

Annual hydrographic variability in Antarctic coastal waters infused with glacial inflow

Maria Osińska¹, Kornelia A. Wójcik-Długoborska², Robert J. Bialik²

¹Institute of Oceanography, University of Gdańsk, Piłsudskiego 46, 81-378 Gdynia, Poland

5 ² Institute of Biochemistry and Biophysics, Polish Academy of Sciences, Pawińskiego 5a, 02-106 Warsaw, Poland

Correspondence to: Robert J. Bialik (rbialik@ibb.waw.pl)

Abstract.

During the thirty-eight months between December 2018 and January 2022, multiparameter hydrographic measurements were taken at thirty-one sites within Admiralty Bay, King George Island, Antarctica. These records consisted of water column
10 measurements (down to 100 m) of temperature, conductivity, turbidity, and pH as well as the dissolved oxygen, dissolved organic matter, chlorophyll A and phycoerythrin contents. The sites were chosen due to their variable distances from glacial fronts and open ocean waters. Fifteen sites were localized within smaller glacial coves, with waters highly impacted by glacial infusions; seven sites were located in the open waters of the main body of Admiralty Bay; and nine were located in intermediate conditions of the Ezcurra Inlet. The final dataset consists of measurements carried out over 142 separate days, with an average
15 3.74 measurements per month. However, data were not collected regularly throughout the year and were collected less frequently during winter, though data were gathered for all but two winter months. On average, each site was investigated 98.2 times. Due to calibration issues, absolute values of optically measured properties occasionally show unrealistic negative values, but the relative distributions of these values remain valid. Variabilities in the measured properties each season and throughout the whole duration of the project reveal regular oscillations as well as possible long-term trends.

20 1 Introduction

When freshwater from glaciers is introduced to the marine environments, it mixes with ambient ocean water masses leading to the formation of new glacially modified water (GMW, (Straneo, 2012). Freshwater export has in this way been shown to influence properties of the coastal ocean, with impacts on the hydrodynamics and thermodynamics (Bendtsen et al., 2015; Chauché et al., 2014). Therefore, there are significant justifications to investigate water quality properties in glacial bays and
25 fjords and to track their variability to potentially predict future changes.

While the majority of studies examining the influence of glacial meltwater on the marine ecosystem have been performed in the Northern Hemisphere, its importance for the functioning of coastal Antarctic waters has long been hypothesized. Nevertheless, widely available data that describe water quality in glacial bays beyond seasonal timescales, at high sampling

resolutions, and that examine multiple variables remains non-existent. In fact, such datasets are scarce for the Arctic and Alaska as well.

To address this deficiency, an intricate investigation campaign was designed with the intention of comprehensively observing the seasonal oscillations and long-term trends in water quality variability of Admiralty Bay (AB), King George Island in Western Antarctica. The goal of this project was to widen the scope of previously gathered observations by expanding the overall duration of monitoring, increasing the frequency and number of measured parameters and to acquire data across all seasons of the year.

2 Research Area


AB is a 177.04 km² cove southeast of King George Island, the largest island of the Southern Shetlands in Western Antarctica. In its interior, AB is subdivided into three distinct areas: Ezcurra, Mackellar and Martel Inlets, which all blend together approximately 11 km from the open ocean waters of the Bransfield Strait, forming the main body of AB (Figure 1). Its coastline has a length of 150 km, of which 102 km consist of rocky coastline and the remaining 38 km consist of ice-water boundaries (Figure 1, yellow lines (Gerrish et al., 2021)). The tidewater glaciers that form these frontiers are the outer regions of two large icefields, the Warsaw and Krakow Icefields. Both icefields are reportedly experiencing unprecedented transformation due to the effects of climate warming (Rückamp et al., 2010; Dziembowski and Bialik, 2022) and are draining into AB through numerous glacial creeks.

To summarize this dataset, it was decided to distinguish different zones within AB to recognize distinctions in water properties that were dependent on proximity to the glacial fronts and open ocean waters. Three zones have been assigned:

- Glacial coves: Distinct smaller bays formed near tidewater glaciers in which marine waters are under the direct influence of glacial meltwater input. Here, three glacial coves were analysed in depth, the cove near Lange Glacier (1.50 km² in area with a 2.81 km long ice-water frontline), Spera Cove (2.45 km² in area with a 4.33 km long ice-water frontline) near Vieville Glacier and Suszczewski Cove near Ecology Glacier (0.69 km² in area with a 0.36 km long ice-water frontline). All three of these basins are undergoing long-term transformation caused by continuously moving and developing glacial fronts. This is visualized in Figure 1, where the light blue line on the glacial cove insets represents the ice-water boundary in 2018 (based on a Sentinel satellite image from Mar 10th, 2018), which is different from the frontline shown on a satellite picture presented in Figure 1 taken in December of 2021 (Sentinel, Dec 29, 2021). The change is especially noticeable in Spera Cove near Vieville glacier, where the ice front has retreated 500 m within three years in some locations.
- Main body of Admiralty Bay: Open bay waters in the main body of the cove, most directly influenced by the open ocean waters of the Bransfield Strait with which it is connected by a 13.45 km wide outlet. Nevertheless, this location is also affected by glacial input, especially in its northern parts.

60 - Ezcurra Inlet (area of 21.32 km²): This is an intermediate area separated from Admiralty Bay waters by a relatively narrow passage (2.40 km wide) and influenced by the surrounding ice coastline (9.58 km long of 32.67 km long coastline).

These areas are shown in Figure 1 and are used as separate, but deeply interrelated, regions for further study. To that end, measurement points were chosen, whose localizations are marked on the map in Figure 1, and their details (location, depth, 65 number of measurements performed at a given point, and, in the case of glacial cove points, distance from the water-ice boundary) are summarized in Table 1.



70 **Figure 1:** Map of Admiralty Bay with measurement points in three distinct zones: main Admiralty Bay (pink), Glacial coves (blue inset boxes) and Ezcurra Inlet (yellow lines based upon (Gerrish et al., 2021). Bright yellow shows current position of the ice-water coastline, in bright blue insets the position of the coastline on Mar 10th, 2018. Source: Sentinel imagery, Dec 29th, 2021.

Table 1: Details of the measurement sites. The depth measurements are based on the YSI EXO sonde depth sensor, depths >100 indicate that in a given site sonde was lowered to cable limits (100 m) and did not reach ocean bottom. Distances from glacial fronts were measured only for sites within smaller Glacial coves adjacent to individual glaciers, since sites in Admiralty Bay and Ezcurra Inlet were influenced by number of glaciers in their vicinity

| site name and zone | latitude | longitude | depth (m) | distance from glacial front (m) (2018-2021) |
|--------------------|----------|-----------|-----------|---------------------------------------------|
| Glacial Coves | | | | |

| | | | | |
|----------------------|----------|----------|------|----------|
| lr1 | -58.4892 | -62.1227 | 19 | 315-322 |
| lr2 | -58.4868 | -62.1195 | >100 | 266-330 |
| lr3 | -58.4845 | -62.1163 | >100 | 275-332 |
| lr4 | -58.4821 | -62.1131 | 23 | 260-343 |
| lr5 | -58.4687 | -62.1120 | 8 | 940-1018 |
| lr6 | -58.4711 | -62.1152 | 66 | 880-951 |
| lr7 | -58.4734 | -62.1184 | >100 | 902-952 |
| lr8 | -58.4758 | -62.1216 | >100 | 868-912 |
| lr9 | -58.4782 | -62.1247 | 3 | 929-932 |
| ve1 | -58.3380 | -62.1361 | 2 | 71-481 |
| ve2 | -58.3429 | -62.1375 | 8 | 359-780 |
| ve3 | -58.3483 | -62.1391 | 29 | 686-1118 |
| ve7 | -58.4613 | -62.1716 | 4 | 455-469 |
| ve8 | -58.4677 | -62.1709 | 2 | 210-232 |
| ve9 | -58.4668 | -62.1734 | 3 | 113-128 |
| <hr/> | | | | |
| Admiralty Bay | | | | |
| lv1 | -58.4624 | -62.1221 | >100 | |
| lv2 | -58.4412 | -62.1251 | >100 | |
| lv3 | -58.3989 | -62.1313 | 71 | |
| lv4 | -58.3777 | -62.1343 | 17 | |
| ve4 | -58.3671 | -62.1445 | 58 | |
| ve5 | -58.4047 | -62.1553 | >100 | |
| ve6 | -58.4424 | -62.1662 | 55 | |
| <hr/> | | | | |
| Ezcurra Inlet | | | | |
| ez1 | -58.6172 | -62.1812 | 56 | |
| ez2 | -58.5994 | -62.1778 | 67 | |
| ez3 | -58.5811 | -62.1750 | 51 | |
| ez4 | -58.5626 | -62.1727 | 61 | |
| ez5 | -58.5441 | -62.1702 | 68 | |
| ez6 | -58.5279 | -62.1655 | 84 | |
| ez7 | -58.5136 | -62.1595 | >100 | |
| ez8 | -58.5012 | -62.1526 | >100 | |
| ez9 | -58.4878 | -62.1462 | >100 | |

75

Measurements in Glacial Coves and Admiralty Bay were taken from December 2018 until January 2022 and in Ezcurra Inlet from October 2019 till January 2022.

Due to proximity to glaciers as well as harsh Antarctic weather sampling in this region was especially strenuous. Each measurement campaign, lasted few hours and has been performed from the decks of small Zodiac boats (Figure 2) that provided little comfort to the crew and getting to assigned sites often involved manoeuvring through moving ice packs and bits of icebergs coming from calving glaciers. Sampling during winter months required working in the dark, in extremely cold temperatures with continuous contact with freezing water.

3 Methodology

3.1 Measured water properties

Measurements were performed with two professional YSI multiparameter EXO sondes (EXO1 and EXO2), designed for simultaneous investigation of multiple water quality properties, used and tested by researchers worldwide (Snazelle, 2015). EXO1 consists of five sensor ports and EXO2 contains seven ports; therefore, the water properties measured varied between the particular campaigns. Of the 3045 measurements collected, 2069 were acquired using the EXO1 sonde, and the remaining 976 were acquired with EXO2 and its larger sensor capacity (details seen in Figure 2).

The list of sensors and properties investigated by them is summarized in Table 2. Some hydrographic properties are derived indirectly from the direct measurements of other water column properties. In these cases, the sondes automatically calculated the additional related values based on universally accepted formulas (Table 2).

Table 2. List of sensors and measured water properties (based upon (YSI Inc, 2017))

| | | Sensor | Measured property | Unit | Accuracy/linearity | Direct measurement (D) or calculated from other measurement (C) |
|------|------|--------------------------|-----------------------|-------|---------------------------------------------------------------------------------------------------|-----------------------------------------------------------------|
| EXO2 | EXO1 | Conductivity/Temperature | Conductivity | µS/cm | 0-100 mS/cm: ±0.5% of reading or 0.001 mS/cm, whichever is greater; 100-200 mS/cm: ±1% of reading | D |
| | | | Specific Conductivity | µS/cm | | C – conductivity adjusted to temp. |
| | | | nLF Conductivity | µS/cm | | C – with temp. compensation |
| | | | Salinity | PSU | | C – based on temp. and conductivity using APHA, 1989 |
| | | | Temperature | °C | ± 0.01° | D |
| | | Depth and Level | Pressure | PSI | ± 0.04 m | D |

| | | | | | |
|----------------------|-------------------------|-----------------------------------|-----------|----------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------|
| not measured by EXO1 | Depth | | m | | C – based on water and atm. pressure |
| | Dissolved Oxygen | Dissolved Oxygen | mg/L | ±1% of reading or ± 0.01 | C –using Stern-Volmer equation from luminescence measurement corrected with temp and atm. pressure |
| | | Dissolved Oxygen Saturation | % | mg/L | |
| | | Dissolved Oxygen Local Saturation | % | | |
| | pH | pH | -, mV | ± 0.01 | C - from electric potential difference |
| | Turbidity | Turbidity | FNU | 0.3 FNU or ±2% of reading, whichever is greater | C – from light scatter |
| | fDOM | Dissolved Organic Matter | QSU, RFU | R ² >0.999 for serial dilution of 300 ppb Quinine Sulphate solution | C – from fluorescence |
| | Total Algae (Chl & BGA) | Chlorophyll A | µg/L, RFU | R ² >0.999 for serial dilution of Rhodamine WT solution from 0-400 µg/L Chl equivalents | C – from fluorescence |
| | | BGA PE (Phycoerythrin) | µg/L, RFU | R ² >0.999 for serial dilution of Rhodamine WT solution from 0-280 µg/L PE equivalents | C – from fluorescence |

95 3.2 Measurement and data handling procedure. Causes of possible data errors and missing values.

Measurements were conducted from the deck of the Zodiac boat (Figure 2). When the boat was in the designated point, the sonde lowered by the cable from the reel to a maximum depth of 100 m. At sites with depth smaller than 100 m (see Table 1 for information on sites' depth) the measurements were performed throughout the whole column of water, until sea bottom was reached. At sites where the depth surpasses 100 m data has been collected only from top 100 m which shows the limitation of this study as data was not obtained from bottom portions of the water column. The sampling rate of the sondes were initially 0.2 Hz up to December 30th, 2019, when the sampling frequency increased to 1 Hz.

The intended descent rate of the instrument was 1 m per second, but since this was manually controlled by the research personnel, the descent rate of the sonde varied significantly. Furthermore, the fact that the measurements were acquired by

different crews may cause some discrepancy in the acquired data. The sensitivity of particular sensors varied, meaning that if
105 the probe was lowered too quickly the measurements taken by some sondes may incorrectly correlate with the depth attributed to those measurements.

Other obstacles were caused by challenging weather and sea conditions. Often, waving and surface currents considerably influenced the position of the boat, making it impossible to remain stationed at the assigned site location for the duration the cast. This can be seen by position data recorded via handheld GPS during sensor deployment and included within the data file.
110 Currents below the surface moved the sonde and cable horizontally from the initial cast position by an unknown extent. On numerous occasions ice prevented scientists from reaching specific sites. This was frequently the case in areas close to glacial fronts, most notably when the water surface froze during winter months and when glacial calving increased in the summer.

All of the sensors were calibrated in accordance with guidelines found in the YSI EXO Manual (YSI Inc, 2017) and replaced
115 after the appropriate time or when malfunctions occurred that could not be otherwise resolved. The depth/pressure level sensor was calibrated at the start of every survey day.

Measured data was firstly recorded in YSI proprietary format in software embedded into all of the sensors. At this stage, data went through real-time data filtering using basic rolling filter as well as adaptive filtering and outlier rejection with default settings of the manufacturer (details see (YSI Inc, 2017)). Gathered data was later downloaded using KorExo software and
120 exported to Matlab where some outlier and extreme values have been extracted due to one of the following reasons:

- Notes from the measurement crew indicated malfunctions or some other issues.
- On sites with depth smaller than 100 m, sonde after reaching the bottom showed unrealistic values from all of the sensors which was caused by the contact with seafloor.
- Other extreme and outlier data was scrutinized individually:
125
 - Continuous abnormal values of a particular sensor during measurement day were deleted indicating sensor malfunction or decalibration.
 - Incidental extreme values recorded within otherwise reasonable datasets were taken out indicating momentary disturbances.

Despite this series of steps whole dataset did not go through any formalized Quality Assessment/Quality Check procedure.
130 Optical sensors for total algae and fDOM showed unrealistic negative values (77.82% of chlorophyll A, 70.87% of phycoerythrin and 60.45% of fDOM readings). This was most probably caused by chosen calibration method using 1-point procedure, based on deionized water as a proxy for 0 fluorescence standard. This methodology has been outlined by the sensor manufacturer (YSI Inc, 2017), but in this environment it has proven insufficient, and suggests necessity of using more robust method of calibration for future measurements. Nevertheless these negative values have been left intact in the data file since
135 they represent the correct variability of the properties; however, their absolute values should be considered carefully, and more attention should be given to the relative units (RFU) for chlorophyll A, phycoerythrin and fDOM.

Turbidity sensor also showed negative values (19.56 % of the readings) but it has been calibrated using 2-point procedure with appropriate standard and its FNU values have been confirmed in Admiralty Bay waters through the laboratory procedure explained in detail by (Wojcik-Długoborska et al., 2022).




Figure 2. Measurements visualisation used sondes and measurement properties. In the background scientists in Zodiac boat during measurements of water properties visibly infused with turbid GMW (source of sonde close-ups: <https://observer.com/products/ysi-exo-series-multiparameter-sonde/>)

4 Results

The results of the measurement campaign discussed above consist of a large and complex dataset describing the variability of the physical, chemical and biological properties in glacially influenced bays. Figure 3 presents a summary of the total number of investigations performed. This shows that even at the sites sampled the least, it was possible to gather data during all seasons. However, most studies were performed during summer across all zones, while the fewest measurements were collected in winter. Interestingly, despite the unpredictable conditions in the Glacial coves, the number of surveys at each site fluctuates around one hundred per location (average 98.2 measurements per site), which is promising for future statistical analysis.




Figure 3. Number of measurements taken at the designated sites in total (a) and by season (calendar) (b) in the period from December 2018 until January 2022

Considering the complete duration of the projects (see Figure 4), it is noticeable how the number of measurement days fluctuated, with increases during the warmer seasons where there was a maximum of seven measurement days per month. In Figure 4(b), we observe that the same tendencies apply to all the zones, and none of them have been more frequently investigated to any degree of significance. The average number of measurements per month was 3.74 in the Glacial coves and 2.91 in Admiralty Bay, with the same number of successful measurement days (111) throughout the whole duration of the project, and 2.42 for Ezcurra Inlet over 92 measurement days.





Figure 4. Number of measurement days per month, (a) in total and (b) per each zone (counted days in which measurements in at least half of zone sites have been performed)

The division of sites into three zones shows how proximity to glacial fronts and open ocean waters alters particular water quality properties. This effect is also notably correlated with seasonal shifts (Figure 5 and 6). In Figure 5 vertical distribution

165 have been presented of all the gathered data. It is apparent that temperature, pH, ODO and phytoplankton pigment values are especially prone to change due to seasonal shifts, whereas salinity and turbidity values stay similar throughout the year. However, Figure 6 in detail illustrates how different properties vary in surface layers in contrast with the whole column of water (limited to 100 m of depth), most notably in salinity and turbidity values, but true for all measured properties except for pH. This shows the impact of both atmospheric forcing and glacial outflow, which, based on buoyant plume model theory

170 (Kimura et al., 2014; Mankoff et al., 2016; Jenkins, 2011) and observations (Chauché et al., 2014; Osińska et al., 2021), is mainly contained in the top layer of the ocean. Therefore, the results provide information on seasonal changes in water properties and glacial-ocean interactions and can be used for validation of previously formulated methods of GMW tracking.



175 **Figure 5. Vertical variability of measured properties divided by zone and season**





Figure 6: Mean values of measured properties dependent on season, divided into the whole column of water and the top 5 m of surface water.

180 The 38-month-long duration of the project allowed for the tracking of seasonal variability across all measured hydrographic properties and showed consistency in all cases (Figure 7). Moreover, this duration permits cautious predictions regarding long-term shifts in water column properties and reveals the impact of climate change or other influential conditions in this region. Using more sophisticated techniques, it is possible to more precisely determine the nature of this variability. The quantities of chlorophyll A, phycoerythrin and fDOM are not presented in Figure 6 since their measurement was significantly less frequent.



185 **Figure 7. Boxplot of monthly properties' mean values (excluding properties measured solely by EXO2 sonde due to their significantly shorter timeseries).**

5 Data availability

Described dataset is freely accessible at the Pangea repository via following doi:
190 <https://doi.pangaea.de/10.1594/PANGAEA.947909> (Osińska et al., 2022), under a non-restrictive license CC BY 4.0..

6 Conclusions

The assembled dataset shared here presents an opportunity for a better understanding of Admiralty Bay water characteristics over the 38-month survey period and can be used in further studies exploring the nature and changes in glacially influenced regions in general. Sheer magnitude of this investigation with 3045 separate measurements acquired on 142 different days
195 validates its importance and inspires optimism regarding future work and application of this data.

The scope of measured parameters (thermodynamic, physical, chemical and biological) paints a wide and precise picture of AB hydrographic variability during all months of the year and may allow for a multidisciplinary analysis of the complex processes that take place in this location. The varied settings of study sites allow for the tracking and identification of GMW and other water masses (Straneo et al., 2011; Chauché et al., 2014). Additionally, this sizable dataset can be used as a tool for
200 better understanding the general hydrodynamics and thermodynamics of glacial bays and fjords and may be employed for the validation of coupled glacier-ocean modelling (Cowton et al., 2015; De Andrés et al., 2021; Bertino and Holland, 2017).

Author Contribution

MO – conceptualization, data curation, formal analysis, investigation, methodology, validation, visualization and writing (original draft preparation), KAW – investigation, methodology and writing (review & editing); RJB – funding acquisition,
205 investigation, project administration, resources, supervision and writing (review & editing)

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgments

This work was supported by the National Science Centre, Poland, Grant No. UMO-2017/25/B/ST10/02092 ‘Quantitative
210 assessment of sediment transport from glaciers of South Shetland Islands on the basis of selected remote sensing methods. Calculations were made possible by software provided by CI TASK (Center of the Tri-City Academic Computer Network) in Gdańsk. Special credits are owed to all invincible members of so called MorMon team, part of Polish Antarctic Station’s crew,

that throughout whole period of the project, in often trying and almost always uncomfortable conditions carried out presented measurements.

215 **References**

- De Andrés, E., Otero, J., Navarro, F. J., and Walczowski, W.: Glacier-plume or glacier-fjord circulation models? A 2-D comparison for Hansbreen-Hansbukta system, Svalbard, *Journal of Glaciology*, 67, 797–810, <https://doi.org/10.1017/jog.2021.27>, 2021.
- APHA: Standard Methods for Examination of Water and Wastewater, 17th ed., Washington D.C., 1989.
- 220 Bendtsen, J., Mortensen, J., Lennert, K., and Rysgaard, S.: Heat sources for glacial ice melt in a west Greenland tidewater outlet glacier fjord: The role of subglacial freshwater discharge, *Geophys Res Lett*, 42, 4089–4095, <https://doi.org/10.1002/2015GL063846>, 2015.
- Bertino, L. and Holland, M. M.: Coupled ice-ocean modeling and predictions, *J Mar Res*, 75, 839–875, <https://doi.org/10.1357/002224017823524017>, 2017.
- 225 Chauché, N., Hubbard, A., Gascard, J. C., Box, J. E., Bates, R., Koppes, M., Sole, A., Christoffersen, P., and Patton, H.: Ice-ocean interaction and calving front morphology at two west Greenland tidewater outlet glaciers, *Cryosphere*, 8, 1457–1468, <https://doi.org/10.5194/tc-8-1457-2014>, 2014.
- Cowton, T., Slater, D., Sole, A., Goldberg, D., and Nienow, P.: Modeling the impact of glacial runoff on fjord circulation and submarine melt rate using a new subgrid-scale parameterization for glacial plumes, *J Geophys Res Oceans*, 120, 796–812, <https://doi.org/10.1002/2014JC010324>, 2015.
- 230 Dziembowski, M. and Bialik, R. J.: The Remotely and Directly Obtained Results of Glaciological Studies on King George Island: A Review, *Remote Sens (Basel)*, 14, 2736, <https://doi.org/10.3390/RS14122736>, 2022.
- Gerrish, L., Fretwell, P., and Cooper, P.: High resolution vector polylines of the Antarctic coastline (7.4) [Data set], <https://doi.org/https://doi.org/10.5285/e46be5bc-ef8e-4fd5-967b-92863fbe2835>, 2021.
- 235 Jenkins, A.: Convection-Driven Melting near the Grounding Lines of Ice Shelves and Tidewater Glaciers, *J Phys Oceanogr*, 41, 2279–2294, <https://doi.org/10.1175/JPO-D-11-03.1>, 2011.
- Kimura, S., Holland, P. R., Jenkins, A., and Piggott, M.: The Effect of Meltwater Plumes on the Melting of a Vertical Glacier Face, *J Phys Oceanogr*, 44, 3099–3117, <https://doi.org/10.1175/JPO-D-13-0219.1>, 2014.
- Mankoff, K. D., Straneo, F., Cenedese, C., Das, S. B., Richards, C. G., and Singh, H.: Structure and dynamics of a subglacial discharge plume in a Greenlandic fjord, *J Geophys Res Oceans*, <https://doi.org/10.1002/2016JC011764>, 2016.
- 240 Osińska, M., Bialik, R. J., and Wójcik-Długoborska, K. A.: Interrelation of quality parameters of surface waters in five tidewater glacier coves of King George Island, Antarctica, *Science of the Total Environment*, 771, 144780, <https://doi.org/10.1016/j.scitotenv.2020.144780>, 2021.

- Osińska, M., Wójcik-Długoborska, K. A., and Bialik, R. J.: Water conductivity, salinity, temperature, turbidity, pH, fluorescent dissolved organic matter (fDOM), optical dissolved oxygen (ODO), chlorophyll a and phycoerythrin measurements in Admiralty Bay, King George Island, from Dec 2018 to Jan 2022, PANGAEA, <https://doi.org/10.1594/PANGAEA.947909>, 2022.
- Rückamp, M., Blindow, N., Suckro, S., Braun, M., and Humbert, A.: Dynamics of the ice cap on King George Island, antarctica: Field measurements and numerical simulations, *Ann Glaciol*, 51, 80–90, <https://doi.org/10.3189/172756410791392817>, 2010.
- Snazelle, T. T.: Evaluation of Xylem EXO water-quality sondes and sensors, U.S. Geological Survey Open-File Report 2015-1063, <https://doi.org/10.3133/OFR20151063>, 2015.
- Straneo, F.: Impact of the large scale ocean circulation on Greenland’s outlet glaciers, *Quaternary International*, 279–280, 472, <https://doi.org/10.1016/j.quaint.2012.08.1584>, 2012.
- Straneo, F., Curry, R. G., Sutherland, D. A., Hamilton, G. S., Cenedese, C., Våge, K., and Stearns, L. A.: Impact of fjord dynamics and glacial runoff on the circulation near Helheim Glacier, *Nat Geosci*, 4, 322–327, <https://doi.org/10.1038/ngeo1109>, 2011.
- Wójcik-Długoborska, K. A., Osińska, M., and Bialik, R. J.: The impact of glacial suspension color on the relationship between its properties and marine water spectral reflectance, *IEEE J Sel Top Appl Earth Obs Remote Sens*, <https://doi.org/10.1109/JSTARS.2022.3166398>, 2022.
- YSI Inc: Exo User Manual, Yellow Springs, 1–154 pp., 2017.