

GRiMeDB: The global river database of methane concentrations and fluxes

Emily H. Stanley¹, Luke C. Loken^{1,2}, Nora J. Casson³, Samantha K. Oliver², Ryan A. Sponseller⁴ Marcus B. Wallin⁵ Liwei Zhang⁶ Gerard Rocher-Ros^{7,8}

5

¹Center for Limnology, University of Wisconsin-Madison, Madison WI, USA 53706

²U.S. Geological Survey, Upper Midwest Science Center, Madison, WI USA 53726

³Department of Geography, University of Winnipeg, Winnipeg, MB Canada R3B 2E9

⁴Department of Ecology and Environmental Science, Umeå University, Umeå, Sweden 90736

10 ⁵Department of Aquatic Sciences and Assessment, Swedish University of Agricultural Sciences, Uppsala, 75007, Sweden

⁶Sino-French Institute for Earth System Science, College of Urban and Environmental Sciences, Peking University, 100871, Beijing, China

⁷Department of Forest Ecology and Management, Swedish University of Agricultural Sciences, 90183 Umeå, Sweden

⁸Integrative Freshwater Ecology Group, Centre for Advanced Studies of Blanes (CEAB-CSIC), 17300 Girona, Spain

15

Correspondence to: Emily H. Stanley (ehstanley@wisc.edu)

Abstract. Despite their small spatial extent, fluvial ecosystems play a significant role in processing and transporting carbon in aquatic networks, which results in substantial emission of methane (CH₄) to the atmosphere. For this reason, considerable effort has been put into identifying patterns and drivers of CH₄ concentrations in streams and rivers and estimating fluxes to the atmosphere across broad spatial scales. Yet progress toward these ends has been slow because of pronounced spatial and temporal variability of lotic CH₄ concentrations and fluxes and by limited data availability across diverse habitats and physicochemical conditions. To address these challenges, we present a comprehensive database of CH₄ concentrations and fluxes for fluvial ecosystems along with broadly relevant and concurrent physical and chemical data. The Global River Methane database (GriMeDB; <https://doi.org/10.6073/pasta/f48cdb77282598052349e969920356ef>, Stanley et al., 2023) includes 24,024 records of CH₄ concentration and 8,205 flux measurements from 5,029 unique sites derived from publications, reports, data repositories, unpublished data sets, and other outlets that became available between 1973 and 2021. Flux observations are reported as diffusive, ebullitive, and total CH₄ fluxes, and GriMeDB also includes 17,655 and 8,409 concurrent measurements of concentrations and 4,444 and 1,521 of fluxes for CO₂ and nitrous oxide (N₂O) respectively. Most observations are date-specific (i.e., not site averages) and many are supported by data for 1 or more of 12 physicochemical variables and 6 site variables. Site variables include codes to characterize marginal channel types (e.g., springs, ditches) and/or presence of human disturbance (e.g., point source inputs, upstream dams). Overall, observations in GRiMeDB encompass the broad range of the climatic, biological, and physical conditions that occur among world river basins, although some geographic gaps remain (arid regions, tropical regions, high latitudes and altitude systems). The global median CH₄ concentration (0.20 μmol L⁻¹) and diffusive flux (0.44 mmol m⁻² d⁻¹) in GRiMeDB are lower than estimates from prior site-averaged compilations, although ranges (0-456 μmol L⁻¹ and -136-4057 mmol m⁻² d⁻¹) and standard deviations (10.69 and 86.4) are greater for this larger and more temporally resolved database. Available flux data are dominated by diffusive measurements despite the recognized importance of ebullitive and plant-mediated CH₄ fluxes. Nonetheless, GriMeDB provides a comprehensive and cohesive resource for examining relationships between CH₄ and environmental drivers, estimating the contribution of fluvial ecosystems to CH₄ emissions, and contextualizing site-based investigations.

1 Introduction

Despite their small areal extent, running-water (fluvial) ecosystems play a significant role in processing and transporting carbon (C) in and through aquatic networks, including the production, consumption, transport, and evasion of carbon dioxide (CO₂) and methane (CH₄). The profound planetary warming effects of CH₄ in the atmosphere, its erratic but accelerating rate of increase over recent years (NOAA, 2022), the significant contributions of natural sources to the growing atmospheric pool (Turner et al., 2019), and improvements in gas measurement technologies have all contributed to a rapid increase in studies of CH₄ dynamics in aquatic environments in general, and fluvial ecosystems in particular. These studies reveal widespread

50 supersaturation of CH₄ in running waters that underlies their larger than expected contribution to the atmospheric pool (Stanley et al., 2016).

Efforts to quantify fluvial CH₄ dynamics at regional, continental, and global scales have been fraught with uncertainty, reflecting the inherent variability of this gas in surface waters combined with a notable limitation in data availability. Sources and sinks of CH₄ are often unevenly distributed over space and time within drainage systems and, as a result, concentrations
55 can vary over 1-3 orders of magnitude over short time periods (sub-daily to sub-weekly; e.g., Natchimuthu et al., 2017; Smith and Böhlke, 2019) or relatively small spatial extents (<10 - <100 m for small streams and large rivers; e.g., Anthony et al., 2012; Crawford et al., 2017; Bretz et al., 2021; Robison et al., 2021). Similarly, several drivers or predictors of CH₄ have been identified in the literature, and these properties also have variable spatial and temporal distributions. Thus, efforts to estimate the total emissions from world rivers have relied on relatively small data sets composed of site-specific values that have been
60 averaged over time, and then used upscaling strategies based on Monte Carlo techniques or extrapolations using predictor variables that have little or no significant statistical relationships with large-scale patterns of gas concentrations or fluxes (Hutchins et al. 2020). Consequently, current global scale estimates of riverine emissions are poorly constrained and highly uncertain (Saunois et al., 2020; Rosentreter et al., 2021).

The combination of rapidly increasing atmospheric concentrations of CH₄, the significant role of fluvial systems in emitting
65 this gas, and, critically, current difficulties in explaining or predicting concentrations and fluxes with reasonable certainty inspired the central goal of this paper: to assemble a comprehensive database of CH₄ concentrations and fluxes for fluvial ecosystems that includes broadly relevant concurrent physical and chemical data. This effort expands upon a prior compilation of CH₄ and CO₂ data (MethDB; Stanley et al., 2015) that was constructed to emphasize among-site differences and included 1,496 concentration records and 532 flux records from 1,080 sites. In this more comprehensive Global River Methane database
70 (GRiMeDB), most data are date-specific (i.e., not averaged over time), the breadth of site types is expanded to include marginal fluvial habitats as well as disturbed and artificial waterways, and CH₄ data are supported by a broad suite of site-specific physical and chemical attributes along with concurrent measurements of CO₂ and N₂O where available. Given the more finely resolved scale of the data and the growth of the field in the past decade, GRiMeDB represents a significant expansion beyond MethDB. Building GRiMeDB with greater detail and breadth of data was done with the intent of increasing opportunities to
75 identify and predict spatial and temporal variation in CH₄, to test hypotheses related to greenhouse gas dynamics, and to reduce uncertainty in future upscaled estimates of gas emissions. In this paper, we (1) provide a detailed description of the components of the database and its construction; (2) summarize some basic patterns of gas concentrations and fluxes from GRiMeDB; and (3) highlight critical data gaps and possible future research opportunities for improving current understanding of CH₄ dynamics in streams and rivers.

80 2 Database components and assembly

GRiMeDB is composed of four tables that contain information related to (1) data sources, (2) sites, (3) gas concentrations and supporting physicochemical data, and (4) gas fluxes. All tables are linked by unique data source identifiers, and all concentration and flux observations are also linked to unique site numbers (Fig. 1). Data included in GRiMeDB were gathered from scientific journals, government reports, public data repositories, theses, dissertations, and unpublished data sets provided by individual investigators. Sources were discovered via searches of bibliographic databases and data repositories (Web of Science, Google Scholar, ProQuest Dissertations & Theses Global, China National Knowledge Infrastructure, Environmental Data Initiative, USGS ScienceBase, Natural Environment Research Council (NERC) Environmental Information Data Centre, Arctic Data Center, PANGAEA, Zenodo) using the keywords: methane and stream* or river* or ditch* or canal*, and searches were repeated numerous times prior to December 2021 for completeness. We also used informal ‘word of mouth’ approaches to discover additional, often unpublished data sets.





Figure 1: General structure of the GRiMe database and connections between its four tables. Information flow began with entering information about each data source into the Sources Table and assigning a unique Source_ID. Site information for each site within a data source was then entered into the Sites table. The site was given a unique Site_ID and linked to its data source by the Source_ID. Source_IDs and Site_IDs were carried over to all concentration and flux observations in their respective tables. Methane (CH₄) observations include site-date combinations with only concentration data (orange), only flux data (green), or both concentration and flux data (brown). Concentrations and available supporting data (described in Sect. 2.3) were entered into the Concentration Table, and each observation was given a unique observation (obs) name. For site-date combinations that had both concentration and flux observations, the Source_ID, Site_ID, observation name, and date information were copied to the Flux Table for data entry. Site-date combinations with flux data only were entered into the Flux Table and given a unique observation name. If a flux observation had associated supporting data, the Source_ID, Site_ID, observation name, and date information were copied to the Concentration Table for supporting data entry. However, if there were no supporting data, matching rows were not added to the Concentration Table.

All potential data sources were first screened to determine their appropriateness for inclusion in GRiMeDB. Several criteria were established *a priori* to ensure the usability of the data and that it was derived from inland running water systems. Coastal sites with

>1 ppt salinity were considered estuarine and thus were excluded. Similarly, sites that were situated in reservoirs or immediately upstream of small dams, dam spillways, beaver ponds, or lake outlets or were subject to experimental manipulation were omitted. We did not enter fluxes derived from chambers attached to collars or inserted into sediments because we could not be certain that such measurements were capturing air-water fluxes. Sources that reported minimum and maximum gas concentrations or fluxes only were not included. Finally, rates expressed on an annual basis were also excluded to avoid introducing uncertainty associated with different upscaling assumptions and methods.

2.1 Sources Table

The Sources Table contains the list of all sources used to build GRiMeDB, a unique identification number (*Source_ID*) for each CH₄ data source, and basic bibliographic information for the data source (*Title*, *Author*, *Source*, publication year [*Pub_year*], and digital object identifiers [*Paper_DOI* or *Data_DOI_primary*] or other persistent identifier; all column titles for this table are defined in Table A1). In several cases, data sources were supplemented with additional supporting information (e.g., associated physicochemical data) from separate sources (described further in Sect. 2.3) or additional or corrected information from authors (Fig. 2). In the latter case, we contacted authors if questions arose regarding their data (e.g., clarification regarding units) and/or to request supporting information or site- or date-specific concentrations or fluxes if published values were aggregated. Inclusion of additional unpublished data from authors is noted in the Sources Table along with a description of the addition or correction. If supporting data from separate published sources were used, the DOI or other persistent identifier for the secondary source was listed in a separate column [*Data_DOI_supporting*].



125 **Figure 2: Workflow for entering data into the Sources Table of GRiMeDB.**

2.2 Sites Table

The Sites Table reports basic information on attributes for all sites where CH₄ was sampled. Each site has a unique identification code (*Site_ID*) and name (usually taken directly from the data source) and is linked to the Sources Table via the *Source_ID* (see Table A2 for detailed descriptions of all columns in the Sites Table). What composes a ‘site’ (i.e., the spatial extent of data collection) varied among data sources and includes (1) discrete sampling points, (2) geomorphically distinct study reaches, and/or (3) larger channel sections, drainage networks, or other geographic units. The second case typically corresponded to reaches such as riffles or pools in small streams. In the third case, multiple points were often sampled within the ‘site’ and data were then presented as averages. The distance between sampling points that had been averaged varied widely, but were typically >1 km, and in some cases exceeded 100 km. Because land use, channel order, slope, etc., can vary substantially across such distances, we included fields to indicate if a site was an aggregation of widespread points (‘aggregated’), and if so, the number of locations in the aggregation (if available). We also limited the resolution of latitude and longitude for these sites to < 3 decimal places. At the opposite extreme, gas sampling at points very close to one another (a ‘high density site’ *sensu* Fig. 3) has the potential to create ambiguities for site delineation and data analysis. To avoid these pitfalls, we combined points with slightly different latitude-longitude values to represent a single site for three specific cases. First, multiple samples collected at different points and/or depths within a channel cross-section were averaged to form a single site. Second, some drainages or regions were surveyed repeatedly (particularly the Congo River basin and streams in Pennsylvania, USA) and it was not always clear if closely-situated (ca. 10-50 m) points from different surveys were intended to be a repeated sampling of the same location or sampling of discrete sites. Some judgment was involved in choosing between these two possibilities, and in a subset of cases, points in close proximity to one another that were sampled on separate dates were treated as a single site. What constituted ‘close proximity’ varied between small streams and large rivers but was always <100 m, and typically <50 m. Finally, three data sources had extremely high sampling densities within discrete reaches (50 - >20,000 samples per reach; Crawford et al., 2016; Call et al., 2018; Loken et al., 2018). Because closely adjacent gas samples can be spatially autocorrelated (Crawford et al., 2017) and including all individual values from these studies would have resulted in their over-representation in the database, individual point measurements were treated as within-reach replicates.




Figure 3: Workflow for entering and checking data for the GRiMeDB Sites Table. ‘Lat-Lon’ is an abbreviation for latitude and longitude.

For a site used in multiple studies, the *Site_ID* was assigned to the earliest paper and a comment was added to the site entry noting its use in other data sources (Fig. 3). Latitude and longitude coordinates were available for most sites; however, in several cases, location information was acquired from authors or estimated from study site figures using Google Earth (© Google Earth 2020). All sites were plotted on Google Earth and inspected (Fig. 3) to identify and correct data errors. If a site’s coordinates were immediately adjacent to, but not on a channel, the coordinates were adjusted to fall on the channel and this

modification was noted in the *Comments* field. If available, additional variables drawn from the data sources were entered to characterize the site, including stream name, basin or region name, elevation, channel slope, Strahler order, basin area, and codes denoting distinct channel or site types (described below). To supplement the available elevation data, we also estimated elevation for all sites except aggregated sites or sites with poorly-resolved coordinates (<3 decimal places for both latitude and longitude) after snapping coordinates to the nearest stream. To determine the adjusted within-channel coordinates, we first downloaded a digital elevation model (DEM) for each site using the function *get_elev_raster()* from the package “elevatr” (version 0.4.2; Hollister et al., 2021) for R statistical software (version 4.2, R Core Team 2021) at a resolution of 6-9 m depending on the location in the globe. Second, the DEM was processed for hydrological correctness using the package “whitebox” (version 1.2.0, Wu, 2020) by filling single cell pits (*fill_single_cell_pits()* function) and breaching depressions (*breach_depressions()* function) to obtain a flow direction model (*d8_pointer()* function). Finally, we calculated a flow accumulation model (*d8_flow_accumulation()* function). If the coordinates reported in the data source had a flow accumulation <10 cells (indicating that they were not located in a preferential flow path), new coordinates were assigned to the cell with the highest flow accumulation within a 50 m radius. If the initial site had a high flow accumulation value (>10 cells), we assumed the site was in a stream channel. Typically, the snapping procedure resulted in very minor changes to a site’s location (median <3 m).

Many studies of CH₄ dynamics have been undertaken to determine if and how specific phenomena such as presence of upstream reservoirs, point source discharges, thermokarst features, or oil and gas extraction potentially affect fluvial CH₄ (and other constituents), usually with an expectation of a net enhancement of concentrations and fluxes. Similarly, other studies have examined sites that may be expected to be enriched in CH₄, but whose fluvial identity might be considered marginal or ambiguous (e.g., springs, floodplain backwaters, ditches, canals). Inclusion of such ‘methane hunting’ studies has the potential to bias the dataset toward higher values (Stanley et al., 2016). Nonetheless we included these studies in GRiMeDB because they provide an opportunity to investigate the consequences of human activity and gain a more comprehensive understanding of fluvial CH₄ dynamics. However, to accommodate future analyses in which use of such data might be unsuitable, or alternatively, when these sites might be the sole focus of a study, we generated a set of channel codes to identify targeted site types (Table 1). Information about four of the codes was not consistently available among data sources and thus their assignment often involved judgment calls. The first case involved determining if the presence of an upstream dam (code DD) was relevant for sites of varying distances downstream. We used a distance of 7 km as a cut-off for this category, although the zone of influence may be far shorter or extend far beyond this distance depending on dam size and operation (Kemenes et al., 2007), respectively. To provide some context for this code, a site’s distance from a dam was acquired from the data source or estimated in Google Earth using the Path tool and reported in the *Comments* field whenever possible. The second case involved straight, symmetrical channels that are common in many agricultural and urban areas. Frequently, it was not known if this unnatural geometry was due to channelization (straightening) of a stream (code CH) or creation of a new channel (ditches and canals; codes DIT and CAN). In the absence of specific information, straight channels were classified as CH. Third, channels draining or passing through wetlands (WS) were

often difficult to identify, particularly given seasonal variation in wetland appearance. Finally, floodplain channels presented a distinct challenge because of the complex nature of these environments and their potential to be classified as either riverine or wetland systems. We used the FP code to indicate habitats that were described as, or appeared to be lentic (i.e., backwaters or connected floodplain lakes) but were persistently connected to the main river channel and thus were part of the fluvial system. Given these ambiguities, we recommend that these four codes be viewed and used with care.

2.3 Concentration Table and Flux Table

The Concentration Table and the Flux Table contain the primary gas data central to GRiMeDB, and the Concentration Table also hosts physical and chemical variables associated with concentration and/or flux observations (see Tables A3 and A4 for the full list of columns and their descriptions). The vast majority of concentration and flux data were extracted from tables within data sources, data repositories, or provided by authors. However, in some cases, values were acquired from figures using graphical digitizing software (WebPlotDigitizer (<https://automeris.io/WebPlotDigitizer/>), GetData (<http://getdata-graph-digitizer.com/>), or DigitizeIt (<https://www.digitizeit.xyz/>)). Plots with log scales or that were difficult to interpret were not digitized. The accuracy and consistency of this method were evaluated by comparing data generated by different individuals digitizing a set of common figures and by comparing digitized results to known results. Agreement between both comparisons was strong (average slope = 0.994, average $R^2 = 0.9996$ for 5 comparisons between individuals digitizing the same dataset, and average slope = 0.998, average $R^2 = 0.997$ for digitized versus actual data for 7 datasets; see Table S1 for further details), demonstrating the reliability of this method of data gathering.

Whenever possible, concentrations and fluxes were entered as values for individual sites on individual days (i.e., not averaged across sites or days) (Fig. 4). Because 1 day represented the lowest level of temporal resolution in GRiMeDB, repeated measurements made on a sub-daily scale were averaged and expressed as a daily value and were not considered to be aggregated over time. If multiple replicates were collected at different times on the same day (e.g., a study of diurnal gas dynamics), this was noted in the *Comments* fields and measurements prior to and after 12:00 a.m. (local time) were entered as separate, consecutive days. Observations resolved to the daily scale can be identified using either a “No” in the *Aggregated_Time* field or by having the same reported starting (*Date_start*) and ending (*Date_end*) dates. If the specific start and end dates were not specified in the data source, we entered the day as the 15th of the month and noted this approximation in the *Comments* field. If available, we also reported minimum and maximum values and standard deviations (SD) for entries that were aggregated over space and/or time. SDs, but not minima and maxima were reported for replicates from non-aggregated sampling when available, except for reach-averaged entries with multiple within-reach measurements and diel studies with multiple within-day values. In these cases, minima and maxima were also included.

Table 1. Codes denoting distinct site or channel attributes or presence of conditions that potentially affect methane (CH₄) concentrations or fluxes. Assignment of codes to a site is based on information provided in the data source and/or physical appearance of a site and a site may have more than one code. Codes are reported in the *Channel_type* field of the Sites Table.

Code	Definition
CAN	Canal or other artificial channel with hardened channel boundaries
CH	Channelized; a channel that has long straight-line sections of uniform width and changes in channel direction are typically distinct angular features rather than curves
DC	Channel in a river delta
DD	Downstream (within 7 km) of a dam. Samples from spillways were excluded.
DIT	Ditch, typically for agricultural drainage, without channel hardening
FP	Site in a floodplain water body connected to the main channel that appears lentic or is described as a floodplain lake or backwater. This category does not include side-channels within floodplains or tributary channels transecting a floodplain
GT	Site below the toe or terminus of a glacier
IMP	Presence of multiple and typically small impoundments in a site's vicinity (e.g., various European rivers, Mississippi River)
PI	Permafrost influenced; this refers specifically to sites at or immediately below thermokarst outflows and not to sites in areas underlain with permafrost
PS	Immediately (<1 km) downstream of a point source discharge
SP	Spring channel; this does not include sites characterized as seeps (features with low flow volume adjacent to channels)
TH	Site receiving inputs of thermogenic CH ₄ , either naturally or as a result of mining, fracking, oil extraction, and other related activities.
WS	Wetland stream; site is in a wetland or immediately downstream from the outlet of a wetland
NORM	Non-targeted site

225

Dealing with concentration data reported as a negative value, zero, or below a detection limit (BDL) is problematic because of inconsistencies in detection limits and reporting practices, and any decision about handling these records introduces some bias (Stow et al., 2018). For example, using a non-numerical format such as BDL or <0.01 is likely to lead to the elimination of these entries during data analysis and thus would introduce a bias against low-value observations. Alternatively, converting any such value to zero would introduce a bias in the opposite direction. As a compromise solution, concentrations recorded as zero in the


230

original data source were entered as zero in GRiMeDB and other below-detection values were entered as -999999. In this latter case, the original data entry format was noted in the *Comments* column. For fluxes, negative and zero values were entered without modification or comment.

235 The Flux Table reports diffusive, ebullitive, and total CH₄ fluxes along with CO₂ and N₂O diffusive fluxes. Given the diverse strategies for measuring each of the three CH₄ flux pathways and potential biases associated with different approaches (Lorke et al., 2015; Chen et al., 2021), values are accompanied by brief categorical descriptions of methods used for each flux type as well as for CO₂ fluxes and the gas exchange coefficient *k*. For a small number of entries, CH₄ fluxes were not directly reported in the data source but information was available (dissolved gas concentration, temperature, and a corresponding *k* value) that allowed us
240 to calculate these fluxes. We also entered BDL values for flux for one data source in which fluxes had been calculated from concentration, but fluxes associated with BDL concentrations had been omitted from the results. Finally, a small number of observations listed diffusive and ebullitive but not total fluxes, so diffusion and ebullition were summed and entered as total flux. In all cases, the added calculations are noted in the *Comments* field.

245 The GRiMeDB Concentration Table includes physicochemical measurements in support of concentration and flux observations (Figs. 1, 4, Table A3). Availability of this supplemental information varied widely among data sources, and was limited to data collected concurrently with gas samples. For data sources with gas fluxes and physicochemical data but not gas concentrations, we created rows in the Concentration Table to capture the supporting data. These records are identified by a “Yes” in the *FluxYesNo* column, *SampleCount* = 0, and NA in the *CH4mean* column. Finally, water temperature was estimated for entries if it
250 was needed to convert gas units and entered in the *WaterTemp_degC_estimated* column. Estimates were typically based on values from adjacent sites or the same site at a similar time (e.g., averages of temperature from the prior and subsequent dates, or from the same month in an adjacent year). Error introduced from these estimates should be small; e.g., ca. <10% of the actual value if the estimated temperature is off by 3°C.

255 Following completion of all data entry, gas and physicochemical variables were converted to ‘new’ standard units (Tables A3, A4). The identities of the new and original units are included in both the Concentration Table and Flux Table for clarity. Elevation was used to estimate atmospheric pressure if needed for unit conversions. We used Henry’s Law, water temperature, and atmospheric pressure to convert dissolved gas values reported in ppm, ppb, μatm, and % saturation (~13% of observations). For observations that reported gas values as percent saturation (<1% of all observations), we also used the global average CH₄, CO₂,
260 and the N₂O atmospheric concentrations from the NOAA Global Monitoring Laboratory (<https://gml.noaa.gov/ccgg/>) for the year 2013, which corresponds to median observation year in the database.




265 **Figure 4: Workflow for entering and checking data for the GRiMeDB Concentration Table and Flux Table.**

2.4 Assessment of representativeness

We assessed the representativeness of sites in GRiMeDB relative to the global distribution of biological, physical, and climatic properties following van den Hoogen et al. (2021). Briefly, we first assigned each site to a corresponding river reach in

270 HydroSHEDS (Linke et al., 2019), which is a global hydrological network database that contains spatial data for a wide array
of hydrological, physiographical, climatic, land cover, geological, edaphic and anthropogenic variables for each river reach.
HydroSHEDS thus provides a multidimensional characterization of global rivers that is well suited for assessing how
representative GRiMeDB sites are in terms of key biophysical and anthropogenic features. After excluding non-numerical
variables (e.g., biome) and variables with monthly values (e.g., monthly precipitation), we performed a principal component
275 analysis (PCA) on all HydroSHEDS subcatchments using all possible combinations of the 54 remaining HydroSHEDS
variables. From this, we selected all principal components (PCs) needed to explain 90% of the variance in the PCA, which
corresponded to 28 PCs and 378 possible bivariate combinations of these PCs. For each unique PC pair, we computed the
convex hull of all sampled sites to determine the distribution of these sites relative to all global river subcatchments for the
specified PCs (Fig. 5). Each HydroSHEDS subcatchment was then assigned a value of 1 or 0 if it fell within or outside the
280 convex hull, respectively. This process was repeated for each of the 378 possible PC combinations. To collapse this
information, we calculated the fraction of cases that a given subcatchment fell within the convex hull for all PC combinations
to obtain a dimensionless summary value ranging from 0 to 1. A subcatchment with a value of 1 for this index of
“representativeness” means that it fell within the convex hull for 100% of the PC combinations, indicating that its overall
characteristics are well captured in the database. It is important to note that this analysis only captures average catchment
285 properties of relatively large river reaches (average subcatchment area: 130 km²). Given the strong local controls on CH₄
concentrations and fluxes, interpretations from this analysis should be made with some caution.



290 **Figure 5: Example of a representative principal components analysis (PCA) hexagon plot based on variability in HydroSHEDS river subcatchment attributes. Hexagon colour indicates the number of subcatchments per hexagon. Subcatchments hosting GRiMeDB sites are plotted in red and contained within the convex hull delineated by red lines. Subcatchments that fall within this polygon are assigned a value of 1 and those outside the perimeter are given a value of 0 to indicate the representativeness of sampled reaches for this pair of PC axes. See Sect. 2.4 for further explanation.**


2.5 Data checking and data analysis

Several approaches were taken to check the accuracy of data in GRiMeDB. This included evaluation of the reliability of digitized data (Sect. 2.3) along with several additional inspection steps. Entries were error checked by a co-author other than the individual who entered the data, including confirmation of site location information, validating units for all variables, and spot- or complete checking of entered gas data (independent units and data check in Fig. 4), depending on dataset length and if data were manually entered or imported directly from a file. Once values had been converted to standard units, all variables were plotted to identify outliers (outlier check; Fig. 4), and extreme values were checked against the original data source. In cases in which errors were present in the original data, if possible, authors were contacted for clarification. In the few rare cases in which issues could not be resolved, the data were excluded. These and all other calculations and analyses were performed in R (version 4.2, R Core Team 2021), using the “dplyr” package (version 1.0.7, Wickham et al., 2021) and “data.table (Dowle and Srinivasin, 2021) for data analysis, “sf” package (version, 1.0, Pebesma, 2018) for spatial data processing, and “ggplot2” (version 3.3.5, Wickham, 2016) and patchwork (Pedersen, 2020) packages for visualization.

3. RESULTS

3.1 Overview of GRiMeDB data

GRiMeDB includes 24,024 records of CH₄ concentration and 8,205 CH₄ flux values from 5,037 unique sites, along with 17,655 and 8,409 concurrent measurements of concentration and 4,444 and 1,521 of flux for CO₂ and N₂O, respectively (Table S2). Although the first concentration and flux values in GRiMeDB were published in 1973 (Lamontagne et al., 1973) and 1987 (de Angelis and Lilley, 1987), respectively, over 70% of all CH₄ concentrations and 80% of flux observations became available after 2015 (the year of publication of MethDB; Fig. 6, Fig. S1). This growth in data availability has occurred predominantly along the spatial axis, as almost two thirds of all sites were added in or after 2015 and over half of all sites in the database have a single concentration and/or flux observation. Conversely, long timeseries are rare, with only 8% of the 5,037 sites having > 10 concentration observations and 4% having >10 diffusive flux records (Fig. 6, Fig. S1). The longest concentration record includes 590 observations distributed over 28 years (Toolik Inlet, *Site_ID* 9025; Kling, 2019a; 2022) while the longest flux record has 82 observations of diffusive flux over 4 years (*Site_ID* 3644; Aho et al. 2021). Further, among the 15 sites with time series >5 years, 12 are situated in either the Toolik Lake region of Alaska, USA (Kling, 2019a; 2019b; 2022) or within the Krycklan watershed in Sweden (Wallin et al., 2018; Wallin, 2021).




325 **Figure 6: Distribution of the number of methane (CH₄) observations per site. Brown bars indicate sites with both concentration (conc) and flux observations. Orange and green bars show sites with only concentration and only flux observations, respectively. Inset: Cumulative observations of CH₄ concentration and flux data based on the year of publication of the data source. The vertical line (2015) indicates the year of MethDB (Stanley et al., 2015) publication. See Fig. S1 for data accumulation and length resolved by CH₄ flux type.**

330 **3.2 Spatial and temporal distribution of data**


Spatially, 40% of all sites and 52% of all CH₄ concentration observations are in North America, followed by Europe (25% of all sites and 26% of all CH₄ concentration values; Table S2). Conversely, there are vast geographic areas with moderate to high channel densities with few or no observations, such as central Canada, Central America, South America beyond the Amazon mainstem area, most of Russia, central and western Asia, New Zealand, and the Malay Archipelago (Fig. 7a).
 335 Geographic limitations in availability of flux data, particularly of ebullition, are pronounced given smaller number of observations and domination of diffusion measurements. Observations of ebullition are absent or limited to 1-2 studies for

Africa, Oceania, Central America, South America, and Russia (Fig. S2). Despite these gaps, there is surprisingly good representation in terms of the range of hydrological, physiographical, climatic, land cover, geological, edaphic, and anthropogenic conditions that exist globally (Fig. 7b). Areas that are poorly represented are characterized by very low channel density associated with arid or polar climates as well as high altitude regions (Greenland, northern Canada, northern Africa, central Australia, Middle Eastern nations, western China, Mongolia, Chile, southern Argentina). Evaluating the distribution or representativeness of sites in terms of system size is difficult given the limited availability of relevant information such as Strahler stream order or basin area, which were reported for only 26% and 28%, respectively, of all sites (Table S2). For sites with these data, counts of observations decline with increasing stream order (Fig. 8) in a log-linear fashion ($R^2 = 0.92$ for concentration and 0.90 for flux; $P < 0.0005$ for both regressions after excluding zero-order counts), consistent with Horton's Law of Stream Numbers (Horton, 1945). Thus, other than the extreme under-representation of zero-order channels, this predictable decline suggests reasonable representation by order. Nonetheless, this result should be interpreted with caution given the scarcity of relevant data. The distribution of counts by basin size follows a similar pattern of under-representation of sites draining very small basins and also indicates a potential over-representation of some large basin sizes (basins of ca. 10,000 km²; Fig. 8).

The distribution of observations among months illustrates seasonal sampling regimes dominated by summer sampling in northern ($>40^\circ$) and southern ($<-20^\circ$) latitudes contrasted by even or erratic sampling at mid-latitudes (Fig. 9). Consistent with the lower representation of southern hemisphere rivers and streams, several months lack concentration and/or flux measurements south of -10° latitude, particularly during winter months. Beyond these gaps, the only months missing data in the northern hemisphere are fluxes in January and February at sites north of 60° latitude and several missing months north of 70° , presumably due to pervasive ice and snow cover.



360 Figure 7: (a) Global distribution of methane (CH_4) observations in the database, colour coded for sites with concentration data only,
 flux data only, or both concentration and flux data. Top and right panels show, respectively, longitudinal and latitudinal patterns of
 the density of CH_4 observations (grey bars) and the density of river area (blue bars). These bars have been aggregated at a 1
 latitudinal or longitudinal degree and rescaled from 0 to 1 for this visualization. River area was obtained from BasinAtlas (Linke et
 al. 2019). (b) Representativeness (dimensionless) of the database based on a wide array of biological, physical, hydrological and land
 365 cover variables (see Sect. 2.4 for details). Values close to 1 indicate a high representativeness, with only 4% of the global river surface
 below a threshold of 0.9. See Fig. S2 for data distribution resolved by CH_4 flux type.



370 **Figure 8: Number of sites with concentration (a, b) or diffusive flux (c, d) observations as a function of stream order (a, c) and basin size (b, d) for the subset of sites with channel order and/or basin size information.**





Figure 9: Number of observations of concentration (left) and flux (right) by month for 10° latitude bands.

375 3.3 CH₄ flux methodology


Records of CH₄ flux are dominated by diffusive flux measurements, which represent 85% of all flux values in the database, with ebullition (8%) and total flux (7%) accounting for the remaining entries (Fig. 10). Not surprisingly, a variety of methods have been used to quantify each flux type, although diffusive flux methods are dominated by calculations based on dissolved gas concentration and a gas exchange coefficient (k) (74% of all observations), while chamber-based methods are most common for quantifying total flux (93% of all observations). Similarly, k is most commonly estimated via physical models ($n = 3188$). Several models have been employed for this calculation, as indicated by >25 different references for k model sources listed in GRiMeDB.



385 **Figure 10: Counts of methane (CH₄) flux observations by type (left), by major methodological categories for each pathway (middle), and for method type used to estimate the gas exchange coefficient k (right). For clarity, the chamber category includes all chamber types and patterns of gas increase in the chamber unless specified; more resolved methodological data are presented in the GRiMeDB Flux Table. See Table A4 for further details about category definitions.**

3.4 Overview of concentration and flux data

390 Concentrations and fluxes of all three gases are characterized by log-normal distributions that vary over several orders of magnitude (Fig. 11) and large coefficients of variation (CVs) for CH₄ and especially N₂O (Table 2). The vast majority (~95%) of CH₄ and CO₂ concentrations appear to be supersaturated, in contrast to N₂O concentrations with 67% of observations above this threshold. Reports of concentrations below detection are scarce for all gases, including N₂O (Table 2).



395


Figure 11: Histograms of gas concentrations and fluxes in GRiMeDB, excluding values reported as below detection or zero; counts of these values are reported in Table 3. Dashed vertical lines in the concentration histograms indicate the 100% saturation concentration based on the median estimated elevation (250 m) and water temperature (12.5) for all sites and atmospheric concentrations of 1.83, 400, and 0.325 ppm for methane (CH₄), carbon dioxide (CO₂), and nitrous oxide (N₂O), respectively.

400 The fraction of observations with zero, below detection, or negative fluxes (5%, 5%, and 19% for diffusive CH₄, CO₂, and N₂O fluxes, respectively), were similar to the frequency of subsaturated concentrations. At the other extreme, the highest CH₄ concentrations (>200 μmol L⁻¹) paradoxically occur in either anthropogenically influenced large rivers of the warm tropics (e.g., Amazon basin: Kemenes et al., 2007; Ganges, Mekong: Begum et al., 2021) or in small boreal headwater streams (e.g., Campeau et al., 2018; Wallin et al, 2018).

405 There were no meaningful univariate relationships between variables that may be used for upscaling (latitude, basin area, and stream order) and mean site concentration or flux (Fig. 12, Table S3). Linear regressions indicated that latitude and flux accounted for a very small percent of the variation in both concentration ($R^2 = 0.006$ and 0.002 , respectively) and flux ($R^2 = 0.036$ and 0.055) among sites. Similarly, concentration and flux among stream orders suggested possible differences for concentration (Kruskal-Wallis $\chi^2 = 47.165$, $df = 8$, $P < 0.001$) and flux ($\chi^2 = 14.777$, $df = 8$, $P = 0.070$). However, results of
 410 corrected pairwise comparisons (using the method of Benjamini and Hochberg, 1995) among orders were ambiguous, suggesting no differences among orders for flux. For concentration, these comparisons indicated possible differences in distributions only between 7th order channels and all other orders, and between 6th vs 1st order sites for concentration. Collectively, these results indicate a lack of a consistent change in CH₄ magnitude across channel orders for flux.

415 **Table 2. Summary statistics for methane (CH₄), carbon dioxide (CO₂), and nitrous oxide (N₂O) concentrations and fluxes. The %BDL (below detection level) column reports the percent of all observations that are below detection limits (including values reported as zero) for concentration. See Table S2 for counts and Table S3 for statistical summaries for all other variables. Standard deviation (SD) and coefficient of variation (CV).**

Metric	Gas (and type)	Mean	Median	Max	Min	SD	CV	%BDL
Concentration (μmol L ⁻¹)								
	CH ₄	1.49	0.20	456	0	10.69	718	3.2
	CO ₂	135	81.7	5,479	0	174.8	130	0.05
	N ₂ O	0.058	0.017	32.9	0	0.602	1,042	0.59
Flux (mmol m ⁻² d ⁻¹)								
	CH ₄ -diffusive	7.31	0.44	4,057	-136	86.4	1,182	
	CH ₄ -ebullitive	5.42	0.28	366	0	24.02	443	
	CH ₄ - total	8.71	0.63	366	-0.05	31.90	366	
	CO ₂	319	128	23,749	-1625	770	242	
	N ₂ O	0.082	0.008	31.3	-11.3	0.981	1,199	



420

Figure 12: Site-average methane (CH₄) concentrations (a-c) and flux (d-f) as function of latitude, basin area, and Strahler stream order. For boxplots, the upper and lower edges of each box are the 25th and 75th percentiles and whiskers are drawn up to 1.5 times the interquartile range.


425 As with relationships between CH₄ and physical site attributes, relationships between CH₄ concentration or flux and water chemistry parameters are also characterized by substantial variability. Representative examples indicate increasing, decreasing, and ambiguous relationships between CH₄ concentrations and fluxes and selected chemical constituents (Fig. 13). One source of the variation in the relationship shown in Fig. 13 may be attributed to differences among sites, as is illustrated for the case of CH₄ concentration versus discharge. The cluster of points in this plot (Fig. 14a) does not suggest an obvious linear relationship between concentration and discharge; however, resolving the data to the site level for sites with multiple observations reveals several significant trends (Fig. 14b). Among 57 sites with >30 observations, 42 had significant relationships ($P < 0.05$) between concentration and discharge and 30 of these 42 trends were negative.

430

Median site concentrations for most categories of targeted channels (Fig. 15) differed from “normal” (NORM) sites (Kruskal-Wallis test $\chi^2 = 460.1$, $df = 12$, $P < 0.0001$ after dropping channel types with <10 observations to improve test reliability).

435


Pairwise Wilcoxon comparisons adjusted to account for multiple comparisons (Benjamini and Hochberg, 1995) indicated that springs (SP) and delta channels (DC) were similar to NORM sites ($P > 0.4$) and impoundment-influence (IMP) sites were marginally different ($P = 0.053$). Concentrations in channels at glacial termini (GT) and floodplain backwaters (FP) were lower ($P < 0.0001$), whereas all other site types had higher site average CH_4 concentrations than NORM sites. Fluxes also varied among channel type (Kruskal-Wallis test $\chi^2 = 126.4$, $df = 8$, $P < 0.0001$ after dropping channel types with < 10 observations), and similar to concentration, fluxes in delta channels (DC) and channelized sites (CH) were similar to NORM channels while all other channel types considered had higher median fluxes.




445

Figure 13: Methane (CH_4) concentration (a-d) and diffusive flux (e-h) versus concurrent measures of dissolved oxygen (O_2 ; $n = 8,529$ and $2,316$ for concentration and flux, respectively), dissolved organic carbon (DOC; $n = 14,441$ and $1,901$), total nitrogen (Total N; $n = 8,378$ and 467) and total phosphorus (Total P; $n = 6,904$ and 240). Three outliers were excluded from the DOC plots, and because of the log scale for CH_4 , negative and zero values have been omitted.

450



455 **Figure 14: Methane (CH₄) concentration versus concurrent measures of discharge for (a) all sites with discharge data and (b) sites with >30 observations (57 sites) with trend lines denoting within-site relationships between concentration and discharge. Each site is represented by a separate colour. Because of the log scale for CH₄, negative and zero values are omitted.**



460 **Figure 15: Boxplots of site-averaged Methane (CH₄) concentration (a) and diffusive flux (b) for channel type categories. Channel**
 categories are defined in Table 1, but briefly are as follows: NORM- non-targeted sites; CAN- canals; CH- channelized streams; DC-
 river delta channels; DD- downstream of dams; DIT- ditches; FP- floodplain backwaters; GT- glacial outflows; IMP- impounded
 465 reaches; PI- permafrost (thermokarst) influenced; PS- point source influenced; SP- springs; TH- thermogenic CH₄ inputs; WS-
 wetland streams. Number of sites per channel type are listed on the right side of each plot. The vertical black line denotes the median
 concentration and flux for non-targeted (NORM) sites. Because a log-scale is used in these plots, zeros and negative values were
 excluded. The actual median for non-targeted sites represented by the vertical line is therefore slightly different than the median
 displayed in the corresponding box plot because of this exclusion. The upper and lower edges of each box are the 25th and 75th
 percentiles, whiskers are drawn up to 1.5 times the interquartile range, and points are plotted if beyond the whiskers.

470 4. Discussion

The rapid increase in availability of aquatic CH₄ (as well as CO₂ and N₂O) data over the past 5-10 years has been remarkable and creates new opportunities for examining patterns and drivers of these gases in lotic ecosystems across broad spatial scales. Similarly, constructing GRiMeDB provided us with an unprecedented opportunity to identify tendencies in when, where, and

475 how CH₄ has been sampled in streams and rivers. Examination of such data collection tendencies can reveal important biases and gaps within a field (Stanley et al. 2019, Gomez-Gener et al. 2021b) and thus points to future research needs and opportunities. Below, we discuss the distribution of sampling efforts and methodological issues, preliminary data analyses, and consider questions that GRiMeDB can help to answer.

4.1 When and where: sampling effort considerations

480 The growth of GHG studies in flowing water systems in the past decade includes a geographic expansion beyond the large body of historic and current work in temperate regions of North America and Europe. In particular, recent research in Africa, Australia, and especially southeast Asia has greatly improved the global coverage of available data. However, studies in arid drainages remain scarce- even beyond what would be expected given their small river surface area. A possible explanation for the limited study of CH₄ in these systems may be the pervasive focus on the contribution of streams and rivers to the global
485 atmospheric CH₄ pool, and the corresponding assumption that aridland systems play a minor role in this context. Yet we suggest that limited study in arid and semi-arid drainages represents a missed opportunity to understand metabolism and carbon cycling in a set of streams and rivers that drain nearly half of the global land surface, are increasingly stressed by growing human water demands (e.g., Sabo et al., 2010; Lian et al., 2021; Stringer et al., 2022), and support ecosystem process rates that are amplified by warm temperatures and highly variable flow regimes (Fisher et al., 1982; Ran et al., 2021). Beyond arid
490 and semi-arid basins, further research emphasis in tropical and high-latitude regions would also be beneficial even given recent improvements in data availability and geographic representation of both areas. Existing data for tropical forests and grasslands are dominated by studies of African rivers (especially the Congo drainage) and the Amazon River system. In fact, observations from tropical areas of the Indomalayan and northern Australasian region represent <3% of all sites, and Central America is represented by a single study. Tropical drainages are frequently characterized by high CH₄ concentrations and fluxes, along
495 with rapid changes in land use and river regulation that are affecting C cycling and GHG dynamics (Park et al., 2018; Flecker et al. 2022). However, understanding or detecting the magnitude and consequences of these anthropogenic changes on fluvial CH₄ is constrained by these current sampling limitations. Finally, while high latitude regions (north of the Arctic Circle) are well represented in GriMeDB with >3,600 concentration observations, more than 80% of these values are derived from studies in the vicinity of the Toolik Field Station in Alaska, USA, and thus do not capture the full biophysical diversity of Arctic
500 biomes (Metcalf et al. 2018). Given that climate change at high latitudes is progressing faster than elsewhere on the planet (IPCC, 2021), and that the global north stores massive quantities of C in soils (Hugelius et al., 2014), more extensive coverage of CH₄ across Arctic drainage systems is warranted.

Although the spatial coverage of CH₄ data has improved markedly over the past decade, expansion across temporal dimensions
505 has lagged. The predominant mode of sample collection has been and continues to be through surveys that yield one or a few observations from individual sites (e.g., Bouillon et al., 2012; Kuhn et al., 2017; Jin et al., 2018; Ho et al., 2022), and studies characterizing seasonal dynamics or responses to a site-specific environmental change are limited. Indeed, long-term (>5 years)

CH₄ datasets in general are extremely rare (Leng et al., 2021); no such data are currently available for fluxes and most long-term concentration records are derived from just a few clustered locations. Determining the consequences of changes in land use or habitat attributes on fluvial CH₄ dynamics have instead relied on space-for-time substitutions (e.g., Smith et al., 2017; Gatti et al., 2018; Woda et al. 2020) rather than on direct observations of change over time. Although this strategy has been successful in revealing variation in GHG dynamics among different site types, current knowledge about how gases vary over time and respond to perturbations is poorly developed because of these data limitations. This deficit may be particularly consequential in the case of climate change, as the broad scope of this phenomenon will inevitably limit the effectiveness of spatial sampling approaches.

The discussion above regarding the ‘when’ and ‘where’ of sampling emphasizes large spatial and relatively long temporal scales, consistent with the extent of GRiMeDB. However, another current deficit in our understanding relates to the degree of heterogeneity of this gas at fine spatial and temporal scales, and thus if current sampling strategies are missing meaningful variation. Recent studies of CO₂ provide a cautionary tale in this context, as failure to account for diurnal variation in this gas results in a consistent under-estimation of fluvial emissions that is quantifiable at regional (Attermeyer et al., 2021) and global (Gómez-Gener et al., 2021b) scales. Similar questions may arise for spatial variation; that is, what is the minimum grain size or appropriate spatial scale for sampling CH₄ in running waters (Crawford et al., 2017; Lupon et al., 2019)? The potential to examine very short-term variation is not possible using GRiMeDB data because of our decision to average within-day measurements given the current small number (ca. 20) of these temporally detailed studies. Assessment of fine-scale spatial variation is also limited because of limited fine-scale sampling in general, as well as by decisions made both by investigators and during database construction. For example, geomorphologically distinct units (e.g., an individual riffle or pool) are often used as a basic sampling unit and results are presented as averages of replicates collected at different points within the study reach (e.g., Hlaváčová et al., 2006; Smith et al., 2017). In general, information about replication was frequently omitted, or if reported, information about variability among replicates was frequently absent. In addition to this limitation, our decision to combine replicates taken at different points in a channel cross-section and within individual channel units that had hundreds to thousands of datapoints to avoid ambiguities for site delineation and data analysis also constrains the opportunity to examine variation at fine spatial scales. However, we anticipate that this situation will change over the next few years, as *in situ* sensors or other devices capable of collecting high-frequency/high density gas measurements become more widely available. Recent papers signal this new frontier and have highlighted the presence (e.g., Lamarche-Gagnon et al., 2019; Smith and Bohlke, 2019; Chen et al., 2021; Taillardat et al., 2022) and absence (e.g., Castro-Morales et al., Chen et al., 2021; 2022; Rovelli et al., 2022; Zhang et al., 2021) of predictable diel variation in CH₄ concentrations and fluxes, and varying degrees of within-reach spatial variability (Crawford et al., 2016; 2017; Call et al., 2018; Bussman et al., 2022).

540

4.2 How: methodological considerations

Measuring dissolved GHG concentrations or fluxes involves multiple steps and calculations. Field and laboratory protocols vary widely in the literature, and methodological variety is particularly conspicuous for flux determination. Ironically, even though many studies of lotic CH₄ dynamics are framed in terms of understanding the contribution of these ecosystems to the rapidly increasing atmospheric CH₄ pool, flux measurements lag far behind those of concentration and the vast majority (ca. 85%) of observations are of the diffusive pathway alone. Further, the most common method for estimating this pathway involve combining dissolved CH₄ concentration with k , the gas exchange coefficient. Quantifying k is notoriously challenging (Hall and Ulseth, 2019) and the large number of approaches for calculating k used among data providers is concerning and undoubtedly introduces substantial uncertainty. A more in-depth consideration of the consequences of different models or strategies for arriving at a k value was beyond the scope of this paper, but inclusion of methodological information should be useful for such an analysis in the future.

Ebullition measurements are notably scarce despite the potential of this pathway to account for a large fraction of total emissions in some streams (e.g., from 30-98% of total CH₄ emissions, as shown in Baulch et al., 2011; Crawford et al., 2014; Chen et al., 2021). The conventional approach to quantifying ebullition involves a combination of capturing bubbles just below the water surface to determine the area and time-specific rate of bubble volume reaching the surface and measuring CH₄ content of recently-erupted bubbles. The episodic nature and extreme spatial heterogeneity of ebullition (Crawford et al., 2014; Spawn et al., 2015; Chen et al., 2021; Robison et al., 2021) require multiple bubble trap replicates that need to be deployed over several days to generate reliable measurements. Given the logistic challenges and labour-intensive work involved, indirect approaches are becoming more common. These approaches typically use the difference between a chamber-based measurement of flux, which is assumed to represent total flux (diffusion + ebullition) and diffusion calculated from dissolved CH₄ and k (i.e., the 'chamber – [concentration + k]' method in Fig. 10) to estimate ebullition (e.g., Campeau et al., 2014; Zhang et al., 2020; Ran et al., 2021). We suggest that this approach should be used cautiously, however. This strategy is arguably inappropriate for situations in which the gas content within a chamber increases in a linear fashion during the measurement period, consistent with the occurrence of diffusive flux alone. Second, it is not clear if it is reasonable to assume that chamber-based measurements capture both diffusion and ebullition, even if a chamber-based flux value is greater than that calculated from dissolved CH₄ concentration. Relatively short chamber deployments are likely to miss or incompletely capture bubble releases, while long-term deployments are vulnerable to sampling artefacts associated with altered concentration gradients within, and/or turbulence around the chamber (Sawakuchi et al., 2014; Lorke et al., 2015). Given these challenges, it is not altogether surprising that comparisons between direct and indirect measurements of ebullition can yield substantially different results (e.g., Yang et al., 2012; Bednařik et al., 2017; Chen et al., 2021).

The final and most profound knowledge gap in the collection of flux data is the absence of measurements of plant-mediated emissions. Plant-mediated fluxes can account for a substantial fraction of total emissions from wetlands and shallow lake

habitats (Bodmer et al., 2021), but the contribution of this pathway is unknown in fluvial systems. Indeed, we did not include
575 plant mediated fluxes in GRiMeDB because we encountered only two papers that had explicitly quantified this pathway in
streams (Sanders et al., 2007; Wilcock and Sorrell, 2008). Although aquatic macrophytes are sparse or absent from many
streams and rivers, they can be abundant in low-gradient, low-disturbance environments (Riis and Biggs, 2003; Gurnell et al.,
2010) where diffusive fluxes would be constrained by low gas exchange rates. Sediment trapping and venting by macrophytes
enhances both methanogenesis and methane emission in these systems (Sanders et al., 2007), but the significance of such
580 processes and the contribution of plant-mediated fluxes at larger spatial scales remain to be determined for fluvial systems
(Bodmer et al., 2021).

4.3 Concentration and flux patterns

Not surprisingly, the massive increase in data availability has led to differences in averages and measures of variability for
585 CH₄ concentrations and fluxes compared to our previous efforts. Median values for all three CH₄ flux pathways in GRiMeDB
are 1.2-2.2 times lower than those reported by Stanley et al. (2016), as well as those from Rosentreter et al. (2021). Conversely,
measures of variability (SD, CV) in GRiMeDB are almost 3-fold greater than previous estimates, undoubtedly due to the far
larger number of observations, the associated expansion of geographic scope and channel types, and the inclusion of temporally
resolved data. It is not yet clear if the sample sizes are sufficient to capture the true global-scale variability of fluvial
590 concentrations and fluxes, and future database updates could be used to examine this possibility.

Despite the slight lowering of median values compared to previous estimates, supersaturated concentrations and positive fluxes
are the norm for CH₄ as well as for CO₂ and N₂O. However, it is likely that CH₄ concentrations and fluxes below detection
limits (BDLs) are under-reported, as is common with environmental data in general (Stow et al., 2018), so these latest estimates
595 may still be slight overestimations of true population medians. Even given the modest number of zero or undetectable CH₄
concentrations in GriMeDB (<2.5%), decisions about handling BDLs can have a small but detectable effect on the estimation
of global averages. For example, if these observations are excluded, median CH₄ concentrations for all other observations
increases from 1.49 to 1.51 μmol L⁻¹. If we keep all of these observations and assign them a value of zero (an unlikely scenario,
but used here to provide a lower limit for this example), then the overall median declines to 1.46 μmol L⁻¹. Although these
600 differences are relatively small, it would likely be consequential for upscaling estimates. At a minimum, we urge GRiMeDB
users to be aware of how these values are handled and encourage future researchers to determine and report detection limits
and include samples that fall below these limits in their results.

A goal of assembling GRiMeDB was to centralize CH₄ data to foster future research efforts. To this end, we included
605 information about habitat conditions that allows the exploration of relationships between CH₄ and potential explanatory
variables and covariates. To demonstrate this opportunity, we provided a limited number of examples of CH₄ versus variables
that have been identified as potential predictors or drivers of CH₄ production, concentration, or flux (Figs. 12-14), and these

plots suggest both the presence and absence of relationships. For example, increasing CH₄ concentrations have been associated with low or decreasing dissolved oxygen and/or increasing organic carbon (e.g., Borges et al., 2018; Jin et al. 2018; Begum et al., 2021) and these relationships are recognizable for concentration but ambiguous for flux across the entirety of the GRiMeDB dataset. Similarly, increased CH₄ production and emissions tend to be elevated in nutrient-rich (eutrophication) lakes (DelSontro et al., 2018) and polluted rivers (Rajkumar et al. 2008; Ho et al., 2022), consistent with positive relationships between CH₄ flux and TN and TP. However, nutrient enrichment in rivers often occurs concurrently with fine sediment and organic matter input; thus it remains to be determined if positive relationships in Figs. 13g and 13h are correlative or reflect a causal mechanism. Finally, increases in discharge have been linked to declines in gas concentration, likely due to source limitation (i.e., dilution) of terrestrial supply (Aho et al., 2021; Gómez- Gómez-Gener et al., 2021a) and/or greater water turbulence, which increases gas exchange and thus reduces supersaturated CH₄ stocks (Billett and Harvey, 2013; Kokic et al. 2018). This relationship is not obvious when all data were considered en masse, but became more apparent when examining within-site dynamics. In contrast to these three confirmatory examples, although latitude and channel size have also been identified as determinants of CH₄ concentrations or used to extrapolate site-specific gas measurements to larger (even global) scales (e.g., Bastviken et al., 2011; Li et al., 2020; Rosentreter et al., 2021), evidence for such relationships is absent from our analysis. Further, even for the former examples that indicated relationships between CH₄ concentration and DO, DOC, or discharge, there is substantial variability present in these relationships, the strength of these predictors is likely to vary across scales, and they explain little of the variability for diffusive fluxes. In short, substantial opportunities exist to identify multivariate relationships between different predictors and CH₄ concentrations and fluxes across different scales, and pursuit of these opportunities should be improved by the substantial increase in data for both gases and potential predictor variables.

The disproportionate contribution of streams and rivers to atmospheric inputs together with the utility of CH₄ as an indicator of anthropogenic influences on drainage systems have inspired several studies that focus on fluvial habitats that are expected to have high concentrations and fluxes. Many of these ‘methane hunting’ studies have demonstrated significant increases in CH₄ concentrations and/or fluxes associated with phenomena such as point source inputs (Alshboul et al., 2016) ditch and canal construction (Peacock et al., 2021), oil and gas extraction (Woda et al. 2020), or passage through wetlands (Taillardat et al., 2022). Such signals persist at the global scale (Fig. 15), highlighting widespread human enhancement of CH₄ emissions from lotic ecosystems. Not all targeted sites are CH₄-rich however. Low concentrations in glacial outflows (GT) likely reflect the effects of cold temperatures and/or low organic carbon availability (Crawford et al., 2015; Burns et al., 2018) while low values at floodplain (FP) sites may be attributable to their more characteristically lentic conditions, which favor higher rates of CH₄ oxidation in the water column. Indeed, oxidation has been shown to represent a significant CH₄ sink in floodplain lakes associated with the Amazon River (Barbosa et al., 2018) and most of the FP sites in GRiMeDB are part of the Amazon system.

As noted in Sect. 3.4, the availability of supporting information is inconsistent, as, for example, only ~25% of data sources provided data on channel order or basin size. However, open-access regional and global geospatial datasets that provide

information about site characteristics (e.g., Linke et al., 2019; Yang et al., 2020) have increased rapidly in the past decade, to the benefit of analyses seeking to link landscape attributes to CH₄ distribution among sites. Recent upscaling efforts analyses (Rosentreter et al., 2021; Liu et al., 2022; Rocher-Ros et al., accepted) have, for example, capitalized on improved estimates of the surface area of world streams and rivers (Allen and Pavelsky, 2018; Yang et al., 2020), while the diverse datasets in
645 HydroSHEDS (Linke et al., 2019) allowed us to evaluate the global representativeness of GRiMeDB sites. As new global-scale datasets become available and become more spatially resolved, we anticipate that their pairing with GRiMeDB data will result in significant improvements in the strength and certainty of data assimilation models, regional to global-scale analyses of CH₄ distribution and drivers, and quantification of fluvial emissions to the atmosphere.

650

5. Data and code availability

GRiMeDB and its associated metadata are available from the the Environmental Data Initiative (Stanley et al., 2023):
<https://doi.org/10.6073/pasta/f48cdb77282598052349e969920356ef>.

655

6. Conclusion

The data gathered in GRiMeDB highlight many new opportunities, both through analysis of CH₄ and supporting data in the database, and by revealing gaps that currently exist across fluvial CH₄ studies. The most conspicuous data limitations include
660 deficits in measurements of non-diffusive flux pathways and underrepresentation of sites from arid, tropical, and arctic biomes. Challenges associated with quantifying ebullition discussed earlier also emphasize the need for more intercomparisons among flux methodologies. Regardless of pathway, flux is a difficult process to quantify and can be highly sensitive to methods or gas exchange model choices, yet there are few comparisons (such as Raymond et al., 2012; Lorke et al., 2015) available to inform these decisions. Finally, we highlight that the expansion of GHG data over the past decade has proceeded largely across
665 spatial rather than temporal dimensions. While this expansion has vastly improved the geographic representativeness of the data, long-term datasets are rare despite their power for generating ecological understanding and informing policy/management in the face of environmental change (Hughes et al., 2017). GHG's are rarely included as routine components of water quality monitoring programs. Thus, we emphasize the compelling need to establish such sampling efforts and perpetuate those few that do exist.

670 Given the rapid growth in both research interest and data in fluvial GHG dynamics, we imagine future updates and expansion of GRiMeDB and we welcome datasets and associated research products (e.g., theses, journal publications, reports, etc.). To facilitate the data acquisition and updating process, a downloadable spreadsheet template and detailed information about its use and submission are available at <https://stanley.limnology.wisc.edu/GRiMe>. Regardless of database updates, we recommend that the minimum basic information to collect along with GHG data that would be most valuable for later analyses include:

675 well-resolved site location data (latitude and longitude); information about site size (Strahler order and/or basin size at the
sampling site), disturbance or modification relevant to GHGs (e.g., categories listed in Table 1); specific sample dates and
times; discharge, dissolved oxygen, and temperature at the time of sample collection; and clear information about units and
method(s) used to measure gas flux. Finally, we strongly encourage data package (sensu Gries et al., 2022) publication in a
680 trustworthy public data repository such as the Environmental Data Initiative that requires metadata to meet FAIR data
principles and increase data findability, accessibility, and re-use (Wilkinson et al., 2016).

Despite highlighting areas of data limitation, it is important to underscore the opportunities that the growth in GHG data
availability- especially of CH₄ data- now provide. Assembly of GRiMeDB was motivated by the goal of having a centralized,
standardized resource to facilitate further studies of CH₄ pattern and process in flowing water systems. Our strategy in
developing this database was to maximize opportunities for identifying patterns and relationships involving this gas in future
685 analyses. Past difficulties with such efforts may well be a product of the common practice of averaging values over time or
among sites and/or of including non-fluvial sites in analyses. Thus, we carefully documented the data and resolved observations
to individual sites and dates whenever possible to match the pronounced spatial and temporal variance of this gas. Similarly,
while we included a range of habitat types in GRiMeDB, unconventional or targeted sites are easily identifiable. Further, we
carefully examined sites to ensure that they were not situated within reservoirs/impoundments or estuaries where distinct
690 processes such as methane oxidation, tidal cycles, or elevated sulphate reductions may obscure or overtake relationships
present in inland flowing water systems. Thus, we are optimistic that analysis of GriMeDB data by itself, or in concert with
other complementary datasets, will provide new and unprecedented opportunities to examine relationships between CH₄ and
environmental drivers or correlates, as well as providing broad contextual information for site-based studies of fluvial carbon
and GHG dynamics.

695

Appendix A. GRiMeDB tables and variables

Table A1. Column titles and description of their content for the GRiMeDB Sources Table.

Column Title	Description
Title	Title of data source
Author	Lead author last name
Source	Identity of the outlet for the data (e.g., journal, data repository, agency that presented the data). For titles with published papers paired with published datasets, the journal is listed in this column
Pub_year	Year of publication, data release, or acquisition of an unpublished dataset
Source_ID	Unique data source identifier
Additional_data	“Yes” in this column indicates that additional data were acquired directly from the author for any field. Additions are described in the Comments field
Comments	Additional information or clarification about the data source
Paper_DOI	DOI or hyperlink for journal article or other publication based on the CH ₄ data
Data_DOI_primary	DOI or hyperlink for CH ₄ data posted in a data repository
Data_DOI_supporting	DOI or hyperlink for separate datasets providing supporting data

Table A2. Column titles and content description for the GRiMeDB Sites Table.

Column Title	Definition
Source_ID	Unique data source identifier from the Sources Table
Site_ID	Unique site identifier
Site_Name	Unique site name
Stream_Name	Stream or river name; taken or modified from the data source or generated <i>de novo</i> when a name was not specified in the data source
Aggregated	Yes or No; “Yes” if CH ₄ data entered are averages from >1 site
N_sites_aggregated	Number of sites that were averaged for aggregated sites
Basin_Region	Name of the larger drainage basin or region that contains the site. This information is included to facilitate site grouping during data analysis
Latitude	Decimal degrees, WGS84 ensemble: EPSG:4326 coordinate system
Longitude	Decimal degrees, WGS84 ensemble: EPSG:4326 coordinate system
Elevation_m	Reported meters above sea level
Slope_m_per_m	Reported channel slope expressed as m m ⁻¹
Strahler_order	Reported Strahler stream order
Basin_size_km ²	Reported basin size in square kilometers
Channel_type	Codes denoting distinct site or channel attributes or presence of specified conditions. See Table 1 for categories and their definitions
Latitude_snapped	Latitude in decimal degrees for site location after snapping to the closest channel for elevation determination
Longitude_snapped	Longitude in decimal degrees for site location after snapping to the closest channel for elevation determination
Elevation_estimated_m	Elevation (meters above sea level) calculated from the DEM. See Sect. 2.2 for details
Comments	Additional information or clarification about the site source

Table A3. Column titles and definitions for the GRiMeDB Concentration Table

Column Title	Definition
Source_ID	Unique paper identifier from the Sources Table
Site_ID	Unique site identifier from the Sites Table
Site_Name	Unique site name from the Sites Table
Conc_Name	Unique name for the sampling event at the site; same as Flux_Name in the Flux Table if both concentration and flux data for the same site-date combination are available
Date_start	First sampling date
Date_end	Last sampling date; this is the same date as the Date_start if data are not aggregated over time
Aggregated_Space	Yes or No; “Yes” if CH ₄ data entered are averages from >1 site
Aggregated_Time	Yes or No; “Yes: if CH ₄ data entered are averages from >1 date
FluxYesNo	Yes or No; “Yes” if there is a corresponding flux measurement associated with this site-date combination
SampleCount	Number of samples or observations corresponding to the mean or median concentration
CH4min	Minimum measured CH ₄ concentration in $\mu\text{mol L}^{-1}$ if data are aggregated spatially or temporally, has multiple within-day measurements (e.g., a diel study), or are from a data-dense spatial study
CH4max	Maximum measured CH ₄ concentration in $\mu\text{mol L}^{-1}$ if data are aggregated spatially or temporally, has multiple within-day measurements (e.g., a diel study), or are from a data-dense spatial study
CH4mean	Mean or sole reported CH ₄ concentration in $\mu\text{mol L}^{-1}$ for the sampling event
CH4_SD	Standard deviation of the mean CH ₄ concentration
CH4median	Median CH ₄ concentration in $\mu\text{mol L}^{-1}$
CO2min	Minimum measured CO ₂ concentration in $\mu\text{mol L}^{-1}$ if data are aggregated spatially or temporally, has multiple within-day measurements (e.g., a diel study), or are from a data-dense spatial study
CO2max	Maximum measured CO ₂ concentration in $\mu\text{mol L}^{-1}$ if data are aggregated spatially or temporally, has multiple within-day measurements (e.g., a diel study), or are from a data-dense spatial study
CO2mean	Mean or sole reported CO ₂ concentration in $\mu\text{mol L}^{-1}$ for the sampling event
CO2_SD	Standard deviation of the mean concentration

Table A3. Continued

CO2median	Median CO ₂ concentration in $\mu\text{mol L}^{-1}$
N2Omin	Minimum measured N ₂ O concentration in $\mu\text{mol L}^{-1}$ if data are aggregated spatially or temporally, has multiple within-day measurements (e.g., a diel study), or are from a data-dense spatial study
N2Omax	Maximum measured N ₂ O concentration in $\mu\text{mol L}^{-1}$ if data are aggregated spatially or temporally, has multiple within-day measurements (e.g., a diel study), or are from a data-dense spatial study
N2Omean	Mean or sole reported N ₂ O concentration in $\mu\text{mol L}^{-1}$ for the concentration for the sampling event
N2O_SD	Standard deviation of the mean N ₂ O concentration
N2Omedian	Median N ₂ O concentration in $\mu\text{mol L}^{-1}$
WaterTemp_degC	Water temperature in degrees C measured concurrently with CH ₄
WaterTemp_degC _estimated	Estimated water temperature in degrees C. This field was populated only for cases in which temperature was needed for gas unit conversion. Most estimates were based on temperatures from adjacent sites, averaging temperatures from prior and proceeding sample dates, or from an adjacent day of the year but from another year.
Cond_uScm	Specific conductance in $\mu\text{S cm}^{-1}$
pH	pH
DO_mgL	Dissolved oxygen in mg L^{-1}
DO_percentsat	Percent saturation of dissolved oxygen
Q	Discharge in $\text{m}^3 \text{s}^{-1}$ measured at the time of sample collection
NO3	NO ₃ or NO ₂ +NO ₃ concentration in $\mu\text{mol L}^{-1}$ measured concurrently with CH ₄
NH4	NH ₄ concentration in $\mu\text{mol L}^{-1}$ measured concurrently with CH ₄
TN	TN or TDN concentration in $\mu\text{mol L}^{-1}$ measured concurrently with CH ₄
SRP	SRP or PO ₄ concentration in $\mu\text{mol L}^{-1}$ measured concurrently with CH ₄
TP	TP or TDP concentration in $\mu\text{mol L}^{-1}$ measured concurrently with CH ₄
DOC	DOC or TOC concentration in $\mu\text{mol L}^{-1}$ measured concurrently with CH ₄
Comments	Any additional relevant information regarding data
new_CH4_unit	Current common units for all CH ₄ concentrations

Table A3. Continued

new_CO2_unit	Current common units for all CO ₂ concentrations
new_N2O_unit	Current common units for all N ₂ O concentrations
new_NO3_unit	Current common units for all NO ₃ or NO ₂ +NO ₃ concentrations
new_NH4_unit	Current common units for all NH ₄ concentrations
new_TN_unit	Current common units for all TN or TDN concentrations
new_SRP_unit	Current common units for all SRP or PO ₄ concentrations
new_TP_unit	Current common units for all TP or TDP concentrations
new_DOC_unit	Current common units for all DOC or TOC concentrations
new_Q_unit	Current common units for all discharge measurements
orig_CH4_unit	Original units for CH ₄ concentration
orig_CO2_unit	Original units for CO ₂ concentration
orig_N2O_unit	Original units for N ₂ O concentration
orig_NO3_unit	Original units for NO ₃ or NO ₂ +NO ₃ concentration
orig_NH4_unit	Original units for NH ₄ concentration
orig_TN_unit	Original units for TN concentration
orig_SRP_unit	Original units for SRP or PO ₄ concentration
orig_TP_unit	Original units for TP concentration
orig_DOC_unit	Original units for DOC concentration
orig_Q_unit	Original units of discharge

710 **Table A4. Column titles and definitions for the GRiMeDB Flux Table**

Column Title	Definition
Source_ID	Unique paper identifier from the Sources Table
Site_ID	Unique site identifier from the Sites Table
Site_Name	Unique site name from the Sites Table
Flux_Name	Unique name for the sampling event at the site; same as Conc_Name in the Concentration Table if both concentration and flux data for the same site-date combination are available
Date_start	First sampling date
Date_end	Last sampling date; this is the same date as the Date_start if data are not aggregated over time
Aggregated_Space	Yes or No; "Yes" if CH ₄ data entered are averages from >1 site
Aggregated_Time	Yes or No; "Yes: if CH ₄ data entered are averages from >1 date
Diffusive_CH4_Flux_Min	Minimum measured CH ₄ diffusive flux in mm m ⁻² d ⁻¹ if data are aggregated or are from diel or data-dense spatial studies
Diffusive_CH4_Flux_Max	Maximum measured CH ₄ diffusive flux in mm m ⁻² d ⁻¹ if data are aggregated or are from diel or data-dense spatial studies
Diffusive_CH4_Flux_Mean	Mean or sole reported CH ₄ diffusive flux in mm m ⁻² d ⁻¹ for the sampling event
Diffusive_CH4_Flux_SD	Standard deviation of the mean CH ₄ diffusive flux
Diffusive_CH4_Flux_Median	Median CH ₄ diffusive flux in mm m ⁻² d ⁻¹
SampleCount_Diffusive	Number of samples or observations corresponding to the mean or median diffusive CH ₄ flux
Diff_Method	Methodological category used to measure diffusive gas flux. Categories (with brief explanations in italics) are: <ul style="list-style-type: none"> chamber (unspecified)- unspecified response <i>use of an unspecified type of chamber (suspended, tethered, or free-floating) and pattern of change gas concentration over time during flux measurements is also not specified</i> chamber (unspecified)- linear response <i>unspecified type of chamber with a linear increase in chamber gas concentration over time or use of a linear model to calculate flux</i> suspended/tethered chamber-unspecified response <i>chamber is restrained to maintain its position and not float downstream during flux measurement</i> suspended/tethered chamber- linear response floating chamber- unspecified response <i>chamber is unrestrained and is able to float downstream during flux measurement</i> floating chamber- linear response conc+k <i>diffusive flux calculated using the equation: flux = k(C_w-C_{eq}), where k = gas exchange coefficient C_w = CH₄ concentration measured in water</i>

Table A4. Continued.

	<i>C_{eq} = CH₄ concentration in water in equilibrium with the atmosphere</i>
	other
	<i>methods other than those described above</i>
Eb_CH4_Flux_Min	Minimum measured CH ₄ ebullitive flux in mm m ⁻² d ⁻¹ if data are aggregated or are from diel or data-dense spatial studies
Eb_CH4_Flux_Max	Maximum measured CH ₄ ebullitive flux in mm m ⁻² d ⁻¹ if data are aggregated or are from diel or data-dense spatial studies
Eb_CH4_Flux_Mean	Mean or sole reported CH ₄ ebullitive flux in mm m ⁻² d ⁻¹ for the sampling event
Eb_CH4_Flux_SD	Standard deviation of the mean CH ₄ ebullitive flux
Eb_CH4_Flux_Median	Median CH ₄ ebullition flux in mm m ⁻² d ⁻¹
SampleCount_Eb	Number of samples or observations corresponding to the mean or median ebullitive CH ₄ flux
	echosounder + bubble analysis
	<i>gas bubble volume determined using echosounder and combined with CH₄ content of recently collected bubbles</i>
	departure from linear increase during measurement
	<i>non-linear change in gas concentrations during chamber-based flux measurements taken as evidence of ebullition; various approaches used to quantify ebullition from these departures</i>
	other
	<i>methods other than those described above</i>
Total_CH4_Flux_Min	Minimum measured total CH ₄ flux in mm m ⁻² d ⁻¹
Total_CH4_Flux_Max	Maximum measured total CH ₄ flux in mm m ⁻² d ⁻¹
Total_CH4_Flux_Mean	Mean or sole reported total CH ₄ flux for the sampling event in mm m ⁻² d ⁻¹
Total_CH4_Flux_SD	Standard deviation of the mean total CH ₄ flux
Total_CH4_Flux_Median	Median measured total CH ₄ flux in mm m ⁻² d ⁻¹
Total_Method	Methodological category used to measure total CH ₄ flux. Categories (with brief explanations in italics) are:
	conc+k and ebullition
	<i>total flux calculated as the sum of separate measurements of diffusion determined by the conc+k method plus ebullition determined from the bubble trap or echosounder approach combined with bubble CH₄ analysis</i>
	floating chamber
	<i>free-floating chamber is assumed to capture diffusive flux and ebullitive flux (if present)</i>
	suspended/tethered chamber
	<i>suspended or tethered chamber is assumed to capture diffusive flux and ebullitive flux (if present)</i>
	chamber and ebullition
	<i>total flux calculated as the sum of separate measurements of diffusion determined using a floating or suspended/tethered chamber plus ebullition determined from the bubble trap or echosounder approach combined with bubble CH₄ analysis</i>

Table A4. Continued

	<p>mass balance <i>total flux represents the difference between all measured inputs to a reach (e.g., dissolved CH₄ from upstream flow, groundwater discharge, and methanogenesis) minus all outputs other than efflux to the atmosphere (e.g., downstream export, methane oxidation)</i></p> <p>other <i>methods other than those described above</i></p>
CO2_Flux_Min	Minimum measured CO ₂ flux in mm m ⁻² d ⁻¹ if data are aggregated or are from diel or data-dense spatial studies
CO2_Flux_Max	Maximum measured CO ₂ flux in mm m ⁻² d ⁻¹ if data are aggregated or are from diel or data-dense spatial studies
CO2_Flux_Mean	Mean or sole reported CO ₂ diffusive flux in mm m ⁻² d ⁻¹ for the sampling event
CO2_Flux_SD	Standard deviation of the mean CO ₂ flux
CO2_Flux_Median	Median CO ₂ flux in mm m ⁻² d ⁻¹
N2O_Flux_Min	Minimum measured N ₂ O flux in mm m ⁻² d ⁻¹ if data are aggregated or are from diel or data-dense spatial studies
N2O_Flux_Max	Maximum measured N ₂ O flux in mm m ⁻² d ⁻¹ if data are aggregated or are from diel or data-dense spatial studies
N2O_Flux_Mean	Mean or sole reported N ₂ O diffusive flux in mm m ⁻² d ⁻¹ for the sampling event
N2O_Flux_Stddev	Standard deviation of the mean N ₂ O flux
N2O_Flux_Median	Median N ₂ O flux in mm m ⁻² d ⁻¹
k_Method	Methodological category used for estimating the gas exchange coefficient, <i>k</i> , Categories (with brief explanations in italics) are:
	<p>physical model <i>k calculated using equations based on physical variables such as channel slope, water velocity, etc.</i></p> <p>chamber + conc <i>k determined by chamber-based measurements of flux, dissolved gas concentration, and re-arrangement of the flux equation</i> $\text{flux} = k(C_w - C_{eq})$<i>to solve for k. Typically, these measurements are made for CO₂, and then k_{CO2} is converted to k_{CH4}</i></p> <p>tracer addition <i>paired conservative and gas tracer additions used to calculate k from concentration declines along a stream reach</i></p> <p>assigned k value <i>use of k values from other dates or sites in the same study or k values considered to be characteristic of the site</i></p> <p>other <i>methods other than those described above</i></p> <p>unknown <i>method to determine k is not described</i></p>
k_ref	k method citation reported in the data source
Comments	Any additional relevant information regarding data entered in this row
new_Diffusive_Flux_unit	Current common units for all diffusive CH ₄ flux data
new_Eb_CH4_Flux_unit	Current common units for all ebullitive CH ₄ flux data

Table A4. Continued.

new_Total_Flux_unit	Current common units for all total CH ₄ flux data
new_CO2_Flux_unit	Current common units for all CO ₂ flux data
new_N2O_Flux_unit	Current common units for all N ₂ O flux data
orig_Diffusive_Flux_unit	Original units for diffusive CH ₄ flux
orig_Eb_CH4_Flux_unit	Original units for ebullitive CH ₄ flux used
orig_Total_Flux_unit	Original units for total CH ₄ flux
orig_CO2_Flux_unit	Original units for CO ₂ flux
orig_N2O_Flux_unit	Original units for N ₂ O flux

Appendix B. Citations for data sources in GRiMeDB, including citations for unpublished datasets. Dates for unpublished dataset correspond to the year the data were provided by data authors.

- 715 Abbott, B. and Jones, J.: Soil respiration, water chemistry, and soil gas data for thermokarst features and undisturbed tundra on the North Slope of Alaska, Arctic Data Center, <https://doi.org/10.18739/A23T9D71C>, 2013.
- Abbott, B. W., Jones, J. B., Godsey, S. E., Larouche, J. R., and Bowden, W. B.: Patterns and persistence of hydrologic carbon and nutrient export from collapsing upland permafrost, *Biogeosciences*, 12, 3725–3740, <https://doi.org/10.5194/bg-12-3725-2015>, 2015.
- 720 Abril, G., Guérin, F., Richard, S., Delmas, R., Galy-Lacaux, C., Gosse, P., Tremblay, A., Varfalvy, L., Dos Santos, M. A., and Matvienko, B.: Carbon dioxide and methane emissions and the carbon budget of a 10-year old tropical reservoir (Petit Saut, French Guiana), *Global Biogeochem. Cycles*, 19, GB4007, <https://doi.org/10.1029/2005GB002457>, 2005.
- Adams, D. D. and Simiyu, G. M.: Greenhouse gas (methane and carbon dioxide) emissions from a tropical river in Kenya: the importance of anthropogenic factors on natural gas flux rates, *SIL Proceedings 1922–2010*, 30, 887-889, <https://doi.org/10.1080/03680770.2009.11902264>, 2009.
- 725 Aho, K. S. and Raymond, P. A.: Differential response of greenhouse gas evasion to storms in forested and wetland streams, *J. Geophys. Res. Biogeosci.*, 124, 649–662, <https://doi.org/10.1029/2018JG004750>, 2019.
- Aho, K., Cawley, K., DelVecchia, A., Stanley, E., and Raymond, P.: Dissolved greenhouse gas concentrations derived from the NEON dissolved gases in surface water data product (DP1.20097.001), Environmental Data Initiative, <https://doi.org/10.6073/pasta/47d7cb6d374b6662cce98e42122169f8>, 2021a
- 730 Aho, K. S., Fair, J. H., Hosen, J. D., Kyzivat, E. D., Logozzo, L. A., Rocher-Ros, G., Weber, L. C., Yoon, B., and Raymond, P. A.: Distinct concentration-discharge dynamics in temperate streams and rivers: CO₂ exhibits chemostasis while CH₄ exhibits source limitation due to temperature control, *Limnol Oceanogr*, 66, 3656–3668, <https://doi.org/10.1002/lno.11906>, 2021b.
- Aho, K., Fair, J., Hosen, J., Kyzivat, E., Logozzo, L., Weber, L., Yoon, B., Zarnetske, J., and Raymond, P.: Dissolved N₂O measurements from the Connecticut River Watershed, Environmental Data Initiative, <https://doi.org/10.6073/pasta/3494ca49fc3283eea5e4fc2f8a24ce3b>, 2021c.
- 735 Aho, K., Hoyle, J., Hosen, J., Kyzivat, E., Logozzo, L., Rocher-Ros, G., Weber, L., Bryan, Y., and Raymond, P.: Dissolved CO₂ and CH₄ concentrations in the Connecticut River Watershed, Environmental Data Initiative, <https://doi.org/10.6073/pasta/af4daec813775b7f426a1db574cbebc7>, 2021d.

- 740 Alshboul, Z., Encinas-Fernández, J., Hofmann, H., and Lorke, A.: Export of dissolved methane and carbon dioxide with effluents from municipal wastewater treatment plants, *Environ. Sci. Technol.*, 50, 5555–5563, <https://doi.org/10.1021/acs.est.5b04923>, 2016.
- Andrews, L. F., Wadnerkar, P. D., White, S. A., Chen, X., Correa, R. E., Jeffrey, L. C., and Santos, I. R.: Hydrological, geochemical and land use drivers of greenhouse gas dynamics in eleven sub-tropical streams, *Aquat. Sci.*, 83, 40, <https://doi.org/10.1007/s00027-021-00791-x>, 2021.
- 745 Anthony, S. E., Prah, F. G., and Peterson, T. D.: Methane dynamics in the Willamette River, Oregon, *Limnol. Oceanogr.*, 57, 1517–1530, <https://doi.org/10.4319/lo.2012.57.5.1517>, 2012.
- Antweiler, R. C., Smith, R. L., Voytek, M. A., Bohlke, J. K., and Dupré, D. H.: Water quality data from two agricultural drainage basins in Northwestern Indiana and northeastern Illinois: III. Biweekly data, 2000-2002, U.S. Geological Survey, <https://pubs.usgs.gov/of/2005/1197/>, 2005a.
- 750 Antweiler, R. C., Smith, R. L., Voytek, M. A., Böhlke, J.-K., and Richards, K. D.: Water quality data from two agricultural drainage basins in northwestern Indiana and northeastern Illinois: I. Lagrangian and synoptic data, 1999-2002, U.S. Geological Survey, <http://pubs.water.usgs.gov/ofr20041317/>, 2005b.
- Arp, C., Kane, D., Hinzman, L., and Stuefer, S.: Hydrographic data, Imnavait Creek Watershed, Alaska, 1985-2017, Arctic Data Center, <https://doi.org/10.18739/A2K649S9D>, 2017.
- 755 Atkins, M. L., Santos, I. R., and Maher, D. T.: Seasonal exports and drivers of dissolved inorganic and organic carbon, carbon dioxide, methane and $\delta^{13}\text{C}$ signatures in a subtropical river network, *Sci. Total Environ.*, 575, 545–563, <https://doi.org/10.1016/j.scitotenv.2016.09.020>, 2017.
- Audet, J., Wallin, M. B., Kyllmar, K., Andersson, S., and Bishop, K.: Nitrous oxide emissions from streams in a Swedish agricultural catchment, *Agric. Ecosyst. Environ.*, 236, 295–303, <https://doi.org/10.1016/j.agee.2016.12.012>, 2017.
- 760 Bagnoud, A., Pramateftaki, P., Bogard, M. J., Battin, T. J., and Peter, H.: Microbial ecology of methanotrophy in streams along a gradient of CH_4 availability, *Front. Microbiol.*, 11, 771, <https://doi.org/10.3389/fmicb.2020.00771>, 2020.
- Baker, M. A., Dahm, C. N., Valett, H. M., Morrice, J. A., Henry, K. S., Campana, M. E., and Wroblecky, G. J.: Spatial and temporal variation in methane distribution at the ground water/surface water interface in headwater catchments, in: 2nd International Conference on Ground Water Ecology, Atlanta, GA, 29-37, <https://www.osti.gov/biblio/37567>, 1994.
- 765

- Ballester, M. V. R. and Santos, J. E. dos: Biogenic gases in tropical floodplain river, *Braz. Arch. Biol. Technol.*, 44, 141–147, <https://doi.org/10.1590/S1516-89132001000200006>, 2001.
- 770 Bange, H. W., Sim, C. H., Bastian, D., Kallert, J., Kock, A., Mujahid, A., and Müller, M.: Nitrous oxide (N₂O) and methane (CH₄) in rivers and estuaries of northwestern Borneo, *Biogeosciences*, 16, 4321–4335, <https://doi.org/10.5194/bg-16-4321-2019>, 2019.
- Banks, E. W., Hatch, M., Smith, S., Underschultz, J., Lamontagne, S., Suckow, A., and Mallants, D.: Multi-tracer and hydrogeophysical investigation of the hydraulic connectivity between coal seam gas formations, shallow groundwater and stream network in a faulted sedimentary basin, *J. Hydrol.*, 578, 124132, <https://doi.org/10.1016/j.jhydrol.2019.124132>, 2019.
- 775 Bansal, S.: PPR stream and river methane. U.S. Geological Survey Northern Prairie Wildlife Research Center, Jamestown, ND, USA, unpublished, 2020.
- Barbosa, P. M., Melack, J. M., Farjalla, V. F., Amaral, J. H. F., Scofield, V., and Forsberg, B. R.: Diffusive methane fluxes from Negro, Solimões and Madeira rivers and fringing lakes in the Amazon basin: Diffusive methane fluxes from rivers and fringing lakes, *Limnol. Oceanogr.*, 61, S221–S237, <https://doi.org/10.1002/lno.10358>, 2016.
- 780 Bastien, J. and Demarty, M.: Spatio-temporal variation of gross CO₂ and CH₄ diffusive emissions from Australian reservoirs and natural aquatic ecosystems, and estimation of net reservoir emissions, *Lakes Reservoirs Res. Manage.* 18, 115–127, <https://doi.org/10.1111/lre.12028>, 2013.
- Baulch, H. M., Dillon, P. J., Maranger, R., and Schiff, S. L.: Diffusive and ebullitive transport of methane and nitrous oxide from streams: Are bubble-mediated fluxes important?, *J. Geophys. Res.*, 116, G04028, <https://doi.org/10.1029/2011JG001656>, 2011.
- 785 Beaulieu, J. J.: Controls on greenhouse gas emissions from headwater streams, Ph.D. thesis, University of Notre Dame, Notre Dame, IN, USA, 205 pp., https://onsearch.library.nd.edu/permalink/f/1phik6l/ndu_aleph002323859, 2007.
- Beaulieu, J. J., Shuster, W. D., and Rebholz, J. A.: Controls on gas transfer velocities in a large river, *J. Geophys. Res.*, 117, G02007, <https://doi.org/10.1029/2011JG001794>, 2012.
- 790 Bednařík, A., Čáp, L., Maier, V., and Rulík, M.: Contribution of methane benthic and atmospheric fluxes of an experimental area (Sitka Stream), *CLEAN Soil Air Water*, 43, 1136–1142, <https://doi.org/10.1002/clen.201300982>, 2015.
- Bednařík, A., Blaser, M., Matoušů, A., Hekera, P., and Rulík, M.: Effect of weir impoundments on methane dynamics in a river, *Sci. Total Environ.*, 584–585, 164–174, <https://doi.org/10.1016/j.scitotenv.2017.01.163>, 2017.

- 795 Begum, M. S., Bogard, M. J., Butman, D. E., Chea, E., Kumar, S., Lu, X., Nayna, O. K., Ran, L., Richey, J. E., Tareq, S. M., Xuan, D. T., Yu, R., and Park, J.: Localized pollution impacts on greenhouse gas dynamics in three anthropogenically modified Asian river systems, *J Geophys. Res. Biogeosci.*, 126, e2020JG006124, <https://doi.org/10.1029/2020JG006124>, 2021a.
- Begum, M. S., Bogard, M. J., Butman, D. E., Chea, E., Kumar, S., Lu, X., Nayna, O., K, Ran, L., Richey, J E, Tareq, S. M, Xuan, D. T., Yu, R., and Park, J.: Spatiotemporal variation in $p\text{CO}_2$, CH_4 , N_2O , DOM, and ancillary water quality measured in the Ganges, Mekong, and Yellow River during 2016 to 2019, PANGAEA, <https://doi.org/10.1594/PANGAEA.926582>, 2021b.
- 800 Billett, M. F. and Harvey, F. H.: Measurements of CO_2 and CH_4 evasion from UK peatland headwater streams, *Biogeochemistry*, 114, 165–181, <https://doi.org/10.1007/s10533-012-9798-9>, 2013.
- Billett, M. F. and Moore, T. R.: Supersaturation and evasion of CO_2 and CH_4 in surface waters at Mer Bleue peatland, Canada, *Hydrol. Process.*, 22, 2044–2054, <https://doi.org/10.1002/hyp.6805>, 2008.
- Bilsley, N.: Processes affecting the spatial and temporal variability of methane in a temperate dammed river system, M.S. thesis, University of California San Diego, San Diego, CA, USA, 94 pp., https://search-library.ucsd.edu/permalink/01UCS_SDI/ld412s/alma991004859049706535, 2012.
- Blackburn, S. R. and Stanley, E. H.: Greenhouse gas fluxes and concentrations and associated habitat data in western Dane County, Wisconsin, USA, streams during the 2018 growing season ver. 2., Environmental Data Initiative, <https://doi.org/10.6073/pasta/eeba6193162bf1189a6b78452686c773>, 2019.
- 810 Blackburn, S. R. and Stanley, E. H.: Floods increase carbon dioxide and methane fluxes in agricultural streams, *Freshwater Biol.*, 66, 62–77, <https://doi.org/10.1111/fwb.13614>, 2021.
- Bodmer, P., Heinz, M., Pusch, M., Singer, G., and Premke, K.: Carbon dynamics and their link to dissolved organic matter quality across contrasting stream ecosystems, *Sci. Total Environ.*, 553, 574–586, <https://doi.org/10.1016/j.scitotenv.2016.02.095>, 2016.
- 815 Bolpagni, R., Laini, A., Mutti, T., Viaroli, P., and Bartoli, M.: Connectivity and habitat typology drive CO_2 and CH_4 fluxes across land-water interfaces in lowland rivers: C fluxes across lowland river interfaces, *Ecohydrology*, 12, e2036, <https://doi.org/10.1002/eco.2036>, 2019.

- 820 Borges, A. V., Abril, G., Darchambeau, F., Teodoru, C. R., Deborde, J., Vidal, L. O., Lambert, T., and Bouillon, S.: Divergent biophysical controls of aquatic CO₂ and CH₄ in the World's two largest rivers, *Sci. Rep.*, 5, 15614, <https://doi.org/10.1038/srep15614>, 2015a.
- Borges, A. V., Darchambeau, F., Teodoru, C. R., Marwick, T. R., Tamooch, F., Geeraert, N., Omengo, F. O., Guérin, F., Lambert, T., Morana, C., Okuku, E., and Bouillon, S.: Globally significant greenhouse-gas emissions from African inland waters, *Nat., Geosci.*, 8, 637–642, <https://doi.org/10.1038/ngeo2486>, 2015b.
- 825 Borges, A. V., Abril, G., and Bouillon, S.: Carbon dynamics and CO₂ and CH₄ outgassing in the Mekong Delta, *Biogeosciences*, 15, 1093–1114, <https://doi.org/10.5194/bg-15-1093-2018>, 2018a.
- Borges, A. V., Darchambeau, F., Lambert, T., Bouillon, S., Morana, C., Brouyère, S., Hakoun, V., Jurado, A., Tseng, H.-C., Descy, J.-P., and Roland, F. A. E.: Effects of agricultural land use on fluvial carbon dioxide, methane and nitrous oxide concentrations in a large European river, the Meuse (Belgium), *Sci. Total Environ.*, 610–611, 342–355, <https://doi.org/10.1016/j.scitotenv.2017.08.047>, 2018b.
- 830 Borges, A. V. and Bouillon, S.: data-base of CO₂, CH₄, N₂O and ancillary data in the Congo River (latest), Zenodo, <https://doi.org/10.5281/ZENODO.3413449>, 2019.
- Bouillon, S., Abril, G., Borges, A. V., Dehairs, F., Govers, G., Hughes, H. J., Merckx, R., Meysman, F. J. R., Nyunja, J., Osburn, C., and Middelburg, J. J.: Distribution, origin and cycling of carbon in the Tana River (Kenya): a dry season basin-scale survey from headwaters to the delta, *Biogeosciences*, 6, 2475–2493, <https://doi.org/10.5194/bg-6-2475-2009>, 2009.
- 835 Bouillon, S., Yambélé, A., Spencer, R. G. M., Gillikin, D. P., Hernes, P. J., Six, J., Merckx, R., and Borges, A. V.: Organic matter sources, fluxes and greenhouse gas exchange in the Oubangui River (Congo River basin), *Biogeosciences*, 9, 2045–2062, <https://doi.org/10.5194/bg-9-2045-2012>, 2012.
- 840 Bouillon, S., Yambélé, A., Gillikin, D. P., Teodoru, C., Darchambeau, F., Lambert, T., and Borges, A. V.: Contrasting biogeochemical characteristics of the Oubangui River and tributaries (Congo River basin), *Sci. Rep.*, 4, 5402, <https://doi.org/10.1038/srep05402>, 2015.
- Bowden, W. B.: Arctic LTER streams chemistry Toolik Field Station, Alaska 1978 to 2019., Environmental Data Initiative, <https://doi.org/10.6073/pasta/3faacd18b63b3bacc5a0dbd6f09660e1>, 2021.

- 845 Bower, S.: Effects of storms on nitrate removal and greenhouse gas emissions from fluvial wetland dominated surface water flow paths. University of New Hampshire., M.S. thesis, University of New Hampshire, Durham, NH, USA, <https://scholars.unh.edu/thesis/1376/>, 2020.
- Bresney, S. R., Moseman-Valtierra, S., and Snyder, N. P.: Observations of greenhouse gases and nitrate concentrations in a Maine river and fringing wetland, Northeast. Nat., 22, 120–143, <https://doi.org/10.1656/045.022.0125>, 2015.
- Bretz, K. A., Jackson, A. R., Rahman, S., Monroe, J. M., and Hotchkiss, E. R.: Integrating ecosystem patch contributions to stream corridor carbon dioxide and methane fluxes, J Geophys Res Biogeosci, 126, e2021JG006313, 850 <https://doi.org/10.1029/2021JG006313>, 2021a.
- Bretz, K. A., Jackson, A. R., Rahman, S., Monroe, J. M., and Hotchkiss, E. R.: Integrating EcosystemPatchContributionsStreamCO2CH4_Data, HydroShare, <https://doi.org/10.4211/hs.ab2b33f27b3b4a0ca9a8ce4b8936753f>, 2021b.
- Brown, R. S. and Hershey, A. E.: Potential effects of the invasive bivalve *Corbicula fluminea* on methane cycling processes 855 in an urban stream, Biogeochemistry, 144, 181–195, <https://doi.org/10.1007/s10533-019-00578-1>, 2019.
- Bugna, G. C., Chanton, J. P., Cable, J. E., Burnett, W. C., and Cable, P. H.: The importance of groundwater discharge to the methane budgets of nearshore and continental shelf waters of the northeastern Gulf of Mexico, Geochim. Cosmochim. Acta, 60, 4735–4746, [https://doi.org/10.1016/S0016-7037\(96\)00290-6](https://doi.org/10.1016/S0016-7037(96)00290-6), 1996.
- Buriánková, I., Brablcová, L., Mach, V., Hýblová, A., Badurová, P., Cupalová, J., Čáp, L., and Rulík, M.: Methanogens and 860 methanotrophs distribution in the hyporheic sediments of a small lowland stream, Fundam. Appl. Limnol., 181, 87–102, <https://doi.org/10.1127/1863-9135/2012/0283>, 2012.
- Burns, R., Wynn, P. M., Barker, P., McNamara, N., Oakley, S., Ostle, N., Stott, A. W., Tuffen, H., Zhou, Z., Tweed, F. S., Chesler, A., and Stuart, M.: Direct isotopic evidence of biogenic methane production and efflux from beneath a temperate glacier, Sci. Rep., 8, 17118, <https://doi.org/10.1038/s41598-018-35253-2>, 2018.
- 865 Burrows, R. M., van de Kamp, J., Bodrossy, L., Venarsky, M., Coates-Marnane, J., Rees, G., Jumppanen, P., and Kennard, M. J.: Methanotroph community structure and processes in an inland river affected by natural gas macro-seeps, FEMS Microbiol. Ecol., 97, fiab130, <https://doi.org/10.1093/femsec/fiab130>, 2021.
- Bussmann, I.: Distribution of methane in the Lena Delta and Buor-Khaya Bay, Russia, Biogeosciences, 10, 4641–4652, <https://doi.org/10.5194/bg-10-4641-2013>, 2013a.

- 870 Bussmann, I.: Methane concentration and isotopic composition ($\delta^{13}\text{C}$) in the waters of the Lena River and the Laptev Sea, in the years 2008, 2009 and 2010, PANGAEA, <https://doi.org/10.1594/PANGAEA.817302>, 2013b.
- Bussmann, I. and Fedorova, I. V.: Dissolved methane concentrations under ice cover in the Lena Delta area, PANGAEA, <https://doi.org/10.1594/PANGAEA.905776>, 2019.
- Bussmann, I., Fedorova, I. V., Juhls, B., Overduin, P. P., and Winkel, M.: Dissolved methane concentrations and oxidation
875 rates in the Lena Delta area, 2016-2018, 2 datasets, PANGAEA, <https://doi.org/10.1594/PANGAEA.920015>, 2020.
- Bussmann, I., Fedorova, I. V., Juhls, B., Overduin, P. P., and Winkel, M.: Methane dynamics in three different Siberian water bodies under winter and summer conditions, Biogeosciences, 18, 2047–2061, <https://doi.org/10.5194/bg-18-2047-2021>, 2021.
- Call, M., Sanders, C. J., Enrich-Prast, A., Sanders, L., Marotta, H., Santos, I. R., and Maher, D. T.: Metadata: Radon-traced pore-water as a potential source of CO_2 and CH_4 to receding black and clear water environments in the Amazon Basin: Black
880 and clear water environments in the Amazon Basin, figshare, <https://figshare.com/s/95945208c33e8800f02c> 2018a.
- Call, M., Sanders, C. J., Enrich-Prast, A., Sanders, L., Marotta, H., Santos, I. R., and Maher, D. T.: Radon-traced pore-water as a potential source of CO_2 and CH_4 to receding black and clear water environments in the Amazon Basin: Black and clear water environments in the Amazon Basin, Limnol. Oceanogr. Lett., 3, 375–383, <https://doi.org/10.1002/lol2.10089>, 2018b.
- Campeau, A.: Dataset for manuscript: Stable carbon isotopes reveal soil-stream DIC linkages in contrasting headwater
885 catchments (Version 1.0), Uppsala University Publications, <http://urn.kb.se/resolve?urn=urn:nbn:se:uu:diva-335488> 2018.
- Campeau, A., Lapierre, J.-F., Vachon, D., and del Giorgio, P. A.: Regional contribution of CO_2 and CH_4 fluxes from the fluvial network in a lowland boreal landscape of Québec: CO_2 and CH_4 emission from boreal rivers, Global Biogeochem. Cycles, 28, 57–69, <https://doi.org/10.1002/2013GB004685>, 2014.
- Campeau, A., Bishop, K. H., Billett, M. F., Garnett, M. H., Laudon, H., Leach, J. A., Nilsson, M. B., Öquist, M. G., and Wallin,
890 M. B.: Aquatic export of young dissolved and gaseous carbon from a pristine boreal fen: Implications for peat carbon stock stability, Global Change Biol, 23, 5523–5536, <https://doi.org/10.1111/gcb.13815>, 2017.
- Campeau, A., Bishop, K., Nilsson, M. B., Klemedtsson, L., Laudon, H., Leith, F. I., Öquist, M., and Wallin, M. B.: Stable carbon isotopes reveal soil-stream DIC linkages in contrasting headwater catchments, J. Geophys. Res. Biogeosci., 123, 149–167, <https://doi.org/10.1002/2017JG004083>, 2018.

- 895 Carey, C. C., Woelmer, W. M., Maze, J. T., and Hounshell, A. G.: Manually-collected discharge data for multiple inflow tributaries entering Falling Creek Reservoir and Beaverdam Reservoir, Vinton, Virginia, USA in 2019, Environmental Data Initiative, <https://doi.org/10.6073/pasta/4d8e7b7bedbc6507b307ba2d5f2cf9a2>, 2019.
- Carey, C. C., Hounshell, A. G., Lofton, M. E., Birgand, F., Bookout, B. J., Corrigan, R. S., Gerling, A. B., McClure, R. P., and Woelmer, W. M.: Discharge time series for the primary inflow tributary entering Falling Creek Reservoir, Vinton, Virginia, USA 2013-2021, Environmental Data Initiative, <https://doi.org/10.6073/pasta/8d22a432aac5560b0f45aa1b21ae4746>, 2021a.
- 900 USA 2013-2021, Environmental Data Initiative, <https://doi.org/10.6073/pasta/8d22a432aac5560b0f45aa1b21ae4746>, 2021a.
- Carey, C. C., Wander, H. L., Woelmer, W. M., Lofton, M. E., Breef-Pilz, A., Doubek, J. P., Gerling, A. B., Hounshell, A. G., McClure, R. P., and Niederlehner, B. R.: Water chemistry time series for Beaverdam Reservoir, Carvins Cove Reservoir, Falling Creek Reservoir, Gatewood Reservoir, and Spring Hollow Reservoir in southwestern Virginia, USA 2013-2020, Environmental Data Initiative, <https://doi.org/10.6073/pasta/8d83ef7ec202eca9192e3da6dd34a4e0>, 2021b.
- 905 Carey, C. C., Hounshell, A. G., McClure, R. P., Gerling, A. B., Lewis, A. S. L., and Niederlehner, B. R.: Time series of dissolved methane and carbon dioxide concentrations for Falling Creek Reservoir and Beaverdam Reservoir in southwestern Virginia, USA during 2015-2021, Environmental Data Initiative, <https://doi.org/10.6073/pasta/2fb836492aace4c13b7962f2718be8e5>, 2022.
- Carpenter, C., Wickland, K. P., Dornblaser, M. M., Clow, D. W., Koch, J. C., and Dantoin, E. D.: Wetland stream water quality data for West Twin Creek, AK, Allequash Creek, WI, and Big Thompson River, CO, 2010-2020: U.S. Geological Survey data release, <https://doi.org/10.5066/p9wys23u>, 2021.
- 910 data for West Twin Creek, AK, Allequash Creek, WI, and Big Thompson River, CO, 2010-2020: U.S. Geological Survey data release, <https://doi.org/10.5066/p9wys23u>, 2021.
- Castro-Morales, K., Canning, A., Körtzinger, A., Göckede, M., Küsel, K., Overholt, W.A., Wichard, T., Redlich, S., Arzberger, S., Kolle, O., and Zimov, N.: Data published in manuscript “Effects of reversal of water flow in an Arctic floodplain river on fluvial emissions of CO₂ and CH₄” by Castro-Morales et al., Zenodo, <https://doi.org/10.5281/zenodo.5758728>, 2021.
- 915 Castro-Morales, K., Canning, A., Körtzinger, A., Göckede, M., Küsel, K., Overholt, W. A., Wichard, T., Redlich, S., Arzberger, S., Kolle, O., and Zimov, N.: Effects of reversal of water flow in an Arctic floodplain river on fluvial emissions of CO₂ and CH₄, J. Geophys. Res. Biogeosci., 127, e2021JG006485, <https://doi.org/10.1029/2021jg006485>, 2022.
- Chamberlain, S. D., Boughton, E. H., and Sparks, J. P.: Underlying ecosystem emissions exceed cattle-emitted methane from subtropical lowland pastures, Ecosystems, 18, 933–945, <https://doi.org/10.1007/s10021-015-9873-x>, 2015.
- 920 Chan, C.-N., Shi, H., Liu, B., and Ran, L.: CO₂ and CH₄ emissions from an arid fluvial network on the Chinese Loess Plateau, Water, 13, 1614, <https://doi.org/10.3390/w13121614>, 2021.

- Chen, C.-T. A., Wang, S.-L., Lu, X.-X., Zhang, S.-R., Lui, H.-K., Tseng, H.-C., Wang, B.-J., and Huang, H.-I.: Hydrogeochemistry and greenhouse gases of the Pearl River, its estuary and beyond, *Quat. Int.*, 186, 79–90, <https://doi.org/10.1016/j.quaint.2007.08.024>, 2008.
- 925 Chen, S., Wang, D., Ding, Y., Yu, Z., Liu, L., Li, Y., Yang, D., Gao, Y., Tian, H., Cai, R., and Chen, Z.: Ebullition controls on CH₄ emissions in an urban, eutrophic river: A potential time-scale bias in determining the aquatic CH₄ flux, *Environ. Sci. Technol.*, 55, 7287–7298, <https://doi.org/10.1021/acs.est.1c00114>, 2021.
- Clarizia, P. E.: Seasonal methane and carbon dioxide emissions along a temperate fluvial wetland dominated river continuum, M.S. thesis, University of New Hampshire, Durham, NH, USA, 40 pp., <https://scholars.unh.edu/thesis/1322/>, 2019.
- 930 Clayer, F., Thrane, J. -E., Brandt, U., Dörsch, P., and Wit, H. A.: Boreal headwater catchment as hot spot of carbon processing from headwater to fjord, *J. Geophys. Res. Biogeosci.*, 126, e2021JG006359, <https://doi.org/10.1029/2021JG006359>, 2021a.
- Clayer, F., Thrane, J. -E., Brandt, U., Dörsch, P., and de Wit, H.: Dataset for “Boreal headwater catchment as hot spot of carbon processing from headwater to fjord” Clayer et al., HydroShare, <https://doi.org/10.4211/hs.b143432cfe72462fac90376631c9a2b3>, 2021b.
- 935 Clilverd, H. M., Jones, J. B., and Kielland, K.: Nitrogen retention in the hyporheic zone of a glacial river in interior Alaska, *Biogeochemistry*, 88, 31–46, <https://doi.org/10.1007/s10533-008-9192-9>, 2008.
- Comer-Warner, S., Krause, S., Gooddy, D. C., Ullah, S., and Wexler, S. K.: Seasonal streambed carbon and nitrogen cycling (including greenhouse gases) in an agriculturally-impacted stream. Measured at Wood Brook UK, 2016-2017, NERC Environmental Information Data Centre, <https://doi.org/10.5285/00601260-285e-4ffa-b381-340b51a7ec50>, 2018.
- 940 Comer-Warner, S. A., Gooddy, D. C., Ullah, S., Glover, L., Percival, A., Kettridge, N., and Krause, S.: Seasonal variability of sediment controls of carbon cycling in an agricultural stream, *Sci. Total Environ.*, 688, 732–741, <https://doi.org/10.1016/j.scitotenv.2019.06.317>, 2019.
- Cotovicz, L. C., Ribeiro, R. P., Régis, C. R., Bernardes, M., Sobrinho, R., Vidal, L. O., Tremmel, D., Knoppers, B. A., and Abril, G.: Greenhouse gas emissions (CO₂ and CH₄) and inorganic carbon behavior in an urban highly polluted tropical coastal lagoon (SE, Brazil), *Environ. Sci. Pollut. Res.*, 28, 38173–38192, <https://doi.org/10.1007/s11356-021-13362-2>, 2021.
- 945 Cowley, K., Looman, A., Maher, D. T., and Fryirs, K.: Geomorphic controls on fluvial carbon exports and emissions from upland swamps in eastern Australia, *Sci. Total Environ.*, 618, 765–776, <https://doi.org/10.1016/j.scitotenv.2017.08.133>, 2018.

- Crawford, J. T.: Carbon gas emissions from headwater streams. University of Wisconsin-Madison, Madison, WI, USA, unpublished, 2014.
- 950 Crawford, J. T. and Stanley, E. H.: Controls on methane concentrations and fluxes in streams draining human-dominated landscapes, *Ecol. Appl.*, 26, 1581–1591, <https://doi.org/10.1890/15-1330>, 2016.
- Crawford, J. and Stanley, E.: Greenhouse gas emissions from streams at North Temperate Lakes LTER 2012, Environmental Data Initiative, <https://doi.org/10.6073/pasta/000c46ce5c2ded2cefbddb596fca3ce2>, 2020.
- Crawford, J. T., Striegl, R. G., Wickland, K. P., Dornblaser, M. M., and Stanley, E. H.: Emissions of carbon dioxide and methane from a headwater stream network of interior Alaska, *J. Geophys. Res. Biogeosci.*, 118, 482–494, <https://doi.org/10.1002/jgrg.20034>, 2013.
- 955 Crawford, J. T., Lottig, N. R., Stanley, E. H., Walker, J. F., Hanson, P. C., Finlay, J. C., and Striegl, R. G.: CO₂ and CH₄ emissions from streams in a lake-rich landscape: Patterns, controls, and regional significance, *Global Biogeochem. Cycles*, 28, 197–210, <https://doi.org/10.1002/2013GB004661>, 2014a.
- 960 Crawford, J. T., Stanley, E. H., Spawn, S. A., Finlay, J. C., Loken, L. C., and Striegl, R. G.: Ebullitive methane emissions from oxygenated wetland streams, *Global Change Biol.*, 20, 3408–3422, <https://doi.org/10.1111/gcb.12614>, 2014b.
- Crawford, J. T., Dornblaser, M. M., Stanley, E. H., Clow, D. W., and Striegl, R. G.: Source limitation of carbon gas emissions in high-elevation mountain streams and lakes, *J. Geophys. Res. Biogeosci.*, 120, 952–964, <https://doi.org/10.1002/2014JG002861>, 2015.
- 965 Crawford, J. T., Loken, L. C., Stanley, E. H., Stets, E. G., Dornblaser, M. M., and Striegl, R. G.: Basin scale controls on CO₂ and CH₄ emissions from the Upper Mississippi River: Mississippi River Greenhouse Gases, *Geophys. Res. Lett.*, 43, 1973–1979, <https://doi.org/10.1002/2015GL067599>, 2016.
- Crawford, J., Stanley, E., and Loken, L.: Ebullitive methane emissions from oxygenated wetland streams at North Temperate Lakes LTER 2013, Environmental Data Initiative, <https://doi.org/10.6073/pasta/7842765e6520b62af6fd4d95c838f177>, 2020.
- 970 Dahm, C. N., Carr, D. L., and Coleman, R. L.: Anaerobic carbon cycling in stream ecosystems, *SIL Proceedings*, 1922–2010, 24, 1600–1604, <https://doi.org/10.1080/03680770.1989.11899028>, 1991.
- Dawson, J. J. C., Billett, M. F., Hope, D., Palmer, S. M., and Deacon, C. M.: Sources and sinks of aquatic carbon in a peatland stream continuum, *Biogeochemistry*, 70, 71–92, <https://doi.org/10.1023/B:BIOG.0000049337.66150.f1>, 2004.

- de Angelis, M. A. and Lilley, M. D.: Methane in surface waters of Oregon estuaries and rivers, *Limnol. Oceanogr.*, 32, 716–722, <https://doi.org/10.4319/lo.1987.32.3.0716>, 1987.
- 975
- de Angelis, M. A. and Scranton, M. I.: Fate of methane in the Hudson River and Estuary, *Global Biogeochem. Cycles*, 7, 509–523, <https://doi.org/10.1029/93GB01636>, 1993.
- Dean, J. F., Billett, M. F., Baxter, R., Dinsmore, K. J., Lessels, J. S., Street, L. E., Subke, J.-A., Tetzlaff, D., Washbourne, I., and Wookey, P. A.: Biogeochemistry of “pristine” freshwater stream and lake systems in the western Canadian Arctic, *Biogeochemistry*, 130, 191–213, <https://doi.org/10.1007/s10533-016-0252-2>, 2016.
- 980
- Dean, J. F., Meisel, O. H., Martyn Rosco, M., Marchesini, L. B., Garnett, M. H., Lenderink, H., van Logtestijn, R., Borges, A. V., Bouillon, S., Lambert, T., Röckmann, T., Maximov, T., Petrov, R., Karsanaev, S., Aerts, R., van Huissteden, J., Vonk, J. E., and Dolman, A. J.: East Siberian Arctic inland waters emit mostly contemporary carbon, *Nat. Commun.*, 11, 1627, <https://doi.org/10.1038/s41467-020-15511-6>, 2020.
- 985
- Deirmendjian, L., Anschutz, P., Morel, C., Mollier, A., Augusto, L., Loustau, D., Cotovicz, L. C., Buquet, D., Lajaunie, K., Chaillou, G., Voltz, B., Charbonnier, C., Poirier, D., and Abril, G.: Importance of the vegetation-groundwater-stream continuum to understand transformation of biogenic carbon in aquatic systems – A case study based on a pine-maize comparison in a lowland sandy watershed (Landes de Gascogne, SW France), *Sci. Total Environ.*, 661, 613–629, <https://doi.org/10.1016/j.scitotenv.2019.01.152>, 2019.
- 990
- DelSontro, T., Beaulieu, J. J., and Downing, J. A.: Greenhouse gas emissions from lakes and impoundments: Upscaling in the face of global change, *Limnol. Oceanogr. Lett.* 3, 64–75, <https://doi.org/10.1002/lol2.10073>, 2018.
- DelSontro, T., Perez, K. K., Sollberger, S., and Wehrli, B.: Methane dynamics downstream of a temperate run-of-the-river reservoir: CH₄ dynamics below a run-of-the-river reservoir, *Limnol. Oceanogr.*, 61, S188–S203, <https://doi.org/10.1002/lno.10387>, 2016.
- 995
- Deng, O., Li, X., Deng, L., Zhang, S., Gao, X., Lan, T., Zhou, W., Tian, D., Xiao, Y., Yang, J., Ou, D., and Luo, L.: Emission of CO₂ and CH₄ from a multi-ditches system in rice cultivation region: Flux, temporal-spatial variation and effect factors, *J. Environ. Manage.*, 270, 110918, <https://doi.org/10.1016/j.jenvman.2020.110918>, 2020.
- Descloux, S., Chanudet, V., Serça, D., and Guérin, F.: Methane and nitrous oxide annual emissions from an old eutrophic temperate reservoir, *Sci. Total Environ.*, 598, 959–972, <https://doi.org/10.1016/j.scitotenv.2017.04.066>, 2017.

- 1000 Deshmukh, C., Guérin, F., Pighini, S., Vongkhamsao, A., Guédant, P., Rode, W., Godon, A., Chanudet, V., Descloux, S., and Serça, D.: Low methane (CH₄) emissions downstream of a monomictic subtropical hydroelectric reservoir (Nam Theun 2, Lao PDR), *Biogeosciences Discuss.*, 12, 11313–11347, <https://doi.org/10.5194/bgd-12-11313-2015>, 2015.
- Dinsmore, K. J., Billett, M. F., Skiba, U. M., Rees, R. M., Drewer, J., and Helfter, C.: Role of the aquatic pathway in the carbon and greenhouse gas budgets of a peatland catchment, *Global Change Biol.*, 16, 2750–2762, <https://doi.org/10.1111/j.1365-2486.2009.02119.x>, 2010.
- 1005 Dinsmore, K. J., Billett, M. F., Dyson, K. E., Harvey, F., Thomson, A. M., Piirainen, S., and Kortelainen, P.: Stream water hydrochemistry as an indicator of carbon flow paths in Finnish peatland catchments during a spring snowmelt event, *Sci. Total Environ.*, 409, 4858–4867, <https://doi.org/10.1016/j.scitotenv.2011.07.063>, 2011.
- Dinsmore, K. J., Billett, M. F., and Dyson, K. E.: Five year record of aquatic carbon and greenhouse gas concentrations from Auchencorth Moss, NERC Environmental Information Data Centre, <https://doi.org/10.5285/3f0820a7-a8c8-4dd7-a058-8db79ba9c7fe>, 2013a.
- 1010 Dinsmore, K. J., Billett, M. F., and Dyson, K. E.: Temperature and precipitation drive temporal variability in aquatic carbon and GHG concentrations and fluxes in a peatland catchment, *Global Change Biol.*, 19, 2133–2148, <https://doi.org/10.1111/gcb.12209>, 2013b.
- 1015 Dinsmore, K. J., Murphey, O., Leith, F., and Carfrae, J.: Aquatic carbon and greenhouse gas concentrations in the Auchencorth Moss catchment following drain blocking, NERC Environmental Information Data Centre, <https://doi.org/10.5285/88ffbf44-0ec0-41d6-9814-04bc3535cd84>, 2016.
- Dodd, A.: Flow regime influences on stream and riparian soil carbon dynamics in the Ozark Highlands and Boston Mountains of Arkansas, Ph.D., University of Arkansas, Fayetteville, AR, USA, 143 pp., <https://scholarworks.uark.edu/etd/2911/>, 2018.
- 1020 Dornblaser, M. M. and Halm, D. R. (Eds.): Water and sediment quality of the Yukon River and its tributaries, from Eagle to St. Marys, Alaska, 2002–2003, U.S. Geological Survey Open-File Report 2006-1228, Reston, VA, 202 pp., <https://doi.org/10.3133/ofr20061228>, 2006.
- Druschke, C. G.: Geomorphic and ecological effects of restoration and flood events on stream in the Driftless Area of Wisconsin, M.S. thesis, University of Wisconsin-Madison, Madison, WI, USA, 59 pp., <https://search.library.wisc.edu/catalog/9913456953802121>, 2021.
- 1025

- Druschke, C. G., Booth, E. G., and Stanley, E. H.: Stream restoration and flood impacts in the Kickapoo River Watershed, Wisconsin, 2019, Environmental Data Initiative, <https://doi.org/10.6073/pasta/4905369228d80920974f555a2dd12229>, 2022.
- Dyson, K. E., Billett, M. F., Dinsmore, K. J., Harvey, F., Thomson, A. M., Piirainen, S., and Kortelainen, P.: Release of aquatic carbon from two peatland catchments in E. Finland during the spring snowmelt period, *Biogeochemistry*, 103, 125–142, 1030 <https://doi.org/10.1007/s10533-010-9452-3>, 2011.
- Einarsdottir, K., Wallin, M. B., and Sobek, S.: High terrestrial carbon load via groundwater to a boreal lake dominated by surface water inflow: Carbon input to a lake via Groundwater, *J. Geophys. Res. Biogeosci.*, 122, 15–29, <https://doi.org/10.1002/2016JG003495>, 2017.
- Emmerton, C. A., St. Louis, V. L., Lehnherr, I., Humphreys, E. R., Rydz, E., and Kosolofski, H. R.: The net exchange of 1035 methane with high Arctic landscapes during the summer growing season, *Biogeosciences*, 11, 3095–3106, <https://doi.org/10.5194/bg-11-3095-2014>, 2014.
- Evans, C. D., Peacock, M., Green, S. M., Holden, J., Chapman, P. J., Lebron, I., Callaghan, N., Grayson, R., and Baird, A. J.: The impact of ditch blocking on fluvial carbon export from a UK blanket bog, *Hydrol. Process.*, 32, 2141–2154, <https://doi.org/10.1002/hyp.13158>, 2018.
- 1040 Federov, Y. A., Khoroshevskaya, V. O., and Tambieva, N. S.: Variations in methane concentrations in the water of the Don River and Taganrog Bay under the effect of natural factors, *Water Resour.*, 30, 89–93, <https://doi.org/10.1023/A:1022059903167>, 2003.
- Flanagan, L. B., Nikkel, D. J., Scherloski, L. M., Tkach, R. E., Smits, K. M., Selinger, L. B., and Rood, S. B.: Multiple 1045 processes contribute to methane emission in a riparian cottonwood forest ecosystem, *New Phytol.*, 229, 1970–1982, <https://doi.org/10.1111/nph.16977>, 2020.
- Flessa, H., Rodionov, A., Guggenberger, G., Fuchs, H., Magdon, P., Shibistova, O., Zrazhevskaya, G., Mikheyeva, N., Kasansky, O. A., and Blodau, C.: Landscape controls of CH₄ fluxes in a catchment of the forest tundra ecotone in northern Siberia, *Global Change Biol.*, 14, 2040–2056, <https://doi.org/10.1111/j.1365-2486.2008.01633.x>, 2008.
- Flury, S. and Ulseth, A. J.: Exploring the sources of unexpected high methane concentrations and fluxes from Alpine headwater 1050 streams, *Geophys. Res. Lett.*, 46, 6614–6625, <https://doi.org/10.1029/2019GL082428>, 2019.
- Foks, S. S., Stackpoole, S. M., Dornblaser, M. M., Whiddon, E. T., Breitmeyer, S. E., Campbell, D. A., Crawford, J. T., Metcalf, E. B., Stets, E. G., Uhle, B. A., Voss, B. M., Wickland, K. P., and Striegl, R. G.: Water quality, quantity, and gas

- fluxes of the Upper Mississippi River basin (WY 2012-2016), U.S. Geological Survey data release, <https://doi.org/10.5066/F7P849S4>, 2018.
- 1055 Foks, S. S., Dornblaser, M. M., Butman, D. E., Campbell, D. A., Koch, J. C., Li, Z., Whiddon, E. T., Wickland, K. P., Striegl, R., Bogard, M. J., Spencer, R. G. M., Textor, S. R., and Johnston, S. E.: Water quality and gas fluxes of Interior Alaska (2014-2018), U.S. Geological Survey data release, <https://doi.org/10.5066/P9C6BDBQ>, 2020.
- Ford, T. E. and Naiman, R. J.: Alteration of carbon cycling by beaver: methane evasion rates from boreal forest streams and rivers, *Can. J. Zool.*, 66, 529–533, <https://doi.org/10.1139/z88-076>, 1988.
- 1060 Galantini, L., Lapierre, J.-F., and Maranger, R.: How are greenhouse gases coupled across seasons in a large temperate river with differential land use?, *Ecosystems*, <https://doi.org/10.1007/s10021-021-00629-5>, 2021.
- Gareis, J. A. L. and Lesack, L. F. W.: Ice-out and freshet fluxes of CO₂ and CH₄ across the air–water interface of the channel network of a great Arctic delta, the Mackenzie, *Polar Res.*, 39, <https://doi.org/10.33265/polar.v39.3528>, 2020.
- Gar’kusha, D. N. and Fedorov, Y. A.: Methane in the water and bottom sediments of the mouth area of the Severnaya Dvina River during the winter time, *Oceanology*, 54, 160–169, <https://doi.org/10.1134/S000143701402009X>, 2014.
- 1065 Gar’kusha, D. N. and Fedorov, Y. A.: Distribution of methane concentration in coastal areas of the Gulf of Petrozavodsk, Lake Onega, *Water Resour*, 42, 331–339, <https://doi.org/10.1134/S0097807815030045>, 2015.
- Gar’kusha, D. N., Fedorov, Y. A., and Khromov, M. I.: Methane in the water and bottom sediments of the mouth area of the Severnaya Dvina River (White Sea), *Oceanology*, 50, 498–512, <https://doi.org/10.1134/S0001437010040065>, 2010a.
- 1070 Gar’kusha, D. N., Fedorov, Y., and Khromov, M. I.: Methane in waters and bottom sediments of the North (Severnaya) Dvina mouth area in 2004-2006, *PANGAEA*, <https://doi.pangaea.de/10.1594/PANGAEA.763838>, 2010b.
- Garnier, J., Vilain, G., Silvestre, M., Billen, G., Jehanno, S., Poirier, D., Martinez, A., Decuq, C., Cellier, P., and Abril, G.: Budget of methane emissions from soils, livestock and the river network at the regional scale of the Seine basin (France), *Biogeochemistry*, 116, 199–214, <https://doi.org/10.1007/s10533-013-9845-1>, 2013.
- 1075 Gatland, J. R., Santos, I. R., Maher, D. T., Duncan, T. M., and Erler, D. V.: Carbon dioxide and methane emissions from an artificially drained coastal wetland during a flood: Implications for wetland global warming potential, *J. Geophys. Res. Biogeosci.*, 119, 1698–1716, <https://doi.org/10.1002/2013JG002544>, 2014.

- 1080 Gatti, R. C., Callaghan, T. V., Rozhkova-Timina, I., Dudko, A., Lim, A., Vorobyev, S. N., Kirpotin, S. N., and Pokrovsky, O. S.: The role of Eurasian beaver (*Castor fiber*) in the storage, emission and deposition of carbon in lakes and rivers of the River Ob flood plain, western Siberia, *Sci. Total Environ.*, 644, 1371–1379, <https://doi.org/10.1016/j.scitotenv.2018.07.042>, 2018.
- Gómez-Gener, L., Obrador, B., von Schiller, D., Marcé, R., Casas-Ruiz, J. P., Proia, L., Acuña, V., Catalán, N., Muñoz, I., and Koschorreck, M.: Hot spots for carbon emissions from Mediterranean fluvial networks during summer drought, *Biogeochemistry*, 125, 409–426, <https://doi.org/10.1007/s10533-015-0139-7>, 2015.
- 1085 Gómez-Gener, L., Gubau, M., von Schiller, D., Marcé, R., and Obrador, B.: Effect of small water retention structures on diffusive CO₂ and CH₄ emissions along a highly impounded river, *Inland Waters*, 8, 449–460, <https://doi.org/10.1080/20442041.2018.1457846>, 2018.
- Gondwe, M. J. and Masamba, W. R. L.: Spatial and temporal dynamics of diffusive methane emissions in the Okavango Delta, northern Botswana, Africa, *Wetlands Ecol. Manage.*, 22, 63–78, <https://doi.org/10.1007/s11273-013-9323-5>, 2014.
- 1090 Gong, X. J., Wang, X. F., Yuan, X. Z., Liu, T. T., and Hou, C. L.: Effects of field town developments on the dissolved and diffusion fluxes of greenhouse gasses in Heishuitan River basin, Chongqing, *Acta Ecol. Sin.*, 39, 8425–8441, <https://www.ecologica.cn/html/2019/22/stxb201810112205.htm>, 2019.
- Gonzalez-Valencia, R. Sepulveda-Jauregui, A., Martinez-Cruz, K. Hoyos-Santillan, J. Dendooven, L. and Thalasso, L.: Methane emissions from Mexican freshwater bodies: correlations with water pollution, *Hydrobiologia*, 721, 9-22, <https://doi.org/10.1007/s10750-013-1632-4>, 2014.
- 1095 Gries, C., Hanson, P. C., O'Brien, M., Servilla, M., Vanderbilt, K. and Waide, R. The Environmental Data Initiative: Connecting the past to the future through data reuse. *Ecol. Evol.*, 13, e9592, <https://doi.org/10.1002/ece3.9592>, 2023.
- Grieve, P. L., Hynek, S. A., Heilweil, V., Sowers, T., Llewellyn, G., Yoxtheimer, D., Solomon, D. K., and Brantley, S. L.: Using environmental tracers and modelling to identify natural and gas well-induced emissions of methane into streams, *Appl. Geochem.*, 91, 107–121, <https://doi.org/10.1016/j.apgeochem.2017.12.022>, 2018.
- 1100 Gu, C., Waldron, S., and Bass, A. M.: Carbon dioxide, methane, and dissolved carbon dynamics in an urbanized river system, *Hydrol., Process.*, 35, <https://doi.org/10.1002/hyp.14360>, 2021.
- Guo, Y., Song, C., Wang, L., Tan, W., Wang, X., Cui, Q., and Wan, Z.: Concentrations, sources, and export of dissolved CH₄ and CO₂ in rivers of the permafrost wetlands, northeast China, *Ecol. Eng.*, 90, 491–497, <https://doi.org/10.1016/j.ecoleng.2015.10.004>, 2016.

- Hall, R. O. Jr. and Ulseth, A. J.: Gas exchange in streams and rivers. *WIREs Water*, 7, e1391, <https://doi.org/10.1002/wat2.1391> 2019
- Halm, D. R. and Dornblaser, M. M. (Eds.): Water and sediment quality in the Yukon River and its tributaries, between Atlin, British Columbia, Canada, and Eagle, Alaska, 2004, U.S. Geological Survey Open-File Report 2007-1197, Reston, VA, 120 pp., <https://doi.org/10.3133/ofr20071197>, 2007.
- 1110
- Hamilton, S. K., Sippel, S., and Melack, J. M.: Oxygen depletion and carbon dioxide and methane production in waters of the Pantanal wetland of Brazil, *Biogeochemistry*, 30, 115–141, <https://doi.org/10.1007/bf00002727>, 1995.
- Han, Y.: Greenhouse gases emission characteristics of rivers in Nanjing and the influencing factors, M.S. thesis, Nanjing University of Information Science and Technology, Nanjing, China, <https://kns.cnki.net/kcms/detail/detail.aspx?dbname=cmfd201401&filename=1013340872.nh>, 2013.
- 1115
- Han, Y., Zhang, G., and Zhao, Y.: Distribution and fluxes of methane in tropical rivers and lagoons of eastern Hainan, *J. Trop. Oceanogr.*, 31, 87–95, <http://www.jto.ac.cn/CN/10.11978/j.issn.1009-5470.2012.02.012>, 2012.
- Hao, X., Ruihong, Y., Zhuangzhuang, Z., Zhen, Q., Xixi, L., Tingxi, L., and Ruizhong, G.: Greenhouse gas emissions from the water–air interface of a grassland river: a case study of the Xilin River, *Sci. Rep.*, 11, 2659, <https://doi.org/10.1038/s41598-021-81658-x>, 2021.
- 1120
- Harms, T. K. and Godsey, S. E.: Collaborative Research: Climate-mediated coupling of hydrology and biogeochemistry in arctic hillslopes, Arctic Data Center, <https://doi.org/10.18739/a23r0ps5p>, 2018.
- Harms, T. K., Rocher-Ros, G., and Godsey, S. E.: Emission of greenhouse gases from water tracks draining Arctic hillslopes, *J. Geophys. Res. Biogeosci.*, 125, e2020JG005889, <https://doi.org/10.1029/2020JG005889>, 2020.
- 1125
- Harrison, J. A., Matson, P. A., and Fendorf, S. E.: Effects of a diel oxygen cycle on nitrogen transformations and greenhouse gas emissions in a eutrophied subtropical stream, *Aquat. Sci.*, 67, 308–315, <https://doi.org/10.1007/s00027-005-0776-3>, 2005.
- Hattenberger, D. M.: Comparing carbon dioxide and methane emissions from restored and unrestored sections of three North Carolina streams, M.S. thesis, North Carolina State University, Raleigh, N.C., USA, 103 pp., <http://www.lib.ncsu.edu/resolver/1840.20/35725>, 2018.
- 1130
- Hayakawa, A., Ikeda, S., Tsushima, R., Ishikawa, Y., and Hidaka, S.: Spatial and temporal variations in nutrients in water and riverbed sediments at the mouths of rivers that enter Lake Hachiro, a shallow eutrophic lake in Japan, *Catena*, 133, 486–494, <https://doi.org/10.1016/j.catena.2015.04.009>, 2015.

- He, B., He, J., Wang, J., Li, J., and Wang, F.: Characteristics of GHG flux from water-air interface along a reclaimed water intake area of the Chaobai River in Shunyi, Beijing, *Atmos. Environ.*, 172, 102–108, 1135 <https://doi.org/10.1016/j.atmosenv.2017.10.060>, 2018.
- Heilweil, V. M., Stolp, B. J., Kimball, B. A., Susong, D. D., Marston, T. M., and Gardner, P. M.: A stream-based methane monitoring approach for evaluating groundwater impacts associated with unconventional gas development, *Groundwater*, 51, 511–524, <https://doi.org/10.1111/gwat.12079>, 2013.
- Heilweil, V. M., Risser, D. W., Conger, R. W., Grieve, P. L., and Hyneck, S. A.: Estimation of methane concentrations and 1140 loads in groundwater discharge to Sugar Run, Lycoming County, Pennsylvania, U.S. Geological Survey Open File Report 2014–1126., Reston, VA, 31 p., <https://doi.org/10.3133/ofr20141126>, 2014.
- Heilweil, V. M., Solomon, D. K., Darrah, T. H., Gilmore, T. E., and Genereux, D. P.: Gas-tracer experiment for evaluating the fate of methane in a coastal plain stream: Degassing versus in-stream oxidation, *Environ. Sci. Technol.*, 50, 10504–10511, <https://doi.org/10.1021/acs.est.6b02224>, 2016.
- 1145 Hendriks, D. M. D., van Huissteden, J., and Dolman, A. J.: Multi-technique assessment of spatial and temporal variability of methane fluxes in a peat meadow, *Agric. For. Meteorol.*, 150, 757–774, <https://doi.org/10.1016/j.agrformet.2009.06.017>, 2010.
- Heppell, C. M. and Binley, A.: Hampshire Avon: Daily discharge, stage and water chemistry data from four tributaries (Sem, Nadder, West Avon, Ebbles), NERC Environmental Information Data Centre, <https://doi.org/10.5285/0dd10858-7b96-41f1-8db5-e7b4c4168af5>, 2016a.
- 1150 Heppell, C. M. and Binley, A.: Hampshire Avon: Vertical head gradient, saturated hydraulic conductivity and pore water chemistry data from six river reaches, NERC Environmental Information Data Centre, <https://doi.org/10.5285/d82a04ce-f04d-40b4-9750-1a2bf7dc29a3>, 2016b.
- Herman-Mercer, N. M.: Water-quality data from the Yukon River Basin in Alaska and Canada, U.S. Geological Survey data release, <https://doi.org/10.5066/f77d2s7b>, 2016.
- 1155 Herreid, A. M., Wymore, A. S., Varner, R. K., Potter, J. D., and McDowell, W. H.: Divergent controls on stream greenhouse gas concentrations across a land-use gradient, *Ecosystems*, <https://doi.org/10.1007/s10021-020-00584-7>, 2020a.
- Herreid, A. M., Wymore, A. S., Varner, R. K., Potter, J. D., and McDowell, W. H.: Divergent controls on stream greenhouse gas concentrations across a land use gradient, *HydroShare*, <https://doi.org/10.4211/hs.2679ce1a6d514b30a54459893557dfe7>, 2020b.

- 1160 Herrero Ortega, S., Romero González-Quijano, C., Casper, P., Singer, G. A., and Gessner, M. O.: Methane emissions from contrasting urban freshwaters: Rates, drivers, and a whole-city footprint, *Global Change Biol.*, 25, 4234–4243, <https://doi.org/10.1111/gcb.14799>, 2019.
- Hlaváčová, E., Rulík, M., and Čáp, L.: Anaerobic microbial metabolism in hyporheic sediment of a gravel bar in a small lowland stream, *River Res. Appl.*, 21, 1003–1011, <https://doi.org/10.1002/rra.866>, 2005.
- 1165 Hlaváčová, E., Rulík, M., Čáp, L., and Mach, V.: Greenhouse gas (CO₂, CH₄, N₂O) emissions to the atmosphere from a small lowland stream in Czech Republic, *Arch. Hydrobiol.*, 165, 339–353, <https://doi.org/10.1127/0003-9136/2006/0165-0339>, 2006.
- Ho, L., Jerves-Cobo, R., Barthel, M., Six, J., Bode, S., Boeckx, P., and Goethals, P.: Dissolved gas concentrations in Cuenca river systems (Ecuador), Environmental Data Initiative, <https://doi.org/10.6073/pasta/545502bf79b3e4e03bad2c5816375b01>,
1170 2021.
- Ho, L., Jerves-Cobo, R., Barthel, M., Six, J., Bode, S., Boeckx, P., and Goethals, P.: Greenhouse gas dynamics in an urbanized river system: influence of water quality and land use, *Environ. Sci. Pollut. Res.*, <https://doi.org/10.1007/s11356-021-18081-2>, 2022.
- Hope, D., Palmer, S. M., Billett, M. F., and Dawson, J. J. C.: Carbon dioxide and methane evasion from a temperate peatland
1175 stream, *Limnol. Oceanogr.*, 46, 847–857, <https://doi.org/10.4319/lo.2001.46.4.0847>, 2001.
- Hope, D., Palmer, S. M., Billett, M. F., and Dawson, J. J. C.: Variations in dissolved CO₂ and CH₄ in a first-order stream and catchment: an investigation of soil-stream linkages, *Hydrol. Process.*, 18, 3255–3275, <https://doi.org/10.1002/hyp.5657>, 2004.
- Hu, B., Wang, D., Zhou, J., Meng, W., Li, C., Sun, Z., Guo, X., and Wang, Z.: Greenhouse gases emission from the sewage draining rivers, *Sci. Total Environ.*, 612, 1454–1462, <https://doi.org/10.1016/j.scitotenv.2017.08.055>, 2018.
- 1180 Hu, J.: Greenhouse gas emissions and spatiotemporal variation from rivers in the Zoige Plateau, M.S. thesis, Northwest Agriculture & Forestry University, Xianyang, China, 56 pp., <https://kns.cnki.net/KCMS/detail/detail.aspx?dbname=CMFD201601&filename=1015333482.nh>, 2015.
- 1185 Huang, Y., Yasarer, L. M. W., Li, Z., Sturm, B. S. M., Zhang, Z., Guo, J., and Shen, Y.: Air–water CO₂ and CH₄ fluxes along a river–reservoir continuum: Case study in the Pengxi River, a tributary of the Yangtze River in the Three Gorges Reservoir, China, *Environ. Monit. Assess.*, 189, 223, <https://doi.org/10.1007/s10661-017-5926-2>, 2017.

- Huotari, J., Nykänen, H., Forsius, M., and Arvola, L.: Effect of catchment characteristics on aquatic carbon export from a boreal catchment and its importance in regional carbon cycling, *Global Change Biol.*, 19, 3607–3620, <https://doi.org/10.1111/gcb.12333>, 2013.
- 1190 Hutchins, R. H. S., Prairie, Y. T., and del Giorgio, P. A.: Large-scale landscape drivers of CO₂, CH₄, DOC, and DIC in boreal river networks, *Global Biogeochem. Cycles*, 33, 125–142, <https://doi.org/10.1029/2018GB006106>, 2019.
- Hutchins, R. H. S., Tank, S. E., Olefeldt, D., Quinton, W. L., Spence, C., Dion, N., Estop-Aragonés, C., and Mengistu, S. G.: Fluvial CO₂ and CH₄ patterns across wildfire-disturbed ecozones of subarctic Canada: Current status and implications for future change, *Global Change Biol.*, 26, 2304–2319, <https://doi.org/10.1111/gcb.14960>, 2020a.
- 1195 Hutchins, R. H. S., Casas-Ruiz, J. P., Prairie, Y. T., and del Giorgio, P. A.: Magnitude and drivers of integrated fluvial network greenhouse gas emissions across the boreal landscape in Québec, *Water Res.*, 173, 115556, <https://doi.org/10.1016/j.watres.2020.115556>, 2020b.
- Hutchins, R. H. S., Prairie, Y. T., and del Giorgio, P. A.: The relative importance of seasonality versus regional and network-specific properties in determining the variability of fluvial CO₂, CH₄ and dissolved organic carbon across boreal Québec, 1200 *Aquat. Sci.*, 83, 72, <https://doi.org/10.1007/s00027-021-00830-7>, 2021.
- Hyvonen, N. P., Huttunen, J. T., Shurpali, N. J., Lind, S. E., Marushchek, M. E., Heito, L., and Martikainen, P. J.: The role of drainage ditches in greenhouse gas emissions and surface leaching losses from a cutaway peatland cultivated with a perennial bioenergy crop, *Boreal Environ. Res.*, <http://hdl.handle.net/10138/229311>, 18, 109–126, 2013.
- Jauhiainen, J. and Silvennoinen H: Diffusion GHG fluxes at tropical peatland drainage canal water surfaces, *Suo*, 63, 93–1205 105, <http://www.suo.fi/article/9882>, 2012.
- Jin, H., Yoon, T. K., Begum, M. S., Lee, E.-J., Oh, N.-H., Kang, N., and Park, J.-H.: Longitudinal discontinuities in riverine greenhouse gas dynamics generated by dams and urban wastewater, *Biogeosciences*, 15, 6349–6369, <https://doi.org/10.5194/bg-15-6349-2018>, 2018.
- Jones, J. B. and Mulholland, P. J.: Influence of drainage basin topography and elevation on carbon dioxide and methane supersaturation of stream water, *Biogeochemistry*, 40, 57–72, <https://doi.org/10.1023/A:1005914121280>, 1998a.
- 1210 Jones, J. B. and Mulholland, P. J.: Methane input and evasion in a hardwood forest stream: Effects of subsurface flow from shallow and deep pathway, *Limnol. Oceanogr.*, 43, 1243–1250, <https://doi.org/10.4319/lo.1998.43.6.1243>, 1998b.

- Jurado, A., Borges, A. V., Pujades, E., Briers, P., Nikolenko, O., Dassargues, A., and Brouyère, S.: Dynamics of greenhouse gases in the river–groundwater interface in a gaining river stretch (Triffoy catchment, Belgium), *Hydrogeol. J.*, 26, 2739–2751, 1215 <https://doi.org/10.1007/s10040-018-1834-y>, 2018.
- Juutinen, S., Väiliranta, M., Kuutti, V., Laine, A. M., Virtanen, T., Seppä, H., Weckström, J., and Tuittila, E.-S.: Short-term and long-term carbon dynamics in a northern peatland-stream-lake continuum: A catchment approach, *J. Geophys. Res. Biogeosci.*, 118, 171–183, <https://doi.org/10.1002/jgrg.20028>, 2013.
- Karlsson, J., Serikova, S., Vorobyev, S. N., Rocher-Ros, G., Denfeld, B. A., and Pokrovsky, O. S.: Carbon emission from 1220 Western Siberian inland waters, Zenodo, <https://doi.org/10.5281/ZENODO.4153050>, 2020.
- Kemenes, A., Forsberg, B. R., and Melack, J. M.: Methane release below a tropical hydroelectric dam, *Geophys. Res. Lett.*, 34, L12809, <https://doi.org/10.1029/2007GL029479>, 2007.
- Kemenes, A., Forsberg, B. R., and Melack, J. M.: LBA-ECO LC-07 Methane and Carbon Dioxide emissions from Balbina Reservoir, Brazil, ORNL DAAC, http://daac.ornl.gov/cgi-bin/dsviewer.pl?ds_id=1143, 2013.
- 1225 Kemenes, A., Forsberg, B. R., and Melack, J. M.: Downstream emissions of CH₄ and CO₂ from hydroelectric reservoirs (Tucuruí, Samuel, and Curuá-Una) in the Amazon basin, *Inland Waters*, 6, 295–302, <https://doi.org/10.1080/TW-6.3.980>, 2016.
- Klaus, M., Geibrink, E., Jonsson, A., Bergström, A.-K., Bastviken, D., Laudon, H., Klaminder, J., and Karlsson, J.: Greenhouse gas emissions from boreal inland waters unchanged after forest harvesting, *Biogeosciences*, 15, 5575–5594, <https://doi.org/10.5194/bg-15-5575-2018>, 2018.
- 1230 Kling, G.: Chemistry from thermokarst impacted soils, lakes, and streams near Toolik Lake Alaska, 2008-2011., Environmental Data Initiative, <https://doi.org/10.6073/pasta/2e55d1587290e642938ac1a6caed6ec6>, 2013.
- Kling, G.: Toolik Inlet discharge data collected in summer 1993, Arctic LTER, Toolik Research Station, Alaska., Environmental Data Initiative, <https://doi.org/10.6073/pasta/ae3cf97a2496946fa8ba0cf964271e56>, 2016a.
- Kling, G.: Toolik Inlet discharge data collected in summer 1994, Arctic LTER, Toolik Research Station, Alaska., 1235 Environmental Data Initiative, <https://doi.org/10.6073/pasta/8cc384d957477d5ad48e926ed26dc89b>, 2016b.
- Kling, G.: Toolik Inlet discharge data collected in summer 1995, Arctic LTER, Toolik Research Station, Alaska., Environmental Data Initiative, <https://doi.org/10.6073/pasta/20e10e53cc8b68cffbe98ed0b234d26a>, 2016c.

- Kling, G.: Toolik Inlet discharge data collected in summer 1996, Arctic LTER, Toolik Research Station, Alaska., Environmental Data Initiative, <https://doi.org/10.6073/pasta/6e9d9bd807d8ec133e91d0e665a1550d>, 2016d.
- 1240 Kling, G.: Toolik Inlet discharge data collected in summer 1997, Arctic LTER, Toolik Research Station, Alaska., Environmental Data Initiative, <https://doi.org/10.6073/pasta/33f027ad109d650964a0a084e5df7b11>, 2016e.
- Kling, G.: Toolik Inlet discharge data collected in summer 1998, Arctic LTER, Toolik Research Station, Alaska., Environmental Data Initiative, <https://doi.org/10.6073/pasta/4b78d41f1462c952140b6d2bd4c5d3e4>, 2016f.
- Kling, G.: Toolik Inlet discharge data collected in summer 1999, Arctic LTER, Toolik Research Station, Alaska.,
1245 Environmental Data Initiative, <https://doi.org/10.6073/pasta/37c5b37970b78525819480aa7e4db43a>, 2016g.
- Kling, G.: Toolik Inlet discharge data collected in summer 2000, Arctic LTER, Toolik Research Station, Alaska., Environmental Data Initiative, <https://doi.org/10.6073/pasta/48d71932248e540223bd5650902dd7a4>, 2016h.
- Kling, G.: Toolik Inlet discharge data collected in summer 2001, Arctic LTER, Toolik Research Station, Alaska., Environmental Data Initiative, <https://doi.org/10.6073/PASTA/4EA8FA2D3B89F4BF2B5DE7B98B6A772C>, 2016i.
- 1250 Kling, G.: Toolik Inlet discharge data collected in summer 2002, Arctic LTER, Toolik Research Station, Alaska., Environmental Data Initiative, <https://doi.org/10.6073/pasta/aa535873109be90a8a1cb133b45dbc67>, 2016j.
- Kling, G.: Toolik Inlet discharge data collected in summer 2003, Arctic LTER, Toolik Research Station, Alaska., Environmental Data Initiative, <https://doi.org/10.6073/pasta/07d2ff982627a2a73343c1785358d0a6>, 2016k.
- Kling, G.: Toolik Inlet discharge data collected in summer 2004, Arctic LTER, Toolik Research Station, Alaska.,
1255 Environmental Data Initiative, <https://doi.org/10.6073/pasta/05f608cdb85f2e558febd0fd399da5cf>, 2016l.
- Kling, G.: Toolik Inlet discharge data collected in summer 2005, Arctic LTER, Toolik Research Station, Alaska., Environmental Data Initiative, <https://doi.org/10.6073/pasta/9dde811179666deedd0ecf911be39f65>, 2016m.
- Kling, G.: Toolik Inlet discharge data collected in summer 2006, Arctic LTER, Toolik Research Station, Alaska., Environmental Data Initiative, <https://doi.org/10.6073/pasta/bd8a06d5dab8691912524db28cc24bcd>, 2016n.
- 1260 Kling, G.: Toolik Inlet discharge data collected in summer 2007, Arctic LTER, Toolik Research Station, Alaska., Environmental Data Initiative, <https://doi.org/10.6073/pasta/3af4cbab73c38f76b2829c3abff8f703>, 2016o.

- Kling, G.: Toolik Inlet discharge data collected in summer 2008, Arctic LTER, Toolik Research Station, Alaska., Environmental Data Initiative, <https://doi.org/10.6073/pasta/48e780b581b1071f19c7e5f4b165035d>, 2016p.
- 1265 Kling, G.: Toolik Inlet discharge data collected in summer 2009, Arctic LTER, Toolik Research Station, Alaska., Environmental Data Initiative, <https://doi.org/10.6073/pasta/94bb7d7a93a46ab5363033de6ee7d603>, 2016q.
- Kling, G.: Tussock Watershed stream discharge, electrical conductivity, and temperature measurements from 1992, Environmental Data Initiative, <https://doi.org/10.6073/pasta/1e224958e278841f9a7a035007c65f21>, 2016r.
- Kling, G.: Tussock Watershed stream discharge, electrical conductivity, and temperature measurements from 1993, Environmental Data Initiative, <https://doi.org/10.6073/pasta/f14f444ce51fa77d5f577db4cddb0564>, 2016s.
- 1270 Kling, G.: Tussock Watershed stream discharge, electrical conductivity, and temperature measurements from 1994, Environmental Data Initiative, <https://doi.org/10.6073/pasta/88124e3e8b4a8bbbd49fbb64d64b62d3>, 2016t.
- Kling, G.: Tussock Watershed stream discharge, electrical conductivity, and temperature measurements from 1995, Environmental Data Initiative, <https://doi.org/10.6073/pasta/7e79c3adc44e965240f1c9d75ea676fb>, 2016u.
- Kling, G.: Tussock Watershed stream discharge, electrical conductivity, and temperature measurements from 1996, Environmental Data Initiative, <https://doi.org/10.6073/pasta/6bd568dba3bfaa58181cfb8abff4d639>, 2016v.
- 1275 Kling, G.: Tussock Watershed stream discharge, electrical conductivity, and temperature measurements from 1997, Environmental Data Initiative, <https://doi.org/10.6073/pasta/4c9e9b2bb4861e73dfeaa6bb5e8fb9cd>, 2016w.
- Kling, G.: Tussock Watershed stream discharge, electrical conductivity, and temperature measurements from 1999, Environmental Data Initiative, <https://doi.org/10.6073/pasta/4b943b5a2de08aca8b7dd48542476f12>, 2016x.
- 1280 Kling, G.: Tussock Watershed stream discharge, electrical conductivity, and temperature measurements from 2000, Environmental Data Initiative, <https://doi.org/10.6073/pasta/53a45c5a110f0af13c5ae0ed3154b8ca>, 2016y.
- Kling, G.: Tussock Watershed stream discharge, electrical conductivity, and temperature measurements from 2002, Environmental Data Initiative, <https://doi.org/10.6073/pasta/5c3e5f2495561903c027c6b06544cf70>, 2016z.
- 1285 Kling, G.: Tussock Watershed stream discharge, electrical conductivity, and temperature measurements from 2003, Environmental Data Initiative, <https://doi.org/10.6073/pasta/b24b8bb901a4b1b825e09c7ab494b39d>, 2016aa.

- Kling, G.: Tussock Watershed stream discharge, electrical conductivity, and temperature measurements from 2004, Environmental Data Initiative, <https://doi.org/10.6073/pasta/459c62f862e1724005eb7d91648bfb44>, 2016ab.
- Kling, G. W.: Biogeochemistry data set for Innavait Creek Weir on the North Slope of Alaska 2002-2018, Environmental Data Initiative, <https://doi.org/10.6073/pasta/733c73c6ebffeac6970b2b0f4dddfe6>, 2019a.
- 1290 Kling, G.: Biogeochemistry data set for soil waters, streams, and lakes near Toolik on the North Slope of Alaska., Environmental Data Initiative, <https://doi.org/10.6073/pasta/574fd24522eee7a0c07fc260ccc0e2fa>, 2019b.
- Kling, G.: Toolik Lake Inlet discharge data collected during summers of 2010 to 2018, Arctic LTER, Toolik Research Station, Alaska., Environmental Data Initiative, <https://doi.org/10.6073/pasta/169d1bae55373c44a368727573ef70eb>, 2019c
- 1295 Kling, G. 2022. Biogeochemistry data set for soil waters, streams, and lakes near Toolik Lake on the North Slope of Alaska, 2012 through 2020 ver. 2, Environmental Data Initiative, <https://doi.org/10.6073/pasta/4e25db9ae9372f5339f2795792814845>, 2022.
- Kling, G. W., Kipphut, G. W., and Miller, M. C.: The flux of CO₂ and CH₄ from lakes and rivers in arctic Alaska, *Hydrobiologia*, 240, 23–36, <https://doi.org/10.1007/BF00013449>, 1992.
- 1300 Köhn, D., Welpelo, C., Günther, A., and Jurasinski, G.: Drainage ditches contribute considerably to the CH₄ budget of a drained and a rewetted temperate fen, *Wetlands*, 41, 71, <https://doi.org/10.1007/s13157-021-01465-y>, 2021.
- Kokic, J., Sahlée, E., Sobek, S., Vachon, D., and Wallin, M. B.: High spatial variability of gas transfer velocity in streams revealed by turbulence measurements, *Inland Waters*, 8, 461–473, <https://doi.org/10.1080/20442041.2018.1500228>, 2018.
- Koné, Y. J. M., Abril, G., Delille, B., and Borges, A. V.: Seasonal variability of methane in the rivers and lagoons of Ivory Coast (West Africa), *Biogeochemistry*, 100, 21–37, <https://doi.org/10.1007/s10533-009-9402-0>, 2010.
- 1305 Krickov, I. V., Serikova, S., Pokrovsky, O. S., Vorobyev, S. N., Lim, A. G., Siewert, M. B., and Karlsson, J.: Dataset for Ob River floodplain, Zenodo, <https://doi.org/10.5281/ZENODO.4563905>, 2021a.
- Krickov, I. V., Serikova, S., Pokrovsky, O. S., Vorobyev, S. N., Lim, A. G., Siewert, M. B., and Karlsson, J.: Sizable carbon emission from the floodplain of Ob River, *Ecol. Indic.*, 131, 108164, <https://doi.org/10.1016/j.ecolind.2021.108164>, 2021b.
- 1310 Kuhn, C., Bettigole, C., Glick, H. B., Seegmiller, L., Oliver, C. D., and Raymond, P.: Patterns in stream greenhouse gas dynamics from mountains to plains in northcentral Wyoming: Aquatic GHG emissions in Wyoming, *J. Geophys. Res. Biogeosci.*, 122, 2173–2190, <https://doi.org/10.1002/2017JG003906>, 2017.

- LaboMaranger: LaboMaranger/RdN_GHG: data available, Zenodo, <https://doi.org/10.5281/ZENODO.4566377>, 2021.
- 1315 Laini, A., Bartoli, M., Castaldi, S., Viaroli, P., Capri, E., and Trevisan, M.: Greenhouse gases (CO₂, CH₄ and N₂O) in lowland springs within an agricultural impacted watershed (Po River Plain, northern Italy), *Chem. Ecol.*, 27, 177–187, <https://doi.org/10.1080/02757540.2010.547489>, 2011.
- 1320 Lamarche-Gagnon, G., Wadham, J. L., Sherwood Lollar, B., Arndt, S., Fietzek, P., Beaton, A. D., Tedstone, A. J., Telling, J., Bagshaw, E. A., Hawkings, J. R., Kohler, T. J., Zarsky, J. D., Mowlem, M. C., Anesio, A. M., and Stibal, M.: Greenland melt drives continuous export of methane from the ice-sheet bed, *Nature*, 565, 73–77, <https://doi.org/10.1038/s41586-018-0800-0>, 2019.
- Lamontagne, R. A., Swinnerton, J. W., Linnenbom, V. J., and Smith, W. D.: Methane concentrations in various marine environments, *J. Geophys. Res.*, 78, 5317–5324, <https://doi.org/10.1029/JC078i024p05317>, 1973.
- Leng, P., Kamjunke, N., Li, F., and Koschorreck, M.: Long-term monitoring CO₂ and CH₄ concentrations from two German streams, figshare, <https://doi.org/10.6084/M9.FIGSHARE.12866945.V1>, 2020.
- 1325 Leng, P., Kamjunke, N., Li, F., and Koschorreck, M.: Temporal patterns of methane emissions from two streams with different riparian connectivity, *J. Geophys. Res. Biogeosci.*, 126, e2020JG006104, <https://doi.org/10.1029/2020jg006104>, 2021.
- Li, L., Yan, R., and Xue, B.: Methane levels of a river network in Wuxi City, China and response to water governance, *Water*, 12, 2617, <https://doi.org/10.3390/w12092617>, 2020a.
- 1330 Li, X., Yao, H., Yu, Y., Cao, Y., and Tang, C.: Greenhouse gases in an urban river: Trend, isotopic evidence for underlying processes, and the impact of in-river structures, *J. Hydrol.*, 591, 125290, <https://doi.org/10.1016/j.jhydrol.2020.125290>, 2020b.
- Liang, X., Zhang, X., Sun, Q., He, C., Chen, X., Liu, X., and Chen, Z.: The role of filamentous algae *Spirogyra* spp. in methane production and emissions in streams, *Aquat. Sci.*, 78, 227–239, <https://doi.org/10.1007/s00027-015-0419-2>, 2016.
- Lilley, M. D., de Angelis, M. A., and Olson, E. J.: Methane concentrations and estimated fluxes from Pacific Northwest rivers, *SIL Communications*, 1953-1996, 25, 187–196, <https://doi.org/10.1080/05384680.1996.11904080>, 1996.
- 1335 Liu, K., Hu, Z., Wei, X., Jiang, Z., Lu, H., and Wang, C.: Research on methane flux production flux in urban black and odourous rivers in summer: Taking Chaoyang Creek in Nanning City as an example, *Earth Environ.*, 43, 415–419, doi: (China): 10.14050/j.cnki1672-9250/2015.04.005, 2015.

- Loken, L., Crawford, J., Stanley, E., Butman, D., and Striegl, R.: Columbia River spatial water chemistry, Environmental Data Initiative, <https://doi.org/10.6073/pasta/e881070c9e8f6b7f774d3c65b27a9f69>, 2018a.
- 1340 Loken, L., Crawford, J., and Stanley, E.: Mississippi River spatial water chemistry Environmental Research Letters datasets, <https://doi.org/10.6073/pasta/c1b9dbd9a96edfb5e39a94cfef2982b9>, 2018b.
- Looman, A., Maher, D. T., Pendall, E., Bass, A., and Santos, I. R.: The carbon dioxide evasion cycle of an intermittent first-order stream: contrasting water–air and soil–air exchange, *Biogeochemistry*, 132, 87–102, <https://doi.org/10.1007/s10533-016-0289-2>, 2017.
- 1345 Looman, A., Santos, I. R., Tait, D. R., Webb, J., Holloway, C., and Maher, D. T.: Dissolved carbon, greenhouse gases, and $\delta^{13}\text{C}$ dynamics in four estuaries across a land use gradient, *Aquat. Sci.*, 81, 22, <https://doi.org/10.1007/s00027-018-0617-9>, 2019.
- Lottig, N. and Stanley, E.: North Temperate Lakes LTER: Northern Highlands stream chemistry survey 2006 ver. 18, Environmental Data Initiative, <https://doi.org/10.6073/pasta/a73255a67041a580601711c9ac761f26>, 2022.
- 1350 Lottig, N. R., Stanley, E. H., and Maxted, J. T.: Assessing the influence of upstream drainage lakes on fluvial organic carbon in a wetland-rich region, *J. Geophys. Res.*, 117, G03011, <https://doi.org/10.1029/2012JG001983>, 2012.
- Ludwig, S., Holmes, R. M., Natali, S., Mann, P., Schade, J., Jimmie, J., Bristol, E., Peter, D., and Dabrowski, J.: Polaris Project 2017: Aquatic isotopes, carbon, and nitrogen, Yukon-Kuskokwim Delta, Alaska, Arctic Data Center, <https://doi.org/10.18739/A20298>, 2018a.
- 1355 Ludwig, S., Holmes, R. M., Natali, S., Schade, J., and Mann, P.: Yukon-Kuskokwim Delta fire: aquatic data, Yukon-Kuskokwim Delta Alaska, 2015-2016, Arctic Data Center, <https://doi.org/10.18739/A2HG45>, 2018b.
- Ludwig, S. M., Natali, S. M., Mann, P. J., Schade, J. D., Holmes, R. M., Powell, M., Fiske, G., and Commane, R.: Using machine learning to predict inland aquatic CO_2 and CH_4 concentrations and the effects of wildfires in the Yukon-Kuskokwim Delta, Alaska, *Global Biogeochem. Cycles*, 36, e2021GB007146, <https://doi.org/10.1029/2021GB007146>, 2022.
- 1360 Ma, P., Li, Y., Qi, L., Tian, X., and Yang, L.: Characteristics of CH_4 emission from Shaying River and its influence factors, *Environ. Sci. Technol. (China)*, 40, 26–30, doi (China): 10.3969/j.issn.1003-6504.2017.02.005, 2017.
- Mach, V., Bednařík, A., Čáp, L., Šipoš, J., and Rulík, M.: Seasonal measurement of greenhouse gas concentrations and emissions along the longitudinal profile of a small stream, *Pol. J. Environ. Stud.*, 25, 2047–2056, <https://doi.org/10.15244/pjoes/61668>, 2016.

- 1365 Maeck, A., DelSontro, T., McGinnis, D. F., Fischer, H., Flury, S., Schmidt, M., Fietzek, P., and Lorke, A.: Sediment trapping by dams creates methane emission hot spots, *Environ. Sci. Technol.*, 47, 8130–8137, <https://doi.org/10.1021/es4003907>, 2013.
- Maier, M.-S., Teodoru, C. R., and Wehrli, B.: Spatio-temporal variations of lateral and atmospheric carbon fluxes from the Danube Delta (dataset): A 2-year dataset of measured concentrations and fluxes, ETH Research Collection, <https://doi.org/10.3929/ethz-b-000416925>, 2020.
- 1370 Maier, M.-S., Teodoru, C. R., and Wehrli, B.: Spatio-temporal variations in lateral and atmospheric carbon fluxes from the Danube Delta, *Biogeosciences*, 18, 1417–1437, <https://doi.org/10.5194/bg-18-1417-2021>, 2021.
- Manning, C. C. M., Preston, V. L., Jones, S. F., Michel, A. P. M., Nicholson, D. P., Duke, P. J., Ahmed, M. M. M., Manganini, K., Else, B. G. T., and Tortell, P. D.: Dissolved methane and nitrous oxide concentrations measured on discrete bottle samples in Cambridge Bay, Nunavut, Canada from 2017, PANGAEA, <https://doi.org/10.1594/PANGAEA.907149>, 2019a.
- 1375 Manning, C. C. M., Preston, V. L., Jones, S. F., Michel, A. P. M., Nicholson, D. P., Duke, P. J., Ahmed, M. M. M., Manganini, K., Else, B. G. T., and Tortell, P. D.: Dissolved methane and nitrous oxide concentrations measured on discrete bottle samples in Cambridge Bay, Nunavut, Canada from 2018, PANGAEA, <https://doi.org/10.1594/PANGAEA.907150>, 2019b.
- Manning, C. C., Preston, V. L., Jones, S. F., Michel, A. P. M., Nicholson, D. P., Duke, P. J., Ahmed, M. M. M., Manganini, K., Else, B. G. T., and Tortell, P. D.: River inflow dominates methane emissions in an Arctic coastal system, *Geophys. Res. Lett.*, 47, e2020GL087669, <https://doi.org/10.1029/2020gl087669>, 2020.
- 1380 Manning, F. C.: Carbon dynamics in oil palm agro-ecosystems, Ph.D. thesis, University of Aberdeen, Aberdeen, UK, 435 pp., https://abdn.alma.exlibrisgroup.com/discovery/delivery/44ABE_INST:44ABE_VU1/12152929810005941, 2019a.
- Manning, F. C., Kho, L. K., Hill, T. C., Cornulier, T., and Teh, Y. A.: Carbon emissions from oil palm plantations on peat soil, *Front. For. Glob. Change*, 2, 37, <https://doi.org/10.3389/ffgc.2019.00037>, 2019b.
- 1385 Marescaux, A., Thieu, V., and Garnier, J.: Carbon dioxide, methane and nitrous oxide emissions from the human-impacted Seine watershed in France, *Sci. Total Environ.*, 643, 247–259, <https://doi.org/10.1016/j.scitotenv.2018.06.151>, 2018.
- Martin, J. B., Pain, A. J., Martin, E. E., Rahman, S., and Ackerman, P.: Comparisons of nutrients exported from Greenlandic glacial and deglaciated watersheds, *Global Biogeochem. Cycles*, 34, e2020GB006661, <https://doi.org/10.1029/2020GB006661>, 2020.

- 1390 Martinez-Cruz, K., Gonzalez-Valencia, R., Sepulveda-Jauregui, A., Plascencia-Hernandez, F., Belmonte-Izquierdo, Y., and Thalasso, F.: Methane emission from aquatic ecosystems of Mexico City, *Aquat. Sci.*, 79, 159–169, <https://doi.org/10.1007/s00027-016-0487-y>, 2017.
- Marwick, T. R., Tamoooh, F., Ogwoka, B., Teodoru, C., Borges, A. V., Darchambeau, F., and Bouillon, S.: Dynamic seasonal nitrogen cycling in response to anthropogenic N loading in a tropical catchment, Athi–Galana–Sabaki River, Kenya, *Biogeosciences*, 11, 443–460, <https://doi.org/10.5194/bg-11-443-2014>, 2014.
- 1395 Marwick, T. R., Tamoooh, F., Ogwoka, B., Borges, A. V., Darchambeau, F., and Bouillon, S.: A comprehensive biogeochemical record and annual flux estimates for the Sabaki River (Kenya), *Biogeosciences*, 15, 1683–1700, <https://doi.org/10.5194/bg-15-1683-2018>, 2018.
- Matoušů, A., Rulík, M., Tušer, M., Bednařík, A., Šimek, K., and Bussmann, I.: Methane dynamics in a large river: a case study of the Elbe River, *Aquat. Sci.*, 81, 12, <https://doi.org/10.1007/s00027-018-0609-9>, 2019.
- 1400 Matveev, A., Blais, M. A., Laurion, I., and Vincent, W. F.: Dissolved methane, carbon dioxide and limnological data from subarctic rivers, northern Québec, Canada, v. 1.0 (2019-2019), *Nordicana D*, <https://doi.org/10.5885/45660CE-8B92339884C146D0>, 2020.
- McGinnis, D. F., Bilsley, N., Schmidt, M., Fietzek, P., Bodmer, P., Premke, K., Lorke, A., and Flury, S.: Deconstructing methane emissions from a small northern European river: Hydrodynamics and temperature as key drivers, *Environ. Sci. Technol.*, 50, 11680–11687, <https://doi.org/10.1021/acs.est.6b03268>, 2016.
- 1405 Mei, D., Ni, M., Liang, X., Hou, L., Wang, F., and He, C.: Filamentous green algae *Spirogyra* regulates methane emissions from eutrophic rivers, *Environ. Sci. Pollut. Res.*, 28, 3660–3671, <https://doi.org/10.1007/s11356-020-10754-8>, 2021.
- Middelburg, J. J., Nieuwenhuize, J., Iversen, N., Høgh, N., de Wilde, H., Helder, W., Seifert, R., and Christof, O.: Methane distribution in European tidal estuaries, *Biogeochemistry*, 59, 95–119, <https://doi.org/10.1023/A:1015515130419>, 2002.
- 1410 Minkinen, K. and Laine, J.: Vegetation heterogeneity and ditches create spatial variability in methane fluxes from peatlands drained for forestry, *Plant Soil*, 285, 289–304, <https://doi.org/10.1007/s11104-006-9016-4>, 2006.
- Minkinen, K., Laine, J., Nykänen, H., and Martikainen, P. J.: Importance of drainage ditches in emissions of methane from mires drained for forestry, *Can. J. For. Res.*, 27, 949–952, <https://doi.org/10.1139/x97-016>, 1997.

- 1415 Morozumi, T., Shingubara, R., Murase, J., Nagai, S., Kobayashi, H., Takano, S., Tei, S., Fan, R., Maximov, T. C., and Sugimoto, A.: Usability of water surface reflectance for the determination of riverine dissolved methane during extreme flooding in northeastern Siberia, *Polar Sci.*, 21, 186–194, <https://doi.org/10.1016/j.polar.2019.01.005>, 2019.
- Morrice, J. A., Dahm, C. N., Valett, H. M., Unnikrishna, P. V., and Campana, M. E.: Terminal electron accepting processes in the alluvial sediments of a headwater stream, *J. N. Amer. Benthol. Soc.*, 19, 593–608, <https://doi.org/10.2307/1468119>, 2000.
- 1420 Mosher, J., Fortner, A., Phillips, J., Bevelhimer, M., Stewart, A., and Troia, M.: Spatial and temporal correlates of greenhouse gas diffusion from a hydropower reservoir in the southern United States, *Water*, 7, 5910–5927, <https://doi.org/10.3390/w7115910>, 2015.
- Mulholland, P. J., Helton, A. M., Poole, G. C., Hall, R. O., Hamilton, S. K., Peterson, B. J., Tank, J. L., Ashkenas, L. R., Cooper, L. W., Dahm, C. N., Dodds, W. K., Findlay, S. E. G., Gregory, S. V., Grimm, N. B., Johnson, S. L., McDowell, W. H., Meyer, J. L., Valett, H. M., Webster, J. R., Arango, C. P., Beaulieu, J. J., Bernot, M. J., Burgin, A. J., Crenshaw, C. L., Johnson, L. T., Niederlehner, B. R., O'Brien, J. M., Potter, J. D., Sheibley, R. W., Sobota, D. J., and Thomas, S. M.: Stream denitrification across biomes and its response to anthropogenic nitrate loading, *Nature*, 452, 202–205, <https://doi.org/10.1038/nature06686>, 2008.
- 1430 Murphy, J. F., Arnold, A., Duerdoth, C. P., Hawczak, A., Pacioglu, O., Pretty, J. L., and Jones, J. I.: Temporal variation in temperature and light availability in the Hampshire Avon, United Kingdom [Macronutrient Cycling], NERC Environmental Information Data Centre, <https://doi.org/10.5285/9B6A6233-85AD-44F4-BA83-4905B8C48713>, 2017.
- Natchimuthu, S., Wallin, M. B., Klemetsson, L., and Bastviken, D.: Spatio-temporal patterns of stream methane and carbon dioxide emissions in a hemiboreal catchment in Southwest Sweden, *Sci. Rep.*, 7, 39729, <https://doi.org/10.1038/srep39729>, 2017.
- 1435 National Ecological Observatory Network (NEON): Chemical properties of surface water (DP1.20093.001): RELEASE-2021 (RELEASE-2021), <https://doi.org/10.48443/05K7-E011>, 2021a.
- National Ecological Observatory Network (NEON): Discharge field collection (DP1.20048.001): RELEASE-2021 (RELEASE-2021), <https://doi.org/10.48443/8TD9-4Z94>, 2021b.
- 1440 Neu, V., Neill, C., and Krusche, A. V.: Gaseous and fluvial carbon export from an Amazon forest watershed, *Biogeochemistry*, 105, 133–147, <https://doi.org/10.1007/s10533-011-9581-3>, 2011.

- Okuku, E. O., Bouillon, S., Tole, M., and Borges, A. V.: Diffusive emissions of methane and nitrous oxide from a cascade of tropical hydropower reservoirs in Kenya, *Lakes & Reserv.*, 24, 127–135, <https://doi.org/10.1111/lre.12264>, 2019.
- 1445 Olid, C., Rodellas, V., Rocher-Ros, G., Garcia-Orellana, J., Diego-Feliu, M., Alorda-Kleinglass, A., Bastviken, D., and Karlsson, J.: Groundwater discharge as a driver of methane emissions from Arctic lakes, *Nat. Commun.*, 13, 3667, <https://doi.org/10.1038/s41467-022-31219-1>, 2022.
- Osudar, R., Liebner, S., Alawi, M., Yang, S., Bussmann, I., and Wagner, D.: Methane turnover and methanotrophic communities in arctic aquatic ecosystems of the Lena Delta, Northeast Siberia, *FEMS Microbiol. Ecol.*, 92, fiw116, <https://doi.org/10.1093/femsec/fiw116>, 2016.
- 1450 Oviedo-Vargas, D., Genereux, D. P., Dierick, D., and Oberbauer, S. F.: The effect of regional groundwater on carbon dioxide and methane emissions from a lowland rainforest stream in Costa Rica, *J. Geophys. Res. Biogeosci.*, 120, 2579–2595, <https://doi.org/10.1002/2015JG003009>, 2015.
- Pain, A., Martin, J., Martin, E., and Rahman, S.: Hydrogeochemistry of Greenlandic proglacial and nonglacial streams, 2017-2018, Arctic Data Center, <https://doi.org/10.18739/A2PC2T94T>, 2019.
- Peacock, M: Wicken Lode data, Swedish Agricultural University, Uppsala, Sweden, unpublished, 2020.
- 1455 Peacock, M.: Forest ditch GHG data, figshare, <https://doi.org/10.6084/m9.figshare.15152253.V3>, 2021a.
- Peacock, M.: Peacock et al, 2021, GCB data figshare, <https://doi.org/10.6084/m9.figshare.14784852.V2>, 2021b.
- Peacock, M., Ridley, L. M., Evans, C. D., and Gauci, V.: Management effects on greenhouse gas dynamics in fen ditches, *Sci. Total Environ.*, 578, 601–612, <https://doi.org/10.1016/j.scitotenv.2016.11.005>, 2017.
- 1460 Peacock, M., Gauci, V., Baird, A. J., Burden, A., Chapman, P. J., Cumming, A., Evans, J. G., Grayson, R. P., Holden, J., Kaduk, J., Morrison, R., Page, S., Pan, G., Ridley, L. M., Williamson, J., Worrall, F., and Evans, C. D.: The full carbon balance of a rewetted cropland fen and a conservation-managed fen, *Agric. Ecosys. Environ.*, 269, 1–12, <https://doi.org/10.1016/j.agee.2018.09.020>, 2019.
- 1465 Peacock, M., Audet, J., Bastviken, D., Cook, S., Evans, C. D., Grinham, A., Holgerson, M. A., Högbom, L., Pickard, A. E., Zielinski, P., and Futter, M. N.: Small artificial waterbodies are widespread and persistent emitters of methane and carbon dioxide, *Global Change Biol.*, 27, 5109–5123, <https://doi.org/10.1111/gcb.15762>, 2021a.

- Peacock, M., Granath, G., Wallin, M. B., Högbom, L., and Futter, M. N.: Significant emissions from forest drainage ditches—An unaccounted term in anthropogenic greenhouse gas inventories?, *J. Geophys. Res. Biogeosci.*, 126, e2021JG006478, <https://doi.org/10.1029/2021JG006478>, 2021b.
- 1470 Pickard, A. E., Skiba, U. M., Carvalho, L., Heal, K. V., Rees, R. M., and Harley, J. F.: River Tay, Scotland, water chemistry and greenhouse gas measurements 2009-2010, NERC Environmental Information Data Centre, <https://doi.org/10.5285/a61da7da-b7ef-40b7-a324-c3711ef81207>, 2019.
- Pickard, A., Dinsmore, K. J., Billett, M. F., and Branagan, M.: Aquatic carbon and greenhouse gas concentrations in headwater streams draining from natural, drained and restored peatland catchments in the Flow Country, Scotland, September 2008-August 2010, NERC Environmental Information Data Centre, <https://doi.org/10.5285/7525088d-e504-456a-bc55-e48d8ca85303>, 2021.
- 1475
- Pickard, A. E., Branagan, M., Billett, M. F., Andersen, R., and Dinsmore, K. J.: Effects of peatland management on aquatic carbon concentrations and fluxes, *Biogeosciences*, 19, 1321–1334, <https://doi.org/10.5194/bg-19-1321-2022>, 2022.
- Pulliam, W. M.: Carbon dioxide and methane exports from a southeastern floodplain swamp, *Ecol. Monogr.*, 63, 29–53, <https://doi.org/10.2307/2937122>, 1993.
- 1480 Qin, X., Li, Y., Wan, Y., Fan, M., Liao, Y., Li, Y., Wang, B., and Gao, Q.: Diffusive flux of CH₄ and N₂O from agricultural river networks: Regression tree and importance analysis, *Sci. Total Environ.*, 717, 137244, <https://doi.org/10.1016/j.scitotenv.2020.137244>, 2020a.
- Qin, X., Li, Y., Wan, Y., Fan, M., Liao, Y., Li, Y., Wang, B., Gao, Q., Wu, H., and Chen, X.: Multiple stable isotopic signatures corroborate the predominance of acetoclastic methanogenesis during CH₄ formation in agricultural river networks, *Agric. Ecosys. Environ.*, 296, 106930, <https://doi.org/10.1016/j.agee.2020.106930>, 2020b.
- 1485
- Qin, Y., Wang, Z., Li, Z., and Yang, B.: CO₂ and CH₄ flux across water-air interface and environmental factors in Pengxi River of the Three Gorges Reservoir, *J. Earth Environ.*, 10, 177–189, doi (China): 10.7515./JEE182071, 2019a.
- Qin, Y., Wang, Z., Li, Z., and Yang, B.: CO₂ and CH₄ partial press and flux across water-air interface in the downstream of the Jinsha River, China, *Appl. Ecol. Environ. Res.*, 17, https://doi.org/10.15666/aeer/1703_58235839, 2019b.
- 1490 Qu, B., Aho, K. S., Li, C., Kang, S., Sillanpää, M., Yan, F., and Raymond, P. A.: Greenhouse gases emissions in rivers of the Tibetan Plateau, *Sci. Rep.*, 7, 16573, <https://doi.org/10.1038/s41598-017-16552-6>, 2017.

- Rajkumar, N. A., Barnes, J., Ramesh, R., Purvaja, R., and Upstill-Goddard, R. C.: Methane and nitrous oxide fluxes in the polluted Adyar River and estuary, SE India, *Mar. Pollut. Bull.*, 56, 2043–2051, <https://doi.org/10.1016/j.marpolbul.2008.08.005>, 2008.
- 1495 Ran, L., Shi, H., and Yang, X.: Magnitude and drivers of CO₂ and CH₄ emissions from an arid/semiarid river catchment on the Chinese Loess Plateau, *J. Hydrol.*, 598, 126260, <https://doi.org/10.1016/j.jhydrol.2021.126260>, 2021.
- Reeburgh, W. S., King, J. Y., Regli, S. K., Kling, G. W., Auerbach, N. A., and Walker, D. A.: A CH₄ emission estimate for the Kuparuk River basin, Alaska, *J. Geophys. Res.*, 103, 29005–29013, <https://doi.org/10.1029/98JD00993>, 1998.
- 1500 Richey, J. E., Devol, A. H., Wofsy, S. C., Victoria, R., and Riberio, M. N. G.: Biogenic gases and the oxidation and reduction of carbon in Amazon River and floodplain waters: Amazon dissolved gases, *Limnol. Oceanogr.*, 33, 551–561, <https://doi.org/10.4319/lo.1988.33.4.0551>, 1988.
- Robison, A. and Wolheim, W.: PIE LTER dissolved methane and water temperature from four headwater streams in Massachusetts and New Hampshire., *Environmental Data Initiative*, <http://dx.doi.org/10.6073/pasta/789147c37923aeccc924ab33b35595eb>, 2021a.
- 1505 Robison, A. and Wollheim, W.: PIE LTER time series of methane, CO₂ and N₂O ebullition measurements at four headwater streams in Massachusetts and New Hampshire., *Environmental Data Initiative*, <https://doi.org/10.6073/pasta/9b7fb5afa7b55f3c198f37ade701a542>, 2021b.
- Robison, A. L., Wollheim, W. M., Turek, B., Bova, C., Snay, C., and Varner, R. K.: Spatial and temporal heterogeneity of methane ebullition in lowland headwater streams and the impact on sampling design, *Limnol. Oceanogr.*, 66, 4063–4076, <https://doi.org/10.1002/lno.11943>, 2021.
- 1510 Rocher-Ros, G.: Miellajokka summer survey, Swedish University of Agricultural Sciences, Umeå, Sweden, unpublished, 2020.
- Rocher-Ros, G.: Impact of reindeer grazing on C emissions and exports from Arctic streams, Swedish University of Agricultural Sciences, Sweden, unpublished, 2021.
- 1515 Roulet, N. T. and Moore, T. R.: The effect of forestry drainage practices on the emission of methane from northern peatlands, *Can. J. For. Res.*, 25, 491–499, <https://doi.org/10.1139/x95-055>, 1995.
- Rovelli, L., Olde, L. A., Heppell, C. M., Binley, A., Yvon-Durocher, G., Glud, R. N., and Trimmer, M.: High-resolution time series of day and night outgassing rates of carbon dioxide and methane for six tributaries of Hampshire River Avon (UK)

- collected with an automated floating chamber in late spring 2015., figshare, <https://doi.org/10.6084/m9.figshare.16545954.v1>,
1520 2021a.
- Rovelli, L., Olde, L. A., Heppell, C. M., Binley, A., Yvon-Durocher, G., Glud, R. N., and Trimmer, M.: Summary data of chamber-based oxygen and methane consumption and production in the streambed and outgassing of carbon dioxide and methane to the atmosphere collected seasonally for six tributaries of Hampshire River Avon (UK) during 2013-2014., figshare, <https://doi.org/10.6084/m9.figshare.16545846.v1>, 2021b.
- 1525 Rovelli, L., Olde, L. A., Heppell, C. M., Binley, A., Yvon-Durocher, G., Glud, R. N., and Trimmer, M.: Contrasting biophysical controls on carbon dioxide and methane outgassing from streams, *J. Geophys. Res. Biogeosci.*, e2021JG006328, 127, <https://doi.org/10.1029/2021JG006328>, 2022.
- Rulík, M., Čáp, L., and Hlaváčová, E.: Methane in the hyporheic zone of a small lowland stream (Sitka, Czech Republic), *Limnologica*, 30, 359–366, [https://doi.org/10.1016/S0075-9511\(00\)80029-8](https://doi.org/10.1016/S0075-9511(00)80029-8), 2000.
- 1530 Sanders, I. A., Heppell, C. M., Cotton, J. A., Wharton, G., Hildrew, A. G., Flowers, E. J., and Trimmer, M.: Emission of methane from chalk streams has potential implications for agricultural practices, *Freshwater Biol.*, 52, 1176–1186, <https://doi.org/10.1111/j.1365-2427.2007.01745.x>, 2007.
- Sansone, F. J., Holmes, M. E., and Popp, B. N.: Methane stable isotopic ratios and concentrations as indicators of methane dynamics in estuaries, *Global Biogeochem. Cycles*, 13, 463–474, <https://doi.org/10.1029/1999gb900012>, 1999.
- 1535 Sawakuchi, H. O., Bastviken, D., Sawakuchi, A. O., Krusche, A. V., Ballester, M. V. R., and Richey, J. E.: Methane emissions from Amazonian rivers and their contribution to the global methane budget, *Global Change Biol.*, 20, 2829–2840, <https://doi.org/10.1111/gcb.12646>, 2014.
- Sawakuchi, H. O., Bastviken, D., Enrich-Prast, A., Ward, N. D., Camargo, P. B., and Richey, J. E.: Low diffusive methane emissions from the main channel of a large Amazonian run-of-the-river reservoir attributed to high methane oxidation, *Front. Environ. Sci.*, 9, 655455, <https://doi.org/10.3389/fenvs.2021.655455>, 2021.
- 1540 Schade, J. D., Bailio, J., and McDowell, W. H.: Greenhouse gas flux from headwater streams in New Hampshire, USA: Patterns and drivers: Greenhouse gas flux from headwater streams, *Limnol. Oceanogr.*, 61, S165–S174, <https://doi.org/10.1002/lno.10337>, 2016.

- Schrier-Uijl, A. P., Kroon, P. S., Leffelaar, P. A., van Huissteden, J. C., Berendse, F., and Veenendaal, E. M.: Methane emissions in two drained peat agro-ecosystems with high and low agricultural intensity, *Plant Soil*, 329, 509–520, <https://doi.org/10.1007/s11104-009-0180-1>, 2010.
- Schrier-Uijl, A. P., Veraart, A. J., Leffelaar, P. A., Berendse, F., and Veenendaal, E. M.: Release of CO₂ and CH₄ from lakes and drainage ditches in temperate wetlands, *Biogeochemistry*, 102, 265–279, <https://doi.org/10.1007/s10533-010-9440-7>, 2011.
- 1545 Schuster, P. F. (Ed.): Water and sediment quality in the Yukon River Basin, Alaska, during Water Year 2001, U.S. Geological Survey, Denver, CO, 120 pp., <https://doi.org/10.3133/ofr03427>, 2003.
- Schuster, P. F. (Ed.): Water and sediment quality in the Yukon River Basin, Alaska, during Water Year 2002, U.S. Geological Survey, Denver, CO, 82 pp., <https://doi.org/10.3133/ofr20051199>, 2006a.
- Schuster, P. F. (Ed.): Water and sediment quality in the Yukon River Basin, Alaska, during Water Year 2003, U.S. Geological Survey, Boulder, CO, 74 pp., <https://doi.org/10.3133/ofr20051397>, 2006b.
- 1555 Schuster, P. F. (Ed.): Water and sediment quality in the Yukon River Basin, Alaska, during Water Year 2004, U.S. Geological Survey, Boulder, CO, 67 pp., <https://doi.org/10.3133/ofr20061258>, 2006c.
- Schuster, P. F. (Ed.): Water and sediment quality in the Yukon River Basin, Alaska, during Water Year 2005, U.S. Geological Survey, Boulder, CO, 65 pp., <https://doi.org/10.3133/ofr20071037>, 2007.
- Schuster, P. F., Maracle, K. B., and Herman-Mercer, N. (Eds.): Water quality in the Yukon River Basin, Alaska, during Water Years 2006 - 2008, U.S. Geological Survey, Reston, VA, 220 pp., <https://doi.org/10.3133/ofr20101241>, 2010.
- Selvam, B. P., Natchimuthu, S., Arunachalam, L., and Bastviken, D.: Methane and carbon dioxide emissions from inland waters in India - implications for large scale greenhouse gas balances, *Global Change Biol.*, 20, 3397–3407, <https://doi.org/10.1111/gcb.12575>, 2014.
- 1560 Shelley, F., Grey, J., and Trimmer, M.: Widespread methanotrophic primary production in lowland chalk rivers, *Proc. R. Soc. B*, 281, 20132854, <https://doi.org/10.1098/rspb.2013.2854>, 2014.
- Shi, W., Chen, Q., Yi, Q., Yu, J., Ji, Y., Hu, L., and Chen, Y.: Carbon Emission from Cascade Reservoirs: Spatial heterogeneity and mechanisms, *Environ. Sci. Technol.*, 51, 12175–12181, <https://doi.org/10.1021/acs.est.7b03590>, 2017.

- Sieczko, A. K., Demeter, K., Singer, G. A., Tritthart, M., Preiner, S., Mayr, M., Meisterl, K., and Peduzzi, P.: Aquatic methane dynamics in a human-impacted river-floodplain of the Danube: Floodplain methane emission, *Limnol. Oceanogr.*, 61, S175–S187, <https://doi.org/10.1002/lno.10346>, 2016.
- Silvennoinen, H., Liikanen, A., Rintala, J., and Martikainen, P. J.: Greenhouse gas fluxes from the eutrophic Temmesjoki River and its Estuary in the Liminganlahti Bay (the Baltic Sea), *Biogeochemistry*, 90, 193–208, <https://doi.org/10.1007/s10533-008-9244-1>, 2008.
- 1575 Smith, L. K., Lewis, Jr., W. M., Chanton, J. P., Cronin, G., and Hamilton, S. K.: Methane emissions from the Orinoco River floodplain, Venezuela, *Biogeochemistry*, 51, 113–140, <https://doi.org/10.1023/A:1006443429909>, 2000.
- Smith, R. L. and Bohlke, J. K.: Methane and nitrous oxide temporal and spatial concentrations in the Iroquois River and Sugar Creek in Northwestern Indiana and Northeastern Illinois, 1999-2003 (ver. 2.0, November 2020), U.S. Geological Survey data release, <https://doi.org/10.5066/f7th8kwz>, 2018.
- 1580 Smith, R. L. and Böhlke, J. K.: Methane and nitrous oxide temporal and spatial variability in two midwestern USA streams containing high nitrate concentrations, *Sci. Total Environ.*, 685, 574–588, <https://doi.org/10.1016/j.scitotenv.2019.05.374>, 2019.
- Smith, R. M., Kaushal, S. S., Beaulieu, J. J., Pennino, M. J., and Welty, C.: Influence of infrastructure on water quality and greenhouse gas dynamics in urban streams, *Biogeosciences*, 14, 2831–2849, <https://doi.org/10.5194/bg-14-2831-2017>, 2017.
- 1585 Soja, G., Kitzler, B., and Soja, A.-M.: Emissions of greenhouse gases from Lake Neusiedl, a shallow steppe lake in Eastern Austria, *Hydrobiologia*, 731, 125–138, <https://doi.org/10.1007/s10750-013-1681-8>, 2014.
- Soued, C. and Prairie, Y. T.: The carbon footprint of a Malaysian tropical reservoir: measured versus modelled estimates highlight the underestimated key role of downstream processes, *Biogeosciences*, 17, 515–527, <https://doi.org/10.5194/bg-17-515-2020>, 2020.
- 1590 Spawn, S., Dunn, S., Fiske, G., Natali, S., Schade, J., and Zimov, N.: Summer methane ebullition from a headwater catchment in Northeastern Siberia, *Inland Waters*, 5, 224–230, <https://doi.org/10.5268/IW-5.3.845>, 2015.
- Stanley, E. H.: BECWA gases, University of Wisconsin-Madison, USA, unpublished, 2020.
- Stets, E. G.: Synoptic carbon gas fluxes from streams, rivers, and lakes in the conterminous US, USGS Landcarbon, 2012 to 2014, U.S. Geological Survey data release, <https://doi.org/10.5066/F72B8W2C>, 2016.

- 1595 Striegl, R. G., Dornblaser, M. M., McDonald, C. P., Rover, J. R., and Stets, E. G.: Carbon dioxide and methane emissions from the Yukon River system, *Global Biogeochem. Cycles*, 26, GB0E05, <https://doi.org/10.1029/2012gb004306>, 2012.
- Sturm, K., Grinham, A., Werner, U., and Yuan, Z.: Sources and sinks of methane and nitrous oxide in the subtropical Brisbane River estuary, South East Queensland, Australia, *Estuarine Coastal Shelf Sci.*, 168, 10–21, <https://doi.org/10.1016/j.ecss.2015.11.002>, 2016.
- 1600 Taillardat, P., Bodmer, P., Deblois, C. P., Ponçot, A., Prijac, A., Riahi, K., Gandois, L., del Giorgio, P. A., Bourgault, M. A., Tremblay, A., and Garneau, M.: Carbon dioxide and methane dynamics in a peatland headwater stream: origin, processes and implications, Zenodo, <https://zenodo.org/record/6073957>, 2021.
- Taillardat, P., Bodmer, P., Deblois, C. P., Ponçot, A., Prijac, A., Riahi, K., Gandois, L., del Giorgio, P. A., Bourgault, M. A., Tremblay, A., and Garneau, M.: Carbon dioxide and methane dynamics in a peatland headwater stream: Origins, processes and implications, *J. Geophys. Res. Biogeosci.*, e2022JG006855, <https://doi.org/10.1029/2022JG006855>, 2022.
- 1605 Taniwaki, R. H.: Methane concentrations and fluxes in agricultural and preserved tropical headwater streams, Environmental Data Initiative, <https://doi.org/10.6073/pasta/82b1aa19ad5b4ef88939303f79c1e74c>, 2022.
- Taniwaki, R. H., Cunha, D. G. F., Bento, C. B., Martinelli, L. A., Stanley, E. H., Filoso, S., Ferreira, M. de S., França, M. V., Ribeiro Júnior, J. W., Schiesari, L. C., and do Carmo, J. B.: Methane concentrations and fluxes in agricultural and preserved tropical headwater streams, *Sci. Total Environ.*, 844, 157238, <https://doi.org/10.1016/j.scitotenv.2022.157238>, 2022.
- 1610 Teodoru, C. R., Nyoni, F. C., Borges, A. V., Darchambeau, F., Nyambe, I., and Bouillon, S.: Dynamics of greenhouse gases (CO₂, CH₄, N₂O) along the Zambezi River and major tributaries, and their importance in the riverine carbon budget, *Biogeosciences*, 12, 2431–2453, <https://doi.org/10.5194/bg-12-2431-2015>, 2015.
- Trimmer, M., Hildrew, A. G., Jackson, M. C., Pretty, J. L., and Grey, J.: Evidence for the role of methane-derived carbon in a free-flowing, lowland river food web, *Limnol. Oceanogr.*, 54, 1541–1547, <https://doi.org/10.4319/lo.2009.54.5.1541>, 2009.
- 1615 Upstill-Goddard, R. C., Barnes, J., Frost, T., Punshon, S., and Owens, N. J. P.: Methane in the southern North Sea: Low-salinity inputs, estuarine removal, and atmospheric flux, *Global Biogeochem. Cycles*, 14, 1205–1217, <https://doi.org/10.1029/1999GB001236>, 2000.
- Upstill-Goddard, R. C., Salter, M. E., Mann, P. J., Barnes, J., Poulsen, J., Dinga, B., Fiske, G. J., and Holmes, R. M.: The riverine source of CH₄ and N₂O from the Republic of Congo, western Congo Basin, *Biogeosciences*, 14, 2267–2281, <https://doi.org/10.5194/bg-14-2267-2017>, 2017.
- 1620

- Vermaat, J. E., Hellmann, F., Dias, A. T. C., Hoorens, B., van Logtestijn, R. S. P., and Aerts, R.: Greenhouse gas fluxes from Dutch peatland water bodies: Importance of the surrounding landscape, *Wetlands*, 31, 493–498, <https://doi.org/10.1007/s13157-011-0170-y>, 2011.
- 1625 Vidon, P. and Serchan, S.: Impact of stream geomorphology on greenhouse gas concentration in a New York Mountain stream, *Water Air Soil Pollut.*, 227, 428, <https://doi.org/10.1007/s11270-016-3131-5>, 2016.
- Villa, J. A., Ju, Y., Smith, G. J., Angle, J. C., Renteria, L., Arntzen, E., Harding, S. F., Stegen, J. C., Wrighton, K. C., and Bohrer, G.: Chamber flux and porewater concentration of CH₄, CO₂ and N₂O, 2018, Columbia River bank at the Hanford site, WA, USA, ESS-DIVE Data Archive, <https://doi.org/10.15485/1595105>, 2020a.
- 1630 Villa, J. A., Smith, G. J., Ju, Y., Renteria, L., Angle, J. C., Arntzen, E., Harding, S. F., Ren, H., Chen, X., Sawyer, A. H., Graham, E. B., Stegen, J. C., Wrighton, K. C., and Bohrer, G.: Methane and nitrous oxide porewater concentrations and surface fluxes of a regulated river, *Sci. Total Environ.*, 715, 136920, <https://doi.org/10.1016/j.scitotenv.2020.136920>, 2020b.
- Vorobyev, S. N., Karlsson, J., Kolesnichenko, Y. Y., Korets, M. A., and Pokrovsky, O. S.: Fluvial carbon dioxide emission from the Lena River basin during the spring flood, *Biogeosciences*, 18, 4919–4936, <https://doi.org/10.5194/bg-18-4919-2021>,
1635 2021.
- Wallin, M. B.: Krycklan catchment time series, Swedish University of Agricultural Sciences, Uppsala, unpublished, 2021.
- Wallin, M. B., Löfgren, S., Erlandsson, M., and Bishop, K.: Representative regional sampling of carbon dioxide and methane concentrations in hemiboreal headwater streams reveal underestimates in less systematic approaches: CO₂ and CH₄ in headwater streams, *Global Biogeochem. Cycles*, 28, 465–479, <https://doi.org/10.1002/2013GB004715>, 2014.
- 1640 Wallin, M. B., Campeau, A., Audet, J., Bastviken, D., Bishop, K., Kocic, J., Laudon, H., Lundin, E., Löfgren, S., Natchimuthu, S., Sobek, S., Teutschbein, C., Weyhenmeyer, G. A., and Grabs, T.: Carbon dioxide and methane emissions of Swedish low-order streams—a national estimate and lessons learnt from more than a decade of observations: Carbon dioxide and methane emissions, *Limnol. Oceanogr. Lett.*, 3, 156–167, <https://doi.org/10.1002/lol2.10061>, 2018.
- Wang, D., Chen, Z., Sun, W., Hu, B., and Xu, S.: Methane and nitrous oxide concentration and emission flux of Yangtze Delta
1645 plain river net, *Sci. China Ser. B-Chem.*, 52, 652–661, <https://doi.org/10.1007/s11426-009-0024-0>, 2009.
- Wang, G., Xia, X., Liu, S., Zhang, L., Zhang, S., Wang, J., Xi, N., and Zhang, Q.: Intense methane ebullition from urban inland waters and its significant contribution to greenhouse gas emissions, *Water Research*, 189, 116654, <https://doi.org/10.1016/j.watres.2020.116654>, 2021.

- Wang, R., Zhang, H., Zhang, W., Zheng, X., Butterbach-Bahl, K., Li, S., and Han, S.: An urban polluted river as a significant hotspot for water–atmosphere exchange of CH₄ and N₂O, *Environ. Pollut.*, 264, 114770, <https://doi.org/10.1016/j.envpol.2020.114770>, 2020.
- Wang, X., He, Y., Chen, H., Yuan, X., Peng, C., Yue, J., Zhang, Q., and Zhou, L.: CH₄ concentrations and fluxes in a subtropical metropolitan river network: Watershed urbanization impacts and environmental controls, *Sci. Total Environ.*, 622–623, 1079–1089, <https://doi.org/10.1016/j.scitotenv.2017.12.054>, 2018.
- 1655 Webb, J. R., Santos, I. R., Maher, D. T., Macdonald, B., Robson, B., Isaac, P., and McHugh, I.: Terrestrial versus aquatic carbon fluxes in a subtropical agricultural floodplain over an annual cycle, *Agric. For. Meteorol.*, 260–261, 262–272, <https://doi.org/10.1016/j.agrformet.2018.06.015>, 2018.
- Wendt, A. K., Sowers, T., Hynek, S., Lemon, J., Beddings, E., Zheng, G., Li, Z., Williams, J. Z., and Brantley, S. L.: Scientist-nonscientist teams explore methane sources in streams near oil/gas development, *J. Contemp. Water Res. Educ.*, 164, 80–111, 1660 <https://doi.org/10.1111/j.1936-704X.2018.03286.x>, 2018.
- Wilcock, R. J. and Sorrell, B. K.: Emissions of greenhouse gases CH₄ and N₂O from low-gradient streams in agriculturally developed catchments, *Water Air Soil Pollut.*, 188, 155–170, <https://doi.org/10.1007/s11270-007-9532-8>, 2008.
- Wilkinson, M., Dumontier, M., Aalbersberg, I., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten J.-W., Bonino da Silva Santos, L., Bourne, P. E., Bouwman, J., Brookes, A. J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, 1665 C. T., Finkers, R., Gonzalez-Beltran, A., Gray, A. J. G., Groth, P., Goble, C., Grethe, J. S., Heringa, J., ‘t Hoen, P. A. C., Hooft, R., Kuhn, T., Kok, R., Kok, J., Lusher, S. J., Martone, M. E., Mons, A., Packer, A. L., Persson, B., Rocca-Serra, P., Roos, M., van Schaik, R., Sansone S.-A., Schultes, E., Sengstag, T., Slater, T., Strawn, G., Swertz, M. A., Thompson, M., van der Lei, J., van Mulligen, E., Veltrerop, J. Waangmeester, A., Wittenburg, P., Wolstencroft, K., Zhao, J. and Mons, B.: The FAIR Guiding Principles for scientific data management and stewardship. *Sci Data*, 3, 160018. [https://doi-
org.ezproxy.library.wisc.edu/10.1038/sdata.2016.18](https://doi-org.ezproxy.library.wisc.edu/10.1038/sdata.2016.18), 2016.
- 1670 Wilkniss, P. E., Lamontagne, R. A., Larson, R. E., and Swinnerton, J. W.: Atmospheric trace gases and land and sea breezes at the Sepik River coast of Papua New Guinea, *J. Geophys. Res.*, 83, 3672-3674, <https://doi.org/10.1029/jc083ic07p03672>, 1978.
- Woda, J., Wen, T., Lemon, J., Marcon, V., Keepports, C. M., Zelt, F., Steffy, L. Y., and Brantley, S. L.: Methane concentrations in streams reveal gas leak discharges in regions of oil, gas, and coal development, *Sci. Total Environ.*, 737, 140105, <https://doi.org/10.1016/j.scitotenv.2020.140105>, 2020.

- Wu, H., Zhao, Q., Gao, Q., Li, Y., Wan, Y., Li, Y., Tian, D., Liao, Y., Fan, M., Ganjurjav, H. Hu, G., Wang, B., Chen, X., and Qin, X.: Human activities inducing high CH₄ diffusive fluxes in an agricultural river catchment in subtropical China, *Sustainability*, 12, 2114, <https://doi.org/10.3390/su12052114>, 2020.
- 1680 Wu, S., Li, S., Zou, Z., Hu, T., Hu, Z., Liu, S., and Zou, J.: High methane emissions largely attributed to ebullitive fluxes from a subtropical river draining a rice paddy watershed in China, *Environ. Sci. Technol.*, 53, 3499–3507, <https://doi.org/10.1021/acs.est.8b05286>, 2019.
- 1685 Wu, W.: Characteristics and influencing factors of greenhouse gas emissions from water bodies of Zhongtianshe River Basin in Tianmu Lake area, M.S. thesis, Nanjing Normal University, Nanjing, China, doi (China): 10.27245/d.cnki.gnjsu.2020.001588, <https://kns.cnki.net/KCMS/detail/detail.aspx?dbname=CMFD202101&filename=1021516788.nh>, 2020.
- Wu, Y., Li, H., Chen, W.: Effects and emission characteristics of greenhouse gases from Wenyu River in summer, *Environ. Sci. Technol. (China)*, doi (China): 10.3969/j.issn.1003-6504.2016.05.002, 39, 8-16, 2016.
- 1690 Wynn, P.: Direct isotopic evidence of biogenic methane production and efflux from beneath a temperate glacier, Lancaster University, <https://doi.org/10.17635/lancaster/researchdata/246>, 2018b.
- Xiao, Q., Hu, Z., Hu, C., Islam, A. R. M. T., Bian, H., Chen, S., Liu, C., and Lee, X.: A highly agricultural river network in Jurong Reservoir watershed as significant CO₂ and CH₄ sources, *Sci. Total Environ.*, 769, 144558, <https://doi.org/10.1016/j.scitotenv.2020.144558>, 2021.
- 1695 Xu, H., Chen, M., Xiao, S., Yu, Z., and Zheng, X.: Temporal and spatial pattern of dissolved methane concentration in the river of a small karst watershed in Western Hubei, China *Environ. Sci.*, <https://kns.cnki.net/kcms/detail/11.2201.x.20210408.0910.002.html>, 2020.
- 1700 Yang, L.: Contrasting methane emissions from upstream and downstream rivers and their associated subtropical reservoir in eastern China, *Sci. Rep.*, 9, 8072, <https://doi.org/10.1038/s41598-019-44470-2>, 2019.
- Yang, L., Li, X., Yan, W., Ma, P., and Wang, J.: CH₄ concentrations and emissions from three rivers in the Chaohu Lake watershed in southeast China, *J. Integr. Agric.*, 11, 665–673, [https://doi.org/10.1016/S2095-3119\(12\)60054-9](https://doi.org/10.1016/S2095-3119(12)60054-9), 2012.
- Yang, S.-S., Chen, I.-C., Ching-Pao, L., Liu, L.-Y., and Chang, C.-H.: Carbon dioxide and methane emissions from Tanswei River in Northern Taiwan, *Atmos. Pollut. Res.*, 6, 52–61, <https://doi.org/10.5094/APR.2015.007>, 2015.

- 1705 Yavitt, J. B., Lang, G. E., and Sexstone, A. J.: Methane fluxes in wetland and forest soils, beaver ponds, and low-order streams of a temperate forest ecosystem, *J. Geophys. Res.*, 95, 22463, <https://doi.org/10.1029/JD095iD13p22463>, 1990.
- Ye, R., Wu, Q., Zhao, Z., Hu, J., Cui, L., and Ding, H.: Concentrations and emissions of dissolved CH₄ and N₂O in the Yarlung Tsangpo River, *Chinese J. Ecol.*, 38, 791–798, doi (China): 10.13292/j.1000-4890.201903.001, 2019.
- Yu, Z., Wang, D., Li, Y., Deng, H., Hu, B., Ye, M., Zhou, X., Da, L., Chen, Z., and Xu, S.: Carbon dioxide and methane dynamics in a human-dominated lowland coastal river network (Shanghai, China): CO₂ and CH₄ in Shanghai River Network, *J. Geophys. Res. Biogeosci.*, 122, 1738–1758, <https://doi.org/10.1002/2017JG003798>, 2017.
- 1710 Zhang, F.: Spatio-temporal dynamics of dissolved concentrations of N₂O and CH₄ and their emission flux in Weihe river of Xinxiang, M.S. thesis, Henan Normal University, 81 pp., Xinxiang, China, <https://kns.cnki.net/KCMS/detail/detail.aspx?dbname=CMFD201701&filename=1016231948.nh>, 2016.
- 1715 Zhang, L., Xia, X., Liu, S., Zhang, S., Li, S., Wang, J., Wang, G., Gao, H., Zhang, Z., Wang, Q., Wen, W., Liu, R., Yang, Z., Stanley, E. H., and Raymond, P. A.: Significant methane ebullition from alpine permafrost rivers on the East Qinghai–Tibet Plateau, *Nat. Geosci.*, 13, 349–354, <https://doi.org/10.1038/s41561-020-0571-8>, 2020.
- Zhang, L., Xia, X., Liu, S., Yang, Z., Stanley, E. H., and Raymond, P. A.: A dataset for methane concentrations and fluxes for alpine permafrost streams and rivers on the East Qinghai-Tibet Plateau, Environmental Data Initiative, <https://doi.org/10.6073/pasta/3e9ed02d7d89a31eba4c36481255084c>, 2021a.
- 1720 <https://doi.org/10.6073/pasta/3e9ed02d7d89a31eba4c36481255084c>, 2021a.
- Zhang, L., Xia, X., Battin, T. J., and Stanley, E. H.: Nitrous oxide dataset for East Qinghai-Tibet Plateau waterways, Environmental Data Initiative, <https://doi.org/10.6073/pasta/ba9340800403c450e7d942d450237dc4>, 2021b.
- Zhang, W., Li, H., Pueppke, S. G., and Pang, J.: Restored riverine wetlands in a headwater stream can simultaneously behave as sinks of N₂O and hotspots of CH₄ production, *Environ. Pollut.*, 284, 117114, <https://doi.org/10.1016/j.envpol.2021.117114>, 2021a.
- 1725 2021a.
- Zhang, W., Li, H., Xiao, Q., and Li, X.: Urban rivers are hotspots of riverine greenhouse gas (N₂O, CH₄, CO₂) emissions in the mixed-landscape chaohu lake basin, *Water Res.*, 189, 116624, <https://doi.org/10.1016/j.watres.2020.116624>, 2021b.
- Zhang, Y., Kang, S., Wei, D., Luo, X., Wang, Z., and Gao, T.: Sink or source? Methane and carbon dioxide emissions from cryoconite holes, subglacial sediments, and proglacial river runoff during intensive glacier melting on the Tibetan Plateau, *Fundam. Res.*, 1, 232–239, <https://doi.org/10.1016/j.fmre.2021.04.005>, 2021.
- 1730 Fundam. Res., 1, 232–239, <https://doi.org/10.1016/j.fmre.2021.04.005>, 2021.

Zhao, J.: Distribution and fluxes of methane and nitrous oxide in the Changjiang (Yangtze River) and its estuary, M.S. thesis, Ocean University of China, Qingdao, China, 81 pp., <https://kns.cnki.net/KCMS/detail/detail.aspx?dbname=CMFD2012&filename=1011230510.nh>, 2011.

1735 Zhao, J., Zhang, G.-L., Wu, Y., and Yang, J.: Distribution and emission of methane from the Changjiang, Environ. Sci., 32, 18–25, doi (China): 10.13227/j.hjcx.2011.01.003, 2011.

Zhao, X.: Study on methane and nitrous oxide emissions from polluted water in Nanjing, M.S. thesis, Nanjing University of Information Science and Technology, Nanjing, China, 55 pp., <https://kns.cnki.net/KCMS/detail/detail.aspx?dbname=CMFD2012&filename=1012369155.nh>, 2012.

1740 Zolkos, S., Tank, S. E., Striegl, R. G., and Kokelj, S. V.: Thermokarst effects on carbon dioxide and methane fluxes in streams on the Peel Plateau (NWT, Canada), J. Geophys. Res. Biogeosci., 124, 1781–1798, <https://doi.org/10.1029/2019JG005038>, 2019.

1745

Supplement.

1750 The supplement related to this article is available at <https://doi.org/10.5194/essd-2022-346-supplement>

Author contributions.

EHS conceived of the project idea, and led data entry, manuscript preparation, and data curation. LCL developed the code used for unit conversions, was responsible for data conversion and QA/QC, and contributed to data visualization, data analysis, and code curation. GRR was responsible for spatial analyses, and contributed to data visualization, code curation, and manuscript preparation. The structure and composition of the manuscript were resulted from collaborative discussions among EHS, GRR, LCL, NJC, SKO, and RAS. All authors contributed to data acquisition, data entry, data checking, and substantial manuscript revising and editing. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

1760

Competing Interests.

The authors declare that they have no conflict of interest.

Acknowledgements.

1765 We are grateful to all the authors who generously shared data or made their data public; our science is better or this transparency. We are also grateful to several of these investigators for patiently and thoughtfully responding to emails from E. Stanley. Corinna Gries provided advice and assistance with data publication. Stanley thanks K. Forest and K. McMahon for guidance during manuscript preparation. Great thanks to M. W. Diebel, S. N.Topp, Y. Yao, K. Castro-Morales, and B. Deemer for their thoughtful and detailed comments that improved this paper.

1770

Financial Support.

Emily H. Stanley acknowledges support from the National Science Foundation #DEB-2025982, NTL LTER. Nora J. Casson was supported in part by the Canada Research Chair program. Gerard Rocher-Ros was supported by the Swedish Research Council (2021-06667) and the Department of Ecology and Environmental Science, Umeå University. Marcus B. Wallin was supported by the Swedish Research Council's VR (2021-04058) and Formas (2019-01105). Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

1780 **References**

- Aho, K. S., Fair, J. H., Hosen, J. D., Kyzivat, E. D., Logozzo, L. A., Rocher-Ros, G., Weber, L. C., Yoon, B., and Raymond, P. A.: Distinct concentration-discharge dynamics in temperate streams and rivers: CO₂ exhibits chemostasis while CH₄ exhibits source limitation due to temperature control, *Limnol. Oceanogr.*, 66, 3656–3668, <https://doi.org/10.1002/lno.11906>, 2021.
- 1785 Allen, G. H. and Pavelsky, T. M.: Global extent of rivers and streams, *Science*, 361, 585–588, <https://doi.org/10.1126/science.aat0636>, 2018.
- Alshboul, Z., Encinas-Fernández, J., Hofmann, H., and Lorke, A.: Export of dissolved methane and carbon dioxide with effluents from municipal wastewater treatment plants, *Environ. Sci. Technol.*, 50, 5555–5563, <https://doi.org/10.1021/acs.est.5b04923>, 2016.
- 1790 Anthony, S. E., Prahl, F. G., and Peterson, T. D.: Methane dynamics in the Willamette River, Oregon, *Limnol. Oceanogr.*, 57, 1517–1530, <https://doi.org/10.4319/lo.2012.57.5.1517>, 2012.
- 1795 Attermeyer, K., Casas-Ruiz, J. P., Fuss, T., Pastor, A., Cauvy-Fraunié, S., Sheath, D., Nydahl, A. C., Doretto, A., Portela, A. P., Doyle, B. C., Simov, N., Gutmann Roberts, C., Niedrist, G. H., Timoner, X., Evtimova, V., Barral-Fraga, L., Bašić, T., Audet, J., Deininger, A., Busst, G., Fenoglio, S., Catalán, N., de Eyto, E., Pilotto, F., Mor, J.-R., Monteiro, J., Fletcher, D., Noss, C., Colls, M., Nagler, M., Liu, L., Romero González-Quijano, C., Romero, F., Pansch, N., Ledesma, J. L. J., Pegg, J., Klaus, M., Freixa, A., Herrero Ortega, S., Mendoza-Lera, C., Bednařík, A., Fonvielle, J. A., Gilbert, P. J., Kenderov, L. A., Rulík, M., and Bodmer, P.: Carbon dioxide fluxes increase from day to night across European streams, *Commun. Earth Environ.*, 2, 118, <https://doi.org/10.1038/s43247-021-00192-w>, 2021.
- 1800 Barbosa, P. M., Farjalla, V. F., Melack, J. M., Amaral, J. H. F., da Silva, J. S., and Forsberg, B. R.: High rates of methane oxidation in an Amazon floodplain lake, *Biogeochemistry*, 137, 351–365, <https://doi.org/10.1007/s10533-018-0425-2>, 2018.
- Bastviken, D., Tranvik, L. J., Downing, J. A., Crill, P. M., and Enrich-Prast, A.: Freshwater Methane emissions offset the continental carbon sink, *Science*, 331, 50–50, <https://doi.org/10.1126/science.1196808>, 2011.
- 1810 Baulch, H. M., Dillon, P. J., Maranger, R., and Schiff, S. L.: Diffusive and ebullitive transport of methane and nitrous oxide from streams: Are bubble-mediated fluxes important?, *J. Geophys. Res.*, 116, G04028, <https://doi.org/10.1029/2011JG001656>, 2011.

- 1815 Bednařík, A., Blaser, M., Matoušů, A., Hekera, P., and Rulík, M.: Effect of weir impoundments on methane dynamics in a river, *Sci. Total Environ.*, 584–585, 164–174, <https://doi.org/10.1016/j.scitotenv.2017.01.163>, 2017.
- 1820 Begum, M. S., Bogard, M. J., Butman, D. E., Chea, E., Kumar, S., Lu, X., Nayna, O. K., Ran, L., Richey, J. E., Tareq, S. M., Xuan, D. T., Yu, R., and Park, J.: Localized pollution impacts on greenhouse gas dynamics in three anthropogenically modified Asian river systems, *J. Geophys. Res. Biogeosci.*, 126, <https://doi.org/10.1029/2020JG006124>, 2021.
- Benjamini, Y. and Hochberg, Y. Controlling the false discovery rate: A practical and powerful approach to multiple testing. *J. R. Statist. Soc. B*, 57, 289–300. <https://doi.org/10.1111/j.2517-6161.1995.tb02031.x>, 1995.
- 1825 Billett, M. F. and Harvey, F. H.: Measurements of CO₂ and CH₄ evasion from UK peatland headwater streams, *Biogeochemistry*, 114, 165–181, <https://doi.org/10.1007/s10533-012-9798-9>, 2013.
- Bodmer, P., Vroom, R., Stepina, T., del Giorgio, P., and Kosten, S.: Methane fluxes of vegetated areas in natural freshwater ecosystems: assessments and global significance, *Earth ArXiv* <https://doi.org/10.31223/X5ND0F>, 5 December 2021.
- 1830 Borges, A. V., Darchambeau, F., Lambert, T., Bouillon, S., Morana, C., Brouyère, S., Hakoun, V., Jurado, A., Tseng, H.-C., Descy, J.-P., and Roland, F. A. E.: Effects of agricultural land use on fluvial carbon dioxide, methane and nitrous oxide concentrations in a large European river, the Meuse (Belgium), *Sci. Total Environ.*, 610–611, 342–355, <https://doi.org/10.1016/j.scitotenv.2017.08.047>, 2018b.
- 1835 Bouillon, S., Yambélé, A., Spencer, R. G. M., Gillikin, D. P., Hernes, P. J., Six, J., Merckx, R., and Borges, A. V.: Organic matter sources, fluxes and greenhouse gas exchange in the Oubangui River (Congo River basin), *Biogeosciences*, 9, 2045–2062, <https://doi.org/10.5194/bg-9-2045-2012>, 2012.
- 1840 Bretz, K. A., Jackson, A. R., Rahman, S., Monroe, J. M., and Hotchkiss, E. R.: Integrating ecosystem patch contributions to stream corridor carbon dioxide and methane fluxes, *J. Geophys. Res. Biogeosci.*, 126, <https://doi.org/10.1029/2021JG006313>, 2021.
- 1845 Burns, R., Wynn, P. M., Barker, P., McNamara, N., Oakley, S., Ostle, N., Stott, A. W., Tuffen, H., Zhou, Z., Tweed, F. S., Chesler, A. and Stuart, M.: Direct isotopic evidence of biogenic methane production and efflux from beneath a temperate glacier, *Sci. Rep.*, 8, 17118. <https://doi.org/10.1038/s41598-018-35253-2>. 2018.

- Bussmann, I., Koedel, U., Schütze, C., Kamjunke, N., and Koschorreck, M.: Spatial variability and hotspots of methane concentrations in a large temperate river, *Front. Environ. Sci.*, 10, 833936, <https://doi.org/10.3389/fenvs.2022.833936>, 2022.
- 1850 Call, M., Sanders, C. J., Enrich-Prast, A., Sanders, L., Marotta, H., Santos, I. R., and Maher, D. T.: Radon-traced pore-water as a potential source of CO₂ and CH₄ to receding black and clear water environments in the Amazon Basin, *Limnol. Oceanogr. Lett.*, 3, 375–383, <https://doi.org/10.1002/lol2.10089>, 2018.
- Campeau, A., Lapierre, J.-F., Vachon, D., and del Giorgio, P. A.: Regional contribution of CO₂ and CH₄ fluxes from the fluvial network in a lowland boreal landscape of Québec, *Global Biogeochem. Cycles*, 28, 57–69, 1855 <https://doi.org/10.1002/2013GB004685>, 2014.
- Campeau, A., Bishop, K., Nilsson, M. B., Klemedtsson, L., Laudon, H., Leith, F. I., Öquist, M., and Wallin, M. B.: Stable carbon isotopes reveal soil-stream DIC linkages in contrasting headwater catchments, *J. Geophys. Res. Biogeosci.*, 123, 149–167, <https://doi.org/10.1002/2017JG004083>, 2018. 1860
- Castro-Morales, K., Canning, A., Körtzinger, A., Göckede, M., Küsel, K., Overholt, W. A., Wichard, T., Redlich, S., Arzberger, S., Kolle, O., and Zimov, N.: Effects of reversal of water flow in an Arctic floodplain river on fluvial emissions of CO₂ and CH₄, *J. Geophys. Res. Biogeosci.*, 127, <https://doi.org/10.1029/2021JG006485>, 2022.
- 1865 Chen, S., Wang, D., Ding, Y., Yu, Z., Liu, L., Li, Y., Yang, D., Gao, Y., Tian, H., Cai, R., and Chen, Z.: Ebullition controls on CH₄ emissions in an urban, eutrophic river: A potential time-scale bias in determining the aquatic CH₄ flux, *Environ. Sci. Technol.*, 55, 7287–7298, <https://doi.org/10.1021/acs.est.1c00114>, 2021.
- 1870 Crawford, J. T., Stanley, E. H., Spawn, S. A., Finlay, J. C., Loken, L. C., and Striegl, R. G.: Ebullitive methane emissions from oxygenated wetland streams, *Global Change Biol.*, 20, 3408–3422, <https://doi.org/10.1111/gcb.12614>, 2014.
- Crawford, J. T., Dornblaser, M. M., Stanley, E. H., Clow, D. W., and Strieg, R. G.: Source limitation of carbon gas emissions in high-elevation mountain streams and lakes, *J. Geophys. Res. Biogeosci.*, 120, 952–64, <https://doi.org/10.1002/2014JG002861>, 2015. 1875
- Crawford, J. T., Loken, L. C., Stanley, E. H., Stets, E. G., Dornblaser, M. M., and Striegl, R. G.: Basin scale controls on CO₂ and CH₄ emissions from the Upper Mississippi River, *Geophys. Res. Lett.*, 43, 1973–1979, <https://doi.org/10.1002/2015GL067599>, 2016.

- 1880 Crawford, J. T., Loken, L. C., West, W. E., Crary, B., Spawn, S. A., Gubbins, N., Jones, S. E., Striegl, R. G., and Stanley, E. H.: Spatial heterogeneity of within-stream methane concentrations, *J. Geophys. Res.-Biogeosci.*, 122, 1036–1048, <https://doi.org/10.1002/2016JG003698>, 2017.
- de Angelis, M. A. and Lilley, M. D.: Methane in surface waters of Oregon estuaries and rivers, *Limnol. Oceanogr.*, 32, 716–
1885 722, <https://doi.org/10.4319/lo.1987.32.3.0716>, 1987.
- Dowle M, and Srinivasan A.: `_data.table`: Extension of ``data.frame``. R package version 1.14.2, <https://CRAN.R-project.org/package=data.table>, 2021.
- Fisher, S. G., Gray, L. J., Grimm, N. B., and Busch, D. E.: Temporal succession in a desert stream ecosystem following flash
1890 flooding, *Ecol.Monogr.*, 52, 93–110, <https://doi.org/10.2307/2937346>, 1982.
- Flecker, A. S., Shi, Q., Almeida, R. M., Angarita, H., Gomes-Selman, J. M., García-Villacorta, R., Sethi, S. A., Thomas, S. A., Poff, N. L., Forsberg, B. R., Heilpern, S. A., Hamilton, S. K., Abad, J. D., Anderson, E. P., Barros, N., Bernal, I. C., Bernstein, R., Cañas, C. M., Dangles, O., Encalada, A. C., Fleischmann, A. S., Goulding, M., Higgins, J., Jézéquel, C., Larson,
1895 E. I., McIntyre, P. B., Melack, J. M., Montoya, M., Oberdorff, T., Paiva, R., Perez, G., Rappazzo, B. H., Steinschneider, S., Torres, S., Varese, M., Walter, M. T., Wu, X., Xue, Y., Zapata-Ríos, X. E., and Gomes, C. P.: Reducing adverse impacts of Amazon hydropower expansion, *Science*, 375, 753–760, <https://doi.org/10.1126/science.abj4017>, 2022.
- Gatti, R. C., Callaghan, T. V., Rozhkova-Timina, I., Dudko, A., Lim, A., Vorobyev, S. N., Kirpotin, S. N., and Pokrovsky,
1900 O. S.: The role of Eurasian beaver (*Castor fiber*) in the storage, emission and deposition of carbon in lakes and rivers of the River Ob flood plain, western Siberia, *Sci. Total Environ.*, 644, 1371–1379, <https://doi.org/10.1016/j.scitotenv.2018.07.042>, 2018.
- Gómez-Gener, L., Hotchkiss, E. R., Laudon, H., and Sponseller, R. A.: Integrating discharge-concentration dynamics across
1905 carbon forms in a boreal landscape, *Water Resour. Res.*, 57, <https://doi.org/10.1029/2020WR028806>, 2021a.
- Gómez-Gener, L., Rocher-Ros, G., Battin, T., Cohen, M. J., Dalmagro, H. J., Dinsmore, K. J., Drake, T. W., Duvert, C., Enrich-Prast, A., Horgby, Å., Johnson, M. S., Kirk, L., Machado-Silva, F., Marzolf, N. S., McDowell, M. J., McDowell, W. H., Miettinen, H., Ojala, A. K., Peter, H., Pumpanen, J., Ran, L., Riveros-Iregui, D. A., Santos, I. R., Six, J., Stanley, E. H.,
1910 Wallin, M. B., White, S. A., and Sponseller, R. A.: Global carbon dioxide efflux from rivers enhanced by high nocturnal emissions, *Nat. Geosci.*, 14, 289–294, <https://doi.org/10.1038/s41561-021-00722-3>, 2021b.

- Gurnell, A. M., O'Hare, J. M., O'Hare, M. T., Dunbar, M. J., and Scarlett, P. M.: An exploration of associations between assemblages of aquatic plant morphotypes and channel geomorphological properties within British rivers, *Geomorphology*, 1915 116, 135–144, <https://doi.org/10.1016/j.geomorph.2009.10.014>, 2010.
- Hlaváčová, E., Rulík, M., Čáp, L., and Mach, V.: Greenhouse gas (CO₂, CH₄, N₂O) emissions to the atmosphere from a small lowland stream in Czech Republic, *Arch. Hydrobiol.*, 165, 339–353, <https://doi.org/10.1127/0003-9136/2006/0165-0339>, 2006.
- 1920 Ho, L., Jerves-Cobo, R., Barthel, M., Six, J., Bode, S., Boeckx, P., and Goethals, P.: Greenhouse gas dynamics in an urbanized river system: influence of water quality and land use, *Environ. Sci. Pollut. Res.*, 29, 37277–37290, <https://doi.org/10.1007/s11356-021-18081-2>, 2022.
- Hollister, J. W., Shah, T., Robitaille A, Beck M, Johnson M. “elevatr”. Access elevation data from various APIs, R package 1925 version 0.4.2, <https://doi.org/10.5281/zenodo.5809645>, 2021.
- Horton, R. E.: Erosional development of streams and their drainage basins; hydrophysical approach to quantitative morphology, *Geol. Soc. America Bull.*, 56, 275, [https://doi.org/10.1130/0016-7606\(1945\)56\[275:EDOSAT\]2.0.CO;2](https://doi.org/10.1130/0016-7606(1945)56[275:EDOSAT]2.0.CO;2), 1945.
- 1930 Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J. W., Schuur, E. A. G., Ping, C.-L., Schirmermeister, L., Grosse, G., Michaelson, G. J., Koven, C. D., O'Donnell, J. A., Elberling, B., Mishra, U., Camill, P., Yu, Z., Palmtag, J., and Kuhry, P.: Estimated stocks of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps, *Biogeosciences*, 11, 6573–6593, <https://doi.org/10.5194/bg-11-6573-2014>, 2014.
- 1935 Hughes, B. B., Beas-Luna, R., Barner, A. K., Brewitt, K., Brumbaugh, D. R., Cerny-Chipman, E. B., Close, S. L., Coblenz, K. E., de Nesnera, K. L., Drobnitch, S. T., Figurski, J. D., Focht, B., Friedman, M., Freiwald, J., Heady, K. K., Heady, W. N., Hettlinger, A., Johnson, A., Karr, K. A., Mahoney, B., Moritsch, M. M., Osterback, A.-M. K., Reimer, J., Robinson, J., Rohrer, T., Rose, J. M., Sabal, M., Segui, L. M., Shen, C., Sullivan, J., Zuercher, R., Raimondi, P. T., Menge, B. A., Grorud-Colvert, K., Novak, M., and Carr, M. H.: Long-term studies contribute disproportionately to ecology and policy, *BioScience*, 67, 271–
1940 281, <https://doi.org/10.1093/biosci/biw185>, 2017.
- Hutchins, R. H. S., Casas-Ruiz, J. P., Prairie, Y. T., and del Giorgio, P. A.: Magnitude and drivers of integrated fluvial network greenhouse gas emissions across the boreal landscape in Québec, *Water Res.*, 173, 115556, <https://doi.org/10.1016/j.watres.2020.115556>, 2020.

- IPCC: Climate Change: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R. and Zhou., B. (eds.). Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, doi:10.1017/9781009157896, 2021.
- 1950
- Jin, H., Yoon, T. K., Begum, M. S., Lee, E.-J., Oh, N.-H., Kang, N., and Park, J.-H.: Longitudinal discontinuities in riverine greenhouse gas dynamics generated by dams and urban wastewater, *Biogeosciences*, 15, 6349–6369, <https://doi.org/10.5194/bg-15-6349-2018>, 2018.
- 1955
- Kemenes, A., Forsberg, B. R., and Melack, J. M.: Methane release below a tropical hydroelectric dam, *Geophys. Res. Lett.*, 34, L12809, <https://doi.org/10.1029/2007gl029479>, 2007.
- Kling, G. W.: Biogeochemistry data set for Innvait Creek Weir on the North Slope of Alaska 2002-2018, Environmental Data Initiative, <https://doi.org/10.6073/pasta/733c73c6ebffeac6970b2b0f4dddfe6>, 2019a.
- 1960
- Kling, G.: Biogeochemistry data set for soil waters, streams, and lakes near Toolik on the North Slope of Alaska., Environmental Data Initiative, <https://doi.org/10.6073/pasta/574fd24522eee7a0c07fc260ccc0e2fa>, 2019b.
- Kling, G. 2022. Biogeochemistry data set for soil waters, streams, and lakes near Toolik Lake on the North Slope of Alaska, 2012 through 2020 ver 2. Environmental Data Initiative, <https://doi.org/10.6073/pasta/4e25db9ae9372f5339f2795792814845>, 2022.
- 1965
- Kokic, J., Sahlée, E., Sobek, S., Vachon, D., and Wallin, M. B.: High spatial variability of gas transfer velocity in streams revealed by turbulence measurements, *Inland Waters*, 8, 461–473, <https://doi.org/10.1080/20442041.2018.1500228>, 2018.
- 1970
- Kuhn, C., Bettigole, C., Glick, H. B., Seegmiller, L., Oliver, C. D., and Raymond, P.: Patterns in stream greenhouse gas dynamics from mountains to plains in northcentral Wyoming, *J. Geophys. Res. Biogeosci.*, 122, 2173–2190, <https://doi.org/10.1002/2017jg003906>, 2017.
- 1975
- Lamarche-Gagnon, G., Wadham, J. L., Sherwood Lollar, B., Arndt, S., Fietzek, P., Beaton, A. D., Tedstone, A. J., Telling, J., Bagshaw, E. A., Hawkings, J. R., Kohler, T. J., Zarsky, J. D., Mowlem, M. C., Anesio, A. M., and Stibal, M.: Greenland melt drives continuous export of methane from the ice-sheet bed, *Nature*, 565, 73–77, <https://doi.org/10.1038/s41586-018-0800-0>, 2019.

- 1980 Leng, P., Kamjunke, N., Li, F., and Koschorreck, M.: Temporal patterns of methane emissions from two streams with different riparian connectivity, *J. Geophys. Res. Biogeosci.*, 126, e2020JG006104, <https://doi.org/10.1029/2020jg006104>, 2021.
- 1985 Li, M., Peng, C., Zhang, K., Xu, L., Wang, J., Yang, Y., Li, P., Liu, Z., and He, N.: Headwater stream ecosystem: an important source of greenhouse gases to the atmosphere. *Water Research*, 190, 116738, <https://doi.org/10.1016/j.watres.2020.116738>, 2021.
- Lian, X., Piao, S., Chen, A., Huntingford, C., Fu, B., Li, L. Z. X., Huang, J., Sheffield, J., Berg, A. M., Keenan, T. F., McVicar, T. R., Wada, Y., Wang, X., Wang, T., Yang, Y., and Roderick, M. L.: Multifaceted characteristics of dryland aridity changes in a warming world, *Nat. Rev. Earth Environ.*, 2, 232–250, <https://doi.org/10.1038/s43017-021-00144-0>, 2021.
- 1990 Lamontagne, R. A., Swinnerton, J. W., Linnenbom, V. J., and Smith, W. D.: Methane concentrations in various marine environments, *J. Geophys. Res.*, 78, 5317–5324, <https://doi.org/10.1029/JC078i024p05317>, 1973.
- Linke, S., Lehner, B., Ouellet Dallaire, C., Ariwi, J., Grill, G., Anand, M., Beames, P., Burchard-Levine, V., Maxwell, S., Moidu, H., Tan, F., and Thieme, M.: Global hydro-environmental sub-basin and river reach characteristics at high spatial resolution, *Sci. Data*, 6, 283, <https://doi.org/10.1038/s41597-019-0300-6>, 2019.
- 1995 Liu, S., Kuhn, C., Amatulli, G., Aho, K., Butman, D. E., Allen, G. H., Lin, P., Pan, M., Yamazaki, D., Brinkerhoff, C., Gleason, C., Xia, X., and Raymond, P. A.: The importance of hydrology in routing terrestrial carbon to the atmosphere via global streams and rivers, *Proc. Natl. Acad. Sci. U.S.A.*, 119, e2106322119, <https://doi.org/10.1073/pnas.2106322119>, 2022.
- 2000 Loken, L. C., Crawford, J. T., Stanley, E. H., Butman, D., and Striegl, R.: Columbia River spatial water chemistry, Environmental Data Initiative, <https://doi.org/10.6073/pasta/e881070c9e8f6b7f774d3c65b27a9f69>, 2018.
- 2005 Lorke, A., Bodmer, P., Noss, C., Alshboul, Z., Koschorreck, M., Somlai-Haase, C., Bastviken, D., Flury, S., McGinnis, D. F., Maeck, A., Müller, D., and Premke, K.: Technical note: drifting versus anchored flux chambers for measuring greenhouse gas emissions from running waters, *Biogeosciences*, 12, 7013–7024, <https://doi.org/10.5194/bg-12-7013-2015>, 2015.
- Lupon, A., Denfeld, B. A., Laudon, H., Leach, J., Karlsson, J., and Sponseller, R. A.: Groundwater inflows control patterns and sources of greenhouse gas emissions from streams, *Limnol. Oceanogr.*, 64, 1545–1557, <https://doi.org/10.1002/lno.11134>, 2019.
- 2010

- 2015 Metcalfe, D. B., Hermans, T. D. G., Ahlstrand, J., Becker, M., Berggren, M., Björk, R. G., Björkman, M. P., Blok, D., Chaudhary, N., Chisholm, C., Classen, A. T., Hasselquist, N. J., Jonsson, M., Kristensen, J. A., Kumordzi, B. B., Lee, H., Mayor, J. R., Prevéy, J., Pantazatou, K., Rousk, J., Sponseller, R. A., Sundqvist, M. K., Tang, J., Uddling, J., Wallin, G., Zhang, W., Ahlström, A., Tenenbaum, D. E., and Abdi, A. M.: Patchy field sampling biases understanding of climate change impacts across the Arctic, *Nat. Ecol. Evol.*, 2, 1443–1448, <https://doi.org/10.1038/s41559-018-0612-5>, 2018.
- NOAA, Increase in atmospheric methane set another record during 2021, <https://www.noaa.gov/news-release/increase-in-atmospheric-methane-set-another-record-during-2021>, last accessed 20 Sept 2022.
- 2020 Park, J.-H., Nayna, O. K., Begum, M. S., Chea, E., Hartmann, J., Keil, R. G., Kumar, S., Lu, X., Ran, L., Richey, J. E., Sarma, V. V. S. S., Tareq, S. M., Xuan, D. T., and Yu, R.: Reviews and syntheses: Anthropogenic perturbations to carbon fluxes in Asian river systems – concepts, emerging trends, and research challenges, *Biogeosciences*, 15, 3049–3069, <https://doi.org/10.5194/bg-15-3049-2018>, 2018.
- 2025 Peacock, M., Audet, J., Bastviken, D., Futter, M. N., Gauci, V., Grinham, A., Harrison, J. A., Kent, M. S., Kosten, S., Lovelock, C. E., Veraart, A. J., and Evans, C. D.: Global importance of methane emissions from drainage ditches and canals, *Environ. Res. Lett.*, 16, 044010, <https://doi.org/10.1088/1748-9326/abeb36>, 2021.
- 2030 Pebesma, E.: Simple features for R: Standardized support for spatial vector data, *The R Journal*, 10, 439, <https://doi.org/10.32614/RJ-2018-009>, 2018.
- Pedersen, T. L. Patchwork: the composer of plots. R package version 1.1.1, <https://patchwork.data-imaginist.com/> 2020.
- 2035 R Core Team: R: A language and environment for statistical computing, 2021.
- Ran, L., Shi, H., and Yang, X.: Magnitude and drivers of CO₂ and CH₄ emissions from an arid/semiarid river catchment on the Chinese Loess Plateau, *J. Hydrol.*, 598, 126260, <https://doi.org/10.1016/j.jhydrol.2021.126260>, 2021.
- 2040 Raymond, P. A., Zappa, C. J., Butman, D., Bott, T. L., Potter, J., Mulholland, P., Laursen, A. E., McDowell, W. H., and Newbold, D.: Scaling the gas transfer velocity and hydraulic geometry in streams and small rivers, *Limnol. Oceanogr. Fluids, Environ.*, 2, 41–53, <https://doi.org/10.1215/21573689-1597669>, 2012.

- 2045 Riis, T. and Biggs, B. J. F.: Hydrologic and hydraulic control of macrophyte establishment and performance in streams, *Limnol. Oceanogr.*, 48, 1488–1497, <https://doi.org/10.4319/lo.2003.48.4.1488>, 2003.
- Robison, A. L., Wollheim, W. M., Turek, B., Bova, C., Snay, C., and Varner, R. K.: Spatial and temporal heterogeneity of methane ebullition in lowland headwater streams and the impact on sampling design, *Limnol. Oceanogr.*, 66, 4063–4076, <https://doi.org/10.1002/lno.11943>, 2021.
- 2050 Rocher-Ros, G., Stanley, E. H., Loken, L. C., Casson, N. J., Raymond, P. A., Liu, S., Amantulli, G., and Sponseller, R. A. Global methane emissions from running waters, accepted.
- Rosentreter, J. A., Borges, A. V., Deemer, B. R., Holgerson, M. A., Liu, S., Song, C., Melack, J., Raymond, P. A., Duarte, C. M., Allen, G. H., Olefeldt, D., Poulter, B., Battin, T. I., and Eyre, B. D.: Half of global methane emissions come from highly variable aquatic ecosystem sources, *Nat. Geosci.*, 14, 225–230, <https://doi.org/10.1038/s41561-021-00715-2>, 2021.
- 2055 Rovelli, L., Olde, L. A., Heppell, C. M., Binley, A., Yvon-Durocher, G., Glud, R. N., and Trimmer, M.: Contrasting biophysical controls on carbon dioxide and methane outgassing from streams, *J. Geophys. Res. Biogeophys.*, 127, <https://doi.org/10.1029/2021JG006328>, 2022.
- 2060 Sabo, J. L., Sinha, T., Bowling, L. C., Schoups, G. H. W., Wallender, W. W., Campana, M. E., Cherkauer, K. A., Fuller, P. L., Graf, W. L., Hopmans, J. W., Kominoski, J. S., Taylor, C., Trimble, S. W., Webb, R. H., and Wohl, E. E.: Reclaiming freshwater sustainability in the Cadillac Desert, *Proc. Natl. Acad. Sci. U.S.A.*, 107, 21263–21269, <https://doi.org/10.1073/pnas.1009734108>, 2010.
- 2065 Sanders, I. A., Heppell, C. M., Cotton, J. A., Wharton, G., Hildrew, A. G., Flowers, E. J., and Trimmer, M.: Emission of methane from chalk streams has potential implications for agricultural practices, *Freshwater Biol.*, 52, 1176–1186, <https://doi.org/10.1111/j.1365-2427.2007.01745.x>, 2007.
- 2070 Saunois, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A., Dlugokencky, E. J., Houweling, S., Patra, P. K., Ciais, P., Arora, V. K., Bastviken, D., Bergamaschi, P., Blake, D. R., Brailsford, G., Bruhwiler, L., Carlson, K. M., Carrol, M., Castaldi, S., Chandra, N., Crevoisier, C., Crill, P. M., Covey, K., Curry, C. L., Etiope, G., Frankenberg, C., Gedney, N., Hegglin, M. I., Höglund-Isaksson, L., Hugelius, G., Ishizawa, M., Ito, A., Janssens-Maenhout, G., Jensen, K. M., Joos, F., Kleinen, T., Krummel, P. B., Langenfelds, R. L., Laruelle, G. G., Liu, L., Machida, T., Maksyutov, S., McDonald, K. C., McNorton, J., Miller, P. A., Melton, J. R., Morino, I., Müller, J., Murguía-Flores, F., Naik, V., Niwa, Y., Noce, S., O'Doherty, S., Parker, R. J., Peng, C., Peng, S., Peters, G. P., Prigent, C., Prinn, R., Ramonet, M., Regnier, P., Riley,

W. J., Rosentreter, J. A., Segers, A., Simpson, I. J., Shi, H., Smith, S. J., Steele, L. P., Thornton, B. F., Tian, H., Tohjima, Y., Tubiello, F. N., Tsuruta, A., Viovy, N., Voulgarakis, A., Weber, T. S., van Weele, M., van der Werf, G. R., Weiss, R. F.,
2080 Worthy, D., Wunch, D., Yin, Y., Yoshida, Y., Zhang, W., Zhang, Z., Zhao, Y., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.:
The global methane budget 2000–2017, *Earth Syst. Sci. Data*, 12, 1561–1623, <https://doi.org/10.5194/essd-12-1561-2020>,
2020.

Sawakuchi, H. O., Bastviken, D., Sawakuchi, A. O., Krusche, A. V., Ballester, M. V. R., and Richey, J. E.: Methane
2085 emissions from Amazonian rivers and their contribution to the global methane budget, *Global Change Biol*, 20, 2829–2840,
<https://doi.org/10.1111/gcb.12646>, 2014.

Smith, R. L. and Böhlke, J. K.: Methane and nitrous oxide temporal and spatial variability in two midwestern USA streams
containing high nitrate concentrations, *Sci. Total Environ.*, 685, 574–588, <https://doi.org/10.1016/j.scitotenv.2019.05.374>,
2090 2019.

Smith, R. M., Kaushal, S. S., Beaulieu, J. J., Pennino, M. J., and Welty, C.: Influence of infrastructure on water quality and
greenhouse gas dynamics in urban streams, *Biogeosciences*, 14, 2831–2849, <https://doi.org/10.5194/bg-14-2831-2017>, 2017.

2095 Spawn, S., Dunn, S., Fiske, G., Natali, S., Schade, J., and Zimov, N.: Summer methane ebullition from a headwater catchment
in Northeastern Siberia, *Inland Waters*, 5, 224–230, <https://doi.org/10.5268/IW-5.3.845>, 2015.

Stanley, E. H., Loken, L. C., Crawford, J. T., Casson, N. J., Oliver, S. K., Gries, C., and Christel, S.: A global database of
methane concentrations and atmospheric fluxes for streams and rivers, Environmental Data Initiative,
<https://doi.org/10.6073/pasta/21f5bd6642e9689baf90262f3c85ac4a>, 2015.

2100

Stanley, E. H., Casson, N. J., Christel, S. T., Crawford, J. T., Loken, L. C., and Oliver, S. K.: The ecology of methane in
streams and rivers: patterns, controls, and global significance, *Ecol. Monogr.*, 86, 146–171, <https://doi.org/10.1890/15-1027>,
2016.

2105 Stanley, E. H., Collins, S. M., Lottig, N. R., Oliver, S. K., Webster, K. E., Cheruvilil, K. S., and Soranno, P. A.: Biases in
lake water quality sampling and implications for macroscale research, *Limnol. Oceanogr.*, 64, 1572–1585,
<https://doi.org/10.1002/lno.11136>, 2019.

Stanley, E. H., Loken, L. C., Rocher-Ros, G., Casson, N. J., Oliver, S. K., Sponseller, R. A., Wallin, M., and Zhang, L.:
2110 GRiMeDB: a comprehensive global database of methane concentrations and fluxes in fluvial ecosystems with supporting

physical and chemical information, ver 2, Environmental Data Initiative, <https://doi.org/10.6073/pasta/f48cdb77282598052349e969920356ef>, 2023.

2115 Stow, C. A., Webster, K. E., Wagner, T., Lottig, N., Soranno, P. A., and Cha, Y.: Small values in big data: The continuing need for appropriate metadata, *Ecol. Inform.*, 45, 26–30, <https://doi.org/10.1016/j.ecoinf.2018.03.002>, 2018.

2120 Stringer, L. C., Mirzabaev, A., Benjaminsen, T. A., Harris, R. M. B., Jafari, M., Lissner, T. K., Stevens, N., and Tirado-von der Pahlen, C.: Climate change impacts on water security in global drylands, *One Earth*, 4, 851–864, <https://doi.org/10.1016/j.oneear.2021.05.010>, 2021.

Taillardat, P., Bodmer, P., Deblois, C. P., Ponçot, A., Prijac, A., Riahi, K., Gandois, L., del Giorgio, P. A., Bourgault, M. A., Tremblay, A., and Garneau, M.: Carbon dioxide and methane dynamics in a peatland headwater stream: Origins, processes and implications, *J. Geophys. Res. Biogeos.*, <https://doi.org/10.1029/2022jg006855>, 2022.

2125 Turner, A. J., Frankenberg, C., and Kort, E. A.: Interpreting contemporary trends in atmospheric methane, *Proc. Natl. Acad. Sci. U.S.A.*, 116, 2805–2813, <https://doi.org/10.1073/pnas.1814297116>, 2019.

van den Hoogen, J., Robmann, N., Routh, D., Lauber, T., van Tiel, N., Danylo, O., and Crowther, T. W.: A geospatial mapping pipeline for ecologists, *BioRxiv*, 2021.07.07.451145. <https://doi.org/10.1101/2021.07.07.451145>. 9 July 2021.

2130 Wallin, M. B.: Krycklan catchment time series, Swedish University of Agricultural Sciences, Uppsala, unpublished, 2021.

2135 Wallin, M. B., Campeau, A., Audet, J., Bastviken, D., Bishop, K., Kokic, J., Laudon, H., Lundin, E., Löfgren, S., Natchimuthu, S., Sobek, S., Teutschbein, C., Weyhenmeyer, G. A., and Grabs, T.: Carbon dioxide and methane emissions of Swedish low-order streams—a national estimate and lessons learnt from more than a decade of observations, *Limnol. Oceanogr. Lett.*, 3, 156–167, <https://doi.org/10.1002/lol2.10061>, 2018.

Wickham, H.: *ggplot2: Elegant Graphics for Data Analysis*, Springer-Verlag, New York, 2016.

2140 Wickham, H., François, R., Henry, L., and Müller, K.: *dplyr: A Grammar of Data Manipulation*, 2021.

Wilcock, R. J. and Sorrell, B. K.: Emissions of greenhouse gases CH₄ and N₂O from low-gradient streams in agriculturally developed catchments, *Water Air Soil Pollut.*, 188, 155–170, <https://doi.org/10.1007/s11270-007-9532-8>, 2008.

- Woda, J., Wen, T., Lemon, J., Marcon, V., Keeports, C. M., Zelt, F., Steffy, L. Y., and Brantley, S. L.: Methane concentrations
2145 in streams reveal gas leak discharges in regions of oil, gas, and coal development, *Sci. Total Environ.*, 737, 140105,
<https://doi.org/10.1016/j.scitotenv.2020.140105>, 2020.
- Wu, Q. whitebox: “WhiteboxTools” R Frontend. v1.2.0, <https://github.com/giswqs/whiteboxR>
- 2150 Zhang, L., Xia, X., Liu, S., Zhang, S., Li, S., Wang, J., Wang, G., Gao, H., Zhang, Z., Wang, Q., Wen, W., Liu, R., Yang, Z.,
Stanley, E. H., and Raymond, P. A.: Significant methane ebullition from alpine permafrost rivers on the East Qinghai–Tibet
Plateau, *Nat. Geosci.*, 13, 349–354, <https://doi.org/10.1038/s41561-020-0571-8>, 2020.
- Yang, L., Li, X., Yan, W., Ma, P., and Wang, J.: CH₄ concentrations and emissions from three rivers in the Chaohu Lake
2155 watershed in southeast China, *Journal of Integrative Agriculture*, 11, 665–673, [https://doi.org/10.1016/S2095-
3119\(12\)60054-9](https://doi.org/10.1016/S2095-3119(12)60054-9), 2012.
- Yang, X., Pavelsky, T. M., Allen, G. H.: The past and future of global river ice, *Nature*. 577, 69–73,
<https://doi.org/10.1038/s41586-019-1848-1>, 2020.
- 2160 Zhang, L., Xia, X., Liu, S., Zhang, S., Li, S., Wang, J., Wang, G., Gao, H., Zhang, Z., Wang, Q., Wen, W., Liu, R., Yang, Z.,
Stanley, E. H., and Raymond, P. A.: Significant methane ebullition from alpine permafrost rivers on the East Qinghai–Tibet
Plateau, *Nat. Geosci.*, 13, 349–354, <https://doi.org/10.1038/s41561-020-0571-8>, 2020.
- 2165 Zhang, W., Li, H., Pueppke, S. G., and Pang, J.: Restored riverine wetlands in a headwater stream can simultaneously
behave as sinks of N₂O and hotspots of CH₄ production, *Environ. Pollut.*, 284, 117114,
<https://doi.org/10.1016/j.envpol.2021.117114>, 2021.