

1 **Inventory of glaciers and perennial snowfields of the**  
2 **~~contiguous~~conterminous USA**  
3

4 Andrew G. Fountain<sup>1</sup>, Bryce Glenn<sup>1</sup>, Christopher Mcneil<sup>2</sup>

5 <sup>1</sup>Department of Geology, Portland State University, 1721 SW Broadway, Portland, OR. 97212  
6 USA, <sup>2</sup>U.S. Geological Survey Alaska Science Center, Anchorage, AK, USA

7 *Correspondence to:* Andrew G. Fountain (andrew@pdx.edu)

8 **Abstract**

9 This report summarizes an updated inventory of glaciers and perennial snowfields of the  
10 ~~contiguous~~conterminous United States. The inventory is based on interpretation of mostly aerial  
11 imagery provided by the National Agricultural Imagery Program, U.S. Department of  
12 Agriculture with some satellite imagery in places where aerial imagery was not suitable. The  
13 inventory includes all perennial snow and ice features ~~greater than~~≥ 0.01 km<sup>2</sup>. Due to aerial  
14 survey schedules and seasonal snow cover, imagery acquired over a number of years were  
15 required. The earliest date is 2013 and the latest is 2020, but more than 73% of the outlines were  
16 acquired from 2015 imagery. The inventory is compiled as shapefiles within a geographic  
17 information system that includes feature classification, area, and location. The inventory  
18 identified 1331 (366.52 ± 14.34 km<sup>2</sup>) glaciers, 1776 (31.0001 ± 9.30 km<sup>2</sup>) perennial snowfields,  
19 and 35 (3.57 km<sup>2</sup> ± no uncertainty) buried-ice features. The data including both the shapefiles  
20 and tabulated results are publicly available at <https://doi.org/10.15760/geology-data.03> (Fountain  
21 & Glenn, 2022).

22 **1. Introduction**

23 Glaciers are an important feature of the landscape for several reasons. Geologically, they modify  
24 the landscape through erosion and deposition (Alley et al., 2019; Benn & Evans, 2010).  
25 Although these processes are typically slow, sudden episodes can occur such as moraine failure  
26 due to fluvial erosion resulting in catastrophic debris flows (Beason et al., 2018; Chiarle et al.,  
27 2007; O'Connor et al., 2001). Hydrologically, glaciers can be viewed as frozen reservoirs of  
28 water that naturally regulate streamflow on seasonal to decadal time scales (Dussaillant et al.,  
29 2019; Fountain & Tangborn, 1985; Moore et al., 2009). ~~Glacial~~Glacier runoff increases during  
30 warm periods and diminishes during cool, wet periods. Thus, ~~glacial~~glacier populated watersheds  
31 have less seasonally variable runoff than ice-free watersheds. Also, ~~glacial~~glacier runoff cools  
32 stream temperatures in the driest and hottest part of the summer after seasonal snowpacks have  
33 vanished (Cadbury et al., 2008; Fellman et al., 2014). ~~As glaciers shrink, they have less ability~~  
34 ~~to buffer seasonal runoff variations and watersheds become more susceptible to drought.~~As  
35 ~~glaciers shrink, they have less ability to buffer seasonal runoff variations and watersheds become~~  
36 ~~more susceptible to drought (Huss & Hock, 2018; Pritchard, 2019).~~ Globally, the loss of  
37 perennial ice from the landscape is a major contributor to sea level rise (Meier, 1984; Parkes &  
38 Marzeion, 2018; Zemp et al., 2019).

39 Glacier inventories have been valuable for assessing glacier contribution to sea level change  
40 (Hock et al., 2009; Pfeffer et al., 2014), and for assessing regional hydrology (~~Yao et al., 2007;~~  
41 ~~Moore et al., 2009).~~ They also provide a baseline for quantifying future glacier changes. Glacier  
42 inventories have been compiled for many regions of the world (Bolch et al., 2010; Smiraglia et  
43 al., 2015; Sun et al., 2018). An exception has been western United States (US), defined here as  
44 those conterminous states west of the 100<sup>th</sup> meridian. Despite a vigorous history of glacier  
45 studies (e.g., Armstrong, 1989; Rasmussen, 2009), glacial geology (e.g. Davis, 1988; Bowerman  
46 and Clark, 2011; Osborn et al., 2012), and regional inventories (e.g. DeVisser & Fountain, 2015;  
47 Fagre et al., 2017; Post et al., 1971) the glacier cover for the entire western US has not been  
48 updated in several decades (Moore et al., 2009; Yao et al., 2007). They also provide a baseline for  
49 quantifying future glacier changes. Updated glacier inventories have been compiled for many  
50 regions of the world (Andreassen et al., 2022; Bolch et al., 2010; Smiraglia et al., 2015; Sun et  
51 al., 2018). An exception has been western United States (US), defined here as those  
52 conterminous states west of the 100<sup>th</sup> meridian. The most recent inventory is (Fountain et al.,  
53 2007, 2017) based on U.S. Geological Survey maps compiled over a 40-year period from the late  
54 1940s to the 1980s. Despite a vigorous history of glacier studies (e.g. Armstrong, 1989;  
55 Rasmussen, 2009)), glacial geology (e.g. Bowerman & Clark, 2011; Davis, 1988; Osborn et al.,  
56 2012)), and regional inventories (e.g. DeVisser & Fountain, 2015; Fagre et al., 2017; Post et al.,  
57 1971) the glacier cover for the entire western US has not been reevaluated.

58 The earliest scientific identification of glacier-populated regions in the western US date to King  
59 (1871) and, more comprehensively, to Russell (1898). The first summary of glacier ~~areas~~  
60 ~~covered area~~ for each state was ~~in 1961~~ (Meier, ~~1961~~). However, the data sources and methods  
61 used to compile the inventories are unknown. Denton (1975) summarized all known glacier  
62 studies in the western US, but did not tabulate glacier ~~areas~~ ~~area~~. Krimmel (2002) updated  
63 Meier's study and provided total glacier area for the various mountain ranges by  
64 ~~summarizing~~ ~~summarizing~~ a variety previous studies published over a 10+ year time span. It is not  
65 clear whether the inventory is complete and no data on individual glaciers are provided. Fountain  
66 et al. (2007, 2017) compiled the first comprehensive inventory of glaciers in the western US. The  
67 data were derived from historical U.S. Geological Survey (USGS) 1:24,000 scale maps compiled  
68 over a 40-year period from the 1940s to the 1980s (Gesch et al., 2002; Usery et al., 2009).  
69 Because the USGS mapping was based on one-time aerial imagery, the misinterpretation of  
70 seasonal snow as perennial was extensive in some regions. The most current study, Selkowitz &  
71 Forster (2016), used Landsat satellite imagery compiled over a four-year period, 2010-2014, and  
72 an automated detection scheme to define perennial snow and ice. However, ~~these early~~  
73 automated schemes are known to misclassify debris-covered ice as ice-free landscape  
74 underestimating glacier area (Earl & Gardner, 2016; Paul et al., 2007; Rabatel et al., 2017).  
75 ~~Recent advances in automated detection have reduced these errors suggesting a more promising~~  
76 ~~future~~ (Lu et al., 2022; Robson et al., 2020).

77 This ~~report~~ ~~paper~~ presents the results of an updated and comprehensive inventory of glaciers and  
78 perennial snowfields of the western US for the purpose of defining their current extent and to  
79 provide of baseline for estimating future changes. ~~The report summarizes~~ ~~We summarize~~ our  
80 methods, uncertainties, tabulated results, and data availability. ~~The data referenced throughout the~~  
81 ~~manuscript are publicly available at <https://doi.org/10.15760/geology-data.03>.~~

## 2. Methods

### 2.1 Data Sources, Classification, Digitizing, and Completeness

The ~~location of the~~ glaciers and perennial snowfields were initially ~~identified by~~ located using a geographic information system (GIS) database from Fountain et al. (2007, [2017](#)). New outlines were manually digitized from three sources of optical imagery. Most of the outlines were digitized from color digital orthographic aerial photographs available from the National Agricultural Imagery Program (NAIP), U.S. Department of Agriculture, Farm Service Agency program ([NAIP, 2017](#)). Since 2009, the imagery is collected on cycles of two to three years. The spatial resolution is at least 1 m (ground sampling distance) with a horizontal accuracy of 6 m of photo-identifiable ground control points (USDA, 2021). NAIP imagery was downloaded from Data Gateway ([https://datagateway.nrcs.usda.gov/GDGHome\\_DirectDownload.aspx](https://datagateway.nrcs.usda.gov/GDGHome_DirectDownload.aspx)). ([NAIP, 2017](#)), ([https://datagateway.nrcs.usda.gov/GDGHome\\_DirectDownload.aspx](https://datagateway.nrcs.usda.gov/GDGHome_DirectDownload.aspx)). Since 2009, the imagery is collected on cycles of two to three years. The aerial imagery was orthorectified using the inertial navigation system - GPS unit in the aircraft. Photo identifiable GPS-survey ground control points were then used to adjust the photo strip. Orthorectified strips, which had  $\geq 30\%$  overlap with adjacent strips, were overlaid with each other and with ground control points to check accuracy. The image strips are then mosaicked together. The spatial resolution was  $\leq 0.6$  m with a horizontal accuracy of  $\leq 6$  m of photo-identifiable ground control points (NAIP, 2017). The NAIP imagery fit the historic USGS glacier outlines remarkably well. In a few cases, NAIP imagery was not suitable due to seasonal snow, deep shadows, or image warping caused by orthophoto rectification, therefore other sources were used including Maxar satellite imagery (Maxar Technologies, Inc) with a spatial resolution of 0.5 – 1 m. In one situation, For 21 perennial snowfields and three glaciers we ~~used~~relied on the most recent snow-free imagery available in Google Earth (Google, Inc), resolution ~ 1m, because no other imagery was suitable. The outlines were digitized in Google Earth and exported to ArcMap (Esri, Inc).

We manually identified all glaciers, ice patches, and perennial snowfields ~~equal or larger than~~  $0.01 \text{ km}^2$ . Glaciers are defined as perennial snow and ice that moves (Cogley et al., 2011). A feature was considered perennial if it was present on the original 1:24,000 USGS topographic maps and present on all Google Earth imagery. Movement was identified by the presence of crevasses. Perennial snowfields and ice patches do not exhibit movement, as indicated by a lack of crevasses observed in the imagery. We do not distinguish between snowfields and ice patches and refer to both as perennial snowfields.

~~Contiguous glacier cover, most commonly on volcanoes, was separated into individual glaciers if they had unique names as indicated on the USGS maps. The orientation of crevasse patterns was used to define flow divides. In the absence of these patterns, shaded relief digital elevation models were used to examine slope changes. These models were derived from aerial lidar data, flown under contract to the USGS (Bard, 2017b, 2017a, 2019; Robinson, 2014) or the Oregon Department of Geology and Mineral Industries (DOGAMI, 2011).~~

121 Contiguous glacier cover, most commonly on volcanoes, was separated into individual glaciers if  
122 they had unique names as indicated on the USGS maps. The orientation of crevasse patterns was  
123 used to define flow divides. In the absence of these patterns, shaded relief maps from digital  
124 elevation models were used. These models were derived from aerial lidar data, flown under  
125 contract to the USGS (Bard, 2017a, 2017b, 2019; Robinson, 2014) or the Oregon Department of  
126 Geology and Mineral Industries (DOGAMI, 2011).

127 We encountered a number of challenges to our classification and delineation of the glaciers and  
128 perennial snowfields. Although crevasses were used to define movement, in a few cases it  
129 appeared that they penetrated through the feature to the bedrock underneath suggesting a  
130 mechanical break up. In these cases, the feature was classified as a snowfield. Debris-cover  
131 made defining the glacier outline for some glaciers on the volcanoes of the Cascade Range. We  
132 relied on local knowledge to help define some boundaries and independent digitization efforts by  
133 the authors and others to provide an uncertainty as explained below. In the high alpine regions of  
134 California, Colorado, and Wyoming, the terminus of some glaciers was hard to define. Rather  
135 than abruptly terminating, the ice seems to thin and smoothly transitions into the surrounding  
136 rock talus (see Wyoming in the appendix-Figure 1). It was unclear whether a thin debris layer  
137 blanketed the ice or cobbles and boulders ~~protruding~~protruded through the thin ice. The  
138 boundary was mapped along the edge of identifiable ice.

139




140  
141 **Figure 1.** An example of a glacier seemingly melting into the talus surrounding the terminus  
142 (upper right). The glacier is flowing from the lower left-hand corner to the upper right-hand  
143 corner. Although not a common problem, one particular difficulty was distinguishingThe glacier  
144 is located in the Wind River Range, WY, INV ID E618081N4774579 and base image is from  
145 the National Agricultural Image Program taken in 2015.

146  
147 In a few situations, we found it difficult to distinguish glaciers from rock glaciers (Brardinoni et  
148 al., 2019). A rock glacier is a mass of rock debris in a matrix of ice that flows (Cogley et al.,  
149 2011). They can be difficult to distinguish from a debris-covered glacier, one that has extensive  
150 rock debris over the ablation zone, that lower part of a glacier with exposed ice in late summer.  
151 We adopted the following topographic classification. If the slope of the apparent glacier/ice  
152 patch/snowfield graded into was similar to the slope of the rock glacier, with no change in sign,  
153 then we considered it part of the rock glacier. (Figure 2a). On the other hand, if the slope  
154 changed sign at the bottom of a topographic depression separates the apparent glacier/snowfield,  
155 such that the topography formed a dip before reaching another topographic high that marks from  
156 the start of a rock glacier, then it was considered independent feature (see Colorado in the  
157 appendix)-Figure 2b). This latter case is similar to the “glacier forefield-connected” rock glacier  
158 as described by (RGIK, 2022).

Formatted: Line spacing: single, Keep with next

160



161

162 **Figure 2.** Examples of glacier versus rock glacier identification. (a) An example of a snowfield  
 163 that is considered part of the rock glacier. Location, Colorado Front Range, 40.827477° N, -  
 164 106.657400° E. Image is from © Google Earth, 9/2014; (b) Tyndall Glacier in the Colorado  
 165 Front Range, 40.305291° N, -105.689602° E, with a rock glacier slightly down valley. Image is  
 166 from © Google Earth 9/2016.

167

168 In a number of cases, we observed buried ice adjacent to a glacier (see Oregon in the  
 169 appendix). Figure 3). Here we use the term ‘buried ice’ to mean dead ice formerly part of a  
 170 flowing glacier, and not the permafrost context of ice embedded within or on top of perennially  
 171 frozen ground. The rocky surface texture of the buried ice was hummocky and very different  
 172 from surrounding bedrock and adjacent ice-, and not a moraine. Occasionally a crack in the  
 173 surface revealed subsurface ice. The feature appeared to be non-moving (dead) ice that is  
 174 covered by debris similar to some of the ice-debris complexes described by Bolch et al. (2019).  
 175 We decided to include these features as a separate classification, ‘buried ice’, because their size  
 176 was large relative to the glacier, they were probably once part of the glacier, and may be  
 177 important local sources of meltwater for streamflow.

178

179

180

181

Formatted: Font: +Body (Calibri), 11 pt, Not Bold, Not Italic

Formatted: Keep with next

Formatted: Font: Not Italic, Font color: Auto

Formatted: Caption

Formatted: Font: Not Italic, Font color: Auto

Formatted: Font: Not Italic, Font color: Auto, Pattern: Clear

Formatted: Font: Not Italic, Font color: Auto

Formatted: Font: Bold, Italic

Formatted: Space After: 8 pt

Formatted: Font color: Custom Color(RGB(34,34,34))



**Figure 3.** Lost Creek Glacier, South Sister, Oregon. Note buried ice and lack of crevasses to the left of the grey-blue ice, suggesting ice that is no longer moving and therefore not part of the dynamic glacier. The white box surrounds an area that has collapsed due to subsurface melt. The inset enlargement shows a cliff edge of exposed dirty ice (white arrow in upper left) indicated by a darker color suggesting wet sediment and a finer texture than the surface debris. The black arrow shows the width of the cleaner ice for scale. Image is from © Google Earth, 8/9/2021.

The glaciers and perennial snowfields outlines were digitized using ArcMap (ESRI Esri, Inc), a geographic information system, at scales varying from 1:300 to 1:2000 depending on image quality and complexity. The projection we used was the native projection of the image, North American Datum of 1983 (NAD83) for NAIP, and World Geodetic System 1984 (WGS84), with the relevant local Universal Transverse Mercator (UTM) zone, for Maxar and Google Earth. When Maxar or Google Earth imagery were used, final outlines were projected into the NAD83 coordinate system. In situations where it was hard to interpret feature geometry, Google Earth was very helpful because its terrain feature provides an oblique perspective that can be tilted and rotated. Each outline was checked independently by the two senior authors of this report, and in some cases by a third collaborator. Google Earth was often used as an additional aid in interpretation because of its tilt and rotation features yielded oblique perspectives. Retaining only those outlines  $\geq 0.01 \text{ km}^2$ , each was checked independently by the two senior authors of this report and in some cases by a third collaborator in order to reduce bias (Leigh et al., 2019). If an outline was revised, then it was returned to its original author for review and correction, and the process iterated until all parties agreed.

Formatted: Font color: Auto

205 **Uncertainty** The uncertainty in glacier area was calculated as one-half the absolute difference  
206 between initial and final revised digitized area (the range) divided by the final area and expressed  
207 as a percentage. For some glaciers where no revision was necessary a 1% uncertainty was  
208 assigned to account for digitizing error, which is known to be relatively small (DeVisser &  
209 Fountain, 2015; Hoffman et al., 2007). For the relatively few glaciers where a small section of  
210 perimeter was masked by deep shadow, seasonal snow patches, rock debris, or poor imagery, a  
211 higher uncertainty was assigned by visually comparing the area in question to the total possible  
212 area of the glacier. The estimated uncertainty, up to 16%, was determined by digitizing the  
213 minimum and maximum perimeters from a sample of glaciers with similar issues. Uncertainty  
214 about the position of a flow divide was considered 5%, due to the topographic ambiguity along a  
215 divide, and estimated from several digitizing efforts. For perennial snowfields a 30% uncertainty is  
216 assigned because the seasonal snow commonly covers the smaller patch of perennial snow and  
217 the seasonal snow varies greatly from year to year. The snowfield uncertainty was arbitrary in  
218 order to note their presence and location, but preclude them from area change calculations  
219 because area differences are typically smaller than the assigned uncertainty.  
220

221  
222 Our initial inventory was then compared sequentially to two other independent inventories to test  
223 for errors of omission or commission. The first comparison was to the Selkowitz and Forster  
224 (2016) inventory (SFI). However, to compare the inventories we had to first reconcile the  
225 different differences in methods of each inventory prior to comparison. Buried-ice features were  
226 eliminated from our inventory because the SFI did not map buried ice. ~~Features removed from~~  
227 ~~the~~ SFI was filtered to only include features  $\leq 0.01 \text{ km}^2$  to match our minimum area  
228 threshold; a small number of glaciers and snowfields features located in Canada; the were  
229 removed; and a few glacier mis-classifications of ponds, lakes and dry lakebeds as glaciers were  
230 removed. Notably, the SFI did not split contiguous ice masses, such as glacier-covered  
231 volcanoes, into individual glaciers, consequently we do not expect the number of features in the  
232 SFI and our inventory to match. Once the two inventories were reconciled, those glaciers and  
233 perennial snowfields unique to one inventory were examined for inclusion in a revised inventory.  
234 Features selected from the SFI were digitized using the same imagery we used for our inventory.  
235

236 ~~The revised inventory was then compared to the 2016 National Land Cover Database (NLCD),~~  
237 ~~which did not map glaciers and perennial snowfields per se, but mapped the distribution of~~  
238 ~~perennial snow and ice (Jin et al., 2019). However, the NLCD used a smaller number of images~~  
239 ~~over time for any one location such that we consider the assessment of 'perennial' has a high~~  
240 ~~uncertainty.~~ The revised inventory was then compared to the 2016 National Land Cover Database  
241 (NLCD, Dewitz, 2019), which did not map glaciers and perennial snowfields per se, but mapped  
242 the distribution of perennial snow and ice (Jin et al., 2019; Wickham et al., 2021). However, the  
243 NLCD used a small number of recent images to assess a 'perennial' presence and therefore  
244 significant errors of commission are expected. Also, the landscape class of snow and ice received  
245 less attention than other classes (e.g. agriculture) such that the timing of imagery acquisition may  
246 be earlier in the summer than optimal and misclassification of clouds as snow and ice may be  
247 present (personal communication C. Homer and J. -Also, the landscape class of snow and ice  
248 received less attention than other classes (e.g. agriculture) such that the timing of imagery  
249 acquisition may be earlier in the summer than optimal and misclassification of clouds as snow



250 ~~and ice may be present (personal communication C. Homer and J. Dewitz, USGS, email~~  
251 ~~December 2015). Again~~The NLCD inventory was compared to the revised inventory and, as  
252 ~~before,~~ the features unique to one inventory were examined for inclusion ~~and those.~~ Those  
253 features selected from the NLCD for inclusion were digitized using the same imagery we used  
254 for our inventory.

Formatted: Font color: Auto

## 255 2.2 Uncertainty


256 Three main sources of uncertainty in the glacier outlines, are georeferencing, digitization, and  
257 interpretation (DeVisser & Fountain, 2015; Sitts et al., 2010). We found georeferencing error to  
258 very small. In any case, the precise location of the outline does not affect its area. Also, the  
259 digitized points are highly correlated such that no deviations from the true outline are caused by  
260 georeferencing. Digitizing error is relatively small, 1%, with good imagery and crisp contrast  
261 between the glacier and ice-free surroundings (DeVisser & Fountain, 2015; Hoffman et al.,  
262 2007). The largest uncertainty is interpretation error caused by poor imagery, shadow, debris  
263 cover, and seasonal snow patches. This uncertainty was calculated in different ways according to  
264 the situation. If the outline was digitized a second (or third) time due different interpretations by  
265 the authors or collaborators the uncertainty is one-half the absolute difference of the between the  
266 largest and smallest digitized areas (the range) divided by the final area and expressed as a  
267 percentage. For the relatively few glaciers where a small section of perimeter was masked by  
268 deep shadow, seasonal snow patches, rock debris, or poor imagery, a higher uncertainty was  
269 assigned by visually estimating the area in question and dividing by the total possible area. In a  
270 few cases the location of a flow divide between glaciers wasn't clear a 5% error is assigned. This  
271 was calculated from the area difference in several test cases where multiple possible flow divides  
272 were digitized. For perennial snowfields, the smaller patch of perennial snow is often covered by  
273 seasonal snow, which varies greatly from year to year. We measured the area of a number of  
274 snowfields over time using late summer historic imagery in Google Earth. Results showed that  
275 the variations in snowfield area could be as much as 30%. We assigned this somewhat arbitrary  
276 uncertainty in order to note snowfield presence and location, but preclude them from area change  
277 calculations because area differences are typically smaller than the assigned uncertainty.

## 280 **3. Results**


281 Our initial inventory identified 2267 glaciers and perennial snowfields totaling 391.95 km<sup>2</sup>.  
282 About 70% (1576) overlapped the features in the SFI. After examining all features unique to  
283 each inventory, we revised our inventory to include 2373 (394.99 km<sup>2</sup>) glaciers and perennial  
284 snowfields. Comparing the revised inventory to the 2016 NLCD resulted in adding another 134  
285 (2.53 km<sup>2</sup>) features, which included 12 (0.38 km<sup>2</sup>) glaciers. The final inventory includes 2542  
286 ~~glacial~~ features composed of 1331 (366.52 km<sup>2</sup>) glaciers, 1176 (31.01 km<sup>2</sup>) perennial snowfields,  
287 and 35 (3.57 km<sup>2</sup>) buried ice deposits (Table 1; Figure 4). Most glaciers and perennial  
288 snowfields, 1554 (62%) were outlined using 2015 NAIP imagery with the remainder outlined  
289 using mostly NAIP imagery from 2013 to 2020. ~~The state of Washington has the greatest number~~  
290 ~~and area of glaciers, perennial snowfields, and buried ice.~~

Formatted: Font color: Auto

Formatted: Space After: 0 pt, Line spacing: single



294  
295



296  
 297 **Figure 1.4.** The spatial distribution and number of glaciers and perennial snowfields, greater  
 298 than 0.01 km<sup>2</sup>, in the western US-United States. Colors indicate the date of aerial and satellite  
 299 imagery used to outline the features. The line is the cumulative total. Base imagery from Esri  
 300 Inc. -Inset is a bar graph and cumulative sum of the number of glaciers and perennial snowfields  
 301 digitized in each image date.  
 302  
 303

304 **Table 1.** The summary of the glacia~~glacier~~ inventory for the American West, exclusive of  
 305 Alaska. Number is the total number of features within each classification (Class), Uncert.~~Max~~  
 306 Area is the largest area uncertainty-of the feature within that class and MaxMean Area is  
 307 maximumthe average area. Note that the uncertainty of 'Buried ice' is unknown.  
 308

State/Region/Class	Number	Total Area km <sup>2</sup>	Max Area km <sup>2</sup>	Mean Area km <sup>2</sup>
--------------------	--------	----------------------------	--------------------------	---------------------------

Deleted Cells  
 Formatted Table

			Une ert km <sup>2</sup>			
<b>California Cascade Range</b>	<b>132</b>	<del>10.63 ±</del> <b>0.61</b>	<del>0.61</del>	<b>1.45</b>	<b>0.08</b>	
	<b>39</b>	<del>5.74 ±</del> <b>0.37</b>	<del>0.37</del>	<b>1.45</b>	<b>0.15</b>	
		Buried ice	5			0.44
Glaciers Perennial snowfields	10	<del>4.61 ±</del> <b>0.17</b>	<del>0.17</del>	1.45	0.46	
	24	<del>0.68 ±</del> <b>0.21</b>	<del>0.21</del>	0.08	0.03	
		<del>4.86 ±</del> <b>0.23</b>	<del>0.23</del>			
<b>Sierra Nevada</b> Buried ice	<b>91</b>	<del>0.23</del> <b>0.13</b>	<del>0.00</del>	<b>0.66</b>	<b>0.05</b>	
	2	<del>4.37 ±</del> <b>0.12</b>	<del>0.12</del>	0.10	0.06	
		64	<del>0.37 ±</del> <b>0.11</b>	<del>0.11</del>	0.66	0.07
Glaciers Perennial snowfields	25	<del>0.03 ±</del> <b>0.00</b>	<del>0.00</del>	0.03	0.01	
	<b>Trinity Alps</b>	<b>2</b>	<del>0.03 ±</del> <b>0.00</b>	<del>0.00</del>	<b>0.02</b>	<b>0.02</b>
		2	<del>0.00</del> <b>0.00</b>	<del>0.00</del>	0.02	0.02
<b>Colorado Elk Mountains</b>	<b>84</b>	<del>2.20 ±</del> <b>0.46</b>	<del>0.46</del>	<b>0.16</b>	<b>0.03</b>	
	5	<del>0.09 ±</del> <b>0.03</b>	<del>0.03</del>	<b>0.0</b>	<b>0.0</b>	
		1	<del>0.01 ±</del> <b>0.00</b>	<del>0.00</del>	0.01	0.01
Glaciers Perennial snowfields	4	<del>0.08 ±</del> <b>0.02</b>	<del>0.02</del>	0.03	0.02	
	<b>Front Range</b>	<b>58</b>	<del>1.73 ±</del> <b>0.33</b>	<del>0.33</del>	<b>0.16</b>	<b>0.03</b>
		13	<del>0.74 ±</del> <b>0.03</b>	<del>0.03</del>	0.16	0.06
Glaciers Perennial snowfields	45		<del>0.99 ±</del> <b>0.30</b>	<del>0.30</del>	0.09	0.02
	<b>Gore Range</b>	<b>7</b>	<del>0.11 ±</del> <b>0.03</b>	<del>0.03</del>	<b>0.02</b>	<b>0.02</b>
		1	<del>0.02 ±</del> <b>0.00</b>	<del>0.00</del>	0.02	0.02
Glaciers Perennial snowfields	6		<del>0.09 ±</del> <b>0.03</b>	<del>0.03</del>	0.02	0.02

Formatted: Centered, Right: -0.07"

Formatted: Left

Formatted: Left

Deleted Cells

Deleted Cells

<b>Medicine Bow Mountains</b>	1	<b>0.04 ± 0.01</b>	<del>0.01</del>	0.04	0.04
Perennial snowfields	1	0.04 ± 0.01	<del>0.01</del>	0.04	0.04
<b>Park Range</b>	6	<b>0.11 ± 0.03</b>		0.0	<del>0.0</del>
Perennial snowfields	6	0.11 ± 0.03		0.0	<del>0.0</del>
<b>San Miguel Mountains</b>	5	<b>0.07 ± 0.02</b>		2	<del>2</del>
Perennial snowfields	5	0.07 ± 0.02		2	<del>2</del>
<b>Sawatch Range</b>	2	<b>0.04 ± 0.01</b>	<del>0.01</del>	0.03	0.02
Perennial snowfields	2	0.04 ± 0.01	<del>0.01</del>	0.03	0.02
<b>Idaho Sawtooth Range</b>	6	<b>0.08 ± 0.02</b>		2	<del>2</del>
Perennial snowfields	6	0.08 ± 0.02		2	<del>2</del>
<b>Montana Beartooth - Absaroka</b>	416	<del>30.26 ± 2.27</del>	<del>2.27</del>	1.45	0.07
Buried ice	1	0.04	-	0.04	0.04
Glaciers	50	4.31 ± 0.12	<del>0.12</del>	0.45	0.09
Perennial snowfields	60	1.72 ± 0.52	<del>0.52</del>	0.22	0.03
<b>Bitterroot Range</b>	4	<b>0.08 ± 0.02</b>	<del>0.02</del>	0.03	0.02
Glaciers	1	<del>0.03</del>	0.03 ± 0.00	0.03	0.03
Perennial snowfields	3	0.05 ± 0.02		0.02	0.02
<b>Cabinet Mountains</b>	9	<b>0.25 ± 0.08</b>		0.08	<del>0.08</del> <del>0.03</del>
Perennial snowfields	9	0.25 ± 0.08		0.08	<del>0.08</del> <del>0.03</del>
<b>Crazy Mountains</b>	13	<b>0.27 ± 0.06</b>	<del>0.06</del>	0.04	0.02
Glaciers	3	0.06 ± 0.00	<del>0.00</del>	0.04	0.02

Deleted Cells

Deleted Cells

Deleted Cells

Formatted: Centered

Deleted Cells

Formatted: Centered

Formatted: Left

Deleted Cells

Deleted Cells

Deleted Cells

Perennial snowfields	10	0.21 ± <del>0.06</del> <u>0.06</u>		0.04	0.02
<b>Lewis Range</b>	<b>230</b>	<b>21.38 ± 1.15</b> <u>1.15</u>	<b>1.45</b>	<b>0.09</b>	
Glaciers	145	19.22 ± <del>0.50</del> <u>0.50</u>	1.45	0.13	
Perennial snowfields	85	2.16 ± <del>0.65</del> <u>0.65</u>	0.09	0.03	
<b>Mission-Swan-Flathead</b>	<b>49</b>	<b>2.20 ± 0.34</b> <u>0.34</u>	<b>0.22</b>	<b>0.04</b>	
Glaciers	11	1.16 ± <del>0.02</del> <u>0.02</u>	0.22	0.11	
Perennial snowfields	38	1.04 ± <del>0.31</del> <u>0.31</u>	0.09	0.03	
<b>Oregon Cascade Range</b>	<b>116</b>	<b>15.38 ± 1.62</b> <u>1.62</u>	<b>1.16</b>	<b>0.13</b>	
<b>Oregon Cascade Range</b>	<b>110</b>	<b>15.24 ± 1.58</b> <u>1.58</u>	<b>1.16</b>	<b>0.14</b>	
Buried ice	7	<del>1.25</del> <u>1.25</u>	-	0.45	0.18
Glaciers	42	11.90 ± <del>0.95</del> <u>0.95</u>	1.16	0.28	
Perennial snowfields	61	2.09 ± <del>0.63</del> <u>0.63</u>	0.15	0.03	
<b>Wallowa Mountains</b>	<b>6</b>	<b>0.14 ± 0.63</b>	<b>0.04</b>	<b>0.04</b>	<b>0.02</b>
Perennial snowfields	6	0.14 ± <del>0.04</del> <u>0.04</u>	0.04	0.04	0.02
<b>Washington Cascade Range-Northern</b>	<b>1481</b>	<b>312.26 ± 16.3</b> <u>16.33</u>	<b>11.24</b>	<b>0.21</b>	
<b>Washington Cascade Range-Northern</b>	<b>1126</b>	<b>186.58 ± 9.64</b> <u>9.64</u>	<b>6.06</b>	<b>0.17</b>	
Buried ice	10	<del>0.50</del> <u>0.50</u>	-	0.15	0.05
Glaciers	706	176.27 ± <del>6.70</del> <u>6.70</u>	6.06	0.25	
Perennial snowfields	410	9.80 ± <del>2.94</del> <u>2.94</u>	0.16	0.02	
<b>Cascade Range-Southern</b>	<b>219</b>	<b>101.66 ± 5.86</b> <u>5.86</u>	<b>11.24</b>	<b>0.46</b>	
Buried ice	10	<del>1.20</del> <u>1.20</u>	-	0.30	0.12
Glaciers	69	95.64 ± <del>4.42</del> <u>4.42</u>	11.24	1.39	

Formatted: Left

Deleted Cells

Deleted Cells

Formatted: Left

Formatted: Left

Perennial snowfields	140	4.82 ± 1.45	1.45	0.33	0.03
<b>Olympic Mountains</b>	<b>136</b>	<b>24.02 ± 0.82</b>	<b>0.82</b>	<b>5.09</b>	<b>0.18</b>
Glacier	106	23.44 ± 0.65	0.65	5.09	0.22
Perennial snowfield	30	0.57 ± 0.17	0.17	0.06	0.02
<b>Wyoming Absaroka Range</b>	<b>307</b>	<b>30.29 ± 2.34</b>	<b>2.34</b>	<b>2.32</b>	<b>0.10</b>
Glacier	10	1.44 ± 0.33	0.33	0.12	0.02
Perennial snowfield	52	0.48 ± 0.05	0.05	0.05	0.02
<b>Bighorn Mountains</b>	<b>8</b>	<b>0.96 ± 0.29</b>	<b>0.29</b>	<b>0.22</b>	<b>0.05</b>
Glacier	3	0.34 ± 0.01	0.01	0.22	0.11
Perennial snowfield	5	0.08 ± 0.02	0.02	0.03	0.02
<b>Teton Range</b>	<b>49</b>	<b>2.04 ± 0.21</b>	<b>0.21</b>	<b>0.23</b>	<b>0.04</b>
Glacier	20	1.46 ± 0.03	0.03	0.23	0.07
Perennial snowfield	29	0.59 ± 0.18	0.18	0.05	0.02
<b>Wind River Range</b>	<b>188</b>	<b>26.39 ± 1.76</b>	<b>1.76</b>	<b>2.32</b>	<b>0.14</b>
Glacier	74	22.42 ± 0.57	0.57	2.32	0.30
Perennial snowfield	114	3.97 ± 1.19	1.19	0.26	0.03
<b>Grand Total</b>	<b>2542</b>	<b>401.10 ± 23.64</b>	<b>23.64</b>	<b>11.24</b>	<b>0.16</b>


Deleted Cells

Deleted Cells

Deleted Cells


309  
310  
311  
312 [Before summarizing the inventory data, a note about the content in Appendix A. It summarizes](#)  
313 [the officially named glaciers that we regard as snowfields or missing; labeling issues found in the](#)  
314 [USGS Geographic Names Information System, the official agency responsible for hosting the](#)  
315 [names and locations of landscape features; and detailed notes, organized by US State, on the](#)  
316 [specific imagery used and challenges encountered digitizing glacier and snowfield outlines.](#)

317  
 318 The glaciers and perennial snowfields are generally small, averaging 0.28 and 0.03 km<sup>2</sup>,  
 319 respectively. Like glaciers elsewhere in the northern hemisphere, most glaciers face north to east.  
 320 (Evans, 2006; Fountain et al., 2017; Schiefer et al., 2007). The distribution of glacier area is  
 321 skewed toward smaller ice masses (Figure 5a). The State of Washington in the Pacific Northwest  
 322 has the largest number of glaciers, ice area and the largest glacier (11.24 km<sup>2</sup> Emmons Glacier)  
 323 of any of the other states (Table 1). Indeed, the glacier cover on Mount Rainier alone (77.37 km<sup>2</sup>)  
 324 is greater than the total sum in all the other states (71.16 km<sup>2</sup>). The elevation distribution of  
 325 glacier-covered area is bimodal with maxima at 2400 m and 3650 m (Figure 5b). The spatial  
 326 distribution of elevations shows a regional climate control with the lowest glaciers and perennial  
 327 snowfields in the maritime climate of the Pacific Northwest of Washington, Oregon, northern  
 328 California, and western Montana and the high elevations located in the continental climate of  
 329 central California, Colorado, Wyoming and southern Montana (Figure 6).  
 330



331 Figure 5. The area and elevation distribution of glaciers in the western U.S., (a) Histogram  
 332 showing the number of glaciers as a function of area. The x-axis intervals are log intervals; (b)  
 333 Elevation distribution of glacier-covered area.  
 334





336  
337 [Figure 6. Elevation distribution of glaciers and perennial snowfields across the western US. Base](#)  
338 [imagery from Esri Inc.](#)

339  
340 The final inventory conflicts with the current database of the Geographic Names Information  
341 System ([US Geological Survey, 2022https://www.usgs.gov/us-board-on-geographic-](https://www.usgs.gov/us-board-on-geographic-names/domestic-names)  
342 [names/domestic-names](https://www.usgs.gov/us-board-on-geographic-names/domestic-names)). The inventory excludes 52 officially named glaciers because 2 have  
343 disappeared, 25 were classified as perennial snowfields, the area of 18 was less than 0.01 km<sup>2</sup>,  
344 and 7 were considered rock glaciers ([Appendix A, Table 2A1](#)). In some cases, a named glacier or  
345 snowfield had split into multiple pieces since the original USGS mapping; all pieces were  
346 assigned the same name in the inventory ([see appendix, sections 6.7 and 6.8](#)). [Appendix A, Table](#)  
347 [A2](#). Several labels that identify the name of the glacier are not clearly associated with a specific  
348 [glacier and these are listed in Table 7.3.](#)

349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393

#### 4. Discussion

The advent of relatively frequent high resolution (< 1 m) optical aerial and satellite imagery available at little or no cost has made compiling and updating glacier inventories a realistic opportunity. Finding suitable imagery spanning only a few years apart provides a near-snapshot of glacier cover. This contrasts strongly with mapping efforts only a few decades ago when aerial-only photographic surveys required decades to cover the western US (Gesch et al., 2002). And the advent of GIS software made digitizing, summarizing, and interrogating digital outlines practical.

We had used the Fountain et al., 2017 historic inventory as a template to locate and update the perimeters of all the glaciers and perennial snowfields. Considering that the inventory was derived from the U.S. Geological Survey 1:24,000 maps, a result of a national effort to remap the entire country at a higher resolution, we were surprised that 240 features (~10%) were missed. These missing features were revealed after comparison with two other independently derived inventories. We had a similar experience in a prior study when comparing two independently derived glacier inventories. Together they suggest that independent efforts are important to compiling a comprehensive inventory.

Multiple checks more accurately define glacier perimeters (Leigh et al., 2019). Different investigators may make different decisions about glacier boundaries and results can differ particularly in debris-covered conditions or along flow divides (Paul et al., 2013). When they agree, it provides some confidence of the interpretation accuracy and where they disagree it provides input for estimating interpretation error.

The total area of glaciers in the western US, 367 km<sup>2</sup>, is a little smaller than that in Austria, 415 km<sup>2</sup>, (Fischer et al., 2015). Like glacier populated regions elsewhere the distribution of glacier area is skewed towards smaller glaciers (e.g. Linsbauer et al., 2012; Mishra et al., 2023; Zalazar et al., 2020). The uncertainty in glacier area is also similar with an overall 5% uncertainty for the total area. Paul et al. (2020) report an uncertainty of 3.3% over a set of 15 glaciers, 4% for 7 glaciers (Zalazar et al., 2020), 2.3% for 15 glaciers (Linsbauer et al., 2021). Our assessment method differs from those cited here in that we estimate the uncertainty for each individual glacier rather than upscaling the uncertainty calculated for a small subsample.

#### 5. Data products and availability

The data are available in three formats. The geospatial data and attribute tables are available in the shapefile (Esri) format and in an open source GeoJSON format. The attribute table is also available as an EXCEL file. These data products can be obtained from <https://doi.org/10.15760/geology-data.03> (Fountain & Glenn, 2022) and from the Global Land Ice Measurements from Space website <http://glims.colorado.edu/glacierdata/>. Maxar imagery

Formatted: List Paragraph

394 was accessed through the USGS and NGA NEXTVIEW license. The Maxar imagery has limited  
395 availability owing to restrictions (proprietary interest). Contact cmcneil@usgs.gov for more  
396 information.

## 397 **6. Conclusions**

399 We have compiled a new and comprehensive inventory of glaciers and perennial snowfields in  
400 the western US from aerial and satellite imagery. Results show that 2542 features are currently  
401 present and include 1331 (366.52 km<sup>2</sup>) glaciers, 1176 (31.01 km<sup>2</sup>) perennial snowfields, and 35  
402 (3.57 km<sup>2</sup>) buried ice deposits. Most of the data were acquired from 2015 NAIP imagery with the  
403 remainder from NAIP imagery and a few satellite images acquired over the period of 2013 to  
404 2020. The state of Washington has the greatest number and area of glaciers and perennial  
405 snowfields. This product updates an older inventory based on USGS 1:24000 maps compiled in  
406 the middle-late 1900's. The new inventory is a significant improvement in accuracy because the  
407 archive of historical imagery in Google Earth greatly aided our efforts to classify glaciers versus  
408 perennial snowfields. Finally, this new inventory provides a baseline for assessing glacier change  
409 in the coterminous US.

## 411 **7. Appendix A**

### 412 **A1 Missing Glaciers**

413 **Table 2-A1** List of officially named glaciers not classified as glaciers and excluded from the  
414 final inventory. Names come from the Geographic Names Information System, (US Geological  
415 Survey, 2022). The 'Reason' column lists why the named glacier is no longer considered a glacier  
416 in our inventory.

Formatted: Font color: Text 1

Formatted: Font color: Text 1

State/Region/Glacier Name	Reason
<b>California</b>	
<b>Sierra Nevada</b>	
Matthes Glaciers	rock glacier
Mount Warlow Glacier	rock glacier
Powell Glacier	rock glacier
<b>Colorado</b>	
<b>Front Range</b>	
Isabelle Glacier	perennial snowfield
Mills Glacier	perennial snowfield
Moomaw Glacier	perennial snowfield
Peck Glacier	perennial snowfield
Rowe Glacier	< 0.01 km <sup>2</sup>
Saint Marys Glacier	< 0.01 km <sup>2</sup>
Taylor Glacier	rock glacier
The Dove	< 0.01 km <sup>2</sup>
<b>Idaho</b>	
<b>Lost River Range</b>	

Borah Glacier	rock glacier
<b>Montana</b>	
<b>Beartooth Mountains-Absaroka Range</b>	
Grasshopper Glacier	rock glacier
<b>Cabinet Mountains</b>	
Blackwell Glacier	perennial snowfield
<b>Crazy Mountains</b>	
Grasshopper Glacier	rock glacier
<b>Lewis Range</b>	
Boulder Glacier	perennial snowfield
<b>Mission-Swan-Flathead Ranges</b>	
Fissure Glacier	< 0.01 km <sup>2</sup>
Gray Wolf Glacier	perennial snowfield
<b>Oregon</b>	
<b>Cascade Range</b>	
Carver Glacier	perennial snowfield
Clark Glacier	perennial snowfield
Irving Glacier	perennial snowfield
Lathrop Glacier	< 0.01 km <sup>2</sup>
Palmer Glacier	perennial snowfield
Skinner Glacier	perennial snowfield
Thayer Glacier	< 0.01 km <sup>2</sup>
<b>Wallowa Mountains</b>	
Benson Glacier	perennial snowfield
<b>Washington</b>	
<b>Cascade Range-Northern</b>	
Lyall Glacier	perennial snowfield
Milk Lake Glacier	disappeared
Snow Creek Glacier	perennial snowfield
Spider Glacier	perennial snowfield
Table Mountain Glacier	< 0.01 km <sup>2</sup>
<b>Cascade Range-Southern</b>	
Ape Glacier	< 0.01 km <sup>2</sup>
Dryer Glacier	perennial snowfield
Forsyth Glacier	< 0.01 km <sup>2</sup>
Meade Glacier	perennial snowfield
Nelson Glacier	< 0.01 km <sup>2</sup>
Packwood Glacier	perennial snowfield
Pinnacle Glacier	< 0.01 km <sup>2</sup>
Pyramid Glaciers	< 0.01 km <sup>2</sup>
Shoestring Glacier	< 0.01 km <sup>2</sup>
Stevens Glacier	perennial snowfield
Talus Glacier	perennial snowfield
Unicorn Glacier	< 0.01 km <sup>2</sup>
Williwakas Glacier	perennial snowfield

**Olympic Mountains**

Anderson Glacier perennial snowfield  
Lillian Glacier < 0.01 km<sup>2</sup>

**Wyoming**

**Absaroka Range**

DuNoir Glacier < 0.01 km<sup>2</sup>

**Teton Range**

Petersen Glacier < 0.01 km<sup>2</sup>  
Teepe Glacier perennial snowfield

**Wind River Range**

Hooker Glacier disappeared  
Harrower Glacier perennial snowfield  
Tiny Glacier < 0.01 km<sup>2</sup>

420  
421  
422  
423  
424  
425  
426  
427  
428

**A2 Glaciers that have split into multiple pieces and current errors in glacier label names**

**Table A2.** List of named glaciers that have split into multiple pieces.  
Names come from the Geographic Names Information System  
(<https://www.usgs.gov/tools/geographic-names-information-system-gnis>). ‘Count’ refers to the  
number of pieces in the updated inventory. ‘Classes’ is the classification of the pieces: glacier,  
perennial snowfield, buried-ice, or a combination.

<u>State/Region/Glacier Name</u>	<u>Count</u>	<u>Classes</u>
<b><u>California</u></b>		
<b><u>Cascade Range</u></b>		
<u>Bolam Glacier</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>
<u>Hotlum Glacier</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>
<u>Whitney Glacier</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>
<u>Wintun Glacier</u>	<u>3</u>	<u>Glaciers and perennial snowfields</u>
<b><u>Sierra Nevada</u></b>		
<u>Goethe Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<u>Lyell Glacier</u>	<u>4</u>	<u>Glaciers and perennial snowfields</u>
<u>Norman Clyde Glacier</u>	<u>3</u>	<u>Glaciers only</u>
<u>Powell Glacier</u>	<u>2</u>	<u>Glacier and Buried-ice</u>
<b><u>Colorado</u></b>		
<b><u>Front Range</u></b>		
<u>Saint Vrain Glaciers</u>	<u>6</u>	<u>Glaciers and perennial snowfields</u>
<b><u>Montana</u></b>		
<b><u>Beartooth Mountains-Absaroka Range</u></b>		
<u>Castle Rock Glacier</u>	<u>3</u>	<u>Glaciers and perennial snowfields</u>
<u>Granite Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<u>Grasshopper Glacier</u>	<u>4</u>	<u>Glaciers and perennial snowfields</u>
<u>Hopper Glacier</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>
<u>Snowbank Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<u>Wolf Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<b><u>Lewis Range</u></b>		

<u>Agassiz Glacier</u>	<u>3</u>	<u>Glaciers only</u>
<u>Blackfoot Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<u>Carter Glaciers</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>
<u>Dixon Glacier</u>	<u>3</u>	<u>Glaciers and perennial snowfields</u>
<u>Harrison Glacier</u>	<u>5</u>	<u>Glaciers and perennial snowfields</u>
<u>Kintla Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<u>Logan Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<u>Shepard Glacier</u>	<u>3</u>	<u>Glaciers only</u>
<u>Siyeh Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<u>Two Ocean Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<u>Whitecrow Glacier</u>	<u>5</u>	<u>Glaciers and perennial snowfields</u>
<b><u>Mission Range-Swan Range-Flathead Range</u></b>		
<u>Swan Glaciers</u>	<u>3</u>	<u>Glaciers and perennial snowfields</u>
<b><u>Oregon</u></b>		
<b><u>Cascade Range</u></b>		
<u>Bend Glacier</u>	<u>3</u>	<u>Glaciers and perennial snowfields</u>
<u>Clark Glacier</u>	<u>2</u>	<u>Perennial snowfields only</u>
<u>Collier Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<u>Diller Glacier</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>
<u>Glisan Glacier</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>
<u>Ladd Glacier</u>	<u>4</u>	<u>Glaciers and perennial snowfields</u>
<u>Langille Glacier</u>	<u>5</u>	<u>Glaciers and perennial snowfields</u>
<u>Newton Clark Glacier</u>	<u>3</u>	<u>Glaciers and perennial snowfields</u>
<u>Palmer Glacier</u>	<u>2</u>	<u>Perennial snowfields only</u>
<u>Prouty Glacier</u>	<u>3</u>	<u>Glaciers and perennial snowfields</u>
<u>Renfrew Glacier</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>
<u>Russell Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<u>Sandy Glacier</u>	<u>4</u>	<u>Glaciers and perennial snowfields</u>
<u>Skinner Glacier</u>	<u>4</u>	<u>Perennial snowfields only</u>
<u>Waldo Glacier</u>	<u>3</u>	<u>Glaciers only</u>
<u>White River Glacier</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>
<u>Whitewater Glacier</u>	<u>3</u>	<u>Glaciers only</u>
<u>Zigzag Glacier</u>	<u>3</u>	<u>Glaciers and perennial snowfields</u>
<b><u>Washington</u></b>		
<b><u>Cascade Range-Northern</u></b>		
<u>Borealis Glacier</u>	<u>4</u>	<u>Glaciers only</u>
<u>Buckner Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<u>Butterfly Glacier</u>	<u>4</u>	<u>Glaciers only</u>
<u>Colchuck Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<u>Company Glacier</u>	<u>3</u>	<u>Glaciers only</u>
<u>Cool Glacier</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>
<u>Dana Glacier</u>	<u>3</u>	<u>Glaciers only</u>
<u>Dark Glacier</u>	<u>3</u>	<u>Glaciers only</u>
<u>Dome Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<u>Douglas Glacier</u>	<u>4</u>	<u>Glaciers and perennial snowfields</u>
<u>Dusty Glacier</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>
<u>East Nooksack Glacier</u>	<u>5</u>	<u>Glaciers only</u>
<u>Entiat Glacier</u>	<u>4</u>	<u>Glaciers and perennial snowfields</u>
<u>Forbidden Glacier</u>	<u>2</u>	<u>Glaciers only</u>

<a href="#"><u>Fremont Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>Goode Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>Hadley Glacier</u></a>	<a href="#"><u>5</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>Hanging Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>Hinman Glacier</u></a>	<a href="#"><u>4</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>Honeycomb Glacier</u></a>	<a href="#"><u>3</u></a>	<a href="#"><u>Glaciers and perennial snowfields</u></a>
<a href="#"><u>Inspiration Glacier</u></a>	<a href="#"><u>3</u></a>	<a href="#"><u>Glaciers and perennial snowfields</u></a>
<a href="#"><u>Isella Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Glaciers and perennial snowfields</u></a>
<a href="#"><u>Jerry Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>Kimtah Glacier</u></a>	<a href="#"><u>3</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>LeConte Glacier</u></a>	<a href="#"><u>7</u></a>	<a href="#"><u>Glaciers and perennial snowfields</u></a>
<a href="#"><u>Lyall Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Perennial snowfields only</u></a>
<a href="#"><u>Mazama Glacier</u></a>	<a href="#"><u>3</u></a>	<a href="#"><u>Glaciers and perennial snowfields</u></a>
<a href="#"><u>McAllister Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>Middle Cascade Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>Neve Glacier</u></a>	<a href="#"><u>3</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>No Name Glacier</u></a>	<a href="#"><u>5</u></a>	<a href="#"><u>Glaciers and perennial snowfields</u></a>
<a href="#"><u>Nohokomeen Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>North Klawatti Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Glaciers and perennial snowfields</u></a>
<a href="#"><u>Pilz Glacier</u></a>	<a href="#"><u>3</u></a>	<a href="#"><u>Glaciers and perennial snowfields</u></a>
<a href="#"><u>Price Glacier</u></a>	<a href="#"><u>4</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>Ptarmigan Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Glaciers and perennial snowfields</u></a>
<a href="#"><u>Queest-alb Glacier (not official)</u></a>	<a href="#"><u>3</u></a>	<a href="#"><u>Glaciers and perennial snowfields</u></a>
<a href="#"><u>Rainbow Glacier</u></a>	<a href="#"><u>3</u></a>	<a href="#"><u>Glaciers and perennial snowfields</u></a>
<a href="#"><u>Redoubt Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>Richardson Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>S Glacier</u></a>	<a href="#"><u>3</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>Sandalee Glacier</u></a>	<a href="#"><u>4</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>Scimitar Glacier</u></a>	<a href="#"><u>3</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>Sholes Glacier</u></a>	<a href="#"><u>4</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>Sitkum Glacier</u></a>	<a href="#"><u>4</u></a>	<a href="#"><u>Glaciers and perennial snowfields</u></a>
<a href="#"><u>Snow Creek Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Perennial snowfields only</u></a>
<a href="#"><u>South Cascade Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>Spider Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>Suiattle Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>Sulphide Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>Thunder Glacier</u></a>	<a href="#"><u>3</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>Thunder Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>White Chuck Glacier</u></a>	<a href="#"><u>5</u></a>	<a href="#"><u>Glaciers and perennial snowfields</u></a>
<a href="#"><u>White Salmon Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>Wyeth Glacier</u></a>	<a href="#"><u>3</u></a>	<a href="#"><u>Glaciers and perennial snowfields</u></a>
<b><u>Cascade Range-Southern</u></b>		
<a href="#"><u>Adams Glacier</u></a>	<a href="#"><u>4</u></a>	<a href="#"><u>Glaciers and perennial snowfields</u></a>
<a href="#"><u>Avalanche Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Glaciers only</u></a>
<a href="#"><u>Conrad Glacier</u></a>	<a href="#"><u>3</u></a>	<a href="#"><u>Glaciers and perennial snowfields</u></a>
<a href="#"><u>Cowlitz Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Glaciers and perennial snowfields</u></a>
<a href="#"><u>Crescent Glacier</u></a>	<a href="#"><u>2</u></a>	<a href="#"><u>Glaciers and perennial snowfields</u></a>
<a href="#"><u>Flett Glacier</u></a>	<a href="#"><u>6</u></a>	<a href="#"><u>Glaciers and perennial snowfields</u></a>
<a href="#"><u>Fryingpan Glacier</u></a>	<a href="#"><u>5</u></a>	<a href="#"><u>Glaciers and perennial snowfields</u></a>

<u>Gotchen Glacier</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>
<u>Kautz Glacier</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>
<u>Klickitat Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<u>Lava Glacier</u>	<u>3</u>	<u>Glaciers and perennial snowfields</u>
<u>McCall Glacier</u>	<u>6</u>	<u>Glaciers and perennial snowfields</u>
<u>Meade Glacier</u>	<u>5</u>	<u>Perennial snowfields only</u>
<u>North Mowich Glacier</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>
<u>Ohanapecosh Glacier</u>	<u>6</u>	<u>Glaciers and perennial snowfields</u>
<u>Paradise Glacier</u>	<u>3</u>	<u>Glaciers and perennial snowfields</u>
<u>Pinnacle Glacier</u>	<u>3</u>	<u>Glaciers and perennial snowfields</u>
<u>Puyallup Glacier</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>
<u>Pyramid Glacier</u>	<u>4</u>	<u>Glaciers and perennial snowfields</u>
<u>Russell Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<u>Sarvant Glaciers</u>	<u>4</u>	<u>Glaciers and perennial snowfields</u>
<u>South Mowich Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<u>South Tahoma Glacier</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>
<u>Success Glacier</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>
<u>Van Trump Glacier</u>	<u>10</u>	<u>Glaciers and perennial snowfields</u>
<u>White Salmon Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<u>Whitman Glacier</u>	<u>5</u>	<u>Glaciers and perennial snowfields</u>
<u>Wilson Glacier</u>	<u>3</u>	<u>Glaciers and perennial snowfields</u>
<b><u>Olympic Mountains</u></b>		
<u>Blue Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<u>Cameron Glaciers</u>	<u>4</u>	<u>Glaciers and perennial snowfields</u>
<u>Carrie Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<u>Eel Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<u>White Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<b><u>Wyoming</u></b>		
<b><u>Teton Range</u></b>		
<u>Middle Teton Glacier</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>
<u>Triple Glaciers</u>	<u>3</u>	<u>Glaciers only</u>
<b><u>Wind River Range</u></b>		
<u>Bull Lake Glacier</u>	<u>3</u>	<u>Glaciers and perennial snowfields</u>
<u>Dinwoody Glacier</u>	<u>2</u>	<u>Glaciers only</u>
<u>Dinwoody Glaciers</u>	<u>3</u>	<u>Glaciers and perennial snowfields</u>
<u>Grasshopper Glacier</u>	<u>3</u>	<u>Glaciers only</u>
<u>Harrower Glacier</u>	<u>2</u>	<u>Perennial snowfields only</u>
<u>Helen Glacier</u>	<u>3</u>	<u>Glaciers only</u>
<u>Lower Fremont Glacier</u>	<u>4</u>	<u>Glaciers and perennial snowfields</u>
<u>Mammoth Glacier</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>
<u>Minor Glacier</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>
<u>Sacagawea Glacier</u>	<u>4</u>	<u>Glaciers and perennial snowfields</u>
<u>Sourdough Glacier</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>
<u>Stroud Glacier</u>	<u>3</u>	<u>Glaciers and perennial snowfields</u>
<u>Twins Glacier</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>
<u>Upper Fremont Glacier</u>	<u>2</u>	<u>Glaciers and perennial snowfields</u>

**Formatted:** Font color: Auto  
**Formatted:** Space After: 8 pt, Line spacing: Multiple  
1.08 li



430 **A3 Labelling errors in the U.S. Geographic Names Information System**  
 431  
 432 **Table A3.** List of officially named glaciers where we identified an issue with the glacier name  
 433 on the 1:24000 U.S. Geological Survey topographical maps (Fountain et al., 2017). Names come  
 434 from the Geographic Names Information System ([https://www.usgs.gov/tools/geographic-](https://www.usgs.gov/tools/geographic-names-information-system-gnis)  
 435 [names-information-system-gnis](https://www.usgs.gov/tools/geographic-names-information-system-gnis)). The 'Issue' column lists the type of issue identified. 'Not  
 436 labeled' indicates the feature was present but not labeled, 'Misidentified' indicates the wrong  
 437 feature was labeled, and 'Label unclear' means the location of the label is not clearly associated  
 438 with a specific glacier.

<u>State/Region/Glacier Name</u>	<u>Issue</u>
<u>Colorado</u>	
<u>Front Range</u>	
<u>Arikaree Glacier</u>	<u>Not labeled</u>
<u>Navajo Glacier</u>	<u>Not labeled</u>
<u>Oregon</u>	
<u>Cascade Range</u>	
<u>Carver Glacier</u>	<u>Misidentified</u>
<u>Milk Creek Glacier</u>	<u>Not labeled</u>
<u>Washington</u>	
<u>Cascade Range-Northern</u>	
<u>S Glacier</u>	<u>Label unclear</u>
<u>Snow Creek Glacier</u>	<u>Label unclear</u>
<u>South Glacier</u>	<u>Not labeled</u>
<u>Cascade Range-Southern</u>	
<u>No Name Glacier</u>	<u>Not labeled</u>
<u>Stevens Glacier</u>	<u>Not labeled</u>
<u>Wyoming</u>	
<u>Wind River Range</u>	
<u>Dinwoody Glaciers</u>	<u>Label unclear</u>
<u>Fremont Glaciers</u>	<u>Label unclear</u>

439  
 440  
 441 **A4 Notes on imagery and interpretation challenges by State.**  
 442

- Formatted: Font: Bold, Font color: Text 1
- Formatted: Indent: Left: 0.25"
- Formatted: List Paragraph

443 **4. Data products and availability**  
 444  
 445 ~~The data are available in three formats. The geospatial data and attribute tables are available in~~  
 446 ~~the shapefile (Esri) format and in an open source GeoJSON format. The attribute table is also~~  
 447 ~~available as an EXCEL file. These data products can be obtained from~~  
 448 ~~[https://doi.org/10.15760/geology\\_data\\_03](https://doi.org/10.15760/geology_data_03) (Fountain and Glenn, 2022) and from the Global Land~~  
 449 ~~Ice Measurements from Space website (to be submitted) <http://glims.colorado.edu/glacierdata/>~~  
 450

451 **5. Summary**  
 452

453 ~~We have compiled a new and comprehensive inventory of glaciers and perennial snowfields in~~  
454 ~~the western US from aerial and satellite imagery. Results show that 2542 glacial features are~~  
455 ~~currently present and include 1231 (366.52 km<sup>2</sup>) glaciers, 1176 (31.01 km<sup>2</sup>) perennial~~  
456 ~~snowfields, and 35 (3.57 km<sup>2</sup>) buried ice deposits. Most of the data were acquired from 2015~~  
457 ~~NAIP imagery with the remainder from NAIP imagery and a few satellite images acquired over~~  
458 ~~the period of 2013 to 2020. The state of Washington has the greatest number and area of glaciers~~  
459 ~~and perennial snowfields. This product updates an older inventory based on USGS 1:24000 maps~~  
460 ~~compiled in the middle-late 1900's. The new inventory is a significant improvement in accuracy~~  
461 ~~because the archive of historical imagery in Google Earth allowed us to classify glaciers versus~~  
462 ~~perennial snowfields. Finally, this new inventory provides a baseline for assessing glacier change~~  
463 ~~in the coterminous US.~~

## 464 ~~6. Appendix~~

465 ~~This appendix, organized by US State, then by mountain range, summarizes the specific~~  
466 ~~imagery used, challenges encountered in feature identification and outline digitization. The~~  
467 ~~most recent suitable NAIP was used in each case. Where such imagery was not suitable~~  
468 ~~Maxar imagery was used. In the Willowa Mountains, Oregon, neither was NAIP suitable nor~~  
469 ~~was Maxar available so images from Google Earth were used. The Selkowitz and Forster~~  
470 ~~(2016) inventory is referred to as the SFI and the National Land Cover Database inventory~~  
471 ~~(Jin et al., 2019) is referred to as the NLCD.~~

### 472 ~~California~~

473 ~~Imagery and DEMs used are listed in Tables A1, A2, A3.~~

474 ~~This appendix, organized by US State, then by mountain range, summarizes the specific~~  
475 ~~imagery used, challenges encountered in feature identification and outline digitization. The~~  
476 ~~Selkowitz and Forster (2016) inventory is referred to as the SFI and the National Land Cover~~  
477 ~~Database inventory (Dewitz, 2019) is referred to as the NLCD.~~

#### 478 ~~A4.1 California~~


479 ~~Imagery and DEMs used are listed in Tables A4, A5, A6.~~

### 480 ~~Cascade Range~~

#### 481 ~~Mount Shasta~~

482 ~~The 2020 black and white Maxar imagery was most useful because of the minimal~~  
483 ~~seasonal snow cover. The 2018 NAIP imagery was helpful in situations where the 2020~~  
484 ~~imagery was obscured by shadow, distortion, or misaligned, and when color was needed~~  
485 ~~to improve interpretation. The 2010 lidar DEM (Robinson, 2014; Table ~~A3A4~~) was used~~  
486 ~~to create a multidirectional hillshade to improve perspective and interpretation (Figure~~  
487 ~~A1).~~

498 The rock debris on the termini of most glaciers and rock debris on some of the upper  
499 parts of the glaciers were challenging to interpret. It was hard to determine whether ice  
500 was present under the debris and whether that ice is part of the active glacier. Spatial  
501 patterns of debris, debris contrasts, and melt streams flowing from the debris were used to  
502 estimate the glacier boundaries.



503  
504 **Figure A1.** Mt. Shasta glaciers in bluish white, perennial snowfields/ice patches in lavender  
505 draped over a 3D rendering created from 2010 lidar (Robinson, 2014).  
506

#### 507 **Sierra Nevada**

509 The 2014 NAIP imagery was the best imagery due to low snow cover. In some cases,  
510 features were difficult to outline because of shadow or image quality. In these cases,  
511 2013/2012 Google Earth imagery were used. Some glaciers were reclassified as rock  
512 glaciers by Trcka (2020). These were re-examined and where we agreed they were  
513 removed from the initial glacier inventory. Defining whether the feature was a glacier or  
514 rock glacier was often difficult, see Colorado section for more discussion.  
515

#### 516 **Trinity Alps**

517 The 2018 imagery was the best for the least snow cover. Justin Garwood (Garwood et al.,  
518 2020) provided outlines for two glaciers, Grizzly and Salmon. The area of the most recent  
519 outline of the Salmon Glacier was  $< 0.01 \text{ km}^2$  and was not included in this inventory. By  
520 2018 all of the other features mapped by the USGS (Fountain et al., 2017) were less than  
521  $0.01 \text{ km}^2$  or had disappeared. ~~An additional feature was added based on the 2016 NLCD~~  
522 ~~(Jin et al. 2019). An additional feature was added based on the 2016 NLCD (Jin et al.,~~  
523 ~~2019).~~

524  
525  
526 **Table A1A4.** List of NAIP imagery used for outlining glaciers and perennial snowfields in  
527 California. 'Date' is the start and end date for flights covering the glaciated portions of the NAIP

528 image. In some cases, flights were completed in a single day.  
 529

Region/Year/Filename	County	Date (Year-M-D)
<b>Cascade Range</b>		
2014		
ortho_1-1_1n_s_ca089_2014_1.sid	Shasta	2014-07-13
ortho_1-1_1n_s_ca093_2014_1.sid	Siskiyou	2014-06-23 to 2014-07-18
2018		
ortho_1-1_hn_s_ca093_2018_1.sid	Siskiyou	2018-07-21 to 2018-09-25
<b>Sierra Nevada</b>		
2014		
ortho_1-1_1n_s_ca019_2014_1.sid	Fresno	2014-07-23 to 2014-08-23
ortho_1-1_1n_s_ca027_2014_1.sid	Inyo	2014-07-23 to 2014-08-23
ortho_1-1_1n_s_ca039_2014_2.sid	Madera	2014-07-18 to 2014-08-15
ortho_1-1_1n_s_ca051_2014_1.sid	Mono	2014-07-17 to 2014-08-15
ortho_1-1_1n_s_ca107_2014_1.sid	Tulare	2014-08-23 to 2014-08-23
<b>Trinity Alps</b>		
2018		
ortho_1-1_hn_s_ca093_2018_1.sid	Siskiyou	2018-07-21 to 2018-09-25

530  
 531  
 532 **Table A2A5.** List of dates of the Maxar imagery used for outlining glaciers and perennial  
 533 snowfields in California.

**Region/ Date (Year-M-D)**

**Cascade Range**

2020-10-05

534  
 535  
 536 **Table A3A6.** List of U.S. Geological Survey digital elevation models used for outlining glaciers  
 537 and perennial snowfields in California.

Filename	Date	Citation	URL
ds852_lidar	2010	Robinson (2014)	<a href="https://pubs.er.usgs.gov/publication/ds852">https://pubs.er.usgs.gov/publication/ds852</a>

538  
 539 **6A4.2 Colorado**  
 540  
 541 The 2015 NAIP was generally free of seasonal snow. Where it persisted at the terminus of a few  
 542 glaciers, images for the same year in Google Earth aided perimeter interpretation. Imagery used  
 543 are listed in Table A4A7.

**Elk Mountains**

No ~~glacial~~ features were mapped in the Elk Mountains by the USGS (Fountain et al., 2017). One glacier and four perennial snowfields were added from the SFI.

### Front Range

The most recent inventory for the Front Range was Hoffman et al. (2007), which used aerial photographs to map the 2001 extent of glaciers. Many features in the Front Range are difficult to classify. The issue is the difference between a glacier or perennial snowfield and a rock glacier. Those that are part of the rock glacier are deleted from the glacier inventory. Those that seem to be separate from rock glaciers are retained. This is a judgement call. From a hydrological point of view, if a snow-ice patch that is part of a rock glacier was counted separately from a rock glacier, it is double counting a water feature.

~~The most challenging situation to interpret occurs when the glacier or perennial snowfield is located up elevation from the rock glacier. If the slope of the snowfield smoothly transitions to the slope of the rock glacier, with no change in sign of the slope, we consider that one feature, a rock glacier (Figure A2). If the terrain dips below the snowfield, changing sign to rise to a topographic high below which the rock glacier clearly emerges, then they are two separate features. The patch does not appear to feed the rock glacier with ice (ice melt maybe, but not ice), because the ice would have to flow uphill to reach the rock glacier (Figure A3).~~

Formatted: Font color: Auto

**Table A4A7.** List of NAIP imagery used for outlining glaciers and perennial snowfields in Colorado. 'Date' is the start and end date for flights covering the glaciated portions of the NAIP image. In some cases, flights were completed in a single day.

Region/Year/Filename	County	Date (Year-M-D)
<b>Elk Mountains</b>		
2015		
ortho_1-1_1n_s_co051_2015_1.sid	Gunnison	2015-09-10 to 2015-09-11
<b>Front Range</b>		
2015		
ortho_1-1_1n_s_co013_2015_1.sid	Boulder	2015-08-25 to 2015-09-20
ortho_1-1_1n_s_co049_2015_1.sid	Grand	2015-08-25 to 2015-09-20
ortho_1-1_1n_s_co057_2015_1.sid	Jackson	2015-09-09
ortho_1-1_1n_s_co069_2015_1.sid	Larimer	2015-08-25 to 2015-09-09
<b>Gore Range</b>		
2015		
ortho_1-1_1n_s_co037_2015_1.sid	Eagle	2015-09-10
<b>Medicine Bow Mountains</b>		
2015		
ortho_1-1_1n_s_co057_2015_1.sid	Jackson	2015-09-09
<b>Park Range</b>		

2015  
 ortho\_1-1\_1n\_s\_co057\_2015\_1.sid Jackson 2015-09-09

**San Miguel Mountains**

2015  
 ortho\_1-1\_1n\_s\_co033\_2015\_1.sid Dolores 2015-09-11  
 ortho\_1-1\_1n\_s\_co091\_2015\_1.sid Ouray 2015-09-11  
 ortho\_1-1\_1n\_s\_co111\_2015\_1.sid San Juan 2015-09-12

**Sawatch Range**

2015  
 ortho\_1-1\_1n\_s\_co037\_2015\_1.sid Eagle 2015-09-10  
 ortho\_1-1\_1n\_s\_co097\_2015\_1.sid Pitkin 2015-09-10 to 2015-09-11

574

575 **6A4.3 Idaho**

576

577 The imagery quality was generally snow free. Of the glacier mapped by the USGS (Fountain et  
 578 al., 2017) only two remain and are classified as perennial snowfields. The Borah Glacier was  
 579 officially named in 2021, ([U.S. Board of Geographic Names](#)), but is < 0.01 km<sup>2</sup>, and is not  
 580 included in the inventory. Table [A5A8](#) lists the imagery used.

581

582

583 **Table A5A8.** List of NAIP imagery used for outlining glaciers and perennial snowfields in  
 584 Idaho. ‘Date’ is the start and end date for flights covering the glaciated portions of the NAIP  
 585 image. In some cases, flights were completed in a single day.

586

<b>Region/Year/Filename</b>	<b>County</b>	<b>Date (Year-M-D)</b>
<b>Sawtooth Range</b>		
2013		
ortho_1-1_hn_s_id015_2013_1.sid	Boise	2013-09-07
2015		
ortho_1-1_1n_s_id013_2015_1.sid	Blaine	2015-07-30
ortho_1-1_1n_s_id015_2015_1.sid	Boise	2015-09-08 to 2015-09-09
2019		
ortho_1-1_hn_s_id037_2019_1.sid	Custer	2019-07-25 to 2019-08-26


587

Formatted: Font: Not Bold



588

589 **Figure A2.** An example of a snowfield that is considered part of the rock glacier. Location,  
590 Colorado Front Range, 40.827477° N, 106.657400° E. Image is from © Google Earth, 9/2014.



Formatted: Keep with next

591  
592

593 **Figure A3.** Tyndall Glacier in the Colorado Front Range, 40.305291° N, -105.689602° E, with a  
594 rock glacier slightly down valley. Image is from © Google Earth 9/2016.

Formatted: Font: +Body (Calibri), 11 pt, Not Bold, Not Italic

Formatted: Font: Not Italic, Font color: Auto

Formatted: Caption

Formatted: Font: Not Italic, Font color: Auto

Formatted: Font: Not Italic, Font color: Auto, Pattern: Clear

Formatted: Font: Not Italic, Font color: Auto

Formatted: Font: Bold, Italic

Formatted: Space After: 8 pt

595

596 **6**  
597 **A4.4 Montana**

598  
599 Image quality varied between mountain ranges due to differences in snow cover. Tables [A6A9](#)  
600 and [A7A10](#) list the imagery used.

601 **Beartooth-Absaroka Range**

602 The 2015 NAIP imagery was the best overall imagery due to the least snow, but Google  
603 Earth was occasionally used as well. Google Earth had imagery dated to 9/11/2015; often  
604 with less seasonal snow than the NAIP imagery. To counter any mismatch in projection,  
605



606 outlines digitized in Google Earth were imported to ArcGIS and projected to match the  
607 NAIP projection.

608 **Bitterroot Range**

609 ~~There were no glacial~~No features were mapped in the Bitterroot Range by the USGS  
610 (Fountain et al., (2017). One glacier and three perennial snowfields were added based on  
611 the NLCD.

612  
613 **Cabinet Range**

614 The USGS mapped four ~~glacial~~ features  $\geq 0.01$  km<sup>2</sup> (Fountain et al., 2017). Inspection of  
615 the 2015 only one was  $\geq 0.01$  km<sup>2</sup>. Seven glaciers and perennial snowfields were added;  
616 five were identified in our initial inventory, the other two were identified by the SFI and  
617 NLCD, respectively. All were less than 0.05 km<sup>2</sup>.

618  
619 **Crazy Mountains**

620 The 2013 NAIP imagery was the best imagery available and included limited seasonal  
621 snow. The 2019 Maxar imagery had too much seasonal snow.

622  
623 **Lewis Range (Glacier National Park)**


624 The most recent published glacier inventory is a 2015 USGS inventory (Fagre et al.,  
625 2017). They outlined the main-body of named-glaciers using 2015 Maxar imagery. We  
626 digitized the outlines of all glaciers and perennial snowfields using 2015 Maxar imagery  
627 where available. Elsewhere, 2015 and 2013 NAIP imagery were used; both years had lots  
628 of seasonal snow cover. Two major glaciers, Blackfoot (Figure ~~A4A2~~) and Harrison  
629 (Figure ~~A5A3~~) glaciers, separated into pieces as it retreated since it was originally  
630 mapped by the USGS (Fountain et al., 2007).

631  
632 **Madison Range**


633 The 2013 NAIP imagery was the only ~~good~~ imagery ~~used~~ due to extensive snow in the  
634 other years. No glaciers or perennial snowfields were found. Of the two features  $\geq 0.01$   
635 km<sup>2</sup> mapped by the USGS (Fountain et al., 2017), the 2013 imagery showed that one  
636 feature is a rock glacier and the other was less than 0.01 km<sup>2</sup>.

637  
638 **Mission-Swan-Flathead Ranges**



639 Based on the least snow cover, the 2013 NAIP was better in the Mission and Flathead  
640 Ranges, and the 2015 NAIP was better in the Swan Range. No glaciers or perennial  
641 snowfields remain in the Flathead Range.



642  
643



644  
645 **Figure A4A2.** The updated (2015) outlines for the Blackfoot Glacier including the main glacier  
646 body (red) and the additional smaller glacier (orange). Base image from the NAIP taken in 2013.



648 **Figure A5A3.** The updated (2015) outlines for Harrison Glacier including the main glacier body  
 649 (red) and the additional smaller glaciers (orange). Base image from the NAIP taken in 2013.

650

651

652 **Table A6A9.** List of NAIP imagery used for outlining glaciers and perennial snowfields in  
 653 Montana. 'Date' is the start and end date for flights covering the glaciated portions of the NAIP  
 654 image. In some cases, flights were completed in a single day.

655

<b>Region/Year/Filename</b>	<b>County</b>	<b>Date (Year-M-D)</b>
<b>Beartooth Mountains-Absaroka Range</b>		
2013		
ortho_1-1_1n_s_mt067_2013_1.sid	Park	2013-08-05 to 2013-09-11
2015		
ortho_1-1_1n_s_mt009_2015_1.sid	Carbon	2015-08-10 to 2015-09-07
ortho_1-1_1n_s_mt067_2015_1.sid	Park	2015-08-19 to 2015-09-11
ortho_1-1_1n_s_mt095_2015_1.sid	Stillwater	2015-08-10 to 2015-09-07
<b>Bitterroot Range</b>		
2013		
ortho_1-1_1n_s_mt001_2013_1.sid	Beaverhead	2013-08-04
2015		
ortho_1-1_1n_s_mt081_2015_2.sid	Ravalli	2015-10-06 to 2015-11-07
<b>Cabinet Mountains</b>		
2015		
ortho_1-1_1n_s_mt053_2015_2.sid	Lincoln	2015-09-11 to 2016-08-15
<b>Crazy Mountains</b>		
2013		
ortho_1-1_1n_s_mt067_2013_1.sid	Park	2013-08-05 to 2013-09-11
ortho_1-1_1n_s_mt097_2013_1.sid	Sweet Grass	2013-08-31 to 2013-09-10
2015		
ortho_1-1_1n_s_mt067_2015_1.sid	Park	2015-08-19 to 2015-09-11
<b>Lewis Range</b>		
2013		
ortho_1-1_1n_s_mt029_2013_1.sid	Flathead	2013-08-21 to 2013-09-01
ortho_1-1_1n_s_mt035_2013_1.sid	Glacier	2013-08-21 to 2013-09-01
2015		
ortho_1-1_1n_s_mt029_2015_2.sid	Flathead	2015-09-30 to 2016-10-21
ortho_1-1_1n_s_mt035_2015_2.sid	Glacier	2015-10-14 to 2016-08-21
<b>Mission Range-Swan-Flathead Ranges</b>		
2013		
ortho_1-1_1n_s_mt029_2013_1.sid	Flathead	2013-08-21 to 2013-09-01
ortho_1-1_1n_s_mt063_2013_1.sid	Missoula	2013-09-01
2015		
ortho_1-1_1n_s_mt047_2015_2.sid	Lake	2015-09-12 to 2016-08-15
ortho_1-1_1n_s_mt063_2015_2.sid	Missoula	2015-09-12 to 2016-08-16

656

657  
658 **Table A7A10.** List of dates of the Maxar imagery used for outlining glaciers and perennial  
659 snowfields in Montana.  
660

**Region/ Date (Year-M-D)**

**Lewis Range**

2015-08-22  
2015-09-01  
2015-09-12  
2015-09-25  
2019-08-20

661  
662 **A4.5 Oregon**

663  
664 Tables ~~A8~~, ~~A9~~**A11**, **A12**, and ~~A10~~**A13** list the imagery and DEM used.

**Cascade Range**

~~Seasonal snow cover was commonly present when this range was imaged by any of the sensors making it difficult to find suitable imagery.~~

Formatted: Indent: Left: 0.5"

Formatted: Font: Not Bold, Not Highlight

Formatted: Not Highlight

**Mount Hood**

The most recent glacier outlines for Mt. Hood were based on 2015 and 2016 Maxar color imagery with interpretation aid using Google Earth. Due to seasonal snow some professional judgement was required in places.

**Mount Jefferson**

~~Seasonal snow was extensive in places.~~ The 2018 NAIP had extensive seasonal snow and ~~was~~ generally only useful near the terminus of some glaciers. Used 2018 Maxar imagery that showed little seasonal snow, but a little cloudy that masked a bit of Whitewater Glacier. Also used Google Earth to help interpret some of the features.

**Three Sisters**

~~Used~~ Maxar 2018 imagery ~~was used~~, but ~~where~~the image ~~was~~ stretching along the feature's headwall ~~and for that segment of the outline~~ 2018 NAIP imagery was used. Two versions of the Maxar imagery for the same day are ~~3~~~~available~~available, one color, one black and white. Color was georectified but suffered stretching along some headwalls. A light early season snowfall occurred before the Maxar image and the snow accumulated in some places just enough to obscure the surface. So, the glacier or snow patch outline was the minimum of the two images with occasional interpolation across the snowy surface to the nearest glacier edge.

~~An example of buried ice on South Sister is shown in Figure A6.~~

Formatted: Font: +Body (Calibri)

**Mount Thielsen**

695 The Lathrop Glacier was named in 1981. At the time of the USGS mapping and now it is  
696 <math><0.01\text{ km}^2</math>, and not counted as part of the inventory. Furthermore, Lathrop Glacier has  
697 been known to disappear in some years and therefore fails the definition of a glacier.  
698

### 699 **Wallowa Mountains**

700 No NAIP imagery was useful and Maxar did not image this region. We used the  
701 8/30/2013 image from Google Earth, which was excellent with little snow. Features were  
702 digitized in Google Earth and then imported into ArcGIS. Because we used NAIP as the  
703 base imagery, we revised the outline from the projection in WGS84 (Google Earth) to  
704 NAD83 UTM Zone 11 (NAIP).  
705



706  
707 **Figure A6.** *Lost Creek Glacier, South Sister, Oregon. Note buried ice and lack of crevasses to*  
708 *the left of the grey blue ice, suggesting ice that is no longer moving and therefore not part of the*  
709 *dynamic glacier. The white box surrounds an area that has collapsed due to subsurface melt.*  
710 *The inset enlargement shows a cliff edge of exposed dirty ice (white arrow) indicated by a*  
711 *darker color suggesting wet sediment and a finer texture than the surface debris. The black*  
712 *arrow shows the width of the cleaner ice for scale. Image is from © Google Earth, 8/9/2021.*  
713

Formatted: Font: Italic, Font color: Text 2

Formatted: Font: Italic, Font color: Text 2

Formatted: Font: Italic

Formatted: Font: Italic, Font color: Text 2

714  
 715 **Table A8A11.** List of NAIP imagery used for outlining glaciers and perennial snowfields in  
 716 Oregon. ‘Date’ is the start and end date for flights covering the glaciated portions of the NAIP  
 717 image. In some cases, flights were completed in a single day.  
 718

Region/Year/Filename	County	Date (Year-M-D)
<b>Cascade Range</b>		
2014		
ortho_1-1_1n_s_or017_2014_1.sid	Deschutes	2014-09-01
ortho_1-1_1n_s_or027_2014_1.sid	Hood River	2014-08-27 to 2014-09-05
ortho_1-1_1n_s_or039_2014_1.sid	Lane	2014-09-01
2016		
ortho_1-1_1n_s_or027_2016_1.sid	Hood River	2016-08-04
2017/2018		
ortho1-1_hn_s_or017_2017_2018_1.sid	Deschutes	2018-07-28
<b>Wallowa Mountains</b>		
2014		
ortho_1-1_1n_s_or063_2014_1.sid	Wallowa	2014-10-05

719  
 720  
 721 **Table A9A12.** List of dates of the Maxar imagery used for outlining glaciers and perennial  
 722 snowfields in Oregon.  
 723

Region/ Date (Year-M-D)
<b>Cascade Range</b>
2015-08-20
2015-09-11
2015-10-05
2016-09-10
2018-09-17
2020-09-20

724  
 725  
 726 **Table A10A13.** List of Oregon Department of Geology and Mineral Industries digital elevation  
 727 models used for outlining glaciers and perennial snowfields in Oregon. <sup>5</sup>  
 728

Filename	Date	URL
2011_OLC_Deschutes	2011	gis.dogami.oregon.gov/maps/lidarviewer/

729  
 730  
 731 **6A4.6 Washington**  
 732

733 The 2015 NAIP imagery was typically excellent with little snow cover, whereas the 2017 NAIP  
 734 had more snow and the 2019 imagery had lots of snow. For most outlines, 2015 NAIP imagery  
 735 was used. In some places, the 2017 NAIP imagery had less snow and was used instead. Maxar



736 imagery was of limited use and often wasn't better than the 2015 or 2017 NAIP. Tables ~~A11,~~  
737 ~~A12, and A13~~A14, A15, A16, list the imagery and DEMs used.

738

739

#### Cascade –Northern

740

The glaciers and perennial snowfields were previously inventoried by (Dick, 2013).

741

742

#### Mount Baker

743

The 2015 NAIP imagery was the best and had little seasonal snow. Google Earth 2009  
744 and 2019 imagery were used to help interpretation. A multidirectional hillshade and 3-  
745 meter contour lines derived from a lidar DEM (Bard, ~~2017b; Table A13~~2017a); were  
746 used to help define flow divides between glaciers, debris covered-ice, and buried ice.

747

There are notable differences between the NAIP imagery and DEM data, particularly in  
748 steep terrain, areas of dark shadow, and debris-covered areas. The DEM helped correct  
749 these positional errors and the benefit of supplying more information on surface texture.

750

Several buried-ice features were identified. The ice appeared to have decoupled from the  
751 active glacier. In a few cases, debris-covered ice is included in the glacier outline because  
752 the ice appears to be directly connected to the glacier, and there was evidence of  
753 movement.

754

755

#### Dragontail Peak

756

The [USGS Geographic Names Information Service \(GNIS\)](#) locates Snow Creek Glacier  
757 at a point on the edge of the southeast glacier (Fountain et al., 2007). In the 2015  
758 imagery, the point is on bedrock, making it unclear which glacier the GNIS is naming.  
759 The USGS identifies both glaciers as Snow Creek Glacier. We labeled both glaciers as  
760 the Snow Creek Glacier.

761

762

#### Glacier Peak

763

For the Glacier Peak region, a multidirectional hillshade and 3-m contour lines derived  
764 from a 2015 lidar DEM (Bard, ~~2017A; Table A13~~ ~~was~~2017b) were used as a guide to  
765 define flow divides.

766

767

#### Hurry-up Peak

768

The point location of the South Glacier provided by the GNIS is over bedrock. We  
769 assume the point refers to the glacier located ~150 m to the north of the point.

770

771

#### Cascade –Southern

772

773

#### Goat Rocks

774

Imagery from 2015 was best, but ~~had~~ more snow than desired. Too much snow ~~was~~  
775 ~~present~~ in 2017 but some ice ~~is~~was exposed. The 2019 imagery was ~~way-too~~ snowy ~~and~~  
776 ~~was considered useless~~ for glacier digitization.

777

778

The outlines are almost entirely based on 2015 imagery, and a few on 2017, where  
779 needed. Used 2009 NAIP imagery to help define the headwalls at the Conrad, McCall,  
780 and Packwood glaciers. Heard (2000) previously mapped the glacier perimeters. The

781

782 maximum extent of the seasonal snow covering the terminal regions was not digitized.  
783 Typically digitized at scales of 1:600 to 1:800. Note that narrow arms of the snowfields  
784 were not typically digitized knowing that they would probably disappear a few days to a  
785 week from the time of imagery.

#### 786 **Mount Adams**

787 No suitable NAIP imagery was found, instead 2019 Maxar imagery was used. In addition  
788 to the Maxar imagery, a multidirectional hillshade and 3-m contour lines derived from a  
789 2016 lidar DEM (Bard 2019, [Table 13A](#)) were used as a guide when delignating flow  
790 divides. Occasionally, 2009 Google Earth imagery was also useful. Extensive snow  
791 covered the mountain when the 2016 lidar was flown masking some of the glacier  
792 termini. However, the DEM was helpful in correcting the imagery where poorly aligned  
793 with the terrain.

794  
795 Multiple buried-ice features were identified near the terminus of several glaciers where  
796 ice appeared to have decoupled from the main active glacier. Large areas below the  
797 glaciers (Mazama, Adams, and Pinnacle) likely have debris-covered ice. We focused on  
798 the features which were likely to contain ice based on meltwater streams exiting near the  
799 features and hummocky terrain which appeared to indicate melt. Ground-based images  
800 from taken between 2014 to 2018 helped decision-making. The images were particularly  
801 helpful in identifying a debris-covered ice cliff at Adams Glacier.

#### 802 **Mount Rainier**

803  
804 In general, the 2019 NAIP and the Maxar (2018-09-25) were used for the outlines ([Table](#)  
805 [A12](#)). Although the GNIS includes the Nisqually Icefall as a separate feature, we  
806 included the icefall as part of the Nisqually Glacier ([Figure A7A4](#)).


#### 807 **Mount St. Helens**

808  
809 We used a GIS layer of geological mapping units that included snow and ice from the  
810 USGS (David Sherrod, [personal USGS written communication](#), 2021) to help guide our  
811 search. The Crater Glacier (INV\_ID E562842N5115499) was heavily debris covered,  
812 and obscured by shadow in some areas.


#### 813 **Olympic Mountains**

814  
815 A 2015 inventory of the region was compiled because more recent imagery (NAIP and  
816 Maxar) were not useful due to seasonal snow. Our updated inventory differs from that  
817 published in Fountain et al. (2017) in two ways. First, they outlined and grouped the  
818 glaciers and perennial snowfields according to watershed rather than individual glacier.  
819 Their goal was to estimate glacier change relative to a previous study by Spicer (1986)  
820 and had to follow Spicer's approach. Second, all outlines were rechecked and compared  
821 to SFI and the NLCD resulting in minor changes.

822  
823  
824  
825  
826



827



828  
 829 **Figure A7A4.** Image of the Nisqually Glacier and Icefall. The orange and red outlines are from  
 830 the updated inventory and the blue outline is from the USGS mapping (Fountain et al., 2007)  
 831 database. The base image is from the NAIP taken in 2019.

832  
 833  
 834 **Table A11A14.** List of NAIP imagery used for outlining glaciers and perennial snowfields in  
 835 Washington. 'Date' is the start and end date for flights covering the glaciated portions of the  
 836 NAIP image. In some cases, flights were completed in a single day. For 2006 the inspection date  
 837 was used, since the start and end dates were not provided.

838

<u>Region/Year/Filename</u>	<u>County</u>	<u>Date (Year-M-D)</u>
Cascade -Northern		

2006			
ortho_1-1_1n_s_wa007_2006_3.sid	Chelan	2006-07-01	
2015			
ortho_1-1_1n_s_wa007_2015_1.sid	Chelan	2015-07-06 to 2015-09-23	
ortho_1-1_1n_s_wa033_2015_1.sid	King	2015-07-06 to 2015-09-27	
ortho_1-1_1n_s_wa037_2015_1.sid	Kittitas	2015-07-06 to 2015-09-23	
ortho_1-1_1n_s_wa047_2015_1.sid	Okanogan	2015-09-09 to 2015-09-11	
ortho_1-1_1n_s_wa057_2015_1.sid	Skagit	2015-07-06 to 2015-09-29	
ortho_1-1_1n_s_wa061_2015_1.sid	Snohomish	2015-07-06 to 2015-09-29	
ortho_1-1_1n_s_wa073_2015_1.sid	Whatcom	2015-09-10 to 2015-09-26	
2017			
ortho_1-1_1n_s_wa007_2017_1.sid	Chelan	2017-10-03 to 2017-10-24	
ortho_1-1_1n_s_wa057_2017_1.sid	Skagit	2017-09-27 to 2017-10-05	
ortho_1-1_1n_s_wa073_2017_1.sid	Whatcom	2017-09-27 to 2017-10-05	
<b>Cascade –Southern</b>			
2015			
ortho_1-1_1n_s_wa041_2015_1.sid	Lewis	2015-07-15 to 2015-07-29	
ortho_1-1_1n_s_wa053_2015_1.sid	Pierce	2015-07-29	
ortho_1-1_1n_s_wa059_2015_1.sid	Skamania	2015-07-15 to 2015-09-12	
ortho_1-1_1n_s_wa077_2015_1.sid	Yakima	2015-07-15 to 2015-07-29	
2019			
ortho_1-1_hn_s_wa053_2019_1.sid	Pierce	2019-08-26	
ortho_1-1_hn_s_wa059_2019_1.sid	Skamania	2019-08-06 to 2019-08-26	
<b>Olympic Mountains</b>			
2015			
ortho_1-1_1n_s_wa009_2015_1.sid	Clallam	2015-07-28 to 2015-09-12	
ortho_1-1_1n_s_wa031_2015_1.sid	Jefferson	2015-07-28 to 2015-09-12	
ortho_1-1_1n_s_wa045_2015_1.sid	Mason	2015-07-28 to 2015-08-19	

839  
840  
841  
842  
843

**Table A12A15.** List of dates of the Maxar imagery used for outlining glaciers and perennial snowfields in Washington.

**Region/ Date (Year-M-D)**

**Cascade Range-Northern**

2018-09-25

**Cascade Range-Southern**

2018-09-25

2019-08-31

**Olympic Mountains**

2015-08-17

2019-09-30

844  
845  
846

**Table A13A16.** List of U.S. Geological Survey digital elevation models used for outlining

847 glaciers and perennial snowfields in Washington. [To access the data both the URL and specific](#)  
848 [identifier are required.](#)

Region	Date	Citation	URL
Mt. Adams	2016	Bard (2019)	<a href="http://www.sciencebase.gov/catalog/item/5bc623b9e4b0fc368ebbe99a">www.sciencebase.gov/catalog/item/5bc623b9e4b0fc368ebbe99a</a>
<del>Mt. Baker</del> Glacier Peak	<del>2014-15</del> 2015	Bard (2017a)	<del><a href="http://www.sciencebase.gov/catalog/item/58518b0ee4b0f99207c4f12c57bf299ee4b0f2f0ceb7534e">www.sciencebase.gov/catalog/item/58518b0ee4b0f99207c4f12c57bf299ee4b0f2f0ceb7534e</a></del> <a href="http://www.sciencebase.gov/catalog/item/57bf299ee4b0f2f0ceb7534e58518b0ee4b0f99207c4f12e">www.sciencebase.gov/catalog/item/57bf299ee4b0f2f0ceb7534e58518b0ee4b0f99207c4f12e</a>
<del>Mt. Baker</del> Glacier Peak	<del>2015</del> 15	Bard (2017b)	<del><a href="http://www.sciencebase.gov/catalog/item/58518b0ee4b0f99207c4f12c57bf299ee4b0f2f0ceb7534e">www.sciencebase.gov/catalog/item/58518b0ee4b0f99207c4f12c57bf299ee4b0f2f0ceb7534e</a></del> <a href="http://www.sciencebase.gov/catalog/item/57bf299ee4b0f2f0ceb7534e58518b0ee4b0f99207c4f12e">www.sciencebase.gov/catalog/item/57bf299ee4b0f2f0ceb7534e58518b0ee4b0f99207c4f12e</a>

Formatted: Font color: Auto

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Don't add space between paragraphs of the same style

849  
850

## 851 6.7A7 Wyoming

852

853

### Wind River Range

854

855 Tables [A14A17](#) and [A15A18](#) list the imagery used. The 2015 NAIP imagery had little  
856 snow in contrast to 2019 imagery. Shadows are common in the 2015 imagery and can be  
857 very dark. Occasionally the 2019 imagery was used to define the glacier-bedrock  
858 headwall boundary. The 2019 Maxar imagery was essentially identical to the NAIP and  
859 because its black and white not as useful. Imagery from 2017 and 2018 were a bit too  
860 snowy around the glacier margin to be useful. The 2018-09-06 Maxar imagery covered  
861 the entire range, with some clouds.

862 In the southern Wind River Range, a new snow dusting was often present, occasionally  
863 making it difficult to outline snowfields and a few glaciers, but mostly snowfields.  
864 Distinguishing seasonal snow from perennial snow was a judgement call. The thin  
865 seasonal If the snow was identified if a slight coloration lightly discolored similar to  
866 underlying rock/soil looking like the color was present or coming from underneath it was  
867 identified as seasonal snow. Also, if many small snow-free patches (a few square meters)  
868 of pockmarked the snow-free surface present or if many rocks protruding protruded  
869 through the snow-, it was considered seasonal. A perennial patch of snow appeared  
870 smooth and white, hiding underlying surface. Thin snow covering glacial cover on glacier  
871 was typically appeared greyish in color, often with banding, much less of a texture and  
872 appeared smoother than the surrounding ice-free landscape.

873 Often, it seems a number of glaciers are thinning in place with a thin layer of debris on  
874 the ice that thickens down valley. The landscape surrounding this relatively smooth  
875 appearance is rumpled like the bedrock terrain further away. Interpretation is difficult.  
876 These are probably shrinking glaciers that are being covered in the debris. The outline is  
877 clear where ice meets bedrock, but in the talus debris area, we digitized along the glacier

878 ice boundary unless some other feature like exposed ice or a crevasse is visible then  
879 included that as part. See INV\_ID E618081N4774579 (Figure A9) for example.

880 At Lower Fremont Glacier, a number of sizable ice patches appear down valley as if a  
881 deposit of buried ice is present. However, there is no obvious connection to the glacier  
882 itself.


883 The GNIS identified a single glacier as the Sacagawea Glacier, and two separate Fremont  
884 Glaciers (Figure A10A5). By 2017 the single glacier had split into four glaciers. We  
885 chose to label the largest glacier and the glacier to the southeast the Sacagawea Glacier.  
886 The other two glaciers were labeled the Fremont Glaciers.



887  
888  
889 **Figure A9.** An example of an unnamed glacier in the Wind River Range, WY, INV\_ID  
890 E618081N4774579 seemingly melting into the talus surrounding the terminus (top of image).  
891 The glacier is flowing from the lower left hand corner to the upper right hand corner. Base  
892 image from the NAIP taken in 2015.

Formatted: Highlight

893  
894



895  
 896 **Figure A10A5.** Image of Fremont Glaciers and Sacagawea Glacier showing the Sacagawea  
 897 outline from the Fountain et al. (2017) database (blue), our updated Fremont Glaciers outlines  
 898 (orange), and updated Sacagawea outlines (red). The base image is from the NAIP, taken in  
 899 2015.

900  
 901  
 902 **Table A14A17.** List of NAIP imagery used for outlining glaciers and perennial snowfields in  
 903 Wyoming. 'Date' is the start and end date for flights covering the glaciated portions of the NAIP  
 904 image. In some cases, flights were completed in a single day. For 2006 the inspection date was  
 905 used, since the start and end dates were not provided.

906

Region/Year/Filename	County	Date (Year-M-D)
Absaroka Range		



2006	ortho_1-2_1n_s_wy029_2006_1.sid	Park	2006-09-02
2015	ortho_1-1_hn_s_wy013_2015_2.sid	Fremont	2015-09-09 to 2015-10-13
	ortho_1-1_hn_s_wy029_2015_2.sid	Park	2015-09-22 to 2015-10-13
<b>Bighorn Mountains, WY</b>			
2015	ortho_1-1_hn_s_wy019_2015_2.sid	Johnson	2015-09-12
<b>Teton Range</b>			
2006	ortho_1-1_1n_s_wy039_2006_1.sid	Teton	2006-09-02
2015	ortho_1-1_hn_s_wy035_2015_2.sid	Sublette	2015-09-09 to 2015-10-13
	ortho_1-1_hn_s_wy039_2015_2.sid	Teton	2015-09-12 to 2015-09-22
2019	ortho_1-1_hn_s_wy039_2019_1.sid	Teton	2019-07-20 to 2015-09-22
<b>Wind River Range</b>			
2006	ortho_1-1_1n_s_wy035_2006_1.sid	Sublette	2006-09-02
2015	ortho_1-1_hn_s_wy013_2015_2.sid	Fremont	2015-09-09 to 2015-10-13
	ortho_1-1_hn_s_wy035_2015_2.sid	Sublette	2015-09-09 to 2015-10-13
2019	ortho_1-1_hn_s_wy013_2019_1.sid	Fremont	2019-07-20 to 2019-08-27
	ortho_1-1_hn_s_wy035_2019_1.sid	Sublette	2019-08-15 to 2019-09-13

907  
908  
909  
910

**Table A15A18.** List of dates of the Maxar imagery used for outlining glaciers and perennial snowfields in Wyoming.

**Region/ Date (Year-M-D)**

Wind River Range  
2018-09-06

911  
912  
913  
914  
915  
916  
917  
918  
919

**6.8 ~~Glaciers that have split into multiple pieces and current errors in glacier label names~~**

~~Table A16 compiles the list of glaciers that have split into multiple pieces since the USGS 1:24000 mapping (Fountain et al., 2017). Table A17 lists current map errors in the labeling of glaciers. The purpose of including this table is to facilitate updating the USGS Geographic Names Information System.~~

920 **Table A16.** List of named glaciers that have split into multiple pieces. Names come from the  
 921 **Geographic Names Information System.** ‘Count’ refers to the number of pieces in the updated  
 922 inventory. ‘Classes’ is the classification of the pieces; glacier, perennial snowfield, buried ice, or  
 923 a combination.

<u>State/Region/Glacier Name</u>	<u>Count</u>	<u>Classes</u>
<b>California</b>		
<b>—Cascade Range</b>		
Dolan Glacier	2	Glaciers and perennial snowfields
Hotlum Glacier	2	Glaciers and perennial snowfields
Whitney Glacier	2	Glaciers and perennial snowfields
Wintun Glacier	2	Glaciers and perennial snowfields
<b>—Sierra Nevada</b>		
Goethe Glacier	2	Glaciers only
Lyonell Glacier	4	Glaciers and perennial snowfields
Norman Clyde Glacier	2	Glaciers only
Powell Glacier	2	Glacier and Buried ice
<b>Colorado</b>		
<b>—Front Range</b>		
Saint Vrain Glaciers	6	Glaciers and perennial snowfields
<b>Montana</b>		
<b>—Beartooth Mountains Absaroka Range</b>		
Castle Rock Glacier	2	Glaciers and perennial snowfields
Granite Glacier	2	Glaciers only
Grasshopper Glacier	4	Glaciers and perennial snowfields
Hopper Glacier	2	Glaciers and perennial snowfields
Snowbank Glacier	2	Glaciers only
Wolf Glacier	2	Glaciers only
<b>—Lewis Range</b>		
Agassiz Glacier	2	Glaciers only
Blackfoot Glacier	2	Glaciers only
Carter Glaciers	2	Glaciers and perennial snowfields
Dixon Glacier	2	Glaciers and perennial snowfields
Harrison Glacier	5	Glaciers and perennial snowfields
Kimble Glacier	2	Glaciers only
Logan Glacier	2	Glaciers only
Shepard Glacier	2	Glaciers only
Siyeh Glacier	2	Glaciers only
Two Ocean Glacier	2	Glaciers only
Whitecrow Glacier	5	Glaciers and perennial snowfields
<b>—Mission Range Swan Range Flathead Range</b>		
Swan Glaciers	2	Glaciers and perennial snowfields
<b>Oregon</b>		
<b>—Cascade Range</b>		
Dend Glacier	2	Glaciers and perennial snowfields
Clark Glacier	2	Perennial snowfields only
Collier Glacier	2	Glaciers only
Diller Glacier	2	Glaciers and perennial snowfields
Glisan Glacier	2	Glaciers and perennial snowfields

Ladd Glacier	4	Glaciers and perennial snowfields
Langille Glacier	5	Glaciers and perennial snowfields
Newton Clark Glacier	2	Glaciers and perennial snowfields
Palmer Glacier	2	Perennial snowfields only
Prouty Glacier	2	Glaciers and perennial snowfields
Renfrew Glacier	2	Glaciers and perennial snowfields
Russell Glacier	2	Glaciers only
Sandy Glacier	4	Glaciers and perennial snowfields
Skinner Glacier	4	Perennial snowfields only
Waldo Glacier	2	Glaciers only
White River Glacier	2	Glaciers and perennial snowfields
Whitewater Glacier	2	Glaciers only
Zigzag Glacier	2	Glaciers and perennial snowfields
<b>Washington</b>		
<b>— Cascade Range Northern</b>		
Borealis Glacier	4	Glaciers only
Buelner Glacier	2	Glaciers only
Butterfly Glacier	4	Glaciers only
Colehuick Glacier	2	Glaciers only
Company Glacier	2	Glaciers only
Cool Glacier	2	Glaciers and perennial snowfields
Dana Glacier	2	Glaciers only
Dark Glacier	2	Glaciers only
Dome Glacier	2	Glaciers only
Douglas Glacier	4	Glaciers and perennial snowfields
Dusty Glacier	2	Glaciers and perennial snowfields
East Nooksack Glacier	5	Glaciers only
Entiat Glacier	4	Glaciers and perennial snowfields
Forbidden Glacier	2	Glaciers only
Fremont Glacier	2	Glaciers only
Goode Glacier	2	Glaciers only
Hadley Glacier	5	Glaciers only
Hanging Glacier	2	Glaciers only
Hinman Glacier	4	Glaciers only
Honeycomb Glacier	2	Glaciers and perennial snowfields
Inspiration Glacier	2	Glaciers and perennial snowfields
Isella Glacier	2	Glaciers and perennial snowfields
Jerry Glacier	2	Glaciers only
Kimtah Glacier	2	Glaciers only
LeConte Glacier	7	Glaciers and perennial snowfields
Lynnall Glacier	2	Perennial snowfields only
Mazama Glacier	2	Glaciers and perennial snowfields
McAllister Glacier	2	Glaciers only
Middle Cascade Glacier	2	Glaciers only
Neve Glacier	2	Glaciers only
No Name Glacier	5	Glaciers and perennial snowfields
Nohokomeen Glacier	2	Glaciers only
North Klawatti Glacier	2	Glaciers and perennial snowfields
Pitz Glacier	2	Glaciers and perennial snowfields

Price Glacier	4	Glaciers only
Parmigan Glacier	2	Glaciers and perennial snowfields
Queest alb Glacier (not official)	2	Glaciers and perennial snowfields
Rainbow Glacier	2	Glaciers and perennial snowfields
Redoubt Glacier	2	Glaciers only
Richardson Glacier	2	Glaciers only
S Glacier	2	Glaciers only
Sandalee Glacier	4	Glaciers only
Scimitar Glacier	2	Glaciers only
Shoher Glacier	4	Glaciers only
Sitkum Glacier	4	Glaciers and perennial snowfields
Snow Creek Glacier	2	Perennial snowfields only
South Cascade Glacier	2	Glaciers only
Spider Glacier	2	Glaciers only
Suiattle Glacier	2	Glaciers only
Sulphide Glacier	2	Glaciers only
Thunder Glacier	2	Glaciers only
Thunder Glacier	2	Glaciers only
White Chuck Glacier	5	Glaciers and perennial snowfields
White Salmon Glacier	2	Glaciers only
Wyeth Glacier	2	Glaciers and perennial snowfields

**— Cascade Range Southern**

Adam Glacier	4	Glaciers and perennial snowfields
Avalanche Glacier	2	Glaciers only
Conrad Glacier	2	Glaciers and perennial snowfields
Cowlitz Glacier	2	Glaciers and perennial snowfields
Crescent Glacier	2	Glaciers and perennial snowfields
Flett Glacier	6	Glaciers and perennial snowfields
Fryingpan Glacier	5	Glaciers and perennial snowfields
Getchen Glacier	2	Glaciers and perennial snowfields
Kautz Glacier	2	Glaciers and perennial snowfields
Klielkitat Glacier	2	Glaciers only
Lava Glacier	2	Glaciers and perennial snowfields
McCall Glacier	6	Glaciers and perennial snowfields
Mondo Glacier	5	Perennial snowfields only
North Mowich Glacier	2	Glaciers and perennial snowfields
Ohanapeosh Glacier	6	Glaciers and perennial snowfields
Paradise Glacier	2	Glaciers and perennial snowfields
Pinnacle Glacier	2	Glaciers and perennial snowfields
Puyallup Glacier	2	Glaciers and perennial snowfields
Pyramid Glacier	4	Glaciers and perennial snowfields
Russell Glacier	2	Glaciers only
Sarvant Glaciers	4	Glaciers and perennial snowfields
South Mowich Glacier	2	Glaciers only
South Tahoma Glacier	2	Glaciers and perennial snowfields
Succors Glacier	2	Glaciers and perennial snowfields
Van Trump Glacier	10	Glaciers and perennial snowfields
White Salmon Glacier	2	Glaciers only
Whitman Glacier	5	Glaciers and perennial snowfields

Wilson Glacier	2	Glaciers and perennial snowfields
<b>Olympic Mountains</b>		
Blue Glacier	2	Glaciers only
Cameron Glacier	4	Glaciers and perennial snowfields
Carrie Glacier	2	Glaciers only
Eel Glacier	2	Glaciers only
White Glacier	2	Glaciers only
<b>Wyoming</b>		
<b>Teton Range</b>		
Middle Teton Glacier	2	Glaciers and perennial snowfields
Triple Glacier	2	Glaciers only
<b>Wind River Range</b>		
Bull Lake Glacier	2	Glaciers and perennial snowfields
Dinwoody Glacier	2	Glaciers only
Dinwoody Glacier	2	Glaciers and perennial snowfields
Grasshopper Glacier	2	Glaciers only
Harrower Glacier	2	Perennial snowfields only
Helen Glacier	2	Glaciers only
Lower Fremont Glacier	4	Glaciers and perennial snowfields
Mammoth Glacier	2	Glaciers and perennial snowfields
Minor Glacier	2	Glaciers and perennial snowfields
Sacagawea Glacier	4	Glaciers and perennial snowfields
Sourdough Glacier	2	Glaciers and perennial snowfields
Stroud Glacier	2	Glaciers and perennial snowfields
Twins Glacier	2	Glaciers and perennial snowfields
Upper Fremont Glacier	2	Glaciers and perennial snowfields

924

925

926 **Table A17.** List of officially named glaciers where we identified an issue with the glacier name  
 927 on the 1:24000 U.S. Geological Survey topographical maps (Fountain et al., 2017). Names come  
 928 from the Geographic Names Information System (U.S. Geological Survey (2022)). The 'Issue'  
 929 column lists the type of issue identified. 'Not labeled' indicates the feature was present but not  
 930 labeled, 'Misidentified' indicates the wrong feature was labeled, and 'Label unclear' indicates it  
 931 is unclear what feature the label is identifying.

<u>State/Region/Glacier Name</u>	<u>Issue</u>
<b>Colorado</b>	
<b>Front Range</b>	
Arikaree Glacier	Not labeled
Navajo Glacier	Not labeled
<b>Oregon</b>	
<b>Cascade Range</b>	
Carver Glacier	Misidentified
Mill Creek Glacier	Not labeled
<b>Washington</b>	
<b>Cascade Range Northern S-Glacier</b>	Label unclear

Formatted: Font color: Auto  
 Formatted: Space After: 8 pt, Line spacing: Multiple 1.08 li

<del>Snow Creek Glacier</del>	<del>Label unclear</del>
<del>South Glacier</del>	<del>Not labeled</del>
<del>Cascade Range Southern</del>	
<del>No Name Glacier</del>	<del>Not labeled</del>
<del>Stevens Glacier</del>	<del>Not labeled</del>
<del>Wyoming</del>	
<del>Wind River Range</del>	
<del>Dinwoody Glaciers</del>	<del>Label unclear</del>
<del>Fremont Glaciers</del>	<del>Label unclear</del>

932

933

934 **9. Author Contributions.**

935

936 Andrew G. Fountain was the principal investigator of the project, he wrote the proposal and  
 937 digitized glacier and snowfield outlines, analyzed the data, and led the writing of this report.  
 938 Bryce Glenn was the GIS expert responsible for the geographic format (e.g. projection,  
 939 attributes, database structure) and quality control. He digitized glacier and snowfield outlines,  
 940 analyzed the data, and helped write the report. Chris McNeil provided some of the imagery.

941

942 **9. Competing Interests.**

943

944 The authors declare that they have no conflict of interest.

945

946 **10. Acknowledgments.**

947

948 Hassan Basagic and Kristina Dick digitized the glaciers of Mt. Shasta and the North Cascade  
 949 National Park, respectively. We greatly appreciate their help and expertise. Nathan Walker of the  
 950 U.S. Forest Service provided a very helpful initial review of this report. We kindly acknowledge  
 951 the funding from the US. Forest Service. ~~Mazar imagery was accessed through the USGS and~~  
 952 ~~NCA NEXTVIEW license. The Mazar imagery has limited availability owing to restrictions~~  
 953 ~~(proprietary interest). Contact cmcneil@usgs.gov for more information.~~ Any use of trade, firm or  
 954 product names is for descriptive purposes only and does not imply endorsement by the U.S.  
 955 Government.

956 **11. References**

957

958

959 Alley, R. B., Cuffey, K. M., & Zoet, L. K. (2019). Glacial erosion: Status and outlook. *Annals of Glaciology*,  
 960 60(80), 1–13. <https://doi.org/10.1017/aog.2019.38>

961 [Andreassen, L. M., Link to external site, this link will open in a new window, Nagy, T., Kjølmoen, B.,](#)  
 962 [Leigh, J. R., & Link to external site, this link will open in a new window. \(2022\). An inventory of Norway's](#)  
 963 [glaciers and ice-marginal lakes from 2018–19 Sentinel-2 data. \*Journal of Glaciology\*, 68\(272\), 1085–1106.](#)  
 964 <https://doi.org/10.1017/jog.2022.20>

Formatted: Font: Bold, Font color: Text 1

Formatted: Indent: Left: 0.25"

Formatted: Normal

Formatted: Font: Not Bold

Formatted: Indent: Left: 0.5", Space After: 0 pt, Line spacing: single

Formatted: Normal

Formatted: Font: +Body (Calibri), 11 pt

965 [Armstrong, R. L. \(1989\). Mass balance history of Blue Glacier, Washington, USA. In J. Oerlemans \(Ed.\),](#)  
966 [Glacier Fluctuations and Climatic Change \(pp. 183–192\). Springer Netherlands.](#)

967 Bard, J. A. (2017a). *High-resolution digital elevation dataset for Glacier Peak and vicinity, Washington,*  
968 *based on lidar surveys of August–November, 2014 and June, 2015* [[Data setdataset](#)]. U.S. Geological  
969 Survey. <https://doi.org/10.5066/F7H41PJG>

970 Bard, J. A. (2017b). *High-resolution digital elevation dataset for Mt Baker and vicinity, Washington,*  
971 *based on lidar surveys of 2015* [[Data setdataset](#)]. U.S. Geological Survey.  
972 <https://doi.org/10.5066/F7WD3XR0>

973 Bard, J. A. (2019). *High-resolution digital elevation dataset for Mount Adams and vicinity, Washington,*  
974 *based on lidar surveys of August–September, 2016* [[Data setdataset](#)]. U.S. Geological Survey.  
975 <https://doi.org/10.5066/P9Z1HF1K>

976 Beason, S. R., Legg, N. T., Kenyon, T. R., Jost, R. P., & Kennard, P. M. (2018). Forecasting and seismic  
977 detection of debris flows in pro-glacial rivers at Mount Rainier National Park, Washington, USA.  
978 *Contained in: Proceedings of the Seventh International Conference on Debris-Flow Hazards Mitigation,*  
979 *Golden, Colorado, USA, June 10–13, 2019, <https://hdl.handle.net/11124/173051>.*  
980 <https://doi.org/10.25676/11124/173232>

981 Benn, D. I., & Evans, D. J. A. (2010). *Glaciers and glaciation* (2nd ed.). Hodder Education.

982 Bolch, T., Menounos, B., & Wheate, R. (2010). Landsat-based inventory of glaciers in western Canada,  
983 1985–2005. *Remote Sensing of Environment, 114*(1), 127–137.  
984 <https://doi.org/10.1016/j.rse.2009.08.015>

985 [Bolch, T., Rohrbach, N., Kutuzov, S., Robson, B. A., & Osmonov, A. \(2019\). Occurrence, evolution and ice](#)  
986 [content of ice-debris complexes in the Ak-Shiirak, Central Tien Shan revealed by geophysical and](#)  
987 [remotely-sensed investigations: Ice-debris complexes in Ak-Shiirak. \*Earth Surface Processes and\*](#)  
988 [Landforms, 44](#)(1), 129–143. <https://doi.org/10.1002/esp.4487>

989 [Bowerman, N. D., & Clark, D. H. \(2011\). Holocene glaciation of the central Sierra Nevada, California.](#)  
990 [\*Quaternary Science Reviews, 30\*](#)(9–10), 1067–1085. <https://doi.org/10.1016/j.quascirev.2010.10.014>

991 Brardinoni, F., Scotti, R., Sailer, R., & Mair, V. (2019). Evaluating sources of uncertainty and variability in  
992 rock glacier inventories. *Earth Surface Processes and Landforms, 44*(12), 2450–2466.  
993 <https://doi.org/10.1002/esp.4674>

994 Cadbury, S. L., Hannah, D. M., Milner, A. M., Pearson, C. P., & Brown, L. E. (2008). Stream temperature  
995 dynamics within a New Zealand glacierized river basin. *River Research and Applications, 24*(1), 68–89.  
996 <https://doi.org/10.1002/rra.1048>

997 Chiarle, M., Iannotti, S., Mortara, G., & Deline, P. (2007). Recent debris flow occurrences associated with  
998 glaciers in the Alps. *Global and Planetary Change, 56*(1–2), 123–136.  
999 <https://doi.org/10.1016/j.gloplacha.2006.07.003>

1000 Cogley, J. G., Hock, R., Rasmussen, L. A., Arendt, A. A., Bauder, A., Braithwaite, R. J., Jansson, P., Kaser,  
1001 G., Möller, M., Nicholson, L. I., & Zemp, M. (2011). *Glossary of Glacier Mass Balance and Related Terms*  
1002 (IHP-VII Technical Documents in Hydrology ~~No.~~-86; IACS Contribution No. 2, p. 114). UNESCO-IHP.

1003 [Davis, P. T. \(1988\). Holocene glacier fluctuations in the American Cordillera. \*Quaternary Science Reviews\*,  
1004 \*7\*\(2\), 129–157.](#)

1005 Denton, G. H. (1975). Conterminous US, Chapter 1. In W. O. Field (Ed.), *Mountain glaciers of the*  
1006 *Northern Hemisphere* (Vol. 1). Corps of Engineers, US Army, Technical Information Analysis Center, Cold  
1007 Regions Research and Engineering Laboratory.

1008 DeVisser, M. H., & Fountain, A. G. (2015). A century of glacier change in the Wind River Range, WY.  
1009 *Geomorphology*, 232, 103–116. <https://doi.org/10.1016/j.geomorph.2014.10.017>

1010 [Dewitz, J. \(2019\). National Land Cover Database \(NLCD\) 2016 Products \(ver. 2.0, July 2020\) \[dataset\].  
1011 U.S. Geological Survey. <https://doi.org/10.5066/P96HHBIE>](#)

1012 Dick, K. (2013). *Glacier Change of the North Cascades, Washington 1900-2009* [M.S.]. Portland State  
1013 University.

1014 Dussaillant, I., Berthier, E., Brun, F., Masiokas, M., Hugonnet, R., Favier, V., Rabatel, A., Pitte, P., & Ruiz,  
1015 L. (2019). Two decades of glacier mass loss along the Andes. *Nature Geoscience*, 12(10), 802–808.  
1016 <https://doi.org/10.1038/s41561-019-0432-5>

1017 Earl, L., & Gardner, A. (2016). A satellite-derived glacier inventory for North Asia. *Annals of Glaciology*,  
1018 57(71), 50–60. <https://doi.org/10.3189/2016AoG71A008>

1019 [Evans, I. S. \(2006\). Local aspect asymmetry of mountain glaciation: A global survey of consistency of  
1020 favoured directions for glacier numbers and altitudes. \*Geomorphology\*, 73\(1–2\), 166–184.  
1021 <https://doi.org/10.1016/j.geomorph.2005.07.009>](#)

1022 Fagre, D. B., McKeon, L. A., Dick, K. A., & Fountain, A. G. (2017). *Glacier margin time series (1966, 1998,*  
1023 *2005, 2015) of the named glaciers of Glacier National Park, MT, USA* [~~Data set~~[dataset](#)]. U.S. Geological  
1024 Survey. <https://doi.org/10.5066/f7p26wb1>

1025 Fellman, J. B., Nagorski, S., Pyare, S., Vermilyea, A. W., Scott, D., & Hood, E. (2014). Stream temperature  
1026 response to variable glacier coverage in coastal watersheds of Southeast Alaska. *Hydrological Processes*,  
1027 28(4), 2062–2073. <https://doi.org/10.1002/hyp.9742>

1028 [Fischer, A., Seiser, B., Stocker Waldhuber, M., Mitterer, C., & Abermann, J. \(2015\). Tracing glacier  
1029 changes in Austria from the Little Ice Age to the present using a lidar-based high-resolution glacier  
1030 inventory in Austria. \*The Cryosphere\*, 9\(2\), 753–766. <https://doi.org/10.5194/tc-9-753-2015>](#)

1031 Fountain, A. G., & Glenn, B. (2022). *Data From: Inventory of Glaciers and Perennial Snowfields of the*  
1032 *Coterminous USA" (2022)* (Data Series ~~No.~~-3; Geology Faculty Datasets). Portland State University.  
1033 <https://doi.org/10.15760/geology-data.03>



1034 Fountain, A. G., Glenn, B., & Basagic, H. J. (2017). The Geography of Glaciers and Perennial Snowfields in  
1035 the American West. *Arctic, Antarctic, and Alpine Research*, 49(3), 391–410.  
1036 <https://doi.org/10.1657/AAAR0017-003>

1037 Fountain, A. G., Hoffman, M. J., Jackson, K., Basagic, H. J., Nylén, T. H., & Percy, D. (2007). *Digital outlines  
1038 and the topography of the Glaciers of the American West* (Open File Report No. 2006–1340; p. 23). US  
1039 Geological Survey. <https://doi.org/10.3133/ofr20061340>

1040 Fountain, A. G., & Tangborn, W. V. (1985). The effect of glaciers on streamflow variations. *Water  
1041 Resources Research*, 21(4), 579–586.

1042 Garwood, J. M., Fountain, A. G., Lindke, K. T., van Hattem, M. G., & Basagic, H. J. (2020). 20th Century  
1043 Retreat and Recent Drought Accelerated Extinction of Mountain Glaciers and Perennial Snowfields in the  
1044 Trinity Alps, California. *Northwest Science*, 94(1), 44. <https://doi.org/10.3955/046.094.0104>

1045 Gesch, D., Oimoen, M., Greenlee, S., Nelson, C., Steuck, M., & Tyler, D. (2002). The national elevation  
1046 dataset. *Photogrammetric Engineering and Remote Sensing*, 68(1), 5–32.

1047 Heard, J. (2000). *Late Pleistocene and Holocene Aged Glacial and Climatic Reconstructions in the Goat  
1048 Rocks Wilderness, Washington, United States*. <https://doi.org/10.15760/etd.557>

1049 Hock, R., de Woul, M., Radić, V., & Dyurgerov, M. (2009). Mountain glaciers and ice caps around  
1050 Antarctica make a large sea-level rise contribution: MOUNTAIN GLACIERS AND ICE CAPS. *Geophysical  
1051 Research Letters*, 36(7), n/a-n/a. <https://doi.org/10.1029/2008GL037020>

1052 Hoffman, M. J., Fountain, A. G., & Achuff, J. M. (2007). 20th-century variations in area of cirque glaciers  
1053 and glacierets, Rocky Mountain National Park, Rocky Mountains, Colorado, USA. *Annals of Glaciology*,  
1054 46(1), 349–354. <https://doi.org/10.3189/172756407782871233>

1055 [Huss, M., & Hock, R. \(2018\). Global-scale hydrological response to future glacier mass loss. \*Nature  
1056 Climate Change\*, 8\(2\), 135–140. <https://doi.org/10.1038/s41558-017-0049-x>](https://doi.org/10.1038/s41558-017-0049-x)

1057 ~~Jin, S., Homer, C. G., Dewitz, J. A., Yang, L., Jin, S., Danielson, P., Dewitz, J., Li, C., Zhu, Z., Xian, G.,  
1058 Coulston, J., Herold, N., & Howard, D., Wickham, J., & Megown, K. (2015). Completion of the  
1059 2011. (2019). Overall Methodology Design for the United States National Land Cover Database for the  
1060 conterminous United States Representing a decade of land cover change information.  
1061 *Photogramm. Eng.-2016 Products. Remote Sens.*, 81(5), 345–354. *Sensing*, 11(24), 2971.  
1062 <https://doi.org/10.3390/rs11242971>~~

1063 King, C. (1871). On the discovery of actual glaciers on the Mountains of the Pacific Slope. *American  
1064 Journal of Science and Arts*, 1(3), 157–167.

1065 Krimmel, R. M. (2002). Glaciers of the conterminous United States. In R. S. Williams, Jr. & J. Ferrigno  
1066 (Eds.), *Satellite image atlas of glaciers of the world: North America* (pp. J329–J381). US Geological Survey  
1067 Professional Paper.

1068 [Leigh, J. R., Stokes, C. R., Carr, R. J., Evans, I. S., Andreassen, L. M., & Evans, D. J. A. \(2019\). Identifying](#)  
1069 [and mapping very small \(<0.5 km<sup>2</sup>\) mountain glaciers on coarse to high-resolution imagery. \*Journal of\*](#)  
1070 [Glaciology, 65\(254\), 873–888. <https://doi.org/10.1017/jog.2019.50>](#)

1071 [Linsbauer, A., Huss, M., Hodel, E., Bauder, A., Fischer, M., Weidmann, Y., Bärtschi, H., & Schmassmann,](#)  
1072 [E. \(2021\). The New Swiss Glacier Inventory SGI2016: From a Topographical to a Glaciological Dataset.](#)  
1073 [Frontiers in Earth Science, 9. <https://www.frontiersin.org/articles/10.3389/feart.2021.704189>](#)

1074 [Linsbauer, A., Paul, F., & Haerberli, W. \(2012\). Modeling glacier thickness distribution and bed](#)  
1075 [topography over entire mountain ranges with GlabTop: Application of a fast and robust approach.](#)  
1076 [Journal of Geophysical Research, 117\(F3\). <https://doi.org/10.1029/2011JF002313>](#)

1077 [Lu, Y., Zhang, Z., Kong, Y., & Hu, K. \(2022\). Integration of optical, SAR and DEM data for automated](#)  
1078 [detection of debris-covered glaciers over the western Nyainqentanglha using a random forest classifier.](#)  
1079 [Cold Regions Science and Technology, 193, 103421. <https://doi.org/10.1016/j.coldregions.2021.103421>](#)

1080 Meier, M. F. (1961). Distribution and variations of glaciers in the United States exclusive of Alaska.  
1081 *International Association of Scientific Hydrology*, 54, 420–429.

1082 Meier, M. F. (1984). Contribution of small glaciers to global sea level. *Science*, 226(4681), 1418–1421.

1083 [Mishra, A., Nainwal, H. C., Bolch, T., Shah, S. S., & Shankar, R. \(2023\). Glacier inventory and glacier](#)  
1084 [changes \(1994–2020\) in the Upper Alaknanda Basin, Central Himalaya. \*Journal of Glaciology\*, 69\(275\),](#)  
1085 [591–606. <https://doi.org/10.1017/jog.2022.87>](#)

1086 Moore, R. D., Fleming, S. W., Menounos, B., Wheate, R., Fountain, A., Stahl, K., Holm, K., & Jakob, M.  
1087 (2009). Glacier change in western North America: Influences on hydrology, geomorphic hazards and  
1088 water quality. *Hydrological Processes*, 23(1), 42–61. <https://doi.org/10.1002/hyp.7162>

1089 [NAIP. \(2017\). National agricultural imagery program \(NAIP\) Information Sheet \(p. 2\) \[Information Sheet\].](#)  
1090 [United States Department of Agriculture.](#)

1091 O'Connor, J. E., Hardison, J. H., & Costa, J. E. (2001). *Debris flows from failures of Neoglacial-age moraine*  
1092 *dams in the Three Sisters and Mount Jefferson wilderness areas, Oregon: A study of the recent debris*  
1093 *flows from moraine-dammed lake releases at central Oregon Cascade Range volcanoes - Mt. Jefferson,*  
1094 *Three Fingered Jack and the Three Sisters / Broken Top.* US Geological Survey.

1095 [Osborn, G., Menounos, B., Ryane, C., Riedel, J. L., Clague, J. J., Koch, J., Clark, D., Scott, K., & Davis, P. T.](#)  
1096 [\(2012\). Latest Pleistocene and Holocene glacier fluctuations on Mount Baker, Washington. \*Quaternary\*](#)  
1097 [Science Reviews, 49, 33–51. <https://doi.org/10.1016/j.quascirev.2012.06.004>](#)

1098 Parkes, D., & Marzeion, B. (2018). Twentieth-century contribution to sea-level rise from uncharted  
1099 glaciers. *Nature*, 563(7732), 551–554. <https://doi.org/10.1038/s41586-018-0687-9>

1100 [Paul, F., Barrand, N. E., Baumann, S., Berthier, E., Bolch, T., Casey, K., Frey, H., Joshi, S. P., Kononov, V.,](#)  
1101 [Le Bris, R., Mölg, N., Nosenko, G., Nuth, C., Pope, A., Racoviteanu, A., Rastner, P., Raup, B., Scharer, K.,](#)  
1102 [Steffen, S., & Winsvold, S. \(2013\). On the accuracy of glacier outlines derived from remote-sensing data.](#)  
1103 [Annals of Glaciology, 54\(63\), 171–182. <https://doi.org/10.3189/2013AoG63A296>](#)

1104 [Paul, F., Kääh, A., & Haeberli, W. \(2007\). Recent glacier changes in the Alps observed by satellite: Consequences for future monitoring strategies. \*Global and Planetary Change\*, 56\(1–2\), 111–122. <https://doi.org/10.1016/j.gloplacha.2006.07.007>](#)

1105

1106

1107 [Paul, F., Rastner, P., Azzoni, R. S., Diolaiuti, G., Fugazza, D., Le Bris, R., Nemec, J., Rabatel, A., Ramusovic, M., Schwaizer, G., & Smiraglia, C. \(2020\). Glacier shrinkage in the Alps continues unabated as revealed by a new glacier inventory from Sentinel-2. \*Earth System Science Data\*, 12\(3\), 1805–1821. <https://doi.org/10.5194/essd-12-1805-2020>](#)

1108

1109

1110

1111 [Pfeffer, W. T., Arendt, A. A., Bliss, A., Bolch, T., Cogley, J. G., Gardner, A. S., Hagen, J.-O., Hock, R., Kaser, G., & Kienholz, C. \(2014\). The Randolph Glacier Inventory: A globally complete inventory of glaciers. \*Journal of Glaciology\*, 60\(221\), 537–552. <https://doi.org/10.3189/2014JoG13J176>](#)

1112

1113

1114 [Post, A., Richardson, D., Tangborn, W. V., & Rossetol, F. \(1971\). \*Inventory of glaciers in the North Cascades, Washington. \(USGS Professional Paper 705-A\)\*. \(p. 26\). US Geological Survey, 705-A.](#)

1115

1116 [Pritchard, H. D. \(2019\). Asia's shrinking glaciers protect large populations from drought stress. \*Nature\*, 569\(7758\), Article 7758. <https://doi.org/10.1038/s41586-019-1240-1>](#)

1117

1118 [Rabatel, A., Sirguey, P., Drolon, V., Maisongrande, P., Arnaud, Y., Berthier, E., Davaze, L., Dedieu, J.-P., & Dumont, M. \(2017\). Annual and Seasonal Glacier-Wide Surface Mass Balance Quantified from Changes in Glacier Surface State: A Review on Existing Methods Using Optical Satellite Imagery. \*Remote Sensing\*, 9\(5\), 507. <https://doi.org/10.3390/rs9050507>](#)

1119

1120

1121

1122 [Rasmussen, L. A. \(2009\). South Cascade Glacier mass balance, 1935–2006. \*Annals of Glaciology\*, 50\(50\), 215–220.](#)

1123

1124 [RGIK. \(2022\). \*Towards standard guidelines for inventorying rock glaciers: Baseline concepts \(version 4.2.2\)\* \(p. 13\). . IPA Action Group Rock glacier inventories and kinematics.](#)

1125

1126 [Robinson, J. E. \(2014\). \*Digital topographic data based on lidar survey of Mount Shasta Volcano, California, July-September 2010\* \(Report No. 852; Data Series\). USGS Publications Warehouse. <https://doi.org/10.3133/ds852>](#)

1127

1128

1129 [Robson, B. A., Bolch, T., MacDonell, S., Hölbling, D., Rastner, P., & Schaffer, N. \(2020\). Automated detection of rock glaciers using deep learning and object-based image analysis. \*Remote Sensing of Environment\*, 250, 112033. <https://doi.org/10.1016/j.rse.2020.112033>](#)

1130

1131

1132 [Russell, I. C. \(1898\). The glaciers of North America. \*The Geographical Journal\*, 12\(6\), 553–564.](#)

1133

1134 [Schiefer, E., Menounos, B., & Wheate, R. \(2007\). Recent volume loss of British Columbian glaciers, Canada: Volume loss of BC glaciers. \*Geophysical Research Letters\*, 34\(16\), L16503. <https://doi.org/10.1029/2007GL030780>](#)

1135

1136 [Selkowitz, D. J., & Forster, R. R. \(2016\). Automated mapping of persistent ice and snow cover across the western U.S. with Landsat. \*ISPRS Journal of Photogrammetry and Remote Sensing\*, 117, 126–140. <https://doi.org/10.1016/j.isprsjprs.2016.04.001>](#)

1137

1138

1139 [Sitts, D. J., Fountain, A. G., & Hoffman, M. J. \(2010\). Twentieth Century Glacier Change on Mount Adams,](#)  
1140 [Washington, USA. \*Northwest Science\*, 84\(4\), 378–385. <https://doi.org/10.3955/046.084.0407>](#)

1141 Smiraglia, C., Azzoni, R. S., D’Agata, C., Maragno, D., Fugazza, D., & Diolaiuti, G. A. (2015). The New  
1142 Italian Glacier Inventory: A didactic tool for a better knowledge of the natural Alpine environment. *J-*  
1143 *Reading-Journal of Research and Didactics in Geography*, 4(1). [http://www.j-](http://www.j-reading.org/index.php/geography/article/view/92)  
1144 [reading.org/index.php/geography/article/view/92](http://www.j-reading.org/index.php/geography/article/view/92)

1145 Spicer, R. (1986). *Glaciers in the Olympic Mountains, Washington—Present distribution and recent*  
1146 *variations*. [M.S.]. University of Washington.

1147 Sun, M., Liu, S., Yao, X., Guo, W., & Xu, J. (2018). Glacier changes in the Qilian Mountains in the past half-  
1148 century: Based on the revised First and Second Chinese Glacier Inventory. *Journal of Geographical*  
1149 *Sciences*, 28(2), 206–220. <https://doi.org/10.1007/s11442-018-1468-y>

1150 Trcka, A. (2020). *Inventory of Rock Glaciers in the American West and Their Topography and Climate*.  
1151 <https://doi.org/10.15760/etd.7509>

1152 ~~U.S. Geological Survey, 2022. [https://www.usgs.gov/tools/geographic-names-information-](https://www.usgs.gov/tools/geographic-names-information-system-gnis)~~  
1153 ~~[system-gnis](https://www.usgs.gov/tools/geographic-names-information-system-gnis). Accessed December 5, 2022~~

1154 Usery, E. L., Varanka, D., & Finn, M. P. (2009). A 125 year history of topographic mapping and GIS in the  
1155 U.S. Geological Survey 1884-2009, part 1: 1884-1980. *ArcNews*, 31(3), 1–1. USGS Publications  
1156 Warehouse.

1157 [Wickham, J., Stehman, S. V., Sorenson, D. G., Gass, L., & Dewitz, J. A. \(2021\). Thematic accuracy](#)  
1158 [assessment of the NLCD 2016 land cover for the conterminous United States. \*Remote Sensing of\*](#)  
1159 [Environment](#), 257, 112357. <https://doi.org/10.1016/j.rse.2021.112357>

1160 [Yao, T., Pu, J., Lu, A., Wang, Y., & Yu, W. \(2007\). Recent glacial retreat and its impact on hydrological](#)  
1161 [processes on the Tibetan Plateau, China, and surrounding regions. \*Arctic, Antarctic, and Alpine Research\*,](#)  
1162 [39\(4\), 642–650.](#)

1163 [Zalazar, L., Ferri, L., Castro, M., Gargantini, H., Gimenez, M., Pitte, P., Ruiz, L., Masiokas, M., Costa, G., &](#)  
1164 [Villalba, R. \(2020\). Spatial distribution and characteristics of Andean ice masses in Argentina: Results](#)  
1165 [from the first National Glacier Inventory. \*Journal of Glaciology\*, 66\(260\), 938–949.](#)  
1166 <https://doi.org/10.1017/jog.2020.55>

1167 Zemp, M., Huss, M., Thibert, E., Eckert, N., McNabb, R., Huber, J., Barandun, M., Machguth, H.,  
1168 Nussbaumer, S. U., Gärtner-Roer, I., Thomson, L., Paul, F., Maussion, F., Kutuzov, S., & Cogley, J. G.  
1169 (2019). Global glacier mass changes and their contributions to sea-level rise from 1961 to 2016. *Nature*,  
1170 568(7752), 382–386. <https://doi.org/10.1038/s41586-019-1071-0>

1171