

1 **Inventory of glaciers and perennial snowfields of the conterminous USA**

2
3 Andrew G. Fountain¹, Bryce Glenn¹, Christopher Mcneil²

4 ¹Department of Geology, Portland State University, 1721 SW Broadway, Portland, OR. 97212
5 USA, ²U.S. Geological Survey Alaska Science Center, Anchorage, AK, USA

6 *Correspondence to:* Andrew G. Fountain (andrew@pdx.edu)

7 **Abstract**

8 This report summarizes an updated inventory of glaciers and perennial snowfields of the
9 conterminous United States. The inventory is based on interpretation of mostly aerial imagery
10 provided by the National Agricultural Imagery Program, U.S. Department of Agriculture with
11 some satellite imagery in places where aerial imagery was not suitable. The inventory includes
12 all perennial snow and ice features $\geq 0.01 \text{ km}^2$. Due to aerial survey schedules and seasonal snow
13 cover, imagery acquired over a number of years were required. The earliest date is 2013 and the
14 latest is 2020, but more than 73% of the outlines were acquired from 2015 imagery. The
15 inventory is compiled as shapefiles within a geographic information system that includes feature
16 classification, area, and location. The inventory identified 1331 ($366.52 \pm 14.34 \text{ km}^2$) glaciers,
17 1776 ($31.01 \pm 9.30 \text{ km}^2$) perennial snowfields, and 35 ($3.57 \text{ km}^2 \pm$ no uncertainty) buried-ice
18 features. The data including both the shapefiles and tabulated results are publicly available at
19 <https://doi.org/10.15760/geology-data.03> (Fountain and Glenn, 2022).

20 **1. Introduction**

21 Glaciers are an important feature of the landscape for several reasons. Geologically, they modify
22 the landscape through erosion and deposition (Alley et al., 2019; Benn and Evans, 2010).
23 Although these processes are typically slow, sudden episodes can occur such as moraine failure
24 due to fluvial erosion resulting in catastrophic debris flows (Beason et al., 2018; Chiarle et al.,
25 2007; O'Connor et al., 2001). Hydrologically, glaciers can be viewed as frozen reservoirs of
26 water that naturally regulate streamflow on seasonal to decadal time scales (Dussailant et al.,
27 2019; Fountain and Tangborn, 1985; Moore et al., 2009). Glacier runoff increases during warm
28 periods and diminishes during cool, wet periods. Thus, glacier populated watersheds have less
29 seasonally variable runoff than ice-free watersheds. Also, glacier runoff cools stream
30 temperatures in the driest and hottest part of the summer after seasonal snowpacks have vanished
31 (Cadbury et al., 2008; Fellman et al., 2014). As glaciers shrink, they have less ability to buffer
32 seasonal runoff variations and watersheds become more susceptible to drought (Huss and Hock,
33 2018; Pritchard, 2019). Globally, the loss of perennial ice from the landscape is a major
34 contributor to sea level rise (Meier, 1984; Parkes and Marzeion, 2018; Zemp et al., 2019).

35 Glacier inventories have been valuable for assessing glacier contribution to sea level change
36 (Hock et al., 2009; Pfeffer et al., 2014), and for assessing regional hydrology (Moore et al., 2009;
37 Yao et al., 2007). They also provide a baseline for quantifying future glacier changes. Updated
38 glacier inventories have been compiled for many regions of the world (Andreassen et al., 2022;
39 Bolch et al., 2010; Smiraglia et al., 2015; Sun et al., 2018). An exception has been western

40 United States (US), defined here as those conterminous states west of the 100th meridian. The
41 most recent inventory is (Fountain et al., 2007, 2017) based on U.S. Geological Survey maps
42 compiled over a 40-year period from the late 1940s to the 1980s. Despite a vigorous history of
43 glacier studies (e.g. Armstrong, 1989; Rasmussen, 2009)), glacial geology (e.g. Bowerman and
44 Clark, 2011; Davis, 1988; Osborn et al., 2012)), and regional inventories (e.g. DeVisser and
45 Fountain, 2015; Fagre et al., 2017; Post et al., 1971) the glacier cover for the entire western US
46 has not been reevaluated.

47 The earliest scientific identification of glacier-populated regions in the western US date to King
48 (1871) and, more comprehensively, to Russell (1898). The first summary of glacier-covered area
49 for each state was Meier (1961). However, the data sources and methods used to compile the
50 inventories are unknown. Denton (1975) summarized all known glacier studies in the western US
51 but did not tabulate glacier area. Krimmel (2002) updated Meier's study and provided total
52 glacier area for the various mountain ranges by summarizing a variety previous studies published
53 over a 10+ year time span. It is not clear whether the inventory is complete and no data on
54 individual glaciers are provided. Fountain et al. (2007, 2017) compiled the first comprehensive
55 inventory of glaciers in the western US. The data were derived from historical U.S. Geological
56 Survey (USGS) 1:24,000 scale maps compiled over a 40-year period from the 1940s to the 1980s
57 (Gesch et al., 2002; Usery et al., 2009). Because the USGS mapping was based on one-time
58 aerial imagery, the misinterpretation of seasonal snow as perennial was extensive in some
59 regions. The most current study, Selkowitz & Forster (2016), used Landsat satellite imagery
60 compiled over a four-year period, 2010-2014, and an automated detection scheme to define
61 perennial snow and ice. However, these early automated schemes are known to misclassify
62 debris-covered ice as ice-free landscape underestimating glacier area (Earl and Gardner, 2016;
63 Paul et al., 2007; Rabatel et al., 2017). Recent advances in automated detection have reduced
64 these errors suggesting a more promising future (Lu et al., 2022; Robson et al., 2020).

65 This paper presents the results of an updated and comprehensive inventory of glaciers and
66 perennial snowfields of the western US for the purpose of defining their current extent and to
67 provide of baseline for estimating future changes. We summarize our methods, uncertainties,
68 tabulated results, and data availability. The data referenced throughout the manuscript are publicly
69 available at <https://doi.org/10.15760/geology-data.03>.

70 **2. Methods**

71

72 **2.1 Data Sources, Classification, Digitizing, and Completeness**

73 The glaciers and perennial snowfields were initially located using a geographic information
74 system (GIS) database from Fountain et al. (2007, 2017). New outlines were manually digitized
75 from three sources of optical imagery. Most of the outlines were digitized from color digital
76 orthographic aerial photographs available from the National Agricultural Imagery Program
77 (NAIP), U.S. Department of Agriculture, Farm Service Agency program (NAIP, 2017),
78 (https://datagateway.nrcs.usda.gov/GDGHome_DirectDownload.aspx). Since 2009, the imagery
79 is collected on cycles of two to three years. The aerial imagery was orthorectified using the
80 inertial navigation system - GPS unit in the aircraft. Photo identifiable GPS-survey ground
81 control points were then used to adjust the photo strip. Orthorectified strips, which had $\geq 30\%$

82 overlap with adjacent strips, were overlaid with each other and with ground control points to
83 check accuracy. The image strips are then mosaicked together. The spatial resolution was ≤ 0.6
84 m with a horizontal accuracy of ≤ 6 m of photo-identifiable ground control points (NAIP, 2017).
85 The NAIP imagery fit the historic USGS glacier outlines remarkably well. In a few cases, NAIP
86 imagery was not suitable due to seasonal snow, deep shadows, or image warping caused by
87 orthophoto rectification, therefore other sources were used including Maxar satellite imagery
88 (Maxar Technologies, Inc) with a spatial resolution of 0.5 – 1 m. For 21 perennial snowfields
89 and three glaciers we relied on the most recent snow-free imagery available in Google Earth
90 (Google, Inc), resolution ~ 1 m, because no other imagery was suitable. The outlines were
91 digitized in Google Earth and exported to ArcMap (Esri, Inc).

92 We manually identified all glaciers, ice patches, and perennial snowfields. Glaciers are defined
93 as perennial snow and ice that moves (Cogley et al., 2011). A feature was considered perennial if
94 it was present on the original 1:24,000 USGS topographic maps and present on all Google Earth
95 imagery. Movement was identified by the presence of crevasses. Perennial snowfields and ice
96 patches do not exhibit movement, as indicated by a lack of crevasses observed in the imagery.
97 We do not distinguish between snowfields and ice patches and refer to both as perennial
98 snowfields.

99 Contiguous glacier cover, most commonly on volcanoes, was separated into individual glaciers if
100 they had unique names as indicated on the USGS maps. The orientation of crevasse patterns was
101 used to define flow divides. In the absence of these patterns, shaded relief maps from digital
102 elevation models were used. These models were derived from aerial lidar data, flown under
103 contract to the USGS (Bard, 2017a, b, 2019; Robinson, 2014) or the Oregon Department of
104 Geology and Mineral Industries (DOGAMI, 2011).

105 We encountered a number of challenges to our classification and delineation of the glaciers and
106 perennial snowfields. Although crevasses were used to define movement, in a few cases it
107 appeared that they penetrated through the feature to the bedrock underneath suggesting a
108 mechanical break up. In these cases, the feature was classified as a snowfield. Debris-cover
109 made defining the glacier outline for some glaciers on the volcanoes of the Cascade Range. We
110 relied on local knowledge to help define some boundaries and independent digitization efforts by
111 the authors and others to provide an uncertainty as explained below. In the high alpine regions of
112 California, Colorado, and Wyoming, the terminus of some glaciers was hard to define. Rather
113 than abruptly terminating, the ice seems to thin and smoothly transitions into the surrounding
114 rock talus (Figure 1). It was unclear whether a thin debris layer blanketed the ice or cobbles and
115 boulders protruded through the thin ice. The boundary was mapped along the edge of identifiable
116 ice.

117



118


119 **Figure 1.** An example of a glacier seemingly melting into the talus surrounding the terminus
120 (upper right). The glacier is flowing from the lower left-hand corner to the upper right-hand
121 corner. The glacier is located in the Wind River Range, WY, INV_ID E618081N4774579 and
122 base image is from the National Agricultural Image Program taken in 2015.

123

124 In a few situations, we found it difficult to distinguish glaciers from rock glaciers (Brardinoni et
125 al., 2019). A rock glacier is a mass of rock debris in a matrix of ice that flows (Cogley et al.,
126 2011). They can be difficult to distinguish from a debris-covered glacier, one that has extensive
127 rock debris over the ablation zone, that lower part of a glacier with exposed ice in late summer.
128 We adopted the following topographic classification. If the slope of the apparent ice
129 patch/snowfield was similar to the slope of the rock glacier then we considered it part of the rock
130 glacier (Figure 2a). On the other hand, if a topographic depression separates the apparent
131 glacier/snowfield from the start of a rock glacier, then it was considered independent feature
132 (Figure 2b). This latter case is similar to the “glacier forefield-connected” rock glacier as
133 described by (RGIK, 2022).

134

135



136

137 **Figure 2.** Examples of glacier versus rock glacier identification. (a) An example of a snowfield
138 that is considered part of the rock glacier. Location, Colorado Front Range, 40.827477° N, -
139 106.657400° E. Image is from © Google Earth, 9/2014; (b) Tyndall Glacier in the Colorado
140 Front Range, 40.305291° N, -105.689602° E, with a rock glacier slightly down valley. Image is
141 from © Google Earth 9/2016.

142


143 In a number of situations, we observed buried ice adjacent to a glacier (Figure 3). Here we use
144 the term ‘buried ice’ to mean dead ice formerly part of a flowing glacier, and not the permafrost
145 context of ice embedded within or on top of perennially frozen ground. The rocky surface texture
146 of the buried ice was hummocky and very different from surrounding bedrock and adjacent ice,
147 and not a moraine. Occasionally a crack in the surface revealed subsurface ice. The feature
148 appeared to be non-moving (dead) ice that is covered by debris similar to some of the ice-debris
149 complexes described by Bolch et al. (2019). We decided to include these features as a separate
150 classification, ‘buried ice’, because their size was large relative to the glacier, they were probably
151 once part of the glacier, and may be important local sources of meltwater for streamflow.

152

153

154

155



156
 157 **Figure 3.** Lost Creek Glacier, South Sister, Oregon. Note buried ice and lack of crevasses to the
 158 left of the grey-blue ice, suggesting ice that is no longer moving and therefore not part of the
 159 dynamic glacier. The white box surrounds an area that has collapsed due to subsurface melt. The
 160 inset enlargement shows a cliff edge of exposed dirty ice (white arrow in upper left) indicated by
 161 a darker color suggesting wet sediment and a finer texture than the surface debris. The black
 162 arrow shows the width of the cleaner ice for scale. Image is from © Google Earth, 8/9/2021.
 163

164 The glaciers and perennial snowfields outlines were digitized using ArcMap (Esri, Inc), a
 165 geographic information system, at scales varying from 1:300 to 1:2000 depending on image
 166 quality and complexity. We used the native projection of the image, North American Datum of
 167 1983 (NAD83) for NAIP, and World Geodetic System 1984 (WGS84) for Maxar and Google
 168 Earth. When Maxar or Google Earth imagery were used, final outlines were projected into the
 169 NAD83 coordinate system. Google Earth was often used as an additional aid in interpretation
 170 because of its tilt and rotation features yielded oblique perspectives. Retaining only those
 171 outlines $\geq 0.01 \text{ km}^2$, each was checked independently by the two senior authors of this report and
 172 in some cases by a third collaborator in order to reduce bias (Leigh et al., 2019). If an outline was
 173 revised, then it was returned to its original author for review and correction, and the process
 174 iterated until all parties agreed.
 175

176 Our initial inventory was then compared sequentially to two other independent inventories to test
 177 for errors of omission or commission. The first comparison was to the Selkowitz and Forster
 178 (2016) inventory (SFI). However, to compare the inventories we had to first reconcile the

179 differences in methods. Buried-ice features were eliminated from our inventory because the SFI
180 did not map buried ice. The SFI was filtered to only include features $\geq 0.01 \text{ km}^2$ to match our
181 minimum area threshold; a small number of features located in Canada were removed; and a few
182 mis-classifications of ponds, lakes and dry lakebeds as glaciers were removed. Notably, the SFI
183 did not split contiguous ice masses, such as glacier-covered volcanoes, into individual glaciers,
184 consequently we do not expect the number of features in the SFI and our inventory to match.
185 Once the two inventories were reconciled, those glaciers and perennial snowfields unique to one
186 inventory were examined for inclusion in a revised inventory. Features selected from the SFI
187 were digitized using the same imagery we used for our inventory.

188
189 The revised inventory was then compared to the 2016 National Land Cover Database
190 (NLCD(Dewitz, 2019), which did not map glaciers and perennial snowfields per se, but mapped
191 the distribution of perennial snow and ice (Jin et al., 2019; Wickham et al., 2021). However, the
192 NLCD used a small number of recent images to assess a ‘perennial’ presence and therefore
193 significant errors of commission are expected. Also, the landscape class of snow and ice received
194 less attention than other classes (e.g. agriculture) such that the timing of imagery acquisition may
195 be earlier in the summer than optimal and misclassification of clouds as snow and ice may be
196 present (personal communication C. Homer and J. Dewitz, USGS, email December 2015). The
197 NLCD inventory was compared to the revised inventory and, as before, the features unique to
198 one inventory were examined for inclusion. Those features selected from the NLCD for inclusion
199 were digitized using the same imagery we used for our inventory.


200 201 2.2 Uncertainty

202
203 Three main sources of uncertainty in the glacier outlines, are georeferencing, digitization, and
204 interpretation (DeVisser and Fountain, 2015; Sitts et al., 2010). We found georeferencing error to
205 very small. In any case, the precise location of the outline does not affect its area. Also, the
206 digitized points are highly correlated such that no deviations from the true outline are caused by
207 georeferencing. Digitizing error is relatively small, 1%, with good imagery and crisp contrast
208 between the glacier and ice-free surroundings (DeVisser and Fountain, 2015; Hoffman et al.,
209 2007). The largest uncertainty is interpretation error caused by poor imagery, shadow, debris
210 cover, and seasonal snow patches. This uncertainty was calculated in different ways according to
211 the situation. If the outline was digitized a second (or third) time due different interpretations by
212 the authors or collaborators the uncertainty is one-half the absolute difference of the between the
213 largest and smallest digitized areas (the range) divided by the final area and expressed as a
214 percentage. For the relatively few glaciers where a small section of perimeter was masked by
215 deep shadow, seasonal snow patches, rock debris, or poor imagery, a higher uncertainty was
216 assigned by visually estimating the area in question and dividing by the total possible area. In a
217 few cases the location of a flow divide between glaciers wasn’t clear a 5% error is assigned. This
218 was calculated from the area difference in several test cases where multiple possible flow divides
219 were digitized. For perennial snowfields, the smaller patch of perennial snow is often covered by
220 seasonal snow, which varies greatly from year to year. We measured the area of a number of
221 snowfields over time using late summer historic imagery in Google Earth. Results showed that
222 the variations in snowfield area could be as much as 30%. We assigned this somewhat arbitrary
223 uncertainty in order to note snowfield presence and location, but preclude them from area change
224 calculations because area differences are typically smaller than the assigned uncertainty.

225
226
227
228
229
230
231
232
233
234
235
236
237

3. Results

Our initial inventory identified 2267 glaciers and perennial snowfields totaling 391.95 km². About 70% (1576) overlapped the features in the SFI. After examining all features unique to each inventory, we revised our inventory to include 2373 (394.99 km²) glaciers and perennial snowfields. Comparing the revised inventory to the 2016 NLCD resulted in adding another 134 (2.53 km²) features, which included 12 (0.38 km²) glaciers. The final inventory includes 2542 features composed of 1331 (366.52 km²) glaciers, 1176 (31.01 km²) perennial snowfields, and 35 (3.57 km²) buried ice deposits (Table 1; Figure 4). Most glaciers and perennial snowfields, 1554 (62%) were outlined using 2015 NAIP imagery with the remainder outlined using mostly NAIP imagery from 2013 to 2020.



238
239

240 **Figure 4.** The spatial distribution and number of glaciers and perennial snowfields, greater than
 241 0.01 km², in the western United States. Colors indicate the date of aerial and satellite imagery
 242 used to outline the features. The line is the cumulative total. Base imagery from Esri Inc. Inset is
 243 a bar graph and cumulative sum of the number of glaciers and perennial snowfields digitized in
 244 each image date.
 245

246 **Table 1.** The summary of the glacier inventory for the American West, exclusive of Alaska.
 247 Number is the total number of features within each classification (Class), ‘Max Area’ is the
 248 largest area of the feature within that class and ‘Mean Area’ is the average area. Note that the
 249 uncertainty of ‘Buried ice’ is unknown.
 250


State/Region/Class	Number	Total Area km²	Max Area km²	Mean Area km²
California	132	10.63 ± 0.61	1.45	0.08
Cascade Range	39	5.74 ± 0.37	1.45	0.15
Buried ice	5	0.44	0.16	0.09
Glaciers	10	4.61 ± 0.17	1.45	0.46
Perennial snowfields	24	0.68 ± 0.21	0.08	0.03
Sierra Nevada	91	4.86 ± 0.23	0.66	0.05
Buried ice	2	0.13	0.10	0.06
Glaciers	64	4.37 ± 0.12	0.66	0.07
Perennial snowfields	25	0.37 ± 0.11	0.03	0.01
Trinity Alps	2	0.03 ± 0.00	0.02	0.02
Glaciers	2	0.03 ± 0.00	0.02	0.02
Colorado	84	2.20 ± 0.46	0.16	0.03
Elk Mountains	5	0.09 ± 0.03	0.03	0.02
Glaciers	1	0.01 ± 0.00	0.01	0.01
Perennial snowfields	4	0.08 ± 0.02	0.03	0.02
Front Range	58	1.73 ± 0.33	0.16	0.03
Glaciers	13	0.74 ± 0.03	0.16	0.06
Perennial snowfields	45	0.99 ± 0.30	0.09	0.02
Gore Range	7	0.11 ± 0.03	0.02	0.02
Glaciers	1	0.02 ± 0.00	0.02	0.02
Perennial snowfields	6	0.09 ± 0.03	0.02	0.02
Medicine Bow Mountains	1	0.04 ± 0.01	0.04	0.04
Perennial snowfields	1	0.04 ± 0.01	0.04	0.04
Park Range	6	0.11 ± 0.03	0.03	0.02
Perennial snowfields	6	0.11 ± 0.03	0.03	0.02
San Miguel Mountains	5	0.07 ± 0.02	0.02	0.01
Perennial snowfields	5	0.07 ± 0.02	0.02	0.01
Sawatch Range	2	0.04 ± 0.01	0.03	0.02
Perennial snowfields	2	0.04 ± 0.01	0.03	0.02
Idaho	6	0.08 ± 0.02	0.02	0.01
Sawtooth Range	6	0.08 ± 0.02	0.02	0.01

Perennial snowfields	6	0.08 ± 0.02	0.02	0.01
Montana	416	30.26 ± 2.27	1.45	0.07
Beartooth -Absaroka	111	6.07 ± 0.64	0.45	0.05
Buried ice	1	0.04	0.04	0.04
Glaciers	50	4.31 ± 0.12	0.45	0.09
Perennial snowfields	60	1.72 ± 0.52	0.22	0.03
Bitterroot Range	4	0.08 ± 0.02	0.03	0.02
Glaciers	1	0.03 ± 0.00	0.03	0.03
Perennial snowfields	3	0.05 ± 0.02	0.02	0.02
Cabinet Mountains	9	0.25 ± 0.08	0.08	0.03
Perennial snowfields	9	0.25 ± 0.08	0.08	0.03
Crazy Mountains	13	0.27 ± 0.06	0.04	0.02
Glaciers	3	0.06 ± 0.00	0.04	0.02
Perennial snowfields	10	0.21 ± 0.06	0.04	0.02
Lewis Range	230	21.38 ± 1.15	1.45	0.09
Glaciers	145	19.22 ± 0.50	1.45	0.13
Perennial snowfields	85	2.16 ± 0.65	0.09	0.03
Mission-Swan-Flathead	49	2.20 ± 0.34	0.22	0.04
Glaciers	11	1.16 ± 0.02	0.22	0.11
Perennial snowfields	38	1.04 ± 0.31	0.09	0.03
Oregon	116	15.38 ± 1.62	1.16	0.13
Cascade Range	110	15.24 ± 1.58	1.16	0.14
Buried ice	7	1.25	0.45	0.18
Glaciers	42	11.90 ± 0.95	1.16	0.28
Perennial snowfields	61	2.09 ± 0.63	0.15	0.03
Wallowa Mountains	6	0.14 ± 0.63	0.04	0.02
Perennial snowfields	6	0.14 ± 0.04	0.04	0.02
Washington	1481	312.26 ± 16.33	11.24	0.21
Cascade Range-Northern	1126	186.58 ± 9.64	6.06	0.17
Buried ice	10	0.50	0.15	0.05
Glaciers	706	176.27 ± 6.70	6.06	0.25
Perennial snowfields	410	9.80 ± 2.94	0.16	0.02
Cascade Range-Southern	219	101.66 ± 5.86	11.24	0.46
Buried ice	10	1.20	0.30	0.12
Glaciers	69	95.64 ± 4.42	11.24	1.39
Perennial snowfields	140	4.82 ± 1.45	0.33	0.03
Olympic Mountains	136	24.02 ± 0.82	5.09	0.18
Glacier	106	23.44 ± 0.65	5.09	0.22
Perennial snowfield	30	0.57 ± 0.17	0.06	0.02
Wyoming	307	30.29 ± 2.34	2.32	0.10
Absaroka Range	62	1.44 ± 0.33	0.12	0.02
Glacier	10	0.48 ± 0.05	0.12	0.05
Perennial snowfield	52	0.96 ± 0.29	0.05	0.02
Bighorn Mountains	8	0.42 ± 0.03	0.22	0.05
Glacier	3	0.34 ± 0.01	0.22	0.11


Perennial snowfield	5	0.08 ± 0.02	0.03	0.02
Teton Range	49	2.04 ± 0.21	0.23	0.04
Glacier	20	1.46 ± 0.03	0.23	0.07
Perennial snowfield	29	0.59 ± 0.18	0.05	0.02
Wind River Range	188	26.39 ± 1.76	2.32	0.14
Glacier	74	22.42 ± 0.57	2.32	0.30
Perennial snowfield	114	3.97 ± 1.19	0.26	0.03
Grand Total	2542	401.10 ± 23.64	11.24	0.16

251
252 Before summarizing the inventory data, a note about the content in Appendix A. It summarizes
253 the officially named glaciers that we regard as snowfields or missing; labeling issues found in the
254 USGS Geographic Names Information System, the official agency responsible for hosting the
255 names and locations of landscape features; and detailed notes, organized by US State, on the
256 specific imagery used and challenges encountered digitizing glacier and snowfield outlines.

257
258 The glaciers and perennial snowfields are generally small, averaging 0.28 and 0.03 km²,
259 respectively. Like glaciers elsewhere in the northern hemisphere, most glaciers face north to east.
260 (Evans, 2006; Fountain et al., 2017; Schiefer et al., 2007). The distribution of glacier area is
261 skewed toward smaller ice masses (Figure 5a). The State of Washington in the Pacific Northwest
262 has the largest number of glaciers, ice area and the largest glacier (11.24 km² Emmons Glacier)
263 of any of the other states (Table 1). Indeed, the glacier cover on Mount Rainier alone (77.37 km²)
264 is greater than the total sum in all the other states (71.16 km²). The elevation distribution of
265 glacier-covered area is bimodal with maxima at 2400 m and 3650 m (Figure 5b). The spatial
266 distribution of elevations shows a regional climate control with the lowest glaciers and perennial
267 snowfields in the maritime climate of the Pacific Northwest of Washington, Oregon, northern
268 California, and western Montana and the high elevations located in the continental climate of
269 central California, Colorado, Wyoming and southern Montana (Figure 6).



271 Figure 5. The area and elevation distribution of glaciers in the western U.S., (a) Histogram
 272 showing the number of glaciers as a function of area. The x-axis intervals are log intervals; (b)
 273 Elevation distribution of glacier-covered area.
 274



276
 277 Figure 6. Elevation distribution of glaciers and perennial snowfields across the western US. Base
 278 imagery from Esri Inc.

279
 280 The final inventory conflicts with the current database of the Geographic Names Information
 281 System (<https://www.usgs.gov/us-board-on-geographic-names/domestic-names>). The inventory
 282 excludes 52 officially named glaciers because 2 have disappeared, 25 were classified as
 283 perennial snowfields, the area of 18 was less than 0.01 km², and 7 were considered rock glaciers
 284 (Appendix A, Table A1). In some cases, a named glacier or snowfield had split into multiple
 285 pieces since the original USGS mapping; all pieces were assigned the same name in the
 286 inventory (Appendix A, Table A2). Several labels that identify the name of the glacier are not
 287 clearly associated with a specific glacier and these are listed in Table 7.3.

288
 289

290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334

4. Discussion

The advent of relatively frequent high resolution (≤ 1 m) optical aerial and satellite imagery available at little or no cost has made compiling and updating glacier inventories a realistic opportunity. Finding suitable imagery spanning only a few years apart provides a near-snapshot of glacier cover. This contrasts strongly with mapping efforts only a few decades ago when aerial-only photographic surveys required decades to cover the western US (Gesch et al., 2002). And the advent of GIS software made digitizing, summarizing, and interrogating digital outlines practical.

We had used the Fountain et al., 2017 historic inventory as a template to locate and update the perimeters of all the glaciers and perennial snowfields. Considering that the inventory was derived from the U.S. Geological Survey 1:24,000 maps, a result of a national effort to remap the entire country at a higher resolution, we were surprised that 240 features (~10%) were missed. These missing features were revealed after comparison with two other independently derived inventories. We had a similar experience in a prior study when comparing two independently derived glacier inventories. Together they suggest that independent efforts are important to compiling a comprehensive inventory.

Multiple checks more accurately define glacier perimeters (Leigh et al., 2019). Different investigators may make different decisions about glacier boundaries and results can differ particularly in debris-covered conditions or along flow divides (Paul et al., 2013). When they agree, it provides some confidence of the interpretation accuracy and where they disagree it provides input for estimating interpretation error.

The total area of glaciers in the western US, 367 km², is a little smaller than that in Austria, 415 km², (Fischer et al., 2015). Like glacier populated regions elsewhere the distribution of glacier area is skewed towards smaller glaciers (e.g. Linsbauer et al., 2012; Mishra et al., 2023; Zalazar et al., 2020). The uncertainty in glacier area is also similar with an overall 5% uncertainty for the total area. Paul et al. (2020) report an uncertainty of 3.3% over a set of 15 glaciers, 4% for 7 glaciers (Zalazar et al., 2020), 2.3% for 15 glaciers (Linsbauer et al., 2021). Our assessment method differs from those cited here in that we estimate the uncertainty for each individual glacier rather than upscaling the uncertainty calculated for a small subsample.

5. Data products and availability

The data are available in three formats. The geospatial data and attribute tables are available in the shapefile (Esri) format and in an open source GeoJSON format. The attribute table is also available as an EXCEL file. These data products can be obtained from <https://doi.org/10.15760/geology-data.03> (Fountain and Glenn, 2022) and from the Global Land Ice Measurements from Space website <http://glims.colorado.edu/glacierdata/>. Maxar imagery was accessed through the USGS and NGA NEXTVIEW license. The Maxar imagery has limited availability owing to restrictions (proprietary interest). Contact cmcneil@usgs.gov for more

335 information.

336

337 **6. Conclusions**

338

339 We have compiled a new and comprehensive inventory of glaciers and perennial snowfields in
340 the western US from aerial and satellite imagery. Results show that 2542 features are currently
341 present and include 1331 (366.52 km²) glaciers, 1176 (31.01 km²) perennial snowfields, and 35
342 (3.57 km²) buried ice deposits. Most of the data were acquired from 2015 NAIP imagery with the
343 remainder from NAIP imagery and a few satellite images acquired over the period of 2013 to
344 2020. The state of Washington has the greatest number and area of glaciers and perennial
345 snowfields. This product updates an older inventory based on USGS 1:24000 maps compiled in
346 the middle-late 1900's. The new inventory is a significant improvement in accuracy because the
347 archive of historical imagery in Google Earth greatly aided our efforts to classify glaciers versus
348 perennial snowfields. Finally, this new inventory provides a baseline for assessing glacier change
349 in the coterminous US.

350

351 **7. Appendix A**

352

353 **A1 Missing Glaciers**

354 **Table A1** List of officially named glaciers not classified as glaciers and excluded from the final
355 inventory. Names come from the Geographic Names Information System, (US Geological
356 Survey, 2022). The 'Reason' column lists why the named glacier is no longer considered a glacier
357 in our inventory.

358

State/Region/Glacier Name	Reason
California	
Sierra Nevada	
Matthes Glaciers	rock glacier
Mount Warlow Glacier	rock glacier
Powell Glacier	rock glacier
Colorado	
Front Range	
Isabelle Glacier	perennial snowfield
Mills Glacier	perennial snowfield
Moomaw Glacier	perennial snowfield
Peck Glacier	perennial snowfield
Rowe Glacier	< 0.01 km ²
Saint Marys Glacier	< 0.01 km ²
Taylor Glacier	rock glacier
The Dove	< 0.01 km ²
Idaho	
Lost River Range	
Borah Glacier	rock glacier
Montana	

Beartooth Mountains-Absaroka Range	
Grasshopper Glacier	rock glacier
Cabinet Mountains	
Blackwell Glacier	perennial snowfield
Crazy Mountains	
Grasshopper Glacier	rock glacier
Lewis Range	
Boulder Glacier	perennial snowfield
Mission-Swan-Flathead Ranges	
Fissure Glacier	< 0.01 km ²
Gray Wolf Glacier	perennial snowfield
<hr/>	
Oregon	
Cascade Range	
Carver Glacier	perennial snowfield
Clark Glacier	perennial snowfield
Irving Glacier	perennial snowfield
Lathrop Glacier	< 0.01 km ²
Palmer Glacier	perennial snowfield
Skinner Glacier	perennial snowfield
Thayer Glacier	< 0.01 km ²
Wallowa Mountains	
Benson Glacier	perennial snowfield
<hr/>	
Washington	
Cascade Range-Northern	
Lyll Glacier	perennial snowfield
Milk Lake Glacier	disappeared
Snow Creek Glacier	perennial snowfield
Spider Glacier	perennial snowfield
Table Mountain Glacier	< 0.01 km ²
Cascade Range-Southern	
Ape Glacier	< 0.01 km ²
Dryer Glacier	perennial snowfield
Forsyth Glacier	< 0.01 km ²
Meade Glacier	perennial snowfield
Nelson Glacier	< 0.01 km ²
Packwood Glacier	perennial snowfield
Pinnacle Glacier	< 0.01 km ²
Pyramid Glaciers	< 0.01 km ²
Shoestring Glacier	< 0.01 km ²
Stevens Glacier	perennial snowfield
Talus Glacier	perennial snowfield
Unicorn Glacier	< 0.01 km ²
Williwakas Glacier	perennial snowfield
Olympic Mountains	
Anderson Glacier	perennial snowfield

Lillian Glacier	< 0.01 km ²
Wyoming	
Absaroka Range	
DuNoir Glacier	< 0.01 km ²
Teton Range	
Petersen Glacier	< 0.01 km ²
Teepe Glacier	perennial snowfield
Wind River Range	
Hooker Glacier	disappeared
Harrower Glacier	perennial snowfield
Tiny Glacier	< 0.01 km ²

359

360

361 **A2 Glaciers that have split into multiple pieces and current errors in glacier label names**

362

363 **Table A2.** List of named glaciers that have split into multiple pieces. Names come from the
 364 Geographic Names Information System ([https://www.usgs.gov/tools/geographic-names-](https://www.usgs.gov/tools/geographic-names-information-system-gnis)
 365 [information-system-gnis](https://www.usgs.gov/tools/geographic-names-information-system-gnis)). ‘Count’ refers to the number of pieces in the updated inventory.
 366 ‘Classes’ is the classification of the pieces; glacier, perennial snowfield, buried-ice, or a
 367 combination.

State/Region/Glacier Name	Count	Classes
California		
Cascade Range		
Bolam Glacier	2	Glaciers and perennial snowfields
Hotlum Glacier	2	Glaciers and perennial snowfields
Whitney Glacier	2	Glaciers and perennial snowfields
Wintun Glacier	3	Glaciers and perennial snowfields
Sierra Nevada		
Goethe Glacier	2	Glaciers only
Lyell Glacier	4	Glaciers and perennial snowfields
Norman Clyde Glacier	3	Glaciers only
Powell Glacier	2	Glacier and Buried-ice
Colorado		
Front Range		
Saint Vrain Glaciers	6	Glaciers and perennial snowfields
Montana		
Beartooth Mountains-Absaroka Range		
Castle Rock Glacier	3	Glaciers and perennial snowfields
Granite Glacier	2	Glaciers only
Grasshopper Glacier	4	Glaciers and perennial snowfields
Hopper Glacier	2	Glaciers and perennial snowfields
Snowbank Glacier	2	Glaciers only
Wolf Glacier	2	Glaciers only
Lewis Range		
Agassiz Glacier	3	Glaciers only
Blackfoot Glacier	2	Glaciers only

Carter Glaciers	2	Glaciers and perennial snowfields
Dixon Glacier	3	Glaciers and perennial snowfields
Harrison Glacier	5	Glaciers and perennial snowfields
Kintla Glacier	2	Glaciers only
Logan Glacier	2	Glaciers only
Shepard Glacier	3	Glaciers only
Siyeh Glacier	2	Glaciers only
Two Ocean Glacier	2	Glaciers only
Whitecrow Glacier	5	Glaciers and perennial snowfields
Mission Range-Swan Range-Flathead Range		
Swan Glaciers	3	Glaciers and perennial snowfields
Oregon		
Cascade Range		
Bend Glacier	3	Glaciers and perennial snowfields
Clark Glacier	2	Perennial snowfields only
Collier Glacier	2	Glaciers only
Diller Glacier	2	Glaciers and perennial snowfields
Glisan Glacier	2	Glaciers and perennial snowfields
Ladd Glacier	4	Glaciers and perennial snowfields
Langille Glacier	5	Glaciers and perennial snowfields
Newton Clark Glacier	3	Glaciers and perennial snowfields
Palmer Glacier	2	Perennial snowfields only
Prouty Glacier	3	Glaciers and perennial snowfields
Renfrew Glacier	2	Glaciers and perennial snowfields
Russell Glacier	2	Glaciers only
Sandy Glacier	4	Glaciers and perennial snowfields
Skinner Glacier	4	Perennial snowfields only
Waldo Glacier	3	Glaciers only
White River Glacier	2	Glaciers and perennial snowfields
Whitewater Glacier	3	Glaciers only
Zigzag Glacier	3	Glaciers and perennial snowfields
Washington		
Cascade Range-Northern		
Borealis Glacier	4	Glaciers only
Buckner Glacier	2	Glaciers only
Butterfly Glacier	4	Glaciers only
Colchuck Glacier	2	Glaciers only
Company Glacier	3	Glaciers only
Cool Glacier	2	Glaciers and perennial snowfields
Dana Glacier	3	Glaciers only
Dark Glacier	3	Glaciers only
Dome Glacier	2	Glaciers only
Douglas Glacier	4	Glaciers and perennial snowfields
Dusty Glacier	2	Glaciers and perennial snowfields
East Nooksack Glacier	5	Glaciers only
Entiat Glacier	4	Glaciers and perennial snowfields
Forbidden Glacier	2	Glaciers only
Fremont Glacier	2	Glaciers only
Goode Glacier	2	Glaciers only

Hadley Glacier	5	Glaciers only
Hanging Glacier	2	Glaciers only
Hinman Glacier	4	Glaciers only
Honeycomb Glacier	3	Glaciers and perennial snowfields
Inspiration Glacier	3	Glaciers and perennial snowfields
Isella Glacier	2	Glaciers and perennial snowfields
Jerry Glacier	2	Glaciers only
Kimtah Glacier	3	Glaciers only
LeConte Glacier	7	Glaciers and perennial snowfields
Lyll Glacier	2	Perennial snowfields only
Mazama Glacier	3	Glaciers and perennial snowfields
McAllister Glacier	2	Glaciers only
Middle Cascade Glacier	2	Glaciers only
Neve Glacier	3	Glaciers only
No Name Glacier	5	Glaciers and perennial snowfields
Nohokomeen Glacier	2	Glaciers only
North Klawatti Glacier	2	Glaciers and perennial snowfields
Pilz Glacier	3	Glaciers and perennial snowfields
Price Glacier	4	Glaciers only
Ptarmigan Glacier	2	Glaciers and perennial snowfields
Queest-alb Glacier (not official)	3	Glaciers and perennial snowfields
Rainbow Glacier	3	Glaciers and perennial snowfields
Redoubt Glacier	2	Glaciers only
Richardson Glacier	2	Glaciers only
S Glacier	3	Glaciers only
Sandalee Glacier	4	Glaciers only
Scimitar Glacier	3	Glaciers only
Sholes Glacier	4	Glaciers only
Sitkum Glacier	4	Glaciers and perennial snowfields
Snow Creek Glacier	2	Perennial snowfields only
South Cascade Glacier	2	Glaciers only
Spider Glacier	2	Glaciers only
Suiattle Glacier	2	Glaciers only
Sulphide Glacier	2	Glaciers only
Thunder Glacier	3	Glaciers only
Thunder Glacier	2	Glaciers only
White Chuck Glacier	5	Glaciers and perennial snowfields
White Salmon Glacier	2	Glaciers only
Wyeth Glacier	3	Glaciers and perennial snowfields
Cascade Range-Southern		
Adams Glacier	4	Glaciers and perennial snowfields
Avalanche Glacier	2	Glaciers only
Conrad Glacier	3	Glaciers and perennial snowfields
Cowlitz Glacier	2	Glaciers and perennial snowfields
Crescent Glacier	2	Glaciers and perennial snowfields
Flett Glacier	6	Glaciers and perennial snowfields
Fryingpan Glacier	5	Glaciers and perennial snowfields
Gotchen Glacier	2	Glaciers and perennial snowfields
Kautz Glacier	2	Glaciers and perennial snowfields

Klickitat Glacier	2	Glaciers only
Lava Glacier	3	Glaciers and perennial snowfields
McCall Glacier	6	Glaciers and perennial snowfields
Meade Glacier	5	Perennial snowfields only
North Mowich Glacier	2	Glaciers and perennial snowfields
Ohanapecosh Glacier	6	Glaciers and perennial snowfields
Paradise Glacier	3	Glaciers and perennial snowfields
Pinnacle Glacier	3	Glaciers and perennial snowfields
Puyallup Glacier	2	Glaciers and perennial snowfields
Pyramid Glacier	4	Glaciers and perennial snowfields
Russell Glacier	2	Glaciers only
Sarvant Glaciers	4	Glaciers and perennial snowfields
South Mowich Glacier	2	Glaciers only
South Tahoma Glacier	2	Glaciers and perennial snowfields
Success Glacier	2	Glaciers and perennial snowfields
Van Trump Glacier	10	Glaciers and perennial snowfields
White Salmon Glacier	2	Glaciers only
Whitman Glacier	5	Glaciers and perennial snowfields
Wilson Glacier	3	Glaciers and perennial snowfields
Olympic Mountains		
Blue Glacier	2	Glaciers only
Cameron Glaciers	4	Glaciers and perennial snowfields
Carrie Glacier	2	Glaciers only
Eel Glacier	2	Glaciers only
White Glacier	2	Glaciers only
Wyoming		
Teton Range		
Middle Teton Glacier	2	Glaciers and perennial snowfields
Triple Glaciers	3	Glaciers only
Wind River Range		
Bull Lake Glacier	3	Glaciers and perennial snowfields
Dinwoody Glacier	2	Glaciers only
Dinwoody Glaciers	3	Glaciers and perennial snowfields
Grasshopper Glacier	3	Glaciers only
Harrower Glacier	2	Perennial snowfields only
Helen Glacier	3	Glaciers only
Lower Fremont Glacier	4	Glaciers and perennial snowfields
Mammoth Glacier	2	Glaciers and perennial snowfields
Minor Glacier	2	Glaciers and perennial snowfields
Sacagawea Glacier	4	Glaciers and perennial snowfields
Sourdough Glacier	2	Glaciers and perennial snowfields
Stroud Glacier	3	Glaciers and perennial snowfields
Twins Glacier	2	Glaciers and perennial snowfields
Upper Fremont Glacier	2	Glaciers and perennial snowfields

368

369

370 **A3 Labelling errors in the U.S. Geographic Names Information System**

371 **Table A3.** List of officially named glaciers where we identified an issue with the glacier name
 372 on the 1:24000 U.S. Geological Survey topographical maps (Fountain et al., 2017). Names come
 373 from the Geographic Names Information System ([https://www.usgs.gov/tools/geographic-](https://www.usgs.gov/tools/geographic-names-information-system-gnis)
 374 [names-information-system-gnis](https://www.usgs.gov/tools/geographic-names-information-system-gnis)). The 'Issue' column lists the type of issue identified. 'Not
 375 labeled' indicates the feature was present but not labeled, 'Misidentified' indicates the wrong
 376 feature was labeled, and 'Label unclear' means the location of the label is not clearly associated
 377 with a specific glacier.

State/Region/Glacier Name	Issue
Colorado	
Front Range	
Arikaree Glacier	Not labeled
Navajo Glacier	Not labeled
Oregon	
Cascade Range	
Carver Glacier	Misidentified
Milk Creek Glacier	Not labeled
Washington	
Cascade Range-Northern	
S Glacier	Label unclear
Snow Creek Glacier	Label unclear
South Glacier	Not labeled
Cascade Range-Southern	
No Name Glacier	Not labeled
Stevens Glacier	Not labeled
Wyoming	
Wind River Range	
Dinwoody Glaciers	Label unclear
Fremont Glaciers	Label unclear

378

379

380 **A4 Notes on imagery and interpretation challenges by State.**

381

382 This appendix, organized by US State, then by mountain range, summarizes the specific
 383 imagery used, challenges encountered in feature identification and outline digitization. The
 384 Selkowitz and Forster (2016) inventory is referred to as the SFI and the National Land Cover
 385 Database inventory (Dewitz, 2019) is referred to as the NLCD.

386

387 **A4.1 California**

388

389 Imagery and DEMs used are listed in Tables A4, A5, A6.

390


391 **Cascade Range**

392

393 **Mount Shasta**

394

395 The 2020 black and white Maxar imagery was most useful because of the minimal
396 seasonal snow cover. The 2018 NAIP imagery was helpful in situations where the 2020
397 imagery was obscured by shadow, distortion, or misaligned, and when color was needed
398 to improve interpretation. The 2010 lidar DEM (Robinson, 2014; Table A4) was used to
399 create a multidirectional hillshade to improve perspective and interpretation (Figure A1).
400 The rock debris on the termini of most glaciers and rock debris on some of the upper
401 parts of the glaciers were challenging to interpret. It was hard to determine whether ice
402 was present under the debris and whether that ice is part of the active glacier. Spatial
403 patterns of debris, debris contrasts, and melt streams flowing from the debris were used to
404 estimate the glacier boundaries.



405
406 **Figure A1.** Mt. Shasta glaciers in bluish white, perennial snowfields/ice patches in lavender
407 draped over a 3D rendering created from 2010 lidar (Robinson, 2014).
408

409 **Sierra Nevada**

411 The 2014 NAIP imagery was the best imagery due to low snow cover. In some cases,
412 features were difficult to outline because of shadow or image quality. In these cases,
413 2013/2012 Google Earth imagery were used. Some glaciers were reclassified as rock
414 glaciers by Trcka (2020). These were re-examined and where we agreed they were
415 removed from the initial glacier inventory. Defining whether the feature was a glacier or
416 rock glacier was often difficult, see Colorado section for more discussion.
417

418 **Trinity Alps**

419 The 2018 imagery was the best for the least snow cover. Justin Garwood (Garwood et al.,
420 2020) provided outlines for two glaciers, Grizzly and Salmon. The area of the most recent
421 outline of the Salmon Glacier was $< 0.01 \text{ km}^2$ and was not included in this inventory. By
422 2018 all of the other features mapped by the USGS (Fountain et al., 2017) were less than
423 0.01 km^2 or had disappeared. An additional feature was added based on the 2016 NLCD
424 (Jin et al., 2019).
425
426

427 **Table A4.** List of NAIP imagery used for outlining glaciers and perennial snowfields in
 428 California. ‘Date’ is the start and end date for flights covering the glaciated portions of the NAIP
 429 image. In some cases, flights were completed in a single day.
 430

Region/Year/Filename	County	Date (Year-M-D)
Cascade Range		
2014		
ortho_1-1_1n_s_ca089_2014_1.sid	Shasta	2014-07-13
ortho_1-1_1n_s_ca093_2014_1.sid	Siskiyou	2014-06-23 to 2014-07-18
2018		
ortho_1-1_hn_s_ca093_2018_1.sid	Siskiyou	2018-07-21 to 2018-09-25
Sierra Nevada		
2014		
ortho_1-1_1n_s_ca019_2014_1.sid	Fresno	2014-07-23 to 2014-08-23
ortho_1-1_1n_s_ca027_2014_1.sid	Inyo	2014-07-23 to 2014-08-23
ortho_1-1_1n_s_ca039_2014_2.sid	Madera	2014-07-18 to 2014-08-15
ortho_1-1_1n_s_ca051_2014_1.sid	Mono	2014-07-17 to 2014-08-15
ortho_1-1_1n_s_ca107_2014_1.sid	Tulare	2014-08-23 to 2014-08-23
Trinity Alps		
2018		
ortho_1-1_hn_s_ca093_2018_1.sid	Siskiyou	2018-07-21 to 2018-09-25

431

432

433 **Table A5.** List of dates of the Maxar imagery used for outlining glaciers and perennial
 434 snowfields in California.

Region/ Date (Year-M-D)

Cascade Range

2020-10-05

435

436

437 **Table A6.** List of U.S. Geological Survey digital elevation models used for outlining glaciers
 438 and perennial snowfields in California.

Filename	Date	Citation	URL
ds852_lidar	2010	Robinson (2014)	https://pubs.er.usgs.gov/publication/ds852

439

A4.2 Colorado

441

442 The 2015 NAIP was generally free of seasonal snow. Where it persisted at the terminus of a few
 443 glaciers, images for the same year in Google Earth aided perimeter interpretation. Imagery used

444 are listed in Table A7.

445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460

Elk Mountains

No features were mapped in the Elk Mountains by the USGS (Fountain et al., 2017). One glacier and four perennial snowfields were added from the SFI.

Front Range

The most recent inventory for the Front Range was Hoffman et al. (2007), which used aerial photographs to map the 2001 extent of glaciers. Many features in the Front Range are difficult to classify. The issue is the difference between a glacier or perennial snowfield and a rock glacier. Those that are part of the rock glacier are deleted from the glacier inventory. Those that seem to be separate from rock glaciers are retained. This is a judgement call. From a hydrological point of view, if a snow-ice patch that is part of a rock glacier was counted separately from a rock glacier, it is double counting a water feature.

461 **Table A7.** List of NAIP imagery used for outlining glaciers and perennial snowfields in
462 Colorado. ‘Date’ is the start and end date for flights covering the glaciated portions of the NAIP
463 image. In some cases, flights were completed in a single day.
464

Region/Year/Filename	County	Date (Year-M-D)
Elk Mountains		
2015		
ortho_1-1_1n_s_co051_2015_1.sid	Gunnison	2015-09-10 to 2015-09-11
Front Range		
2015		
ortho_1-1_1n_s_co013_2015_1.sid	Boulder	2015-08-25 to 2015-09-20
ortho_1-1_1n_s_co049_2015_1.sid	Grand	2015-08-25 to 2015-09-20
ortho_1-1_1n_s_co057_2015_1.sid	Jackson	2015-09-09
ortho_1-1_1n_s_co069_2015_1.sid	Larimer	2015-08-25 to 2015-09-09
Gore Range		
2015		
ortho_1-1_1n_s_co037_2015_1.sid	Eagle	2015-09-10
Medicine Bow Mountains		
2015		
ortho_1-1_1n_s_co057_2015_1.sid	Jackson	2015-09-09
Park Range		
2015		
ortho_1-1_1n_s_co057_2015_1.sid	Jackson	2015-09-09
San Miguel Mountains		
2015		
ortho_1-1_1n_s_co033_2015_1.sid	Dolores	2015-09-11
ortho_1-1_1n_s_co091_2015_1.sid	Ouray	2015-09-11
ortho_1-1_1n_s_co111_2015_1.sid	San Juan	2015-09-12

Sawatch Range

2015

ortho_1-1_1n_s_co037_2015_1.sid	Eagle	2015-09-10
ortho_1-1_1n_s_co097_2015_1.sid	Pitkin	2015-09-10 to 2015-09-11

465

466 A4.3 Idaho

467

468 The imagery quality was generally snow free. Of the glacier mapped by the USGS (Fountain et
469 al., 2017) only two remain and are classified as perennial snowfields. The Borah Glacier was
470 officially named in 2021 (U.S. Board of Geographic Names), but is < 0.01 km², and is not
471 included in the inventory. Table A8 lists the imagery used.

472

473

474 **Table A8.** List of NAIP imagery used for outlining glaciers and perennial snowfields in Idaho.
475 'Date' is the start and end date for flights covering the glaciated portions of the NAIP image. In
476 some cases, flights were completed in a single day.

477

Region/Year/Filename	County	Date (Year-M-D)
Sawtooth Range		
2013		
ortho_1-1_hn_s_id015_2013_1.sid	Boise	2013-09-07
2015		
ortho_1-1_1n_s_id013_2015_1.sid	Blaine	2015-07-30
ortho_1-1_1n_s_id015_2015_1.sid	Boise	2015-09-08 to 2015-09-09
2019		
ortho_1-1_hn_s_id037_2019_1.sid	Custer	2019-07-25 to 2019-08-26

478

479

480 A4.4 Montana

481

482 Image quality varied between mountain ranges due to differences in snow cover. Tables A9 and
483 A10 list the imagery used.

484

485 **Beartooth-Absaroka Range**

486 The 2015 NAIP imagery was the best overall imagery due to the least snow, but Google
487 Earth was occasionally used as well. Google Earth had imagery dated to 9/11/2015; often
488 with less seasonal snow than the NAIP imagery. To counter any mismatch in projection,
489 outlines digitized in Google Earth were imported to ArcGIS and projected to match the
490 NAIP projection.

491

492 **Bitterroot Range**

492 No features were mapped in the Bitterroot Range by the USGS (Fountain et al., (2017)).
493 One glacier and three perennial snowfields were added based on the NLCD.

494

495 **Cabinet Range**

496 The USGS mapped four features $\geq 0.01 \text{ km}^2$ (Fountain et al., 2017). Inspection of the
497 2015 only one was $\geq 0.01 \text{ km}^2$. Seven glaciers and perennial snowfields were added; five
498 were identified in our initial inventory, the other two were identified by the SFI and
499 NLCD, respectively. All were less than 0.05 km^2 .

500

501 **Crazy Mountains**

502 The 2013 NAIP imagery was the best imagery available and included limited seasonal
503 snow. The 2019 Maxar imagery had too much seasonal snow.

504

505 **Lewis Range (Glacier National Park)**

506 The most recent published glacier inventory is a 2015 USGS inventory (Fagre et al.,
507 2017). They outlined the main-body of named-glaciers using 2015 Maxar imagery. We
508 digitized the outlines of all glaciers and perennial snowfields using 2015 Maxar imagery
509 where available. Elsewhere, 2015 and 2013 NAIP imagery were used; both years had lots
510 of seasonal snow cover. Two major glaciers, Blackfoot (Figure A2) and Harrison (Figure
511 A3) glaciers, separated into pieces as it retreated since it was originally mapped by the
512 USGS (Fountain et al., 2007).

513

514 **Madison Range**


515 The 2013 NAIP imagery was the only imagery used due to extensive snow in the other
516 years. No glaciers or perennial snowfields were found. Of the two features $\geq 0.01 \text{ km}^2$
517 mapped by the USGS (Fountain et al., 2017), the 2013 imagery showed that one feature
518 is a rock glacier and the other was less than 0.01 km^2 .

519

520 **Mission-Swan-Flathead Ranges**


521 Based on the least snow cover, the 2013 NAIP was better in the Mission and Flathead
522 Ranges, and the 2015 NAIP was better in the Swan Range. No glaciers or perennial
523 snowfields remain in the Flathead Range.

524



525

526 **Figure A2.** The updated (2015) outlines for the Blackfoot Glacier including the main glacier
527 body (red) and the additional smaller glacier (orange). Base image from the NAIP taken in 2013.



528

529 **Figure A3.** The updated (2015) outlines for Harrison Glacier including the main glacier body
 530 (red) and the additional smaller glaciers (orange). Base image from the NAIP taken in 2013.

531

532

533 **Table A9.** List of NAIP imagery used for outlining glaciers and perennial snowfields in
 534 Montana. 'Date' is the start and end date for flights covering the glaciated portions of the NAIP
 535 image. In some cases, flights were completed in a single day.

536

Region/Year/Filename	County	Date (Year-M-D)
Beartooth Mountains-Absaroka Range		
2013		
ortho_1-1_1n_s_mt067_2013_1.sid	Park	2013-08-05 to 2013-09-11

2015	ortho_1-1_1n_s_mt009_2015_1.sid	Carbon	2015-08-10 to 2015-09-07
	ortho_1-1_1n_s_mt067_2015_1.sid	Park	2015-08-19 to 2015-09-11
	ortho_1-1_1n_s_mt095_2015_1.sid	Stillwater	2015-08-10 to 2015-09-07
Bitterroot Range			
2013	ortho_1-1_1n_s_mt001_2013_1.sid	Beaverhead	2013-08-04
2015	ortho_1-1_1n_s_mt081_2015_2.sid	Ravalli	2015-10-06 to 2015-11-07
Cabinet Mountains			
2015	ortho_1-1_1n_s_mt053_2015_2.sid	Lincoln	2015-09-11 to 2016-08-15
Crazy Mountains			
2013	ortho_1-1_1n_s_mt067_2013_1.sid	Park	2013-08-05 to 2013-09-11
	ortho_1-1_1n_s_mt097_2013_1.sid	Sweet Grass	2013-08-31 to 2013-09-10
2015	ortho_1-1_1n_s_mt067_2015_1.sid	Park	2015-08-19 to 2015-09-11
Lewis Range			
2013	ortho_1-1_1n_s_mt029_2013_1.sid	Flathead	2013-08-21 to 2013-09-01
	ortho_1-1_1n_s_mt035_2013_1.sid	Glacier	2013-08-21 to 2013-09-01
2015	ortho_1-1_1n_s_mt029_2015_2.sid	Flathead	2015-09-30 to 2016-10-21
	ortho_1-1_1n_s_mt035_2015_2.sid	Glacier	2015-10-14 to 2016-08-21
Mission Range-Swan-Flathead Ranges			
2013	ortho_1-1_1n_s_mt029_2013_1.sid	Flathead	2013-08-21 to 2013-09-01
	ortho_1-1_1n_s_mt063_2013_1.sid	Missoula	2013-09-01
2015	ortho_1-1_1n_s_mt047_2015_2.sid	Lake	2015-09-12 to 2016-08-15
	ortho_1-1_1n_s_mt063_2015_2.sid	Missoula	2015-09-12 to 2016-08-16

537
538
539
540
541

Table A10. List of dates of the Maxar imagery used for outlining glaciers and perennial snowfields in Montana.

<u>Region/ Date (Year-M-D)</u>
Lewis Range
2015-08-22
2015-09-01
2015-09-12
2015-09-25
2019-08-20

542 **A4.5 Oregon**

543

544 Tables A11, A12, and A13 list the imagery and DEM used.

545

546 **Cascade Range**

547

548 Seasonal snow cover was commonly present when this range was imaged by any of the
549 sensors making it difficult to find suitable imagery.

550

551 **Mount Hood**

552 The most recent glacier outlines for Mt. Hood were based on 2015 and 2016 Maxar color
553 imagery with interpretation aid using Google Earth. Due to seasonal snow some
554 professional judgement was required in places.

555

556 **Mount Jefferson**

557 The 2018 NAIP had extensive seasonal snow and was generally only useful near the
558 terminus of some glaciers. Used 2018 Maxar imagery that showed little seasonal snow,
559 but a little cloudy that masked a bit of Whitewater Glacier. Also used Google Earth to
560 help interpret some of the features.

561

562 **Three Sisters**

563 Maxar 2018 imagery was used, but the image was stretching along the feature's headwall
564 and for that segment of the outline 2018 NAIP imagery was used. Two versions of the
565 Maxar imagery for the same day are available, one color, one black and white. Color was
566 georectified but suffered stretching along some headwalls. A light early season snowfall
567 occurred before the Maxar image and the snow accumulated in some places just enough
568 to obscure the surface. So, the glacier or snow patch outline was the minimum of the two
569 images with occasional interpolation across the snowy surface to the nearest glacier edge.

570

571 **Mount Thielsen**

572 The Lathrop Glacier was named in 1981. At the time of the USGS mapping and now it is
573 $<0.01 \text{ km}^2$, and not counted as part of the inventory. Furthermore, Lathrop Glacier has
574 been known to disappear in some years and therefore fails the definition of a glacier.

575

576 **Wallowa Mountains**

577 No NAIP imagery was useful and Maxar did not image this region. We used the
578 8/30/2013 image from Google Earth, which was excellent with little snow. Features were
579 digitized in Google Earth and then imported into ArcGIS. Because we used NAIP as the
580 base imagery, we revised the outline from the projection in WGS84 (Google Earth) to
581 NAD83 UTM Zone 11 (NAIP).

582

583

584 **Table A11.** List of NAIP imagery used for outlining glaciers and perennial snowfields in
585 Oregon. 'Date' is the start and end date for flights covering the glaciated portions of the NAIP

586 image. In some cases, flights were completed in a single day.
 587

Region/Year/Filename	County	Date (Year-M-D)
Cascade Range		
2014		
ortho_1-1_1n_s_or017_2014_1.sid	Deschutes	2014-09-01
ortho_1-1_1n_s_or027_2014_1.sid	Hood River	2014-08-27 to 2014-09-05
ortho_1-1_1n_s_or039_2014_1.sid	Lane	2014-09-01
2016		
ortho_1-1_1n_s_or027_2016_1.sid	Hood River	2016-08-04
2017/2018		
ortho1-1_hn_s_or017_2017_2018_1.sid	Deschutes	2018-07-28
Wallowa Mountains		
2014		
ortho_1-1_1n_s_or063_2014_1.sid	Wallowa	2014-10-05

588
 589
 590 **Table A12.** List of dates of the Maxar imagery used for outlining glaciers and perennial
 591 snowfields in Oregon.
 592

Region/ Date (Year-M-D)
Cascade Range
2015-08-20
2015-09-11
2015-10-05
2016-09-10
2018-09-17
2020-09-20

593
 594
 595 **Table A13.** List of Oregon Department of Geology and Mineral Industries digital elevation
 596 models used for outlining glaciers and perennial snowfields in Oregon.
 597

Filename	Date	URL
2011_OLC_Deschutes	2011	gis.dogami.oregon.gov/maps/lidarviewer/

598
 599
 600 **A4.6 Washington**
 601
 602 The 2015 NAIP imagery was typically excellent with little snow cover, whereas the 2017 NAIP
 603 had more snow and the 2019 imagery had lots of snow. For most outlines, 2015 NAIP imagery
 604 was used. In some places, the 2017 NAIP imagery had less snow and was used instead. Maxar
 605 imagery was of limited use and often wasn't better than the 2015 or 2017 NAIP. Tables A14,
 606 A15, A16, list the imagery and DEMs used.
 607

608 **Cascade –Northern**

609 The glaciers and perennial snowfields were previously inventoried by (Dick, 2013).

610

611 **Mount Baker**

612 The 2015 NAIP imagery was the best and had little seasonal snow. Google Earth 2009
613 and 2019 imagery were used to help interpretation. A multidirectional hillshade and 3-
614 meter contour lines derived from a lidar DEM (Bard, 2017a); were used to help define
615 flow divides between glaciers, debris covered-ice, and buried ice. There are notable
616 differences between the NAIP imagery and DEM data, particularly in steep terrain, areas
617 of dark shadow, and debris-covered areas. The DEM helped correct these positional
618 errors and the benefit of supplying more information on surface texture.

619

620 Several buried-ice features were identified. The ice appeared to have decoupled from the
621 active glacier. In a few cases, debris-covered ice is included in the glacier outline because
622 the ice appears to be directly connected to the glacier, and there was evidence of
623 movement.

624

625 **Dragontail Peak**

626 The USGS Geographic Names Information Service (GNIS) locates Snow Creek Glacier
627 at a point on the edge of the southeast glacier (Fountain et al., 2007). In the 2015
628 imagery, the point is on bedrock, making it unclear which glacier the GNIS is naming.
629 The USGS identifies both glaciers as Snow Creek Glacier. We labeled both glaciers as
630 the Snow Creek Glacier.

631

632 **Glacier Peak**

633 For the Glacier Peak region, a multidirectional hillshade and 3-m contour lines derived
634 from a 2015 lidar DEM (Bard, 2017b) were used as a guide to define flow divides.

635

636 **Hurry-up Peak**

637 The point location of the South Glacier provided by the GNIS is over bedrock. We
638 assume the point refers to the glacier located ~150 m to the north of the point.

639

640 **Cascade –Southern**

641

642 **Goat Rocks**

643 Imagery from 2015 was best, but had more snow than desired. Too much snow was
644 present in 2017 but some ice was exposed. The 2019 imagery was too snowy for glacier
645 digitization.

646

647 The outlines are almost entirely based on 2015 imagery, and a few on 2017, where
648 needed. Used 2009 NAIP imagery to help define the headwalls at the Conrad, McCall,
649 and Packwood glaciers. Heard (2000) previously mapped the glacier perimeters. The
650 maximum extent of the seasonal snow covering the terminal regions was not digitized.
651 Typically digitized at scales of 1:600 to 1:800. Note that narrow arms of the snowfields
652 were not typically digitized knowing that they would probably disappear a few days to a

653 week from the time of imagery.

654

655 **Mount Adams**

656 No suitable NAIP imagery was found, instead 2019 Maxar imagery was used. In addition
657 to the Maxar imagery, a multidirectional hillshade and 3-m contour lines derived from a
658 2016 lidar DEM (Bard 2019) were used as a guide when delignating flow divides.

659 Occasionally, 2009 Google Earth imagery was also useful. Extensive snow covered the
660 mountain when the 2016 lidar was flown masking some of the glacier termini. However,
661 the DEM was helpful in correcting the imagery where poorly aligned with the terrain.

662

663 Multiple buried-ice features were identified near the terminus of several glaciers where
664 ice appeared to have decoupled from the main active glacier. Large areas below the
665 glaciers (Mazama, Adams, and Pinnacle) likely have debris-covered ice. We focused on
666 the features which were likely to contain ice based on meltwater streams exiting near the
667 features and hummocky terrain which appeared to indicate melt. Ground-based images
668 from taken between 2014 to 2018 helped decision-making. The images were particularly
669 helpful in identifying a debris-covered ice cliff at Adams Glacier.

670

671 **Mount Rainier**

672 In general, the 2019 NAIP and the Maxar (2018-09-25) were used for the outlines.

673 Although the GNIS includes the Nisqually Icefall as a separate feature, we included the
674 icefall as part of the Nisqually Glacier (Figure A4).

675

676 **Mount St. Helens**

677 We used a GIS layer of geological mapping units that included snow and ice from the
678 USGS (David Sherrod, USGS written communication, 2021) to help guide our search.

679 The Crater Glacier (INV_ID E562842N5115499) was heavily debris covered, and
680 obscured by shadow in some areas.

681

682 **Olympic Mountains**


683

684 A 2015 inventory of the region was compiled because more recent imagery (NAIP and
685 Maxar) were not useful due to seasonal snow. Our updated inventory differs from that
686 published in Fountain et al. (2017) in two ways. First, they outlined and grouped the
687 glaciers and perennial snowfields according to watershed rather than individual glacier.
688 Their goal was to estimate glacier change relative to a previous study by Spicer (1986)
689 and had to follow Spicer's approach. Second, all outlines were rechecked and compared
690 to SFI and the NLCD resulting in minor changes.

691

692

693



694

695 **Figure A4.** Image of the Nisqually Glacier and Icefall. The orange and red outlines are from the
 696 updated inventory and the blue outline is from the USGS mapping (Fountain et al., 2007)
 697 database. The base image is from the NAIP taken in 2019.

698

699

700 **Table A14.** List of NAIP imagery used for outlining glaciers and perennial snowfields in
 701 Washington. ‘Date’ is the start and end date for flights covering the glaciated portions of the
 702 NAIP image. In some cases, flights were completed in a single day. For 2006 the inspection date
 703 was used, since the start and end dates were not provided.

704

Region/Year/Filename	County	Date (Year-M-D)
Cascade –Northern		

2006		
ortho_1-1_1n_s_wa007_2006_3.sid	Chelan	2006-07-01
2015		
ortho_1-1_1n_s_wa007_2015_1.sid	Chelan	2015-07-06 to 2015-09-23
ortho_1-1_1n_s_wa033_2015_1.sid	King	2015-07-06 to 2015-09-27
ortho_1-1_1n_s_wa037_2015_1.sid	Kittitas	2015-07-06 to 2015-09-23
ortho_1-1_1n_s_wa047_2015_1.sid	Okanogan	2015-09-09 to 2015-09-11
ortho_1-1_1n_s_wa057_2015_1.sid	Skagit	2015-07-06 to 2015-09-29
ortho_1-1_1n_s_wa061_2015_1.sid	Snohomish	2015-07-06 to 2015-09-29
ortho_1-1_1n_s_wa073_2015_1.sid	Whatcom	2015-09-10 to 2015-09-26
2017		
ortho_1-1_1n_s_wa007_2017_1.sid	Chelan	2017-10-03 to 2017-10-24
ortho_1-1_1n_s_wa057_2017_1.sid	Skagit	2017-09-27 to 2017-10-05
ortho_1-1_1n_s_wa073_2017_1.sid	Whatcom	2017-09-27 to 2017-10-05
Cascade –Southern		
2015		
ortho_1-1_1n_s_wa041_2015_1.sid	Lewis	2015-07-15 to 2015-07-29
ortho_1-1_1n_s_wa053_2015_1.sid	Pierce	2015-07-29
ortho_1-1_1n_s_wa059_2015_1.sid	Skamania	2015-07-15 to 2015-09-12
ortho_1-1_1n_s_wa077_2015_1.sid	Yakima	2015-07-15 to 2015-07-29
2019		
ortho_1-1_hn_s_wa053_2019_1.sid	Pierce	2019-08-26
ortho_1-1_hn_s_wa059_2019_1.sid	Skamania	2019-08-06 to 2019-08-26
Olympic Mountains		
2015		
ortho_1-1_1n_s_wa009_2015_1.sid	Clallam	2015-07-28 to 2015-09-12
ortho_1-1_1n_s_wa031_2015_1.sid	Jefferson	2015-07-28 to 2015-09-12
ortho_1-1_1n_s_wa045_2015_1.sid	Mason	2015-07-28 to 2015-08-19

705
706
707
708
709

Table A15. List of dates of the Maxar imagery used for outlining glaciers and perennial snowfields in Washington.

Region/ Date (Year-M-D)

Cascade Range-Northern

2018-09-25

Cascade Range-Southern

2018-09-25

2019-08-31

Olympic Mountains

2015-08-17

2019-09-30

710
711
712

Table A16. List of U.S. Geological Survey digital elevation models used for outlining glaciers

713 and perennial snowfields in Washington. To access the data both the URL and specific identifier
714 are required.

Region	Date	Citation	URL www.sciencebase.gov/catalog/item/
Mt. Adams	2016	Bard (2019)	5bc623b9e4b0fc368ebbe99a
Mt. Baker	2015	Bard (2017a)	58518b0ee4b0f99207c4f12c
Glacier Peak	2014-15	Bard (2017b)	57bf299ee4b0f2f0ceb7534e

715

716

717 **A7 Wyoming**

718

719 **Wind River Range**

720

721 Tables A17 and A18 list the imagery used. The 2015 NAIP imagery had little snow in
722 contrast to 2019 imagery. Shadows are common in the 2015 imagery and can be very
723 dark. Occasionally the 2019 imagery was used to define the glacier-bedrock headwall
724 boundary. The 2019 Maxar imagery was essentially identical to the NAIP and because its
725 black and white not as useful. Imagery from 2017 and 2018 were a bit too snowy around
726 the glacier margin to be useful. The 2018-09-06 Maxar imagery covered the entire range,
727 with some clouds.


728 In the southern Wind River Range, a new snow dusting was often present, occasionally
729 making it difficult to outline snowfields and a few glaciers. Distinguishing seasonal snow
730 from perennial snow was a judgement call. If the snow was slightly discolored similar to
731 underlying rock/soil looking like the color was coming from underneath it was identified
732 as seasonal snow. Also, if many snow-free patches (a few square meters) pockmarked the
733 snow or if many rocks protruded through the snow, it was considered seasonal. A
734 perennial patch of snow appeared smooth and white, hiding underlying surface. Thin
735 snow cover on glacier ice appeared greyish in color and appeared smoother than the
736 surrounding ice-free landscape.

737 At Lower Fremont Glacier, a number of sizable ice patches appear down valley as if a
738 deposit of buried ice is present. However, there is no obvious connection to the glacier
739 itself.

740 The GNIS identified a single glacier as the Sacagawea Glacier, and two separate Fremont
741 Glaciers (Figure A5). By 2017 the single glacier had split into four glaciers. We chose to
742 label the largest glacier and the glacier to the southeast the Sacagawea Glacier. The other
743 two glaciers were labeled the Fremont Glaciers.

744

745



747

748 **Figure A5.** Image of Fremont Glaciers and Sacagawea Glacier showing the Sacagawea outline
 749 from the Fountain et al. (2017) database (blue), our updated Fremont Glaciers outlines (orange),
 750 and updated Sacagawea outlines (red). The base image is from the NAIP, taken in 2015.

751

752

753 **Table A17.** List of NAIP imagery used for outlining glaciers and perennial snowfields in
 754 Wyoming. ‘Date’ is the start and end date for flights covering the glaciated portions of the NAIP
 755 image. In some cases, flights were completed in a single day. For 2006 the inspection date was
 756 used, since the start and end dates were not provided.

757

Region/Year/Filename	County	Date (Year-M-D)
Absaroka Range		

2006	ortho_1-2_1n_s_wy029_2006_1.sid	Park	2006-09-02
2015	ortho_1-1_hn_s_wy013_2015_2.sid	Fremont	2015-09-09 to 2015-10-13
	ortho_1-1_hn_s_wy029_2015_2.sid	Park	2015-09-22 to 2015-10-13
Bighorn Mountains, WY			
2015	ortho_1-1_hn_s_wy019_2015_2.sid	Johnson	2015-09-12
Teton Range			
2006	ortho_1-1_1n_s_wy039_2006_1.sid	Teton	2006-09-02
2015	ortho_1-1_hn_s_wy035_2015_2.sid	Sublette	2015-09-09 to 2015-10-13
	ortho_1-1_hn_s_wy039_2015_2.sid	Teton	2015-09-12 to 2015-09-22
2019	ortho_1-1_hn_s_wy039_2019_1.sid	Teton	2019-07-20 to 2015-09-22
Wind River Range			
2006	ortho_1-1_1n_s_wy035_2006_1.sid	Sublette	2006-09-02
2015	ortho_1-1_hn_s_wy013_2015_2.sid	Fremont	2015-09-09 to 2015-10-13
	ortho_1-1_hn_s_wy035_2015_2.sid	Sublette	2015-09-09 to 2015-10-13
2019	ortho_1-1_hn_s_wy013_2019_1.sid	Fremont	2019-07-20 to 2019-08-27
	ortho_1-1_hn_s_wy035_2019_1.sid	Sublette	2019-08-15 to 2019-09-13

758

759

760 **Table A18.** List of dates of the Maxar imagery used for outlining glaciers and perennial
761 snowfields in Wyoming.

Region/ Date (Year-M-D)

Wind River Range

2018-09-06

762

763

764 **8. Author Contributions.**

765

766 Andrew G. Fountain was the principal investigator of the project, he wrote the proposal and
767 digitized glacier and snowfield outlines, analyzed the data, and led the writing of this report.
768 Bryce Glenn was the GIS expert responsible for the geographic format (e.g. projection,
769 attributes, database structure) and quality control. He digitized glacier and snowfield outlines,
770 analyzed the data, and helped write the report. Chris McNeil provided some of the imagery.
771

772 **9. Competing Interests.**

773

774 The authors declare that they have no conflict of interest.

775

776 **10. Acknowledgments.**

777

778 Hassan Basagic and Kristina Dick digitized the glaciers of Mt. Shasta and the North Cascade
779 National Park, respectively. We greatly appreciate their help and expertise. Nathan Walker of the
780 U.S. Forest Service provided a very helpful initial review of this report. We kindly acknowledge
781 the funding from the US. Forest Service. Any use of trade, firm or product names is for
782 descriptive purposes only and does not imply endorsement by the U.S. Government.

783 **11. References**

784

785 Alley, R. B., Cuffey, K. M., and Zoet, L. K.: Glacial erosion: status and outlook, *Ann. Glaciol.*, 60, 1–13,
786 <https://doi.org/10.1017/aog.2019.38>, 2019.

787 Andreassen, L. M., Nagy, T., Kjølmoen, B., and Leigh, J. R.: An inventory of Norway’s glaciers and ice-
788 marginal lakes from 2018–19 Sentinel-2 data, *J. Glaciol.*, 68, 1085–1106,
789 <https://doi.org/10.1017/jog.2022.20>, 2022.

790 Armstrong, R. L.: Mass balance history of Blue Glacier, Washington, USA, in: *Glacier Fluctuations and*
791 *Climatic Change*, edited by: Oerlemans, J., Springer Netherlands, 183–192, 1989.

792 Bard, J. A.: High-resolution digital elevation dataset for Glacier Peak and vicinity, Washington, based on
793 lidar surveys of August–November, 2014 and June, 2015, <https://doi.org/10.5066/F7H41PJJ>, 2017a.

794 Bard, J. A.: High-resolution digital elevation dataset for Mt Baker and vicinity, Washington, based on
795 lidar surveys of 2015, <https://doi.org/10.5066/F7WD3XR0>, 2017b.

796 Bard, J. A.: High-resolution digital elevation dataset for Mount Adams and vicinity, Washington, based
797 on lidar surveys of August–September, 2016, <https://doi.org/10.5066/P9Z1HF1K>, 2019.

798 Beason, S. R., Legg, N. T., Kenyon, T. R., Jost, R. P., and Kennard, P. M.: Forecasting and seismic detection
799 of debris flows in pro-glacial rivers at Mount Rainier National Park, Washington, USA, *Contain. Proc.*
800 *Seventh Int. Conf. Debris-Flow Hazards Mitig. Gold. Colo. USA June 10-13 2019*
801 [Httpshdlhandlenet11124173051](httpshdlhandlenet11124173051), <https://doi.org/10.25676/11124/173232>, 2018.

802 Benn, D. I. and Evans, D. J. A.: *Glaciers and glaciation*, 2nd ed., Hodder Education, London, 802 pp., 2010.

803 Bolch, T., Menounos, B., and Wheate, R.: Landsat-based inventory of glaciers in western Canada, 1985–
804 2005, *Remote Sens. Environ.*, 114, 127–137, <https://doi.org/10.1016/j.rse.2009.08.015>, 2010.

805 Bolch, T., Rohrbach, N., Kutuzov, S., Robson, B. A., and Osmonov, A.: Occurrence, evolution and ice
806 content of ice-debris complexes in the Ak-Shiirak, Central Tien Shan revealed by geophysical and
807 remotely-sensed investigations: Ice-debris complexes in Ak-Shiirak, *Earth Surf. Process. Landf.*, 44, 129–
808 143, <https://doi.org/10.1002/esp.4487>, 2019.

809 Bowerman, N. D. and Clark, D. H.: Holocene glaciation of the central Sierra Nevada, California, *Quat. Sci.*
810 *Rev.*, 30, 1067–1085, <https://doi.org/10.1016/j.quascirev.2010.10.014>, 2011.

811 Brardinoni, F., Scotti, R., Sailer, R., and Mair, V.: Evaluating sources of uncertainty and variability in rock
812 glacier inventories, *Earth Surf. Process. Landf.*, 44, 2450–2466, <https://doi.org/10.1002/esp.4674>, 2019.

813 Cadbury, S. L., Hannah, D. M., Milner, A. M., Pearson, C. P., and Brown, L. E.: Stream temperature
814 dynamics within a New Zealand glacierized river basin, *River Res. Appl.*, 24, 68–89,
815 <https://doi.org/10.1002/rra.1048>, 2008.

816 Chiarle, M., Iannotti, S., Mortara, G., and Deline, P.: Recent debris flow occurrences associated with
817 glaciers in the Alps, *Glob. Planet. Change*, 56, 123–136,
818 <https://doi.org/10.1016/j.gloplacha.2006.07.003>, 2007.

819 Cogley, J. G., Hock, R., Rasmussen, L. A., Arendt, A. A., Bauder, A., Braithwaite, R. J., Jansson, P., Kaser,
820 G., Möller, M., Nicholson, L. I., and Zemp, M.: Glossary of Glacier Mass Balance and Related Terms,
821 UNESCO-IHP, Paris, 2011.

822 Davis, P. T.: Holocene glacier fluctuations in the American Cordillera, *Quat. Sci. Rev.*, 7, 129–157, 1988.

823 Denton, G. H.: Conterminous US, Chapter 1, in: *Mountain glaciers of the Northern Hemisphere*, vol. 1,
824 edited by: Field, W. O., Corps of Engineers, US Army, Technical Information Analysis Center, Cold
825 Regions Research and Engineering Laboratory, 1975.

826 DeVisser, M. H. and Fountain, A. G.: A century of glacier change in the Wind River Range, WY,
827 *Geomorphology*, 232, 103–116, <https://doi.org/10.1016/j.geomorph.2014.10.017>, 2015.

828 Dewitz, J.: National Land Cover Database (NLCD) 2016 Products (ver. 2.0, July 2020),
829 <https://doi.org/10.5066/P96HHBIE>, 2019.

830 Dick, K.: *Glacier Change of the North Cascades, Washington 1900-2009*, M.S., Portland State University,
831 Portland, OR, 127 pp., 2013.

832 Dussailant, I., Berthier, E., Brun, F., Masiokas, M., Hugonnet, R., Favier, V., Rabatel, A., Pitte, P., and
833 Ruiz, L.: Two decades of glacier mass loss along the Andes, *Nat. Geosci.*, 12, 802–808,
834 <https://doi.org/10.1038/s41561-019-0432-5>, 2019.

835 Earl, L. and Gardner, A.: A satellite-derived glacier inventory for North Asia, *Ann. Glaciol.*, 57, 50–60,
836 <https://doi.org/10.3189/2016AoG71A008>, 2016.

837 Evans, I. S.: Local aspect asymmetry of mountain glaciation: A global survey of consistency of favoured
838 directions for glacier numbers and altitudes, *Geomorphology*, 73, 166–184,
839 <https://doi.org/10.1016/j.geomorph.2005.07.009>, 2006.

840 Fagre, D. B., McKeon, L. A., Dick, K. A., and Fountain, A. G.: Glacier margin time series (1966, 1998, 2005,
841 2015) of the named glaciers of Glacier National Park, MT, USA, <https://doi.org/10.5066/f7p26wb1>, 2017.

842 Fellman, J. B., Nagorski, S., Pyare, S., Vermilyea, A. W., Scott, D., and Hood, E.: Stream temperature
843 response to variable glacier coverage in coastal watersheds of Southeast Alaska, *Hydrol. Process.*, 28,
844 2062–2073, <https://doi.org/10.1002/hyp.9742>, 2014.

845 Fischer, A., Seiser, B., Stocker Waldhuber, M., Mitterer, C., and Abermann, J.: Tracing glacier changes in
846 Austria from the Little Ice Age to the present using a lidar-based high-resolution glacier inventory in
847 Austria, *The Cryosphere*, 9, 753–766, <https://doi.org/10.5194/tc-9-753-2015>, 2015.

848 Fountain, A. G. and Glenn, B.: Data From: Inventory of Glaciers and Perennial Snowfields of the
849 Coterminous USA" (2022), Portland State University, Portland, OR, 2022.

850 Fountain, A. G. and Tangborn, W. V.: The effect of glaciers on streamflow variations, *Water Resour. Res.*,
851 21, 579–586, 1985.

852 Fountain, A. G., Hoffman, M. J., Jackson, K., Basagic, H. J., Nylén, T. H., and Percy, D.: Digital outlines and
853 the topography of the Glaciers of the American West, US Geological Survey, 2007.

854 Fountain, A. G., Glenn, B., and Basagic, H. J.: The Geography of Glaciers and Perennial Snowfields in the
855 American West, *Arct. Antarct. Alp. Res.*, 49, 391–410, <https://doi.org/10.1657/AAAR0017-003>, 2017.

856 Garwood, J. M., Fountain, A. G., Lindke, K. T., van Hattem, M. G., and Basagic, H. J.: 20th Century Retreat
857 and Recent Drought Accelerated Extinction of Mountain Glaciers and Perennial Snowfields in the Trinity
858 Alps, California, *Northwest Sci.*, 94, 44, <https://doi.org/10.3955/046.094.0104>, 2020.

859 Gesch, D., Oimoen, M., Greenlee, S., Nelson, C., Steuck, M., and Tyler, D.: The national elevation dataset,
860 *Photogramm. Eng. Remote Sens.*, 68, 5–32, 2002.

861 Heard, J.: Late Pleistocene and Holocene Aged Glacial and Climatic Reconstructions in the Goat Rocks
862 Wilderness, Washington, United States, <https://doi.org/10.15760/etd.557>, 2000.

863 Hock, R., de Woul, M., Radić, V., and Dyurgerov, M.: Mountain glaciers and ice caps around Antarctica
864 make a large sea-level rise contribution: MOUNTAIN GLACIERS AND ICE CAPS, *Geophys. Res. Lett.*, 36,
865 n/a-n/a, <https://doi.org/10.1029/2008GL037020>, 2009.

866 Hoffman, M. J., Fountain, A. G., and Achuff, J. M.: 20th-century variations in area of cirque glaciers and
867 glacierets, Rocky Mountain National Park, Rocky Mountains, Colorado, USA, *Ann. Glaciol.*, 46, 349–354,
868 <https://doi.org/10.3189/172756407782871233>, 2007.

869 Huss, M. and Hock, R.: Global-scale hydrological response to future glacier mass loss, *Nat. Clim. Change*,
870 8, 135–140, <https://doi.org/10.1038/s41558-017-0049-x>, 2018.

871 Jin, S., Homer, C., Yang, L., Danielson, P., Dewitz, J., Li, C., Zhu, Z., Xian, G., and Howard, D.: Overall
872 Methodology Design for the United States National Land Cover Database 2016 Products, *Remote Sens.*,
873 11, 2971, <https://doi.org/10.3390/rs11242971>, 2019.

874 King, C.: On the discovery of actual glaciers on the Mountains of the Pacific Slope, *Am. J. Sci. Arts*, 1,
875 157–167, 1871.

876 Krimmel, R. M.: Glaciers of the conterminous United States, in: Satellite image atlas of glaciers of the
877 world: North America, edited by: Williams, Jr., R. S. and Ferrigno, J., US Geological Survey Professional
878 Paper, Washington, DC, J329–J381, 2002.

879 Leigh, J. R., Stokes, C. R., Carr, R. J., Evans, I. S., Andreassen, L. M., and Evans, D. J. A.: Identifying and
880 mapping very small (<0.5 km²) mountain glaciers on coarse to high-resolution imagery, *J. Glaciol.*, 65,
881 873–888, <https://doi.org/10.1017/jog.2019.50>, 2019.

882 Linsbauer, A., Paul, F., and Haerberli, W.: Modeling glacier thickness distribution and bed topography
883 over entire mountain ranges with GlabTop: Application of a fast and robust approach, *J. Geophys. Res.*,
884 117, <https://doi.org/10.1029/2011JF002313>, 2012.

885 Linsbauer, A., Huss, M., Hodel, E., Bauder, A., Fischer, M., Weidmann, Y., Bärtschi, H., and Schmassmann,
886 E.: The New Swiss Glacier Inventory SGI2016: From a Topographical to a Glaciological Dataset, *Front.*
887 *Earth Sci.*, 9, 2021.

888 Lu, Y., Zhang, Z., Kong, Y., and Hu, K.: Integration of optical, SAR and DEM data for automated detection
889 of debris-covered glaciers over the western Nyainqentanglha using a random forest classifier, *Cold Reg.*
890 *Sci. Technol.*, 193, 103421, <https://doi.org/10.1016/j.coldregions.2021.103421>, 2022.

891 Meier, M. F.: Distribution and variations of glaciers in the United States exclusive of Alaska, *Int. Assoc.*
892 *Sci. Hydrol.*, 54, 420–429, 1961.

893 Meier, M. F.: Contribution of small glaciers to global sea level, *Science*, 226, 1418–1421, 1984.

894 Mishra, A., Nainwal, H. C., Bolch, T., Shah, S. S., and Shankar, R.: Glacier inventory and glacier changes
895 (1994–2020) in the Upper Alaknanda Basin, Central Himalaya, *J. Glaciol.*, 69, 591–606,
896 <https://doi.org/10.1017/jog.2022.87>, 2023.

897 Moore, R. D., Fleming, S. W., Menounos, B., Wheate, R., Fountain, A., Stahl, K., Holm, K., and Jakob, M.:
898 Glacier change in western North America: influences on hydrology, geomorphic hazards and water
899 quality, *Hydrol. Process.*, 23, 42–61, <https://doi.org/10.1002/hyp.7162>, 2009.

900 NAIP: National agricultural imagery program (NAIP) Information Sheet, United States Department of
901 Agriculture, 2017.

902 O’Connor, J. E., Hardison, J. H., and Costa, J. E.: Debris flows from failures of Neoglacial-age moraine
903 dams in the Three Sisters and Mount Jefferson wilderness areas, Oregon: a study of the recent debris
904 flows from moraine-dammed lake releases at central Oregon Cascade Range volcanoes - Mt. Jefferson,
905 Three Fingered Jack and the Three Sisters / Broken Top, US Geological Survey, Reston, VA, 93 pp., 2001.

906 Osborn, G., Menounos, B., Ryane, C., Riedel, J. L., Clague, J. J., Koch, J., Clark, D., Scott, K., and Davis, P.
907 T.: Latest Pleistocene and Holocene glacier fluctuations on Mount Baker, Washington, *Quat. Sci. Rev.*,
908 49, 33–51, <https://doi.org/10.1016/j.quascirev.2012.06.004>, 2012.

909 Parkes, D. and Marzeion, B.: Twentieth-century contribution to sea-level rise from uncharted glaciers,
910 *Nature*, 563, 551–554, <https://doi.org/10.1038/s41586-018-0687-9>, 2018.

911 Paul, F., Kääb, A., and Haeberli, W.: Recent glacier changes in the Alps observed by satellite:
912 Consequences for future monitoring strategies, *Glob. Planet. Change*, 56, 111–122,
913 <https://doi.org/10.1016/j.gloplacha.2006.07.007>, 2007.

914 Paul, F., Barrand, N. E., Baumann, S., Berthier, E., Bolch, T., Casey, K., Frey, H., Joshi, S. P., Konovalov, V.,
915 Le Bris, R., Mölg, N., Nosenko, G., Nuth, C., Pope, A., Racoviteanu, A., Rastner, P., Raup, B., Scharrer, K.,
916 Steffen, S., and Winsvold, S.: On the accuracy of glacier outlines derived from remote-sensing data, *Ann.*
917 *Glaciol.*, 54, 171–182, <https://doi.org/10.3189/2013AoG63A296>, 2013.

918 Paul, F., Rastner, P., Azzoni, R. S., Diolaiuti, G., Fugazza, D., Le Bris, R., Nemeč, J., Rabatel, A., Ramusovic,
919 M., Schwaizer, G., and Smiraglia, C.: Glacier shrinkage in the Alps continues unabated as revealed by a
920 new glacier inventory from Sentinel-2, *Earth Syst. Sci. Data*, 12, 1805–1821,
921 <https://doi.org/10.5194/essd-12-1805-2020>, 2020.

922 Pfeffer, W. T., Arendt, A. A., Bliss, A., Bolch, T., Cogley, J. G., Gardner, A. S., Hagen, J.-O., Hock, R., Kaser,
923 G., and Kienholz, C.: The Randolph Glacier Inventory: a globally complete inventory of glaciers, *J. Glaciol.*,
924 60, 537–552, <https://doi.org/10.3189/2014JoG13J176>, 2014.

925 Post, A., Richardson, D., Tangborn, W. V., and Rosselot, F.: Inventory of glaciers in the North Cascades,
926 Washington. (USGS Professional Paper 705-A)., US Geological Survey, 705-A, Washington, DC, 1971.

927 Pritchard, H. D.: Asia’s shrinking glaciers protect large populations from drought stress, *Nature*, 569,
928 649–654, <https://doi.org/10.1038/s41586-019-1240-1>, 2019.

929 Rabatel, A., Sirguey, P., Drolon, V., Maisongrande, P., Arnaud, Y., Berthier, E., Davaze, L., Dedieu, J.-P.,
930 and Dumont, M.: Annual and Seasonal Glacier-Wide Surface Mass Balance Quantified from Changes in
931 Glacier Surface State: A Review on Existing Methods Using Optical Satellite Imagery, *Remote Sens.*, 9,
932 507, <https://doi.org/10.3390/rs9050507>, 2017.

933 Rasmussen, L. A.: South Cascade Glacier mass balance, 1935–2006, *Ann. Glaciol.*, 50, 215–220, 2009.

934 RGIK: Towards standard guidelines for inventorying rock glaciers: baseline concepts (version 4.2.2), . IPA
935 Action Group Rock glacier inventories and kinematics, 2022.

936 Robinson, J. E.: Digital topographic data based on lidar survey of Mount Shasta Volcano, California, July-
937 September 2010, Reston, VA, <https://doi.org/10.3133/ds852>, 2014.

938 Robson, B. A., Bolch, T., MacDonell, S., Hölbling, D., Rastner, P., and Schaffer, N.: Automated detection
939 of rock glaciers using deep learning and object-based image analysis, *Remote Sens. Environ.*, 250,
940 112033, <https://doi.org/10.1016/j.rse.2020.112033>, 2020.

941 Russell, I. C.: The glaciers of North America, *Geogr. J.*, 12, 553–564, 1898.

942 Schiefer, E., Menounos, B., and Wheate, R.: Recent volume loss of British Columbian glaciers, Canada:
943 Volume loss of BC glaciers, *Geophys. Res. Lett.*, 34, L16503, <https://doi.org/10.1029/2007GL030780>,
944 2007.

- 945 Selkowitz, D. J. and Forster, R. R.: Automated mapping of persistent ice and snow cover across the
946 western U.S. with Landsat, *ISPRS J. Photogramm. Remote Sens.*, 117, 126–140,
947 <https://doi.org/10.1016/j.isprsjrs.2016.04.001>, 2016.
- 948 Sitts, D. J., Fountain, A. G., and Hoffman, M. J.: Twentieth Century Glacier Change on Mount Adams,
949 Washington, USA, *Northwest Sci.*, 84, 378–385, <https://doi.org/10.3955/046.084.0407>, 2010.
- 950 Smiraglia, C., Azzoni, R. S., D’Agata, C., Maragno, D., Fugazza, D., and Diolaiuti, G. A.: The New Italian
951 Glacier Inventory: a didactic tool for a better knowledge of the natural Alpine environment, *J-Read.-J.*
952 *Res. Didact. Geogr.*, 4, 2015.
- 953 Spicer, R.: *Glaciers in the Olympic Mountains, Washington -- Present distribution and recent variations.*,
954 M.S., University of Washington, Seattle, WA, 158 pp., 1986.
- 955 Sun, M., Liu, S., Yao, X., Guo, W., and Xu, J.: Glacier changes in the Qilian Mountains in the past half-
956 century: Based on the revised First and Second Chinese Glacier Inventory, *J. Geogr. Sci.*, 28, 206–220,
957 <https://doi.org/10.1007/s11442-018-1468-y>, 2018.
- 958 Trcka, A.: *Inventory of Rock Glaciers in the American West and Their Topography and Climate*,
959 <https://doi.org/10.15760/etd.7509>, 2020.
- 960 Usery, E. L., Varanka, D., and Finn, M. P.: A 125 year history of topographic mapping and GIS in the U.S.
961 Geological Survey 1884-2009, part 1: 1884-1980, *ArcNews*, 31, 1–1, 2009.
- 962 Wickham, J., Stehman, S. V., Sorenson, D. G., Gass, L., and Dewitz, J. A.: Thematic accuracy assessment
963 of the NLCD 2016 land cover for the conterminous United States, *Remote Sens. Environ.*, 257, 112357,
964 <https://doi.org/10.1016/j.rse.2021.112357>, 2021.
- 965 Yao, T., Pu, J., Lu, A., Wang, Y., and Yu, W.: Recent glacial retreat and its impact on hydrological
966 processes on the Tibetan Plateau, China, and surrounding regions, *Arct. Antarct. Alp. Res.*, 39, 642–650,
967 2007.
- 968 Zalazar, L., Ferri, L., Castro, M., Gargantini, H., Gimenez, M., Pitte, P., Ruiz, L., Masiokas, M., Costa, G.,
969 and Villalba, R.: Spatial distribution and characteristics of Andean ice masses in Argentina: results from
970 the first National Glacier Inventory, *J. Glaciol.*, 66, 938–949, <https://doi.org/10.1017/jog.2020.55>, 2020.
- 971 Zemp, M., Huss, M., Thibert, E., Eckert, N., McNabb, R., Huber, J., Barandun, M., Machguth, H.,
972 Nussbaumer, S. U., Gärtner-Roer, I., Thomson, L., Paul, F., Maussion, F., Kutuzov, S., and Cogley, J. G.:
973 Global glacier mass changes and their contributions to sea-level rise from 1961 to 2016, *Nature*, 568,
974 382–386, <https://doi.org/10.1038/s41586-019-1071-0>, 2019.
- 975