



1 **Inventory of glaciers and perennial snowfields of the coterminous USA**

2

3 Andrew G. Fountain¹, Bryce Glenn¹, Christopher Mcneil²

4 ¹Department of Geology, Portland State University, 1721 SW Broadway, Portland, OR. 97212
5 USA, ²U.S. Geological Survey Alaska Science Center, Anchorage, AK, USA

6 *Correspondence to:* Andrew G. Fountain (andrew@pdx.edu)

7 **Abstract**

8 This report summarizes an updated inventory of glaciers and perennial snowfields of the
9 contiguous United States. The inventory is based on interpretation of mostly aerial imagery
10 provided by the National Agricultural Imagery Program, U.S. Department of Agriculture with
11 some satellite imagery in places where aerial imagery was not suitable. The inventory includes
12 all perennial snow and ice features greater than 0.01 km². Due to aerial survey schedules and
13 seasonal snow cover, imagery acquired over a number of years were required. The earliest date is
14 2013 and the latest is 2020, but more than 73% of the outlines were acquired from 2015 imagery.
15 The inventory is compiled as shapefiles within a geographic information system that includes
16 feature classification, area, and location. The inventory identified 1331 (366.52 km²) glaciers,
17 1776 (31.00 km²) perennial snowfields, and 35 (3.57 km²) buried-ice features. The data
18 including both the shapefiles and tabulated results are publicly available at
19 <https://doi.org/10.15760/geology-data.03> (Fountain & Glenn, 2022).

20 **1. Introduction**

21 Glaciers are an important feature of the landscape for several reasons. Geologically, they modify
22 the landscape through erosion and deposition (Alley et al., 2019; Benn & Evans, 2010).
23 Although these processes are typically slow, sudden episodes can occur such as moraine failure
24 due to fluvial erosion resulting in catastrophic debris flows (Beason et al., 2018; Chiarle et al.,
25 2007; O'Connor et al., 2001). Hydrologically, glaciers can be viewed as frozen reservoirs of
26 water that naturally regulate streamflow on seasonal to decadal time scales (Dussaillant et al.,
27 2019; Fountain & Tangborn, 1985; Moore et al., 2009). Glacial runoff increases during warm
28 periods and diminishes during cool, wet periods. Thus, glacial watersheds have less seasonally
29 variable runoff than ice-free watersheds. Also, glacial runoff cools stream temperatures in the
30 driest and hottest part of the summer after seasonal snowpacks have vanished (Cadbury et al.,
31 2008; Fellman et al., 2014). As glaciers shrink, they have less ability to buffer seasonal runoff
32 variations and watersheds become more susceptible to drought. Globally, the loss of perennial
33 ice from the landscape is a major contributor to sea level rise (Meier, 1984; Parkes & Marzeion,
34 2018; Zemp et al., 2019).

35 Glacier inventories have been valuable for assessing glacier contribution to sea level change
36 (Hock et al., 2009; Pfeffer et al., 2014), and for assessing regional hydrology (Yao et al., 2007;
37 Moore et al., 2009). They also provide a baseline for quantifying future glacier changes. Glacier
38 inventories have been compiled for many regions of the world (Bolch et al., 2010; Smiraglia et
39 al., 2015; Sun et al., 2018). An exception has been western United States (US), defined here as



40 those conterminous states west of the 100th meridian. Despite a vigorous history of glacier
41 studies (e.g., Armstrong, 1989; Rasmussen, 2009), glacial geology (e.g. Davis, 1988; Bowerman
42 and Clark, 2011; Osborn et al., 2012), and regional inventories (e.g. DeVisser & Fountain, 2015;
43 Fagre et al., 2017; Post et al., 1971) the glacier cover for the entire western US has not been
44 updated in several decades.

45 The earliest scientific identification of glacier-populated regions in the western US date to King
46 (1871) and, more comprehensively, to Russell (1898). The first summary of glacier areas for
47 each state was in 1961 (Meier, 1961). However, the data sources and methods used to compile
48 the inventories are unknown. Denton (1975) summarized all known glacier studies in the western
49 US, but did not tabulate glacier areas. Krimmel (2002) updated Meier's study and provided total
50 glacier area for the various mountain ranges by summarizing a variety previous studies published
51 over a 10+ year time span. It is not clear whether the inventory is complete and no data on
52 individual glaciers are provided. Fountain et al. (2007, 2017) compiled the first comprehensive
53 inventory of glaciers in the western US. The data were derived from historical U.S. Geological
54 Survey (USGS) 1:24,000 scale maps compiled over a 40-year period from the 1940s to the 1980s
55 (Gesch et al., 2002; Usery et al., 2009). Because the USGS mapping was based on one-time
56 aerial imagery, the misinterpretation of seasonal snow as perennial was extensive in some
57 regions. The most current study, Selkowitz & Forster (2016), used Landsat satellite imagery
58 compiled over a four-year period, 2010-2014, and an automated detection scheme to define
59 perennial snow and ice. However, automated schemes are known to misclassify debris-covered
60 ice as ice-free landscape underestimating glacier area (Earl & Gardner, 2016; Paul et al., 2007;
61 Rabatel et al., 2017).

62 This report presents the results of an updated and comprehensive inventory of glaciers and
63 perennial snowfields of the western US for the purpose of defining their current extent and to
64 provide of baseline for estimating future changes. The report summarizes our methods,
65 uncertainties, tabulated results, and data availability.

66 **2. Methods**

67
68 The location of the glaciers and perennial snowfields were initially identified by a geographic
69 information system (GIS) database from Fountain et al. (2007). New outlines were manually
70 digitized from color digital orthographic aerial photographs available from the National
71 Agricultural Imagery Program (NAIP), U.S. Department of Agriculture, Farm Service Agency
72 program (NAIP, 2017). Since 2009, the imagery is collected on cycles of two to three years. The
73 spatial resolution is at least 1 m (ground sampling distance) with a horizontal accuracy of 6 m of
74 photo-identifiable ground control points (USDA, 2021). NAIP imagery was downloaded from
75 Data Gateway (https://datagateway.nrcs.usda.gov/GDGHome_DirectDownload.aspx). In a few
76 cases, NAIP imagery was not suitable due to seasonal snow, deep shadows, or image warping
77 caused by orthophoto rectification, therefore other sources were used including Maxar satellite
78 imagery (Maxar Technologies, Inc) with a spatial resolution of 0.5 – 1 m. In one situation, we
79 used the most recent imagery available in Google Earth (Google, Inc), resolution ~ 1m, because
80 no other imagery was suitable.



81 We manually identified all glaciers, ice patches, and perennial snowfields equal or larger than
82 0.01 km². Glaciers are defined as perennial snow and ice that moves (Cogley et al., 2011). A
83 feature was considered perennial if it was present on the original 1:24,000 USGS topographic
84 maps and present on all Google Earth imagery. Movement was identified by the presence of
85 crevasses. Perennial snowfields and ice patches do not exhibit movement, as indicated by a lack
86 of crevasses observed in the imagery. We do not distinguish between snowfields and ice patches
87 and refer to both as perennial snowfields.

88 Contiguous glacier cover, most commonly on volcanoes, was separated into individual glaciers if
89 they had unique names as indicated on the USGS maps. The orientation of crevasse patterns was
90 used to define flow divides. In the absence of these patterns, shaded relief digital elevation
91 models were used to examine slope changes. These models were derived from aerial lidar data,
92 flown under contract to the USGS (Bard, 2017b, 2017a, 2019; Robinson, 2014) or the Oregon
93 Department of Geology and Mineral Industries (DOGAMI, 2011).

94 We encountered a number of challenges to our classification and delineation of the glaciers and
95 perennial snowfields. Although crevasses were used to define movement, in a few cases it
96 appeared that they penetrated through the feature to the bedrock underneath suggesting a
97 mechanical break up. In these cases, the feature was classified as a snowfield. In the high alpine
98 regions of California, Colorado, and Wyoming, the terminus of some glaciers was hard to define.
99 Rather than abruptly terminating, the ice seems to thin and smoothly transitions into the
100 surrounding rock talus (see Wyoming in the appendix). It was unclear whether a thin debris layer
101 blanketed the ice or cobbles and boulders protruding through the thin ice. The boundary was
102 mapped along the edge of identifiable ice.

103 Although not a common problem, one particular difficulty was distinguishing glaciers from rock
104 glaciers (Brardinoni et al., 2019). A rock glacier is a mass of rock debris in a matrix of ice that
105 flows (Cogley et al., 2011). They can be difficult to distinguish from a debris-covered glacier,
106 one that has extensive rock debris over the ablation zone, that lower part of a glacier with
107 exposed ice in late summer. We adopted the following topographic classification. If the slope of
108 the apparent glacier/snowfield graded into the slope of the rock glacier, with no change in sign,
109 then we considered it part of the rock glacier. On the other hand, if the slope changed sign at the
110 bottom of the apparent glacier/snowfield, such that the topography formed a dip before reaching
111 another topographic high that marks the start of a rock glacier, then it was considered
112 independent feature (see Colorado in the appendix).

113 In a number of cases we observed buried ice adjacent to a glacier (see Oregon in the appendix).
114 The texture was hummocky and very different from surrounding bedrock and adjacent ice.
115 Occasionally a crack in the surface revealed subsurface ice. We decided to include these features
116 as a separate classification, 'buried ice', because their size was large relative to the glacier, they
117 were probably once part of the glacier, and may be important local sources of meltwater for
118 streamflow.

119
120 The glaciers and perennial snowfields outlines were digitized using ArcMap (ESRI, Inc), a
121 geographic information system, at scales varying from 1:300 to 1:2000 depending on image



122 quality and complexity. The projection used was the native projection of the image, North
123 American Datum of 1983 (NAD83) for NAIP, and World Geodetic System 1984 (WGS84), with
124 the relevant local Universal Transverse Mercator (UTM) zone, for Maxar and Google Earth.
125 When Maxar or Google Earth imagery were used, final outlines were projected into the NAD83
126 coordinate system. In situations where it was hard to interpret feature geometry, Google Earth
127 was very helpful because its terrain feature provides an oblique perspective that can be tilted and
128 rotated. Each outline was checked independently by the two senior authors of this report, and in
129 some cases by a third collaborator. If an outline was revised, then it was returned to its original
130 author for review and correction, and the process iterated until all parties agreed.

131

132 **Uncertainty** The uncertainty in glacier area was calculated as one-half the absolute difference
133 between initial and final revised digitized area (the range) divided by the final area and expressed
134 as a percentage. For some glaciers where no revision was necessary a 1% uncertainty was
135 assigned to account for digitizing error, which is known to be relatively small (DeVisser &
136 Fountain, 2015; Hoffman et al., 2007). For the relatively few glaciers where a small section of
137 perimeter was masked by deep shadow, seasonal snow patches, rock debris, or poor imagery, a
138 higher uncertainty was assigned by visually comparing the area in question to the total possible
139 area of the glacier. The estimated uncertainty, up to 16%, was determined by digitizing the
140 minimum and maximum perimeters from a sample of glaciers with similar issues. Uncertainty
141 about the position of a flow divide was considered 5%, due to the topographic ambiguity along a
142 divide, and estimated from several digitizing efforts. For perennial snowfields a 30% uncertainty is
143 assigned because the seasonal snow commonly covers the smaller patch of perennial snow and
144 the seasonal snow varies greatly from year to year. The snowfield uncertainty was arbitrary in
145 order to note their presence and location, but preclude them from area change calculations
146 because area differences are typically smaller than the assigned uncertainty.

147

148 Our initial inventory was then compared sequentially to two other independent inventories to test
149 for errors of omission or commission. The first comparison was to the Selkowitz and Forster
150 (2016) inventory (SFI). However, we had to reconcile the different methods of each inventory
151 prior to comparison. Buried-ice features were eliminated from our inventory because the SFI did
152 not map buried ice. Features removed from the SFI include, features $< 0.01 \text{ km}^2$ to match our
153 minimum area threshold; a small number of glaciers and snowfields located in Canada; the few
154 glacier classifications of ponds, lakes and dry lakebeds. Notably, the SFI did not split contiguous
155 ice masses, such as glacier-covered volcanoes, into individual glaciers, consequently we do not
156 expect the number of features in the SFI and our inventory to match. Once the two inventories
157 were reconciled, those glaciers and perennial snowfields unique to one inventory were examined
158 for inclusion in a revised inventory. Features selected from the SFI were digitized using the same
159 imagery we used for our inventory.

160

161 The revised inventory was then compared to the 2016 National Land Cover Database (NLCD),
162 which did not map glaciers and perennial snowfields per se, but mapped the distribution of
163 perennial snow and ice (Jin al., 2019). However, the NLCD used a smaller number of images
164 over time for any one location such that we consider the assessment of ‘perennial’ has a high
165 uncertainty. Also, the landscape class of snow and ice received less attention than other classes
166 (e.g. agriculture) such that the timing of imagery acquisition may be earlier in the summer than



167 optimal and misclassification of clouds as snow and ice may be present (personal communication
168 C. Homer and J. Dewitz, 2015). Again, the features unique to one inventory were examined for
169 inclusion and those features selected from the NLCD were digitized using the same imagery we
170 used for our inventory.


171

172 **3. Results**

173

174 Our initial inventory identified 2267 glaciers and perennial snowfields totaling 391.95 km².
175 About 70% (1576) overlapped the features in the SFI. After examining all features unique to
176 each inventory, we revised our inventory to include 2373 (394.99 km²) glaciers and perennial
177 snowfields. Comparing the revised inventory to the 2016 NLCD resulted in adding another 134
178 (2.53 km²) features, which included 12 (0.38 km²) glaciers. The final inventory includes 2542
179 glacial features composed of 1331 (366.52 km²) glaciers, 1176 (31.01 km²) perennial snowfields,
180 and 35 (3.57 km²) buried ice deposits (Table 1; Figure 1). Most glaciers and perennial
181 snowfields, 1554 (62%) were outlined using 2015 NAIP imagery with the remainder outlined
182 using mostly NAIP imagery from 2013 to 2020. The state of Washington has the greatest number
183 and area of glaciers, perennial snowfields, and buried ice.

184



185
 186
 187
 188
 189

Figure 1. The spatial distribution and number of glaciers and perennial snowfields, greater than 0.01 km², in the western US. Colors indicate the date of aerial and satellite imagery used to outline the features. The line is the cumulative total. Base imagery from Esri Inc.



190 **Table 1.** The summary of the glacial inventory for the American West, exclusive of Alaska.
 191 Number is the total number of features within each classification (Class), ‘Uncert.’ is the area
 192 uncertainty, and ‘Max Area’ is maximum area. Note that the uncertainty of ‘Buried ice’ is
 193 unknown.
 194

State/Region/Class	Number	Total Area km²	Total Uncert km²	Max Area km²	Mean Area km²
California	132	10.63	0.61	1.45	0.08
Cascade Range	39	5.74	0.37	1.45	0.15
Buried ice	5	0.44	-----	0.16	0.09
Glaciers	10	4.61	0.17	1.45	0.46
Perennial snowfields	24	0.68	0.21	0.08	0.03
Sierra Nevada	91	4.86	0.23	0.66	0.05
Buried ice	2	0.13	0.00	0.10	0.06
Glaciers	64	4.37	0.12	0.66	0.07
Perennial snowfields	25	0.37	0.11	0.03	0.01
Trinity Alps	2	0.03	0.00	0.02	0.02
Glaciers	2	0.03	0.00	0.02	0.02
Colorado	84	2.20	0.46	0.16	0.03
Elk Mountains	5	0.09	0.03	0.03	0.02
Glaciers	1	0.01	0.00	0.01	0.01
Perennial snowfields	4	0.08	0.02	0.03	0.02
Front Range	58	1.73	0.33	0.16	0.03
Glaciers	13	0.74	0.03	0.16	0.06
Perennial snowfields	45	0.99	0.30	0.09	0.02
Gore Range	7	0.11	0.03	0.02	0.02
Glaciers	1	0.02	0.00	0.02	0.02
Perennial snowfields	6	0.09	0.03	0.02	0.02
Medicine Bow Mountains	1	0.04	0.01	0.04	0.04
Perennial snowfields	1	0.04	0.01	0.04	0.04
Park Range	6	0.11	0.03	0.03	0.02
Perennial snowfields	6	0.11	0.03	0.03	0.02
San Miguel Mountains	5	0.07	0.02	0.02	0.01
Perennial snowfields	5	0.07	0.02	0.02	0.01
Sawatch Range	2	0.04	0.01	0.03	0.02
Perennial snowfields	2	0.04	0.01	0.03	0.02
Idaho	6	0.08	0.02	0.02	0.01
Sawtooth Range	6	0.08	0.02	0.02	0.01
Perennial snowfields	6	0.08	0.02	0.02	0.01
Montana	416	30.26	2.27	1.45	0.07
Beartooth -Absaroka	111	6.07	0.64	0.45	0.05
Buried ice	1	0.04	-----	0.04	0.04
Glaciers	50	4.31	0.12	0.45	0.09
Perennial snowfields	60	1.72	0.52	0.22	0.03



Bitterroot Range	4	0.08	0.02	0.03	0.02
Glaciers	1	0.03	0.00	0.03	0.03
Perennial snowfields	3	0.05	0.02	0.02	0.02
Cabinet Mountains	9	0.25	0.08	0.08	0.03
Perennial snowfields	9	0.25	0.08	0.08	0.03
Crazy Mountains	13	0.27	0.06	0.04	0.02
Glaciers	3	0.06	0.00	0.04	0.02
Perennial snowfields	10	0.21	0.06	0.04	0.02
Lewis Range	230	21.38	1.15	1.45	0.09
Glaciers	145	19.22	0.50	1.45	0.13
Perennial snowfields	85	2.16	0.65	0.09	0.03
Mission-Swan-Flathead	49	2.20	0.34	0.22	0.04
Glaciers	11	1.16	0.02	0.22	0.11
Perennial snowfields	38	1.04	0.31	0.09	0.03
Oregon	116	15.38	1.62	1.16	0.13
Cascade Range	110	15.24	1.58	1.16	0.14
Buried ice	7	1.25	-----	0.45	0.18
Glaciers	42	11.90	0.95	1.16	0.28
Perennial snowfields	61	2.09	0.63	0.15	0.03
Wallowa Mountains	6	0.14	0.04	0.04	0.02
Perennial snowfields	6	0.14	0.04	0.04	0.02
Washington	1481	312.26	16.33	11.24	0.21
Cascade Range-Northern	1126	186.58	9.64	6.06	0.17
Buried ice	10	0.50	-----	0.15	0.05
Glaciers	706	176.27	6.70	6.06	0.25
Perennial snowfields	410	9.80	2.94	0.16	0.02
Cascade Range-Southern	219	101.66	5.86	11.24	0.46
Buried ice	10	1.20	-----	0.30	0.12
Glaciers	69	95.64	4.42	11.24	1.39
Perennial snowfields	140	4.82	1.45	0.33	0.03
Olympic Mountains	136	24.02	0.82	5.09	0.18
Glacier	106	23.44	0.65	5.09	0.22
Perennial snowfield	30	0.57	0.17	0.06	0.02
Wyoming	307	30.29	2.34	2.32	0.10
Absaroka Range	62	1.44	0.33	0.12	0.02
Glacier	10	0.48	0.05	0.12	0.05
Perennial snowfield	52	0.96	0.29	0.05	0.02
Bighorn Mountains	8	0.42	0.03	0.22	0.05
Glacier	3	0.34	0.01	0.22	0.11
Perennial snowfield	5	0.08	0.02	0.03	0.02
Teton Range	49	2.04	0.21	0.23	0.04
Glacier	20	1.46	0.03	0.23	0.07
Perennial snowfield	29	0.59	0.18	0.05	0.02
Wind River Range	188	26.39	1.76	2.32	0.14
Glacier	74	22.42	0.57	2.32	0.30



Perennial snowfield	114	3.97	1.19	0.26	0.03
Grand Total	2542	401.10	23.64	11.24	0.16

195
 196
 197
 198
 199
 200
 201
 202
 203
 204
 205
 206
 207
 208

The final inventory conflicts with the current database of the Geographic Names Information System (US Geological Survey, 2022). The inventory excludes 52 officially named glaciers because 2 have disappeared, 25 were classified as perennial snowfields, the area of 18 was less than 0.01 km², and 7 were considered rock glaciers (Table 2). In some cases, a named glacier or snowfield had split into multiple pieces since the original USGS mapping; all pieces were assigned the same name in the inventory (see appendix, sections 6.7 and 6.8).

Table 2. List of officially named glaciers not classified as glaciers and excluded from the final inventory. Names come from the Geographic Names Information System, (US Geological Survey, 2022). The 'Reason' column lists why the named glacier is no longer considered a glacier in our inventory.

State/Region/Glacier Name	Reason
California	
Sierra Nevada	
Matthes Glaciers	rock glacier
Mount Warlow Glacier	rock glacier
Powell Glacier	rock glacier
Colorado	
Front Range	
Isabelle Glacier	perennial snowfield
Mills Glacier	perennial snowfield
Moomaw Glacier	perennial snowfield
Peck Glacier	perennial snowfield
Rowe Glacier	< 0.01 km ²
Saint Marys Glacier	< 0.01 km ²
Taylor Glacier	rock glacier
The Dove	< 0.01 km ²
Idaho	
Lost River Range	
Borah Glacier	rock glacier
Montana	
Beartooth Mountains-Absaroka Range	
Grasshopper Glacier	rock glacier
Cabinet Mountains	
Blackwell Glacier	perennial snowfield
Crazy Mountains	
Grasshopper Glacier	rock glacier
Lewis Range	
Boulder Glacier	perennial snowfield



Mission-Swan-Flathead Ranges	
Fissure Glacier	< 0.01 km ²
Gray Wolf Glacier	perennial snowfield
<hr/>	
Oregon	
Cascade Range	
Carver Glacier	perennial snowfield
Clark Glacier	perennial snowfield
Irving Glacier	perennial snowfield
Lathrop Glacier	< 0.01 km ²
Palmer Glacier	perennial snowfield
Skinner Glacier	perennial snowfield
Thayer Glacier	< 0.01 km ²
Wallowa Mountains	
Benson Glacier	perennial snowfield
<hr/>	
Washington	
Cascade Range-Northern	
Lyall Glacier	perennial snowfield
Milk Lake Glacier	disappeared
Snow Creek Glacier	perennial snowfield
Spider Glacier	perennial snowfield
Table Mountain Glacier	< 0.01 km ²
Cascade Range-Southern	
Ape Glacier	< 0.01 km ²
Dryer Glacier	perennial snowfield
Forsyth Glacier	< 0.01 km ²
Meade Glacier	perennial snowfield
Nelson Glacier	< 0.01 km ²
Packwood Glacier	perennial snowfield
Pinnacle Glacier	< 0.01 km ²
Pyramid Glaciers	< 0.01 km ²
Shoestring Glacier	< 0.01 km ²
Stevens Glacier	perennial snowfield
Talus Glacier	perennial snowfield
Unicorn Glacier	< 0.01 km ²
Williwakas Glacier	perennial snowfield
Olympic Mountains	
Anderson Glacier	perennial snowfield
Lillian Glacier	< 0.01 km ²
<hr/>	
Wyoming	
Absaroka Range	
DuNoir Glacier	< 0.01 km ²
Teton Range	
Petersen Glacier	< 0.01 km ²
Teepe Glacier	perennial snowfield
Wind River Range	



Hooker Glacier	disappeared
Harrower Glacier	perennial snowfield
Tiny Glacier	< 0.01 km ²

209

210

211 **4. Data products and availability**

212

213 The data are available in three formats. The geospatial data and attribute tables are available in
214 the shapefile (Esri) format and in an open source GeoJSON format. The attribute table is also
215 available as an EXCEL file. These data products can be obtained from
216 <https://doi.org/10.15760/geology-data.03> (Fountain and Glenn, 2022) and from the Global Land
217 Ice Measurements from Space website (to be submitted) <http://glims.colorado.edu/glacierdata/>

218

219 **5. Summary**

220

221 We have compiled a new and comprehensive inventory of glaciers and perennial snowfields in
222 the western US from aerial and satellite imagery. Results show that 2542 glacial features are
223 currently present and include 1331 (366.52 km²) glaciers, 1176 (31.01 km²) perennial
224 snowfields, and 35 (3.57 km²) buried ice deposits. Most of the data were acquired from 2015
225 NAIP imagery with the remainder from NAIP imagery and a few satellite images acquired over
226 the period of 2013 to 2020. The state of Washington has the greatest number and area of glaciers
227 and perennial snowfields. This product updates an older inventory based on USGS 1:24000 maps
228 compiled in the middle-late 1900's. The new inventory is a significant improvement in accuracy
229 because the archive of historical imagery in Google Earth allowed us to classify glaciers versus
230 perennial snowfields. Finally, this new inventory provides a baseline for assessing glacier change
231 in the coterminous US.

232

233 **6. Appendix**

234

235 This appendix, organized by US State, then by mountain range, summarizes the specific
236 imagery used, challenges encountered in feature identification and outline digitization. The
237 most recent suitable NAIP was used in each case. Where such imagery was not suitable
238 Maxar imagery was used. In the Wallowa Mountains, Oregon, neither was NAIP suitable nor
239 was Maxar available so images from Google Earth were used. The Selkowitz and Forster
240 (2016) inventory is referred to as the SFI and the National Land Cover Database inventory
241 (Jin et al., 2019) is referred to as the NLCD.

242

243 **California**

244

245 Imagery and DEMs used are listed in Tables A1, A2, A3.

246

247 **Cascade Range**


248

249 **Mount Shasta**

250



251 The 2020 black and white Maxar imagery was most useful because of the minimal
252 seasonal snow cover. The 2018 NAIP imagery was helpful in situations where the 2020
253 imagery was obscured by shadow, distortion, or misaligned, and when color was needed
254 to improve interpretation. The 2010 lidar DEM (Robinson, 2014; Table A3) was used to
255 create a multidirectional hillshade to improve perspective and interpretation (Figure A1).
256 The rock debris on the termini of most glaciers and rock debris on some of the upper
257 parts of the glaciers were challenging to interpret. It was hard to determine whether ice
258 was present under the debris and whether that ice is part of the active glacier. Spatial
259 patterns of debris, debris contrasts, and melt streams flowing from the debris were used to
260 estimate the glacier boundaries.



261

262 **Figure A1.** Mt. Shasta glaciers in bluish white, perennial snowfields/ice patches in lavender
263 draped over a 3D rendering created from 2010 lidar (Robinson, 2014).
264
265

266 **Sierra Nevada**

267 The 2014 NAIP imagery was the best imagery due to low snow cover. In some cases,
268 features were difficult to outline because of shadow or image quality. In these cases,
269 2013/2012 Google Earth imagery were used. Some glaciers were reclassified as rock
270 glaciers by Trcka (2020). These were re-examined and where we agreed they were
271 removed from the initial glacier inventory. Defining whether the feature was a glacier or
272 rock glacier was often difficult, see Colorado section for more discussion.
273

274 **Trinity Alps**

275 The 2018 imagery was the best for the least snow cover. Justin Garwood (Garwood et al.,
276 2020) provided outlines for two glaciers, Grizzly and Salmon. The area of the most recent
277 outline of the Salmon Glacier was $< 0.01 \text{ km}^2$ and was not included in this inventory. By
278 2018 all of the other features mapped by the USGS (Fountain et al., 2017) were less than
279 0.01 km^2 or had disappeared. An additional feature was added based on the 2016 NLCD
280 (Jin et al. 2019).
281
282



283 **Table A1.** List of NAIP imagery used for outlining glaciers and perennial snowfields in
 284 California. ‘Date’ is the start and end date for flights covering the glaciated portions of the NAIP
 285 image. In some cases, flights were completed in a single day.
 286

Region/Year/Filename	County	Date (Year-M-D)
Cascade Range		
2014		
ortho_1-1_1n_s_ca089_2014_1.sid	Shasta	2014-07-13
ortho_1-1_1n_s_ca093_2014_1.sid	Siskiyou	2014-06-23 to 2014-07-18
2018		
ortho_1-1_hn_s_ca093_2018_1.sid	Siskiyou	2018-07-21 to 2018-09-25
Sierra Nevada		
2014		
ortho_1-1_1n_s_ca019_2014_1.sid	Fresno	2014-07-23 to 2014-08-23
ortho_1-1_1n_s_ca027_2014_1.sid	Inyo	2014-07-23 to 2014-08-23
ortho_1-1_1n_s_ca039_2014_2.sid	Madera	2014-07-18 to 2014-08-15
ortho_1-1_1n_s_ca051_2014_1.sid	Mono	2014-07-17 to 2014-08-15
ortho_1-1_1n_s_ca107_2014_1.sid	Tulare	2014-08-23 to 2014-08-23
Trinity Alps		
2018		
ortho_1-1_hn_s_ca093_2018_1.sid	Siskiyou	2018-07-21 to 2018-09-25

287
 288
 289
 290

Table A2. List of dates of the Maxar imagery used for outlining glaciers and perennial snowfields in California.

Region/ Date (Year-M-D)
Cascade Range
2020-10-05

291
 292
 293
 294

Table A3. List of U.S. Geological Survey digital elevation models used for outlining glaciers and perennial snowfields in California.

Filename	Date	Citation	URL
ds852_lidar	2010	Robinson (2014)	https://pubs.er.usgs.gov/publication/ds852

295
 296
 297
 298
 299

6.2 Colorado

The 2015 NAIP was generally free of seasonal snow. Where it persisted at the terminus of a few glaciers, images for the same year in Google Earth aided perimeter interpretation. Imagery used



300 are listed in Table A4.

301

302

Elk Mountains

303

No glacial features were mapped in the Elk Mountains by the USGS (Fountain et al., 2017). One glacier and four perennial snowfields were added from the SFI.

304

305

306

Front Range

307

The most recent inventory for the Front Range was Hoffman et al. (2007), which used aerial photographs to map the 2001 extent of glaciers. Many features in the Front Range are difficult to classify. The issue is the difference between a glacier or perennial snowfield and a rock glacier. Those that are part of the rock glacier are deleted from the glacier inventory. Those that seem to be separate from rock glaciers are retained. This is a judgement call. From a hydrological point of view, if a snow-ice patch that is part of a rock glacier was counted separately from a rock glacier, it is double counting a water feature.

308

309

310

311

312

313

314

315

316

The most challenging situation to interpret occurs when the glacier or perennial snowfield is located up-elevation from the rock glacier. If the slope of the snowfield smoothly transitions to the slope of the rock glacier, with no change in sign of the slope, we consider that one feature, a rock glacier (Figure A2). If the terrain dips below the snowfield, changing sign to rise to a topographic high below which the rock glacier clearly emerges, then they are two separate features. The patch does not appear to feed the rock glacier with ice (ice melt maybe, but not ice), because the ice would have to flow uphill to reach the rock glacier (Figure A3).

317

318

319

320

321

322

323

324

325

326

Table A4. List of NAIP imagery used for outlining glaciers and perennial snowfields in Colorado. ‘Date’ is the start and end date for flights covering the glaciated portions of the NAIP image. In some cases, flights were completed in a single day.

327

328

Region/Year/Filename	County	Date (Year-M-D)
Elk Mountains		
2015		
ortho_1-1_1n_s_co051_2015_1.sid	Gunnison	2015-09-10 to 2015-09-11
Front Range		
2015		
ortho_1-1_1n_s_co013_2015_1.sid	Boulder	2015-08-25 to 2015-09-20
ortho_1-1_1n_s_co049_2015_1.sid	Grand	2015-08-25 to 2015-09-20
ortho_1-1_1n_s_co057_2015_1.sid	Jackson	2015-09-09
ortho_1-1_1n_s_co069_2015_1.sid	Larimer	2015-08-25 to 2015-09-09
Gore Range		
2015		
ortho_1-1_1n_s_co037_2015_1.sid	Eagle	2015-09-10
Medicine Bow Mountains		
2015		
ortho_1-1_1n_s_co057_2015_1.sid	Jackson	2015-09-09



Park Range

2015
 ortho_1-1_1n_s_co057_2015_1.sid Jackson 2015-09-09

San Miguel Mountains

2015
 ortho_1-1_1n_s_co033_2015_1.sid Dolores 2015-09-11
 ortho_1-1_1n_s_co091_2015_1.sid Ouray 2015-09-11
 ortho_1-1_1n_s_co111_2015_1.sid San Juan 2015-09-12

Sawatch Range

2015
 ortho_1-1_1n_s_co037_2015_1.sid Eagle 2015-09-10
 ortho_1-1_1n_s_co097_2015_1.sid Pitkin 2015-09-10 to 2015-09-11

329

330 **6.3 Idaho**

331

332 The imagery quality was generally snow free. Of the glacier mapped by the USGS (Fountain et
 333 al., 2017) only two remain and are classified as perennial snowfields. The Borah Glacier was
 334 officially named in 2021, but is < 0.01 km², and is not included in the inventory. Table A5 lists
 335 the imagery used.


336

337

338 **Table A5.** List of NAIP imagery used for outlining glaciers and perennial snowfields in Idaho.
 339 'Date' is the start and end date for flights covering the glaciated portions of the NAIP image. In
 340 some cases, flights were completed in a single day.


Region/Year/File name	County	Date (Year-M-D)
Sawtooth Range		
2013		
ortho_1-1_hn_s_id015_2013_1.sid	Boise	2013-09-07
2015		
ortho_1-1_1n_s_id013_2015_1.sid	Blaine	2015-07-30
ortho_1-1_1n_s_id015_2015_1.sid	Boise	2015-09-08 to 2015-09-09
2019		
ortho_1-1_hn_s_id037_2019_1.sid	Custer	2019-07-25 to 2019-08-26

341



342

343 **Figure A2.** An example of a snowfield that is considered part of the rock glacier. Location,
344 Colorado Front Range, 40.827477° N, -106.657400° E. Image is from © Google Earth, 9/2014.



345
346

347 **Figure A3.** Tyndall Glacier in the Colorado Front Range, 40.305291° N, -105.689602° E, with a
348 rock glacier slightly down valley. Image is from © Google Earth 9/2016.

349

350 **6.4 Montana**

351

352 Image quality varied between mountain ranges due to differences in snow cover. Tables A6 and
353 A7 list the imagery used.

354

355 **Beartooth-Absaroka Range**

356

357 The 2015 NAIP imagery was the best overall imagery due to the least snow, but Google
358 Earth was occasionally used as well. Google Earth had imagery dated to 9/11/2015; often
359 with less seasonal snow than the NAIP imagery. To counter any mismatch in projection,
360 outlines digitized in Google Earth were imported to ArcGIS and projected to match the
361 NAIP projection.

361

Bitterroot Range



362 There were no glacial features were mapped in the Bitterroot Range by the USGS
363 (Fountain et al., (2017). One glacier and three perennial snowfields were added based on
364 the NLCD.

365

366

Cabinet Range

367 The USGS mapped four glacial features $\geq 0.01 \text{ km}^2$ (Fountain et al., 2017). Inspection of
368 the 2015 only one was $\geq 0.01 \text{ km}^2$. Seven glaciers and perennial snowfields were added;
369 five were identified in our initial inventory, the other two were identified by the SFI and
370 NLCD, respectively. All were less than 0.05 km^2 .

371

372

Crazy Mountains

373 The 2013 NAIP imagery was the best imagery available and included limited seasonal
374 snow. The 2019 Maxar imagery had too much seasonal snow.

375

376

Lewis Range (Glacier National Park)

377 The most recent published glacier inventory is a 2015 USGS inventory (Fagre et al.,
378 2017). They outlined the main-body of named-glaciers using 2015 Maxar imagery. We
379 digitized the outlines of all glaciers and perennial snowfields using 2015 Maxar imagery
380 where available. Elsewhere, 2015 and 2013 NAIP imagery were used; both years had lots
381 of seasonal snow cover. Two major glaciers, Blackfoot (Figure A4) and Harrison (Figure
382 A5) glaciers, separated into pieces as it retreated since it was originally mapped by the
383 USGS (Fountain et al., 2007).

384

385

Madison Range


386 The 2013 NAIP imagery was the only good imagery due to extensive snow in the other
387 years. No glaciers or perennial snowfields were found. Of the two features $\geq 0.01 \text{ km}^2$
388 mapped by the USGS (Fountain et al., 2017), the 2013 imagery showed that one feature
389 is a rock glacier and the other was less than 0.01 km^2 .

390

391


Mission-Swan-Flathead Ranges

392 Based on the least snow cover, the 2013 NAIP was better in the Mission and Flathead
393 Ranges, and the 2015 NAIP was better in the Swan Range. No glaciers or perennial
394 snowfields remain in the Flathead Range.



395

396 **Figure A4.** The updated (2015) outlines for the Blackfoot Glacier including the main glacier
397 body (red) and the additional smaller glacier (orange). Base image from the NAIP taken in 2013.



398

399 **Figure A5.** The updated (2015) outlines for Harrison Glacier including the main glacier body
 400 (red) and the additional smaller glaciers (orange). Base image from the NAIP taken in 2013.

401

402

403 **Table A6.** List of NAIP imagery used for outlining glaciers and perennial snowfields in
 404 Montana. ‘Date’ is the start and end date for flights covering the glaciated portions of the NAIP
 405 image. In some cases, flights were completed in a single day.

Region/Year/Filename	County	Date (Year-M-D)
Beartooth Mountains-Absaroka Range		
2013		
ortho_1-1_1n_s_mt067_2013_1.sid	Park	2013-08-05 to 2013-09-11
2015		
ortho_1-1_1n_s_mt009_2015_1.sid	Carbon	2015-08-10 to 2015-09-07



ortho_1-1_1n_s_mt067_2015_1.sid	Park	2015-08-19 to 2015-09-11
ortho_1-1_1n_s_mt095_2015_1.sid	Stillwater	2015-08-10 to 2015-09-07
Bitterroot Range		
2013		
ortho_1-1_1n_s_mt001_2013_1.sid	Beaverhead	2013-08-04
2015		
ortho_1-1_1n_s_mt081_2015_2.sid	Ravalli	2015-10-06 to 2015-11-07
Cabinet Mountains		
2015		
ortho_1-1_1n_s_mt053_2015_2.sid	Lincoln	2015-09-11 to 2016-08-15
Crazy Mountains		
2013		
ortho_1-1_1n_s_mt067_2013_1.sid	Park	2013-08-05 to 2013-09-11
ortho_1-1_1n_s_mt097_2013_1.sid	Sweet Grass	2013-08-31 to 2013-09-10
2015		
ortho_1-1_1n_s_mt067_2015_1.sid	Park	2015-08-19 to 2015-09-11
Lewis Range		
2013		
ortho_1-1_1n_s_mt029_2013_1.sid	Flathead	2013-08-21 to 2013-09-01
ortho_1-1_1n_s_mt035_2013_1.sid	Glacier	2013-08-21 to 2013-09-01
2015		
ortho_1-1_1n_s_mt029_2015_2.sid	Flathead	2015-09-30 to 2016-10-21
ortho_1-1_1n_s_mt035_2015_2.sid	Glacier	2015-10-14 to 2016-08-21
Mission Range-Swan-Flathead Ranges		
2013		
ortho_1-1_1n_s_mt029_2013_1.sid	Flathead	2013-08-21 to 2013-09-01
ortho_1-1_1n_s_mt063_2013_1.sid	Missoula	2013-09-01
2015		
ortho_1-1_1n_s_mt047_2015_2.sid	Lake	2015-09-12 to 2016-08-15
ortho_1-1_1n_s_mt063_2015_2.sid	Missoula	2015-09-12 to 2016-08-16

406
 407
 408
 409
 410

Table A7. List of dates of the Maxar imagery used for outlining glaciers and perennial snowfields in Montana.

Region/ Date (Year-M-D)

Lewis Range

2015-08-22
 2015-09-01
 2015-09-12
 2015-09-25
 2019-08-20

411



412 **6.5 Oregon**

413

414 Tables A8, A9, and A10 list the imagery and DEM used.

415

416 **Cascade Range**

417

418 **Mount Hood**

419 The most recent glacier outlines for Mt. Hood were based on 2015 and 2016 Maxar color
420 imagery with interpretation aid using Google Earth. Due to seasonal snow some
421 professional judgement was required in places.

422

423 **Mount Jefferson**

424 Seasonal snow was extensive in places. The 2018 NAIP had extensive seasonal snow and
425 generally only useful near the terminus of some glaciers. Used 2018 Maxar imagery that
426 showed little seasonal snow, but a little cloudy that masked a bit of Whitewater Glacier.
427 Also used Google Earth to help interpret some of the features.

428

429 **Three Sisters**

430 Used Maxar 2018 imagery, but where image stretching along the feature's headwall 2018
431 NAIP imagery was used. Two versions of the Maxar imagery for the same day are
432 3available, one color, one black and white. Color was georectified but suffered stretching
433 along some headwalls. A light early season snowfall occurred before the Maxar image
434 and the snow accumulated in some places just enough to obscure the surface. So, the
435 glacier or snow patch outline was the minimum of the two images with occasional
436 interpolation across the snowy surface to the nearest glacier edge.

437

438 An example of buried ice on South Sister is shown in Figure A6.

439

440 **Mount Thielsen**

441 The Lathrop Glacier was named in 1981. At the time of the USGS mapping and now it is
442 $<0.01 \text{ km}^2$, and not counted as part of the inventory. Furthermore, Lathrop Glacier has
443 been known to disappear in some years and therefore fails the definition of a glacier.

444

445 **Wallowa Mountains**

446 No NAIP imagery was useful and Maxar did not image this region. We used the
447 8/30/2013 image from Google Earth, which was excellent with little snow. Features were
448 digitized in Google Earth and then imported into ArcGIS. Because we used NAIP as the
449 base imagery, we revised the outline from the projection in WGS84 (Google Earth) to
450 NAD83 UTM Zone 11 (NAIP).

451



452

453 **Figure A6.** Lost Creek Glacier, South Sister, Oregon. Note buried ice and lack of crevasses to
 454 the left of the grey-blue ice, suggesting ice that is no longer moving and therefore not part of the
 455 dynamic glacier. The white box surrounds an area that has collapsed due to subsurface melt. The
 456 inset enlargement shows a cliff edge of exposed dirty ice (white arrow) indicated by a darker
 457 color suggesting wet sediment and a finer texture than the surface debris. The black arrow shows
 458 the width of the cleaner ice for scale. Image is from © Google Earth, 8/9/2021.

459

460

461 **Table A8.** List of NAIP imagery used for outlining glaciers and perennial snowfields in Oregon.
 462 ‘Date’ is the start and end date for flights covering the glaciated portions of the NAIP image. In
 463 some cases, flights were completed in a single day.

Region/Year/Filename	County	Date (Year-M-D)
Cascade Range		
2014		
ortho_1-1_1n_s_or017_2014_1.sid	Deschutes	2014-09-01
ortho_1-1_1n_s_or027_2014_1.sid	Hood River	2014-08-27 to 2014-09-05
ortho_1-1_1n_s_or039_2014_1.sid	Lane	2014-09-01
2016		



ortho_1-1_1n_s_or027_2016_1.sid	Hood River	2016-08-04
2017/2018		
ortho1-1_hn_s_or017_2017_2018_1.sid	Deschutes	2018-07-28
Wallowa Mountains		
2014		
ortho_1-1_1n_s_or063_2014_1.sid	Wallowa	2014-10-05

464

465

466 **Table A9.** List of dates of the Maxar imagery used for outlining glaciers and perennial
 467 snowfields in Oregon.

Region/ Date (Year-M-D)
Cascade Range
2015-08-20
2015-09-11
2015-10-05
2016-09-10
2018-09-17
2020-09-20

468

469

470 **Table A10.** List of Oregon Department of Geology and Mineral Industries digital elevation
 471 models used for outlining glaciers and perennial snowfields in Oregon. ‘

Filename	Date	URL
2011_OLC_Deschutes	2011	gis.dogami.oregon.gov/maps/lidarviewer/

472

473

474 **6.6 Washington**

475

476 The 2015 NAIP imagery was typically excellent with little snow cover, whereas the 2017 NAIP
 477 had more snow and the 2019 imagery had lots of snow. For most outlines, 2015 NAIP imagery
 478 was used. In some places, the 2017 NAIP imagery had less snow and was used instead. Maxar
 479 imagery was of limited use and often wasn’t better than the 2015 or 2017 NAIP. Tables A11,
 480 A12, and A13 list the imagery and DEMs used.

481

482 **Cascade –Northern**

483 The glaciers and perennial snowfields were previously inventoried by (Dick, 2013).

484

485 **Mount Baker**

486 The 2015 NAIP imagery was the best and had little seasonal snow. Google Earth 2009
 487 and 2019 imagery were used to help interpretation. A multidirectional hillshade and 3-
 488 meter contour lines derived from a lidar DEM (Bard, 2017b; Table A13); were used to
 489 help define flow divides between glaciers, debris covered-ice, and buried ice. There are
 490 notable differences between the NAIP imagery and DEM data, particularly in steep
 491 terrain, areas of dark shadow, and debris-covered areas. The DEM helped correct these
 492 positional errors and the benefit of supplying more information on surface texture.



493 Several buried-ice features were identified. The ice appeared to have decoupled from the
494 active glacier. In a few cases, debris-covered ice is included in the glacier outline because
495 the ice appears to be directly connected to the glacier, and there was evidence of
496 movement.

497 **Dragontail Peak**

499 The GNIS locates Snow Creek Glacier at a point on the edge of the southeast glacier
500 (Fountain et al., 2007). In the 2015 imagery, the point is on bedrock, making it unclear
501 which glacier the GNIS is naming. The USGS identifies both glaciers as Snow Creek
502 Glacier. We labeled both glaciers as the Snow Creek Glacier.

504 **Glacier Peak**

505 For the Glacier Peak region, a multidirectional hillshade and 3-m contour lines derived
506 from a 2015 lidar DEM (Bard, 2017A; Table A13) was used as a guide to define flow
507 divides.

509 **Hurry-up Peak**

510 The point location of the South Glacier provided by the GNIS is over bedrock. We
511 assume the point refers to the glacier located ~150 m to the north of the point.

513 **Cascade –Southern**

515 **Goat Rocks**

516 Imagery from 2015 was best, but more snow than desired. Too much snow in 2017 but
517 some ice is exposed. The 2019 imagery was way too snowy and was considered useless
518 for glacier digitization.

520 The outlines are almost entirely based on 2015 imagery, and a few on 2017, where
521 needed. Used 2009 NAIP imagery to help define the headwalls at the Conrad, McCall,
522 and Packwood glaciers. Heard (2000) previously mapped the glacier perimeters. The
523 maximum extent of the seasonal snow covering the terminal regions was not digitized.
524 Typically digitized at scales of 1:600 to 1:800. Note that narrow arms of the snowfields
525 were not typically digitized knowing that they would probably disappear a few days to a
526 week from the time of imagery.

528 **Mount Adams**

529 No suitable NAIP imagery was found, instead 2019 Maxar imagery was used. In addition
530 to the Maxar imagery, a multidirectional hillshade and 3-m contour lines derived from a
531 2016 lidar DEM (Bard 2019, Table 13A) were used as a guide when delineating flow
532 divides. Occasionally, 2009 Google Earth imagery was also useful. Extensive snow
533 covered the mountain when the 2016 lidar was flown masking some of the glacier
534 termini. However, the DEM was helpful in correcting the imagery where poorly aligned
535 with the terrain.

537 Multiple buried-ice features were identified near the terminus of several glaciers where



538 ice appeared to have decoupled from the main active glacier. Large areas below the
539 glaciers (Mazama, Adams, and Pinnacle) likely have debris-covered ice. We focused on
540 the features which were likely to contain ice based on meltwater streams exiting near the
541 features and hummocky terrain which appeared to indicate melt. Ground-based images
542 from taken between 2014 to 2018 helped decision-making. The images were particularly
543 helpful in identifying a debris-covered ice cliff at Adams Glacier.

544

545

Mount Rainier

546

In general, the 2019 NAIP and the Maxar (2018-09-25) were used for the outlines (Table
547 A12). Although the GNIS includes the Nisqually Icefall as a separate feature, we
548 included the icefall as part of the Nisqually Glacier (Figure A7).

549

550

Mount St. Helens

551

We used a GIS layer of geological mapping units that included snow and ice from the
552 USGS (David Sherrod, personal communication, 2021) to help guide our search. The
553 Crater Glacier (INV_ID E562842N5115499) was heavily debris covered, and obscured
554 by shadow in some areas.

555

556

Olympic Mountains

557


558

A 2015 inventory of the region was compiled because more recent imagery (NAIP and
559 Maxar) were not useful due to seasonal snow. Our updated inventory differs from that
560 published in Fountain et al. (2017) in two ways. First, they outlined and grouped the
561 glaciers and perennial snowfields according to watershed rather than individual glacier.
562 Their goal was to estimate glacier change relative to a previous study by Spicer (1986)
563 and had to follow Spicer's approach. Second, all outlines were rechecked and compared
564 to SFI and the NLCD resulting in minor changes.

565

566

567



568

569 **Figure A7.** Image of the Nisqually Glacier and Icefall. The orange and red outlines are from the
 570 updated inventory and the blue outline is from the USGS mapping (Fountain et al., 2007)
 571 database. The base image is from the NAIP taken in 2019.

572

573

574 **Table A11.** List of NAIP imagery used for outlining glaciers and perennial snowfields in
 575 Washington. ‘Date’ is the start and end date for flights covering the glaciated portions of the
 576 NAIP image. In some cases, flights were completed in a single day. For 2006 the inspection date
 577 was used, since the start and end dates were not provided.

578

Region/Year/Filename	County	Date (Year-M-D)
Cascade –Northern 2006		



ortho_1-1_1n_s_wa007_2006_3.sid	Chelan	2006-07-01
2015		
ortho_1-1_1n_s_wa007_2015_1.sid	Chelan	2015-07-06 to 2015-09-23
ortho_1-1_1n_s_wa033_2015_1.sid	King	2015-07-06 to 2015-09-27
ortho_1-1_1n_s_wa037_2015_1.sid	Kittitas	2015-07-06 to 2015-09-23
ortho_1-1_1n_s_wa047_2015_1.sid	Okanogan	2015-09-09 to 2015-09-11
ortho_1-1_1n_s_wa057_2015_1.sid	Skagit	2015-07-06 to 2015-09-29
ortho_1-1_1n_s_wa061_2015_1.sid	Snohomish	2015-07-06 to 2015-09-29
ortho_1-1_1n_s_wa073_2015_1.sid	Whatcom	2015-09-10 to 2015-09-26
2017		
ortho_1-1_1n_s_wa007_2017_1.sid	Chelan	2017-10-03 to 2017-10-24
ortho_1-1_1n_s_wa057_2017_1.sid	Skagit	2017-09-27 to 2017-10-05
ortho_1-1_1n_s_wa073_2017_1.sid	Whatcom	2017-09-27 to 2017-10-05
Cascade –Southern		
2015		
ortho_1-1_1n_s_wa041_2015_1.sid	Lewis	2015-07-15 to 2015-07-29
ortho_1-1_1n_s_wa053_2015_1.sid	Pierce	2015-07-29
ortho_1-1_1n_s_wa059_2015_1.sid	Skamania	2015-07-15 to 2015-09-12
ortho_1-1_1n_s_wa077_2015_1.sid	Yakima	2015-07-15 to 2015-07-29
2019		
ortho_1-1_hn_s_wa053_2019_1.sid	Pierce	2019-08-26
ortho_1-1_hn_s_wa059_2019_1.sid	Skamania	2019-08-06 to 2019-08-26
Olympic Mountains		
2015		
ortho_1-1_1n_s_wa009_2015_1.sid	Clallam	2015-07-28 to 2015-09-12
ortho_1-1_1n_s_wa031_2015_1.sid	Jefferson	2015-07-28 to 2015-09-12
ortho_1-1_1n_s_wa045_2015_1.sid	Mason	2015-07-28 to 2015-08-19

579
 580
 581
 582
 583

Table A12. List of dates of the Maxar imagery used for outlining glaciers and perennial snowfields in Washington.

Region/ Date (Year-M-D)

Cascade Range-Northern

2018-09-25

Cascade Range-Southern

2018-09-25

2019-08-31

Olympic Mountains

2015-08-17

2019-09-30

584
 585
 586
 587

Table A13. List of U.S. Geological Survey digital elevation models used for outlining glaciers and perennial snowfields in Washington.



Region	Date	Citation	URL www.sciencebase.gov/catalog/item/
Mt. Adams	2016	Bard (2019)	5bc623b9e4b0fc368ebbe99a
Glacier Peak	2014-15	Bard (2017a)	57bf299ee4b0f2f0ceb7534e
Mt. Baker	2015	Bard (2017b)	58518b0ee4b0f99207c4f12c

588

589

590

6.7 Wyoming

591

592

Wind River Range

593

594

595

596

597

598

599

Tables A14 and A15 list the imagery used. The 2015 NAIP imagery had little snow in contrast to 2019 imagery. Shadows are common in the 2015 imagery and can be very dark. Occasionally the 2019 imagery was used to define the glacier-bedrock headwall boundary. The 2019 Maxar imagery was essentially identical to the NAIP and because its black and white not as useful. Imagery from 2017 and 2018 were a bit too snowy around the glacier margin to be useful. The 2018-09-06 Maxar imagery covered the entire range, with some clouds.

600

601

602

603

604

605

606

607

608

In the southern Wind River Range, a new snow dusting was often present, occasionally making it difficult to outline snowfields and glaciers, but mostly snowfields. Distinguishing seasonal snow from perennial snow was a judgement call. The thin seasonal snow was identified if a slight coloration similar to underlying rock/soil was present or many small patches (a few square meters) of snow-free surface present or many rocks protruding through the snow. A perennial patch of snow appeared smooth and white, hiding underlying surface. Thin snow covering glacial ice was typically greyish in color, often with banding, much less of a texture than surrounding ice-free landscape.

609

610

611

612

613

614

615

Often, it seems a number of glaciers are thinning in place with a thin layer of debris on the ice that thickens down valley. The landscape surrounding this relatively smooth appearance is rumpled like the bedrock terrain further away. Interpretation is difficult. These are probably shrinking glaciers that are being covered in the debris. The outline is clear where ice meets bedrock, but in the talus debris area, we digitized along the glacier ice boundary unless some other feature like exposed ice or a crevasse is visible then included that as part. See INV_ID E618081N4774579 (Figure A9) for example.

616

617

618

At Lower Fremont Glacier, a number of sizable ice patches appear down valley as if a deposit of buried ice is present. However, there is no obvious connection to the glacier itself.

619

620


The GNIS identified a single glacier as the Sacagawea Glacier, and two separate Fremont Glaciers (Figure A10). By 2017 the single glacier had split into four glaciers. We chose



621 to label the largest glacier and the glacier to the southeast the Sacagawea Glacier. The
622 other two glaciers were labeled the Fremont Glaciers.



623
624 **Figure A9.** An example of an unnamed glacier in the Wind River Range, WY, INV_ID
625 E618081N4774579 seemingly melting into the talus surrounding the terminus (top of image).
626 The glacier is flowing from the lower left-hand corner to the upper right-hand corner. Base
627 image from the NAIP taken in 2015.
628
629



630

631 **Figure A10.** Image of Fremont Glaciers and Sacagawea Glacier showing the Sacagawea outline
 632 from the Fountain et al. (2017) database (blue), our updated Fremont Glaciers outlines (orange),
 633 and updated Sacagawea outlines (red). The base image is from the NAIP, taken in 2015.

634

635

636 **Table A14.** List of NAIP imagery used for outlining glaciers and perennial snowfields in
 637 Wyoming. ‘Date’ is the start and end date for flights covering the glaciated portions of the NAIP
 638 image. In some cases, flights were completed in a single day. For 2006 the inspection date was
 639 used, since the start and end dates were not provided.

640

Region/Year/Filename	County	Date (Year-M-D)
Absaroka Range		
2006		



ortho_1-2_1n_s_wy029_2006_1.sid	Park	2006-09-02
2015		
ortho_1-1_hn_s_wy013_2015_2.sid	Fremont	2015-09-09 to 2015-10-13
ortho_1-1_hn_s_wy029_2015_2.sid	Park	2015-09-22 to 2015-10-13
Bighorn Mountains, WY		
2015		
ortho_1-1_hn_s_wy019_2015_2.sid	Johnson	2015-09-12
Teton Range		
2006		
ortho_1-1_1n_s_wy039_2006_1.sid	Teton	2006-09-02
2015		
ortho_1-1_hn_s_wy035_2015_2.sid	Sublette	2015-09-09 to 2015-10-13
ortho_1-1_hn_s_wy039_2015_2.sid	Teton	2015-09-12 to 2015-09-22
2019		
ortho_1-1_hn_s_wy039_2019_1.sid	Teton	2019-07-20 to 2015-09-22
Wind River Range		
2006		
ortho_1-1_1n_s_wy035_2006_1.sid	Sublette	2006-09-02
2015		
ortho_1-1_hn_s_wy013_2015_2.sid	Fremont	2015-09-09 to 2015-10-13
ortho_1-1_hn_s_wy035_2015_2.sid	Sublette	2015-09-09 to 2015-10-13
2019		
ortho_1-1_hn_s_wy013_2019_1.sid	Fremont	2019-07-20 to 2019-08-27
ortho_1-1_hn_s_wy035_2019_1.sid	Sublette	2019-08-15 to 2019-09-13

641

642

643 **Table A15.** List of dates of the Maxar imagery used for outlining glaciers and perennial

644 snowfields in Wyoming.

Region/ Date (Year-M-D)

Wind River Range

2018-09-06

645

646

6.8 Glaciers that have split into multiple pieces and current errors in glacier label names

647

648

649 Table A16 compiles the list of glaciers that have split into multiple pieces since the USGS
 650 1:24000 mapping (Fountain et al., 2017). Table A17 lists current map errors in the labeling of
 651 glaciers. The purpose of including this table is to facilitate updating the USGS Geographic
 652 Names Information System.

653

654 **Table A16.** List of named glaciers that have split into multiple pieces. Names come from the
 655 Geographic Names Information System. ‘Count’ refers to the number of pieces in the updated



656 inventory. ‘Classes’ is the classification of the pieces; glacier, perennial snowfield, buried-ice, or
 657 a combination.

State/Region/Glacier Name	Count	Classes
California		
Cascade Range		
Bolam Glacier	2	Glaciers and perennial snowfields
Hotlum Glacier	2	Glaciers and perennial snowfields
Whitney Glacier	2	Glaciers and perennial snowfields
Wintun Glacier	3	Glaciers and perennial snowfields
Sierra Nevada		
Goethe Glacier	2	Glaciers only
Lyell Glacier	4	Glaciers and perennial snowfields
Norman Clyde Glacier	3	Glaciers only
Powell Glacier	2	Glacier and Buried-ice
Colorado		
Front Range		
Saint Vrain Glaciers	6	Glaciers and perennial snowfields
Montana		
Beartooth Mountains-Absaroka Range		
Castle Rock Glacier	3	Glaciers and perennial snowfields
Granite Glacier	2	Glaciers only
Grasshopper Glacier	4	Glaciers and perennial snowfields
Hopper Glacier	2	Glaciers and perennial snowfields
Snowbank Glacier	2	Glaciers only
Wolf Glacier	2	Glaciers only
Lewis Range		
Agassiz Glacier	3	Glaciers only
Blackfoot Glacier	2	Glaciers only
Carter Glaciers	2	Glaciers and perennial snowfields
Dixon Glacier	3	Glaciers and perennial snowfields
Harrison Glacier	5	Glaciers and perennial snowfields
Kintla Glacier	2	Glaciers only
Logan Glacier	2	Glaciers only
Shepard Glacier	3	Glaciers only
Siyeh Glacier	2	Glaciers only
Two Ocean Glacier	2	Glaciers only
Whitecrow Glacier	5	Glaciers and perennial snowfields
Mission Range-Swan Range-Flathead Range		
Swan Glaciers	3	Glaciers and perennial snowfields
Oregon		
Cascade Range		
Bend Glacier	3	Glaciers and perennial snowfields
Clark Glacier	2	Perennial snowfields only
Collier Glacier	2	Glaciers only
Diller Glacier	2	Glaciers and perennial snowfields
Glisan Glacier	2	Glaciers and perennial snowfields
Ladd Glacier	4	Glaciers and perennial snowfields
Langille Glacier	5	Glaciers and perennial snowfields



Newton Clark Glacier	3	Glaciers and perennial snowfields
Palmer Glacier	2	Perennial snowfields only
Prouty Glacier	3	Glaciers and perennial snowfields
Renfrew Glacier	2	Glaciers and perennial snowfields
Russell Glacier	2	Glaciers only
Sandy Glacier	4	Glaciers and perennial snowfields
Skinner Glacier	4	Perennial snowfields only
Waldo Glacier	3	Glaciers only
White River Glacier	2	Glaciers and perennial snowfields
Whitewater Glacier	3	Glaciers only
Zigzag Glacier	3	Glaciers and perennial snowfields
Washington		
Cascade Range-Northern		
Borealis Glacier	4	Glaciers only
Buckner Glacier	2	Glaciers only
Butterfly Glacier	4	Glaciers only
Colchuck Glacier	2	Glaciers only
Company Glacier	3	Glaciers only
Cool Glacier	2	Glaciers and perennial snowfields
Dana Glacier	3	Glaciers only
Dark Glacier	3	Glaciers only
Dome Glacier	2	Glaciers only
Douglas Glacier	4	Glaciers and perennial snowfields
Dusty Glacier	2	Glaciers and perennial snowfields
East Nooksack Glacier	5	Glaciers only
Entiat Glacier	4	Glaciers and perennial snowfields
Forbidden Glacier	2	Glaciers only
Fremont Glacier	2	Glaciers only
Goode Glacier	2	Glaciers only
Hadley Glacier	5	Glaciers only
Hanging Glacier	2	Glaciers only
Hinman Glacier	4	Glaciers only
Honeycomb Glacier	3	Glaciers and perennial snowfields
Inspiration Glacier	3	Glaciers and perennial snowfields
Isella Glacier	2	Glaciers and perennial snowfields
Jerry Glacier	2	Glaciers only
Kimtah Glacier	3	Glaciers only
LeConte Glacier	7	Glaciers and perennial snowfields
Lyllall Glacier	2	Perennial snowfields only
Mazama Glacier	3	Glaciers and perennial snowfields
McAllister Glacier	2	Glaciers only
Middle Cascade Glacier	2	Glaciers only
Neve Glacier	3	Glaciers only
No Name Glacier	5	Glaciers and perennial snowfields
Nohokomeen Glacier	2	Glaciers only
North Klawatti Glacier	2	Glaciers and perennial snowfields
Pilz Glacier	3	Glaciers and perennial snowfields
Price Glacier	4	Glaciers only
Ptarmigan Glacier	2	Glaciers and perennial snowfields



Queest-alb Glacier (not official)	3	Glaciers and perennial snowfields
Rainbow Glacier	3	Glaciers and perennial snowfields
Redoubt Glacier	2	Glaciers only
Richardson Glacier	2	Glaciers only
S Glacier	3	Glaciers only
Sandalee Glacier	4	Glaciers only
Scimitar Glacier	3	Glaciers only
Sholes Glacier	4	Glaciers only
Sitkum Glacier	4	Glaciers and perennial snowfields
Snow Creek Glacier	2	Perennial snowfields only
South Cascade Glacier	2	Glaciers only
Spider Glacier	2	Glaciers only
Suiattle Glacier	2	Glaciers only
Sulphide Glacier	2	Glaciers only
Thunder Glacier	3	Glaciers only
Thunder Glacier	2	Glaciers only
White Chuck Glacier	5	Glaciers and perennial snowfields
White Salmon Glacier	2	Glaciers only
Wyeth Glacier	3	Glaciers and perennial snowfields
Cascade Range-Southern		
Adams Glacier	4	Glaciers and perennial snowfields
Avalanche Glacier	2	Glaciers only
Conrad Glacier	3	Glaciers and perennial snowfields
Cowlitz Glacier	2	Glaciers and perennial snowfields
Crescent Glacier	2	Glaciers and perennial snowfields
Flett Glacier	6	Glaciers and perennial snowfields
Fryingpan Glacier	5	Glaciers and perennial snowfields
Gotchen Glacier	2	Glaciers and perennial snowfields
Kautz Glacier	2	Glaciers and perennial snowfields
Klickitat Glacier	2	Glaciers only
Lava Glacier	3	Glaciers and perennial snowfields
McCall Glacier	6	Glaciers and perennial snowfields
Meade Glacier	5	Perennial snowfields only
North Mowich Glacier	2	Glaciers and perennial snowfields
Ohanapecosh Glacier	6	Glaciers and perennial snowfields
Paradise Glacier	3	Glaciers and perennial snowfields
Pinnacle Glacier	3	Glaciers and perennial snowfields
Puyallup Glacier	2	Glaciers and perennial snowfields
Pyramid Glacier	4	Glaciers and perennial snowfields
Russell Glacier	2	Glaciers only
Sarvant Glaciers	4	Glaciers and perennial snowfields
South Mowich Glacier	2	Glaciers only
South Tahoma Glacier	2	Glaciers and perennial snowfields
Success Glacier	2	Glaciers and perennial snowfields
Van Trump Glacier	10	Glaciers and perennial snowfields
White Salmon Glacier	2	Glaciers only
Whitman Glacier	5	Glaciers and perennial snowfields
Wilson Glacier	3	Glaciers and perennial snowfields
Olympic Mountains		



Blue Glacier	2	Glaciers only
Cameron Glaciers	4	Glaciers and perennial snowfields
Carrie Glacier	2	Glaciers only
Eel Glacier	2	Glaciers only
White Glacier	2	Glaciers only
Wyoming		
Teton Range		
Middle Teton Glacier	2	Glaciers and perennial snowfields
Triple Glaciers	3	Glaciers only
Wind River Range		
Bull Lake Glacier	3	Glaciers and perennial snowfields
Dinwoody Glacier	2	Glaciers only
Dinwoody Glaciers	3	Glaciers and perennial snowfields
Grasshopper Glacier	3	Glaciers only
Harrower Glacier	2	Perennial snowfields only
Helen Glacier	3	Glaciers only
Lower Fremont Glacier	4	Glaciers and perennial snowfields
Mammoth Glacier	2	Glaciers and perennial snowfields
Minor Glacier	2	Glaciers and perennial snowfields
Sacagawea Glacier	4	Glaciers and perennial snowfields
Sourdough Glacier	2	Glaciers and perennial snowfields
Stroud Glacier	3	Glaciers and perennial snowfields
Twins Glacier	2	Glaciers and perennial snowfields
Upper Fremont Glacier	2	Glaciers and perennial snowfields

658

659

660 **Table A17.** List of officially named glaciers where we identified an issue with the glacier name
 661 on the 1:24000 U.S. Geological Survey topographical maps (Fountain et al., 2017). Names come
 662 from the Geographic Names Information System (U.S. Geological Survey (2022)). The 'Issue'
 663 column lists the type of issue identified. 'Not labeled' indicates the feature was present but not
 664 labeled, 'Misidentified' indicates the wrong feature was labeled, and 'Label unclear' indicates it
 665 is unclear what feature the label is identifying.

State/Region/Glacier Name	Issue
Colorado	
Front Range	
Arikaree Glacier	Not labeled
Navajo Glacier	Not labeled
Oregon	
Cascade Range	
Carver Glacier	Misidentified
Milk Creek Glacier	Not labeled
Washington	
Cascade Range-Northern	
S Glacier	Label unclear
Snow Creek Glacier	Label unclear
South Glacier	Not labeled
Cascade Range-Southern	



No Name Glacier	Not labeled
Stevens Glacier	Not labeled
<hr/>	
Wyoming	
Wind River Range	
Dinwoody Glaciers	Label unclear
Fremont Glaciers	Label unclear

666

667

668 **Author Contributions.** Andrew G. Fountain was the principal investigator of the project, he
669 wrote the proposal and digitized glacier and snowfield outlines, analyzed the data, and led the
670 writing of this report. Bryce Glenn was the GIS expert responsible for the geographic format
671 (e.g. projection, attributes, database structure) and quality control. He digitized glacier and
672 snowfield outlines, analyzed the data, and helped write the report. Chris McNeil provided some
673 of the imagery.

674

675 **Competing Interests.** The authors declare that they have no conflict of interest.

676

677 **Acknowledgments.** Hassan Basagic and Kristina Dick digitized the glaciers of Mt. Shasta and
678 the North Cascade National Park, respectively. We greatly appreciate their help and expertise.
679 Nathan Walker of the U.S. Forest Service provided a very helpful initial review of this report.
680 We kindly acknowledge the funding from the US. Forest Service. Mazar imagery was accessed
681 through the USGS and NGA NEXTVIEW license. The Maxar imagery has limited availability
682 owing to restrictions (proprietary interest). Contact cmcneil@usgs.gov for more information.
683 Any use of trade, firm or product names is for descriptive purposes only and does not imply
684 endorsement by the U.S. Government.

685 **References**

686

687 Alley, R. B., Cuffey, K. M., & Zoet, L. K. (2019). Glacial erosion: Status and outlook. *Annals of*
688 *Glaciology*, 60(80), 1–13. <https://doi.org/10.1017/aog.2019.38>

689 Bard, J. A. (2017a). *High-resolution digital elevation dataset for Glacier Peak and vicinity,*
690 *Washington, based on lidar surveys of August–November, 2014 and June, 2015* [Data set]. U.S.
691 Geological Survey. <https://doi.org/10.5066/F7H41PJG>

692 Bard, J. A. (2017b). *High-resolution digital elevation dataset for Mt Baker and vicinity,*
693 *Washington, based on lidar surveys of 2015* [Data set]. U.S. Geological Survey.
694 <https://doi.org/10.5066/F7WD3XR0>

695 Bard, J. A. (2019). *High-resolution digital elevation dataset for Mount Adams and vicinity,*
696 *Washington, based on lidar surveys of August–September, 2016* [Data set]. U.S. Geological
697 Survey. <https://doi.org/10.5066/P9Z1HF1K>

698 Beason, S. R., Legg, N. T., Kenyon, T. R., Jost, R. P., & Kennard, P. M. (2018). Forecasting and
699 seismic detection of debris flows in pro-glacial rivers at Mount Rainier National Park,



- 700 Washington, USA. *Contained in: Proceedings of the Seventh International Conference on*
701 *Debris-Flow Hazards Mitigation, Golden, Colorado, USA, June 10-13, 2019,*
702 *Https://Hdl.Handle.Net/11124/173051.* <https://doi.org/10.25676/11124/173232>
- 703 Benn, D. I., & Evans, D. J. A. (2010). *Glaciers and glaciation* (2nd ed.). Hodder Education.
- 704 Bolch, T., Menounos, B., & Wheate, R. (2010). Landsat-based inventory of glaciers in western
705 Canada, 1985–2005. *Remote Sensing of Environment*, 114(1), 127–137.
706 <https://doi.org/10.1016/j.rse.2009.08.015>
- 707 Brardinoni, F., Scotti, R., Sailer, R., & Mair, V. (2019). Evaluating sources of uncertainty and
708 variability in rock glacier inventories. *Earth Surface Processes and Landforms*, 44(12), 2450–
709 2466. <https://doi.org/10.1002/esp.4674>
- 710 Cadbury, S. L., Hannah, D. M., Milner, A. M., Pearson, C. P., & Brown, L. E. (2008). Stream
711 temperature dynamics within a New Zealand glacierized river basin. *River Research and*
712 *Applications*, 24(1), 68–89. <https://doi.org/10.1002/rra.1048>
- 713 Chiarle, M., Iannotti, S., Mortara, G., & Deline, P. (2007). Recent debris flow occurrences
714 associated with glaciers in the Alps. *Global and Planetary Change*, 56(1–2), 123–136.
715 <https://doi.org/10.1016/j.gloplacha.2006.07.003>
- 716 Cogley, J. G., Hock, R., Rasmussen, L. A., Arendt, A. A., Bauder, A., Braithwaite, R. J.,
717 Jansson, P., Kaser, G., Möller, M., Nicholson, L. I., & Zemp, M. (2011). *Glossary of Glacier*
718 *Mass Balance and Related Terms* (IHP-VII Technical Documents in Hydrology No. 86; IACS
719 Contribution No. 2, p. 114). UNESCO-IHP.
- 720 Denton, G. H. (1975). Conterminous US, Chapter 1. In W. O. Field (Ed.), *Mountain glaciers of*
721 *the Northern Hemisphere* (Vol. 1). Corps of Engineers, US Army, Technical Information
722 Analysis Center, Cold Regions Research and Engineering Laboratory.
- 723 DeVisser, M. H., & Fountain, A. G. (2015). A century of glacier change in the Wind River
724 Range, WY. *Geomorphology*, 232, 103–116. <https://doi.org/10.1016/j.geomorph.2014.10.017>
- 725 Dick, K. (2013). *Glacier Change of the North Cascades, Washington 1900-2009* [M.S.].
726 Portland State University.
- 727 Dussailant, I., Berthier, E., Brun, F., Masiokas, M., Hugonnet, R., Favier, V., Rabatel, A., Pitte,
728 P., & Ruiz, L. (2019). Two decades of glacier mass loss along the Andes. *Nature Geoscience*,
729 12(10), 802–808. <https://doi.org/10.1038/s41561-019-0432-5>
- 730 Earl, L., & Gardner, A. (2016). A satellite-derived glacier inventory for North Asia. *Annals of*
731 *Glaciology*, 57(71), 50–60. <https://doi.org/10.3189/2016AoG71A008>
- 732 Fagre, D. B., McKeon, L. A., Dick, K. A., & Fountain, A. G. (2017). *Glacier margin time series*
733 *(1966, 1998, 2005, 2015) of the named glaciers of Glacier National Park, MT, USA* [Data set].
734 U.S. Geological Survey. <https://doi.org/10.5066/f7p26wb1>



- 735 Fellman, J. B., Nagorski, S., Pyare, S., Vermilyea, A. W., Scott, D., & Hood, E. (2014). Stream
736 temperature response to variable glacier coverage in coastal watersheds of Southeast Alaska.
737 *Hydrological Processes*, 28(4), 2062–2073. <https://doi.org/10.1002/hyp.9742>
- 738 Fountain, A. G., & Glenn, B. (2022). *Data From: Inventory of Glaciers and Perennial*
739 *Snowfields of the Coterminous USA" (2022)* (Data Series No. 3; Geology Faculty Datasets).
740 Portland State University. <https://doi.org/10.15760/geology-data.03>
- 741 Fountain, A. G., Glenn, B., & Basagic, H. J. (2017). The Geography of Glaciers and Perennial
742 Snowfields in the American West. *Arctic, Antarctic, and Alpine Research*, 49(3), 391–410.
743 <https://doi.org/10.1657/AAAR0017-003>
- 744 Fountain, A. G., Hoffman, M. J., Jackson, K., Basagic, H. J., Nylén, T. H., & Percy, D. (2007).
745 *Digital outlines and the topography of the Glaciers of the American West* (Open File Report No.
746 2006–1340; p. 23). US Geological Survey. <https://doi.org/10.3133/ofr20061340>
- 747 Fountain, A. G., & Tangborn, W. V. (1985). The effect of glaciers on streamflow variations.
748 *Water Resources Research*, 21(4), 579–586.
- 749 Garwood, J. M., Fountain, A. G., Lindke, K. T., van Hattem, M. G., & Basagic, H. J. (2020).
750 20th Century Retreat and Recent Drought Accelerated Extinction of Mountain Glaciers and
751 Perennial Snowfields in the Trinity Alps, California. *Northwest Science*, 94(1), 44.
752 <https://doi.org/10.3955/046.094.0104>
- 753 Gesch, D., Oimoen, M., Greenlee, S., Nelson, C., Steuck, M., & Tyler, D. (2002). The national
754 elevation dataset. *Photogrammetric Engineering and Remote Sensing*, 68(1), 5–32.
- 755 Heard, J. (2000). *Late Pleistocene and Holocene Aged Glacial and Climatic Reconstructions in*
756 *the Goat Rocks Wilderness, Washington, United States*. <https://doi.org/10.15760/etd.557>
- 757 Hock, R., de Woul, M., Radić, V., & Dyurgerov, M. (2009). Mountain glaciers and ice caps
758 around Antarctica make a large sea-level rise contribution: MOUNTAIN GLACIERS AND ICE
759 CAPS. *Geophysical Research Letters*, 36(7), n/a-n/a. <https://doi.org/10.1029/2008GL037020>
- 760 Hoffman, M. J., Fountain, A. G., & Achuff, J. M. (2007). 20th-century variations in area of
761 cirque glaciers and glacierets, Rocky Mountain National Park, Rocky Mountains, Colorado,
762 USA. *Annals of Glaciology*, 46(1), 349–354. <https://doi.org/10.3189/172756407782871233>
- 763 Homer, C. G., Dewitz, J. A., Yang, L., Jin, S., Danielson, P., Xian, G., Coulston, J., Herold, N.
764 D., Wickham, J., & Megown, K. (2015). Completion of the 2011 National Land Cover Database
765 for the conterminous United States-Representing a decade of land cover change information.
766 *Photogramm. Eng. Remote Sens.*, 81(5), 345–354.
- 767 King, C. (1871). On the discovery of actual glaciers on the Mountains of the Pacific Slope.
768 *American Journal of Science and Arts*, 1(3), 157–167.



- 769 Krimmel, R. M. (2002). Glaciers of the conterminous United States. In R. S. Williams, Jr. & J.
770 Ferrigno (Eds.), *Satellite image atlas of glaciers of the world: North America* (pp. J329–J381).
771 US Geological Survey Professional Paper.
- 772 Meier, M. F. (1961). Distribution and variations of glaciers in the United States exclusive of
773 Alaska. *International Association of Scientific Hydrology*, 54, 420–429.
- 774 Meier, M. F. (1984). Contribution of small glaciers to global sea level. *Science*, 226(4681),
775 1418–1421.
- 776 Moore, R. D., Fleming, S. W., Menounos, B., Wheate, R., Fountain, A., Stahl, K., Holm, K., &
777 Jakob, M. (2009). Glacier change in western North America: Influences on hydrology,
778 geomorphic hazards and water quality. *Hydrological Processes*, 23(1), 42–61.
779 <https://doi.org/10.1002/hyp.7162>
- 780 O'Connor, J. E., Hardison, J. H., & Costa, J. E. (2001). *Debris flows from failures of Neoglacial-*
781 *age moraine dams in the Three Sisters and Mount Jefferson wilderness areas, Oregon: A study*
782 *of the recent debris flows from moraine-dammed lake releases at central Oregon Cascade Range*
783 *volcanoes - Mt. Jefferson, Three Fingered Jack and the Three Sisters / Broken Top*. US
784 Geological Survey.
- 785 Parkes, D., & Marzeion, B. (2018). Twentieth-century contribution to sea-level rise from
786 uncharted glaciers. *Nature*, 563(7732), 551–554. <https://doi.org/10.1038/s41586-018-0687-9>
- 787 Paul, F., Kääb, A., & Haeberli, W. (2007). Recent glacier changes in the Alps observed by
788 satellite: Consequences for future monitoring strategies. *Global and Planetary Change*, 56(1–2),
789 111–122. <https://doi.org/10.1016/j.gloplacha.2006.07.007>
- 790 Pfeffer, W. T., Arendt, A. A., Bliss, A., Bolch, T., Cogley, J. G., Gardner, A. S., Hagen, J.-O.,
791 Hock, R., Kaser, G., & Kienholz, C. (2014). The Randolph Glacier Inventory: A globally
792 complete inventory of glaciers. *Journal of Glaciology*, 60(221), 537–552.
793 <https://doi.org/10.3189/2014JG13J176>
- 794 Post, A., Richardson, D., Tangborn, W. V., & Rosselot, F. (1971). *Inventory of glaciers in the*
795 *North Cascades, Washington. (USGS Professional Paper 705-A)*. (p. 26). US Geological
796 Survey, 705-A.
- 797 Rabatel, A., Sirguey, P., Drolon, V., Maisongrande, P., Arnaud, Y., Berthier, E., Davaze, L.,
798 Dedieu, J.-P., & Dumont, M. (2017). Annual and Seasonal Glacier-Wide Surface Mass Balance
799 Quantified from Changes in Glacier Surface State: A Review on Existing Methods Using Optical
800 Satellite Imagery. *Remote Sensing*, 9(5), 507. <https://doi.org/10.3390/rs9050507>
- 801 Robinson, J. E. (2014). *Digital topographic data based on lidar survey of Mount Shasta*
802 *Volcano, California, July-September 2010* (Report No. 852; Data Series). USGS Publications
803 Warehouse. <https://doi.org/10.3133/ds852>
- 804 Russell, I. C. (1898). The glaciers of North America. *The Geographical Journal*, 12(6), 553–564.



- 805 Selkowitz, D. J., & Forster, R. R. (2016). Automated mapping of persistent ice and snow cover
806 across the western U.S. with Landsat. *ISPRS Journal of Photogrammetry and Remote Sensing*,
807 *117*, 126–140. <https://doi.org/10.1016/j.isprsjprs.2016.04.001>
- 808 Smiraglia, C., Azzoni, R. S., D'Agata, C., Maragno, D., Fugazza, D., & Diolaiuti, G. A. (2015).
809 The New Italian Glacier Inventory: A didactic tool for a better knowledge of the natural Alpine
810 environment. *J-Reading-Journal of Research and Didactics in Geography*, *4*(1). <http://www.j-reading.org/index.php/geography/article/view/92>
- 812 Spicer, R. (1986). *Glaciers in the Olympic Mountains, Washington—Present distribution and*
813 *recent variations*. [M.S.]. University of Washington.
- 814 Sun, M., Liu, S., Yao, X., Guo, W., & Xu, J. (2018). Glacier changes in the Qilian Mountains in
815 the past half-century: Based on the revised First and Second Chinese Glacier Inventory. *Journal*
816 *of Geographical Sciences*, *28*(2), 206–220. <https://doi.org/10.1007/s11442-018-1468-y>
- 817 Trcka, A. (2020). *Inventory of Rock Glaciers in the American West and Their Topography and*
818 *Climate*. <https://doi.org/10.15760/etd.7509>
- 819 U.S. Geological Survey, 2022. [https://www.usgs.gov/tools/geographic-names-information-](https://www.usgs.gov/tools/geographic-names-information-system-gnis)
820 [system-gnis](https://www.usgs.gov/tools/geographic-names-information-system-gnis). Accessed December 5, 2022
- 821 Usery, E. L., Varanka, D., & Finn, M. P. (2009). A 125 year history of topographic mapping and
822 GIS in the U.S. Geological Survey 1884-2009, part 1: 1884-1980. *ArcNews*, *31*(3), 1–1. USGS
823 Publications Warehouse.
- 824 Zemp, M., Huss, M., Thibert, E., Eckert, N., McNabb, R., Huber, J., Barandun, M., Machguth,
825 H., Nussbaumer, S. U., Gärtner-Roer, I., Thomson, L., Paul, F., Maussion, F., Kutuzov, S., &
826 Cogley, J. G. (2019). Global glacier mass changes and their contributions to sea-level rise from
827 1961 to 2016. *Nature*, *568*(7752), 382–386. <https://doi.org/10.1038/s41586-019-1071-0>
- 828