

# A global Lagrangian eddy dataset based on satellite altimetry

Tongya Liu<sup>1,2</sup> and Ryan Abernathey<sup>3</sup>

<sup>1</sup>State Key Laboratory of Satellite Ocean Environment Dynamics, Second Institute of Oceanography, Ministry of Natural Resources, Hangzhou, China

<sup>2</sup>Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai), Zhuhai, China

<sup>3</sup>Lamont-Doherty Earth Observatory, Columbia University, New York, NY, USA

**Correspondence:** Tongya Liu (liutongya@sio.org.cn)

## 1 **Abstract.**

2 The methods used to identify coherent ocean eddies are either Eulerian or Lagrangian in nature, and nearly all existing eddy  
3 datasets are based on the Eulerian method. In this study, millions of Lagrangian particles are advected by satellite-derived  
4 surface geostrophic velocities over the period of 1993–2019. Using the method of Lagrangian-averaged vorticity deviation  
5 (LAVD), we present a global Lagrangian eddy dataset (GLED v1.0, Liu and Abernathey, 2022, <https://doi.org/10.5281/zenodo.7349753>  
6 ). This open-source dataset contains not only the general features (eddy center position, equivalent radius, rotation property,  
7 etc.) of eddies with lifetimes of 30, 90, and 180 days, but also the trajectories of particles trapped by coherent eddies over the  
8 lifetime. We present the statistical features of Lagrangian eddies and compare them with those of the most widely used sea  
9 surface height (SSH) eddies, focusing on generation sites, size, and propagation speed. A remarkable feature is that Lagrangian  
10 eddies is generally smaller than SSH eddies, with a radius ratio of about 0.5. Also, the validation using Argo floats indicates  
11 that coherent eddies from GLED v1.0 exist in the real ocean and have the ability to transport water parcels. Our eddy dataset  
12 provides an additional option for oceanographers to understand the interaction between coherent eddies and other physical or  
13 biochemical processes in the Earth system.

## 14 **1 Introduction**

15 Mesoscale eddies, defined as rotating structures ranging typically from tens to hundreds of kilometers and lasting for several  
16 weeks to months, are ubiquitous in the global ocean (Fu et al., 2010; Chelton et al., 2011b, hereinafter CS11). And these eddies  
17 can trap, transport, and stir tracers such as heat, salt, and biochemical components in the ocean, thereby playing significant  
18 roles in nutrient distribution (Chelton et al., 2011a; Frenger et al., 2015), altering large-scale ocean circulation patterns (Aber-  
19 nathey and Marshall, 2013; Liu et al., 2022b), affecting marine ecosystems (Mahadevan, 2016; McGillicuddy Jr, 2016), and  
20 modulating climate variability (Busecke and Abernathey, 2019; Li et al., 2022b). Isolated mesoscale eddies in the ocean are  
21 generally considered as coherent structures with a material barrier that can trap the fluid within the eddy interior (Haller, 2015).  
22 Therefore, understanding the eddy structure and the degree of material transport by eddies are key issues for more accurate  
23 parameterization of mesoscale eddies in coarse-resolution marine ecosystem or climate models. To achieve this goal, herein  
24 we seek to produce a global coherent eddy dataset based on satellite observations.

25 Many methods have been proposed to identify mesoscale eddies from numerous oceanic databases such as satellite maps,  
26 numerical simulation products, and Argo floats. These existing methods generally fall into two categories: Eulerian and La-  
27 grangian (Haller, 2015; Abernathey and Haller, 2018). The core idea of Eulerian methods is to detect the eddy boundary based  
28 on certain physical or geometrical contours from the instantaneous flow field, and then track these boundaries at neighbor-  
29 ing times. Frequently used Eulerian eddy boundaries includes contours of Okubo-Weiss parameter, sea surface height (SSH;  
30 CS11), potential vorticity (Zhang et al., 2014), velocity streamlines (Nencioli et al., 2010), etc. By contrast, rather than depend-  
31 ing on instantaneous images, Lagrangian methods examine trajectories of water parcels over a finite time interval to identify  
32 the skeletons of coherent structures. Different techniques such as finite-time Lyapunov exponents (Shadden et al., 2005), finite-  
33 scale Lyapunov exponents (d’Ovidio et al., 2009), and Lagrangian-averaged vorticity deviation (LAVD; Haller et al., 2016)  
34 have been proposed for eddy detection. Both Eulerian and Lagrangian methods have advantages and disadvantages.

35 The most significant advantage of Eulerian methods is their operational simplicity: if continuous images of flow fields are  
36 available, then searching for eddy centers and boundaries becomes relatively straightforward once the dynamical criterion is  
37 determined. This feature means that Eulerian methods are used extensively, especially for SSH eddies (following geostrophic  
38 equilibrium) derived from the sea level anomaly (SLA). And the development of satellite observations facilitates eddy identifi-  
39 cation on a global scale. Using 16 years of altimetry maps with weekly intervals, the first mesoscale eddy dataset was produced  
40 (CS11) and the general features of mesoscale eddies were analyzed statistically. Later, Faghmous et al. (2015) presented a  
41 global SSH eddy dataset over the period of 1993–2014 using the daily altimetry product and a SLA-based method similar to  
42 that used in CS11. Until 2016, the eddy census of CS11 was updated routinely by a research team at Oregon State University,  
43 then in 2017 its operation was transferred to CLS/CNES, and it is now distributed by AVISO as the Mesoscale Eddy Trajectory  
44 Atlas (META). Several versions of this dataset—from META1.0exp to META3.1exp—are available to users, and Pegliasco et al.  
45 (2022) described the improvements from one release to the next. In addition, Dong et al. (2022) constructed a multi-parameter  
46 eddy dataset based on the velocity vector field from satellite observations. These Eulerian eddy datasets have been used widely  
47 to study the interaction between mesoscale eddies and other processes of the Earth system.

48 Mesoscale eddies are generally believed to be able to trap and transport the interior fluid when the nonlinearity parameter  
49  $U/c$  is greater than 1, where  $U$  is the azimuthal eddy speed and  $c$  is the eddy propagation speed. Statistics suggest that more  
50 than 90% of observed SSH eddies satisfy this criterion (CS11). By assuming no effective water exchange between the eddy  
51 interior and background flows, many studies have conducted estimates of heat, salt, and mass transports by Eulerian eddies  
52 on regional and global scales (Dong et al., 2014; Zhang et al., 2014; Frenger et al., 2015; He et al., 2018). Among them,  
53 the most appealing result shows that the westward zonal eddy mass transport in the subtropical gyre can reach 30–40 Sv,  
54 which is surprisingly comparable to the wind-driven gyre transport (Zhang et al., 2014). However, many recent works provide  
55 clear evidence that Eulerian methods strongly overestimate the degree of material transport by mesoscale eddies. Horizontally,  
56 observations and numerical simulations both suggest that Eulerian eddies are far from coherent structures because there is  
57 strong and persistent water exchange across the Eulerian eddy boundary (such as the SSH contour) during the eddy lifespan  
58 (Beron-Vera et al., 2013; Wang et al., 2016; Liu et al., 2019, 2022a). The contribution of coherent structures to the total eddy  
59 transport is very limited, and most eddy transport is induced by incoherent motions such as swirling and filamentation outside

60 the eddy cores (Wang et al., 2015; Abernathey and Haller, 2018; Zhang et al., 2019; Xia et al., 2022). In addition,  $U/c$  has  
61 been shown to be an ineffective indicator of eddy coherent transport because the leakage magnitude of initially trapped water is  
62 generally significant and does not depend on this parameter (Liu et al., 2022a). The overestimation of coherent eddy transport  
63 might be attributed to the common shortcomings of Eulerian methods (see discussion in Haller, 2015; Abernathey and Haller,  
64 2018). The essential issue is that Eulerian eddy boundaries detected at neighboring times do not necessarily trap the same fluid,  
65 and this can be rectified under the Lagrangian framework.

66 Lagrangian coherent structures have been identified successfully using different techniques. And these eddies can truly trap  
67 and transport materials for a certain distance without obvious leakage. However, few studies employ Lagrangian eddies to  
68 estimate eddy material transport for the following potential reasons. First, compared with the contour searching of Eulerian  
69 methods, Lagrangian algorithms are much more complicated for calculating some physical parameters (e.g., LAVD; details  
70 in Section 2) over a time interval. Second, flow fields with high spatial and temporal resolutions are needed to drive millions  
71 of Lagrangian particles, which brings huge calculation and storage pressures. Third, the definition method determines that  
72 Lagrangian eddies have a preset duration, rather than a free duration like Eulerian eddies, and identifying Lagrangian eddies  
73 with different lifetimes is also computationally expensive.

74 Recently, Abernathey and Haller (2018) used satellite-derived geostrophic velocities in the eastern Pacific to advect La-  
75 grangian particles, and they used the LAVD method to identify rotationally coherent Lagrangian vortices (RCLVs, also called  
76 Lagrangian eddies) over a period of 25 years, which is the first large-scale application of objective Lagrangian eddy detec-  
77 tion. Based on numerical model outputs, Xia et al. (2022) used the three-dimensional LAVD method to detect global coherent  
78 eddies, and they estimated the coherent transport across each latitude or longitude to be only about 1 Sv. Tian et al. (2022)  
79 also applied the LAVD method to global eddy detection and presented a 90-day RCLVs dataset, but they adopted a very tight  
80 threshold to define the eddy boundary (Tarshish et al., 2018), which would greatly underestimate the size of Lagrangian eddies  
81 (see Figure 4).

82 Nearly all public global eddy datasets are based on the Eulerian framework, and identifying coherent eddies is not an easy  
83 task. Therefore, it is necessary to develop a global Lagrangian eddy dataset based on observational data. So far, we have  
84 conducted a series of works towards this goal, including regional eddy identification (Abernathey and Haller, 2018; Liu et al.,  
85 2022a), parameter sensitivity tests (Tarshish et al., 2018), and numerical experiments (Sinha et al., 2019; Liu et al., 2019; Zhang  
86 et al., 2019). In this study, we extend the work of Abernathey and Haller (2018) to the global ocean to identify coherent eddies  
87 using the LAVD method, and we generate a Lagrangian eddy dataset based on altimetry observations. This dataset provides not  
88 only general features (eddy center position, equivalent radius, rotation property, etc.) of eddies with lifespans of 30, 90, and 180  
89 days but also the trajectory of particles trapped by coherent eddy boundaries over the lifetime, and to the best of our knowledge  
90 this is the first attempt at a public eddy dataset. Also, we compare this dataset to the latest SSH eddy dataset (META3.1exp)  
91 to understand the statistical differences between the two types of eddies. Our eddy dataset provides an additional option for  
92 oceanographers in studying the interactions between coherent eddies and other physical or biochemical processes.

93 Although some studies have revealed several vertical features of mesoscale eddies, such as regional variability (Zhang  
94 et al., 2013), surface and subsurface-intensified types (Dilmahamod et al., 2018), and eddy vertical tilt (Li et al., 2022a), our

95 understanding of the three-dimensional structure of mesoscale eddies is still limited due to the lack of subsurface observations.  
96 The prevailing assumption is that mesoscale eddies are approximately in geostrophic balance, so this study mainly concentrates  
97 on two-dimensional coherent eddies based on geostrophic currents. We will discuss how unsolved motions affect coherent  
98 eddies later. We encourage users of our product to be mindful of the limitations of the underlying satellite-derived geostrophic  
99 velocity fields used to derive our coherent eddies.

100 This paper is organized as follows. Section 2 presents the complete process of generating the global Lagrangian eddy dataset.  
101 Section 3 illustrates the basic information of the dataset, the statistical features of coherent eddies, the comparison with SSH  
102 eddies, and the dataset validation. Section 4 introduces the availability of the eddy dataset and related algorithms. Finally,  
103 Section 5 provides the discussion and conclusions.

## 104 **2 Generation of eddy dataset**

### 105 **2.1 Satellite altimetry**


106 Because observational data for the subsurface flow field are quite rare, we consider only two-dimensional coherent eddies from  
107 the near-surface geostrophic velocity field  $\mathbf{v}_g = (u, v)$  that can be derived according to the geostrophic relation

$$108 \quad \hat{\mathbf{k}} \times \mathbf{v}_g = -\frac{g}{f} \nabla \eta, \quad (1)$$

109 where  $g$  is the acceleration due to gravity,  $f$  is the Coriolis parameter,  $\hat{\mathbf{k}}$  is the unit vertical vector pointing upward, and  $\eta$  is  
110 the SSH. In this study, we use the satellite altimetry product (SEALEVEL\_GLO\_PHY\_L4\_REP\_OBSERVATIONS\_008\_047)  
111 distributed by the Copernicus Marine Environment Monitoring Service. This dataset merges along-track measurements from  
112 several altimeter missions and interpolates them to a  $1/4^\circ$  latitude-longitude grid. It provides daily variables including the  
113 SLA, the absolute dynamic topography (ADT, equivalent to SSH), and the precomputed geostrophic velocities based on (1).  
114 Note that velocities in the equatorial region (within  $\pm 5^\circ$ ) are estimated based on a higher-order vorticity balance (Lagerloef  
115 et al., 1999) since the geostrophy is not satisfied. We choose the time period of 27 years, from 1 January 1993 to 30 December  
116 2019. In addition, following the procedure described by Abernathey and Marshall (2013), a small correction to the geostrophic  
117 velocities is applied to eliminate the divergence due to the meridional change of  $f$  and to perform no-normal-flow boundary  
118 conditions at the coastlines. Compared with noncorrected flow fields, this correction has an insignificant effect on the coher-  
119 ent eddy identification in the open ocean (Abernathey and Haller, 2018). Although the geostrophic current is an incomplete  
120 representation of the full flow in the real ocean, it is by far the leading-order component at the scales of interest in this study.

### 121 **2.2 Particle advection**


122 The first step in generating the global Lagrangian eddy dataset is to advect particles using surface geostrophic velocities (Figure  
123 1). The satellite altimetry product with a  $1/4^\circ$  grid resolution can well resolve  $\sim 200$  km length structures in the equatorial  
124 region,  $\sim 50$  km length structures at the mid-latitudes, and  $\sim 25$  km length structures at high latitudes (Ballarotta et al., 2019).



**Figure 1.** Flowchart of eddy dataset generation based on satellite observations.

125 To reflect properly the fine structure of material transport barriers and Lagrangian eddies, it is necessary to employ an extremely  
 126 dense mesh of Lagrangian particles with higher resolution than the forcing velocity field (Haller et al., 2016; Abernathey and  
 127 Haller, 2018). However, we should not pursue high resolution particle excessively because of the consequent computational  
 128 and storage burdens. Sensitivity tests by Abernathey and Haller (2018) suggest that a particle spacing of  $1/32^\circ$  is necessary to  
 129 identify RCLVs accurately, and in the present study we use the same resolution and release Lagrangian particles over the global  
 130 ocean (between  $0^\circ$  and  $360^\circ$  longitude and  $80^\circ\text{S}$  and  $80^\circ\text{N}$  latitude; Figure 2a), a total of 39 848 999 points. To our knowledge,  
 131 this is the highest resolution to date for a Lagrangian particle mesh applied at global scale. Note that the points on land are  
 132 masked because they never move. It is important to note that using ultrahigh-resolution particles does not necessarily improve  
 133 the resolution of the flow field as geostrophic currents are inherently unable to resolve small-scale/high-frequency processes,  
 134 such as submesoscale flows, tides, and inertia-gravity waves. The real benefit is to avoid the discontinuous areas in the LAVD  
 135 fields induced by coarse particle seeding, which allows us to obtain the clear structure of mesoscale coherent eddies.

136 The MITgcm (Adcroft et al., 2018), an open-source ocean general circulation model, is used to solve the kinematic equation  
 137 for Lagrangian particles  $d\mathbf{X}/dt = \mathbf{u}$ , where  $\mathbf{X} = (X, Y)$  is the position vector and  $\mathbf{u}$  is a two-dimensional velocity field.  
 138 The model can typically operate in either online or offline mode. Here, we employ the offline mode in which the internal  
 139 dynamical kernel is turned off and velocity fields are read from preset files with a frequency of 1 day. The FLT package is  
 140 enabled to track Lagrangian particles via implementing fourth-order Runge-Kutta integration. Compared with other tools for



**Figure 2.** (a) Initial latitudes of released Lagrangian particles. (b) Zonal and (c) meridional displacements of particles (in degree) after 180 days.

141 particle tracking, MITgcm provides a convenient configuration for parallel computing on a high-performance cluster, making  
 142 the global calculation more efficient.

143 From January 1993 to June 2019, the Lagrangian particle mesh is initialized on the first day of every month, and these par-  
 144 ticles are advected forward for 180 days, amounting to 318 180-day runs in total. In the zonal direction, the periodic boundary  
 145 condition is used to allow particles crossing zero longitude. Figures 2b and 2c show zonal and meridional displacements of  
 146 particles in a random time interval, which clearly display some main currents (e.g. western boundary currents, zonal tropical  
 147 currents, and Antarctic Circumpolar Current) and eddy-like structures. In each model run, the relative vorticity is calculated  
 148 on the Eulerian grid and interpolated to Lagrangian particle positions. To reduce the storage pressure, the relative vorticity and  
 149 the particle trajectory are output every 10 days, with the total volume still exceeding 20 TB.

### 150 2.3 Lagrangian eddy identification

151 Consider a two-dimensional coherent eddy, all fluid parcels along its material boundary should have the same average angular  
 152 speed when rotating around the eddy core, which is analogous to solid body rotation. Based on this physical intuition, Haller  
 153 et al. (2016) proposed an objective vorticity-based method to identify the material boundary of a coherent eddy by searching  
 154 for the outermost closed contour of the LAVD. In a two-dimensional flow, given a finite time interval  $(t_0, t_1)$ , the LAVD is  
 155 defined as the average of the vorticity deviation along the Lagrangian particle trajectory, that is,

$$156 \text{LAVD}_{t_0}^{t_1}(x_0, y_0) = \frac{1}{t_1 - t_0} \int_{t_0}^{t_1} |\zeta'[X(x_0, y_0, t), Y(x_0, y_0, t), t]| dt, \quad (2)$$


157 where  $(X, Y)$  is the position for the particle released initially at point  $(x_0, y_0)$  and  $\zeta'$  is the instantaneous relative vorticity  
 158 deviation from the spatial average over the whole domain. The LAVD (always positive) examines the average magnitude of  
 159 local rotation for each Lagrangian particle over the time interval. A larger (smaller) LAVD value implies that the particle rotates  
 160 faster (slower), with the local maximum representing the eddy center and the eddy boundary being the outermost closed LAVD  
 161 curve encircling the center. This definition determines that all particles inside the boundary must rotate around the eddy core  
 162 during the time interval, which is essentially different from Eulerian methods based on instantaneous fields.

163 The algorithm employed for detecting RCLVs has been described in previous studies (Abernathey and Haller, 2018; Tarshish  
 164 et al., 2018; Liu et al., 2019; Zhang et al., 2019; Liu et al., 2022a). Once a local LAVD maximum is determined, we search  
 165 outward for closed LAVD curves. There might be multiple closed contours around a center, which are all objective options for  
 166 the Lagrangian eddy boundary that is expected to be a convex but allowing small deviations. To confine the boundary choice,  
 167 two parameters are introduced here: the convexity deficiency (CD, Haller et al., 2016) and the coherency index (CI, Tarshish  
 168 et al., 2018). The CD is defined as the ratio of the area difference between the contour and its convex hull to the total contour's  
 169 area (see Figure 7 in Tarshish et al., 2018), which means that the closer CD is to zero, the closer the eddy boundary is to being  
 170 a convex curve. The CI examines the change in spatial compactness of particles inside the contour over a time interval, which  
 171 is expressed as

$$172 \text{CI} = \frac{\sigma^2(t_0) - \sigma^2(t_1)}{\sigma^2(t_0)}, \quad (3)$$

173 where  $\sigma^2(t) = \langle |\mathbf{X}(t) - \langle \mathbf{X}(t) \rangle|^2 \rangle$ ,  $\langle \rangle$  indicates an average over all particles and  $||$  is the standard Euclidean distance.  
 174 Theoretically, the CI is less than 1 in value, and with decreasing CI, the eddy particle tends to rapidly disperse and develop  
 175 filaments. The RCLV boundary is determined when the outermost contour satisfies both the CD and CI thresholds.

176 In this study, the combination of  $CD < 0.1$  and  $CI > -1$  is adopted according to the sensitivity analysis by Tarshish et al.  
 177 (2018). Their results indicate that CD values of 0.01, 0.1, and 0.25 are three representative thresholds for strictly coherent,  
 178 moderately coherent, and leaky vortices, respectively, as is shown in Figure 3a and 3b. Although a small amount of filaments  
 179 exists, the RCLV defined by  $CD < 0.1$  can basically trap the initial water parcels and maintain the coherent structure over  
 180 the lifetime. It is clear that, the thresholds of 0.25 and 0.01 (adopted by Tian et al., 2022) will greatly overestimate and




**Figure 3.** (a) A random example of the 30-day LAVD field (color map) and three identified RCLV boundaries using three different CD values. The green dashed thick line is the eddy boundary defined by the SLA contour. (b) Initial and final particle positions trapped by three boundaries. The dashed black line is the eddy center trajectory and the black dot is the initial eddy center.

181 underestimate, respectively, the size of the coherent eddy. This parameter combination is a moderate threshold for identifying  
 182 coherent eddies and has been employed successfully in our previous studies (Liu et al., 2019, 2022a). In addition, we repeatedly  
 183 conduct the test of RCLV identification in the random regions and time periods, and as shown in Figure 4, the determined  
 184 parameters perform well in identifying RCLVs with lifetimes of 30 and 90 days.

185 Except for the ability to trap and transport tracers, one of the most significant differences between Eulerian and Lagrangian  
 186 eddies is the fact that the LAVD is defined over a specific, fixed finite time interval. Eulerian eddy tracking, in contrast, can  
 187 detect eddies of arbitrary lifetimes (of course, without any guarantee of material coherence). Computational pressure dictates  
 188 that it is impossible to release Lagrangian particles at any time and identify Lagrangian eddies with an open lifespan, and  
 189 to date there is no clear solution to reconcile this difference between the Eulerian and Lagrangian frameworks. In this study,  
 190 we choose three typical lifetimes to identify Lagrangian eddies, i.e., 30, 90, and 180 days. Coherent eddies with lifetimes  
 191 longer than 180 days are not considered because their number is quite limited based on our results (Figure 6) and those of  
 192 Abernathey and Haller (2018). (While eddies of different lifetimes in a specific location may overlap, we cannot say that they  
 193 are the “same” eddy because they will, in general, have different material boundaries.) After identifying boundaries for all  
 194 eddies over 27 years from 954 LAVD fields, the related eddy parameters (such as radius and movement speed) are calculated,  
 195 then we conduct quality control to discard eddies with a radius smaller than 25 km and to check that all the eddy parameters  
 196 fall within reasonable ranges. At this point, the Global Lagrangian Eddy Dataset (GLED v1.0, Liu and Abernathey, 2022,  
 197 <https://doi.org/10.5281/zenodo.7349753>) has been generated based on satellite observations.





**Figure 4.** (a) 30-day LAVD (in  $s^{-1}$ ) field in the northern hemisphere and 90-day LAVD field in the southern hemisphere calculating from 1 October 2016. (b) Identified boundaries (red contours) of 90-day RCLVs in the west of Australia. (c) Initial (red dots) and final (blue dots) positions of 90-day RCLVs, with black lines representing the eddy center trajectories. (d) Identified boundaries (red contours) of 30-day RCLVs in the Gulf Stream region. (e) Initial (red dots) and final (blue dots) positions of 30-day RCLVs, with black lines representing the eddy center trajectories.

## 198 3 Results

### 199 3.1 Description of eddy dataset

200 GLED v1.0 contains two components. First, the general features of coherent eddies are provided in the directory named  
 201 *eddyinfo*. The information about 30-day, 90-day, and 180-day eddies is stored separately in three JSON files, which contain the  
 202 following attributes:

- 203 • *id*: an eddy's unique ID composed by identification date, lifetime, and eddy number in the corresponding detection  
 204 interval;
- 205 • *date\_start*: generation date of the eddy;
- 206 • *duration*: eddy lifespan (in days);
- 207 • *radius*: equivalent radius (in kilometers) that is derived from the area enclosed by the eddy boundary;

- 208 • *cyc*: eddy rotation type (1 for anticyclonic, -1 for cyclonic);
- 209 • *center\_lon*, *center\_lat*: the longitude (in degrees North) and latitude (in degrees East) of the eddy center with a frequency  
210 of 10 days;
- 211 • *dx*, *dy*: zonal and meridional displacements (in kilometers) of the eddy over the eddy duration;
- 212 • *speed\_x*, *speed\_y*: averaged zonal and meridional propagation speeds (in meter per second) of the eddy, which equal the  
213 displacements divided by the eddy duration;
- 214 • *vort*: domain-averaged relative vorticity within the eddy boundary (in per second);
- 215 • *lavd*: domain-averaged LAVD value within the eddy boundary (in per second);


216 Researchers can filter the eddy data based on their studying regions, time periods, or other conditions. For example, if investi-  
217 gating the statistical behaviours of coherent eddies generated around the Kuroshio extension region ( $25 - 35^{\circ}\text{N}$ ,  $140 - 150^{\circ}\text{E}$ ),  
218 then 2445 30-day, 210 90-day, and 17 180-day eddies over 27 years will be selected for conducting the related analysis.

219 Second, the trajectories of all Lagrangian particles inside the eddy boundary are provided in the directory named *eddytraj*,  
220 which to the best of our knowledge is the first attempt at an open-source eddy dataset. We use an NC file with a three-  
221 dimensional array to store the particle positions every 10 days for each eddy, with the array dimensions being particle initial  
222 longitude, particle initial latitude, and time. Each NC file is named by its unique eddy ID, and the grid number of the two  
223 position dimensions is adjusted according to the eddy size. We randomly load six data records to show the particle positions  
224 during the eddy lifetime (Figure 5), and we find that these eddies all perform well in maintaining the coherent structure. An  
225 interesting phenomenon is that the eddy in Figure 5a is not initially located around a closed SLA contour, but a coherent  
226 structure does exist. This type of coherent eddies are all neglected when using the Eulerian method (Liu et al., 2019). Another  
227 typical feature is that the coherent eddy is much smaller than the outermost closed SLA contour (Figure 5b), indicating that this  
228 SSH eddy is highly leaky and far from a coherent structure. The second component of GLED v1.0 clearly displays the detailed  
229 process of material transport by coherent eddies, which is significant for understanding further the influence of coherent eddies  
230 in the distribution of oceanic tracers, especially some biogeochemical tracers such as chlorophyll (Gaube and McGillicuddy Jr,  
231 2017) and nutrients (Hughes and Miller, 2017).

### 232 3.2 General features of global coherent eddies

233 To assess GLED v1.0, in this subsection we calculate some statistics of global Lagrangian eddies and compare them with  
234 those of a new SSH eddy product (META3.1exp, publicly available at [https://www.aviso.altimetry.fr/en/data/products/value-  
235 added-products/global-mesoscale-eddy-trajectory-product.html](https://www.aviso.altimetry.fr/en/data/products/value-added-products/global-mesoscale-eddy-trajectory-product.html)). This dataset updates the detection algorithm and the tracking  
236 scheme, and changes the input sea level field from SLA to ADT (Pegliasco et al., 2022), but it is essentially the same as the  
237 eddy product proposed by CS11, falling into the Eulerian category.


238 From January 1993 to December 2019, META3.1exp provides 619 510, 166 426, and 44 329 SSH eddies with radii larger  
239 than 25 km and lifetimes longer than 30 days, 90 days, and 180 days, respectively. Our dataset contains many more short-lived



**Figure 5.** Positions of particles (colored dots) inside the eddy boundary every 10 days for randomly selected (a, d) 30-day, (b, e) 90-day, and (c, f) 180-day RCLVs. The unique eddy ID is marked in the title of each panel. Different colors denote the different times, and the blue dots represent the initial positions. The SLA fields are overlaid using black contours with solid lines for positive values and dashed lines for negative values.

240 but fewer long-lived coherent eddies, with the numbers of 30-, 90-, and 180-day RCLVs in GLED v1.0 being 1 095 356, 116  
 241 656, and 13 643, respectively. Census statistics of the numbers for RCLVs and SSH eddies originating in  $2^\circ \times 2^\circ$  grids over 27  
 242 years are shown in Figure 6. For RCLVs with the three lifetimes, the peak values of eddy number are generally located close to  
 243 the eastern boundaries of ocean basins, much higher than that in the western-boundary current regions (Figures 6a, 6c, and 6e).  
 244 This spatial feature is not in agreement with the previous analysis by CS11 and the pattern based on META3.1exp, which both  
 245 show SSH eddies to be distributed broadly in the mid-latitude regions between  $10^\circ\text{N/S}$  and  $60^\circ\text{N/S}$  with no obvious east-west  
 246 asymmetry (Figures 6b, 6d, and 6f). Compared with SSH eddies with lifetimes longer than 180 days that can be observed nearly  
 247 everywhere in the global ocean except for the tropics, the number of 180-day RCLVs is quite limited and they are concentrated  
 248 in the southwest of Australia and the interior ocean of the Atlantic.


249 To understand intuitively the differences between RCLVs and SSH eddies, we choose two regions—one in the northeast  
 250 Pacific and the other in the Antarctic Circumpolar Current (ACC)—to display the location and size features of eddies on a  
 251 random date (Figure 7). These two regions are selected because they represent weak and strong eddy kinetic energy (EKE)  
 252 scenarios. The global EKE map exhibits that the northeast Pacific is less energetic (Whalen et al., 2018) and is typically  
 253 considered as a "desert" of long-lived eddies (CS11), but numerous short-lived SSH eddies and RCLVs are distributed widely  
 254 (Figure 7a). The most noteworthy feature is that RCLVs are generally smaller in size than SSH eddies and not necessarily  
 255 enclosed by the SSH contour. Based on their relative positions to SSH eddies, RCLVs can be classified into two categories (Liu  
 256 et al., 2019): overlapping and non-overlapping. The latter are quite different from traditional geostrophic eddies and appears



**Figure 6.** The geographic distribution of eddy generation number in  $2^\circ \times 2^\circ$  grids for (a, c, e) RCLVs and (b, d, f) SSH eddies over 27 years. Three time intervals (30, 90, and 180 days) are considered and the grid without eddies is masked.

257 frequently, deserving further investigation of their structure and evolution. Another feature is that many RCLVs propagate  
 258 eastward over the lifespan in this region, which has not been noticed before. As one of the most energetic regions, the ACC  
 259 region is rich in SSH eddies with large radii and amplitudes (Figure 7b), but few of them have a coherent core, indicating that  
 260 these SSH eddies cannot maintain a coherent structure for as little as 30 days. We identify only 39 30-day RCLVs in region 2,  
 261 much fewer than the number (124) in region 1 with the same size. The reduced number of coherent eddies along the main path  
 262 of the jet-like current can also be seen clearly in the Gulf Stream and the Kuroshio Extension regions.

263 We now examine the statistics of eddy radius, zonal propagation speed, and meridional propagation speed for all RCLVs  
 264 and SSH eddies in  $10^\circ$  latitude bins, which are shown using the box plot in Figure 8. Outside of the tropical region, both types  
 265 of eddies basically decrease in size with latitude, reflecting the dependence of the Rossby deformation radius on the Coriolis  
 266 parameter (Chelton et al., 1998), but the averaged RCLV radius is only half of the SSH eddy radius, which is consistent with  
 267 the regional examples shown in Figure 7 and our previous analysis in the eastern Pacific (Abernathey and Haller, 2018). In  
 268 the tropics, the RCLV radius is only about 40 km because there are numerous non-overlapping RCLVs with small size (not  
 269 shown). In addition, it is observed that RCLVs and SSH eddies have similar westward propagation speeds, consistent with the




**Figure 7.** Locations of 30-day RCLVs (blue dots) and SSH eddies(>30 days, black contours) in (a) region 1 (blue box in Figure 6a) and (b) region 2 (red box in Figure 6a) on 1 January 2009. The red lines are the center trajectories of the RCLVs, and the color map represents the SLA field.

270 phase speed of long Rossby wave (Killworth et al., 1997), except for the tropical region where some RCLVs move eastward  
 271 with the background tropical flows. For the meridional propagation speed, its magnitude is usually lower an order than that of  
 272 the zonal speed, and both types of eddies have similar patterns, with the difference emerging in  $30^{\circ}\text{S}$ – $0^{\circ}$  where there are many  
 273 RCLVs along the eastern boundary (see Figure 6a).

### 274 3.3 Dataset validation

275 In this subsection, Argo floats are used to verify the existence of identified coherent eddies and their ability to trap water parcels.  
 276 Since its initial deployment in the early 2000s, the Argo profiling float array has expanded to encompass all the world's oceans  
 277 with more than 3000 active floats. The standard Argo float is designed to conduct a 10-day cycle, during which it measures  
 278 conductivity, temperature, and pressure at various depths. At the start of one cycle, the float sinks to a parking depth of 1000  
 279 m and drifts along with ocean currents for about 9 days. Then, it descends to a depth of 2000 m and rises to the surface  
 280 while continuously measuring the ocean's properties. Upon reaching the surface, the float transmits its data to satellites before  
 281 starting another cycle. The Argo data is available from the global data centers (<ftp://ftp.ifremer.fr/ifremer/argo>) and the position  
 282 information of floats from 2000 to 2019 is considered here.


283 Due to the horizontal structure of mesoscale eddies being basically independent of the depth (Zhang et al., 2013), despite  
 284 the eddy intensity having the vertical variability (Dilmahamad et al., 2018), it is believed that coherent eddies defined from



**Figure 8.** Statistics of (a) radius, (b) zonal propagation speed, and (c) meridional propagation speed for RCLVs and SSH eddies. The box plot shows statistics of all eddies in  $10^\circ$  bins. The box and the black whisker span the 25th to 75th and 10th to 90th percentiles of the distribution, respectively. The black line in the box indicates the median. The means of all eddies in a bin are shown using dashed lines, blue for RCLVs and red for SSH eddies.

285 surface geostrophic currents can maintain the coherent structure above a certain depth. Numerical simulations show that the  
 286 averaged depth of coherent eddies does not exceed 500 m (Xia et al., 2022). However, the composite analysis based on Argo  
 287 floats suggests that the density anomaly of mesoscale eddies in the west of Australia can penetrate deeper than 1000 m (see  
 288 Figure 2 in He et al., 2021), which surpasses the parking depth of Argo floats. And there are many long-lived RCLVs near this  
 289 area (Figure 6). Therefore, we search for Argo floats that are initially trapped by 90-day and 180-day RCLVs from GLED v1.0  
 290 in the southern Indian Ocean ( $20 - 60^\circ\text{S}$ ,  $45 - 145^\circ\text{E}$ , black box in Figure 6c) and examine if these floats can be carried for a  
 291 long range. A larger region than that in He et al. (2021) is used here in order to expand the sample size.

292 Figures 9a-9c show a case in which an Argo float is trapped by a 180-day RCLV for its entire lifespan. At the initial  
 293 time, the float is located close to the eddy center and moves westward along with the eddy for more than 400 km. We then  
 294 perform statistical analysis for 1001 90-day RCLVs and 270 180-day RCLVs. The calculation of the time-based distance



**Figure 9.** (a, b, c) A case showing that an Argo float is trapped by a 180-day RCLV (red dots), with the SLA field overlaid in black contours. The blue line tracks the eddy center's trajectory, while the yellow line tracks the Argo float's trajectory. (d, e) The distance between eddy centers and Argo floats that are initially trapped during the eddy lifetime for 90-day and 180-day RCLVs. Thin gray lines indicate the ensemble and the thick blue (red) line is the mean of 90-day (180-day) RCLVs. The green dashed line is the average radius of RCLVs. (f) The distribution probability of Argo floats with respect to the normalized eddy radius at the final time for 90-day (blue line) and 180-day (red line) RCLVs.

295 between the eddy centers and trapped Argo floats is carried out. Figures 9d and 9e show that the majority of the floats can be  
 296 continuously carried by RCLVs, with only a small portion escaping rapidly. The final distribution probability of Argo floats  
 297 within a specified distance normalized by eddy radius is calculated (Figure 9f). It is observed that for both 90-day and 180-day  
 298 RCLVs, approximately 70% of the floats are still located within 3 times the eddy radius. Given the vertical tilt of mesoscale  
 299 eddies (Li et al., 2022a), it is reasonable to conclude that these floats have been moving with coherent eddies since their initial  
 300 trapping. The reason that some Argo floats escape from RCLVs might be because these eddies do not reach the parking depth  
 301 of Argo floats. It is speculated that there would have been a stronger correlation if Argo floats were deployed at shallower  
 302 depths. In general, these findings suggest that coherent eddies from GLED v1.0 exist in the real ocean and have the ability to  
 303 transport water parcels.

#### 304 **4 Code and datasets availability**

305 Our dataset GLED v1.0 is available at <https://doi.org/10.5281/zenodo.7349753> (Liu and Abernathey, 2022). It is convenient  
306 to load the data using Python, Matlab, or other programming languages. Detailed examples for reading and analyzing the data  
307 using Python can be found in a GitHub repository (<https://github.com/liutongya/GLED>), in which we also provide the related  
308 algorithms to reproduce the generation of GLED v1.0. Users can apply these algorithms to regional or global identification of  
309 coherent eddies with different lifespans based on velocity fields from observations or numerical simulations.

#### 310 **5 Conclusions and discussion**

311 Methods employed to identify oceanic mesoscale eddies can be classified into Eulerian and Lagrangian frameworks, and nearly  
312 all public global eddy dataset are based on the Eulerian framework (e.g., CS11) because of its operational simplicity. Eulerian  
313 eddies are generally treated as coherent structures that can transport tracers such as heat, salt, and nutrients, and they have  
314 been used widely to evaluate the material transport by eddies (e.g., Zhang et al., 2014), but recent studies under the Lagrangian  
315 framework have provided clear evidence that (i) Eulerian eddies are far from being coherent studies and (ii) using Eulerian  
316 methods will greatly overestimate the degree of real coherent transport (e.g., Abernathey and Haller, 2018; Liu et al., 2019). To  
317 provide an additional option for oceanographers in studying mesoscale eddies, in this study, we proposed a global Lagrangian  
318 eddy dataset (GLED v1.0) based on satellite observations.

319 Millions of Lagrangian particles with a resolution of  $1/32^\circ$  were advected by satellite-derived surface geostrophic velocities  
320 for 180 days from the first day of every month over the period from January 1993 to June 2019. Using the LAVD method  
321 proposed by Haller et al. (2016), we identified coherent eddies (RCLVs) with lifetimes of 30, 90, and 180 days to generate  
322 GLED v1.0. This open-source dataset contains not only general features of coherent eddies (center position, equivalent radius,  
323 rotation property, etc.), but also the trajectories of particles trapped by coherent eddy boundaries over the lifetime. To the best  
324 of our knowledge, this is the first attempt to date to provide the position of Lagrangian particles advected by geostrophic flows  
325 in an eddy dataset.

326 We compared the statistical features of RCLVs in GLED v1.0 with those of SSH eddies in META3.1exp. Unlike SSH eddies  
327 that are broadly distributed in the global ocean basins, RCLVs tend to be generated close to the eastern boundaries, and the  
328 RCLV numbers along the main paths of western-boundary currents and the ACC are very limited. The zonal and meridional  
329 propagation speeds of RCLVs are found to be qualitatively similar to those of SSH eddies in most regions, but RCLVs are  
330 much smaller than SSH eddies with a radius ratio of about 0.5. In addition, we conducted the dataset validation based on Argo  
331 floats and found that about 70% of Argo floats that are initially trapped can always be carried by 90-day and 180-day RCLVs  
332 during the lifetime.

333 Although the size of coherent eddies is limited, it does not necessarily mean that the role of mesoscale eddies in the material  
334 transport is insignificant. It is speculated that the contribution of coherent structures to the overall eddy transport is small, and  
335 the incoherent motions such as stirring and filamentation on the periphery of mesoscale eddies might make a leading-order  
336 contribution (Hausmann and Czaja, 2012; Abernathey and Haller, 2018). Therefore, more attention is required to understand



337 material transport by the filamentary structures, and the global particle trajectories produced by this study might be useful for  
338 studying the motion behaviour outside coherent cores. In addition, given the huge difference in estimating the global eddy  
339 mass transport between Zhang et al. (2014) and Xia et al. (2022), it is worth conducting a more accurate estimate of the eddy  
340 coherent contribution based on our dataset and other observations.

341 Because of the computation and storage pressures, GLED v1.0 only provides RCLVs identified over three time intervals.  
342 And it is still unclear how to reconcile the difference between the free Eulerian lifetime and the fixed Lagrangian lifetime. In  
343 order to better satisfy the users' needs, as well as the eddy information in the dataset, we provide the related algorithms to  
344 reproduce our results completely, from driving Lagrangian particles to RCLV identification. Users should feel free to modify  
345 the configuration (e.g., the date of releasing particles and the identification time interval) according to their own research.

346 Although we have produced a useful eddy dataset under the Lagrangian framework, one should note that not all studies  
347 must use Lagrangian eddies. Eulerian eddies are still convenient and meaningful when the coherent structure is not the main  
348 concern. Researchers should select the suitable method and dataset based on their objectives. This present study offers relief  
349 from the dilemma that the Eulerian eddy dataset is nearly the only option for studying mesoscale eddies.

350 One limitation of the present dataset is that RCLVs are based on surface geostrophic velocities, which might introduce  
351 potential errors due to limited spatial and temporal sampling of satellite data. Lacorata et al. (2019) evaluated the Lagrangian  
352 simulations based on satellite-derived currents with respect to real drifter trajectories. They found that surface currents from  
353 satellite observations have overall good Lagrangian skills for large-scale transport and dispersion numerical simulations, but  
354 the finite-resolution flow field tends to underestimate relative dispersion at scales smaller than 100 km. The differences between  
355 simulated and real drifter trajectories might come from the fact that geostrophic currents fail to capture small-scale processes  
356 and vertical motions, such as submesoscale currents and inertia-gravity waves. Since the present study also used satellite data  
357 to drive particles, we did not expect to see better performance in the direct comparison between simulated particles and real  
358 near-surface drifters. This is the reason why we verify the accuracy of the dataset using Argo floats, whose motions are mainly  
359 determined by geostrophic flows. In addition, Sinha et al. (2019) investigated particle evolution driven by hourly-, daily-,  
360 and weekly-averaged velocities from a  $1/48^\circ$  numerical simulation, and several cases showed that small-scale/high-frequency  
361 motions from hourly and daily velocities can make the coherent structure identified from weekly velocities leaky and cause  
362 strong vertical motions of particles. Recent works highlighted the role of small motions in material transport, but the extent to  
363 which these motions affect coherent structures is still an open question.

364 Here, we propose several potential application scenarios of GLED v1.0. First, it can be used to understand the structure and  
365 physical dynamics around mesoscale coherent eddies, including their interactions with multi-scale oceanic and atmospheric  
366 processes. Second, it can be used to estimate the coherent eddy transport of heat, salt, and nutrients, which can provide more  
367 accurate parameterization in climate and ecosystem models. Third, it can be used to explore the behavior and distribution of  
368 marine organisms and how they are influenced by coherent eddies. In addition, we need to remind users to be careful when  
369 using the particle trajectory in regions where submesoscale processes are active.

370 Although limitations exist, the satellite-derived geostrophic flow field is still the only large-scale velocity observation that  
371 resolves mesoscale structures, and our dataset has been verified to be reasonable at the geostrophic scale. We see this study as

372 an important step toward fully understanding the features of mesoscale coherent structures, and we expect to update this dataset  
373 to version 2.0 once the observational data from the Surface Water and Ocean Topography mission become available. It would  
374 be quite meaningful to explore differences between the two versions, which will lead to new insights regarding multi-scale  
375 interactions and more accurate parameterization of eddy transport in numerical models.

376 *Video supplement.* The video supplement is available at <https://vimeo.com/773609039>.

377 *Author contributions.* RA proposed the idea and launched this project. TL and RA developed the related algorithm. TL conducted the offline  
378 particle advection and data analysis. TL organized the eddy dataset. TL and RA wrote the manuscript.

379 *Competing interests.* The authors declare that they have no conflict of interest.

380 *Disclaimer.* Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional  
381 affiliations.

382 *Acknowledgements.* This project has been supported by the National Natural Science Foundation of China (42106008, 42227901). We thank  
383 Nathaniel Tarshish, Anirban Sinha, Wenda Zhang, and Ci Zhang for their early involvement to push this project forward. We thank two  
384 anonymous reviewers for their helpful and constructive comments.

## 385 **References**

- 386 Abernathey, R. and Haller, G.: Transport by lagrangian vortices in the eastern pacific, *Journal of Physical Oceanography*, 48, 667–685, 2018.
- 387 Abernathey, R. P. and Marshall, J.: Global surface eddy diffusivities derived from satellite altimetry, *Journal of Geophysical Research: Oceans*, 118, 901–916, 2013.
- 389 Adcroft, A., Campin, J.-M., Doddridge, S. D., Evangelinos, C., Ferreira, D., Follows, M., Forget, G., Hill, H., Jahn, O., Klymak, J., et al.: MITgcm documentation, Release checkpoint67a-12-gbf23121, 19, 2018.
- 391 Ballarotta, M., Ubelmann, C., Pujol, M.-I., Taburet, G., Fournier, F., Legeais, J.-F., Faugère, Y., Delepouille, A., Chelton, D., Dibarboure, G., et al.: On the resolutions of ocean altimetry maps, *Ocean Science*, 15, 1091–1109, 2019.
- 393 Beron-Vera, F. J., Wang, Y., Olascoaga, M. J., Goni, G. J., and Haller, G.: Objective detection of oceanic eddies and the Agulhas leakage, *Journal of Physical Oceanography*, 43, 1426–1438, 2013.
- 395 Busecke, J. J. and Abernathey, R. P.: Ocean mesoscale mixing linked to climate variability, *Science Advances*, 5, eaav5014, 2019.
- 396 Chelton, D. B., DeSzoeki, R. A., Schlax, M. G., El Naggar, K., and Siwertz, N.: Geographical variability of the first baroclinic Rossby radius of deformation, *Journal of Physical Oceanography*, 28, 433–460, 1998.
- 398 Chelton, D. B., Gaube, P., Schlax, M. G., Early, J. J., and Samelson, R. M.: The influence of nonlinear mesoscale eddies on near-surface oceanic chlorophyll, *Science*, 334, 328–332, 2011a.
- 400 Chelton, D. B., Schlax, M. G., and Samelson, R. M.: Global observations of nonlinear mesoscale eddies, *Progress in oceanography*, 91, 167–216, 2011b.
- 402 Dilmahamod, A. F., Aguiar-González, B., Penven, P., Reason, C., De Ruijter, W., Malan, N., and Hermes, J.: SIDDIES corridor: a major east-west pathway of long-lived surface and subsurface eddies crossing the subtropical South Indian Ocean, *Journal of Geophysical Research: Oceans*, 123, 5406–5425, 2018.
- 405 Dong, C., McWilliams, J. C., Liu, Y., and Chen, D.: Global heat and salt transports by eddy movement, *Nature communications*, 5, 1–6, 2014.
- 407 Dong, C., Liu, L., Nencioli, F., Bethel, B. J., Liu, Y., Xu, G., Ma, J., Ji, J., Sun, W., Shan, H., et al.: The near-global ocean mesoscale eddy atmospheric-oceanic-biological interaction observational dataset, *Scientific Data*, 9, 1–13, 2022.
- 409 d’Ovidio, F., Isern-Fontanet, J., López, C., Hernández-García, E., and García-Ladona, E.: Comparison between Eulerian diagnostics and finite-size Lyapunov exponents computed from altimetry in the Algerian basin, *Deep Sea Research Part I: Oceanographic Research Papers*, 56, 15–31, 2009.
- 412 Faghmous, J. H., Frenger, I., Yao, Y., Warmka, R., Lindell, A., and Kumar, V.: A daily global mesoscale ocean eddy dataset from satellite altimetry, *Scientific data*, 2, 1–16, 2015.
- 414 Frenger, I., Münnich, M., Gruber, N., and Knutti, R.: Southern Ocean eddy phenomenology, *Journal of Geophysical Research: Oceans*, 120, 7413–7449, 2015.
- 416 Fu, L.-L., Chelton, D. B., Le Traon, P.-Y., and Morrow, R.: Eddy dynamics from satellite altimetry, *Oceanography*, 23, 14–25, 2010.
- 417 Gaube, P. and McGillicuddy Jr, D. J.: The influence of Gulf Stream eddies and meanders on near-surface chlorophyll, *Deep Sea Research Part I: Oceanographic Research Papers*, 122, 1–16, 2017.
- 419 Haller, G.: Lagrangian coherent structures, *Annual Review of Fluid Mechanics*, 47, 137–162, 2015.
- 420 Haller, G., Hadjighasem, A., Farazmand, M., and Huhn, F.: Defining coherent vortices objectively from the vorticity, *Journal of Fluid Mechanics*, 795, 136–173, 2016.

422 Hausmann, U. and Czaja, A.: The observed signature of mesoscale eddies in sea surface temperature and the associated heat transport, *Deep*  
423 *Sea Research Part I: Oceanographic Research Papers*, 70, 60–72, 2012.

424 He, Q., Zhan, H., Cai, S., He, Y., Huang, G., and Zhan, W.: A new assessment of mesoscale eddies in the South China Sea: Surface features,  
425 three-dimensional structures, and thermohaline transports, *Journal of Geophysical Research: Oceans*, 123, 4906–4929, 2018.

426 He, Y., Feng, M., Xie, J., He, Q., Liu, J., Xu, J., Chen, Z., Zhang, Y., and Cai, S.: Revisit the vertical structure of the eddies and eddy-induced  
427 transport in the Leeuwin Current system, *Journal of Geophysical Research: Oceans*, 126, e2020JC016556, 2021.

428 Hughes, C. W. and Miller, P. I.: Rapid water transport by long-lasting modon eddy pairs in the southern midlatitude oceans, *Geophysical*  
429 *Research Letters*, 44, 12–375, 2017.

430 Killworth, P. D., Chelton, D. B., and de Szoeke, R. A.: The speed of observed and theoretical long extratropical planetary waves, *Journal of*  
431 *Physical Oceanography*, 27, 1946–1966, 1997.

432 Lacorata, G., Corrado, R., Falcini, F., and Santoleri, R.: FSLE analysis and validation of Lagrangian simulations based on satellite-derived  
433 GlobCurrent velocity data, *Remote sensing of environment*, 221, 136–143, 2019.

434 Lagerloef, G. S., Mitchum, G. T., Lukas, R. B., and Niiler, P. P.: Tropical Pacific near-surface currents estimated from altimeter, wind, and  
435 drifter data, *Journal of Geophysical Research: Oceans*, 104, 23 313–23 326, 1999.

436 Li, H., Xu, F., and Wang, G.: Global mapping of mesoscale eddy vertical tilt, *Journal of Geophysical Research: Oceans*, p. e2022JC019131,  
437 2022a.

438 Li, J., Roughan, M., and Kerry, C.: Drivers of ocean warming in the western boundary currents of the Southern Hemisphere, *Nature Climate*  
439 *Change*, 12, 901–909, 2022b.

440 Liu, T. and Abernathey, R.: A global Lagrangian eddy dataset based on satellite altimetry (GLED v1.0),  
441 <https://doi.org/10.5281/zenodo.7349753>, 2022.

442 Liu, T., Abernathey, R., Sinha, A., and Chen, D.: Quantifying Eulerian eddy leakiness in an idealized model, *Journal of Geophysical Research:*  
443 *Oceans*, 124, 8869–8886, 2019.

444 Liu, T., He, Y., Zhai, X., and Liu, X.: Diagnostics of coherent eddy transport in the South China Sea based on satellite observations, *Remote*  
445 *Sensing*, 2022a.

446 Liu, T., Ou, H.-W., Liu, X., and Chen, D.: On the role of eddy mixing in the subtropical ocean circulation, *Frontiers in Marine Science*, 9,  
447 832 992, 2022b.

448 Mahadevan, A.: The impact of submesoscale physics on primary productivity of plankton, *Annual review of marine science*, 8, 161–184,  
449 2016.

450 McGillicuddy Jr, D. J.: Mechanisms of physical-biological-biogeochemical interaction at the oceanic mesoscale, *Annual Review of Marine*  
451 *Science*, 8, 125–159, 2016.

452 Nencioli, F., Dong, C., Dickey, T., Washburn, L., and McWilliams, J. C.: A vector geometry–based eddy detection algorithm and its appli-  
453 cation to a high-resolution numerical model product and high-frequency radar surface velocities in the Southern California Bight, *Journal*  
454 *of atmospheric and oceanic technology*, 27, 564–579, 2010.

455 Pegliasco, C., Delepouille, A., Mason, E., Morrow, R., Faugère, Y., and Dibarboure, G.: META3. 1exp: a new global mesoscale eddy trajectory  
456 atlas derived from altimetry, *Earth System Science Data*, 14, 1087–1107, 2022.

457 Shadden, S. C., Lekien, F., and Marsden, J. E.: Definition and properties of Lagrangian coherent structures from finite-time Lyapunov  
458 exponents in two-dimensional aperiodic flows, *Physica D: Nonlinear Phenomena*, 212, 271–304, 2005.

459 Sinha, A., Balwada, D., Tarshish, N., and Abernathey, R.: Modulation of lateral transport by submesoscale flows and inertia-gravity waves,  
460 *Journal of Advances in Modeling Earth Systems*, 11, 1039–1065, 2019.

461 Tarshish, N., Abernathey, R., Zhang, C., Dufour, C. O., Frenger, I., and Griffies, S. M.: Identifying Lagrangian coherent vortices in a  
462 mesoscale ocean model, *Ocean Modelling*, 130, 15–28, 2018.

463 Tian, F., Wang, M., Liu, X., He, Q., and Chen, G.: SLA-based orthogonal parallel detection of global rotationally coherent Lagrangian  
464 vortices, *Journal of Atmospheric and Oceanic Technology*, 2022.

465 Wang, Y., Olascoaga, M. J., and Beron-Vera, F. J.: Coherent water transport across the South Atlantic, *Geophysical Research Letters*, 42,  
466 4072–4079, 2015.

467 Wang, Y., Beron-Vera, F. J., and Olascoaga, M. J.: The life cycle of a coherent Lagrangian Agulhas ring, *Journal of Geophysical Research:*  
468 *Oceans*, 121, 3944–3954, 2016.

469 Whalen, C. B., MacKinnon, J. A., and Talley, L. D.: Large-scale impacts of the mesoscale environment on mixing from wind-driven internal  
470 waves, *Nature Geoscience*, 11, 842–847, 2018.

471 Xia, Q., Li, G., and Dong, C.: Global oceanic mass transport by coherent eddies, *Journal of Physical Oceanography*, 2022.

472 Zhang, W., Wolfe, C. L., and Abernathey, R.: Role of Coherent Eddies in Potential Vorticity Transport in Two-layer Quasigeostrophic  
473 Turbulence, arXiv preprint arXiv:1911.01520, 2019.

474 Zhang, Z., Zhang, Y., Wang, W., and Huang, R. X.: Universal structure of mesoscale eddies in the ocean, *Geophysical Research Letters*, 40,  
475 3677–3681, 2013.

476 Zhang, Z., Wang, W., and Qiu, B.: Oceanic mass transport by mesoscale eddies, *Science*, 345, 322–324, 2014.