

Long-term ash dispersal dataset of the Sakurajima Taisho eruption for ashfall disaster countermeasure

Haris Rahadiano^{1,2}, Hirokazu Tatano², Masato Iguchi³, Hiroshi L. Tanaka⁴, Tetsuya Takemi², Sudip Roy⁵

¹Graduate School of Informatics, Kyoto University, Kyoto, 606-8501, Japan

²Disaster Prevention Research Institute, Kyoto University, Uji, 611-0011, Japan

³Sakurajima Volcano Research Center, Disaster Prevention Research Institute, Kyoto University, Sakurajima, 851-1419, Japan

⁴Center for Computational Sciences, Division of Global Environmental Science, University of Tsukuba, Ibaraki, 305-8577, Japan

⁵Department of Computer Science and Engineering, Indian Institute of Technology Roorkee, Roorkee, Uttarakhand, 247-667, India

Correspondence to: Haris Rahadiano (haris.rahadiano.88n@st.kyoto-u.ac.jp)

Abstract. A large volcanic eruption can generate large amounts of ash which affect the socio-economic activities of surrounding areas, affecting airline transportation, socio-economics activities, and human health. Accumulated ashfall has devastating impacts on areas surrounding the volcano and in other regions, and eruption scale and weather conditions may escalate ashfall hazards to wider areas. It is crucial to discover places with a high probability of exposure to ashfall deposition. Here, as a reference for ashfall disaster countermeasures, we present a dataset containing the estimated distributions of the ashfall deposit and airborne ash concentration, obtained from a simulation of ash dispersal following a large-scale explosive volcanic eruption. We selected the Taisho (1914) eruption of the Sakurajima volcano, as our case study. This was the strongest eruption in Japan in the last century, and our study provides a baseline for a worst-case scenario. We employed One Eruption Scenario (OES) approach by replicating the actual event under various extended weather conditions to show how it would affect contemporary Japan. We generated an ash dispersal dataset by simulating the ash transport of the Taisho eruption scenario using a volcanic ash dispersal model and meteorological reanalysis data for 64 years (1958–2021). We explain the dataset production and provide the dataset in multiple formats for broader audiences. We examine the validity of the dataset, its limitations, and its uncertainties. Countermeasure strategies can be derived from this dataset, to reduce ashfall risk. The dataset is available at the DesignSafe-CI Data Depot: <https://www.designsafe-ci.org/data/browser/public/designsafe.storage.published/PRJ-2848v2> or through the DOI: <https://www.doi.org/10.17603/ds2-vw5f-t920> by selecting Version 2 (Rahadiano and Tatano, 2020).

1 Introduction

A volcanic eruption emits dangerous pollutants including tephra in the form of very fine ash. These affect human lives over vast areas (Bonadonna et al., 2021).

Deleted: per

Deleted: Term Ash ...sh Dispersal ...ispersal Dataset ...ata{ ...

Deleted: can affect the socio-economic activities of surrounding ...

Moved (insertion) [5]

Deleted: Therefore, i...t is crucial to discover places with a high ...

Deleted: of

Deleted: consideringfollowing a large-scale explosive volcanic ...

Moved up [5]: Accumulated ashfall has devastating impacts on both surrounding areas of the volcano and other regions, affecting airline transportation, socio-economics activities, and human health. Therefore, it is crucial to discover places with a high probability of exposure to ashfall deposition.

Deleted: This knowledge can help assess the additional risk in the infrastructures, human lives, and economic impacts to make a better volcanic eruption response plan. We generated the ...n ash disper ...

Deleted: a

Deleted: in order

Deleted: further reduce ashfall rs...sk. ...

Deleted: is one of the events that

Deleted: several dangerous pollutants including tephra in the fo ...

Deleted: . Large eruptions eject enormous amount of

Deleted: including tephra in the form of very fine ash

Deleted: and other eruptive materials into the air. In the form of very fine ash, tTephra hampers the residents near the volcano and citizens in the farther areas ... ADDIN ZOTERO_ITEM ...

Deleted: ¶

135 When a large volcanic eruption occurs, volcanic ashes can travel far from the volcano source, disrupting socio-economic activities in many ways. Fast-travelling ash endangers a larger population and significantly disrupts airline transportation (Folch et al., 2012; Peng and Peterson, 2012; Tanaka and Iguchi, 2019). Settled volcanic ash (i.e., ashfall) directly affects human health and livelihoods, destroys vegetation, crops, and pastures, and physically damages infrastructure, by clogging drainage systems, contaminating water supplies, disrupting traffic, and damaging vehicles on roads (Barsotti et al., 2010; Ayris and Delmelle, 2012; Wilson et al., 2012; Damby et al., 2013; Poulidis et al., 2018). The cumulative weight of ashfall on the roof can collapse buildings and short-circuit electricity inside them (Zuccaro et al., 2013; Hampton et al., 2015). Evidence from large eruption events (e.g., St. Helens in 1980, Mt. Pinatubo in 1991, Eyjafjallajökull in 2010) demonstrates the cataclysmic consequences of ashfall in wide areas (Tilling et al., 1990; Fero et al., 2008, 2009; Folch et al., 2012; Peng and Peterson, 2012). Ashfall hazards are uncertain, and depend on the eruption magnitude and intensity and the wind condition at the time of eruption. Because socio-economic activities have dynamically changing exposures to ashfall hazards, we believe that estimating the risk of finding effective countermeasures is critical.

145 Currently, densely populated and modernised cities require volcanic risk reduction strategies (Miyaji et al., 2011) to lessen the impacts of ashfall. Assessing the risks to infrastructure, human lives, and economic systems allows better response plans to be developed. Hazard and risk assessment for ashfall mostly relies on quantifying accumulated ashfall on the ground, then extending the analysis by combining appropriate social data (e.g., population, or building data, or others). Many researchers used this approach to determine the probability of impacts of future eruption events (Jenkins et al., 2014, 2015, 2018; Wilson et al., 2012; Biass et al., 2017). An ashfall risk assessment requires long-term weather data for statistical analysis and simulations of typical and exceptional conditions (Hattori et al., 2013, 2016). Long-term ashfall deposit data encompassing vast impact areas during an extended period is an important requisite for conducting such a study. This paper presents a generated dataset of ash dispersal over a vast region of Japan for 64 years (1958–2021). Using the Taisho eruption of the

155 Sakurajima volcano, the largest and the strongest eruption in Japan during the last century (Todde et al., 2017; Poulidis et al., 2018) as a case study, we simulated a volcanic ash transport and dispersion model (VATDM) to demonstrate an ash dispersal process over entire Japan. We adapt an OES strategy as a baseline for a worst-case scenario (Bonadonna, 2006; Barsotti et al., 2018). We used long-term weather reanalysis data to provide an extensive dataset to give input to volcanic disaster response plans.

160 Sakurajima volcano was selected as an area of interest because of its high explosivity and potential for large-scale explosive eruptions within the next 20–25 years, due to the recurrence of magma supply rates as similar phenomena that led to the last eruption (Yamasato et al., 2013; Hickey et al., 2016; Biass et al., 2017; Poulidis et al., 2018). To address such an urgent issue, the authorities have been preparing a disaster response plans utilising volcano hazard maps. These hazard maps contain information about the impacts of historical eruptions, potential locations of ash deposition in municipalities, and volcanic alerts and warnings, along with guidelines on evacuation directions and procedures (Kagoshima City, 2010; Kyushu Regional Development Bureau, 2017).

- Deleted:** happensoccurs, volcanic ashes could ...an travel far a (...)
- Moved (insertion) [23]**
- Deleted:** .. for example, damaging critical infrastructures and buildings, and causing health problems among the people (Wilson et al., 2012; Zuccaro et al., 2013)
- Deleted:** es
- Deleted:** are
- Deleted:** ing human health and livelihoods; ... destroying (...)
- Deleted:** In addition, ash plumes from the volcanoes are a safety hazard that can severely damage commercial airlines (Folch et al., 2012; Peng and Peterson, 2012; Tanaka and Iguchi, 2019).
- Moved (insertion) [24]**
- Deleted:** the latest large eruption events (e.g., St. Helens in 198 (...)
- Deleted:** The 2010 eruption of Eyjafjallajökull has amplifies the evidence. During this event, Volcanic volcanic ashes flew to thousands of kilometres away, forcing almost entire all airports in 25 countries to close for a week-long and costing a loss of billions of dollars in revenue for airlines (Folch et al., 2012; Peng and Peterson, 2012). Apart from the eruption scale, the weather conditions, mainly winds, primarily contribute to escalating ashfall catastrophes to a broader range. Ergo, aA
- Deleted:** Those events illustrates that aAshfall hazards have ... (...)
- Deleted:** based on
- Deleted:** due to (1)...nd depend on its ...he eruption magnitude (...)
- Deleted:** The presentCurrently, densely populated and moderni (...)
- Deleted:** to provide vital input to for the development of a more comprehensive ashfall risk assessment and emergency management
- Deleted:** case ...f the Sakurajima volcano, the largest and the (...)
- Deleted:** We selected We select
- Deleted:** was selected
- Deleted:** due tobecause of its high explosivity and its ...otential (...)
- Deleted:** ¶
As supporting evidence, three large eruptions in of Sakurajima volcano have occurred within the last six centuries: the Bunmei eruption in 1471, the An'ei eruption in 1779, and the Taisho eruption in 1914. The latter was considered the biggest largest and the strongest eruption in Japan during the last century (Todde et al., 2017; Poulidis et al., 2018). Furthermore, Hickey et al. (2016) showed that similar phenomena of magma supply rates that led to the Taisho eruption are currently happeningoccurring, suggesting (...)
- Deleted:** Addressing (...)
- Deleted:** Kagoshima Ccity municipal government...uthorities h (...)
- Deleted:** the
- Deleted:** risk management and evacuation plan by providing an (...)
- Deleted:** s necessary...information about the impacts of histori (...)
- Deleted:** with
- Deleted:** ndlong with a (...)

365 Recently, some studies assessed the impacts of continuous Vulcanian activities and scenarios for an explosive eruption of Sakurajima volcano, providing insights into how volcanic ash could disperse and affect livelihoods (Biasi et al., 2017; Poulidis et al., 2018). However, those studies only discussed the proximal impacts in short to medium terms, with no further analysis available for impacts on distal locations. Furthermore, previous large-scale explosive eruptions brought calamities across almost all of Japan. Economic losses amounted to billions because obstructed air transportation was one of the more
 370 significant repercussions. These losses were not realised by the authorities during the Taisho eruption event that occurred more than a century ago. Moreover, the Japanese airspace would suffer worse implications if a similar explosive eruption occurred in the near present. At least 40% of the air traffic in the Japanese airspace would be affected if Sakurajima volcano had a large-scale explosive eruption in the present day, almost four times higher than the number of people affected by the devastating Typhoon Jebi back in 2018 (Takebayashi et al., 2021). It is urgent to learn how large would be the catastrophic
 375 effect on contemporary Japan if the same event occurred in present times. Therefore, it is crucial to develop a risk analysis for a potentially broader range of ashfall, because when a future ashfall arrives in major socio-economic centres of Japan, such as Fukuoka, Osaka, and Tokyo, a large population will be exposed.
The primary aim of this study was to develop a dataset for input to an ashfall risk assessment for development of an emergency management plan. The dataset will be useful to simulate ashfall impacts from the Taisho eruption to
 380 contemporary Japan, understand the ash dispersal over an extensive region following a large-scale explosive eruption at Sakurajima volcano, and discover the condition that exacerbates the consequences. The size of the dataset, and its data on ground deposits and airborne concentrations, will enable users to prepare responses by creating conditional ashfall risk maps, enhancing early warning systems, planning for ash clean-up operations, and preparing for closures of air and land
 385 transportation and for long-term long-distance evacuation. This paper is structured as follows: Section 1 introduces the general background and motivation for this study. Section 2 presents the process and impacts of ashfall dispersal from the Taisho eruption in 1914, derived from previous studies and historical reports of the eruption. Section 3 describes the methodology for producing the dataset, and Section 4 describes the format and gives examples of usage of the dataset. Section 5 addresses the validation and limitations of the dataset. Finally, Section 6 concludes the paper. Unless otherwise
specified, all time dimensions in this study use the Japan Standard Time (JST, +09UTC).

390 2 Ashfall dispersal during the Taisho eruption

The Taisho eruption has been extensively studied by researchers all over the world, mainly because of the following three
factors:

1. Detailed historical reports in English were compiled and available shortly after the eruption (Omori, 1914; Koto, 1916).
2. The current unrest of the Sakurajima volcano (Iguchi, 2016; Iguchi et al., 2020; Poulidis et al., 2017, 2018).

Deleted:

Deleted: Furthermore, the Osumi Office of the Rivers and National Highways produced maps for the potential ash deposition from a large eruption case in municipalities surrounding the Sakurajima volcano (Kyushu Regional Development Bureau, 2017).

Deleted: Lately/Recently, some research ...studies has ...sessed (...)

Deleted:

Deleted: the ...revious large-scale explosive eruptions brought (...)

Deleted: s

Deleted: modern times regarding ashfall disaster countermeasures

Deleted: forto the development of ...n more comprehensive (...)

Deleted: Using the Taisho eruption case, we simulated the volcanic ash transport and dispersion model (VATDM) to demonstrate an ash dispersal process in large areas (entire Japan). We approached this issue by adapting one eruption scenario (OES) strategy, specifically as a baseline for the worst-case scenario (Bonadonna, 2006; Barsotti et al., 2018). We conducted this simulation in over an extended period (1958–2021, 64 years) with the a long-term weather reanalysis data to compensate for the incomplete wind profiles at the time of the eruption and to further validate the dataset.

Deleted: It

Deleted: Our study is an attempt

Deleted: characterise

Deleted: and

Deleted: variety of ...sh dispersal over an extensive region (...)

Deleted: . Furthermore, it is important to recognise the behaviour of ash transport, starting from discovering the condition of reproducing the actual ashfall distribution to finding

Deleted: led to much worse

Deleted: We produced this dataset for various purposes and usage, such as estimating the volcanic ashfall risk by analysing both the dispersal and deposition patterns of volcanic ashfall over a long period; developing the hazard maps for the disaster risk management process; and seeking a better comprehension of the tendency from of ashfall distribution from a large eruption over an large extensive region.

Deleted: Given (The size of the dataset, the available informati (...)

Deleted: ¶

Deleted: of for this study, followed by... Section 2, (...)

Deleted: Ddispersal during the Taisho Eruption (...)

Deleted: is one of the eruption events that has been extensively (...)

Deleted: The availability of dDetailed historical reports in (...)

Deleted: concurrent unrest activities (...)

3. The probability of a large-scale explosive eruption in the next decades that would be likely to resemble the past event (Hickey et al., 2016).

This section focuses on how volcanic ash spread across Japan, especially to distal locations, during the Taisho eruption. The complete evidence and chronology of the Taisho eruption are provided in the chronicles cited above and the studies conducted by Kobayashi (1982); Yasui et al. (2006, 2007); and Todde et al. (2017). Unless specified otherwise, the explanation here refers to the historical chronicles and the study by Todde et al. (2017). The Taisho eruption was a large-scale explosive eruption with a Volcanic Explosivity Index (VEI) (Newhall and Self, 1982) of four. An enormous volume of volcanic materials was ejected from the two main active vents during the explosive phases of the eruption, which began on 12 January 1914 10:00:00 and lasted 48 h. The tephra mostly ejected from the western vent (Nabeyama) and mixed with those from the eastern vent (Yokoyama), producing plumes estimated to be between 10 and 18 km high. The plume height, upper altitude westerlies, and surface winds influenced the pattern of ash dispersal, and were significant drivers of the ash deposition over a very wide area.

Most tephra dispersed eastward during the first several days of the eruption, leaving Kagoshima City with only minor ash deposits. In contrast, locations close to the eastern vent had ashfall deposit of four metres deep. Three days after the eruption started, ashfall deposits were reported in Kyushu, Shikoku, western Japan, and Sendai in northern Japan. Village offices, tobacco plantations, and local meteorological observatories kept a record of the exact time of ash arrival and sightings, showing that volcanic ash reached the major cities of Fukuoka at 08:00:00 and Osaka at midnight on 13 January 1914. Tokyo on the morning of 14 January 1914, and finally Sendai in the afternoon of the same day. Heavy ash fallout (approximately $1.5 \times 10^6 \text{ km}^2$) was reported at Ogasawara (Bonin Islands), approximately 1222 km away from the vent. Dispersal of very fine ashes during the first 13 h of eruption before it reached the climactic phase may have caused this phenomenon. Meanwhile, the activity of the eruption in the following days was largely minor in intensity, and ashfall continued within 200 km from the volcano until 19 January 1914.

Unfortunately, although an abundance of Japan-wide ashfall reports, no measurements of ashfall extent could be obtained. Most recent research focused on the proximal impacts, and there remains little analysis of the distal impacts. To the best of our knowledge, only one study attempted to draw all ashfall-affected areas completely. Dispersal maps at proximal regions have recently updated (Todde et al., 2017; Mita et al., 2018), but the map of wider exposure (Fig. 1) has not been modernised. The maps showing the eruption impacts in the southern Kyushu region contain complete information regarding the ground measurement of ash deposits but, the map for all Japan did not. The information provided in Fig. 1 is limited but vital, as at least the distance travelled by the ashfall is known.

Deleted: high probability of a large-scale explosive

Deleted: s

Deleted: paper section will ...ocues on how volcanic

Deleted: eruptive

Deleted: its the explosive phases of the eruption. ... The explos

Deleted: gigantic

Deleted: km and 18 km high. The plume height, upper altitude

Deleted: considerable

Deleted: plume height

Deleted: a

Deleted: the significant drivers of the vast ...sh deposition proc

Deleted:

Deleted: on the first several days of the eruption

Deleted: ¶

Deleted: the locations close to the eastern vent had ashfall depe

Deleted: there is an abundance of ashfall reports across ...apan

Field Code Changed

Formatted: No underline




Figure 1. Map of ash dispersal over all Japan following the Taisho eruption, (Omori, 1914).

3 Methods of data generation.

3.1 Simulation of selected eruption scenario

We used a fixed volcanological scenario over an extended contemporary period recorded in the meteorological reanalysis dataset to capture the extended daily variability of ash dispersal from the Taisho eruption. The PUFF model is an ash tracking model developed during the Redoubt volcano eruption in 1989, and is known for its high resolution and accuracy results (Tanaka, 1994; Searcy et al., 1998; Scollo et al., 2011; Folch, 2012). We ran the PUFF model for 23,376 days from 1 January 1958 to 31 December 2021. This model considers ash as a collection of a finite number of virtual particles and computes their motion over time. It works best within the first 48 h of the eruption (Searcy et al., 1998; Tanaka, 1994). The PUFF model is a Lagrangian-based model with several advantages over other (Eulerian) approaches. Eight of the nine Volcanic Ash Advisory Centers (VAACs) worldwide, including Tokyo VAAC, operate models based on the Lagrangian approach (Folch, 2012; Lin et al., 2012). A recent study used the PUFF model to simulate the explosive eruptions of Sakurajima volcano 2017 and 2018 to assess its utility in forecasting real-time ash fallout, and produced satisfactory results.

Moved (insertion) [25]


Formatted: Caption, Centered

Deleted: The ash dispersal over all Japan following map of

Moved down [6]: Todde et al. (2017) presented a simple and straightforward isochrones map derived from the chronicles in Fig.

Moved up [23]: Fast-travelling ash endangers a larger population and significantly disrupts airline transportation.

Deleted: 2



Deleted:

Moved (insertion) [6]

Deleted: The wide exposure map, shown in Fig. 12, did not get the same interest for modernisation compared to the dispersal maps at

Deleted: Fast-travelling ash endangers a larger population and significantly disrupts airline transportation. Evidence from the lat

Deleted: D

Deleted: Generation

Deleted:

Deleted: Methods

Deleted: methods

Deleted: on

Deleted: Selected selected Eruption ...ruption Scenario

Deleted: time period recorded in the meteorological reanalysis

Deleted: ¶

Deleted:

Deleted: We run the PUFF model for cases of 23376 days from 1 January 1958 to 31 December 2021.

Deleted: ¶

Deleted: ¶
The PUFF model is an ash tracking model developed during the

985 Simulations agreed well with measurements recorded by instruments at several points near the volcano (Tanaka and Iguchi, 2019). Here, we construct the model using many random variables $r_i(t)$, where $i = 1 \sim M$ and M is the total number of particles. $r_i(t) = (x, y, z)$ represents the position vector for the i^{th} particle at time t with its origin at the volcano vent. Using the discrete-time increment $\Delta t = 300$ s, the governing equation can be written as:

$$\begin{cases} r_i(0) = S_i, & i = 1 \sim M, \text{ for } t = 0, \\ r_i(t + \Delta t) = r_i(t) + V\Delta t + Z\Delta t + G\Delta t, & i = 1 \sim M, \text{ for } t > 0 \end{cases} \quad (1)$$

990 where S_i is the initial location of all the particles at the vent, $V = (u, v)$ is a vector for the wind velocity moving the particles and $Z = (c_h, c_h, c_v)$ is a vector for the diffusion velocity containing the diffusion speeds generated by Gaussian random numbers. $G = (0, 0, -w_t)$ is the gravitational fallout velocity obtained by approximating the extended Stokes law for various particle sizes (Tanaka, 1994; Tanaka and Yamamoto, 2002). The movement of particles steers the diffusion Z Z -direction, and the size of the particles affects the gravitational fallout G (Searcy et al., 1998; Tanaka, 1994). The diffusion speed c (c_h or c_v) was obtained by the random walk process related to the diffusion coefficient K as $c = \sqrt{2K/\Delta t}$ (Tanaka et al., 2016; Tanaka and Iguchi, 2019). Tanaka and Yamamoto (2002) conducted several diffusion tests with various values of K , and compared the results with satellite images of actual dispersals from several volcanic eruptions in the past. Based on this research, we assigned a suitable horizontal diffusion coefficient $K_h = 150 \text{ m}^2\text{s}^{-1}$ and vertical diffusion coefficient $K_v = 1.5 \text{ m}^2\text{s}^{-1}$ (Table 1).

1000 **Table 1. Input values for the simulations used in this study.**

Eruption Source Parameters	Value
Mean Estimated Eruption Mass ($\times 10^{10}\text{kg}$)	1.1
Mass Eruption Rate, ε (kg/h)	5.8×10^6 (Min) - 2.7×10^{10} (Max)
Horizontal Diffusion Coefficient, K_h (m/s)	150
Vertical Diffusion Coefficient, K_v (m/s)	1.5
Log-Scale Mean Grain Size (mm)	1.0

1005 These diffusion coefficients are consistent with the in situ observations documented by Eliasson et al. (2014) and adjusted for the Sakurajima volcano. The values used here follow those reported by Tanaka et al. (2016, 2019) but are smaller than those reported by Fero et al. (2008, 2009) and Kratzmann et al. (2010). To further investigate the ash dispersal process to distal locations in all Japan, we extended the simulation period of the PUFF model to 96 h after the eruption began. Following Eq. (1), by modelling the source location of the volcano vent as $S(x, y, z)$, we can adjust the number of particles released at each time step to obtain the optimal statistical information from the model.

Deleted: Both s

Deleted: the

Deleted: installed

Deleted: by

Deleted: ¶

Deleted: a large number of

Deleted: an

Deleted: is

Deleted: L

Deleted: diverse

Deleted: such

Deleted: the

Deleted: as mentioned in

Formatted: Caption

Formatted: Justified

Formatted Table

Formatted: Font: Not Bold

Formatted: Justified

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Justified

Formatted: Font: Not Bold

Formatted: Justified

Formatted: Font: Not Bold

Formatted: Justified

Deleted: ¶

Formatted: Font: Not Italic

Deleted: in this study

Deleted: previous studies

Deleted: the study conducted

Deleted: In order t

Deleted: entire

Deleted: ours

Deleted: started

Deleted: ¶

Deleted:

Deleted: draw

For each time step, we assigned a Gaussian random number for each particle (M_0) and scaled it according to the emission rate, making the value of the total number of particles (M) changes linearly with the emission rate (Tanaka et al., 2016). We assigned the initial number of particles to 5000 due to constraints in the available computational power. The size of volcanic ash varies from fine ash to boulders. Large particles tend to settle out quickly so airborne particle size decrease over time. Therefore, each ash particle will have a different grain size (Bonadonna et al., 1998), and we assume that each particle has an initial grain size following a logarithmic Gaussian distribution with a standard deviation of 2.0 centred at -3.0. Thus, the average particle size was approximately 1.0 mm on a log scale, with 68% of the particles within the range of 10 μm - 10 cm (Table 1). Given the initial vertical velocity of the emission with specified damping (e-folding) time τ_0 , the particles are distributed randomly in a vertical manner from the vent z_1 to the plume's peak z_2 continually. Then, using time integration for the vertical velocity in the momentum equation, we obtained the final form as $z(0) = z_1$ and $z(t) = z_2$ during time step Δt (Tanaka et al., 2016).




Figure 2. Plume height progression during the Taisho eruption on 12-13 January 1914 with information on both subsequent events and materials ejected. PF means pyroclastic flows that occurred from 13 January 1914 (Takebayashi et al., 2021).

3.2 Estimation of mass eruption rate based on plume height transitions

We used the estimated emission rate of Iguchi (2014), based on a previous study (Kobayashi, 1982). We consider the maximum estimated plume height as 17,890 m around midnight on 13 January 1914 which is close to the range of plume heights in the historical chronicles (Sect. 2; Fig. 2). We obtained the value of the mass eruption rate (ϵ) and the mean eruption mass (Table 1) with the help of temporal change in plume height. The obtained values agreed well with a recent observation (Todde et al., 2017). The pressure of a reservoir that ejects gas through a nozzle ($P(t)$) at time t decays exponentially with time (Nishimura, 1998) following Eq. (2):

$$P(t) = \Delta P_0 \left[1 + \frac{\gamma-1}{2} \left(\frac{2}{\gamma+1} \right)^{\frac{\gamma+1}{2(\gamma-1)}} v_0 \frac{A}{V_{\text{Vol}}} t \right]^{\frac{-2\gamma}{\gamma-1}} \quad (2)$$

Deleted:
Deleted: ¶
Deleted: For each time step, we assigned a Gaussian random ...
Deleted: .
Deleted: ¶
Deleted: 2.0
Deleted: 3.0.
Deleted: is approximately 1.0 mm on a log scale, with 68% of t ...
Deleted:
Deleted: by
Deleted: the
Deleted: ¶

Deleted: -
Deleted: ¶
Deleted: on
Deleted: Mass mass Eruption ...ruption Rate ...ate based o ...
Deleted: s
Deleted: Transition
Deleted: In this study, wWe used the estimated emission rate fi ...
Deleted: ion
Deleted: . 2; Fig. 2). The temporal change of the plume height ...
Deleted: see Table 1) with the help of temporal change in plum ...

where Vol is the volume of the reservoir, A is the cross-sectional area of the nozzle, v_0 is the initial ejection velocity, P_0 is the initial pressure, and γ is the specific heat ratio (values within the range 1.01–1.4). Because the Taisho eruption is a Plinian eruption, the plume height (H) is proportional to a quarter of the power of the heat discharge rate ($\frac{dQ}{dt}$) (Morton et al., 1956). Assuming that the volume of the reservoir is much larger than the ejected gas volume, the mass eruption rate (ϵ) is equivalent to the decay rate of the pressure ($\frac{dP}{dt}$), which can be estimated from the changes in plume height as follows:

$$H \propto \left(\frac{dQ}{dt}\right)^{1/4} \propto (\epsilon)^{1/4} = k \left(\frac{dP}{dt}\right)^{1/4} \quad (3)$$

3.3 Long-term wind field

Long-term weather data from meteorological reanalysis are necessary inputs to long-term ash transport simulation, as the wind strongly affects the sedimentary properties of volcanic ashfall (Hattori et al., 2013, 2016; Poulidis et al., 2018). One available long-period historical dataset is the JRA-55 reanalysis dataset, which includes the global meteorological reanalysis dataset from 1958 to the present, developed by the Japan Meteorological Agency (JMA) to study long-term variations in atmospheric and climate phenomena. The JRA-55 employs a reduced Gaussian horizontal grid system and applies a vertically conservative semi-Lagrangian advection scheme (Kobayashi et al., 2015). We utilised the JRA-55 data for the PUFF model, considering the horizontal wind grid to be $1.25^\circ \times 1.25^\circ$. We employed 16 vertical layers from 1000 to 10 hPa atmospheric pressure, and we used this data at four time points (00, 06, 12, and 18 UTC) daily. The 3D wind field consisted of zonal wind (U), meridional wind (V), and geo-potential height (gph), which were interpolated to the position of each ash particle in space and time using the cubic spline method.

The ash dispersal from the Taisho eruption was controlled by the westerlies (250 hPa) and surface winds (1000 hPa) (See Sect. 2). The most frequent directions at upper altitude winds are from W and SW, which are strongest during winter and weakest in summer. During winter and spring, winds mostly come from west, and the southwesterlies are more prominent in summer and autumn. Meanwhile, weaker winds dominate on the surface with distinct directions during cold seasons (autumn and winter), which blow from WSW-SW and NNE-NE direction, and warm seasons (spring and summer), where the winds are just southwesterly. Wind conditions depend on season (Fig. 3) so ash deposition depends on season of the eruption (Biass et al., 2017; Poulidis and Takemi, 2017; Poulidis et al., 2018).


3.4 Recording ashfall location and thickness

This study considers the ashfall deposits on the ground as particles with non-positive altitudes, with their location marked as a longitude-latitude pair. For each simulation, as the computation progressed, the particles with a negative value in the altitude dimension were subset to other files before being compiled when the simulation was completed. During this process, we captured the rest of the flowing particles as airborne ash concentrations, copied them to different files, and saved them using time marking every 90 min.

Deleted: W
Deleted: –
Deleted: Since
Deleted: i

Formatted: Justified, Line spacing: 1.5 lines

Deleted: ¶



Deleted: Term

Deleted: Wind

Deleted: Field

Deleted: In order to conduct the long-term ash transport

Deleted: the

Deleted: values

Deleted: .

Deleted: characteristics

Deleted: the

Deleted:

Deleted: of the

Deleted: (Kobayashi et al., 2015)

Deleted: . This data contain

Deleted: date


Deleted: a

Deleted: time


Deleted: hPa

Deleted: different

Deleted: in space and time



9



Deleted:

Formatted: Caption

Figure 3. Ten wind roses illustrating wind conditions inferred from the JRA-55 Reanalysis dataset (Kobavashi et al., 2015) for upper altitude winds (first column) and surface winds (second column), (a) and (f), 1958–2021; (b) and (g) winter, (c) and (h) spring, (d) and (i) summer, and (e) and (j) autumn. Roses indicate the probability of the provenance and sequential colours indicate the speed gradients.

We separated the measurement of the ashfall deposit and airborne ash concentrations into two files using this mechanism. Furthermore, we measured the ashfall thickness (ashfall depth) by dividing the surface area by the ashfall density according to its particle size. First, we allocated the ashfall particles to grids according to their location. Then, we set all ashfall particles in those grids according to their average values. We considered all the adjacent grids and the pre-assigned mass values multiplied by the number of particles to obtain the total mass of the ashfall deposit. It is possible to obtain the mass of each particle based on the particle size. However, because we deliberately designate the total initial number of particles as a constant number and smaller than the actual number in the real case, the total mass of all particles is less than the actual mass, and the difference may increase as the scale of the eruption increases. Therefore, considering the particle size under total mass conservation, the virtual mass of each particle is necessary (Shimbori et al., 2009). When the eruption reached its peak, each ash particle contained approximately 450 tons of virtual mass. This method alleviated the uncertainty that appears when assigning a small number of particles in the simulation (Scollo et al., 2011). After assigning the virtual mass to each particle in one grid, the thickness of the ashfall χ_{ij} at grid i, j obtained from the following equation:

$$\chi_{ij} = \sum_n \frac{\tilde{m}(D_n)}{\Delta x \Delta y} \quad (4)$$

where \tilde{m} is the total virtual mass for the total n particles within the grid i, j of area $\Delta x \Delta y$ corresponding to the particle size D_n (Shimbori et al., 2009). All computations (running the simulation, processing wind fields, and computing the total ashfall accumulation on the ground) were performed by a workstation with 32GB RAM, dual 16-cores Intel Xeon E5-2620V4 2.10 GHz processors, and 4GB NVIDIA Quadro P1000 GPU.

4. Dataset overview and usage examples

4.1 Data description and availability

The dataset contains three data series:

1. Ashfall deposit measurements from 128.5° E to 148.6° E, and 30.0° N to 45.9° N in 0.1°×0.1° grid (approximately 10 km²) (1958–2021).
2. Ashfall deposit measurements from 129° E to 132° E, and 30.5° N to 32.5° N in 0.01°×0.01° grid (approximately 1 km²) (1958–2021).
3. The time series of the total airborne ash concentrations in the air every 90 min (1958–2021).

To ensure maximum availability for diverse users of the dataset, we prepared it in two formats: a space-separated ASCII table in comma-separated value (CSV) format (Table 2) and a multi-dimensional array structure in the Network Common Data Format (NetCDF).

Formatted

Deleted: 4

Deleted: he ten wind roses illustrate...ng the ...ind conditio

Deleted: surface winds (first column) and surface winds

Deleted: The first row (

Deleted: b

Deleted:) portrays all wind conditions for the entire ...958--

Deleted: 0

Deleted: ,; while the rest shows the influence of seasonality.

Deleted: c) and (gd

Deleted: are the wind conditions in

Deleted: spring, (ce... and (f

Deleted: are the wind conditions in

Deleted: summer, (g...) and (h

Deleted: are the wind conditions in

Deleted: autumn, and (i

Deleted: are the wind conditions in

Deleted: winter

Deleted: correspond toindicate the probability of the

Deleted: 3.4 Recording Ashfall ashfall Location location and

Deleted:

Deleted: considering

Deleted: ¶

Deleted: get

Deleted: ¶

Deleted:

Deleted: since because we deliberately designate the total initia

Deleted: We assume that one ash particle contained 450 tons of

Deleted: would contained approximately 450 tons of virtual ma

Deleted:

Deleted:

Deleted: is donemethod to ...lleviated the uncertainty that

Field Code Changed

Formatted

Deleted: from running the simulation, processing wind fields, a

Deleted: Overview ...view and Availability

Formatted

Deleted: Each ash dispersal simulation produces

Deleted: (~(approximately 10 km²) for an extended period

Deleted: ~1 km²) for an extended period ...1958--

Deleted: collection of the total airborne ash concentrations in

Deleted: ¶

Deleted: the diverse users of the dataset, we prepared the

415 NetCDF is useful for supporting access to diverse types of scientific data, and its files are self-describing, network-
transparent, directly accessible, and extensible (Unidata, 2021). We developed the NetCDF files using the *Xarray* library in
Python using the NETCDF4 package (Hoyer and Hamman, 2017). Owing to the high dimensionality of the airborne ash
concentrations, we only provided these data in the NetCDF format. Ashfall deposit data are available in CSV and NetCDF
420 formats. The name of each file was set to the simulation date in ISO-8601 format (*YYYYMMDD<<.csv, .nc>>*), and files
were stored inside a directory named by year (*YYYY*) within each region (Japan and Kagoshima). Each dataset contains an
ordered set of location markings (longitude, latitude) and ashfall depth (in cm), which are stored in two decimal floating-
point formats. The location markings are the coordinates of the ashfall deposit in NetCDF data format. The airborne ash
concentrations are the total number of particles at a specific location and time. The tracking period of the ash particles began
90 min after the simulation and was completed after 96 h. As the three-dimensional shape of the affected area and the total
425 number of particles change throughout the time dimension, we assigned *date time* as the primary coordinate. The dataset
directory consists of two folders based on the data observations: *Airborne* and *Deposit*. The *Deposit* folder contains two child
directories based on the data format and observation range, which break down to every year from 1958 to 2021. The
Airborne dataset has a straightforward directory structure (year, *YYYY*) because it consists of one data format without a
separate observational range. Detailed structure of the dataset for both the ground deposits and airborne concentrations is
430 provided in the Appendix A.

Table 2: Tabular view of sample deposit data for two regions: Japan and Kagoshima. Each column is separated by a comma,
corresponding to CSV format.

<i>Deposit/CSV/Japan/<YYYY>/<YYYYMMDD.csv></i>			<i>Deposit/CSV/Kagoshima/<YYYY>/<YYYYMMDD.csv></i>		
latitude	longitude	deposit	latitude	longitude	deposit
30.00	128.50	0.00	30.00	129.00	0.00
30.00	128.60	0.10	30.00	129.01	0.10
...
30.10	128.50	0.20	30.01	129.00	0.20
30.10	128.60	0.30	30.01	129.01	0.30
<i>%.2f</i>	<i>%.2f</i>	<i>%.2f</i>	<i>%.2f</i>	<i>%.2f</i>	<i>%.2f</i>

435 The dataset is available at the DesignSafe-CI Data Depot (<https://www.designsafe-ci.org>) hosted at the Texas Advanced
Computing Center (TACC). This data depot is provided by the Natural Hazards Engineering Research Infrastructure
(NHERI), which provides the natural hazards engineering community and researchers with state-of-the-art
cyberinfrastructure (Rathje et al., 2017).

Deleted:

Deleted: Meanwhile, the a

Deleted: We set t

Deleted: dataset

Deleted: respective

Deleted: In addition, e

Deleted: set of

Deleted: is

Deleted: in the NetCDF data format

Deleted: data

Deleted: In contrast to the deposit data, t

Deleted: collection of

Deleted: provides

Deleted: started

Deleted: uses

Formatted: Caption

Deleted: he t

Deleted: the

Deleted: both

Deleted: the

Formatted Table

Deleted: In order tTo ensure maximum availability for the diverse users of the dataset, we prepared the dataset in two formats. We stored the dataset as a space-separated ASCII value table in comma-separated value (CSV) format (see Table 2) and a multi-dimensional array structure in the Network Common Data Format (NetCDF). NetCDF is useful for supporting access to diverse kinds types of scientific data, and its files are self-describing, network-transparent, directly accessible, and extensible (Unidata, 2021). We developed the NetCDF files using the *xarray Xarray* library in Python using the NETCDF4 package (Hoyer and Hamman, 2017). Due Owing to the high dimensionality of the airborne ash concentrations, we only provided these data in the NetCDF format. Meanwhile, the ashfall deposit data are available in CSV and NetCDF formats. We set the name of each dataset file as the simulation date in ISO-8601 format (*YYYYMMDD<<.csv, .nc>>*), stored inside a directory ...

Deleted:

Deleted: ¶

Deleted:

Deleted: As the three-dimensional shape of the affected area an ...

Deleted: ¶

Deleted:

Deleted: on

Deleted: in

Deleted: comprehensive

The dataset can be accessed directly at <https://www.designsafe-ci.org/data/browser/public/designsafe.storage.published/PRJ-2848v2> or through the DOI: <https://www.doi.org/10.17603/ds2-vw5f-t920> (Rahadiano and Tatano, 2020). When accessing the dataset from the DOI, the link will direct to Version 1 instead of Version 2; please select an appropriate version for the latest updates. Users can access the dataset without prior registration and may choose any data to download. Data can be downloaded as a single file or a collection of multiple files.

4.2 Usage examples of the dataset

The dataset comprises ash deposits on the ground and airborne concentrations of a typical large-scale eruption at Sakurajima volcano (Sect. 2) for an extended period. The data can be used to prepare a response for an ashfall disaster. Here, we demonstrate how to use these data to create conditional ashfall hazard maps. The information in such maps serve as a reference for designing an early warning system, planning ash clean-up operations, managing closures of on air and land transportation, and preparing long-term long-distance evacuation strategies. The maps describe the probability distribution of exceeding a selected hazard threshold based on the statistical distribution of the wind profiles (Bonadonna, 2006).




Figure 4. Probability distribution of ash dispersal with the threshold for airport closure risk (≥ 0.02 cm). Airport locations are marked by black aeroplane icons and the busiest airports by larger red icons. The colour gradient shows the probability of ash deposits exceeding the threshold. Human geography base (ESRI, 2022) provided the basemap for this figure.

- Deleted:
- Deleted: the old version (
- Deleted:)
- Deleted: the new version (
- Deleted:).
- Deleted: Therefore,
- Deleted: V
- Deleted: 2 of the dataset
- Deleted: directly
- Deleted: any
- Deleted: can
- Deleted: The d
- Deleted: as
- Deleted: by selecting
- Deleted: at once
- Deleted: Please select the Version 2 of the dataset for the latest (...)
- Commented [HR1]: Added section explaining the usage of th (...)
- Deleted: ¶
- Formatted (...)
- Deleted: in
- Deleted: see
- Deleted: Given the size of the dataset, the users can utilise t
- Deleted: make
- Deleted: preparation
- Deleted: can
- Deleted: and closure
- Deleted: ement
- Deleted: even
- Deleted: for
- Moved (insertion) [26]
- Deleted: The maps describe the probability distribution of (...)
- Formatted (...)
- Deleted: 5
- Deleted: The p
- Deleted: on the map
- Deleted: shown larger in
- Deleted: the difference in
- Deleted: is used as the basemap for this figure
- Deleted: is used as
- Deleted: The information in such maps can become serve as (...)





Figure 5. Seasonal probability distribution of ash dispersal with the threshold for airport closure risk (≥ 0.02 cm). The panels show the probability distribution of ash dispersal during (a) winter, (b) summer, (c) spring and (d) autumn. Airport locations are marked by black aeroplane icons and the busiest airports by larger red icons. The colour gradient shows the difference in the probability of ash deposits exceeding the threshold. Human geography base (ESRI, 2022) provided the basemap for this figure.

Formatted: Caption

Moved up [26]: The maps describe the probability distribution of exceeding a selected hazard threshold based on the statistical distribution of the wind profiles (Bonadonna, 2006).


Deleted: Several researchers used such maps to illustrate hazard assessments of a specific volcanic eruption scenario, identify the possible impacts of the assumed maximum expected events, and quantify the probability of first-order impacts on built environments (Bonadonna, 2006; Biass et al., 2014, 2016).

Several researchers used such maps to illustrate hazard assessments of a specific volcanic eruption scenario, identify the possible impacts of the assumed maximum expected events, and quantify the probability of first-order impacts on built environments (Bonadonna, 2006; Biass et al., 2014, 2016). One example is a conditional ashfall hazard map showing airport closure risk of all of Japan for all the wind profiles used in the simulation (Fig. 4). The threshold, or hazardous limit, of all hazard maps in this subsection is defined the Cabinet Office's Working Group on Wide-Area Ashfall Countermeasure at the time of Large Eruptions, Government of Japan (2018). We used a 0.02-cm threshold for ash on the airport's runway that would disrupt the take-offs and landing sequence of aeroplanes (Fig. 4 and Fig. 5). The busiest airports in Japan are Fukuoka, Kansai, Haneda, and Narita (Ministry of Land, Infrastructure, and Tourism, 2022).



- Deleted:** Figure 4 reveals the
- Deleted:** account
- Deleted:** for the
- Deleted:** f
- Deleted:** the
- Deleted:** , and all airport locations are obtained from the
- Moved down [27]:** Ministry of Land, Infrastructure, and Tourism (2022).
- Deleted:** In Fig. 4 and Fig. 5, w
- Deleted:** W
- Deleted:** d
- Deleted:**
- Deleted:** as the hazardous limit
- Deleted:** ,
- Deleted:** as defined by the Cabinet Office's Working Group on Wide-Area Ashfall Countermeasure at the time of Large Eruptions, Government of Japan (2018). A
- Deleted:** and mark all airport locations are marked bywith aeroplane icons on the map in black, and t
- Deleted:** (
- Deleted:**) are shown larger in red icons
- Deleted:** (Ministry of Land, Infrastructure, and Tourism, 2022)
- Moved (insertion) [27]**
- Deleted:** Ministry of Land, Infrastructure, and Tourism (2022)
- Field Code Changed**
- Formatted:** No underline
- Deleted:** Airport Management Status,

Figure 6. Probability distribution of ash dispersal with the threshold for road closure risk (≥ 3 cm). Black and white stripes mark railways, and highways and other roads are coloured lines. The colour gradient shows the probability of ash deposits exceeding the threshold. Human geography base (ESRI, 2022) provided the basemap for this figure.



Formatted: Caption

Deleted: .

Deleted: The figure illustrates how In average, a large-scale eruption in Sakurajima volcano would affect a lot of airports from the south to the east, with the busiest airports would have more than a 5% probability of closure after an eruption., Such disruptions possibly displacing will displace a numerous passengers and inflicting tremendous economic losses. The dataset can also be used to portray more prevalent information by showing the seasonal inclination on the airport closure risk, as shown in Fig. 56. For example, a large eruption during summer would bring the greatest impact in terms of the number of airports, and an eruption at autumn would disrupt a high number of traffics of Japan's busiest airports. Furthermore, the airborne concentrations data can further derive an further extended analysis on the timing of ash arrival in some airports that would present knowledge on how to do protection or even an evacuation of aircraft if there are sufficient time windows...

Formatted: Caption

Deleted:

Figure 7. Seasonal probability distribution of ash dispersal with the threshold for road closure risk (≥ 3 cm). The panels show the probability distribution of ash dispersal during (a) winter, (b) summer, (c) spring and (d) autumn. Black and white stripes mark railways on the map, and highways and other roads are coloured lines. The colour gradient shows the probability of ash deposits exceeding the threshold. Human geography base (ESRI, 2022) provided the basemap for this figure.

In average, a large-scale eruption in Sakurajima volcano would affect many airports from south to east, with the busiest airports having more than 5% probability of closure after an eruption, possibly displacing numerous passengers and inflicting economic losses. The dataset can also be used to portray the seasonal variations in the airport closure risk (Fig. 5). For example, a large eruption during summer would have greatest on the number of airports, and an eruption during autumn would also disrupt traffic at Japan's busiest airports. Further analysis of the airborne concentration data can inform the timing of ash arrival in some airports that would inform how to protect or evacuate aircraft if there was sufficient time. The variety of geographical dimensions offered by the dataset also allows investigation of the immediate consequences of ground ashfall deposits urban areas surrounding the Sakurajima volcano. Analogous conditional ashfall hazard maps can be made for the risk of disruption to land transportation network (Fig. 6 and Fig. 7). We used a 3-cm threshold as the hazard limit for closing roads and railways, when they would be impassable by cars and trains. The figures illustrate how a large-scale eruption of Sakurajima volcano could severely affect the eastern parts of Kagoshima City (located downwind of the volcano) with more than 25% probability, regardless of eruption time. The data also show how the seasonal patterns determine the extent of the ash deposits in the western parts of South Kyushu, with a large-scale eruption in summer would be escalating the damage to almost the entire Kagoshima Prefecture. The projected intensity of the ashfall hazards to the transportation networks provide insights for preparing responses and countermeasures. These preparations include scheduling roads and highway blockages, designing evacuation strategies, and managing the ash cleaning process shortly after the eruption.

5 Validation and Limitations

5.1 Validation

Simulation results have been validated using methods that include comparison to post-event reports, matching the ash cloud trajectory with satellite images, and adjusting the model's input to find the best-fit values (Webley and Mastin, 2009; Folch, 2012); for example, PUFF model simulations of the 1980 eruption of St. Helens, 1991 eruption of Hudson volcano, 1991 eruption of Mt. Pinatubo in the Philippines, 2001 eruption of Mt. Etna in Italy, 2006 eruption of Mt. Augustine, and all eruptions of North Pacific volcanoes from 1970–2010 (Fero et al., 2008, 2009; Kratzmann et al., 2010; Webley et al., 2008; Scollo et al., 2011; Webley et al., 2012). Several past studies validated ash dispersal simulations using the PUFF model for smaller-scale eruption of the Sakurajima volcano (Tanaka and Iguchi, 2019; Tanaka et al., 2020). The PUFF model has never been validated for historical large-scale eruptions of Sakurajima volcano, with its great plume height, eruption volume, and range of ash deposition. Ideally, if precise wind data at the time of the selected eruption are available, we can replicate the eruption; therefore, simulation results with similar wind patterns to the eruption would resemble the historical reports. The dataset presented here offered an opportunity to validate the simulation of ash dispersal from a large-scale eruption of

Formatted: Caption

Deleted: 8

Deleted: in coloured lines. The colour gradient shows the

Deleted: (ESRI, 2022)

Deleted: The dataset can also be used to portray more prevalent information by showing the seasonal inclination on airport closure risk, as portrayed shown in Fig. 6. For example, a large eruption during summer would bring the greatest impact in terms of the number of airports, and an eruption at autumn would disrupt a high number of traffics of Japan's busiest airports. Furthermore, the airborne concentrations data can derive further extended analysis on the timing of ash arrival in some airports that would present knowledge on how to do protection or even an evacuation of aircraft if there are sufficient time windows. The variety of geographical dimensions offered by the dataset also expands the opportunity to investigate the immediate consequences of ground ashfall deposits in the proximal urban areas surrounding Sakurajima volcano.¶

Figure 7 displays the conditional ashfall hazard maps accounting for the airport closure risk ofn entireall of Japan for all the wind profiles used in the simulation. We use a 3 cm threshold as the hazardous limit for the closures of roads and railways, as it would make them impassable by cars and trains, as defined by the Cabinet Office's Working Group on Wide-Area Ashfall Countermeasure at the time of Large Eruptions, Government of Japan (2018). The figure illustrates how a large-scale eruption in Sakurajima volcano can severely affect the eastern parts of Kagoshima Ceity, located downwind of the volcano regardless of eruption time, with more than 25% probability, regardless of eruption time. The data also shows how the seasonal patterns determine the extent of the ash deposits in the western parts of South Kyushu, with a summer eruption would escalating the damages to almost the entire Kagoshima Pprefecture, as depicted in Figure. 8. ¶

Knowing tThe intensity of the ashfall hazards to the transportation networks can giveprovide necessary insights tofor preparing responses and countermeasures. These preparations, which include scheduling roads and highway blockages, making evacuation

Deleted: -----Section Break (Continuous)-----

Deleted: Dataset Validation and Limitations

Deleted: Common ...alidations

Deleted:

Deleted: Techniques and Limited Information on Ground Truth of the Taisho Eruption

Deleted: result

Deleted: Section 3.1 briefly describes the strength and the well-known performance of the PUFF model for ash dispersal simulati

Formatted: No underline

Deleted: . Such methods have been used successfully for validating the simulation using the; for example, PUFF model

Formatted: No underline

Deleted: In addition, sSeveral past researcher...udies have

Deleted: Quality the ,using a degree of accuracy of ash dispersal simulation on a large-scale eruptions¶


Deleted: Ideally, if the ...recise wind data at the time of the

Sakurajima volcano over all Japan. We directly compared selected simulation results with the available reports of the Taisho eruption. However, making a direct comparison with the ground truth data was challenging owing to the following:

1. Lack of complete ground reports. Available reports from cities and tobacco plantations across Japan mentioned subjective ash deposit observations without a quantitative measurement, e.g., “frost-like ash” “house roof become white” and “slight deposit”.
2. Lack of complete wind data at the time of eruption. Only surface observations are available at limited observation points, and it is not possible to completely replicate the ash transport process.

Despite these limitations, we attempted to utilise an innovative approach to use all available information to validate the dataset. The ash dispersal map (Fig. 1) provided locations of ash deposition across all of Japan, and the detailed weather conditions that control the ash dispersal process are available. We utilised these data for validation, as follows:

1. We manifested the ground data in the same dimension as the simulation output by digitally redrawing the ash distribution map for all of Japan, provided by Omori (1914), using geographic information software (GIS). The resultant ground observation data resembled the original ash distribution of the Taisho eruption (Fig. 8).
2. We found wind pattern in the wind data of the JRA-55 Reanalysis that are equivalent to the day of the eruption. We expected that the similarity in weather features would also match similar wind profile across multiple atmospheric levels. We subset the number of simulation results for validation checking with the data produced in the previous step.



3.

- Deleted:** a wide space (all of ...apan)... We employed the ...
- Deleted:** edfor all of. laredwith
- Deleted:** in this study, ...aking a direct comparison with the ...
- Deleted:** The lLack of complete ground reports. Section 2 ...
- Deleted:** S yet
- Formatted:** List Paragraph, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.25" + Indent at: 0.5"
- Deleted:** The ecomplete wind data at the time of the ...ruptio ...
- Deleted:** ds
- Deleted:** ¶
- Deleted:** present a validation result using... all available ...
- Deleted:** these information
- Deleted:** iese information to proceed with the...ata for validatio ...
- Deleted:** Mmanifesting...d the ground data in the same ...
- Deleted:** The equivalent
- Deleted:** on the wind data of the JRA-55 Reanalysis that are ...
- Formatted:** List Paragraph, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.25" + Indent at: 0.5"
- Deleted:** Therefore, we utilised to thateise information to proceed with the validation process, eliminationsattemptedbyas follows:

Figure 8. Transformed ash dispersal map using ArcGIS software with airport icons for impacts reference. Human geography base (ESRI, 2022) provided the basemap for this figure.

Appendix B provides the detailed explanation of each step. The simulation result of the chosen date was matched with the new ground data using a binary contingency matrix (Table 3). The evaluation focused on measuring how the ash dispersal map produced from the simulation on selected dates would cover an equivalent region of interest to the ground data. We limited the evaluation to all ash deposits on the land and omitted ash deposits on the sea. We quantified the hit rate score *Hit* of the simulation matching in the binary contingency table as follows:

$$Hit = \frac{TP+TN}{TP+TN+FP+FN} \quad (5)$$

For brevity, we chose only the top date of the semantic textual similarity (STS) scores list in Appendix B (5 March 1997) in the matching process. The simulation for the chosen date reached satisfactory performance with a high *Hit* score (0.832), indicating that the selected date with weather conditions similar to those of the Taisho eruption had a reasonably similar ash dispersal pattern. Figure 9 exhibits further evidence of similarities in the ash dispersal map for the selected simulation results.

Table 3. Binary contingency table for the validation.


Actual Ash Precipitation of the Taisho Eruption	Selected Simulation Results	
	Affected	Unaffected
Affected	TP	FN
Unaffected	FP	TN

TP true positive; TN true negative; FP false positive; FN false negative

5.2 Dataset limitations

Generally, results from the simulation of volcanic ash dispersal and transport models depend on the initial values of parameters of eruption source and wind conditions during the eruption. These parameters are critical in determining the impacts of ashfall on proximal and distal locations (Bonadonna et al., 2012; Folch, 2012; Macedonio et al., 2016; Mastin et al., 2009; Webley and Mastin, 2009). We used deterministic values of plume height, erupted mass, and mass eruption rate as the eruption source parameters. We utilised the calculation by Iguchi (2014), incorporating the study by Morton et al. (1956) for the relationship between the observed plume height in the historical reports and studies (Kobayashi, 1982; Koto, 1916; Omori, 1914; Todde et al., 2017) to produce the mass eruption rate. The change in plume height over time was positively correlated with the total eruption mass and eruption rate. However, the method introduces a significant bias in the mean height value. The eruption rate is roughly proportional to the fourth power of height.

Formatted: Caption, No bullets or numbering



Deleted:

Deleted: Manifesting the ground data in the same dimension with the simulation output by digitally redrawing the ash distribution map for entire all of Japan, provided by Omori

Deleted: a shows placing the to do their

Deleted: ¶

Deleted: simply the data because of a reason for the ¶ omitted proceed with a normalyha in the and continue to, entire is for show the approximately situated C eigned with in C

Deleted: We show the detailed explanation of each step

Deleted:

Deleted: in Appendix B. Figure 8 portrays the transformed ash

Formatted: Superscript

Deleted: Table 3. The binary contingency table for the validation process. ¶

Actual Ash Precipitation of the Taisho Eruption

Moved (insertion) [12]

Formatted Table

Moved (insertion) [13]

Formatted Table

Formatted: Indent: Left: 1", First line: 0.5"

Deleted: To make it brief, we chose only the top date of the STS scores list in Table B2 (5 March 1997) in the matching process.




Figure 9. Comparison of ash distribution on the selected date (5 March 1997) according to the original ash dispersal map and simulation results. Human geography base (ESRI, 2022) provided the basemap for this figure.

Therefore, error in assigning the precise height will significantly affect its conversion to the eruption rate; and therefore, to the outcome of the simulation (Folch et al., 2012; Mastin et al., 2009). We assigned the parameters of the largest eruption size, or plume height of the last eruption as a baseline of a conservative safety factor, although we recognise that the variability could change the significance of the implied hazard. The values used here agreed fairly well with the past observations, and the bias in the eruption source parameter should be small and not significantly affect the ash dispersal mechanism. However, variations in these parameters result in different eruption rates and total mass, resulting in different ashfall footprints. Future vent location and eruption style, size, and duration significantly affect the ashfall in proximal and distal areas (Bonadonna et al., 2012; Mastin et al., 2009; Selva et al., 2018). Furthermore, some parameters, such as the diffusion coefficient, settling particle velocity law (see Table 1), orographic effects, and the lack of topographical data, need to be set arbitrarily causing discrepancies when using different variations depending on the model (Macedonio et al., 2016; Poulidis et al., 2017, 2018; Scollo et al., 2011). Finally, the distributions of total grain size and particle density and shape, and the chosen aggregation method also affect the accuracy of model outputs (Bonadonna et al., 2012; Folch et al., 2010, 2012; Mastin et al., 2009; Selva et al., 2018). Here, the aim of the simulation was to characterise the Taisho eruption, eliminating variations in those variables because we used a fixed scenario as the worst case. Using fixed values for the eruption source parameters to precisely replicate the transition in the Taisho eruption is a limitation which restricts the possibilities of future events.

Deleted: 10

Deleted: The cComparison of the ...sh distribution on the ...

Formatted: Caption

Deleted: (ESRI, 2022)

Deleted: Meanwhile, *FN* denotes the false negative, in which particular grid points are marked as unaffected in the simulation results but appear to be affected in the ground data. In reverse, *FP* denotes the false positive, in which particular grid points are marked as affected in the simulation results but appear to be unaffected in the ground data. Therefore, the hit rate score *Hit* is the simulation result correctly dispersed ashes to the same grid points as the ground data, divided by all defined grid points from the original ash dispersal map. To make it brief, we chose only the top date of the STS scores list in Table B2 (5 March 1997) in the matching process. The simulation results of the chosen date reached satisfactory performance with a high *Hit* score (0.832).

Moved up [1.3]: Meanwhile, *FN* denotes the false negative, in which particular grid points are marked as unaffected in the simulation results but appear to be affected in the ground data. In reverse, *FP* denotes the false positive, in which particular grid points are marked as affected in the simulation results but appear to be unaffected in the ground data. Therefore, the hit ...

Deleted: ¶

Deleted: This result means that the selected date with weat ...

Deleted: Figure 9 further exhibit the evidence of the ...

Deleted: 5.2 Finding the Identical Weather Characteristics¶

Deleted: 5.2 Dataset limitations¶

Deleted:

Deleted:

Deleted: s or plume height of the last eruption as a baseline of t ...

Deleted:

Deleted: we used in this study...ere agreed fairly well with the ...

Deleted: ¶

Deleted:

Deleted: on those ...hese parameters could ...result in different ...

Deleted:

Deleted: ¶

Deleted: Furthermore, depending on the models, the...some ...

Deleted:

Deleted: we approach ...he aim of the ...imulation with the m ...

Deleted: ¶

Deleted:

Deleted: We acknowledged the limitations of this study that ...

Deleted:

Deleted: Following the result mentioned earlier by

Hickey et al. (2016) illustrate how the current uplift situation is similar to that of the preceding eruption event, so our results could provide valuable input to authorities and stakeholders. The dataset provides a guideline for understanding how such a large explosive eruption would affect contemporary Japan. Inaccuracies also occur due to the effects of winds and atmospheric humidity, and other factors (Folch et al., 2012; Mastin et al., 2009). Meteorological conditions determine the ashfall dispersion trajectory and deposit location (Macedonio et al., 2016). Reanalysis datasets are favourable because they have better accuracy than the forecast dataset (Folch et al., 2012). The variability in the meteorological databases (e.g., ERA-Interim, JRA-55, NARR, etc.) is relatively insignificant (Selva et al., 2018). This dataset, intended to provide guidance for a worst case scenario, may remain helpful in extending the present study toward ashfall risk analysis and decision-making. We argue that the dataset here is vital to support future researchers and emergency managers in analysing and planning responses to crises that may occur under a variety of conditions. These risk assignments are rough estimates applying to the most likely future eruption size and type at Sakurajima volcano. There might be inconsistencies in the dataset that introduce bias in the ashfall hazard and risk analyses, so future users should consider all these sources of uncertainty.

Deleted: Hickey et al. (2016) that ...illustrates... how the curre...

Deleted:

Deleted: presented here will be ...rovides a guideline to ...

Deleted:

Deleted: Moreover, i...naccuracies also can ...ccur from ...ue ...

Deleted: ¶

Deleted: Therefore, rReanalysis datasets are favourable to use ...

Deleted:

Deleted: We understood that t...here might be ...

6 Conclusion

Ash dispersal products from large eruptions have devastating impacts on various sectors, in areas near the volcano and in distal locations. Emergency managers and crisis response planners need to acknowledge the compounded effect of ashfall deposits on the infrastructures to manage losses appropriately. This study presents a dataset that is useful for research and planning, focusing on the context of the entire country. This paper describes the process of generating a dataset based on the recent large eruption of Sakurajima volcano (the Taisho eruption) by simulating the eruption process over an extended period from 1958 to 2021. We validated the simulation of ash dispersal over all Japan following a large-scale eruption of Sakurajima volcano within some limiting constraints. Finally, although it contains some degree of inaccuracies, the dataset can be used for further studies in ashfall risk analysis and decision-making, and will support future researchers and emergency managers in devising disaster responses for different conditions.

Deleted: The a...sh dispersal products from large eruptions ...

Deleted: ¶

Deleted:

Deleted: e paper describes the complete generation ...rocess of ...

Deleted: Swestoestat thethe sohave beenFurthermore, we conducted the a validation procedure by identifying the similarity in the weather pattern at the time of the eruption and proving that similar weather conditions will bring identical ashfall distribution, thus verifying the dataset.

Deleted:

Deleted: we also acknowledged the limitation and uncertainties the dataset could bring. A...lthough it contains some degree of ...

Commented [HR2]: Moved figures to appendices based on Referee #1 recommendation

Deleted: the NetCDF format are...s shown in Fig. A1 for...

Appendix A

The structure of the dataset in NetCDF format is shown in Fig. A1 (ground deposits data) and Fig. A2 (airborne concentrations data).

Left: all Japan		Right: Kagoshima region	
Coordinates:	<ul style="list-style-type: none"> latitude (float32) 30.0 30.1 30.2 ... 45.7 45.8 45.9 longitude (float32) 128.5 128.6 128.7 ... 148.5 148.6 eruption_date (datetime64[ns]) ... 	Coordinates:	<ul style="list-style-type: none"> latitude (float32) 30.0 30.01 30.02 ... 32.99 33.0 longitude (float32) 129.0 129.01 ... 131.99 132.0 eruption_date (datetime64[ns]) ...
Data variables:	<ul style="list-style-type: none"> deposit (float32) ... 	Data variables:	<ul style="list-style-type: none"> deposit (float32) ...
Attributes:	<ul style="list-style-type: none"> Name: Long-period Ashfall Deposition Dataset (1958-2021) from the Sakurajima Taisho Eruption. Type: Daily Ashfall Deposit for Japan region from simulation on 1958/01/01 10:00 JST (01UTC). Description: This dataset presents the ashfall deposit thickness in cm for Japan. The dataset generated from a numerical simulation of the 1914 Taisho eruption in Sakurajima using PUFF model and JRA-55 Reanalysis data. Edition: 1 Authors: Haris Rahadianto Publisher: Disaster Prevention Research Institute (DPRI), Kyoto University, Japan License: Open Data Commons Attribution License (ODC-By) 	Attributes:	<ul style="list-style-type: none"> Name: Long-period Ashfall Deposition Dataset (1958-2021) from the Sakurajima Taisho Eruption. Type: Daily Ashfall Deposit for Kagoshima region from simulation on 1958/01/01 10:00 JST (01UTC). Description: This dataset presents the ashfall deposit thickness in cm for Japan. The dataset generated from a numerical simulation of the 1914 Taisho eruption in Sakurajima using PUFF model and JRA-55 Reanalysis data. Edition: 1 Authors: Haris Rahadianto Publisher: Disaster Prevention Research Institute (DPRI), Kyoto University, Japan License: Open Data Commons Attribution License (ODC-By)

Figure A1. Structures of sample deposit data in NetCDF format. Left: all Japan, Right: Kagoshima region.

Left: all Japan		Right: Kagoshima region	
Coordinates:	<ul style="list-style-type: none"> date_time (datetime64[ns]) 1958-01-01T11:30:00 ... 1958-01-... eruption_date (datetime64[ns]) ... 	Coordinates:	<ul style="list-style-type: none"> date_time (datetime64[ns]) 1958-01-01T11:30:00 ... 1958-01-... eruption_date (datetime64[ns]) ...
Data variables:	<ul style="list-style-type: none"> longitude (float32) ... latitude (float32) ... altitude (float32) ... ash_particles (float32) ... 	Data variables:	<ul style="list-style-type: none"> longitude (float32) ... latitude (float32) ... altitude (float32) ... ash_particles (float32) ...
Attributes:	<ul style="list-style-type: none"> Name: Long-period Ashfall Deposition Dataset (1958-2021) from the Sakurajima Taisho Eruption. Type: Airborne Ashes for 96 hours from 1958/01/01 10:00 JST (01UTC), recorded at every 90 minutes. Description: This dataset presents the total number of airborne ash particles from the starting simulation date. The dataset generated from a numerical simulation of the 1914 Taisho eruption in Sakurajima using PUFF model and JRA-55 Reanalysis data. Edition: 1 Authors: Haris Rahadianto Publisher: Disaster Prevention Research Institute (DPRI), Kyoto University, Japan License: Open Data Commons Attribution License (ODC-By) 	Attributes:	<ul style="list-style-type: none"> Name: Long-period Ashfall Deposition Dataset (1958-2021) from the Sakurajima Taisho Eruption. Type: Airborne Ashes for 96 hours from 1958/01/01 10:00 JST (01UTC), recorded at every 90 minutes. Description: This dataset presents the total number of airborne ash particles from the starting simulation date. The dataset generated from a numerical simulation of the 1914 Taisho eruption in Sakurajima using PUFF model and JRA-55 Reanalysis data. Edition: 1 Authors: Haris Rahadianto Publisher: Disaster Prevention Research Institute (DPRI), Kyoto University, Japan License: Open Data Commons Attribution License (ODC-By)

Figure A2. Structures of sample airborne data in NetCDF format.

Appendix B

A diagram explaining the flow of the validation procedure for this study is portrayed in Fig. B1.

B.1 Transforming the original ash precipitation map

To transform ground data to the same area dimension as the model output to enable its directly comparison with simulation results, we digitised the 1914 ash dispersal map of Fig. 1 by placing binary markings to indicate the affected regions; i.e., 1 for red-striped grids and 0 for the rest.

- Deleted: The s
- Deleted: the
- Deleted: for both regions in the
- Deleted: NetCDF format

- Deleted: The s
- Deleted: the
- Deleted: the
- Commented [HR3]: Moved the details of the validation from Sect 5 to here
- Deleted: The



Figure B1. Flowchart of validation of the dataset. (1) Digitising the original ash precipitation map to become new ground data, and (2) selecting a simulation result for matching with ground observation data.

All the binary values have a pair of longitude-latitude values to represent their positions on the map (Fig. B1). We added current airport locations to emphasise the impacts of very fine ashfall across all Japan. We used ArcGIS software with a WGS 84 projection (ESRI, 2017).

B.2 Finding the identical weather characteristics

The weather at the time of eruption determines the ash fallout trajectory. At the time of the Taisho eruption the weather was unusual, which explains why the ashfall moved unpredictably to distal locations. Previous analyses have not adjusted for the anomalous weather conditions that day (e.g., Biass et al., (2017), among many others). The Taisho eruption occurred in winter when mid-latitude cyclones were active. Stationary planetary waves generated by the land-sea contrasts and the flow over the Himalayas reinforce the tropospheric jet stream and the baroclinic wave. In Japan, the reinforced jet streams cause the westerlies at 250 hPa pressure to exceed 70 ms^{-1} . The winds in the tropospheric jet stream blow from the west throughout the year; they are strongest in winter and weakest in summer (Wallace and Hobbs, 2006). The Meteorological Research Institute of the JMA (JMA-MRI) further clarifies how the westerly wind traverses above the Sakurajima volcano all year long from west to east. The westerlies reach their peak velocity in January and then continue to weaken until their lowest speed in summer, before rising again (Maeda et al., 2012). This cycle runs for the entire year and is a determining factor for the ash dispersal pattern.

Deleted: The f

Deleted: the

Deleted: procedure using

Deleted: produced in this paper

Deleted: the process of d

Deleted: the process of s

Deleted: the

Formatted: Normal

Deleted: B.1 Transforming the original ash precipitation map¶

In the first step, as shown in Fig. B1, we transformed the ash dispersal map in Fig. 2 using ArcGIS software with a WGS 84 projection (ESRI, 2017). In addition, we added the current airport locations to signify the impacts of the very fine ashfall on entire Japan. We conducted the map transformation by placing binary markings to indicate the affected regions, i.e., 1 for red-stripped grids and 0 for the rest. All the binary values have a pair of longitude-latitude values to represent their positions on the map. This procedure is necessary, as we finally have a ground data in the same area dimension as the model output to directly compare it with the selected simulation result.

Deleted: The main component of the validation part is to match the simulation result with the new ground data. However, we cannot simply check the data with any results from the simulation because of the unusual weather at the time of the eruption. Due to the absence of a complete wind profile at the time of the eruption, we investigated the weather conditions on that day. Understanding the weather components at the time of eruption is crucial for determining the behaviour of the ash fallout trajectory. This feature of the Taisho eruption is often omitted, although it explains why the ashfall from the eruption moves farther to Japanese mainland in distal locations instead of the Pacific Ocean as the winds in the upper altitude during winter suggested (see Fig. 3b).¶ Several prior researchers did not proceed with their analysis of this anomaly, which has proven to be influential concerning the impacts in distal municipalities (e.g., Biass et al., (2017), among many others).

Deleted: ¶

Deleted: ¶

Deleted: With the s

Deleted: ,

Deleted: these phenomena

Deleted: to be the strongest

Deleted: one of the

Deleted: cid

Deleted: s




Figure B2. Seasonal progression of the westerlies over Japan at 12 km from the surface (Maeda et al., 2012). The red dashed square indicates the region covering Sakurajima volcano and Kagoshima city. The vertical axis shows the latitude of the wind position. The horizontal axis shows monthly changes of the wind velocity (left) and the longitude of the winds (right). The thick black arrow from left to right indicates the location of maximum velocity of westerlies based on seasonal progression. The straight blue lines indicate all regions in Japan affected by the westerlies. The small numbers inside the left image indicate the wind contour.

Figure B2 shows a time-latitude cross-section of the east-west wind at the 200 hPa (approximately 12 km from the surface), averaged at 130°E to 140°E based on the 30-year survey (Maeda et al., 2012; Mita et al., 2018). The Sakurajima volcano and its vicinity are situated around 31° N (red dashed square), including Kagoshima City. Therefore, westerlies would usually bring most of the ashfall eastwards from the eruption. Instead, chronicles reported that ash traversed to the north, west, and southwest (Omori, 1914; Todde et al., 2017). This pattern contrasted with the preceding eruption (An'ei eruption – November 1779) that followed the pattern of the westerlies (Tsukui, 2011). Investigation of past patterns of ash dispersal from Sakurajima volcano also reaffirm the differences arising from seasonal features, agreeing that during winter, tephra should mostly go eastward, and mostly fall in the Pacific (Biass et al., 2017; Poulidis et al., 2018; Mita et al., 2018).

When the Taisho eruption occurred, the vicinity of Kagoshima City and Sakurajima volcano experienced a maximum atmospheric pressure during the first 4 days, providing a unique characteristic of the Taisho eruption. From 9 January 1914 to the morning of 13 January 1914 atmospheric pressure in Japan was consistently recorded as unusually high, resulting in stable sea waves and low-velocity winds, indicating calm and clear weather in most places (Omori, 1914). Unexpectedly, the weak winds helped many people living near the volcano to evacuate by boat (Kitagawa, 2015). The eruption seemed to play a significant role in increasing the temperature in Kagoshima, but apparently the opposite was true (Omori, 1914). The leading cause of the high temperature at the time of eruption time was a sudden increase in southerly winds, as observed in Okinawa, 373 km from the vent (Omori, 1914). Furthermore, the chronicles of the tri-daily general weather conditions during the eruption reported that similar weather phenomena also occurred on 17 January 1914 (Omori, 1914). This vital information helped us derive a method for validating our simulation results. We concluded that clear, sunny weather with weak winds played a major role in transporting ashfall to all Japan.

Formatted: Caption

Deleted: degree

Deleted: westerlies

Deleted: for the

Deleted: image

Deleted: on the

Deleted: image

Deleted: ing

Deleted: s

Deleted:

Deleted: 5

Deleted: the

Deleted: to the east

Deleted: es

Deleted: ¶

Deleted: Research i

Deleted: g

Deleted: the

Deleted: patterns

Deleted: s

Deleted: due to

Deleted: with most of it

Deleted: ing

Deleted:

Deleted: four

Deleted: ¶

Deleted:

Deleted: n

Deleted: ,

Deleted: d

Deleted: , it turned out to be

Deleted:

Deleted: explained

Deleted: and found

Deleted: ¶

Deleted: Following such reports, w

Deleted: entire

Deleted: ,

Such conditions although rare (less than 11%, as winds during winter are usually strong (see Fig. 3e)), can recur. We assumed that the wind pattern would be similar on days with identical weather as irregular as the day of the Taisho eruption.

We obtained wind data at various atmospheric pressure heights that matched the wind data during the eruption. Thus, the simulation on the dates with similar weather is expected to produce a similar ash distribution to all of Japan, comparable with the transformed ash dispersal map in Fig. 8. The Taisho eruption occurred in winter, so we searched for winter day(s) with weather conditions like the eruption day (12 January 1914). We searched the available weather information (from the chronicles and historical weather information provided by the JMA on their website) for clear sunny days with weak winds with high atmospheric pressure (anti-cyclonic) and higher temperatures during December, January, February, and March from 1 January 1958 to 31 December 2021. These features are quite distinct and rarely occur in Japanese winter. However, the Japanese authority only maintained surface weather data (temperature, pressure, humidity, precipitation, and wind speed) in a major meteorological observatory located in the capital of selected prefectures. These past data consist of daily and hourly observations. No other measurements were recorded at different atmospheric pressure levels. The surface weather chart on the day of the eruption (12 January 1914) and all simulation dates is available. Therefore, we utilised a surface weather chart for a clearer description of weather phenomena on a particular day. Weather chart consist of symbols and features that describe a particular meteorological pattern in a specific space-time dimension. Weather charts were created by plotting measurements including mean sea-level pressure, wind barbs, and cloud cover, and identifying synoptic-scale features such as weather fronts. A meteorologist usually performs an isobaric analysis of the maps by constructing lines of equal mean sea level pressure (Wallace and Hobbs, 2006).

The weather chart at 06:00:00 on 12 January 1914 (Fig. B3) displays the high-pressure condition centred over central Honshu and covering almost all Japan. These local maxima in the pressure field explain the common weather characteristics in that specific region. The wind becomes stronger when the pressure gradient between the high- and low-pressure systems increases. Land friction weakens the wind coming from high-pressure systems. Hence, high-pressure systems typically result in weaker winds with clear skies and relatively warm weather (Wallace and Hobbs, 2006); thus, the interpretation of the weather chart corroborates the weather reports of the eruption day. Accordingly, searching for similarities in surface weather charts is appropriate for identifying identical weather conditions.

B.2.1 Weather report commentary on the surface weather charts

To search for identical weather by comparing features in pairs of weather charts for similarity we initially employed an image similarity method. However, this was very difficult, as the weather chart on the eruption day differed greatly from the modern weather chart used in Japan. The current daily weather chart differed from the weather chart on 12 January 1914 in the measurement time and scale of barometric pressure, dimension and projection of the map, and area included in the map. The format of daily weather charts in Japan has changed several times after World War II. In 1883 the JMA began producing daily surface weather charts for the Asia-Pacific region, which consisted of Japan and Japanese-occupied areas pre-World War II.

Deleted: and s

Deleted: is

Deleted:)

Deleted: characteristically

Deleted: From this observation, w

Deleted: il

Deleted: also

Deleted: characteristics

Deleted: Using this assumption, w

Deleted: further

Deleted: ing

Deleted: ¶

Owing to seasonal differences, finding dates with similar weather only uses winter dates, because t

Deleted: as well. In detail,

Deleted: and found the

Deleted: The

Commented [A4]: I omitted this text because the H cannot be seen in Fig. B3.

Deleted: search process only focused on winter days (December, January, February, and March) from 1 January 1958 to 31 December 2021. The important main features to query are clear sunny days, weak winds with high atmospheric pressure (anti-cyclonic), and higher temperatures, which are quite distinct and rarely occur in Japanese winter. These features reiterate the peculiarity of the Taisho eruption and its diverging ash distribution. Weather comparisons were conducted using available weather information from the chronicles and historical weather information provided by the JMA on their website. However, the past meteorological data are limited because the Japanese authority only maintained surface weather condition (temperature, pressure, humidity, precipitation, and wind speed) in a major meteorological observatory located in the capital of selected prefectures. These past data consist of daily and hourly observations. Apart from surface observations, there are no other available measurement recorded at different atmospheric pressure levels. Other available information is the surface weather chart on the day of the eruption (12 January 1914) and all simulation dates. Therefore, we decided to utilise a surface weather chart instead of surface observations, as it provides a clearer description of a weather phenomenon on a particular day. ¶
A weather chart, also known as a weather map, usually consists of symbols and features that describe a particular meteorological pattern in a specific space-time dimension. Weather charts were created by plotting the relevant measurements such as mean sea-level pressure, wind barbs, and cloud cover. This plotting can help find synoptic-scale features, such as weather fronts. A meteorologist usually performs an isobaric analysis of these maps, which involves the construction of lines of equal mean sea level pressure (Wallace and Hobbs, 2006).




Figure B3. Weather chart at 06:00:00 on 12 January 1914, the time of the Taisho eruption, with additional markings of pressure measurement converted to the latest convention (mmHg → hPa). Modified from Omori (1914).

Since August 1958, the JMA has provided daily weather charts at various upper levels (500, 700, and 850 hPa) for the Asia-Pacific and 500 hPa for the Northern Hemisphere region. Since March 1996, the JMA added a daily surface weather chart for the Northern Hemisphere and a 300 hPa level for the Asia-Pacific region. Finally, in February 1999 the JMA updated all the weather charts and added a specific surface weather chart for the regions of Japan. These changes were accompanied by changes in the weather chart format to that currently in daily use. Owing to the ever-changing formats of past weather charts, identifying image similarity was time-consuming because of the need for multiple training data for every format to decide whether the two images are similar, and was not feasible. Instead we utilised other information available in the daily weather charts, namely the general weather report commentary (天気概況). Experts in meteorology developed this commentary as a guide to weather conditions during a particular day. Figure B4a portrays the weather report commentary on 12 January 1914. All historical daily weather charts from 1883 to March 1938, have a written commentary explaining the weather conditions. However, this feature was omitted from April 1938 until December 1995. The JMA resumed the commentary feature in the monthly compilation of historical daily weather charts from January 1996 to date (Database of Weather Charts for Hundred Years, 2022) (Fig. B4b) (Daily Weather Chart, 2022).

Formatted: Caption

Deleted: The weather chart on the Taisho eruption with additional markings of pressure measurement converted to the latest convention (mmHg → hPa). Modified from Omori (1914).

Deleted:

Deleted: Figure B3 shows the weather chart at 06:00:00 on 12 January 1914, with the modern convention of pressure measurements. This weather chart displays how the high-pressure condition covers almost the entire Japan centring the central Honshu. These local maxima in the pressure field, denoted by a big "H" letter, can simply explain the common weather characteristics in that specific region. The wind becomes stronger when the pressure gradient between the high- and low-pressure systems increases. Land friction weakens the wind coming out from high-pressure systems. Hence, high-pressure systems typically result in weaker winds with clear skies and relatively warm weather (Wallace and Hobbs, 2006). These implications corroborate the weather reports during the eruption day. Accordingly, using the similarity on the surface weather chart is appropriate for identifying identical weather conditions.

Deleted: B.2.1 Weather report commentary on the surface weather charts

Our initial attempt to search for identical weather by using weather chart similarity is by employed an image similarity method to compare features drawn in a pair of weather charts. However, this attempt turned out to be very difficult, as the weather chart on the eruption day was much different from the modern weather chart currently used in Japan.

The measurement time and scale of barometric pressure, the dimension and projection of the map, and the area included in the map were all different between the current daily weather chart and the weather chart on 12 January 1914.

The format of the daily weather charts in Japan has changed several times after World War II. Starting in 1883, the JMA produced daily surface weather charts for the Asia-Pacific region, consisting of Japan and Japanese-occupied areas pre-World War II. Since August 1958, the JMA has provided daily weather charts at various upper levels (500 hPa, 700 hPa, and 850 hPa) for the Asia-Pacific and 500 hPa for the Northern Hemisphere region. Then, starting in March 1996, the JMA added a daily surface weather chart for the Northern Hemisphere and a 300 hPa level for the Asia-Pacific region. Finally, in February 1999, the JMA updated all the weather charts and added a specific surface weather chart for the regions of Japan. These changes were also accompanied by changes in the weather chart format, and the current daily weather charts follows the 1999 format. Thus, owing to the ever-changing formats of past weather charts and the modern ones, performing image similarity is not feasible and is a time-consuming task because we need to have multiple training data for every format to decide whether the two images are similar. We moved to another option which includes employing other information available in the daily weather chart. Another piece of information available inside the weather chart is the general weather report commentary (天気概況). Experts in meteorology developed this commentary as a guide to weather conditions during a particular day. Figure B4a portrays the weather report commentary on 12 January 1914 located at the bottom left of the second page.




Figure B4. General weather report commentary on the weather condition inside the weather chart. (a) Old weather chart included the commentary on the bottom of the second page on the day of the Taisho eruption (Database of Weather Charts for Hundred Years, 2022). (b) Modern weather chart: the first page of the daily weather chart compilation for December 2020. The commentary is placed at the bottom of the monthly compilation of daily weather charts (Japan Meteorological Agency, 2022).

Formatted: Caption

Deleted: The general weather report commentary on the weather condition inside the weather chart. (a) The old weather chart put the commentary on the bottom of the second page on the day of the Taisho eruption (Database of Weather Charts for Hundred Years, 2022). (b) The modern weather chart mentions the commentary on the bottom of the monthly compilation of daily weather charts (Japan Meteorological Agency, 2022).

The daily commentary usually includes the important weather characteristics such as high- or low-pressure centres or fronts, extreme weather phenomena such as typhoons, heavy rains, or heavy snow, and other special events such as disasters. This description enabled us to find weather similar to the day of the eruption during the dates used in the simulation (1996–2021, 3120 days). Several initial conversions were required to overcome a minor distinction between the old and modern commentaries, namely:

1. The old commentary on the weather chart on the day of the eruption was based on old Japanese syntax.
2. The old commentary addressed the weather in Japan and in Japanese-occupied areas.

We converted the old commentary to contemporary Japanese syntax (Table B1) and removed all observation areas outside the modern Japan region. The old commentary consists of three parts: the location of high- or low-pressure centres, the weather on the main islands of Japan (Honshu, Kyushu, and Hokkaido), and in other parts of the Asia-Pacific under Japanese occupation. This commentary agrees with the meteorological reports provided in the chronicles (Omori, 1914). We focused on the weather on the main islands of Japan, and converted the old commentary to a form similar to that in the current daily weather charts.

3035 The new phrases in the last column give more brief comments that match with the contemporary style in the modern weather charts while still maintaining the important features (high-pressure all over Japan, clear/sunny weather in the main islands, Kyushu is cloudy, and Hokkaido is either clear or cloudy with strong south-westerly winds). Then, we compared these phrases in contemporary Japanese with all the commentaries in the modern weather chart (from 1 January 1996 to 31 March 2021) provided by the JMA in winter (from January 1996 to March 2021), in the historical archive collected by the National Institute of Informatics (Database of Weather Charts for Hundred Years, 2022). These commentaries explain the weather conditions on a particular day in a random sequence; that is, not in a similar order to the processed commentary phrases (from 12 January 1914). Finally, we utilised the natural language processing (NLP) method to find semantic textual similarity between all commentaries with sentence embedding using SentenceBERT (Reimers and Gurevych, 2019).

3045 **Table B1: Conversion of the general weather commentary in the weather chart of 12 January 1914.**

Original Form	Contemporary Form	Final Form
高気圧東に移り、中心本邦中部に在りて、773mmを示し。揚子江低気圧は目下漢口付近に存す。本邦にては、天気おおむね良好なれども、九州にてはすでに曇天となり、北海道にては南西風強く晴曇相半す。満州および朝鮮にてはおおむね曇天となれり、琉球および台湾にてはこれ反して快晴なり。	高気圧は東に移動し、日本の中心にあり、気圧計の値は773mmを示すしている。長江にある低気圧は、ちょうど武漢付近に存在する。日本では、天気は晴れているが、九州地方では既に曇りとなっており、北海道では南西の風が強く晴れたり曇ったりしている。満州と朝鮮はおおむね曇り、琉球(沖縄)と台湾では、反して快晴。	日本付近は移動性高気圧に覆われている。日本列島は晴れ。一方、九州は曇り。北海道は南西風が吹き、晴れや、曇り。

B.2.2 Semantic textual similarity of weather report commentary using SentenceBERT (SBERT)

To find similar weather, both commentaries should have a meaning that is semantically similar, regardless of the sentence structures and words used to form the sentence. The vast improvements in machine learning (ML) and artificial intelligence (AI) applied to NLP tasks allowed us to find the similarity between two sentences using semantic textual similarity (STS) by embedding all sentences first. Word embedding represents words in the form of real-valued vectors which encode the meaning of a word such that the words that are closer in the vector space will be similar in meaning (Jurafsky and James, 2000). Sentence embedding is the sum of individual word embedding (Cer et al., 2017). STS scores semantic similarity on a continuous scale (e.g., a vehicle and a car are more similar than a wave and a car) rather than binary (e.g., a vehicle and a car are the same, and a wave and a car are not the same). STS provides a unified framework that allows for the extrinsic evaluation of multiple semantic components.

Deleted: All historical daily weather charts, from 1883 up to March 1938, have a commentary written on a weather chart explaining the weather conditions. However, this feature was omitted from April 1938 until December 1995. The JMA resumed adding a general weather report commentary feature to the daily weather chart from January 1996 to date, under the weather chart images in the monthly compilation of historical daily weather charts (Database of Weather Charts for Hundred Years, 2022). Figure B4b shows the first page of the daily weather charts compilation for December 2020 from the JMA as a reference (Daily Weather Chart, 2022). The commentary usually includes the important weather characteristics on that day, such as the existence of high- or low-pressure centres or fronts, the description of extreme weather phenomena such as typhoons, heavy rains, or heavy snow, and other special events on the day, such as disasters. This description allows us to find similar weather between the dates used in the simulation and the day of the eruption, even though we further decreased the number of dates used in the searching process (1996-2021, 3120 days). However, before proceeding with the searching process, we need to perform several conversions because there is a minor distinction between the old commentary and the modern ones, which are:

<#>The old commentary on the weather chart on the day of the eruption was based on old Japanese syntax.
 <#>The old commentary addressed both the weather in Japan and the weather in a Japanese-occupied areas.
 We solve these slight differences by simply converting the old commentary to contemporary Japanese syntax and removing all observation areas outside the modern Japan region. The first column in Table B1 denotes the original form written in the Old Japanese syntax. The second column shows the transformation from the Old to the contemporary Japanese. The old commentary consists of three parts: explaining the location of high- or low-pressure centres, the

Moved (insertion) [17]
Deleted: The conversion of the general weather commentary in the weather chart on 12 January 1914.

Formatted: Caption

Formatted Table

Deleted: This commentary agrees with the meteorological report provided in the chronicles (Omori, 1914). Therefore, we omitted the last phrases and focused only on the weather on the main islands of Japan. Then, further converted it to a form similar to the commentary in the current daily weather charts. The new phrases in the last column give more brief comments that match

Moved up [17]: This commentary agrees with the provided in the chronicles (Omori, 1914). Therefore, we omitted the last phrases and focused only on the weather on the main islands of Japan. Then, further converted it to a form similar to the commentary in the current daily weather charts. The new phrases in the last column give more brief comments that match with the contemporary style in the modern weather charts while

Deleted: B.2.2 Semantic textual similarity on weather report commentary using SentenceBERT (SBERT)
 To find similar weather, both commentaries should have an equivalent meaning that is semantically similar, regardless of the sentence structures and words used to form the sentence. The vast improvements in machine learning (ML) and artificial intelligence (AI), especially in NLP tasks, allow us to find the similarity between

This framework includes word sense disambiguation and induction, lexical substitution, semantic role labelling, multi-word expression detection and handling, anaphora and co-reference resolution, time and date resolution, named-entity handling, under-specification, hedging, semantic scoping, and discourse analysis (Agirre et al., 2012). Here, we considered a pair of sentences to be similar (in this case, the weather condition) if they were located close to each other, as both sentences are mapped to a dense vector space (Reimers and Gurevych, 2019, 2020). Google led the current state-of-the-art NLP development with its technology, Bidirectional Encoder Representations from Transformers (BERT), which can handle various NLP tasks accurately, including question answering, sentence classification, and sentence-pair regression (Devlin et al., 2019). However, BERT is not optimal for handling tasks such as large-scale semantic similarity comparison, clustering, and information retrieval via semantic search (Reimers and Gurevych, 2020). To alleviate this issue, Reimers and Gurevych (2019) developed SentenceBERT (SBERT), which is a modification of the BERT network using Siamese and Triplet networks that can derive semantically meaningful sentence embedding.

The Siamese network architecture enables input sentences to be fixed-sized vectors. We used SBERT to find weather conditions similar to those of 12 January 1914 (query) from all the corpora available (general weather report commentary on daily weather charts 1996–2021) (Fig. B5) We defined the corpora as a collection of commentaries, which may contain one or more sentences explaining the weather conditions in a non-uniform sequence. SBERT embeds each sentence in both query and corpus with models pre-trained specifically to handle the Japanese language. The result of the sentence embedding is vectors of length 768 values, specific to each sentence, representing the sentence in vector space (Reimers and Gurevych, 2019).



Figure B5. Flowchart for finding dates with weather report commentaries similar to the day of eruption, using a semantic search, based on semantic textual similarity between the old and the modern commentaries, by SentenceBERT (SBERT).

We employed pre-trained models for the Japanese language available from the Hugging Face model hub (Hugging Face, 2022). Sentence embedding resources for the Japanese language are limited, and we used an appropriate pre-trained model exclusively developed for Japanese sentences, *sbert-base-ja* (Abe, 2021) (Colorful Scoop, 2022); CC-BY-SA 3.0, Colorful Scoop, 2021). We encoded queries and corpora into the vector space then conducted a semantic search to find the commentary with most similarity between the query and corpora.

Deleted: This framework includes word sense disambiguation and induction, lexical substitution, semantic role labelling, multi-word expression detection and handling, anaphora and co-reference resolution, time and date resolution, named-entity handling, under-specification, hedging, semantic scoping and discourse analysis (Agirre et al., 2012).

Here, we can consider a pair of sentences to be similar (in this case, the weather condition) if they are located close to each other, as both sentences are mapped to a dense vector space (Reimers and Gurevych, 2019, 2020). Google has led the current state-of-the-art NLP development with its technology, Bidirectional Encoder Representations from Transformers (BERT), which can handle various NLP tasks accurately, including question answering, sentence classification, and sentence-pair regression (Devlin et al., 2019). However, BERT is not optimal for handling several tasks, such as large-scale semantic similarity comparison, clustering, and information retrieval via semantic search (Reimers and Gurevych, 2020).

To alleviate this issue, Reimers and Gurevych (2019) developed SentenceBERT (SBERT), which is a modification of the BERT network using Siamese and Triplet networks that can derive semantically meaningful sentence embedding.

The Siamese network architecture enables input sentences to be fixed-sized vectors. Thus, we use SBERT to find a similar weather conditions on 12 January 1914 (query) from all the corpora available (general weather report commentary on daily weather chart 1996–2021). Figure B5 explains all steps to find similar weather conditions through a semantic search based on STS between the old and the modern commentaries. We define the corpora as a collection of a commentaries, which may contain one or more sentences explaining the weather conditions in a non-uniform sequence. SBERT embeds each sentence on both query and corpus with pre-trained models, specifically to handle the Japanese language. The result of the sentence embedding is vectors with a length of 768 values, specific to each sentence, representing the sentence in the vector space (Reimers and Gurevych, 2019).

Formatted: Caption

Deleted: The flowchart for finding the date with a similar weather report commentary to the day of eruption using SBERT.

Deleted:

Semantic search offers an advantage of finding synonymous sentences, even when the sentences are constructed using different words and order, as the search process focuses on understanding the query's content. We performed a symmetric semantic search because the query and corpus have similar lengths and numbers of sentences (Reimers and Gurevych, 2019). We used cosine similarity to compute the similarity scores between the query and all corpora. Cosine similarity is a popular similarity metric used to identify a pair of sentences with overlapping meanings, as these sentences are likely to be located next to each other in the vector space (Cer et al., 2017; Reimers and Gurevych, 2019; Arsov et al., 2019).

$$\text{sim}_v(q, c) = \frac{v(q) \cdot v(c)}{\|v(q)\|_2 \|v(c)\|_2} \quad (6)$$

Given a pair of sentence vectors $v(q)$ and $v(c)$, cosine similarity sim_v is computed by the dot product and the magnitude between two vectors, where q and c are the components of vector $v(q)$ and vector $v(c)$, respectively. Here, q is each sentence in the queries, which is the feature we want to find, and c is each sentence in all the commentaries collected from 1996 to 2021.

Table B2. Top 10 weather similarities scores of the selected dates.

Date	STS Score
5 March 1997	0.9635
24 February 2006	0.9575
2 February 2011	0.9551
11 December 1996	0.9532
16 March 2001	0.9514
30 March 2009	0.9508
28 January 2009	0.9507
29 December 1996	0.9503
19 January 2019	0.9497
14 January 2005	0.9496

STS: semantic textual similarity

The results of the comparison are between {0} for the most dissimilar sentences, and {1} if a pair of sentences is very similar (Singhal, 2001). Similar weather conditions were found by ranking all corpora based on their similarity scores (their similarity with the query). We selected only the highest result from the top 10 results of the search process (Table B2). Table B3 presents the weather commentary on the top result of the semantic search.

Deleted: We employed pre-trained models available for the Japanese language from the Hugging Face model hub (Hugging Face, 2022). However, sentence embedding resources for the Japanese language are limited, and we found only one appropriate pre-trained model exclusively developed for Japanese sentences, *sbert-base-ja* (Abe, 2021). This pre-trained model is from Colorful Scoop, a non-profit entity developing a conversational AI (Colorful Scoop, 2022). This company produced its own BERT Japanese model, which is trained using the Japanese Wikipedia corpus and released in 2021 under the CC-BY-SA 3.0 licence. The *sbert-base-ja* pre-trained model is an extension of their original model and is trained using the Japanese translation of the Stanford Natural Language Inference (JSNLI) dataset released by Kyoto University under CC-BY-SA 4.0 licence (Yoshikushi et al., 2020), with 535,000 sentence pairs, utilising AdamW optimiser with a 2×10^{-5} learning rate (Abe, 2021). Once we encode both queries and corpora into the vector space, we conduct a semantic search to find the most similar commentary between the query and corpora. The benefit of using semantic search is that we can find synonymous sentences, although constructed using completely different words and order, as the search process focuses on understanding the query's content. Specifically, we performed a symmetric semantic search because both the query and corpus have similar lengths and numbers of sentences (Reimers and Gurevych, 2019). We used cosine similarity to compute the similarity scores between the query and all corpora.

Deleted: ¶
Cosine similarity is a popular similarity metric used to identify a pair of sentences with an overlapping meanings, as these sentences are likely to be located next to each other in the vector space (Cer et al., 2017; Reimers and Gurevych, 2019; Arsov et al., 2019).

Deleted: 5

Deleted: Given a pair of sentence vectors $v(q)$ and $v(c)$, cosine similarity sim_v is computed by the dot product and the magnitude between two vectors, where q and c are the components of vector $v(q)$ and vector $v(c)$, respectively. Here, q is each sentence in the queries, which is the feature we want to find, and c is each sentence in all the commentaries collected from 1996 to 2021.

Deleted: The results of the comparison are between {0} for the most dissimilar sentences, and {1} if a pair of sentences is very similar (Singhal, 2001). The last step is to find similar weather conditions by ranking all corpora based on their similarity scores with the query. In brief, we selected only the top highest result from the top 10 results obtained from the search process, as shown in Table B2.

Deleted: he t

Formatted Table

Deleted: ¶

Table B3: Top result of the semantic search with its weather description from the commentary.

Date	STS Score	Commentary (ja)	Commentary (en)
1997/03/05	0.963	全国的に平年より5~10°C高い。日本付近は移動性高気圧に覆われるが、大陸から気圧の谷が進んでくる。北海道で曇りの他は似中晴れ。夜になると日本海側は曇り。山陰や九州北部では雨。	5~10°C above normal nationwide. A mobile high-pressure system will cover the area around Japan, but a trough of pressure will move in from the continent. Mostly cloudy in Hokkaido and sunny throughout. Cloudy over the Sea of Japan at night. Rain in the San-in region and northern Kyushu.

STS: semantic textual similarity. The commentary in English is a direct translation using DeepL translation software (DeepL Translator, 2022), with sentences in bold indicating weather conditions similar to the Taisho eruption.




Figure B6. Surface weather chart for Asia Pacific region at 00 UTC 5 March 1997. Weather chart image was obtained from the Japan Meteorological Agency (Database of Weather Charts for Hundred Years, 2022; Daily Weather Chart, 2022).

B.2.3 Weather on the selected date

The date with the highest STS scores (5 March 1997) agreed well with the weather features mentioned in the chronicles. The weather on the selected date was generally clear and sunny because of the high-pressure covering Japan's main islands, with the exception of some rain that the chronicles did not mention, demonstrating it may not be possible to observe exactly similar weather. This slight difference may slightly affect the simulated distribution of ashfall.

Deleted: he t

Moved down [19]: Table B3 describes the general commentary on the top results of the semantic search. The commentary in English is a direct translation using DeepL translation software (DeepL Translator, 2022), with sentences in bold indicating weather conditions similar to the Taisho eruption.

Deleted:

Deleted: process

Formatted: Caption

Formatted Table

Formatted: Caption

Moved (insertion) [18]

Deleted: Table B3: The top result of the semantic search process with its weather description from the commentary.

Formatted: Justified

Deleted: Surface weather chart for Asia Pacific region at 00 UTC 5 March 1997. Weather chart image is obtained from the JMA (Database of Weather Charts for Hundred Years, 2022; Daily Weather Chart, 2022).

Deleted: The results of the comparison are between {0} for the most dissimilar sentences, and {1} if a pair of sentences is very similar (Singhal, 2001). The last step is to find similar weather conditions by ranking all corpora based on their similarity scores with the query. In brief, we selected only the top highest result from the top 10 results obtained from the search process, as shown in Table B2. Table B3 describes the general commentary on the top results of the semantic search. The commentary in English is a direct translation using DeepL translation software (DeepL Translator, 2022), with sentences in bold indicating weather conditions similar to the Taisho eruption.

Formatted: Top: 0.39", Bottom: 0.93", Width: 8.27", Footer distance from edge: 0.51"

Moved (insertion) [19]

Deleted: condition

Figure B6 depicts the weather chart for the chosen date, illustrating more apparent features analogous to the weather chart published during the eruption. The chosen date and the eruption date exhibit the same weather: high-pressure covering all Japan with a low-pressure centred in the north. On the selected date, the high-pressure centre moved from the south to cover Japan at night. The general weather report commentary, accompanied by a daily weather chart, provide critical information that clarified the weather conditions on the respective dates. Another feature, not mentioned in the commentary, was the wind conditions on the selected date. The chronicles reveal that the wind was weak, making it possible to evacuate by boat to escape the eruption (Omori, 1914; Kitagawa, 2015).

Author contributions

HR performed most of the work to produce the dataset, namely processing data and simulations, analysing results, and writing this manuscript. HT also conceptualised the project and supervised the simulations and analyses. MI produced the Taisho eruption data used as the eruption source parameters used in the model to generate the dataset. HLT is the creator of the PUFF model and provided the original FORTRAN codes of the model and the default values of the parameters used in the simulation. TT verified the weather similarity process by incorporating the general weather report commentary inside the weather chart. SR assisted in developing a semantic search methodology to find the semantic similarities through the commentary.

Competing Interest

The authors declare that they have no conflict of interest.

Acknowledgements

We acknowledge the works of the Sakurajima Volcano Research Center (SVRC), Disaster Prevention Research Institute, Kyoto University in Kagoshima, Japan, which provided the Taisho eruption data used in this study. We thank the JMA for supplying the JRA-55 Reanalysis data from 1958 to 2020. The JRA-55 data were obtained from the archive of the Center for Computational Sciences (CCS), University of Tsukuba, Ibaraki, Japan. Furthermore, the authors are grateful for the help from Si Ha, Ryo Hirako, and Toraemon Matsumoto, who assisted in the map digitalisation process of the original ash dispersal map and the translation process for the old weather charts. HR acknowledges fellowship and financial support from the Tokio Marine Kagami Memorial Foundation for this work. We would like to thank Editage (www.editage.com) for English language editing.

Deleted: The date with the highest STS scores (5 March 1997) agreed well with the weather features mentioned in the chronicles. The weather on selected date was generally clear and sunny because of the high-pressure covering Japan's main islands. With the exception of some rain that the chronicles did not mention. This difference indicate that we may not observe the exactly similar weather completely. This slight difference may affect the ashfall distribution produced by the simulation slightly. Figure B6 depicts the weather chart for the chosen date, illustrating more apparent features analogous to the weather chart published during the eruption. Both the chosen date and the eruption date exhibit the same weather characteristics, as high-pressure covering entire Japan regions, added with an influence of a low-pressure centre in the north. On the selected date, the high-pressure centre moved from the south and then covered Japan at night. The weather charts presented in Fig. B6 further substantiate the continuous format transition of surface weather analysis in Japan. The general weather report commentary, accompanied by the establishment of a daily weather chart, is proven to be critical information that clarifies the weather conditions on the respective date.¶ Another feature, not mentioned in the commentary was the wind conditions during the selected dates. The chronicles briefly revealed that the wind was weak, making it possible to evacuate using boats from the island to escape the eruption (Omori, 1914; Kitagawa, 2015). ¶ To further analyse the weather similarities between the selected date and 12 January 1914, we drew the condition of the wind using a wind rose graph as depicted in Fig. B7.¶

Deleted:

Deleted: The graph consists of winds observations for every 6 h from the start of the simulation (00 UTC on the selected date) until the simulation process finished (00 UTC four days after the eruption start). We chose the winds observations both at the 1000 hPa pressure level (equivalent to the surface pressure) and 250 hPa pressure level (equivalent to the location of the westerlies) based on the findings explained in Sect. 2. The surface winds characteristics differed in terms of directions but had a comparable velocity between the two selected dates. The winds surface winds during ¶

Deleted: Contributions

Deleted: HR did the most work to produce the dataset by doing the data processing and simulations, analysing the results, and writing this manuscript. In addition, HT conceptualised the project and supervised both the simulations and the result analysis. MI produced the Taisho eruption data used as eruption source parameters used in the model to generate the dataset. HLT is the creator of the PUFF model, provided the original FORTRAN codes of the model and the default value of the parameters used in the simulation. TT verified the weather similarity process incorporat¶

Deleted: We acknowledge the works of Sakurajima Volcano Research Center (SVRC), Disaster Prevention Research Institute, Kyoto University in Kagoshima, Japan, that provided the Taisho eruption data used in this work. In addition, we thank the JMA for supplying the JRA-55 Reanalysis data from 1958 to 2020. The JRA-55 data were obtained from the archive of the Center for Computational Sciences (CCS), University of Tsukuba, Ibaraki, Japan. Furthermore, the authors were grateful for the help from Si Ha, Ryo Hirako, and Toraemon Matsumoto, who assisted in the ¶

References

Abe, N.: Sentence BERT Base Japanese Model, Python, Colorful Scoop, 2021.

3565 Agirre, E., Cer, D., Diab, M., and Gonzalez-Agirre, A.: SemEval-2012 Task 6: A pilot on semantic textual similarity, in: Proceedings of the Sixth International Workshop on Semantic Evaluation (SemEval 2012), SEM 2012: The First Joint Conference on Lexical and Computational Semantics, Montréal, Canada, 385–393, 2012.

Arsov, N., Dukovski, M., Evkoski, B., and Cvetkovski, S.: A Measure of Similarity in Textual Data Using Spearman’s Rank Correlation Coefficient. ArXiv [eprint], arXiv:1911.11750, 2019.

3570 Ayris, P. M. and Delmelle, P.: The immediate environmental effects of tephra emission, Bull. Volcanol., 74, 1905–1936, <https://doi.org/10.1007/s00445-012-0654-5>, 2012.

Barsotti, S., Andronico, D., Neri, A., Del Carlo, P., Baxter, P. J., Aspinall, W. P., and Hincks, T.: Quantitative assessment of volcanic ash hazards for health and infrastructure at Mt. Etna (Italy) by numerical simulation, J. Volcanol. Geotherm. Res., 192, 85–96, <https://doi.org/10.1016/j.jvolgeores.2010.02.011>, 2010.

3575 Barsotti, S., Di Rienzo, D. I., Thordarson, T., Björnsson, B. B., and Karlsdóttir, S.: Assessing impact to infrastructures due to tephra fallout from Óræfajökull volcano (Iceland) by using a scenario-based approach and a numerical model, Front. Earth Sci., 6, 196, <https://doi.org/10.3389/feart.2018.00196>, 2018.

Biass, S., Scaini, C., Bonadonna, C., Folch, A., Smith, K., and Höskuldsson, A.: A multi-scale risk assessment for tephra fallout and airborne concentration from multiple Icelandic volcanoes – Part 1: Hazard assessment, Nat. Hazards Earth Syst. Sci., 14, 2265–2287, <https://doi.org/10.5194/nhess-14-2265-2014>, 2014.

3580 Biass, S., Bonadonna, C., di Traglia, F., Pistolesi, M., Rosi, M., and Lestuzzi, P.: Probabilistic evaluation of the physical impact of future tephra fallout events for the Island of Vulcano, Italy, Bull. Volcanol, 78, 37, <https://doi.org/10.1007/s00445-016-1028-1>, 2016.

3585 Biass, S., Todde, A., Cioni, R., Pistolesi, M., Geshi, N., and Bonadonna, C.: Potential impacts of tephra fallout from a large-scale explosive eruption at Sakurajima volcano, Japan, Bull. Volcanol., 79, <https://doi.org/10.1007/s00445-017-1153-5>, 2017.

Bonadonna, C.: Probabilistic modelling of tephra dispersion, in: Statistics in Volcanology, Special Publications of IAVCEI, 1, edited by: Mader, H. M., Coles, S. G., Connor, C. B., and Connor, L. J., The Geological Society of London on behalf of The International Association of Volcanology and Chemistry of the Earth’s Interior, 243–259, <https://doi.org/10.1144/IAVCEI001.19>, 2006.

3590 Bonadonna, C., Ernst, G. G. J., and Sparks, R. S. J.: Thickness variations and volume estimates of tephra fall deposits: The importance of particle Reynolds number, J. Volcanol. Geotherm. Res., 81, 173–187, [https://doi.org/10.1016/S0377-0273\(98\)00007-9](https://doi.org/10.1016/S0377-0273(98)00007-9), 1998.

3595 Bonadonna, C., Folch, A., Loughlin, S., and Puempel, H.: Future developments in modelling and monitoring of volcanic ash clouds: Outcomes from the first IAVCEI-WMO workshop on Ash Dispersal Forecast and Civil Aviation, Bull. Volcanol., 74, 1–10, <https://doi.org/10.1007/s00445-011-0508-6>, 2012.

Bonadonna, C., Biass, S., Menoni, S., and Gregg, C. E.: Assessment of risk associated with tephra-related hazards, in: *Forecasting and Planning for Volcanic Hazards, Risks, and Disasters, 2*, edited by: Schroeder, J. F., volume edited by, Papale, P., Elsevier, 329–378, <https://doi.org/10.1016/B978-0-12-818082-2.00008-1>, 2021.

3600 Bursik, M., Jones, M., Carn, S., Dean, K., Patra, A., Pavolonis, M., Pitman, E. B., Singh, T., Singla, P., Webley, P., Bjornsson, H., and Ripepe, M.: Estimation and propagation of volcanic source parameter uncertainty in an ash transport and dispersal model: Application to the Eyjafjallajökull plume of 14–16 April 2010, *Bull. Volcanol.*, 74, 2321–2338, <https://doi.org/10.1007/s00445-012-0665-2>, 2012.

3605 *Central Disaster Management Council, Executive Committee for Disaster Prevention: Impacts of Ashfall and Countermeasures in the Tokyo Metropolitan Area: Mt. Fuji Eruption Case (in Japanese)*, Working Group on Ashfall Countermeasures for Large-Scale Eruptions, Cabinet Office, Government of Japan, 2020.

Cer, D., Diab, M., Agirre, E., Lopez-Gazpio, I., and Specia, L.: SemEval-2017 Task 1: Semantic textual similarity – Multilingual and cross-lingual focused evaluation, in: *Proceedings of the 11th International Workshop on Semantic Evaluation (SemEval-2017)*, 11th International Workshop on Semantic Evaluation (SemEval-2017), Vancouver, Canada, 1–14, <https://doi.org/10.18653/v1/S17-2001>, 2017.

3610 *Colorful Scoop*: <https://colorfulcoop.com/>, last access: 27 January 2022.

Daily weather chart: <https://www.data.jma.go.jp/fcd/yoho/hibiten/>, last access: 3 January 2022.

3615 Damby, D. E., Horwell, C. J., Baxter, P. J., Delmelle, P., Donaldson, K., Dunster, C., Fubini, B., Murphy, F. A., Natrass, C., Sweeney, S., Tetley, T. D., and Tomatis, M.: The respiratory health hazard of tephra from the 2010 Centennial eruption of Merapi with implications for occupational mining of deposits, *J. Volcanol. Geotherm. Res.*, 261, 376–387, <https://doi.org/10.1016/j.jvolgeores.2012.09.001>, 2013.

Database of weather charts for hundred years: <http://agora.ex.nii.ac.jp/digital-typhoon/weather-chart/>, last access: 3 January 2022.

Environmental Systems Research Institute (ESRI): ArcGIS Pro 2.7, 2017.

3620 *Environmental Systems Research Institute (ESRI): Human geography base [basemap]*, <https://www.arcgis.com/home/item.html?id=2afe5b807fa74006be6363fd243ffb30>, 2022.

Deep, L. translator: <https://www.deepl.com/translator/>, last access: 27 January 2022.

Devlin, J., Chang, M.-W., Lee, K., and Toutanova, K.: BERT: Pretraining of Deep Bidirectional Transformers for Language Understanding, *ArXiv [eprint]*, arXiv:1810.04805, 2019.

3625 *Eliasson, J., Yoshitani, J., Weber, K., Yasuda, N., Iguchi, M., and Vogel, A.: Airborne measurement in the ash plume from mount Sakurajima: Analysis of gravitational effects on dispersion and fallout*, *Int. J. Atmos. Sci.*, 2014, 1–16, <https://doi.org/10.1155/2014/372135>, 2014.

Fero, J., Carey, S. N., and Merrill, J. T.: Simulation of the 1980 eruption of Mount St. Helens using the ash-tracking model PUFF, *J. Volcanol. Geotherm. Res.*, 175, 355–366, <https://doi.org/10.1016/j.jvolgeores.2008.03.029>, 2008.

- 3630 [Fero, J., Carey, S. N., and Merrill, J. T.: Simulating the dispersal of tephra from the 1991 Pinatubo eruption: Implications for the formation of widespread ash layers, *J. Volcanol. Geotherm. Res.*, 186, 120–131, <https://doi.org/10.1016/j.jvolgeores.2009.03.011>, 2009.](#)
- [Folch, A.: A review of tephra transport and dispersal models: Evolution, current status, and future perspectives, *J. Volcanol. Geotherm. Res.*, 235–236, 96–115, <https://doi.org/10.1016/j.jvolgeores.2012.05.020>, 2012.](#)
- 3635 [Folch, A., Costa, A., Durant, A., and Macedonio, G.: A model for wet aggregation of ash particles in volcanic plumes and clouds: 2. Model application, *J. Geophys. Res.*, 115, B09202, <https://doi.org/10.1029/2009JB007176>, 2010.](#)
- [Folch, A., Costa, A., and Basart, S.: Validation of the FALL3D ash dispersion model using observations of the 2010 Eyjafjallajökull volcanic ash clouds, *Atmos. Environ.*, 48, 165–183, <https://doi.org/10.1016/j.atmosenv.2011.06.072>, 2012.](#)
- 3640 [Hampton, S. J., Cole, J. W., Wilson, G., Wilson, T. M., and Broom, S.: Volcanic ashfall accumulation and loading on gutters and pitched roofs from laboratory empirical experiments: Implications for risk assessment, *J. Volcanol. Geotherm. Res.*, 304, 237–252, <https://doi.org/10.1016/j.jvolgeores.2015.08.012>, 2015.](#)
- [Hattori, Y., Suto, H., Toshida, K., and Hirakuchi, H.: Development of Estimation Method for Tephra Transport and Dispersal Characteristics with Numerical Simulation Technique \(part 1\) – Meteorological Effects on Ash Fallout with Shinmoe-dake Eruption \(in Japanese\), Central Research Institute of Electric Power Industry \(CRIEPI\), Japan, 2013.](#)
- 3645 [Hattori, Y., Suto, H., Toshida, K., and Hirakuchi, H.: Development of Estimation Method for Tephra Transport and Dispersal Characteristics with Numerical Simulation Technique \(part 2\) – A Method of Selecting Meteorological Conditions and the Effects on Ash Deposition and Concentration in Air for Kanto-Area \(in Japanese\), Central Research Institute of Electric Power Industry \(CRIEPI\), Japan, 2016.](#)
- [Hickey, J., Gottsmann, J., Nakamichi, H., and Iguchi, M.: Thermomechanical controls on magma supply and volcanic deformation: Application to Aira caldera, Japan, *Sci Rep. Sci. Rep.*, 6, 32691, <https://doi.org/10.1038/srep32691>, 2016.](#)
- 3650 [Hoyer, S. and Hamman, J. J.: xarray: N-D labeled Arrays and Datasets in Python, *J. Open Res. Softw.*, 5, <https://doi.org/10.5334/jors.148>, 2017.](#)
- [Face, H.: <https://huggingface.co/>, last access: 27 January 2022.](#)
- [Iguchi, M.: Sakurajima Taisho Eruption, 100th Anniversary of the Disaster Prevention in Sakurajima, Kyoto University, Kyoto, Japan, 2014.](#)
- 3655 [Iguchi, M.: Method for real-time evaluation of discharge rate of volcanic ash – Case study on intermittent eruptions at the Sakurajima volcano, Japan, *J. Disaster Res.*, 11, 4–14, <https://doi.org/10.20965/jdr.2016.p0004>, 2016.](#)
- [Iguchi, M., Nakamichi, H., and Tameguri, T.: Integrated study on forecasting volcanic hazards of Sakurajima volcano, Japan, *J. Disaster Res.*, 15, 174–186, <https://doi.org/10.20965/jdr.2020.p0174>, 2020.](#)
- 3660 [Japan's Daily Weather Chart, Japan Meteorological Agency: <https://www.data.jma.go.jp/fcd/yoho/hibiten/>, last access: 3 January 2022.](#)
- [Jenkins, S. F., Spence, R. J. S., Fonseca, J. F. B. D., Solidum, R. U., and Wilson, T. M.: Volcanic risk assessment: Quantifying physical vulnerability in the built environment, *J. Volcanol. Geotherm. Res.*, 276, 105–120, <https://doi.org/10.1016/j.jvolgeores.2014.03.002>, 2014.](#)

- 3665 Jenkins, S. F., Wilson, T. M., Magill, C., Miller, V., Stewart, C., Blong, R., Marzocchi, W., Boulton, M., Bonadonna, C., and Costa, A.: Volcanic ash fall hazard and risk, in: Global Volcanic Hazards and Risk, edited by: Loughlin, S. C., Sparks, S., Brown, S. K., Jenkins, S. F., and Vye-Brown, C., Cambridge University Press, Cambridge, 173–222, <https://doi.org/10.1017/CBO9781316276273.005>, 2015.
- Jenkins, S. F., Magill, C. R., and Blong, R. J.: Evaluating relative tephra fall hazard and risk in the Asia-Pacific region, *Geosphere*, 14, 492–509, <https://doi.org/10.1130/GES01549.1>, 2018.
- 3670 Jurafsky, D. and James, M. H.: *Speech and Language Processing: An Introduction to Natural Language Processing, Computational Linguistics, and Speech Recognition*, Prentice Hall, Upper Saddle River, NJ, 2000.
- Kagoshima City, Crisis Management Division: Sakurajima Volcano Hazard Map, Kagoshima Municipal Government, Kagoshima, Japan, 2010.
- 3675 Kitagawa, K.: Living with an active volcano: Informal and community learning for preparedness in south of Japan, in: Observing the Volcano World, edited by: Fearnley, C. J., Bird, D. K., Haynes, K., McGuire, W. J., and Jolly, G., Springer International Publishing, Cham, 677–689, https://doi.org/10.1007/11157_2015_12, 2015.
- Kobayashi, S., Ota, Y., Harada, Y., Ebita, A., Moriya, M., Onoda, H., Onogi, K., Kamahori, H., Kobayashi, C., Endo, H., Miyaoka, K., and Takahashi, K.: The JRA-55 reanalysis: General specifications and basic characteristics, *J. Meteorol. Soc. Jpn.*, 93, 5–48, <https://doi.org/10.2151/jmsj.2015-001>, 2015.
- 3680 Kobayashi, T.: Geology of Sakurajima volcano: A review, *Bull. Volcanol. Soc. Jpn.*, 27, 277–292, 1982.
- Koto, B.: The great eruption of Sakura-Jima, *J. Coll. Sci. Imperial Univ. Tokyo*, 38, 1–237, 1916.
- Kratzmann, D. J., Carey, S. N., Fero, J., Scasso, R. A., and Naranjo, J.-A.: Simulations of tephra dispersal from the 1991 explosive eruptions of Hudson volcano, Chile, *Journal of Volcanology and Geothermal Research*, 190, 337–352, <https://doi.org/10.1016/j.jvolgeores.2009.11.021>, 2010.
- 3685 Kyushu Regional Development Bureau and Osumi Office of the Rivers and National Highways: Sakurajima volcano wide area disaster prevention map (in Japanese), Minist. Land Infrastruct. Transp. Tourism Sakurajima, Kagoshima, Japan, 2017.
- Lin, J., Brunner, D., Gerbig, C., Stohl, A., Luhar, A., and Webley, P.: Lagrangian modeling of the atmosphere: An introduction, in: *Lagrangian Modeling of the Atmosphere*, American Geophysical Union, Washington, DC, USA, 1–15, 2012.
- 3690 Macedonio, G., Costa, A., Scollo, S., and Neri, A.: Effects of eruption source parameter variation and meteorological dataset on tephra fallout hazard assessment: Example from Vesuvius (Italy), *J. Appl. Volcanol.*, 5, 5, <https://doi.org/10.1186/s13617-016-0045-2>, 2016.
- 3695 Madankan, R., Pouget, S., Singla, P., Bursik, M., Dehn, J., Jones, M., Patra, A., Pavolonis, M., Pitman, E. B., Singh, T., and Webley, P.: Computation of probabilistic hazard maps and source parameter estimation for volcanic ash transport and dispersion, *J. Comput. Phys.*, 271, 39–59, <https://doi.org/10.1016/j.jcp.2013.11.032>, 2014.
- Maeda, S., Hirahara, S., Hagiya, S., Murai, H., and Oikawa, Y., 05 March 2012: Forecasting Changes in Seasonal Progress in Future Climate (based on the changes of Westerly Wind) (in Japanese): <http://wind.gp.tohoku.ac.jp/yamase/reports/meeting5.html>, Climate Information Division, Global Environment and Ocean Department, Japan Meteorological Agency.

- 3700 Mastin, L. G., Guffanti, M., Servranckx, R., Webley, P., Barsotti, S., Dean, K., Durant, A., Ewert, J. W., Neri, A., Rose, W. I., Schneider, D., Siebert, L., Stunder, B., Swanson, G., Tupper, A., Volentik, A., and Waythomas, C. F.: A multidisciplinary effort to assign realistic source parameters to models of volcanic ash-cloud transport and dispersion during eruptions, *J. Volcanol. Geotherm. Res.*, 186, 10–21, <https://doi.org/10.1016/j.jvolgeores.2009.01.008>, 2009.
- 3705 Mita, K., Ishimine, Y., Iwamatsu, A., Kamikozuru, H., Koga, M., Takeda, N., Nakamura, T., Matsumoto, S., Iguchi, M., Baba, M., Yamaguchi, K., Shimokawa, E., and Maki, M.: Tentative Reports on the Traffic Countermeasures for a Large-Scale Eruption of Sakurajima Volcano. Case Study: Ash Deposit on Kagoshima City (in Japanese), *Earthquake and Volcanic Disaster Prevention Center, Kagoshima University*, 2018.
- Miyaji, N., Kan'no, A., Kanamaru, T., and Mannen, K.: High-resolution reconstruction of the Hiei eruption (AD 1707) of Fuji volcano, Japan, *J. Volcanol. Geotherm. Res.*, 207, 113–129, <https://doi.org/10.1016/j.jvolgeores.2011.06.013>, 2011.
- 3710 Morton, B. R., Taylor, S. G., and Turner, J. S.: Turbulent gravitational convection from maintained and instantaneous sources: Proceedings of the R. Soc. Lond. A Mat., 234, 1–23, <https://doi.org/10.1098/rspa.1956.0011>, 1956.
- Newhall, C. G. and Self, S.: The volcanic explosivity index (VEI) an estimate of explosive magnitude for historical volcanism, *J. Geophys. Res.*, 87, 1231, <https://doi.org/10.1029/JC087iC02p01231>, 1982.
- 3715 Nishimura, T.: Source mechanisms of volcanic explosion earthquakes: Single force and implosive sources, *J. Volcanol. Geotherm. Res.*, 86, 97–106, [https://doi.org/10.1016/S0377-0273\(98\)00088-2](https://doi.org/10.1016/S0377-0273(98)00088-2), 1998.
- Omori, F.: The Sakura-Jima eruption and earthquakes, *Bulletin of the Imperial Earthquake Investigation Committee, University of Tokyo*, 8, 1–630, 1914.
- Peng, J. and Peterson, R.: Attracting structures in volcanic ash transport, *Atmos. Environ.*, 48, 230–239, <https://doi.org/10.1016/j.atmosenv.2011.05.053>, 2012.
- 3720 Peterson, R. and Dean, K.: Analysis and Optimization of a Lagrangian Volcanic Ash Particle Tracking Model called Puff, *AGU Fall Meet. Admin.*, 2002.
- Poulidis, A. P. and Takemi, T.: A 1998–2013 climatology of Kyushu, Japan: Seasonal variations of stability and rainfall, *Int. J. Climatol.*, 37, 1843–1858, <https://doi.org/10.1002/joc.4817>, 2017.
- 3725 Poulidis, A. P., Takemi, T., Iguchi, M., and Renfrew, I. A.: Orographic effects on the transport and deposition of volcanic ash: A case study of Mount Sakurajima, Japan, *J. Geophys. Res. Atmos.*, 122, 9332–9350, <https://doi.org/10.1002/2017JD026595>, 2017.
- Poulidis, A. P., Takemi, T., Shimizu, A., Iguchi, M., and Jenkins, S. F.: Statistical analysis of dispersal and deposition patterns of volcanic emissions from MT. Sakurajima, Japan, *Atmos. Environ.*, 179, 305–320, <https://doi.org/10.1016/j.atmosenv.2018.02.021>, 2018.
- 3730 Rahadianto, H. and Tatano, H.: 62 Years Simulated Sakurajima Taisho Eruption Ashfall Deposit Data (1958-2019), *Disaster Prevention Research Institute, Kyoto University, Japan [dataset]*, <https://doi.org/10.17603/ds2-vw5f-t920>, 2020.
- 3735 Rathje, E. M., Dawson, C., Padgett, J. E., Pinelli, J.-P., Stanzione, D., Adair, A., Arduino, P., Brandenberg, S. J., Cockerill, T., Dey, C., Esteva, M., Haan, F. L., Hanlon, M., Kareem, A., Lowes, L., Mock, S., and Mosqueda, G.: DesignSafe: New cyberinfrastructure for natural hazards engineering, *Nat. Hazards Rev.*, 18, 06017001, [https://doi.org/10.1061/\(ASCE\)NH.1527-6996.0000246](https://doi.org/10.1061/(ASCE)NH.1527-6996.0000246), 2017.

Reimers, N. and Gurevych, I.: Sentence-BERT: Sentence Embeddings Using Siamese BERT-Networks, ArXiv [eprint], arXiv:1908.10084, 2019.

Reimers, N. and Gurevych, I.: Making Monolingual Sentence Embeddings Multilingual Using Knowledge Distillation, ArXiv [eprint], arXiv:2004.09813, 2020.

3740 Scollo, S., Prestifilippo, M., Coltelli, M., Peterson, R. A., and Spata, G.: A statistical approach to evaluate the tephra deposit and ash concentration from PUFF model forecasts, *J. Volcanol. Geotherm. Res.*, 200, 129–142, <https://doi.org/10.1016/j.jvolgeores.2010.12.004>, 2011.

Searcy, C., Dean, K., and Stringer, W.: PUFF: A high-resolution volcanic ash tracking model, *J. Volcanol. Geotherm. Res.*, 80, 1–16, [https://doi.org/10.1016/S0377-0273\(97\)00037-1](https://doi.org/10.1016/S0377-0273(97)00037-1), 1998.

3745 Selva, J., Costa, A., De Natale, G., Di Vito, M. A., Isaia, R., and Macedonio, G.: Sensitivity test and ensemble hazard assessment for tephra fallout at Campi Flegrei, Italy, *J. Volcanol. Geotherm. Res.*, 351, 1–28, <https://doi.org/10.1016/j.jvolgeores.2017.11.024>, 2018.

3750 Shimbori, T., Aikawa, Y., Fukui, K., Hashimoto, A., Seino, N., and Yamasato, H.: Quantitative tephra fall prediction with the JMA mesoscale tracer transport model for volcanic ash: A case study of the eruption at Asama volcano in 2009 (in Japanese), *Pap. Meteorol. Geophys.*, 61, 13–29, <https://doi.org/10.2467/mripapers.61.13>, 2009.

Singhal, A.: *Modern Information Retrieval: A Brief Overview*, 9, 2001.

Takebayashi, M., Onishi, M., and Iguchi, M.: Large volcanic eruptions and their influence on air transport: The case of Japan, *J. Air Transp. Manag.*, 97, 102136, <https://doi.org/10.1016/j.jairtraman.2021.102136>, 2021.

3755 Tanaka, H. L.: Development of A prediction scheme for volcanic ash fall from redoubt volcano, Alaska, in: *Proceedings of the First International Symposium on Volcanic Ash and Aviation Safety, First International Symposium on Volcanic Ash and Aviation Safety*, 283–291, <https://doi.org/10.3133/b2047>, 1994.

Tanaka, H. L. and Iguchi, M.: Numerical simulations of volcanic ash plume dispersal for Sakura-Jima using real-time emission rate estimation, *J. Disaster Res.*, 14, 160–172, <https://doi.org/10.20965/jdr.2019.p0160>, 2019.

3760 Tanaka, H. L. and Yamamoto, K.: Numerical simulation of volcanic plume dispersal from usu volcano in Japan on 31 March 2000 using PUFF model, *Earth Planets Space*, 54, 743–752, <https://doi.org/10.1186/BF03351727>, 2002.

Tanaka, H. L., Iguchi, M., and Nakada, S.: Numerical simulations of volcanic ash plume dispersal from Kelud volcano in Indonesia on February 13, 2014, *J. Disaster Res.*, 11, 31–42, <https://doi.org/10.20965/jdr.2016.p0031>, 2016.

Tilling, R. I., Topinka, L. J., Swanson, D. A., and Survey (U.S.), G.: *Eruptions of Mount St. Helens: Past, Present, and Future*, US Department of the Interior, Geological Survey, 1990.

3765 Todde, A., Cioni, R., Pistolesi, M., Geshi, N., and Bonadonna, C.: The 1914 Taisho eruption of Sakurajima volcano: Stratigraphy and dynamics of the largest explosive event in Japan during the twentieth century, *Bull. Volcanol.*, 79, 72, <https://doi.org/10.1007/s00445-017-1154-4>, 2017.

Tsukui, M.: Ash-fall distribution of 1779 an'ei eruption, Sakurajima volcano: Revealed by historical documents (in Japanese), *Bull. Volcanological Soc. Jpn.*, 56, 89–94, https://doi.org/10.18940/kazan.56.2-3_89, 2011.

3770 [Unidata: NetCDF, UCAR/Unidata, https://doi.org/10.5065/D6H70CW6, 2021.](https://doi.org/10.5065/D6H70CW6)

[Wallace, J. M. and Hobbs, P. V.: Atmospheric Science: An Introductory Survey, 2nd ed, Academic Press, Amsterdam, Paris, 2006.](#)

[Webley, P. and Mastin, L.: Improved prediction and tracking of volcanic ash clouds, J. Volcanol. Geotherm. Res., 186, 1–9, https://doi.org/10.1016/j.jvolgeores.2008.10.022, 2009.](https://doi.org/10.1016/j.jvolgeores.2008.10.022)

3775 [Wilson, T. M., Stewart, C., Sword-Daniels, V., Leonard, G. S., Johnston, D. M., Cole, J. W., Wardman, J., Wilson, G., and Barnard, S. T.: Volcanic ash impacts on critical infrastructure, Phys. Chem. Earth Parts A B C, 45–46, 5–23, https://doi.org/10.1016/j.pce.2011.06.006, 2012.](https://doi.org/10.1016/j.pce.2011.06.006)

[Yamasato, H., Funasaki, J., and Takagi, Y.: The Japan Meteorological Agency's Volcanic Disaster Mitigation Initiatives, National Research Institute for Earth Science and Disaster Prevention, Japan, 2013.](#)

3780 [Yasui, M., Takahashi, M., Ishihara, K., and Miki, D.: Records on the 1914–1915 eruption of Sakurajima volcano, Japan \(in Japanese\), Bull. Volcanological Soc. Jpn, 41, 75–107, 2006.](#)

[Yasui, M., Takahashi, M., Ishihara, K., and Miki, D.: Eruptive Style and Its Temporal Variation through the 1914–1915 Eruption of Sakurajima Volcano, 52, Southern Kyushu, Japan \(in Japanese\), Bulletin of The Volcanological Society of Japan, 161–186, https://doi.org/10.18940/kazan.52.3_161, 2007.](https://doi.org/10.18940/kazan.52.3_161)

3785 [Yoshikushi, T., Kawahara, D., and Kurohara, S.: Multilingualization of a Natural Language Inference Dataset Using Machine Translation, Kyoto University, 2020.](#)

[Zuccaro, G., Leone, M. F., Del Cogliano, D., and Sgroi, A.: Economic impact of explosive volcanic eruptions: A simulation-based assessment model applied to Campania region volcanoes, J. Volcanol. Geotherm. Res., 266, 1–15, https://doi.org/10.1016/j.jvolgeores.2013.09.002, 2013.](https://doi.org/10.1016/j.jvolgeores.2013.09.002)

- Deleted:** ¶
 Biass, S., Scaini, C., Bonadonna, C., Folch, A., Smith, K., and Höskuldsson, A.: A multi-scale risk assessment for tephra fallout and airborne concentration from multiple Icelandic volcanoes – Part 1: Hazard assessment, Nat. Hazards Earth Syst. Sci., 14, 2265–2287, <https://doi.org/10.5194/nhess-14-2265-2014>, 2014.¶
 Biass, S., Bonadonna, C., di Traglia, F., Pistolesi, M., Rosi, M., and Lestuzzi, P.: Probabilistic evaluation of the physical impact of future tephra fallout events for the Island of Vulcano, Italy, Bull. Volcanol, 78, 37, <https://doi.org/10.1007/s00445-016-1028-1>, 2016. Central Disaster Management Council, Executive Committee for Disaster Prevention: Impacts of Ashfall and Countermeasures in the Tokyo Metropolitan Area: Mt. Fuji Eruption Case (in Japanese), Working Group on Ashfall Countermeasures for Large-Scale Eruptions, Cabinet Office, Government of Japan, 2020.¶
 Environmental Systems Research Institute (ESRI): ArcGIS Pro 2.7, 2017.¶
 Environmental Systems Research Institute (ESRI): Human geography base [basemap], <https://www.arcgis.com/home/item.html?id=2afe5b807fa74006b>...
- Deleted: Figures Caption**¶
 Figure 1: The arrival time of ashfall during the Taisho eruption event throughout Japan. The blue dashed line indicates the ...
- Formatted:** Justified
- Moved down [7]:** The probability distribution of ash dispersal with the threshold for airport closure risk (≥ 0.02 cm). Airport locations are marked by black aeroplane icons of ...
- Moved down [14]:** The flowchart of the validation procedure using the dataset produced in this paper. (1) the process of digitising the original ash precipitation map to become new ...
- Moved up [18]:** Date
- Moved (insertion) [1]** ...
- Moved up [1]:** The weather chart on the Taisho eruption with additional markings of pressure measurement converted to the latest convention (mmHg \rightarrow hPa). Modified from Omori (1914).
- Moved down [11]:** The comparison of the ash distribution coverage between the original ash dispersal map and the simulation results on the selected date (5 March 1997). The red ...
- Moved up [2]:** The general weather report commentary on the weather condition inside the weather chart. (a) The old weather chart put the commentary on the bottom of the second page on ...
- Moved (insertion) [4]**
- Moved up [3]:** (a) Surface weather chart for Asia Pacific region at 00 UTC 5 March 1997, and (b) Surface weather chart for Asia Pacific region at 00 UTC 24 February 2006. Both ...
- Formatted Table**
- Moved up [12]:** Actual Ash Precipitation of the Taisho Eruption
- Moved (insertion) [7]** ...
- Formatted Table**
- Formatted** ...