

# 1 The UKSCAPE-G2G river flow and soil moisture datasets: 2 Grid-to-Grid model estimates for the UK for historical and 3 potential future climates

4 Alison L Kay, Victoria A Bell, Helen N Davies, Rosanna A Lane, Alison C

5 Rudd

6 UK Centre for Ecology & Hydrology, Wallingford, UK, OX10 8BB

7 Correspondence to: A.L. Kay ([alkay@ceh.ac.uk](mailto:alkay@ceh.ac.uk))

## 8 Abstract

9 Appropriate adaptation planning is contingent upon information about the potential  
10 future impacts of climate change, and hydrological impact assessments are of  
11 particular importance. The UKSCAPE-G2G datasets were produced, as part of the  
12 NERC UK-SCAPE programme, to contribute to this information requirement. They  
13 use the Grid-to-Grid (G2G) national-scale hydrological model configured for both  
14 Great Britain and Northern Ireland (and the parts of the Republic of Ireland that drain  
15 to rivers in NI). Six separate datasets are provided, for two sets of driving data —  
16 one observation-based (1980–2011) and one climate projection-based (1980–2080)  
17 — for both river flows and soil moisture on 1 km x 1 km grids across GB and NI. The  
18 river flow datasets include grids of monthly mean flow, annual maxima of daily mean  
19 flow, and annual minima of 7-day mean flow ( $\text{m}^3\text{s}^{-1}$ ). The soil moisture datasets are  
20 grids of monthly mean soil moisture content (m water / m soil), which should be  
21 interpreted as depth-integrated values for the whole soil column. The climate  
22 projection-based datasets are produced using data from the 12-member 12km  
23 regional climate model ensemble of the latest UK climate projections (UKCP18),  
24 which uses RCP8.5 emissions. The production of the datasets is described, along  
25 with details of the file format, and how the data should be used. Example maps are  
26 provided, as well as simple UK-wide analyses of the various outputs. These suggest  
27 potential future decreases in summer flows, annual minimum 7-day flows, and  
28 summer/autumn soil moisture, along with possible future increases in winter flows  
29 and annual maximum flows. References are given for published papers providing  
30 more detailed spatial analyses, and some further potential uses of the data are  
31 suggested.

## 32 Keywords

33 Climate change; hydrological impacts; rainfall-runoff; UK Climate Projections 2018;  
34 UKCP18

## 35 1 Introduction

36 Information on the potential future impacts of climate change is crucial to enable  
37 appropriate adaptation planning, and impacts on the hydrological cycle and river  
38 flows are one of the main ways by which climate change will affect both society and  
39 the natural environment. UK-SCAPE (UK Status, Change And Projections of the  
40 Environment; [ukscape.ceh.ac.uk](http://ukscape.ceh.ac.uk)) is a five-year programme funded by the Natural  
41 Environment Research Council (NERC) as part of a National Capability Science  
42 Single Centre award, and the main aim of Work Package 2.2 of UK-SCAPE is to

43 deliver data and analyses showing how future climate change could influence water  
44 quantity. The hydrological datasets presented here were produced as part of UK-  
45 SCAPE WP2.2.

46 The datasets consist of 1 km x 1 km gridded outputs from a national-scale  
47 hydrological model (Grid-to-Grid), and include both river flows and soil moisture, for  
48 Great Britain (GB) and Northern Ireland (NI). The model has been driven with  
49 observation-based data, and with an ensemble of Regional Climate Model (RCM)  
50 data from the latest climate projections for the UK (UKCP18; Lowe et al. 2018). A  
51 summary of the six available datasets, including references, is provided in Table 1.  
52 The datasets have been used within UK-SCAPE WP2.2 to support analyses of  
53 potential future changes in river flows and soil moisture (Kay 2021, Lane & Kay  
54 2021, Kay et al. 2021a, Kay et al. 2022a), but could also be used to support other  
55 hydrological research and wider studies such as ecological and agricultural  
56 modelling.

57 Section 2 describes how the datasets were produced, including the hydrological  
58 model and the driving data applied. Section 3 presents some high-level analyses of  
59 the datasets, and describes the results of more detailed analyses presented in other  
60 published papers. Section 4 discusses potential uses and caveats, with conclusions  
61 in Section 5.

62

63 **Table 1 Summary of the six UKSCAPE-G2G datasets.**

---

<b>Observation-driven</b>	
River flow	GB: <a href="https://doi.org/10.5285/2f835517-253e-4697-b774-ab6ff2c0d3da">https://doi.org/10.5285/2f835517-253e-4697-b774-ab6ff2c0d3da</a> (Kay et al. 2021c) NI: <a href="https://doi.org/10.5285/f5fc1041-e284-4763-b8b7-8643c319b2d0">https://doi.org/10.5285/f5fc1041-e284-4763-b8b7-8643c319b2d0</a> (Kay et al. 2021d)
Soil moisture	GB and NI: <a href="https://doi.org/10.5285/c9a85f7c-45e2-4201-af82-4c833b3f2c5f">https://doi.org/10.5285/c9a85f7c-45e2-4201-af82-4c833b3f2c5f</a> (Kay et al. 2021e)
<b>Climate projection-driven</b>	
River flow	GB: <a href="https://doi.org/10.5285/18be3704-0a6d-4917-aa2e-bf38927321c5">https://doi.org/10.5285/18be3704-0a6d-4917-aa2e-bf38927321c5</a> (Kay et al. 2022b) NI: <a href="https://doi.org/10.5285/76057b0a-b18f-496f-891c-d5b22bd0b291">https://doi.org/10.5285/76057b0a-b18f-496f-891c-d5b22bd0b291</a> (Kay et al. 2022c)
Soil moisture	GB and NI: <a href="https://doi.org/10.5285/f7142ced-f6ff-486b-af33-44fb8f763cde">https://doi.org/10.5285/f7142ced-f6ff-486b-af33-44fb8f763cde</a> (Kay et al. 2022d)

---

64

## 65 **2 Data production methods**

### 66 **2.1 The hydrological model**

67 The Grid-to-Grid (G2G) is a national-scale grid-based hydrological model which  
68 typically operates on a 1km x 1km grid at a 15-minute time-step (Bell et al. 2009),  
69 with an optional snow module (Bell et al. 2016). It was originally configured to cover  
70 Great Britain (GB), on a spatial domain aligned with the GB national grid, but more  
71 recently a version was configured to cover Northern Ireland (NI) and areas in the  
72 Republic of Ireland (RoI) that drain into NI, also on a domain aligned with the GB  
73 national grid (Kay et al. 2021a). The G2G is configured using spatial datasets (e.g.  
74 soil types, land-cover, flow directions), in preference to parameter identification via

75 calibration; where model parameters are required (such as the wave speeds used in  
76 lateral routing) nationally-applicable values are applied (Bell et al. 2009).

77 G2G has been shown to perform well for a wide range of catchments across GB and  
78 NI, including for the modelling of high flows / floods and low flows / droughts (Bell et  
79 al. 2009, 2016; Rudd et al. 2017; Formetta et al. 2018; Kay et al. 2021a,b). This is  
80 particularly the case for catchments with more natural flow regimes, as the model  
81 does not routinely account for artificial influences like abstractions/discharges  
82 (Rameshwaran et al. 2022). While the effect of urban/suburban land-cover on runoff  
83 is accounted for, the effect of lake/reservoir storage and regulation is generally  
84 neglected at the national scale; lake grid-cells are treated as though they were rivers.  
85 This has a minimal effect across most of GB; the largest lake in Scotland, Loch  
86 Lomond, has an area of  $\sim 71\text{km}^2$ , and the largest lake in England, Windermere, has  
87 an area of  $\sim 15\text{km}^2$ . But in NI the dominant presence of Lough Neagh ( $\sim 390\text{km}^2$ )  
88 limits model performance downstream (the Lower Bann river; Kay et al. 2021a), and  
89 Lough Erne in the south-west of NI is also relatively large (Upper and Lower Lough  
90 Erne have a combined area of  $\sim 144\text{km}^2$ ).

## 91 **2.2 Observation-based driving data**

92 Gridded time-series of precipitation and potential evaporation (PE) are required to  
93 drive the G2G, plus temperature for the snow module. The observation-based driving  
94 data are applied as follows:

- 95 • Daily 1km grids of precipitation (CEH-GEAR; Tanguy et al. 2016) are divided  
96 equally over each model time-step within a day. For use in NI they are first re-  
97 projected from the Irish national grid to the GB national grid.
- 98 • Monthly 40km grids of PE for short grass (MORECS; Hough and Jones 1997) are  
99 copied down to the 1km grid, then divided equally over each model time-step  
100 within a month. For use in NI they are first re-projected from the Irish national grid  
101 to the GB national grid. The data do not cover all the required parts of the UK, so  
102 have been extended where necessary (i.e. some coastal areas and some parts of  
103 the RoI that drain into NI) by copying from the nearest cell with data.
- 104 • Daily 1km grids of min and max temperature (HadUK-Grid; Met Office 2019) are  
105 interpolated through the day using a sine curve (Kay and Crooks 2014). The data  
106 do not cover the required parts of the RoI, so have been infilled from the nearest  
107 cell with data, using a lapse rate with elevation data (Morris and Flavin 1990).

## 108 **2.3 Climate projection-based driving data**

109 The climate change simulations use data from the UKCP18 Regional projections  
110 (Met Office Hadley Centre 2018). These comprise a 12-member perturbed  
111 parameter ensemble (PPE) of the Hadley Centre  $\sim 12\text{km}$  Regional Climate Model  
112 (RCM) nested in an equivalent PPE of their  $\sim 60\text{km}$  Global Climate model (GCM)  
113 (Murphy et al. 2018). Ensemble member 01 represents the standard  
114 parameterisation, with members 02-15 representing a range of credible variations in  
115 parameters (note that there are no RCM equivalents for GCM PPE members 02, 03  
116 and 14). The data cover Dec 1980–Nov 2080 under just one emissions scenario,  
117 RCP8.5 (Riahi et al. 2011), and have a 360-day year (twelve 30-day months). The  
118 data are available re-projected from the native climate model grid onto a 12km grid  
119 aligned with the GB national grid – the latter are used here.

120 The climate projection data are applied as follows:

- 121 • Daily 12km grids of precipitation are directly available from the UKCP18 Regional  
122 projections. These are first adjusted for bias using 12km grids of monthly  
123 correction factors derived by comparing baseline values against CEH-GEAR data  
124 averaged up to the 12km resolution (Kay 2021, Kay et al. 2021a). They are then  
125 spatially downscaled to 1km using patterns of average annual rainfall (1961–  
126 1990; Bell et al. 2007), and divided equally over each model time-step within a  
127 day (as for observed data).
- 128 • Daily 12km grids of PE are not directly available from the UKCP18 Regional  
129 projections. Instead, they are calculated from other meteorological variables in a  
130 way which closely replicates MORECS (as in Robinson et al. (2021, 2022), but  
131 using the bias-adjusted precipitation in the interception component). PE is only  
132 estimated for 12km 'land' RCM boxes; where PE is required for boxes classed as  
133 'sea' in the RCM, it is copied from the nearest 12km 'land' box. The method also  
134 includes increased stomatal resistance under future higher atmospheric CO<sub>2</sub>  
135 concentrations (Rudd and Kay 2016, Guillod et al. 2018). The 12km PE are  
136 copied down to the 1km grid, then divided equally over each model time-step  
137 within a month (as for observed data).
- 138 • Daily 12km grids of min and max temperature are directly available from the  
139 UKCP18 Regional projections. These are downscaled to 1km using a lapse rate  
140 with elevation data, and interpolated through the day using a sine curve (as for  
141 observed data).

## 142 **2.4 Hydrological model runs and outputs**

143 The observation-based simulation (hereafter 'SIMOBS') is initialised using a states  
144 file saved at the end of a prior observation-based run (Jan 1970–Nov 1980). The  
145 same state initialisation file is used for each RCM-based simulation (hereafter  
146 'SIMRCM').

147 Model outputs consist of 1km x 1km gridded time-series of

- 148 • monthly mean river flow (m<sup>3</sup>s<sup>-1</sup>);
- 149 • annual maxima (AMAX) of daily mean river flow (m<sup>3</sup>s<sup>-1</sup>), for water years  
150 (October–September);
- 151 • annual minima (AMIN) of 7-day mean river flow (m<sup>3</sup>s<sup>-1</sup>), for years spanning  
152 December–November; and
- 153 • monthly mean soil moisture content (m water / m soil).

154 While it is possible to output 1km x 1km gridded daily time-series from G2G, these  
155 are not typically produced as they are very large files (especially if long time periods  
156 are covered, as is the case for the SIMRCM runs). Instead, the AMAX and AMIN  
157 flows are calculated and saved during the model run, to enable analyses of high and  
158 low flows without saving daily gridded flows. The AMAX of daily mean flows are  
159 extracted for water years (Oct–Sep), to try to avoid extraction of the same high flow  
160 event from consecutive years. AMIN extraction would usually use calendar years,  
161 but Dec-Nov is used here to match the climate model data running from December  
162 1980 to November 2080, whilst still trying to avoid extraction of the same low flow  
163 event from consecutive years.

164 The flow variables are provided for all non-sea and non-tidal 1km cells with a  
165 catchment drainage area of at least 50km<sup>2</sup>, while the soil moisture is provided for all  
166 non-sea 1km cells. G2G soil moisture estimates are provided as monthly averages  
167 of daily mean soil moisture in the unsaturated zone, which can be interpreted as

168 volumetric soil moisture content,  $\theta$ , where  $0 \leq \theta \leq 1$ . In G2G soil depth can vary from  
 169 a few centimetres to several metres, and soil moisture estimates should be  
 170 interpreted as depth-integrated values for the whole soil column.

171 **2.5 Format of the gridded datasets**

172 The 1km x 1km gridded data are provided as a NetCDF4 file for each variable,  
 173 following UKCEH gridded dataset conventions. The file naming convention is  
 174 described in Table 2 for the observation-based datasets and Table 3 for the climate  
 175 projection-based datasets.

176

177 **Table 2 The file naming convention for the observation-based datasets.**

Data	Names of NetCDF files	Years available
monthly mean river flow	G2G_GB_mmflow_obs_1980_2011.nc	Dec 1980–
	G2G_NI_mmflow_obs_1980_2011.nc	Nov 2011
annual maxima of daily mean river flow	G2G_GB_amaxflow_obs_1980_2011.nc	Oct 1981–
	G2G_NI_amaxflow_obs_1980_2011.nc	Sep 2011
annual minima of 7-day mean river flow	G2G_GB_aminflow_obs_1980_2011.nc	Dec 1980–
	G2G_NI_aminflow_obs_1980_2011.nc	Nov 2011
monthly mean soil moisture content	G2G_GB_mmsoil_obs_1980_2011.nc	Dec 1980–
	G2G_NI_mmsoil_obs_1980_2011.nc	Nov 2011

178

179 **Table 3 The file naming convention for the climate projection-based datasets.**

Data	Names of NetCDF files	Years available
monthly mean river flow	G2G_GB_mmflow_UKCP18RCM_ensnum_1980_2080.nc	Dec 1980–
	G2G_NI_mmflow_UKCP18RCM_ensnum_1980_2080.nc	Nov 2080
annual maxima of daily mean river flow, and dates of occurrence	G2G_GB_amaxflow_UKCP18RCM_ensnum_1980_2080.nc	Oct 1981–
	G2G_NI_amaxflow_UKCP18RCM_ensnum_1980_2080.nc	Sep 2080
annual minima of 7-day mean river flow, and dates of occurrence	G2G_GB_aminflow_UKCP18RCM_ensnum_1980_2080.nc	Dec 1980–
	G2G_NI_aminflow_UKCP18RCM_ensnum_1980_2080.nc	Nov 2080
monthly mean soil moisture content	G2G_GB_mmsoil_UKCP18RCM_ensnum_1980_2080.nc	Dec 1980–
	G2G_NI_mmsoil_UKCP18RCM_ensnum_1980_2080.nc	Nov 2080

ensnum is the number of the ensemble member (01, 04, 05, 06, 07, 08, 09, 10, 11, 12, 13, 15)

180

181 For the observation-based datasets, the time stamp in the NetCDF files is “days  
 182 since 1900-01-01”, and the monthly mean river flows and monthly mean soil  
 183 moisture are nominally assigned to the first day of the month. The annual  
 184 maximum/minimum flow values are nominally assigned to the start year of the 12-  
 185 month period over which they are calculated, e.g. the annual maximum flow  
 186 assigned to 1981 is for 1/10/1981–30/9/1982 (water years), while the annual  
 187 minimum flow assigned to 1981 is for 1/12/1981–30/11/1982 (Dec–Nov years). The  
 188 ‘time\_bnds’ variable gives the start and end dates of the time period over which the  
 189 annual maximum or minimum flow are extracted.

190 For the climate projection-based datasets, the data have 30-day months due to the  
 191 “360\_day” calendar of the Hadley Centre climate model. The files are otherwise as  
 192 above, except that the dates of occurrence of the annual maximum and minimum



193 flows are also provided, as additional variables in the 'amaxflow' and 'aminflow'  
194 NetCDF files respectively.

195 Table 4 summarises the spatial domains covered by the GB and NI datasets. River  
196 flows are only provided for non-sea and non-tidal river cells with a catchment area of  
197 at least 50km<sup>2</sup>, and set to missing elsewhere. Soil moisture estimates are provided  
198 for all non-sea cells, and set to missing elsewhere.

199

200 **Table 4 Summary of domain sizes and extents, including the OSGB co-ordinates for**  
201 **the lower left and upper right corners (m).**

	GB	NI
Domain size	700 km × 1000 km	187 km × 170 km
Lower left corner	(0,0)	(-7000,440000)
Upper right corner	(700000,1000000)	(180000,610000)

202


203 To aid use of the datasets, further data files are provided for both GB and NI,  
204 including catchment area grids, grids identifying majority lake cells, and grids  
205 identifying the approximate locations of river flow gauging stations (Table 5). The  
206 catchment area grids are mapped in Figure 1, while the majority lake cells and  
207 gauging station locations are mapped in Figure 2 (note that although GB and NI are  
208 mapped together, the data for GB and NI are provided separately). At the gauging  
209 station locations the G2G flow estimates can be compared to observed river flows.

210

211 **Table 5 The additional data files for GB and NI.**

<b>Data</b>	<b>File names</b>	<b>Description</b>
Catchment area grid	UKSCAPE_G2G_GB_CatchmentAreaGrid.nc UKSCAPE_G2G_NI_CatchmentAreaGrid.nc	Digitally-derived catchment area (km <sup>2</sup> ) draining to every 1km x 1km grid box.
Majority lake cells grid	UKSCAPE_G2G_GB_SoilMoisture_LakeGrid.nc UKSCAPE_G2G_NI_SoilMoisture_LakeGrid.nc  UKSCAPE_G2G_NI_LakeGrid.nc	Cells with greater than 85% of area covered by water (according to 25m data from Land Cover Map 2015, Rowland et al. 2017). These grids can be applied to exclude use of soil moisture data in majority lake cells.  1=land, 2=lake, and -9999=sea. As above but cells with greater than 70% of area covered by water, plus some manual additions of cells for Lough Erne to avoid more than one change from river to lake to river for each flow pathway. This grid can be applied to exclude use of river flow data in lake cells in NI.
Gauging station location grid	UKSCAPE_G2G_GB_NRFASStationIDGrid.nc UKSCAPE_G2G_NI_NRFASStationIDGrid.nc	Best locations corresponding to 1038 gauging stations in GB and 43 gauging stations in NI, referenced by NRFA station number (nrfa.ceh.ac.uk). NRFA station number at gauging station locations, 0=land, and -9999=sea.
Gauging station info	UKSCAPE_GB_NRFASStationIDs.csv UKSCAPE_NI_NRFASStationIDs.csv	Information on stations included in location grids. Information for 18 additional stations is included in the GB file; these are each located in the same 1km cell as one of the stations in the grid (as detailed in the comments column of the csv file).

212




214

215

216

**Figure 1 Map showing the catchment area grids for GB and NI (see Table 5).**





218

219

220

221

222

**Figure 2** Map showing the majority lake cells and gauging station locations for GB and NI (see Table 5), along with the main rivers (catchment area  $\geq 50\text{km}^2$ ; blue lines). Note that the 'majority lake cells (>70% water)' cover small areas around Lough Neagh and Lough Erne in Northern Ireland only.

## 223 2.6 How to use the datasets

224 River flows from the observation-based simulation can be compared to observed  
225 (gauged) river flows (for example from the National River Flow Archive, NRFA;  
226 [nrfa.ceh.ac.uk](http://nrfa.ceh.ac.uk)), and to facilitate such comparisons files identifying gauging station  
227 locations on the 1km G2G model grid for GB and NI are provided (see Table 5).  
228 However, it should be borne in mind that G2G provides natural flow estimates, so  
229 comparisons in catchments affected by artificial influences like abstractions and  
230 discharges may not be as good as those in catchments with relatively natural flow  
231 regimes (Rameshwaran et al. 2022). Also, although the gauging station locations  
232 have been identified as the G2G cell closest in terms of geographical location and  
233 catchment area, and checks have been undertaken to ensure that the G2G flows are  
234 for the correct river rather than a nearby river with a similar catchment area, in some  
235 cases the derived catchment area draining to the 1km x 1km cell will be different to  
236 the “observed” NRFA catchment area. This problem can particularly affect smaller  
237 catchments, for which discretisation to a 1km x 1km grid can lead to proportionally  
238 larger errors, although flow data provided here are in any case limited to catchments  
239 with drainage areas of at least 50km<sup>2</sup>. The catchment area grids (Table 5) can be  
240 used to check the drainage area of gauged catchments, and could also be used to  
241 identify the most appropriate 1km x 1km cell corresponding to any particular  
242 ungauged catchment of interest.

243 Users should be aware that the effect of water bodies such as lakes and reservoirs is  
244 not accounted for within the model; any impact of lake storage and regulation on  
245 downstream river flows has been neglected, and lake grid-cells are treated as  
246 though they were rivers. The data files thus include ‘river flows’ and ‘soil moisture’ for  
247 1km cells located within lakes. Additional files identify majority lake cells in GB and  
248 NI, so that these can be excluded from analyses if desired (see Table 5).

249 The historical portion of the climate projection-based river flow and soil moisture  
250 datasets can be compared to the observation-based datasets, or to observed data.  
251 However, comparisons in either case should only be made statistically (over multi-  
252 decadal periods), not directly (time point by time point), as there will be no  
253 equivalence between observed weather features and those in the RCM PPE at the  
254 same date. An example of such a comparison is presented in Supp. Fig. 4 of Kay et  
255 al. 2021a, where mean monthly flows, flood frequency curves and low flow frequency  
256 curves are compared for the baseline SIMRCM ensemble and SIMOBS, for 8  
257 catchments in NI. Comparison of climate projection-based simulations to the  
258 observation-based simulation will indicate how both natural variability and  
259 (remaining) biases in the climate projection data affect the hydrological model  
260 simulations for the baseline period, while comparison to observed data themselves  
261 will be additionally affected by the accuracy of the G2G model.

262 The climate projection-based datasets for baseline (historical) and future periods can  
263 be compared statistically, to investigate the potential future impacts of climate  
264 change on river flows (e.g. Kay 2021, Lane & Kay 2021, Kay et al.2021a) and soil  
265 moisture (e.g. Kay et al. 2022a). Analyses should use the full ensemble; each  
266 member should be considered as a different but plausible realisation. Comparison  
267 between periods should use the same ensemble member for each period, not a  
268 baseline from one member and a future from another member.

269 The observation-based datasets for GB can be considered updates to MaRIUS-  
270 G2G-MORECS-monthly flow and soil moisture data (Bell et al. 2018a). The main

271 differences are the shorter simulation period here (Dec 1980–Nov 2011 vs 1960–  
272 2015 or 1891–2015), the inclusion here of the optional snow module, some changes  
273 to the land-sea mask, some changes related to infilling of missing soil type data, and  
274 minor changes to the discretised river flow network to improve the G2G model  
275 catchment areas (thus the additional spatial datasets provided here may differ in  
276 places to those provided with the MaRIUS dataset).


277 The climate projection-based datasets for GB are analogous to the MaRIUS-G2G-  
278 WAH2-monthly flow and soil moisture data (Bell et al. 2018b), which were driven by  
279 weather@home climate model data (Guillod et al. 2017). The main differences, as  
280 well as the factors listed above for the observation-based datasets, are the climate  
281 model version, the smaller ensemble size here (12 members vs 100 members), and  
282 the provision here of transient rather than time-slice data (Dec 1980–Nov 2080 vs  
283 1900–2006, 2020–2049 and 2070–2099). Note that, as far as the authors are aware,  
284 there has been no comparison between the weather@home and UKCP18 RCM  
285 climate datasets.

## 286 **3 Results**

### 287 **3.1 Monthly mean river flows**

288 Maps of example monthly mean river flows across GB and NI from SIMOBS and two  
289 SIMRCM ensemble members (Figure 3) illustrate the accumulation of water as it  
290 flows downstream, with typically higher flows for downstream locations with larger  
291 catchment areas. The example maps also show the generally lower flows in summer  
292 (July) compared to winter (January). Note that, although GB and NI are mapped  
293 together, the data for GB and NI are provided separately.






294

295 **Figure 3 Maps of monthly mean river flows ( $\text{m}^3\text{s}^{-1}$ ) for January and July 1982, from**  
 296 **SIMOBS (left) and two SIMRCM ensemble members (01 – centre, and 15 – right). Also**  
 297 **shown are Lough Neagh and Lough Erne in NI (bright blue shading).**

298

299 Time-series plots of UK-mean annual mean river flows from SIMOBS and the  
 300 SIMRCM ensemble show good correspondence (Figure 4). There is a relatively  
 301 small but highly statistically significant decrease in the SIMRCM ensemble mean  
 302 flow over Dec 1980–Nov 2080 ( $-0.695 \text{ m}^3\text{s}^{-1} / 100 \text{ years}$ ) (Figure 4). Six of the 12  
 303 individual ensemble members show decreases significant at the 10% level, while  
 304 four show non-significant decreases and two show non-significant increases. Plots of  
 305 the monthly climatology of UK-mean river flows for the first and last 30 years (Dec  
 306 1980–Nov 2010 and Dec 2050–Nov 2080) show a clear reduction in flows during  
 307 summer and early autumn, but a possible increase in winter (Figure 4).

308



309

310 **Figure 4 Time-series of UK-mean annual mean river flows (top), and the baseline (Dec**  
 311 **1980–Nov 2010) and future (Dec 2050–Nov 2080) monthly climatology of UK-mean**  
 312 **river flows (bottom), for SIMOBS and the SIMRCM ensemble. The shading in the**  
 313 **bottom plot shows the SIMRCM ensemble range for each period.**


314

315 Kay (2021) used the GB SIMRCM monthly mean river flow data to investigate  
 316 potential future changes in seasonal mean flows, for two future time-slices (2020–  
 317 2050 and 2050–2080). This suggested large decreases in summer mean flows  
 318 everywhere, but possible increases in winter mean flows, especially in the north and  
 319 west. A similar analysis using the NI SIMRCM monthly mean river flow data (Kay et  
 320 al. 2021a) suggested decreases in spring–autumn mean flows, especially in  
 321 summer, but possible increases in winter mean flows.

### 322 3.2 Extreme river flows

323 Maps of example GB and NI AMAX of daily mean river flows and AMIN of 7-day  
 324 mean river flows from SIMOBS and two SIMRCM ensemble members (Figure 5)  
 325 show less spatial variation than those of monthly mean river flows (when plotted on  
 326 the same scale). Note that, although GB and NI are mapped together, the data for  
 327 GB and NI are provided separately.

328



329

330

331

332

333

334

**Figure 5 Maps of AMAX of daily mean river flows for Oct 1982–Sep 1983 ( $\text{m}^3\text{s}^{-1}$ ; top) and AMIN of 7-day mean river flows for Dec 1982–Nov 1983 ( $\text{m}^3\text{s}^{-1}$ ; bottom), from SIMOBS (left) and two SIMRCM ensemble members (01 – centre, and 15 – right). Also shown are Lough Neagh and Lough Erne in NI (bright blue shading).**

335

336

337

338

339

340

341


342

343

344

Time-series plots of UK-mean AMAX and AMIN river flows from SIMOBS and the SIMRCM ensemble show good correspondence (Figure 6). The SIMRCM ensemble mean AMAX flows show a statistically significant increase over Oct 1981–Sep 2080 ( $8.51 \text{ m}^3\text{s}^{-1} / 100 \text{ years}$ ) (Figure 6). Nine of the 12 individual ensemble members show increases in AMAX flows significant at the 10% level, while one shows non-significant increases and two show non-significant decreases. The SIMRCM ensemble mean AMIN flows show a highly statistically significant decrease over Dec 1980–Nov 2080 ( $-0.670 \text{ m}^3\text{s}^{-1} / 100 \text{ years}$ ) (Figure 6), and all 12 individual ensemble members show decreases significant at the 10% level.





345  
 346 **Figure 6 Time-series of UK-mean AMAX of daily mean river flows (top) and AMIN of 7-**  
 347 **day mean river flows (bottom), for SIMOBS and the SIMRCM ensemble.**

348


349 Lane & Kay (2021) used the GB SIMRCM AMAX and AMIN river flow data to  
 350 investigate potential future changes in high/low flows by 2050–2080. All ensemble  
 351 members showed large reductions in 10-year return period low flows. The direction  
 352 of change for 10-year return period high flows was more uncertain, but increases of  
 353 over 9% were possible in most areas. Simultaneous worsening of both high and low  
 354 flow extremes was projected in the west. A similar analysis using the NI SIMRCM  
 355 AMAX and AMIN flow data (Kay et al. 2021a) suggested large reductions in 10-year  
 356 return period low flows everywhere, and large increases in 10-year return period high  
 357 flows for some locations and ensemble members. Analyses of the GB and NI dates  
 358 of occurrence of SIMRCM AMAX and AMIN showed few significant changes in  
 359 timing (Lane & Kay 2021, Kay et al. 2021a).

360

### 361 **3.3 Soil moisture**

362 Maps of example GB and NI monthly mean soil moisture content from SIMOBS and  
 363 two SIMRCM ensemble members (Figure 7) show the spatial variation, which is  
 364 generally related to the variation in soil types. The example maps also show the  
 365 generally drier soils in summer (July) compared to winter (January), and show  
 366 differences between the two selected SIMRCM ensemble members. Note that,

367 although GB and NI are mapped together, the data for GB and NI are provided  
368 separately.  
369




370  
371 **Figure 7 Maps of monthly mean soil moisture content (m water / m soil) for January**  
372 **and July 1982, from SIMOBS (left) and two SIMRCM ensemble members (01 – centre,**  
373 **and 15 – right).**

374

375 Time-series plots of UK-mean annual mean soil moisture content from SIMOBS and  
376 the SIMRCM ensemble show good correspondence (Figure 8). The SIMRCM  
377 ensemble mean soil moisture content shows a highly statistically significant  
378 decrease over Dec 1980–Nov 2080 (-0.035 / 100 years) (Figure 8), and all 12  
379 individual ensemble members show decreases significant at the 10% level. Plots of  
380 the monthly climatology of UK-mean soil moisture content for the first and last 30  
381 years (Dec 1980–Nov 2010 and Dec 2050–Nov 2080) show a clear reduction in  
382 summer and autumn (Figure 8).

383



384  
 385 **Figure 8** Time-series of UK-mean annual mean soil moisture content (m water / m soil;  
 386 top), and the baseline (Dec 1980–Nov 2010) and future (Dec 2050–Nov 2080) monthly  
 387 climatology of UK-mean soil moisture content (m water / m soil; bottom), for SIMOBS  
 388 and the SIMRCM ensemble. The shading in the bottom plot shows the SIMRCM  
 389 ensemble range for each period.

390

391 Kay et al. (2022a) used the GB and NI SIMRCM monthly mean soil moisture data to  
 392 investigate potential future changes in occurrence of indicative soil moisture  
 393 extremes and changes in typical wetting and drying dates of soils by 2050–2080  
 394 across the UK. This also suggested large increases in the spatial occurrence of low  
 395 soil moisture levels, and later soil wetting dates. Changes to soil drying dates were  
 396 less apparent.

## 397 4 Discussion

398 Ensemble data from the historical period of the climate projection-driven datasets  
 399 show good correspondence with the observation-driven datasets, for both river flows  
 400 (Figure 4 and Figure 6) and soil moisture (Figure 8). More detailed performance  
 401 analyses are provided elsewhere (Kay 2021, Lane & Kay 2021, Kay et al. 2021a,  
 402 Kay et al. 2022a).

403 The climate projection-driven river flow and soil moisture datasets suggest potential  
 404 future decreases in summer flows, annual minimum 7-day flows, and  
 405 summer/autumn soil moisture, along with possible future increases in winter flows  
 406 and annual maximum flows. More detailed analyses, presented elsewhere, illustrate



407 the variation in these general trends, both spatially and by ensemble member (Kay  
408 2021, Lane & Kay 2021, Kay et al. 2021a, Kay et al. 2022a). These changes are  
409 consistent with the climate projections, which give wetter winters and drier and hotter  
410 summers (Murphy et al. Fig. 5.2), and increased summer PE (Robinson et al. 2022).

411 A study of trends in historical gauged flows from the UK Benchmark Network  
412 (Harrigan et al. 2017) shows a tendency for an increase in winter mean flows and  
413 high flow indices over the past 50 years, although with significant natural decadal  
414 variability (clear so-called flood-rich and flood-poor periods). The datasets here  
415 suggest that this overall trend could continue into the future, and they could  
416 potentially be used to further investigate natural variability. The analysis of Harrigan  
417 et al. (2017) shows less consistent changes in summer mean flows or low flow  
418 indices, with catchments in the south/east often showing decreases, but catchments  
419 in the north/west typically showing increases. The datasets here suggest that more  
420 consistent decreases could be seen everywhere in future.

421 However, the fact that the UKCP18 Regional climate projections applied here only  
422 use one GCM/RCM needs to be borne in mind. Other climate models tend to give  
423 smaller decreases (or increases) in summer precipitation than the UKCP18 Regional  
424 projections (Murphy et al. 2018 Fig. 5.2), so are likely to give lower reductions in  
425 summer flows and soil moisture. Similarly, other climate models give a wider range  
426 of changes in winter precipitation than the UKCP18 Regional projections (Murphy et  
427 al. 2018 Fig. 5.2), so could give larger or smaller increases in winter flows. The  
428 UKCP Local projections, produced by nesting a ~2.2km convection-permitting model  
429 in each RCM PPE member (Kendon et al. 2021), also give some differences in  
430 climatic changes compared to the RCM (Kendon et al. 2021), and consequently  
431 some differences in hydrological impacts (Kay 2022). In addition, the use of a high  
432 emissions scenario (RCP8.5) for the UKCP18 Regional projections is likely to lead to  
433 more extreme changes than would occur for lower emissions (e.g. Arnell et al. 2014),  
434 although RCP8.5 should not be considered implausible (Schwalm et al. 2020).

435 Further sources of uncertainty in the datasets include the observation-based PE and  
436 the calculation of RCM PE. MORECS 40km monthly PE data are used for the  
437 observation-based hydrological model runs; lower spatial and temporal resolution  
438 than the other driving data (Section 2.2). A dataset of 1km daily PE has since been  
439 derived (Brown et al. 2022) using HadUK-Grid data (Met Office 2019), although  
440 some the variables required had to be interpolated from monthly to daily. The RCM  
441 PE used here includes the effect of stomatal closure under higher CO<sub>2</sub>  
442 concentrations but does not include a potential leaf area increase due to carbon  
443 fertilisation (Rudd and Kay 2016; Robinson et al.2022).

444 Potential future changes in land cover are also excluded, as are any artificial  
445 influences on river flows. Also, only one hydrological model has been applied; a  
446 catchment-based dataset of simulated river flows from the 'Enhanced future Flows  
447 and Groundwater' (eFlaG) project (Hannaford et al. 2022b) uses similar driving data  
448 from the UKCP18 Regional projections for three hydrological models (including  
449 G2G), so could be used to look at potential uncertainty from hydrological model  
450 structure (Hannaford et al. 2022a). Note that eFlaG used HadUK-Grid rainfall, both  
451 for observation-based runs and for bias correction of UKCP18 RCM data, whereas  
452 CEH-GEAR rainfall were used here since data are included for the parts of the  
453 Republic of Ireland draining into Northern Ireland, enabling gridded flow simulation  
454 across Northern Ireland. The ability to provide full and coherent coverage, of gauged

455 and ungauged locations is a particular strength of national-scale grid-based models  
456 like G2G, in contrast to catchment-based models.

## 457 **5 Conclusions**

458 The datasets presented here provide consistent spatial simulations of river flows and  
459 soil moisture for the whole of the UK, driven by both observed data and by an  
460 ensemble of regional climate model data from the latest UK climate projections,  
461 UKCP18. These enable direct studies of historical and potential future river flows and  
462 soil moisture, but they can also be used to provide inputs for further studies, for  
463 example to simulate water quality (e.g. Hutchins et al. 2016), crop yields (e.g. Cai et  
464 al. 2009), irrigated agriculture economic risk (Salmoral et al. 2019), or ecological  
465 impacts (e.g. Bussi et al. 2016).

466 An online (anonymous) stakeholder survey was carried out for UK-SCAPE WP2.2 in  
467 late 2021 (Kay et al. 2022e). This asked a set of questions divided into three broad  
468 classes; 'job role and level of experience', 'data of interest', and 'data format/access  
469 preferences'. The responses on 'data of interest' showed that there is a lot of interest  
470 in water quantity, including both river flows and soil moisture, and a lot of interest in  
471 potential future changes in river flows, although slightly less so for changes in soil  
472 moisture. Furthermore, the responses on 'data format/access preferences' showed  
473 that the greatest 1<sup>st</sup> preference was for grids covering sub-regions or the whole  
474 country, although perhaps unsurprising this varied by job role (Academic,  
475 Government/Regulator, Practitioner/ Consultant), which likely influences how data  
476 are used. A large proportion of respondents were also happy downloading the full  
477 dataset as NetCDF files from the EIDC. The datasets described here thus provide for  
478 a significant stakeholder demand, although there is always more that could be done.  
479 Further developments could include a web-tool allowing interactive data exploration  
480 and plotting.

## 481 **Acknowledgements**

482 This work was supported by the Natural Environment Research Council award  
483 number NE/R016429/1 as part of the UK-SCAPE programme delivering National  
484 Capability. Thanks to Emma Robinson (UKCEH) for work on the estimation of PE  
485 from climate model data.

## 486 **Data Availability**

487 The six datasets described in this manuscript are available from the Environmental  
488 Information Data Centre (EIDC); see Table 1.

## 489 **References**

490 Arnell NW, Charlton MB, Lowe JA (2014). The effect of climate policy on the impacts  
491 of climate change on river flows in the UK. *J Hydrol*, 510, 424-435.

492 Bell VA, Kay AL, Davies HN, Jones RG (2016). An assessment of the possible  
493 impacts of climate change on snow and peak river flows across Britain. *Clim  
494 Change*, 136(3), 539–553.

495 Bell VA, Kay AL et al. (2007). Development of a high resolution grid-based river flow  
496 model for use with regional climate model output. *Hydrol Earth Syst Sci*, 11, 532–  
497 549.

498 Bell VA, Kay AL et al. (2009). Use of soil data in a grid-based hydrological model to  
499 estimate spatial variation in changing flood risk across the UK. *J Hydrol*, 377, 335–  
500 350.

501 Bell VA, Rudd AC, Kay AL, Davies HN (2018a). Grid-to-Grid model estimates of  
502 monthly mean flow and soil moisture for Great Britain (1960 to 2015): observed  
503 driving data [MaRIUS-G2G-MORECS-monthly]. NERC Environmental Information  
504 Data Centre. doi:10.5285/e911196a-b371-47b1-968c-661eb600d83b.

505 Bell VA, Rudd AC, Kay AL, Davies HN (2018b). Grid-to-Grid model estimates of  
506 monthly mean flow and soil moisture for Great Britain: weather@home2 (climate  
507 model) driving data [MaRIUS-G2G-WAH2-monthly]. NERC Environmental  
508 Information Data Centre. doi:10.5285/3b90962e-6fc8-4251-853e-b9683e37f790.

509 Brown, M.J., Robinson, E.L., Kay, A.L., Chapman, R., Bell, V.A., Blyth, E.M. (2022).  
510 Potential evapotranspiration derived from HadUK-Grid 1km gridded climate  
511 observations 1969-2021 (Hydro-PE HadUK-Grid). NERC EDS Environmental  
512 Information Data Centre. doi:10.5285/9275ab7e-6e93-42bc-8e72-59c98d409deb.

513 Bussi G, Whitehead PG, Bowes MJ et al. (2016). Impacts of climate change, land-  
514 use change and phosphorus reduction on phytoplankton in the River Thames (UK),  
515 *Sci Total Environ*, 572, 1507-1519.

516 Cai X, Wang D, Laurent R (2009). Impact of Climate Change on Crop Yield: A Case  
517 Study of Rainfed Corn in Central Illinois. *J Appl Meteorol Climatol*, 4(9), 1868–1881.

518 Formetta, G., Prodocimi, I., Stewart, E., Bell, V. (2018). Estimating the index flood  
519 with continuous hydrological models: an application in Great Britain. *Hydrol Res*, 49,  
520 123–133.

521 Guillod BP, Jones RG, Dadson SJ et al. (2018). A large set of potential past, present  
522 and future hydro-meteorological time series for the UK. *Hydrol. Earth Syst. Sci.*,  
523 22(1), 611–634.

524 Guillod BP, Jones RG, Kay AL et al. (2017). Managing the Risks, Impacts and  
525 Uncertainties of drought and water Scarcity (MaRIUS) project: Large set of potential  
526 past and future climate time series for the UK from the weather@home2 model.  
527 Centre for Environmental Data Analysis.  
528 doi:10.5285/0cea8d7aca57427fae92241348ae9b03.

529 Hannaford J, Mackay JD, Ascot M et al. (2022a). eFLaG: enhanced future FLoWS  
530 and Groundwater. A national dataset of hydrological projections based on UKCP18.  
531 *Earth Syst Sci Data Discuss*, doi:10.5194/essd-2022-40.

532 Hannaford J, Mackay J, Ascot M, et al. (2022b). Hydrological projections for the UK,  
533 based on UK Climate Projections 2018 (UKCP18) data, from the Enhanced Future  
534 Flows and Groundwater (eFLaG) project. NERC EDS Environmental Information  
535 Data Centre. doi:10.5285/1bb90673-ad37-4679-90b9-0126109639a9.

536 Harrigan S, Hannaford J et al. (2018). Designation and trend analysis of the updated  
537 UK Benchmark Network of river flow stations: the UKBN2 dataset. *Hydrology*  
538 *Research*, 49, 552–567.



539 Hough M, Jones RJA (1997). The United Kingdom Meteorological Office rainfall and  
540 evaporation calculation system: MORECS version 2.0– an overview. *Hydro Earth*  
541 *Syst Sci*, 1(2), 227–239.

542 Hutchins MG, Williams RJ, Prudhomme C et al. (2016). Projections of future  
543 deterioration in UK river quality are hampered by climatic uncertainty under extreme  
544 conditions, *Hydrological Sciences Journal*, 61:16, 2818-2833.

545 Kay AL (2021). Simulation of river flow in Britain under climate change: baseline  
546 performance and future seasonal changes. *Hydro Process*. 35(4), e14137, doi:  
547 10.1002/hyp.14137.

548 Kay, A.L. (2022). Differences in hydrological impacts using regional climate model  
549 and nested convection-permitting model data. *Climatic Change*, 173(1-2), 11,  
550 doi:10.1007/s10584-022-03405-z.

551 Kay AL, Crooks SM (2014). An investigation of the effect of transient climate change  
552 on snowmelt, flood frequency and timing in northern Britain. *International Journal of*  
553 *Climatology*, 34(12), 3368–3381.

554 Kay AL, Davies HN, Lane RA, Rudd AC, Bell VA (2021a). Grid-based simulation of  
555 river flows in Northern Ireland: model performance and future flow changes. *Journal*  
556 *of Hydrology: Regional Studies*, 38, 100967.

557 Kay AL, Griffin A, Rudd AC, Chapman RM, Bell VA, Arnell NW (2021b). Climate  
558 change effects on indicators of high and low river flow across Great Britain.  
559 *Advances in Water Resources*, 151, 103909.

560 Kay AL, Lane RA, Bell VA (2022a). Grid-based simulation of soil moisture in the UK:  
561 future changes in extremes and wetting and drying dates. *Environmental Research*  
562 *Letters*, 17(7), 074029.

563 Kay AL, Rudd AC, Davies HN, Lane RA, Bell VA (2021c). Grid-to-Grid model  
564 estimates of river flow for Great Britain driven by observed data (1980 to 2011).  
565 NERC Environmental Information Data Centre. doi:10.5285/2f835517-253e-4697-  
566 b774-ab6ff2c0d3da.

567 Kay AL, Rudd AC, Davies HN, Lane RA, Bell VA (2021d). Grid-to-Grid model  
568 estimates of river flow for Northern Ireland driven by observed data (1980 to 2011).  
569 NERC Environmental Information Data Centre. doi:10.5285/f5fc1041-e284-4763-  
570 b8b7-8643c319b2d0.

571 Kay AL, Rudd AC, Davies HN, Lane RA, Bell VA (2021e). Grid-to-Grid model  
572 estimates of soil moisture for Great Britain and Northern Ireland driven by observed  
573 data (1980 to 2011). NERC Environmental Information Data Centre.  
574 doi:10.5285/c9a85f7c-45e2-4201-af82-4c833b3f2c5f.

575 Kay AL, Rudd AC, Davies HN, Lane RA, Bell VA (2022b). Grid-to-Grid model  
576 estimates of river flow for Great Britain driven by UK Climate Projections 2018  
577 (UKCP18) Regional (12km) data (1980 to 2080) v2. NERC Environmental  
578 Information Data Centre. doi:10.5285/18be3704-0a6d-4917-aa2e-bf38927321c5.

579 Kay AL, Rudd AC, Davies HN, Lane RA, Bell VA (2022c). Grid-to-Grid model  
580 estimates of river flow for Northern Ireland driven by UK Climate Projections 2018  
581 (UKCP18) Regional (12km) data (1980 to 2080) v2. NERC Environmental  
582 Information Data Centre. doi:10.5285/76057b0a-b18f-496f-891c-d5b22bd0b291.

583 Kay AL, Rudd AC, Davies HN, Lane RA, Bell VA (2022d). Grid-to-Grid model  
584 estimates of soil moisture for Great Britain and Northern Ireland driven by UK  
585 Climate Projections 2018 (UKCP18) Regional (12km) data (1980 to 2080). NERC  
586 Environmental Information Data Centre. doi:10.5285/f7142ced-f6ff-486b-af33-  
587 44fb8f763cde.

588 Kay AL, Spencer M, Bell VA (2022e). UK-SCAPE WP2.2: Water Futures.  
589 Stakeholder questionnaire results. Wallingford, UK Centre for Ecology & Hydrology,  
590 14pp. nora.nerc.ac.uk/id/eprint/531705/

591 Kendon E, Short C, Pope J, Chan S, Wilkinson J, Tucker S, Bett P, Harris G (2021).  
592 Update to UKCP Local (2.2km) projections. Met Office Hadley Centre, Exeter, UK.

593 Lane RA, Kay AL (2021). Climate change impact on the magnitude and timing of  
594 hydrological extremes across Great Britain. *Frontiers in Water*, 3:684982,  
595 doi:10.3389/frwa.2021.684982.

596 Lowe JA, Bernie D, Bett P et al. (2018). UKCP18 Science Overview report. Exeter,  
597 UK: Met Office Hadley Centre.

598 Met Office Hadley Centre (2018). UKCP18 Regional Projections on a 12km grid over  
599 the UK for 1980-2080. CEDA, September 2019.  
600 catalogue.ceda.ac.uk/uuid/589211abeb844070a95d061c8cc7f604.

601 Met Office, Hollis D, McCarthy M et al. (2019). HadUK-Grid Gridded Climate  
602 Observations on a 1km grid over the UK, v1.0.0.0 (1862-2017). Centre for  
603 Environmental Data Analysis, 14 November 2019.  
604 doi:10.5285/2a62652a4fe6412693123dd6328f6dc8.

605 Morris DG, Flavin RW (1990). A digital terrain model for hydrology. In: Proceedings  
606 of the 4th International Symposium on Spatial Data Handling, Zurich, Switzerland,  
607 23–27 July 1990, 250–262.

608 Murphy, J.M., Harris, G.R., Sexton, D.M.H. et al. (2018). UKCP18 Land Projections:  
609 Science Report. Met Office Hadley Centre, Exeter, UK.

610 Rameshwaran P, Bell VA, Brown MJ, Davies HN, Kay AL, Rudd AC, Sefton C  
611 (2022). Use of abstraction and discharge data to improve the performance of a  
612 national-scale hydrological model. *Water Resources Research*, 5(1),  
613 e2021WR029787.

614 Riahi K, Krey V et al. (2011). RCP-8.5: exploring the consequence of high emission  
615 trajectories. *Clim Change*, 109, 33–57.

616 Robinson EL, Kay AL, Brown M, Chapman R, Bell V, Blyth EM (2021). Potential  
617 evapotranspiration derived from the UK Climate Projections 2018 Regional Climate  
618 Model ensemble 1980-2080 (Hydro-PE UKCP18 RCM). NERC Environmental  
619 Information Data Centre. doi:10.5285/eb5d9dc4-13bb-44c7-9bf8-c5980fcf52a4.

620 Robinson EL, Kay AL, Brown M, Chapman R, Bell V, Blyth E (2022). Hydro-PE:  
621 gridded datasets of historical and future Penman-Monteith potential evaporation for  
622 the United Kingdom. *Earth Syst. Sci. Data Discuss.*, doi:10.5194/essd-2022-288.

623 Rowland CS, Morton RD, Carrasco L et al. (2017) Land Cover Map 2015 (25m  
624 raster, N. Ireland). NERC EIDC, doi:10.5285/47f053a0-e34f-4534-a843-  
625 76f0a0998a2f.

- 626 Rudd AC, Bell VA, Kay AL (2017). National-scale analysis of simulated hydrological  
627 droughts (1891-2015). *J Hydrol*, 550, 368-385.
- 628 Rudd AC, Kay AL (2016). Use of very high resolution climate model data for  
629 hydrological modelling: estimation of potential evaporation. *Hydrology Research*,  
630 47(3), 660–670, doi:10.2166/nh.2015.028.
- 631 Salmoral, G., Rey, D. Rudd, A., de Margon, P., Holman I. (2019). A Probabilistic Risk  
632 Assessment of the National Economic Impacts of Regulatory Drought Management  
633 on Irrigated Agriculture. *Earth's Future*, 7(2), 178-196.
- 634 Schwalm, C.R., Glendon, S., Duffy, P.B. (2020). RCP8.5 tracks cumulative CO2  
635 emissions. *Proc Nat Acad. Sci*, 117, 19656-19657.
- 636 Tanguy M, Dixon H et al. (2016). Gridded estimates of daily and monthly areal  
637 rainfall for the United Kingdom (1890-2015) [CEH-GEAR]. NERC EIDC.  
638 doi:10.5285/33604ea0-c238-4488-813d-0ad9ab7c51ca.
- 639