

1 **The HTAP\_v3 emission mosaic: merging regional and global monthly emissions (2000-**  
2 **2018) to support air quality modelling and policies**

3  
4 Monica Crippa<sup>1</sup>, Diego Guizzardi<sup>1</sup>, Tim Butler<sup>2</sup>, Terry Keating<sup>3</sup>, Rosa Wu<sup>16</sup>, Jacek Kaminski<sup>4</sup>,  
5 Jeroen Kuenen<sup>5</sup>, Junichi Kurokawa<sup>6</sup>, Satoru Chatani<sup>15</sup>, Tazuko Morikawa<sup>7</sup>, George Pouliot<sup>17</sup>,  
6 Jacinthe Racine<sup>8</sup>, Michael D. Moran<sup>16</sup>, Zbigniew Klimont<sup>9</sup>, Patrick M. Manseau<sup>8</sup>, Rabab  
7 Mashayekhi<sup>8</sup>, Barron H. Henderson<sup>17</sup>, Steven J. Smith<sup>10</sup>, Harrison Suchyta<sup>10</sup>, Marilena  
8 Muntean<sup>1</sup>, Efsio Solazzo<sup>11</sup>, Manjola Banja<sup>1</sup>, Edwin Schaaf<sup>1</sup>, Federico Pagani<sup>12</sup>, Jung-Hun  
9 Woo<sup>13,14</sup>, Jinseok Kim<sup>14</sup>, Fabio Monforti-Ferrario<sup>1</sup>, Enrico Pisoni<sup>1</sup>, Junhua Zhang<sup>16</sup>, David  
10 Niemi<sup>8</sup>, Mourad Sassi<sup>8</sup>, Tabish Ansari<sup>2</sup>, Kristen Foley<sup>17</sup>

11  
12 <sup>1</sup>European Commission, Joint Research Centre (JRC), Ispra, Italy

13 <sup>2</sup>Institute for Advanced Sustainability Studies (IASS), Potsdam, 14467, Germany

14 <sup>3</sup>U.S. Environmental Protection Agency, Washington DC 20460, USA

15 <sup>4</sup>Institute of Environmental Protection, National Research Institute, Poland

16 <sup>5</sup>Department of Climate, Air and Sustainability, TNO, Utrecht, The Netherlands

17 <sup>6</sup>Asia Center for Air Pollution Research (ACAP), 1182 Sowa, Nishi-ku, Niigata-shi 950-2144,  
18 Japan

19 <sup>7</sup>Japan Automobile Research Institute (JARI)

20 <sup>8</sup>Air Quality Policy-Issue Response Section, Canadian Centre for Meteorological and  
21 Environmental Prediction, Environment and Climate Change Canada (ECCC), Dorval,  
22 Quebec, Canada

23 <sup>9</sup>International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria

24 <sup>10</sup>Joint Global Change Research Institute, Pacific Northwest National Lab, College Park, MD,  
25 USA

26 <sup>11</sup>UniSystems Company, Milan (Italy)

27 <sup>12</sup>GFT Italia S.r.l., VIA SILE, 20139 Milano, Italy

28 <sup>13</sup>Department of Civil and Environmental Engineering, Konkuk University, Seoul, South Korea

29 <sup>14</sup>Department of Technology Fusion Engineering, Konkuk University, Seoul, South Korea

30 <sup>15</sup>National Institute for Environmental Studies (NIES), Tsukuba, 305-8506, Japan

31 <sup>16</sup>Air Quality Research Division, Environment and Climate Change Canada, Toronto, Ontario,  
32 Canada

33 <sup>17</sup>U.S. Environmental Protection Agency, North Carolina, USA

34 Correspondence to: [Marilena.muntean@ec.europa.eu](mailto:Marilena.muntean@ec.europa.eu), [monica.crippa@ext.ec.europa.eu](mailto:monica.crippa@ext.ec.europa.eu)

35 **Abstract.** This study, performed under the umbrella of the Task Force on Hemispheric  
36 Transport of Air Pollution (TF-HTAP), responds to the need of the global and regional  
37 atmospheric modelling community of having a mosaic emission inventory of air pollutants that  
38 conforms to specific requirements: global coverage, long time series, spatially distributed

1 emissions with high time resolution, and a high sectoral resolution. The mosaic approach of  
2 integrating official regional emission inventories based on locally reported data, with a global  
3 inventory based on a globally consistent methodology, allows modellers to perform simulations  
4 of a high scientific quality while also ensuring that the results remain relevant to policymakers.

5 HTAP\_v3, an ad-hoc global mosaic of anthropogenic inventories, has been developed by  
6 integrating official inventories over specific areas (North America, Europe, Asia including  
7 Japan and Korea) with the independent Emissions Database for Global Atmospheric Research  
8 (EDGAR) inventory for the remaining world regions. The results are spatially and temporally  
9 distributed emissions of SO<sub>2</sub>, NO<sub>x</sub>, CO, NMVOC, NH<sub>3</sub>, PM<sub>10</sub>, PM<sub>2.5</sub>, Black Carbon (BC), and  
10 Organic Carbon (OC), with a spatial resolution of 0.1 x 0.1 degree and time intervals of months  
11 and years covering the period 2000-2018 (<https://doi.org/10.5281/zenodo.7516361>,  
12 [https://edgar.jrc.ec.europa.eu/dataset\\_htap\\_v3](https://edgar.jrc.ec.europa.eu/dataset_htap_v3)). The emissions are further disaggregated to 16  
13 anthropogenic emitting sectors. This paper describes the methodology applied to develop such  
14 an emission mosaic, reports on source allocation, differences among existing inventories, and  
15 best practices for the mosaic compilation. One of the key strengths of the HTAP\_v3 emission  
16 mosaic is its temporal coverage, enabling the analysis of emission trends over the past two  
17 decades. The development of a global emission mosaic over such long time series represents a  
18 unique product for global air quality modelling and for better-informed policy making,  
19 reflecting the community effort expended by the TF-HTAP to disentangle the complexity of  
20 transboundary transport of air pollution.

## 21 **1 Introduction**

22 Common international efforts have procured an agreement to reduce global air pollutant  
23 emissions. For this purpose, the United Nations Economic Commission for Europe (UNECE)  
24 Convention on Long Range Transboundary Air Pollution (CLRTAP) and the Task Force on  
25 Hemispheric Transport of Air Pollution (TF-HTAP) have been instrumental in developing the  
26 understanding of intercontinental transport of air pollution and thus contributing to the  
27 reduction of key pollutants in Europe and North America.

28 The success of CLRTAP is based on meeting strict reduction targets for pollutant releases.  
29 Therefore, evaluating the resulting implications of these reductions requires an ongoing  
30 improvement of global emission inventories in terms of emission updating and of  
31 methodological refinements. These aspects are instrumental to gain understanding of  
32 transboundary air pollution processes and drivers and to measure the effectiveness of emissions  
33 reduction and air quality mitigation policies. New guidance is available to achieve further  
34 emission reductions across all emitting sectors. For example, the 2019 establishment of the  
35 Task Force for International Cooperation on Air Pollution, which is intended to promote  
36 international collaboration for preventing and reducing air pollution and improving air quality  
37 globally (UNECE, 2021). As part of the ongoing effort by CLRTAP to reduce emissions and  
38 to set out more effective and accountable mitigation measures, the 2005 Gothenburg Protocol  
39 (UNECE, 2012) has been revised, including the review of the obligations in relation to  
40 emission reductions and mitigation measures (e.g., black carbon and ammonia) and the review  
41 of the progress towards achieving the environmental and health objectives of the Protocol.

42  
43 The Task Force on Hemispheric Transport of Air Pollution (TF-HTAP) of the Convention has  
44 a mandate to promote the scientific understanding of the intercontinental transport of air  
45 pollution to and from the UNECE area (<https://unece.org/geographical-scope>), to quantify its  
46 impacts on human health, vegetation and climate, and to identify emission mitigation options  
47 that will shape future global policies.

1 This paper describes and discusses a consistent global emission inventory of air pollutants  
2 emitted by anthropogenic activities. This important database has been developed to assess the  
3 contribution of anthropogenic air pollution emission sources within and outside the UNECE-  
4 area through atmospheric modelling. This inventory has been compiled based on officially  
5 reported emissions, and an independent global inventory where officially reported emissions  
6 are not used. This harmonised emissions “mosaic” dataset, hereafter referred to as the  
7 HTAP\_v3, contains annual and monthly:

- 8 - emission time series (from 2000 to 2018) of SO<sub>2</sub>, NO<sub>x</sub> (expressed as NO<sub>2</sub> mass unit),  
9 CO, NMVOC, NH<sub>3</sub>, PM<sub>10</sub>, PM<sub>2.5</sub>, BC, OC by emitting sector and country, and
- 10 - spatially distributed emissions on a global grid with spatial spacing of 0.1x0.1 degree.

11 The creation of a global emission mosaic requires the harmonisation of several data sources,  
12 detailed analysis of contributing sectors for the different input inventories, development of data  
13 quality control procedures, and a robust and consistent gap-filling methodology when lacking  
14 information. The development of HTAP\_v3 builds upon the previous experience of the  
15 HTAPv1 (Janssens-Maenhout et al., 2012) and HTAPv2.2 (Janssens-Maenhout et al., 2015)  
16 global inventories. HTAP\_v3, as requested by the TF-HTAP modelling community, provides  
17 a more refined sectoral disaggregation compared to the previous HTAP emission mosaics. It  
18 also includes tools ([https://edgar.jrc.ec.europa.eu/htap\\_tool/](https://edgar.jrc.ec.europa.eu/htap_tool/)) that allow the extraction of  
19 emission data over selected domains (detailed later in section 4).

20 This paper describes the development the HTAP\_v3 database as a global anthropogenic air  
21 pollutant emissions inventory mosaic for the period 2000-2018. The HTAP\_v3 mosaic has  
22 been composed by integrating official, spatially distributed emissions data from CAMS-REG-  
23 v5.1 (Kuenen et al., 2022), US EPA (U.S. Environmental Protection Agency, 2021b, a),  
24 Environment and Climate Change Canada (ECCC) (NPRI, 2017), REAS, CAPSS-KU, and  
25 JAPAN (<https://www.env.go.jp/air/osen/pm/info.html>) (Kurokawa and Ohara, 2020; Chatani  
26 et al., 2018; Chatani et al., 2020) inventories. As the information gathered from the official  
27 reporting covers only part of the globe, HTAP\_v3 has been completed using emissions from  
28 the Emissions Database for Global Atmospheric Research (EDGAR) version 6.1  
29 ([https://edgar.jrc.ec.europa.eu/dataset\\_ap61](https://edgar.jrc.ec.europa.eu/dataset_ap61)).

30 One of the key strengths of the HTAP\_v3 emission mosaic is the temporal coverage of the  
31 emissions, spanning the 2000-2018 period, enabling the analysis of emission trends over the  
32 past two decades. The development of a global emission mosaic over such long time series  
33 represents a unique product for air quality modelling and for better-informed policy making,  
34 reflecting the effort of the TF-HTAP community to improve understanding of the  
35 transboundary transport of air pollution. The year 2000 was chosen as the start year since it  
36 often represents the year from which complete datasets of annual air pollutant emissions can  
37 be generated. It also represents a turning point for several emerging economies (e.g., China)  
38 and the strengthening of mitigation measures in historically developed regions (e.g., EU, USA,  
39 etc.).

40 The two previous HTAP emission mosaics had limited temporal coverage. HTAPv1 covered  
41 the period 2000-20005 with annual resolution ([https://edgar.jrc.ec.europa.eu/dataset\\_htap\\_v1](https://edgar.jrc.ec.europa.eu/dataset_htap_v1),  
42 (Janssens-Maenhout et al., 2012)), while HTAPv2.2 covered two recent years (2008 and 2010),  
43 but with monthly resolution (Janssens-Maenhout et al., 2015)  
44 ([https://edgar.jrc.ec.europa.eu/dataset\\_htap\\_v2](https://edgar.jrc.ec.europa.eu/dataset_htap_v2)). However, the needs of the TF-HTAP  
45 modelling community are continuously evolving to both foster forward-looking air quality  
46 science and produce more fit-for-purpose analyses in support of efficient policy making.  
47 HTAP\_v3 therefore not only covers the time period of the previous HTAP phases, but also

1 extends it forward by almost a decade, to provide the most up-to-date picture of global air  
2 pollutant emission trends. Another distinguishing feature of the HTAPv3 mosaic is a  
3 considerably higher sectoral resolution than previous iterations of the HTAP mosaic  
4 inventories (section 2.2), enabling more policy-relevant use of the inventory.

5 The methodology and data sources for the HTAP\_v3 emission mosaic are described in section  
6 2. The long-time coverage of two decades, allows comprehensive trend analysis (see section  
7 3), the HTAP\_v3 data format and data-set access are presented in section 4 and conclusions  
8 are provided in section 5.

## 10 **2 HTAP\_v3 emission mosaic overview: data sources, coverage, and methodology**

### 11 **2.1 Data input**

12 The HTAP\_v3 mosaic is a database of monthly- and sector-specific global air pollutant  
13 emission gridmaps developed by integrating spatially explicit regional information from recent  
14 officially-reported national or regional emission inventories. Data from six main regional  
15 inventories were integrated into HTAP\_v3, which covered only North America, Europe, and a  
16 portion of Asia (including Japan, China, India, and South Korea) (Fig.1). The geographical  
17 domain covered by each of these inventories is depicted in Fig. 1, while further details on each  
18 contributing inventory are presented in section 2.3. The emissions for all other countries,  
19 international shipping and aviation (international and domestic) have been retrieved from the  
20 Emissions Database for Global Atmospheric Research (EDGARv6.1,  
21 [https://edgar.jrc.ec.europa.eu/dataset\\_ap61](https://edgar.jrc.ec.europa.eu/dataset_ap61)) as represented by the grey areas in Fig.1.  
22 Depending on the pollutant, more than half of global emissions are provided by region-specific  
23 inventories, while the remaining contribution is derived from the EDGAR global inventory as  
24 reported in the bar graph of Fig.1, where the share of each individual inventory to global  
25 emissions is represented. For all pollutants, the Asian domain is contributing most to global  
26 emissions, hence the importance of having accurate emission inventories for this region.

27 Recent literature studies (Puliafito et al., 2021; Huneus et al., 2020; Álamos et al., 2022; Keita  
28 et al., 2021; MEIC, 2022) document additional regional/local inventories which may contribute  
29 to future updates of HTAP\_v3, in particular extending the mosaic compilation to regions in the  
30 Southern Hemisphere. Considering relative hemispheric emission levels as well as the  
31 atmospheric dynamics happening in the Northern Hemisphere and regulating the  
32 transboundary transport of air pollution, the current HTAP\_v3 mosaic should still satisfy the  
33 needs of the atmospheric modelling community, although improvements using latest available  
34 inventories for Africa and South America may also be considered for future updates.

35 Table 1 provides an overview of all data providers, in terms of geographical and temporal  
36 coverage, data format, and sectoral and pollutant data availability. Table 2 defines the  
37 HTAP\_v3 sectors and corresponding IPCC codes. Table 3 further details the sector-pollutant  
38 data availability for each inventory and the gap-filling approach required for some sectors and  
39 pollutants.

### 40 **2.2 Pollutant, spatial, temporal and sectoral coverage**

41 The HTAP\_v3 emission mosaic helps to address the transboundary role of air pollutants by  
42 providing a key input for atmospheric modellers and supporting the evaluation of  
43 environmental impact analyses for poor air quality. For this reason, HTAP\_v3 provides global  
44 0.1 x 0.1 degree emission gridmaps for all air pollutants and specifically for acidifying and

1 eutrophying gases (such as SO<sub>2</sub>, NH<sub>3</sub>, NO<sub>x</sub>), ozone precursors (NMVOC, CO, NO<sub>x</sub>), and  
2 primary particulate matter (PM<sub>10</sub>, PM<sub>2.5</sub>, BC, OC).

3 Emissions from each officially-reported inventory were submitted to HTAP on 0.1 x 0.1 degree  
4 regional gridmaps. Spatial allocation was performed to these gridmaps for each sector by each  
5 inventory group using the best available set of subsector spatial surrogate fields used by each  
6 group (e.g., <https://www.emascenter.org/sa-tools>). EDGARv6.1 global gridmaps are also on a  
7 0.1 x 0.1 degree grid.

8 Compared to the two previous HTAP emission mosaics, HTAP\_v3 input emission gridmaps  
9 were provided with monthly time distributions to better reflect the regional seasonality of sector  
10 specific emissions (e.g., household, power generation, and agricultural activities). Information  
11 on emission peaks over certain months of the year is also a useful information for the  
12 development of territorial policies to mitigate localised emission sources in space and time  
13 (e.g., emissions from residential heating over winter months, agricultural residue burning, etc.).

14 The HTAP\_v3 mosaic provides emissions for gaseous and particulate matter air pollutants  
15 arising from all anthropogenic emitting sectors except for wildfires and savannah burning,  
16 which represent major sources of particulate matter and CO emissions. Wildfires and savannah  
17 burning are not included in the current mosaic since community efforts are ongoing to tackle  
18 these sources specifically. Modellers can find these additional sources on several publicly  
19 available global wildfire emission datasets compiled based on the best available scientific  
20 knowledge, such as the Global Fire Emission Database (GFED,  
21 <https://www.globalfiredata.org/>) or the Global Wildfire Information System (GWIS,  
22 <https://gwis.jrc.ec.europa.eu/>). When using satellite retrieved emissions from fires, they should  
23 be treated with caution to avoid double counting the emissions released by e.g. agricultural  
24 crop residue burning activities.

25 HTAP\_v3 provides emissions at higher sectoral disaggregation than previous HTAP  
26 experiments<sup>1</sup> to better understand drivers of emission trends and the effectiveness of sector-  
27 specific policy implementation. Emissions from 16 sectors are provided by the HTAP\_v3  
28 mosaic, namely: International Shipping; Domestic Shipping; Domestic Aviation; International  
29 Aviation; Energy; Industry; Fugitives; Solvent Use; Road Transport; Brake and Tyre Wear;  
30 Other Ground Transport; Residential; Waste; Agricultural Waste Burning; Livestock; and  
31 Agricultural Crops. Further details on the sector definitions as well as their correspondence  
32 with the IPCC codes (IPCC, 1996, 2006) are provided in Table 2. The selection of the number  
33 of sectors was constrained by the sectoral disaggregation of the input inventories (see Table  
34 S1). Table 3 provides the complete overview of the emission data provided by each inventory  
35 group indicating the pollutants covered for each sector and eventual gap-filling information  
36 included using the EDGARv6.1 data. Table 4 reports a summary of the main features of the 3  
37 HTAP emission mosaics, showing the advancements achieved with this work. The high sector  
38 disaggregation available within the HTAP\_v3 mosaic gives needed flexibility to modellers to  
39 include or exclude emission sub-sectors in their simulations, in particular when integrating the  
40 anthropogenic emissions provided by HTAP\_v3 with other components (e.g. natural  
41 emissions, forest fires, etc.). However, we recommend particular caution when using a natural

---

<sup>1</sup>HTAPv1 covered 10 broad emission sectors (Aircraft, Ships, Energy, Industry Processes, Ground Transport, Residential, Solvents, Agriculture, Agriculture Waste Burning, and Waste), while even broader sectoral emissions were provided in HTAPv2.2 (Air, Ships, Energy, Industry, Transport, Residential (including waste), and Agriculture (only for NH<sub>3</sub>)).

1 emissions model such as MEGAN (Model of Emissions of Gases and Aerosols from Nature,  
2 <https://www2.acom.ucar.edu/modeling/model-emissions-gases-and-aerosols-nature-megan>),  
3 which includes the estimation of NMVOC emissions from crops and soil NO<sub>x</sub> emissions  
4 (including agricultural soils) that are also provided by the HTAP\_v3 mosaic.

## 5 6 **2.3 Inventory overviews**

7 In the following sub-sections, details are provided on each officially-reported inventory used  
8 to construct the HTAP\_v3 emission mosaic.

### 9 **2.3.1 CAMS-REG-v5.1 inventory**

10 The CAMS-REG-v5.1 emission inventory was developed to support air pollutant and  
11 greenhouse gas modelling activities at the European scale. The inventory builds largely on the  
12 official reported data to the UN Framework Convention on Climate Change (UNFCCC) for  
13 greenhouse gases (for CO<sub>2</sub> and CH<sub>4</sub>), and the Convention on Long-Range Transboundary Air  
14 Pollution (CLRTAP) for air pollutants. For the latter, data are collected for NO<sub>x</sub>, SO<sub>2</sub>, CO,  
15 NMVOC, NH<sub>3</sub>, PM<sub>10</sub> and PM<sub>2.5</sub>, including all major air pollutants. For each of these pollutants,  
16 the emission data are collected at the sector level at which these are reported for the time series  
17 2000-2018 for each year and country. The CAMS-REG inventory covers UNECE-Europe,  
18 extending eastward until 60°E, therefore including the European part of Russia. For some non-  
19 EU countries, the reported data are found to be partially available or not available at all. In  
20 other cases, the quality of the reported data is found to be insufficient, i.e. with important data  
21 gaps or following different formats or methods. In this case, emission data from the IIASA  
22 GAINS model instead (IIASA, 2018) are used. This model is the main tool used to underpin  
23 pan-European and EU level air quality policies such as the UNECE Convention on Long Range  
24 Transboundary Air Pollution (UNECE, 2012) and the EU National Emission reduction  
25 Commitments Directive (European Commission, 2016).

26 After collecting all the emission data from reporting and GAINS, the source sectors are  
27 harmonised, distinguishing around 250 different subsectors. For each detailed sector, a  
28 speciation is applied to the PM<sub>2.5</sub> and PM<sub>10</sub> emissions, distinguishing elemental carbon  
29 (representing BC in the HTAP\_v3 inventory), organic carbon and other non-carbonaceous  
30 emissions for both the coarse (2.5-10 μm) and fine (<2.5 μm) mode.

31 A consistent spatial resolution is applied across the entire domain, where a specific proxy is  
32 selected for each subsector to spatially distribute emissions, including for instance the use of  
33 point source emissions, e.g., from the European Pollutant Release and Transfer Register (E-  
34 PRTR), complemented with additional data from the reporting of EU Large Combustion Plants  
35 (European Commission, 2001) and the Platts/WEPP commercial database for power plants  
36 (Platts, 2017). Road transport emissions are spatially disaggregated using information from  
37 OSM (Open Street Map, 2017), combined with information on traffic intensity in specific road  
38 segments from OTM (OpenTransportMap, 2017). Agricultural livestock emissions are  
39 spatially distributed using global gridded livestock numbers (FAO, 2010). Furthermore,  
40 CORINE land cover (Copernicus Land Monitoring Service, 2016) and population density are  
41 other key spatial distribution proxies.

42 After having spatially distributed the data, the ~ 250 different source categories are aggregated  
43 to fit with the HTAP\_v3 sector classification (Table S1). CAMS-REG-v5.1 is an update of an  
44 earlier version, CAMS-REG-v4.2 and based on the 2020 submissions to cover the years 2000-  
45 2018. A detailed description of the CAMS-REG-v4.2 inventory is provided in Kuenen et al.  
46 (2022).

1 The data are provided as gridded annual totals at a resolution of 0.05°x0.1° (lat-lon). Along  
2 with the grids, additional information is available including height profiles as well as temporal  
3 profiles to break down the annual emissions into hourly data (monthly profiles, day-of-the-  
4 week profiles and hourly profiles for each day). Furthermore, the CAMS-REG inventory  
5 provides dedicated speciation profiles for NMVOC per year, country and sector.

### 6 **2.3.2 US EPA inventory**

7 Emissions estimates for the United States were based primarily on estimates produced for the  
8 EPA's Air QUALity Time Series Project (EQUATES), which generated a consistent set of  
9 modelled emissions, meteorology, air quality, and pollutant deposition for the United States  
10 spanning the years 2002 through 2017 (<https://www.epa.gov/cmaq/equates>). For each sector, a  
11 consistent methodology was used to estimate emissions for each year in the 16-year period, in  
12 contrast to the evolving methodologies applied in the triennial U.S. National Emissions  
13 Inventories (NEIs) produced over that span. The HTAPv3 time series was extended back one  
14 year to 2001 and forward one year to 2018 using country, sector, and pollutant specific trends  
15 from EDGARv6.1.

16  
17 Emissions estimates were calculated for more than 8000 Source Classification Codes grouped  
18 into 101 sectors and then aggregated to the 16 HTAP\_v3 emission sectors. The 2017 NEI (U.S.  
19 Environmental Protection Agency, 2021b) served as the base year for the time series. For each  
20 sector, emissions estimates were generated for previous years using one of four methods: 1)  
21 applying new methods to create consistent emissions for all years, 2) scaling the 2017 NEI  
22 estimates using annual sector-specific activity data and technology information at the county  
23 level, 3) using annual emissions calculated consistently in previous NEIs and interpolating to  
24 fill missing years, and 4) assuming emissions were constant at 2017 levels. The assumption of  
25 constant emissions was applied to a very limited number of sources. Foley et al. (2023)  
26 provides a detailed explanation of the assumptions used for each sector.

27 Emissions from electric generating units were estimated for individual facilities, combining  
28 available hourly emissions data for units with continuous emissions monitors (CEMs) and  
29 applying regional fuel-specific profiles to units without CEMS. On-road transport and non-  
30 road mobile emissions were estimated using emission factors from the MOVES v3 model (U.S.  
31 Environmental Protection Agency, 2021a). A complete MOVES simulation was completed  
32 only for the NEI years with national adjustment factors applied for years plus or minus one  
33 from the NEI year. For California, emission factors for all on-road sources for all years were  
34 based on the California Air Resources Board Emission Factor Model (EMFAC)  
35 (<https://ww2.arb.ca.gov/our-work/programs/mobile-source-emissions-inventory/>). New non-  
36 road emissions estimates for Texas were provided by the Texas Commission on Environmental  
37 Quality. Emissions from oil and gas exploration and production were calculated using point  
38 source specific data and the EPA Oil and Gas Tool (U.S. Environmental Protection Agency,  
39 2021b), incorporating year-specific spatial, temporal, and speciation profiles. Residential wood  
40 combustion estimates were developed with an updated methodology incorporated into the 2017  
41 NEI and scaled backward to previous years using a national activity as a scaling factor. Solvent  
42 emissions were estimated using the Volatile Chemical Product (VCPy) framework of Seltzer  
43 et al. (2021). Emissions from livestock waste were calculated with revised annual animal  
44 counts to address missing data and methodological changes over the period. Emissions for

1 agricultural burning were developed using a new suite of activity data with the same  
2 methodology and input data sets from 2002 onwards. County-level estimates were only  
3 available for 2002 because activity data based on satellite information was not yet available.  
4 Emissions for forest wildfires, prescribed burns, grass and rangeland fires were also calculated  
5 in EQUATES but not included in the HTAP\_v3 data. For EQUATES, fugitive dust emissions  
6 (e.g., unpaved road dust, coal pile dust, dust from agricultural tilling) were reduced to account  
7 for precipitation and snow cover by grid cell. For use in HTAP\_v3, however, no  
8 meteorological adjustments (which decrease annual PM<sub>10</sub> emissions by about 75% on average)  
9 were applied to fugitive dust emissions. Wind-blown fugitive dust emissions are not included  
10 in the estimates for other regions in the HTAP\_v3 mosaic

11 Non-point source emissions were allocated spatially based on a suite of activity surrogates (e.g.  
12 population, total road miles, housing, etc.), many of which are sector specific. The spatial  
13 allocation factors were calculated for the 0.1 degree grid used by EDGARv6.1 with no  
14 intermediate re-gridding. The spatial allocation factors for all sectors were held constant for  
15 the entire time series except for oil and gas sectors which were year-specific. Depending on  
16 the sector, either 2017-based or 2014-based surrogates were developed for the same sectors as  
17 in the EQUATES.

18 Emissions from the US EPA inventory were provided from 2002-2017 (Table 1). Emissions  
19 for the year 2018 were estimated applying country, sector and pollutant specific trends from  
20 EDGAR, as well as for years 2000 and 2001 to complete the entire time series. Table S1  
21 provides an overview about the US EPA inventory sector mapping to the HTAP\_v3 sectors.

### 22 **2.3.3 Environment and Climate Change Canada (ECCC) inventory**

23 The Canadian emissions inventory data were obtained from 2018-released edition of Canada's  
24 Air Pollutant Emissions Inventory (APEI) originally compiled by the Pollutant Inventories and  
25 Reporting Division (PIRD) of Environment and Climate Change Canada (ECCC) (APEI,  
26 2018). This inventory contains a comprehensive and detailed estimate of annual emissions of  
27 seven criteria air pollutants (SO<sub>2</sub>, NO<sub>x</sub>, CO, NMVOC, NH<sub>3</sub>, PM<sub>10</sub>, PM<sub>2.5</sub>) at the national and  
28 provincial/territorial level for each year for the period from 1990 to 2016. The APEI inventory  
29 was developed based on a bottom-up approach for facility-level data reported to the National  
30 Pollutant Release Inventory (NPRI) (NPRI, 2017), as well as an in-house top-down emission  
31 estimates based on source-specific activity data and emissions factors. In general,  
32 methodologies used to estimate Canadian emissions are consistent with those developed by the  
33 U.S. EPA (EPA, 2009) or those recommended in the European emission inventory guidebook  
34 (EMEP/EEA, 2013). These methods are often further adjusted by PIRD to reflect the Canadian  
35 climate, fuels, technologies and practices.

36 To prepare emissions in the desired HTAP classification, the APEI sector emissions were first  
37 mapped to the United Nations Economic Commission for Europe (UNECE) Nomenclature for  
38 Reporting (NFR) categories, which involved dividing the sector emissions into their  
39 combustion and process components. The NFR categories were then mapped to the HTAP 16  
40 sector categories provided in the sector disaggregation scheme guide. Table S1 provides an  
41 overview of ECCC sector mapping to the HTAP\_v3 sectors.

42 The HTAP-grouped APEI inventory emissions files were further processed by the Air Quality  
43 Policy-Issue Response (REQA) Section of ECCC to prepare the air-quality-modelling version  
44 of inventory files in the standard format (i.e., FF10 format) supported by the U.S EPA  
45 emissions processing framework. To process emissions into gridded, speciated and total  
46 monthly values, a widely-used emissions processing system called the Sparse Matrix Operator



1 Kernel Emissions (SMOKE) model, version 4.7 (UNC, 2019) was used. As part of the  
2 preparation for SMOKE processing, a gridded latitude-longitude North American domain at  
3 0.1 x 0.1 degree resolution was defined with 920 columns and 450 rows covering an area of -  
4 142W to -50W and 40N to 85N. The point-source emissions in the APEI include latitude and  
5 longitude information so those sources were accurately situated in the appropriate grid cell in  
6 the Canadian HTAP gridded domain. However, to allocate provincial-level non-point source  
7 emissions into this domain, a set of gridded spatial surrogate fields was generated for each  
8 province from statistical proxies, such as population, road network, dwellings, crop  
9 distributions, etc. Over 80 different surrogate ratio files were created using the 2011 Canadian  
10 census data obtained from Statistics Canada website ([https://www12.statcan.gc.ca/census-  
11 recensement/2011/index-eng.cfm](https://www12.statcan.gc.ca/census-recensement/2011/index-eng.cfm)) and other datasets, such as the Canadian National Road  
12 Network (<https://open.canada.ca/data/en/dataset/3d282116-e556-400c-9306-ca1a3cada77f>).

13 To map the original APEI inventory species to the HTAP's desired list of species, PM  
14 speciation profiles from the SPECIATE version 4.5 database (EPA, 2016) were used to  
15 calculate source-type-specific EC and OC emissions. As a final step in SMOKE processing,  
16 the monthly emissions values were estimated using a set of sector-specific temporal profiles  
17 developed and recommended by the U.S. EPA (Sassi, 2021). For the point sources the NPRI  
18 annually reported monthly emissions proportions were applied. Emissions for the years 2017  
19 and 2018 were calculated by applying sector- and pollutant-specific trends from EDGAR.

#### 20 **2.3.4 REASv3.2.1 inventory**

21 The Regional Emission inventory in ASia (REAS) series have been developed for providing  
22 historical trends of emissions in the Asian region including East, Southeast, and South Asia.  
23 REASv3.2.1, the version used in HTAP\_v3, runs from 1950 to 2015. REASv3.2.1 includes  
24 emissions of SO<sub>2</sub>, NO<sub>x</sub>, CO, NMVOCs, NH<sub>3</sub>, CO<sub>2</sub>, PM<sub>10</sub>, PM<sub>2.5</sub>, BC, and OC from major  
25 anthropogenic sources: fuel combustion in power plant, industry, transport, and domestic  
26 sectors; industrial processes; agricultural activities; evaporation; and others.

27 Emissions from stationary fuel combustion and non-combustion sources are traditionally  
28 calculated using activity data and emission factors, including the effects of control  
29 technologies. For fuel consumption, the amount of energy consumption for each fuel type and  
30 sector was obtained from the International Energy Agency World Energy Balances for most  
31 countries and province-level tables in the China Energy Statistical Yearbook were used for  
32 China. Other activity data such as the amount of emissions produced from industrial processes  
33 were obtained from related international and national statistics. For emission factors, those  
34 without effects of abatement measures were set and then, effects of control measures were  
35 considered based on temporal variations of their introduction rates. Default emission factors  
36 and settings of country- and region-specific emission factors and removal efficiencies were  
37 obtained from scientific literature studies as described in Kurokawa and Ohara (2020) and  
38 references therein.

39 Emissions from road transport were calculated using vehicle numbers, annual distance  
40 travelled, and emission factors for each vehicle type. The number of registered vehicles were  
41 obtained from national statistics in each country and the World Road Statistics. For emission  
42 factors, year-to-year variation were considered by following procedures: (1) Emission factors  
43 of each vehicle type in a base year were estimated; (2) Trends of the emission factors for each  
44 vehicle type were estimated considering the timing of road vehicle regulations in each country  
45 and the ratios of vehicle production years; (3) Emission factors of each vehicle type during the  
46 target period were calculated using those of base years and the corresponding trends.

1 In REASv3.2.1, only large power plants were treated as point sources. For emissions from  
2 cement, iron, and steel plants, grid allocation factors were developed based on positions,  
3 production capacities, and start and retire years for large plants. Gridded emission data of  
4 EDGARv4.3.2 were used for grid allocation factors for the road transport sector. Rural, urban,  
5 and total population data were used to allocation emissions from the residential sector. For  
6 other sources, total population were used for proxy data.

7 For temporal distribution, if data for monthly generated power and production amounts of  
8 industrial products were available, monthly emissions were estimated by allocating annual  
9 emissions to each month using the monthly data as proxy. For the residential sector, monthly  
10 variation of emissions was estimated using surface temperature in each grid cell. If there is no  
11 appropriate proxy data, annual emissions were distributed to each month based on number of  
12 dates in each month.

13 Monthly gridded emission data sets at  $0.25^{\circ} \times 0.25^{\circ}$  resolution for major sectors and emission  
14 table data for major sectors and fuel types in each country and region during 1950-2015 are  
15 available in text format from a data download site of REAS (<https://www.nies.go.jp/REAS/>).  
16 Table S1 provides an overview about the REASv3.2.1 sector mapping to the HTAP\_v3 sectors.

17 More details of the methodology of REASv3.2.1 are available in Kurokawa and Ohara (2020)  
18 and its supplement. (Note that REASv3.2.1 is the version after error corrections of REASv3.2  
19 of Kurokawa and Ohara (2020)). Details of the error corrections are described in the data  
20 download site of REAS.) Table S1 provides an overview about the REASv3.2.1 sector mapping  
21 to the HTAP\_v3 sectors.

22 The MEIC inventory (<http://meicmodel.org>, 2021) is not currently included in the HTAP\_v3  
23 mosaic. Since the REAS inventory only includes emissions until 2015, the REAS-based  
24 HTAP\_v3 mosaic is only complete until this year. Emissions beyond 2015 were extrapolated  
25 using trends derived from a combination of MEIC and EDGAR. To extend the Chinese  
26 emission estimates to most recent years, MEIC data were used to adjust sector and pollutant  
27 specific trend for China for the years 2016 and 2017 (refer to Table S2 for the mapping sectors  
28 of MEIC and HTAP\_v3). Then, the 2018 data were calculated based on the 2015-2017 trend.  
29 For all the other countries belonging to the REAS domain, the emissions were extended beyond  
30 2015 applying the sector-, country-, and pollutant-specific trends from EDGAR.

### 31 **2.3.5 CAPSS-KU inventory**

32 In the Republic of Korea, the National Air Emission Inventory and Research Center (NAIR)  
33 estimates annual emissions of the air pollutants CO, NO<sub>x</sub>, SO<sub>x</sub>, TSP, PM<sub>10</sub>, PM<sub>2.5</sub>, BC, VOCs,  
34 and NH<sub>3</sub> via the Clean Air Policy Support System (CAPSS). The CAPSS inventory is divided  
35 into four source-sector levels (high, medium, low and detailed) based on the European  
36 Environment Agency's (EEA) CORE Inventory of AIR emissions (EMEP/CORINAIR). For  
37 activity data, various national- and regional-level statistical data collected from 150 domestic  
38 institutions are used. For large point sources, emissions are estimated directly using real-time  
39 stack measurements. For small point, area and mobile sources, indirect calculation methods  
40 using activity data, emission factors, and control efficiency are used.

41 Even though CAPSS (Clean Air Policy Support System) has been estimating annual emissions  
42 since 1999, some inconsistencies exist in the time series because of the data and methodological  
43 changes over the period. For example, emissions of PM<sub>2.5</sub> were initiated from the year 2011  
44 and not from 1999. Therefore, in the CAPSS emission inventory, PM<sub>2.5</sub> emissions were  
45 calculated from 2011, and post-2011 the PM<sub>10</sub> to PM<sub>2.5</sub> emission ratio was used to calculate the  
46 emissions from 2000 to 2010. These limitations make it difficult to compare and analyse

1 emissions inter-annually. To overcome these limitations, re-analysis of the annual emissions  
2 of pollutants was conducted using upgrades of the CAPSS inventory, such as missing source  
3 addition and emission factor updates.

4 The biomass combustion and fugitive dust sector emissions from 2000 to 2014 were estimated  
5 and added in the inventory, which are newly calculated emission sources from 2015. As for the  
6 on-road mobile sector, new emission factors using 2016 driving conditions were applied from  
7 the year 2000 to 2015. Since the emissions from the combustion of imported anthracite coal  
8 were calculated only from 2007, the coal use statistics of imported anthracite from 2000 to  
9 2006 were collected to estimate emissions for those years.

10 After all the adjustments, a historically re-constructed emissions inventory using the latest  
11 emission estimation method and data was developed. Table S1 provides an overview about the  
12 CAPSS sector mapping to the HTAP\_v3 sectors.

### 13 **2.3.6 JAPAN inventory (PM2.5EI and J-STREAM)**

14 The Japanese emission inventory contributing to the HTAP\_v3 mosaic is jointly developed by  
15 the Ministry of the Environment, Japan (MOEJ) for emissions arising from mobile sources and  
16 by the National Institute of Environmental Studies (NIES) for estimating emissions from fixed  
17 sources.

18 The mobile source emissions data for the HTAP\_5.1, 5.2, and 5.4 sectors are based on the air  
19 pollutant emission inventory named “PM2.5 Emission Inventory (PM2.5EI),  
20 <https://www.env.go.jp/air/osen/pm/info.html>). PM2.5EI has been developed for the years 2012  
21 and 2015, while for 2018 is currently under development. Almost all anthropogenic sources  
22 are covered, but emissions from vehicles are estimated in particular detail based on JATOP  
23 (Shibata, 2021). The emission factor of automobiles is constructed by MOEJ as a function of  
24 the average vehicle speed over several kilometres in a driving cycle that simulates driving on  
25 a real road. Emission factors are organized by 7 types of vehicles, 2 fuel types, 5 air pollutants,  
26 and regulation years, and have been implemented since 1997 as a project of MOEJ. By using  
27 these emission factors and giving the average vehicle speed on the road to be estimated, it is  
28 possible to estimate the air pollutant emissions per kilometre per vehicle. The hourly average  
29 vehicle speed of trunk roads, which account for 70% of Japan's traffic volume, is obtained at  
30 intervals of several kilometres nationwide every five years, so the latest data for the target year  
31 is used. For narrow roads, the average vehicle speed by prefecture measured by probe  
32 information is applied. It is 20 km/h in Tokyo, but slightly faster in other prefectures. Starting  
33 emission is defined as the difference between the exhaust amount in the completely cold state  
34 and the warm state in the same driving cycle and is estimated by the times the engine started  
35 in a day. Chassis dynamometer tests are performed in a well-prepared environment, so for more  
36 realistic emissions estimates, temperature correction factor, humidity correction factor,  
37 deterioration factor, DPF regeneration factor, and soak time correction factor are used. In  
38 addition to running and starting emissions, evaporative emissions from gasoline vehicles and  
39 non-exhaust particles such as road dust (including brake wear particles) and tire wear particles  
40 are combined to provide a vehicle emissions database with a spatial resolution of approximately  
41 1 km × 1 km (30° latitude, 45° longitude), and a temporal resolution of an hour by month,  
42 including weekdays and holidays.

1 Off-road vehicle emissions are estimated separately for 17 types of construction machinery,  
2 industrial machinery (forklifts), and 5 types of agricultural machinery. In all cases, emission  
3 factors by type and regulatory year per workload are used, as researched by the MOEJ.  
4 Although not as precise as automobiles, the off-road database is provided with the same  
5 temporal and spatial resolution as the automobile database.

6 Emissions from stationary sources in Japan are derived from the emission inventory developed  
7 in the Japan's Study for Reference Air Quality Modelling (J-STREAM) model intercomparison  
8 project (Chatani et al., 2018; Chatani et al., 2020). In this emission inventory, emissions from  
9 stationary combustion sources are estimated by multiplying emission factors and activities  
10 including energy consumption, which is available in the comprehensive energy statistics. Large  
11 stationary sources specified by the air pollution control law need to report emissions to the  
12 government every three years. The emission factors and their annual variations were derived  
13 from the emissions reported by over 100,000 sources (Chatani et al., 2020). For fugitive VOC  
14 emissions, MOEJ maintains a special emission inventory to check progress on regulations and  
15 voluntary actions targeting 30% reduction of fugitive VOC emissions starting from 2000. VOC  
16 emissions estimated in this emission inventory are used. Emissions from agricultural sources  
17 are consistent with the emissions estimated in the national greenhouse gas emission inventory  
18 (Center for Global Environmental Research et al., 2022). Emissions of all the stationary sources  
19 are divided into prefecture, city, and grid (approximately 1 x 1 km) levels based on spatial  
20 proxies specific to each source. Emissions for the year 2018 were estimated applying sector-  
21 and pollutant-specific trends from EDGAR. Table S1 provides an overview about the Japanese  
22 inventory sector mapping to the HTAP\_v3 sectors.

## 23 **2.4 Gap-filling methodology with EDGARv6.1**

24 EDGAR is a globally consistent emission inventory of air pollutant and greenhouse gases  
25 developed and maintained by the Joint Research Centre of the European Commission  
26 (<https://edgar.jrc.ec.europa.eu/>). The EDGAR methodology used to compute GHG and air  
27 pollutant emissions has been described in detail in several publications (Janssens-Maenhout et  
28 al., 2019; Crippa et al., 2018) and summarised here after. In EDGAR, air pollutant emissions  
29 are computed making use of international statistics as activity data (e.g., International Energy  
30 Balance data, Food and Agriculture Organisation statistics, US Commodity statistics, etc.),  
31 region- and/or country-specific emission factors by pollutant/sector and technology and  
32 abatement measures, following Eq. 1:

$$33 \quad EM_{i(C,t,x)} = \sum_{j,k} AD_{i(C,t)} * TECH_{i,j(C,t)} * EOP_{i,j,k(C,t)} * EF_{i(C,t,x)} * (1 - RED)_{i,j,k(C,t,x)}$$

34 (Eq. 1)

35 where EM are the emissions from a given sector *i* in a country *C* accumulated during a year *t*  
36 for a chemical compound *x*, AD the country-specific activity data quantifying the human  
37 activity for sector *i*, TECH the mix of *j* technologies (varying between 0 and 1), EOP the mix  
38 of *k* (end-of-pipe) abatement measures (varying between 0 and 1) installed with a share *k* for  
39 each technology *j*, and EF the uncontrolled emission factor for each sector *i* and technology *j*  
40 with relative reduction (RED) by abatement measure *k*. Emission factors are typically derived  
41 from the EMEP/EEA Guidebooks (EMEP/EEA, 2013, 2019, 2016), the AP-42(EPA, 2009)  
42 inventory and scientific literature.

1 Annual country and sector specific air pollutant emissions are then disaggregated into monthly  
2 values (Crippa et al., 2020) and subsequently spatially distributed making use of detailed proxy  
3 data (Janssens-Maenhout et al., 2019; Crippa et al., 2021).

4 As the most comprehensive and globally consistent emission database, the latest update of the  
5 EDGAR air pollutant emissions inventory, EDGARv6.1  
6 ([https://edgar.jrc.ec.europa.eu/dataset\\_ap61](https://edgar.jrc.ec.europa.eu/dataset_ap61)), is used in the HTAP\_v3 mosaic to complete  
7 missing information from the officially reported inventories, as reported in Table 3.  
8 EDGARv6.1 includes important updates to estimate air pollutant emissions such as the  
9 improvement of road transport emission estimates for many world regions, the inclusion of  
10 agricultural NMVOC emissions, revised monthly emission profiles (Crippa et al., 2020) and  
11 updated spatial proxies to distribute national emissions by sector over the globe (Crippa et al.,  
12 2021). EDGARv6.1 also includes new international shipping proxies and their monthly  
13 distribution based on the STEAM model (Jalkanen et al., 2012; Johansson et al., 2017). In the  
14 Supplementary Material (section S2), the assessment of EDGAR emission data is reported in  
15 comparison with global and regional inventories.

## 16 **3 Results**

### 17 **3.1 Annual time series analysis: trends and regional and sectoral contributions**

18 Having a consistent set of global annual emission inventories for a two-decade period allows  
19 the investigation of global emissions trends for the inventory pollutants and regional and  
20 sectoral contributions. Figure 2 presents annual time series (2000–2018) of the global emissions  
21 of the nine air pollutants included in the HTAP\_v3 mosaic separated into the actual  
22 contributions of 12 regions. Figure 3 shows the corresponding relative contributions of (a) 16  
23 sectors and (b) 12 regions to the 2018 global emissions of these same pollutants. We can then  
24 discuss each pollutant in turn.

25 Global SO<sub>2</sub> emissions declined from 99.4 to 72.9 Mt over the past two decades. This decreasing  
26 pattern is found for several world regions with the fastest decline in Eastern Asia, where after  
27 the year 2005 SO<sub>2</sub> emissions began to decrease steadily. This is consistent with the use of  
28 cleaner fuels with lower sulphur content and the implementation of desulphurisation techniques  
29 in power plants and industrial facilities in China in accordance with the 11th Five-Year Plan  
30 (FYP, 2006–2010 (Planning Commission, 2008)) and the 12th Five-Year Plan (FYP, 2011–  
31 2015 (Hu, 2016)) (Sun et al., 2018). Similarly, industrialised regions, such as North America  
32 and Europe, are characterised by a continuous decreasing trend in SO<sub>2</sub> emission, which had  
33 started well before the year 2000 due to the implementation of environmental and air quality  
34 legislation (EEA, 2022). Increasing SO<sub>2</sub> emissions, on the other hand, are found for Southern  
35 Asia (+112% compared to 2000), South-East Asia and developing Pacific (+62%), and Africa  
36 (+40%). These increases mostly arise from the energy, industry, and (partly) residential  
37 sectors, and reflect the need for emerging and developing economies to mitigate these  
38 emissions. Emissions estimated using satellite retrievals and model inversions confirm the  
39 trends provided by the HTAP\_v3 mosaic (Liu et al., 2018). SO<sub>2</sub> is mostly emitted by power  
40 generation and industrial activities, which in 2018 represent 42% and 26%, respectively, of the  
41 global total. Despite measures in some specific sea areas to mitigate sulphur emissions, globally  
42 they have been rising steadily with increasing activity. International shipping represents 13.8%  
43 of global SO<sub>2</sub> emissions in 2018, and it is 41% higher compared to the 2000 levels (Fig. 3).

44 Global NO<sub>x</sub> emissions increased from 110.4 Mt in 2000 to 117.4 Mt in 2018 as a result of the  
45 increase in energy- and industry-related activities for most of the world regions (in particular  
46 over the Asian domain). The strongest decreases are found for North America (-63%), Europe  
47 (-42%), Asia-Pacific Developed (-32%) and to a lower extent for Eurasia (-6%). Comparable

1 spatio-temporal patterns are found by satellite OMI data and ground based measurements of  
2 NO<sub>2</sub> concentrations (Jamali et al., 2020). NO<sub>x</sub> is mainly produced at high combustion  
3 temperatures (e.g., power and industrial activities, 38% of the global total), but also by  
4 transportation (27% of the global total) and international shipping (14% of the global total).

5 CO is mostly emitted by incomplete combustion processes from residential combustion,  
6 transportation and the burning of agricultural residues. Globally, CO emissions showed little  
7 change over the past two decades (502.7 Mt in 2000 vs. 499.8 in 2018), but different regional  
8 trends are present. Historically industrialised regions have reduced their emissions over the  
9 years (-42% in Europe and -62% in North America), while CO emissions increased in Africa  
10 by 45% and in Southern Asia by 49%. Road transport CO emissions halved over the past two  
11 decades (-55%), while the emissions from all other sectors increased. These results are  
12 consistent with MOPITT satellite retrievals, which mostly show the same trends over the  
13 different regional domains over the past decades (Yin et al., 2015).

14 NMVOC emissions increased from 115.2 Mt in 2000 to 146 Mt in 2018. These emissions are  
15 mostly associated with the use of solvents (25% of the 2018 global total), fugitive emissions  
16 (23%), road transportation (including both combustion and evaporative emissions, 15%) and  
17 small-scale combustion activities (19%). The most prominent increases in the emissions at the  
18 global level are found for the solvents sector (+81%). In 2018, NMVOC emissions from  
19 solvents were 3.7 and 3.5 times higher than in 2000 in China and India, respectively, while a  
20 rather stable trend in found for US and Europe.

21 Global NH<sub>3</sub> emissions increased from 47.4 Mt in 2000 to 58.9 Mt in 2018 due to enhanced  
22 emissions from agricultural activities. In particular, NH<sub>3</sub> emissions strongly increased in Africa  
23 (+60.5%), South-East Asia and developing Pacific (48.3%), Southern Asia (+38.7%), and Latin  
24 America and Caribbean (+41.1%).

25 Particulate matter emissions showed little change over the past two decades at the global level,  
26 whereas regional emission increases are found for Africa (e.g., +47.0% for PM<sub>10</sub>), Latin  
27 America and Caribbean (+39.1%), Middle East (48.3%), and Southern Asia (+56%), mostly  
28 associated with increases in agricultural waste burning and the livestock, energy, and waste  
29 sectors. By contrast, Eastern Asia (-39.5%), Europe (-24.3%), and Asia-Pacific Developed (-  
30 36.8%) significantly decreased their PM<sub>10</sub> emissions over the past two decades due to the  
31 continuous implementation of reduction and abatement measures for the energy, industry, road  
32 transport and residential sectors ((Crippa et al., 2016). As shown in Fig. 3, the relative  
33 contribution of North America to global PM<sub>10</sub> is quite high compared to other substances due  
34 to fugitive dust emissions (e.g., unpaved road dust, coal pile dust, dust from agricultural tilling)  
35 which have not been adjusted for meteorological conditions (e.g., rain, snow) and near-source  
36 settling and mitigation (e.g., tree wind breaks) because these removal mechanisms are better  
37 addressed by the chemical transport models. Additional uncertainty may be therefore  
38 introduced for these emissions, depending on the modelling assumptions of each official  
39 inventory. Similarly, particulate matter speciation into its carbonaceous components is often  
40 challenging and subjected to higher level of uncertainty, for instance because different  
41 definitions are used for PM in inventories, including condensable emissions or not (Denier van  
42 der Gon et al., 2015). Attempts to improve the accuracy of such emissions (e.g. BC and OC  
43 emissions over the European domain) are ongoing.

44 Figure 3a shows more than 50% difference at the global level between PM<sub>2.5</sub> emissions and the  
45 sum of its carbonaceous components (BC and OC), which however varies depending on the  
46 region and sector. The largest difference between PM<sub>2.5</sub> and the sum of BC and OC is generally  
47 found for the energy and industrial sectors, where due to the high temperatures BC and OC are

1 largely burned. Within this sector, the non carbonaceous fraction of PM<sub>2.5</sub> represents around  
2 75% in Europe, 78% in the USA and up to more than 95% over Asian countries (e.g. China  
3 and India). This PM fraction is represented by other minerals, ash (mostly when burning coal)  
4 and sulphate. Road transport is also a sector showing large differences between PM<sub>2.5</sub> and the  
5 sum of BC and OC, with around 40% difference for Europe, around 90% difference for USA  
6 and lower values for India and China (around 15%). This component may be associated with  
7 other minerals. For the residential sector, this difference is generally lower and around 25%  
8 (for Europe and Asian countries), while around 37% in the USA and is possibly associated  
9 with other minerals and ash due to coal combustion. Shipping is also a sector where a large  
10 component of PM<sub>2.5</sub> (around 70%) is not associated with carbonaceous fractions but to  
11 sulphate. In particular, regions within the Sulphur Emission Control Area (SECA) show lower  
12 contributions from sulphates (e.g. Europe and USA) with an overall contribution of 5-10%.  
13 Another source of uncertainty which may contribute to enhancing the difference between PM<sub>2.5</sub>  
14 and the sum of BC and OC is associated on how different inventories consider condensable  
15 particulate matter.

### 16 **3.2 Emission maps**

17 Spatially distributed emission data describe where emissions take place, as input for local,  
18 regional and global air quality modelling. As noted in section 2.2, nationally aggregated air  
19 pollutant emissions are spatially distributed over the corresponding national territory using  
20 spatial proxy data which are believed to provide a relatively good representation of where  
21 emissions takes place. Depending on the emitting sector, air pollutants can be associated with  
22 the spatial distributions of point sources (e.g., in the case of power plant or industrial activities),  
23 road networks (e.g., for transportation related emissions), settlement areas (e.g., for small-scale  
24 combustion emissions), crop and livestock distribution maps, ship tracks etc. Using reliable  
25 and up-to-date spatial information to distribute national emissions is therefore relevant,  
26 although challenging. Multiple assumptions are often made by inventory compilers when  
27 developing their inventories, which may result in differences when analysing spatially  
28 distributed emissions provided by different inventory compilers over the same geographical  
29 domain.

30 One key goal of the HTAP\_v3 mosaic is to collate in one inventory the most accurate spatially-  
31 distributed emissions for all air pollutants at the global level, based on the best available local  
32 information. Point sources related with emissions from power plant and industrial facilities  
33 represent one the most critical spatial information to be retrieved, and their misallocation can  
34 significantly affect the characterisation of local air quality. This challenge is also present in the  
35 HTAP\_v3 mosaic. For example, the REASv3.2.1 inventory is still using limited information  
36 to distribute emissions from these two sectors especially for industrial plants. Depending on  
37 the region, point source information could be limited compared to datasets used in inventories  
38 of North America and Europe. To overcome this issue, the participation of national emission  
39 inventory developers not only from China, but also India and other Asian countries is  
40 recommended. The impact can be seen in Fig. 4, which shows the global map of SO<sub>2</sub> emissions  
41 in 2018 based on the HTAP\_v3 mosaic compilation, where information about the magnitude  
42 and the type of emission sources for the different regions can be retrieved. The energy and  
43 industry sectors contribute a large fraction of SO<sub>2</sub> emissions (Fig. 3a), but the spatial  
44 distribution of these emissions is qualitatively different in North America and Europe than in  
45 Asia (i.e., more “spotty”, less smooth and widely distributed). Ship tracks cover the entire  
46 geographical marine domain, consistent with emissions from the STEAM model (Jalkanen et  
47 al., 2012; Johansson et al., 2017) included in the EDGARv6.1 database, although showing  
48 marked emissions over the Mediterranean Sea, Asian domain, Middle East and North

1 American coasts. Furthermore, emissions from power plant and industrial activities, as well as  
2 small-scale combustion are prominent over the Asian domain, Eastern Europe, and some  
3 African regions.

4 Sector-specific case studies are presented in the maps of Figs. 5-8. Figure 5 shows the  
5 comparison of annual NO<sub>x</sub> emissions for the year 2000 and 2018. The road transport sector is  
6 a key source of NO<sub>x</sub> emissions (cf. Fig. 3a), and this contribution is reflected in the visible  
7 presence of road networks in the maps. Decreasing emissions are found for industrialised  
8 regions (USA, Europe, Japan) thanks to the introduction of increasingly restrictive legislation  
9 on vehicle emissions since the 1990s, whereas a steep increase is found for emerging  
10 economies and in particular India, China, and the Asian domain. Figure 6 shows the different  
11 spatial allocation of PM<sub>10</sub> emissions from the residential sector during the month of January  
12 2018, with higher emission intensities evident in the Northern Hemisphere (cold season) and  
13 the lower values in the Southern Hemisphere (warm season). Figures 7 and 8 show the spatio-  
14 temporal allocation of agriculture-related emissions, and specifically, PM<sub>10</sub> emissions from  
15 agricultural waste burning and NH<sub>3</sub> emissions from agricultural soil activities.

### 16 **3.3 Monthly temporal distribution**

#### 17 **3.3.1 Monthly variability by region**

18 The magnitude of air pollutant emissions varies by month because of the seasonality of  
19 different anthropogenic activities and their geographical location (e.g., Northern vs. Southern  
20 Hemisphere regions). Figures 9 and 10 (and S3.1, S3.2 and S3.3) show the monthly distribution  
21 of regional emissions for those pollutants and sectors for which higher variability is expected.  
22 The year 2015 was chosen since it is the last year for which all of the official data providers  
23 have data. Figure 9 shows monthly NH<sub>3</sub> emissions by region from three agricultural activities  
24 (agricultural waste burning, livestock, and crops). These sectors display the largest variability  
25 by month, reflecting the seasonal cycle and the region-specific agricultural practices, such as  
26 fertilisation, crop residue burning, manure and pasture management, animal population  
27 changes, etc. In Figure 10, NO<sub>x</sub> emissions from residential activities show a particular monthly  
28 distribution, with the highest emissions occurring during the cold months shifted for the  
29 Northern and Southern Hemispheres. By contrast, regions in the equatorial zone do not show  
30 a marked monthly profile even for residential activities. The energy sector also follows  
31 monthly-seasonal cycles related to the demand for power generation, which is also correlated  
32 with ambient temperature and local day length. Transport-related emissions do not show a large  
33 variation by month, whereas daily and weekly cycles for transport-related emissions, which are  
34 typically more relevant, are beyond the temporal resolution of this work.

35 Although a spatio-temporal variability of the HTAP\_v3 emissions is found in these figures, a  
36 more in-depth analysis reveals that with the exception of few regions and sectors (e.g., Canada,  
37 USA and regions gap-filled with EDGAR), no inter-annual variability of the monthly profiles  
38 is present, meaning that the majority of official inventories assume the same monthly  
39 distribution of the emissions for the past two decades (refer to Figs. S3.4-S3.9). This is different  
40 from the approach used for example by EDGAR (Crippa et al., 2020), ECCC for Canada, and  
41 U.S. EPA for the USA, where year-dependent monthly profiles are used for specific sectors, in  
42 particular for residential, power generation, and agricultural activities. Further analysis has  
43 shown that for the European domain regional rather than country-specific monthly profiles are  
44 applied. Therefore, for Europe new state-of-the-art profiles have been made available under  
45 the CAMS programme by Guevara et al. (2021).



### 1 **3.3.2 Spatially-distributed monthly emissions**

2 An important added value of HTAP\_v3 comes from the availability of monthly gridmaps that  
3 reflect the seasonality of the emissions for different world regions. Access to spatially  
4 distributed monthly emissions is essential to design effective mitigation actions, providing  
5 information on hot spots of emissions and critical periods of the year when emissions are  
6 highest.

7 Figure 11 shows mid-season PM<sub>2.5</sub> monthly emissions arising from the residential sector in  
8 2018. The global map shows higher emissions in the Northern Hemisphere during January,  
9 while the opposite pattern is found for the Southern Hemisphere in July. Agriculture is an  
10 important activity characterised by strong seasonal patterns, as shown in Figs. 12 and 13. Figure  
11 12 shows PM<sub>10</sub> monthly emission maps from agricultural residue burning in 2018 from  
12 HTAP\_v3, highlighting higher emissions over certain months of the year related with specific  
13 burning practices of agricultural residues for different world regions. For example, during the  
14 month of April, intense burning of crop residues is found in Africa (Nigeria, Ethiopia, Sudan,  
15 South Africa, etc.), South America (Brazil, Argentina, Colombia, etc.), Northern India, and  
16 South-Eastern Asia (e.g., Vietnam, Thailand, Indonesia, Philippines, etc.). Figure 13 represents  
17 the yearly variability of NH<sub>3</sub> emissions from agricultural soils activities, mostly related with  
18 fertilisation. During the month of March and April, intense agricultural soils activities are found  
19 over Europe and North America compared to other months, while during the month of October  
20 the highest emissions are for this sector are found in China, India, several countries of the Asian  
21 domain, but also in USA, Australia, and Latin America. These results are consistent with  
22 satellite based observations performed using Cross-track Infrared Sounder (Shephard et al.,  
23 2020).

### 24 **3.4 Vertical distribution of the emissions**

#### 25 **3.4.1 Aircraft emissions**

26 In EDGAR6.1 the emissions are provided at three effective altitude levels (landing/take-off,  
27 ascent/descent, and cruising). The spatial proxy for the aviation sector is derived from  
28 International Civil Aviation Organization (ICAO, 2015) which specifies a typical flight pattern  
29 with landing/take-off cycle within few km of the airport, followed by climb-out/descending  
30 phase during the first 100 km and the last 100km of a flight and finally the remaining part from  
31 101 km until the last 101 km as the cruise phase. Routes and airport locations are taken from  
32 the Airline Route Mapper of ICAO (2015). In HTAP\_v3, aircraft emissions are provided as  
33 domestic and international, but with no information about altitude ranges. We recommend  
34 modellers to use the corresponding EDGARv6.1 data  
35 ([https://edgar.jrc.ec.europa.eu/dataset\\_ap61](https://edgar.jrc.ec.europa.eu/dataset_ap61)) including the vertical distribution of the  
36 emissions.

#### 37 **3.4.2 Speciation of NMVOC emissions**

38 For emission data to be useful for modellers, total NMVOC emissions must be decomposed  
39 into emissions of individual NMVOC species. As the chemical mechanisms used by models  
40 can differ with respect to the NMVOC species they include, it is not practical to provide an  
41 NMVOC speciation which is usable by all models. Instead, a speciation is provided here for  
42 the set of 25 NMVOCs defined by Huang et al. (2017) and the corresponding data are made  
43 available on the HTAP\_v3 website. The absolute values of 25-category speciated NMVOC  
44 emissions were obtained for all countries for the 28 EDGAR sectors from here:  
45 [https://edgar.jrc.ec.europa.eu/dataset\\_ap432\\_VOC\\_spec](https://edgar.jrc.ec.europa.eu/dataset_ap432_VOC_spec). The absolute NMVOC emissions of  
46 each species from each sector in this dataset were remapped to the HTAP\_v3 sectors following  
47

1 the mapping from Table 2, then converted to a speciation by dividing by the total emissions of  
2 each individual species for the four world regions defined by Huang et al. (2017): Asia; Europe;  
3 North America; and Other. The resulting NMVOC speciation is provided in the supplementary  
4 material to this paper for the 25 NMVOC species, 4 world regions, and 15 emitting NMVOC  
5 sectors<sup>2</sup> following the HTAP\_v3 sector classification (including 13 sectors defined over the 4  
6 world regions, and the two international sectors: international shipping and international  
7 aviation). The list of countries comprising each region is also provided in the supplement.

### 8 **3.5 Qualitative assessment of the uncertainty of a global emission mosaic**

9 Assessing the uncertainty of a global emission mosaic is challenging since it consists of several  
10 bottom-up inventories and by definition it prevents a consistent global uncertainty calculation.  
11 Each emission inventory feeding the HTAP\_v3 mosaic is characterized by its own uncertainty  
12 which is documented by the corresponding literature describing each dataset (see Table 2 and  
13 section 2.3) and which should be cited by the users of the mosaic for a quantitative assessment  
14 of regional uncertainties. However, the mosaic compilation process may also introduce  
15 additional uncertainties compared to the input datasets. In order to limit these additional  
16 uncertainties, we made the following considerations:

17 -for each emission inventory both the national totals and gridded data by sector were gathered.  
18 This process allows the mosaic compilers not to introduce additional uncertainty compared to  
19 the original input regional datasets. In fact, additional uncertainties may arise from the  
20 extraction of the national totals from spatially distributed data (e.g. country border issues which  
21 were one limitation of previous editions of the HTAP mosaics). Therefore, when regional  
22 trends are described by region and pollutant (see section 3), no additional source of uncertainty  
23 has to be considered from the mosaic compilation approach.

24 -the sector definition and mapping has been accurately developed following the IPCC  
25 categories and when no data was available for a certain combination of sector and pollutant a  
26 gapfilling procedure is applied using the EDGAR database. Also in this case no additional  
27 uncertainty should be considered compared to the input datasets.

28 -any additional uncertainty introduced by the temporal disaggregation can be deemed as  
29 negligible since each inventory already provided monthly resolution emission gridmaps and  
30 time series.

31 In this work we also provide a qualitative indication of the emission variability by HTAP sector  
32 and pollutant at the global level. Table S6 summarises the variability of global HTAP\_v3  
33 emissions by sector for the boundary years of this mosaic (2000 and 2018) compared to the  
34 global EDGARv6.1 data. EDGAR emissions are considered as the reference global emission  
35 inventory against which comparing the HTAP\_v3 estimates although these two global products  
36 are not fully independent. The variability of the global emissions is calculated as the relative  
37 difference of the estimates of the two inventories, i.e.  $(\text{EDGARv6.1-HTAP\_v3})/\text{HTAP\_v3}$ .  
38 Emission variabilities are also classified as low (L,  $L < 15\%$ ), low medium (LM,  
39  $15\% < LM < 50\%$ ), upper medium (UM,  $50\% < UM < 100\%$ ), high (H,  $H > 100\%$ ), based on the  
40 EMEP/EEA Guidebook (2019) information. The largest variability is found domestic shipping  
41 emissions (CO and NMVOC), energy (OC, BC), agricultural crops (PM), road transport (PM,  
42 NMVOC) and industry (NH<sub>3</sub>, NMVOC). In absence of a full uncertainty assessment the

---

<sup>2</sup> No speciation profile is provided for the 'tyre and brake wear sector' not being a source of NMVOC emissions.

1 variability can be used as proxy of structural uncertainty, keeping in mind that variability could  
2 be biased towards overconfidence, thus underestimating the uncertainty. Furthermore, the  
3 uncertainty of the spatial proxies has not been assessed and maybe subject of future activity  
4 updates.

5

#### 6 **4 Data availability**

7 The HTAP\_v3 emission mosaic data can be freely accessed and cited using  
8 <https://doi.org/10.5281/zenodo.7516361>. All data can be also accessed through the EDGAR  
9 website at the following link: [https://edgar.jrc.ec.europa.eu/dataset\\_htap\\_v3](https://edgar.jrc.ec.europa.eu/dataset_htap_v3).

10 Data are made available in the following formats:

- 11 • Monthly gridmaps of emissions (in Mg/month) at 0.1x0.1degree resolution: there is one  
12 .NetCDF file per year and substance that includes the emissions for each sector for the  
13 12 months.
- 14 • Monthly gridmaps of emission fluxes (in kg/m<sup>2</sup>/s) at 0.1x0.1degree resolution: there is  
15 one .NetCDF file per year and substance that includes the emission fluxes for each  
16 sector the emission fluxes for the 12 months.
- 17 • Annual gridmaps of emissions (in Mg/year at 0.1x0.1degree resolution: there is one  
18 .NetCDF file per year and substance that includes the emissions for each sector.
- 19 • Annual gridmaps of emission fluxes (in kg/m<sup>2</sup>/s) at 0.1x0.1degree resolution: there is  
20 one .NetCDF file per year and substance that includes the emission fluxes for each  
21 sector.

22 The full set of HTAP\_v3 data is quite large, requiring substantial network bandwidth and time  
23 for download, and substantial storage space. To make it easier for users to query and use the  
24 data, additional products are available. For global modellers who may not require such high  
25 spatial resolution, gridmaps at 0.5x0.5 degree resolution are made available following the  
26 abovementioned specifications of the higher spatial resolution data. Furthermore, to allow  
27 regional modellers to download only the data for the regions they need, the JRC EDGAR group  
28 has also developed an interface to allow the users of the HTAP\_v3 mosaic to extract emission  
29 data over arbitrarily specified geographical domains. The HTAP tool is accessible after creation  
30 of an ECAS account (<https://webgate.ec.europa.eu/cas/login>) and it is available at:  
31 [https://edgar.jrc.ec.europa.eu/htap\\_tool/](https://edgar.jrc.ec.europa.eu/htap_tool/).

32

#### 33 **5 Conclusions**

34 The global air pollution mosaic inventory HTAP\_v3 presented and discussed in this paper is a  
35 state-of-the-art database for addressing the present status and the recent evolution of a set of  
36 policy-relevant air pollutants. The inventory is made by the harmonization and blending of six  
37 regional inventories, gapfilled using the most recent release of EDGAR (EDGARv6.1). By  
38 directly incorporating the best available local information, including the spatial distribution of  
39 emissions, the HTAP\_v3 mosaic inventory can be used for policy-relevant studies at both  
40 regional and global levels. As such, the HTAP\_v3 mosaic inventory provides a complement to  
41 globally consistent emission inventories such as EDGAR. The global and regional trends of air  
42 pollutant emissions in the HTAP\_v3 mosaic are comparable with other commonly available  
43 global emission datasets.

1 By providing consistent times series for almost two decades, HTAP\_v3 allows an evaluation  
2 of the impact and success of the pollution control measures deployed across various regions of  
3 the world since 2000. Similarly, its finer sectoral resolution is suitable for understanding how  
4 and where technological changes have resulted in emissions reductions, suggesting possible  
5 pathways for strengthening appropriate policy actions.

6 All these features make HTAP\_v3 a database of interest for policy makers active in the air  
7 quality regulatory efforts. HTAP\_v3 provides a picture of a world where most pollutant  
8 emissions are following a steady or decreasing path. However, several areas of the world show  
9 an increasing emission trend, with wide portions of the world remaining subjected to  
10 unsatisfactory levels of ambient air quality.

11 When using the HTAP\_v3 emission mosaic, users should consider the following limitations,  
12 for example when combining the HTAP\_v3 data with other emission input needed to run  
13 atmospheric models:

14 - agricultural waste burning emissions should be treated with caution to avoid double-counting  
15 when combined with existing biomass burning emission inventories;

16 - NMVOC and NO<sub>x</sub> emissions from agricultural soils should be treated with caution to avoid  
17 double-counting when combining the HTAP\_v3 data with a natural emissions model such as  
18 MEGAN (Model of Emissions of Gases and Aerosols from Nature);

19 - the speciation of NO<sub>x</sub> emissions into its components (NO, NO<sub>2</sub>, HONO) is not provided by  
20 the global HTAP\_v3 mosaic and it is beyond the scope of the current work since the regional  
21 inventories report total NO<sub>x</sub> with no speciation. Standard practice in global models is to emit  
22 all anthropogenic NO<sub>x</sub> as NO, while we expect that regional modelling groups will have access  
23 to appropriate best practices for their particular regions. In particular for road transport, the  
24 partitioning of NO<sub>x</sub> emissions between NO, NO<sub>2</sub>, and HONO is highly region-dependent and  
25 it is based on the fleet composition (e.g., number of diesel vehicles relative to gasoline vehicles)  
26 and technology level (e.g., the level of exhaust after treatment).

27 Thanks to the continuous improvement of local and regional emission inventories, recent  
28 literature shows new datasets that report regional information over areas of the world not  
29 covered by local inventories in the current HTAP\_v3 mosaic (e.g. Argentina (Puliafito et al.  
30 2021), Africa (Keita et al., 2021) and the MEIC inventory  
31 ([http://meicmodel.org.cn/?page\\_id=1772&lang=en](http://meicmodel.org.cn/?page_id=1772&lang=en))). Future updates to this mosaic may also  
32 integrate reliable and up to data information over South America or Africa as time and  
33 resources permit.

34 Similar to its predecessor (e.g. HTAP\_v2.2 mosaic inventory), we expect that this new  
35 HTAP\_v3 mosaic inventory will be used as a basis for global assessments of long-range,  
36 transboundary transport of air pollution under the Task Force on Hemispheric Transport of Air  
37 Pollution, while also providing a convenient and useful information for regional modellers  
38 seeking the best available regional emissions with a consistent gap-filling methodology.

39

#### 40 **Author contributions.**

41 MC and DG developed the mosaic gathering input from all data providers. The co-chairs of the  
42 TF-HTAP (TK, TB, RW and JaKa) fostered the dialogue with international institutions

1 contributing to this work with their data. PM, RM, JR, JZ, DN, MS, MDM, RW provided data  
2 for Canada, JuKu, SC, TM provided data for Japan, JeKu provided data for Europe, J-HW, JK  
3 provided data for Korea, TK, GP provided data for USA, JiKi provided data for Asia. The JRC  
4 EDGAR group (MC, ES, DG, EP, MM, FM, ES, MB, FP) lead the drafting of the publication  
5 with input from colleagues contributing to the HTAP\_v3 mosaic. SJS and HS performed  
6 detailed data comparison among available emission inventories. TA calculated and provided  
7 the NMVOC speciation fractions for all the sectors for the four regions.

8

9 **Competing interests.** The authors declare that they have no conflicts of interest nor competing  
10 interests.

### 11 **Acknowledgements.**

12 The authors would like to thank all HTAP\_v3 data providers for the fruitful cooperation. The  
13 views expressed in this publication are those of the authors and do not necessarily reflect the  
14 views or policies of the European Commission.

15 J-STREAM emission inventory for Japan was developed by Environment Research and  
16 Technology Development Fund (JPMEERF20165001 and JPMEERF20215005) of the  
17 Environmental Restoration and Conservation Agency Provided by the Ministry of  
18 Environment of Japan, and FRIEND (Fine Particle Research Initiative in East Asia Considering  
19 National Differences) Project through the National Research Foundation of Korea (NRF)  
20 funded by the Ministry of Science and ICT (2020M3G1A1114622).

### 21 **6 References**

22

23 Álamos, N., Huneus, N., Opazo, M., Osses, M., Puja, S., Pantoja, N., Denier van der Gon, H., Schueftan,  
24 A., Reyes, R., and Calvo, R.: High-resolution inventory of atmospheric emissions from transport,  
25 industrial, energy, mining and residential activities in Chile, *Earth Syst. Sci. Data*, 14, 361-379,  
26 10.5194/essd-14-361-2022, 2022.

27 APEI: Air Pollutant Emissions Inventory: overview, Canada.ca,  
28 <https://www.canada.ca/content/dam/eccc/images/apEI/apEI-2018-en.pdf>, 2018.

29 Center for Global Environmental Research, Earth System Division, and National Institute for  
30 Environmental Studies: National Greenhouse Gas Inventory Report of JAPAN 2022, Greenhouse Gas  
31 Inventory Office of Japan and Ministry of the Environment, Japan (eds.), 2022.

32 Chatani, S., Shimadera, H., Itahashi, S., Yamaji, K. J. A. C., and Physics: Comprehensive analyses of  
33 source sensitivities and apportionments of PM 2.5 and ozone over Japan via multiple numerical  
34 techniques, *Atmos. Chem. Phys.*, 20, 10311-10329, doi.org/10.5194/acp-20-10311-2020, 2020.

35 Chatani, S., Yamaji, K., Sakurai, T., Itahashi, S., Shimadera, H., Kitayama, K., and Hayami, H. J. A.:  
36 Overview of model inter-comparison in Japan's study for reference air quality modeling (J-STREAM),  
37 *Atmosphere*, 9, 19, <https://doi.org/10.3390/atmos9010019>, 2018.

38 Copernicus Land Monitoring Service: CORINE Land Cover 2012, available at:  
39 <https://land.copernicus.eu/pan-european/corine-land-cover/clc-2012> (last access: 13 August 2018),  
40 2016.

- 1 Crippa, M., Janssens-Maenhout, G., Dentener, F., Guizzardi, D., Sindelarova, K., Muntean, M.,  
2 Van Dingenen, R., and Granier, C.: Forty years of improvements in European air quality: regional  
3 policy-industry interactions with global impacts, *Atmos. Chem. Phys.*, 16, 3825-3841, 10.5194/acp-16-  
4 3825-2016, 2016.
- 5 Crippa, M., Solazzo, E., Huang, G., Guizzardi, D., Koffi, E., Muntean, M., Schieberle, C., Friedrich, R.,  
6 and Janssens-Maenhout, G.: High resolution temporal profiles in the Emissions Database for Global  
7 Atmospheric Research, *Scientific Data*, 7, 121, 10.1038/s41597-020-0462-2, 2020.
- 8 Crippa, M., Guizzardi, D., Pisoni, E., Solazzo, E., Guion, A., Muntean, M., Florczyk, A., Schiavina, M.,  
9 Melchiorri, M., and Hutfilter, A. F.: Global anthropogenic emissions in urban areas: patterns, trends,  
10 and challenges, *Environmental Research Letters*, 16, 074033, 10.1088/1748-9326/ac00e2, 2021.
- 11 Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Dentener, F., van Aardenne, J. A., Monni, S.,  
12 Doering, U., Olivier, J. G. J., Pagliari, V., and Janssens-Maenhout, G.: Gridded emissions of air pollutants  
13 for the period 1970–2012 within EDGAR v4.3.2, *Earth Syst. Sci. Data*, 10, 1987-2013, 10.5194/essd-10-  
14 1987-2018, 2018.
- 15 Denier van der Gon, H. A. C., Bergström, R., Fountoukis, C., Johansson, C., Pandis, S. N., Simpson, D.,  
16 and Visschedijk, A. J. H.: Particulate emissions from residential wood combustion in Europe – revised  
17 estimates and an evaluation, *Atmos. Chem. Phys.*, 15, 6503-6519, 10.5194/acp-15-6503-2015, 2015.
- 18 EEA: Emission trends of sulphur oxides, [https://www.eea.europa.eu/data-and-maps/daviz/emission-](https://www.eea.europa.eu/data-and-maps/daviz/emission-trends-of-sulphur-oxides#tab-chart_1)  
19 [trends-of-sulphur-oxides#tab-chart\\_1](https://www.eea.europa.eu/data-and-maps/daviz/emission-trends-of-sulphur-oxides#tab-chart_1), Last access: October 2022, 2022.
- 20 EMEP/EEA: EMEP/EEA Air Pollutant Emission Inventory Guidebook 2013. Technical Guidance to  
21 Prepare National Emission Inventories, Luxembourg: Publications Office of the European Union,  
22 Technical Report No. 12/2013, <http://www.eea.europa.eu/publications/emep-eeaguidebook-2013>,  
23 2013.
- 24 EMEP/EEA: EMEP/EEA air pollutant emission inventory guidebook 2016, EEA Report No 21/2016,  
25 2016.
- 26 EMEP/EEA: EMEP/EEA air pollutant emission inventory guidebook 2019, European Environment  
27 Agency, 2019, Luxembourg, Publications Office of the European Union, ISBN 978-92-9480-098-5, ISSN  
28 1977-8449, doi:10.2800/293657, 2019, European Environment Agency, Luxembourg, Publications  
29 Office of the European Union, ISBN 978-92-9480-098-5, ISSN 1977-8449, doi:10.2800/293657, 2019.
- 30 EPA: AP-42: Compilation of Air Emissions Factors, [https://www.epa.gov/air-emissions-factors-and-](https://www.epa.gov/air-emissions-factors-and-quantification/ap-42-compilation-air-emissions-factors)  
31 [quantification/ap-42-compilation-air-emissions-factors](https://www.epa.gov/air-emissions-factors-and-quantification/ap-42-compilation-air-emissions-factors), 2009.
- 32 EPA: Final report SPECIATE version 4.5 database development documentation, Report EPA/600/R-  
33 16/294 (Abt Associates, 2016), SPECIATE 4.5: Speciation Database Development Documentation -  
34 September 30, 2016 (epa.gov) 2016.
- 35 European Commission: Directive 2001/80/EC of the European Parliament and of the Council of 23  
36 October 2001 on the limitation of emissions of certain pollutants into the air from large combustion  
37 plants, available at: <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=celex%3A32001L0080> (last  
38 access: 1 July 2021), 2001.
- 39 European Commission: Directive (EU) 2016/2284 on the reduction of national emissions of certain  
40 atmospheric pollutants, available at:

1 [content/EN/TXT/?uri=uriserv:OJ.L .2016.344.01.0001.01.ENG&toc=OJ:L:2016:344:TOC](https://www.fao.org/land-water/land-land-governance/land-resources-planning-toolbox/category/details/en/c/1236449/) (last access: 1  
2 July 2021), 2016.

3 FAO: Gridded Livestock of the World (GLW3), available at: [http://www.fao.org/land-water/land-land-](http://www.fao.org/land-water/land-land-governance/land-resources-planning-toolbox/category/details/en/c/1236449/)  
4 [governance/land-resources-planning-toolbox/category/details/en/c/1236449/](http://www.fao.org/land-water/land-land-governance/land-resources-planning-toolbox/category/details/en/c/1236449/) (last access: 10  
5 August 2018), 2010.

6 Foley, K. M., Pouliot, G. A., Eyth, A., Aldridge, M. F., Allen, C., Appel, K. W., Bash, J. O., Beardsley, M.,  
7 Beidler, J., Choi, D., Farkas, C., Gilliam, R. C., Godfrey, J., Henderson, B. H., Hogrefe, C., Koplitz, S. N.,  
8 Mason, R., Mathur, R., Misenis, C., Possiel, N., Pye, H. O. T., Reynolds, L., Roark, M., Roberts, S.,  
9 Schwede, D. B., Seltzer, K. M., Sonntag, D., Talgo, K., Toro, C., Vukovich, J., Xing, J., and Adams, E.:  
10 2002–2017 anthropogenic emissions data for air quality modeling over the United States, Data in Brief,  
11 47, 109022, <https://doi.org/10.1016/j.dib.2023.109022>, 2023.

12 Guevara, M., Jorba, O., Tena, C., Denier van der Gon, H., Kuenen, J., Elguindi, N., Darras, S., Granier,  
13 C., and Pérez García-Pando, C.: Copernicus Atmosphere Monitoring Service TEMPORal profiles (CAMS-  
14 TEMPO): global and European emission temporal profile maps for atmospheric chemistry modelling,  
15 Earth Syst. Sci. Data, 13, 367-404, 10.5194/essd-13-367-2021, 2021.

16 Hu, A.-G.: The Five-Year Plan: A new tool for energy saving and emissions reduction in China, J  
17 Advances in Climate Change Research, 7, 222-228, <https://doi.org/10.1016/j.accre.2016.12.005>,  
18 2016.

19 Huang, G., Brook, R., Crippa, M., Janssens-Maenhout, G., Schieberle, C., Dore, C., Guizzardi, D.,  
20 Muntean, M., Schaaf, E., and Friedrich, R.: Speciation of anthropogenic emissions of non-methane  
21 volatile organic compounds: a global gridded data set for 1970–2012, Atmos. Chem. Phys., 17, 7683-  
22 7701, <https://doi.org/10.5194/acp-17-7683-2017>, 2017.

23 Huneus, N., van Der Gon, H. D., Castesana, P., Menares, C., Granier, C., Granier, L., Alonso, M., de  
24 Fatima Andrade, M., Dawidowski, L., and Gallardo, L. J. A. E.: Evaluation of anthropogenic air pollutant  
25 emission inventories for South America at national and city scale, 235, 117606, 2020.

26 ICAO: International Civil Aviation Organisation data: Airline Route Mapper data,  
27 <http://arm.64hosts.com/>, 2015.

28 IIASA: Progress towards the achievement of the EU's Air Quality and Emission Objectives, available at:  
29 [https://ec.europa.eu/environment/air/pdf/clean\\_air\\_outlook\\_overview\\_report.pdf](https://ec.europa.eu/environment/air/pdf/clean_air_outlook_overview_report.pdf) (last access: 1  
30 July 2021) 2018.

31 IPCC: Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories: Reporting Instructions.,  
32 <https://www.ipcc-nggip.iges.or.jp/public/gl/guidelin/ch1ri.pdf>, 1996.

33 IPCC: Guidelines for National Greenhouse Gas Inventory, Volume 5: Waste. Internet: [http://www.ipcc-](http://www.ipcc-nggip.iges.or.jp/public/2006gl/vol5.html)  
34 [nggip.iges.or.jp/public/2006gl/vol5.html](http://www.ipcc-nggip.iges.or.jp/public/2006gl/vol5.html), 2006.

35 Jalkanen, J. P., Johansson, L., Kukkonen, J., Brink, A., Kalli, J., and Stipa, T.: Extension of an assessment  
36 model of ship traffic exhaust emissions for particulate matter and carbon monoxide, Atmos. Chem.  
37 Phys., 12, 2641-2659, 10.5194/acp-12-2641-2012, 2012.

38 Jamali, S., Klingmyr, D., and Tagesson, T.: Global-Scale Patterns and Trends in Tropospheric NO2  
39 Concentrations, 2005–2018, Remote Sensing, 12, 3526, <https://doi.org/10.3390/rs12213526>, 2020.

- 1 Janssens-Maenhout, G., Dentener, F., Van Aardenne, J., Monni, S., Pagliari, V., Orlandini, L., Klimont,  
2 Z., Kurokawa, J.-i., Akimoto, H., and Ohara, T.: EDGAR-HTAP: a harmonized gridded air pollution  
3 emission dataset based on national inventories, European Commission Publications Office, Ispra,  
4 Italy, <https://data.europa.eu/doi/10.2788/14069>, 25229, 40, 2012.
- 5 Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Dentener, F., Bergamaschi,  
6 P., Pagliari, V., Olivier, J. G. J., Peters, J. A. H. W., van Aardenne, J. A., Monni, S., Doering, U., Petrescu,  
7 A. M. R., Solazzo, E., and Oreggioni, G. D.: EDGAR v4.3.2 Global Atlas of the three major greenhouse  
8 gas emissions for the period 1970–2012, *Earth Syst. Sci. Data*, 11, 959-1002, 10.5194/essd-11-959-  
9 2019, 2019.
- 10 Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Dentener, F., Muntean, M., Pouliot, G., Keating, T.,  
11 Zhang, Q., Kurokawa, J., Wankmüller, R., Denier van der Gon, H., Kuenen, J. J. P., Klimont, Z., Frost, G.,  
12 Darras, S., Koffi, B., and Li, M.: HTAP\_v2.2: a mosaic of regional and global emission grid maps for 2008  
13 and 2010 to study hemispheric transport of air pollution, *Atmos. Chem. Phys.*, 15, 11411-11432,  
14 10.5194/acp-15-11411-2015, 2015.
- 15 Johansson, L., Jalkanen, J.-P., and Kukkonen, J.: Global assessment of shipping emissions in 2015 on a  
16 high spatial and temporal resolution, *Atmospheric Environment*, 167, 403-415,  
17 <https://doi.org/10.1016/j.atmosenv.2017.08.042>, 2017.
- 18 Keita, S., Lioussé, C., Assamoi, E. M., Doumbia, T., N'Datchoh, E. T., Gnamien, S., Elguindi, N., Granier,  
19 C., and Yoboué, V.: African anthropogenic emissions inventory for gases and particles from 1990 to  
20 2015, *Earth Syst. Sci. Data*, 13, 3691-3705, 10.5194/essd-13-3691-2021, 2021.
- 21 Kuenen, J., Dellaert, S., Visschedijk, A., Jalkanen, J. P., Super, I., and Denier van der Gon, H.: CAMS-  
22 REG-v4: a state-of-the-art high-resolution European emission inventory for air quality modelling, *Earth  
23 Syst. Sci. Data*, 14, 491-515, 10.5194/essd-14-491-2022, 2022.
- 24 Kurokawa, J. and Ohara, T.: Long-term historical trends in air pollutant emissions in Asia: Regional  
25 Emission inventory in ASia (REAS) version 3, *Atmospheric Chemistry and Physics*, 20, 12761-12793,  
26 <https://doi.org/10.5194/acp-20-12761-2020>, 2020.
- 27 Liu, F., Choi, S., Li, C., Fioletov, V. E., McLinden, C. A., Joiner, J., Krotkov, N. A., Bian, H., Janssens-  
28 Maenhout, G., Darmenov, A. S., and da Silva, A. M.: A new global anthropogenic SO<sub>2</sub> emission  
29 inventory for the last decade: a mosaic of satellite-derived and bottom-up emissions, *Atmos. Chem.  
30 Phys.*, 18, 16571-16586, 10.5194/acp-18-16571-2018, 2018.
- 31 MEIC: [http://meicmodel.org.cn/?page\\_id=1772&lang=en](http://meicmodel.org.cn/?page_id=1772&lang=en), 2022.
- 32 NPRI: National Pollutant Release Inventory Summary Report 2016, Cat. No.: En81-14E-PDF ISSN: 2369-  
33 7733, [https://publications.gc.ca/collections/collection\\_2017/eccc/En81-14-2016-eng.pdf](https://publications.gc.ca/collections/collection_2017/eccc/En81-14-2016-eng.pdf), 2017.
- 34 Open Street Map: Open Street Map, available at: <https://www.openstreetmap.org/> (last access: 8  
35 March 2019), 2017.
- 36 OpenTransportMap: Open Transport Map, available at: <http://opentransportmap.info/> (last access: 8  
37 March 2019), 2017.
- 38 Planning Commission: Eleventh Five Year Plan 2007-12, [http://www.gov.cn/zhengce/content/2008-  
39 03/28/content\\_4877.htm](http://www.gov.cn/zhengce/content/2008-03/28/content_4877.htm), Volume-I. Inclusive Growth, 2008.



1 Platts: World Electric Power Plants database, available at:  
2 <https://www.spglobal.com/platts/en/commodities/electric-power> (last access: 21 February 2018),  
3 2017.

4 Puliafito, S. E., Bolaño-Ortiz, T. R., Fernandez, R. P., Berná, L. L., Pascual-Flores, R. M., Urquiza, J., López-  
5 Noreña, A. I., and Tames, M. F.: High-resolution seasonal and decadal inventory of anthropogenic gas-  
6 phase and particle emissions for Argentina, *Earth Syst. Sci. Data*, 13, 5027-5069, 10.5194/essd-13-  
7 5027-2021, 2021.

8 Sassi, M., Zhang, J., Moran, M.D.: 2015 SMOKE-Ready Canadian Air Pollutant Emission Inventory (APEI)  
9 Package, version 1 [Data set, Zenodo] <https://doi.org/10.5281/zenodo.4883639>, 2021.

10 Seltzer, K. M., Pennington, E., Rao, V., Murphy, B. N., Strum, M., Isaacs, K. K., and Pye, H. O. T.: Reactive  
11 organic carbon emissions from volatile chemical products, *Atmos. Chem. Phys.*, 21, 5079-5100,  
12 10.5194/acp-21-5079-2021, 2021.

13 Shephard, M. W., Dammers, E., Cady-Pereira, K. E., Kharol, S. K., Thompson, J., Gainariu-Matz, Y.,  
14 Zhang, J., McLinden, C. A., Kovachik, A., Moran, M., Bittman, S., Sioris, C. E., Griffin, D., Alvarado, M.  
15 J., Lonsdale, C., Savic-Jovcic, V., and Zheng, Q.: Ammonia measurements from space with the Cross-  
16 track Infrared Sounder: characteristics and applications, *Atmos. Chem. Phys.*, 20, 2277-2302,  
17 10.5194/acp-20-2277-2020, 2020.

18 Shibata, Y., and Morikawa, T.: Review of the JCAP/JATOP Air Quality Model Study in Japan,  
19 *Atmosphere*, 12(8), 943, <https://www.mdpi.com/2073-4433/12/8/943>, 2021.

20 Sun, W., Shao, M., Granier, C., Liu, Y., Ye, C. S., and Zheng, J. Y.: Long-Term Trends of Anthropogenic  
21 SO<sub>2</sub>, NO<sub>x</sub>, CO, and NMVOCs Emissions in China, *Earth's Future*, 6, 1112-1133,  
22 <https://doi.org/10.1029/2018EF000822>, 2018.

23 U.S. Environmental Protection Agency: Overview of EPA's MOtor Vehicle Emission Simulator  
24 (MOVES3), (2021) EPA-420-R-21-004, <https://nepis.epa.gov/Exe/ZyPDF.cgi?Dockkey=P1011KV2.pdf>  
25 (accessed 21 December 2021)., 2021a.


26 U.S. Environmental Protection Agency: 2017 National Emissions Inventory (NEI) Technical Support  
27 Document, [https://www.epa.gov/air-emissions-inventories/2017-national-emissions-inventory-nei-](https://www.epa.gov/air-emissions-inventories/2017-national-emissions-inventory-nei-data#doc)  
28 [data#doc](https://www.epa.gov/air-emissions-inventories/2017-national-emissions-inventory-nei-data#doc) (accessed 21 December 2021), 2021b.

29 UNC: Community Modeling and Analysis System CMAS, SMOKE v4.7 User's Man,  
30 <https://www.cmascenter.org/smoke/>, 2019.


31 UNECE: 1999 Protocol to Abate Acidification, Eutrophication and Ground-level Ozone to the  
32 Convention on Long-range Transboundary Air Pollution, as amended on 4 May 2012, available at:  
33 [https://www.unece.org/fileadmin/DAM/env/documents/2013/air/eb/ECE.EB.AIR.114\\_ENG.pdf](https://www.unece.org/fileadmin/DAM/env/documents/2013/air/eb/ECE.EB.AIR.114_ENG.pdf) (last  
34 access: 1 July 2021), 2012.

35 UNECE: Annual Report: Connecting countries, recovering better, driving progress, improving lives,  
36 Information Service, United Nations Economic Commission for Europe, Palais des Nations, CH-1211  
37 Geneva10, Switzerland, [https://unece.org/sites/default/files/2022-](https://unece.org/sites/default/files/2022-04/UNECE%20Annual%20Report%202021%20Web%20%281%29%200.pdf)  
38 [04/UNECE%20Annual%20Report 2021 Web%20%281%29 0.pdf](https://unece.org/sites/default/files/2022-04/UNECE%20Annual%20Report%202021%20Web%20%281%29%200.pdf), 2021.

39 Yin, Y., Chevallier, F., Ciais, P., Broquet, G., Fortems-Cheiney, A., Pison, I., and Saunois, M.: Decadal  
40 trends in global CO emissions as seen by MOPITT, *Atmos. Chem. Phys.*, 15, 13433-13451, 10.5194/acp-  
41 15-13433-2015, 2015.



1




2

3 **Figure 1 – Overview of the HTAP\_v3 mosaic data providers. Data from officially**  
 4 **reported emission gridmaps were collected from the US Environmental Protection**  
 5 **Agency, Environment and Climate Change Canada, CAMS-REG-v5.1 for Europe,**  
 6 **REASv3.2.1 for most of the Asian domain, CAPSS-KU for Korea and JAPAN (PM2.5EI**  
 7 **and J-STREAM) for Japan. The share of the total emissions covered by each data**  
 8 **provider is reported in the bar chart at the bottom.**

9

10





1

2 **Figure 2 – Time series of gaseous and particulate matter pollutants from HTAP\_v3 by**  
 3 **aggregated regions. Regional grouping follows the Intergovernmental Panel on Climate**  
 4 **Change Sixth Assessment Report (IPCC AR6) definitions. Table S3 provides information**  
 5 **on the regional belonging of each country to the IPCC AR6 regions.**

6


7

8



3 **Figure 3 - Sectoral (panel a) and regional (panel b) breakdown of air pollutant emissions**  
 4 **from HTAP\_v3 for the year 2018. At the top of each bar in panel a, total emissions for**  
 5 **each pollutant are reported (in Mt).**






1

2 **Figure 4 – HTAP\_v3 mosaic: SO<sub>2</sub> emission gridmaps for the year 2018.**


3

4










1

2 **Figure 5 – HTAP\_v3 mosaic: NO<sub>x</sub> emission gridmaps in 2000 (top panel) and 2018**  
 3 **(bottom panel).**


1



2


3 **Figure 6 – HTAP\_v3 mosaic: PM<sub>2.5</sub> emissions from residential activities in January 2018.**






1

2 **Figure 7 – HTAP\_v3 mosaic: PM<sub>10</sub> emissions from agricultural waste burning in January**  
 3 **2018.**




1  
2  
3  
4

**Figure 8 – HTAP\_v3 mosaic: NH<sub>3</sub> emissions from agricultural soils activities in January 2018.**



1

2 **Figure 9 – Monthly variability of NH<sub>3</sub> emissions for agriculture related activities for the**  
 3 **different world regions in 2015.**




4

5 **Figure 10 – Monthly variability of NO<sub>x</sub> emissions for relevant emission sectors for the**  
 6 **different world regions in 2015.**

7




1



2


3 **Figure 11 – PM<sub>2.5</sub> monthly emission maps from the residential sector in 2018 from**  
4 **HTAP\_v3.**

5



6

7 **Figure 12 – PM<sub>10</sub> monthly emission maps from agricultural residues burning in 2018 from**  
8 **HTAP\_v3.**







1  
2  
3  
4  
5  
6  
7

**Figure 13 – NH<sub>3</sub> monthly emission maps from agricultural soils in 2018 from HTAP\_v3.**




**Table 1 – Overview of data input to the HTAP\_v3 emission mosaic.**

Data source	CAMS-REG-v5.1	US EPA	ECCEC	REASv3.2.1
Type of data source	Country inventories as emission time series by sector and country and emission gridmaps as .csv files.	Country inventory	Country inventory as emission time series by sector and country and emission gridmaps as .NetCDF files.	Country inventories as emission gridmaps as text files.
Sectors coverage	All sectors, excluding international shipping and aviation (international and domestic).	All sectors, excluding international shipping and aviation (international and domestic).	All sectors, excluding agricultural waste burning, international shipping and aviation (international and domestic).	All sectors, excluding brake and tyre wear, domestic shipping, waste, agricultural waste burning, international shipping and aviation (international and domestic).

<b>Temporal coverage</b>	2000-2018	2002-2017	2000-2016	2000-2015+trends from MEIC over China for 2016, 2017, 2018
<b>Temporal resolution</b>	Annual emission gridmaps + monthly profiles	Monthly emission gridmaps	Monthly emission gridmaps	Monthly emission gridmaps
<b>Spatial resolution</b>	0.1°x0.1°	0.1°x0.1°	0.1°x0.1°	0.1°x0.1° (The original spatial resolution of REASv3.2.1 is 0.25°x0.25°. Assuming that emissions are equally distributed in the 0.25° cell, REASv3.2.1 data were converted to 0.1° cell and provided to HTAP v3)
<b>Substances</b>	SO <sub>2</sub> , NO <sub>x</sub> , CO, NMVOC, NH <sub>3</sub> , PM <sub>10</sub> , PM <sub>2.5</sub> , BC, OC	SO <sub>2</sub> , NO <sub>x</sub> , CO, NMVOC, NH <sub>3</sub> , PM <sub>10</sub> , PM <sub>2.5</sub> , BC, OC	SO <sub>2</sub> , NO <sub>x</sub> , CO, NMVOC, NH <sub>3</sub> , PM <sub>10</sub> , PM <sub>2.5</sub> , BC, OC	SO <sub>2</sub> , NO <sub>x</sub> , CO, NMVOC, NH <sub>3</sub> , PM <sub>10</sub> , PM <sub>2.5</sub> , BC, OC
<b>Geocoverage</b>				
<b>References</b>				<a href="http://meicmodel.org">http://meicmodel.org</a>

1

<b>Data source</b>	<b>CAPSS-KU</b>	<b>JAPAN (PM2.5EI and J-STREAM)</b>	<b>EDGARv6.1</b>
<b>Type of data source</b>	Country inventory as emission time series by sector and country and emission gridmaps as .NetCDF files.	Country inventory as emission time series by sector and country and emission gridmaps as .NetCDF files.	Country inventory as emission time series by sector and country and emission gridmaps as .NetCDF files.
<b>Sectors coverage</b>	All sectors, excluding international shipping and aviation (international and domestic).	All sectors, excluding international shipping, domestic shipping and aviation	All sectors, including international shipping and aviation (international and domestic)

		(international and domestic).	
<b>Temporal coverage</b>	2000-2018	2000-2017	2000-2018
<b>Temporal resolution</b>	Annual emission gridmaps + monthly profiles	Monthly emission gridmaps	Monthly emission gridmaps
<b>Spatial resolution</b>	0.1°x0.1°	0.1°x0.1°	0.1°x0.1°
<b>Substances</b>	SO <sub>2</sub> , NO <sub>x</sub> , CO, NMVOC, NH <sub>3</sub> , PM <sub>10</sub> , PM <sub>2.5</sub> , BC, OC	SO <sub>2</sub> , NO <sub>x</sub> , CO, NMVOC, NH <sub>3</sub> , PM <sub>10</sub> , PM <sub>2.5</sub> , BC, OC	SO <sub>2</sub> , NO <sub>x</sub> , CO, NMVOC, NH <sub>3</sub> , PM <sub>10</sub> , PM <sub>2.5</sub> , BC, OC
<b>Geocoverage</b>			
<b>References</b>			<a href="https://edgar.jrc.ec.europa.eu/dataset_ap61">https://edgar.jrc.ec.europa.eu/dataset_ap61</a>

1 Table 2 – Definition of HTAP\_v3 sectors and correspondence to IPCC codes.

HTAP_v3 main sectors	HTAP_v3 detailed sectors	Sector description	IPCC 1996 codes	IPCC 2006 codes
HTAP_1 : International Shipping	HTAP_1: International Shipping	International water-borne navigation.	1C2	1.A.3.d.i
HTAP_2 : Aviation	HTAP_2.1: Domestic Aviation	Civil Aviation.	1A3aii	1.A.3.a.ii
	HTAP_2.2: International Aviation	International Aviation.	1A3ai	1.A.3.a.i
HTAP_3 : Energy	HTAP_3: Energy	Power generation.	1A1a	1.A.1.a
HTAP_4 : Industry	HTAP_4.1: Industry	Industrial non-power large-scale combustion emissions and emissions of industrial processes. It includes: manufacturing, mining, metal, cement, chemical and fossil fuel fires.	1A2 + 2 + 5B	1A2 + 2 (excluding 2.D.3 + 2.E + 2.F + 2.G) + 7A
	HTAP_4.2: Fugitive	It includes oil and gas exploration and production and transmission, including evaporative emissions (mainly NMVOC).	1B + 1A1b + 1A1ci + 1A1cii + 1A5biii	1.B + 1.A.1.b + 1.A.1.c.i + 1.A.1.c.i.i + 1.A.5.b.i.i.i
	HTAP_4.3: Solvents	Solvents and product use.	3	2D3 + 2E + 2F + 2G
HTAP_5 : Ground Transport	HTAP_5.1: Road Transport	Road Transport, combustion and evaporative emissions only.	1A3b (excluding resuspension)	1.A.3.b (excluding resuspension)
	HTAP_5.2: Brake and Tyre wear	Re-suspended dust from pavements or tyre and brake wear from road transport.	1A3b (resuspension only)	1.A.3.b (resuspension only)
	HTAP_5.3: Domestic shipping	Domestic shipping: inland waterways + domestic shipping.	1A3d2	1.A.3.d.ii
	HTAP_5.4: Other ground transport	Ground transport by pipelines and other ground transport of mobile machinery.	1A3c + 1A3e	1.A.3.c + 1.A.3.e.ii
HTAP_6 : Residential	HTAP_6: Residential	Small-scale combustion, including heating, cooling, lighting, cooking and auxiliary engines, to equip residential,	1A4 + 1A5	1.A.4 + 1.A.5



		commercial buildings, service institutes, and agricultural facilities and fisheries.		
<b>HTAP_7 : Waste</b>	<b>HTAP_7: Waste</b>	Solid waste disposal and wastewater treatment.	6	4
<b>HTAP_8 : Agriculture</b>	<b>HTAP_8.1: Agricultural waste burning</b>	Agricultural waste burning (excluding Savannah burning).	4F	3.C.1.b
	<b>HTAP_8.2: Agriculture livestock</b>	Livestock emissions, including manure management.	4B	3.A.2
	<b>HTAP_8.3: Agriculture crops</b>	Emissions from crops, fertilisers, and all agricultural soils activities.	4C + 4D	3.C.2 + 3.C.3 + 3.C.4 + 3.C.7

1

2

3

4 **Table 3 – Overview of pollutant and sector provided by each inventory in HTAP\_v3. Cells**  
5 **with N/A indicate that the emissions for those sectors were not provided and/or used in**  
6 **HTAP\_v3 for a specific inventory, while gapfilled with the corresponding information**  
7 **from EDGARv6.1. The other cells represent the data availability for each sector and**  
8 **inventory. The color codes used for the pollutants refer to the data source: black color**  
9 **represents pollutant emissions provided by a specific inventory, red color indicates**  
10 **emissions gapfilled using EDGARv6.1 and violet color indicates combinations of sectors-**  
11 **pollutants available for specific regional inventories but not in EDGAR, which typically**  
12 **represent minor sources of emissions included in officially reported inventories. These**  
13 **minor sources are included in the HTAP\_v3 mosaic.**

14

<b>Data provider</b>	<b>REA Sv3.2 .1</b>	<b>CAP SS-KU</b>	<b>JAPAN</b>	<b>ECCE</b>	<b>US EPA</b>	<b>CAMS-REG-v5.1</b>	<b>EDGAR v6.1</b>
<b>HTAP_1: International Shipping</b>	N/A	N/A	N/A	N/A	N/A	N/A	All substances
<b>HTAP_2.1: Domestic Aviation</b>	N/A	N/A	N/A	N/A	N/A	N/A	All substances
<b>HTAP_2.2: International Aviation</b>	N/A	N/A	N/A	N/A	N/A	N/A	All substances
<b>HTAP_3: Energy</b>	All substances	All substances	BC, OC, NOx, NH3,	All substances	All substances	All substances	All substances

			CO, PM2.5, PM10, NMVO C, SO2				
<b>HTAP_4.1: Industry</b>	All subst ances	All subst ances	BC, OC, NOx, NH3, CO, PM2.5, PM10, NMVO C, SO2	All substanc es	All substanc es	All substanc es	All substanc es
<b>HTAP_4.2: Fugitive</b>	All subst ances	BC, OC, NOx, NH3, CO, PM2. 5, PM10 , NMV OC, SO2	BC, OC, NOx, NH3, CO, PM2.5, PM10, NMVO C, SO2	All substanc es	All substanc es	All substanc es	All substanc es
<b>HTAP_4.3: Solvents</b>	NMV OC, NH3, PM10 , PM2. 5	NMV OC, NH3, PM10 , PM2. 5	NMVO C, NH3, PM10, PM2.5	NMVO C, NH3, PM10, PM2.5	CO, NOx, OC, NMVOC , NH3, PM10, PM2.5, SO2	NOx, NH3, CO, PM2.5, PM10, NMVO C, SO2	All substanc es
<b>HTAP_5.1: Road Transport</b>	All subst ances	All subst ances	All substanc es	All substanc es	All substanc es	All substanc es	All substanc es
<b>HTAP_5.2: Brake and Tyre wear</b>	N/A	BC, OC, NOx, NH3, CO, PM2. 5, PM10 , NMV OC, SO2	BC,OC, PM2.5,P M10	BC,OC, PM2.5,P M10	BC,OC, PM2.5,P M10, NMVOC	BC,OC, PM2.5,P M10	BC,OC, PM2.5,P M10

<b>HTAP_5.3: Domestic shipping</b>	N/A	All substances	N/A	All substances	All substances	BC, OC, NOx, NH3, CO, PM2.5, PM10, NMVOC, SO2	All substances
<b>HTAP_5.4: Other ground transport</b>	All substances	All substances	BC, OC, NOx, NH3, CO, PM2.5, PM10, NMVOC, SO2	All substances	All substances	All substances	All substances
<b>HTAP_6: Residential</b>	All substances	All substances	BC, OC, NOx, NH3, CO, PM2.5, PM10, NMVOC, SO2	All substances	All substances	All substances	All substances
<b>HTAP_7: Waste</b>	N/A	All substances	All substances	All substances	All substances	All substances	All substances
<b>HTAP_8.1: Agricultural waste burning</b>	N/A	All substances	All substances	N/A	All substances	All substances	All substances
<b>HTAP_8.2: Agriculture livestock</b>	NH3, NMVOC, NOx, PM10, PM2.5	NH3, NMVOC, NOx, PM10, PM2.5, BC, OC	NH3, NMVOC, NOx, PM10, PM2.5	NH3, NMVOC, NOx, PM10, PM2.5, BC, OC	NH3, NMVOC, NOx, PM10, PM2.5, BC, OC	NH3, NMVOC, NOx, PM10, PM2.5, OC	NH3, NMVOC, NOx, PM10, PM2.5
<b>HTAP_8.3: Agriculture crops</b>	NH3, NOx, PM10, PM2.5	NH3, NOx, PM10, PM2.5, BC, OC	NH3, NOx, PM10, PM2.5	NH3, NOx, PM10, PM2.5, BC, NMVOC, OC	NH3, NOx, PM10, PM2.5, BC, CO, NMVOC, SO2	NH3, NOx, PM10, PM2.5, SO2, CO, OC, NMVOC	NH3, NOx, PM10, PM2.5

1 Table 4 – Main features of the different HTAP mosaics.

	<b>HTAP v1</b>	<b>HTAP v2.2</b>	<b>HTAP v3</b>
<b>Time coverage</b>	2000-2005	2008 and 2010	2000-2018
<b>Time resolution</b>	yearly	yearly and monthly	yearly and monthly
<b>Substances</b>	CH <sub>4</sub> , NMVOC, CO, SO <sub>2</sub> , NO <sub>x</sub> , NH <sub>3</sub> , PM <sub>10</sub> , PM <sub>2.5</sub> , BC, OC	SO <sub>2</sub> , NO <sub>x</sub> , CO, NMVOC, NH <sub>3</sub> (only for agriculture), PM <sub>10</sub> , PM <sub>2.5</sub> , BC, OC	SO <sub>2</sub> , NO <sub>x</sub> , CO, NMVOC, NH <sub>3</sub> , PM <sub>10</sub> , PM <sub>2.5</sub> , BC, OC
<b>Sectors</b>	Aircraft, Ships, Energy, Industry Processes, Ground Transport, Residential, Solvents, Agriculture, Agriculture Waste Burning, and Waste	Air, Ships, Energy, Industry, Transport, Residential (including waste), and Agriculture (only for NH <sub>3</sub> )	International Shipping, Domestic Shipping, Domestic Aviation, International Aviation, Energy, Industry, Fugitives, Solvent Use, Road Transport, Brake and Tyre Wear, Other Ground Transport, Residential, Waste, Agricultural Waste Burning, Livestock, and Agricultural Crops
<b>Geographical coverage</b>	Globe	Globe	Globe
<b>Spatial resolution</b>	0.1°x0.1°	0.1°x0.1°	0.1°x0.1°
<b>Input datasets</b>	UNFCCC, REAS, GAINS, EMEP, EPA, EDGARv4.1	US EPA, Environment Canada, MICS, TNO/EMEP Europe (MACC II), MICS Asia III+, REAS2.1, EDGARv4.3	CAMS-REG-v5.1, REASv3.2.1, US EPA, ECCC, CAPSS-KU, JAPAN (PM2.5EI and J-STREAM), EDGARv6.1

<b>Reference</b>	Janssens-Maenhout et al., 2012	Janssens-Maenhout et al., 2015	This work
------------------	--------------------------------	--------------------------------	-----------