

Multitemporal glacier inventory revealing four decades of glacier changes in the Ladakh region

Mohd Soheb¹, Alagappan Ramanathan¹, Anshuman Bhardwaj², Millie Coleman^{2,3}, Brice R. Rea², Matteo Spagnolo², Shaktiman Singh², Lydia Sam²

¹ School of Environmental Sciences, Jawaharlal Nehru University, India.

² School of Geosciences, University of Aberdeen, United Kingdom.

³ School of Natural and Built Environment, Queen's University Belfast, United Kingdom.

Correspondence to: Mohd Soheb (sohaib.achaa@gmail.com)

Abstract. Multi-temporal inventories of glacierised regions provide an improved understanding of water resource availability. In this study, we present a Landsat-based multi-temporal inventory of glaciers in four Upper Indus sub-basins and three internal drainage basins in the Ladakh region for the years 1977, 1994, 2009 and 2019. The study records data on 2257 glaciers (of individual size $>0.5 \text{ km}^2$) covering an area of $\sim 7923 \pm 106 \text{ km}^2$ which is equivalent to $\sim 30\%$ of the total glacier population and $\sim 89\%$ of the total glacierised area of the region. Glacier area ranged between 0.5 ± 0.02 and $862 \pm 16 \text{ km}^2$, while glacier length ranged between 0.4 ± 0.02 and $73 \pm 0.54 \text{ km}$. Shayok Basin has the largest glacierised area and glacier population, while Tsokar has the least. Results show that the highest concentration of glaciers is found in the higher elevation zones, between 5000 and 6000 m a.s.l, with most of the glaciers facing towards the NW-NE quadrant. The error assessment shows that the uncertainty, based on the buffer-based approach, ranges between 2.6 and 5.1% for glacier area, and 1.5 and 2.6% for glacier length with a mean uncertainty of 3.2 and 1.8%, respectively. This multitemporal inventory is in good agreement with previous studies undertaken in parts of the Ladakh region. The new glacier database for the Ladakh region will be valuable for policy-making bodies, and future glaciological and hydrological studies. The data can be viewed and downloaded from PANGAEA, <https://doi.org/10.1594/PANGAEA.940994> (Soheb et al., 2022).

1. Introduction:

The Himalaya is the largest storehouse of snow and ice outside the Polar Regions. This large reserve of water plays a crucial role in the hydro-economy of the region (Bolch, 2019; Frey et al., 2014; Maurer et al., 2019; Pritchard, 2019). Any change to the Himalayan cryosphere would have a direct impact on the hydrology, further influencing the communities downstream whose livelihood and economy relies on, and are supported by, the major river systems e.g., the Brahmaputra, Ganges and Indus, among others. In high altitude arid regions like Ladakh, where the majority of glaciers are small and restricted to higher altitudes, meltwater serves as an important driver of the economy, especially in years with low winter precipitation when glacier melt becomes the major (or only) source of water (Schmidt and

Nüsser, 2012, 2017). Recent studies have reported that Himalayan glaciers are retreating at an alarming rate (Azam et al., 2021; Bolch, 2019; Kääb et al., 2015; Maurer et al., 2019; Pritchard, 2019; Shean et al., 2020, among others) with glaciers of the Western Himalayas showing less shrinkage than the glaciers of the central and eastern parts (Azam et al., 2021; Shukla et al., 2020; Singh et al., 2016). Glaciers in the nearby Karakoram region display long-term irregular behaviour with frequent glacier advances/surges and minimal shrinkage, which is yet to be fully understood (Azam et al., 2021; Bhambri et al., 2013; Bolch et al., 2012; Kulkarni, 2010; Liu et al., 2006; Minora et al., 2013; Negi et al., 2021). Glaciers of the Karakoram region experienced an increase in area post-2000, due to surge-type glaciers. In just the upper Shayok valley, as many as 18 glaciers, occupying more than one-third of the glacierised area, showed surge-type behaviour (Bhambri et al., 2011, 2013; Negi et al., 2021). However, not all regions of Ladakh have been analysed at the same level of spatio-temporal detail. In particular, our knowledge of glacier dynamics and their response to climate change is still incomplete in the cold-arid, high-altitude Ladakh region (~105,476 km²) comprising both, the Himalayan and Karakoram ranges. Few studies have focused on the glaciers of this region (e.g. Bhambri et al., 2011, 2013; Chudley et al., 2017; Negi et al., 2021; Nüsser et al., 2012; Schmidt & Nüsser, 2012, 2017; Shukla et al., 2020).

The advent of remote sensing technologies has permitted the mapping and measuring of various glacier attributes even in the absence of sufficient in-situ observations (Bhardwaj et al., 2015). Glacierised area estimations have often relied on global and regional glacier inventories such as the Randolph Glacier Inventory (RGI), Global Land Ice Measurements from Space (GLIMS), Geological Survey of India (GSI) inventory and Space Application Centre India (SAC) inventory, among others (Chinese Glacier Inventory (CGI), Glacier Area Mapping for Discharge from the Asian Mountains (GAMDAM), International Centre for Integrated Mountain Development (ICIMOD)). However, given the large scale of these inventories, automated techniques are employed, in most of the cases, to map and calculate glacier extent with differing levels of success. Additionally, the varying quality of satellite imagery acquired from different time periods are sometimes necessitated in high mountain areas, such as Ladakh. Together, these two factors can lead to over- or under-estimation of glacier areas leading to erroneous information on temporal change. Moreover, there is no multi-temporal glacier inventory available for the entire Ladakh region, which can inform us on the changes in the natural frozen water reserves which have put the water security of this entire cold-arid region under significant stress during recent years. The residents of Ladakh have witnessed a decrease in agricultural yields, the main driver of economic development of the region, due to a decrease in water resources (Barrett and Bosak, 2018). The water scarcity together with an increase in tourism footprint (four times more tourists (327,366) in 2018 than 2010, a number that is more than the entire population of Ladakh) has led to a shift in livelihood from agriculture to other commercial activities (Müller et al., 2020), though even the latter relies heavily on water resources. In order to cope with water scarcity, some people of Ladakh have developed new water management techniques, commonly known as ‘ice reservoirs’ or ‘ice stupas’, to supplement agricultural activities (Nüsser, et al., 2019a,b).

This study presents a new multi-temporal glacier inventory for the Union Territory of Ladakh, India, covering 42 years of change between 1977 and 2019. This new dataset and analyses of glacier distribution will help to improve understanding of the glacier dynamics and the impact of ongoing climate change on water resources in the Ladakh region, where glaciers are the only source of water in the dry season. The inventories are entirely based on Landsat images acquired mostly during late-summer with additional quality control provided through high-resolution

PlanetScope and Google Earth imagery. We further establish a comparison with the existing inventories and data available in recent studies from the region. The dataset produced in this study can be viewed and downloaded from: *PANGAEA*, <https://doi.org/10.1594/PANGAEA.940994> (Soheb et al., 2022)

2. Study Area:

This study focuses on glaciers in the Upper Indus Basin (UIB) upstream of Skardu and three internal drainage/endorheic basins (IDBs) within Ladakh, namely Tsokar, Tsomoriri and Pangong Basins. The geographic extent of the study area lies within a latitude of 31.1° to 35.6° N and a longitude of 75.1° to 81.8° E and covers a vast region of the Karakoram and Western Himalayan ranges. UIB has an area of ~105,476 km², of which ~8302 km² (8%) is glacierised by ~6300 glaciers spanning elevations between ~3400 m and ~7500 m a.s.l. (as per RGI 6.0). IDBs of Tsokar (1036 km²), Tsomoriri (5462 km²) and Pangong (21,206 km²) house ~30, 345 and 812 glaciers, comprising a glacierised area of ~7 (0.6%), 185 (3.4%) and 437 (2.1%) km², respectively (as per RGI 6.0). The glaciers of IDBs are at a comparatively higher elevation, spanning from ~4800 to ~6800 m a.s.l. Meltwater from these glaciers drains into the lakes within each basin. Pangong Lake (a saline lake), situated at an elevation of ~4241 m a.s.l., is the largest with an area of ~703 km². Both Tsomoriri (freshwater lake at ~4522 m a.s.l.) and Tsokar (saline lake at ~4531 m a.s.l.) Lakes are designated Ramsar sites which occupy areas of ~140 and ~15 km², respectively. Since the majority of the investigated area (UIB and IDBs combined) falls within Ladakh, the combined area of UIB and IDBs will be referred to as “Ladakh region” hereafter.




Figure 1: Location map of the study area: the boundaries of studied Upper Indus Basin and internal drainage basins are outlined in black and red on the digital elevation model (DEM) and in the inset map. Inset map shows the study area with respect to the Himalayan and Karakoram region. Black dots and stars represent the respective basins' major settlements and field investigated glaciers. The background image (ASTER GDEM) courtesy: NASA/METI/AIST/Japan Spacesystems and U.S./Japan ASTER Science Team, <https://earthdata.nasa.gov/>.

The Ladakh region has a cold-arid climate due to the rain shadow and elevation effects of the Himalaya and Karakoram mountains (Schmidt and Nüsser, 2017). Mean annual air temperature and annual precipitation range between 0 to 10 °C and 20 to 145 mm, respectively (Hersbach et al., 2020; Figure 2). This region is inhabited by ~700,000 people (as per Census of India 2011, Census of China 2020), most of which are directly, or indirectly, dependent on snow and glacier meltwater to support hydropower generation, irrigation and domestic needs.




Figure 2: Mean annual (a, c) and monthly (b, d) temperature and precipitation at Shiquanhe and Leh stations.

3. Data and methods

3.1. Data

This study utilises multiple Landsat level-1 precision and terrain (L1TP) corrected scenes (63 scenes in total) from four different periods: 1977 \pm 5 (hereafter 1977), 1994 \pm 1 (hereafter 1994), 2009 and 2019 \pm 1 (hereafter 2019). Scenes from the 1970s are majorly (12 out of 17) from the year 1976 and 1977 however due to higher cloud cover and less availability of imagery during the earlier Landsat period, five scenes from 1972, 1979 and 1980 were also included to aid the digitization of glaciers (Table S1). Images from the late in the ablation season (July-October), having least snow and cloud cover (<30% overall, and not over the glacierised parts), were selected and used for glacier identification and boundary delineation. Advanced Space borne Thermal Emission and Reflection Radiometer Global Digital Elevation Model (ASTER GDEM) scenes were also used for basin delineation and calculating slope, aspect and elevation metrics of the glaciers. Glacier digitisation, basin delineation and calculation of area were all performed in ArcGIS 10.4. Details of the imagery used in this study are presented in (Table 1 and Table S1).

Table 1: Information on the satellite imagery used in this study (Detailed info. in Table S1).

Dataset	Year of Acquisition	Spatial Resolution	No. of image used	Source	Purpose
Landsat MSS	1977 \pm 5	60m	17	https://earthexplorer.usgs.gov/	Glacier area mapping
Landsat TM	1994 \pm 1, 2009	30m	14, 18		

Landsat OLI	2019±1	15m	14		
ASTER GDEM	2000-2013	30m	17	https://earthdata.nasa.gov/	Topography and basin delineation

3.2. Basin delineation

Basin delineation was carried using ASTER GDEM V003 and the Hydrology tool in ArcGIS. The input DEM was first analysed to fill-in all sinks with careful consideration of the potential for basin area over-estimation (Khan et al., 2014). UIB was delineated using a pour point selected at the Indus River in Skardu as we aimed to assess all the tributary basins of the Ladakh region. The UIB obtained by this approach was further divided into second-order tributary basins, i.e., Shayok, Suru, Zaskar and Leh Basins. A small portion of the leftover area from UIB after second-order tributary basin delineation was merged into the Leh Basin in order to investigate the UIB upstream of Skardu. Delineation of the three endorheic basins (IDBs) that lie partially or completely in the Ladakh region, i.e., Tsokar, Tsomoriri and Pangong Basins, was also carried out using the same method with the help of respective lakes as a pour point. The digitisation of the three lakes (Tsokar, Tsomoriri and Pangong Lake) was carried out manually for the years 1977, 1994, 2009 and 2019 using Landsat imagery.

3.3. Glacier mapping

Glaciers were mapped using a two-way approach, closely following the Global Land Ice Measurements from Space (GLIMS) guidelines (Paul et al., 2009): 1) automatic mapping of the clean glacier and 2) manually correcting the glacier outlines and digitisation of debris cover. First, a band ratio approach between NIR (Near Infrared) and SWIR (Shortwave Infrared) (as suggested by Paul et al., 2002, 2015; Racoviteanu et al., 2009; Bhardwaj et al 2015; Schmidt & Nüsser, 2017; Smith et al., 2015; Winsvold et al., 2014, 2016) with a threshold of 2.0 ($NIR/SWIR > 2 = \text{ice/snow}$) was used on 2019 Landsat OLI images to delineate the clean part of glaciers. A median filter of kernel size 3 x 3 was applied to remove the isolated and small pixels outside the glacier area. The NIR and SWIR band ratio approach is good at distinguishing glacier pixels from water features with similar spectral reflectance values (Racoviteanu et al., 2009; Zhang et al., 2019). This approach failed in areas with high snow/cloud cover, shadows, frozen channels/lakes and debris cover. The snow/cloud cover and frozen lakes/stream problem were addressed by selecting Landsat scenes from the ablation period (July-October) with the cloud cover < 30%. The issue with the snow-covered regions in accumulation zones, where the delineation was the most challenging, was resolved using the best available imagery of any time between 1977 and 2019 because glaciers are not expected to change their shape significantly in the higher accumulation zones. One of the major issues was the debris covered glaciers, which had to be manually digitised, with the support of high-resolution Google Earth and PlanetScope imagery from 2019 ±2. The result was then used as a basis for manual digitisation of debris covered glaciers in other years where high-resolution images are not available. In most cases, identification of the glacier terminus was made with certain contextual characteristics at the snout, e.g., the emergence of meltwater streams, proglacial lakes, ice walls, end moraines etc. (Figure S1).

The glacier outlines from 2019 were used as a starting point for the subsequent digitization of glacier areas in 2009, 1994 and 1977. Glacier length was measured using a semi-automatic approach, by employing the DEM to identify a

central flow line for each mapped glacier (Ji et al., 2017; Le Bris and Paul, 2013). Further manual corrections were undertaken to account for the flow lines of glaciers that have multiple tributaries and multiple highest/lowest points. Furthermore, some mapping errors are still expected to be present in this inventory due to a possible misinterpretation of glacier features, and the quantification of such errors are difficult owing to the lack of reliable reference in-situ data in the Ladakh region. Such errors were minimized by keeping a fixed map-scale of 1:10,000 in most cases, and undertaking a quality check on glacier outlines using high-resolution images. In case of MSS images and smaller glaciers, a map-scale of 1:25,000 was also used whenever required.

Other specific glacier attributes were also extracted including new glacier Ids, Global Land Ice Measurements from Space (GLIMS)-Ids, Randolph Glacier Inventory (RGI 6.0)-Ids, coordinates (latitude and longitude), elevation (maximum, mean and minimum), aspect (mean), slope (mean), area, length (maximum), area uncertainty and length uncertainty.

3.4. Uncertainty

This study involves the use of satellite imagery to extract various glacier parameters. It is therefore subject to uncertainties which may arise mainly from four different sources: (1) the quality of the image (with potential issues due to seasonal snow, shadows and cloud cover), (2) sensor characteristics (spatial/spectral resolution), (3) interpretation of glacial features and methodology used, and (4) post-processing techniques (Le Bris and Paul, 2013; Paul et al., 2013, 2017; Racoviteanu et al., 2009, 2019). Error due to sources 1, 3, and 4 are generally minor and can be visually identified and corrected (section 3.3), but an exact quantification is difficult due to the lack of reference data available from the region (Racoviteanu et al., 2009; Shukla et al., 2020). Type 4 errors are significant and have an impact on both glacier area and length estimation. Therefore, we applied a buffer-based assessment to glacier areas with the buffer width set to one-pixel for debris covered and a half-pixel for clean ice (Bolch et al., 2010; Granshaw & Fountain, 2006; Mölg et al., 2018; Paul et al., 2017; Racoviteanu et al., 2009; Shukla et al., 2020; Tielidze & Wheate, 2018), given that the level 1TP Landsat images were corrected to sub-pixel geometric accuracy (Bhambri et al., 2013). A buffer-based method provides the maximum and minimum estimates of uncertainty with respect to glacier size, where the values vary with size of the glacier and spatial resolution of the imagery used. Thus, it is more specific to the dataset and most recommended when there is no reliable reference data available (Paul et al., 2017; Racoviteanu et al., 2009; Shukla et al., 2020). The same approach was also followed to estimate the uncertainties in lake areas with one-pixel as the buffer width.

The associated uncertainty for smaller glaciers ($<0.5 \text{ km}^2$) amounts to ~12-25%. Therefore, all the glaciers with an area of less than 0.5 km^2 , which comprise ~70% and ~10% of the total glacier count and glacierised area respectively, are not included in this study. For the remaining glaciers, the uncertainty in glacier area ranged between ± 2.1 and $\pm 7.2\%$ depending on the spatial resolution of the satellite imagery and the individual glacier size. The highest uncertainty was for the year 1977 due to the coarser spatial resolution of Landsat MSS data when applied to the smallest glaciers ($0.5\text{-}1 \text{ km}^2$). For most of the glaciers, lengths are assumed to be accurate to ± 1 pixel at the terminus (Le Bris and Paul, 2013). Therefore, a buffer of one-pixel was set to determine the uncertainty in glacier length. The

length uncertainty ranged between ± 1.5 and $\pm 2.6\%$ with maximum uncertainty observed for the smallest glacier category ($0.5\text{--}1\text{ km}^2$). The methods yielded an overall uncertainty of 4.2, 1.8 and 1.5% for glacier area, glacier length and lake area, respectively (Table S2).

Uncertainties related to other attributes (mean elevation, mean slope and mean aspect) of the inventory are difficult to estimate due to the use of the ASTER GDEM product in this study, which was developed using a collage of archived scenes acquired between 2000 and 2013. In addition, the local undulations and surface change over time will have only marginal effects on parameters (elevation, slope and aspect) that are averaged over the entire glacier as averaging compensates for most of the changes (Frey and Paul, 2012). However, for parameters like maximum and minimum elevations, where one cell is used and no averaging is applied, the uncertainty is $\sim \pm 9\text{m}$, as the vertical accuracy of ASTER GDEM is $\pm 8.55\text{m}$ for glacierised areas of high Asia (Yao et al., 2020) and $\pm 8.86\text{m}$ elsewhere (Mukherjee et al., 2013).

4. Results

4.1. General statistics

In total, 2257 glaciers ($>0.5\text{ km}^2$) were compiled in the current inventory (Table 2), with a total glacierised area of $\sim 8511 \pm 430$, 8173 ± 215 , 8096 ± 214 and $7923 \pm 106\text{ km}^2$ for the years 1977, 1994, 2009 and 2019, respectively. The glacierised area corresponds to $\sim 6\%$ of the Ladakh region with individual areas ranging between 0.5 ± 0.02 and $862 \pm 16\text{ km}^2$. Glacier length in the Ladakh region varies between 0.4 ± 0.02 and $73 \pm 0.54\text{ km}$ with a mean length of $2.9 \pm 0.05\text{ km}$. About 90% of the glaciers are shorter than 5km in length while only 6% of glaciers have a length of $< 1\text{ km}$. Larger glaciers are mainly located in the Shayok and Zaskar Basins with the Siachen Glacier being the largest ($862 \pm 16\text{ km}^2$), longest ($73 \pm 0.54\text{ km}$) and covers the greatest elevation range of $\sim 3616\text{m}$ ($3702\text{--}7318\text{m a.s.l.}$). The major lakes in each endorheic basins of Pangong, Tsokar and Tsomoriri occupy an area of 3, 2 and 2.5%, respectively. The lake areas for the year 1977, 1994, 2009 and 2019 were 610 ± 14 , 619 ± 8 , 669 ± 8 and $705 \pm 8\text{ km}^2$ for Pangong, 13.5 ± 0.9 , 17 ± 0.7 , 18.3 ± 0.7 and $18.8 \pm 0.6\text{ km}^2$ for Tsokar and 140 ± 2.6 , 141 ± 1.3 , 141 ± 1.3 and $141 \pm 1.1\text{ km}^2$, respectively.

4.2. Glacier distribution in the Ladakh region

Glacierised areas and population in the Ladakh region vary across basins. Shayok Basin has the largest distribution of glacierised area and population (74% and 56%), whereas the Tsokar Basin has the least (0.04% and 0.1%), respectively (Table 2). Based on size distribution, the glacier area category of $1\text{--}5\text{ km}^2$ comprises the highest area (28% of the total), while the category of $50\text{--}100\text{ km}^2$ occupies the least glacierised area (9%) of the region. Most glaciers ($\sim 90\%$ of the total) in the Ladakh region have an area of $< 5\text{ km}^2$ but occupy only 37% of the total glacierised area. The population and area of glaciers in each area class are different in each basin but the proportion of glaciers, smaller than 5 km^2 , is greater than 87% in all basins. Glaciers larger than 100 km^2 ($n=7$, $< 1\%$ of the total) are only present in the Shayok Basin and occupy ~ 24 and 32% of the total glacierised area of Ladakh and Shayok Basin, respectively.

4.3. Glacier hypsometry, slope and aspect

Figure 3 (iii and iv) shows the glacier elevations and hypsometry with 100m elevation intervals of seven basins of the Ladakh region. The highest and lowest glacier elevation are 7740 and 3249m a.s.l., both in the Shayok Basin. Whereas mean elevation of the glacier ranges between 4345-6355m a.s.l. (Figure 3iii). Small glaciers mainly occupy the higher elevations above 5500, and vice versa. The majority (73%, 5810 km²) of the glacierised area is distributed in the 5000-6000 m a.s.l. elevation range, while only 14% is located below 5000m, and 13% above 6000m a.s.l. (Figure 3iv). The mean slope of these glaciers ranges between 8 and 46°, and is found to decrease with increasing glacier area. Glaciers with an area greater than 100 km² (n- 7, <1% of the total) have the lowest mean slope of 13° whereas, higher mean slopes (23°) are found for smaller glaciers (43% of the total). Overall, the mean glacier slope is ~21° (Figure 3v). Around 74% (1665) of the glaciers face the northern quadrant (NW-NE) amounting to ~50% (3940 km²) of the glacierised area. While 9, 5, 3, 3 and 4% of the glaciers face East, South-East, South, South-West and West which constitute 24, 6, 8, 6 and 6% of the glacierised area, respectively. However, the orientation and respective area coverage of glaciers vary within individual basins (Figure 3i, ii).

Table 2: Basin-wide glacier information of Ladakh region based on present study for the year 2019.

Basin	Basin area km ²	Total Area > 0.5 km ²		Area 0.5-1 km ²		Area 1-5 km ²		Area 5-10 km ²		Area 10-50 km ²		Area 50-100 km ²		Area > 100 km ²	
		Count	Area km ²	Count	Area km ²	Count	Area km ²	Count	Area km ²	Count	Area km ²	Count	Area km ²	Count	Area km ²
All	132180	2257	7923	980	694	1053	2206	124	853	84	1617	10	674	7	1879
Shayok	33579	1268 (56%)	5864 (74%)	495 (51%)	351 (51%)	609 (58%)	1304 (59%)	88 (71%)	621 (73%)	60 (71%)	1151 (71%)	8 (80%)	559 (83%)	7 (100%)	1879 (100%)
Leh	46579	247 (11%)	334 (4%)	147 (15%)	105 (16%)	95 (9%)	191 (9%)	4 (3%)	26 (3%)	1 (1%)	12 (1%)	0	0	0	0
Suru	10502	201 (9%)	498 (6%)	81 (8%)	59 (9%)	100 (9%)	212 (10%)	12 (10%)	69 (8%)	8 (10%)	159 (10%)	0	0	0	0
Zaskar	14817	256 (12%)	775 (10%)	116 (12%)	82 (12%)	111 (11%)	235 (11%)	15 (12%)	108 (13%)	12 (14%)	235 (15%)	2 (20%)	115 (17%)	0	0
Tsokar	1036	3 (0.1%)	3.5 (0.04%)	2 (0.2%)	1.5 (0.2%)	1 (0.1%)	2 (0.1%)	0	0	0	0	0	0	0	0
Tsomoriri	5462	94 (4%)	135 (2%)	47 (5%)	22 (3%)	46 (4%)	95 (4%)	1 (1%)	7 (1%)	0	0	0	0	0	0
Pangong	21206	190 (8%)	315 (4%)	92 (9%)	63 (9%)	91 (9%)	168 (8%)	4 (3%)	22 (2%)	3 (4%)	60 (4%)	0	0	0	0




Figure 3: General statistics of the glaciers in the Ladakh region: orientation of glaciers (i) and associated area distribution (ii), Maximum, minimum and mean elevation of glaciers (iii), hypsometry of glacierised area (iv) and slope against glacier area (v) and elevation (vi).

5. Discussion

5.1. The produced dataset and limitations

The multitemporal inventory of glaciers ($>0.5 \text{ km}^2$) in the Ladakh region for the years 1977, 1994, 2009 and 2019 is available at PANGAEA portal (<https://doi.org/10.1594/PANGAEA.940994>; Soheb et al., 2022). The dataset is

provided in two different GIS-ready file formats, i.e., GeoPackages (*.gpkg) and Shapefiles (*.dbf, *.prj, *.sbn, *.sbx, *.shp, *.shx) to support a wider end users. GeoPackage is a relatively new and open-source file format which is now being widely used and supported, whereas Shapefile format is one of the most widely used proprietary but open file formats for vector datasets, supported by open-source GIS tools such as QGIS. The outlines of glaciers, basins and lakes are all referenced to the WGS 84 / UTM zone 43N datum. For each region, there is one file for basin outlines, and four files for glacier and lake (if present) outlines for 1977, 1994, 2009 and 2019. Each glacier outline file contains glacier Ids (New glacier Ids, Randolph Glacier Inventory 6.0 Ids, and Global Land Ice Measurements from Space initiative Ids), coordinates (latitude and longitude), elevation (maximum, mean and minimum), aspect (mean), slope (mean), area, length (maximum), area uncertainty and length uncertainty. Whereas, the Lake Outline file contains coordinates, area, elevation and area uncertainty.

When using this dataset it is important to understand the key limitations of such regional-scale glacier inventories. Some of the key user limitations of the dataset are: (1) Glaciers smaller than 0.5 km² (which comprise ~70% and ~10% of the total glacier population and glacierised area, respectively) were not included in this inventory due to the higher uncertainty (~12-25%) associated with these glacier outlines; (2) Inventories produced in this study are entirely based on the medium resolution Landsat imagery, in the same way as other global or regional-scale glacier inventories. Although the uncertainty associated with these inventories do not considerably impact regional-scale analyses, care should be taken while using these data for a small subset of glaciers. It should also be noted that it is not feasible to produce multitemporal inventories regionally using high-resolution datasets due to the paucity and high costs of such high-resolution datasets; (3) The inventories of 1977±5, 1994±1 and 2019±1 are produced using images with a range of acquisition dates due to the lack of data continuity within a particular year (more details in section 3.1); and (4) The time periods chosen in this study are based on the availability of datasets and sufficient temporal gaps between the datasets to allow multitemporal glacio-hydrological analyses.

5.2. Significance of the present inventory

The glacier inventory presented here has several improvements compared to the existing regional and global inventories. Firstly, it covers the glaciers (> 0.5 km²; n = 2257; ~7923 ±106 km²) for the entire Ladakh region with manual correction and quality control undertaken using freely available high-resolution images. The analyses were further extended to estimate the distribution of ice masses at the sub-basin scale. Secondly, the temporal aspect of the glacierised area will aid hydrological and glaciological modelling aimed at understanding past and future system evolution. Finally, the new inventory will aid both the scientific community studying the glaciers and water resources of the Ladakh region, and the administration of the Union Territory of Ladakh, Government of India in developing efficient mitigation and adaptation strategies by improving the projections of change on timescales relevant to policy makers.

5.3. Comparison of inventories in the Ladakh region

Differences in estimates of the glacierised areas are meaningful as they can lead to an over or under estimation of the available water resources. Therefore, correctly estimating glacier area over time is necessary for understanding glacier dynamics, future response to climate forcing and the water resources they provide. Table 3 presents a comparison between the present inventory and the Randolph Glacier Inventory (RGI) 6.0 (Pfeffer et al., 2014), the International Centre for Integrated Mountain Development (ICIMOD) inventory (Bajracharya et al., 2011, 2019; Williams, 2013) and the Glacier Area Mapping for Discharge in Asian Mountains (GAMDAM) inventory (Guo et al., 2015; Nuimura et al., 2015; Sakai, 2019), for the Ladakh region. The comparison involves glacier outlines for 2009 from the present study and excludes glaciers smaller than 0.5 km² from regional inventories to achieve the closest match temporally and for glacier size categories. This should be taken as a first order comparison, given the fact that the uncertainties have been estimated with different approaches for the different inventories. Specifically the uncertainty estimated for the GAMDAM and ICIMOD inventories differs only slightly to the one applied here, given that they used a normalized standard deviation approach on the datasets produced by several operators on the same subset of glaciers (Bajracharya et al., 2011, 2019; Guo et al., 2015; Nuimura et al., 2015; Sakai, 2019). Whereas, in case of RGI 6.0 inventory, the uncertainty estimation approach differs significantly from the one presented here, because their errors were calculated on a collection of glaciers due to the vast quantity of data acquired from multiple sources and approaches used to produced them (Pfeffer et al., 2014). Figure 4 presents a comparison of the only three inventories (present, RGI 6.0 and ICIMOD) for the five field-investigated glaciers of Ladakh region because RGI and GAMDAM inventories share the same outlines for these glaciers.

The comparison showed a higher glacierised area in the RGI/GAMDAM inventories and lower in the ICIMOD inventory (Table 3) than the present inventory, with most of the differences contributed by the basins having the higher glacierised areas (Shayok and Zaskar) and from the larger glaciers (>10 km²). Such inconsistencies among the inventories are a product of several factors, e.g. 1) absence of change in glaciers over time due to the use of imagery with a wide range of acquisition years (Figure 4 a, c, d); 2) misinterpretation of the glacier terminus due to icing, debris, snow and cloud cover (Nagai et al., 2016), and 3) the methodology used. The smaller difference between the present and the ICIMOD inventory is mainly due to the adoption of a similar technique (i.e., a semi-automated approach) and the shorter time frame of the analysis that generated the ICIMOD inventory (i.e., 2002-2009).

Table 3: Basin and class wise comparison of the glacierised area between the present study and other inventories (RGI 6.0, ICIMOD and GAMDAM).

Region	Present Study	RGI 6.0			GAMDAM			ICIMOD		
		Area		Difference		Area		Area		Difference
		km ²	km ²	km ²	%	km ²	km ²	km ²	km ²	%
Shayok	5938	6999	1061	15	6616	678	10	5456	-482	-9
Zaskar	808	880	72	8	932	124	13	819	11	1
Suru	532	525	-7	-1	564	32	6	506	-26	-5
Leh	354	342	-12	-3	356	2	1	322	-32	-10
Tsokar	4	4.4	1	15	4.3	1	13	4.1	0	9
Tsomoriri	141	142	1	1	143	2	2	116	-25	-21

Pangong	320	320	0	0	335	15	4	-	-	-
Area Class										
0.5-1	758	774	16	2	803	45	6	662	-96	-7
1-5.	2284	2437	153	6	2385	101	4	1958	-326	-12
5-10.	862	961	99	10	925	63	7	766	-96	-10
10-50.	1628	1959	331	17	1824	196	11	1356	-272	-20
50-100	678	730	52	7	599	-79	-13	592	-86	-15
>100	1887	2351	464	20	2412	525	22	1887	0	0
Total	8096	9212	1116	14	8950	854	11	7223	-533	-7




Figure 4: Comparison of inventories on the field investigated glaciers of the Ladakh region: a) Parkachik glacier, Suru Basin; b) Pensila glacier, Suru Basin; c) Lato glacier, Leh Basin; d) Khardung glacier, Shayok Basin; e) Stok glacier, Leh Basin. The background image (Landsat) courtesy: U.S. Geological Survey/NASA, <https://earthexplorer.usgs.gov/>.

5.4. Comparison with recent studies

The data from the recent spatio-temporal change studies from different sub-regions of Ladakh (Figure 5) are not in the public domain, except from Shukla et al., 2020. Hence, it is not possible to use these to validate our results.

Therefore, our comparison mostly focuses on the rate of change for some of the individual glaciers (n=21, Figure 5) from the literature and the bulk properties of a set of glaciers in different regions (Table S3, Figure 6). Our results agree well with the studies conducted by others (Bhambri et al., 2013; Chudley et al., 2017; Garg et al., 2022; Garg et al., 2021; Negi et al., 2021; Schmidt & Nüsser, 2012, 2017; Shukla et al., 2020) on individual glaciers of various sizes as well as on a set of glaciers, respectively (Figure 6, Table S3). However, the results differ significantly only on some glaciers and especially in a part of the Shayok Basin (e.g. Kumdan (D), Aktash (E) and Thusa glaciers(I)). In the Shayok Basin surge-type glaciers are common (Bhambri et al., 2013, 2017), the difference in analysis period between the present and other studies is the likely cause of the difference in glacier area statistics. Figure S2 presents the dynamics of the Kumdan and Aktash glaciers as an example of surge type glacier of this region.

No significant difference was observed in rate of change of glacierised areas between the present study and other studies in the Leh, Tsomoriri, Zaskar and Suru Basins. In contrast, the number of glaciers and glacierised area vary among these studies (present and others) but paint a similar picture of relatively lower retreat in the Shayok Basin (Bhambri et al., 2013; Negi et al., 2021), higher in Leh, Tsokar, Tsomoriri (Chudley et al., 2017; Schmidt & Nüsser, 2012, 2017) and moderate in Zaskar and Suru Basins (Garg et al., 2022; Garg et al., 2022; Shukla et al., 2020).




Figure 5: Presents the spatial extent of different studies undertaken in Ladakh region. Black stars represent the individual glaciers. The background image (ASTER GDEM) courtesy: NASA/METI/AIST/Japan Spacesystems and U.S./Japan ASTER Science Team, <https://earthdata.nasa.gov/>.




Figure 6: Comparison between the present study and other studies undertaken in different basins of Ladakh region over different time periods

6. Data availability

The entire dataset of the Landsat based multitemporal inventory of glaciers, larger than 0.5 km², in Ladakh region for the year 1977, 1994, 2009 and 2019 is available at: PANGAEA, <https://doi.org/10.1594/PANGAEA.940994> (Soheb et al., 2022).

7. Conclusions

We compiled a new glacier inventory of the Ladakh region for 1977, 1994, 2009 and 2019 based on 63 Landsat (MSS, TM and OLL) images, with least cloud/snow cover, acquired during the summer time (July-October). The inventory includes 2257 glaciers, larger than 0.5 km², covering an area of $\sim 7923 \pm 106$ km² which is $\sim 14\%$ and $\sim 11\%$ less than the RGI 6.0 and the GAMDAM, and 7% more than the ICIMOD inventories. The glacierised area accounts for $\sim 6\%$ of the Ladakh region with individual glacier areas ranging between 0.5 ± 0.02 and 862 ± 16 km². About 90% of the glacier population are smaller than 5 km² but combined they occupy only 37% of the glacierised area. The seven largest glaciers, larger than 100 km², account for ~ 1879 km² or 24% of the total. The Shayok Basin and glacier area category 1-5 km² hosts the highest number of glacier population and glacierised area; whereas, Tsokar Basin accounts for the least. More than 70% of the glaciers are in the north-facing quadrant (NW-NE) and are concentrated in the higher elevation zones, between 5000 and 6000 m a.s.l. The error assessment shows that the uncertainty, based on a buffer-based approach, ranges between 2.6 and 5.1% for glacier area, and 1.5 and 2.6% for glacier length with a mean uncertainty of 3.2 and 1.8%, respectively. The uncertainty varies depending on the quality of the images and size of the glaciers. Our results also show a good agreement with other studies undertaken in parts of the Ladakh region for individual glaciers (n=21) and bulk properties of a set of glaciers.

The new multi-temporal inventory presented here will assist in planning the management of water resources, and for guiding scientific research focusing on glacier mass balance, hydrology and glacier change within the region. The detailed information and multi-temporal nature of this inventory will also aid in improving the existing global and regional glacier inventories especially in the cold-arid Ladakh region where the majority of the population is highly dependent on glacier-derived melt water resources for domestic, irrigation and hydropower generation needs.

Author contribution

MSo, AR, AB conceptualized and designed the study. MSo, AB and MC did the analysis. MSo wrote and AR, AB, MSp, BR, SS and LS edited the manuscript. All the authors have equally contributed to interpretation of the results.

Competing interests:

The authors declare that they have no conflict of interest.

Acknowledgements

The authors are thankful to the School of Environmental Sciences, Jawaharlal Nehru University, New Delhi, India, for the lab facilities and the United States Geological Survey for the Landsat and ASTER imageries. The authors also thank Planet Labs and Google for the high resolution PlanetScope and Google Earth imageries. We are also thankful to the Scottish Funding Council and the University Of Aberdeen, United Kingdom for financially supporting our work.

References

- Azam, M. F., Kargel, J. S., Shea, J. M., Nepal, S., Haritashya, U. K., Srivastava, S., Maussion, F., Qazi, N., Chevallier, P., Dimri, A. P., Kulkarni, A. V., Cogley, J. G., and Bahuguna, I. M.: Glaciohydrology of the Himalaya-Karakoram, *Science* 373, eabf3668, <https://doi.org/10.1126/science.abf3668>, 2021.
- Bajracharya, S. R. and Shrestha, B. R., International Centre for Integrated Mountain Development, and Sweden (Eds.): The status of glaciers in the Hindu Kush-Himalayan region, International Centre for Integrated Mountain Development, Kathmandu, 127 pp., 2011.
- Bajracharya, S. R., Maharjan, S. B., and Shrestha, F.: Glaciers in the Indus Basin, in: Indus River Basin, Elsevier, 123–144, <https://doi.org/10.1016/B978-0-12-812782-7.00006-0>, 2019.
- Barrett, K. and Bosak, K.: The Role of Place in Adapting to Climate Change: A Case Study from Ladakh, Western Himalayas, *Sustainability*, 10, 898, <https://doi.org/10.3390/su10040898>, 2018.
- Bhambri, R., Bolch, T., Chaujar, R. K., and Kulshreshtha, S. C.: Glacier changes in the Garhwal Himalaya, India, from 1968 to 2006 based on remote sensing, *J. Glaciol.*, 57, 543–556, <https://doi.org/10.3189/002214311796905604>, 2011.
- Bhambri, R., Bolch, T., Kawishwar, P., Dobhal, D. P., Srivastava, D., and Pratap, B.: Heterogeneity in glacier response in the upper Shyok valley, northeast Karakoram, *The Cryosphere*, 7, 1385–1398, <https://doi.org/10.5194/tc-7-1385-2013>, 2013.
- Bhambri, R., Hewitt, K., Kawishwar, P., and Pratap, B.: Surge-type and surge-modified glaciers in the Karakoram, *Sci. Rep.*, 7, 15391, <https://doi.org/10.1038/s41598-017-15473-8>, 2017.
- Bhardwaj, A., Joshi, P., Snehmani., Sam, L., Singh, M. K., Singh, S., and Kumar, R.: Applicability of Landsat 8 data for characterizing glacier facies and supraglacial debris, *Int. J. Appl. Earth Obs. Geoinformation*, 38, 51–64, <https://doi.org/10.1016/j.jag.2014.12.011>, 2015.
- Bolch, T.: Past and Future Glacier Changes in the Indus River Basin, in: Indus River Basin, Elsevier, 85–97, <https://doi.org/10.1016/B978-0-12-812782-7.00004-7>, 2019.
- Bolch, T., Menounos, B., and Wheate, R.: Landsat-based inventory of glaciers in western Canada, 1985–2005, *Remote Sens. Environ.*, 114, 127–137, <https://doi.org/10.1016/j.rse.2009.08.015>, 2010.
- Bolch, T., Kulkarni, A., Kaab, A., Huggel, C., Paul, F., Cogley, J. G., Frey, H., Kargel, J. S., Fujita, K., Scheel, M., Bajracharya, S., and Stoffel, M.: The State and Fate of Himalayan Glaciers, *Science*, 336, 310–314, <https://doi.org/10.1126/science.1215828>, 2012.
- Census of India (2011), Jammu and Kashmir, Series 02- Part XII A-B, District Census Handbook, Leh and Kargil. India - Census of India 2011 - Jammu & Kashmir - Series 02 - Part XII A - District Census Handbook, Leh (censusindia.gov.in), last assess: 30 September 2021.
- Chudley, T. R., Miles, E. S., and Willis, I. C.: Glacier characteristics and retreat between 1991 and 2014 in the Ladakh Range, Jammu and Kashmir, *Remote Sens. Lett.*, 8, 518–527, <https://doi.org/10.1080/2150704X.2017.1295480>, 2017.

415 Frey, H. and Paul, F.: On the suitability of the SRTM DEM and ASTER GDEM for the compilation of
 416 topographic parameters in glacier inventories, *Int. J. Appl. Earth Obs. Geoinformation*, 18, 480–490,
 417 <https://doi.org/10.1016/j.jag.2011.09.020>, 2012.

418 Frey, H., Machguth, H., Huss, M., Huggel, C., Bajracharya, S., Bolch, T., Kulkarni, A., Linsbauer, A.,
 419 Salzmann, N., and Stoffel, M.: Estimating the volume of glaciers in the Himalayan-Karakoram region
 420 using different methods, *The Cryosphere*, 8, 2313–2333, <https://doi.org/10.5194/tc-8-2313-2014>, 2014.

421 Garg, P. K., Garg, S., Yousuf, B., Shukla, A., Kumar, V., and Mehta, M.: Stagnation of the Pensilungpa
 422 glacier, western Himalaya, India: causes and implications, *J. Glaciol.*, 68, 221–235,
 423 <https://doi.org/10.1017/jog.2021.84>, 2022a.

424 Garg, S., Shukla, A., Garg, P. K., Yousuf, B., and Shukla, U. K.: Surface evolution and dynamics of the
 425 Kangriz glacier, western Himalaya in past 50 years, *Cold Reg. Sci. Technol.*, 196, 103496,
 426 <https://doi.org/10.1016/j.coldregions.2022.103496>, 2022b.

427 Granshaw, F. D. and G. Fountain, A.: Glacier change (1958–1998) in the North Cascades National Park
 428 Complex, Washington, USA, *J. Glaciol.*, 52, 251–256, <https://doi.org/10.3189/172756506781828782>,
 429 2006.

430 Guo, W., Liu, S., Xu, J., Wu, L., Shangguan, D., Yao, X., Wei, J., Bao, W., Yu, P., Liu, Q., and Jiang, Z.: The
 431 second Chinese glacier inventory: data, methods and results, *J. Glaciol.*, 61, 357–372,
 432 <https://doi.org/10.3189/2015JoG14J209>, 2015.

433 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C.,
 434 Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati,
 435 G., Bidlot, J., Bonavita, M., Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J.,
 436 Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley,
 437 S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and
 438 Thépaut, J.: The ERA5 global reanalysis, *Q. J. R. Meteorol. Soc.*, 146, 1999–2049,
 439 <https://doi.org/10.1002/qj.3803>, 2020.

440 Ji, Q., Yang, T., He, Y., Qin, Y., Dong, J., and Hu, F.: A simple method to extract glacier length based on
 441 Digital Elevation Model and glacier boundaries for simple basin type glacier, *J. Mt. Sci.*, 14, 1776–1790,
 442 <https://doi.org/10.1007/s11629-016-4243-5>, 2017.

443 Kääb, A., Treichler, D., Nuth, C., and Berthier, E.: Brief Communication: Contending estimates of
 444 2003–2008 glacier mass balance over the Pamir–Karakoram–Himalaya, *The Cryosphere*, 9,
 445 557–564, <https://doi.org/10.5194/tc-9-557-2015>, 2015.

446 Khan, A., Richards, K. S., Parker, G. T., McRobie, A., and Mukhopadhyay, B.: How large is the Upper Indus
 447 Basin? The pitfalls of auto-delineation using DEMs, *J. Hydrol.*, 509, 442–453,
 448 <https://doi.org/10.1016/j.jhydrol.2013.11.028>, 2014.

449 Kulkarni, A. V.: Monitoring Himalayan cryosphere using remote sensing techniques, *J. India Inst. Sci.* 90,
 450 457–469, 2010.

451 Le Bris, R. and Paul, F.: An automatic method to create flow lines for determination of glacier length: A
 452 pilot study with Alaskan glaciers, *Comput. Geosci.*, 52, 234–245,
 453 <https://doi.org/10.1016/j.cageo.2012.10.014>, 2013.

454 Liu, S., Ding, Y., Shangguan, D., Zhang, Y., Li, J., Han, H., Wang, J., and Xie, C.: Glacier retreat as a result of
 455 climate warming and increased precipitation in the Tarim river basin, northwest China, *Ann. Glaciol.*, 43,
 456 91–96, <https://doi.org/10.3189/172756406781812168>, 2006.

457 Maurer, J. M., Schaefer, J. M., Rupper, S., and Corley, A.: Acceleration of ice loss across the Himalayas
 458 over the past 40 years, *Sci. Adv.*, 5, eaav7266, <https://doi.org/10.1126/sciadv.aav7266>, 2019.

459 Minora, U., Bocchiola, D., D’Agata, C., Maragno, D., Mayer, C., Lambrecht, A., Mosconi, B., Vuillermoz, E.,
 460 Senese, A., Compostella, C., Smiraglia, C., and Diolaiuti, G.: 2001–2010 glacier changes in the Central
 461 Karakoram National Park: a contribution to evaluate the magnitude and rate of the "Karakoram
 462 anomaly", *The Cryosphere Discuss.*, 7, 2891–2941, <https://doi.org/10.5194/tcd-7-2891-2013>, 2013.

463 Mölg, N., Bolch, T., Rastner, P., Strozzi, T., and Paul, F.: A consistent glacier inventory for Karakoram and
 464 Pamir derived from Landsat data: distribution of debris cover and mapping challenges, *Earth Syst. Sci.*
 465 *Data*, 10, 1807–1827, <https://doi.org/10.5194/essd-10-1807-2018>, 2018.

466 Mukherjee, S., Joshi, P. K., Mukherjee, S., Ghosh, A., Garg, R. D., and Mukhopadhyay, A.: Evaluation of
 467 vertical accuracy of open source Digital Elevation Model (DEM), *Int. J. Appl. Earth Obs. Geoinformation*,
 468 21, 205–217, <https://doi.org/10.1016/j.jag.2012.09.004>, 2013.

469 Müller, J., Dame, J., and Nüsser, M.: Urban Mountain Waterscapes: The Transformation of Hydro-Social
 470 Relations in the Trans-Himalayan Town Leh, Ladakh, India, *Water*, 12, 1698,
 471 <https://doi.org/10.3390/w12061698>, 2020.

472 National Bureau of Statistics. Communiqué of the seventh National Population Census.
 473 <http://www.stats.gov.cn/tjsj/>, last access: 30 September 2021.

474 Nagai, H., Fujita, K., Sakai, A., Nuimura, T., and Tadono, T.: Comparison of multiple glacier inventories
 475 with a new inventory derived from high-resolution ALOS imagery in the Bhutan Himalaya, *The*
 476 *Cryosphere*, 10, 65–85, <https://doi.org/10.5194/tc-10-65-2016>, 2016.

477 Negi, H. S., Kumar, A., Kanda, N., Thakur, N. K., and Singh, K. K.: Status of glaciers and climate change of
 478 East Karakoram in early twenty-first century, *Sci. Total Environ.*, 753, 141914,
 479 <https://doi.org/10.1016/j.scitotenv.2020.141914>, 2021.

480 Nuimura, T., Sakai, A., Taniguchi, K., Nagai, H., Lamsal, D., Tsutaki, S., Kozawa, A., Hoshina, Y., Takenaka,
 481 S., Omiya, S., Tsunematsu, K., Tshering, P., and Fujita, K.: The GAMDAM glacier inventory: a quality-
 482 controlled inventory of Asian glaciers, *The Cryosphere*, 9, 849–864, [https://doi.org/10.5194/tc-9-849-](https://doi.org/10.5194/tc-9-849-2015)
 483 2015, 2015.

484 Nüsser, M., Schmidt, S., and Dame, J.: Irrigation and Development in the Upper Indus Basin:
 485 Characteristics and Recent Changes of a Socio-hydrological System in Central Ladakh, India, *Mt. Res.*
 486 *Dev.*, 32, 51–61, <https://doi.org/10.1659/MRD-JOURNAL-D-11-00091.1>, 2012.

487 Nüsser, M., Dame, J., Parveen, S., Kraus, B., Baghel, R., and Schmidt, S.: Cryosphere-Fed Irrigation
 488 Networks in the Northwestern Himalaya: Precarious Livelihoods and Adaptation Strategies Under the
 489 Impact of Climate Change, *Mt. Res. Dev.*, 39, <https://doi.org/10.1659/MRD-JOURNAL-D-18-00072.1>,
 490 2019a.

491 Nüsser, M., Dame, J., Kraus, B., Baghel, R., and Schmidt, S.: Socio-hydrology of “artificial glaciers” in
 492 Ladakh, India: assessing adaptive strategies in a changing cryosphere, *Reg. Environ. Change*, 19, 1327–
 493 1337, <https://doi.org/10.1007/s10113-018-1372-0>, 2019b.

494 Paul, F., Barry, R. G., Cogley, J. G., Frey, H., Haeberli, W., Ohmura, A., Ommanney, C. S. L., Raup, B.,
 495 Rivera, A., and Zemp, M.: Recommendations for the compilation of glacier inventory data from digital
 496 sources, *Ann. Glaciol.*, 50, 119–126, <https://doi.org/10.3189/172756410790595778>, 2009.

497 Paul, F., Barrand, N. E., Baumann, S., Berthier, E., Bolch, T., Casey, K., Frey, H., Joshi, S. P., Konovalov, V.,
 498 Le Bris, R., Mölg, N., Nosenko, G., Nuth, C., Pope, A., Racoviteanu, A., Rastner, P., Raup, B., Scharrer, K.,
 499 Steffen, S., and Winsvold, S.: On the accuracy of glacier outlines derived from remote-sensing data, *Ann.*
 500 *Glaciol.*, 54, 171–182, <https://doi.org/10.3189/2013AoG63A296>, 2013.

501 Paul, F., Bolch, T., Briggs, K., Kääb, A., McMillan, M., McNabb, R., Nagler, T., Nuth, C., Rastner, P., Strozzi,
 502 T., and Wuite, J.: Error sources and guidelines for quality assessment of glacier area, elevation change,
 503 and velocity products derived from satellite data in the Glaciers_cci project, *Remote Sens. Environ.*, 203,
 504 256–275, <https://doi.org/10.1016/j.rse.2017.08.038>, 2017.

505 Pfeiffer, W. T., Arendt, A. A., Bliss, A., Bolch, T., Cogley, J. G., Gardner, A. S., Hagen, J.-O., Hock, R., Kaser,
 506 G., Kienholz, C., Miles, E. S., Moholdt, G., Mölg, N., Paul, F., Radić, V., Rastner, P., Raup, B. H., Rich, J.,
 507 Sharp, M. J., and The Randolph Consortium: The Randolph Glacier Inventory: a globally complete
 508 inventory of glaciers, *J. Glaciol.*, 60, 537–552, <https://doi.org/10.3189/2014JoG13J176>, 2014.

509 Pritchard, H. D.: Asia’s shrinking glaciers protect large populations from drought stress, *Nature*, 569,
 510 649–654, <https://doi.org/10.1038/s41586-019-1240-1>, 2019.

511 Racoviteanu, A. E., Paul, F., Raup, B., Khalsa, S. J. S., and Armstrong, R.: Challenges and
 512 recommendations in mapping of glacier parameters from space: results of the 2008 Global Land Ice
 513 Measurements from Space (GLIMS) workshop, Boulder, Colorado, USA, *Ann. Glaciol.*, 50, 53–69,
 514 <https://doi.org/10.3189/172756410790595804>, 2009.

515 Racoviteanu, A. E., Rittger, K., and Armstrong, R.: An Automated Approach for Estimating Snowline
 516 Altitudes in the Karakoram and Eastern Himalaya From Remote Sensing, *Front. Earth Sci.*, 7, 220,
 517 <https://doi.org/10.3389/feart.2019.00220>, 2019.

518 Sakai, A.: Brief communication: Updated GAMDAM glacier inventory over high-mountain Asia, *The*
 519 *Cryosphere*, 13, 2043–2049, <https://doi.org/10.5194/tc-13-2043-2019>, 2019.

520 Schmidt, S. and Nüsser, M.: Changes of High Altitude Glaciers from 1969 to 2010 in the Trans-Himalayan
 521 Kang Yatze Massif, Ladakh, Northwest India, *Arct. Antarct. Alp. Res.*, 44, 107–121,
 522 <https://doi.org/10.1657/1938-4246-44.1.107>, 2012.

523 Schmidt, S. and Nüsser, M.: Changes of High Altitude Glaciers in the Trans-Himalaya of Ladakh over the
 524 Past Five Decades (1969–2016), *Geosciences*, 7, 27, <https://doi.org/10.3390/geosciences7020027>, 2017.

525 Shean, D. E., Bhushan, S., Montesano, P., Rounce, D. R., Arendt, A., and Osmanoglu, B.: A Systematic,
 526 Regional Assessment of High Mountain Asia Glacier Mass Balance, *Front. Earth Sci.*, 7, 363,
 527 <https://doi.org/10.3389/feart.2019.00363>, 2020.

528 Shukla, A., Garg, S., Mehta, M., Kumar, V., and Shukla, U. K.: Temporal inventory of glaciers in the Suru
 529 sub-basin, western Himalaya: impacts of regional climate variability, *Earth Syst. Sci. Data*, 12, 1245–
 530 1265, <https://doi.org/10.5194/essd-12-1245-2020>, 2020.

531 Singh, S., Kumar, R., Bhardwaj, A., Sam, L., Shekhar, M., Singh, A., Kumar, R., and Gupta, A.: Changing
 532 climate and glacio-hydrology in Indian Himalayan Region: a review, *WIREs Clim. Change*, 7, 393–410,
 533 <https://doi.org/10.1002/wcc.393>, 2016.

534 Soheb, M; Ramanathan, A; Bhardwaj, A; Coleman, M; Spagnolo, M; Rea, B. R; Singh, S; Sam, L: Landsat-
 535 based multitemporal glacier inventory data of over four decades (1977-2019) for Ladakh
 536 region. PANGAEA [data set], <https://doi.pangaea.de/10.1594/PANGAEA.940994>, 2022.

537 Tielidze, L. G. and Wheate, R. D.: The Greater Caucasus Glacier Inventory (Russia, Georgia and
 538 Azerbaijan), *The Cryosphere*, 12, 81–94, <https://doi.org/10.5194/tc-12-81-2018>, 2018.

539 Williams, M. W.: The Status of Glaciers in the Hindu Kush–Himalayan Region, *Mt. Res. Dev.*, 33, 114,
 540 <https://doi.org/10.1659/mrd.mm113>, 2013.

541 Winsvold, S. H., Kaab, A., and Nuth, C.: Regional Glacier Mapping Using Optical Satellite Data Time
 542 Series, *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.*, 9, 3698–3711,
 543 <https://doi.org/10.1109/JSTARS.2016.2527063>, 2016.

544 Yao, J., Chao-lu, Y., and Ping, F.: Evaluation of the Accuracy of SRTM3 and ASTER GDEM in the Tibetan
 545 Plateau Mountain Ranges, *E3S Web Conf.*, 206, 01027, <https://doi.org/10.1051/e3sconf/202020601027>,
 546 2020.

547 Zhang, M., Wang, X., Shi, C., and Yan, D.: Automated Glacier Extraction Index by Optimization of
 548 Red/SWIR and NIR /SWIR Ratio Index for Glacier Mapping Using Landsat Imagery, *Water*, 11, 1223,
 549 <https://doi.org/10.3390/w11061223>, 2019.

550