

PalVol v1: A proxy-based semi-stochastic ensemble reconstruction of volcanic stratospheric sulfur injection for the last glacial cycle (140,000 – 50 BP)

5 Julie Christin Schindlbeck-Belo^{1*}, Matthew Toohey², Marion Jegen¹, Steffen Kutterolf¹, Kira Rehfeld³

¹GEOMAR Helmholtz Centre for Ocean Research Kiel, Wischhofstr.1-3, 24148 Kiel, Germany

²Institute of Space and Atmospheric Studies, Department of Physics & Engineering Physics, University of Saskatchewan, S7N 5A2 Saskatoon, Canada

10 ³Department of Geosciences and Department of Physics, Geo- und Umweltforschungszentrum, Schnarrenbergstr. 94/96, Tübingen University, 72076 Tübingen, Germany

Correspondence to: Julie C. Belo (jbelo@geomar.de)

15 **Abstract.**

Perturbations in stratospheric aerosol due to explosive volcanic eruptions are a primary contributor to natural climate variability. Observations of stratospheric aerosol are available for the past decades, and information from ice cores has been used to derive estimates of stratospheric sulfur injections and aerosol optical depth over the Holocene (approximately 10,000 BP to present) and into the last glacial period, extending back to 60,000 BP. Tephra records of past volcanism, compared to
20 ice cores, are less complete, but extend much further into the past. To support model studies of the potential impacts of explosive volcanism on climate variability over across timescales, we present here an ensemble reconstruction of volcanic stratospheric sulfur injection (VSSI) over the last 140,000 years that is based primarily on terrestrial and marine tephra records. VSSI values are computed as a simple function of eruption magnitude, based on VSSI estimates from ice cores and satellite observations for identified eruptions. To correct for the incompleteness of the tephra record we include stochastically generated
25 synthetic eruptions, assuming a constant background eruption frequency from the ice core Holocene record. While the reconstruction often differs from ice core estimates for specific eruptions due to uncertainties in the data used and reconstruction method, it shows good agreement with an ice core based VSSI reconstruction in terms of millennial-scale cumulative VSSI variations over the Holocene. The PalVol reconstruction provides a new basis to test the contributions of forced vs. unforced natural variability to the spectrum of climate, and the mechanisms leading to abrupt transitions in the
30 palaeoclimate record with low-to-high complexity climate models. The PalVol volcanic forcing reconstruction is available at <https://doi.org/10.26050/WDCC/PalVolv1> (Toohey, Schindlbeck-Belo, 2023).

1 Introduction

Explosive volcanic eruptions transfer massive quantities of material from the solid Earth into the atmosphere. Eruptive plumes contain large amounts of solid material, as well as gaseous compounds including water vapor, carbon dioxide and sulfur-containing species (mostly SO₂), often being combined into aerosols. The solid volcanic fragments are fragmented magma that is ejected by an eruption and are called tephra (mostly ash: $\phi < 2\text{mm}$, and lapilli: $\phi 2$ to 64mm). They generally fall back to the surface of the Earth rather quickly, producing a tephra layer with decreasing thickness and grain size with increasing distance to the volcano. Such deposits persist as a record of past volcanic eruptions that can be seen in outcrops, or in sediment cores extracted from marine and lacustrine environments (e.g., Kutterolf et al., 2016; Schindlbeck et al., 2016).

Gaseous emissions from eruptions can persist in the atmosphere much longer than solid emissions. While the emission of H₂O and CO₂ from a single eruption is generally insignificant compared to the atmosphere burden of these species, volcanic emissions of sulfur-containing species can produce significant increases in atmospheric sulfur (Robock, 2000). Under atmospheric conditions, sulfur-containing species produce sulfate aerosol particles, small droplets of sulfate (SO₄²⁻) and water in liquid form that are dispersed in a gaseous matrix (Robock, 2000). Sulfur emissions to the troposphere (from volcanic eruptions or anthropogenic activities) produce sulfate aerosols that have an atmospheric lifetime of days to weeks, as they grow in size and are eventually “rained out” (Hamill, 1997). Sulfate aerosols in the cold and dry lower stratosphere do not generally grow as large as those in the troposphere, and as a result persist in the stratosphere for months to years, over which time they are transported around the globe. These sulfate aerosols have important impacts on atmospheric radiative transfer, by scattering solar radiation and absorbing longwave radiation, with the net effect of a decrease in downwelling net radiation at the Earth’s surface, which leads to a cooling of surface temperatures (Robock, 2000).

Post-volcanic large-scale (global or hemispheric) cooling has been observed after recent eruptions (e.g., El Chichón in 1982, Mt. Pinatubo in 1991), and is apparent in millennial scale climate reconstructions, for example the 1815 Tambora eruption in Indonesia (e.g., Rampino and Self, 1982, 1993). It has been shown that negative radiative forcing from volcanic aerosol perturbations is the primary driver of natural climate variability over the past thousand to two thousand years (Sigl et al., 2015; Schurer et al., 2013). Additionally, Kobashi et al. (2017) showed that also during the Holocene, volcanic eruptions played an important role in centennial to millennial temperature variability on Greenland. Representing the intermittent natural forcing in climate model experiments for the late Holocene and the Glacial period improves modeled variability on timescales from decades to centuries (Ellerhoff et al., 2022). And Abbott et al. (2022) provide a 400-year reconstruction of volcanic forcing from 13.2-12.8 ka BP, which supports model simulations for the Younger Dryas inception.

The role of volcanic eruptions in longer-term climate variability has long been speculated but remains poorly understood. Strong volcanic eruptions have been linked to multi-decadal periods of cooler than usual surface temperatures, for example during the Little Ice Age (Owens et al., 2017; Miller et al., 2012; Zanchettin et al., 2013; Schurer et al., 2014; Timmreck et al., 2021) and the so-called “Late Antiquity Little Ice Age (LALIA, Büntgen et al., 2020, Toohey et al., 2016). The manifestation of volcanic radiative forcing as multi-decadal temperature anomalies has been suggested to be related to the thermal inertia of

the Earth's oceans, which dampen the initial temperature response, but also prolong it through the accumulation of negative energy input in the deep ocean (Gupta et al., 2018). Indeed, ocean sea surface temperature changes over the 801-1800 CE period are well explained by the time series of volcanic eruptions (McGregor et al., 2015). There are climate modeling studies that suggest that strong volcanic radiative forcing can perturb ocean circulation modes (Swingedouw et al., 2017), which may produce long term perturbations to surface climate (Miller et al., 2014; Schleussner and Feulner, 2013; Schleussner et al., 2015; Otterå et al., 2010, Zhong et al., 2011). Indeed, at the global scale, clusters of eruptions have been linked to global mean temperature variations over the Common Era (Pages2k-Consortium, 2019). But other work has suggested a potential match in the timings of large eruptions or clusters of eruptions with the sudden transitions in climate between stadials and interstadials (Baldini et al., 2015; Bay et al., 2006; Lohmann and Svensson, 2022), although the robustness of this connection is limited by uncertainties in eruption magnitudes and timings.


Ice core sulfur (Huybers and Langmuir, 2009) and tephra chronologies (e.g., Praetorius et al., 2016; Sigvaldason et al., 1992) both attest to a marked increase in eruption frequencies during and after the last glaciation, especially in the northern hemisphere (NH) mid-to-high latitudes. On longer timescales, variations in eruption frequency from deep marine cores suggest periodicities on Milankovic time scales (Kutterolf et al., 2013, 2019; Schindlbeck et al., 2018a), suggesting a connection between the climate changes brought about by variations in orbital parameters and volcanic eruption frequencies. A leading theory is that mass transfer between ice sheets and the ocean due to changing temperatures leads to changes in the pressure on the Earth's crust which can modulate crustal stress fields and enhance the possibility of associated eruptive events by providing pathways for the magma to rise (Kutterolf et al., 2013).

Recent ice core-based reconstructions of volcanic sulfur emissions confirm the increase in explosive eruptive frequency after the last deglaciation, and its likely impact on stratospheric aerosol levels (Sigl et al., 2022). Ice core data presented by Lin et al. (2022) extends into the last glacial period, and suggests that during glacial conditions, the frequency of large eruptions with significant VSSI was similar to that after the deglaciation.

The emissions of subaerial eruptions are eventually deposited to the Earth's surface and in some cases preserved providing us with records of past volcanic activity. Tephra layers from past eruptions are often discernable in terrestrial outcrops, but the records are often incomplete since younger deposits overlay the stratigraphy or the deposits are already heavily weathered, eroded, or covered by vegetation. Ash and tephra, either transported by fallout or density flow processes, are well-preserved in the marine sediments, since wide areas of the seafloor are relatively unaffected by erosion or bioturbation (Freundt et al., 2021). This makes marine sediments outstanding archives for previous eruptions. By piston or gravity coring (~up to 20 m depth) or drilling (several hundreds of meters) the marine sediment records can be recovered and the marine ash layers provide a stratigraphically controlled record. While drilling is time-consuming and expensive, and the lateral coverage thus limited to a few sites, multiple shorter piston or gravity cores can be taken in a certain region providing good coverage, which enables detection of lateral stratigraphic changes or local erosion. Due to their length, however, they are often limited to the last two glacial cycles. In general, marine tephra records from sediment cores can cover several millions of years depending on the

used coring/drilling technique. There are geographical gaps, or regions with sparse coverage of cores and by far not all cores
100 have been studied for their tephra inventory in detail.


Ice cores (Fig. 1) provide a good archive for past volcanic eruptions, as the volcanic sulfate aerosols and in some cases ash
particles are deposited to the surface and incorporated into the glacial ice. Chemical analyses of ice cores therefore provide
time series, which are especially valuable when the dating of the ice cores is of high quality (Hammer, 1977; Gao et al., 2008;
Sigl et al., 2015). Past volcanic eruptions are evidenced by strong increases in sulfur/sulfate content of the ice, or in the
105 electrical conductivity of the ice, as the analysis traverses the ice core and thus reaches backwards in time. By synchronizing
multiple ice cores from Greenland and Antarctica, recent efforts have produced estimates of the VSSI from volcanic eruptions
covering the past 2,500 years (Toohey and Sigl, 2017), the Holocene (Cole-Dai et al., 2021; Sigl et al., 2015), and the late
glacial period and deglaciation (Lin et al., 2022).



110 **Figure 1: Global overview map created using GeoMapApp (<http://www.geomapapp.org>; GMRT-Global Multi-Resolution Topography) (Ryan et al., 2009). Red triangles mark global distribution of volcanoes from the LaMEVE database (Crossweller et al., 2012). Black squares mark regions with marine sediment cores that have been used in this publication. Red circles show positions of ice cores in Greenland and Antarctica used by Cole-Dai et al. (2021) and Sigl et al. (2015, 2022 and references therein).**

115 Due to limitations with ice cores—absolute length, thinning, and synchronization, there is a limit to the length of time ice cores can be used to reconstruct past volcanism. Tephra records from sediment cores, on the other hand, extend much further back in time. However, they have their own limitations: incompleteness, dating uncertainty, not as direct a measure of the stratospheric sulfate aerosol load as ice core sulfate records are. There is a strong temporal trend in the dataset, which is

120 described by a decreasing number of detected events going back in time (Fig. 2). Brown et al. (2014) for example emphasize
in the LaMEVE database, which covers the last 1.8 Ma, about 40% of the detected eruptions included occurred during the
Holocene (the past 11 ka), and they conclude that the decrease going back in time is mainly due to under-reporting of eruptions.
The time trend in underreporting is found to depend on the magnitude of eruption, with the frequency of smaller eruptions
($4.0 \leq M < 5.0$) falling off much faster than that for larger eruptions ($M > 6$) (Fig. 2). Despite these limitations, far reaching
125 tephra records do provide valuable information about specific events (e.g., Pinatubo, Toba) and changes in eruption frequency
with time. For example, analysis of tephra records long enough to cover several glacial cycles, show that eruption frequencies
vary on periods representative of Milankovitch cycles, supporting the claims that the Earth's climate influences eruption
frequencies on long time scales through changes in sea level and associated crustal stresses (Paterne et al., 1990; Rampino et
al., 1979; Kutterolf et al., 2013, 2019; Schindlbeck et al., 2018a).



130 **Figure 2: Scatter plot showing the distribution of eruptions by magnitude over the last 140,000 years. Data comprises all eruptions
from the LaMEVE database as well as additional eruptions from marine sediment cores as described in Section 2.1.**

Here we present a new time series of volcanic stratospheric sulfur injection called “PalVol” (Toohey and Schindlbeck-Belo,
2023) covering the last 140 ka BP that is based on terrestrial and marine tephra records, and includes stochastically generated
135 “synthetic” eruptions to correct for the incompleteness of tephra records, in particular regarding small to medium eruptions
($M < 6$). We provide an ensemble of time series, with each realization providing different timings of events found in the tephra
record according to the uncertainties in the dating of the events, as well as different timings of the synthetic events. Our aim is
not to provide an accurate reconstruction of the actual timing and magnitude of eruptions over the last glacial cycle. This is
potentially even impossible. Rather, our aim is to provide a plausible set of such time series, each of which might approximate
140 a possible true history given the information available and some basic assumptions. While we do not guarantee the accuracy
of the timing and magnitude of specific eruptions, we do aim to produce a time series which represents our best estimate of
the stochastic forcing provided by volcanic eruptions as well as some accuracy in terms of millennial-scale variability in
volcanic forcing.

The paper is organized as follows: in Section 2, the data and methods used to produce the ensemble VSSI time series product are introduced. In Section 3 the ensemble VSSI product is compared to existing ice core-based reconstructions. Discussion and conclusions are included in Section 4.

2 Data and methods

2.1 Tephra Data


The majority of tephra data used in this study is extracted from the LaMEVE database (Crosweller et al., 2012; LaMEVE Version 3), a global compilation of large magnitude eruptions with VEI ≥ 4 (Volcanic Explosivity Index), and/or magnitude 4 that are known from terrestrial deposits and outcrops. The LaMEVE database is publicly available and summarizes information regarding (1) erupted mass/volume and therefore magnitude/VEI, (2) eruption dates as well as the applied dating techniques and uncertainties, (3) the source volcano, (4) eruption parameters (e.g., column height), and (5) rock types. We focused on the time interval from 130,700 years BP to 2014 AD (comprising 1,501 events).

The eruption data taken from LaMEVE is complemented and corrected by records from a suite of marine cores and recent studies (Fig. 1; appendices tables A1, A2), including cores and samples from Central America, offshore the Izu-Bonin volcanic arc, offshore the Cape Verdes, offshore the Kamchatka peninsula, and from the Hellenic Arc. Eruptive volumes and magnitudes for eruptions with no published values have been calculated applying a single isopach approach following the methods described in Schindlbeck et al. (2018b).

The tephra record is characterized by increasing incompleteness back in time (Papale, 2018; Kiyosugi et al., 2015; Brown et al., 2014) (Fig. 2). The reasons for missing eruptions in terrestrial tephra records are mostly erosion and alteration, or burial by younger deposits (e.g., Lavigne et al., 2004; Pollard et al., 2003). However, generally it is thought that the incompleteness is more apparent for smaller magnitude eruptions (e.g., $M=4$) compared to larger magnitudes (e.g., $M \geq 5$) (Brown et al., 2014) since preservation in marine sediments and on land is strongly increasing with the thicknesses of deposits (Wetzel, 2009; Freundt et al., 2021) (Fig. 2). In the marine environment, increasing time for bioturbation and diagenesis (the time in which bio-organisms and chemical interaction with formation waters can actively modify the ash layers physically and chemically) as well as the depositional environment play a role for incompleteness in the records (Hopkins et al., 2020; Freundt et al., 2021). For long records also the plate motions (except for ocean island volcanoes) might be responsible for a decreasing amount of recorded eruptions in the past since most of the eruptive records are coupled with convergent margins and associated arc volcanism (e.g., Schindlbeck et al., 2015). The deeper and older the ash layers in the marine sediments are, the greater the distance between the volcanic source and deposition location at the time of the eruption due to plate motion (e.g., Schindlbeck et al. 2015). This mechanism could and therefore reduce the possibility that smaller eruptions are recorded in the marine sediment archive from convergent margins.

Marine or lacustrine ash layers as well as terrestrial deposits can be either directly or indirectly dated. The techniques comprise radiometric techniques (e.g., radiocarbon (^{14}C), $^{40}\text{Ar}/^{39}\text{Ar}$ mineral dating, zircon dating), orbital tuning of oxygen isotope

curves, sedimentation rates, (Drexler et al., 1980; Kutterolf et al., 2008), which lead to age uncertainties of approximately 1-10% of the estimated age (Fig. 3). The most precise dates are obtained for tephra that can be associated with historical observations of an eruption or with ice core or tree ring signals. The current literature has updated several eruption ages, which have not been included in PalVol v1, but which will be included in the next version. These comprise especially ages obtained by ice core or dendrochronological studies (e.g., Bárðarbunga 877 CE (Plunkett et al., 2023); White River Ash 853 CE (Mackay et al., 2022); Ilopango 431 CE (Smith et al., 2020); Okmok II 43 BCE (McConnell et al., 2020); Aniakchak II 1628 BCE (Pearson et al., 2022); Laacher See 13,006 BP (Reinig et al., 2021)).



185 **Figure 3: Dating uncertainties over time for eruptions included in the LaMEVE data base over the X-Y period, color-coded according to the different dating techniques as provided by Croweller et al. (2012), Cisneros et al. (2021a,b). Gray dashed lines indicate the 1% and 10% error range.**

Magnitude (M) is the preferred measure of eruption size used in the LaMEVE database (Croweller et al., 2012), and is the quantity used here to estimate VSSI (Walker, 1980; Newhall and Self, 1982). Magnitude is a function of erupted mass, which is typically derived from estimates of erupted volumes, classically calculated by drawing isopach maps (areas of equal thickness of the ash layer; e.g., Bonadonna and Houghton, 2005; Fierstein and Nathenson, 1992; Pyle, 1995a) and integrating over area. For eruptions used in this study which are not included in LaMEVE, magnitudes are taken from the respective publications. For the volcanic eruptions described in Derkachev et al. (2020) no volume estimates were available and we therefore calculated minimum volumes applying a single isopach approach (Schindlbeck et al., 2018b) based on the given thickness data. The estimated uncertainty in mass estimates (and thus magnitude) are of the order of 10 to 20% (e.g., Kutterolf et al., 2021b; Klawonn et al., 2014), and depends on the sample size (available outcrops on land and core density in the sea), and uncertainties in rock densities used for converting volumes into masses.

2.2 Ice core-based VSSI

VSSI time series derived from polar ice cores are used in this work, both as a basis for the PalVol VSSI reconstruction methods (see Section 2.3, 2.4) and to validate the new reconstruction through statistical comparisons (Section 3).

For the -500 to 1900 CE period (#256 eruptions), VSSI is taken from the eVolv2k reconstruction (Toohey and Sigl, 2017). These estimates are based on a set of continuous sulfate records from a suite of ice cores from Greenland and Antarctica. As in earlier ice core-based reconstructions (e.g., Gao et al., 2008), tropical eruptions are identified by simultaneous deposition in both Antarctica and Greenland, and signals present in only one hemisphere are assumed to result from extratropical eruptions.

205 VSSI is estimated by applying empirically-derived transfer functions to the ice sheet average sulfate flux values (Toohey and Sigl, 2017, Gao et al., 2008).

Volcanic stratospheric sulfur injection estimates for the Holocene (from 9,500 BCE or 11,500 years BP to 1900 CE; comprising 1,496 eruptions) are available from the HolVol v1.1 reconstruction (Sigl et al., 2022). The method of VSSI reconstruction is very similar to that of eVolv2k, although the number of ice cores used is necessarily smaller since fewer cores cover the full

210 Holocene. Despite the lower number of samples, the HolVol reconstruction shows good agreement with the 2,500-year long eVolv2k record, strengthening confidence in its accuracy (Sigl et al., 2022). Uncertainty in the timing of eruptions in HolVol is estimated to be ± 1 to 5 years on average over the last 2,500 years and better than ± 5 to 15 years for the rest of the Holocene. Reported uncertainties in the VSSI values in HolVol for explosive (i.e., non-effusive) eruptions are typically between approximately 20 and 40% (Sigl et al., 2021).

215 Ice core derived VSSI estimates have been recently reconstructed for the 60-9 ka BP period, which covers the late glacial period as well as the Early Holocene (Lin et al., 2022). Due to the thinning of ice sheets with age, the reconstruction of Lin et al. (2022) is limited to eruptions which produced strong deposition to both Greenland and Antarctica. The reconstruction provides estimates of stratospheric sulfate loading for 85 eruptions with bipolar deposition. To convert the sulfate aerosol mass estimates of Lin et al. (2022) to units of mass sulfur, we divide by a factor of 3 to account for the ratio of molar masses for

220 sulfur (32 g/mol) to sulfate (96 g/mol).

2.3 Deriving VSSI from tephra

VSSI is estimated assuming a linear relationship between VSSI and erupted volume, i.e., a power law relationship between VSSI and eruption magnitude, as used by Pyle et al. (1996) and Metzner et al. (2014). Here, we derive a fit of VSSI to magnitude using ice core-derived VSSI (Section 2.2) and tephra-based magnitudes (Section 2.1) for identified eruptions, as

225 well as recent eruptions for which estimates of sulfur emission are available from satellite instruments (Carn, 2022) (Fig. 4). appendix Table A1 lists eruption data which is used to derive a relationship between VSSI and eruption magnitude. For the period 1980 to 2014, we use sulfur emission estimates from satellite instruments compiled by Carn (2022). Emitted sulfur is matched to the eruptions listed in LaMEVE for this period. For the period before the satellite era, we rely on VSSI estimates and eruption attributions included in the eVolv2k reconstruction (Toohey and Sigl, 2017), supplemented with a few events

230 from the HolVol v1 reconstruction (Sigl et al., 2022). Compared to prior investigations of the VSSI-vs-magnitude relationship (Pyle et al., 1996; Metzner et al., 2014), this data compilation includes more satellite-based estimates, as well including ice core-based estimates, extending the coverage to larger magnitude eruptions, particularly due to the inclusion of recently attributed ice core signals of Okmok (McConnel et al., 2020) and Crater Lake (Sigl et al., 2022) (Fig. 4).

235 2.4. Semi-stochastic VSSI time series generation

In order to correct for the incompleteness of the tephra-based eruption time series, we manufacture a synthetic eruption time series with the same statistical characteristics as an input data set, with timing and magnitudes of eruptions randomized. To construct our supplementary synthetic eruption time series, we draw on the efficient methodology of Bethke et al. (2017), but using as a basis the recent HolVol ice core-based reconstruction over the years -4000 to 1900 CE. The algorithm of Bethke et al. (2017) stochastically produces a new eruption time series based on the eruption magnitude-frequency distribution of the input time series. Our time range is chosen so as to include a large period to enhance the statistical basis of the frequency distribution, while excluding the early to mid-Holocene (11 to 6 kaBP), for which the eruption frequency is amplified compared to the mid-to-late Holocene, at least in part to eruptionpo. When an eruption is randomly generated, characteristics of the eruption from HolVol, including VSSI, eruption region (tropical, NH extratropical or southern hemisphere (SH) extratropical), and month are copied into the constructed synthetic time series. For eruptions that are unidentified in the HolVol base data (which is the majority of events), eruption parameters month and precise latitude (within three latitude ranges) are unknown, and set to default values. In our synthetic eruptions, the eruption month is randomized uniformly across the calendar year, and the eruption latitude is randomized within the identified tropical, NH and SH bands using the probability density of the LaMEVE data set between 10 ka BP and the present. We repeat this process 100 times to produce 100 synthetic eruption time series.

To produce an ensemble of final VSSI time series therefore requires merging each synthetic time series--based on the statistics of the HolVol ice core data--with the evidence from the tephra record. Merging the two eliminates the decreasing eruption frequency backwards in time present in the tephra record, assuming that this characteristic of the tephra record is a product of incompleteness. It also assumes that the true eruption frequency distribution is approximately static with time. For each of the 100 ensemble members, the synthetic and tephra records are merged so that each event in the tephra record is inserted into the synthetic record while also removing from the synthetic record a synthetic event with closest matching magnitude within a window of 500 years. To represent the dating uncertainty in the tephra-based events in the ensemble of forcing time series, and also avoid clumping of eruptions around intervals of 1000's of years due to the limited resolution of the reported dates of the tephra-based events, thus potentially creating an artificial millennial-scale periodicity in the radiative forcing, we add a random, normally distributed perturbation to the reported date of the tephra-based eruptions, based on the estimated dating uncertainty (see Section 2.1 and figure 3). Thus, the dates of tephra-based eruptions in our reconstruction match the original dates for an eruption within the reported uncertainty. The dating perturbation is performed separately for each of the 100 ensemble members, so that the date assigned differs in each realization of the data product, and the spread in dates between realizations depends on the dating uncertainty of the tephra event.

265 **2.5. Aerosol optical properties**

The impact of volcanic aerosol radiative forcing is implemented in climate models in different levels of complexity. The simplest models take as input variations in the top of atmosphere radiative flux anomalies (W/m^2), which represent the net effect of scattering and absorption of radiation by stratospheric aerosol (e.g., the energy balance model used in Pages-2k-Consortium, 2019). Comprehensive climate models, on the other hand, may require detailed optical properties of the aerosol, as a function of latitude, height, wavelength and time, to be used in the radiative calculations of the model (as e.g., for the Max Planck Institute Earth System Model used in Bader et al., 2019).

To produce timeseries of the radiative impacts of past eruptions, including the detailed optical properties required by comprehensive models, we use the Easy Volcanic Aerosol (EVA) forcing generator (Toohey et al., 2016). This simple model takes as input the eruption timing, VSSI estimates, and eruption latitude, and produces aerosol extinction, single scattering albedo and scattering asymmetry factor as a function of latitude, altitude, wavelength and time. These variables are the result of a simple 3-box model of stratospheric transport, scaling approximations between aerosol mass and AOD at $0.55 \mu\text{m}$, and Mie theory which describes the scattering of radiation by spheres. The overall impact of stratospheric aerosol on the Earth's energy balance is roughly proportional to the AOD in the visible part of the electromagnetic spectrum, so it is common to illustrate the volcanic forcing simply as the AOD at $0.55 \mu\text{m}$, either as a function of latitude, or as a global (area weighted) mean.

2.6. A note on date formatting

Throughout the manuscript we use two conventions with regarding to dates. For periods extending no further than around 2,500 years into the past, we use a variation of the ISO8601 format, which is very similar to the usual “Common Era” (CE) system, differing though in the sense that the ISO8601 system includes a year 0, while the Common Era system does not. The two systems therefore differ by one year for years before 1 CE, for example, 44 BCE would correspond to the year -43 in the ISO8601 system.

For dates further in the past, we use the widely used “Before Present” (BP) system, which indicates the number of years before 1950.

290 **3. Results**

3.1 VSSI-to-eruption magnitude relationship


The relationship between eruption magnitude and VSSI as observed by satellite instruments over the most recent decades and estimated from ice cores over the last 2500 years and Holocene is shown in Figure 4. Satellite observations offer predominantly information on the VSSI from M=4 to M=5 eruptions: the only eruptions larger than M=5 that have been observed directly are

295 the 1991 Pinatubo ($M=6.1$) and Cerro Hudson ($M=5.8$) (Crossweller et al., 2012 and references therein). For $M=4$ eruptions, the VSSI observed by satellites covers a range from 0 to approximately 1 TgS. The largest values of VSSI observed from satellites are associated with the eruptions of Pinatubo ($M=6.1$, $VSSI=7.6$ TgS) and El Chichón ($M=5.1$, $VSSI=4.0$ TgS). Eruptions from the ice core reconstructions extend the range of eruption magnitudes included in the analysis, we use here 24 eruptions, of which 21 have magnitude greater or equal to 5.0.

300 The VSSI values from satellite instruments and ice core reconstructions taken together show a proportionality with eruption magnitude: larger eruption magnitudes lead generally to larger VSSI values. Following Pyle et al. (1996), we fit a power law relation to the data to obtain a best fit relationship for VSSI in TgS: $VSSI = (1.67 \times 10^{-5}) \times 6.27^M$

This fit is very similar to fits produced by Pyle et al. (1996) based solely on satellite data from the period 1979–1993, and also somewhat similar to the fit presented by Metzner et al. (2014), who based their fit on petrologically obtained sulfur emission
305 values from Central American eruptions (see Figure 4).

There is clearly a significant amount of scatter around the line of best fit. This scatter can be the result of uncertainties in the eruption magnitude estimates as well as the VSSI values, but is likely dominated by actual variability in sulfur emission between different eruptions with similar magnitude (e.g., Andres et al., 1993; Sigurdsson, 1990), and on the proportion of emitted sulfur that is injected into the stratosphere. The strong apparent scatter for $M=4$ eruptions in Figure 4 represents a
310 modest scatter in term of absolute numbers, with VSSI ranging from approximately 0 to 1 TgS, reflecting variations in the sulfur output as well as the modulating effect of plume heights which in many cases are not high enough to bring sulfur into the stratosphere. For $M \geq 5$, the proportionality between VSSI and magnitude is more compact, the majority of events falling within approximately an order of magnitude around the best fit line. Notable outliers for which the VSSI is larger than the best fit line relationship include Laki (1783), for which the Greenland VSSI estimate has been suggested to potentially include a
315 significant amount of tropospheric aerosol (Lanciki et al., 2012), and Hekla (1766), another Icelandic eruption for which the same scenario may hold. In contrast, a VSSI much smaller than expected, based on the best fit relationship, is seen for the Millennium eruption of Changbaishan (940 CE), which has been discussed previously and may be due to some combination of sulfur-poor magma (Horn and Schmincke, 2000) and short stratospheric lifetime due to injection height or seasonal atmospheric dynamics (Iacovino et al., 2016). Accordingly, VSSI estimated from eruption magnitudes should be understood
320 to have significant uncertainty for any individual eruption.



325 **Figure 4: Volcanic stratospheric sulfur injection (VSSI) derived from ice cores (Toohey and Sigl, 2017; Sigl et al., 2021) and satellite observations (Carn, 2022) as a function of eruption magnitude from the LaMEVE database (Crosweller et al., 2012). A least squares power law best fit is shown in black, compared to similar fits from Pyle et al. (1996) and Metzner et al. (2014). A 1-sigma uncertainty range to the fit is shown as gray shading.**

3.2 Common Era

330 During the Common Era (0-2000 CE) the eruption rate as recorded in the tephra records is larger than that of the ice core records during the chosen base period (-4000 to 1400 CE). Therefore, we can base the PalVol reconstruction purely on the tephra records without any synthetic events that could otherwise be included to compensate for undersampling by the tephra (Fig. 5).

335 The timeseries of VSSI calculated from the tephra magnitude estimates shows reasonable agreement with the ice core-based eVolV2k reconstruction (Fig. 5). The Rinjani (Samalas) eruption of 1257 CE, which produced the largest VSSI in eVolV2k (59.4 TgS) is well reproduced in the tephra time series, with an estimated VSSI of 69.4 TgS. The large eruptions of Tambora (1815) and Krakatau (1883) are apparent in both time series, albeit larger by a factor of approximately two in the tephra-based estimates compared to eVolV2k. The tephra record includes a large VSSI associated with the Changbaishan eruption of 942, which is not found in ice cores, which is likely due to the sulfur-poor content of the erupted magma (Horn and Schmincke, 2000). Other prominent events in the 1st millennium CE of the tephra VSSI time series, including TBJ (450±30 CE, VSSI=38.9

340 TgS), Taupo (230 ± 16 CE, $VSSI=57.2$ TgS) and Ambrym (190 ± 135 , $VSSI=47.2$ TgS) are likely to be responsible for ice core signals within the dating uncertainty of the tephra events.





Figure 5: VSSI estimates for the Common Era from (a) tephra and (b) ice cores (eVolv2k, Toohey and Sigl, 2017). (c) Centennial total VSSI for both data sets, with the correlation of the two centennial time series.

345 Given the large uncertainties in VSSI estimated from tephra (and the non-negligible uncertainties in the same value from ice
cores), we would not and should not expect the VSSI values for individual eruptions to agree to a precision better than an order
of magnitude. When multiple eruptions are averaged together, we expect the errors to compensate to some degree, and
cumulative VSSI values to become more reliable. Centennial total VSSI is shown for the tephra reconstruction and eVolv2k
in Figure 5c. The correlation between the two centennial time series is notable, with an overall correlation coefficient of
350 $R=0.50$. The correlation is especially good in the 2nd millennium, with both reconstructions showing elevated mean VSSI
amounts for the 13th and 19th centuries, as well as more modest elevated values for the 15th and 17th centuries. While the
tephra time series misses the elevated VSSI values of the 6th Century, there is some agreement in the elevated VSSIs during
the 2nd and 3rd centuries.

3.3 Holocene

355 VSSI reconstructed from tephra is shown in Figure 6 for the Holocene period (roughly 11 to 0 ka BP) and compared to the HolVol ice core-based reconstruction. Both data sets show an increase in large magnitude VSSI in the early Holocene, around 11 to 7 ka BP. Particularly, both data sets include four quite large events between 9 and 7 ka BP, with the tephra-based values reaching approximately 100 TgS while the ice core values reach up to over 150 TgS.



360 **Figure 6: Comparing tephra and ice core-based VSSI time series over the Holocene. Upper panel: VSSI time series derived from tephra records. Lower panel: HoVol ice core-based VSSI time series (Sigl et al., 2022). Gray shading shows the period used to base eruption statistics for the synthetic time series.**


In a next step we compare the statistical characteristics of the tephra data, the PalVol reconstruction ensemble, and the HoVol
365 ice core-based VSSI reconstruction over the period of the Holocene on millennial timescales. Comparing first the number of events in the tephra and ice core data sets (Fig. 7a) we see that the number of tephra events drops rapidly with increasing age, from ~275 events/ka from 0-1 ka BP to less than 100 events/ka before 2 ka BP, to less than 50/ka in the early Holocene, consistent with prior analyses of the LaMEVE database (Brown et al. 2014).

From roughly 0 to 9 ka BP, the ice core number of events holds approximately steady at around 100 events per ka, which
370 increases during the deglaciation period, reaching a maximum of ~160 events per ka around 11 ka BP. In the most recent 2 ka, the frequency of tephra events is larger than the frequency of ice core events. This is likely due in part to overcounting of tephra events, due to overestimation of the magnitude of events (increasing the absolute number of $VEI \geq 4$ events included in LaMEVE). We speculate that another source of discrepancy may be undercounting of $VEI \geq 4$ eruptions by ice cores, since not all eruptions may leave traces in ice cores if not explosive enough or sulfur poor. Before 0 CE, the incompleteness of the

375 tephra record is clear compared to the ice cores, and the degree of incompleteness increases roughly linearly until around 6 ka BP. There may be a weak local maximum in tephra events around 9 to 10 ka BP CE in agreement with the ice core increase here, but the difference in tephra events between this period and the local minimum at 7 to 8 ka BP is rather small.

Next, we compare the cumulative VSSI per millennium derived directly from the tephra data compared to the HoIVol VSSI database (Fig. 7b). The tephra-derived VSSI per ka is, in all millennia before 2 ka BP, smaller than the HoIVol values, which
 380 is perhaps unsurprising given the much smaller number of events in the tephra database compared to the ice cores. In contrast, over the Common Era (0-2 ka BP), the tephra VSSI/ka is larger or roughly equal to the ice core VSSI/ka.

Over the Holocene, the average VSSI per millennium from HoIVol is 638 TgS, while that from the pure tephra data is 231 TgS per millennium.



385 **Figure 7: Millennial-scale variations in eruptive characteristics over the Holocene in ice core (HoIVol), tephra and PalVol time series. (a) Number of volcanic events per millennium in each dataset, (b) cumulative VSSI per millennium, (c) average VSSI per event, (d) histogram of correlation coefficients calculated between the ensemble of semi-synthetic millennial cumulative VSSI time series with the HoIVol time series. Each bar indicates the fraction of all ensemble members with correlation coefficients between the values defining the edges of the bar along the horizontal axis. In (d), the correlation between the tephra time series and HoIVol is indicated by the orange cross. In panels a-c, gray lines indicate values for each of the 100 stochastic realizations, and the black line the ensemble mean.**

390

395 Despite clear differences between the tephra and ice core VSSI reconstructions for specific events, and a bias with tephra
showing lower values for most millennia of the Holocene, there is correlation between the two datasets in terms of millennial
VSSI over the Holocene, with both showing larger VSSI/ka values in the early Holocene (11 to 7 ka BP) compared to the mid
Holocene (2 to 7 ka BP). The correlation coefficient between the millennial VSSI totals for the tephra and ice core-based
datasets over the 10 to 2 ka BP CE period is $R=0.46$. This correlation comes about despite the strong undercounting of the
tephra database in the early Holocene. Evidently, although the number of events captured by the tephra data sets is small, the
400 events that are counted tend to be the larger eruptions, which contribute the most to the VSSI millennial sums. Figure 7c shows
the average VSSI per event as a function of millennium, which shows a similar structure for both the tephra and ice core data,
with larger average VSSI per event in the early Holocene compared to the mid and late Holocene. This implies that the increase
in VSSI/ka in the early Holocene arises from an increase in the frequency of large eruptions, which is evident in both the ice
core and tephra data.

405 Now we compare the semi-stochastic PalVol VSSI reconstruction, constructed by adding stochastically generated events to
the tephra data, to the statistics of the HolVol reconstruction. By construction, the PalVol reconstruction includes a relatively
constant number of events per millennium, around 100 events/ka (Fig. 7a), in agreement with the frequency of events in the
HolVol reconstruction over the chosen base period -4000 to 1900 CE (roughly 6 to 0 ka BP). The number of events/ka for
each individual ensemble member of the reconstruction will vary around this number based on the stochastic event generation,
410 and the inter-millennial variance matches well with the variance of the HolVol reconstruction over the base period. The PalVol
reconstruction clearly does not include the increase in events/ka in the early Holocene that is seen in HolVol (Fig. 7a). This is
by construction, since we do not adjust the eruption frequency probability with time.

Nonetheless, in Figure 7b, we see that after adding the stochastically generated events to create the PalVol forcing
reconstruction, the agreement of the millennial distribution of VSSI/ka with the HolVol reconstruction improves compared to
415 the pure tephra time series. First, the overall bias is reduced (but not eliminated), with an average VSSI/ka of 505 TgS for
PalVol compared to 639 TgS for HolVol. Secondly, since more stochastically generated events are added to the early Holocene
compared to the late Holocene to make up for the larger bias in event frequency (see Fig. 7a), the VSSI/ka is boosted more in
the early Holocene. This improves the correlation: the mean PalVol VSSI/ka time series shows a correlation of $R=0.73$ with
HolVol. A histogram of the correlation coefficient between each individual PalVol VSSI time series with HolVol (Fig. 7d)
420 shows the largest proportion of ensemble members have a correlation of 0.7-0.8. We conclude that over the Holocene, the
addition of stochastic events to the tephra data improves resulting time series in comparison to ice core derived reconstructions,
both by reducing the low bias in the tephra data, and improving the inter-millennial variability of VSSI.

3.3 Last glacial cycle

The PalVol VSSI reconstruction is compared to the ice core-based VSSI reconstruction of Lin et al. (2022) in Figure 8. The
425 Lin et al. (2022) events used here constitute 85 eruptions with matched bipolar signals with deposition of >20 kg/km² in

Antarctica and $>10 \text{ kg/km}^2$ in Greenland between 60–9 ka BP. Tephra events in this period have a median dating uncertainty of 2300 years (900-4,600 years 0.25-0.75 interquartile range), therefore, we don't expect clear temporal matches to the ice core events for the reported dates of tephra events. Despite this, we find a decent agreement in the VSSI frequency distribution of the largest events in both time series for the period overall. For example, we find 11 events with VSSI $>100 \text{ TgS}$ in the Lin et al. (2022) time series, while in the PalVol record for the same 60-9 ka BP period, we have 10 such events. The largest VSSI signal in the PalVol reconstruction over this period is associated with the 27 ka BP Taupo Oruanui eruption with $M=8.1$, leading to an estimated VSSI of 480 TgS, a factor of almost 5 times greater than the ice core derived value of 127 TgS. This represents an overestimate compared to the ice core-based reconstruction for this event, which is not overly surprising given the uncertainty in the tephra-based reconstruction method.

435





Figure 8: Volcanic stratospheric sulfur injection from PalVol (top) and the ice core-based reconstruction of Lin et al., 2022, for the 60-9 ka BP period. Stochastically generated PalVol VSSI (in gray) are from a single ensemble member, while tephra-based estimates (red) are shown at the reported dates (not randomized).

440 Reconstructed VSSI is shown in Figure 9 for the full PalVol reconstruction period. Individual eruptions are shown in panels a and b, for which the decreasing number of detected eruptions with age, and the resulting increasing number of synthetic eruptions included is apparent. The four largest VSSI estimates in PalVol exceed the range plotted on Figure 9: the corresponding VSSI values are listed in Table 1 as part of the top twenty VSSI estimates. The eruption of Toba is the largest eruption of the past 140,000 years, and results in an estimated 3,000 TgS VSSI using our method, with a wide estimated uncertainty range spanning 310 to 29,000 TgS. Prior estimates of Toba's sulfate emission fall within a very broad range, multiple studies were compiled by Oppenheimer (2002) to define a range of 35 to 3,300 TgS. More recently, Costa et al. (2014)

445

estimated Toba's sulfur emission as 850-1,750 TgS, which falls within our uncertainty range, while Crick et al., (2021) estimated a range of 72-233 TgS, which falls outside our uncertainty range. It must be stressed that our estimates of VSSI for the strongest eruptions is based on an extrapolation of the VSSI-to-magnitude relationship beyond what has been observed, and non-linearities in the physical processes (e.g., plume collapse) are not considered here.



455 **Figure 9: Tephra-based volcanic stratospheric sulfur injection (VSSI) reconstruction. (a) VSSI based on the pure tephra record. (b) A semi-synthetic VSSI time series, based on merging a stochastic synthetic record with the tephra reconstruction. (c) Millennial VSSI rates for the tephra, TephraSynth and HolVol reconstructions.**

The Los Chocoyos eruption of Atitlan, ~75 ka BP, has the same magnitude as the Taupo eruption, and therefore the same estimated VSSI of 480 TgS with a range of 57 to 4,000 TgS. This value is very similar to the 343 TgS estimated by Metzner et al. (2014), although the method used there was very similar to that used here.

460 Our estimate for the Changbaishan eruption (946 CE) is 130 TgS, while Horn and Schmincke (2000) estimated a release of 5.7 TgS, and Iacovino et al. (2016) a release of 45 TgS. A lack of strong ice core sulfate signals around the documented time

of the eruption has been used as evidence that the sulfur emission from Changbaishan must have been quite minor (Oppenheimer et al., 2017), and likely much lower than the estimate here based only on the eruption magnitude.


465 **Table 1: Estimated VSSI for the top 20 largest eruptions of the past 140,000 years based on the LaMEVE database.**

Volcano	Year	Year BP	Magnitude	VSSI (TgS)	Min VSSI (TgS)	Max VSSI (TgS)
Toba	-73050	75000	9.1	3000	310	29000
Taupo	-23410	25360	8.1	480	57	4000
Atitlán	-73050	75000	8.1	480	57	4000
Aira	-27205	29155	7.9	330	40	2700
Asosan	-85050	87000	7.7	230	29	1800
Campi Flegrei	-37900	39850	7.5	160	20	1200
Ata	-106050	108000	7.5	160	20	1200
Pacaya	-122050	124000	7.5	160	20	1200
Changbaishan	946	1004	7.4	130	17	1000
Long Island	-17328	19278	7.4	130	17	1000
Opala	-42534	44484	7.4	130	17	1000
Okataina	-45550	47500	7.4	130	17	1000
Maninjau	-50050	52000	7.4	130	17	1000
Asosan	-121050	123000	7.4	130	17	1000
Santorini	-1610	3560	7.3	110	14	840
Kikai	-5284	7234	7.3	110	14	840
Kurile Lake	-6437	8387	7.3	110	14	840

Gorely	-37031	38981	7.3	110	14	840
Nemo Peak	-43050	45000	7.3	110	14	840
Shikotsu	-43883	45833	7.3	110	14	840

Figure 9c shows the millennial average VSSI for the PalVol reconstruction as well as the pure tephra data and the HolVol ice core reconstruction. Each PalVol ensemble member is shown in gray: these reconstructions generally show millennial average VSSI ranging from approximately 0.3 to 0.7 TgS/year, consistent with the mean value and variability from HolVol. Exceptions occur in millennia which contain the strongest eruptions, for which the millennial average VSSI can increase by a factor of 2 or more. The ensemble average PalVol millennial VSSI is approximately constant with age except for the perturbations due to the largest eruptions, and some small increases due to enhanced numbers of detected strong eruptions, for example in the 55-40 ka BP period.

One realization of PalVol VSSI time series is visualized in Figure 10, with VSSI magnitudes plotted as “bubbles” as a function of ka BP and latitude. Many characteristics of the LaMEVE database are apparent: the decreasing eruption sampling with increasing age, the dependence of this sampling with latitude, with a generally more complete sampling in the NH mid latitudes. This figure illustrates the distribution of the tephra-based and synthetic eruptions in time and latitude, and specifically the latitude distribution of the synthetic events based on the latitudinal distribution of the LaMEVE eruptions.



480 **Figure 10: The PalVol global VSSI product, with the coordinates of each circle representing the time and latitude of an eruption, and the circle size the VSSI in TgS. Circle colors signify the type of event, either detected events compiled in LaMEVE (red) or additional records from marine sediment cores (purple), or the stochastic synthetic events used here to fill the record (gray).**

Data availability

The PalVol volcanic stratospheric sulfur injection stratospheric aerosol optical depth datasets described herein are available
485 through the World Data Center for Climate in netCDF format (<https://doi.org/10.26050/WDCC/PalVolv1>; Toohey and Schindlbeck-Belo, 2023).

4. Conclusions and Discussion

We have produced PalVol VSSI reconstructions that cover the last glacial cycle (the last 140 kyrs). The ensembles incorporate tephra data of past eruptions, as well as stochastically generated events to attempt to correct for the undersampling of events
490 with increasing age. VSSI values are derived from the eruption magnitude, which is itself estimated based on the thickness of tephra layers in the surrounding of a volcano. Importantly, the VSSI values for individual eruptions have significant uncertainties, as we (and prior works; e.g., Andres et al., 1993; Sigurdsson, 1990) showed that the amount of sulfur released by an eruption can vary by orders of magnitude for any given eruption magnitude (Fig. 4). Especially for M=4 eruptions the VSSI varies significantly since some reach the stratosphere and some do not (Fig. 4), which depends on their actual plume
495 height, but also on latitude. Uncertainties in VSSI are included in our reconstruction as upper and lower bounds on the VSSI for each eruption, based on the uncertainty in our derived magnitude to VSSI relationship. We note as well that this relationship may not be constant in time, and indeed changes may occur, e.g., to changes in the height of the tropopause or in stratospheric circulation from glacial conditions to interglacials. This would be similar to predicted changes in future volcanic radiative forcing due to changes in the tropopause height due to climate change (Aubry et al., 2016). This is something we did not take
500 into account in our reconstructions. Nonetheless, although the VSSI for any specific eruption is quite uncertain, we expect that due to the observed relationship between VSSI and magnitude, this will be compensated when averaged over a sufficiently large set of eruptions, and therefore the tephra data may contain information on variations in VSSI on long time scales.

Furthermore, the PalVol reconstruction shows good agreement with the ice core based HolVol VSSI time series in terms of millennial variations in cumulative VSSI, but not particularly in terms of the timing and magnitude of individual events.
505 This agreement arises from the shared increase in cumulative VSSI in the early Holocene compared to the mid and late Holocene. In both data sets, this arises from the increased frequency of relatively large eruptions. Evidently, while the tephra time series is incomplete during this period, the sampling of large events is good enough to detect the increase in the amount of large eruptions, which translate into increases in millennial cumulative VSSI.

We assumed a constant eruption frequency distribution, but there is evidence that variations in eruption frequency are driven
510 by changes in mass distribution of ice sheets and respective sea level changes. The PalVol reconstruction includes a small

increase in VSSI/ka in the early Holocene, qualitatively consistent with observations from the ice cores, but with an amplitude which is actually smaller. Therefore, we propose that future iteration of stochastic forcing could take into account such variations, for example by using sea level reconstructions as a basis to estimate variations in eruption frequency.

515 The PalVol reconstruction takes the form of an ensemble of realizations, where each realization differs in terms of timing and size of the stochastically generated events, and the timing of the events taken from the tephra corresponding to the uncertainty in the tephra dating. While any single realization is unlikely to be a very accurate reconstruction of the true history of VSSI, it is quite possible that some realizations include aspects that are realistic. Therefore, the ensemble of VSSI time series represents a probability distribution of the probable forcing from volcanic eruptions over this period, given the information we currently have from proxies.

520 We emphasize that users wanting the most accurate reconstruction of VSSI over the last glacial cycle could consider using a merged product, for example by concatenating the HolVol ice core time series with PalVol for the period, which occurs before the beginning of HolVol. Future improvements to the PalVol reconstruction will be possible with the addition of new tephra data, improvements in the dating, and magnitude estimates of volcanic events.

Appendices

525 Table A1: Compilation of matched eruption magnitudes from LaMEVE with volcanic stratospheric sulfur injection (VSSI) estimates from satellite measurements or ice core-based reconstructions. Volcano. eruption year and magnitude taken from the LaMEVE database. VSSI in TgS is taken from 3 sources. indicated by last column: 1: from satellite observations compiled by Carn et al. (2022); 2: the eVol2k ice core reconstruction (Toohey and Sigl, 2017); 3: the HolVol ice core reconstruction (Sigl et al., 2022).

530

Volcano	Eruption Year	Magnitude	VSSI (TgS)	Source
Puyehue-Cordón Caulle	2011	4.8	0.1	1
Eyjafjallajökull	2010	4	0.0025	1
Merapi	2010	4	0.15	1
Sarychev Peak	2009	4	0.6	1
Chaitén	2008	4.2	0.007	1
Reventador	2002	4.6	0.042	1
Láscar	1993	4.4	0.225	1
Spurr	1992	4	0.4	1

Hudson. Cerro	1991	5.8	2	1
Pinatubo	1991	6.1	7.597	1
Kelut	1990	4.1	0.075	1
Redoubt	1989	4.3	0.07	1
Augustine	1986	4	0.01	1
Chikurachki	1986	4	0.375	1
Kliuchevskoi	1986	4.5	0.005	1
Chichón. El	1982	5.1	4.045	1
Galunggung	1982	4.8	0.3725	1
Alaid	1981	4.7	0.55	1
Pagan	1981	4.4	0.16	1
St. Helens	1980	4.8	0.455	1
Okataina	1886	5.3	0.74	2
Krakatau	1883	6.5	9.34	2
Askja	1875	4.9	0.67	2
Makian	1861	4.7	4.53	2
Cosigüina	1835	5.8	9.48	2
Tambora	1812	7	28.08	2
Grímsvötn	1783	5	20.81	2
Hekla	1766	4.3	2.52	2
Katla	1755	5	1.18	2
Shikotsu	1739	5.6	3.44	2
Katla	1721	5	0.81	2
Gamkonora	1673	5	4.67	2
Shikotsu	1667	5.4	3.48	2
Huaynaputina	1600	6.1	18.85	2
Bárdarbunga	1477	5.4	5.12	2
Rinjani	1257	7	59.42	2
Katla	934	5.7	16.23	2
Bárdarbunga	870	5.4	3.99	2

Churchill	847	6.1	2.48	2
Ilopango	450	6.7	10.91	3
Okmok	-76	6.7	33.91	3
Aniakchak	-1645	6.9	22.77	3
Crater Lake	-5677	6.8	107.9	3
Khangar	-5699	6.3	30.05	3

Table A2: Additional data from a suite of marine cores and recent studies including cores samples from Central America, offshore the Izu-Bonin volcanic arc, offshore the Cape Verdes, offshore the Kamchatka peninsula.

Sample name	Age [Ma]	Source	Tephra Volume [km ³]	DRE [km ³]	Magnitude
Schindlbeck et al., 2016					
344-U1413B-1H-1-9-11	0.0011	Rincon de la Vieja tephra	0.85	0.40	4.92
334-U1378B-3H-6 29-30	0.027	Rincon	2.56	1.19	5.40
344-U1413A-1H-4-16-18	0.033	Poas	2.04	0.95	5.20
334-U1378B-4H-5 51-53	0.036	Ilopango, TB4			
344-U1381C-2H-1-97-99	0.06	Fontana			
344-U1381C-2H-1-105-107	0.06	Las Sierras	2.02	0.94	5.30
170-1039B-2H-3-62-68	0.087	Poas Lapilli Tuff	1.67	0.78	5.20
334-U1378B-5H-4 106-108	0.076	Poas Plantanar	0.32	0.15	4.50
344-U1381C-2H-3-39-41	0.09	Las Sierras	5.05	2.36	5.40
334-U1378B-6H-4-93-95	0.116	Apoyo	7.66	3.58	5.50
Derkachev et al., 2020	Age [ka]	Source	Tephra Volume [km³]	DRE [km³]	Magnitude
WP1†	8.7	Karymskyc caldera	0.63	0.29	4.80
WPL1	~20	?	0.16	0.07	4.10
WP2	~28	EVF	9.44	4.41	5.90
WP3	~36	EVF	0.63	0.29	4.80

WP4	~39	Gorely	7.69	3.59	5.70
WPL2	~76	Opala?	0.70	0.33	4.80
WP5	~80	Gorely	10.49	4.90	6.00
WPL3	~88	?	0.39	0.18	4.60
WP6	~97	Opala?	2.24	1.04	5.40
WP7	~103	EVF	14.31	6.68	6.20
WP8	~107	EVF	1.26	0.59	5.10
WP9	~125	Gorely?	14.68	6.86	6.20
*volumes are calculated for a 30° single isopach following the methods described in Schindlbeck et al., 2018a					
	Age [Ma]	Source			Magnitude
Schindlbeck et al., 2018a					
350-U1436A-1H-1-48-50	0.0083	Sumisu Knoll?			4.00
350-U1436A-1H-1-84-86	0.0145	Sumisu/Agoashima?			4.30
350-U1436A-1H-1-124-126	0.0217	Sumisu/Agoashima?			4.60
350-U1436A-1H-2-24-27	0.03	Sumisu Knoll?			4.00
350-U1436A-1H-2-42-44	0.0337	Sumisu Knoll?			4.50
350-U1436A-1H-3-79-81	0.0725	Agoashima?			5.00
350-U1436A-2H-1-25-27	0.088	Hachijojima?			4.40
350-U1436A-2H-2-16-18	0.1039	Hachijojima/Sumisu/Agoas			4.60
350-U1436A-2H-2-46-48	0.1058	Agoashima/Hajijojima?			4.90
350-U1436A-2H-2-68-70	0.1072	Sumisu Knoll?			4.20
350-U1436A-2H-3-90-92	0.118	Hachijojima/Torishima?			4.90
350-U1437B-1H-2-37-38	0.0193	Sumisu Knoll?			5.50
350-U1437B-1H-3-18-20	0.0263	Sumisu Knoll?			5.00
350-U1437B-2H-4-46-48	0.1098	Torishima/Hachijojima? Zoned			5.20
350-U1437B-2H-4-72-74	0.1107	Torishima/Hachijojima?			5.20
350-U1437B-2H-4-128-130	0.1127	Torishima/Hachijojima?			5.50

350-U1437B-2H-5-117-119	0.1232	Sumisu Knoll?			4.70
350-U1437B-2H-5-144-145	0.1264	Minami Hachijo?			4.70
350-U1437B-2H-6-25-27	0.1301	Myojin Knoll			4.60
Eisele et al., 2015	Age [ka]	Source		DRE [km³]	Magnitude
C1	17	Cadamosto		0.21	4.30
C2	18-20	Fogo			5.00
C3	21-28	Fogo			5.00
C4	24	Fogo		1.44	5.40
C5	26-28	Fogo			5.00
C6	37	Fogo			5.00
C7	40	Cadamosto		0.12	4.20
C8	50-53	Fogo			5.00
C9	61	Fogo		0.97	5.30
C10	88	Fogo		0.95	5.30

Author contributions

535 JCSB, MT, MJ, SK, KR: conceptualization; JCSB, MT: writing the original draft; MJ, SK, KR: writing- reviewing and editing; JCSB, MT: data and storage; MT: data analyses; MT, KR, SK: funding acquisition

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

540 Research has been funded by the PalMod project (www.palmod.de, subproject no. 01LP1926E). This work benefitted greatly as a result of the authors' participation in the Past Global Changes (PAGES) Volcanic Impacts on Climate and Society (VICS) working group which in turn received support from the Swiss Academy of Sciences and the Chinese Academy of Sciences. We would like to thank the DKRZ for their help with publishing the data.

References

- 545 Abbott, P. M., Niemeier, U., Timmreck, C., Riede, F., McConnell, J. R., Severi, M., Fischer, H., Svensson, A., Toohey, M., Reinig, F., and Sigl, M.: Volcanic climate forcing preceding the inception of the Younger Dryas: Implications for tracing the Laacher See eruption, *Quaternary Sci Rev*, 274, 2021.
- Andres, R. J., Rose, W. I., Stoiber, R. E., Williams, S. N., Matías, O., and Morales, R.: A summary of sulfur dioxide emission rate measurements from Guatemalan volcanoes, *Bull. Volcanol.*, 55, 379-388, 10.1007/BF00301150, 1993.
- 550 Aubry, T. J., Jellinek, A. M., Degruyter, W., Bonadonna, C., Radić, V., Clyne, M., and Quainoo, A.: Impact of global warming on the rise of volcanic plumes and implications for future volcanic aerosol forcing, *J. Geophys. Res.: Atmospheres*, 121, 13,326-313,351, 2016.
- Bader, J., Jungclaus, J., Krivova, N., Lorenz, S., Maycock, A., Raddatz, T., Schmidt, H., Toohey, M., Wu, C.-J., and Claussen, M.: Global temperature modes shed light on the Holocene temperature conundrum, *Nat. Comm.*, 11, 4726, 10.1038/s41467-020-18478-6, 2020.
- 555 Baldini, J. U., Brown, R. J., and McElwaine, J. N.: Was millennial scale climate change during the Last Glacial triggered by explosive volcanism?, *Sci. Rep.*, 5, 1-9, 2015.
- Bay, R., Bramall, N., Price, P., Clow, G., Hawley, R., Udisti, R., and Castellano, E.: Globally synchronous ice core volcanic tracers and abrupt cooling during the last glacial period, *J. Geophys. Res.: Atmos.*, 111, 2006.
- 560 Bethke, I., Outten, S., Otterå, O. H., Hawkins, E., Wagner, S., Sigl, M., and Thorne, P.: Potential volcanic impacts on future climate variability, *Nat. Clim. Change*, 7, 799-805, 2017.
- Bonadonna, C. and Houghton, B. F.: Total grain-size distribution and volume of tephra-fall deposits, *Bull. Volcanol.*, 67, 441-446, 2005.
- Brown, S. K., Crossweller, H. S., Sparks, R. S. J., Cottrell, E., Deligne, N. I., Guerrero, N. O., Hobbs, L., Kiyosugi, K., Loughlin, S. C., and Siebert, L.: Characterisation of the Quaternary eruption record: analysis of the Large Magnitude Explosive Volcanic Eruptions (LaMEVE) database, *J. Appl. Volcanol.*, 3, 1-22, 2014.
- 565 Büntgen, U., Arseneault, D., Boucher, É., Churakova, O. V., Gennaretti, F., Crivellaro, A., Hughes, M. K., Kirilyanov, A. V., Klippel, L., and Krusic, P. J.: Prominent role of volcanism in Common Era climate variability and human history, *Dendrochronologia*, 64, 125757, 2020.
- 570 Carn, S.: Multi-Satellite Volcanic Sulfur Dioxide L4 Long-Term Global Database V4, Greenbelt, MD, USA, Goddard Earth Science Data and Information Services Center (GES DISC), Accessed: March 18, 2022, 10.5067/MEASURES/SO2/DATA405, 2022.
- Cisneros de León, A., Schindlbeck-Belo, J. C., Kutterolf, S., Danišik, M., Schmitt, A. K., Freundt, A., Pérez, W., Harvey, J. C., Wang, K. L., and Lee, H. Y.: A history of violence: magma incubation, timing and tephra distribution of the Los Chocoyos supereruption (Atitlán Caldera, Guatemala), *J. Quat. Sci.*, 36, 169-179, 2021a.
- 575

- Cisneros de León, A., Schmitt, A., Kutterolf, S., Schindlbeck-Belo, J. C., Hernández, W., Sims, K., Garrison, J., Kant, L., Weber, B., and Wang, K. L.: Zircon and melt extraction from a long-lived and vertically extensive magma system underneath Ilopango Caldera (El Salvador), *Geochem., Geophys., Geosys.*, 22, 2021b.
- 580 Cole-Dai, J., Ferris, D. G., Kennedy, J. A., Sigl, M., McConnell, J. R., Fudge, T. J., Taylor, K. C., Souney, J. M., Geng, L., and Maselli, O. J.: Comprehensive Record of Volcanic Eruptions in the Holocene (11,000 years) from the WAIS Divide, Antarctica ice core, *J. Geophys. Res. Atmos.*, 126, doi: 10.1029/2020JD032855., 2021.
- Costa, A., Smith, V. C., Macedonio, G., and Matthews, N. E.: The magnitude and impact of the Youngest Toba Tuff super-eruption, *Front. Earth Sci.*, 2, 16, 2014.
- Crick, L., Burke, A., Hutchison, W., Kohno, M., Moore, K. A., Savarino, J., Doyle, E. A., Mahony, S., Kipfstuhl, S., Rae, J. W. B., Steele, R. C. J., Sparks, R. S. J., and Wolff, E. W.: New insights into the ~ 74ka Toba eruption from sulfur isotopes of 585 polar ice cores, *Clim. Past*, 17, 2119-2137, 10.5194/cp-17-2119-2021, 2021.
- Crosweller, H. S., Arora, B., Brown, S. K., Cottrell, E., Deligne, N. I., Guerrero, N. O., Hobbs, L., Kiyosugi, K., Loughlin, S. C., and Lowndes, J.: Global database on large magnitude explosive volcanic eruptions (LaMEVE), *J. Appl. Volcanol.*, 1, 1-13, 2012.
- 590 Derkachev, A. N., Gorbarenko, S. A., Ponomareva, V. V., Portnyagin, M. V., Malakhova, G. I., and Liu, Y.: Middle to late Pleistocene record of explosive volcanic eruptions in marine sediments offshore Kamchatka (Meiji Rise, NW Pacific), *J. Quat. Sci.*, 35, 362-379, 2020.
- Drexler, J. W., Rose jr., W. I., Sparks, R. S. J., and Ledbetter, M. T.: The Los Chocoyos Ash, Guatemala: A Major Stratigraphic Marker in Middle America and in the Three Ocean Basins, *Quat. Res.*, 13, 327-345, 1980.
- 595 Eisele, S., Reißig, S., Freundt, A., Kutterolf, S., Nürnberg, D., Wang, K., and Kwasnitschka, T.: Pleistocene to Holocene offshore tephrostratigraphy of highly explosive eruptions from the southwestern Cape Verde Archipelago, *Mar. Geol.*, 369, 233-250, 2015.
- Ellerhoff, B., Kirschner, M. J., Ziegler, E., Holloway, M. D., Sime, L., and Rehfeld, K.: Contrasting state-dependent effects of natural forcing on global and local climate variability, *Geophys. Res. Lett.*, e2022GL098335, 2022.
- 600 Fierstein, J. and Nathenson, M.: Another look at the calculation of fallout tephra volumes, *Bull. Volcanol.*, 54, 156-167, 1992.
- Freundt, A., Schindlbeck-Belo, J. C., Kutterolf, S., and Hopkins, J. L.: Tephra layers in the marine environment: A review of properties and emplacement processes, *Geol. Soc., London, Spec. Pub.*, 520, <https://doi.org/10.1144/SP520-2021-50>, 2021.
- Gao, C., Robock, A., and Ammann, C.: Volcanic forcing of climate over the past 1500 years: An improved ice core-based index for climate models, *J. Geophys. Res.: Atmos.*, 113, 2008.
- 605 Gupta, M. and Marshall, J.: The climate response to multiple volcanic eruptions mediated by ocean heat uptake: Damping processes and accumulation potential, *J. Clim.*, 31, 8669-8687, 2018.
- Hamill, P., Jensen, E. J., Russell, P. B. and Bauman, J. J.: The Life Cycle of Stratospheric Aerosol Particles., *Bull. Am. Meteorol. Soc.*, 78(7), doi:10.1175/1520-0477(1997)078<1395:TLCOSA>2.0.CO;2, 1997.

- Hammer, C.: Dating of Greenland ice cores by microparticle concentration analyses, *Isotopes and Impurities in Snow and Ice*, 610 297-301, 1977.
- Hopkins, J. L., Wysoczanski, R. J., Orpin, A. R., Howarth, J. D., Strachan, L. J., Lunenburg, R., McKeown, M., Ganguly, A., Twort, E., and Camp, S.: Deposition and preservation of tephra in marine sediments at the active Hikurangi subduction margin, *Quat. Sci. Rev.*, 247, 106500, 2020.
- Horn, S. and Schmincke, H.-U.: Volatile emission during the eruption of Baitoushan Volcano (China/North Korea) ca. 969 615 AD, *Bull. Volcanol.*, 61, 537-555, 2000.
- Huybers, P. and Langmuir, C.: Feedback between deglaciation, volcanism, and atmospheric CO₂, *Earth Planet. Sci. Lett.*, 286, 479-491, 2009.
- Iacovino, K., Ju-Song, K., Sisson, T., Lowenstern, J., Kuk-Hun, R., Jong-Nam, J., Kun-Ho, S., Song-Hwan, H., Oppenheimer, C., and Hammond, J. O.: Quantifying gas emissions from the “Millennium eruption” of Paektu volcano, democratic People’s 620 Republic of Korea/China, *Sci. Adv.*, 2, e1600913, 2016.
- Kiyosugi, K., Connor, C., Sparks, R. S. J., Crossweller, H. S., Brown, S. K., Siebert, L., Wang, T., and Takarada, S.: How many explosive eruptions are missing from the geologic record? Analysis of the quaternary record of large magnitude explosive eruptions in Japan, *J. Appl. Volcanol.*, 4, 1-15, 2015.
- Klawonn, M., Houghton, B. F., Swanson, D. A., Fagents, S. A., Wessel, P., and Wolfe, C. J.: Constraining explosive volcanism: 625 subjective choices during estimates of eruption magnitude, *Bull. Volcanol.*, 76, 1-6, 2014.
- Kobashi, T., Menviel, L., Jeltsch-Thömmes, A., Vinther, B. M., Box, J. E., Muscheler, R., Nakaegawa, T., Pfister, P. L., Döring, M., Leuenberger, M., Wanner, H., and Ohmura, A.: Volcanic influence on centennial to millennial Holocene Greenland temperature change, *Sci. Rep.*, 7, 1441, 10.1038/s41598-017-01451-7, 2017.
- Kutterolf S., Schindlbeck, J.C., Anselmetti, F.S., Ariztegui, D., Brenner, M., Curtis, J.H., Schmidt, D., Hodell, D.A., Müller, 630 A.D., Pérez, L., Pérez, W., Schwalb, A., Frische, M., Wang, K-L.: A 400-ka tephrochronological framework for Central America from Lake Petén Itzá (Guatemala) sediments, *Quat. Sci. Rev.*, 150, 200-220, doi:10.1016/j.quascirev.2016.08.023, 2016.
- Kutterolf, S., Schindlbeck, J. C., Jegen, M., Freundt, A., and Straub, S. M.: Milankovitch frequencies in tephra records at volcanic arcs: The relation of kyr-scale cyclic variations in volcanism to global climate changes, *Quat. Sci. Rev.*, 204, 1-16, 635 2019.
- Kutterolf, S., Jegen, M., Mitrovica, J. X., Kwasnitschka, T., Freundt, A., and Huybers, P.: A detection of Milankovitch frequencies in global volcanic activity, *Geology*, 41, 227-230, 10.1130/g33419.1, 2013.
- Kutterolf, S., Freundt, A., Pérez, W., Mörz, T., Schacht, U., Wehrmann, H., and Schmincke, H.-U.: The Pacific offshore record of Plinian arc volcanism in Central America, part 1: Along-arc correlations, *Geochem. Geophys. Geosyst.*, 9, 640 doi:10.1029/2007GC001631, 2008.
- Kutterolf, S., Freundt, A., Druitt, T., McPhie, J., Nomikou, P., Pank, K., Schindlbeck-Belo, J. C., Hansteen, T., and Allen, S.: The medial offshore record of explosive volcanism along the central to eastern Aegean Volcanic Arc: 2. Tephra ages and

- volumes, eruption magnitudes and marine sedimentation rate variations, *Geochem. Geophys. Geosyst.*, 22, e2021GC010011, 2021a.
- 645 Kutterolf, S., Freundt, A., Hansteen, T. H., Dettbarn, R., Hampel, F., Sievers, C., Wittig, C., Allen, S., Druitt, T. H., McPhie, J., Nomikou, P., Pank, K., Schindlbeck-Belo, J., Wang, K.-L., Lee, H.-Y., and Friedrichs, B.: The medial offshore record of explosive volcanism along the central to eastern Aegean Volcanic Arc, part 1: Tephrostratigraphic correlations, *Geochem. Geophys. Geosyst.*, 22, e2021GC010010, doi:10.1029/2021GC010010, 2021b.
- Lanciki, A., Cole-Dai, J., Thiemens, M. H. and Savarino, J.: Sulfur isotope evidence of little or no stratospheric impact by the
650 1783 Laki volcanic eruption, *Geophys. Res. Lett.*, 39, L01806, doi:10.1029/2011GL050075, 2012.
- Lavigne, F.: Rate of sediment yield following small-scale volcanic eruptions: a quantitative assessment at the Merapi and Semeru stratovolcanoes, Java, Indonesia, *Earth Surface Processes and Landforms: The Journal of the British Geomorphological Research Group*, 29, 1045-1058, 2004.
- Lin, J., Svensson, A., Hvidberg, C. S., Lohmann, J., Kristiansen, S., Dahl-Jensen, D., Steffensen, J. P., Rasmussen, S. O.,
655 Cook, E., and Kjær, H. A.: Magnitude, frequency and climate forcing of global volcanism during the last glacial period as seen in Greenland and Antarctic ice cores (60–9 ka), *Clim. Past*, 18, 485-506, 2022.
- Lohmann, J. and Svensson, A.: Ice core evidence for major volcanic eruptions at the onset of Dansgaard–Oeschger warming events, *Clim. Past*, 18, 2021-2043, 10.5194/cp-18-2021-2022, 2022.
- Mackay, H., Plunkett, G., Jensen, B. J. L., Aubry, T. J., Corona, C., Kim, W. M., Toohey, M., Sigl, M., Stoffel, M., Anchukaitis,
660 K. J., Raible, C., Bolton, M. S. M., Manning, J. G., Newfield, T. P., Di Cosmo, N., Ludlow, F., Kostick, C., Yang, Z., Coyle McClung, L., Amesbury, M., Monteath, A., Hughes, P. D. M., Langdon, P. G., Charman, D., Booth, R., Davies, K. L., Blundell, A., and Swindles, G. T.: The 852/3 CE Mount Churchill eruption: examining the potential climatic and societal impacts and the timing of the Medieval Climate Anomaly in the North Atlantic region, *Clim. Past*, 18, 1475-1508, 2022.
- McConnell, J. R., Sigl, M., Plunkett, G., Burke, A., Kim, W. M., Raible, C. C., Wilson, A. I., Manning, J. G., Ludlow, F.,
665 Chellman, N. J., Innes, H. M., Yang, Z., Larsen, J. F., Schaefer, J. R., Kipfstuhl, S., Mojtabavi, S., Wilhelms, F., Opel, T., Meyer, H., and Steffensen, J. P.: Extreme climate after massive eruption of Alaska’s Okmok volcano in 43 BCE and effects on the late Roman Republic and Ptolemaic Kingdom, *Proceedings of the National Academy of Sciences*, 117, 15443-15449, doi:10.1073/pnas.2002722117, 2020.
- McGregor, H. V., Evans, M. N., Goosse, H., Leduc, G., Martrat, B., Addison, J. A., Mortyn, P. G., Oppo, D. W., Seidenkrantz,
670 M.-S., and Sicre, M.-A.: Robust global ocean cooling trend for the pre-industrial Common Era, *Nat. Geosci.*, 8, 671-677, 2015.
- Metzner, D., Kutterolf, S., Toohey, M., Timmreck, C., Niemeier, U., Freundt, A., and Krüger, K.: Radiative forcing and climate impact resulting from SO₂ injections based on a 200,000-year record of Plinian eruptions along the Central American Volcanic Arc, *Int. J. Earth Sci.*, 103, 2063-2079, 2014.
- Miller, G. H., Geirsdóttir, Á., Zhong, Y., Larsen, D. J., Otto-Bliesner, B. L., Holland, M. M., Bailey, D. A., Refsnider, K. A.,
675 Lehman, S. J., and Southon, J. R.: Abrupt onset of the Little Ice Age triggered by volcanism and sustained by sea-ice/ocean feedbacks, *Geophys. Res. Lett.*, 39, 2012.

- Miller, R. L., Knippertz, P., Pérez García-Pando, C., Perlwitz, J. P., and Tegen, I.: Impact of dust radiative forcing upon climate, in: *Mineral dust*, Springer, 327-357, 2014.
- Oppenheimer, C.: Limited global change due to the largest known Quaternary eruption, Toba \approx 74kyr BP?, *Quat. Sci. Rev.*, 680 21, 1593-1609, [https://doi.org/10.1016/S0277-3791\(01\)00154-8](https://doi.org/10.1016/S0277-3791(01)00154-8), 2002.
- Oppenheimer, C., Wacker, L., Xu, J., Galván, J. D., Stoffel, M., Guillet, S., Corona, C., Sigl, M., Di Cosmo, N., and Hajdas, I.: Multi-proxy dating the ‘Millennium Eruption’ of Changbaishan to late 946 CE, *Quat. Sci. Rev.*, 158, 164-171, 2017.
- Otterå, O. H., Bentsen, M., Drange, H., and Suo, L.: External forcing as a metronome for Atlantic multidecadal variability, *Nat. Geosci.*, 3, 688-694, 2010.
- 685 Owens, M. J., Lockwood, M., Hawkins, E., Usoskin, I., Jones, G. S., Barnard, L., Schurer, A., and Fasullo, J.: The Maunder minimum and the Little Ice Age: an update from recent reconstructions and climate simulations, *J. Space Weather Space Clim.*, 7, A33, 2017.
- Pages2k-Consortium, 2019: Consistent multidecadal variability in global temperature reconstructions and simulations over the Common Era, *Nat. Geosci.*, 12, 643-649, [10.1038/s41561-019-0400-0](https://doi.org/10.1038/s41561-019-0400-0), 2019.
- 690 Papale, P.: Global time-size distribution of volcanic eruptions on Earth, *Sci. Rep.*, 8, 1-11, 2018.
- Paterne, M., Labeyrie, J., Guichard, F., Mazaud, A., and Maitre, F.: Fluctuations of the Campanian explosive volcanic activity (South Italy) during the past 190,000 years, as determined by marine tephrochronology, *Earth and Planetary Science Letters*, 98, 166-174, 1990.
- Pearson, C., Sigl, M., Burke, A., Davies, S., Kurbatov, A., Severi, M., Cole-Dai, J., Innes, H., Albert, P. G., and Helmick, M.: 695 Geochemical ice-core constraints on the timing and climatic impact of Aniakchak II (1628 BCE) and Thera (Minoan) volcanic eruptions, *PNAS Nexus*, doi: [10.1093/pnasnexus/pgac048](https://doi.org/10.1093/pnasnexus/pgac048), 2022.
- Plunkett, G., Sigl, M., McConnell, J. R., Pilcher, J. R., and Chellman, N. J.: The significance of volcanic ash in Greenland ice cores during the Common Era, *Quaternary Sci Rev*, 301, 107936, 2023.
- Pollard, A., Blockley, S., and Ward, K.: Chemical alteration of tephra in the depositional environment: theoretical stability 700 modelling, *Journal of Quaternary Science*, 18, 385-394, 2003.
- Praetorius, S., Mix, A., Jensen, B., Froese, D., Milne, G., Wolhowe, M., Addison, J., and Prah, F.: Interaction between climate, volcanism, and isostatic rebound in Southeast Alaska during the last deglaciation, *Earth Planet. Sci. Lett.*, 452, 79-89, <https://doi.org/10.1016/j.epsl.2016.07.033>, 2016.
- Pyle, D. M.: Assessment of the minimum volume of tephra fall deposits, *J. Volc. Geo. Res.*, 69, 379-382, 1995.
- 705 Pyle, D. M., Beattie, P. D., and Bluth, G. J. S.: Sulphur emissions to the stratosphere from explosive volcanic eruptions, *Bull. Volcanol.*, 57, 663-671, 1996.
- Rampino, M. R. and Self, S.: Historic eruptions of Tambora (1815), Krakatau (1883), and Agung (1963), their stratospheric aerosols, and climate impact, *Quat. Res.*, 18, 127-143, 1982.
- Rampino, M. R. and Self, S.: Climate-Volcanism Feedback and the Toba Eruption of \sim 74,000 Years Ago, *Quat. Res.*, 40, 269- 710 280, <https://doi.org/10.1006/qres.1993.1081>, 1993.

- Rampino, M. R., Self, S., and Fairbridge, R. W.: Can rapid climatic change cause volcanic eruptions?, *Science*, 206, 826 - 829, 1979.
- Reinig, F., Wacker, L., Jöris, O., Oppenheimer, C., Guidobaldi, G., Nievergelt, D., Adolphi, F., Cherubini, P., Engels, S., Esper, J., Land, A., Lane, C., Pfanz, H., Remmele, S., Sigl, M., Sookdeo, A., and Büntgen, U.: Precise date for the Laacher
715 See eruption synchronizes the Younger Dryas, *Nature*, 595, 66- 69, 2021.
- Robock, A.: Volcanic eruptions and climate., *Reviews of Geophysics*, 38, 191-219, 2000.
- Ryan, W. B. F., Carbotte, S. M., Coplan, J. O., O'Hara, S., Melkonian, A., Arko, R., Weissel, R. A., Ferrini, V., Goodwillie, A., Nitsche, F., Bonczkowski, J., and Zemsky, R.: Global Multi-Resolution Topography synthesis, *Geochem., Geophys., Geosys.*, 10, Q03014, 10.1029/2008gc002332, 2009.
- 720 Schindlbeck, J. C., Kutterolf, S., Straub, S. M., Andrews, G. D., Wang, K. L., and Mleneck-Vautravers, M. J.: One Million Years tephra record at IODP Sites U 1436 and U 1437: Insights into explosive volcanism from the Japan and Izu arcs, *Isl. Arc*, 27, e12244, 2018a.
- Schindlbeck, J. C., Jegen, M., Freundt, A., Kutterolf, S., Straub, S. M., Mleneck-Vautravers, M. J., and McManus, J. F.: 100-kyr cyclicity in volcanic ash emplacement: evidence from a 1.1 Myr tephra record from the NW Pacific, *Sci.Rep.*, 8, 4440,
725 10.1038/s41598-018-22595-0, 2018b.
- Schindlbeck, J. C., Kutterolf, S., Freundt, A., Alvarado, G., Wang, K. L., Straub, S., Hemming, S., Frische, M., and Woodhead, J.: Late Cenozoic tephrostratigraphy offshore the southern Central American Volcanic Arc: 1. Tephra ages and provenance, *Geochem., Geophys., Geosys.*, 17, 4641-4668, 2016.
- Schindlbeck, J. C., Kutterolf, S., Freundt, A., Straub, S. M., Wang, K.-L., Jegen, M., Hemming, S. R., Baxter, A. T., and
730 Sandoval, M. I.: The Miocene Galápagos ash layer record of Integrated Ocean Drilling Program Legs 334 and 344: Ocean-island explosive volcanism during plume-ridge interaction, *Geology*, 43, 599-602, 10.1130/g36645.1, 2015.
- Schleussner, C.-F. and Feulner, G.: A volcanically triggered regime shift in the subpolar North Atlantic Ocean as a possible origin of the Little Ice Age, *Clim. Past*, 9, 1321-1330, 2013.
- Schleussner, C.-F., Divine, D., Donges, J. F., Miettinen, A., and Donner, R. V.: Indications for a North Atlantic ocean
735 circulation regime shift at the onset of the Little Ice Age, *Clim. Dyn.*, 45, 3623-3633, 2015.
- Schurer, A. P., Tett, S. F., and Hegerl, G. C.: Small influence of solar variability on climate over the past millennium, *Nat. Geosci.*, 7, 104-108, 2014.
- Schurer, A. P., Hegerl, G. C., Mann, M. E., Tett, S. F., and Phipps, S. J.: Separating forced from chaotic climate variability over the past millennium, *J. Clim.*, 26, 6954-6973, 2013.
- 740 Sigl, M., Toohey, M., McConnell, J. R., Cole-Dai, J., and Severi, M.: HolVol: Reconstructed volcanic stratospheric sulfur injections and aerosol optical depth for the Holocene (9500 BCE to 1900 CE), PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.928646>, 2021.

- Sigl, M., Toohey, M., McConnell, J. R., Cole-Dai, J., and Severi, M.: Volcanic stratospheric sulfur injections and aerosol optical depth during the Holocene (past 11500 years) from a bipolar ice-core array, *Earth Syst. Sci. Data*, 14, 3167-3196, 10.5194/essd-14-3167-2022, 2022.
- 745 Sigl, M., Winstrup, M., McConnell, J. R., Welten, K. C., Plunkett, G., Ludlow, F., Büntgen, U., Caffee, M., Chellman, N., and Dahl-Jensen, D.: Timing and climate forcing of volcanic eruptions for the past 2,500 years, *Nature*, 523, 543-549, 2015.
- Sigurdsson, H.: Evidence of volcanic loading of the atmosphere and climate response, *Glob. Planet. Change*, 3, 277-289, [https://doi.org/10.1016/0921-8181\(90\)90024-7](https://doi.org/10.1016/0921-8181(90)90024-7), 1990.
- 750 Sigvaldason, G. E., Annertz, K., and Nilsson, M.: Effect of Glacial loading/deloading on volcanism: postglacial volcanic production rate of the Dyngjufjöll area, central Iceland., *Bull. Volcanol.*, 54, 385 - 392, 1992.
- Smith, V. C., Costa, A., Aguirre-Diaz, G., Pedrazzi, D., Scifo, A., Plunkett, G., Poret, M., Tournigand, P. Y., Miles, D., Dee, M. W., McConnell, J. R., Sunye-Puchol, I., Harris, P. D., Sigl, M., Pilcher, J. R., Chellman, N., and Gutierrez, E.: The magnitude and impact of the 431 CE Tierra Blanca Joven eruption of Ilopango, El Salvador, *P Natl Acad Sci USA*, 117, 26061-26068, 2020.
- 755 Swingedouw, D., Mignot, J., Ortega, P., Khodri, M., Menegoz, M., Cassou, C., and Hanquiez, V.: Impact of explosive volcanic eruptions on the main climate variability modes, *Glob. Planet. Change*, 150, 24-45, 2017.
- Timmreck, C., Toohey, M., Zanchettin, D., Brönnimann, S., Lundstad, E., and Wilson, R.: The unidentified eruption of 1809: a climatic cold case, *Clim. Past*, 17, 1455-1482, 10.5194/cp-17-1455-2021, 2021.
- 760 Toohey, M. and Sigl, M.: Volcanic stratospheric sulfur injections and aerosol optical depth from 500 BCE to 1900 CE, *Earth Syst. Sci. Data*, 9, 809-831, 2017.
- Toohey, M., Krüger, K., Sigl, M., Stordal, F., and Svensen, H.: Climatic and societal impacts of a volcanic double event at the dawn of the Middle Ages, *Clim. Change*, 136, 401-412, 2016.
- Toohey, M., Schindlbeck-Belo, J. C.: PalVol: A proxy-based semi-stochastic ensemble reconstruction of volcanic stratospheric aerosol for the last glacial cycle (Version 1). World Data Center for Climate (WDCC) at DKRZ. <https://doi.org/10.26050/WDCC/PalVolv1>, 2023
- 765 Wetzel, A.: The preservation potential of ash layers in the deep-sea: The example of the 1991-Pinatubo ash in the South China Sea, *Sedimentology*, 56, 1992-2009, 2009.
- Zanchettin, D., Bothe, O., Graf, H. F., Lorenz, S. J., Luterbacher, J., Timmreck, C., and Jungclaus, J. H.: Background conditions influence the decadal climate response to strong volcanic eruptions, *J. Geophys. Res.: Atmos.*, 118, 4090-4106, 2013.
- 770 Zhong, Y., Miller, G., Otto-Bliesner, B., Holland, M., Bailey, D., Schneider, D., and Geirsdottir, A.: Centennial-scale climate change from decadal-paced explosive volcanism: a coupled sea ice-ocean mechanism, *Clim. Dyn.*, 37, 2373-2387, 2011.