

# CMEMS-LSCE: A global 0.25-degree, monthly reconstruction of the surface ocean carbonate system

Thi-Tuyet-Trang Chau<sup>1</sup>, Marion Gehlen<sup>1</sup>, Nicolas Metzler<sup>2</sup>, and Frédéric Chevallier<sup>1</sup>

<sup>1</sup>Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France

<sup>2</sup>Laboratoire LOCEAN (IPSL), Sorbonne Université, CNRS-IRD-MNHN, Paris, F-75005, France

**Correspondence:** Thi-Tuyet-Trang CHAU (trang.chau@lsce.ipsl.fr, thi.tuyet.trang.chau@gmail.com)

**Abstract.** Observation-based data reconstructions of global surface ocean carbonate system variables play an essential role in monitoring the recent status of ocean carbon uptake and ocean acidification as well as their impacts on marine organisms and ecosystems. So far ongoing efforts are directed towards exploring new approaches to describe the complete marine carbonate system and to better recover its fine-scale features. In this respect, our research activities within the Copernicus Marine Environment Monitoring Service (CMEMS) aim at developing a sustainable production chain of observation-derived global ocean carbonate system datasets at high space-time resolution. As the start of the long-term objective, this study introduces a new global 0.25° monthly reconstruction, namely CMEMS-LSCE, for the period 1985-2021. The CMEMS-LSCE reconstruction derives datasets of six carbonate system variables including surface ocean partial pressure of CO<sub>2</sub> ( $p\text{CO}_2$ ), total alkalinity ( $A_T$ ), total dissolved inorganic carbon (DIC), surface ocean pH, and saturation states with respect to aragonite ( $\Omega_{ar}$ ) and calcite ( $\Omega_{ca}$ ). Reconstructing  $p\text{CO}_2$  relies on an ensemble of neural network models mapping gridded observation-based data provided by the Surface Ocean CO<sub>2</sub> Atlas (SOCAT). Surface ocean  $A_T$  is estimated with a multiple linear regression approach, and the remaining carbonate variables are resolved by CO<sub>2</sub> system speciation given the reconstructed  $p\text{CO}_2$  and  $A_T$ .  $1\sigma$ -uncertainty associated with these estimates is also provided. Here,  $\sigma$  stands for either ensemble standard deviation of  $p\text{CO}_2$  estimates or total uncertainty for each of the five other variables propagated through the processing chain with input data uncertainty. We demonstrate that the 0.25°-resolution  $p\text{CO}_2$  product outperforms a coarser spatial resolution (1°) thanks to a higher data coverage nearshore and a better description of horizontal and temporal variations in  $p\text{CO}_2$  across diverse ocean basins, particularly in the coastal-open-ocean continuum. Product qualification with observation-based data confirms reliable reconstructions with root-of-mean-square-deviation from observations less than 8%, 4%, and 1% relative to the global mean of  $p\text{CO}_2$ ,  $A_T$  (DIC), and pH. The global average  $1\sigma$ -uncertainty is below 5% and 8% for  $p\text{CO}_2$  and  $\Omega_{ar}$  ( $\Omega_{ca}$ ), 2% for  $A_T$  and DIC, and 0.4% for pH relative to their global mean values. Both model-observation misfit and model uncertainty indicate that coastal data reproduction still needs further improvement, wherein high temporal and horizontal gradients of carbonate variables and representative uncertainty from data sampling would be taken into account in priority. This study also presents a potential use case of the CMEMS-LSCE carbonate data product in tracking the recent state of ocean acidification.

## 1 Introduction

25 Between 1750 and 2019, the ocean took up an estimated 25% (or  $170 \pm 20$  PgC) of total cumulated anthropogenic  $\text{CO}_2$  ( $685 \pm 75$  PgC) emitted to the atmosphere (IPCC AR6 - the Sixth Assessment Report of the United Nations Intergovernmental Panel on Climate Change, Canadell et al., 2021). While the uptake of anthropogenic  $\text{CO}_2$  mitigates global warming it also profoundly modifies seawater chemistry in a suite of well-understood reactions (Orr et al., 2005) leading to an increase in hydrogen ion concentration ( $[\text{H}^+]$ ), as well as a decrease in carbonate ion concentration ( $[\text{CO}_3^{2-}]$ ) and in the saturation state of seawater ( $\Omega$ )  
30 with respect to calcium carbonate minerals ( $\text{CaCO}_3$ ). The increase in hydrogen ion concentration ( $[\text{H}^+]$ ) is commonly reported as a decrease in  $\text{pH}$  ( $\text{pH} = -\log[\text{H}^+]$ ) and referred to as ocean acidification.

Changes in carbonate chemistry impact calcifying plankton and benthos as a direct result of decreasing seawater saturation state with respect to  $\text{CaCO}_3$  (Fabry et al., 2008; Thomsen et al., 2015). Ocean acidification also modifies the production of marine trace gases exchanged at the air-sea interface (Hopkins et al., 2020), the availability of nutrients fueling primary  
35 production (Doney et al., 2009), as well as the speciation of pollutants (Millero et al., 2009; Hoffmann et al., 2012). These chemical changes interact with warming and ocean deoxygenation to drive major changes in marine ecosystems (Doney et al., 2020) and to alter global biogeochemical cycles with the potential for feeding back on radiative forcing (Gehlen et al., 2011; Hopkins et al., 2020). The likelihood for major disruptive impacts of ocean acidification on marine ecosystems, if future  
40  $\text{CO}_2$  emissions were to go unabated, is reflected by the Sustainable Development Goal 14.3 (SDG 14.3) - "Reduce Ocean Acidification: minimize and address impacts of ocean acidification" (<https://www.globalgoals.org/14-life-below-water>, last access: 20/03/2023). Albeit not specifically mentioned, moving towards SDG 14.3 implies the understanding of historical and contemporary carbonate chemistry, its mean state, trends and variability.

~~Earth system models have been widely used to track changes in ocean  $\text{pH}$  over the historical period and to project its future evolution under different  $\text{CO}_2$  emission pathways (Bopp et al., 2013; Gattuso et al., 2015; Kwiatkowski et al., 2020; Cooley et al., 2022; Ji  
45 -The present-day global surface ocean  $\text{pH}$  is roughly 0.1  $\text{pH}$  units less than at the beginning of the industrial era (Gattuso et al., 2015; Jiang et al., 2015) corresponding to an increase in hydrogen ion concentration of 26% (Doney, 2010). By the end of the 21<sup>st</sup> century, the  $\text{pH}$  is projected to decrease by  $0.16 \pm 0.002$   $\text{pH}$  units in response to the IPCC AR6 low emission scenario (SSP1-2.6), respectively by  $0.44 \pm 0.005$   $\text{pH}$  units in response to the IPCC AR6 high emission pathway (SSP5-8.5) relative to 1870–1899 (Kwiatkowski et al., 2020). Understanding impacts on marine biota requires to move towards finer spatial and temporal scales  
50 than resolved by the current generation of Earth system models (Torres et al., 2021), as well as to expand the analysis from  $\text{pH}$  to other carbonate system variables such as the saturation state with respect to calcium carbonate minerals and the buffer capacity. The development and implementation of environmental management strategies equally rely on understanding and attributing the variability of the carbonate system from diurnal to decadal time scales to underlying physical-chemical-biological processes.~~

55 In situ time series have played an important role in monitoring ocean acidification over the last decades (Bates et al., 2014; Lauvset et al., 2015; Sutton et al., 2019; Pérez et al., 2021; Leseurre et al., 2022; Skjelvan et al., 2022). At these sites, seawater  $\text{pH}$  ( $\Omega$ ) has been either directly measured or calculated from measurements of other carbonate system variables. These variables

include surface ocean partial pressure of CO<sub>2</sub> ( $p\text{CO}_2$ ), total alkalinity ( $A_T$ ), and dissolved inorganic carbon (DIC). While changes in time series of carbonate system variables well reflect impacts of enhanced anthropogenic CO<sub>2</sub> uptake on ocean chemistry at a local scale (Steinberg et al., 2001; González-Dávila and Santana-Casiano, 2009; Dore et al., 2009; Bates et al., 2014; Pérez et al., 2021), the reliable upscaling to large ocean regions or entire basins requires a significant extension of the existing observing network (Lauvset et al., 2015; Bakker et al., 2016; Sutton et al., 2019; Lauvset et al., 2022a).

Time series data are completed by bottle data from international cruises. These data are synthesized by the Global Ocean Data Analysis Project v2.2022 (GLODAPv2.2022) and include about 1.4 million measurements of surface-to-interior ocean  $p\text{H}$ ,  $A_T$ , DIC, and other parameters (Lauvset et al., 2022a, <https://www.glodap.info/>, last access: 30/9/2022)(Lauvset et al., 2022b, <https://www.glo>. Likewise, underway measurements of near-surface CO<sub>2</sub> fugacity, i.e.,  $p\text{CO}_2$  corrected for non-ideal gas behavior, are compiled in the Surface Ocean CO<sub>2</sub> Atlas (SOCAT) since its first release in 2011 (Pfeil et al., 2013). That latest version SOCATv2022 yields approximately 33.7 million high-quality controlled data (Bakker et al., 2022, <http://www.socat.info/>, last access: 17/6/2022). Despite millions of observations available, data coverage is still modest, e.g., CO<sub>2</sub> fugacity samples over the global ocean cover less than 2% of its surface for each month in the last three decades (Bakker et al., 2016; Hauck et al., 2020). Mapping methods have thus become an essential tool in ocean carbon cycle research allowing to interpolate or extrapolate these sparse measurements into space-time varying fields of carbonate system variables(e.g., Rödenbeck et al., 2015) and used for global carbon budget estimates (Friedlingstein et al., 2022).

Recent years have seen the rapid development of machine learning approaches to map global surface ocean  $p\text{CO}_2$  (see Rödenbeck et al., 2013; Landschützer et al., 2016; Denvil-Sommer et al., 2019; Gregor et al., 2019; Chau et al., 2022b, for instance). Thanks to these efforts, the carbon cycle community can now draw on an ensemble of reconstructions for the observation-based assessment of the ocean carbon sink (Friedlingstein et al., 2022). However, only a few global observation-based reconstructions are available for  $p\text{H}$ ,  $A_T$ , DIC, and  $\Omega$  with respect to calcite and aragonite (see Gregor and Gruber, 2021, for a review). The reconstruction of global distributions of these variables is hampered by an insufficient amount of direct measurements (Bakker et al., 2016; Lauvset et al., 2022a). Alternatively, the complete carbonate system can be obtained by speciation given the information of any couple of  $p\text{CO}_2$ ,  $p\text{H}$ ,  $A_T$  or DIC together with chemical (e.g., phosphate, silicate, nitrate) and physical variables (e.g., temperature, salinity), as well as corresponding dissociation constants (Park, 1969; Lewis and Wallace, 1998; Dickson et al., 2007).

Regardless of the developments in different observation-based estimation methods, Takahashi et al. (2014), Iida et al. (2021), and Gregor and Gruber (2021) propose global climatologies or monthly varying fields of all variables of the carbonate system, i.e.,  $p\text{CO}_2$ ,  $p\text{H}$ ,  $A_T$ , DIC, and  $\Omega$ . These data products have a spatial resolution of 1° (~100km × 100km) or even coarser. Nevertheless, the variations of carbonate system variables over the coastal regions where their instantaneous gradients are driven by smaller-scale features like ocean upwelling, wind turbulence, eddies, water runoff, and sharp biological productivity (Jones et al., 2012; Bakker et al., 2016; Laruelle et al., 2017) are poorly described at such spatial resolutions. Here we improve on existing studies by providing a global 0.25°, monthly observation-based surface ocean carbonate system product consisting of datasets of six core surface ocean carbonate system variables of the marine carbonate system (see Table 1 and Appendix A for definitions) and their associated 1 $\sigma$ -uncertainty. This high-resolution data product covers the years from 1985 to 2021.

Laboratoire des Sciences du Climat et de l'Environnement (LSCE) is in charge of the product within the European Copernicus Marine Environment Monitoring Service (CMEMS). Our product is referred to as CMEMS-LSCE hereafter.

95 The reconstruction of surface ocean carbonate system variables starts with the reconstruction of surface ocean  $p\text{CO}_2$  and  $A_T$  in each regular grid of  $1\text{month} \times 0.25^\circ \times 0.25^\circ$  in the period 1985-2021 (444 months in total). Next, variables  $p\text{H}$ ,  $\text{DIC}$ , and  $\Omega$  are derived by speciation. Advantages of the combination of  $p\text{CO}_2$  and  $A_T$  over others for the speciation of the carbonate system are: (1)  $p\text{CO}_2$  is the most extensively measured parameter, (2)  $A_T$  can be accurately predicted from salinity, temperature, and nutrient concentrations, and (3) the combination of these two prior variables results in the slightest uncertainty of  $p\text{H}$  estimates  
100 (Zeebe and Wolf-Gladrow, 2001; Lauvset and Gruber, 2014; Takahashi et al., 2014; Orr et al., 2018). The three main successive modules used in the CMEMS-LSCE production chain are summarized as follows.

**Table 1.** CMEMS-LSCE carbonate system variables.

Standard names	Notations	Units
1. Partial pressure of $\text{CO}_2$ in surface seawater	$p\text{CO}_2$	$\mu\text{atm}$
2. Total alkalinity in surface seawater	$A_T$	$\mu\text{mol kg}^{-1}$
3. Surface ocean dissolved inorganic carbon	$\text{DIC}$	$\mu\text{mol kg}^{-1}$
4. Surface seawater $p\text{H}$ reported on total scale	$p\text{H}$	-
5. Saturation state for surface seawater with respect to aragonite	$\Omega_{ar}$	-
6. Saturation state for surface seawater with respect to calcite	$\Omega_{ca}$	-

i) *Reconstruction of  $p\text{CO}_2$*  (Sect. 3.1): ~~a modified version of~~ the CMEMS-LSCE-FFNN **ensemble-based** approach (Chau et al., 2022b) is **applied-modified** to map gridded datasets of SOCATv2022  $\text{CO}_2$  fugacity and predictors in order to reconstruct  $p\text{CO}_2$  at a **spatial-finer spatial scale** resolution of  $0.25^\circ$  for every month in the period 1985-2021 (444 months in  
105 total). The ~~CMEMS-LSCE-FFNN works on an ensemble of 100~~ **spatial resolution of new** feed-forward neural networks (FFNNs) **is 16-fold higher than the original**. By design, ~~100-member ensemble model outputs allow the ensemble of model outputs allows~~ to yield the best model estimate (i.e., ensemble mean) and model uncertainty (i.e., ensemble standard deviation) ~~for surface ocean  $p\text{CO}_2$  in each grid at each  $0.25^\circ$ -grid cell and each month. The primary modification of this study's version and the original CMEMS-LSCE-FFNN (Chau et al., 2022b) is an increase of 16-fold in model~~  
110 **spatial-resolution**. Global monthly reconstructions of  $p\text{CO}_2$  proposed by this study complement the previous climatological product by Landschützer et al. (2020), i.e., a combination of the two existing datasets covering respectively the open ocean at  $1^\circ$  (Landschützer et al., 2016) and the coastal sector at  $0.25^\circ$  (Laruelle et al., 2017).

ii) *Reconstruction of  $A_T$*  (Sect. 3.2): locally interpolated alkalinity regression (LIAR; Carter et al., 2016, 2018) is chosen to estimate ~~total alkalinity on regular grids of  $1\text{month} \times 0.25^\circ \times 0.25^\circ$  monthly~~ **total alkalinity** over the global surface  
115 ocean ~~for the years 1985-2021. LIAR works with multiple linear regression models, each representing a combination of predictor variables. The best linear model, which has the lowest prediction error among the others, is retained for the final estimation of  $A_T$ .~~ Various reconstruction methods for  $A_T$  exist (see Carter et al., 2016; Broullón et al., 2019; Gregor and Gruber, 2021, for a review), but we choose LIAR due to its global applicability, simplicity in setting, and



accuracy compared to other published approaches (Carter et al., 2018; Gregor and Gruber, 2021). Importantly, LIAR  
120 allows determining reconstruction uncertainty propagated from multiple sources of input uncertainties at desired model  
resolutions.

iii) *Reconstruction of pH, DIC, and saturation states with respect to aragonite ( $\Omega_{ar}$ ) and calcite ( $\Omega_{ca}$ )* (Sect. 3.3): CO2SYS  
(Lewis and Wallace, 1998; Van Heuven et al., 2011) is a standard software used for the speciation of carbonate parameters  
in the marine CO<sub>2</sub> system (see Olsen et al., 2016; Bresnahan et al., 2021; Gregor and Gruber, 2021; Woosley, 2021, for  
125 a few). ~~The CO2SYS speciation is built on a set of equilibrium equations (Dickson et al., 2007; Dickson, 2010). Given  
the reconstructed  $p\text{CO}_2$  and  $A_T$ , non-CO<sub>2</sub> acid-base constituents, physical variables, and equilibrium constants, this  
method allows solving pH, DIC,  $\Omega_{ar}$ , and  $\Omega_{ca}$  at the same input resolutions.~~ A complementary of the CO2SYS software  
developed by Orr et al. (2018) is used to quantify the uncertainty associated with these carbonate system variables. All  
the input data uncertainties are propagated through the CO2SYS processing chain.

130 The global monthly, 0.25°-resolution datasets of  ~~$p\text{CO}_2$ ,  $A_T$ , pH, DIC,  $\Omega_{ar}$ , and  $\Omega_{ca}$~~  **surface carbonate variables** are intensi-  
sively evaluated against different observation-based products independent from our model fitting at a global scale to in situ  
locations (Table 2). In Section 4, multiple metrics are proposed for product ~~analyzes~~ **analyses** and assessments. Results are  
presented in section 5 with emphasis on the evaluation of the best reconstruction and associated model uncertainty for each  
variable (Sect. 5). This section also highlights the advantages obtained with an increase in spatial resolution and presents an  
135 application of the CMEMS-LSCE product in tracking ocean acidification over the last three decades. Section 6 summarizes  
key results, discusses the potential for future model upgrades, and introduces possible product use cases. **The high-resolution  
data product described in this manuscript (netCDF format) can be accessed via repository under data DOI: 10.14768/a2f0891b-  
763a-49e9-af1b-78ed78b16982.**

## 2 Data used and reprocessing

### 140 2.1 Input data products for surface ocean carbonate system reconstructions

Many observation-based products are used as predictors of our target carbonate system variables (Table 1). Global ocean  
maps of sea surface temperature (SST), **sea surface** salinity (SSS), ~~height~~ **sea surface height above geoid** (SSH), chlorophyll-  
*a* (Chl-*a*) come from the Copernicus Marine Environment Monitoring Service (CMEMS: Good et al., 2020; Nardelli et al.,  
2016; Droghei et al., 2018; Maritorena et al., 2010). Mixed layer depth (MLD) fields belong to Estimating the Circulation  
and Climate of the Ocean project Phase II (ECCO2, Menemenlis et al., 2008). CO<sub>2</sub> mole fractions ( $x\text{CO}_2$ ) are derived from  
145 the CO<sub>2</sub> atmospheric inversion of the Copernicus Atmosphere Monitoring Service (CAM5, Chevallier et al., 2005, 2010;  
Chevallier, 2013). Surface ocean concentrations of nitrate (NO<sub>3</sub>), silicate (SiO<sub>2</sub>), and phosphate (PO<sub>4</sub>) are extracted from the  
World Ocean Atlas 2018 (WOA18, Garcia et al., 2019). The climatological  $p\text{CO}_2$  ( $p\text{CO}_2^{\text{clim}}$ ) product is provided by Lamont  
Doherty Earth Observatory (LDEO, Takahashi et al., 2009). Details of these products including resource access, data coverage,  
150 and resolutions are presented in Table 1.

**Table 1.** Input data used in the reconstructions of CMEMS-LSCE carbonate system variables over the global ocean in 1985-2021.

Variables	Notations	Units	Products	Resolutions	References
1. CO <sub>2</sub> fugacity	$f\text{CO}_2$	$\mu\text{atm}$	Surface Ocean CO <sub>2</sub> Atlas version 2022 (SOCATv2022, 1985-2021)	monthly, 1° (open ocean) and 0.25° (coastal ocean)	Bakker et al. (2022)
2. Sea surface temperature	SST	°C	CMEMS SST_GLO_SST_L4_REP_OBSERVATIONS_010_011 and SST_GLO_SST_L4_NRT_OBSERVATIONS_010_001 (1985-2021)	daily, 0.05°	Good et al. (2020)
3. Sea surface salinity	SSS	PSU	CMEMS MULTIOBS_GLO_PHY_S_SURFACE_MYNRT_015_013 (1993-2021)	monthly, 0.25°	Nardelli et al. (2016); Droghei et al. (2018)
4. Sea surface height	SSH	m	CMEMS SEALEVEL_GLO_PHY_L4_MY_008_047 and SEALEVEL_GLO_PHY_L4_NRT_OBSERVATIONS_008_046 (1993-2021)	daily, 0.25°	CLS-TOULOUSE
5. Mixed layer depth	MLD	m	Estimating the Circulation and Climate of the Ocean project Phase II (ECCO2, 1992-2021)	daily, 0.25°	Menemenlis et al. (2008)
6. Chlorophyll- <i>a</i>	CHL- <i>a</i>	$\text{mg m}^{-3}$	CMEMS OCEANCOLOUR_GLO_CHL_L4_REP_OBSERVATIONS_009_082 and OCEANCOLOUR_GLO_CHL_L4_NRT_OBSERVATIONS_009_033 (1998-2021)	daily, 0.25°	GLOCOLOUR, Maritorea et al. (2010)
7. CO <sub>2</sub> mole fraction	$x\text{CO}_2$	ppm	CO <sub>2</sub> atmospheric inversion from the Copernicus Atmosphere Monitoring Service (CAMS, 1985-2021)	3-hourly, 1.9° × 3.75°	Chevallier et al. (2005, 2010); Chevallier (2013)
8. $p\text{CO}_2$ climatology	$p\text{CO}_2^{\text{clim}}$	$\mu\text{atm}$	Lamont Doherty Earth Observatory (LDEO, climatology)	monthly, 4° × 5°	Takahashi et al. (2009)
9. Nitrate 10. Silicate 11. Phosphate	NO <sub>3</sub> SiO <sub>2</sub> PO <sub>4</sub>	$\mu\text{mol kg}^{-1}$	World Ocean Atlas 2018 (WOA18, climatologies)	monthly, 1°	Garcia et al. (2019)

\* Last access was on 15/4/2022 for all input databases except for SOCATv2022 data (17/6/2022) and WOA18 data (30/7/2022).

\*\* Data products 1-8 are used in the  $p\text{CO}_2$  reconstruction. Products 2-3 and 9-11 are used to compute  $A_T$ , DIC,  $p\text{H}$ ,  $\Omega_{ar}$ , and  $\Omega_{ca}$ .

\*\*\* Global values of product uncertainty (data<sup>†</sup> or analysis errors expressed as  $\sigma$ ) have been reported for  $f\text{CO}_2^{\ddagger}$  (< 5  $\mu\text{atm}$ ), SST (0.15 °C), SSS (0.2 PSU), SSH (0.02 m), MLD (-), CHL-*a* (0.03  $\text{mg m}^{-3}$ ),  $x\text{CO}_2$  (-),  $p\text{CO}_2^{\text{clim}}$  (10  $\mu\text{atm}$ ), NO<sub>3</sub> (1.8  $\mu\text{mol kg}^{-1}$ ), SiO<sub>2</sub> (3.6  $\mu\text{mol kg}^{-1}$ ), PO<sub>4</sub> (0.12  $\mu\text{mol kg}^{-1}$ ).

155 With the exception of  $x\text{CO}_2$ , nutrient concentrations, and  $p\text{CO}_2^{\text{clim}}$ , these input data products have original resolutions equivalent to or even finer than a spatial resolution of 0.25° and a temporal resolution of monthly. When mismatches in data resolutions appear, input data products are interpolated to fit the pre-defined model resolutions. The datasets of SST and  $x\text{CO}_2$  - the two key variables driving global  $p\text{CO}_2$  changes (Bates et al., 2014; Gruber et al., 2019; Landschützer et al., 2019; Chau et al., 2022b; Friedlingstein et al., 2022) - cover the full learning period and the whole globe as expected. The other predictor data are not available before the 1990s, when new types of satellite measurements started, and one of them (i.e., Chl-*a*) does not cover the high latitudes of the winter hemisphere. We therefore gap-fill the time series in an ad hoc manner, as in previous studies (Landschützer et al., 2016; Gregor et al., 2019; Chau et al., 2022b). Monthly climatologies of SSS, Chl-*a*, and MLD computed on the available data are used for each missing year. Likewise, climatologies plus linear trends of SSH following global warming effects serve for the pre-1993 period. Missing Chl-*a* data in the high latitudes of the winter hemisphere are replaced by the minimum concentration of Chl-*a* over the available data for the same grid cell (~0.01  $\text{mg m}^{-3}$ ). WOA18 nutrients and LDEO  $p\text{CO}_2^{\text{clim}}$  are already climatologies per se and we apply them for all the analysis years 1985-2021. The

160

sine function is applied to convert latitude while both the sine and cosine are used to transform longitude to conserve their periodical behaviors.

165 CO<sub>2</sub> fugacity from [Surface Ocean CO<sub>2</sub> Atlas version 2022 \(SOCATv2022, Bakker et al., 2022\)](#) SOCATv2022 (Bakker et al., 2022) is used as the target data in our monthly  $p\text{CO}_2$  reconstructions. The SOCAT project collects and qualifies underway observations via international vessels, moorings, or autonomous platforms. It grids the observations at spatial resolutions of 1° or 0.25° resulting in the two major SOCAT gridded data products. The temporal resolution of these two products is monthly. While the 1°-data product (SOCATv2022r100) covers the global ocean, the 0.25° covers solely the coastal regions. The SOCAT coastal areas is within 400 km from the shoreline (Sabine et al., 2013; Bakker et al., 2016); see Fig. A1a for an illustration. To merge the two resolutions, we first duplicate the 1°-open-ocean SOCATv2022 data ( $\sim 2 \times 10^5$  data points) over its sixteen 0.25° sub-cells. This 0.25°-open-ocean data are then combined with the 0.25°-coastal-ocean SOCATv2022 data ( $\sim 4 \times 10^5$  data points) to generate a global monthly 0.25° ocean data product fed to our reconstruction model of  $p\text{CO}_2$  (Sect. 3.1). The merged SOCATv2022 product at monthly, 0.25° resolutions is referred to as SOCATv2022r025 hereafter. The assumption of open-ocean data homogeneity of  $p\text{CO}_2$  within 1°-grid boxes ( $\sim 100 \text{ km} \times 100 \text{ km}$ ) does not degrade the reconstruction skill over the global open ocean (see Sect. 5 for results) where  $p\text{CO}_2$  observations are spatially auto-correlated within a global median distance of  $400 \pm 250 \text{ km}$  (Jones et al., 2012). The data distribution of SOCATv2022 CO<sub>2</sub> fugacity before and after combining is shown in Fig. A2 and Table 3.

## 2.2 Product qualification and comparison

180 The monthly, 0.25°-resolution reconstructions of carbonate system variables are qualified with gridded observation-based datasets and in-situ time series which are not used in our model fitting (Table 2).

- The SOCAT data in each reconstruction month are excluded from the model fitting, which avoids overfitting and ensures fairness in the model evaluation (Chau et al., 2022b). The global monthly, 0.25°-resolution CMEMS-LSCE-FFNN  $p\text{CO}_2$  fields [at a spatial resolution of 0.25°](#) can therefore be evaluated against the  $p\text{CO}_2$  data converted from SOCATv2022 CO<sub>2</sub> fugacity [\(Eq. A2\)](#) (Körtzinger, 1999) at the same resolution. Doing this, CMEMS-LSCE-FFNN  $p\text{CO}_2$  is assessed with more than  $32 \times 10^5$  open-ocean data and  $4 \times 10^5$  coastal-ocean data (Table 3). The SOCATv2022 [measurements data](#) have low random uncertainty (2-5  $\mu\text{atm}$ ) but the spatio-temporal sampling bias from the month and grid centers is significant (Bakker et al., 2016). The 0.25°-data reconstruction is also compared to its previous version with a spatial resolution of 1° (Chau et al., 2022a, b).
- The monthly, 0.25° reconstructions of  $A_T$ , DIC, and  $p\text{H}$  are qualified based on [Global Ocean Data Analysis Project bottle data version 2.2022 \(GLODAPv2.2022, Lauvset et al., 2022a\)](#) GLODAPv2.2022 data (Lauvset et al., 2022b). GLODAP provides non-gridded datasets of ocean carbon variables which have been compiled and bias-corrected from water samples taken at various depths. The measurement uncertainty is  $4 \mu\text{mol kg}^{-1}$  in  $A_T$  and DIC and between 0.01 – 0.02 in  $p\text{H}$  (Lauvset et al., 2022a). Only direct measurements at depths shallower than 10 m and with a flag of 2 (best quality control) are selected for this evaluation. Measurements in each box of 1month  $\times$  10m  $\times$  0.25°  $\times$  0.25° are averaged to

obtain representative data of surface  $A_T$ , DIC, and  $pH$  at  $0.25^\circ$ -grid cells for months in the period 1985-2021. This results in roughly  $16 \times 10^3$  data points for  $A_T$  and DIC over the global ocean (Table 4). Only half of that amount stems from direct  $pH$  measurements. Another half, referred to as indirect measurements (i.e.,  $pH$  calculated with  $A_T$  and DIC), is excluded from this data evaluation. Over 30% of ~~these GLODAP~~ data are distributed along the coasts. ~~The number of the GLODAPv2.2022 gridded data (Table 4) is much less than the SOCATv2022 gridded data (Table 3).~~

- In situ time series of direct measurements of carbonate system variables ( $pCO_2$ ,  $A_T$ , DIC, and  $pH$ ) are used to qualify our product at local scale (Table 2).

a) Sutton et al. (2019) present data over multiple sites equipped with autonomous moorings measuring surface ocean  $pCO_2$  and  $pH$  ~~from the open ocean to the continental shelves~~ since 2004. These ~~time series were used to qualify the CMEMS-LSCE-FFNN reconstruction in Chau et al. (2022b). This study only revisits eight coastal sites (Table A2) where both stations cover a wide range of oceanic conditions from the tropics to high latitudes (Fig. A1b). More than half of 42 stations distribute over the continental shelves, and many of them observe  $pCO_2$  and  $pH$  have been measured and the  $1^\circ$ -reconstruction poorly constrains most of these measurements (see later in Sect. 3.1). The eight stations are located along the US coast, the Gulf of Mexico, and in a Caribbean coral reef (Fig. A1b and Table A2 in the regime of coral reefs (Tables A2 and A3). Measurement uncertainty is up to  $2 \mu atm$  reported for  $pCO_2$  and 0.02 for  $pH$ .~~

b) For  $A_T$  and DIC, we consider ~~four time series~~: (1) eight time series accessible to provide insights into changes in the surface ocean carbonate system over the recent decades (Bates et al., 2014; Coppola et al., 2020; Gregor and Gruber, 2021; Péro et al., 2022). Two of them belong to the biogeochemical observing systems located in the subtropical Atlantic: Bermuda Atlantic Time Series (BATS, Michaels and Knap, 1996; Steinberg et al., 2001), (2) Atmospheric Flux Dynamics Time Series in the Mediterranean (DYFAMED, Coppola et al., 2021), (3) and European Station for Time-Series in the Ocean Canary islands (ESTOC, González-Dávila and Santana-Casiano, 2009), and (4) Hawaii Ocean Time-series (HOT, Dore et al., 2009). The other two provide direct measurements in the same ocean basin but in the subpolar region: Irminger Sea and Iceland Sea (IRMINGER and ICELAND, Olafsson et al., 2010). Further mentions time series distributed in specific conditions including a high-Arctic fjord in Svalbard (AWIPEV, Fischer et al., 2017; Gattuso et al., 2023) and the Mediterranean basin (DYFAMED, Coppola et al., 2021). Hawaii Ocean Time-series (HOT, Dore et al., 2009) and OISO time series in the North Pacific (Fig. A1b). These long-term time series provide insights into changes in the surface ocean carbonate system over the recent decades (Bates et al., 2014; Coppola et al., 2020; Gruber et al., 2022). The subpolar Southern Ocean (KERFIX, Metzl and Lo Monaco, 1998) allow to complete the reconstruction qualification in different ocean basins. Measurement uncertainty (f.i., from replication experiments) reported at these sites is below  $3 \mu mol kg^{-1}$ . Except for HOT and ESTOC ~~stations provide surface ocean observations of  $A_T$  and DIC, and BATS and DYFAMED data (surface ocean measurements)~~, data over all the stations are extracted at

seawater depth shallower than 10 m. A monthly average is applied for all the mentioned time series in order to be compatible with output from the CMEMS-LSCE chain of models.

**Table 2.** Data sources used in product evaluation and comparison. Values in brackets of each variable present measurement-based data uncertainty or analysis<sup>‡</sup> uncertainty.

	Product	Data type	Evaluation variables	Reference
Global ocean	1. Surface Ocean CO <sub>2</sub> Atlas version 2022 (SOCATv2022, 1985-2021), last access: 17/6/2022	observation-based gridding, resolution: 1° (global ocean) and 0.25° (coastal ocean), monthly	$p\text{CO}_2$ (< 5 $\mu\text{atm}$ )	Bakker et al. (2022)
	2. CMEMS global ocean surface carbon product (MULTI-OBS_GLO_BIO_CARBON_SURFACE_REP_015_008, 1985-2021), last access: 05/12/2022	SOCAT-based reconstruction, resolution: 1°, monthly	$p\text{CO}_2^{\ddagger}$ (8 $\mu\text{atm}$ )	Chau et al. (2022a, b)
Time series stations	3. Global Ocean Data Analysis Project bottle data version 2.2022 (GLODAPv2.2022, 1985-2021), last access: 30/9/2022	observation	$A_T$ (4 $\mu\text{mol kg}^{-1}$ ) DIC (4 $\mu\text{mol kg}^{-1}$ ) $p\text{H}$ (0.01 – 0.02)	Lauvset et al. (2022a, b)
	4. Autonomous time series from surface buoys since 2004 (see details in Table A2), last access: 15/10/2022	observation	$p\text{CO}_2$ (2 $\mu\text{atm}$ ) $p\text{H}$ (0.02)	Sutton et al. (2019)
	5. Bermuda Atlantic Time Series (BATS: 31.7°N-64.2°W, 1988-2021), last access: 30/10/2022	observation	$A_T$ (3 $\mu\text{mol kg}^{-1}$ ) DIC (1 $\mu\text{mol kg}^{-1}$ )	Michaels and Knapp (1996); Steinberg et al. (2001)
	6. Atmospheric Flux Dynamics Time Series in the Mediterranean (DYFAMED: 43.5°N-7.9°E, 1998-2017), last access: 23/03/2023	observation	$A_T$ , DIC	Coppola et al. (2020, 2021)
	7. European Station for Time-Series in the Ocean Canary islands (ES-TOC: 29.2°N-15.5°W, 1995-2009), last access: 30/10/2022	observation	$A_T$ , DIC	González-Dávila and Santana-Casiano (2009)
	8. Hawaii Ocean Time-series (HOT: 22.5°N-158.1°W, 1988-2020), last access: 30/10/2022	observation	$A_T$ , DIC	Dore et al. (2009)
	9. Underwater observatory in Spitsbergen (Svalbard) (AWIPEV: 78.93°N-11.92°E, 2015-2020), last access: 20/07/2023	observation	$A_T$ (2.6 $\mu\text{mol kg}^{-1}$ ) DIC (3 $\mu\text{mol kg}^{-1}$ )	Fischer et al. (2017); Gattuso et al. (2023)
	10. Irminger Sea and the Iceland Sea time series (IRMINGER: 64.33°N-28.00°W, 2014-2021; ICELAND: 68.00°N-12.67°W, 2014-2021), last access: 20/07/2023	observation	$A_T$ (-) DIC (2 $\mu\text{mol kg}^{-1}$ )	Olafsson et al. (2010)
	11. Southern Ocean time series under the OISO project (KERFIX: 50.67°S-68.42°E, 1992-2018), last access: 20/07/2023	observation	$A_T$ (3 $\mu\text{mol kg}^{-1}$ ) DIC (3 $\mu\text{mol kg}^{-1}$ )	Metzl and Lo Monaco (1998)

### 3 Reconstruction methods

#### 3.1 Ensemble $p\text{CO}_2$ mapping feed-forward neural networks

The CMEMS-LSCE-FFNN (Chau et al., 2022b) is based on an ensemble of 100 feed-forward neural network (FFNN) models mapping SOCAT  $\text{CO}_2$  fugacity ( $f\text{CO}_2$ ) and predictor variables (Eq. 1).

$$235 \quad f\text{CO}_2 = \text{FFNN}(\text{SST}, \text{SSS}, \text{SSH}, \text{Chl} - a, \text{MLD}, x\text{CO}_2, f\text{CO}_2^{\text{clim}}, \text{latitude}, \text{longitude}) \quad (1)$$

The predictors of  $f\text{CO}_2$  include sea surface temperature (SST), salinity (SSS), surface height (SSH), chlorophyll- $a$  (Chl- $a$ ), mixed layer depth (MLD),  $\text{CO}_2$  mole fraction ( $x\text{CO}_2$ ),  $f\text{CO}_2$  climatologies ( $f\text{CO}_2^{\text{clim}}$ ), and the geographical coordinates (latitude and longitude). The datasets of SOCAT  $f\text{CO}_2$  and predictors are first reprocessed to match model fitting requirements (Sect. 2.1). After excluding the data in the reconstruction month, the data within the 3-month window are randomly separated  
240 into FFNN training and validation subsets with a ratio of 2 : 1. The subsampling process is repeated for each 100 FFNN runs that results in 100 different datasets for model fitting. The excluded SOCATv2022 datasets are used in model evaluation. The CMEMS-LSCE-FFNN approach was originally developed for  $p\text{CO}_2$  reconstructions at monthly,  $1^\circ$  resolutions where  $p\text{CO}_2$  is converted from  $f\text{CO}_2$  following the formulation by Körtzinger (1999). The model best estimate and its uncertainty are defined as the ensemble mean ( $\mu$ ) and ensemble spread ( $\sigma$ ) of 100 model outputs of  $p\text{CO}_2$ .

245 This study slightly modifies the CMEMS-LSCE-FFNN ensemble approach by Chau et al. (2022b) to achieve  $p\text{CO}_2$  reconstructions at monthly,  $0.25^\circ$  resolutions. Some of the input datasets presented here (Table 1) are different from those presented in Chau et al. (2022b) (Table S1). The up-to-date input datasets have higher resolutions and a better coverage over the coastal ocean as well as in the high latitudes. Furthermore, the new CMEMS data resources offer space-time varying uncertainty fields which are important in quantifying carbonate system variable uncertainties.

250 For comparable evaluations in this study, we execute 100-member ensembles of FFNN models at spatial resolutions of both  $1^\circ$  (FFNNr100) and  $0.25^\circ$  (FFNNr025) using the same lot of input data resources (Table 1). Remind that the training data of  $f\text{CO}_2$  is extracted from the SOCATv2022r100 product for FFNNr100 while it comes from the SOCATv2022r025 product (i.e., the merged product of the  $1^\circ$ -open-ocean dataset and the  $0.25^\circ$ -coastal-ocean dataset) for FFNNr025. All input datasets are reprocessed with respect to each model resolution (Sect. 2.1). Sect. 3.1 compares these two CMEMS-LSCE-FFNN versions  
255 and highlights the skill of the finer resolution data product.

#### 3.2 Locally interpolated alkalinity regression

Locally interpolated alkalinity regression (LIAR; Carter et al., 2016, 2018) is an ensemble-based regression method developed for the global reconstruction of total alkalinity ( $A_T$ ). Regression coefficients were learned on GLODAPv2 data (Olsen et al., 2016) binned within regular windows of  $5^\circ \times 5^\circ$ . For prediction, the LIAR software interpolates between the regression  
260 coefficients to arbitrary resolutions specified by the users. This study employs eight LIAR models (Carter et al., 2018, Table 2) for calculating  $A_T$  at monthly,  $0.25^\circ$  resolutions. Each model represents a combination of predictor variables (see the full

presentation in Eq. 2),

$$A_T = \text{LIAR}(\text{SSS}, \text{SST}, \text{NO}_3, \text{SiO}_2). \quad (2)$$

The eight regression models include salinity (SSS) - the predominant predictor of  $A_T$  - and some combinations of temperature (SST), nitrate ( $\text{NO}_3$ ), and silicate ( $\text{SiO}_2$ ). The model which has the smallest propagation uncertainty is chosen to provide the best estimate of  $A_T$ .

Global monthly total alkalinity and  $1\sigma$ -uncertainty are estimated with given input data from the monthly CMEMS SSS and SST fields and from the WOA18 datasets of nutrient concentrations (Table 1). Uncertainty of the  $A_T$  field is estimated systematically through input uncertainty propagation along the processing chain (Carter et al., 2018). Here we define the input uncertainty of predictors in terms of standard deviations ( $1\sigma$ ). Input uncertainty fields associated to the monthly CMEMS SSS and SST are products' analysis errors (see e.g., Fig. A7) while uncertainties of the WOA18  $\text{NO}_3$  and  $\text{SiO}_2$  climatologies are set to 15% of data values per cell. The 15% quantity refers to the median percentage of standard analysis errors against climatological means of nutrient concentrations (see product standard errors in Table 6, Garcia et al., 2019). The WOA18 standard analysis errors are defined as misfits between their interpolated data and GLODAPv2 bottle data (Olsen et al., 2016). Spatial distribution of the error percentage of the WOA18 nutrient concentrations at the ocean surface is illustrated in Fig. A8.

### 3.3 Carbonate system speciation

The CO2SYS speciation software was first developed by Lewis and Wallace (1998) to determine carbonate system parameters in the marine  $\text{CO}_2$  system based on a set of equilibrium equations (Dickson et al., 2007). Here we use the speciation program written by Van Heuven et al. (2011) and its extension with uncertainty propagation proposed by Orr et al. (2018). To obtain a complete description of the ocean carbonate system, the CO2SYS initialization requires the following input conditions:

- i) values of any couple of the parameters  $p\text{CO}_2$ ,  $A_T$ , DIC, and  $p\text{H}$ ,
- ii) temperature and pressure,
- iii) total concentrations of all the non- $\text{CO}_2$  acid-base systems,
- iv) equilibrium constants used to describe seawater acid-base chemistry.

The (iii)-condition involves total concentrations of both conservative and non-conservative constituents in the non- $\text{CO}_2$  acid-base systems. The amount of conservative constituents such as borate, fluoride, and sulfate in surface seawater is estimated with salinity. The total concentration of non-conservative constituents (nutrients) is computed approximately with silicate ( $\text{SiO}_2$ ), and phosphate ( $\text{PO}_4$ ). Further information of the carbonate system speciation can be found in Dickson et al. (2007) and Dickson (2010).

With the reconstructions of  $p\text{CO}_2$  and  $A_T$  (Sects 3.1 and 3.2), the CO2SYS speciation software is used to derive  $p\text{H}$ , DIC,  $\Omega_{ar}$ , and  $\Omega_{ca}$ , and determine their uncertainty over the ocean surface at a resolution of  $0.25^\circ$ . Equation 3 expresses all input-output variables of CO2SYS for this study. Note that the estimates for other carbonate system variables such as hydrogen ion



( $H^+$ ) concentration and Revelle Factor (RF) - a measure of the carbonate buffer capacity- are also available (Figs. A4 and A6) but beyond the scope of our data evaluation.

$$295 \quad pH, DIC, \Omega_{ar}, \Omega_{ca} = CO2SYS(pCO_2, A_T, SST, SSS, P, SiO_2, PO_4, constants) \quad (3)$$

The FFNN best estimate (ensemble mean) of  $pCO_2$  reconstructions (Sect. 3.1) and the LIAR outputs of  $A_T$  (Sect. 3.2) are used as the prior inputs of the CO2SYS at each grid cell for every month in the period 1985-2021. We take the same data products of SST, SSS, and nutrient concentrations as for the previous reconstructions (Table 1). Pressure (P) is assumed to be 0 dbar at the ocean surface. For equilibrium constants, we choose the best empirical values recommended by Dickson et al. (2007) and  
 300 Dickson (2010). These settings include (1) the dissociation constants  $K_1$  and  $K_2$  from Lueker et al. (2000) and  $K_{HSO_4}$  from Dickson (1990) in combination with the total boron-ratio-salinity formulation by Uppstrom (1974).

The uncertainty of the CO2SYS variables is estimated by error propagation (Orr et al., 2018). Inputs for the CO2SYS error propagation include the reconstruction uncertainty of  $pCO_2$  (FFNN ensemble standard deviation) and of  $A_T$  (LIAR error propagation). The uncertainty of SST, SSS, and nutrient concentrations are set to the same values as in the previous section  
 305 (Sect. 3.2). Equilibrium constants' standard errors are default values (see Table 1, Orr et al., 2018). As for FFNN and LIAR, uncertainty values of each carbonate system variable are computed for each month in 1985-2021 and at each  $0.25^\circ$ -grid box over the global surface ocean.

## 4 Evaluation metrics

### 4.1 Model best estimate and uncertainty quantification

310 The 100 FFNN models result in an ensemble of 100 estimates of global monthly,  $0.25^\circ$  surface ocean  $pCO_2$  fields (Sect. 3.1). Specify any  $t = 1 : 444$  (month),  $i = 1 : 180$  (latitude), and  $j = 1 : 360$  (longitude), the best estimate ( $\mu_{tij}$ ) and uncertainty ( $\sigma_{tij}$ ) at time  $t$  and grid cell  $ij$  are deduced from 100 FFNN  $pCO_2$  estimates ( $X(t, i, j, m)$ ) $_{m=1}^{m=100}$  as follows.

$$\mu_{tij} = \frac{\sum_{m=1}^{m=100} X(t, i, j, m)}{100}, \quad (4a)$$

$$\sigma_{tij} = \sqrt{\frac{\sum_{m=1}^{m=100} [X(t, i, j, m) - \mu_{tij}]^2}{100}}. \quad (4b)$$

315 For  $pH$ ,  $A_T$ ,  $DIC$ ,  $\Omega_{ar}$ , and  $\Omega_{ca}$ , the best estimates and associated uncertainties ( $\mu_{tij}$  and  $\sigma_{tij}$ ) are obtained directly from the LIAR and CO2SYS speciation tools and their error propagation (Sects. 3.2 and 3.3).

To assign representatives of  $\mu$  and  $\sigma$  estimates for carbonate system variables at a specific space-time window, we define statistics with respect to each of the three following cases:

i) a representative over a period of time ( $T$  months)

$$320 \quad \mu_{ij} = \frac{\sum_t \mu_{tij}}{T}, \quad (5a)$$

$$\sigma_{ij} = \sqrt{\frac{\sum_t \sigma_{tij}^2}{T}}. \quad (5b)$$

ii) a representative over a region (e.g., ocean basins and sub-basins, the global ocean)

$$\mu_t = \frac{\sum_{ij} \mu_{tij} \times A_{ij}}{\sum_{ij} A_{ij}}, \quad (6a)$$

$$\sigma_t = \sqrt{\frac{\sum_{ij} \sigma_{tij}^2 \times A_{ij}}{\sum_{ij} A_{ij}}}. \quad (6b)$$

325 iii) a representative over a period of time and a region

$$\mu = \frac{\sum_{t,ij} \mu_{tij} \times A_{ij}}{T \times \sum_{ij} A_{ij}}, \quad (7a)$$

$$\sigma = \sqrt{\frac{\sum_{t,ij} \sigma_{tij}^2 \times A_{ij}}{T \times \sum_{ij} A_{ij}}}. \quad (7b)$$

where  $t$  is in a time period with length  $T$  and  $A_{ij}$  is the area of each grid cell in a desired region. It is noteworthy that the statistics in Eqs. (5b)-(7b) are not the standard deviation associated to the mean quantities in Eqs. (5a)-(7a), but they stand for the best representative of uncertainty estimates over an ocean basin and/or time period. These statistics also support for the comparison with model-observation deviation (e.g., Eq. 10) which is typically used in the calculation of standard uncertainty proposed in the previous studies (Jiang et al., 2019; Iida et al., 2021; Gregor and Gruber, 2021). Subscripts in the notations of the best model estimates ( $\mu$ ) and model uncertainties ( $\sigma$ ) in Eqs. (4)-(7) are dropped out for general situations.

The best model estimate ( $\mu$ ) is assessed against model uncertainty ( $\sigma$ ) through  $\sigma$ -to- $\mu$  ratio (%)

$$335 \quad R(\mu, \sigma) = 100\% \frac{\sigma}{|\mu|}. \quad (8)$$

The  $\sigma$ -to- $\mu$  ratio allows evaluating the significance of the model estimate. Model estimates of carbonate variables are reliable with  $R(\sigma, \mu)$  values less than 20% (Rose, 2013).

## 4.2 Model performance in comparison with evaluation data

Assume that  $\mu_{tij}$  and  $O_{tij}$  are the best model estimate and an observation (or its gridded data) available at time  $t$  and grid cell  $ij$ , and  $\mu$  and  $O$  are respectively their means over the total number of evaluation data ( $N$ ). Model skills are assessed against observation data (Table 2) with the following metrics:

- mean model-observation differences (Bias)

$$\text{Bias} = \frac{\sum_{t,ij} (\mu_{tij} - O_{tij})}{N}, \quad (9)$$

- root-of-mean-square-deviation (RMSD)

$$345 \quad \text{RMSD} = \sqrt{\frac{\sum_{t,ij} (\mu_{tij} - O_{tij})^2}{N}}, \quad (10)$$


- coefficient of determination ( $r^2$ )

$$r^2 = \frac{\left[ \sum_{t,ij} (\mu_{tij} - \mu) \times (O_{tij} - O) \right]^2}{\sum_{t,ij} (\mu_{tij} - \mu)^2 \times \sum_{t,ij} (O_{tij} - O)^2}. \quad (11)$$

## 5 Results

### 5.1 Surface ocean $p\text{CO}_2$

350 This section presents the reconstruction of surface ocean  $p\text{CO}_2$  at monthly and  $0.25^\circ$  resolutions. The reconstruction skill is evaluated against SOCATv2022 data from global to ocean basin scale and at the level of grid cells (Sect. 2.2). We compare the novel reconstruction at a higher spatial resolution to the one at a coarser spatial resolution (Chau et al., 2022b). Emphasis is put on the skill to reproduce spatial and temporal variations of  $p\text{CO}_2$  across a variety of coastal regions and time series stations.



**Figure 1.** CMEMS-LSCE-FFNN  $p\text{CO}_2$  over the global ocean at a spatial resolution of  $0.25^\circ$ . Temporal means of the model best estimate and  $1\sigma$ -uncertainty per grid cell over 1985-2021 are calculated by using Eq. (5).

**Table 3. Skill scores-Evaluation statistics** for monthly CMEMS-LSCE-FFNN reconstructions of  $p\text{CO}_2$  at  $1^\circ$  (r100) and  $0.25^\circ$  (r025) spatial resolutions computed over the period 1985-2021. r100  $\rightarrow$  025 and r025  $\rightarrow$  100 are referred to the versions ~~upsealed or~~ downscaled or ~~upscaled~~ from the original CMEMS-LSCE-FFNN  $p\text{CO}_2$  at  $1^\circ$  and  $0.25^\circ$  resolutions. SOCATv2022 gridded data independent from CMEMS-LSCE-FFNN training are used as benchmarks for model evaluation (see text for details). Statistics including total numbers of data, RMSD (Eq. 10), and  $r^2$  (Eq. 11) are reported for both the open ocean (O) and coastal region (C). \* marks results with respect to the primary product proposed in this study.

Basins			Number of data		RMSD [ $\mu\text{atm}$ ]				$r^2$			
			r100	r025*	r100	r025 $\rightarrow$ 100	r025*	r100 $\rightarrow$ 025	r100	r025 $\rightarrow$ 100	r025*	r100 $\rightarrow$ 025
0.	Globe	(O)	207174	3317273	<del>14.32</del> 14.3	<del>14.08</del> 14.1	<del>14.29</del> 14.3	<del>14.38</del> 14.4	0.83	0.83	0.83	0.83
		(C)	101007	431758	<del>26.61</del> 26.6	<del>26.48</del> 26.5	<del>27.55</del> 27.6	<del>28.50</del> 28.5	0.72	0.72	0.74	0.72
1.	Arctic	(O)	537	8589	<del>27.93</del> 27.9	<del>27.43</del> 27.4	<del>28.04</del> 28.0	<del>28.06</del> 28.1	0.69	0.69	0.67	0.67
		(C)	5897	25844	<del>38.74</del> 38.7	<del>38.56</del> 38.6	<del>41.46</del> 41.5	<del>43.17</del> 43.2	0.55	0.56	0.55	0.52
2.	Atlantic	(O)	54797	876116	<del>13.76</del> 13.8	<del>13.57</del> 13.6	<del>13.69</del> 13.7	<del>13.78</del> 13.8	0.81	0.81	0.81	0.81
		(C)	49770	227665	<del>24.99</del> 25.0	<del>24.78</del> 24.8	<del>25.17</del> 25.2	<del>26.05</del> 26.1	0.76	0.76	0.77	0.77
3.	Pacific	(O)	120604	1932981	<del>14.59</del> 14.6	<del>14.30</del> 14.3	<del>14.54</del> 14.5	<del>14.67</del> 14.7	0.85	0.85	0.85	0.85
		(C)	26847	104269	<del>26.79</del> 26.8	<del>26.90</del> 26.9	<del>28.46</del> 28.5	<del>28.95</del> 29.0	0.71	0.71	0.69	0.67
4.	Indian Ocean	(O)	4485	71719	<del>10.34</del> 10.3	<del>10.17</del> 10.2	<del>10.26</del> 10.3	<del>10.34</del> 10.3	0.88	0.88	0.88	0.88
		(C)	1522	6187	<del>23.50</del> 23.5	<del>22.82</del> 22.8	<del>25.40</del> 25.4	<del>26.51</del> 26.5	0.69	0.71	0.69	0.69
5.	Southern Ocean	(O)	26751	427868	<del>14.42</del> 14.4	<del>14.29</del> 14.3	<del>14.52</del> 14.5	<del>14.43</del> 14.4	0.69	0.69	0.69	0.69
		(C)	16971	67793	<del>26.01</del> 26.0	<del>25.80</del> 25.8	<del>27.35</del> 27.4	<del>28.80</del> 28.8	0.61	0.61	0.64	0.59

Figure 1 presents global maps at  $0.25^\circ$ -resolution of long-term averages of  $p\text{CO}_2$  and corresponding uncertainty estimates. Reconstructed  $p\text{CO}_2$  distributions reveal well documented large scale structures. Values are high over upwelling regions (e.g., Equatorial Pacific, California Boundary Current, Western Arabian Sea). Low  $p\text{CO}_2$  is associated with increased  $\text{CO}_2$  solubility in cold high latitudes seawater (e.g., Arctic), strong biological production (e.g., China Sea), or the combination of both (e.g., subpolar Northern Atlantic, Southern Ocean between  $35 - 50^\circ\text{S}$ ). Spatial structures appear coherent from small to large spatial scales, both along the coast and moving towards the open ocean (see also in Figs. 2-4). The combination of a down-scaled version of open-ocean and higher-resolution coastal SOCATv2022 data (Sect. 2.1) yields  $p\text{CO}_2$  distributions without discontinuities. The uncertainty map (Fig 1b) represents the confidence level in surface ocean  $p\text{CO}_2$  estimates (Fig 1a). Predominantly low uncertainty estimates ( $\sigma < 5 \mu\text{atm}$ ) indicate the global stability of the ensemble reconstruction. Exceptions are found in many coastal regions, open-ocean areas with sparse data coverage (e.g., Southern Pacific, Indian Ocean), or regions with substantially high or low surface ocean  $p\text{CO}_2$  (e.g., Arctic, eastern equatorial Pacific). However,  $p\text{CO}_2$  is reconstructed with a high degree of confidence over most of the global ocean with a  $\sigma$ -to- $\mu$  ratio (Eq. 8) below 10% (Fig. A9a).

Skill scores of the monthly,  $0.25^\circ$ -resolution reconstruction are presented in Table 3 (columns marked by an asterisk). The global RMSD (Eq. 10) between the best reconstruction and SOCATv2022r025  $p\text{CO}_2$  over the entire period is ~~14.29~~14.3  $\mu\text{atm}$  for the open ocean and ~~27.55~~27.6  $\mu\text{atm}$  for the coastal ocean. These two model errors are lower than 4% and 8% of the global mean  $p\text{CO}_2$  (Table 6). Moreover, variability present in observation-based data is reproduced by the CMEMS-LSCE-FFNN with high values of  $r^2$  (open ocean: 0.83, coast: 0.74). The reconstruction quality is similar among major ocean basins. Spatial distributions of SOCATv2022 data, bias, and RMSD are shown in Figs. A2-bd and A3-bdfh. Estimation skills are low in the

ocean basins with sparse data coverage and significant space-time variability of  $p\text{CO}_2$  (e.g., Arctic, eastern Equatorial Pacific, land-ocean continuum).

Table 3 also presents statistics for the monthly FFNN products of surface ocean  $p\text{CO}_2$  at spatial resolutions of  $0.25^\circ$  (r025) and  $1^\circ$  (r100) together with their variants (r100  $\rightarrow$  025 and r025  $\rightarrow$  100). The latter are respectively extrapolation and interpolation versions of the original r100 and r025 datasets, i.e.,. We used the Climate Data Operators (CDO) remap operator to regrid FFNN model outputs regridded to a finer or coarser spatial resolution. For compatibility, we compare statistics between:

- i) FFNN(r025) and FFNN(r100  $\rightarrow$  025) by using SOCATv2022r025 as evaluation data,
- ii) FFNN(r025  $\rightarrow$  100) and FFNN(r100) by using SOCATv2022r100 as evaluation data.


The FFNN(r025) central to this study yields a lower RMSD and a higher correlation to the SOCAT data than the FFNN(r100  $\rightarrow$  025). As expected, the improvement in reconstruction skill with higher model resolution is larger over coastal regions than in the open ocean. The FFNN(r025) product after interpolating to a coarser resolution, i.e., FFNN(r025  $\rightarrow$  100), agrees with the original  $1^\circ$ -resolution data product over all the ocean.

The motivation to increase the spatial resolution of the reconstruction is to improve the representation of horizontal gradients of  $p\text{CO}_2$  at fine scales. Figures 2-4 exemplify spatial distributions for the two reconstructions (r025 and r100) over the coastal-open-ocean continuum. Ten distinct oceanic regions are considered (see Fig. A1a and Table A1 for the ten locations), which can be classified into three groups:


- permanent Eastern Boundary current upwelling systems with relatively high  $p\text{CO}_2$  (California Current System - CCS, Humboldt Current System - HCS, Canary Current System - CnCS, and Benguela Current System - BCS),
- regions characterized by low  $p\text{CO}_2$  values driven by cold water temperatures and strong biological production (Labrador Sea, Western South Atlantic, Northern Europe, and Sea of Japan),
- other regions either under the influence of strong river runoff (Amazon mouth) or monsoon-driven upwelling (Western Arabian Sea).

The legend of Figs 2-4 includes regional RMSD and  $r^2$  computed between the best estimates of two models and coastal-ocean SOCATv2022r025 data. The coarser spatial resolution product is co-located at the same  $0.25^\circ$ -grid cells for this analysis. These figures illustrate important discrepancies in  $p\text{CO}_2$  data density between coastal regions with poorly monitored regions (e.g., HCS, BCS, Amazon mouth) contrasting with areas with higher data coverage (e.g., Northern Europe, Sea of Japan).

Over 7 out of the 10 analysed regions the reconstruction at monthly,  $0.25^\circ$  resolutions yields RMSDs below 10% of the global mean of coastal-ocean  $p\text{CO}_2$  estimates (Table 6) and  $r^2$  values higher than 0.3; e.g., Northern Europe (RMSD = 33.90  $\mu\text{atm}$ ,  $r^2 = 0.80$ ), Sea of Japan (RMSD = 20.84  $\mu\text{atm}$ ,  $r^2 = 0.70$ ), and CnCS (RMSD = 30.36  $\mu\text{atm}$ ,  $r^2 = 0.35$ ). The CMEMS-LSCE-FFNN model projections of  $p\text{CO}_2$  lack skill over the HCS (RMSD = 54.54  $\mu\text{atm}$ ,  $r^2 = 0.29$ ), the region under influence of the Amazon river (RMSD = 45.93  $\mu\text{atm}$ ,  $r^2 = 0.37$ ), and the Western Arabian Sea (RMSD = 45.31  $\mu\text{atm}$ ,  $r^2 = 0.38$ ).




**Figure 2.** Comparison of CMEMS-LSCE-FFNN mapping  $p\text{CO}_2$  at  $1^\circ$  (r100) and  $0.25^\circ$  (r025) resolutions over 4 permanent upwelling regions associated with the Eastern Boundary Currents (California, [PeruHumboldt](#), Canary, and Benguela; see Figure A1-ABGH for geographical locations). For each region, spatial distributions of  $p\text{CO}_2$  ( $\mu$ ) and uncertainty ( $\sigma$ ) estimates, and coastal-ocean RMSD of  $p\text{CO}_2$  averaged over 1985-2021 (Eqs. 5 and 10) are shown. Metrics presented in the legend for each of the 3rd row include the number of coastal-ocean SOCATv2022 data ( $N$ ), regional RMSD (Eq. 10) and  $r^2$  (Eq. 11).



**Figure 3.** Comparison of CMEMS-LSCE-FFNN mapping  $p\text{CO}_2$  at  $1^\circ$  (r100) and  $0.25^\circ$  (r025) resolutions over 4 regions characterized with low  $p\text{CO}_2$  values (Labrador, Western South America Atlantic, Northern Europe, and Japan; see Figure A1-CEFJ for geographical locations). For each region, spatial distributions of  $p\text{CO}_2$  ( $\mu$ ) and uncertainty ( $\sigma$ ) estimates, and coastal-ocean RMSD of  $p\text{CO}_2$  averaged over 1985-2021 (Eqs. 5 and 10) are shown. Metrics present in the legend for each of the 3rd row include the number of coastal-ocean SOCATv2022 data ( $N$ ), regional RMSD (Eq. 10) and  $r^2$  (Eq. 11).

405  $\mu\text{atm}$ ,  $r^2 = 0.47$ ,  $r^2 = 0.49$ ). In nearshore sectors of these coastal areas,  $p\text{CO}_2$  estimates are also subject to a substantial amount of uncertainty ( $\sigma > 20 \mu\text{atm}$ ). The lack of model skill reflects the combination of low data density and strong  $p\text{CO}_2$  gradients






**Figure 4.** Comparison of CMEMS-LSCE-FFNN mapping  $p\text{CO}_2$  at  $1^\circ$  (r100) and  $0.25^\circ$  (r025) resolutions over the mouth of the river Amazon and the Western Arabian Sea (see Fig. A1-DI for geographical locations). For each region, spatial distributions of  $p\text{CO}_2$  ( $\mu$ ) and uncertainty ( $\sigma$ ) estimates, and coastal-ocean RMSD of  $p\text{CO}_2$  averaged over 1985-2021 (Eqs. 5 and 10) are shown. Metrics present in the legend for each of the 3rd row include the number of coastal-ocean SOCATv2022 data ( $N$ ), regional RMSD (Eq. 10) and  $r^2$  (Eq. 11).

driven by multiple underlying physical and biogeochemical processes. The HCS, for instance, is characterized by the highest  $p\text{CO}_2$  levels (Fig. 2) among the four Eastern Boundary Current Systems, with interannual variability amplified with the El Niño–Southern Oscillation (ENSO) events (Feely et al., 1999; Landschützer et al., 2016). Similarly, high  $p\text{CO}_2$  levels with substantial seasonal variability are observed over the Western Arabian Sea (Fig. 4b), the key driver being monsoonal upwelling (Sabine et al., 2002; Sarma et al., 2013) (Sabine et al., 2002; Sarma et al., 2013, 2023). In contrast to the two aforementioned coastal regions, high  $\text{CO}_2$  undersaturation as well as strong  $p\text{CO}_2$  gradients (Fig. 4a) are found in the area under the influence of Amazon river discharge (Olivier et al., 2022). Extreme values and large variability of  $p\text{CO}_2$  challenge any approach to estimate  $p\text{CO}_2$  data over these regions (Ibáñez et al., 2015; Bakker et al., 2016; Landschützer et al., 2020).

The two FFNN reconstructions (r025 and r100) share similarities in overall structures of  $p\text{CO}_2$  over the coastal-open-ocean continuum (Figs. 2-4). However, the higher spatial resolution outperforms its lower resolution counterpart in reproducing fine-scale features of  $p\text{CO}_2$  in the transition from nearshore regions to the adjacent open ocean. The increase in model spatial resolution translates into a greater spatial coverage of the continental shelves such as Labrador Sea, Northern Europe, and Sea of Japan (Fig. 3), and thus an increase in the number of data over the coastal domain. The increase in spatial resolution allows



**Figure 5.** Time series of surface ocean  $p\text{CO}_2$  ( $\mu\text{atm}$ ) at coastal observing stations (Table A2 and Fig. A1b): model best estimate (curve),  $1\sigma$ -uncertainty (envelope), and monthly average of in situ observations (point). The reconstructed data at  $1^\circ$  (r100) and  $0.25^\circ$  (r025) resolutions are co-located to in situ observations provided by Sutton et al. (2019). Means of the best estimate and  $1\sigma$ -uncertainty ( $\mu \pm \sigma$ ) calculated over the observing time span are shown in brackets. Statistics include number of months with observations ( $N$ ), Bias, RMSD, and  $r^2$  computed for the two reconstructions.  $\sigma_{p\text{CO}_2}^t$  stands for temporal standard deviation from monthly averages of  $p\text{CO}_2$  observations.

a gain in prediction probability of  $p\text{CO}_2$  variations on the order of roughly 2% over the Eastern Boundary Currents to 8%–7% over the Western South Atlantic (Figs. 2-3b).

Reconstruction skill of seasonal to inter-annual variability of  $p\text{CO}_2$  at local scale, the reconstruction of in situ  $p\text{CO}_2$  is further assessed at eight coastal monitoring sites (Sutton et al., 2019) and illustrated in (Sutton et al., 2019) over the open ocean is at


the high order of confidence (Table A3). Low RMSD (between 7.8 and 23.5  $\mu\text{atm}$ ) and sustainably high  $r^2$  (from 0.45 to 0.98) dominate evaluation statistics over the 18 open-ocean stations. Obviously, CMEMS-LSCE-FFNN has less skill in the coastal sector and model-observation deviation varies depending on a wide range of  $p\text{CO}_2$  conditions. However, coastal-ocean RMSD can be smaller than 10% of station climatology (e.g., KILONALU, KANEOHE, ALAWAI) and the reproduction availability of temporal variations of  $p\text{CO}_2$  possibly exceeds 70% (e.g., SEAK, KODIAK, DABOB). Through Fig. 5 (see Sect. 2.2, we further assess seasonal to inter-annual variability reproduced at the eight coastal sites (see Fig. A1b and Table A2 for data description and Fig. A1b for station locations) station location) where measurements are available for both  $p\text{CO}_2$  and pH (analyzed in Section 5.3) and they are poorly constrained by the  $1^\circ$ -reconstruction (Chau et al., 2022b). The temporal variability of  $p\text{CO}_2$  reported for these time series sites reflects a combination of processes (Sutton et al., 2019), e.g., California Current System (CAPEARAGO and CCE2), western coastal upwelling (CAPEELIZABETH), eutrophication enhancing respiration of  $\text{CO}_2$  (FIRSTLANDING), and multiple stressors on coral reef environments (CHEECAROCKS, GREYREEF). Results from reconstructions at two spatial resolutions are compared:  $1^\circ$  (100, black curve) and  $0.25^\circ$  (r025, color curve). As shown in Fig. 5 (scattered points for observations) time series of coastal  $p\text{CO}_2$  are still short. The longest time series covers 127 months of  $p\text{CO}_2$  monitoring since 2010 (CCE2) while the shortest one contributes 17 months with observations (FIRSTLANDING).

Analyzing the eight station time series, we have found that data have been sampled within a few days with an average offset of about a week from the month center. At these coastal sites, the temporal standard deviation from monthly averages of  $p\text{CO}_2$  ( $\sigma_{p\text{CO}_2}^t$ ) exceeds measurement-analytical errors (2  $\mu\text{atm}$ , Sutton et al., 2019).  $\sigma_{p\text{CO}_2}^t$  ranges from 20.12 to 20.1  $\mu\text{atm}$  at GREYR-EFF to values as large as 65.6  $\mu\text{atm}$  at CAPEARAGO or 69.98 to 70  $\mu\text{atm}$  at FIRSTLANDING. The monthly average of  $p\text{CO}_2$  might not be adequately represented by discrete samples at sites with a large temporal standard deviation of  $p\text{CO}_2$ . The misfit between the monthly reconstruction and discrete observations is exacerbated in dynamical coastal environments and might explain in part the large RMSD of reconstructions of monthly coastal  $p\text{CO}_2$  (e.g., GREYREEF: 38.34 to 38.3  $\mu\text{atm}$ , CAPEARAGO: 79.86 to 79.9  $\mu\text{atm}$ , FIRSTLANDING: 77.32 to 77.3  $\mu\text{atm}$ ) for the r025 reconstruction. The RMSD is mostly lower for the FFNN reconstruction at  $0.25^\circ$  resolution compared to the FFNN at  $1^\circ$  resolution by 2.11 to 2.2  $\mu\text{atm}$  (CCE2) to 23.32 to 23.2  $\mu\text{atm}$  (COASTALMS). Similarly,  $r^2$  increases between 7%-23% at higher resolution. Overall, seasonal to interannual variations of coastal-ocean  $p\text{CO}_2$  are better reproduced in the reconstruction at  $0.25^\circ$  resolution (Fig. 5).

## 5.2 Total alkalinity and dissolved inorganic carbon

This section presents and analyzes global ocean surface reconstructions of total alkalinity ( $A_T$ ) and dissolved inorganic carbon (DIC) at monthly,  $0.25^\circ$  resolutions over 1985-2021. GLODAPv2.2022 bottle data (Sect. 2.2) serve as reference data for model evaluation. Model reconstruction skill is further assessed at the four-eight Eulerian time series sites: AWIPEV, BATS, DYFAMED, ESTOC, and HOTS-HOT, ICELAND, IRMINGER, and KERFIX (Table 2).


Figure 6 shows spatial distributions of the climatological mean and uncertainty (Eq. 5) for  $A_T$  and DIC. Despite being in part influenced by common biological and physical processes, both properties have contrasting distributions due to the strong correlation between surface ocean  $A_T$  and salinity (Lee et al., 2006; Broullón et al., 2019), as well as the contribution of air-sea gas exchange and biological productivity on surface ocean DIC levels (Feely et al., 2001; Takahashi et al., 2014). Over subtropical



**Figure 6.** CMEMS-LSCE  $A_T$  and DIC over the global ocean at a spatial resolution of  $0.25^\circ$ . Temporal means of the model best estimate and  $1\sigma$ -uncertainty per grid cell over 1985-2021 are calculated by using Eq. (5).

Atlantic gyres and the Mediterranean Sea, oceanic areas with net evaporation,  $A_T$  exceeds  $2400 \mu\text{mol kg}^{-1}$ . Total alkalinity falls below  $2150 \mu\text{mol kg}^{-1}$  in regions where precipitation, river freshwater runoff, or seasonal sea-ice melting dilute surface water salinity (e.g., subpolar North Pacific, Arctic, and equatorial river outflows). The distribution of DIC is relatively uniform between the Atlantic, Pacific, and Indian Ocean basins, but shows pronounced latitudinal gradients. High concentrations of DIC are found throughout the Southern Ocean ( $\text{DIC} > 2100 \mu\text{mol kg}^{-1}$ ) where strong upwelling brings up subsurface water enriched in  $\text{CO}_2$  and nutrients. The inefficient utilization of nutrients in this high nutrient low chlorophyll region limits the biological drawdown of DIC allowing the massive DIC input to be spread horizontally by westerlies (Key et al., 2004; Menviel et al., 2018). Levels of DIC below  $1900 \mu\text{mol kg}^{-1}$  are reconstructed over the Equatorial Pacific, the Equatorial Eastern Atlantic, the Eastern Indian Ocean, and coastal areas on the Arctic Ocean. While low DIC levels associated with Equatorial upwelling reflect gas exchanges across the air-sea interface and enhanced biological production, the interaction between physical and biogeochemical processes at work in the Indian Ocean are less well understood (Takahashi et al., 2014). Low DIC levels found close to river mouths reflect outgassing of  $\text{CO}_2$  across the salinity gradient, as well as enhanced biological uptake fueled by river nutrient inputs. Representation uncertainty (Fig. 6-cd) associated with monthly alkalinity and DIC reconstructions is lower than  $20 \mu\text{mol kg}^{-1}$  throughout the open ocean. The open-ocean  $\sigma$ -to- $\mu$  ratio (Eq. 8) ranges between  $0.5 - 1.5\%$  which is relatively small (Fig. A9-cd). **The largest model uncertainty-DIC uncertainty is computed through  $\text{CO}_2\text{SYS}$  error propagation with reconstruction uncertainties of  $p\text{CO}_2$  and  $A_T$  set as inputs. The largest values ( $\sigma > 30 \mu\text{mol kg}^{-1}$ ) is computed appear**

nearshore and surrounding oceanic islands, a feature (Fig. 6d). A similar feature is found on the field of  $A_T$  (Fig. 6c) inherited from input uncertainty associated with the CMEMS salinity product (Fig. A7a).




**Figure 7.** Monthly time series of  $A_T$  and DIC at BATS, DYFAMED, ESTOC and HOT stations (Table 2 and Fig. A1b): model best estimate (curve), 1σ-uncertainty (envelope), and monthly average of surface (0-10 m) observations (point). Means of the best estimate and 1σ-uncertainty ( $\mu \pm \sigma$ ) calculated over the observing time span are shown in brackets if accessible. Statistics include number of months with observations ( $N$ ), Bias, RMSD, and  $r^2$ .  $\sigma_{A_T}^t$  [ $\sigma_{DIC}^t$ ] stands for temporal standard deviation from monthly averages of  $A_T$  [DIC] observations.

475 We qualify monthly, 0.25° reconstructions of  $A_T$  and DIC with [measurements from GLODAPv2.2022 data](#) (Lauvset et al., 2022a) for the 37-year period (Table 4 and Fig. 8). The global open-ocean reconstruction scores a RMSD of ~~22.09~~22.1 μmol kg<sup>-1</sup> and a  $r^2$  of 0.9 in  $A_T$ . Similar numbers are found for DIC (RMSD=~~22.67~~22.7 μmol kg<sup>-1</sup> and  $r^2 = 0.9$ ). The model scores the good fit in the open Indian Ocean with RMSD smaller than 15.5 μmol kg<sup>-1</sup> and  $r^2$  above 0.92 for both variables. The reconstruction deviates from GLODAP data in the western North Atlantic, subpolar North Pacific, tropics, and nearby major

480 rivers (Fig. 8-abcd).





**Figure 8.** Spatial distribution of reconstruction skills for  $A_T$ , DIC, and  $pH$  over 1985-2021. Mean model-data difference (Bias) and root-of-mean square-deviation (RMSD) between the reconstruction and GLODAPv2.2022 surface data (0-10 m) at a spatial resolution of  $0.25^\circ$ . The size of grid cells is scaled upon a better visualization.

$A_T$  and DIC are underestimated in the continental shelves of north Alaska and the northeastern Atlantic, the Mediterranean Sea, South China Sea, and nearby river plumes (Fig. 8-ac). The Arctic yields the poorest estimations among all the ocean basins with a global RMSD over  $100 \mu\text{mol kg}^{-1}$  (Table 4). The prediction probability of variability in  $A_T$  [DIC] is relatively large for the open ocean 79% [71%], but rather unsatisfying over the coastal ocean (46% [40%]). Extrapolating these carbonate variables  
 485 towards the shore remains challenging with much higher errors and uncertainty estimates obtained over the continental shelf compared to the open-ocean reconstruction (Table 4, Figs. 6-cd and . 8-abcd). The coastal-ocean errors are on the order of 10% of the global mean values of  $A_T$  and DIC (Table 6).

The reconstruction of  $A_T$  distributions relies on LIAR coefficients fit with GLODAPv2 data (Olsen et al., 2016) covering the years before 2015. These data are also part of the latest version GLODAPv2.2022 (Lauvset et al., 2022a). They do there-  
 490 fore not correspond to an independent dataset for the evaluation data of the CMEMS-LSCE reconstruction. To overcome this

**Table 4.** Skill scores computed between CMEMS-LSCE and GLODAPv2.v2022 in  $A_T$ , DIC, and  $pH$  over the period 1985-2021. Total numbers of data, RMSD (Eq. 10), and  $r^2$  (Eq. 11) are reported for both the open ocean (O) and coastal region (C). Basin identification is shown in Fig. A1.

Basins			Number of data		$A_T$ [ $\mu\text{mol kg}^{-1}$ ]		DIC [ $\mu\text{mol kg}^{-1}$ ]		pH [-]	
					RMSD	$r^2$	RMSD	$r^2$	RMSD	$r^2$
0.	Globe	(O)	10269	(5411)	<del>22.09</del> 22.1	0.90	<del>22.67</del> 22.7	0.90	0.022	0.70
		(C)	6309	(2080)	<del>82.01</del> 82.0	0.72	<del>72.39</del> 72.4	0.62	0.060	0.45
1.	Arctic	(O)	103	(26)	<del>107.09</del> 107.1	0.79	<del>113.28</del> 113.3	0.71	0.106	0.32
		(C)	1635	(300)	<del>148.71</del> 148.7	0.46	<del>126.77</del> 126.8	0.4	0.107	0.48
2.	Atlantic	(O)	2785	(932)	<del>30.10</del> 30.1	0.74	<del>28.66</del> 28.7	0.72	0.028	0.58
		(C)	2422	(941)	<del>44.50</del> 44.5	0.71	<del>39.09</del> 39.1	0.69	0.046	0.45
3.	Pacific	(O)	4539	(3222)	<del>13.61</del> 13.6	0.92	<del>15.95</del> 16.0	0.92	0.019	0.74
		(C)	1380	(639)	<del>28.43</del> 28.4	0.76	<del>44.36</del> 44.4	0.45	0.057	0.34
4.	Indian Ocean	(O)	1177	(551)	<del>15.05</del> 15.1	0.92	<del>13.79</del> 13.8	0.96	0.012	0.90
		(C)	328	(62)	<del>16.56</del> 16.6	0.92	<del>21.97</del> 22.0	0.90	0.013	0.82
5.	Southern Ocean	(O)	1665	(680)	<del>10.96</del> 11.0	0.64	<del>13.21</del> 13.2	0.92	0.019	0.68
		(C)	544	(138)	<del>22.53</del> 22.5	0.50	<del>24.48</del> 24.5	0.77	0.023	0.65

limitation accomplish a cross-validation, reconstructions of  $A_T$  and DIC are compared to observations for Eulerian-eight time series stations: AWIPEV, BATS, DYFAMED, ESTOC, and HOTHOT, ICELAND, IRMINGER, and KERFIX (see Table 2 and Fig. A1b for data sources and station locations). Table A4 presents the evaluation statistics for all the stations and Figure 7 illustrates the comparison between monthly time series of  $A_T$  and DIC extracted from the CMEMS-LSCE datasets and measurements at these long-term monitoring sites. The four stations stand out as the four sustained long-term observation-time series for carbonate system variables monitoring sites. More than 270 [80] months in the years 1988-2021 [1995-2009 and 1998-2017] include measurements of  $A_T$  and DIC at BATS and HOT [ESTOC and DYFAMED]. As shown in Table A4, the Arctic site (AWIPEV) provides 52-month data in 2015-2020 while the three other stations sparsely observed  $A_T$  and DIC at the surface layer resulting in fewer than 30 monthly mean data in 2014-2021 (IRMINGER and ICELAND) and 1992-2018 (KERFIX). The reconstructed time series fit monthly averages of in situ measurements well. Mean estimates of  $A_T$  [DIC] over the observing period are about ~~2305~~2283 [1983]  $\mu\text{mol kg}^{-1}$  at ~~HOT~~KERFIX [HOT] to 2420 [~~2129~~2219]  $\mu\text{mol kg}^{-1}$  at DYFAMED [AWIPEV]. At all the stations (DYFAMED and AWIPEV excepted) and for the two variables, model-observation misfit is small (Bias  $< 10 < 11$   $\mu\text{mol kg}^{-1}$ , RMSD  $< 13 < 14$   $\mu\text{mol kg}^{-1}$ ) relative to the aforementioned mean estimates (Table A4). The highest offset between the CMEMS-LSCE estimation and observations for all the stations is found at DYFAMED ( $A_T$ :  $-145.1$   $\mu\text{mol kg}^{-1}$ , DIC:  $-124.69-124.7$   $\mu\text{mol kg}^{-1}$ ). DYFAMED provides long-term time series of  $A_T$  and DIC measurements in the Northwestern Mediterranean Sea (Fig. A1b). Salinity and alkalinity have substantial values due to the net evaporation (Coppola et al., 2020). The average of  $A_T$  in the Mediterranean Sea exceeds that for the global ocean by 10%



(Palmiéri et al., 2015). These characteristics set the Mediterranean Sea aside from the ocean basins. **Although the bias between reanalysed SSS and observations (Fig. A10) is relatively small**, LIAR (Carter et al., 2018) was trained on GLODAPv2 (Olsen et al., 2016) including **a-only** few observations in this area. The distinct relationship between alkalinity and salinity prevailing in the Mediterranean Sea is likely not reproduced by LIAR leading to an underestimation of  $A_T$  and a systematic bias to DIC at DYFAMED (Fig. 7). ESTOC is located close to the North Atlantic east coast and under the influence of the Canary Current System (CCS, Fig. A1). Spatial gradients and temporal variability are higher in the CCS (Fig. 2c) compared to BATS and HOT which are both located in the center of subtropical gyres. **The lowest prediction skill of temporal variability is obtained for ESTOC** **Despite showing good estimates of  $A_T$  and DIC in RMSD at ESTOC, temporal variability of observations are reconstructed with the lowest  $r^2$  (Table A4)**. Particularly, seasonality to multi-year variations in DIC are predicted at  $r^2 = 0.47$  for ESTOC compared to  $r^2 > 0.7$  for **AWIPEV, ICELAND, IRMINGER**, BATS and HOT. Over all the stations, the model underestimates temporal changes of  $A_T$  (Fig. 7a; BATS:  $r^2 = 0.33$ , DYFAMED:  $r^2 = 0.12$ , ESTOC:  $r^2 = 0.03$ , HOT:  $r^2 = 0.32$ ) which can be attributed to the large discrepancy in variability between in situ measurements and the CMEMS time series of salinity (Fig. A10a; BATS:  $r^2 = 0.33$ , DYFAMED:  $r^2 = 0.19$ , ESTOC:  $r^2 = 0.03$ , HOT:  $r^2 = 0.35$ ). Model uncertainty ( $1\sigma$ -envelop) of monthly  $A_T$  and DIC estimates (Fig. 7a) is also inflated somewhat proportional to the CMEMS salinity product uncertainty (Fig. A10a).


### 5.3 Surface ocean pH and saturation state with respect to carbonate minerals

Surface ocean pH and saturation states with respect to aragonite ( $\Omega_{ar}$ ) and calcite ( $\Omega_{ca}$ ) are critical indicators used to measure ocean acidification. This section first presents an overall evaluation of these variables. We then introduce estimates involved in the monitoring of ocean acidification in 1985-2021 as an essential application of the CMEMS-LSCE surface ocean carbon product.

#### 5.3.1 General analysis and evaluation

The spatial distribution of surface ocean pH reported on total hydrogen ion ( $H^+$ ) scale is shown in Fig. 9 (the corresponding figure for  $H^+$ , Fig. A4, is included in the supplementary). Both temporal means of the best model estimate and  $1\sigma$ -uncertainty of pH share spatial patterns with  $pCO_2$  (Fig. 1). Variables pH and  $pCO_2$  correlate closely through equilibrium relationships of dissolved  $CO_2$  in seawater: an increase in  $pCO_2$  generally corresponds to a decrease in pH. The distribution of the climatological mean of pH displays a gradient with latitude between 8.03 and 8.11 pH units across most of the basins (Fig. 9a). Values of pH below 8 are associated with the upwelling of  $CO_2$ -rich waters (e.g., Eastern Equatorial Pacific, Western Arabian Sea). pH exceeds 8.15 in sub- and polar cold surface water and in the regions with high biological productivity (e.g. Labrador Sea, Nordic Seas, Southern Ocean between  $35^\circ S$ - $50^\circ S$ ).

The saturation state of surface ocean waters with respect to calcium carbonate minerals aragonite and calcite is defined as the ratio of the product of the concentrations of calcium ions ( $Ca^{2+}$ ) and carbonate ions ( $CO_3^{2-}$ ) to the solubility of the respective calcium carbonate mineral ( $CaCO_3$ ) in surface seawater (Eq. ??). Aragonite being the more soluble polymorph, its degree of saturation ( $\Omega_{ar}$ ) is smaller than that of calcite ( $\Omega_{ca}$ ) (Mucci, 1983). With the exception of this offset, the spatial distributions



**Figure 9.** CMEMS-LSCE  $pH$  and  $\Omega_{ar}$  over the global ocean at a spatial resolution of  $0.25^\circ$ . Temporal means of the model best estimate and  $1\sigma$ -uncertainty per grid cell over 1985-2021 are calculated by using Eq. (5).

of their climatological means share common spatial patterns over the global ocean (Figs. 9b and A5a). Surface seawater is generally supersaturated, i.e.,  $\Omega_{ar}$  and  $\Omega_{ca}$  greater than 1. The magnitude of surface ocean calcium carbonate saturation state varies with latitude. Values as large as 3.7-4.5 [5-7] for aragonite [calcite] are reconstructed in subtropical and tropical regions.  $\Omega_{ar}$  and  $\Omega_{ca}$  decrease toward the poles. In the Southern Ocean, surface seawater enriched in  $CO_2$  from vertical mixing has  $\Omega_{ar}$  [ $\Omega_{ca}$ ] values in the range of 1.5-2.1 [2-3.4]. Low saturation states are also computed in the Arctic and for waters of upwelling regimes (Fig. 9b). Locally  $\Omega_{ar}$  drops below 1.3, and even fall under the  $CaCO_3$  dissolution threshold of 1 (Gattuso and Hansson, 2011) in the Arctic water runoff and Baltic sea.

The uncertainty ( $1\sigma$ ) of  $pH$ ,  $\Omega_{ar}$ , and  $\Omega_{ca}$  propagated the speciation of the  $CO_2$  system takes into account the ensemble spread of  $pCO_2$  estimates and analysis errors of other variables (Sect. 3.3). Monthly  $pH$  uncertainty estimates fall in the 95% confidence interval of [0.008, 0.036] with a global mean value of 0.011. These estimates are in close agreement with the global uncertainty between 0.01-0.022  $pH$  units calculated by Jiang et al. (2019), Iida et al. (2021), and Gregor and Gruber (2021).  $pH$  uncertainty is typically larger than 0.03 in the Arctic and in coastal regions (Figs. 9c). In contrast, the reconstructions of  $\Omega_{ar}$  and  $\Omega_{ca}$  are subject to high uncertainty ( $\sigma > 0.175$ ) between  $30^\circ S$ - $30^\circ N$  (Fig. 9d and A5b). Regarding the  $\sigma$ -to- $\mu$  ratio, mean uncertainty estimates per cell for the saturation states in the (sub-) tropical band are relatively small compared to the mean of the best monthly estimates (Figs. A9-ef). The Arctic and the coastal oceans remain the regions with largest reconstruction uncertainties for  $\Omega_{ar}$  and  $\Omega_{ca}$ , as well as for  $pCO_2$  and  $pH$  (Figs. A9-ab). Excluding these regions,  $R(\sigma, \mu)$  (Eq. 8) is less than 0.3% for  $pH$  and 8% for  $\Omega_{ar}$  and  $\Omega_{ca}$ .


The monthly CMEMS-LSCE reconstruction at  $0.25^\circ$  resolution is assessed against  $pH$  measurements from GLODAPv2.2022 bottle data (Table 2). For the period 1985-2021, the global RMSD amounts to 0.022 [0.060]  $pH$  units and  $r^2$  scores at 0.70 [0.45] over the open [coastal] ocean (Table 4). Model bias lies within  $[-0.01, 0.01]$   $pH$  units and RMSD is below 0.02  $pH$  units over the open ocean, except for high latitudes over  $60^\circ$  (Figs. 8-ef). At local scale, the ~~eight coastal~~ time series from Sutton et al. (2019) are used for further evaluation (Tables A2 and A3). There exists much ~~less-fewer~~ evaluation data for  $pH$  than for  $pCO_2$ , e.g., only 2 months of monitoring  $pH$  at COASTALLA and FIRSTLANDING (Table A3). ~~Monthly time series of CMEMS-LSCE or devoid of  $pH$  are coherent with these measurements at equatorial observing systems. CMEMS-LSCE reconstructs  $pH$  measurements (Tables A2 and A3). Measurement uncertainty of  $pH$  at these coastal sites is reported to be around 0.02 over the open ocean with rather high scores, e.g., at BOBOA (RMSD = 0.011 and  $r^2 = 0.71$ ) and KEO (RMSD = 0.014 and  $r^2 = 0.86$ ). Referring to the eight coastal sites evaluated for  $pH$  units- $CO_2$  in Section 5.1, RMSD can be as small as 0.035 and 0.04  $pH$  units at CCE2 and GRAYSREEF while it is over 0.05  $pH$  units at the other stations (e.g., COASTALLA: 0.068, CAPEARAGO: 0.069). Similar to the  $pCO_2$  time series (Fig. 5, Sect. 5.1),  $pH$  has been monitored with low sampling frequency (roughly a few days in the tracking month) and the temporal sampling deviation of instantaneous observations from monthly averages ( $\sigma_{pH}^t$ ) is significant. This temporal sampling uncertainty of  $pH$  contributes to the mismatch between model estimates and observations. For example,  $\sigma_{pH}^t$  amounts to 0.048  $pH$  units at CCE2 and 0.020  $pH$  units at GRAYSREEF, and reaches highest values of 0.078  $pH$  units at COASTALLA and 0.086  $pH$  units at CAPEARAGO. Although model-observation misfit and model uncertainty remain high over the coastal sector (see also Figs. 8-ef and 9c), their estimates do not surpass 1% of the global mean  $pH$  (8.082). The reconstructed  $pH$  time series reproduce measurement variability with relatively high correlation,  $r^2$  in ~~[0.21, 0.69]~~[0.21, 0.94], that reinforces the reliability of CMEMS-LSCE  $pH$  data.~~

### 5.3.2 Ocean acidification: key features from global to local scales


The monthly,  $0.25^\circ$  CMEMS-LSCE datasets of  $pH$ ,  $\Omega_{ar}$ , and  $\Omega_{ca}$  are at the basis of of two CMEMS ocean indicators monitoring surface ocean acidification from 1985 to 2021: (1) annual global means and (2) global trend maps.

In Fig. 10, we present annual global means of surface ocean  $pH$  and saturation states for aragonite ( $\Omega_{ar}$ ). An illustration of calcite ( $\Omega_{ca}$ ) is provided in the Appendix (Fig. A13a). For each variable, the calculation of annual global area-weighted means of best estimates (line) and  $1\sigma$ -uncertainties (envelope) follows Eq. (7). The trends reported in the legend result from linear least-squares regression on annual global means of 100-ensembles of the carbonate system variables. These ensembles are generated with Gaussian distribution having the mean and variance as best model estimate  $\mu$  and squared uncertainty ( $\sigma^2$ ) at monthly time steps and  $0.25^\circ$ -grid cells, respectively.  $pH$  decreases from  $8.110 \pm 0.017$  in 1985 to  $8.049 \pm 0.014$  in 2021 with a descend rate of  $-0.017 \pm 0.004$  decade $^{-1}$ . Similar trends are found for the surface ocean saturation states with respect to calcium carbonate minerals. The global mean estimates of  $\Omega_{ar}$  [ $\Omega_{ca}$ ] amount to  $3.141 \pm 0.198$  [ $4.807 \pm 0.302$ ] and  $2.862 \pm 0.174$  [ $4.372 \pm 0.266$ ] for the open and coastal oceans. The saturation state declines at a rate of  $-0.080 \pm 0.029$  decade $^{-1}$  with respect to aragonite while the reduction is steeper for calcite ( $-0.114 \pm 0.045$  decade $^{-1}$ ).

Global trend maps of surface ocean  $pH$ ,  $\Omega_{ar}$ , and  $\Omega_{ca}$  over the entire period are illustrated in Figs. 11 and A13b. Linear least-squares regression is used to estimate secular trends at every  $0.25^\circ$ -grid cell. The linear fits of each variable against time rely on



**Figure 10.** Yearly global area-weighted mean of surface seawater  $pH$  reported on total scale (a) and surface ocean saturation states with respect to aragonite (b). Global means of the best estimate ( $\mu$ , plain line) and of uncertainty ( $\sigma$ , envelop) are computed with Eq. (7a). Trend and uncertainty in the legend are computed with linear regressions on the 100-member ensemble of yearly global means for each variable.



**Figure 11.** Global trend maps of surface seawater  $pH$  reported on total scale (a) and surface ocean saturation states with respect to aragonite (b). Linear trend of CMEMS-LSCE  $pH$  and  $\Omega_{ar}$  is estimated per  $0.25^\circ$ -grid cell over 1985-2021. Cross-hatching covers the regions where with uncertainty over 10% [20%] of  $pH$  [ $\Omega_{ar}$ ] trend estimates.

the 100-member ensemble generated with the best estimates and propagated uncertainties of  $pH$ ,  $\Omega_{ar}$ , and  $\Omega_{ca}$  (see Figs. A14 for examples). Regression slope and residual standard deviation estimates are defined as linear trend and uncertainty of  $pH$ ,  $\Omega_{ar}$ , and  $\Omega_{ca}$ . Hatched area represents  $pH$  [ $\Omega_{ar}$  and  $\Omega_{ca}$ ] trend estimates ( $\mu$ ) with highest uncertainties ( $\sigma$ ), i.e.,  $\sigma$ -to- $\mu$  ratio (Eq. 8) above 10% [20%]. These regions include a portion of the Arctic, Antarctic, equatorial Pacific, and coastal ocean (Figs. 11, A11, and A12). 95% of  $pH$  trend estimates over the global ocean is in the range of  $[-0.022, -0.012]$  decade $^{-1}$  (Fig. 11a). In the broad open ocean of the tropics and subtropics,  $pH$  has been declining around  $-0.018$  decade $^{-1}$  to  $-0.012$  decade $^{-1}$ . Faster decrease rates are found in the Indian Ocean and Southern Ocean with values between  $-0.022$  and  $-0.018$  decade $^{-1}$ . Fastest reductions are computed for the eastern equatorial Pacific and the Arctic with rates exceeding  $-0.025$  decade $^{-1}$ . A similar magnitude of  $pH$  trends over these regions is also found in (Lauvset et al., 2015; Leseurre et al., 2022; Ma et al.,

2023). The spatial distribution of saturation states with respect to calcium carbonate minerals generally shows the opposite latitudinal pattern (Figs. 11b and A13b). The magnitude of  $\Omega_{ar}$  [ $\Omega_{ca}$ ] trends over the 30°S-30°N band can be as large as  $-0.086 \text{ decade}^{-1}$  [ $-0.134 \text{ decade}^{-1}$ ] to the greatest extent of  $-0.186 \text{ decade}^{-1}$  [ $-0.275 \text{ decade}^{-1}$ ] (e.g., eastern equatorial Pacific). Trends of  $\Omega_{ar}$  and  $\Omega_{ca}$  computed in polar and subpolar northern hemisphere regions are not significant.

**Table 5.** Secular trend estimates of  $pH$  and  $\Omega_{ar}$  at seven time-series stations (Bates et al., 2014). Trend and uncertainty estimates are reported as  $\mu \pm \sigma$ . Monthly time series in the CMEMS-LSCE datasets are extracted at the grid box nearest to each station location (Fig. A1b). For the first three stations, this study calculates linear trends starting in the year 1985. Brackets show values computed over the full period 1985-2021.

Stations	Coordinates	Time span	$pH$ trend [ $\text{decade}^{-1}$ ]		$\Omega_{ar}$ trend [ $\text{decade}^{-1}$ ]	
			Bates et al. (2014)	This study	Bates et al. (2014)	This study
1. Iceland Sea	68.00°N 12.66°W	1983-2012	$-0.014 \pm 0.005$	$-0.010 \pm 0.001$ ( $-0.014 \pm 0.001$ )	$-0.018 \pm 0.027$	$-0.013 \pm 0.011$ ( $-0.025 \pm 0.008$ )
2. Irminger Sea	64.30°N 28.00°W	1983-2012	$-0.026 \pm 0.006$	$-0.014 \pm 0.001$ ( $-0.016 \pm 0.001$ )	$-0.080 \pm 0.040$	$-0.006 \pm 0.011$ ( $-0.039 \pm 0.009$ )
3. BATS	32.00°N 64.00°W	1983-2012	$-0.017 \pm 0.001$	$-0.014 \pm 0.001$ ( $-0.016 \pm 0.001$ )	$-0.095 \pm 0.007$	$-0.079 \pm 0.016$ ( $-0.074 \pm 0.010$ )
4. ESTOC	29.04°N 15.50°W	1995-2012	$-0.018 \pm 0.002$	$-0.018 \pm 0.002$ ( $-0.019 \pm 0.001$ )	$-0.115 \pm 0.023$	$-0.103 \pm 0.031$ ( $-0.089 \pm 0.011$ )
5. HOT*	22.75°N 158.00°W	1988-2012	$-0.016 \pm 0.001$	$-0.016 \pm 0.001$ ( $-0.019 \pm 0.001$ )	$-0.084 \pm 0.011$	$-0.100 \pm 0.020$ ( $-0.102 \pm 0.011$ )
6. CARIACO*	10.50°N 64.66°W	1995-2012	$-0.025 \pm 0.004$	$-0.017 \pm 0.003$ ( $-0.018 \pm 0.001$ )	$-0.066 \pm 0.028$	$-0.059 \pm 0.053$ ( $-0.099 \pm 0.018$ )
7. Munida	45.70°S 171.50°E	1998-2012	$-0.013 \pm 0.003$	$-0.017 \pm 0.002$ ( $-0.017 \pm 0.001$ )	$-0.085 \pm 0.026$	$-0.088 \pm 0.032$ ( $-0.070 \pm 0.009$ )

\*Stations with direct observations of  $pH$ .

605 Trend estimates derived from reconstructions of  $pH$  and  $\Omega_{ar}$  are evaluated at seven time series stations (Bates et al., 2014)  
in Table 5. Time series locations are shown in Fig. A1b. With the exception of CARIACO and HOT ~~excepted~~ for which  $pH$   
measurements are available, long-term trend estimates by Bates et al. (2014) rely on time series of  $pH$  and  $\Omega_{ar}$  calculated  
via speciation from measurements of  $A_T$  and DIC. A-100-member ~~ensemble-ensembles~~ of monthly time series of  $pH$  and  
 $\Omega_{ar}$  are extracted from the 0.25°-grid box nearest to each monitoring station. Linear least-squares regression is then used  
610 to infer estimates of their secular trends and associated uncertainties (see Fig. A14 for illustration). Trend estimates derived  
from CMEMS-LSCE reconstructions at HOT, BATS, ESTOC, and Munida are in line with previous studies for both  $pH$   
and  $\Omega_{ar}$  (Dore et al., 2009; González-Dávila and Santana-Casiano, 2009; Bates et al., 2014). The magnitude of the trend  
estimate at Irminger Sea for 1985-2012 ( $pH$ :  $-0.014 \pm 0.001 \text{ decade}^{-1}$ ,  $\Omega_{ar}$ :  $-0.006 \pm 0.011 \text{ decade}^{-1}$ ) is smaller than that  
determined by Bates et al. (2014). However, the CMEMS-LSCE  $pH$  trend is consistent with the estimate by Pérez et al. (2021)



615  $(-0.017 \pm 0.002 \text{ decade}^{-1})$ . Moreover,  $1\sigma$ -uncertainty reported for both  $pH$  and  $\Omega_{ar}$  trend estimates by Bates et al. (2014) is large at this station ( $pH$ :  $-0.025 \pm 0.006 \text{ decade}^{-1}$ ,  $\Omega_{ar}$ :  $-0.080 \pm 0.040 \text{ decade}^{-1}$ ) highlighting the associated uncertainty. Long-term trends of  $pH$  and  $\Omega_{ar}$  are also under-estimated at the Iceland Sea monitoring site, but the bias is not as large as at Irminger Sea (Table 5). Low data sampling frequency at these two stations (Table 1, Bates et al., 2014) could be on account of trend estimate deviation. At CARIACO, the CMEMS-LSCE time series yields a decrease in  $\Omega_{ar}$  of  $-0.059 \pm 0.053 \text{ decade}^{-1}$ ,  
620 relatively close to Bates et al. (2014) ( $-0.066 \pm 0.028 \text{ decade}^{-1}$ ). The decrease in  $pH$  derived from CMEMS-LSCE is, however, larger than in Bates et al. (2014).

## 6 Summary Conclusions and Discussion

This study presents the CMEMS-LSCE product, a dataset of six carbonate system variables (Table ??6) covering the global surface ocean at a spatial resolution of  $0.25^\circ$  for every month in the period 1985-2021 (444 months). Datasets of individual  
625 carbonate system variables are built on the combination of the three methods. First, we adapt an ensemble of 100 feed-forward neural network models (CMEMS-LSCE-FFNN, Chau et al., 2022b) to estimate surface ocean partial pressure of  $CO_2$  ( $pCO_2$ ) at the pre-defined data resolution. Second, the high-resolution total alkalinity ( $A_T$ ) reconstruction is obtained by using locally interpolated alkalinity regression (LIAR, Carter et al., 2016, 2018). Finally, surface ocean  $pH$ , total dissolved inorganic carbon (DIC), and saturation states with respect to aragonite ( $\Omega_{ar}$ ) and calcite ( $\Omega_{ca}$ ) are calculated with the carbonate system  
630 speciation software (CO2SYS, Lewis and Wallace, 1998; Van Heuven et al., 2011; Orr et al., 2018), given the global monthly reconstructions of  $pCO_2$  and  $A_T$  and other environmental input data (Sect. 3). Results are 2D-fields of the best estimate and associated uncertainty ( $1\sigma$ ) of carbonate system variables available at each grid box of  $1\text{month} \times 0.25^\circ \times 0.25^\circ$ .  $1\sigma$ -uncertainty is referred to as the ensemble standard deviation of 100 FFNN outputs for  $pCO_2$  while it is propagated through the processing chain of LIAR and CO2SYS taking into account different uncertainty sources of input parameters for other variables.

635 Multiple observation-based datasets, which are not used for the CMEMS-LSCE reconstructions at monthly and  $0.25^\circ$  resolutions, serve as benchmarks in the assessments of product quality from global to local scales (e.g., Tables 3, 4, and A3; Figs. 2-5 and 7-8). A summary of the primary statistics for all the six carbonate variables is presented Table 6. Over the full period 1985-2021, CMEMS-LSCE yields global RMSDs of ~~14.29~~14.3  $\mu\text{atm}$  and ~~27.55~~27.6  $\mu\text{atm}$  in comparison with SOCATv2022  $pCO_2$  for the open and coastal oceans, respectively. Temporal variability of observation-based data is well reproduced with  $r^2$   
640 of 0.83 for the open ocean and 0.74 for the coastal domain. In comparison to CMEMS-LSCE at monthly and  $1^\circ$  resolutions (Chau et al., 2022b), the reconstructions over coastal areas are improved at higher resolution(Figs. 2-4). Furthermore, the. The monthly,  $0.25^\circ$  reconstruction outperforms its  $1^\circ$  counterpart in reproducing horizontal and temporal gradients of  $pCO_2$  over a variety of oceanic regions as well as at nearshore time series stations (Figs. 2-5). Evaluations with GLODAPv2022 bottle data and time series stations results in good reconstruction skills for  $A_T$ , DIC, and  $pH$  at monthly and  $0.25^\circ$  resolutions (Tables 4  
645 and A3, Figs. 7 and 8). At the global scale, the open-ocean reconstruction scores a RMSD smaller than  $23 \mu\text{mol kg}^{-1}$  and a  $r^2$  of 0.9 in  $A_T$  and DIC. The model-observation deviation is higher in the coastal zone. However, it does not exceed 5% of the global mean values and  $r^2$  is above 0.6 for both coastal  $A_T$  and DIC. Regarding  $pH$ , the CMEMS-LSCE reconstruction

provides estimates with  $\text{RMSD} = 0.022$  [0.060] and  $r^2 = 0.7$  [0.45] over the open [coastal] ocean. From the statistics in Tables 3 and 4, the Indian Ocean and the Southern Ocean have poor data density (Fig. A2) but generally show the best global reconstruction among the ocean basins. Thus, model evaluation with different numbers of observation data might not reflect a fair comparison of skill scores (e.g., RMSD and  $r^2$ ) between regions. Data density is much higher in the Arctic, Atlantic, and Pacific than in the Indian and Southern Oceans. The increased data density reveals stronger spatio-temporal variability, for instance, related to coastal dynamics or upwelling than resolved in the two latter basins. RMSD and  $r^2$  computed on the lower data variability result in better model scores.

The spatial distribution of long-term mean  $1\sigma$ -uncertainty estimates (Figs. 1b, 6cd, and 9cd) indicates higher confidence levels for open-ocean estimates than over the coastal sector. The evaluation of temporal mean  $1\sigma$ -uncertainty estimates relative to climatological mean values  $\mu$  (Figs. 1a, 6ab, and 9ab) results in  $\sigma$ -to- $\mu$  ratio (Eq. 8) below 5% and 8% for  $p\text{CO}_2$  and  $\Omega_{ar}$ , 2% for  $A_T$  and DIC, and 0.4% for  $p\text{H}$  over the open ocean (Fig. A9). The  $\sigma$ -to- $\mu$  ratio reaches values as high as 10% to 20% for  $p\text{CO}_2$  and  $\Omega_{ar}$  in the coastal domain. The global mean of open-ocean  $1\sigma$ -uncertainty estimates (Eq. 7a) for CMEMS-LSCE  $p\text{CO}_2$  (8.488.5  $\mu\text{atm}$ ),  $A_T$  (16.6616.7  $\mu\text{mol kg}^{-1}$ ), DIC (15.7515.8  $\mu\text{mol kg}^{-1}$ ),  $p\text{H}$  (0.011), and  $\Omega_{ar}$  (0.180) are in line with those reported by previous studies despite being derived from different statistics. For instance, Iida et al. (2021) calculated  $1\sigma$ -uncertainty based on the median absolute deviation of regression model fits from open-ocean observations. Their approach yielded global  $\sigma$ -averages of 17.8  $\mu\text{atm}$ , 11.5  $\mu\text{mol kg}^{-1}$ , 0.018, and 0.110 for  $p\text{CO}_2$ , normalized DIC,  $p\text{H}$ , and  $\Omega_{ar}$ , respectively. In Gregor and Gruber (2021), the authors propagated the sum squared errors (global RMSD and measurement uncertainties) of  $p\text{CO}_2$  (15  $\mu\text{atm}$ ) and  $A_T$  (22  $\mu\text{mol kg}^{-1}$ ) obtaining global uncertainty estimates of 19  $\mu\text{mol kg}^{-1}$  in DIC and 0.022 in  $p\text{H}$ . Mean uncertainty estimates over the coastal region are on the order of twofold that computed for the open ocean for these four variables (Table 6), corroborating results by Gregor and Gruber (2021) (Fig. 7).

**Table 6.** Summary in global evaluation statistics for CMEMS-LSCE surface ocean carbonate system datasets at monthly,  $0.25^\circ$  resolutions over the period 1985-2021.  $\mu$  and  $\sigma$  stand for the global area-weighted means of monthly best estimates and  $1\sigma$ -uncertainties for each variable (Eq. 7). RMSD (Eq. 10) and  $r^2$  (Eq. 11) are computed with SOCATv2022 for  $p\text{CO}_2$  and GLODAPv2.2022 for  $p\text{H}$ ,  $A_T$ , and DIC. The division between the coastal (C) and open (O) oceans is at 400 km on a distance from the shore line (Fig. A1a).

Variables	Standard names	Units	Sector	$\mu$	$\sigma$	RMSD	$r^2$
1. $p\text{CO}_2$	Partial pressure of $\text{CO}_2$ in surface seawater	$\mu\text{atm}$	(O)	364.48364.5	8.48-8.5	14.29-14.3	0.83
			(C)	359.35359.4	17.1017.1	27.55-27.6	0.74
2. $A_T$	Total alkalinity in surface seawater	$\mu\text{mol kg}^{-1}$	(O)	2305.782305.8	16.66-16.7	22.09-22.1	0.90
			(C)	2263.022263.0	38.36-38.4	82.01-82.0	0.72
3. DIC	Surface ocean dissolved inorganic carbon	$\mu\text{mol kg}^{-1}$	(O)	2031.122031.1	15.75-15.8	22.67-22.7	0.90
			(C)	2008.652008.7	33.40-33.4	72.39-72.4	0.62
4. $p\text{H}$	Surface seawater $p\text{H}$ reported on total scale	-	(O)	8.082	0.011	0.022	0.70
			(C)	8.082	0.021	0.060	0.45
5. $\Omega_{ar}$	Saturation state for surface seawater with respect to aragonite	-	(O)	3.059	0.180	-	-
			(C)	2.864	0.206	-	-
6. $\Omega_{ca}$	Saturation state for surface seawater with respect to calcite	-	(O)	4.674	0.275	-	-
			(C)	4.384	0.314	-	-



Our high-resolution carbon data product opens the door to various analyses of the marine carbonate system from global to local scale. This study exemplifies an application of the data for monitoring ocean acidification over recent years. The monitoring indicators derived from the monthly,  $0.25^\circ$  surface ocean CMEMS-LSCE product consist of (1) yearly global means of surface ocean  $pH$  and saturation states with respect aragonite  $\Omega_{ar}$  and calcite  $\Omega_{ca}$  and (2) global maps of multi-annual trends of surface ocean  $pH$ ,  $\Omega_{ar}$ , and  $\Omega_{ca}$  (Figs. 10, 11, and A13). In 1985, the global mean surface ocean  $pH$  was  $8.110 \pm 0.017$ . It was  $8.049 \pm 0.014$  in 2021 (Fig. 10a). Over the same 37-year time period,  $\Omega_{ar}$  decreased from  $3.141 \pm 0.198$  to  $2.862 \pm 0.174$  (Fig. 10b). The rate of decline of surface ocean  $pH$  and  $\Omega_{ar}$  was respectively  $-0.017 \pm 0.004 \text{ decade}^{-1}$  and  $-0.080 \pm 0.029 \text{ decade}^{-1}$  since 1985 (see also results for  $\Omega_{ca}$  in Sect. 5.3.2). Estimates of  $pH$  trend lie between  $[-0.022, -0.012] \text{ decade}^{-1}$  across most of the open ocean (Fig. 11a). In general, surface ocean  $pH$  decreased more rapidly in the Indian Ocean and Southern Ocean than the tropics and subtropics. These findings are in close agreement with the suggestions by Lauvset et al. (2015) and Ma et al. (2023) but future studies would need to include analyses of underlying drivers to provide insight into regional differences in  $pH$  changes. By contrast, the greatest reduction in surface ocean saturation states (Fig. 11b) was computed for the two latter regions. The global trend maps of  $pH$  and  $\Omega_{ar}$  highlight the Eastern Equatorial Pacific as one of the vulnerable regions with respect to ocean acidification. In this area, the decline rate of  $pH$  exceeds  $-0.025 \text{ decade}^{-1}$  and  $-0.186 \text{ decade}^{-1}$  for  $\Omega_{ar}$ . The comparison of multi-annual trends of  $pH$  and  $\Omega_{ar}$  at time series stations (Table 5 and Fig. A14) highlighted the consistency between CMEMS-LSCE estimates and previous studies (Dore et al., 2009; González-Dávila and Santana-Casiano, 2009; Bates et al., 2014; Pérez et al., 2021). For most of these sites, the trends evaluated for 1985-2021 are greater than those relative to the sub-period before the year 2012. The faster rate of ocean acidification over the full period compared to the pre-2012 probably reflects a steeper acceleration in ocean uptake of anthropogenic  $CO_2$  in the last decade. **The global maps of CMEMS-LSCE  $pH$ ,  $\Omega$ , and their trend estimates would be potential indicators for ocean acidification along with the SDG 14.3.1 - "Average marine acidity ( $pH$ ) measured at agreed suite of representative sampling stations" (<https://sdgs.un.org/goals/goal14>: last access 31/07/2023).**

The production chain of CMEMS-LSCE carbonate system variables will be maintained and further improvements with the aim to reduce model-observation misfit and improve the quantification of model uncertainty are on the way forward. Being at the core of the chain, model upgrades of CMEMS-LSCE-FFNN will be tackled first. At the time, SOCAT does not provide open-ocean data of  $CO_2$  fugacity gridded at monthly,  $0.25^\circ$  resolutions. Our ensemble-based approach draws thus on two SOCATv2022 data sources: a "downscaled" version of the  $1^\circ$ -open-ocean data and the  $0.25^\circ$ -coastal-ocean data (see Sect. 2.1). Open-ocean SOCAT datasets gridded at finer regular resolutions (if accessible) will be updated to gain more accuracy in our model fitting. Selections of data products for predictors needed for model input are equally important. For instance, the CMEMS SSS product used here results in a globally good reconstruction of total alkalinity (Table 4). However, the temporal variability in CMEMS SSS data does not match that in observations (Fig. A10) and this feature is retained in time series of total alkalinity (Fig. 7). Despite best efforts in determining overall product uncertainty in estimates of carbonate system variables, part of input uncertainty is still not taken into account or only partially quantified due to lack of time-space varying uncertainty fields associated with predictor variables (e.g. SSH, Chl-a, MLD, nutrient concentrations). Moreover, temporal sampling bias in  $pCO_2$  and  $pH$  is likely to contribute to deviations between observations and model output (Fig. 5 and Table A3). The

total ~~measurement-analytical~~ error uncertainty should be considered with great care during reconstruction and model output evaluation.


705 The CMEMS-LSCE approach leads as the first series of long-term reconstructions of  $p\text{CO}_2$ ,  $p\text{H}$ ,  $A_T$ ,  $\text{DIC}$ ,  $\Omega_{ca}$  and  $\Omega_{ar}$  extending seamlessly from the global open ocean to coastal regions at monthly,  $0.25^\circ$  resolutions. Future use cases recommended for this high-resolution product include (1) estimation of monthly to interannual variations, long-term trends of carbonate system variables, as well as of air-sea  $\text{CO}_2$  exchanges at the surface layer from local scale to large ocean basins, (2) analyses in interactions between these variables and effects of other physical and biogeochemical factors on ocean acidification and  
710 changes in the marine carbonate system, (3) assessments of horizontal and temporal gradients of carbonate system variables in the coastal-open ocean continuum, (4) evaluation or combination with other model- or observation-based products (e.g., Biogeochemistry Argo, Southern Ocean Carbon and Climate Observations and Modeling), and (5) improvements in coastal reconstructions based on observation system simulation experiments (e.g., with finer spatio-temporal model resolutions). The CMEMS-FFNN surface ocean carbon product at monthly,  $0.25^\circ$  resolutions will be accessible through the CMEMS data portal  
715 (see Sect. 7).

## 7 Data availability

The CMEMS-LSCE datasets (netCDF format) of six carbonate system variables have been delivered to the European Copernicus Marine Environment Monitoring Service (CMEMS, Product ID: MULTIOBS\_GLO\_BIO\_CARBON\_SURFACE\_REP\_015\_008, DOI: 10.48670/moi-00047). Since November 2022, the product with monthly and  $1^\circ$  resolutions is available at the CMEMS  
720 portal (Chau et al., 2022a, b). The CMEMS-LSCE data product at monthly and  $0.25^\circ$  resolutions proposed in this study will replace its coarser resolution version in due course. For the time being, the high-resolution data product described in this manuscript can be accessed via repository under data DOI: 10.14768/a2f0891b-763a-49e9-af1b-78ed78b16982 (Chau et al., 2023).

## Appendix A: Definitions of ocean carbonate system variables

725 ~~Chemical reactions of dissolved  $\text{CO}_2$  in seawater follow a series of the following equilibria,~~



where (g) and (aq) stand for a gas or the species in an aqueous solution.  $\text{CO}_2(\text{aq})$  refers to the combination of aqueous  $\text{CO}_2$   
730 and its weak acid  $\text{H}_2\text{CO}_3$ .  $\text{HCO}_3^-$  and  $\text{CO}_3^{2-}$  are bicarbonate and carbonate ions.

~~Definitions of the essential variables involved in the carbonate system equilibria are on the list below (see in Dickson et al., 2007; Dickso~~

~~-~~

i) Surface ocean  $p\text{CO}_2$  is partial pressure of  $\text{CO}_2$  in air which is in equilibrium with that in water sample. It is not the same as surface ocean fugacity of  $\text{CO}_2$  ( $f\text{CO}_2$ ).  $p\text{CO}_2$  can be converted from  $f\text{CO}_2$  via

$$735 \quad p\text{CO}_2 = f\text{CO}_2 \exp\left(-P \frac{B + 2\delta}{RT^*}\right).$$

where  $P$  is total atmospheric pressure at surface water,  $T^*$  is absolute temperature,  $R$  is the gas constant, and  $B$  and  $\delta$  are cross-virial coefficients (Körtzinger, 1999).

ii) Seawater  $p\text{H}$  is a negative logarithmic scale of total concentration of hydrogen ions ( $\text{H}^+$ ) in aqueous solution. Total  $\text{H}^+$  is the sum of concentrations of free  $\text{H}^+$  and  $\text{HSO}_4^-$  ions. The  $p\text{H}$  scale typically ranges from 0 to 14.  $p\text{H} = 7$  is the threshold specifying whether a water sample is in acidic (i.e.,  $p\text{H} < 7$ ) or basic (i.e.,  $p\text{H} > 7$ ) conditions.

740

iii) Total alkalinity ( $A_T$ ) measures the capacity of seawater against acidification. By definition,  $A_T$  is total concentration of dissolved alkaline substances corresponding to the ability in  $\text{H}^+$  attracting over  $\text{H}^+$  releasing. The major contributions to alkalinity includes bicarbonate ( $\text{HCO}_3^-$ ), carbonate ( $\text{CO}_3^{2-}$ ), and hydroxide ( $\text{OH}^-$ ) ions. Total alkalinity can be approximated with

$$745 \quad A_T = [\text{HCO}_3^-] + 2[\text{CO}_3^{2-}] + [\text{OH}^-] - [\text{H}^+].$$

iv) Total dissolved inorganic carbon (DIC) is the sum in concentrations of the three primary aqueous species in seawater,

$$\text{DIC} = [\text{HCO}_3^-] + [\text{CO}_3^{2-}] + [\text{CO}_2(\text{aq})].$$

v) Calcium carbonate saturation state ( $\Omega$ ) is defined as follows,

$$\Omega = \frac{[\text{Ca}^{2+}][\text{CO}_3^{2-}]}{K_{sp}},$$

750 where  $[\text{Ca}^{2+}]$  is the concentration of dissolved calcium ions and  $K_{sp}$  is the solubility of calcium carbonate in seawater.  $\text{CaCO}_3$  has two principal minerals: aragonite and calcite. Aragonite, which is more soluble than calcite ( $\Omega_{ar} < \Omega_{ca}$ ), is produced by many marine shells and skeletons including corals, pteropods, clams, and mussels. A  $\Omega_{ar}$  value greater than 1, i.e., preferable conditions in shell formation, indicates supersaturated seawater with respect to aragonite, and vice versa.

755 vi) Revelle factor (RF) measures the buffer capacity for the carbonate system in seawater that decreases as  $p\text{H}$  increases. Revelle factor is expressed by the ratio between instantaneous changes of dissolved  $\text{CO}_2$   $\left(\frac{[\Delta\text{CO}_2(\text{aq})]}{[\text{CO}_2(\text{aq})]}\right)$  and of DIC  $\left(\frac{[\Delta\text{DIC}]}{[\text{DIC}]}\right)$  in seawater,


$$\text{RF} = \frac{[\Delta\text{CO}_2(\text{aq})]}{[\text{CO}_2(\text{aq})]} \left(\frac{[\Delta\text{DIC}]}{[\text{DIC}]}\right)^{-1}.$$

*Author contributions.* TTTC, FC, and MG developed the CMEMS-LSCE-FFNN model at a quarter-degree resolution. TTTC has prepared  
760 script codes and executed the experiments with support from FC in setting and running the model at the HPC resources of TGCC. TTTC,  
MG, NM, and FC shaped the first manuscript version. All the authors contribute to the manuscript revision.

*Competing interests.* The author and co-authors have declared that they have no competing interests.

*Acknowledgements.* This research has been supported by the MOB TAC project of the European Copernicus Marine Environment Moni-  
765 HPC resources of TGCC under the allocation A0110102201 made by GENCI. The Surface Ocean CO<sub>2</sub> Atlas (SOCAT, [www.socat.info](http://www.socat.info), last  
access: 20 March 2023) is an international effort endorsed by the International Ocean Carbon Coordination Project (IOCCP), the Surface  
Ocean Lower Atmosphere Study (SOLAS) and the Integrated Marine Biogeochemistry and Ecosystem Research program (IMBER), to de-  
770 [deliver a uniformly quality-controlled surface ocean CO<sub>2</sub> database. The Global Ocean Data Analysis Project \(GLODAP, \[www.glodap.info\]\(http://www.glodap.info\), last  
access: 21 August 2023\) provides access to ocean surface-to-bottom quality controlled data of carbonate system variables collected through  
international cruises. We thank Anna Conchon for her help in testing and wrapping LIAR and CO2SYS Matlab toolboxes. We are grateful  
for constructive comments from two anonymous reviews to refine the manuscript.](#)


## Appendix



**Figure A1.** a) Ocean basins (<https://github.com/RECCAP2-ocean/RECCAP2-shared-resources/tree/master/data/regions>, last access: 11/7/2022): coastal mask (grey, approximately 400 km from the [shore line](#)shoreline), feature regions [analyzed](#)–[analysed](#) in this study (cyan box, Table A1); b) Location of time series stations recording in situ observations used in data evaluation (Table 2): blue stars ([Bates et al., 2014](#)) for ocean acidification ([Bates et al., 2014](#)), black star ([Coppola et al., 2021](#)) stars for  $A_T$  and DIC ([Metzl and Lo Monaco, 1998](#); [Coppola et al., 2021](#); [Gattuso et al., 2023](#)), and other coloured scattered objects ([Sutton et al., 2019](#)) for  $pCO_2$  and  $pH$  ([Sutton et al., 2019](#)). Asterisk (\*) marks the two stations with also  $A_T$  and DIC observations ([Olafsson et al., 2010](#)) available for assessments.


**Table A1.** Information of feature regions [analyzed](#)–[analysed](#) in this study (Fig. A1a - cyan boxes).

Notations	Regions	Coordinates	
		Latitude	Longitude
A	California Current System	25°N-45°N	130°W-110°W
B	Humboldt Current System	30°S-0°	90°W-70°W
C	Labrador Sea	45°N-65°N	70°W-45°W
D	Amazon river mouth	5°S-15°N	60°W-40°W
E	Western South Atlantic	60°S-40°S	70°W-50°W
F	Northern Europe	50°N-70°N	10°W-25°E
G	Canary Current System	5°N-30°N	30°W-10°W
H	Benguela Current System	35°S-15°S	5°E-20°E
I	Western Arabian Sea	5°N-24°N	45°E-65°E
J	Sea of Japan	30°N-50°N	120°E-150°E




**Figure A2.** Spatial distribution of total months in 1985-2021 containing SOCATv2022 gridded data. Left:  $1^\circ$ -data product (r100), right:  $0.25^\circ$ -data product (r025). Open-ocean data (O) in each  $0.25^\circ$ -grid box is created by setting conservatively the open-ocean SOCATv2022 data at the  $1^\circ$ -grid box containing it. The coastal-ocean SOCATv2022 data (C) are assigned within 400 km from the shoreline (Fig. A1a).






**Figure A3.** Global maps of mean model-data difference (Bias, abcd) and root-of-mean-square-deviation (RMSD, efgh) between the reconstruction and SOCATv2022  $p\text{CO}_2$  [ $\mu\text{atm}$ ] over 1985-2021. Left: CMEM-LSCE-FFNN with a resolution of  $1^\circ$  (r100), right: CMEM-LSCE-FFNN with a resolution of  $0.25^\circ$  (r025). Open-ocean data (O) in each  $0.25^\circ$ -grid box used for evaluation is created by setting conservatively the open-ocean SOCATv2022 data value at the  $1^\circ$ -grid box containing it. Coastal-ocean data (C) are extracted from each of the two SOCATv2022 gridded data products.







**Figure A4.** CMEMS-LSCE  $H^+$  over the global ocean at a spatial resolution of  $0.25^\circ$ . Temporal means of the model best estimate and  $1\sigma$ -uncertainty per grid cell over 1985-2021 are calculated by using Eq. (5).




**Figure A5.** CMEMS-LSCE  $\Omega_{ca}$  over the global ocean at a spatial resolution of  $0.25^\circ$ . Temporal means of the model best estimate and  $1\sigma$ -uncertainty per grid cell over 1985-2021 are calculated by using Eq. (5).




**Figure A6.** CMEMS-LSCE Revelle Factor (RF) over the global ocean at a spatial resolution of  $0.25^\circ$ . Temporal means of the model best estimate per grid cell over 1985-2021 are calculated by using Eq. (5).




**Figure A7.** Spatial distribution of CMEMS SSS and SST product uncertainty over the global ocean at a spatial resolution of  $0.25^\circ$ .  $1\sigma$ -uncertainty is computed per grid cell by using Eq. (5) over 1985-2021.



**Figure A8.** Median percentage of analysis error uncertainty against of climatological mean of surface WOA18 nutrient data: phosphate ( $\text{PO}_4$ ), nitrate ( $\text{NO}_3$ ), and silicate ( $\text{SiO}_2$ ).




**Figure A9.** Spatial distribution of  $R(\sigma, \mu)$  [%] (Eq 8), i.e., the ratio of model uncertainty ( $\sigma$ ) against model best estimate ( $\mu$ ).




**Figure A10.** Monthly time series of SSS and SST at BATS, DYFAMED, ESTOC, and HOT stations (Table 2 and Fig. A1b): **model best estimate** CMEMS data (curve),  $1\sigma$ -uncertainty (envelope), and monthly average of observations (point). Means of **the best estimate** CMEMS data and  $1\sigma$ -uncertainty ( $\mu \pm \sigma$ ) calculated over the observing time span are shown in brackets if accessible. Statistics include **the number** of months with observations ( $N$ ), Bias, RMSD, and  $r^2$ .  $\sigma_{SSS}^t$  [ $\sigma_{SST}^t$ ] stands for temporal standard deviation from monthly averages of SSS and SST observations. Temporal variations in SSS observations are poorly described in the CMEMS SSS time series (Table 1) used in CMEMS-LSCE reconstructions.






**Figure A11.** a) Global surface seawater  $pH$  trend over the period 1985-2021, b)  $1\sigma$ -uncertainties associated to trend estimates, c)  $\sigma$ -to- $\mu$  ratio  $R(\sigma, \mu)$  [%] (Eq. 8) between uncertainty estimates (b) and the best trend estimates (a), d) mask applied over the regions where  $R(\mu, \sigma) > 10\%$ .




**Figure A12.** a) Global surface seawater  $\Omega_{ar}$  trend over the period 1985-2021, b)  $1\sigma$ -uncertainties associated to trend estimates, c)  $\sigma$ -to- $\mu$  ratio  $R(\sigma, \mu)$  [%] (Eq. 8) between uncertainty estimates (b) and the best trend estimates (a), d) mask applied over the regions where  $R(\mu, \sigma) > 20\%$ .



**Figure A13.** a) Yearly global area-weighted mean of surface ocean saturation states with respect to calcite ( $\Omega_{ca}$ ): Global means of the best estimate ( $\mu$ , plain line) and uncertainty ( $\sigma$ , envelop) are computed with Eq. (7a). b) Global trend maps of  $\Omega_{ca}$  over 1985-2021: Cross-hatching covers the regions where with uncertainty of a trend estimate over 20% of the trend value.





**Figure A14.** Linear trend estimates learned on 100-member ensemble (grey points) of yearly mean time series of  $pH$  and  $\Omega_{ar}$  at different stations (Bates et al., 2014).  $\mu \pm \sigma$  present linear slope and residual standard deviation. Black or blue lines stand for linear fits over the full or sub-period in 1985-2021 (see Table 5 for comparison).

**Table A2.** Information of moored time series of ~~coastal-surface-ocean-surface-ocean~~  $p\text{CO}_2$  and  $p\text{H}$  observations (Sutton et al., 2019).

Stations	Abbreviations	Coordinates	Date range
Ala Wai Water Quality Buoy Pacific island	ALAWAI	21.3°N, 157.9°W	06/2008-12/2020
Bay of Bengal Ocean Indian Ocean	BOBOA	15.0°N, 90.0°W	11/2013-12/2017
Bermuda Testbed Mooring	BTM	31.5°N, 64.2°W	10/2005f-12/2006
Cape Arago	CAPEARAGO	43.3°N, 124.5°W	06/2017-12/2020
Cape Elizabeth	CAPEELIZABETH	47.4°N, 124.7°W	06/2006-05/2020
California Current Ecosystem 2	CCE2	34.3°N, 120.8°W	01/2010-06/2021
Chá b'a Buoy in the Northwest Enhanced Moored Observatory and Olympic Coast NMS	CHABA	47.9°N, 126°W	07/2010-09/2020
Chuuk Lagoon Ocean Acidification Mooring	CHUUK	7.5°N, 151.9°E	11/2011-12/2017
Cheeca Rocks Ocean Acidification Mooring in Florida Keys National Marine Sanctuary	CHEECAROCKS	24.9°N, 80.6°W	12/2011-12/2021
Coastal Louisiana buoy	COASTALLA	28.5°N, 90.3°W	07/2017-08/2020
Central Gulf of Mexico Ocean Observing System Station 01	COASTALMS	30.0°N, 88.6°W	05/2009-05/2017
Crescent Reef Bermuda Buoy	CRESCENTREEF	32.4°N, 64.8°W	11/2010-12/2014
Coral Reef Instrumented Monitoring Platform 1	CRIMP1	21.4°N, 157.8°W	12/2005-12/2007
Coral Reef Instrumented Monitoring Platform 2	CRIMP2	21.5°N, 157.8°W	06/2008-12/2019
Chesapeake Bay Interpretive Buoy System Ocean Acidification Buoy at First Landing	FIRSTLANDING	37.0°N, 76.1°W	04/2018-09/2020
NANOOS ORCA buoy in Dabob Bay	DABOB	47.8°N, 122.8°W	06/2011-01/2016
Gulf of Alaska Ocean Acidification Mooring	GAKOA	59.9°N, 149.4°W	05/2011-12/2017
NDBC Buoy 41008 in Gray's Reef National Marine Sanctuary	GRAYSREEF	31.4°N, 80.9°W	07/2006-08/2018
Coastal Western Gulf of Maine Mooring	GULFOFMAINE	43.0°N, 70.5°W	07/2006-06/2021
Hog Reef Bermuda Buoy	HOGREEF	32.5°N, 64.8°W	12/2010-12/2017
North Atlantic Ocean Acidification Mooring	ICELAND	68.0°N, 12.7°W	08/2013-06/2017
Kaneohe Bay Ocean Acidification Offshore Observatory	KANEOHE	21.5°N, 157.8°W	09/2011-12/2019
Kuroshio Extension Observatory	KEO	32.3°N, 144.6°E	09/2007-12/2019
Kilo Nalu Water Quality Buoy at South Shore Oahu	KILONALU	21.3°N, 157.9°W	08/2008-12/2018
Kodiak Alaska Ocean Acidification Mooring	KODIAK	57.7°N, 152.3°W	03/2013-12/2015
La Parguera Ocean Acidification Mooring	LAPARGUERA	18.0°N, 67.1°W	01/2009-12/2018
Southeastern Bering Sea Mooring Site 2	M2	56.5°N, 164.0°W	05/2013-09/2017
Newport Hydrographic Line Station 10 Ocean Acidification Mooring	NH10	44.9°N, 124.8°W	04/2014-12/2016
Ocean Station Papa	PAPA	50.1°N, 144.8°W	06/2007-12/2019
Southeast Alaska Ocean Acidification Mooring	SEAK	56.3°N, 134.7°W	03/2013-12/2015
Southern Ocean Flux Station	SOFS	46.8°S, 142.0°E	11/2011-12/2020
Stratus	STRATUS	19.7°S, 85.6°W	10/2006-12/2017
National Data Buoy Center (NDBC) Tropical Atmosphere Ocean	TAO110W	0.0°N, 110.0°W	09/2009-12/2018
NDBC Tropical Atmosphere Ocean	TAO125W	0.0°N, 125.0°W	05/2004-12/2020
NDBC Tropical Atmosphere Ocean	TAO140W	0.0°N, 140.0°W	05/2004-4/2019
NDBC Tropical Atmosphere Ocean	TAO155W	0.0°N, 155.0°W	01/2010-07/2020
NDBC Tropical Atmosphere Ocean	TAO165E	0.0°N, 165.0°E	02/2010-12/2019
NDBC Tropical Atmosphere Ocean	TAO170W	0.0°N, 170.0°W	07/2005-12/2017
NDBC Tropical Atmosphere Ocean	TAO8S165E	8.0°S, 165.0°E	06/2009-11/2011
ORCA buoy at Twanoh in Hood Canal	TWANOH	47.4°N, 123.0°W	08/2009-12/2020
Woods Hole Oceanographic Institution Hawaii Ocean Time-series Station	WHOTS	22.7°N, 158.0°W	12/2004-12/2018

**Table A3.** Statistics computed between CMEMS-LSCE datasets (0.25°) and time series of  $p\text{CO}_2$  and  $p\text{H}$  measurements (Sutton et al., 2019) at open-ocean (O) and coastal stations (C): total numbers of monthly mean observations (N), temporal standard deviation of observations from their monthly averages ( $\sigma^t$ ), RMSD (Eq. 10), and  $r^2$  (Eq. 11). See Table A2 and Fig. A1b for stations' information and locations.

Stations	$p\text{CO}_2$ [ $\mu\text{atm}$ ]				$p\text{H}$ [-]			
	N	$\sigma^t$	RMSD	$r^2$	N	$\sigma^t$	RMSD	$r^2$
<del>1</del> -O1. BOBOA	42	9.9	9.6	0.69	19	0.010	0.011	0.71
O2. BTM	15	7.2	11.4	0.98	0	—	—	—
O3. CHUUK	66	10.9	23.5	0.60	33	0.014	0.033	0.78
O5. CRESCENTREEF	41	13.6	45.0	0.93	0	—	—	—
O5. HOGREEF	60	23.3	47.8	0.85	0	—	—	—
O6. ICELAND	24	8.1	16.9	0.68	4	0.007	0.092	0.94
O7. KEO	130	9.0	10.1	0.92	48	0.010	0.014	0.86
O8. PAPA	135	5.7	8.7	0.60	103	0.007	0.017	0.40
O9. SOFS	61	7.1	7.8	0.81	0	—	—	—
O10. STRATUS	116	6.3	9.2	0.80	10	0.004	0.042	0.43
O11. TAO110W	67	22.8	17.7	0.79	0	—	—	—
O12. TAO125W	124	16.5	16.0	0.43	0	—	—	—
O13. TAO140W	92	11.4	12.0	0.44	0	—	—	—
O14. TAO155W	45	11.3	12.2	0.75	0	—	—	—
O15. TAO165E	39	9.1	19.8	0.76	0	—	—	—
O16. TAO170W	87	9.0	17.9	0.45	0	—	—	—
O17. TAO8S165E	29	7.1	10.5	0.49	0	—	—	—
O18. WHOTS	143	4.7	8.0	0.80	23	0.004	0.01	0.57
C1. ALAWAI	112	20.3	22.8	0.32	0	—	—	—
C2. CAPEARAGO	33	<del>65.60</del> 65.6	<del>79.86</del> 79.9	0.19	31	0.086	0.069	0.22
<del>2</del> -C3. CAPEELIZABETH	92	<del>42.54</del> 42.5	<del>41.44</del> 41.4	0.52	11	0.061	0.057	0.69
<del>3</del> -C4. CCE2	127	<del>45.31</del> 45.3	<del>32.44</del> 32.4	0.16	58	0.048	0.035	0.24
<del>4</del> -C5. CHABA	75	51.2	65.3	0.44	42	0.056	0.064	0.58
C6. CHEECAROCKS	73	<del>44.05</del> 44.1	<del>62.42</del> 62.4	0.25	40	0.038	0.066	0.21
<del>5</del> -C7. COASTALLA	22	<del>59.15</del> 59.2	<del>57.41</del> 57.4	0.52	2	0.078	0.068	—
<del>6</del> -C8. COASTALMS	41	<del>43.04</del> 43.0	<del>42.50</del> 42.5	0.51	15	0.062	0.065	0.25
<del>7</del> -C9. CRIMP1	23	26.4	95.6	0.62	0	—	—	—
C10. CRIMP2	119	85.4	90.7	0.58	0	—	—	—
C11. DABOB	24	55.0	89.3	0.74	0	—	—	—
C12. FIRSTLANDING	17	<del>69.98</del> 70.0	<del>77.32</del> 77.3	0.49	2	0.061	0.042	—
<del>8</del> -C13. GAKOA	64	20.3	66.0	0.73	0	—	—	—
C14. GRAYSREEF	96	<del>20.12</del> 20.1	<del>38.34</del> 38.3	0.65	49	0.020	0.040	0.66
C15. GULFOFMAINE	144	26.7	31.2	0.54	77	0.029	0.042	0.27
C16. KANEOHE	49	20.9	23.5	0.36	35	0.025	0.034	0.25
C17. KILONALU	69	14.3	11.6	0.56	0	—	—	—
C18. KODIAK	34	31.7	62.1	0.78	0	—	—	—
C19. LAPARGUERA	103	12.9	41.6	0.41	48	0.012	0.038	0.33
C20. M2	23	25.4	55.8	0.24	0	—	—	—
C21. NH10	25	64.6	31.4	0.46	15	0.058	0.048	0.42
C22. SEAK	31	53.4	141.3	0.82	0	—	—	—
C23. TWANOH	57	126.2	199.9	0.33	0	—	—	—

**Table A4.** Statistics computed between CMEMS-LSCE datasets ( $0.25^\circ$ ) and time series of  $A_T$  and DIC measurements (0–10 m depth): total numbers of monthly mean observations (N), temporal standard deviation of observations from their monthly averages ( $\sigma^t$ ), RMSD (Eq. 10), and  $r^2$  (Eq. 11). See Table 2 and Fig. A1b for stations’ information and locations.

Stations	$A_T$ [ $\mu\text{mol kg}^{-1}$ ]				DIC [ $\mu\text{mol kg}^{-1}$ ]			
	N	$\sigma^t$	RMSD	$r^2$	N	$\sigma^t$	RMSD	$r^2$
1. AWIPEV	52	15.8	32.0	0.33	52	17.7	29.6	0.71
2. BATS	303	2.4	9.6	0.34	351	1.9	8.2	0.82
3. DYFAMED	84	1.3	145.7	0.12	84	2.0	124.7	0.61
4. ESTOC	298	1.3	8.3	0.03	108	1.4	11.2	0.47
5. HOT	298	0.8	12.5	0.32	298	0.6	10.6	0.70
6. ICELAND	27	2.5	13.4	0.24	27	0.8	13.7	0.79
7. IRMINGER	29	1.8	8.4	0.30	23	1.7	14.6	0.84
8. KERFIX	23	1.0	7.4	0.22	23	3.4	14.0	0.52

## References

- Bakker, D., Alin, S., Becker, M., Bittig, H., Castaño-Primo, R., Feely, R. A., Gritzalis, T., Kadono, K., Kozyr, A., Lauvset, S. K., Metzl, N., Munro, D., Nakaoka, S.-i., Nojiri, Y., O'Brien, K., Olsen, A., Pfeil, B., Pierrot, D., Steinhoff, T., Sullivan, K., Sutton, A., Sweeney, C., Tilbrook, B., Wada, C., Wanninkhof, R., Wranne, A. W., et al.: SOCAT version 2022 for quantification of ocean CO<sub>2</sub> uptake, [https://www.socat.info/wp-content/uploads/2022/06/2022\\_Poster\\_SOCATv2022\\_release.pdf](https://www.socat.info/wp-content/uploads/2022/06/2022_Poster_SOCATv2022_release.pdf), 2022.
- Bakker, D. C. E., Pfeil, B., Landa, C. S., Metzl, N., O'Brien, K. M., Olsen, A., Smith, K., Cosca, C., Harasawa, S., Jones, S. D., Nakaoka, S., Nojiri, Y., Schuster, U., Steinhoff, T., Sweeney, C., Takahashi, T., Tilbrook, B., Wada, C., Wanninkhof, R., Alin, S. R., Balestrini, C. F., Barbero, L., Bates, N. R., Bianchi, A. A., Bonou, F., Boutin, J., Bozec, Y., Burger, E. F., Cai, W.-J., Castle, R. D., Chen, L., Chierici, M., Currie, K., Evans, W., Featherstone, C., Feely, R. A., Fransson, A., Goyet, C., Greenwood, N., Gregor, L., Hankin, S., Hardman-Mountford, N. J., Harlay, J., Hauck, J., Hoppema, M., Humphreys, M. P., Hunt, C. W., Huss, B., Ibáñez, J. S. P., Johannessen, T., Keeling, R., Kitidis, V., Körtzinger, A., Kozyr, A., Krasakopoulou, E., Kuwata, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lo Monaco, C., Manke, A., Mathis, J. T., Merlivat, L., Millero, F. J., Monteiro, P. M. S., Munro, D. R., Murata, A., Newberger, T., Omar, A. M., Ono, T., Paterson, K., Pearce, D., Pierrot, D., Robbins, L. L., Saito, S., Salisbury, J., Schlitzer, R., Schneider, B., Schweitzer, R., Sieger, R., Skjelvan, I., Sullivan, K. F., Sutherland, S. C., Sutton, A. J., Tadokoro, K., Telszewski, M., Tuma, M., van Heuven, S. M. A. C., Vandemark, D., Ward, B., Watson, A. J., and Xu, S.: A multi-decade record of high-quality *f*CO<sub>2</sub> data in version 3 of the Surface Ocean CO<sub>2</sub> Atlas (SOCAT), *Earth System Science Data*, 8, 383–413, <https://doi.org/10.5194/essd-8-383-2016>, 2016.
- Bates, N. R., Astor, Y. M., Church, M. J., Currie, K., Dore, J. E., González-Dávila, M., Lorenzoni, L., Muller-Karger, F., Olafsson, J., and Santana-Casiano, J. M.: A time-series view of changing surface ocean chemistry due to ocean uptake of anthropogenic CO<sub>2</sub> and ocean acidification, *Oceanography*, 27, 126–141, 2014.
- Bopp, L., Resplandy, L., Orr, J. C., Doney, S. C., Dunne, J. P., Gehlen, M., Halloran, P., Heinze, C., Ilyina, T., Séférian, R., Tjiputra, J., and Vichi, M.: Multiple stressors of ocean ecosystems in the 21st century: projections with CMIP5 models, *Biogeosciences*, 10, 6225–6245, 2013.
- Bresnahan, P. J., Takeshita, Y., Wirth, T., Martz, T. R., Cyronak, T., Albright, R., Wolfe, K., Warren, J. K., and Mertz, K.: Autonomous in situ calibration of ion-sensitive field effect transistor pH sensors, *Limnology and Oceanography: Methods*, 19, 132–144, 2021.
- Broullón, D., Pérez, F. F., Velo, A., Hoppema, M., Olsen, A., Takahashi, T., Key, R. M., Tanhua, T., González-Dávila, M., Jeansson, E., et al.: A global monthly climatology of total alkalinity: a neural network approach, *Earth System Science Data*, 11, 1109–1127, 2019.
- Canadell, J. G., Monteiro, P. M., Costa, M. H., Da Cunha, L. C., Cox, P. M., Alexey, V., Henson, S., Ishii, M., Jaccard, S., Koven, C., et al.: Global carbon and other biogeochemical cycles and feedbacks In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, <https://doi.org/10.1017/9781009157896.007>, 2021.
- Carter, B. R., Williams, N. L., Gray, A. R., and Feely, R. A.: Locally interpolated alkalinity regression for global alkalinity estimation, *Limnology and Oceanography: Methods*, 14, 268–277, <https://doi.org/https://doi.org/10.1002/lom3.10087>, 2016.
- Carter, B. R., Feely, R. A., Williams, N. L., Dickson, A. G., Fong, M. B., and Takeshita, Y.: Updated methods for global locally interpolated estimation of alkalinity, pH, and nitrate, *Limnology and Oceanography: Methods*, 16, 119–131, <https://doi.org/https://doi.org/10.1002/lom3.10232>, 2018.
- Chau, T. T. T., Gehlen, M., and Chevallier, F.: Global ocean surface carbon product, Research Report CMEMS-MOB-QUID-015-008, Le Laboratoire des Sciences du Climat et de l'Environnement, <https://doi.org/10.48670/moi-00047>, quality Information Document, 2022a.

- 810 Chau, T. T. T., Gehlen, M., and Chevallier, F.: A seamless ensemble-based reconstruction of surface ocean  $p\text{CO}_2$  and air–sea  $\text{CO}_2$  fluxes over the global coastal and open oceans, *Biogeosciences*, 19, 1087–1109, <https://doi.org/10.5194/bg-19-1087-2022>, 2022b.
- Chau, T. T. T., Gehlen, M., Metzl, N., and Chevallier, F.: CMEMS-LSCE: A global 0.25-degree, monthly reconstruction of the surface ocean carbonate system, ESPRI/IPSL [dataset], <https://doi.org/10.14768/a2f0891b-763a-49e9-af1b-78ed78b16982>, 2023.
- Chevallier, F.: On the parallelization of atmospheric inversions of  $\text{CO}_2$  surface fluxes within a variational framework, *Geoscientific Model Development*, 6, 783–790, <https://doi.org/10.5194/gmd-6-783-2013>, 2013.
- 815 Chevallier, F., Fisher, M., Peylin, P., Serrar, S., Bousquet, P., Bréon, F.-M., Chédin, A., and Ciais, P.: Inferring  $\text{CO}_2$  sources and sinks from satellite observations: Method and application to TOVS data, *Journal of Geophysical Research: Atmospheres*, 110, <https://doi.org/https://doi.org/10.1029/2005JD006390>, 2005.
- Chevallier, F., Ciais, P., Conway, T. J., Aalto, T., Anderson, B. E., Bousquet, P., Brunke, E. G., Ciattaglia, L., Esaki, Y., Fröhlich, M., Gomez, A., Gomez-Pelaez, A. J., Haszpra, L., Krummel, P. B., Langenfelds, R. L., Leuenberger, M., Machida, T., Maignan, F., Matsueda, H., Morguí, J. A., Mukai, H., Nakazawa, T., Peylin, P., Ramonet, M., Rivier, L., Sawa, Y., Schmidt, M., Steele, L. P., Vay, S. A., Vermeulen, A. T., Wofsy, S., and Worthy, D.:  $\text{CO}_2$  surface fluxes at grid point scale estimated from a global 21 year reanalysis of atmospheric measurements, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/https://doi.org/10.1029/2010JD013887>, 2010.
- 820 [Cooley, S., Schoeman, D., Bopp, L., Boyd, P., Donner, S., Ito, S.-i., Kiessling, W., Martinetto, P., Ojea, E., Racault, M.-F., et al.: Oceans and coastal ecosystems and their services, in: IPCC AR6 WGII, Cambridge University Press, 2022.](#)
- Coppola, L., Boutin, J., Gattuso, J.-P., Lefevre, D., and Metzl, N.: The Carbonate System in the Ligurian Sea, chap. 4, pp. 79–103, John Wiley & Sons, Ltd, <https://doi.org/https://doi.org/10.1002/9781119706960.ch4>, 2020.
- Coppola, L., Diamond, R. E., and Carval, T.: Dyfamed observatory data. SEANOE [dataset], <https://doi.org/0.17882/43749>, 2021.
- Denvil-Sommer, A., Gehlen, M., Vrac, M., and Mejia, C.: LSCE-FFNN-v1: a two-step neural network model for the reconstruction of surface ocean  $p\text{CO}_2$  over the global ocean, *Geoscientific Model Development*, 12, 2091–2105, <https://doi.org/10.5194/gmd-12-2091-2019>, 2019.
- 830 Dickson, A. G.: Standard potential of the reaction  $-\text{AgCl}(\text{s}) + 1/2\text{H}_2(\text{g}) = \text{Ag}(\text{s}) + \text{HCl}(\text{aq})$  and the standard acidity constant of the ion  $\text{HSO}_4^-$  in synthetic sea-water from 273.15-K to 318.15-K, *Journal of Chemical Thermodynamics*, 22, 113–127, [https://doi.org/doi:10.1016/0021-9614\(90\)90074-z](https://doi.org/doi:10.1016/0021-9614(90)90074-z), 1990.
- Dickson, A. G.: The carbon dioxide system in seawater: equilibrium chemistry and measurements in Guide to best practices for ocean acidification research and data reporting, [https://www.pmel.noaa.gov/co2/files/dickson\\_thecarbondioxidesysteminseawater\\_equilibriumchemistryandmeasurementspp17-40.pdf](https://www.pmel.noaa.gov/co2/files/dickson_thecarbondioxidesysteminseawater_equilibriumchemistryandmeasurementspp17-40.pdf), 2010.
- 835 Dickson, A. G., Sabine, C. L., and Christian, J. R.: Guide to best practices for ocean  $\text{CO}_2$  measurements., North Pacific Marine Science Organization, 2007.
- [Doney, S. C.: The growing human footprint on coastal and open-ocean biogeochemistry, science, 328, 1512–1516, 2010.](#)
- 840 Doney, S. C., Fabry, V. J., Feely, R. A., and Kleypas, J. A.: Ocean acidification: the other  $\text{CO}_2$  problem, *Annual review of marine science*, 1, 169–192, 2009.
- Doney, S. C., Busch, D. S., Cooley, S. R., and Kroeker, K. J.: The impacts of ocean acidification on marine ecosystems and reliant human communities, *Annual Review of Environment and Resources*, 45, 2020.
- Dore, J. E., Lukas, R., Sadler, D. W., Church, M. J., and Karl, D. M.: Physical and biogeochemical modulation of ocean acidification in the central North Pacific, *Proceedings of the National Academy of Sciences*, 106, 12 235–12 240, 2009.
- 845 Droghei, R., Buongiorno Nardelli, B., and Santoleri, R.: A new global sea surface salinity and density dataset from multivariate observations (1993–2016), *Frontiers in Marine Science*, 5, 84, 2018.

- Fabry, V. J., Seibel, B. A., Feely, R. A., and Orr, J. C.: Impacts of ocean acidification on marine fauna and ecosystem processes, *ICES Journal of Marine Science*, 65, 414–432, 2008.
- 850 Feely, R. A., Wanninkhof, R., Takahashi, T., and Tans, P.: Influence of El Niño on the equatorial Pacific contribution to atmospheric CO<sub>2</sub> accumulation, *Nature*, 398, 597–601, 1999.
- Feely, R. A., Sabine, C. L., Takahashi, T., and Wanninkhof, R.: Uptake And Storage Of Carbon Dioxide In The Ocean: The Global CO<sub>2</sub> Survey, *OCEANOGRAPHY-WASHINGTON DC-OCEANOGRAPHY SOCIETY-*, 14, 18–32, 2001.
- Fischer, P., Schwanitz, M., Loth, R., Posner, U., Brand, M., and Schröder, F.: First year of practical experiences of the new Arctic AWIPEV-COSYNA cabled Underwater Observatory in Kongsfjorden, Spitsbergen, *Ocean Science*, 13, 259–272, <https://doi.org/10.5194/os-13-259-2017>, 2017.
- 855 Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Gregor, L., Hauck, J., Le Quéré, C., Luijckx, I. T., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Alkama, R., Arneeth, A., Arora, V. K., Bates, N. R., Becker, M., Bellouin, N., Bittig, H. C., Bopp, L., Chevallier, F., Chini, L. P., Cronin, M., Evans, W., Falk, S., Feely, R. A., Gasser, T., Gehlen, M., Gkritzalis, T., Gloege, L., Grassi, G., Gruber, N., Gürses, O., Harris, I., Hefner, M., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Jain, A. K., Jersild, A., Kadono, K., Kato, E., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Landschützer, P., Lefèvre, N., Lindsay, K., Liu, J., Liu, Z., Marland, G., Mayot, N., McGrath, M. J., Metzl, N., Monacci, N. M., Munro, D. R., Nakaoka, S.-I., Niwa, Y., O’Brien, K., Ono, T., Palmer, P. I., Pan, N., Pierrot, D., Pocock, K., Poulter, B., Resplandy, L., Robertson, E., Rödenbeck, C., Rodriguez, C., Rosan, T. M., Schwinger, J., Séférian, R., Shutler, J. D., Skjelvan, I., Steinhoff, T., Sun, Q., Sutton, 860 A. J., Sweeney, C., Takao, S., Tanhua, T., Tans, P. P., Tian, X., Tian, H., Tilbrook, B., Tsujino, H., Tubiello, F., van der Werf, G. R., Walker, A. P., Wanninkhof, R., Whitehead, C., Willstrand Wranne, A., Wright, R., Yuan, W., Yue, C., Yue, X., Zaehle, S., Zeng, J., and Zheng, B.: Global Carbon Budget 2022, *Earth System Science Data*, 14, 4811–4900, <https://doi.org/10.5194/essd-14-4811-2022>, 2022.
- Garcia, H., Weathers, K., Paver, C., Smolyar, I., Boyer, T., Locarnini, M., Zweng, M., Mishonov, A., Baranova, O., Seidov, D., et al.: World ocean atlas 2018. Vol. 4: Dissolved inorganic nutrients (phosphate, nitrate and nitrate+ nitrite, silicate), <https://archimer.ifremer.fr/doc/00651/76336/>, 2019.
- 870 Gattuso, J.-P. and Hansson, L.: *Ocean acidification*, Oxford university press, 2011.
- Gattuso, J.-P., Magnan, A., Billé, R., Cheung, W. W. L., Howes, E. L., Joos, F., Allemand, D., Bopp, L., Cooley, S. R., Eakin, C. M., Hoegh-Guldberg, O., Kelly, R. P., Pörtner, H. O., Rogers, A. D., Baxter, J. M., Laffoley, D., Osborn, D., Rankovic, A., Rochette, J., Sumaila, U. R., TreyerAlliouane, S., and Turley, C.: ~~Contrasting futures for ocean and society from different anthropogenic CO<sub>2</sub> emissions scenarios~~, *Science* Fischer, P.: High-frequency, 349, aac4722, , 2015.-year-round time series of the carbonate chemistry in a high-Arctic fjord (Svalbard), *Earth System Science Data*, 15, 2809–2825, <https://doi.org/10.5194/essd-15-2809-2023>, 2023.
- 875
- Gehlen, M., Gruber, N., Gangstø, R., Bopp, L., and Oschlies, A.: Biogeochemical consequences of ocean acidification and feedbacks to the earth system, *Ocean acidification*, 1, 230–248, 2011.
- 880 González-Dávila, M. and Santana-Casiano, J.: Sea surface and atmospheric fco2 data measured during the estoc time series cruises from 1995-2009, CDIAC, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tennessee. doi, 10, 2009.
- Good, S., Fiedler, E., Mao, C., Martin, M. J., Maycock, A., Reid, R., Roberts-Jones, J., Searle, T., Waters, J., While, J., and Worsfold, M.: The Current Configuration of the OSTIA System for Operational Production of Foundation Sea Surface Temperature and Ice Concentration Analyses, *Remote Sensing*, 12, <https://www.mdpi.com/2072-4292/12/4/720>, 2020.
- 885



- Gregor, L. and Gruber, N.: OceanSODA-ETHZ: a global gridded data set of the surface ocean carbonate system for seasonal to decadal studies of ocean acidification, *Earth System Science Data*, 13, 777–808, <https://doi.org/10.5194/essd-13-777-2021>, 2021.
- Gregor, L., Lebehot, A. D., Kok, S., and Scheel Monteiro, P. M.: A comparative assessment of the uncertainties of global surface ocean CO<sub>2</sub> estimates using a machine-learning ensemble (CSIR-ML6 version 2019a) – have we hit the wall?, *Geoscientific Model Development*, 12, 5113–5136, <https://doi.org/10.5194/gmd-12-5113-2019>, 2019.
- Gruber, N., Clement, D., Carter, B. R., Feely, R. A., Van Heuven, S., Hoppema, M., Ishii, M., Key, R. M., Kozyr, A., Lauvset, S. K., et al.: The oceanic sink for anthropogenic CO<sub>2</sub> from 1994 to 2007, *Science*, 363, 1193–1199, 2019.
- Hauck, J., Zeising, M., Le Quéré, C., Gruber, N., Bakker, D. C., Bopp, L., Chau, T. T. T., Gürses, Ö., Ilyina, T., Landschützer, P., Lenton, A., Resplandy, L., Rödenbeck, C., Schwinger, J., and Séférian, R.: Consistency and Challenges in the Ocean Carbon Sink Estimate for the Global Carbon Budget, *Frontiers in Marine Science*, 7, 852, <https://doi.org/10.3389/fmars.2020.571720>, 2020.
- Hoffmann, L. J., Breitbarth, E., Boyd, P. W., and Hunter, K. A.: Influence of ocean warming and acidification on trace metal biogeochemistry, *Marine Ecology Progress Series*, 470, 191–205, 2012.
- Hopkins, F. E., Suntharalingam, P., Gehlen, M., Andrews, O., Archer, S. D., Bopp, L., Buitenhuis, E., Dadou, I., Duce, R., Goris, N., et al.: The impacts of ocean acidification on marine trace gases and the implications for atmospheric chemistry and climate, *Proceedings of the Royal Society A*, 476, 20190769, 2020.
- Ibáñez, J. S. P., Diverrès, D., Araujo, M., and Lefèvre, N.: Seasonal and interannual variability of sea-air CO<sub>2</sub> fluxes in the tropical Atlantic affected by the Amazon River plume, *Global Biogeochemical Cycles*, 29, 1640–1655, 2015.
- Iida, Y., Takatani, Y., Kojima, A., and Ishii, M.: Global trends of ocean CO<sub>2</sub> sink and ocean acidification: an observation-based reconstruction of surface ocean inorganic carbon variables, *Journal of Oceanography*, 77, 323–358, 2021.
- Jiang, L.-Q., Carter, B. R., Feely, R. A., Lauvset, S. K., and Olsen, A.: Surface ocean pH and buffer capacity: past, present and future, *Scientific reports*, 9, 1–11, 2019.
- Jiang, L.-Q., Dunne, J., Carter, B. R., Tjiputra, J. F., Terhaar, J., Sharp, J. D., Olsen, A., Alin, S., Bakker, D. C. E., Feely, R. A., Gattuso, J.-P., Hogan, P., Ilyina, T., Lange, N., Lauvset, S. K., Lewis, E. R., Lovato, T., Palmieri, J., Santana-Falcón, Y., Schwinger, J., Séférian, R., Strand, G., Swart, N., Tanhua, T., Tsujino, H., Wanninkhof, R., Watanabe, M., Yamamoto, A., and Ziehn, T.: Global Surface Ocean Acidification Indicators From 1750 to 2100, *Journal of Advances in Modeling Earth Systems*, 15, e2022MS003563, 2023.
- Jones, S., Le Quéré, C., and Rödenbeck, C.: Autocorrelation characteristics of surface ocean pCO<sub>2</sub> and air-sea CO<sub>2</sub> fluxes, *Global biogeochemical cycles*, 26, 2012.
- Key, R. M., Kozyr, A., Sabine, C. L., Lee, K., Wanninkhof, R., Bullister, J. L., Feely, R. A., Millero, F. J., Mordy, C., and Peng, T.-H.: A global ocean carbon climatology: Results from Global Data Analysis Project (GLODAP), *Global Biogeochemical Cycles*, 18, <https://doi.org/https://doi.org/10.1029/2004GB002247>, 2004.
- Kwiatkowski, L., Torres, O., Bopp, L., Aumont, O., Chamberlain, M., Christian, J. R., Dunne, J. P., Gehlen, M., Ilyina, T., John, J. G., Lenton, A., Li, H., Lovenduski, N. S., Orr, J. C., Palmieri, J., Santana-Falcón, Y., Schwinger, J., Séférian, R., Stock, C. A., Tagliabue, A., Takano, Y., Tjiputra, J., Toyama, K., Tsujino, H., Watanabe, M., Yamamoto, A., Yool, A., and Ziehn, T.: Twenty-first century ocean warming, acidification, deoxygenation, and upper-ocean nutrient and primary production decline from CMIP6 model projections, *Biogeosciences*, 17, 3439–3470, 2020.
- Körtzinger, A.: Determination of carbon dioxide partial pressure (pCO<sub>2</sub>), chap. 9, pp. 149–158, John Wiley & Sons, Ltd, <https://doi.org/https://doi.org/10.1002/9783527613984.ch9>, 1999.

- Landschützer, P., Gruber, N., and Bakker, D. C.: Decadal variations and trends of the global ocean carbon sink, *Global Biogeochemical Cycles*, 30, 1396–1417, 2016.
- 925 Landschützer, P., Ilyina, T., and Lovenduski, N. S.: Detecting Regional Modes of Variability in Observation-Based Surface Ocean  $p\text{CO}_2$ , *Geophysical Research Letters*, 46, 2670–2679, 2019.
- Landschützer, P., Laruelle, G. G., Roobaert, A., and Regnier, P.: A uniform  $p\text{CO}_2$  climatology combining open and coastal oceans, *Earth System Science Data*, 12, 2537–2553, <https://doi.org/10.5194/essd-12-2537-2020>, 2020.
- Laruelle, G. G., Landschützer, P., Gruber, N., Tison, J.-L., Delille, B., and Regnier, P.: Global high-resolution monthly  $p\text{CO}_2$  climatology for  
930 the coastal ocean derived from neural network interpolation, *Biogeosciences*, 14, 4545–4561, 2017.
- Lauvset, S. K. and Gruber, N.: Long-term trends in surface ocean pH in the North Atlantic, *Marine Chemistry*, 162, 71–76, 2014.
- Lauvset, S. K., Gruber, N., Landschützer, P., Olsen, A., and Tjiputra, J.: Trends and drivers in global surface ocean pH over the past 3 decades, *Biogeosciences*, 12, 1285–1298, <https://doi.org/10.5194/bg-12-1285-2015>, 2015.
- 935 Lauvset, S. K., Lange, N., Tanhua, T., Bittig, H. C., Olsen, A., Kozyr, A., Alin, S., Álvarez, M., Azetsu-Scott, K., Barbero, L., Becker, S., Brown, P. J., Carter, B. R., da Cunha, L. C., Feely, R. A., Hoppema, M., Humphreys, M. P., Ishii, M., Jeansson, E., Jiang, L.-Q., Jones, S. D., Lo Monaco, C., Murata, A., Müller, J. D., Pérez, F. F., Pfeil, B., Schirnack, C., Steinfeldt, R., Suzuki, T., Tilbrook, B., Ulfso, A., Velo, A., Woosley, R. J., and Key, R. M.: GLODAPv2.2022: the latest version of the global interior ocean biogeochemical data product, *Earth System Science Data*, 14, 5543–5572, <https://doi.org/10.5194/essd-14-5543-2022>, ~~2022~~-2022a.
- 940 Lauvset, S. K., Lange, N., Tanhua, T., Bittig, H. C., Olsen, A., Kozyr, A., Alin, S., Álvarez, M., Azetsu-Scott, K., Barbero, L., Becker, S., Brown, P. J., Carter, B. R., da Cunha, L. C., Feely, R. A., Hoppema, M., Humphreys, M. P., Ishii, M., Jeansson, E., Jiang, L.-Q., Jones, S. D., Lo Monaco, C., Murata, A., Müller, J. D., Pérez, F. F., Pfeil, B., Schirnack, C., Steinfeldt, R., Suzuki, T., Tilbrook, B., Ulfso, A., Velo, A., Woosley, R. J., and Key, R. M.: Global Ocean Data Analysis Project version 2.2022 (GLODAPv2.2022) (NCEI Accession 0257247), NOAA National Centers for Environmental Information [dataset], <https://doi.org/10.25921/1f4w-0t92>, 2022b.
- 945 Lee, K., Tong, L. T., Millero, F. J., Sabine, C. L., Dickson, A. G., Goyet, C., Park, G.-H., Wanninkhof, R., Feely, R. A., and Key, R. M.: Global relationships of total alkalinity with salinity and temperature in surface waters of the world's oceans, *Geophysical research letters*, 33, 2006.
- Leseurre, C., Lo Monaco, C., Reverdin, G., Metzl, N., Fin, J., Mignon, C., and Benito, L.: Summer trends and drivers of sea surface  $f\text{CO}_2$  and pH changes observed in the southern Indian Ocean over the last two decades (1998–2019), *Biogeosciences*, 19, 2599–2625,  
950 <https://doi.org/10.5194/bg-19-2599-2022>, 2022.
- Lewis, E. and Wallace, D.: CO2SYS-Program developed for the CO2 system calculations, Carbon Dioxide Inf Anal Center Report ORNL/CDIAC-105, <https://doi.org/https://doi.org/10.15485/1464255>, 1998.
- Lueker, T. J., Dickson, A. G., and Keeling, C. D.: Ocean  $p\text{CO}_2$  calculated from dissolved inorganic carbon, alkalinity, and equations for K1 and K2: validation based on laboratory measurements of  $\text{CO}_2$  in gas and seawater at equilibrium, *Marine chemistry*, 70, 105–119, 2000.
- 955 Ma, D., Gregor, L., and Gruber, N.: Four Decades of Trends and Drivers of Global Surface Ocean Acidification, *Global Biogeochemical Cycles*, 37, e2023GB007765, <https://doi.org/https://doi.org/10.1029/2023GB007765>, e2023GB007765 2023GB007765, 2023.
- Maritorea, S., d'Andon, O. H. F., Mangin, A., and Siegel, D. A.: Merged satellite ocean color data products using a bio-optical model: Characteristics, benefits and issues, *Remote Sensing of Environment*, 114, 1791–1804, <https://doi.org/https://doi.org/10.1016/j.rse.2010.04.002>, 2010.

- 960 Menemenlis, D., Campin, J., Heimbach, P., Hill, C., Lee, T., Nguyen, A., Schodlok, M., and Zhang, H.: ECCO2: High Resolution Global Ocean and Sea Ice Data Synthesis, 2008, OS31C-1292, 2008.
- Menviel, L., Spence, P., Yu, J., Chamberlain, M., Matear, R., Meissner, K., and England, M. H.: Southern Hemisphere westerlies as a driver of the early deglacial atmospheric CO<sub>2</sub> rise, *Nature Communications*, 9, 1–12, 2018.
- Metzl, N. and Lo Monaco, C.: OISO - Ocean Indien Service d'Observation [dataset], <https://doi.org/10.18142/228>, 1998.
- 965 Michaels, A. F. and Knap, A. H.: Overview of the US JGOFS Bermuda Atlantic Time-series Study and the Hydrostation S program, *Deep Sea Research Part II: Topical Studies in Oceanography*, 43, 157–198, 1996.
- Millero, F. J., Woosley, R., Ditrolio, B., and Waters, J.: Effect of ocean acidification on the speciation of metals in seawater, *Oceanography*, 22, 72–85, 2009.
- Mucci, A.: The solubility of calcite and aragonite in seawater at various salinities, temperatures, and one atmosphere total pressure, *American Journal of Science*, 283, 780–799, 1983.
- 970 Nardelli, B. B., Droghei, R., and Santoleri, R.: Multi-dimensional interpolation of SMOS sea surface salinity with surface temperature and in situ salinity data, *Remote Sensing of Environment*, 180, 392–402, 2016.
- Olafsson, J., Olafsdottir, S., Benoit-Cattin, A., and Takahashi, T.: The Irminger Sea and the Iceland Sea time series measurements of sea water carbon and nutrient chemistry 1983–2008, *Earth System Science Data*, 2, 99–104, 2010.
- 975 Olivier, L., Boutin, J., Reverdin, G., Lefèvre, N., Landschützer, P., Speich, S., Karstensen, J., Labaste, M., Noisel, C., Ritschel, M., Steinhoff, T., and Wanninkhof, R.: Wintertime process study of the North Brazil Current rings reveals the region as a larger sink for CO<sub>2</sub> than expected, *Biogeosciences*, 19, 2969–2988, <https://doi.org/10.5194/bg-19-2969-2022>, 2022.
- Olsen, A., Key, R. M., Van Heuven, S., Lauvset, S. K., Velo, A., Lin, X., Schirnick, C., Kozyr, A., Tanhua, T., Hoppema, M., et al.: The Global Ocean Data Analysis Project version 2 (GLODAPv2)—an internally consistent data product for the world ocean, *Earth System Science Data*, 8, 297–323, 2016.
- 980 Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., et al.: Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms, *Nature*, 437, 681–686, 2005.
- Orr, J. C., Epitalon, J.-M., Dickson, A. G., and Gattuso, J.-P.: Routine uncertainty propagation for the marine carbon dioxide system, *Marine Chemistry*, 207, 84–107, <https://doi.org/https://doi.org/10.1016/j.marchem.2018.10.006>, 2018.
- 985 Palmiéri, J., Orr, J., Dutay, J.-C., Béranger, K., Schneider, A., Beuvier, J., and Somot, S.: Simulated anthropogenic CO<sub>2</sub> storage and acidification of the Mediterranean Sea, *Biogeosciences*, 12, 781–802, 2015.
- Park, P. K.: OCEANIC CO<sub>2</sub> SYSTEM: AN EVALUATION OF TEN METHODS OF INVESTIGATION1, *Limnology and Oceanography*, 14, 179–186, <https://doi.org/https://doi.org/10.4319/lo.1969.14.2.0179>, 1969.
- Pérez, F. F., Olafsson, J., Ólafsdóttir, S. R., Fontela, M., and Takahashi, T.: Contrasting drivers and trends of ocean acidification in the subarctic Atlantic, *Scientific Reports*, 11, 1–16, 2021.
- 990 Pfeil, B., Olsen, A., Bakker, D. C., Hankin, S., Koyuk, H., Kozyr, A., Malczyk, J., Manke, A., Metzl, N., Sabine, C. L., et al.: A uniform, quality-controlled Surface Ocean CO<sub>2</sub> Atlas (SOCAT), *Earth System Science Data*, 5, 125–143, 2013.
- Rödenbeck, C., Keeling, R. F., Bakker, D. C., Metzl, N., Olsen, A., Sabine, C., and Heimann, M.: Global surface-ocean pCO<sub>2</sub> and sea-air CO<sub>2</sub> flux variability from an observation-driven ocean mixed-layer scheme, *Ocean Science*, 9, 193–216, 2013.
- 995 Rödenbeck, C., Bakker, D. C. E., Gruber, N., Iida, Y., Jacobson, A. R., Jones, S., Landschützer, P., Metzl, N., Nakaoka, S., Olsen, A., Park, G.-H., Peylin, P., Rodgers, K. B., Sasse, T. P., Schuster, U., Shutler, J. D., Valsala, V., Wanninkhof, R., and Zeng, J.: Data-based estimates

Rose, A.: Vision: human and electronic, Springer Science & Business Media, 2013.

1000 Sabine, C. L., Key, R. M., Feely, R. A., and Greeley, D.: Inorganic carbon in the Indian Ocean: Distribution and dissolution processes, *Global Biogeochemical Cycles*, 16, 15–1, 2002.

Sabine, C. L., Hankin, S., Koyuk, H., Bakker, D. C. E., Pfeil, B., Olsen, A., Metzl, N., Kozyr, A., Fassbender, A., Manke, A., Malczyk, J., Akl, J., Alin, S. R., Bellerby, R. G. J., Borges, A., Boutin, J., Brown, P. J., Cai, W.-J., Chavez, F. P., Chen, A., Cosca, C., Feely, R. A., González-Dávila, M., Goyet, C., Hardman-Mountford, N., Heinze, C., Hoppema, M., Hunt, C. W., Hydes, D., Ishii, M., Johannessen, T., 1005 Key, R. M., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lourantou, A., Merlivat, L., Midorikawa, T., Mintrop, L., Miyazaki, C., Murata, A., Nakadate, A., Nakano, Y., Nakaoka, S., Nojiri, Y., Omar, A. M., Padin, X. A., Park, G.-H., Paterson, K., Perez, F. F., Pierrot, D., Poisson, A., Ríos, A. F., Salisbury, J., Santana-Casiano, J. M., Sarma, V. V. S. S., Schlitzer, R., Schneider, B., Schuster, U., Sieger, R., Skjelvan, I., Steinhoff, T., Suzuki, T., Takahashi, T., Tedesco, K., Telszewski, M., Thomas, H., Tilbrook, B., Vandemark, D., Veness, T., Watson, A. J., Weiss, R., Wong, C. S., and Yoshikawa-Inoue, H.: Surface Ocean CO<sub>2</sub> Atlas (SOCAT) gridded 1010 data products, *Earth System Science Data*, 5, 145–153, <https://doi.org/10.5194/essd-5-145-2013>, 2013.

Sarma, V. V. S. S., Lenton, A., Law, R. M., Metzl, N., Patra, P. K., Doney, S., Lima, I. D., Dlugokencky, E., Ramonet, M., and Valsala, V.: Sea–air CO<sub>2</sub> fluxes in the Indian Ocean between 1990 and 2009, *Biogeosciences*, 10, 7035–7052, <https://doi.org/10.5194/bg-10-7035-2013>, 2013.

Sarma, V. V. S. S., Sridevi, B., Metzl, N., Patra, P. K., Lachkar, Z., Chakraborty, K., Goyet, C., Levy, M., Mehari, M., and Chandra, N.: Air-Sea 1015 Fluxes of CO<sub>2</sub> in the Indian Ocean Between 1985 and 2018: A Synthesis Based on Observation-Based Surface CO<sub>2</sub>, Hindcast and Atmospheric Inversion Models, *Global Biogeochemical Cycles*, 37, e2023GB007694, <https://doi.org/https://doi.org/10.1029/2023GB007694>, 2023.

Skjelvan, I., Lauvset, S. K., Johannessen, T., Gundersen, K., and Skagseth, Ø.: Decadal trends in ocean acidification from the Ocean Weather Station M in the Norwegian Sea, *Journal of Marine Systems*, 234, 103775, <https://doi.org/10.1016/j.jmarsys.2022.103775>, 2022.

1020 Steinberg, D. K., Carlson, C. A., Bates, N. R., Johnson, R. J., Michaels, A. F., and Knap, A. H.: Overview of the US JGOFS Bermuda Atlantic Time-series Study (BATS): a decade-scale look at ocean biology and biogeochemistry, *Deep Sea Research Part II: Topical Studies in Oceanography*, 48, 1405–1447, 2001.

Sutton, A. J., Feely, R. A., Maenner-Jones, S., Musielwicz, S., Osborne, J., Dietrich, C., Monacci, N., Cross, J., Bott, R., Kozyr, A., et al.: 1025 Autonomous seawater pCO<sub>2</sub> and pH time series from 40 surface buoys and the emergence of anthropogenic trends, *Earth System Science Data*, 11, 421–439, 2019.

Takahashi, T., Sutherland, S. C., Wanninkhof, R., Sweeney, C., Feely, R. A., Chipman, D. W., Hales, B., Friederich, G., Chavez, F., Sabine, C., Watson, A., Bakker, D. C., Schuster, U., Metzl, N., Yoshikawa-Inoue, H., Ishii, M., Midorikawa, T., Nojiri, Y., Körtzinger, A., Steinhoff, T., Hoppema, M., Olafsson, J., Arnarson, T. S., Tilbrook, B., Johannessen, T., Olsen, A., Bellerby, R., Wong, C., Delille, B., Bates, N., and de Baar, H. J.: Climatological mean and decadal change in surface ocean pCO<sub>2</sub>, and net sea–air CO<sub>2</sub> flux over the global oceans, *Deep 1030 Sea Research Part II: Topical Studies in Oceanography*, 56, 554–577, <https://doi.org/https://doi.org/10.1016/j.dsr2.2008.12.009>, 2009.

Takahashi, T., Sutherland, S. C., Chipman, D. W., Goddard, J. G., Ho, C., Newberger, T., Sweeney, C., and Munro, D.: Climatological distributions of pH, pCO<sub>2</sub>, total CO<sub>2</sub>, alkalinity, and CaCO<sub>3</sub> saturation in the global surface ocean, and temporal changes at selected locations, *Marine Chemistry*, 164, 95–125, 2014.

- Thomsen, J., Haynert, K., Wegner, K. M., and Melzner, F.: Impact of seawater carbonate chemistry on the calcification of marine bivalves, *Biogeosciences*, 12, 4209–4220, 2015.
- ~~Torres, O., Kwiatkowski, L., Sutton, A. J., Dorey, N., and Orr, J. C.: Characterizing mean and extreme diurnal variability of ocean CO<sub>2</sub> system variables across marine environments, *Geophysical Research Letters*, 48, e2020GL090228, 2021.~~
- Uppstrom, L.: The boron/chlorinity ratio of deep-sea water from the Pacific Ocean, *Deep Sea Res.*, 21, 161–162, 1974.
- Van Heuven, S., Pierrot, D., Rae, J., Lewis, E., and Wallace, D.: CO<sub>2</sub>SY<sub>S</sub> v 1.1, MATLAB program developed for CO<sub>2</sub> system calculations, ORNL/CDIAC-105b. Oak Ridge, TN: Oak Ridge National Laboratory, [https://cdiac.ess-dive.lbl.gov/ftp/co2sys/CO2SYS\\_calc\\_MATLAB\\_v1.1/](https://cdiac.ess-dive.lbl.gov/ftp/co2sys/CO2SYS_calc_MATLAB_v1.1/), 2011.
- Woosley, R. J.: Evaluation of the temperature dependence of dissociation constants for the marine carbon system using pH and certified reference materials, *Marine Chemistry*, 229, 103 914, 2021.
- Zeebe, R. E. and Wolf-Gladrow, D.: CO<sub>2</sub> in seawater: equilibrium, kinetics, isotopes, 65, Gulf Professional Publishing, 2001.