

28 **Abstract**

29 The climatic response of mountain permafrost and glaciers located in high-elevation mountain areas has major
30 implications for the stability of mountain slopes and related geomorphological hazards, water storage and
31 supply, and preservation of paleoclimatic archives. Despite a good knowledge of physical processes that
32 govern the climatic response of mountain permafrost and glaciers, there is a lack of observational datasets from
33 summit areas. This represents a crucial gap in knowledge and a serious limit for model-based projections of
34 future behaviour of permafrost and glaciers.

35 A new observational dataset is available for the summit area of Mt. Ortles, which is the highest summit of
36 South Tyrol, Italy. This paper presents a series of air, englacial, soil surface and rock wall temperatures
37 collected between 2010 and 2016. Details are provided regarding instrument types and characteristics, field
38 methods, data quality control and assessment. The obtained data series are available through an open data
39 repository.

40 In the observed period, the mean annual air temperature at 3830 m a.s.l. was between -7.8 and -8.6°C . The
41 most shallow layers of snow and firn (down to a depth of about 10 m) froze during winter. However melt water
42 percolation restored isothermal conditions during the ablation season and the entire firn layer was found at the
43 melting pressure point. Glacier ice is cold, however only from about 30 m depth. Englacial temperature
44 decreases with depth reaching a minimum of almost -3°C close to the bedrock, at 75 m depth. A small glacier
45 located at 3470 m a.s.l., close to the summit of Mt. Ortles, was also found in cold conditions down to a depth
46 of 9.5 m. The mean annual ground surface temperature was negative for all but one monitored sites, indicating
47 cold ground conditions and the existence of permafrost in nearly all debris-mantled slopes of the summit.
48 Similarly, the mean annual rock wall temperature was negative at most monitored sites, except the lowest one
49 at 3030 m a.s.l. This suggests that the rock faces of the summit are affected by permafrost at all exposures.

50

51

52

53

54

55

56 **1. Introduction**

57 High-elevation mountain areas are complex systems influenced by physical processes occurring in the
58 atmosphere, cryosphere and lithosphere. These processes closely interact and govern the energy and mass
59 balance and climatic response of mountain permafrost and glaciers located at high elevation. Their response
60 to climatic changes has important consequences for i) the stability of mountain slopes and related
61 geomorphological hazards (Huggel et al., 2015; Knight and Harrison, 2023), ii) the thermal regime, water
62 storage and stability of mountain glaciers (Deline et al., 2015), iii) the hydrological balance and water supply
63 from glacierized catchments (Irvine-Fynn and Hubbard, 2017), and iv) the formation and preservation of
64 paleoclimatic archives, such as glacier geochemical records (Gabrielli et al., 2010).

65 The ongoing atmospheric warming is leading to a deep transformation of these high-elevation systems, which
66 react sensitively to climatic changes. Indeed, the thermal state of the cryosphere is strongly influenced by
67 variations in air temperature, which regulates its energy and mass balance and dynamic behaviour (Harris et
68 al., 2009; Cicoira et al., 2019; Deline et al., 2015). Projections of future global climate indicate further warming
69 in absence of mitigation policies such as the reduction of greenhouse gas emission (IPCC, 2022). For this
70 reason, the current impacts on high-elevation mountain areas are expected to continue and possibly accelerate.
71 Direct observations of the thermal state and response of high-elevation mountain areas are of great importance.
72 Even though the physical processes that govern the energy and mass balance and climatic response of mountain
73 permafrost and glaciers are known, model-based projections of their future behaviour are subject to large
74 uncertainty. This is because the observational datasets required for model calibration and validation are
75 particularly scarce for these summit areas, where model inputs and results are often poorly constrained and
76 extrapolated, in absence of direct observations (Charbonneau et al., 1981; Machguth et al., 2008; Carturan et
77 al., 2012, Zolles et al., 2019; Kinnard et al., 2022).

78 Thermal observations in high-elevation mountain areas are also of great value for i) improving knowledge on
79 the air temperature variability (e.g. the so-called elevation-dependent warming, Pepin et al., 2015 and 2022),
80 or the glacier cooling effect (Braithwaite et al., 2002; Carturan et al., 2015; Troxler et al., 2020; Shaw et al.,
81 2023), ii) better understanding the relationship between climatic proxies and meteorological variables (e.g. ice
82 cores, Bohleber et al., 2013), iii) evaluating/improving models (e.g. permafrost distribution models, Boekli et

83 al., 2012), iv) biological and biogeochemical studies (e.g. Rathore et al., 2018), and v) setting baseline
84 conditions for future studies and trend analyses.

85 In this paper we present a novel six-year dataset of air, rock, soil surface and englacial temperatures collected
86 between 2010 and 2016 on the summit of Mt. Ortles (46.508° N, 10.541° E, 3905 m a.s.l.), in the eastern Italian
87 Alps. These observations were carried out in the framework of the Ortles Project (ortles.org; Gabrielli et al.,
88 2016), which is an international research project, coordinated by the Byrd Polar and Climate Research Center,
89 The Ohio State University (USA) and the Hydrographic Office of the Autonomous Province of Bolzano, with
90 the aim of extracting ice cores from the Alto dell'Ortles Glacier (Oberer Ortlerferner) to be used for
91 paleoclimatic and paleoenvironmental investigations.

92 Here we provide a full description of the experimental site, data-collection methods and equipment, raw data
93 processing and final datasets.

94

95 **2. Site description**

96 Mount Ortles (46.508° N, 10.541° E) is located in the Ortles-Cevedale Mountain Group, which is the largest
97 glacierized area in the Italian Alps (Carturan et al., 2013) and it is the highest peak (3905 m a.s.l.) of South
98 Tyrol (Fig. 1). From a lithological point of view, the summit of Mt. Ortles is mainly composed by dolomites,
99 alternated with dark-stratified limestones and paraconglomeratic limestone levels and breccias. Local outcrops
100 of phyllites rich in quartz and orthogneiss can be found at the base of the mountain (Montrasio et al., 2012).

101 The northern part of the Ortles-Cevedale mountain range is characterised by a continental climate, with scarce
102 annual precipitation (500 mm in the lower valley), which falls mostly in summer. Towards the south, there is
103 an increasing Mediterranean influence and the annual precipitation maxima are in spring and autumn, with
104 cumulative amounts of 900 mm in the lower valleys. In the glaciated areas in the middle of the mountain range,
105 at 3000-3200 m a.s.l., the mean annual precipitation has been estimated between 1400 and 1500 mm (Carturan,
106 2010; Carturan et al., 2012). Using mass balance observations in the period from 2009 to 2016 it is possible to
107 estimate 1300-1400 mm of annual precipitation on top of Mt. Ortles. However, the snow accumulation and
108 glacier mass balance are highly affected by wind and estimating the actual precipitation is almost impossible
109 at this site. In comparison, the closest weather station (Solda, 1905 m a.s.l.) averaged an annual precipitation
110 of 1090 mm between 1989 and 2019.


111 The mean annual isotherm of 0°C is located at about 2500 m a.s.l. At the elevation of the glaciers (above 3000
112 m) the snow cover shows a typical annual cycle, so the accumulation season lasts between October and May,
113 and the ablation season between June and September. On the glaciers, however, snowfalls are frequent during
114 summer, especially above 3300-3500 m. Liquid precipitation is rare on top of Mt. Ortles, but some rain events
115 have been observed in the last 15 years.

116 Glaciers, glacierets and snowfields cover the Mt. Ortles flanks. Here we describe the Alto dell'Ortles and
117 Hintergrat glaciers, which are the two ice bodies investigated in the Ortles Project. The summit area is almost
118 entirely covered by the Alto dell'Ortles Glacier (Oberer Ortlerferner), which is the highest glacier of South
119 Tyrol, ranging in altitude between 3018 and 3905 m a.s.l. and covering an area of 1.04 km² (year 2008). The
120 observed glacier thickness is about 75 m (Gabrielli et al., 2012) and the vertical ice profile encompasses the
121 last ~7 kyr (Gabrielli et al., 2016). This glacier is polythermal, with temperate firn and cold ice underneath
122 (Gabrielli et al., 2012). From geomorphological evidence (trimlines and moraines) it is possible to estimate a
123 maximum Little Ice Age (14th - 19th centuries) area of 2.09 km² for this glacier, and a 50% area loss since then.
124 Between 1984 and 2005 the (geodetic) mass balance of the glacier was closer to equilibrium (-0.18 m w.e. y⁻¹
125 ¹) when compared to the majority of glaciers in the Ortles-Cevedale Group in the same period (mean balance
126 rate of -0.69 m w.e. y⁻¹, Carturan et al., 2013). A small glacier, named Hintergrat, covers part of the eastern
127 rocky ridge of Mt. Ortles. The area of this glacier is 0.09 km² and its elevation ranges between 3340 and 3580
128 m a.s.l. This glacier is mostly in cold thermal conditions and its front hangs over the Fine del Mondo Glacier
129 (Ende der Welt Ferner) underneath.

130 Mountain permafrost is widespread on Mt. Ortles, according to the permafrost distribution modelled by
131 Boeckli et al. (2012), that indicates 'permafrost in nearly all conditions' above 2600 m for areas with northern
132 exposure, and above 2900 m for areas with southern exposure.

133 Before the Ortles Project, no specific investigation existed on the air, englacial and permafrost temperatures
134 of this mountain.

135



136

137 Figure 1. Geographic location of Mt. Ortles and of sites instrumented for air, rock, englacial and soil surface
 138 temperature measurements. Close ups of the a) Payer Refuge, b) Lombardi Bivouac, c) Mt. Ortles summit
 139 area, and d) Hintergrat ridge are reported in the panes on the right. The background hillshaded DEM (2017
 140 LiDAR survey) is from <http://geocatalogo.retecivica.bz.it/>.

141


142 3. Data description

143 The temperature datasets presented in this work were obtained by installing stand-alone dataloggers connected
 144 to one or several temperature sensors. Due to the remoteness of the study site, dataloggers were powered by
 145 lithium or lead-acid batteries, which in the case of the Automatic Weather Station (AWS, Section 3.1) were
 146 recharged daily by solar panels. Periodic field visits, mostly performed from June to September, were used for
 147 instrumentation maintenance and data download using a laptop. No real-time transmission of data was setup.
 148 The dataset is characterised by a good time coverage and a few gaps (Fig. 2), indicating the suitability of the
 149 selection of the equipment and field procedures for installation and maintenance. The most significant temporal
 150 gaps affect soil surface temperature datasets due to the impossibility of accessing dataloggers in late summer
 151 of 2011. Other minor gaps were caused by temporary malfunctions or by damaged equipment, e.g., the rupture

152 of the fan-aspirated radiation shield at the AWS between February and August 2013, which forced us to treat
 153 this period as a data gap. We did not undertake gap-filling, in order to keep the data recorded in the field as
 154 unchanged as possible.

155 Details of measuring equipment and installations are provided in the following sections. Further details about
 156 the instruments are provided in Table 1 and a topographic description of instrumented sites is reported in Table
 157 E1, in the Appendix. Figures showing examples of data series for each variable are provided in the following
 158 subsections.

159



160

161 Figure 2. Monthly coverage of temperature measurements available from 2010 to 2016 on Mt. Ortles. Partial
 162 coverage indicates the occurrence of data gaps for specific months.

163

164

165 Table 1. Sensor characteristics, setup and period of operation for air, englacial, soil surface and rock wall
 166 temperature measurements on Mt. Ortles.

167

Measured variable	Sensor	Radiation shield	Period of operation		Initial height (m)	Unit	Interval	Integration method and interval	Accuracy
			from	to					
Air temperature data (AWS)									

Air Temperature	Vaisala HMP155A	R. M. Young 43502 fan-aspirated radiation shield	Sep 2011	Jun 2015	+4	°C	15 min	avg 1 h	$\pm(0.226 - 0.0028 \cdot T)$ °C from -80 to 20°C, $\pm(0.055 + 0.0057 \cdot T)$ °C from 20 to 60 °C
Air Temperature	Vaisala HMP155A	Campbell Scientific MET 21 radiation shield with natural ventilation	Sep 2012	Jun 2015	+4	°C	15 min	avg 1 h	$\pm(0.226 - 0.0028 \cdot T)$ °C from -80 to 20°C, $\pm(0.055 + 0.0057 \cdot T)$ °C from 20 to 60 °C
Air Temperature (Backup and comparison)	Gemini TGP-4020	R. M. Young 43502 fan-aspirated radiation shield	Sep 2011	Jun 2015	+4	°C	1 h	avg 1 h	$\pm(0.2 - 0.005 \cdot T)$ °C from -40 to 0°C, ± 0.2 °C from 0°C to 40°C
Air Temperature (Backup and comparison)	Gemini TGP-4020	R.M. Young 41303-5 radiation shield with natural ventilation	Sep 2011	Jun 2015	+4	°C	1 h	avg 1 h	$\pm(0.2 - 0.005 \cdot T)$ °C from -40 to 0°C, ± 0.2 °C from 0°C to 40°C
Air Temperature (Backup and comparison)	Onset Hobo H8 Pro Temp	Davis 7714 radiation shield with natural ventilation	Sep 2012	Sep 2016	+4	°C	1 h	avg 1 h	$\pm(0.63 - 0.022 \cdot T)$ °C from -40 to 0°C, ± 0.63 °C from 0°C to 40°C

Englacial temperature data

Snow and firn temperature at the AWS site	Geoprecision thermistor string (15 sensors)		Jun 2012	Sep 2016	-0.6 / -1.6 / -2.6 / -3.6 / -4.6 / -5.6 / -6.6 / -7.6 / -8.6 / -9.6 / -10.6 / -11.6 / -12.6 / -13.6 / -14.6	°C	2 h	instant 2 h	± 0.5 °C from -10°C to +85°C
Englacial temperature at the borehole drilling site (short string)	Sommer thermistor string (11 sensors)		Nov 2011	Aug 2015	0 / -1 / -2 / -3 / -4 / -6 / -8 / -10 / -15 / -20 / -25	°C	1 h	instant 1h	± 0.1 °C from 0°C to 70°C
Englacial temperature at the borehole drilling site (long string)	Sommer thermistor string (4 sensors)		Nov 2011	Apr 2013	-15 / -35 / -55 / -75	°C	1 h	instant 1h	± 0.1 °C from 0°C to 70°C
Englacial temperature at the Hintergrat Glacier	Sommer thermistor string (5 sensors)		Nov 2011	Aug 2016	-1.5 / -3.5 / -5.5 / -7.5 / -9.5	°C	1 h	instant 1h	± 0.1 °C from 0°C to 70°C

Soil surface temperature data

Soil surface temperature at Lombardi bivouac - ORTL_05	Gemini TGP-4020		Sep 2010	Sep 2016	-0.15	°C	1 h	avg 1h	$\pm(0.2 - 0.005 \cdot T)$ °C from -40 to 0°C, ± 0.2 °C from 0°C to 40°C
Soil surface temperature at Lombardi bivouac - ORTL_06	Gemini TGP-4020		Sep 2010	Sep 2016	-0.12	°C	1 h	avg 1h	$\pm(0.2 - 0.005 \cdot T)$ °C from -40 to 0°C, ± 0.2 °C from 0°C to 40°C
Soil surface temperature at Payer refuge - ORTL_07	Gemini TGP-4020		Sep 2010	Aug 2016	-0.05	°C	1 h	avg 1h	$\pm(0.2 - 0.005 \cdot T)$ °C from -40 to 0°C, ± 0.2 °C from 0°C to 40°C
Soil surface temperature at Payer refuge - ORTL_08	Gemini TGP-4020		Sep 2010	Aug 2016	-0.05	°C	1 h	avg 1h	$\pm(0.2 - 0.005 \cdot T)$ °C from -40 to 0°C, ± 0.2 °C from 0°C to 40°C
Soil surface temperature at Hintergrat ridge - ORTL_10	Gemini TGP-4020		Sep 2010	Aug 2016	-0.05	°C	1 h	avg 1h	$\pm(0.2 - 0.005 \cdot T)$ °C from -40 to 0°C, ± 0.2 °C from 0°C to 40°C
Soil surface temperature at Hintergrat ridge - ORTL_11	Gemini TGP-4020		Sep 2010	Aug 2013	-0.05	°C	1 h	avg 1h	$\pm(0.2 - 0.005 \cdot T)$ °C from -40 to 0°C, ± 0.2 °C from 0°C to 40°C

Rock wall temperature data

Rock wall temperature at Mt. Ortles summit - CIMA_ALTO	Geoprecision thermistor string (3 sensors)	Sep 2011	Nov 2013	-0.10 / -0.30 / -0.55	°C	1 h	instant 1h	±0.5 °C from -30 to -5 °C, ±0.1 °C from -5 to +40 °C
Rock wall temperature Mt. Ortles summit - CIMA_VERTICALE	Geoprecision thermistor string (3 sensors)	Sep 2011	Sep 2016	-0.10 / -0.30 / -0.55	°C	1 h	instant 1h	±0.5 °C from -30 to -5 °C, ±0.1 °C from -5 to +40 °C
Rock wall temperature at Vorgipfel - ANTICIMA_SUD	Geoprecision thermistor string (3 sensors)	Sep 2011	Sep 2016	-0.10 / -0.30 / -0.55	°C	1 h	instant 1h	±0.5 °C from -30 to -5 °C, ±0.1 °C from -5 to +40 °C
Rock wall temperature at Lombardi bivouac - BIV_LOMBARDI	Geoprecision thermistor string (3 sensors)	Sep 2011	Sep 2016	-0.10 / -0.30 / -0.55	°C	1 h	instant 1h	±0.5 °C from -30 to -5 °C, ±0.1 °C from -5 to +40 °C
Rock wall temperature at Hintergrat - HINTERGRAT	Geoprecision thermistor string (3 sensors)	Oct 2011	Aug 2014	-0.10 / -0.30 / -0.55	°C	1 h	instant 1h	±0.5 °C from -30 to -5 °C, ±0.1 °C from -5 to +40 °C
Rock wall temperature at Payer refuge - PAYER	Geoprecision thermistor string (3 sensors)	Nov 2011	Aug 2016	-0.10 / -0.30 / -0.55	°C	1 h	instant 1h	±0.5 °C from -30 to -5 °C, ±0.1 °C from -5 to +40 °C

168

169

170

171 **3.1 Air temperature data**

172 On 30 September 2011 an Automatic Weather Station (AWS, Fig. 3) was installed on the upper accumulation
173 area of Alto dell'Ortles Glacier, at an elevation of 3830 m a.s.l. The AWS was equipped with a Campbell
174 Scientific CR-1000 datalogger, solar panels and sensors for air temperature and relative humidity (Vaisala
175 HMP155A), wind speed and direction (R. M. Young 05103), incoming and outgoing shortwave and longwave
176 radiation (Delta Ohm LP Pyra 05 and LP PIRG 01), and snow depth (Campbell Scientific SR50A). The
177 equipment was supported by an aluminum tower (composed of 2-m modules), anchored in the firn at 2 m depth
178 and supported by wooden boards. After the installation, the tower extended 4 m from the surface. The sensors
179 and solar panels were fixed on top of the tower, whereas the datalogger/battery housing box was fixed at the
180 bottom (Figs. F1 and F2).

181



182

183 Figure 3. The automatic weather station installed in the upper accumulation area of Alto dell'Ortles Glacier.

184 The Mt. Ortles summit (3905 m a.s.l.) is visible in the background.


185

186 The Vaisala HMP155A sensor was installed inside a R. M. Young 43502 fan-aspirated radiation shield. Two
187 standalone Gemini TGP-4020 dataloggers equipped with PB-5003-1 thermistor probes were also installed for
188 comparison and backup purposes, one in the same fan-aspirated radiation shield of the Vaisala HMP155A
189 sensor and one inside a 6-plate R.M. Young 41303-5 radiation shield with natural ventilation. In September
190 2012, we installed an additional HMP155A temperature sensor inside a 15-plate Campbell Scientific MET 21
191 radiation shield with natural ventilation, and an Onset Hobo H8 Pro Temp datalogger housed in an 8-plate
192 Davis 7714 radiation shield with natural ventilation (Figs. F3 and F4). Sensor specifications are reported in
193 Table 1. The Vaisala HMP155A data were recorded as 15-minute mean values, whereas the Gemini TGP-4020
194 dataloggers recorded hourly minimum and maximum temperature, and the Hobo H8 Pro Temp datalogger
195 recorded hourly instantaneous temperature. All the temperature records have been converted into hourly

196 averages, averaging 15-minute means for the Vaisala HMP155A sensors, minimum and maximum hourly
197 temperature for the Gemini TGP-4020 dataloggers, and instantaneous temperature at the beginning and at the
198 end of each hour for the Hobo H8 Pro Temp datalogger, assuming a linear variation of temperature during
199 each hour.

200 All the air temperature sensors were installed at the same level (± 20 cm). The height of the air temperature
201 sensors above the glacier surface changed with the snow accumulation over time (mean height = 241 cm, 5th
202 percentile = 61 cm, 95th percentile = 407 cm). To prevent burial by snow accumulation, the tower was elongated
203 annually by adding a 2-m module. Figure 4 shows the height of the sensors above the glacier surface, as
204 reconstructed from the snow depth data and maintenance logs (Table D1).

205



206

207 Figure 4. Air temperature sensors height above the snow surface.

208

209


210 Despite the remote and harsh environment, the AWS worked properly without major interruptions. In June
211 2015 it was removed, but the Hobo H8 Pro Temp logger was left on site and recorded additional 15 months of

212 data. The main issue linked to the specific environment of installation was ice and snow accretion combined
213 with strong winds, which damaged the fan-aspirated radiations shield in February 2013. The obtained data are
214 shown in Fig. 5.

215

216

Air temperature (AWS) [°C]



217

218 Figure 5. Daily mean air temperature series measured by different sensors installed at the Automatic Weather
219 Station on Mt. Ortles.

220

221 3.2 Englacial temperature data

222 Englacial temperature measurements were collected at three different sites on Mt. Ortles: i) at the AWS site
223 (3830 m), ii) at the borehole drilling site (3859 m), and iii) at the Hintergrat glacier (3476 m). The obtained
224 englacial temperature data are shown in Fig. 6.

225

226 3.2.1 Snow and firn temperature data at the AWS site

227 On the 18th of June 2012 a 20-meter thermistor string manufactured by Geoprecision GmbH (Germany) was
228 installed 10 m east of the AWS (Fig. F1). The thermistor string was composed of a Dallas M-Log5W
229 datalogger, powered by a 3.6 V lithium battery, and connected to 15 digital Dallas temperature sensors spaced
230 one meter from each other. The string was lowered into a 14.6 m hole drilled using a steam ice drill. The initial
231 depth of temperature sensors ranged between 0.6 and 14.6 m, and increased afterwards up to about 6 m due to
232 the accumulation of snow. The logger was housed inside a plastic box on the glacier surface, subsequently
233 buried in the snow. Instantaneous temperature data were recorded with a 2-hour frequency.
234 The data were retrieved by means of a laptop using a USB dongle connected wireless (radio transmission) to
235 the logger, below the glacier surface. We were able to retrieve temperature data with the logger buried below
236 a maximum of ~6 m of snow and firn. The thermistor string worked properly without interruptions and without
237 requiring maintenance or battery replacement. Sensor specifications are reported in Table 1.

238

239 3.2.2 Englacial temperature at the borehole drilling site

240 The site where the Ortles Project deep ice cores were extracted is a small col (3859 m; 10°32'34", 46°30'25")
241 between the summit of Mt. Ortles and the Anticima Sud/Vorgipfel (3845 m, Figs. 1 and G1). The ice is 75 m
242 thick at this site as indicated by geophysical sensing prospecting and confirmed by ice core drilling operations
243 (Gabielli et al., 2016). Two thermistor strings were installed in borehole number 3 on the 5th of October 2011,
244 immediately after the completion of the drilling operations (Fig. G3a). The strings were composed of an MDL
245 8/3 datalogger, manufactured by Sommer GmbH & Co KG (Austria). The logger was connected to 44031
246 thermistors, manufactured by ThermX (USA).

247 A first (short) thermistor string was 35 m long, and was equipped with temperature sensors at 0, 1, 2, 3, 4, 6,
248 8, 10, 15, 20, 25 m (initial depth). The other (long) string was 100 m, with temperature sensors at 15, 35, 55
249 and 75 m (initial depth). Burial depth of sensor increased over time due to net snow and firn accumulation.

250 Dataloggers and exceeding portions of strings were housed inside a metal box and arranged on a winding
251 system (Fig. G3), making it possible to extend the thermistor strings and to arise the box at the glacier surface
252 periodically. Field maintenance was also required to replace batteries and download the stored data.
253 Instantaneous temperature data were recorded with a 1-hour frequency.

254 The short string worked properly until removal in summer 2015, with the exception of a two-month gap in
255 summer 2013. The long string stopped working in April 2013, possibly due to ice dynamics and deformation
256 of the borehole at a depth below 25-30 m. Sensor specifications are reported in Table 1.

257


258 3.2.3 Englacial temperature at the Hintergrat Glacier

259 On the 14th of October 2011 a thermistor string was installed at 3476 m a.s.l. on top of the Hintergrat Glacier
260 (Fig. G2). The string was manufactured by Sommer GmbH & Co KG (Austria), with the same components as
261 those installed at the ice core drilling site (Section 3.2.2, Fig. G4). It was lowered into a hole drilled using a
262 steam ice drill down to a depth of 9.6 m. We did not reach the glacier bottom at this site. The temperature
263 sensors were placed at 1.5, 3.5, 5.5, 7.5 and 9.5 m below the glacier surface. This lower site is subject to net
264 ablation, therefore in this case the initial depth decreased through time and the sensor at 1.5 m depth came to
265 the surface in summer 2013. After 2013, this sensor's data were consequently discarded.

266

267

268



269

270 Figure 6. Englacial temperature measured at various sites and various depths on the Alto dell'Ortles Glacier.

271

272


273 3.3 Soil surface temperature data

274 The thermal regime of the soil surface at six deglaciated sites on Mt. Ortles was monitored using standalone

275 temperature dataloggers over the period between September 2010 and September 2016. We used Gemini TGP-

276 4020 dataloggers, powered by 3.6 V lithium batteries, and equipped with PB-5001 probes, which were placed

277 5-15 cm below the soil surface (Figs. H1 and H2). Mean temperature data were recorded at hourly intervals.
278 Periodic maintenance was required to download the data and replace exhausted batteries.
279 The monitored sites range in elevation between 2899 and 3466 m a.s.l. The dataloggers were placed in pairs
280 at three main locations (Figs. 1 and H3): refuge Payer (ORTL_07 and ORTL_08), bivouac Lombardi
281 (ORTL_05 and ORTL_06) and Hintergrat ridge (ORTL_10 and ORTL_11).
282 The data series extend from late summer 2010 to late summer 2016, with a gap between autumn 2011 and late
283 summer 2012 and for ORTL_05 and ORTL_06 also between July and August 2015, due to the impossibility
284 of accessing the dataloggers for maintenance. ORTL_11 was buried under snow and firn after 2013 and has
285 never been recovered. The obtained soil surface temperature data are displayed in Fig. 7.
286



287
288 Figure 7. Soil surface temperature data measured at various locations on Mt. Ortles.
289

290

291

292 **3.4 Rock wall temperature data**

293 The sub-surface temperature of rock walls located at six sites on Mt. Ortles was monitored starting in late
294 summer and autumn 2011 (Fig. 2). Very steep/almost vertical rock walls with different exposures and
295 elevations were selected for monitoring (Figs. 1 and I3, Table E1). Two sites were established next to the Mt.
296 Ortles summit (3900 and 3880 m, facing East), one at the Vorgipfel (3844 m, facing South), one at the
297 Hintergrat (3370 m, facing North-East), one at the bivouac Lombardi (3351, facing West) and one at the refuge
298 Payer (3030 m, facing East).


299 Rock temperature data were acquired using Geoprecision Dallas M-Log5W dataloggers, powered by a 3.6 V
300 lithium battery, and connected to three digital Dallas temperature sensors installed at 0.1, 0.3 and 0.55 m depth,
301 into holes drilled with a hammer drill (Fig. I1). Instantaneous temperature data were stored at hourly intervals
302 and downloaded with a remote connection using a wireless USB dongle and a laptop (Fig. I2).

303 The datalogger placed at the Hintergrat was damaged by hikers in late July 2012 but was repaired in late
304 August 2012. It remained operational until August 2014 when it was removed due to another badly damage.

305 One of the loggers installed at the Mt. Ortles summit ("CIMA_ALTO") stopped working in November 2013
306 due to battery failure while the other dataloggers worked properly until the end of the monitoring period, in
307 late summer 2016. Sensor specifications are reported in Table 1. The obtained rock wall temperature data are
308 displayed in Fig. 8.

309

Rock wall temperature [°C]



310

311 Figure 8. Rock wall temperature measurements at various sites and three depths on Mt. Ortles.

312

313 4. Data quality control and assessment

314 The temperature datasets presented in this work were carefully inspected to detect possible problems affecting
315 raw measurements and to ensure the highest possible accuracy (Table 1). Data quality controls allowed
316 assigning a quality flag to each temperature record, as described in Table B1.

317 Air temperature sensors were exposed to harsh conditions, without protection from snow, rock or firn/ice as in
318 the case of sensors used to measure the englacial, soil and rock wall temperature. For this reason, the air
319 temperature sensors were subjected to possible damage by strong winds and lightning, ice and snow accretion,
320 and burial by snow in case of abundant snowfall. In addition, they were subjected to the typical issues affecting
321 air temperature measurements, arising from low wind speed and high solar radiation, worsened by high surface
322 albedo, which generally lead to errors due to heating during daytime (WMO, 2021). Sensor drifting should
323 also be taken into account as a potential problem.

324 In order to spot problematic periods we carried out a sensor-by-sensor intercomparison, calculating hourly and
325 daily temperature differences among pairs of sensors, including data from two neighbouring weather stations
326 (Madriccio, 2825 m a.s.l., and Cima Beltovo, 3328 m a.s.l., Weather Service - Autonomous Province of
327 Bozen/Bolzano) for additional confirmation. We compared temperature difference series with maintenance
328 logs, to understand the sources of malfunctions and anomalies, and to assign data quality flags to air
329 temperature series. Data recorded during malfunctions were handled as data gaps and removed from the
330 published series. Anomalies in periods of heavy snowfalls, which caused snow/ice accretion and a rapid
331 decrease of sensor height, are flagged with a specific code (Table B1).

332 Englacial and rock wall temperature were checked in the same way by calculating hourly and daily differences
333 among sensors located at the same site, and checking irregularities (i.e. sudden jumps in temperature
334 differences) in combination with field observations during maintenances. We have detected no malfunctions,
335 but it is possible that after maintenance operations (detailed in Table D1 to D4) a short period of a few hours
336 or a few days was required to reach a new equilibrium at englacial temperature monitoring sites. We have
337 highlighted these maintenance operations with potential impacts on measured temperature using a data quality
338 code, reported in Table B1.

339 Soil surface temperature data displayed no obvious anomalies and were checked in the 'zero-curtain' phase,
340 that is the 0°C plateau during the snowmelt phase. Only ORTL_07 required a correction of measured
341 temperature in 2014 (offset applied = -0.35°C) and 2015 (offset applied = -1.1°C), to correct discrepancies
342 larger than sensor accuracies reported in Table 1. We have highlighted these adjustments with a quality flag in
343 the corresponding data files (Table B1).

344

345 **5. Data availability**

346 The datasets from this study are publicly available at <https://doi.org/10.5281/zenodo.7879969> (Carturan et al.,
347 2023). The data files are stored in Microsoft Excel .xlsx format. Detailed information on the file content and
348 structure are reported in the Appendix of this manuscript (Tables A1, B1 and C1).

349

350

351

352 **6. Summary of observations and research outlook**

353 The datasets collected on Mt. Ortle enable description of its thermal state within a time window of six years
 354 (2011-2016). This period is long enough to provide a picture of modern average conditions and interannual
 355 variability, and it is useful as a baseline for possible future studies aimed at detecting changes and trends in
 356 monitored variables.

357 In the period from 02-10-2011 to 01-09-2016, the mean daily air temperature ranged between -30.1°C (12-02-
 358 2012) and 6.1°C (03-08-2013), averaging -8.3°C. These statistics have been extracted from a merged time
 359 series, which combines the sensors that have the longest time coverage (Fig. 2), i.e. the GEMINI TGP-4020
 360 with natural ventilation before 15-06-2015 and the ONSET Hobo H8 (natural ventilation) from 15-06-2015.
 361 The air temperature reached hourly extremes of -33.3°C (09-02-2012 at 24:00 UTC) and 10.1°C (20-08-2012
 362 at 10:00 UTC). These extremes must be viewed with caution due to the high sensitivity of short-term
 363 temperature fluctuations to possible errors, mainly due to low natural ventilation. The aspirated shield installed
 364 at the weather station proved subject to damage and malfunction. However, it was operational at the time when
 365 the two extreme values were recorded, providing identical minimum temperature of -33.3°C, and a maximum
 366 of 8.0°C at 13:00 of 20-08-2012. In the common period of operation (overlaps are shown in Fig. 5), the average
 367 difference in mean daily air temperature among pairs of installed sensors did not exceed 0.60°C in absolute
 368 value (Table 2).

369

370 Table 2. Average difference in mean daily air temperature among pairs of installed sensors (f.a. = fan-aspirated;
 371 n.v. = natural ventilation). Column headings represent the first term of the difference calculation.

372

	Vaisala HMP155A (f.a.)	Vaisala HMP155A (n.v.)	Gemini TGP- 4020 (f.a.)	Gemini TGP- 4020 (n.v.)	Onset Hobo H8 Pro Temp (n.v.)
Vaisala HMP155A (f.a.)		-0.28	0.11	0.18	-0.02
Vaisala HMP155A (n.v.)			0.41	0.60	0.32
Gemini TGP-4020 (f.a.)				0.08	-0.15

Gemini TGP-4020
(n.v.)

-0.28

373

374

375 The mean daily air temperature was above 0°C for 39 days in 2012, 48 in 2013, 15 in 2014, 44 in 2015, and
376 31 in 2016. These results highlight a significant interannual variability in the length of the melt season at this
377 high-elevation site. The mean annual air temperature averaged -8.3°C, ranging between -8.6°C in 2012 and
378 2013, and -7.8°C in 2016 (Table 3).

379

380 Table 3. Mean annual air temperature recorded at the automatic weather station on Mt. Ortles, with five
381 different sensors (f.a. = fan-aspirated; n.v. = natural ventilation). The mean temperature is reported only for
382 years with less than one month missing of data, and is calculated between the 1st of September and the 31st of
383 August.

384

Sensor	Vaisala HMP155A (f.a.)	Vaisala HMP155A (n.v.)	Gemini TGP- 4020 (f.a.)	Gemini TGP- 4020 (n.v.)	Onset Hobo H8 Pro Temp (n.v.)	Merged (n.v.)
Year						
2012	-8.9		-8.7	-8.6		-8.6
2013				-8.6	-9.0	-8.6
2014	-8.4		-8.2	-8.1	-8.4	-8.1
2015					-8.6	-8.3
2016					-8.1	-7.8

385

386 Englacial temperature measurements reveal warm firn and isothermal summer conditions down to a depth of
387 25 m on the upper part of the Alto dell'Ortles Glacier. The summer was cold (and snowy) enough only in 2014
388 to preserve below-zero temperature in firn and snow down to a depth of about 15 m at the AWS and 10 m at
389 the drilling site. On the other hand deep borehole data available until 10/04/2013 confirm that glacier ice below
390 the firn-ice transition is cold throughout the year, as detected during ice coring operations at the drilling site in
391 October 2011 (Gabielli et al. 2012). The ice temperature decreased with depth reaching a minimum of -3°C
392 at the glacier bed, at 75 m below the surface, and does not change significantly throughout the year.

393 The Hintergrat glacier is also composed of cold ice, which is subject to net surface ablation at the string
 394 installation site. Indeed, the sensor at 1.5 m initial depth was exposed at the surface in August 2013. A 1.2 m
 395 layer of firn formed in 2014, but underwent complete ablation by the end of the following summer.

396 Soil surface temperature measurements, and in particular the mean annual ground surface temperature
 397 (MAGST, Table 4), suggest the existence of permafrost on most of the monitored sites (Guglielmin et al.,
 398 2003; Ballantyne, 2018), with the exception of the ORTL07 site which is at 2994 m a.s.l, close to the Payer
 399 refuge (Fig. 1). The results of ORTL_10 and ORTL_11 can be compared to analogous observations
 400 (unpublished) carried out in the same period on Mt Vioz (3520 m a.s.l., 14 km south of Mt. Ortles), using the
 401 same devices and field techniques at two sites with similar elevation and exposure. On Mt. Vioz the MAGST
 402 was -2.1°C for the site with southern exposure and -2.9°C for the site with eastern exposure, indicating slightly
 403 colder soil surface thermal conditions.

404

405 Table 4. Mean annual ground surface temperature (MAGST) recorded at six different sites on Mt. Ortles. Site
 406 locations are reported in Fig. 1. MAGST is reported only for years with less than one month of missing data,
 407 and is calculated between the 1st of September and the 31st of August.

408

Sensor	ORTL_05	ORTL_06	ORTL_07	ORTL_08	ORTL_10	ORTL_11
Year						
2011	-2.6	-2.4	1.3	-1.2	-1.2	-2.4
2012						
2013	-3.5	-2.7	1.1	-1.0	-0.7	-2.1
2014	-3.2	-2.1	0.9	-0.5	-1.5	
2015			1.9	-0.3	-1.0	
2016	-3.4	-1.6	0.7	-1.6	-1.9	

409

410

411

412 Rock wall temperature provided results that are in line with soil surface temperature measurements. The
 413 warmest site was close to the Payer refuge, with mean annual rock surface temperature (MARST) above the
 414 freezing level (Table 5). All the other monitored rock walls displayed below-freezing MARST and similar
 415 behaviour, with the exception of CIMA_ALTO, close to the Mt. Ortles summit, where rock temperature
 416 fluctuations appear to be dampened by snow accumulation between September and May (Fig. 8).

417 The collected data are being analysed for the interpretation of the ice core drilled in the framework of the Ortles
 418 Project. In particular, air and englacial temperature data are used for developing and validating a model that
 419 aims at reproducing the formation of the isotopic record in snow and firn.

420 Together with rock wall and soil surface temperature, these datasets represent unique observations at such
 421 elevation in the eastern European Alps, and may contribute to the study and understanding of specific aspects
 422 of the climatic sensitivity of the alpine cryosphere. For example, they can be used for the development of
 423 permafrost distribution and degradation models, air temperature simulations over glacierized areas (including
 424 the so-called glacier cooling effect), snow and glacier mass balance models, glacio-hydrological forecasting
 425 systems, or dynamic glacier models that take into account the thermal state of glaciers and its variability.

426

427 Table 5. Mean annual rock surface temperature (MARST) recorded at six different site on Mt. Ortles. Site
 428 locations are reported in Fig. 1. MARST is reported only for years with less than one month of missing data,
 429 and is calculated between the 1st of September and the 31st of August.

430

LOCATION	CIMA_ALTO			CIMA_VERTICALE			ANTICIMA_SUD			BIVACCO_LOMBARDI			HINTERGRAT			PAYER			
	0.10	0.30	0.55	0.10	0.30	0.55	0.10	0.30	0.55	0.10	0.30	0.55	0.10	0.30	0.55	0.10	0.30	0.55	
Depth (m)																			
Year																			
2012	-2.1	-2.3	-2.6	-2.6	-2.5	-2.5	-1.8	-2.2	-2.7	-2.4	-2.3	-2.4							
2013	-2.0	-2.3	-2.5	-3.1	-3.1	-3.0	-3.1	-3.5	-3.9	-2.8	-2.7	-2.7	-2.2	-2.1	-2.2	1.4	1.2	0.9	
2014				-2.9	-2.8	-2.8	-2.5	-2.8	-3.3	-2.8	-2.7	-2.7				1.8	1.5	1.2	
2015				-2.1	-2.1	-2.1	-2.2	-2.6	-3.1	-1.5	-1.4	-1.6				2.7	2.4	2.1	
2016				-2.4	-2.3	-2.3	-2.2	-2.5	-3.0	-3.2	-3.1	-3.1				1.9	1.6	1.3	

431

432

433

434

435

436

APPENDICES

437

438 **APPENDIX A: Variables in data files**

439 Table A1. Column names for variables reported in data files.

Variable	Column name
Air temperature (fan-aspirated Vaisala HMP155A)	Air_T_HMP_Asp
Air temperature (natural-ventilation Vaisala HMP155A)	Air_T_HMP_Nat
Air temperature (fan-aspirated Gemini TGP-4020)	Air_T_TGP_Asp
Air temperature (natural-ventilation Gemini TGP-4020)	Air_T_TGP_Nat
Air temperature (natural-ventilation Onset Hobo H8 Pro Temp)	Air_T_H8_Nat
Englacial temperature at the AWS site	AWS_En (depth m)
Englacial temperature at the borehole drilling site (short string)	BH_En_SS (depth m)
Englacial temperature at the borehole drilling site (long string)	BH_En_LS (depth m)
Englacial temperature at the Hintergrat Glacier	HG_En (depth m)
Soil surface temperature at bivouac Lombardi - ORTL_05	GST_ORTL05
Soil surface temperature at bivouac Lombardi - ORTL_06	GST_ORTL06
Soil surface temperature at refuge Payer - ORTL_07	GST_ORTL07
Soil surface temperature at refuge Payer - ORTL_08	GST_ORTL08
Soil surface temperature at Hintergrat ridge - ORTL_10	GST_ORTL10
Soil surface temperature at Hintergrat ridge - ORTL_11	GST_ORTL11
Rock wall temperature at Mt. Ortles summit - CIMA_ALTO	Rw_ALTO (depth m)
Rock wall temperature Mt. Ortles summit - CIMA_VERTICALE	Rw_VERTICALE (depth m)
Rock wall temperature at Vorgipfel - ANTICIMA_SUD	Rw_ANTICIMA (depth m)
Rock wall temperature at bivouac Lombardi - BIV_LOMBARDI	Rw_LOMBARDI (depth m)
Rock wall temperature at Hintergrat - HINTERGRAT	Rw_HINTERGRAT (depth m)
Rock wall temperature at refuge Payer - PAYER	Rw_PAYER (depth m)

440

441

442 **APPENDIX B: Quality flags for data files**

443 Table B1. Quality code flags reported in data files, their meaning and explanations.

Quality code flag (“_FI” inflection in column names)	Meaning	Explanation
1	Good data	No issues detected during quality checks
0	No data	Data missing or removed (malfunctioning, physically implausible, sensor/device damaged, sensor underneath snow)
2	Maintenance	Data are affected by field maintenance of instrumentation
3	Ice/snow accretion	The air temperature data are affected by ice or snow accretion
4	Small height of the sensor	The air temperature sensor is less than 1 m above the snow surface
5 (offset)	Sensor offset	Offset applied to correct soil surface temperature data, based on the zero-curtain phase during snow melt (offset value in brackets)

444

445

446

447 **APPENDIX C: Data files structure**

448 Table C1. Structure of data files. For sensors at different depth below the surface, the depth in m is reported
 449 after the variable name, in brackets.

Date and hour (UTC)	Variable name (depth m)	Quality flag code
DD/MM/YYYY HH:MM	value	code

454

455

456 **APPENDIX D: Maintenance logs**

457 Table D1. Field operations and maintenance for the air temperature sensors mounted at the AWS on Mt. Orles.

Date	Field operations
01/10/2011	AWS setup and datalogger launch
18/06/2012	Datalogger download, check of sensor status and functioning
07/09/2012	Datalogger download, check of sensor status and functioning, 2 m increase in height of support tower, installation of two additional sensors (Vaisala HMP155A with natural ventilation, and Onset Hobo H8 Pro Temp)
01/07/2013	Datalogger download, check of sensor status and functioning. The fan-aspirated radiation shield was found damaged and not working

03/09/2013	Datalogger download, check of sensor status and functioning, 2 m increase in height of support tower. The fan-aspirated radiation shield was repaired and resumed working properly
03/07/2014	Datalogger download, check of sensor status and functioning
23/09/2014	2 m elongation of support tower. Sensors have been partially buried by snow between 2 and 23/09/2014
29/06/2015	Datalogger download, check of sensor status and functioning. Support tower lengthened by 2 m. Sensors have been partially buried by snow between 15 and 29/06/2015. Removal of all sensors except the Onset Hobo H8 Pro Temp
31/08/2015	Onset Hobo H8 Pro Temp download, check of sensor status and functioning
02/09/2016	Onset Hobo H8 Pro Temp download, check of sensor status and functioning. Sensor removed

458

459

460 Table D2. Field operations and maintenance for the englacial temperature sensors installed on Mt. Ortles.

Date	Field operations
17/11/2011	Installation and launch of the Sommer thermistor strings at the borehole drilling site and at the Hintergrat Glacier.
18/06/2012	Installation and launch of the Geoprecision thermistor string at the AWS site. Download and maintenance (battery replacement) of the Sommer thermistor strings at the borehole drilling site
28/08/2012	Download and maintenance (battery replacement) of the Sommer thermistor string at the Hintergrat Glacier. Sensor at 7.5 m initial depth stopped working on 18/08/2012
07/09/2012	Download of the Geoprecision thermistor string at the AWS site. Download and maintenance (battery replacement) of the Sommer thermistor strings at the borehole drilling site
01/07/2013	Download of the Geoprecision thermistor string at the AWS site. Download and maintenance (battery replacement, logger raised to the surface) of the Sommer thermistor strings at the borehole drilling site
23/08/2013	Download and maintenance (battery replacement) of the Sommer thermistor string at the Hintergrat Glacier. The sensor at 1.5 m initial depth was above the glacier surface
03/09/2013	Download of the Geoprecision thermistor string at the AWS site. Download and maintenance (battery replacement) of the Sommer thermistor strings at the borehole drilling site. Long string stopped working on 10/04/2013
03/07/2014	Download and maintenance (battery replacement, logger replacement) of the short Sommer thermistor string at borehole drilling site. Removal of the datalogger of the long Sommer thermistor string at borehole drilling site (no longer working).
28/08/2014	Download and maintenance (battery replacement) of the Sommer thermistor string at the Hintergrat Glacier. Sensor at 5.5 m initial depth stopped working on 17/08/2014
23/09/2014	Download and maintenance (battery replacement, logger raised to the surface) of the short Sommer thermistor string at the borehole drilling site
18/10/2014	Download of the Geoprecision thermistor string at the AWS site
29/06/2015	Download of the Geoprecision thermistor string at the AWS site
27/08/2015	Download and maintenance (battery replacement) of the Sommer thermistor string at the Hintergrat Glacier
31/08/2015	Download and removal of the short Sommer thermistor string at the borehole drilling site
23/08/2016	Download and removal of the Sommer thermistor string at the Hintergrat Glacier
02/09/2016	Download of the Geoprecision thermistor string at the AWS site

461

462

463 Table D3. Field operations and maintenance for the soil surface temperature sensors installed on Mt. Ortles.

Date	Field operations
02/09/2010	Installation and launch of the ORTL_05, ORTL_06, ORTL_07 and ORTL_08 Gemini TGP-4020
23/09/2010	Installation and launch of the ORTL_10 and ORTL_11 Gemini TGP-4020
28/08/2012	Download and maintenance (battery replacement, logger re-launch) at the ORTL_07, ORTL_08, ORTL_10 and ORTL_11
07/09/2012	Download and maintenance (battery replacement, logger re-launch) at the ORTL_05 and ORTL_06
23/08/2013	Download and maintenance (battery replacement, logger re-launch) at the ORTL_07, ORTL_08, ORTL_10 and ORTL_11
03/09/2013	Download and maintenance (battery replacement, logger re-launch) at the ORTL_05 and ORTL_06
28/08/2014	Download and maintenance (battery replacement, logger re-launch) at the ORTL_07, ORTL_08 and ORTL_10. ORTL_05, ORTL_06 and ORTL_11 not found, buried by snow
27/08/2015	Download and maintenance (battery replacement, logger re-launch) at the ORTL_07, ORTL_08 and ORTL_10
31/08/2015	Download and maintenance (battery replacement, logger re-launch) at the ORTL_05 and ORTL_06
23/08/2016	Download and removal of the ORTL_07, ORTL_08 and ORTL_10
02/09/2016	Download and removal of the ORTL_05 and ORTL_06

464

465

466 Table D4. Field operations and maintenance for the rock wall temperature sensors installed on Mt. Ortles.

Date	Field operations
30/08/2011	Installation and launch of the Geoprecision thermistor strings at CIMA_ALTO, CIMA_VERTICALE, ANTICIMA_SUD, BIV_LOMBARDI
14/10/2011	Installation and launch of the Geoprecision thermistor string at HINTERGRAT
28/11/2011	Installation and launch of the Geoprecision thermistor string at PAYER
28/08/2012	Download of thermistor strings at HINTERGRAT and PAYER. Repair of the Geoprecision thermistor string at HINTERGRAT
07/09/2012	Download of thermistor strings at CIMA_ALTO, CIMA_VERTICALE, ANTICIMA_SUD, BIV_LOMBARDI
23/08/2013	Download of thermistor strings at HINTERGRAT and PAYER
03/09/2013	Download of thermistor strings at CIMA_ALTO, CIMA_VERTICALE, ANTICIMA_SUD, BIV_LOMBARDI
28/08/2014	Download of thermistor string at PAYER
01/09/2014	Download of thermistor strings at HINTERGRAT, damaged, removed
27/08/2015	Download of thermistor string at PAYER
31/08/2015	Download of thermistor strings at CIMA_ALTO, CIMA_VERTICALE, ANTICIMA_SUD, BIV_LOMBARDI. CIMA_ALTO damaged, removed
23/08/2016	Download and removal of the thermistor string at PAYER
02/09/2016	Download and removal of the thermistor strings at CIMA_VERTICALE, ANTICIMA_SUD, BIV_LOMBARDI

467

468

469 **APPENDIX E: Characteristics of measurement sites**

470 Table E1. Topographic and geomorphological characteristics of sites instrumented for temperature
 471 measurements.

Measured variable	Elevation (m a.s.l.)	East coordinate UTM WGS84 (m)	North coordinate UTM WGS84 (m)	Aspect	Slope (degrees)	Site description
Air Temperature (automatic weather station)	3830	618254	5151614	NW	11	Upper accumulation area of Alto dell'Ortles Glacier
Snow and firn temperature at the AWS site	3830	618260	5151619	NW	11	Upper accumulation area of Alto dell'Ortles Glacier
Englacial temperature at the borehole drilling site (short and long strings)	3859	618373	5151536	W	7	Upper accumulation area of Alto dell'Ortles Glacier
Englacial temperature at the Hintergrat Glacier	3476	619435	5151395	N	12	Hintergrat Glacier
Soil surface temperature at Lombardi bivouac - ORTL_05	3351	618202	5152846	SW	7	Northern ridge of Mt. Ortles, bedrock covered by a thin layer of debris (fine gravel, sand)
Soil surface temperature at Lombardi bivouac - ORTL_06	3371	618284	5152772	N	22	Northern ridge of Mt. Ortles, recently deglaciated bedrock covered by a discontinuous layer of loose debris (fine gravel, sand)
Soil surface temperature at Payer - refuge ORTL_07	2994	618361	5153936	N	22	Northern ridge of Mt. Ortles, bedrock covered by a thick layer of debris (pebbles, gravel, sand) with sparse vegetation
Soil surface temperature at Payer - refuge ORTL_08	2899	618287	5154105	W	36	Northern ridge of Mt. Ortles, bedrock covered by coarse debris with isolated areas of thinner debris (fine sand and silt).
Soil surface temperature at Hintergrat ridge ORTL_10	3460	619628	5151341	S	22	Eastern ridge of Mt. Ortles, bedrock covered by a layer of debris (fine gravel, sand).
Soil surface temperature at Hintergrat ridge ORTL_11	3466	619491	5151374	SE	11	Eastern ridge of Mt. Ortles, bedrock covered by a thin layer of coarse debris (gravel, sand), close to the edge of the Hintergrat Glacier
Rock wall temperature at Mt. Ortles summit - CIMA_ALTO	3900	618512	5151691	E	70	70 m south of Mt. Ortles summit (3905 m), in a sub-vertical rock face about 30 m below the crest edge
Rock wall temperature Mt. Ortles summit - CIMA_VERTICALE	3880	618512	5151691	E	90	70 m south of Mt. Ortles summit (3905 m), in a vertical rock face about 50 m below the ridge, 20 m below CIMA_ALTO
Rock wall temperature at Vorgipfel - ANTICIMA_SUD	3810	618327	5151269	S	90	Vertical rock face, about 10 m below the upper rock wall edge
Rock wall temperature at Lombardi bivouac - BIV_LOMBARDI	3351	618213	5152784	W	70	Northern ridge of Mt. Ortles, sub-vertical rock wall, about 30 m below the crest edge
Rock wall temperature at Hintergrat - HINTERGRAT	3370	619710	5151334	NE	90	Eastern ridge of Mt. Ortles, vertical rock wall, about 10 m below the crest edge
Rock wall temperature at Payer refuge - PAYER	3030	618372	5153812	SE	90	Northern ridge of Mt. Ortles, vertical rock wall, about 20 m below the crest edge

472
473 **APPENDIX F: Description of the measuring equipment for air temperature**

474



475
476 Figure F1. The automatic weather station (AWS) installed on Mt. Ortles, whose summit is visible in the
477 background. The stake behind the weather station indicates the site of the Geoprecision thermistor string. Photo
478 taken on 7 September 2012, after the lengthening of the support tower.

479




480

481 Figure F2. The wooden boards placed at the bottom of the support aluminum tower, at 2 m depth in the firn,

482

during the AWS installation. Photo taken on 30 September 2011.




483
 484 Figure F3. Detail of the AWS seen from the west: a) R. M. Young 05103 anemometers, p) Delta Ohm LP
 485 PIRG 01 pyrgeometers, r) Delta Ohm LP Pyra 05 radiometers, s) Campbell Scientific SR50A snow depth
 486 sensors, v1) Vaisala HMP155A inside the R. M. Young 43502 fan-aspirated radiation shield, g1) Gemini TGP-
 487 4020 datalogger inside the R. M. Young 43502 fan-aspirated radiation shield, v2) Vaisala HMP155A inside
 488 the 15-plates Campbell Scientific MET 21 radiation shield with natural ventilation, o) Onset Hobo H8 Pro

489 Temp datalogger inside the 8-plates Davis 7714 radiation shield with natural ventilation. Photo taken on 7
490 September 2012.

491

492



493

494 Figure F4. Detail of the AWS seen from the east: a) R. M. Young 05103 anemometers, p) Delta Ohm LP PIRG

495 01 pyrgeometers, r) Delta Ohm LP Pyra 05 radiometers, s) Campbell Scientific SR50A snow depth sensors,

496 v1) Vaisala HMP155A inside the R. M. Young 43502 fan-aspirated radiation shield, g2) Gemini TGP-4020
497 datalogger inside the 6-plates R.M. Young 41303-5 radiation shield with natural ventilation, o) Onset Hobo
498 H8 Pro Temp datalogger inside the 8-plates Davis 7714 radiation shield with natural ventilation. Photo taken
499 on 7 September 2012.

500

501

502

503

504 **APPENDIX G: Description of the measuring equipment for englacial temperature**

505



506

507 Figure G1. The drilling site seen from the summit of Mt. Ortles. The Vorgipfel-Anticima Sud is visible in the
508 background. Photo taken on 1 October 2011 during the ice drilling operations, and before setting up the drilling
509 site thermistor strings for englacial temperature measurements.

510


511
512
513
514
515
516



517
518 Figure G2. The Hintergrat Glacier seen from the south east (aerial photo taken on 28 August 2012). The black
519 asterisk indicates the location of the borehole equipped with the thermistor string.

520
521
522
523
524
525
526
527


528
529



530
531
532
533
534
535
536
537
538
539
540
541
542
543
544

Figure G3. a) Lowering of a thermistor string into the borehole n. 3 at the drilling site. b) The winding systems of the two thermistor strings installed at the drilling site. c) The two metal boxes containing the thermistor string data loggers and the batteries. d) Final arrangement of the box housing the winding systems and the data loggers. The summit of Mt. Ortles is visible in the background. Photos taken on 17 November 2011.

545
546



547

548 Figure G4. a) Borehole drilling at the Hintergrat Glacier using a steam ice drill. b) The box containing the
549 winding system, the thermistor string data logger and the batteries. c) Final arrangement of the box housing
550 the winding systems and the data logger. Photos taken on 17 November 2011 and 27 August 2015.

551

552 APPENDIX H: Description of the measuring equipment for soil surface temperature

553



554

555 Figure H1. The soil surface temperature datalogger (Gemini TGP-4020) installed at the ORTL_05 site, close
556 to the Lombardi bivouac. The white ellipse indicates the PB-5001 external probe placed underneath the debris
557 surface. Photo taken on 2 September 2010.

558

559

560



561

562 Figure H2. Data download and logger maintenance at the ORTL_10 soil surface temperature site on 28
563 August 2012.

564


565

566

567

568


569



570
571
572
573
574
575
576

Figure H3. Location of the six sites equipped with dataloggers for soil surface temperature measurement on Mt. Ortles.

578



579

580 Figure I1. The rock wall temperature datalogger (Geoprecision thermistor string) installed close to the Payer
581 refuge. The datalogger is anchored to the rock wall and is connected to three temperature sensors placed at 0.1,
582 0.3 and 0.55 m depth inside a horizontal hole drilled in the rock wall. Photo taken on 28 November 2011.

583



584
585
586
587
588
589
590

Figure I2. Launching of the CIMA ALTO data logger. A wireless USB dongle secures the wireless connection to the laptop used for launching the logger and for downloading the temperature data. Photo taken on 30 August 2011.



591
592

593 Figure I3. Location of the six sites equipped with data loggers for rock wall temperature measurement on Mt.
594 Ortles (BIV_LOMBARDI and HINTERGRAT from Google Earth Pro 7.3 (2022)).

595

596

597 **Acknowledgements**

598 This work is a contribution to the Ortles Project, a program supported by two NSF awards no. 1060115 & no. 1461422
599 to The Ohio State University and by the Ripartizione Protezione Antincendi e Civile of the Autonomous province of
600 Bolzano in collaboration with the Ripartizione Opere idrauliche e Ripartizione Foreste of the Autonomous province of
601 Bolzano and the Stelvio National Park. This is Ortles Project publication 11 (www.ortles.org). The research was funded
602 by the Italian MIUR Project (PRIN 2010-11), "Response of morphoclimatic system dynamics to global changes and
603 related geomorphological hazards" (local and national coordinators G. Dalla Fontana and C. Baroni). The authors are
604 grateful to all the students, technicians and scientists who contributed to the field activities in the period from 2009 to
605 2016, the Alpine guides of the Alpinschule of Solda, the helicopter companies Airway, Air Service Center, Star Work
606 Sky and the Hotel Franzenshöhe for logistical support. The authors acknowledge the support of Vinicio Carraro (TeSAF
607 Department of the University of Padova) for the setup of the automatic weather station, and of Umberto Morra di Cella
608 and Paolo Pogliotti (ARPA Val d'Aosta) for the setup of the rock wall temperature dataloggers. The soil surface
609 temperature measured on Mt. Vioz were kindly provided by the Servizio Geologico of the Autonomous province of
610 Trento (Matteo Zumiani).

611

612

613 **References**

614

615 Ballantyne, C. K.: Periglacial geomorphology, John Wiley & Sons Ltd., Hoboken, NJ, USA, 472 pp., ISBN:
616 978-1-405-10006-9, 2018.

617

618 Boeckli, L., Brenning, A., Gruber, S., and Noetzli, J.: Permafrost distribution in the European Alps: Calculation
619 and evaluation of an index map and summary statistics, *The Cryosphere*, 6, 807–820,
620 <https://doi.org/10.5194/tc-6-807-2012>, 2012.

621

622 Bohleber, P., Wagenbach, D., Schöner, W., and Böhm, R.: To what extent do water isotope records from low
623 accumulation Alpine ice cores reproduce instrumental temperature series?, *Tellus B*, 65, 1–17,
624 <https://doi.org/10.3402/tellusb.v65i0.20148>, 2013.

625

626 Braithwaite, R. J., Zhang, Y., and Raper, S. C. B.: Temperature sensitivity of the mass balance of mountain
627 glaciers and icecaps as a climatological characteristic, *Z. Gletsch.kd. Glazialgeol.*, 38, 35–61 pp., 2002.

628

629 Carturan, L.: Climate change effects on the cryosphere and hydrology of a high-altitude watershed, Ph.D.
630 thesis, TeSAF – University of Padova, 187 pp., 2010.

631

632 Carturan, L., Cazorzi, F., and Dalla Fontana, G.: Distributed mass-balance modelling on two neighbouring
633 glaciers in Ortles-Cevedale Italy, from 2004 to 2009, *J. Glaciol.*, 58, 467–486,
634 <https://doi.org/10.3189/2012JoG11J111>, 2012a.

635

636 Carturan, L., Dalla Fontana, G., and Borga, M.: Estimation of winter precipitation in a high-altitude catchment
637 of the Eastern Italian Alps: Validation by means of glacier mass balance observations, *Geogr. Fis. Din. Quat.*,
638 35, 37–48, <https://doi.org/10.4461/GFDQ.2012.35.4>, 2012b.

639

640 Carturan, L., Filippi, R., Seppi, R., Gabrielli, P., Notarnicola, C., Bertoldi, L., Paul, F., Rastner, P., Cazorzi,
641 F., Dinale, R., and Dalla Fontana, G.: Area and volume loss of the glaciers in the Ortles-Cevedale group
642 (Eastern Italian Alps): controls and imbalance of the remaining glaciers, *The Cryosphere*, 7, 1339–1359,
643 <https://doi.org/10.5194/tc-7-1339-2013>, 2013.

644

645 Carturan, L., Cazorzi, F., De Blasi, F., and Dalla Fontana, G.: Air temperature variability over three glaciers
646 in the Ortles–Cevedale (Italian Alps): Effects of glacier fragmentation, comparison of calculation methods,
647 and impacts on mass balance modeling, *The Cryosphere*, 9(3), 1129– 1146, [https://doi.org/10.5194/tc-9-1129-](https://doi.org/10.5194/tc-9-1129-2015)
648 2015, 2015.

649

650 Carturan, L., De Blasi, F., Dinale, R., Dragà, G., Gabrielli, P., Mair, V., Seppi, R., Tonidandel, D., Zanoner,
651 T., Zandrini, T. L., and Dalla Fontana, G.: Data from air, englacial and permafrost temperature measurements
652 on Mt. Ortles (Eastern European Alps) (Version 1.0) [Data set]. Zenodo.
653 <https://doi.org/10.5281/zenodo.7879969>, 2023.

654

655 Charbonneau, R., Lardeau, J. P., and Obled, C.: Problems of modelling a high mountainous drainage basin
656 with predominant snow yields, *Hydrol. Sci. B.*, 26, 345–361, <https://doi.org/10.1080/02626668109490899>,
657 1981.

658

659 Cicoira, A., Beutel, J., Faillettaz, J., Gärtner-Roer, I., and Vieli, A.: Resolving the influence of temperature
660 forcing through heat conduction on rock glacier dynamics: A numerical modelling approach, *The Cryosphere*,
661 13, 927–942, <https://doi.org/10.5194/tc-13-927-2019>, 2019.

662

663 Deline, P., Gruber, S., Delaloye, R., Fischer, L., Geertsema, M., Giardino, M., Hasler, A., Kirkbride, M.,
664 Krautblatter, M., Magnin, F., McColl, S., Ravanel, L., and Schoeneich, P.: Ice Loss and Slope Stability in
665 High-Mountain Regions, *Snow and Ice-Related Hazards, Risks, and Disasters*, Academic Press, 521–561,
666 <https://doi.org/10.1016/B978-0-12-394849-6.00015-9>, 2015.

667

668 Gabrielli, P., Carturan, L., Gabrieli, J., Dinale, R., Krainer, K., Hausmann, H., Davis, M., Zagorodnov, V.,
669 Seppi, R., Barbante, C., Fontana, G. D., and Thompson, L. G.: Atmospheric warming threatens the untapped
670 glacial archive of Ortles mountain, South Tyrol, *J. Glaciol.*, 56, 843–853,
671 <https://doi.org/10.3189/002214310794457263>, 2010.

672

673 Gabrielli, P., Barbante, C., Carturan, L., Cozzi, G., Fontana, G. D., Dinale, R., Dragà, G., Gabrieli, J.,
674 Kehrwald, N., Mair, V., Thompson, L. G., and Tonidandel, D.: Discovery of cold ice in a new drilling site in
675 the eastern European Alps, *Geogr. Fis. Din. Quat.*, 35, 101–105, <https://doi.org/10.4461/GFDQ.2012.35.10>,
676 2012.

677

678 Gabrielli, P., Barbante, C., Bertagna, G., Bertó, M., Binder, D., Carton, A., Carturan, L., Cazorzi, F., Cozzi,
679 G., Dalla Fontana, G., Zanoner, T., and Zennaro, P.: Age of the Mt. Ortles ice cores, the Tyrolean Iceman and
680 glaciation of the highest summit of South Tyrol since the Northern Hemisphere Climatic Optimum, *The*
681 *Cryosphere*, 10, 2779–2797, <https://doi.org/10.5194/tc-10-2779-2016>, 2016.

682

683 Guglielmin, M., Aldighieri, B., and Testa, B.: PERMACLIM: A model for the distribution of mountain
684 permafrost, based on climatic observations, *Geomorphology*, 51, 245–257, [https://doi.org/10.1016/S0169-555X\(02\)00221-0](https://doi.org/10.1016/S0169-555X(02)00221-0), 2003.

686

687 Harris, C., Arenson, L. U., Christiansen, H. H., Etzelmüller, B., Frauenfelder, R., Gruber, S., Haeberli, W.,
688 Hauck, C., Hölzle, M., Humlum, O., Springman, S. M., and Vonder Mühl, D.: Permafrost and climate in
689 Europe: Monitoring and modelling thermal, geomorphological and geotechnical responses, *Earth Sci. Rev.*,
690 92, 117–171, <https://doi.org/10.1016/j.earscirev.2008.12.002>, 2009.

691

692 Huggel, C., Carey, M., Clague, J. J., and Kääh, A.: The high-mountain cryosphere: Environmental changes
693 and human risks, Cambridge University Press, 1–371 pp., <https://doi.org/10.1017/CBO9781107588653>, 2015.

694

695 Irvine-Fynn, T. and Hubbard, B.: Glacier Hydrology and Runoff, In *International Encyclopedia of Geography:*
696 *People, the Earth, Environment and Technology*, Elsevier,
697 <https://doi.org/https://doi.org/10.1002/9781118786352.wbieg0709>, 2017.

698

699 Kinnard, C., Larouche, O., Demuth, M. N., and Menounos, B.: Modelling glacier mass balance and climate
700 sensitivity in the context of sparse observations: application to Saskatchewan Glacier, western Canada, *The*
701 *Cryosphere*, 16, 3071–3099, <https://doi.org/10.5194/tc-16-3071-2022>, 2022.

702

703 Knight, J. and Harrison, S.: The sensitivity and evolutionary trajectory of the mountain cryosphere:
704 Implications for mountain geomorphic systems and hazards, *Land Degrad. Dev.*,
705 <https://doi.org/10.1002/ldr.4630>, 2023.

706

707 Langsdorf, S., Löschke, S., Möller, V., Okem, A., Officer, S., Rama, B., Belling, D., Dieck, W., Götze, S.,
708 Kersher, T., Mangele, P., Maus, B., Mühle, A., Nabiyeva, K., Nicolai, M., Niebuhr, A., Petzold, J., Prentzler,
709 E., Savolainen, J., Scheuffele, H., Weisfeld, S., Weyer, N., Pörtner, H.-O., Roberts, D. C., Tignor, M.,

710 Poloczanska, E. S., Mintenbeck, K., Alegría, A., Craig, M., Langsdorf, S., Löschke, S., Möller, V., Okem, A.,
711 and Rama, B.: IPCC, 2022: Climate Change 2022: Impacts, Adaptation and Vulnerability. Working Group II
712 Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge
713 University Press, Cambridge, UK and New York, NY, USA, 3056 pp.,
714 <https://doi.org/10.1017/9781009325844>, 2022.

715

716 Machguth, H., Purves, R. S., Oerlemans, J., Hoelzle, M., and Paul, F.: Exploring uncertainty in glacier mass
717 balance modelling with Monte Carlo simulation, *The Cryosphere*, 2, 191–204, [https://doi.org/10.5194/tc-2-](https://doi.org/10.5194/tc-2-191-2008)
718 191-2008, 2008.

719

720 Montrasio A, Berra F, Cariboni M, Ceriani M, Deichmann N, Ferliga C, Gregnanin A, Guerra S, Guglielmin
721 M, Jadoul F, Longhin M, Mair V, Mazzoccola D, Sciesa E, and Zappone A: Note illustrative della Carta
722 Geologica d'Italia - Foglio 024 Bormio, ISPRA, Roma, Italy, 16–26 pp., 2012.

723

724 Pepin, N., Bradley, R. S., Diaz, H. F., Baraer, M., Caceres, E. B., Forsythe, N., Fowler, H., Greenwood, G.,
725 Hashmi, M. Z., Liu, X. D., Williamson, S. N., and Yang, D. Q.: Elevation-dependent warming in mountain
726 regions of the world, *Nat. Clim. Change*, 5, 424–430, <https://doi.org/10.1038/nclimate2563>, 2015.

727

728 Pepin, N. C., Arnone, E., Gobiet, A., Haslinger, K., Kotlarski, S., Notarnicola, C., Palazzi, E., Seibert, P.,
729 Serafin, S., Schöner, W., Terzago, S., Thornton, J. M., Vuille, M., and Adler, C.: Climate Changes and Their
730 Elevational Patterns in the Mountains of the World, *Rev. Geophys.*, 60,
731 <https://doi.org/10.1029/2020RG000730>, 2022.

732

733 Shaw, T. E., Buri, P., McCarthy, M., Miles, E. S., Ayala, Á., and Pellicciotti, F.: The Decaying Near-Surface
734 Boundary Layer of a Retreating Alpine Glacier, *Geophys. Res. Lett.*, 50,
735 <https://doi.org/10.1029/2023gl103043>, 2023.

736

737 Rathore, N., Thakur, D., and Chawla, A.: Seasonal variations coupled with elevation gradient drives significant
738 changes in eco-physiological and biogeochemical traits of a high altitude evergreen broadleaf shrub,
739 *Rhododendron anthopogon*, *Plant Physiol. Bioch.*, 132, 708–719,
740 <https://doi.org/10.1016/J.PLAPHY.2018.08.009>, 2018.

741

742 Troxler, P., Ayala, Á., Shaw, T. E., Nolan, M., Brock, B. W., and Pellicciotti, F.: Modelling spatial patterns of
743 near-surface air temperature over a decade of melt seasons on McCall Glacier, Alaska, *J. Glaciol.*, 1–15(257),
744 386– 400, <https://doi.org/10.1017/jog.2020.12>, 2020.

745

746 World Meteorological Organization: Guide to Instruments and Methods of Observation, Volume I,
747 Measurement of Meteorological Variables (WMO No. 8), Geneva, Switzerland, 2021.

748

749 Zolles, T., Maussion, F., Galos, S. P., Gurgiser, W., and Nicholson, L.: Robust uncertainty assessment of the
750 spatio-temporal transferability of glacier mass and energy balance models, *The Cryosphere*, 13, 469–489,
751 <https://doi.org/10.5194/tc-13-469-2019>, 2019.