

Indicators of Global Climate Change 2022: Annual update of large-scale indicators of the state of the climate system and the human influence

5 Piers M. Forster¹, Christopher J. Smith^{1,2}, Tristram Walsh³, William F. Lamb^{4,1}, Robin Lamboll⁵, Mathias Hauser⁶, Aurélien Ribes⁷, Debbie Rosen¹, Nathan Gillett⁸, Matthew D. Palmer^{9,10}, Joeri Rogelj⁵, Karina von Schuckmann¹¹, Sonia I. Seneviratne⁶, Blair Trewin¹², Xuebin Zhang⁸, Myles Allen³, Robbie Andrew¹³, Arlene Birt¹⁴, Alex Borger¹⁵, Tim Boyer¹⁶, Jiddu A. Broersma¹⁵, Lijing Cheng¹⁷, Frank Dentener¹⁸, Pierre Friedlingstein^{19,20}, José M. Gutiérrez²¹, Johannes Gütschow²², Bradley Hall²², Masayoshi Ishii²⁴, Stuart Jenkins³, Xin Lan^{22,40}, June-Yi Lee²⁵, Colin Morice⁹, Christopher Kadow²⁶, John Kennedy²⁷, Rachel Killick⁹, Jan C. Minx^{4,1}, Vaishali Naik²⁸, Glen P. Peters¹³, Anna Pirani²⁹, Julia Pongratz^{30,39}, Carl-Friedrich Schleussner³¹, Sophie Szopa³², Peter Thorne³³, Robert Rohde³⁴, Maisa Rojas Corradi³⁵, Dominik Schumacher⁶, Russell Vose³⁶, Kirsten Zickfeld³⁷, Valerie Masson-Delmotte³², Panmao Zhai³⁸

¹Priestley Centre, University of Leeds, Leeds, LS2 9JT, UK

15 ²International Institute for Applied Systems Analysis (IIASA), Austria

³Environmental Change Institute, University of Oxford, UK

⁴Mercator Research Institute on Global Commons and Climate Change (MCC), Berlin, Germany

⁵Centre for Environmental Policy, Imperial College London, UK

20 ⁶Institute for Atmospheric and Climate Science, Department of Environmental Systems Science, ETH Zurich, Zurich, Switzerland

⁷Université de Toulouse, Météo France, CNRS, France

⁸Environment and Climate Change Canada, Canada

⁹Met, Office Hadley Centre, UK

¹⁰School of Earth Sciences, University of Bristol, UK

25 ¹¹Mercator Ocean international, Toulouse, France

¹²Bureau of Meteorology, Australia

¹³CICERO, Center for International Climate Research, Norway

¹⁴Backgroundstories.com, Minneapolis College of Art and Design

¹⁵ClimateChangeTracker.org

30 ¹⁶NOAA, National Centers for Environmental Information, Silver Spring, MD, USA

¹⁷Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China

¹⁸European Commission, & Joint Research Centre, Institute for Environment and Sustainability, Ispra, Italy

¹⁹Faculty of Environment, Science and Economy, University of Exeter, UK

35 ²⁰Laboratoire de Météorologie Dynamique/Institut Pierre-Simon Laplace, CNRS, Ecole Normale Supérieure/Université PSL, Paris, France

²¹Instituto de Física de Cantabria (CSIC-University of Cantabria), Spain

²²Climate Resource, Australia/Germany

²³NOAA, Global Monitoring Laboratory, Boulder, CO, USA

²⁴Meteorological Research Institute, Tsukuba, Japan

40 ²⁵Research Center for Climate Sciences, Pusan National University and Center for Climate Physics, Institute for Basic Science, Pusan, Republic of Korea

²⁶German Climate Computing Center (DKRZ)

Deleted: Lamb^{4,1}, Matthew D. Palmer^{9,10}, Joeri Rogelj⁵, Palmer^{5,6}, Karina von Schuckmann¹¹, Sonia I. Seneviratne⁶, Schuckmann⁷, Blair Trewin¹², Xuebin Myles Allen³, Robbie Andrew¹³, Andrew⁷, Arlene Birt¹⁴, Birt¹⁰, Alex Borger¹⁵, Borger¹¹, Tim Boyer¹⁶, Boyer¹², Jiddu A. Broersma¹⁵, Broersma¹¹, Lijing Cheng¹⁷, Cheng¹³, Frank Dentener¹⁸, Dentener¹⁴, Pierre Friedlingstein^{19,20}, Friedlingstein^{15,16}, Nathan Gillett¹⁷, José M. Gutiérrez²¹, Gutiérrez¹⁸, Johannes Gütschow²², Gütschow¹⁹, Mathias Hauser²⁰, Bradley Hall²², Hall²¹, Masayoshi Ishii²⁴, Ishii²², Stuart Jenkins³, Robin Lamboll²³, Xin Lan^{22,40}, Lan²¹, June-Yi Lee²⁵, Lee²⁴, Colin Morice⁹, Morice⁵, Christopher Kadow²⁶, Kadow²⁶, John Kennedy²⁷, Kennedy²⁶, Rachel Killick⁹, Killick², Jan C. Minx^{4,1}, Vaishali Naik²⁸, Naik²⁷, Glen P. Peters¹³, Peters⁹, Anna Pirani²⁹, Pirani²⁸, Julia Pongratz^{30,39}, Pongratz²⁹, Aurélien Ribes³⁰, Joeri Rogelj²³, Debbie Rosen¹, arl-Rosen¹, arl-Friedrich Schleussner³¹, Sonia I. Seneviratne²⁰, Sophie Szopa³², Peter Thorne³³, Robert Rohde³⁴, Maisa Rojas Corradi³⁵, Dominik Schumacher⁶, Schumacher²⁰, Russell Vose³⁶, Kirsten Zickfeld³⁷, Xuebin Zhang¹⁷, Valerie Masson-Delmotte³⁸, Panmao Zhai³⁸, Zhai³⁹ (... 11)

Deleted: ¹Met

Deleted: ⁶School

Deleted: ⁷Mercator

Deleted: ⁸Bureau

Deleted: ⁹CICERO

Deleted: ¹⁰Backgroundstories

Deleted: ¹¹ClimateChangeTracker

Deleted: ¹²NOAA

Deleted: ¹³Institute

Deleted: ¹⁴European

Deleted: ¹⁵Faculty

Deleted: ¹⁶Laboratoire

Deleted: ¹⁷Environment and Climate Change Canada, Canada⁴

¹⁸Instituto

Deleted: ¹⁹Climate

Deleted: ²⁰Institute for Atmospheric and Climate Science, Department of Environmental Systems Science, ETH Zurich, Zurich, Switzerland⁴

²¹NOAA

Deleted: ²²Meteorological

Deleted: ²³Centre for Environmental Policy, Imperial College London, UK⁴

²⁴Research

Deleted: ²⁵German

145 ²⁷No affiliation, independent
²⁸NOAA, GFDL, Princeton, New Jersey, USA
²⁹Université Paris-Saclay, France; CMCC, Italy; Università Cà Foscari, Italy
³⁰University of Munich, Germany
³¹Climate Analytics, Berlin, Germany and Geography Department and IRI THESys, Humboldt-Universität zu Berlin, Berlin, Germany
150 ³²Université Paris-Saclay, CNRS, CEA, UVSQ, Laboratoire des Sciences du Climat et de l'Environnement, 91191, Gif-sur-Yvette, France
³³ICARUS Climate Research Centre, Maynooth University, Maynooth, Ireland
³⁴Berkeley Earth, Berkeley, CA, USA
³⁵University of Chile, Santiago, Chile
155 ³⁶NOAA's National Centers for Environmental Information (NCEI), Asheville, NC, USA
³⁷Simon Fraser University, Vancouver, Canada
³⁸Chinese Academy of Meteorological Sciences, Beijing, China
³⁹Max Planck Institute for Meteorology, Hamburg, Germany
⁴⁰CIRES, University of Colorado Boulder, USA

Deleted: ²⁶No
Deleted: ²⁷NOAA
Deleted: ²⁸Université
Deleted: ²⁹University
Deleted: and Max Planck Institute for Meteorology, Hamburg, Germany...³⁰Université de Toulouse, Météo France, CNRS, France
³⁰Université de Toulouse, Météo France, CNRS, France
Deleted: ³⁰Université de Toulouse, Météo France, CNRS, France
Deleted: ³²Laboratoire
Deleted: IPSL, Paris
Deleted: ³⁸Laboratoire des Sciences du Climat et de l'Environnement (LSCE) / Institut Pierre Simon Laplace (IPSL), CEA-CNRS-UVSQ (UMR8212), Université Paris-Saclay, Gif-sur-Yvette, France
³⁹Chinese

160 *Correspondence to:* Piers. M. Forster (p.m.forster@leeds.ac.uk)

Abstract. Intergovernmental Panel on Climate Change (IPCC) assessments are the trusted source of scientific evidence for climate negotiations taking place under the United Nations Framework Convention on Climate Change (UNFCCC), including the first global stocktake under the Paris Agreement that will conclude at COP28 in December 2023. Evidence-based decision making needs to be informed by up-to-date and timely information on key indicators of the state of the climate system and of the human influence on the global climate system. However, successive IPCC reports are published at intervals of 5-10 years, creating potential for an information gap between report cycles.

We follow methods as close as possible to those used in the IPCC Sixth Assessment Report (AR6) Working Group One (WGI) report. We update monitoring datasets to produce updated estimates for key climate indicators related to forcing of the climate system: emissions of greenhouse gases and short-lived climate forcers, greenhouse gas concentrations, radiative forcing, surface temperature changes, the Earth's energy imbalance, warming attributed to human activities, the remaining carbon budget and estimates of global temperature extremes. The purpose of this effort, grounded in an open data, open science approach, is to make annually updated reliable global climate indicators available in the public domain (<https://doi.org/10.5281/zenodo.7883758>, Smith et al., 2023). As they are traceable to IPCC report methods, they can be trusted by all parties involved in UNFCCC negotiations and help convey wider understanding of the latest knowledge of the climate system and its direction of travel.

Deleted: base this update on the assessment
Deleted: , updating the
Deleted: and
Deleted: including emissions
Deleted: and consistent with

The indicators show that human induced warming reached 1.14 [0.9 to 1.4] °C averaged over the 2013-2022 decade, and 1.26 [1.0 to 1.6] °C in 2022. Over the 2013-2022 period, Human induced warming has been increasing at an unprecedented rate of

Deleted: period
Deleted: is

over 0.2 °C per decade. This high rate of warming is caused by a combination of greenhouse gas emissions being at an all-time high of 54 ± 5.3 GtCO₂e over the last decade, as well as reductions in the strength of aerosol cooling. Despite this, there is evidence that increases in greenhouse gas emissions have slowed, a continued series of these annual updates over this critical decade can track real world changes in direction for the human influence on climate.

1 Introduction

Increased greenhouse gas concentrations combined with reductions in aerosol pollution have led to rapid increases in human induced effective radiative forcing, which has in turn led to atmosphere, land, cryosphere and ocean warming (Gulev et al., 2021). This in turn has led to an intensification of many weather and climate extremes, particularly more frequent and more intense hot extremes, and heavy precipitation across most regions of the world (Seneviratne et al., 2021). Given the speed of recent change, and the need for evidence-based decision-making, this Indicators of Global Climate Change (IGCC) update assembles the latest scientific understanding on the current state of the climate system, how it is evolving and the human influence to support policymakers whilst the next IPCC assessment is under preparation. This first annual update is focused on indicators related to heating of the climate system, building from greenhouse gas emissions towards estimates of human-induced warming and the remaining carbon budget. In future years, this effort could be expanded to encompass other indicators, including global precipitation changes and related extremes.

This update complements other international efforts under the auspices of the Global Climate Observing System (GCOS) and the World Meteorological Organization (WMO). Annual state of the climate reports are released by WMO which use much of the same data analysed here for surface temperature and energy budget trends. The Bulletin of American Meteorological Society (BAMS) releases annual State of the Climate reports covering many essential variables including temperature and greenhouse gas concentrations. However, these reports focus on statistics from the previous year and make slightly different choices over datasets and analysis compared to the IPCC (see Sect. 5). The Global Carbon Project publishes updated carbon dioxide datasets which are used directly in this report. There is no similarly structured activity that provides all the necessary datasets to update annually the assessment of human influence on global surface temperature.

The update is based on methodologies for key climate indicators assessed by the IPCC Sixth Assessment Report (AR6) of the physical science basis of climate change (WGI report; IPCC, 2021a) as well as Chapter 2 of the WGIII report (Dhakal et al., 2022), and is aligned with the efforts initiated in AR6 to implement FAIR principles for reproducibility and reusability (Pirani et al., 2022; Iturbide et al., 2022). IPCC reports make a much wider assessment of the science and methodologies - we do not attempt to reproduce the comprehensive nature of these IPCC assessments here.

Deleted: 57

Deleted: 6

Deleted: are signs

Deleted: emission levels are starting to stabilise, and we can hope that ...

Deleted: might

Deleted: a

Deleted: -

Deleted: change of

Deleted: over this critical decade.

Deleted: and

Deleted: is proposed to assemble

Deleted: ¶

Deleted: analyses and integrates

Deleted: , especially those

Deleted: Essential Climate Variables programme

Deleted: Organization's

Deleted:) Global Atmospheric Watch programme.

Deleted: release

Deleted: .

Deleted: and methane emission

Deleted: current report

Deleted: updates

Deleted: make an annual

Deleted: the

Deleted: - and this is the goal here

Deleted: of

Deleted: (

Deleted: .

Deleted: . 2022). We trace methodologies to these IPCC reports, but as calibrations are revised and science moves forward, we update methods where necessary. However, we do this as transparently as possible to distinguish methodological differences from physical evolution....

270 The IPCC Special Report on Global Warming of 1.5°C (SR1.5), published in 2018, provided an assessment of the level of human-induced warming and cumulative emissions to date (Allen et al., 2018) and the remaining carbon budget (Rogelj et al., 2018) to support the evidence base on how the world is progressing in terms of meeting aspects of the Paris Agreement. The AR6 WGI Report, published in 2021, assessed past, current and future changes of these and other key global climate indicators, as well as undertaking an assessment of the Earth’s energy budget. It also updated its approach for estimating human-induced warming and global warming level. In AR6 WGI and here, reaching a level of global warming is defined as the global surface temperature change, averaged over a 20-year period, exceeding a particular level of global warming, e.g. 1.5°C global warming. Given the current rates of change and the likelihood of reaching 1.5°C of global warming in the first half of the 2030s (Lee et al., 2021; Lee et al., 2023; Riahi et al., 2022), it is important to have robust, trusted, and also timely climate indicators in the public domain to form an evidence base for effective science-based decision making.

280 When making their assessments, authors of IPCC reports assess published literature, but also apply established published analysis methods to assessed datasets, such as that produced by the latest climate model intercomparison projects (Lee et al., 2021). The authors combine and analyze both model and observational data as part of their expert assessment, making assessments of the trustworthiness and error characteristics of different datasets. It is this synthetic analysis by IPCC authors that derives the estimates of key climate indicators. Wherever possible, these same assessed methodological approaches are implemented, here to provide the updates with variations clearly flagged and documented. The same approach, using the same datasets (updated by 2 years) and methods as employed in WGI, was used in the AR6 SYR (2023) report to provide an updated assessment of the latest atmospheric well mixed greenhouse gas concentrations (up to 2021) and decadal average change in global surface temperature (+1.15°C [1.00°C–1.25°C] in 2013-2022 for global surface temperature). However, the assessment of human-induced warming was not updated (and therefore only covers warming up to the decade 2010-2019), nor was the remaining carbon budget updated, so the related information in the 2023 SYR report remained based on data up to the end of 290 2019.

The indicators in this first annual update give important insights into the magnitude and the pace of global warming. This paper provides the basis for a dashboard of climate indicators grounded in IPCC methodologies and directly traceable to reports published as part of the AR6 cycle. We employ datasets that can be updated on a regular basis between the publication of IPCC reports. Note that there are other similar initiatives underway to update other AR6 cycle products; for example, the evolution of the WGI Interactive Atlas (Gutiérrez et al., 2021) is being developed under the Copernicus Climate Change Service (C3S) and has potential connections and synergies with this initiative that will be explored in the future.

Deleted: The COP21 Paris Agreement of 2015 expressly sets out to limit global warming levels through greenhouse gas emission reduction commitments in Articles 2 and 4 respectively. Article 2.1.a sets the goal of holding global temperature increase to well below 2°C above pre-industrial levels, and pursuing efforts to limit the increase to 1.5°C; Article 4.1 states the aim for global greenhouse gas emissions (GHGs) to peak as soon as possible, and to reach a balance between anthropogenic emissions by sources and removals by sinks of GHGs in the second half of the century. Article 2 also sets out clear targets for adaptation (Article 2.1b) and implementation (Article 2.1c). Establishing policies to effectively support efforts to meet these aims and commitments requires reliable indicators of both the state of the climate system and the human influence on climate.

Both COP26 (Glasgow) and COP27 (Sharm El-Sheikh) also “recognized[s] the importance of the best available science for effective climate action” (UNFCCC, 2022a,b). COP27 in 2022 reiterated its invitation to Parties to consider further actions to reduce by 2030 non-carbon dioxide greenhouse gas emissions, including methane. A global stocktake to be held every five years, starting in 2023, has been established under the Paris Agreement to evaluate the collective progress of countries’ actions in the implementation of the Paris Agreement and its long-term goals. The IPCC assessment of the physical science basis provides a wide range of information with relevance for the global stocktake, complementing other products from AR6. The now complete AR6 cycle updated GHG emissions and concentrations, the current state of the climate, near-term and long-term projections of global warming and of the climate system, the attribution of extreme events, and remaining carbon budgets.

The 2015 COP21 Decision invited the IPCC to prepare a special report on the impacts of 1.5°C and related greenhouse gas emission pathways to help inform its work (UNFCCC, 2015). The resulting

Deleted: Agreement’s Article 2

Deleted: 2021b

Deleted: Cross-Chapter Box 1.1, Table 1 in Chapter 1 of the AR6 report (Chen et al., 2021) maps how the material assessed by WGI may be relevant for the global stocktake.

Deleted: 2021b

Deleted: It is

Deleted: that we

Deleted: implementing

Deleted: In this work, we focus on providing updated estimates of key global ...

Deleted: of the state of climate: the Earth’s heat inventory, human-induced warming and the remaining carbon budget. To do this requires updates to emissions, greenhouse gas concentrations, effective radiative forcing, energy imbalance and surface temp (... [2])

Deleted: to

Deleted: Gutierrez

Deleted: .

Our longer-term ambition is to rigorously track both **climate** system change and methodological improvements between IPCC report cycles, thereby building consistency and awareness. An example of why tracking methodological change is important was the **updated estimate for historic warming (the increase in global surface temperature from 1850-1900 to 1986-2005). This was 0.08 [-0.01 to 0.12] °C higher in the AR6 than in the fifth assessment report (AR5) and SR1.5.** Datasets and methods of evaluating global temperature changes altered between the **AR5** and AR6, leading to a small shift in the historical temperature. This was reflected in changes between AR5 and AR6, whereas SR1.5 **mostly relied on methodologies from AR5** (see AR6 WGI Cross Chapter Box 2.3, Gulev et al., 2021). Annual updates provide **indications** of possible future methodological shifts that subsequent IPCC reports may make as science advances, and can detail their impact on perceived trends.

We adopt the Global Carbon Budget ethos of a community-wide inclusive effort that synthesises work from across a large and diverse global scientific community in a timely fashion (Friedlingstein et al., 2022a). Like the Global Carbon Budget, this initiative arises from the international science community to establish a knowledge base to support policy debate and action to **meet the Paris Agreement temperature goal.**

The update is organised as follows: Emissions (Sect. 2) and GHG concentrations (Sect. 3) are used to develop updated estimates of effective radiative forcing (Sect. 4). Observations of global surface temperature change (Sect. 5) and Earth's energy imbalance (Sect. 6) are key global indicators of a warming world. The global **surface** temperature change is formally attributed to human activity in Sect. 7, which tracks human-induced warming. Section 8 updates the remaining carbon budget to policy-relevant temperature thresholds. Section 9 **gives an example of** global-scale indicators **associated with climate extremes of maximum land surface temperatures.**

An important purpose of the exercise is to make these indicators widely available and understood. Plans for a web dashboard are discussed in Sect. 10, code and data availability in Sect. 11 and conclusions presented in Sect. 12. Data is available at <https://doi.org/10.5281/zenodo.7883758> (Smith et al., 2023).

380 2. Emissions

Historic emissions from human activity were assessed in both AR6 WGI and WGIII. Chapter 5 of WGI assessed CO₂ and CH₄ emissions in the context of the carbon cycle (Canadell et al., 2021). Chapter 6 of WGI assessed emissions in the context of understanding the climate and air quality **impacts** of short-lived climate forcers (Szopa et al., 2021). Chapter 2 of WGIII,

Deleted: We track how specific global climate indicators have changed since reported in AR6. It breaks the change down into components from the march of time and any revisions to methods and/or data. For this first report, the methodologies are either identical or very close to those employed in AR6, keeping our results as consistent as possible. There are places where we need to depart as indicated in the text. These occasions are when a particular dataset is not updateable through 2022 and an alternative approach has had to be used.

Deleted: real

Deleted: ,

Deleted: shift in the historical baseline used for

Deleted: temperatures between SR1.5 and AR6.

Deleted: fifth (AR5) assessment report

Deleted: baseline

Deleted: chose to broadly follow AR5 methods

Deleted: 1

Deleted: forewarning

Deleted: 2022

Deleted: slow down and ultimately stop the increase of greenhouse gases in the atmosphere. The update is focussed on building from emissions towards estimates of human-induced warming and the remaining carbon budget.

Deleted:

Deleted:

Deleted:

Deleted:

Deleted: estimates preliminary

Deleted: of

Deleted: extremes. These changes in

Deleted: are not directly related to the other indicators but are included to showcase the possibility of extending the indicator set in future years.

Deleted: ,

Deleted: . (

Deleted: .

Deleted: impact

published one year later (Dhakal et al., 2022), looked at the sectoral sources of emissions and gave the most up to date understanding of the current level of emissions. This section bases its methods and data on those employed in this WGIII chapter.

Deleted:)

Deleted: based

425 2.1 Methods of estimating greenhouse gas emissions changes

Like in AR6 WGIII, net GHG emissions in this paper refer to releases of GHG from anthropogenic sources minus removals by anthropogenic sinks, for those species of gases that are reported under the common reporting format of the UNFCCC. This includes CO₂ emissions from fossil fuels and industry (CO₂-FFI), net CO₂ emissions from land use, land use change and forestry (CO₂-LULUCF), CH₄, N₂O, and fluorinated gas (F-gas) emissions. CO₂-FFI mainly comprises fossil-fuel combustion emissions, as well as emissions from industrial processes such as cement production. This excludes biomass and biofuel use by industry. CO₂-LULUCF is mainly driven by deforestation, but also includes anthropogenic removals on land from afforestation and reforestation, emissions from logging and forest degradation, emissions and removals in shifting cultivation cycles, as well as emissions and removals from other land-use change and land management activities, including peat burning and drainage. The non-CO₂ GHGs, CH₄, N₂O and F-gas emissions - are linked to the fossil-fuel extraction, agriculture, industry and waste sectors.

Deleted:),

Deleted:),

Deleted: ,

Deleted: - also known as land use change emissions -

Deleted: such as

Deleted: or forestry

Deleted: GHG emissions

Deleted: gases

Deleted: and

Global regulatory conventions have led to a two-fold categorisation of F-gas emissions (also known as halogenated gases). Under UNFCCC accounting, countries record emissions of hydrofluorocarbons (HFCs), perfluorocarbons (PFCs), sulphur hexafluoride (SF₆), and nitrogen trifluoride (NF₃) - hereinafter “UNFCCC F-gases”. However, national inventories tend to exclude halons, chlorofluorocarbons (CFCs) and hydrochlorofluorocarbons (HCFCs) - hereinafter “ODS (Ozone Depleting Substances) F-gases” - as they have been initially regulated under the Montreal protocol and its amendments. In line with the WGIII assessment, ODS-F-gases and other substances, including ozone and aerosols, are not included in our GHG emissions reporting, but are included in subsequent assessments of concentrations, effective radiative forcing, human-induced warming, carbon budgets and climate impacts in line with the WGI assessment.

Deleted: assessment

Deleted: concentration

There are also varying conventions used to quantify CO₂-LULUCF fluxes. These include the use of bookkeeping models, dynamic global vegetation models (DVGMs), and the national inventory approach (Pongratz et al. 2021). Each differs in terms of their applied system boundaries and definitions, and are not directly comparable. However, efforts to “translate” between bookkeeping estimates and national inventories using DVGMs have demonstrated a degree of consistency between the varying approaches (Friedlingstein et al., 2022a; Grassi et al., 2023).

465 Each category of GHG emissions included here is covered by varying primary sources and datasets. Although many datasets cover individual categories, few extend across multiple categories, and only a minority have frequent and timely update schedules. Notable datasets include the Global Carbon Budget (GCB; Friedlingstein et al., 2022b), which covers CO₂-FFI and CO₂-LULUCF; the Emissions Database for Global Atmospheric Research (EDGAR; Crippa et al., 2022) and the Potsdam Real-time Integrated Model for probabilistic Assessment of emissions Paths (PRIMAP-hist; Gütschow et al., 2016; Gütschow and Pflüger 2023), which cover CO₂-FFI, CH₄, N₂O and UNFCCC F-gases; and the Community Emissions Data system (CEDS; O'Rourke et al., 2021), which covers CO₂-FFI, CH₄, and N₂O. As detailed below not all these datasets were employed in this update.

In AR6 WGIII, total net GHG emissions were calculated as the sum of CO₂-FFI, CH₄, N₂O and UNFCCC F-gases from EDGAR, and net CO₂-LULUCF emissions from the GCB. Net CO₂-LULUCF emissions followed the GCB convention and were derived from the average of three bookkeeping models (Hansis et al., 2015; Houghton and Nassikas, 2017; Gasser et al., 2020). Version 6 of EDGAR was used (with a fast-track methodology applied for the final year of data - 2019), alongside the 2020 version of the GCB (Friedlingstein et al., 2020). CO₂-equivalent emissions were calculated using global warming potentials with a 100-year time horizon from AR6 WGI Chapter 7 (Forster et al., 2021). Uncertainty ranges were based on a comparative assessment of available data and expert judgement, corresponding to a 90% confidence interval (Minx et al., 2021): ±8% for CO₂-FFI, ±70% for CO₂-LULUCF, ±30% for CH₄, and F-gases, and ±60% for N₂O. (Note that the GCB assesses one standard deviation uncertainty for CO₂-FFI as ±5%, and of ±2.6 GtCO₂ for CO₂-LULUCF; Friedlingstein et al., 2022a). The total uncertainty was summed in quadrature assuming independence of estimates per species / source. Reflecting these uncertainties, AR6 WGIII reported emissions to two significant figures only. Uncertainties in GWP100 metrics were not applied (Minx et al., 2021).

This analysis tracks the same compilation of GHGs as in AR6 WGIII. We follow the same approach for estimating uncertainties and CO₂-equivalent emissions. We also use the same type of data sources, but make important changes to the specific selection of data sources to further improve the quality of the data as suggested in the knowledge gap discussion of the WGIII report (Dhakal et al., 2022). Instead of using EDGAR data (which is now available as version 7), we use GCB data for CO₂-FFI, PRIMAP-hist data for CH₄, and N₂O, and atmospheric concentrations with best-estimate lifetimes for UNFCCC F-gas emissions (Hodnebrog et al., 2020). As in AR6 WGIII we use GCB for net CO₂-LULUCF emissions, taking the average of three bookkeeping models.

Deleted: , with significant differences in uncertainties.

Deleted: 2022

Deleted: 3

Deleted: CH₄ and N₂O emissions from biomass combustion from the Global Fire Emissions Database (GFED; Van Der Werf 2017) were added to EDGAR. ...

Deleted: .

Deleted: uncertainties

Deleted: uses an absolute uncertainty range

Deleted: , both corresponding to a 95% confidence interval

Deleted: 2022

Deleted: aggregate

Deleted: for GHG emissions was computed as the square root of the sum of squared uncertainties for each gas.

Deleted: manuscript

Deleted: their

Deleted: .

Deleted: 2020b

Deleted: GFED for biomass combustion CH₄ and N₂O emissions, and ...

Deleted: 3

There are three reasons for these specific data choices. First, national greenhouse gas emissions inventories tend to use improved, higher-tier methods for estimating emissions fluxes than global inventories such as EDGAR or CEDS (Dhakal et al., 2022; Minx et al., 2021). As GCB and PRIMAP-hist integrate the most recent national inventory submissions to the UNFCCC, selecting these databases makes best use of **country-level improvements in data gathering infrastructures**. Second, comprehensive reporting of F-gas emissions has remained challenging in national inventories **and may exclude some military applications** (see Minx et al., 2021; Dhakal et al., 2022). However, **F-gases are entirely anthropogenic substances, and their concentrations can be measured effectively and reliably in the atmosphere**. We therefore follow the AR6 WGI approach in making use of direct atmospheric observations. Third, the choice of GCB data for CO₂-FFI means we can integrate its projection of that year's CO₂ emissions at the time of publication (i.e., for 2022). No other dataset except GCB provides projections of CO₂ emissions on this timeframe. At this point in the publication cycle (mid-year), the other chosen sources provide data points with a two-year time lag (i.e., for 2021). While these data choices inform our overall assessment of GHG emissions, we provide a comparison across datasets for each emissions category, as well as between our estimates and an estimate derived from AR6 WGIII-like databases (i.e., EDGAR for CO₂-FFI and non-CO₂ GHG emissions, GCB for CO₂-LULUCF).

Deleted: the investments countries have made into

Deleted: as global F-gas concentrations

Deleted: they

530 2.2 Updated global greenhouse gas emissions

Total global GHG emissions reached **55 ± 5.2 GtCO₂e** in 2021. The main contributing sources were CO₂-FFI (37 ± 3 GtCO₂), CO₂-LULUCF (3.9 ± 2.8 GtCO₂), CH₄ (8.9 ± 2.7 GtCO₂e), N₂O (2.9 ± 1.8 GtCO₂e) and F-gas emissions (2 ± 0.59 GtCO₂e). GHG emissions rebounded in 2021, following a single year decline during the COVID-19 induced lockdowns of 2020. Prior to this event in 2019, emissions were **55 ± 5.4 GtCO₂e** - i.e. almost the same level as in 2021. Initial projections indicate that CO₂ emissions from fossil fuel and industry and land use change remained **similar** in 2022, at 37 ± 3 GtCO₂ and 3.9 ± 2.8 GtCO₂, respectively (Friedlingstein et al., 2022a). Note that ODS-F-gases such as chlorofluorocarbons and hydrochlorofluorocarbons are excluded from national GHG emissions inventories. For consistency with AR6, they are also excluded here. Including them here would increase total global GHG emissions by 1.6Gt GtCO₂e in 2021.

Deleted: 57

Deleted: 6

Deleted: 11 ± 3.

Deleted: 3.3 ±

Deleted: significant

Deleted: 57

Deleted: 7

Deleted: stable

Deleted: 2022

Average GHG emissions for the decade 2012-2021 were **54 ± 5.3 GtCO₂e**. Average decadal GHG emissions have increased steadily since the 1970s across all major groups of GHG, driven primarily by increasing CO₂ emissions from fossil fuel and industry, but also rising emissions of CH₄ and N₂O. UNFCCC F-gas emissions have grown more rapidly than other greenhouse gases reported under the UNFCCC, but from low levels. By contrast, ODS F-gas emissions have declined substantially since the 1990s. Both the magnitude and trend of CO₂ emissions from land use change remain highly uncertain, with the latest data indicating **an average** net flux between 4-5 GtCO₂/yr for the past few decades.

Deleted: 56

Deleted: 6


Deleted: a relatively stable

AR6 WGIII reported total net anthropogenic emissions of 59 ± 6.6 GtCO₂e in 2019, and decadal average emissions of 56 ± 6.0 GtCO₂e from 2010-2019. By comparison, our estimates here for the AR6 period sum to 55 ± 5.4 GtCO₂e in 2019, and 53 ± 5.3 GtCO₂e for the same decade (2010-2019). The difference between these figures, including the reduced relative uncertainty range, is partly driven by the substantial revision in GCB CO₂-LULUCF estimates between the 2020 version (used in AR6 WGIII) of 6.6 GtCO₂ and the 2022 version (used here) of 4.6 GtCO₂. The main reason for this downward revision comes from updated estimates of agricultural areas by the FAO and uses multi-annual land-cover maps from satellite remote sensing, leading to lower emissions from cropland expansion, particularly in the tropical regions. It is important to note that this change is not a reflection of changed and improved methodology per se, but an update of the resulting estimation, due to updates in the available input data. Second, there are relatively small changes resulting from improvements in datasets since AR6, with the direction of changes depending on the considered gases. CH₄ accounts for the largest of these at -1.8 GtCO₂e in 2019, which is related to the switch from EDGAR in AR6 to PRIMAP-hist in this study. EDGAR estimates considerably higher CH₄ emissions - from fugitive fossil sources, as well as the livestock, rice cultivation and waste sectors - compared to country reported data using higher tier methods, as compiled in PRIMAP-hist. Generally, uncertainty in these sectors is relatively high as calculations are based on activity data and assumed emissions factors which are hard to determine and vary greatly over countries. Differences in the remaining gases for 2019 are relatively small in magnitude (increases: N₂O (+0.18 GtCO₂e), UNFCCC-F-gases (+ 0.48 GtCO₂e); and decreases: CO₂-FFI (-0.8 GtCO₂e)). Overall, excluding the change due to CO₂-LULUCF and CH₄, they impact the total GHG emissions estimate by -0.14 GtCO₂e.


New literature not available at the time of the AR6 suggests that increases in atmospheric methane concentrations are also driven by methane emissions from wetland changes resulting from climate change (e.g. Basu et al., 2022; Peng et al., 2022; Nisbet et al., 2023; Zhang et al., 2023). Such carbon cycle feedbacks are not taken into account here, as we focus on estimates of emissions resulting directly from human activities.

- Deleted: 57
- Deleted: 7
- Deleted: 55
- Deleted: 6
- Deleted: is primarily
- Deleted: :
- Deleted: an
- Deleted: estimation
- Deleted: adjustment
- Deleted: methodology
- Deleted: changes
- Deleted: smaller differences in the estimates from the described use of better datasets
- Deleted: However, as shown in Figure 1 below
- Deleted: and the direction of these differences depend on the gases considered
- Deleted: 5
- Deleted: CH₄ (-0.001 GtCO₂e),
- Deleted: , this only impacts
- Deleted: 19


(a) Global total greenhouse gas emissions




(b) Global CO₂ emissions from fossil fuel & industry (FFI)




(c) Global CO₂ emissions from land use change (LULUCF)




(d) Global CH₄ emissions




(e) Global N₂O emissions




(f) Global F-gas emissions




(a) Global total greenhouse gas emissions



(c) Global CO₂ emissions from land use change (LULUCF)



(e) Global N₂O emissions



Deleted:

Deleted: Not shown in panels d and e are biomass combustion emissions from GFED (Van Der Werf 2017), which are included in the aggregate estimate in panel a.

605

Figure 1: Annual global anthropogenic greenhouse gas emissions by source, 1970-2021. Refer to Sect. 2.1 for a list of datasets. Starred datasets (*) indicate the sources used to compile global total greenhouse gas emissions in panel a. CO₂ equivalent emissions in panels a and f are calculated using GWPs with a 100-year time horizon from the AR6 WGI Chapter 7 (Forster et al., 2021). F-gas emissions in panel a) comprise only UNFCCC F-gas emissions (see Sect. 2.1 for a list of species).

610

Table 1: Global anthropogenic greenhouse gas emissions by source and decade.

| Gt CO ₂ e | 1970-1979 | 1980-1989 | 1990-1999 | 2000-2009 | 2010-2019 | 2020-2021 | 2021 | 2022 (projection) |
|-------------------------|-----------|-----------|-----------|-----------|-----------|-----------|---------|-------------------|
| GHG | 30±4 | 35±4.4 | 39±4.9 | 45±5.1 | 53±5.3 | 54±5.3 | 55±5.2 | |
| CO ₂ -FFI | 17±1.4 | 20±1.6 | 24±1.9 | 29±2.3 | 36±2.8 | 36±2.9 | 37±3 | 37±3 |
| CO ₂ -LULUCF | 4.4±3.1 | 4.8±3.4 | 5.3±3.7 | 5±3.5 | 4.7±3.3 | 4.5±3.2 | 3.9±2.8 | 3.9±2.8 |
| CH ₄ | 6.2±1.9 | 6.6±2 | 7.3±2.2 | 8±2.4 | 8.6±2.6 | 8.7±2.6 | 8.9±2.7 | |
| N ₂ O | 1.9±1.1 | 2.1±1.3 | 2.2±1.3 | 2.4±1.5 | 2.7±1.6 | 2.8±1.7 | 2.9±1.8 | |
| UNFCCC F-gases | 0.58±0.17 | 0.78±0.23 | 0.77±0.23 | 1±0.3 | 1.5±0.46 | 1.7±0.5 | 2±0.59 | |

620 Notes: All numbers refer to decadal averages, except for annual estimates in 2021 and 2022. CO₂ equivalent emissions are calculated using GWP with a 100-year time horizon from AR6 WGI Chapter 7 (Forster et al., 2021). Projections of non-CO₂ GHG emissions in 2022 remain unavailable at the time of publication. Uncertainties are ±8% for CO₂-FFI, ±70% for CO₂-LULUCF, ±30% for CH₄ and F-gases, and ±60% for N₂O, corresponding to a 90% confidence interval. ODS F-gases are excluded, as noted in Sect. 2.1.

2.3 Non-methane short lived climate forcers

625 In addition to GHG emissions, we provide an update of anthropogenic emissions of non-methane short-lived climate forcers (SLCFs) (SO₂, BC, OC, NO_x, VOCs, CO and NH₃). HFCs are considered in Sect. 2.2. Updating emissions of many short-lived climate forcing agents to 2022 based on established datasets is not possible as compiling global data can take several years. Yet, as SLCF emissions are needed in this paper to update effective radiative forcing (ERF) estimates through 2022, updated emission datasets, where they are available, are combined with projected data to make SLCF emission time series complete.

630

Inserted Cells

Deleted: 31...4.42 (... [3])

Deleted: 36...4.96 (... [4])

Deleted: 41

Deleted: 47

Deleted: 56...5.36 (... [5])

Deleted: 57...5.26 (... [6])

Inserted Cells

Inserted Cells

Deleted: LUCF

Deleted: 9±...± (... [7])

Deleted: 6...2.23 (... [8])

Deleted: 4...2.45 (... [9])

Deleted: 9...2.67 (... [10])

Deleted: 10±3.1

Deleted: 11±3.

Deleted Cells

Inserted Cells

Inserted Cells

Deleted: 2...9±1.13 (... [11])

Deleted: 4

Deleted: 6...1.65 (... [12])

Deleted: 3.1±1

Deleted: 3.3±2

Deleted Cells

Inserted Cells

Inserted Cells

Deleted: 7...0.465 (... [13])

Inserted Cells

680 As in Dhakal et al. (2022), sectoral emissions of SLCFs are derived from two sources. For fossil fuel, industrial, waste and agricultural sectors, we use the CEDS dataset that provided SLCF emissions for CMIP6 (Hoesly et al., 2018). CEDS provides global emissions totals from 1750 to 2019 in its most recent version (O'Rourke et al., 2021). No CEDS emissions data is yet available beyond 2019. As a first estimate, the SLCF emissions time series are extrapolated to 2022 using the “two-year blip” scenario (Forster et al., 2020) of global emissions suppressed by the economic slowdown due to COVID-19. These projections are proxy estimates from Google and Apple mobility data over 2020, and assume a slow return to pre-pandemic emissions activity levels by 2022. Other near-real time emissions estimates covering the COVID-19 pandemic era tend to show less of an emissions reduction than the two-year blip scenario (Guevara et al., 2023). It should be stressed that accurate quantification of SLCF emissions during this period is not possible.

Deleted: emission

690 We do not explicitly account for the introduction of strict fuel sulphur controls brought in by the International Maritime Organization on 1 January 2020, which was expected to reduce SO₂ emissions from the global shipping sector by 8.5 Tg against a pre-COVID baseline (around 10% of 2019 total SO₂ emissions). SO₂ reductions from shipping are partly accounted for in the proxy activity dataset, and including a specific shipping adjustment may double-count emissions reductions.

695 For biomass burning SLCF emissions we follow AR6 WGIII (Dhakal et al., 2022) and use the Global Fire Emissions Dataset (GFED; Randerson et al., 2017) for 1997 to 2022, with the dataset extended back to 1750 for CMIP6 (van Marle et al., 2017). Estimates from 2017 to 2022 are provisional. The potential for both sources of emissions data to be updated in future versions exist, particularly in light of a forthcoming update to CEDS and quantification of shipping sector SO₂ reductions. Other natural emissions, which are important for gauging some SLCF concentrations, are considered as constant in the context of calculating concentrations and ERF.

Deleted:

Deleted: .

700 Estimated emissions used here are based on a combination of GFED emissions for biomass-burning emissions and CEDS up until 2019 extended with the “two-year blip” scenario for fossil, agricultural, industrial and waste sectors. Under this scenario, emissions of all SLCFs are reduced in 2022 relative to 2019 (Table 2). As described in Sect. 4, this has implications for several categories of anthropogenic radiative forcing. Trends in SLCFs emissions are spatially heterogeneous (Szopa et al., 2021) with strong shifts in the geographical distribution of emissions over the 2010-2019 decade. Very different lockdown measures have been applied for COVID around the world resulting in various length and intensity of activity reductions and effect on air pollutant emissions (Sokhi et al., 2021). SLCF emissions have been seen to return to their pre-COVID levels by 2022 in some regions, sometimes with rebound effect, but not in all (Putaud et al., 2023, Lonsdale and Sun, 2023) but quantification at the global scale is not yet available.

Deleted:

Uncertainties ~~associated with these emission estimates~~ are difficult to quantify. From the non-biomass burning sectors they are estimated to be smallest for SO₂ (±14%), largest for ~~black carbon (BC)~~ (a factor of two), and intermediate for other species (Smith et al., 2011; Bond et al., 2013; Hoesly et al., 2018). Uncertainties are also likely to increase ~~both~~ backwards in time (Hoesly et al., 2018), and ~~again~~ in the most recent years. The estimates of non-biomass burning emissions for 2020, 2021 and 720 2022 are highly uncertain owing to the use of proxy activity data, ~~scenario extension, and the impact of sulphur controls in the shipping sector.~~ Future updates of CEDS are expected to include uncertainties (Hoesly et al., 2018). Even though trends over recent years are uncertain, the general decline in ~~some~~ SLCF emissions derived is supported by aerosol optical depth measurements (e.g. Quaas et al., 2022).

Deleted: in the emissions underlying data

Deleted: and

Deleted: . We do not provide a formal assessment of emissions uncertainty here as uncertainties in underlying datasets are not routinely quantified.

Table 2: Emissions of the major SLCFs in 1750, 2019 and 2022

| Compound Species | 1750 emissions (Tg yr ⁻¹) | 2019 emissions (Tg yr ⁻¹) | 2022 emissions (Tg yr ⁻¹) |
|--|--|--|--|
| Sulphur dioxide (SO ₂) + sulphate (SO ₄) | 0.3 | 85.9 | 76.9 |
| Black carbon (BC) | 2.1 | 7.8 | 6.7 |
| Organic carbon (OC) | 15.4 | 34.7 | 26.0 |
| Ammonia (NH ₃) | 6.6 | 66.5 | 65.3 |
| Oxides of nitrogen (NO _x) | 19.4 | 142.9 | 131.8 |
| Volatile organic compounds (VOCs) | 60.6 | 227.2 | 189.6 |
| Carbon monoxide (CO) | 348.4 | 937.8 | 764.1 |

Notes. Emissions of SO₂ + SO₄ use SO₂ molecular weights. Emissions of NO_x use NO₂ molecular weights. VOCs are for the total mass.

3 Well-mixed greenhouse gas concentrations

AR6 Working Group I assessed well mixed GHG concentrations in Chapter 2 (Gulev et al., 2021) and additionally provided a dataset of concentrations of 52 well-mixed GHGs from 1750 to 2019 in its Annex III (IPCC, 2021c). Footnotes in AR6 SYR updated CO₂, CH₄ and N₂O concentrations to 2021 (Lee et al., 2023). In this update we extended the record to 2022 for all 52 gases.

Ozone is an important greenhouse gas with strong regional variation both in the stratosphere and troposphere (Szopa et al., 2021). Its ERF arising from its regional distribution is assessed in Sect. 4, but following AR6 convention is not included with the GHGs discussed here. Other non-methane SLCFs are heterogeneously distributed in the atmosphere and are also not typically reported in terms of a globally averaged concentration. Globally averaged concentrations for these are normally model derived, supplemented by local monitoring networks and satellite data (Szopa et al., 2021).

Deleted: 1

755 As in AR6, CO₂ concentrations are taken from the NOAA Global Monitoring Laboratory (GML) and updated through 2022
(Lan et al., 2023a). Although, here CO₂ is reported on the updated WMO-CO₂-X2019 scale, whereas in AR6, values were
reported on the WMO-CO₂-X2007 scale. This improved calibration increases CO₂ concentrations by around 0.2 ppm (Hall et
al., 2021). In AR6, CH₄ and N₂O were reported as the average from NOAA and AGAGE global networks. For 2022 as updated
AGAGE data is not currently available, we used only NOAA data [Lan et al., 2023b], and multiplied N₂O by 1.0007 to be
760 consistent with a NOAA-AGAGE average. NOAA CH₄ in 2022 was used without adjustment since the NOAA and AGAGE
global means CH₄ are consistent within 2 ppb. Mixing ratio uncertainties for 2022 are assumed to be similar to 2019, and we
adopt the same uncertainties as assessed in AR6 WGI.

Many halogenated greenhouse gases are reported on a global mean basis from NOAA and/or AGAGE until 2020 or 2021 (SF₆
765 is available in the NOAA dataset up to 2022). Where both NOAA and AGAGE data are used for the same gas, we take a mean
of the two datasets. Where both networks are used and the last full year of data availability is different, the difference between
the dataset mean and the dataset with the longer time series in this last year is used as an additive offset to the dataset with the
longer time series. Some obvious inconsistencies are removed such as sudden changes in concentrations when missing data is
reported as zero.

770 Some of the more minor halogenated gases are not part of the NOAA or AGAGE operational network and are currently only
reported in literature sources until 2019, or possibly 2015 (Droste et al., 2020; Laube et al., 2014; Schoenenberger et al., 2015;
Simmonds et al., 2017; Vollmer et al., 2018). Concentrations of gases where 2022 data is not yet available are extrapolated
forwards to 2022 using the average growth rate over the last 5 years of available data. These assumptions have an imperceptible
775 effect on the total ERF assessed in Sect. 4, whereas excluding these gases would have an impact.

780 The global surface mean mixing ratios of CO₂, CH₄ and N₂O in 2022 were 417.1 [± 0.4] ppm, 1911.9 [± 3.3] ppb and 335.9 [± 0.4] ppb. Concentrations of all three major GHGs have increased from 2019 values reported in AR6 WGI, which were 410.1 [± 0.36] ppm for CO₂, 1866.3 [± 3.2] ppb for CH₄ and 332.1 [± 0.7] ppb for N₂O. CO₂ concentrations in 2019 are updated to 410.3 ppm using the new WMO-CO₂-X2019 scale adopted here. Concentrations of most categories of halogenated GHGs have increased from 2019 to 2022, from 109.4 to 114.2 ppt on a CF₄-equivalent scale for PFCs; 237.1 ppt to 287.2 ppt on an HFC-134a-equivalent scale for HFCs; 9.9 ppt to 11.0 ppt for SF₆ and 2.1 to 2.8 ppt for NF₃. Only Montreal Protocol halogenated GHGs have decreased in concentration, from 1031.9 ppt in 2019 to 1016.6 ppt in 2022 on a CFC-12-equivalent scale, demonstrating the continued success of the Montreal Protocol. Although even here, concentrations of some minor CFCs are rising (see also Western et al. 2023). In this update we employ AR6 derived uncertainty estimates and do not perform a new assessment. Table 1 in Sect. 3 of the Supplementary Material shows specific updated concentrations for all the GHGs considered.

4 Effective Radiative Forcing (ERF)

- Deleted: 4
- Deleted: (table 3).
- Deleted: Note, AR6 SYR quoted updated values but to less precision. These ...
- Deleted: not given
- Deleted: to avoid the perception of an inconsistency.
- Deleted: ,
- Deleted: eq
- Deleted: ,
- Deleted: eq
- Deleted: ,
- Deleted: ¶
- Table 3: Annual mean concentrations of well-mixed greenhouse gases in 2022, 2019, 1850 and 1750. Except for CO₂, CH₄ and N₂O,...
- Deleted: are in parts per trillion by volume [ppt].

805

Moved (insertion) [1]

ERFs were principally assessed in Chapter 7 of AR6 Working Group I (Forster et al., 2021). Chapter 7 focussed on assessing ERF from changes in atmospheric concentrations, it also supported estimates of ERF in Chapter 6 that attributed forcing to specific precursor emissions (Szopa et al., 2021) and also generated the time history of ERF shown in AR6 WGI Figure 2.10 and discussed in Chapter 2 (Gulev et al., 2021). Only the concentration based estimates are updated this year. The emission

810

Moved (insertion) [2]

based estimates relied on specific chemistry climate model integrations and a consistent method of applying updates to these would need to be developed in the future.

815

Each IPCC report has successively updated both the method of calculation and the time history of different warming and cooling contributions, measured as ERFs. Both types of updates have contributed to a significantly changed forcing estimate between successive reports. For example, Forster et al., (2021) updated the methodology to exclude land-surface temperature related adjustments from the forcing calculation, which generally increased estimates. At the same time GHG levels increased and the time history of aerosol forcing was revised, overall leading to a higher total ERF estimate in AR6 compared to AR5. These IPCC updates flow from an assessment of varied literature and also rely on updates to concentrations and/or emissions.

820

There is no published regularly updated total ERF indicator outside of the IPCC process, although the European Copernicus programme has trialed such a product (Bellouin et al., 2020). For radiative forcing, NOAA annually updates estimates for the main GHGs, calculating radiative forcing (RF) using the set of formulas to estimate RFs from concentrations (Montzka, 2022). Updated RF formulas were employed in AR6 (Forster et al., 2021) and these updated expressions are also employed here in Supplementary Material, Sect. 4.

825

Moved (insertion) [3]

The ERF calculation follows the methodology used in AR6 WGI (Smith et al., 2021). For each category of forcing, a 100,000 member probabilistic Monte Carlo ensemble is sampled to span the assessed uncertainty range in each forcing. All uncertainties are reported as 5-95% ranges and provided in square brackets. The only significant methodological change compared to AR6 is for the volcanic ERF estimate. Firstly, The preindustrial baseline data has been improved by switching to a new longer record of stratospheric aerosol optical depth before 1750 (Sigl et al, 2022). Secondly, choices have also been made to include the January 2022 eruption of Hunga Tonga-Hunga Ha'apai as an exceptional positive ERF perturbation from the increase in stratospheric water vapour (Millan et al., 2022; Sellito et al., 2022; Jenkins et al., 2023). The methods are all detailed in Supplementary Material, Sect. 4.

830

Moved (insertion) [4]

835 The summary results for the anthropogenic constituents of ERF and solar irradiance in 2022 relative to 1750 are shown in Figure 2a. In Table 3, these are summarised alongside the equivalent ERFs from AR6 (1750-2019) and AR5 (1750-2011). Figure 2b shows the time evolution of ERF from 1750 to 2022.

Moved (insertion) [5]

Moved (insertion) [6]

840 Total anthropogenic ERF has increased to 2.91 [2.19 to 3.63] W m⁻² in 2022 relative to 1750, compared to 2.72 [1.96 to 3.48] W m⁻² for 2019 relative to 1750 in AR6. The main contributions to this increase are from increases in greenhouse gas concentrations and a reduction in the magnitude of aerosol forcing. Decadal trends in ERF have increased markedly and are now over 0.6 W m⁻² per decade. These are discussed further in the discussion and conclusions (Sect. 12).

845 The ERF from well-mixed GHGs is 3.45 [3.14 to 3.75] W m⁻² for 1750-2022, of which 2.25 W m⁻² is from CO₂, 0.56 W m⁻² from CH₄, 0.22 W m⁻² from N₂O and 0.41 W m⁻² from halogenated gases. This is an increase from 3.32 [3.03 to 3.61] W m⁻² for 1750-2019 in AR6. ERFs from CO₂, CH₄ and N₂O have all increased since the AR6 WG1 assessment for 1750-2019 owing to increases in atmospheric concentrations.

850 **The total aerosol ERF (sum of the ERF from aerosol radiation interactions (ERF_{ari}) and aerosol cloud interactions (ERF_{aci})) for 1750-2022 is -0.98 [-1.58 to -0.40] W m⁻² compared to -1.06 [-1.71 to -0.41] W m⁻² assessed for 1750-2019 in AR6 WG1.**

Moved (insertion) [7]

855 This continues a trend of weakening aerosol forcing due to reductions in precursor emissions. The majority of this reduction is from ERF_{aci} which is determined to be -0.77 [-1.33 to -0.23] W m⁻² compared to -0.84 [-1.45 to -0.25] W m⁻² in AR6 for 1750-2019. ERF_{ari} for 1750-2022 is -0.21 [-0.42 to 0.00] W m⁻², marginally weaker than the -0.22 [-0.47 to 0.04] W m⁻² assessed for 1750-2019 in AR6 WG1 (Forster et al., 2021). The largest contributions to ERF_{ari} are from SO₂ (primary source of sulphate aerosol; -0.21 W m⁻²), BC (+0.12 W m⁻²), OC (-0.04 W m⁻²) and NH₃ (primary source of nitrate aerosol; -0.03 W m⁻²). ERF_{ari} is not weakening as fast as ERF_{aci} due to reductions in the warming influence of BC cancelling out some of the reduced sulphate cooling. ERF_{ari} also includes terms from CH₄, N₂O and NH₃ which are small but have all increased.

Moved (insertion) [8]

860 Ozone ERF is determined as 0.48 [0.24 to 0.72] W m⁻² for 1750-2022, similar to the AR6 assessment of 0.47 [0.24 to 0.71] W m⁻² for 1750-2019. Land use forcing and stratospheric water vapour from methane oxidation are unchanged (to two decimal places) since AR6. The decline in BC emissions from 2019 to 2022 has reduced ERF from light absorbing particles on snow and ice from 0.08 [0.00 to 0.18] W m⁻² for 1750-2019 to 0.06 [0.00 to 0.14] W m⁻² for 1750-2022. We determine from provisional data that aviation activity in 2022 had not yet returned to pre-COVID levels. Therefore, ERF from contrails and contrail-induced cirrus is lower than AR6, at 0.05 [0.02 to 0.09] W m⁻² in 2022 compared to 0.06 [0.02 to 0.10] W m⁻² in 2019.

865

The headline assessment of solar ERF is unchanged, at 0.01 [-0.06 to +0.08] W m⁻² from pre-industrial to the 2009-2019 solar cycle mean. Separate to the assessment of solar forcing over complete solar cycles, we provide a single year 2022 solar ERF of 0.06 [-0.02 to +0.14] W m⁻². This is higher than the single year estimate of solar ERF for 2019 (a solar minimum) of -0.02 [-0.08 to 0.06] W m⁻².


870

For volcanic ERF, updating of the pre-industrial dataset for stratospheric aerosol optical depth (sAOD) increased the sAOD over 500 BCE to 1749 CE, resulting in a larger difference to post-1750 sAOD and resulting in a volcanic ERF difference of +0.015 W m⁻² compared to AR6 (see Sect. 4 in the Supplementary Material). In addition, the earlier Holocene was more volcanically active than the period after 500 BCE, further increasing the mean sAOD baseline. Taking the longer baseline period into account in the new preindustrial dataset, post-1750 ERF is further increased by 0.031 W m⁻².

875

The net effect is that volcanic forcing after 1750 has increased by +0.046 W m⁻² compared to AR6 due to dataset updates and by account of the fact that the post-1750 period was less volcanically active on average than the early Holocene which is now used in the ERF calculation.

Moved (insertion) [9]



880




Figure 2: Effective radiative forcing from 1750-2022. (a) 1750-2022 change in ERF, showing best estimates (bars) and 5-95% uncertainty ranges (lines) from major anthropogenic components to ERF, total anthropogenic ERF, and solar forcing. (b) Time evolution of ERF from 1750 to 2022. Best estimates from major anthropogenic categories are shown along with solar and volcanic forcing (thin coloured lines), total (thin black line) and anthropogenic total (thick black line). 5-95% uncertainty in the anthropogenic forcing is shown in shaded grey. Note solar forcing in 2022 is a single-year estimate.

Table 3: Contributions to anthropogenic effective radiative forcing (ERF) for 1750-2022 assessed in this section.

| Forcer | 1750-2022 | 1750-2019 (AR6) | 1750-2011(AR5) | Reason for change from AR6 |
|------------------|-------------------|-------------------|-------------------|---------------------------------|
| | W m ⁻² | W m ⁻² | W m ⁻² | |
| CO ₂ | 2.25 | 2.16 | 1.82 | |
| | [1.98 to 2.52] | [1.90 to 2.41] | [1.63 to 2.01] | Increases in GHG concentrations |
| CH ₄ | 0.56 | 0.54 | 0.48 | |
| | [0.45 to 0.67] | [0.43 to 0.65] | [0.43 to 0.53] | |
| N ₂ O | 0.22 | 0.21 | 0.17 | |
| | [0.19 to 0.25] | [0.18 to 0.24] | [0.14 to 0.20] | |

Moved (insertion) [10]

Moved down [11]: Greenhouse gas

Deleted: 1850

Deleted: 1750

Deleted: CO₂ [ppm] ... [14]

Deleted: NF₃

Deleted: 7

Deleted: 0.0

Deleted: 0.0

Merged Cells

Deleted: 11.

Deleted: SF₆

Deleted: 9.9

Deleted: 0.0

Deleted: SO₂F₂

Deleted: 2.8

Deleted: 2.5

Deleted: 0.0

Deleted: HFCs as HFC-134a-eq ... [15]

| | | | | |
|---|-----------------|-----------------|-----------------|---|
| Halogenated GHGs | 0.41 | 0.41 | 0.36 | |
| | [0.33 to 0.49] | [0.33 to 0.49] | [0.32 to 0.40] | |
| Ozone | 0.48 | 0.47 | 0.35 | Changes in precursor emissions and chemically active GHGs; net effect almost cancels. |
| | [0.24 to 0.72] | [0.24 to 0.71] | [0.21 to 0.67] | |
| Stratospheric water vapour | 0.05 | 0.05 | 0.07 | |
| | [0.00 to 0.10] | [0.00 to 0.10] | [0.02 to 0.12] | |
| Aerosol-radiation interactions | -0.21 | -0.22 | -0.45 | Reduction in aerosol and aerosol precursor emissions. |
| | [-0.42 to 0.00] | [-0.47 to 0.04] | [-0.95 to 0.05] | |
| Aerosol-cloud interactions | -0.77 | -0.84 | -0.45 | |
| | [-1.33 to 0.23] | [-1.45 to 0.25] | [-1.2 to 0.0] | |
| Land use | -0.20 | -0.20 | -0.15 | |
| | [-0.30 to 0.10] | [-0.30 to 0.10] | [-0.25 to 0.05] | |
| Light-absorbing particles on snow and ice | 0.06 | 0.08 | 0.04 | Reduction in BC emissions. |
| | [0.00 to 0.14] | [0.00 to 0.18] | [0.02 to 0.09] | |
| Contrails and aviation-induced cirrus | 0.05 | 0.06 | 0.05 | As of 2022, global aviation activity has not yet returned to pre-COVID19 levels. |
| | [0.02 to 0.09] | [0.02 to 0.10] | [0.02 to 0.15] | |
| Total anthropogenic | 2.91 | 2.72 | 2.3 | Increase in GHG concentrations and reduction in aerosol emissions. |
| | [2.19 to 3.63] | [1.96 to 3.48] | [1.1 to 3.3] | |
| Solar irradiance | 0.01 | 0.01 | 0.05 | |
| | [-0.06 to 0.08] | [-0.06 to 0.08] | [0.0 to 0.10] | |

All values are in $W m^{-2}$ and 5-95% ranges are in square brackets. As a comparison, the equivalent assessments from AR6 (1750-2019) and AR5 (1750-2011; Myhre et al., 2013b) are shown. Solar ERF is included and unchanged from AR6, based on the most

- Deleted: HFC-23
- Deleted: 36.1
- Deleted: 32.5
- Deleted: 0.0
- Deleted: HFC-32 ... [16]
- Deleted: .
- Deleted: HFC-143a
- Deleted: 28.9
- Deleted: 0.0
- Deleted: HFC-152a
- Deleted: 7.5
- Deleted: 7.2
- Deleted: 0.0
- Merged Cells ... [17]
- Deleted: HFC-227ea
- Deleted: 2.1
- Deleted: 1.6
- Deleted: 0.0
- Deleted: HFC-236fa ... [18]
- Deleted: HFC-365mfc
- Deleted: 2
- Deleted: 1
- Deleted: 0.0
- Merged Cells ... [19]
- Deleted: HFC-43-10mee
- Deleted: 3
- Deleted: 3
- Deleted: 0.0
- Deleted: 34.
- Deleted: PFCs as CF₄-eq
- Deleted: 114.2
- Deleted: 109.4
- Deleted: 34.0
- Deleted: 34.
- Deleted: CF₄
- Deleted: 88.4
- Deleted: 85.6
- Deleted: 34.0
- Deleted: C₂F₆ ... [20]
- Deleted: 4
- Deleted: 3
- Deleted: CFC-13
- Deleted: 0.0
- Deleted: 0.0
- Deleted: CFC-113
- Deleted: 68.2
- Deleted: 69.8
- Deleted: 0.0
- Moved up [1]: [y](#)
- Moved up [2]: 2021). Only the concentration based estimates are
- Moved up [3]: [f](#)
- Moved up [4]:). For each category of forcing, a 100,000 member
- Moved up [5]: The summary results for the anthropogenic
- Moved up [6]: these are summarised alongside the equivalent

recent solar cycle (2009-2019) thus differing from the single-year estimate in Fig. 2a. Volcanic ERF is excluded due to the sporadic nature of eruptions.

5. Global surface temperature

AR6 WGI Chapter 2 assessed the 2001-2020 globally averaged surface temperature change above an 1850-1900 baseline to be 0.99 [0.84 to 1.10] °C and 1.09 [0.95 to 1.20] °C for 2011-2020 (Gulev et al., 2021). Updated estimates to 2022 were also given in AR6 SYR (Lee et al., 2023). The AR6 SYR estimates match those given here. We describe the update in detail and provide further quantification and comparisons.

There are choices around the methods used to aggregate surface temperatures into a global average, how to correct for systematic errors in measurements, methods of infilling missing data, and whether surface measurements or atmospheric temperatures just above the surface are used. These choices, and others, affect temperature change estimates and contribute to uncertainty (IPCC AR6 WGI Chapter 2, Cross Chapter Box 2.3, Gulev et al., 2021). The methods chosen here closely follow AR6 WGI and are presented in Supplementary Material, Sect. 5. Confidence intervals are taken from AR6 as only one of the employed datasets regularly updates ensembles (see Supplementary Material, Sect. 5).

Based on the updates available as of February 2023 (which were reported in the AR6 SYR), the change in global surface temperature from 1850-1900 to 2013-2022, using the same underlying data sets and methodology as AR6, is 1.15 [1.00-1.25] °C, an increase of 0.06 °C within two years from the 2011-2020 value reported in the AR6 Working Group I report (Table 4). The change from 1850-1900 to 2003-2022 was 1.03 [0.87-1.13] °C, 0.04 °C higher than the earlier value reported in AR6 WGI. These changes are broadly consistent with typical warming rates over the last few decades, which were assessed in AR6 as 0.76 °C over the 1980-2020 period (using ordinary-least-square linear trends), or 0.019 °C per year (Gulev et al., 2021). They are also broadly consistent with projected warming rates from 2001-2020 to 2021-2040 reported in AR6, which are in the order of 0.025 °C per year under most scenarios (Lee et al., 2021).

Table 4: Estimates of global surface temperature change from 1850-1900 [very likely ranges] for IPCC AR6 and the present study.

| Time Period | Temperature change from 1850-1900 (°C) | |
|-------------|--|------------|
| | IPCC AR6 | This study |
| | | |

Moved (insertion) [12]

Deleted: Table 4: Contributions to anthropogenic effective radiative forcing (ERF) for 1750-2022 assessed in this section.

Moved (insertion) [13]

Moved (insertion) [14]

Deleted: Forcer

Deleted: 1750-2022
W m²

Deleted: 1750-2019 (AR6)
W m²

Deleted: 1750-2011(AR5)
W m²

Deleted: Reason for


Merged Cells

Deleted: AR6

| | | |
|-------------------------------------|---|---|
| <u>Global, most recent 10 years</u> | <u>1.09 [0.95 to 1.20]</u> <u>(to 2011-2020)</u> | <u>1.15 [1.00 to 1.25]</u> <u>(to 2013-2022)</u> |
| <u>Global, most recent 20 years</u> | <u>0.99 [0.84 to 1.10]</u> <u>(to 2001-2020)</u> | <u>1.03 [0.87 to 1.13]</u> <u>(to 2003-2022)</u> |
| <u>Land, most recent 10 years</u> | <u>1.59 [1.34 to 1.83]</u> <u>(to 2011-2020)</u> | <u>1.65 [1.36 to 1.90]</u> <u>(to 2013-2022)</u> |
| <u>Ocean, most recent 10 years</u> | <u>0.88 [0.68 to 1.01]</u> <u>(to 2011-2020)</u> | <u>0.93 [0.73 to 1.04]</u> <u>(to 2013-2022)</u> |

- Deleted: 2.25 []
- Deleted: 98
- Deleted: 2.16 [1.90 to 2.41]
- Deleted: 82 []
- Deleted: 63
- Deleted: Increases in GHG concentrations
- Deleted Cells
- Deleted: CO₂
- Deleted: 2.01
- Deleted: 2.52
- Deleted: 56 []
- Deleted: 45
- Deleted: 54 [0.43]
- Deleted: 0.48 [0.43 to 0.53]
- Deleted: CH₄
- Deleted: 0.67
- Deleted: 0.65
- Deleted: N₂O
- Deleted: 22 []
- Deleted: 19
- Deleted: 21 []
- Deleted: 18
- Deleted: 0.17 [0.14 to 0.20]
- Deleted Cells
- Deleted: 0.25
- Deleted: 0.24
- Deleted: Halogenated GHGs

... [26]



475 **Figure 3. Annual (thin line) and decadal (thick line) means of Global surface temperature (expressed as a change from the 1850-1900 reference period).**

480 **Note that the temperatures for single years include considerable variability and are influenced by natural forcings such as the El Niño-Southern Oscillation and sporadic volcanic eruptions that might either cool or warm the climate for short periods (Jenkins et al., 2023). At current warming rates individual years may exceed warming of 1.5°C several years before a long-term mean exceeds this level (Trewin, 2022).**

Deleted: All values are in $W m^{-2}$ and 5-95% ranges are in square brackets. As a comparison, the equivalent assessments from AR6 (1750-2019) and AR5 (1750-2011; Myhre et al., 2013b) are shown. Solar ERF is included and unchanged from AR6, based on the most recent solar cycle (2009-2019) thus differing from the single-year estimate in Fig. 2a. Volcanic ERF is excluded due to the sporadic nature of eruptions.¶ 5.

Moved up [12]: ¶ There are choices around the methods used to aggregate surface temperatures into a global average, how to correct for systematic errors in measurements, methods of infilling missing data, and whether surface measurements or atmospheric temperatures just above the surface are used. These choices, and others, affect temperature change estimates and contribute to uncertainty (IPCC AR6 WGI Chapter 2, Cross Chapter Box 2).

Moved down [15]: 2021).

Moved up [13]: ¶ Based on the updates available as of February 2023 (which were reported in the AR6 SYR), the change in

Moved up [14]: . These changes are broadly consistent with typical warming rates over the last few decades, which were assessed in AR6 as 0.76 °C over the 1980-2020 period (using ordinary-least-square linear trends), or 0.019 °C per year (Gulev et al., 2021). They are also broadly consistent with projected warming rates from 2001-2020 to 2021-2040 reported in AR6, which are in the order of 0.025 °C per year under most scenarios (Lee et al.,

Deleted: AR6 WGI Chapter 2 assessed the 2001-2020 globally averaged surface temperature change above an 1850-1900 baseline to be 0.99 [0.84 to 1.10] °C and 1.09 [0.95 to 1.20] °C for 2011-2020 (Gulev et al., 2021). Updated estimates to 2022 were also given in AR6 SYR (Lee et al., 2023). The AR6 SYR estimates match those given here. We describe the update in detail and provide further quantification and comparisons.¶

Additionally to IPCC reports, the World Meteorological Organisation (WMO) has developed a set of indicators for global climate monitoring (Trewin et al., 2020) and annually publishes 'state of the climate' reports with a global surface temperature estimate for the year of publication. For example, their 2022 "State of the Global Climate Report" gave an estimated global surface temperature (... [27])

Deleted: I, Gulev et al.,

Deleted: ¶ Surface temperature information on land and sea is available with low latency through WMO distribution channels, with monthly station data from a substantial number of stations reported within a few days of the end of the month. These are consolidated into (... [28])

Deleted: GMST from 1850-1900 to 2013-2022, using the same underlying data sets and methodology as AR6, is 1.15 [1.00-1.25] °C, an increase of 0.06 °C within two years from the 2011-2020 value reported in the AR6 Working Group I report. The change from 1850-1900 to 2003-2022 was 1.03 [0.87-1.13] °C, 0.04 °C higher than the earlier value reported in the AR6 Working Group I report

6. Earth Energy Imbalance

630 The Earth energy imbalance (EEI) assessed in Chapter 7 of AR6 WGI (Forster et al., 2021), provides a measure of accumulated additional energy (heating) in the climate system, and hence plays a critical role in our understanding of climate change. It represents the difference between the radiative forcing acting to warm the climate and Earth's radiative response, which acts to oppose this warming. On annual and longer timescales, the Earth heat inventory changes associated with EEI are dominated by the changes in global ocean heat content (OHC), which accounts for about 90% of global heating since the 1970s (Forster et al., 2021). This planetary heating results in changes to the Earth system such as sea level rise, ocean warming, ice loss, rise in temperature and water vapour in the atmosphere, and permafrost thawing (e.g., Cheng et al., 2022; von Schuckmann et al., 2023a), with adverse impacts for ecosystems and human systems (Douville et al., 2021; IPCC, 2022).

640 On decadal timescales, changes in global surface temperatures (Sect. 5) can become decoupled from EEI by ocean heat rearrangement processes (e.g., Palmer and McNeill, 2014; Allison et al., 2020). Therefore, the increase in the Earth heat inventory provides a more robust indicator of the rate of global change on interannual-to-decadal timescales (Cheng et al., 2019; Forster et al., 2021; von Schuckmann et al., 2023a). AR6 found increased confidence in the assessment of changes in the Earth heat inventory compared to previous IPCC reports due to observational advances and closure of the energy and global sea level budgets (Forster et al., 2021; Fox-Kemper et al., 2021).

645 AR6 estimated with high confidence that EEI increased from 0.50 [0.32-0.69] W m⁻² during the period 1971-2006 to 0.79 [0.52-1.06] W m⁻² during the period 2006-2018 [very likely range] (Forster et al., 2021). The contributions to increases in the Earth heat inventory throughout 1971-2018 remained fairly stable: 91% for the full-depth ocean; 5% for the land; 3% for the cryosphere and about 1% for the atmosphere (Forster et al., 2021). The increase in EEI (Figure 4) has also been reported by (Cheng et al. 2019; von Schuckmann et al., 2020; 2023a; Loeb et al., 2021; Hakuba et al., 2021; Kramer et al., 2021; Raghuraman et al., 2021). Drivers for the most recent period (i.e., past two decades) are both the increases in effective radiative forcing (Sect. 4) as well as climate feedbacks, such as cloud and sea ice changes. The degree of contribution from the different drivers are uncertain and still under active investigation.

655 While changes in EEI have been effectively monitored at the top-of-atmosphere by satellites since the mid-2000s, we rely on estimates of OHC change to determine the absolute magnitude of EEI, and its evolution on inter-annual to multi-decadal time series. The AR6 assessment of ocean heat content change for the 0-2000 m layer was based on global annual mean time series from five ocean heat content datasets: IAP (Cheng et al., 2017); Domingues et al., (2008); EN4 (Good et al., 2013); Ishii et al.,

Moved (insertion) [16]

Deleted: 2021b),¶

Moved (insertion) [17]

Moved (insertion) [18]

Moved (insertion) [19]

Moved (insertion) [20]

Moved (insertion) [21]

Moved (insertion) [22]

Moved (insertion) [23]

Moved (insertion) [15]

Moved (insertion) [24]

Moved (insertion) [25]

660 (2017); NCEI (Levitus et al., 2012). Four of these datasets routinely provide updated OHC time series for the BAMS State of
the Climate report, and all are used for the GCOS Earth heat inventory (von Schuckmann et al., 2020; 2023a) and the annual
WMO global state of the climate. The uncertainty assessment for the 0-2000 m layer used the ensemble method described by
Palmer et al. (2021) that separately accounts for *parametric* and *structural* uncertainty. The >2000 m OHC change and
associated uncertainty was assessed based on trend analysis of the available hydrographic data following Purkey and Johnson
665 (2010). All five of the datasets used for the 0-2000 m OHC assessment are now updated at least annually and should in principle
support an AR6 assessment time series update within the first few months of each year. There is potential to increase the
observational ensemble used in the assessment by supplementing this set with additional data products that are also available
annually for future updates. There is also a potential to update the uncertainty estimate after a more comprehensive
understanding of the error sources.

670 Estimates of EEI should also account for the other elements of the Earth heat inventory, i.e., the atmospheric warming, the
latent heat of global ice loss, and heating of the continental land surface (Forster et al., 2021; Cuesta-Valero et al., 2021; 2022;
Steiner et al., 2020; Nitzbon et al., 2022a; Vanderkelen et al., 2020; Adusumilli et al., 2022). Some of these components of the
Earth heat inventory are routinely updated by a community-based initiative reported in von Schuckmann et al. (2020; 2023a).

675 However, in the absence of annual updates to all heat inventory components, a pragmatic approach is to use recent OHC
change as a proxy for EEI, scaling the value up as required based on historical partitioning between Earth system components.

We carry out an update to the AR6 estimate of changes in the Earth heat inventory based on updated observational *time series
for the period 1971-2020*. Time series of heating associated with loss of ice and warming of the atmosphere and continental
680 land surface are obtained from the recent Global Climate Observing System (GCOS) initiative (von Schuckmann et al., 2023b;
Adusumilli et al., 2022; Cuesta-Valero et al., 2023; Vanderkelen and Thiery, 2022; Nitzbon et al., 2022b; Kirchengast et al.,
2022). We use the original AR6 time series ensemble OHC time series for the period 1971-2018 and then switch to a smaller
four-member ensemble for the period 2019-2022. We “splice” the two sets of time series by adding an offset as needed to
ensure that the 2018 values are identical. The AR6 heating rates and uncertainties for the ocean below 2000 m are assumed to
685 be time-constant. The time-evolution of the Earth heat inventory is determined as a simple summation of time series of:
*atmospheric heating; continental land heating; heating of the cryosphere; and heating of the ocean over three depth layers: 0-
700 m, 700-2000 m, and below 2000 m. While von Schuckmann et al. (2023a) have also quantified heating of permafrost and
inland lakes and reservoirs, these additional terms are very small and are omitted here for consistency with AR6 (Forster et al.,
2021).*

690

Moved (insertion) [26]

Moved (insertion) [27]

Moved (insertion) [28]

Moved (insertion) [29]

Moved (insertion) [30]

A full propagation of uncertainties across all heat inventory components depends on the specific choice of time-period and different estimates are not directly comparable. Therefore we take a simple pragmatic approach, using the total ocean heat content uncertainty as a proxy for the total uncertainty, since this term is two orders of magnitude larger than the other terms (Forster et al., 2021). In order to provide estimates of the EEI up to the year 2022, we scale up the values of OHC change in 2021 and 2022 to reflect the about 90% contribution of the ocean to changes in the Earth heat inventory. The EEI is then simply computed as the difference in global energy inventory over each period, converted to units of $W m^{-2}$ using the surface area of the Earth and the elapsed time. The uncertainties in the global energy inventory for the end-point years are assumed to be independent and added in quadrature, following the approach used in AR6 (Forster et al., 2021).

Table 5: Estimates of the Earth energy imbalance (EEI) for AR6 and the present study.

| Time Period | Earth energy imbalance ($W m^{-2}$) | |
|-------------|---------------------------------------|---------------------|
| | IPCC AR6 | This study |
| 1971-2018 | 0.57 [0.43 to 0.72] | 0.57 [0.43 to 0.72] |
| 1971-2006 | 0.50 [0.32 to 0.69] | 0.50 [0.31 to 0.68] |
| 2006-2018 | 0.79 [0.52 to 1.06] | 0.79 [0.52 to 1.07] |
| 1975-2022 | - | 0.65 [0.48 to 0.81] |
| 2010-2022 | - | 0.89 [0.63 to 1.15] |

In our updated analysis, we find successive increases in EEI for each 20-year period since 1973, with an estimated value of 0.44 [0.05 to 0.83] $W m^{-2}$ during 1973-1992 that almost doubled to 0.82 [0.60 to 1.04] $W m^{-2}$ during 2003-2022 (Figure 4b). In addition, there is some evidence that the warming signal is propagating into the deeper ocean over time, as seen by a robust increase of deep (700-2000m) ocean warming since the 1990s (Cheng et al. 2019, 2022). The model simulations qualitatively agree with the observational evidence (e.g. Gleckler et al., 2016; Cheng et al. 2019), further suggesting more than half of the

Moved (insertion) [31]

Deleted: surface temperature change from 1850-1900 [v... [29]

Moved (insertion) [32]

Deleted: Temperature change from 1850-1900 (°C)

Deleted: Global, most recent 10 years

Deleted: 1.09 [...57 [0.4395...to 0.72][1.20] [30]

Deleted: 1.15 [1.00...to 0.721.25 [31]

Deleted: (to 2013-2022)

Deleted: Global, most recent 20 years

Deleted: 99...[0.3284...to 0.69][1.10] [32]

Deleted: 1.03 [...50 [0.3187...to 0.681.13 [33]

Deleted: (to 2003-2022)

Deleted: Land, most recent 10 years

Deleted: 1.59 [1.34...to 1.06][83] [34]

Deleted: 1.65 [1.36...to 1.0790 [35]

Deleted: (to 2013-2022)

Deleted: Ocean, most recent 10 years


Deleted: 0.88 [0.68 to 1.01] [36]

Deleted: 93...[0.4873...to 0.811.04 [37]

Deleted: (to 2013-2022)

Moved up [16]: 6. Earth Energy Imbalance [38]

Moved up [17]: This planetary heating results in change [39]



Deleted: [38]

Deleted: , 2021).

Deleted: IPCC, 2022). [40]

Moved up [19]: Palmer and McNeal, 2014; Allison et al., 2020).

Deleted: Forster et al.,

Moved up [20]: 2021; von Schuckmann et al., 2023a).

Deleted: AR6 found increased confidence in the assessment [41]

Moved up [21]: 2021; Fox-Kemper et al., 2021). [42]

Deleted: AR6 estimated with high confidence that EEI inc [42]

Moved up [22]: 0.69] $W m^{-2}$ during the period 1971-2006 to 0.79

Deleted: von Schuckmann et al.,

Moved up [23]: 2020; 2023a; Loeb et al., 2021; Hakuba et al.,

Deleted: 2021), and drivers for the most recent period (i.e. [43]

Moved up [24]: [44]

Deleted: top-of-atmosphere by satellites since the mid-200 [44]

OHC increase since the late 1800s occurs after the 1990s. For 1973-1992 the contribution by ocean vertical layer was 66%, 28% and 1% for 0-700 m, 700-2000 m and >2000 m, respectively. During 2013-2022 the corresponding layer contributions were 50%, 33% and 8%.

The update of the AR6 assessment periods to end in 2022 results in systematic increases of EEL of 0.08 W m⁻² for 1975-2022 relative to 1971-2018 and 0.10 W m⁻² for 2010-2022 relative to 2006-2018 (Table 5).




Figure 4: a) Observed changes in the Earth heat inventory for the period 1971-2020 with component contributions as indicated in the figure legend; b) Estimates of the Earth energy imbalance for IPCC AR6 assessment periods, for consecutive twenty-year periods, and the most recent decade. Shaded regions indicate the *very likely* range (90% confidence interval). Data use and approach are based on the AR6 methods, and further described in Sect. 6.

7 Human-induced global warming

Human-induced warming, also known as anthropogenic warming, refers to the component of observed global surface temperature increase over a specific period (for instance, from 1850-1900 as a proxy for pre-industrial climate to the last decade) attributable to

Moved (insertion) [33]

Deleted: Table 6: Estimates of the Earth energy imbalance (EEI) for AR6 and the present study⁴

Moved (insertion) [34]

110 both the direct and indirect effects of human activities, which are typically grouped as follows: well-mixed greenhouse-gases (consisting of CO₂, CH₄, N₂O, and F-gases), and other-human forcings (consisting of aerosol radiation interaction, aerosol cloud interaction, black carbon on snow, contrails, ozone, stratospheric H₂O, and land use) (Eyring et al., 2021). While *total warming*, the actual observed temperature change potentially resulting from both natural climate variability (internal variability of the climate system, and the climate response to natural forcing) and human influences, is the quantity directly related to climate impacts and therefore relevant for adaptation, mitigation efforts focus on human-induced warming as the more relevant indicator for tracking progress against climate stabilisation targets. Further, as the attribution analysis allows to disentangle human-caused warming from possible contributions from solar and volcanic forcing and internal variability (e.g. related to El Niño/La Niña events), it avoids misperception about short-term fluctuations in temperature

115 An assessment of human-induced warming was therefore provided in two reports within the IPCC's 6th assessment cycle: first in SR1.5 in 2018 (Chapter 1 Sect. 1.2.1.3 and Figure 1.2 (Allen et al., 2018), summarised in SPM A.1 and Figure SPM.1 (IPCC, 2018)) and second in AR6 in 2021 (WGI Chapter 3 Sect. 3.3.1.1.2 and Figure 3.8 (Eyring et al., 2021), summarised in WGI SPM A.1.3 and Figure SPM.2 (IPCC, 2021b)).

Moved (insertion) [35]

Moved (insertion) [36]

Moved (insertion) [37]

7.1.1 Warming Period Definitions in the IPCC Sixth Assessment Cycle

120 AR6 defined the current human-induced warming relative to the 1850-1900 baseline as the decade-average of the previous 10-year period (see AR6 WGI Chapter 3). This paper provides an update of the 2010-2019 period used in the AR6 to the 2013–2022 decade. SR1.5 defined current human-induced warming as the average of a 30-year period centred on the current-year assuming the recent rate of warming continues (see SR1.5 Chapter 1). This definition is currently almost identical to the present-day single-year value of human-induced warming, differing by about 0.01°C (see results in Sect. 7.4); the attribution assessment in SR1.5 was therefore provided as single-year warming. This section also updates the SR1.5 single-year approach by providing a year 2022 value.

7.1.2 Estimates of global surface temperature: GMST and GSAT

130 AR6 WGI (Chapter 2 Cross-Chapter Box 2.3, Gulev et al., 2021) described how global mean surface air temperature (GSAT), as is typically diagnosed from climate models, is physically distinct from the global mean surface temperature (GMST) estimated from observations, which generally combine measurements of near-surface temperature over land, and in some cases over ice, with measurements of sea surface temperature over the ocean

135 Based on conflicting lines of evidence from climate models, which show stronger warming of GSAT compared to GMST, and observations, which tend to show the opposite, Gulev et al. (2021) assessed with *high confidence* that long-term trends in the two indicators differ by less than 10%, but that there is *low confidence* in the sign of the difference in trends. Therefore, with *medium confidence*, in AR6 WGI Chapter 3 (Eyring et al., 2021), the best estimates and *likely* ranges for attributable warming expressed in terms of GMST were assessed to be equal to those for GSAT, with the consequence that the AR6 warming attribution results can be interpreted as both GMST and GSAT. While, based on the WGI Chapter 2 (Gulev et al., 2021) assessment, WGI Chapter 3 (Eyring et al., 2021) treated estimates of attributable warming in GSAT and GMST from the literature together, without any rescaling, we note that climate-model based estimates of attributable warming in GSAT are

Moved (insertion) [38]

Moved (insertion) [39]

Moved (insertion) [40]

140 expected to be systematically higher than corresponding estimates of attributable warming in GMST (see e.g. Cowtan et al., 2015; Richardson et al., 2018; Beusch et al., 2020; Gillett et al., 2021). Therefore, given an opportunity to update these analyses from AR6, it is more consistent, and more comparable with observations of GMST, to report attributable changes in GMST using all three methods (described in Sect. 7.2). The SR1.5 assessment of attributable warming was given in terms of GMST, which is continued here. In line with Sect. 2 and AR6 WGI, we adopt GMST as the estimate of global surface temperature.

Moved (insertion) [41]

Moved (insertion) [42]

145 7.2 Methods

Both SR1.5 and AR6 drew on evidence from a range of literature for their assessments of human-induced warming, before selecting results from a smaller subset to produce a quantified estimate. While both the SR1.5 and AR6 assessments used the latest Global Warming Index (GWI) results (Haustein et al., 2017), AR6 also incorporated results from two other methods, Regularised Optimal Fingerprinting (ROF) (as in Gillett et al., 2021) and Kriging for Climate Change (KCC) (as in Ribes et al., 2021). In AR6, all three methods gave results consistent not only with each other, but also results from AR6 WGI Chapter 7 (see WGI Chapter 7 Supplementary Material (Smith et al., 2021; Figure 3.8 of AR6 WGI Chapter 3 (Eyring et al., 2021; and Supplementary Material Sect. 7 Figure 1), though the results from Chapter 7 were not included in the AR6 final calculation because they were not statistically independent. Of the methods used, two (Gillett et al., 2021; Ribes et al., 2021) relied on CMIP6 DAMIP (Gillett et al., 2016) simulations which ended in 2020, and hence require modifications to update to the most recent years. The other two methods (Haustein et al., 2017; Smith et al., 2021) are updatable, and can also be made consistent with other aspects of the AR6 assessment and methods. The three methods used in the final assessment of contributions to warming in AR6 are used again with revisions for this annual update, and are presented in Supplementary Material, Sect. 7 with any updates to their approaches described in Sect. 7.2.

Moved (insertion) [43]

Moved (insertion) [44]

Moved (insertion) [45]

155 7.3 Updated estimates of human-caused warming to date

160 7.3.1 Updated estimate using the AR6 WGI methodology

Factoring in results from all three methods, AR6 WGI Chapter 3 (Erving et al., 2021) defined the *likely* (66% - 100% probability interval) range for each warming component as the smallest 0.1°C-precision range that enveloped the 5th to 95th percentile ranges of each method.

165 In addition, a best estimate was provided for the Human-induced (Ant) warming component, calculated as the mean of the 50th percentile values for each method. Best estimates were not provided in AR6 for the other components (well-mixed greenhouse gases (GHG), other human forcings (OHF), natural forcings (Nat)), with their values in AR6 WGI Figure SPM.2(b) simply being given as the midpoint between the lower and upper bound of the *likely* range, and therefore not directly comparable with the central values given for human-induced and observed warming. In order to make a meaningful and

Moved (insertion) [46]

Moved (insertion) [47]

consistent comparison, and provide meaningful insight into interannual changes, an improvement is made in this update: the multi-method-mean best estimate approach is extended for all warming components.

7.3.2 Updated estimate using the SR1.5 methodology applied to the AR6 WGI datasets

While a variety of literature was drawn upon for the assessment of human-induced warming in SR1.5 Chapter 1 (Allen et al., 2018), only one method, the Global Warming Index (GWI), was used to provide a quantitative assessment of the 2017, 'present-day', level of human-induced warming. The latest results for this method were provided in Haustein et al. 2017, which gave a central estimate for human-induced warming in 2017 of 1.01°C with 5-95% range of (0.87°C to 1.22°C).

SR1.5 then accounted for methodological uncertainty by rounding this value to 0.1°C precision for its final assessment of 1.0°C and assessing the 0.8°C to 1.2°C range as a *likely* range. No assessment of the contributions from other components was provided due to limitations in the GWI approach at the time.

Moved (insertion) [48]

While it is possible to continue the SR1.5 assessment approach of using a single method (GWI) rounded to 0.1°C-precision, for the purpose of providing annual updates this is insufficient; (i) 0.1°C-precision is too coarse to capture meaningful inter-annual changes to the level of present-day warming, (ii) using different selections of methods prevents meaningful comparison between the results for *decadal mean* and *present-day* warming calculations, and (iii) using the mean of multiple methods increases the robustness of the results. These points are simultaneously addressed in this update by adopting the latest multi-method assessment approach, as established in WGI AR6, for both the AR6 *decadal mean* warming update and the SR1.5 *present-day single-year* warming update.

Further, where SR1.5 only provided an assessment for human-induced warming, updates in available attribution methods since SR1.5 mean that it is now also possible to provide a fully-consistent assessment for all warming components. As with the attribution assessment in SR1.5, this update reports values in Table 6(b) for *single-year present-day* attributable warming, (as discussed in Sect. 7.1.1), with a comparison to results calculated using the SR1.5 trend based definition also provided below in Sect. 7.4.

Moved (insertion) [49]

7.4 Results

Results are summarised in Table 6 and Figure 5. WGI AR6 results for 2010-2019 are quoted in Table 6(a), compared with a repeat calculation using updated methods and datasets, and finally updated for the 2013-2022 period. Results from SR1.5 are quoted in Table 6(b) for the 2017

level of human-induced warming, compared with a repeat calculation using the updated selection of methods and datasets (see Sect. 7.2) and the WGI AR6 multi-method assessment approach (see Sect. 7.3.2), and finally updated for 2022. Method-specific contributions to the assessment results, along with timeseries, are given in Supplementary Material Sect. 7.

Moved (insertion) [50]

Moved (insertion) [51]

The repeat calculations for attributable warming in 2010-2019 exhibit good correspondence with the results in WGI AR6 for the same period, (see also Supplementary Material Sect. 7), with an exact correspondence in the best estimate and *likely* range of human-induced warming (Ant).

The repeat calculation for the level of attributable anthropogenic warming in 2017 is about 0.1°C larger than the estimate provided in SR1.5 for the same period, resulting from changes in methods and observational data (see above). The updated results for warming contributions in 2022 are also higher than in 2017 due to five additional years of anthropogenic forcing. A repeat assessment using the SR1.5 trend-based definition (see Sect. 7.1.1) leads to results that are very similar to the single-year results reported in Table 6(b), with 0.02°C differences at most.

The attribution assessment in WGI AR6 concluded that, averaged for the 2010-2019 period, all observed warming was human-induced, with solar and volcanic drivers and internal climate variability estimated not to make a contribution. This conclusion remains the same for the 2013-2022 period. Generally, whatever methodology is used, the best estimate of the human-caused warming to date is (within small uncertainties) equal to the observed warming to date.

Table 6: Updates to assessments in the 6th IPCC assessment cycle of warming attributable to multiple influences.

| Definition | Estimates of warming attributable to multiple influences, in °C, relative to the 1850–1900 baseline period Results are given as best estimates, with the likely range in brackets, and reported as Global Mean Surface Temperature. | | | | | | | |
|------------|--|----------------------|---|--|---|---|---|--|
| | (a) IPCC AR6 Attributable Warming Update Average value for previous 10-year period | | (b) IPCC SR1.5 Attributable Warming Update Value for single-year period | | (iii) 2013-2022 Updated value using updated methods and datasets | (i) 2017 Quoted from SR1.5 Chapter 1 Sect. 1.2.1.3 | (ii) 2017 Repeat calculation to using the updated methods and datasets | (iii) 2022 Updated value using updated methods and datasets |
| Period | (i) 2010-2019 Quoted from AR6 Chapter 3 Sect. 3.3.1.1.2 Table 3.1 | | (ii) 2010-2019 Repeat calculation using the updated methods and datasets | | | | | |
| Component | | | | | | | | |
| Observed | 1.06 (0.88 to 1.21) | 1.07 (0.89 to 1.22)* | 1.15 (1.00 to 1.25)* | | | | | |

- Moved up [32]: Earth energy imbalance (W m⁻²)
- Deleted: Time Period
- Deleted Cells
- Split Cells
- Deleted: Square brackets are [very likely ranges]
- Deleted: This study
- Moved (insertion) [65]
- Moved (insertion) [66]
- Moved (insertion) [67]
- Moved (insertion) [69]
- Moved (insertion) [68]
- Deleted: 1971-2018
- Deleted: 57 [0.43
- Deleted: 0.72]
- Inserted Cells
- Inserted Cells
- Inserted Cells
- Inserted Cells
- Deleted: 57 [0.43
- Deleted: 0.72]

| | | | | | |
|------------------------------------|---|---|---|-------------------|---|
| Anthropogenic | 1.07 (0.8 to 1.3) <small>1.07 (0.8 to 1.3)</small> | 1.14 (0.9 to 1.4) <small>1.14 (0.9 to 1.4)</small> | 1.0 (0.8 to 1.2) <small>1.0 (0.8 to 1.2)</small> | 1.13 (0.9 to 1.3) | 1.26 (1.0 to 1.6) |
| Well-mixed greenhouse gases | 1.40** (1.0 to 2.0) <small>1.40 (1.0 to 2.0)</small> | 1.33 (1.0 to 1.8) <small>1.33 (1.0 to 1.8)</small> | 1.40 (1.1 to 1.8) <small>1.40 (1.1 to 1.8)</small> | N/A (1.1 to 1.8) | 1.38 (1.1 to 2.0) <small>1.38 (1.1 to 2.0)</small> |
| Other human forcings | -0.32** (-0.8 to 0.0) <small>-0.32 (-0.8 to 0.0)</small> | -0.26 (-0.7 to 0.1) <small>-0.26 (-0.7 to 0.1)</small> | -0.25 (-0.7 to 0.1) <small>-0.25 (-0.7 to 0.1)</small> | N/A (-0.7 to 0.1) | -0.24 (-0.7 to 0.1) <small>-0.24 (-0.7 to 0.1)</small> |
| Natural forcings | 0.03** (-0.1 to 0.1) <small>0.03 (-0.1 to 0.1)</small> | 0.05 (-0.1 to 0.1) <small>0.05 (-0.1 to 0.1)</small> | 0.04 (0.0 to 0.2) <small>0.04 (0.0 to 0.2)</small> | N/A (-0.1 to 0.1) | 0.03 (-0.1 to 0.1) <small>0.03 (-0.1 to 0.1)</small> |

Results from the 6th IPCC assessment cycle, for both AR6 and SR1.5, are quoted in columns labelled (i), and are compared with repeat calculations in columns labelled (ii) for the same period using the updated methods and datasets to see how methodological and dataset updates alone would change previous assessments. Assessments for the updated periods are reported in columns labelled (iii). Table 6.1(a): * Updated GMST observations, quoted from Sect. 5 of this update, are marked with an asterisk, with very likely ranges given in brackets. ** In AR6 WGI, best estimate values were not provided for warming attributable to well-mixed greenhouse gases, other human forcings, and natural forcings, (though they did receive a likely range, as discussed in Sect. 7.3.1); for comparison, best estimates (marked with two asterisks) have been retrospectively calculated in an identical way to the best estimate that AR6 provided for anthropogenic warming.

Moved (insertion) [70]

Deleted: 1971-2006

Deleted: 50 [0.32...to 1.3)0.69] ... [47]

Deleted: .50 [...831...to 1.2)0.68] ... [50]

Inserted Cells ... [48]

Inserted Cells ... [49]

Inserted Cells ... [51]

Moved (insertion) [52]

Deleted: 2006-2018

Deleted: .79 [0.52...to 2.0)1.06] ... [52]

Deleted: .79 [0.52...to 1.8)07] ... [53]

Inserted Cells ... [54]

Inserted Cells ... [55]

Inserted Cells ... [56]

Deleted: -

Deleted Cells ... [57]

Inserted Cells ... [59]

Inserted Cells ... [60]

Inserted Cells ... [61]

Deleted: 1975-2022

Deleted: 65 [...848...to 0.0)81] ... [58]

Deleted: 2010-2022

Deleted: -

Inserted Cells ... [63]

Inserted Cells ... [64]

Deleted: 89 [...163...to 0.1).15] ... [62]

Moved up [33]: For 1973-1992 the contribution by ocean vertical

Moved up [34]: Figure 4: a) Observed changes in the Earth

Moved up [35]: . An assessment of human-induced warming was

Moved up [36]: 2018), summarised in SPM A.1 and Figure

Moved up [37]: 2021), summarised in WGI SPM A.1.3 and

Moved up [38]: . Based on conflicting lines of evidence from

Moved up [39]: the best estimates and likely ranges for

Moved up [40]: treated estimates of attributable warming in

Moved up [41]:). The SR1.5 assessment of attributable warming

Moved up [42]: .

Moved up [43]: Of the methods used, two (Gillett et al.,

Moved up [44]: relied on CMIP6 DAMIP (Gillett et al., ... [73]

Moved up [46]: In addition, a best estimate was provided for the

Moved up [47]: In order to make a meaningful and consistent

Moved up [48]: SR1.5 then accounted for methodological

Moved up [49]: Further, where SR1.5 only provided an

Moved up [50]: level of human-induced warming, compared

Moved up [51]: .

Deleted: ... [65]

Deleted: 6). ... [66]

Deleted: 7.1 Introduction ... [67]

Deleted: .

Deleted: .

Deleted: AR6 ... [68]

Deleted: (2021) assessed that long-term trends in the two i ... [69]

Deleted: . (2021))

Deleted: 7.2), though we also show the sensitivity of result ... [70]

Deleted: Reporting attributable warming updates in terms ... [71]

Deleted: 2017), AR6 also incorporated results from two of ... [72]




Figure 5: Updated assessed contributions to observed warming relative to 1850-1900, cf. AR6 WGI SPM.2. Results for all time periods in this figure are calculated using updated datasets and methods. The 2010-2019 decade-average assessed results repeat the AR6 2010-2019 assessment, and the 2017 single-year assessed results repeat the SR1.5 2017 assessment. For each double bar the lighter and darker shading refers to the earlier and later period respectively. The 2013-2022 decade-average, and 2022 single-year results are the updated assessments for AR6 and SR1.5 respectively. Panel (a) shows updated observed global warming from Sect. 5, expressed as total GMST, due to both anthropogenic and natural influences. Whiskers give the very likely range. Panel (b) and Panel (c) show updated assessed contributions to warming, expressed as global mean surface temperature, from natural forcings and total human-induced forcings, which in turn consists of contributions from well-mixed greenhouse-gases, and other human forcings. Whiskers give the likely range.

Moved (insertion) [54]

Moved (insertion) [55]

880 **8 Remaining Carbon Budget**

AR6 assessed the remaining carbon budget (RCB) in Chapter 5 of its WGI report (Canadell et al., 2021) for 1.5°C, 1.7°C and 2°C thresholds (see Table 7). They were also reported in its Summary for Policy Makers (Table SPM2, IPCC, 2021b). These are updated in this section using the same method with transparently described updates.

Moved (insertion) [56]

885 AR5 (IPCC, 2013) assessed that global surface temperature increase is close to linearly proportional to the total amount of cumulative CO₂ emissions (Collins et al., 2013). The most recent AR6 report reaffirmed this assessment (Canadell et al., 2021). This near-linear relationship implies that for keeping global warming below a specified temperature level, one can estimate the total amount of CO₂ that can ever be emitted, also known as the carbon budget. When expressed relative to a recent reference period, this is referred to as the remaining carbon budget (Rogelj et al., 2018).

890 The RCB is estimated by application of the WGI AR6 method described in Rogelj et al. (2019), which involves the combination of the assessment of five factors: (i) the most recent decade of human-induced warming (ii) the transient climate response to cumulative emissions of CO₂ (TCRE), (iii) the zero emissions commitment (ZEC), (iv) the temperature contribution of non-CO₂ emissions, and (v) an adjustment term for Earth system feedbacks that are otherwise not captured through the other factors. AR6 WGI reassessed all five terms (Canadell et al., 2021). The incorporation of factor (v) was further considered by Lamboll and Rogelj (2022).

Moved (insertion) [57]

895 Of these factors, only factor (i) (human-induced warming) lends itself to a regular and systematic annual update, the decade-long period, 2010-2019, was used in WGI AR6. Up-to-date historical CO₂ emissions from the middle of this period until the start of the RCB are required to have an as up-to-date RCB estimate as possible.

900 Other factors can be updated, but depend on new evidence and insights being published rather than an additional year of observational data becoming available. Factor (iv) (temperature contribution of non-CO₂ emissions) depends both on the available mitigation scenario evidence and the assessment of non-CO₂ warming. Additional scenario evidence has become available through the publication of the scenario database supporting the AR6 WGIII report (Byers et al., 2022) which is taken into account in this update.

Moved (insertion) [58]

Moved (insertion) [59]

910 The RCB for 1.5°C, 1.7°C and 2°C warming levels are re-assessed based on the most recent available data. Estimated RCBs are reported below. They are expressed both relative to 2020 to compare to AR6 and relative to the start of 2023 for estimates based on the 2013-2022 human-induced warming update. Note that between the start of 2020 and the end of 2022, about 122

Moved (insertion) [60]

915 GtCO₂ have been emitted (Sect. 2). Based on the variation in non-CO₂ emissions across the scenarios in AR6 WGIII scenario
database, the estimated RCB values can be higher or lower by around 200 GtCO₂, depending on how deeply non-CO₂ emissions
are reduced. The impact of non-CO₂ emissions on warming includes both the warming effects of other greenhouse gases such
as methane and the cooling effects of aerosols such as sulphates. The impacts of these are assessed using a climate emulator
(MAGICC, Meinshausen et al., 2011), which was updated to more accurately capture recent updates from the AR6 WGIII
report, but whose results were not captured in the AR6 WGI carbon budget estimates. This emulator update increased the
estimate of the importance of aerosols, which are expected to decline with time in low emissions pathways (Rogelj et al.,
2014), causing a net warming, and decreasing the remaining carbon budget. The AR6 WGI version of MAGICC is used here.
920 If instead, the FaIR emulator were used, this would give reduced non-CO₂ warming and a larger carbon budget (Lamboll and
Rogelj, 2022).

Table 7: Updated estimates of the Remaining Carbon Budget for 1.5°C, 1.7°C and 2.0°C, for five levels of likelihood, considering only uncertainty in TCRE.

| Historical cumulative CO ₂ emissions (1850 -2019) AR6 WGI Table SPM.2 | | 2390 (± 240; likely (66%-100% probability) range) | | | | | | |
|--|-----------|---|-----|-----|-----|-----|--|--|
| Remaining carbon budgets Case / update | Base year | Estimated remaining carbon budgets from the beginning of base year (GtCO ₂) | | | | | | |
| | | 17% | 33% | 50% | 67% | 83% | | |
| Likelihood of limiting global warming to temperature limit | | | | | | | | |
| 1.5°C from AR6 WGI | 2020 | 900 | 650 | 500 | 400 | 300 | | |
| + AR6 emulator update | 2020 | 750 | 500 | 400 | 300 | 200 | | |
| + as above with AR6 scenario update | 2020 | 750 | 500 | 400 | 300 | 200 | | |
| + as above with warming update (2013-2022) (best estimate) | 2023 | 500 | 300 | 250 | 150 | 100 | | |

Moved (insertion) [61]

Deleted: Estimates of warming attributable to multiple influences, in °C, relative to the 1850–1900 baseline period
 Results are given as best estimates, with the likely range in brackets, and reported as Global Mean Surface Temperature.

Inserted Cells

Moved (insertion) [63]

Moved (insertion) [64]

Deleted: (a) IPCC AR6 Attributable Warming Update
 Average value for previous 10-year period

Deleted: (b) IPCC SR1.5 Attributable Warming Update
 Present-day value for single-year period

Moved (insertion) [62]

Deleted: Definition

Moved up [65]: (i) 2010-2019
 Quoted from AR6 Chapter 3 Sect. 3.3.1.1.2 Table 3.1

Moved up [66]: (ii) 2010-2019
 Repeat calculation using the updated methods and datasets

Moved up [67]: (iii) 2013-2022
 Updated value using updated methods and datasets

Moved up [68]: Repeat calculation to using the updated methods and datasets

Moved up [69]: (iii) 2022
 Updated value using updated methods and datasets

Deleted: (i) 2017
 Quoted from SR1.5 Chapter 1 Sect. 1.2.1.3

Deleted: (ii) 2017

Split Cells

Deleted: Period

Deleted: Component

Deleted: 1.06 (0.88 to 1.21)

Deleted: 1.07 (0.89 to 1.22) *

Deleted: 1.15 (1.00 to 1.25) *

Deleted: Observed

| | | | | | | | |
|--|------|------|------|------|------|-----|-----|
| 1.7°C from AR6 WGI | 2020 | 1450 | 1050 | 850 | 700 | 550 | |
| + AR6 emulator update | 2020 | | 1250 | 900 | 700 | 600 | 450 |
| + as above with AR6 scenario update | 2020 | | 1300 | 950 | 750 | 600 | 500 |
| + as above with warming update (2013-2022) (best estimate) | 2023 | | 1100 | 800 | 600 | 500 | 350 |
| 2°C from AR6 WGI | 2020 | 2300 | 1700 | 1350 | 1150 | 900 | |
| + AR6 emulator update | 2020 | 2050 | 1500 | 1200 | 1000 | 800 | |
| + as above with AR6 WGIII scenario update | 2020 | 2200 | 1650 | 1300 | 1100 | 900 | |
| + as above with warming update (2013-2022) (best estimate) | 2023 | 2000 | 1450 | 1150 | 950 | 800 | |

Estimates start from AR6 WGI estimates (first row for each warming level), updated with the latest scenario information from AR6 WGIII (second row for each warming level), and an update of the anthropogenic historical warming which is estimated for the 2013-2022 period (third row for each warming level). Estimates are expressed relative to the start of year 2020 or 2023. The probability includes only the uncertainty in how the Earth immediately responds to carbon, not long-term committed warming or uncertainty in other emissions. All values are rounded to the nearest 50 GtCO₂.

Updated RCB estimates presented in Table 7

for 1.5°C, 1.7°C and 2.0°C of global warming are smaller than AR6, and geophysical and other uncertainties therefore have become larger in relative terms. This is a feature that will have to be kept in mind when communicating budgets. The estimates presented here differ from those presented in the annual Global Carbon Budget (GCB) publications (Friedlingstein et al., 2022a). The GCB updates have previously started from the AR6 WGI estimate and subtracted the latest estimates of historical CO₂ emissions. The RCB estimates presented here take into account the same updates in historical CO₂ emissions from the GCB as well as the latest available quantification of human-induced warming to date and a reassessment of non-CO₂ warming contributions.

If the single year human-induced warming until 2022 (Sect. 7) was used directly in the RCB calculation, this would lead to similar remaining carbon budgets estimates to those from the decadal average approach used here; the 50% likelihood estimates

Moved up [70]: 1.07 (0.8 to 1.3)

Deleted: 1.07 (0.8 to 1.3)

Deleted: 1.14 (0.9 to 1.4)

Deleted: 1.0 (0.8 to 1.2)

Deleted: 1.13 (0.9 to 1.4)

Deleted: 1.26 (1.0 to 1.6)

Deleted: Anthropogenic

Moved up [52]: Well-mixed greenhouse gases

Deleted: 1.40** (1.0 to 2.0)

Deleted: 1.33 (1.0 to 1.8)

Deleted: 1.40 (1.1 to 1.8)

Deleted: N/A

Deleted: 1.38 (1.1 to 1.8)

Deleted: 1.49 (1.1 to 2.0)

Deleted: -0.32** (-0.8 to 0.0)

Deleted: -0.26 (-0.7 to 0.1)

Deleted: -0.25 (-0.7 to 0.1)

Deleted: N/A

Deleted: -0.25 (-0.7 to 0.1)

Deleted: -0.24 (-0.7 to 0.1)

Deleted: Other human forcings

Deleted: 0.03** (-0.1 to 0.1)

Deleted: 0.05 (-0.1 to 0.1)

Deleted: 0.04 (0.0 to 0.1)

Deleted: N/A

Deleted: 0.04 (-0.1 to 0.2)

Deleted: 0.03 (-0.1 to 0.1)

Deleted: Natural forcings

Moved up [53]: best estimate values were not provided for

Moved up [54]: Results for all time periods in this figure are

Moved up [55]: and 2022 single-year results are the updated

Moved up [56]: . They were also reported in its Summary for

Moved up [57]: , (ii) the transient climate response to cumulative


Moved up [58]: 

Moved up [59]: 2022) which is taken into account in this

Moved up [60]: Note that between the start of 2020 and the end

Moved up [61]: version of MAGICC is used here. If instead, the

Deleted: Results from the 6th IPCC assessment cycle, f(... [81]



would be unchanged although other likelihoods alter somewhat because the spread due to TCRE uncertainty starts 5 years later. However, we choose to only show the decadal calculation as this was assessed to be the best estimate for human-induced warming and the method adopted in AR6 WGI.

Moved (insertion) [73]

The RCB for limiting warming to 1.5°C is becoming very small. It is important, however, to correctly interpret this information. RCB estimates take into account projected reductions in non-CO₂ emissions that are aligned with a global transition to net zero CO₂ emissions. These estimates assume median reductions in non-CO₂ emissions between 2020-2050 of CH₄ (50%) N₂O (25%) and SO₂ (77%). If these non-CO₂ greenhouse gas emission reductions are not achieved the RCB would be smaller (see Supplementary Material Sect. 8). Note that the 50% RCB is expected to be exhausted a few years before the 1.5°C global warming level is reached.

9. Examples of climate and weather extremes : maximum temperature over land

Climate and weather extremes belong to the most visible human-induced climate changes. Within AR6 WGI, a full chapter was dedicated to the assessment of past and projected changes in extremes on continents (Seneviratne et al., 2021), and the chapter on ocean, cryosphere and sea level changes also provided assessments on changes in marine heatwaves (Fox-Kemper et al., 2021). Global indicators related to climate extremes include averaged changes in climate extremes, e.g., the mean increase of annual minimum and maximum temperatures on land (AR6 WGI Chapter 11, Figure 11.2, Seneviratne et al., 2021) or the area affected by certain types of extremes (AR6 WGI Chapter 11, Box 11.1, Figure 1, Seneviratne et al., 2021; Sippel et al., 2015). In contrast to global surface temperature, extreme indicators are less established. They are therefore expected to be subject to improvements, reflecting advances in understanding and better data collection. Indeed, such efforts are planned within the World Climate Research Programme (WCRP) Grand Challenge on Weather and Climate Extremes, which will likely inform the next iteration of this study.

Moved (insertion) [74]

Moved (insertion) [75]

Moved (insertion) [76]

As part of this first update, we provide an upgraded version of the analysis in Figure 11.2 from Seneviratne et al., 2021. Like the analysis of global mean temperature, the choice of data sets is based on a compromise on the length of the data record, the data availability, near-real time updates and long-term support. As the indicator (in its current form) averages over all available land grid points, the spatial coverage should be high to obtain a meaningful average, which further limits the choice of datasets. The HadEX3 dataset (Dunn et al., 2020), which is used for Figure 11.2 in Seneviratne et al. (2021), is static and does not cover years after 2018. We therefore additionally include the Berkeley Earth Surface Temperature dataset (building off Rohde et al., 2013), and the fifth generation ECMWF atmospheric reanalysis of the global climate (ERA5; Hersbach et al., 2020). Berkeley

Moved (insertion) [77]

Earth data currently enable an analysis of annual indices up to 2021 while ERA5 is updated daily with a latency of about 5 days (and the final release occurs after 2–3 months).

315 Our proposed climate indicator of changes in temperature extremes consists of land-averaged annual maximum temperatures (TXx), (excluding Antarctica). For HadEX3 we select the years 1961–2018, to exclude years with insufficient data coverage, and require at least 90% temporal completeness, thus applying the same criteria as for Figure 11.2 (Seneviratne et al., 2021). Berkeley Earth provides daily maximum temperatures and we require more than 99% data availability for each individual year and grid, such that years with more than four missing days are removed. Based on this criterion, Berkeley Earth covers at least
320 95% of the global land area from 1955 onwards. ERA5, on the other hand, has full spatiotemporal coverage by design, and hence the entire currently available time period of 1950 to 2022 is used. The annual maximum temperature is then computed for each grid cell, and a global area-weighted average is calculated for all grid cells with at least 90% temporal completeness in the respective available period (1955–2021 and 1961–2018 for Berkeley Earth and HadEX3, while ERA5 is again not affected by this criterion). We thus enforce high data availability to adequately calculate global land-averaged TXx across all
325 three datasets, but their coverage is not identical which introduces minor deviations in the estimated global land averages. The resulting TXx timeseries are then computed as anomalies with respect to a baseline period of 1961–1990.

To express the TXx as anomalies with respect to 1850-1900 we add an offset to all three datasets. The offset is based on the Berkeley Earth data and is derived from the linear regression of land-mean TXx to the annual mean global mean air temperature over the period 1955 to 2020. The offset is then calculated as the slope of the linear regression times the global mean temperature difference between the reference periods 1850-1900 and 1961-1990 (see Supplementary Material, Figure 4).

Moved (insertion) [78]

335 Our climate has warmed rapidly in the last few decades, which also manifests in changes in the occurrence and intensity of climate and weather extremes. We visualise this with land-averaged annual maximum temperatures (TXx) from three different datasets (ERA5, Berkeley Earth and HadEX3), expressed as anomalies with respect to the pre-industrial baseline period of 1850–1900 (Figure 6). From about 1980 onwards, all employed datasets point to a strong TXx increase, which coincides with the transition from global dimming, associated with aerosol increases, to brightening, associated with decreases (Wild et al., 2005). Together with strongly increasing greenhouse gas emissions (Sect. 2), this explains why human-induced climate change has emerged at an even greater pace in the last four decades than previously. For example, land-averaged annual maximum
340 temperatures have warmed by more than 0.5 °C in the past 10 years (1.72 °C with respect to pre-industrial conditions) compared to the first decade of the millennium (1.22 °C; Table 8). Since the offset relative to our pre-industrial baseline period is calculated relative to 1961–1990, within the latter period, temperature anomalies align by construction but can diverge afterwards. In an extensive comparison of climate extreme indices across several reanalyses and observational products, Dunn

Moved (insertion) [79]

Moved (insertion) [80]

et al. (2022), point to an overall strong correspondence between temperature extreme indices across reanalysis and observational products, with ERA5 exhibiting especially high correlations to HadEX3 among all regularly updated datasets. This suggests that both our choice of datasets and approach to calculate anomalies does not affect our conclusion — the intensity of heatwaves across all land areas has unequivocally increased since pre-industrial times.




Figure 6: Time series of observed temperature anomalies for land average annual maximum temperature (TXx) for ERA5 (1950–2022), Berkeley Earth (1955–2021), and HadEX3 (1961–2018), with respect to 1850–1900. Note that the datasets have different spatial coverage and are not coverage-matched. All anomalies are calculated relative to 1961–1990 and an offset of 0.53°C is added to obtain TXx values relative to 1850–1900. Note that while the HadEX3 numbers are the same as shown in Seneviratne et al. (2021) Figure 11.2, these numbers were not specifically assessed.

Table 8: Anomalies of land average annual maximum temperature (TXx) for recent decades based on HadEX3 and ERA5.

| Period | Anomaly w.r.t. 1961-1990 (°C) | | Anomaly w.r.t. 1850-1900 (°C) | | |
|-----------|-------------------------------|------|-------------------------------|--|--|
| | HadEX3 | ERA5 | ERA5 | | |
| 2000-2009 | 0.72 | 0.69 | 1.23 | | |

- Moved (insertion) [81]
- Moved up [62]: Case / update
- Moved up [63]: Base year
- Moved up [64]: Estimated remaining carbon budgets from the beginning of base year (GtCO₂)
- Deleted: Likelihood of limiting global warming to temperature limit.
- Deleted: 17%
- Deleted: 33%
- Deleted: 50%
- Deleted: 67%
- Deleted: 83%
- Deleted Cells
- Deleted Cells
- Deleted Cells

| | | | | | | |
|-----------|------|-----------|------|------|--|--|
| 2009-2018 | 1.01 | 1.02 | 1.55 | | | |
| 2010-2019 | | | 1.11 | 1.64 | | |
| | | 2011-2020 | 1.12 | 1.65 | | |
| 2012-2021 | 1.18 | 1.74 | | | | |

The anomalies with respect to 1850-1900 are derived by adding an offset of 0.53°C. Note that while the HadEX3 numbers are the same as shown in Seneviratne et al. (2021) Figure 11.2, these numbers were not specifically assessed.

10. Dashboard data visualisations

The Climate Change Tracker (), a platform hosting a range of publicly available climate data, aims to provide a range of audiences with a reliable, user-friendly means of tracking and understanding climate change and its progression.

Building on the existing platform, a bespoke “dashboard” places a number of the updated IPCC-consistent indicators of climate change set out above in the public domain. This bespoke dashboard is primarily aimed at policymakers involved in UNFCCC negotiations, but the ultimate intention is to reach and inform a much wider audience.

The dashboard initially focuses on three key indicator sets: greenhouse gas emissions (Sect. 2); human-induced global warming (Sect. 7); and the remaining global carbon budget (Sect. 8), bringing together and presenting up-to-date information crucial to effective climate decision-making in a findable, accessible, traceable and reproducible way. In addition, the Climate Change Tracker provides standardised application programming interfaces (APIs), dashboards and charts to embed in third-party apps and websites. All data is traceable to the github repository employed for this paper (Sect. 11).

In time, and with feedback from the user community, the initial set of indicators displayed by the dashboard may be expanded to include others alongside their rates of change.

11. Code and data availability

The carbon budget calculation is available from . The code and data used to produce other indicators is available in repositories under . All data is available from (Smith et al., 2023). Data is provided under a CC-BY 4.0 Licence.

Inserted Cells ... [89]

Deleted: 5°C from AR6 WGI

Deleted: 2020

Deleted: 900

Deleted: 650

Deleted Cells ... [90]

Deleted: 500

Deleted Cells ... [91]

Deleted: 400

Deleted Cells ... [92]

Deleted: 300

Deleted Cells ... [93]

Moved down [82]: 2020

Deleted: + AR6 emulator update

Deleted: 750

Deleted: 500

Deleted: 400

Deleted: 300

Deleted: 200

Deleted: + AR6 scenario update

Deleted Cells ... [94]

Deleted: 750

Deleted: 500

Deleted: 400

Deleted: 300

Deleted: 200

Deleted: + warming update (2013-2022) (best esti... [95]

Inserted Cells ... [96]

Inserted Cells ... [97]

Deleted: 7°C from AR6 WGI

Deleted: 2020

Deleted: 1450

Deleted: 1050

Deleted: 850

Deleted: 700

Deleted: 550

Deleted Cells ... [98]

Deleted Cells ... [99]

Deleted Cells ... [100]

Deleted Cells ... [101]

Deleted Cells ... [102]

Moved up [71]: . The probability includes only the uncertainty

Moved up [72]: for 1.5°C, 1.7°C and 2.0°C of global warming

Moved up [73]: 1

Moved up [74]: Climate and weather extremes belong to the most

Moved up [75]: related to climate extremes include averaged

Deleted: + AR6 emulator update ... [103]

Deleted: updated estimate relative to the 2013-2022 pe... [104]

Deleted: Updated RCB estimates presented in Table 8

Deleted: . 2022). The GCB updates have previously start... [105]

Deleted: This means that reductions in aerosol cooling w... [106]

Deleted: Global metrics

Deleted: . Indeed, such efforts are planned within the We... [107]

Moved up [76]: Grand Challenge on Weather and Climate

Moved up [77]: 2013). and the fifth generation ECMWF

770 **12. Discussion and conclusions**

The first year of the Global Climate Change (IGCC) initiative has built from the AR6 report cycle to provide a comprehensive update of the climate change indicators required to estimate the human induced warming and the remaining carbon budget. Table 9 presents a summary of the headline figures from each section compared to that given in the AR6 assessment. The main substantive data change since AR6 is that land-use CO₂ emissions have been revised down by around 2 GtCO₂. However, as CO₂ ERF and human induced warming estimates depend on concentrations, not emissions, this does not affect most of the other findings. Note it does slightly increase the remaining carbon budget, but this is only by 5 GtCO₂, less than the 50 GtCO₂ rounding precision.

Moved (insertion) [86]

775

Table 9: Summary of headline results and methodological updates from the Indicators of Global Climate Change (IGCC) initiative.

| Climate Indicator | AR6 2021 assessment | This assessment | 2023 | Explanation of changes | Methodological updates |
|---|--|--|---|--|---|
| Greenhouse gas emissions | AR6 WGIII Chapter 2: Dhakal et al. (2022); see also Minx et al. (2021) | 2010-2019 average: 53 ± 5.6 GtCO ₂ e 2012-2021 average: 54 ± 5.3 GtCO ₂ e | 56 ± 6 GtCO ₂ e* | The change from AR6 is due to a systematic downward revision in CO ₂ -LULUCF and CH ₄ estimates. Real-world emissions have slightly increased. Average emissions in the past decade grew at a slower rate than in the previous decade. Note following convention, ODS F-gases are excluded from the total. | CO ₂ -LULUCF emissions revised down. PRIMAP-hist used in place of EDGAR for CH ₄ and N ₂ O emissions. Atmospheric measurements taken for F-gas emissions. These changes reduce estimates by around 3 GtCO ₂ e (Sect. 2) |
| Greenhouse gas concentrations | AR6 WGI Chapter 2: Gulev et al. (2021) | 2019: CO ₂ 410.1 [± 0.36] ppm CH ₄ 1866.3 [± 3.2] ppb | 2022: CO ₂ 417.1 [± 0.4] ppm CH ₄ 1911.9 [± 3.3] ppb | Continued and increasing emissions. | Updates based on NOAA data as AGAGE not yet available for 2022. To make an AR6-like product, N ₂ O scaled to approximate NOAA-AGAGE average (Sect. 3) |
| Effective radiative forcing change since 1750 | | 2019: 2.91 [2.19 to 3.63] W m ⁻² | 2022: 2.91 [2.19 to 3.63] W m ⁻² | Overall substantial increase and high decadal rate of change. | Minor update in aerosol precursor method for improved future |

Inserted Cells
 Deleted: Period
 Deleted: Anomaly w.r.t. 1961-1990 (°C)
 Deleted: Anomaly w.r.t. 1850-1900 (°C)
 Inserted Cells

Moved (insertion) [11]

Deleted: ERA5
 Deleted: ERA5
 Deleted: HadEX3
 Deleted: Period (... [110])
 Inserted Cells

Deleted: 0.72
 Deleted: 0.69
 Deleted: 1.23

| | | | | | |
|---|--|--|---|--|---|
| <u>AR6 WGI Chapter 7: Forster et al. (2021)</u> | | <u>2.72 [1.96 to 3.48] W m⁻²</u> | | <u>arising from increases in greenhouse gas concentrations and reductions in aerosol precursors</u> | <u>estimates - had no impact at quoted accuracy level (Sect. 4)</u> |
| <u>Global mean surface temperature change above 1850-1900</u> | | <u>2011-2020 average: 1.09 [0.95 to 1.20] °C</u> | <u>2013-2022 average: 1.15 [1.00-1.25] °C</u> | <u>An increase of 0.06 °C within two years, indicating a high decadal rate of change.</u> | <u>Methods match AR6 (Sect. 5)</u> |
| <u>AR6 WGI Chapter 2: Gulev et al. (2021)</u> | | <u>1.09 [0.95 to 1.20] °C</u> | <u>1.15 [1.00-1.25] °C</u> | <u>indicating a high decadal rate of change.</u> | |
| <u>Earth's energy imbalance</u> | | <u>2006-2018 average: 0.79 [0.52-1.06] W m⁻²</u> | <u>2010-2022. average: 0.89 [0.63 to 1.15] W m⁻²</u> | <u>Substantial increase in energy imbalance estimated based on increased rate of ocean heating.</u> | <u>Ocean heat content timeseries extended from 2018 to 2022 using 4 of the 5 AR6 datasets. Other heat inventory terms updated following von Schuckmann et al (2023). Ocean heat content uncertainty is used as a proxy for total uncertainty. Further details in Sect. 6.</u> |
| <u>AR6 WGI Chapter 7: Forster et al. (2021)</u> | | <u>0.79 [0.52-1.06] W m⁻²</u> | <u>0.89 [0.63 to 1.15] W m⁻²</u> | | |
| <u>Human induced global warming since preindustrial</u> | <u>2010-2019 average: 1.07 [0.8 to 1.3] °C</u> | <u>2013-2022 average: 1.14 [0.9 to 1.4] °C</u> | <u>An increase of 0.07 °C within three years, indicating a high decadal rate of change.</u> | <u>The three methods for the basis of the AR6 assessment are retained, but each has new input data (Sect. 7)</u> | |
| <u>AR6 WGI Chapter 3: Eyring et al. (2021)</u> | <u>1.07 [0.8 to 1.3] °C</u> | <u>1.14 [0.9 to 1.4] °C</u> | <u>indicating a high decadal rate of change.</u> | | |
| <u>Remaining carbon budget for 50% likelihood of limiting global warming to 1.5°C</u> | <u>From the start of 2020: 250 GtCO₂ (depends on stronger or weaker</u> | <u>The 1.5°C budget is becoming very small. The RCB can exhaust before the 1.5°C</u> | <u>Methods match AR6 (Sect. 8).</u> | | |

Inserted Cells

Moved (insertion) [82]

Deleted: 1.01

Deleted: 2009-2018

Deleted: 02

Deleted: 1.55

Inserted Cells

Inserted Cells

Deleted: 1.11

Deleted: 1.64

Deleted: -

Deleted Cells

Deleted: 2011-

Deleted: 1.65

| | | | | |
|---|--|---|----------------------------|----------------------------|
| AR6 WGI Chapter 5: Canadell et al. (2021) | accompanying non-CO ₂ mitigation) | threshold is reached due to having to allow for future non-CO ₂ warming. | | |
| Land average maximum temperature change compared to preindustrial | | | 2009-2018 average: 1.55 °C | 2013-2022 average: 1.74 °C |
| AR6 WGI Chapter 11: Seneviratne et al., 2021 | | | | |
| https://climatechangetracker.org/ https://github.com/RIamboll/AR6CarbonBudgetCalc https://github.com/ClimateIndicator https://doi.org/10.5281/zenodo.7883758 | | | | |

Figure 7 summarises contributions to warming, repeating Figure 2.1 of the AR6 Synthesis Report (Lee et al., 2023). It highlights changes since the assessment period in ARG WGI. Table 9 also summarises methodological updates.

- Deleted:** -
- Deleted:** 1.12
- Inserted Cells**
- Inserted Cells**
- Deleted:** 2012-2021
- Deleted:** 18
- Deleted:** 71
- Deleted:** -
- Deleted Cells**
- Moved up [83]:** In time, and with feedback from the user community,
- Moved up [84]:** 11. Code and data availability The carbon budget calculation is available from
- Deleted:** The anomalies with respect to 1850-1900 are derived by adding an offset of 0.53°C. Note that while the HadEX3 numbers are the same as shown in Seneviratne et al. (2021) Figure 11.2, these numbers were not specifically assessed.
- 10. Dashboard data visualisations** Software engineer and data scientist authors (Borger and Broersma) have created The Climate Change Tracker (
- Deleted:**) a “dashboard” of publicly available climate data. This builds off their experience of financial industry products: providing real time intuitive access to complex information. The aim of this tracker is to present a range of audiences with a reliable, user-friendly platform for tracking and understanding climate change and its progression. Building on the existing platform, we place updated IPCC-consistent indicators of climate change into the public domain via a bespoke “dashboard” aimed primarily at policymakers involved in UNFCCC negotiations, but also intended to reach and inform a much wider audience.
- The policy-facing dashboard initially focuses on three key indicator sets: greenhouse gas emissions (Sect. 2); human-induced global warming (Sect. 7); and the remaining global carbon budget (Sect. 8). The climate change indicator dashboard will bring together and present up-to-date information crucial to effective climate decision-making in a findable, accessible, traceable and reproducible... [111]
- Deleted:** this set of indicators may be expanded to look at other indicators as well as rates of change. However, the current ... [112]
- Deleted:** . The code and data used to produce other indicators is available in repositories under
- Moved up [85]:** 2023). Data is provided under a CC-BY 4.0 Licence.
- Moved up [86]:** . However, as CO₂ ERF and human induced warming estimates depend on concentrations, not emissions, this does
- Deleted:** . All data is available from
- Deleted:** .
- Deleted:** The table

Since AR6 WGI (2021), humans had added to global warming





Figure 7. The causal chain from emissions to resulting warming of the climate system. Emissions of GHG have increased rapidly over recent decades (panel a). These emissions have led to increases in the atmospheric concentrations of several GHGs including the three major well-mixed GHGs (panel b). The global surface temperature (shown as annual anomalies from an 1850–1900 baseline) has increased by around 1.15°C since 1850–1900 (panel c). The human-induced warming estimate over the last decade is a close match to the observed warming (panel d). Whiskers show 5% to 95% ranges. Figure is modified from AR6 SYR with a zoom on the period 2000 to 2022 for the upper two panels (Figure 2.1, Lee et al., 2023).

Since AR6 WGI (2021), humans had added to global warming



Deleted:

It is hoped that this update can support the science community in its collection and provision of reliable and timely global climate data. In future years we are particularly interested in improving SLCF updating methods to get a more accurate estimate of short-term ERF changes. The work also highlights the importance of high-quality metadata to document changes in methodological approaches over time. In future years we hope to improve the robustness of the indicators presented here but also extend the breadth of indicators reported through coordinated research activities. For example, we could begin to make use of new satellite data inversion techniques to infer recent emissions. We are particularly interested in exploring how we might update indicators of regional climate extremes and their attribution, which are particularly relevant for supporting actions on adaptation and loss and damage.

Generally, scientists and scientific organisations such as WMO and IPCC have an important role as “watchdogs” to critically inform evidence-based decision making. This annual update traced to IPCC methods can provide a reliable, timely source of trustworthy information. As well as helping inform decisions, we can use the update to track changes in dataset homogeneity between their use in one IPCC report and the next. We can also provide information and testing to motivate updates in methods that future IPCC reports might choose to employ.

Figure 8 shows decadal trends for the attributed warming and ERF. The most recent trends were unprecedented at the time of AR6 and have increased further since then (red markers), showing that human activities are consistently causing global warming recently of more than 0.2 °C per decade. As nations and businesses forge climate policies and take meaningful action, the latest available evidence shows that global actions are not yet at the scale to manifest a substantive shift in the direction of global human influence on the Earth’s energy imbalance and the resulting global warming. Indeed, our results point to the opposite: the evidence shows continued increase in cumulative CO₂ emissions, increased emissions of other GHGs, and gains in air quality at the expense of the loss of the cooling effect from aerosols. Both AR6 WGI and WGIII reports highlighted the benefits of short-term reductions in methane emissions to counter the loss of aerosol cooling and further improve air quality - however, at the global scale, methane emissions are at their highest level and rising (see Table 1). Policy makers, civil society and the scientific community require monitoring data and analyses from rigorous, robust assessments available on a regular basis. These results illustrate how assessments such as ours provide a strong “reality check” based on science and real-world data.

Deleted: These

Deleted: our assessment

Deleted: travel for

Deleted: high levels of greenhouse gas

Deleted: combined with improvements

Deleted: , are reducing

Deleted: level

Deleted: aerosol

Deleted: - leading to an unprecedented rate of human-induced warming....

Deleted:





Figure 8: Decadal trends in human-induced warming - left axis, and anthropogenic effective radiative forcing (ERF) - right axis. These are computed from the [GlobalWarming Index](#) human-induced warming estimate shown in [Supplementary Material Sect. 7](#), and Figure 2b respectively. The red points mark three additional years since the AR6 [time series](#) for these indicators ended in 2019.

This is a critical decade: [human-induced global](#) warming rates are at their highest historical level and 1.5 °C global warming might be expected to be reached or [exceeded](#) within the next 10 years [in the absence of cooling from major volcanic eruptions](#) ([Lee et al., 2021](#)). Yet this is also the decade that global greenhouse gas emissions could be expected to peak and begin to substantially decline. The indicators of global climate change presented here show that the Earth's energy imbalance has increased to around 0.9 W m⁻², averaged over the last 12 years. [This also has implications for the committed response of slow components in the climate system \(glaciers, deep ocean, ice sheets\) and committed long-term sea level rise but this is not part of the update here. However, rapid and stringent GHG emission decreases could halve warming rates over the next 20 years.](#)



Deleted:

Deleted: GWI

Deleted: Appendix B, Figure B2

Deleted: timeseries

Deleted: passed

Deleted: .

Deleted: This means that there are large energy flows into

Deleted: and rates of human induced warming will remain high as greenhouse gas emissions remain high. Nevertheless, these warming rates do not need to be locked in as

(McKenna et al., 2021). Table 1 shows that although global GHG emissions are at a long term high, their rate of increases have slowed. A continued series of these annual updates over this critical decade can track real world changes in direction for the human influence on climate.

<https://github.com/ClimateIndicator/anthropogenic-warming-assessment/>13. Acknowledgements

Piers Forster, Debbie Rosen, Joeri Rogelj and Robin Lamboll were supported by the EU Horizon 2020 Research and Innovation Programme grant no.820829 (CONSTRAIN). Chris Smith was supported by a NERC/IIASA collaborative research fellowship (NE/T009381/1). Matthew D. Palmer, Colin Morice and Rachel Killick were supported by the Met Office Hadley Centre Climate Programme funded by BEIS. William F. Lamb and Jan C. Minx were supported by the ERC-2020-SyG "GENIE" (grant ID 951542). Pierre Friedlingstein, Glen P. Peters and Robbie M. Andrew were supported by EU Horizon 2020 Research and Innovation Programme grant no. 821003 (4C). HadEX3 [3.0.4] data were obtained from <https://www.metoffice.gov.uk/hadobs/hadex3/> on 05.04.2023 and are © British Crown Copyright, Met Office, 2022, provided under an Open Government Licence <http://www.nationalarchives.gov.uk/doc/open-government-licence/version/2/>.

14. Author contributions

PMF, CJS, MA, PF, JR, MRC and AP developed the concept of an annual update in discussions with the wider IPCC community over many years. CJS led the work of the data repositories. A. Borger and JAB led the website development with visualisation support from DR, JMG and A. Birt. VMD, PZ, SS, JM, C-FS, SIS, VN, AP, J-YL, NG, FD, GP, BT, MSP, MRC, JR, PF, MA and PT provided important IPCC and UNFCCC framing. PMF coordinated the production of the manuscript with support from DR. WFL led Sect. 2 with contributions from CJS, JM, PF, GP, JG, JP and RA. CJS led Sects. 3 and 4 with contributions from BH, FD, SS, VN and XL. BT led Sect. 5 with contributions from PT, CM, CK, JK, RR, RV and LC. Kvs and MDP led Sect. 6 with contributions from LC, MI, TB and RK. TW led Sect. 7 with contributions and calculations from AR, NG and MR. JR led Sect. 8 with contributions from RL and KZ. Sect. 9 was led by SIS and XC with calculations by MH and DS. All authors either edited or commented on the manuscript.

15. Competing interests

The authors declare no competing interests.

Deleted: .

Deleted: greenhouse gas

Deleted: they are beginning to stabilise, giving some hope that

Deleted: time the indicators of global climate change presented here

Deleted: a

Deleted: -

Deleted: change

Deleted: Appendix A Climate and weather extremes - offset calculation

Berkeley Earth — Land mean TXx




Figure A1: Calculation of land mean annual maximum temperature (TXx) offset between 1850-1900 and 1961-1990. A linear regression of TXx as a function of global mean temperature from Berkeley Earth is fitted to data from 1955-2020. The TXx offset of 0.53 °C is then obtained by multiplying the slope of the linear regression (1.25 °C / °C) with the global mean temperature difference between 1850-1900 and 1961-1990 (0.43°C).

... [113]

Deleted: .

Deleted: RL

Deleted: JR

References

- Adusumilli, S., Straneo, F., Hendricks, S., Korosov, A., Lavergne, T., Lawrence, I., Marzeion, B., Otosaka, I., Schweiger, A., Shepherd, A., Slater, D., Slater, T., Timmermanns, M.-L., and Zemp, M.: GCOS EHI 1960-2020 Cryosphere Heat Content, https://doi.org/10.26050/WDC/GCOS_EHI_1960-2020_CRHC, 2022.
- <https://doi.org/10.1007/s00382-003-0313-9> Allen, M. R., O. P. Dube, W. Solecki, F. Aragón-Durand, W. Cramer, S. Humphreys, M. Kainuma, J. Kala, N. Mahowald, Y. Mulugetta, R. Perez, M. Wairiu, and K. Zickfeld, 2018: Framing and Context. In: Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty [Masson-Delmotte, V., P. Zhai, H.-O. Pörtner, D. Roberts, J. Skea, P.R. Shukla, A. Pirani, W. Moufouma-Okia, C. Péan, R. Pidcock, S. Connors, J.B.R. Matthews, Y. Chen, X. Zhou, M.I. Gomis, E. Lonnoy, T. Maycock, M. Tignor, and T. Waterfield (eds.)], Cambridge University Press, Cambridge, UK and New York, NY, USA, 49-92, <https://doi.org/10.1017/9781009157940.003>, 2018.
- Allison, L. C., Palmer, M. D., Allan, R. P., Hermanson, L., Liu, C., and Smith, D. M.: Observations of planetary heating since the 1980s from multiple independent datasets, *Environ. Res. Commun.*, 2, 101001, <https://doi.org/10.1088/2515-7620/abbb39>, 2020.
- <https://doi.org/10.1039/c2cp20017a> Basu, S., Jan, X., Dlugokencky, E., Michel, S., Schwietzke, S., Miller, J. B., Bruhwiler, L., Oh, Y., Tans, P. P., Apudula, F., Gatti, L. V., Jordan, A., Necki, J., Sasakawa, M., Morimoto, S., Di Iorio, T., Lee, H., Arduini, J., and Manca, G.: Estimating emissions of methane consistent with atmospheric measurements of methane and $\delta^{13}\text{C}$ of methane. *Atmos. Chem. Phys.*, 22, 15351–15377, 2022.
- Bellouin, N., Davies, W., Shine, K. P., Quaas, J., Mülmenstädt, J., Forster, P. M., Smith, C., Lee, L., Regayre, L., Brasseur, G., Sudarchikova, N., Bouarar, I., Boucher, O., and Myhre, G.: Radiative forcing of climate change from the Copernicus reanalysis of atmospheric composition, *Earth Syst. Sci. Data*, 12, 1649–1677, <https://doi.org/10.5194/essd-12-1649-2020>, 2020.
- Beusch, L., Gudmundsson, L., and Seneviratne, S. I.: Crossbreeding CMIP6 Earth System Models With an Emulator for Regionally Optimized Land Temperature Projections, *Geophys. Res. Lett.*, 47, <https://doi.org/10.1029/2019GL086812>, 2020.
- Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment, *J. Geophys. Res.-Atmos.*, 118, 5380–5552, <https://doi.org/10.1002/jgrd.50171>, 2013.

Deleted: 15.

Deleted: Allen, M. R. and Stott, P. A.: Estimating signal amplitudes in optimal fingerprinting, part I: theory, *Climate Dynamics*, 21, 477–491,

Deleted: , 2003.

Moved (insertion) [87]

Moved (insertion) [88]

Moved (insertion) [89]

Moved (insertion) [90]

Byers, E., Krey, V., Kriegler, E., Riahi, K., Schaeffer, R., Kikstra, J., Lamboll, R., Nicholls, Z., Sandstad, M., Smith, C., van der Wijst, K., Lecocq, F., Portugal-Pereira, J., Saheb, Y., Stromann, A., Winkler, H., Auer, C., Brutschin, E., Lepault, C., Müller-Casseres, E., Gidden, M., Huppmann, D., Kolp, P., Marangoni, G., Werning, M., Calvin, K., Guivarch, C., Hasegawa, T., Peters, G., Steinberger, J., Tavoni, M., van Vuuren, D., Al-Khourdajie, A., Forster, P., Lewis, J., Meinshausen, M., Rogelj, J., Samset, B., and Skeie, R.: AR6 Scenarios Database, <https://doi.org/10.5281/ZENODO.5886912>, 2022.

Canadell, J.G., P. M. S. Monteiro, M. H. Costa, L. Cotrim da Cunha, P. M. Cox, A.V. Eliseev, S. Henson, M. Ishii, S. Jaccard, C. Koven, A. Lohila, P. K. Patra, S. Piao, J. Rogelj, S. Syampungani, S. Zaehle, and K. Zickfeld: Global Carbon and other Biogeochemical Cycles and Feedbacks. In Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 673–816, <https://doi.org/10.1017/9781009157896.007>, 2021.

Cheng, L., Trenberth, K. E., Fasullo, J., Boyer, T., Abraham, J., and Zhu, J.: Improved estimates of ocean heat content from 1960 to 2015, *Sci. Adv.*, 3, e1601545, <https://doi.org/10.1126/sciadv.1601545>, 2017.

<https://climateactiontracker.org/global/cat-emissions-gaps/> Cheng, L., Abraham, J., Hausfather, Z., and Trenberth, K. E.: How fast are the oceans warming?. *Science*, 363, 128–129, , 2019.

Cheng, L., Von Schuckmann, K., Abraham, J. P., Trenberth, K. E., Mann, M. E., Zanna, L., England, M. H., Zika, J. D., Fasullo, J. T., Yu, Y., Pan, Y., Zhu, J., Newsom, E. R., Bronselaer, B., and Lin, X.: Past and future ocean warming, *Nat. Rev. Earth. Environ.*, 3, 776–794, , 2022.

Collins, M., Knutti, R., Arblaster, J., Dufresne, J.-L., Fichfet, T., Friedlingstein, P., Gao, X., Gutowski, W.J., Johns, T., Krinner, G., Shongwe, M., Tebaldi, C., Weaver, A.J. & Wehner, M.: Long-term Climate Change: Projections, Commitments and Irreversibility. In: V.B. Stocker T.F., .D. Qin, G.K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia & P.M. Midgley (eds.). Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge, United Kingdom and New York, NY, USA, Cambridge University Press. pp. 1029–1136, 2013.

Cowtan, K. <https://doi.org/10.1002/qj.2297>, Hausfather, Z., Hawkins, E., Jacobs, P., Mann, M. E., Miller, S. K., Steinman, B. A., Stolpe, M. B., and Way, R. G.: Robust comparison of climate models with observations using blended land air and ocean sea surface temperatures, *Geophys. Res. Lett.*, 42, 6526–6534, , 2015.

Crippa, M., Guizzardi, D., Banja, M., Solazzo, E., Muntean, M., Schaaf, E., Pagani, F., Monforti-Ferrario, F., Olivier, J. G. J., Quadrelli, R., Riskez Martin, A. , Taghavi-Moharamli, P., Grassi, G., Rossi, S., Oom, D., Branco, A., San-Miguel, J.,

- Moved down [91]:** K.
- Moved down [92]:** Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.
- Moved down [93]:** Lonnoy, J.
- Moved down [94]:** Matthews, T.
- Moved down [95]:** Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp.
- Deleted:** Chen, D., M. Rojas, B. H. Samset, K. Cobb, A. Diongue Niang, P. Edwards, S. Emori, S. H. Faria, E. Hawkins, P. Hope, P. Huybrechts, M. Meinshausen, S.
- Deleted:** Mustafa, G.-K. Plattner, and A.-M. Tréguier. 2021: Framing, Context, and Methods. In Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I.
- Deleted:** B. R.
- Deleted:** K.
- Deleted:** 147–286,
- Deleted:** , 2021.¶
- Moved (insertion) [91]**
- Deleted:** Climate Action Tracker, CAT Emissions Gap:
- Deleted:** , last access: 26 April 2023.¶
- Moved (insertion) [96]**
- Moved (insertion) [97]**
- Moved (insertion) [98]**
- Deleted:** . and Way, R. G.: Coverage bias in the HadCRUT4 temperature series and its impact on recent temperature trends, *Q.J.R. Meteorol. Soc.*, 140, 1935–1944,
- Deleted:** , 2014.¶
Cowtan, K
- Deleted:** <https://doi.org/10.1002/2015GL064888>,

Vignati, E.: CO₂ emissions of all world countries: JRC/IEA/PBL 2022 report, Publications Office, LU, <https://doi.org/10.2760/07904>, 2022.

Cuesta-Valero, F. J., García-García, A., Beltrami, H., González-Rouco, J. F., and García-Bustamante, E.: Long-term global ground heat flux and continental heat storage from geothermal data, *Clim. Past*, 17, 451–468, <https://doi.org/10.5194/cp-17-451-2021>, 2021.

Cuesta-Valero, F. J., Beltrami, H., García-García, A., Krinner, G., Langer, M., MacDougall, A. H., Nitzbon, J., Peng, J., von Schuckmann, K., Seneviratne, S. I., Smith, N., Thiery, W., Vanderkelen, I., and Wu, T.: Continental heat storage: Contributions from ground, inland waters, and permafrost thawing, *Earth Syst. Dynam. Discuss.* [preprint], <https://doi.org/10.5194/esd-2022-32>, 2022.

Cuesta-Valero, F. J., Beltrami, H., García-García, A., Krinner, G., Langer, M., MacDougall, A., Nitzbon, J., Peng, J., von Schuckmann, K., Seneviratne, S., Thiery, W., Vanderkelen, I., Wu, T.: GCOS EHI 1960-2020 Continental Heat Content (Version 2), World Data Center for Climate (WDCC) at DKRZ, https://doi.org/10.26050/WDCC/GCOS_EHI_1960-2020_CoHC_v2, 2023.

Dhakal, S., J. C. Minx, F. L. Toth, A. Abdel-Aziz, M. J. Figueroa Meza, K. Hubacek, I. G. C. Jonckheere, Yong-Gun Kim, G. F. Nemet, S. Pachauri, X. C. Tan, T. Wiedmann: Emissions Trends and Drivers. In IPCC, 2022: Climate Change 2022: Mitigation of Climate Change. Contribution of Working Group III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [P.R. Shukla, J. Skea, R. Slade, A. Al Khourdajie, R. van Diemen, D. McCollum, M. Pathak, S. Some, P. Vyas, R. Fradera, M. Belkacemi, A. Hasija, G. Lisboa, S. Luz, J. Malley, (eds.)]. Cambridge University Press, Cambridge, UK and New York, NY, USA, <https://doi.org/10.1017/9781009157926.004>, 2022.

Domingues, C. M., Church, J. A., White, N. J., Gleckler, P. J., Wijffels, S. E., Barker, P. M., and Dunn, J. R.: Improved estimates of upper-ocean warming and multi-decadal sea-level rise, *Nature*, 453, 1090–1093, <https://doi.org/10.1038/nature07080>, 2008.

Douville, H., K. Raghavan, J. Renwick, R. P. Allan, P. A. Arias, M. Barlow, R. Cerezo-Mota, A. Cherchi, T. Y. Gan, J. Gergis, D. Jiang, A. Khan, W. Pokam Mba, D. Rosenfeld, J. Tierney, and O. Zolina: *Water Cycle Changes*. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekci, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1055–1210, . 2021.

Droste, E. S., Adcock, K. E., Ashfold, M. J., Chou, C., Fleming, Z., Fraser, P. J., Gooch, L. J., Hind, A. J., Langenfelds, R. L., Leedham Elvidge, E. C., Mohd Hanif, N., O'Doherty, S., Oram, D. E., Ou-Yang, C.-F., Panagi, M., Reeves, C. E., Sturges, W.

- Moved down [99]: . M.,
- Moved down [100]: H.,
- Deleted: Dessler, A. E., Schoeberl, M. R., Wang, T., Davis, S
- Deleted: Rosenlof, K.
- Deleted: and Vernier, J.-P.: Variations of stratospheric water vapor over the past three decades, *J. Geophys. Res.-Atmos.*, 119, 12588–12598,
- Deleted: , 2014.¶
- Moved down [101]: S.,
- Moved up [97]: E.,
- Moved down [102]: P.,
- Moved up [96]: L.,
- Moved up [89]: Chem. Phys.,
- Moved up [90]: Phys.,
- Deleted: Dhomse, S.
- Deleted: Mann, G. W., Antuña Marrero, J. C., Shallcross, S.
- Deleted: Chipperfield, M.
- Deleted: Carslaw, K. S., Marshall, L., Abraham, N.
- Deleted: and Johnson, C. E.: Evaluating the simulated radiative forcings, aerosol properties, and stratospheric warmings from the 1963 Mt Agung, 1982 El Chichón, and 1991 Mt Pinatubo volcanic aerosol clouds, *Atmos.*
- Deleted: 20, 13627–13654,
- Deleted: , 2020.¶
- Moved (insertion) [100]
- Moved (insertion) [103]
- Moved (insertion) [92]
- Moved (insertion) [93]
- Moved (insertion) [94]
- Moved (insertion) [95]

T., and Laube, J. C.: Trends and emissions of six perfluorocarbons in the Northern Hemisphere and Southern Hemisphere, *Atmos. Chem. Phys.*, 20, 4787–4807, <https://doi.org/10.5194/acp-20-4787-2020>, 2020.

Dunn, R. J. H., Alexander, L. V., Donat, M. G., Zhang, X., Bador, M., Herold, N., Lippmann, T., Allan, R., Aguilar, E., Barry, A. A., Brunet, M., Caesar, J., Chagnaud, G., Cheng, V., Cinco, T., Durre, I., Guzman, R., Htay, T. M., Wan Ibadullah, W. M., Bin Ibrahim, M. K. I., Khoshkam, M., Kruger, A., Kubota, H., Leng, T. W., Lim, G., Li-Sha, L., Marengo, J., Mbatha, S., McGree, S., Menne, M., Milagros Skansi, M., Ngwenya, S., Nkrumah, F., Oonariya, C., Pabon-Caicedo, J. D., Panthou, G., Pham, C., Rahimzadeh, F., Ramos, A., Salgado, E., Salinger, J., Sané, Y., Sopaheluwakan, A., Srivastava, A., Sun, Y., Timbal, B., Trachow, N., Trewin, B., Schrier, G., Vazquez-Aguirre, J., Vasquez, R., Villarroel, C., Vincent, L., Vischel, T., Vose, R., and Bin Hj Yussof, M. N.: Development of an updated global land in situ-based data set of temperature and precipitation extremes: HadEX3, *J. Geophys. Res.-Atmos.*, 125, e2019JD032263, <https://doi.org/10.1029/2019JD032263>, 2020.

Dunn, R. J. H., Donat, M. G., and Alexander, L. V.: Comparing extremes indices in recent observational and reanalysis products, *Front. Clim.*, 4, 98905, <https://doi.org/10.3389/fclim.2022.989505>, 2022.

<https://doi.org/10.1002/2016GL071930> Eyring, V., N. P. Gillett, K.M. Achuta Rao, R. Barimalala, M. Barreiro Parrillo, N. Bellouin, C. Cassou, P. J. Durack, Y. Kosaka, S. McGregor, S. Min, O. Morgenstern, and Y. Sun: Human Influence on the Climate System. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 423–552, <http://doi:10.1017/9781009157896.005>, 2021.

Forster, P. M., Forster, H. I., Evans, M. J., Gidden, M. J., Jones, C. D., Keller, C. A., Lamboll, R. D., Le Quéré, C., Rogelj, J., Rosen, D., Schleussner, C. F., Richardson, T. B., Smith, C. J. and Turnock, S. T.: Current and future global climate impacts resulting from COVID-19, *Nature Clim. Chang.*, 10, 913–919, <https://doi.org/10.1038/s41558-020-0883-0>, 2020.

Forster, P., T. Storelvmo, K. Armour, W. Collins, J.-L. Dufresne, D. Frame, D.J. Lunt, T. Mauritsen, M.D. Palmer, M. Watanabe, M. Wild, and H. Zhang, 2021: The Earth's Energy Budget, Climate Feedbacks, and Climate Sensitivity. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 923–1054, <https://doi.org/10.1017/9781009157896.009>, 2021.

Fox-Kemper, B., Fox-Kemper, B., H. T. Hewitt, C. Xiao, G. Aðalgeirsdóttir, S.S. Drijfhout, T. L. Edwards, N. R. Golledge, M. Hemer, R. E. Kopp, G. Krinner, A. Mix, D. Notz, S. Nowicki, I. S. Nurhati, L. Ruiz, J.-B. Sallée, A. B. A. Slangen, and Y.

Deleted: Etmiman, M., Myhre, G., Highwood, E. J., and Shine, K. P.: Radiative forcing of carbon dioxide, methane, and nitrous oxide: A significant revision of the methane radiative forcing, *Geophys. Res. Lett.*, 43, 12614–12623,

Deleted: , 2016.

Deleted: doi:10.1017/9781009157896.009,

- Yu: Ocean, Cryosphere and Sea Level Change. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1211–1362, <https://doi.org/10.1017/9781009157896.011>, 2021.
- 1200 Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Le Quéré, C., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S., Aragão, L. E. O. C., Arneth, A., Arora, V., Bates, N. R., Becker, M., Benoit-Cattin, A., Bittig, H. C., Bopp, L., Bultan, S., Chandra, N., Chevallier, F., Chini, L. P., Evans, W., Florentie, L., Forster, P. M., Gasser, T., Gehlen, M., Gilfillan, D., Gkritzalis, T., Gregor, L., Gruber, N., Harris, I., Hartung, K., Haverd, V., Houghton, R. A., Ilyina, T., Jain, A. K., Joetzjer, E., Kadono, K., Kato, E., Kitidis, V., Korsbakken, J. I.,
- 1205 Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Liu, Z., Lombardozi, D., Marland, G., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., Niwa, Y., O’Brien, K., Ono, T., Palmer, P. I., Pierrot, D., Poulter, B., Resplandy, L., Robertson, E., Rödenbeck, C., Schwinger, J., Séférian, R., Skjelvan, I., Smith, A. J. P., Sutton, A. J., Tanhua, T., Tans, P. P., Tian, H., Tilbrook, B., van der Werf, G., Vuichard, N., Walker, A. P., Wanninkhof, R., Watson, A. J., Willis, D., Wiltshire, A. J., Yuan, W., Yue, X., and Zaehle, S.: Global carbon budget 2020, *Earth Syst. Sci. Data*, 12, 3269–3340, <https://doi.org/10.5194/essd-12-3269-2020>, 2020.
- 1210 Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Gregor, L., Hauck, J., Le Quéré, C., Luijkx, I. T., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Alkama, R., Arneth, A., Arora, V. K., Bates, N. R., Becker, M., Bellouin, N., Bittig, H. C., Bopp, L., Chevallier, F., Chini, L. P., Cronin, M., Evans, W., Falk, S., Feely, R. A., Gasser, T., Gehlen, M., Gkritzalis, T., Gloege, L., Grassi, G., Gruber, N.,
- 1215 Gürses, Ö., Harris, I., Hefner, M., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Jain, A. K., Jersild, A., Kadono, K., Kato, E., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Landschützer, P., Lefèvre, N., Lindsay, K., Liu, J., Liu, Z., Marland, G., Mayot, N., McGrath, M. J., Metzl, N., Monacci, N. M., Munro, D. R., Nakaoka, S.-I., Niwa, Y., O’Brien, K., Ono, T., Palmer, P. I., Pan, N., Pierrot, D., Pockock, K., Poulter, B., Resplandy, L., Robertson, E., Rödenbeck, C., Rodriguez, C., Rosan, T. M., Schwinger, J., Séférian, R., Shutler, J. D., Skjelvan, I., Steinhoff, T., Sun, Q., Sutton, A. J., Sweeney, C.,
- 1220 Takao, S., Tanhua, T., Tans, P. P., Tian, X., Tian, H., Tilbrook, B., Tsujino, H., Tubiello, F., van der Werf, G. R., Walker, A. P., Wanninkhof, R., Whitehead, C., Willstrand Wranne, A., et al.: Global Carbon Budget 2022, *Earth Syst. Sci. Data*, 14, 4811–4900, <https://doi.org/10.5194/essd-14-4811-2022>, 2022a, <https://doi.org/10.1029/2005JD006019> ~~Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Gregor, L., Hauck, L., Le Quéré, C., Luijkx, I. T., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell,~~

Deleted: , 2022

Moved (insertion) [102]

Moved (insertion) [99]

Deleted: Fueglistaler, S. and Haynes, P. H.: Control of interannual and longer-term variability of stratospheric water vapor, *J. Geophys. Res.*, 110, D24108,

Deleted: , 2005.†

Moved (insertion) [104]

230 [J. G., Ciais, P., Jackson, R. B., Alin, S., Alkama, R., Arneeth, A., Arora, V. K., Bates, N. R., Becker, M., Bellouin, N., Bittig, H. C., Bopp, L., Chevallier, F., Chini, L. P., Cronin, M., Evans, W., Falk, S., Feely, R. A., Gasser, T., Gehlen, M., Gkritzalis, T., Gloege, L., Grassi, G., Gruber, N., Gürses, Ö., Harris, I., Hefner, M., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Jain, A. T., Jersild, A., Kadono, K., Kato, E., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. J., Landschützer, P., Lefèvre, N., Lindsay, Keith., Liu, J., Marland, G., Mayot, N., McGrath, M. J., Metz, N., Monacci, N. M., Munro, D. R., Nakaoka, S.-I., Niwa, Y., O'Brien, K., Ono, T., Palmer, P. I., Pan, N., Pierrot, D., Pocock, K., Poulter, B., Resplandy, L., Robertson, E., Rödenbeck, C., Rodriguez, C., Rosan, T. M., Schwinger, J., Séférian, R., Shutler, J. D., Skjelvan, I., Steinhoff, T., Sun, Q., Sutton, A. J., Sweeney, C., Takao, S., Tanhua, T., Tans, P. P., Tian, X., Tian, H., Tilbrook, B., Tsujino, H., Tubiello, F., van der Werf, G. R., Walker, A. P., Wanninkhof, R., Whitehead, C., Wranne, A., Wright, R. M., Yuan, W., Yue, C., Yue, X., Zaehle, S., Zeng, J., Zheng, B. and Zhu, L.:](#) Supplemental data of the Global Carbon Budget 2022, ICOS-ERIC Carbon Portal [data set], . 2022b.

235 Gasser, T., Crepin, L., Quilcaille, Y., Houghton, R. A., Ciais, P., and Obersteiner, M.: Historical CO2 emissions from land use and land cover change and their uncertainty, *Biogeosciences*, 17, 4075–4101, <https://doi.org/10.5194/bg-17-4075-2020>, 2020.

240 Gillett, N. P., Shiogama, H., Funke, B., Hegerl, G., Knutti, R., Matthes, K., Santer, B. D., Stone, D., and Tebaldi, C.: The Detection and Attribution Model Intercomparison Project (DAMIP v1.0) contribution to CMIP6, *Geosci. Model. Dev.*, 9, 3685–3697, <https://doi.org/10.5194/gmd-9-3685-2016>, 2016.

245 Gillett, N.P., Kirchmeier-Young, M., Ribes, A., Shiogama, H., Hegerl, G.C., Knutti, R., Gastineau, G., John, J.G., Li, L., Nazarenko, L., Rosenbloom, N., Seland, Ø., Wu, T., Yukimoto, S., and Ziehn, T.: Constraining human contributions to observed warming since the pre-industrial period, *Nat. Clim. Chang.*, 11, 207–212, <https://doi.org/10.1038/s41558-020-00965-9>, 2021.

250 Gleckler, P. J., Durack, P. J., Stouffer, R. J., Johnson, G. C., and Forest, C. E.: Industrial-era global ocean heat uptake doubles in recent decades, *Nat. Clim. Chang.*, 6, 394–398, <https://doi.org/10.1038/nclimate2915>, 2016.

255 <https://www.globalfiredata.org/> Good, S. A., Martin, M. J., and Rayner, N. A.: EN4: Quality controlled ocean temperature and salinity profiles and monthly objective analyses with uncertainty estimates, THE EN4 DATA SET, *J. Geophys. Res.-Oceans*, 118, 6704–6716, <https://doi.org/10.1002/2013JC009067>, 2013.

260 [Grassi, G., Schwingshackl, C., Gasser, T., Houghton, R. A., Sitch, S., Canadell, J. G., Cescatti, A., Ciais, P., Federici, S., Friedlingstein, P., Kurz, W. A., Sanz Sanchez, M. J., Abad Viñas, R., Alkama, R., Bultan, S., Ceccherini, G., Falk, S., Kato, E., Kennedy, D., Knauer, J., Korosuo, A., Melo, J., McGrath, M. J., Nabel, J. E. M. S., Poulter, B., Romanovskaya, A. A., Rossi, S., Tian, H., Walker, A. P., Yuan, W., Yue, X., and Pongratz, J.:](#) Harmonising the land-use flux estimates of global models and national inventories for 2000–2020, *Earth Syst. Sci. Data*, 15, 1093–1114, . 2023.

Moved (insertion) [101]

Moved (insertion) [105]

Moved (insertion) [106]

Moved (insertion) [107]

Moved (insertion) [108]

Moved (insertion) [109]

Moved (insertion) [110]

Moved (insertion) [111]

Moved (insertion) [112]

Moved (insertion) [113]

Moved (insertion) [114]

Deleted: .

Moved (insertion) [115]

Deleted: Nature Climate Change,

Deleted: Global fire emissions database,

Deleted: , last accessed 24 April 2023.

Moved (insertion) [116]

Moved (insertion) [117]

Moved (insertion) [118]

Moved (insertion) [119]

Moved (insertion) [120]

- 4265 Guevara, M., Petetin, H., Jorba, O., Denier van der Gon, H., Kuenen, J., Super, I., Granier, C., Doumbia, T., Ciais, P., Liu, Z., Lamboll, R. D., Schindlbacher, S., Matthews, B., and Pérez García-Pando, C.: Towards near-real time air pollutant and greenhouse gas emissions: lessons learned from multiple estimates during the COVID-19 Pandemic, *EGUsphere* [preprint], 2023, 1–36, <https://doi.org/10.5194/egusphere-2023-186>, 2023.
- Gulev, S. K., P. W. Thorne, J. Ahn, F. J. Dentener, C. M. Domingues, S. Gerland, D. Gong, D. S. Kaufman, H. C. Nnamchi, 4270 J. Quaas, J.A. Rivera, S. Sathyendranath, S.L. Smith, B. Trewin, K. von Schuckmann, and R. S. Vose: Changing State of the Climate System. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*[Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom 4275 and New York, NY, USA, pp. 287–422, <https://doi.org/10.1017/9781009157896.004>, 2021.
- Gutiérrez, J. M., R. G. Jones, G. T. Narisma, L. M. Alves, M. Amjad, I. V. Gorodetskaya, M. Grose, N. A. B. Klutse, S. Krakovska, J. Li, D. Martínez-Castro, L. O. Mearns, S. H. Mernild, T. Ngo-Duc, B. van den Hurk, and J.-H. Yoon: Atlas. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. 4280 Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1927–2058, <https://doi.org/10.1017/9781009157896.021>, 2021. Note: The companion Interactive Atlas is available at <http://interactive-atlas.ipcc.ch>
- Gütschow, J., Jeffery, M. L., Gieseke, R., Gebel, R., Stevens, D., Krapp, M., and Rocha, M.: The PRIMAP-hist national 4285 historical emissions time series, *Earth Syst. Sci. Data*, 8, 571–603, <https://doi.org/10.5194/essd-8-571-2016>, 2016.
- Gütschow, J., and Pflüger, M.: The PRIMAP-hist national historical emissions time series (1750-2021) v2.4.1 (2.4.1), Zenodo [data set], <https://doi.org/10.5281/zenodo.7585420>, 2023.
- Hakuba, M. Z., Frederikse, T., and Landerer, F. W.: Earth's energy imbalance from the ocean perspective (2005–2019), *Geophys Res Lett*, 48, e2021GL093624, <https://doi.org/10.1029/2021GL093624>, 2021.
- 4290 Hall, B. D., Crotwell, A. M., Kitzis, D. R., Mefford, T., Miller, B. R., Schibig, M. F., and Tans, P. P.: Revision of the World Meteorological Organization Global Atmosphere Watch (WMO/GAW) CO₂ calibration scale, *Atmos. Meas. Tech.*, 14, 3015–3032, <https://doi.org/10.5194/amt-14-3015-2021>, 2021.
- Hansis, E., Davis, S. J., and Pongratz, J.: Relevance of methodological choices for accounting of land use change carbon fluxes, *Global Biogeochem. Cy.*, 29, 1230–1246, <https://doi.org/10.1002/2014GB004997>, 2015.

4295 Haustein, K., Allen, M. R., Forster, P. M., Otto, F. E. L., Mitchell, D. M., Matthews, H. D., and Frame, D. J.: A real-time Global Warming Index, *Sci Rep*, 7, 15417, <https://doi.org/10.1038/s41598-017-14828-5>, 2017.

Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., 4300 Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Q. J. R. Meteorol. Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.

<https://doi.org/10.1038/s41612-020-00150-x>Hodnebrog, Ø., Aamaas, B., Fuglestedt, J. S., Marston, G., Myhre, G., Nielsen, C. J., Sandstad, M., Shine, K. P., and Wallington, T. J.: Updated Global Warming Potentials and Radiative Efficiencies 4305 of Halocarbons and Other Weak Atmospheric Absorbers, *Rev. Geophys.*, 58, e2019RG000691, <https://doi.org/10.1029/2019RG000691>, 2020.

Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M., Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J.-I., Li, M., Liu, L., Lu, Z., Moura, M. C. P., O'Rourke, P. R., and Zhang, Q.: Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community 4310 Emissions Data System (CEDS), *Geosci. Model. Dev.*, 11, 369–408, <https://doi.org/10.5194/gmd-11-369-2018>, 2018.

Houghton, R. A., and Nassikas, A. A.: Global and regional fluxes of carbon from land use and land cover change 1850–2015, *Global Biogeochem. Cy.*, 31, 456–472, <https://doi.org/10.1002/2016GB005546>, 2017.

<https://doi.org/10.1029/2010JD015065><https://www.iata.org/en/iata-repository/publications/economic-reports/airline-industry-economic-performance---june-2022---report><https://doi.org/10.1787/558987b9-en>IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report 4315 of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp, <https://doi:10.1017/CBO9781107415324>, 2013.

IPCC: Summary for Policymakers. In: Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming 4320 of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty [Masson-Delmotte, V., P. Zhai, H.-O. Pörtner, D. Roberts, J. Skea, P.R. Shukla, A. Pirani, W. Moufouma-Okia, C. Péan, R. Pidcock, S. Connors, J.B.R. Matthews, Y. Chen, X. Zhou, M.I. Gomis, E. Lonnoy, T. Maycock, M. Tignor, and T. Waterfield (eds.)]. Cambridge University Press, Cambridge, UK and New York, NY, USA, pp. 3-24, <https://doi.org/10.1017/9781009157940.001>, 4325 2018.

Deleted: Hodnebrog, Ø., Myhre, G., Kramer, R. J., Shine, K. P., Andrews, T., Faluvegi, G., Kasoar, M., Kirkevåg, A., Lamarque, J.-F., Mülmenstädt, J., Olivé, D., Samsat, B. H., Shindell, D., Smith, C. J., Takemura, T., and Voulgarakis, A.: The effect of rapid adjustments to halocarbons and N2O on radiative forcing, *npj Clim. Atmos. Sci.*, 3, 43,

Deleted: , 2020a.¶

Deleted: , 2020b

Moved up [88]: F.,

Moved up [120]: H.,

Moved up [106]: M.,

Moved up [118]: A.,

Moved up [116]: G.,

Deleted: Hurst, D.

Deleted: Oltmans, S. J., Vömel, H., Rosenlof, K.

Deleted: Davis, S.

Deleted: Ray, E.

Deleted: Hall, E.

Deleted: and Jordan, A. F.: Stratospheric water vapor trends over Boulder, Colorado: Analysis of the 30 year Boulder record, *J. Geophys. Res.-Atmos.*, 116,

Deleted: , 2011.¶

IATA: Global Outlook for Air Transport: Times of Turbulence, IATA,

Deleted: , 2022. ¶
IEA: World oil statistics (Edition 2021), IEA Oil Information Statistics (database),

Deleted: , 2022 (accessed on 24 April 2023).¶

1355 IPCC: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896>, 2021a.

1360 IPCC: Summary for Policymakers, in: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp.3–32, <https://doi.org/10.1017/9781009157896.001>, 2021b.

1365 IPCC: Annex III: Tables of historical and projected well-mixed greenhouse gas mixing ratios and effective radiative forcing of all climate forcers [Dentener F.J., B. Hall, C. Smith (eds.)]. In Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 2139–2152, <https://doi.org/10.1017/9781009157896.017>, 2021c.

1370 IPCC: Climate Change 2022: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A. Alegría, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, B. Rama (eds.)]. Cambridge University Press. Cambridge University Press, Cambridge, UK and New York, NY, USA, 3056 pp., <https://doi.org/10.1017/9781009325844>, 2022.

1375 Ishii, M., Fukuda, Y., Hirahara, S., Yasui, S., Suzuki, T., and Sato, K.: Accuracy of Global Upper Ocean Heat Content Estimation Expected from Present Observational Data Sets, SOLA, 13, 163–167, <https://doi.org/10.2151/sola.2017-030>, 2017.

1380 Iturbide, M., Fernández, J., Gutiérrez, J. M., Pirani, A., Huard, D., Al Khourdajie, A., Baño-Medina, J., Bedia, J., Casanueva, A., Cimadevilla, E., Cofiño, A. S., De Felice, M., Diez-Sierra, J., García-Diez, M., Goldie, J., Herrera, D. A., Herrera, S., Manzanas, R., Milovac, J., Radhakrishnan, A., San-Martín, D., Spinuso, A., Thyng, K. M., Trenham, C., and Yelekçi, Ö.: Implementation of FAIR principles in the IPCC: the WGI AR6 Atlas repository, Sci Data, 9, 629, <https://doi.org/10.1038/s41597-022-01739-y>, 2022.

Jenkins, S., Smith, C., Allen, M., and Grainger, R.: Tonga eruption increases chance of temporary surface temperature anomaly above 1.5 °C, Nature Clim. Chang., 13, 127–129, <https://doi.org/10.1038/s41558-022-01568-2>, 2023.

Deleted: .

<https://doi.org/10.5194/acp-9-6109-2009><https://doi.org/10.5194/gmd-10-4005-2017> Kadow, C., Hall, D. M., and Ulbrich, U.: Artificial intelligence reconstructs missing climate information, *Nat. Geosci.*, 13, 408–413, <https://doi.org/10.1038/s41561-020-0582-5>, 2020.

<https://doi.org/10.5194/acp-21-5015-2021> Kirchengast, G., Gorfer, M., Mayer, M., Steiner, A. K., and Haimberger, L.: GCOS EHI 1960-2020 Atmospheric Heat Content, https://doi.org/10.26050/WDCC/GCOS_EHI_1960-2020_AHC, 2022.

<https://doi.org/10.5194/essd-12-2607-2020> Kramer, R. J., He, H., Soden, B. J., Oreopoulos, L., Myhre, G., Forster, P. M., and Smith, C. J., Observational evidence of increasing global radiative forcing, *Geophys. Res. Lett.*, 48, e2020GL091585, <https://doi.org/10.1029/2020GL091585>, 2021.

Lamboll, R. D. and Rogelj, J.: Code for estimation of remaining carbon budget in IPCC AR6 WGI, Zenodo [code], <https://doi.org/10.5281/zenodo.6373365>, 2022.

Lan, X., Tans, P. and Thoning, K.W.: Trends in globally-averaged CO2 determined from NOAA Global Monitoring Laboratory measurements, Version 2023-04, <https://doi.org/10.15138/9N0H-ZH07>, 2023a.

Lan, X., Thoning, K. W., and Dlugokencky, E.J.: Trends in globally-averaged CH4 N2O, and SF6 determined from NOAA Global Monitoring Laboratory measurements, Version 2023-04, <https://doi.org/10.15138/P8XG-AA10>, 2023b.

Laube, J., Newland, M., Hogan, C., Brenninkmeijer, A.M., Fraser, P.J., Martinerie, P., Oram, D.E., Reeves, C.E., Röckmann, T., Schwander, J., Witrant, E., Sturges, W.T.: Newly detected ozone-depleting substances in the atmosphere. *Nature Geosci.*, 7, 266–269, <https://doi.org/10.1038/ngeo2109>, 2014.

<https://doi.org/10.5194/gmd-14-3007-2021>, <https://doi.org/10.1016/j.atmosenv.2020.117834> Lee, J.-Y., J. Marotzke, G. Bala, L. Cao, S. Corti, J.P. Dunne, F. Engelbrecht, E. Fischer, J.C. Fyfe, C. Jones, A. Maycock, J. Mutemi, O. Ndiaye, S. Panickal, and T. Zhou: Future Global Climate: Scenario-Based Projections and Near-Term Information. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 553–672, <https://doi.org/10.1017/9781009157896.006>, 2021.

Lee, H., K. Calvin, D. Dasgupta, G. Krinner, A. Mukherji, P. Thorne, C. Trisos, J. Romero, P. Aldunce, K. Barrett, G. Blanco, W.W.L. Cheung, S.L. Connors, F. Denton, A. Diongue-Niang, D. Dodman, M. Garschagen, O. Geden, B. Hayward, C. Jones, F. Jotzo, T. Krug, R. Lasco, J.-Y. Lee, V. Masson-Delmotte, M. Meinshausen, K. Mintenbeck, A. Mokssit, F.E.L. Otto, M. Pathak, A. Pirani, E. Poloczanska, H.-O. Pörtner, A. Revi, D.C. Roberts, J. Roy, A.C. Ruane, J. Skea, P.R. Shukla, R. Slade, A. Slangen, Y. Sokona, A.A. Sörensson, M. Tignor, D. van Vuuren, Y.-M. Wei, H. Winkler, P. Zhai, and Z. Zommers:

- Moved up [110]: M.
- Moved up [105]: , Chini, L.
- Moved up [114]: F.,
- Moved up [107]: , Hurtt, G.
- Moved up [109]: N.,
- Moved up [111]: J.,
- Moved up [108]: L.,
- Moved up [119]: J.,
- Deleted: Joshi, M.
- Deleted: and Jones, G. S.: The climatic effects of the direct injection of water vapour into the stratosphere by large volcanic eruptions, *Atmos. Chem. Phys.*, 9, 6109–6118,
- Deleted: , 2009. [114]
- Deleted: P., Egorova, T., Evans, M., González-Rouco, J.
- Deleted: Goosse, H
- Deleted: C., Joos, F., Kaplan, J. O., Khodri, M., Klein Gc (... [115]
- Deleted: Lorenz, S.
- Deleted: Luterbacher, J., Man, W., Maycock, A. C., Meit (... [116]
- Deleted: Phipps, S. J., Pongratz, J., Rozanov, E., Schmid (... [117]
- Deleted: Yeo, K. L., Zanchettin, D., Zhang, Q., and Zorit (... [118]
- Deleted: , 2017.
- Deleted: Keeble, J., Hassler, B., Banerjee, A., Checa-Gar (... [119]
- Deleted: , 2021.
- Deleted: Kovilakam, M., Thomason, L. W., Ernest, N., R (... [120]
- Deleted: , 2020.
- Moved up [87]: Lan, X.,
- Moved down [121]: . D.,
- Moved up [112]: L.,
- Moved down [122]: . E.,
- Moved up [98]: Environ.,
- Deleted: Leach, N. J., Jenkins, S., Nicholls, Z., Smith, C. (... [121]
- Deleted: , 2021. [122]
- Deleted: Lim, L.
- Deleted: Lund, M. T., Millar, R. J., Owen, B., Penner, J
- Deleted: Pitari, G., Prather, M. J., Sausen, R., and Wilco (... [123]
- Deleted: 244, 117834,
- Deleted: , 2021a.
- Deleted: 2021b

4495 Synthesis Report of the IPCC Sixth Assessment Report (AR6): Summary for Policymakers. Intergovernmental Panel on
Climate Change [accepted], available at <https://www.ipcc.ch/report/ar6/syr/>. 2023.

Lenssen, N. J. L., Schmidt, G. A., Hansen, J. E., Menne, M. J., Persin, A., Ruedy, R., and Zyss, D.: Improvements in the
GISTEMP Uncertainty Model, *J. Geophys. Res.-Atmos.*, 124, 6307–6326, <https://doi.org/10.1029/2018JD029522>,
2019.

Levitus, S., Antonov, J. I., Boyer, T. P., Baranova, O. K., Garcia, H. E., Locarnini, R. A., Mishonov, A. V., Reagan, J. R.,
4500 Seidov, D., Yarosh, E. S., and Zweng, M. M.: World ocean heat content and thermosteric sea level change (0–2000 m), 1955–
2010, *Geophys. Res. Lett.*, 39, <https://doi.org/10.1029/2012GL051106>, 2012.

Loeb, N. G., Johnson, G. C., Thorsen, T. J., Lyman, J. M., Rose, F. G., Kato, S.: Satellite and ocean data reveal marked increase
in Earth’s heating rate. *Geophys. Res. Lett.*, 48, e2021GL093047, <https://doi.org/10.1029/2021GL093047>, 2021.

Lonsdale, C. R. and Sun, K.: Nitrogen oxides emissions from selected cities in North America, Europe, and East Asia observed
4505 by TROPOMI before and after the COVID-19 pandemic, *EGU sphere* [preprint], 2023, 1–30,
<https://doi.org/10.5194/egusphere-2023-346>, 2023.

van Marle, M. J. E., Kloster, S., Magi, B. I., Marlon, J. R., Daniau, A.-L., Field, R. D., Arneeth, A., Forrest, M., Hantson, S.,
Kehrwald, N. M., Knorr, W., Lasslop, G., Li, F., Mangeon, S., Yue, C., Kaiser, J. W., and van der Werf, G. R.: Historic global
biomass burning emissions for CMIP6 (BB4CMIP) based on merging satellite observations with proxies and fire models
4510 (1750–2015), *Geosci. Model Dev.*, 10, 3329–3357, <https://doi.org/10.5194/gmd-10-3329-2017>, 2017.
<https://doi.org/10.5194/gmd-10-2247-2017> McKenna, C. M., Maycock, A. C., Forster, P. M., Smith, C. J., and
Tokarska, K. B.: Stringent mitigation substantially reduces risk of unprecedented near-term warming rates, *Nature Climate
Change*, 11, 126–131, <https://doi.org/10.1038/s41558-020-00957-9>, 2021.

Meinshausen, M., Raper, S. C. B., and Wigley, T. M. L.: Emulating coupled atmosphere-ocean and carbon cycle models with
4515 a simpler model, *MAGICC6 – Part I: Model description and calibration*, *Atmos. Chem. Phys.*, 11, 1417–1456,
<https://doi.org/10.5194/acp-11-1417-2011>, 2011.
<https://doi.org/10.5194/gmd-13-3571-2020> Millán, L., Santee, M. L., Lambert, A., Livesey, N. J., Werner, F.,
Schwartz, M. J., Pumphrey, H. C., Manney, G. L., Wang, Y., Su, H., Wu, L., Read, W. G., and Froidevaux, L.: The Hunga
Tonga-Hunga Ha’apai Hydration of the Stratosphere, *Geophys. Res. Lett.*, 49, e2022GL099381,
4520 <https://doi.org/10.1029/2022GL099381>, 2022.
<https://doi.org/10.5194/acp-17-7213-2017> Minx, J. C., Lamb, W. F., Andrew, R. M., Canadell, J. G., Crippa, M.,
Döbbling, N., Forster, P. M., Guizzardi, D., Olivier, J., Peters, G. P., Pongratz, J., Reisinger, A., Rigby, M., Saunio, M.,
Smith, S. J., Solazzo, E., and Tian, H.: A comprehensive and synthetic dataset for global, regional, and national greenhouse

Deleted: .

Moved up [113]: T.,

Deleted: Matthes, K., Funke, B., Andersson, M. E., Barnard, L., Beer, J., Charbonneau, P., Chilverd, M. A., Dudok de Wit, T., Haberreiter, M., Hendry, A., Jackman, C. H., Kretzschmar, M., Kruschke, T., Kunze, M., Langematz, U., Marsh, D. R., Maycock, A. C., Miosos, S., Rodger, C. J., Scaife, A. A., Seppälä, A., Shangquan, M., Sinnhuber, M., Tourpali, K., Usoskin, I., van de Kamp, M., Verronen, P.

Deleted: and Versick, S.: Solar forcing for CMIP6 (v3.2), *Geosci. Model Dev.*, 10, 2247–2302,

Deleted: , 2017.¶

Moved up [117]: R.

Moved up [104]: ., Canadell, J.

Deleted: Meinshausen, M., Nicholls, Z.

Deleted: J., Lewis, J., Gidden, M. J., Vogel, E., Freund, M., Beyerle, U., Gessner, C., Nauels, A., Bauer, N

Deleted: G., Daniel, J. S., John, A., Krummel, P. B., Luderer, G., Meinshausen, N., Montzka, S. A., Rayner, P. J., Reimann, S., Smith, S. J., van den Berg, M., Velders, G. J. M., Vollmer, M. K., and Wang, R. H. J.: The shared socio-economic pathway (SSP) greenhouse gas concentrations and their extensions to 2500, *Geosci. Model Dev.*, 13, 3571–3605,

Deleted: , 2020.¶

Deleted: Millar, R. J., Nicholls, Z. R., Friedlingstein, P., and Allen, M. R.: A modified impulse-response representation of the global near-surface air temperature and atmospheric concentration response to carbon dioxide emissions, *Atmos. Chem. Phys.*, 17, 7213–7228,

Deleted: , 2017.¶

gas emissions by sector 1970–2018 with an extension to 2019, *Earth Syst. Sci. Data*, 13, 5213–5252, <https://doi.org/10.5194/essd-13-5213-2021>, 2021.

4555 Montzka, S: The NOAA Annual Greenhouse Gas Index (AGGI), <https://gml.noaa.gov/aggi/aggi.html>, 2022. <https://doi.org/10.1029/2011JD017187>, <https://doi.org/10.1029/2019JD032361>, 2021.

Myhre, G., Samset, B. H., Schulz, M., Balkanski, Y., Bauer, S., Bernsten, T. K., Bian, H., Bellouin, N., Chin, M., Diehl, T., Easter, R. C., Feichter, J., Ghan, S. J., Hauglustaine, D., Iversen, T., Kinne, S., Kirkevåg, A., Lamarque, J.-F., Lin, G., Liu, X., Lund, M. T., Luo, G., Ma, X., van Noije, T., Penner, J. E., Rasch, P. J., Ruiz, A., Seland, Ø., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., Wang, P., Wang, Z., Xu, L., Yu, H., Yu, F., Yoon, J.-H., Zhang, K., Zhang, H., and Zhou, C.: Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations, *Atmos. Chem. Phys.*, 13, 1853–1877, <https://doi.org/10.5194/acp-13-1853-2013>, 2013a.

Myhre, G., D. Shindell, F.-M. Bréon, W. Collins, J. Fuglestedt, J. Huang, D. Koch, J.-F. Lamarque, D. Lee, B. Mendoza, T. Nakajima, A. Robock, G. Stephens, T. Takemura and H. Zhang: Anthropogenic and Natural Radiative Forcing. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/CBO9781107415324.018>, 2013.

4560 Nisbet, E. G., Manning, M. R., Dlugokencky, E. J., Michel, S. E., Lan, X., Roeckmann, T., Gon, H. A. D. V. D., Palmer, P., Oh, Y., Fisher, R., Lowry, D., France, J. L., and White, J. W. C.: Atmospheric methane: Comparison between methane's record in 2006–2022 and during glacial terminations, *Preprints*, <https://doi.org/10.22541/essoar.167689502.25042797/v1>, 2023.

Nitzbon, J., Krinner, G., Deimling, T. S. von, Werner, M., and Langer, M.: Quantifying the Permafrost Heat Sink in Earth's Climate System, *ESS Open Archive [preprint]*, <https://doi.org/10.1002/essoar.10511600.1>, 2022a.

Nitzbon, J., Krinner, G., Langer, M.: GCOS EHI 1960–2020 Permafrost Heat Content, World Data Center for Climate (WDCC) at DKRZ, https://doi.org/10.26050/WDCC/GCOS_EHI_1960-2020_PHC, 2022b.

4575 O'Rourke, Patrick R, Smith, Steven J, Mott, Andrea, Ahsan, Hamza, McDuffie, Erin E, Crippa, Monica, Klimont, Zbigniew, McDonald, Brian, Wang, Shuxiao, Nicholson, Matthew B, Feng, Leyang, & Hoesly, Rachel M.: CEDS v_2021_04_21 Release Emission Data (v_2021_02_05), Zenodo [data set], <https://doi.org/10.5281/zenodo.4741285>, 2021. [5, 917–920. https://doi.org/10.1038/nclimate2716](https://doi.org/10.1038/nclimate2716), 2015.

4580 Palmer, M. D. and McNeill, D. J.: Internal variability of Earth's energy budget simulated by CMIP5 climate models, *Environ. Res. Lett.*, 9, 034016, <https://doi.org/10.1088/1748-9326/9/3/034016>, 2014.

Deleted: Morice, C. P., Kennedy, J. J., Rayner, N. A., and Jones, P. D.: Quantifying uncertainties in global and regional temperature change using an ensemble of observational estimates: The HadCRUT4 data set: THE HADCRUT4 DATASET, *J. Geophys. Res.*, 117, n/a-n/a.

Deleted: , 2012.
Morice, C. P., Kennedy, J. J., Rayner, N. A., Winn, J. P., Hogan, E., Killick, R. E., Dunn, R. J. H., Osborn, T. J., Jones, P. D., and Simpson, I. R.: An Updated Assessment of Near-Surface Temperature Change From 1850: The HadCRUT5 Data Set, *J. Geophys. Res.-Atmos.*, 126, e2019JD032361,

Deleted: 2013b

Moved (insertion) [122]

Moved (insertion) [121]

Moved down [123]: L.,

Moved up [115]: Chang.,

Deleted: Otto, F. E.

Deleted: Frame, D. J., Otto, A., and Allen, M. R.: Embracing uncertainty in climate change policy, *Nature Clim.*

Palmer, M. D., Domingues, C. M., Slangen, A. B. A., and Dias, F. B.: An ensemble approach to quantify global mean sea-level rise over the 20th century from tide gauge reconstructions, *Environ. Res. Lett.*, 16, 044043, <https://doi.org/10.1088/1748-9326/abdaec>, 2021.

[Peng, S., Lin, X., Thompson, R. L., Xi, Y., Liu, G., Hauglustaine, D., Lan, X., Poulter, B., Ramonet, M., Saunio, M., Yin, Y., Zhang, Z., Zheng, B., and Ciais, P.: Wetland emission and atmospheric sink changes explain methane growth in 2020, *Nature*, 612, 477–482, . 2022.](#)

Pirani, A., Alegria, A., Khourdajie, A. A., Gunawan, W., Gutiérrez, J. M., Holsman, K., Huard, D., Juckes, M., Kawamiya, M., Klutse, N., Krey, V., Matthews, R., Milward, A., Pascoe, C., Van Der Shrier, G., Spinuso, A., Stockhause, M., and Xiaoshi Xing: The implementation of FAIR data principles in the IPCC AR6 assessment process, <https://doi.org/10.5281/ZENODO.6504469>, 2022.

[Pongratz, J., Schwingshackl, C., Bultan, S., Obermeier, W., Havermann, F., and Guo, S.: Land Use Effects on Climate: Current State, Recent Progress, and Emerging Topics, *Curr Clim Change Rep.* 7, 99–120, . 2021.](#)

Purkey, S.G. and Johnson, G.C., Warming of Global Abyssal and Deep Southern Ocean Waters between the 1990s and 2000s: Contributions to Global Heat and Sea Level Rise Budgets, *J. Climate*, 23, 23, 6336–6351, <https://doi.org/10.1175/2010JCLI3682.1>, 2010.

Putaud, J.-P., Pisoni, E., Mangold, A., Hueglin, C., Sciare, J., Pikridas, M., Savvides, C., Ondracek, J., Mbengue, S., Wiedensohler, A., Weinhold, K., Merkel, M., Poulain, L., van Pinxteren, D., Herrmann, H., Massling, A., Nordstroem, C., Alastuey, A., Reche, C., Pérez, N., Castillo, S., Sorribas, M., Adame, J. A., Petaja, T., Lehtipalo, K., Niemi, J., Riffault, V., de Brito, J. F., Colette, A., Favez, O., Petit, J.-E., Gros, V., Gini, M. I., Vratolis, S., Eleftheriadis, K., Diapouli, E., Denier van der Gon, H., Yttri, K. E., and Aas, W.: Impact of 2020 COVID-19 lockdowns on particulate air pollution across Europe, *EGUsphere* [preprint], . 2023.

[Qasmi, S. and Ribes, A.: Reducing uncertainty in local temperature projections, *Sci. Adv.*, 8, eabo6872, https://doi.org/10.1126/sciadv.abo6872, 2022.](#)

Quaas, J., Jia, H., Smith, C., Albright, A. L., Aas, W., Bellouin, N., Boucher, O., Doutriaux-Boucher, M., Forster, P. M., Grosvenor, D., Jenkins, S., Klimont, Z., Loeb, N. G., Ma, X., Naik, V., Paulot, F., Stier, P., Wild, M., Myhre, G., and Schulz, M.: Robust evidence for reversal of the trend in aerosol effective climate forcing, *Atmos. Chem. Phys.*, 22, 12221–12239, <https://doi.org/10.5194/acp-22-12221-2022>, 2022.

Raghuraman, S.P., Paynter, D. and Ramaswamy, V.: Anthropogenic forcing and response yield observed positive trend in Earth's energy imbalance, *Nat. Commun.* 12, 4577, <https://doi.org/10.1038/s41467-021-24544-4>, 2021.

[Randerson, J. T., van der Werf, G. R., Giglio, L., Collatz, G. J., and Kasibhatla, P. S.: Global Fire Emissions Database, Version 4.1 \(GFEDv4\), ORNL Distributed Active Archive Center \[dataset\], . 2017](#)

Deleted: <https://doi.org/10.5194/egusphere->

Deleted: -434, 2023

Moved (insertion) [123]

Riahi, K., Schaeffer, J. Arango, K. Calvin, C. Guivarch, T. Hasegawa, K. Jiang, E. Kriegler, R. Matthews, G.P. Peters, A. Rao, S. Robertson, A.M. Sebbit, J. Steinberger, M. Tavoni, D.P. van Vuuren, 2022: Mitigation pathways compatible with long-term goals. In IPCC, 2022: Climate Change 2022: Mitigation of Climate Change. Contribution of Working Group III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [P.R. Shukla, J. Skea, R. Slade, A. Al Khourdajie, R. van Diemen, D. McCollum, M. Pathak, S. Some, P. Vyas, R. Fradera, M. Belkacemi, A. Hasija, G. Lisboa, S. Luz, J. Malley, (eds.)]. Cambridge University Press, Cambridge, UK and New York, NY, USA, <https://doi.org/10.1017/9781009157926.005>, 2022.

Ribes, A., Qasmi, S., and Gillett, N. P.: Making climate projections conditional on historical observations, *Sci. Adv.*, 7, eabc0671, <https://doi.org/10.1126/sciadv.abc0671>, 2021.

Deleted: Ribes, A., Planton, S., and Terray, L.: Application of regularised optimal fingerprinting to attribution. Part I: method, properties and idealised analysis, *Clim. Dyn.*, 41, 2817–2836, <https://doi.org/10.1007/s00382-013-1735-7>, 2013.

Richardson, M., Cowtan, K., and Millar, R. J.: Global temperature definition affects achievement of long-term climate goals, *Environ. Res. Lett.*, 13, 054004, <https://doi.org/10.1088/1748-9326/aab305>, 2018.

Rogelj, J., D. Shindell, K. Jiang, S. Fifita, P. Forster, V. Ginzburg, C. Handa, H. Khesghi, S. Kobayashi, E. Kriegler, L. Mundaca, R. Séférian, and M. V. Vilariño: Mitigation Pathways Compatible with 1.5°C in the Context of Sustainable Development. In: *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty* [Masson-Delmotte, V., P. Zhai, H.-O. Pörtner, D. Roberts, J. Skea, P.R. Shukla, A. Pirani, W. Moufouma-Okia, C. Péan, R. Pidcock, S. Connors, J. B. R. Matthews, Y. Chen, X. Zhou, M. I. Gomis, E. Lonnoy, T. Maycock, M. Tignor, and T. Waterfield (eds.)]. Cambridge University Press, Cambridge, UK and New York, NY, USA, pp. 93-174, <https://doi.org/10.1017/9781009157940.004>, 2018.

Rogelj, J., Forster, P. M., Kriegler, E., Smith, C. J., and Séférian, R.: Estimating and tracking the remaining carbon budget for stringent climate targets, *Nature*, 571, 335–342, <https://doi.org/10.1038/s41586-019-1368-z>, 2019.

Rogelj, J., Rao, S., McCollum, D. L., Pachauri, S., Klimont, Z., Krey, V., and Riahi, K.: Air-pollution emission ranges consistent with the representative concentration pathways, *Nature Clim. Chang.*, 4 (6), 446–450, , 2014.

Rohde, R., Muller, R., Jacobsen, R., Perlmutter, S., Rosenfeld, A., et al.: Berkeley Earth Temperature Averaging Process Geoinfor. Geostat., An Overview 1:2., , 2013. <https://berkeleyearth.org/wp-content/uploads/2022/12/Methods-GIGS-1-103.pdf>

Deleted: ,

Deleted: ,

Deleted: ,

Deleted: ,

Deleted: ,

Deleted: . (2013)

Deleted: ,

Deleted: ,

Deleted: available at:

Rohde, R. A. and Hausfather, Z.: The Berkeley Earth Land/Ocean Temperature Record, *Earth Sys. Sci. Data*, 12, 3469–3479, <https://doi.org/10.5194/essd-12-3469-2020>, 2020.

Schoenenberger, F., Vollmer, M.K., Rigby, M., Hill, M., Fraser, P.J., Krummel, P.B., Langenfelds, R.L., Rhee, T.S., Peter, T., Reimann, S.: First observations, trends, and emissions of HCFC-31 (CH2ClF) in the global atmosphere, *Geophys. Res. Lett.*, 42, 7817–7824, <https://doi.org/10.1002/2015GL064709>, 2015.

von Schuckmann, K., Cheng, L., Palmer, M. D., Hansen, J., Tassone, C., Aich, V., Adusumilli, S., Beltrami, H., Boyer, T., Cuesta-Valero, F. J., Desbruyères, D., Domingues, C., García-García, A., Gentine, P., Gilson, J., Gorfer, M., Haimberger, L., Ishii, M., Johnson, G. C., Killick, R., King, B. A., Kirchengast, G., Kolodziejczyk, N., Lyman, J., Marzeion, B., Mayer, M., Monier, M., Monselesan, D. P., Purkey, S., Roemmich, D., Schweiger, A., Seneviratne, S. I., Shepherd, A., Slater, D. A.,
1680 Steiner, A. K., Straneo, F., Timmermans, M.-L., and Wijffels, S. E.: Heat stored in the Earth system: where does the energy go?, *Earth Syst. Sci. Data*, 12, 2013–2041, <https://doi.org/10.5194/essd-12-2013-2020>, 2020.

von Schuckmann, K., Minière, A., Gues, F., Cuesta-Valero, F. J., Kirchengast, G., Adusumilli, S., Straneo, F., Ablain, M., Allan, R. P., Barker, P. M., Beltrami, H., Blazquez, A., Boyer, T., Cheng, L., Church, J., Desbruyeres, D., Dolman, H., Domingues, C. M., García-García, A., Giglio, D., Gilson, J. E., Gorfer, M., Haimberger, L., Hakuba, M. Z., Hendricks, S.,
1685 Hosoda, S., Johnson, G. C., Killick, R., King, B., Kolodziejczyk, N., Korosov, A., Krinner, G., Kuusela, M., Landerer, F. W., Langer, M., Lavergne, T., Lawrence, I., Li, Y., Lyman, J., Marti, F., Marzeion, B., Mayer, M., MacDougall, A. H., McDougall, T., Monselesan, D. P., Nitzbon, J., Otsuka, I., Peng, J., Purkey, S., Roemmich, D., Sato, K., Sato, K., Savita, A., Schweiger, A., Shepherd, A., Seneviratne, S. I., Simons, L., Slater, D. A., Slater, T., Steiner, A. K., Suga, T., Szekely, T., Thiery, W., Timmermans, M.-L., Vanderkelen, I., Wijffels, S. E., Wu, T., and Zemp, M.: Heat stored in the Earth system 1960–2020:
1690 where does the energy go?, *Earth System Science Data*, 15, 1675–1709, <https://doi.org/10.5194/essd-15-1675-2023>, 2023a.

von Schuckmann, K., Minière, A., Gues, F., Cuesta-Valero, F. J., Kirchengast, G., Adusumilli, S., Straneo, F., Ablain, M., Allan, R. P., Barker, P. M., Beltrami, H., Blazquez, A., Boyer, T., Cheng, L., Church, J., Desbruyeres, D., Dolman, H., Domingues, C. M., García-García, A., Giglio, D., Gilson, J. E., Gorfer, M., Haimberger, L., Hakuba, M. Z., Hendricks, S.,
1695 Hosoda, S., Johnson, G. C., Killick, R., King, B., Kolodziejczyk, N., Korosov, A., Krinner, G., Kuusela, M., Landerer, F. W., Langer, M., Lavergne, T., Lawrence, I., Li, Y., Lyman, J., Marti, F., Marzeion, B., Mayer, M., MacDougall, A. H., McDougall, T., Monselesan, D. P., Nitzbon, J., Otsuka, I., Peng, J., Purkey, S., Roemmich, D., Sato, K., Sato, K., Savita, A., Schweiger, A., Shepherd, A., Seneviratne, S. I., Simons, L., Slater, D. A., Slater, T., Steiner, A. K., Suga, T., Szekely, T., Thiery, W., Timmermans, M.-L., Vanderkelen, I., Wijffels, S. E., Wu, T., and Zemp, M.: GCOS EHI 1960–2020 Earth Heat Inventory
1700 Ocean Heat Content (Version 2), https://doi.org/10.26050/WDCC/GCOS_EHI_1960-2020_OHC_v2, 2023b.

Sellitto, P., Podglajen, A., Belhadji, R., Boichu, M., Carboni, E., Cuesta, J., Duchamp, C., Kloss, C., Siddans, R., Bègue, N., Blarel, L., Jegou, F., Khaykin, S., Renard, J.-B., and Legras, B.: The unexpected radiative impact of the Hunga Tonga eruption of 15th January 2022, *Commun. Earth Environ.*, 3, 288, <https://doi.org/10.1038/s43247-022-00618-z>, 2022.

Seneviratne, S.I., X. Zhang, M. Adnan, W. Badi, C. Dereczynski, A. Di Luca, S. Ghosh, I. Iskandar, J. Kossin, S. Lewis, F.
1705 Otto, I. Pinto, M. Satoh, S. M. Vicente-Serrano, M. Wehner, and B. Zhou: Weather and Climate Extreme Events in a Changing Climate. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment*

Deleted: .

Deleted: .

Deleted: .

Deleted: .

- Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1513–1766, doi:10.1017/9781009157896.013.1513–1766, 2021.
- 4715 <https://doi.org/10.1017/9781009157896.013>, 2021.
- Sigl, M., Toohey, M., McConnell, J. R., Cole-Dai, J., and Severi, M.: Volcanic stratospheric sulfur injections and aerosol optical depth during the Holocene (past 11,500 years) from a bipolar ice-core array, *Earth Syst. Sci. Data*, 14, 3167–3196, <https://doi.org/10.5194/essd-14-3167-2022>, 2022.
- Simmonds, P. G., Rigby, M., McCulloch, A., O'Doherty, S., Young, D., Mühle, J., Krummel, P. B., Steele, P., Fraser, P. J., Manning, A. J., Weiss, R. F., Salameh, P. K., Harth, C. M., Wang, R. H. J., and Prinn, R. G.: Changing trends and emissions of hydrochlorofluorocarbons (HCFCs) and their hydrofluorocarbon (HFCs) replacements, *Atmos. Chem. Phys.*, 17, 4641–4655, <https://doi.org/10.5194/acp-17-4641-2017>, 2017.
- 4720 Sippel, S., Zscheischler, J., Heimann, M., Otto, F. E. L., Peters, J., and Mahecha, M. D.: Quantifying changes in climate variability and extremes: Pitfalls and their overcoming, *Geophys. Res. Lett.*, 42, 9990–9998, <https://doi.org/10.1002/2015GL066307>, 2015.
- 4725 <https://doi.org/10.1038/s41612-020-00131-0>, 2020.
- [Skeie, R. B., Myhre, G., Hodnebrog, Ø., Cameron-Smith, P. J., Deushi, M., Hegglin, M. I., Horowitz, L. W., Kramer, R. J., Michou, M., Mills, M. J., Olivié, D. J. L., Connor, F. M. O., Paynter, D., Samset, B. H., Sellar, A., Shindell, D., Takemura, T., Tilmes, S., and Wu, T.: Historical total ozone radiative forcing derived from CMIP6 simulations, *npj Clim. Atmos. Sci.*, 3, 32, <https://doi.org/10.1038/s41612-020-00131-0>, 2020.](https://doi.org/10.1038/s41612-020-00131-0)
- 4730 [Smith, C. J., Kramer, R. J., Myhre, G., Forster, P. M., Soden, B. J., Andrews, T., Boucher, O., Faluvegi, G., Fläschner, D., Hodnebrog, Ø., Kasoar, M., Kharin, V., Kirkevåg, A., Lamarque, J.-F., Mülmenstädt, J., Olivié, D., Richardson, T., Samset, B. H., Shindell, D., Stier, P., Takemura, T., Voulgarakis, A., and Watson-Parris, D.: Understanding Rapid Adjustments to Diverse Forcing Agents, *Geophys. Res. Lett.*, 45, 12,023–12,031, <https://doi.org/10.1029/2018GL079826>, 2018a.](https://doi.org/10.1029/2018GL079826)
- 4735 [Smith, C. J., Forster, P. M., Allen, M., Leach, N., Millar, R. J., Passerello, G. A., and Regayre, L. A.: FAIR v1.3: A simple emissions-based impulse response and carbon cycle model, *Geoscientific Model Development*, 11, 2273–2297, <https://doi.org/10.5194/gmd-11-2273-2018>, 2018b.](https://doi.org/10.5194/gmd-11-2273-2018)
- [Smith, C. J., Harris, G. R., Palmer, M. D., Bellouin, N., Collins, W., Myhre, G., Schulz, M., Golaz, J.-C., Ringer, M., Storelvmo, T., and Forster, P. M.: Energy Budget Constraints on the Time History of Aerosol Forcing and Climate Sensitivity, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033622, <https://doi.org/10.1029/2020JD033622>, 2021a.](https://doi.org/10.1029/2020JD033622)
- 4740 Smith, C., Nicholls, Z. R. J., Armour, K., Collins, W., Forster, P., Meinshausen, M., Palmer, M. D., and Watanabe, M.: The Earth's Energy Budget, Climate Feedbacks, and Climate Sensitivity Supplementary Material, in: *Climate Change 2021: The*

- Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., 2021.
- 4750 ~~Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)~~, available from <https://www.ipcc.ch/>, 2021c.
- Smith, C., Walsh, T., Forster, P.M., Gillett, N., Hauser, M., Lamb, W., Lamboll, R., Palmer, M., Ribes, A., Schumacher, D., Seneviratne, S., Trewin, B., and von Schuckmann, K.: Indicators of Global Climate Change 2022 (v2023.05.24), Zenodo. <https://doi.org/10.5281/zenodo.7883758>, 2023.
- Smith, S. J., van Aardenne, J., Klimont, Z., Andres, R. J., Volke, A., and Delgado Arias, S.: Anthropogenic sulfur dioxide emissions: 1850–2005, *Atmos. Chem. and Phys.*, 11, 1101–1116, <https://doi.org/10.5194/acp-11-1101-2011>, 2011.
- Sokhi, R. S., Singh, V., Querol, X., Finardi, S., Targino, A. C., Andrade, M. de F., Pavlovic, R., Garland, R. M., Massagué, J., Kong, S., Baklanov, A., Ren, L., Tarasova, O., Carmichael, G., Peuch, V.-H., Anand, V., Arbilla, G., Badali, K., Beig, G., Belalcazar, L. C., Bolignano, A., Brimblecombe, P., Camacho, P., Casallas, A., Charland, J.-P., Choi, J., Chourdakis, E., Coll, I., Collins, M., Cyrus, J., Silva, C. M. da, Giosa, A. D. D., Leo, A. D., Ferro, C., Gavidia-Calderon, M., Gayen, A., Ginzburg, A., Godefroy, F., Gonzalez, Y. A., Guevara-Luna, M., Haque, S. M., Havenga, H., Herod, D., Hörrak, U., Hussein, T., Ibarra, S., Jaimes, M., Kaasik, M., Khaiwal, R., Kim, J., Kousa, A., Kukkonen, J., Kulmala, M., Kuula, J., Violette, N. L., Lanzani, G., Liu, X., MacDougall, S., Manseau, P. M., Marchegiani, G., McDonald, B., Mishra, S. V., Molina, L. T., Mooibroek, D., Mor, S., Moussiopoulos, N., Murena, F., Niemi, J. V., Noe, S., Nogueira, T., Norman, M., Pérez-Camaño, J. L., Petäjä, T., Piketh, S., Rathod, A., Reid, K., Retama, A., Rivera, O., Rojas, N. Y., Rojas-Quincho, J. P., José, R. S., Sánchez, O., Seguel, R. J., Sillanpää, S., Su, Y., Tapper, N., Terrazas, A., Timonen, H., Toscano, D., Tsegas, G., Velders, G. J. M., Vlachokostas, C., Schneidmesser, E. von, VPM, R., Yadav, R., Zalakeviciute, R., and Zavala, M.: A global observational analysis to understand changes in air quality during exceptionally low anthropogenic emission conditions, *Environment International*, 157, 106818, <https://doi.org/10.1016/j.envint.2021.106818>, 2021.
- Steiner, A. K., Ladstädter, F., Randel, W. J., Maycock, A. C., Fu, Q., Claud, C., Gleisner, H., Haimberger, L., Ho, S.-P., Keckhut, P., Leblanc, T., Mears, C., Polvani, L. M., Santer, B. D., Schmidt, T., Sofieva, V., Wing, R., and Zou, C.-Z.: Observed Temperature Changes in the Troposphere and Stratosphere from 1979 to 2018, *J. Climate*, 33, 8165–8194, <https://doi.org/10.1175/JCLI-D-19-0998.1>, 2020.
- 4770 ~~Stevenson, D. S., Young, P. J., Naik, V., Lamarque, J.-F., Shindell, D. T., Voulgarakis, A., Skeie, R. B., Dalsoren, S. B., Myhre, G., Bernsten, T. K., Folberth, G. A., Rumbold, S. T., Collins, W. J., MacKenzie, I. A., Doherty, R. M., Zeng, G., van Noije, T. P. C., Strunk, A., Bergmann, D., Cameron-Smith, P., Plummer, D. A., Strode, S. A., Horowitz, L., Lee, Y. H., Szopa,~~

Deleted: 2021b

Moved up [103]: . In Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I.

Deleted: Smith, C., Z. R. J. Nicholls, K. Armour, W. Collins, P. Forster, M. Meinshausen, M. D. Palmer, and M. Watanabe: The Earth's Energy Budget, Climate Feedbacks, and Climate Sensitivity Supplementary Material

Deleted: (01).

785 [S., Sudo, K., Nagashima, T., Josse, B., Cionni, I., Righi, M., Eyring, V., Conley, A., Bowman, K. W., Wild, O., and Archibald, A.: Tropospheric ozone changes, radiative forcing and attribution to emissions in the Atmospheric Chemistry and Climate Model Intercomparison Project \(ACCMIP\), *Atmos. Chem. Phys.*, 13, 3063–3085, <https://doi.org/10.5194/acp-13-3063-2013>, 2013.](#)

790 [Sun, W., Li, Q., Huang, B., Cheng, J., Song, Z., Li, H., Dong, W., Zhai, P., and Jones, P.: The Assessment of Global Surface Temperature Change from 1850s: The C-LSAT2.0 Ensemble and the CMST-Interim Datasets, *Advances in Atmospheric Sciences*, 38, 875–888, <https://doi.org/10.1007/s00376-021-1012-3>, 2021.](#)

795 Szopa, S., V. Naik, B. Adhikary, P. Artaxo, T. Bernsten, W.D. Collins, S. Fuzzi, L. Gallardo, A. Kiendler-Scharr, Z. Klimont, H. Liao, N. Unger, and P. Zanis: Short-Lived Climate Forcers. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 817–922, <https://doi.org/10.1017/9781009157896.008>, 2021.

800 [Taha, G., Loughman, R., Zhu, T., Thomason, L., Kar, J., Rieger, L., and Bourassa, A.: OMPS LP Version 2.0 multi-wavelength aerosol extinction coefficient retrieval algorithm, *Atmos. Meas. Tech.*, 14, 1015–1036, <https://doi.org/10.5194/amt-14-1015-2021>, 2021.](#)

805 [Taylor, K. E., Crucifix, M., Braconnot, P., Hewitt, C. D., Doutriaux, C., Broccoli, A. J., Mitchell, J. F. B., and Webb, M. J.: Estimating Shortwave Radiative Forcing and Response in Climate Models, *J. Climate*, 20, 2530–2543, <https://doi.org/10.1175/JCLI4143.1>, 2007.](#)

[Thomason, L. W., Ernest, N., Millán, L., Rieger, L., Bourassa, A., Vernier, J.-P., Manney, G., Luo, B., Arfeuille, F., and Peter, T.: A global space-based stratospheric aerosol \hack\breakclimatology: 1979–2016, *Earth System Science Data*, 10, 469–492, <https://doi.org/10.5194/essd-10-469-2018>, 2018.](#)

810 [Thornhill, G. D., Collins, W. J., Kramer, R. J., Olivíe, D., Skeie, R. B., O'Connor, F. M., Abraham, N. L., Checa-Garcia, R., Bauer, S. E., Deushi, M., Emmons, L. K., Forster, P. M., Horowitz, L. W., Johnson, B., Keeble, J., Lamarque, J.-F., Michou, M., Mills, M. J., Mulcahy, J. P., Myhre, G., Nabat, P., Naik, V., Oshima, N., Schulz, M., Smith, C. J., Takemura, T., Tilmes, S., Wu, T., Zeng, G., and Zhang, J.: Effective radiative forcing from emissions of reactive gases and aerosols – a multi-model comparison, *Atmos. Chem. Phys.*, 21, 853–874, <https://doi.org/10.5194/acp-21-853-2021>, 2021a.](#)

[Thornhill, G., Collins, W., Olivíe, D., Skeie, R. B., Archibald, A., Bauer, S., Checa-Garcia, R., Fiedler, S., Folberth, G., Giermundsen, A., Horowitz, L., Lamarque, J.-F., Michou, M., Mulcahy, J., Nabat, P., Naik, V., O'Connor, F. M., Paulot, F., Schulz, M., Scott, C. E., Séférian, R., Smith, C., Takemura, T., Tilmes, S., Tsigaridis, K., and Weber, J.: Climate-driven](#)

815 [chemistry and aerosol feedbacks in CMIP6 Earth system models, Atmos. Chem. Phys., 21, 1105–1126,
<https://doi.org/10.5194/acp-21-1105-2021>, 2021b.](#)
[Toohey, M. and Sigl, M.: Volcanic stratospheric sulfur injections and aerosol optical depth from 500\BCE to 1900\CE, Earth
Syst. Sci. Data, 9, 809–831, <https://doi.org/10.5194/essd-9-809-2017>, 2017.](#)

Trewin, B.: Assessing Internal Variability of Global Mean Surface Temperature From Observational Data and Implications
1820 for Reaching Key Thresholds, Journal of Geophysical Research: Atmospheres, 127, e2022JD036747,
<https://doi.org/10.1029/2022JD036747>, 2022.

[Trewin, B., Cazenave, A., Howell, S., Huss, M., Isensee, K., Palmer, M. D., Tarasova, O., and Vermeulen, A.: Headline
Indicators for Global Climate Monitoring, Bulletin of the American Meteorological Society, 102, E20–E37,
<https://doi.org/10.1175/BAMS-D-19-0196.1>, 2021.](#)

825 [UNFCCC: Report of the Conference of the Parties on its twenty-first session, held in Paris from 30 November to 13 December
2015, Addendum, Part two: Action taken by the Conference of the Parties at its, twenty-first session, Decisions adopted by the
Conference of the Parties, <https://unfccc.int/resource/docs/2015/cop21/eng/10a01.pdf>, 2015.](#)
[UNFCCC: Report of the Conference of the Parties serving as the meeting of the Parties to the Paris Agreement on its third
session, held in Glasgow from 31 October to 13 November 2021. Addendum Part two: Action taken by the Conference of the
830 Parties serving as the meeting of the Parties to the Paris Agreement at its third session,
\[https://unfccc.int/sites/default/files/resource/CMA2021_10_Add3_E.pdf\]\(https://unfccc.int/sites/default/files/resource/CMA2021_10_Add3_E.pdf\), 2022a.](#)
[UNFCCC: Decision -/CP.27, Sharm-El-Sheikh Implementation Plan,
\[https://unfccc.int/sites/default/files/resource/cop27_auv_2_cover%20decision.pdf\]\(https://unfccc.int/sites/default/files/resource/cop27_auv_2_cover%20decision.pdf\) 2022b.](#)

Van Der Werf, G. R., Randerson, J. T., Giglio, L., Van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., Van Marle, M. J.
1835 E., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997–2016,
Earth Syst. Sci. Data, 9, 697–720, <https://doi.org/10.5194/essd-9-697-2017>, 2017.

Vanderkelen, I., van Lipzig, N. P. M., Lawrence, D. M., Droppers, B., Golub, M., Gosling, S. N., Janssen, A. B. G., Marcé,
R., Schmied, H. M., Perroud, M., Pierson, D., Pokhrel, Y., Satoh, Y., Schewe, J., Seneviratne, S. I., Stepanenko, V. M., Tan,
Z., Woolway, R. I., and Thiery, W.: Global Heat Uptake by Inland Waters, Geophysical Research Letters, 47, e2020GL087867,
1840 <https://doi.org/10.1029/2020GL087867>, 2020.

Vanderkelen, I. and Thiery, W.: GCOS EHI 1960-2020 Inland Water Heat Content,
https://doi.org/10.26050/WDCC/GCOS_EHI_1960-2020_IWHC, 2022.

Vollmer, M. K., Young, D., Trudinger, C. M., Mühle, J., Henne, S., Rigby, M., Park, S., Li, S., Guillevic, M., Mitrevski, B.,
Harth, C. M., Miller, B. R., Reimann, S., Yao, B., Steele, L. P., Wyss, S. A., Lunder, C. R., Arduini, J., McCulloch, A., Wu,
1845 S., Rhee, T. S., Wang, R. H. J., Salameh, P. K., Hermansen, O., Hill, M., Langenfelds, R. L., Ivy, D., O'Doherty, S., Krummel,

P. B., Maione, M., Etheridge, D. M., Zhou, L., Fraser, P. J., Prinn, R. G., Weiss, R. F., and Simmonds, P. G.: Atmospheric histories and emissions of chlorofluorocarbons CFC-13 (CClF₃), ΣCFC-114 (C₂Cl₂F₄), and CFC-115 (C₂ClF₅), Atmos. Chem. Phys., 18, 979–1002, <https://doi.org/10.5194/acp-18-979-2018>, 2018.

Vose, R. S., Huang, B., Yin, X., Arndt, D., Easterling, D. R., Lawrimore, J. H., Menne, M. J., Sanchez-Lugo, A., and Zhang, H. M.: Implementing Full Spatial Coverage in NOAA's Global Temperature Analysis, Geophys. Res. Lett., 48, e2020GL090873, <https://doi.org/10.1029/2020GL090873>, 2021.

Western, L. M., Vollmer, M. K., Krummel, P. B., Adcock, K. E., Fraser, P. J., Harth, C. M., Langenfelds, R. L., Montzka, S. A., Mühle, J., O'Doherty, S., Oram, D. E., Reimann, S., Rigby, M., Vimont, I., Weiss, R. F., Young, D., and Laube, J. C.: Global increase of ozone-depleting chlorofluorocarbons from 2010 to 2020, Nat. Geosci., 16, 309–313, <https://doi.org/10.1038/s41561-023-01147-w>, 2023.

Wild, M., Gilgen, H., Roesch, A., Ohmura, A., Long, C. N., Dutton, E. G., Forgan, B., Kallis, A., Russak, V., and Tsvetkov, A.: From Dimming to Brightening: Decadal Changes in Solar Radiation at Earth's Surface, Science, 308, 847–850, <https://doi.org/10.1126/science.1103215>, 2005.

https://library.wmo.int/doc_num.php?explnum_id=11593,

<https://doi.org/10.1002/2014JD021710><https://doi.org/10.5194/egusphere-2023-689>Zhang, Z., Poulter, B., Feldman, A.F., Ying, Q., Ciais, P., Peng, S. and Xin, L.: Recent intensification of wetland methane feedback, Nat. Clim. Chang. 13, 430–433, . 2023.

Deleted: WMO: State of the Global Climate 2022, World Meteorological Organisation,

Deleted: 2023. [Zelinka, M. D., Andrews, T., Forster, P. M., and Taylor, K. E.: Quantifying components of aerosol-cloud-radiation interactions in climate models, J. Geophys. Res.-Atmos., 119, 7599–7615,](#)

Deleted: 2014. [Zelinka, M. D., Smith, C. J., Qin, Y., and Taylor, K. E.: Aerosol Effective Radiative Forcings in CMIP Models, EGU sphere \[preprint\],](#)

Deleted: , 2023.

Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00



Page 1: [1] Deleted Piers Forster 25/05/2023 07:42:00

Page 11: [3] Deleted Piers Forster 25/05/2023 07:42:00



Page 11: [4] Deleted Piers Forster 25/05/2023 07:42:00



Page 11: [4] Deleted Piers Forster 25/05/2023 07:42:00



Page 11: [5] Deleted Piers Forster 25/05/2023 07:42:00



Page 11: [5] Deleted Piers Forster 25/05/2023 07:42:00



Page 11: [6] Deleted Piers Forster 25/05/2023 07:42:00



Page 11: [6] Deleted Piers Forster 25/05/2023 07:42:00



Page 11: [7] Deleted Piers Forster 25/05/2023 07:42:00



Page 11: [7] Deleted Piers Forster 25/05/2023 07:42:00



Page 11: [8] Deleted Piers Forster 25/05/2023 07:42:00



Page 11: [8] Deleted Piers Forster 25/05/2023 07:42:00



Page 11: [9] Deleted Piers Forster 25/05/2023 07:42:00



Page 11: [9] Deleted Piers Forster 25/05/2023 07:42:00



Page 11: [10] Deleted Piers Forster 25/05/2023 07:42:00



Page 11: [10] Deleted Piers Forster 25/05/2023 07:42:00



Page 11: [11] Deleted Piers Forster 25/05/2023 07:42:00

Page 11: [12] Deleted Piers Forster 25/05/2023 07:42:00

Page 11: [13] Deleted Piers Forster 25/05/2023 07:42:00

Page 11: [13] Deleted Piers Forster 25/05/2023 07:42:00

Page 19: [14] Deleted Piers Forster 25/05/2023 07:42:00

Page 19: [15] Deleted Piers Forster 25/05/2023 07:42:00

Page 20: [16] Deleted Piers Forster 25/05/2023 07:42:00

Page 20: [17] Merged Cells Piers Forster 25/05/2023 07:42:00

Merged Cells

Page 20: [18] Deleted Piers Forster 25/05/2023 07:42:00

Page 20: [19] Merged Cells Piers Forster 25/05/2023 07:42:00

Merged Cells

Page 20: [20] Deleted Piers Forster 25/05/2023 07:42:00

Page 20: [21] Deleted Piers Forster 25/05/2023 07:42:00

Page 20: [22] Deleted Piers Forster 25/05/2023 07:42:00

Page 20: [23] Deleted Piers Forster 25/05/2023 07:42:00

Page 20: [24] Deleted Piers Forster 25/05/2023 07:42:00

Page 20: [25] Deleted Piers Forster 25/05/2023 07:42:00

Page 22: [26] Deleted Piers Forster 25/05/2023 07:42:00

Page 23: [27] Deleted Piers Forster 25/05/2023 07:42:00

Page 23: [28] Deleted Piers Forster 25/05/2023 07:42:00

Page 26: [29] Deleted Piers Forster 25/05/2023 07:42:00

Page 26: [29] Deleted Piers Forster 25/05/2023 07:42:00

▼.....
Page 26: [31] Deleted Piers Forster 25/05/2023 07:42:00

▼.....
Page 26: [31] Deleted Piers Forster 25/05/2023 07:42:00

▼.....
Page 26: [32] Deleted Piers Forster 25/05/2023 07:42:00

▼.....
Page 26: [32] Deleted Piers Forster 25/05/2023 07:42:00

▼.....
Page 26: [32] Deleted Piers Forster 25/05/2023 07:42:00

▼.....
Page 26: [33] Deleted Piers Forster 25/05/2023 07:42:00

▼.....
Page 26: [33] Deleted Piers Forster 25/05/2023 07:42:00

▼.....
Page 26: [33] Deleted Piers Forster 25/05/2023 07:42:00

▼.....
Page 26: [34] Deleted Piers Forster 25/05/2023 07:42:00

▼.....
Page 26: [34] Deleted Piers Forster 25/05/2023 07:42:00

▼.....
Page 26: [35] Deleted Piers Forster 25/05/2023 07:42:00

▼.....
Page 26: [35] Deleted Piers Forster 25/05/2023 07:42:00

▼.....
Page 26: [36] Deleted Piers Forster 25/05/2023 07:42:00

▼.....
Page 26: [37] Deleted Piers Forster 25/05/2023 07:42:00

▼.....
Page 26: [37] Deleted Piers Forster 25/05/2023 07:42:00

Page 26: [39] Moved to page 24 (Move #17) Piers Forster 25/05/2023 07:42:00



Page 26: [40] Deleted Piers Forster 25/05/2023 07:42:00



Page 26: [41] Deleted Piers Forster 25/05/2023 07:42:00



Page 26: [42] Deleted Piers Forster 25/05/2023 07:42:00



Page 26: [43] Deleted Piers Forster 25/05/2023 07:42:00



Page 26: [44] Deleted Piers Forster 25/05/2023 07:42:00



Page 26: [45] Deleted Piers Forster 25/05/2023 07:42:00



Page 26: [46] Deleted Piers Forster 25/05/2023 07:42:00



Page 32: [47] Deleted Piers Forster 25/05/2023 07:42:00



Page 32: [47] Deleted Piers Forster 25/05/2023 07:42:00



Page 32: [48] Inserted Cells Piers Forster 25/05/2023 07:42:00

Inserted Cells

Page 32: [49] Inserted Cells Piers Forster 25/05/2023 07:42:00

Inserted Cells

Page 32: [50] Deleted Piers Forster 25/05/2023 07:42:00



Page 32: [50] Deleted Piers Forster 25/05/2023 07:42:00



Page 32: [50] Deleted Piers Forster 25/05/2023 07:42:00



Page 32: [51] Inserted Cells Piers Forster 25/05/2023 07:42:00

Inserted Cells

Page 32: [53] Deleted **Piers Forster** **25/05/2023 07:42:00**



Page 32: [53] Deleted **Piers Forster** **25/05/2023 07:42:00**



Page 32: [54] Inserted Cells **Piers Forster** **25/05/2023 07:42:00**

Inserted Cells

Page 32: [55] Inserted Cells **Piers Forster** **25/05/2023 07:42:00**

Inserted Cells

Page 32: [56] Inserted Cells **Piers Forster** **25/05/2023 07:42:00**

Inserted Cells

Page 32: [57] Deleted Cells **Piers Forster** **25/05/2023 07:42:00**

Deleted Cells

Page 32: [58] Deleted **Piers Forster** **25/05/2023 07:42:00**



Page 32: [58] Deleted **Piers Forster** **25/05/2023 07:42:00**



Page 32: [58] Deleted **Piers Forster** **25/05/2023 07:42:00**



Page 32: [59] Inserted Cells **Piers Forster** **25/05/2023 07:42:00**

Inserted Cells

Page 32: [60] Inserted Cells **Piers Forster** **25/05/2023 07:42:00**

Inserted Cells

Page 32: [61] Inserted Cells **Piers Forster** **25/05/2023 07:42:00**

Inserted Cells

Page 32: [62] Deleted **Piers Forster** **25/05/2023 07:42:00**



Page 32: [62] Deleted **Piers Forster** **25/05/2023 07:42:00**



Page 32: [62] Deleted **Piers Forster** **25/05/2023 07:42:00**



Page 32: [63] Inserted Cells **Piers Forster** **25/05/2023 07:42:00**

Inserted Cells

Page 32: [66] Deleted Piers Forster 25/05/2023 07:42:00

Page 32: [67] Deleted Piers Forster 25/05/2023 07:42:00

Page 32: [68] Deleted Piers Forster 25/05/2023 07:42:00

Page 32: [69] Deleted Piers Forster 25/05/2023 07:42:00

Page 32: [70] Deleted Piers Forster 25/05/2023 07:42:00

Page 32: [71] Deleted Piers Forster 25/05/2023 07:42:00

Page 32: [72] Deleted Piers Forster 25/05/2023 07:42:00

Page 32: [73] Moved to page 29 (Move #44) Piers Forster 25/05/2023 07:42:00

Page 32: [73] Moved to page 29 (Move #44) Piers Forster 25/05/2023 07:42:00

Page 32: [74] Deleted Piers Forster 25/05/2023 07:42:00

Page 32: [75] Deleted Piers Forster 25/05/2023 07:42:00

Page 32: [76] Deleted Piers Forster 25/05/2023 07:42:00

Page 32: [77] Deleted Piers Forster 25/05/2023 07:42:00

Page 32: [78] Deleted Piers Forster 25/05/2023 07:42:00

Page 32: [79] Deleted Piers Forster 25/05/2023 07:42:00

Page 32: [80] Deleted Piers Forster 25/05/2023 07:42:00

Page 36: [81] Deleted Piers Forster 25/05/2023 07:42:00

Page 36: [84] Deleted Piers Forster 25/05/2023 07:42:00

▼
Page 36: [85] Deleted Piers Forster 25/05/2023 07:42:00

▼
Page 36: [86] Deleted Piers Forster 25/05/2023 07:42:00

▼
Page 36: [87] Deleted Piers Forster 25/05/2023 07:42:00

▼
Page 36: [88] Deleted Piers Forster 25/05/2023 07:42:00

▼
Page 40: [89] Inserted Cells Piers Forster 25/05/2023 07:42:00

Inserted Cells

Page 40: [90] Deleted Cells Piers Forster 25/05/2023 07:42:00

Deleted Cells

Page 40: [91] Deleted Cells Piers Forster 25/05/2023 07:42:00

Deleted Cells

Page 40: [92] Deleted Cells Piers Forster 25/05/2023 07:42:00

Deleted Cells

Page 40: [93] Deleted Cells Piers Forster 25/05/2023 07:42:00

Deleted Cells

Page 40: [94] Deleted Cells Piers Forster 25/05/2023 07:42:00

Deleted Cells

Page 40: [95] Deleted Piers Forster 25/05/2023 07:42:00

Page 40: [96] Inserted Cells Piers Forster 25/05/2023 07:42:00

Inserted Cells

Page 40: [97] Inserted Cells Piers Forster 25/05/2023 07:42:00

Inserted Cells

Page 40: [98] Deleted Cells Piers Forster 25/05/2023 07:42:00

Deleted Cells

Page 40: [99] Deleted Cells Piers Forster 25/05/2023 07:42:00

Deleted Cells

Page 40: [100] Deleted Cells Piers Forster 25/05/2023 07:42:00

Deleted Cells

Page 40: [101] Deleted Cells Piers Forster 25/05/2023 07:42:00

Page 40: [104] Deleted Piers Forster 25/05/2023 07:42:00

Page 40: [105] Deleted Piers Forster 25/05/2023 07:42:00

Page 40: [106] Deleted Piers Forster 25/05/2023 07:42:00

Page 40: [107] Deleted Piers Forster 25/05/2023 07:42:00

Page 40: [108] Deleted Piers Forster 25/05/2023 07:42:00

Page 40: [108] Deleted Piers Forster 25/05/2023 07:42:00

Page 40: [109] Deleted Piers Forster 25/05/2023 07:42:00

Page 41: [110] Deleted Piers Forster 25/05/2023 07:42:00

| | | | |
|--|--------------------------------------|--------------------------------------|--|
| Greenhouse gas concentrations AR6 WGI Chapter 2: Gulev et al. (2021) | <u>2019:</u> | <u>2022:</u> | Updates based on NOAA data as AGAGE not yet available for 2022. To make an AR6-like product, N ₂ O scaled to approximate NOAA-AGAGE average (Sect. 3) |
| | CO ₂ , 410.1 ± 0.36 ppm | CO ₂ , 417.1 ± 0.4 ppm | |
| | CH ₄ , 1866.3 ± 3.2 ppb | CH ₄ , 1911.9 ± 3.3 ppb | |
| | N ₂ O, 332.1 ± 0.7 ppb | N ₂ O, 335.9 ± 0.4 ppb | |

Page 43: [111] Deleted Piers Forster 25/05/2023 07:42:00

Page 43: [112] Deleted Piers Forster 25/05/2023 07:42:00

Page 47: [113] Deleted Piers Forster 25/05/2023 07:42:00

Page 57: [114] Deleted Piers Forster 25/05/2023 07:42:00

Page 57: [115] Deleted Piers Forster 25/05/2023 07:42:00

Page 57: [116] Deleted Piers Forster 25/05/2023 07:42:00

Page 57: [117] Deleted Piers Forster 25/05/2023 07:42:00

▼.....
Page 57: [120] Deleted Piers Forster 25/05/2023 07:42:00

▼.....
Page 57: [121] Deleted Piers Forster 25/05/2023 07:42:00

▼.....
Page 57: [122] Deleted Piers Forster 25/05/2023 07:42:00

▼.....
Page 57: [123] Deleted Piers Forster 25/05/2023 07:42:00