

High resolution aerosol data from the top 3.8 ka of the EGRIP ice core

Tobias Erhardt^{1,2,*}, Camilla M. Jensen^{1,*}, Florian Adolphi^{1,2}, Helle Astrid Kjær³, Remi Dallmayr², Birthe Twarloh², Melanie Behrens², Motohiro Hirabayashi⁴, Kaori Fukuda⁴, Jun Ogata⁴, François Burgay^{5,6}, Federico Scoto^{5,7}, Ilaria Crotti⁵, Azzurra Spagnesi^{5,8}, Niccoló Maffezzoli^{5,8}, Delia Segato^{5,8}, Chiara Paleari⁹, Florian Mekhaldi⁹, Raimund Muscheler⁹, Sophie Darfeuil¹⁰, and Hubertus Fischer¹

¹Climate and Environmental Physics, Physics Institute, and Oeschger Center for Climate Change Research, University of Bern, Bern, Switzerland

²Alfred Wegener Institute, Helmholtz Center for Polar and Marine Science, Bremerhaven, Germany

³Physics of Ice, Climate and Earth, Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark

⁴National Institute of Polar Research, Research Organization of Information and Systems, Tokyo, Japan

⁵Ca' Foscari University of Venice, Department of Environmental Sciences, Informatics and Statistics, Venice, Italy

⁶Paul Scherrer Institute, Laboratory of Environmental Chemistry (LUC), Villigen PSI, Switzerland

⁷National Research Council of Italy, Institute of Atmospheric Sciences and Climate (ISAC-CNR), Lecce, Italy

⁸National Research Council of Italy, Institute of Polar Sciences, Venice, Italy

⁹Department of Geology, Lund University, Lund, Sweden

¹⁰University Grenoble Alpes, IRD, CNRS, INRAE, Grenoble INP, IGE, Grenoble, France

*These authors contributed equally to this work.

Correspondence: Tobias Erhardt (tobias.erhardt@awi.de)

Abstract. Here we present the high-resolution CFA data from the top 479 m of the East Greenland Ice coring Project (EGRIP) ice core covering the past 3.8 thousand years. The data consists of 1 mm-depth-resolution profiles of calcium, sodium, ammonium, nitrate and electrolytic conductivity as well as decadal averages of these profiles. [The nominally 1-mm data represents an oversampling of the record as the true resolution is limited by the analytical setup to approximately 1 cm.](#) Alongside the data we provide a description of the measurement setup, procedures, the relevant references for the specific methods as well as an assessment of the precision of the measurements, the sample to depth assignment and the depth and temporal resolution of the data set. ~~The nominally 1-mm data represents an oversampling of the record as the true resolution is limited by the analytical setup to approximately 1 cm.~~ The error of absolute depth assignment of the data may be on the order of ± 2 cm, however relative depth offsets between the records of the individual species is only on the order of 1 mm. The presented data has sub-annual resolution over the entire depth range and has already formed part of the data for an annually layer-counted time scale for the EGRIP ice core used to improve and revise the multi-core Greenland ice-core chronology (GICC05) to a new version, GICC21 (Sinnl et al., 2022). The data is available in full 1-mm resolution and decadal averages on PANGAEA (<https://doi.org/10.1594/PANGAEA.945293> (Erhardt et al., 2022b))

1 Introduction

15 Ice cores from polar regions and their proxy records have allowed us to study the past climate and its variability in great detail. Concentrations of aerosol constituents in the ice are among the widely used ice-core proxy records. Deposited onto the ice sheet they are well preserved in the ice matrix as either soluble or insoluble impurities accessible through various analysis techniques. They reflect the past aerosol deposition to the ice sheets at high temporal resolution, often allowing us to resolve the seasonal variability of the aerosols species if measured at high enough depth resolution. The high temporal resolution in
20 turn can enable the generation of annual-layer counted age models from the aerosol records (e.g. [Sinnl et al., 2022](#)). Aerosol records of both shallow and deep ice cores are routinely measured at high depth resolution using continuous melting and analysis setups, i.e., using so-called continuous flow analysis (CFA) (e.g. Sigg et al., 1994; Kaufmann et al., 2008). These setups have overcome some of the analytical challenges/limitations that the very low impurity concentrations and the desired high depth resolution pose and have become the [gold-de facto](#) standard for long, high-resolution aerosol records.

25 Typically, ice cores that aim to provide climate reconstructions far back in time are drilled away from fast-flowing ice, ideally on the domes and ice-divides of an ice sheet, where effects of glacial flow on the records are minimal. The East Greenland ice coring project ice core (EGRIP) with its location inside of the North East Greenland ice stream (NEGIS) (Vallelonga et al., 2014; Hvidberg et al., 2020) is far from this ideal. However, the goal of this project was not only a long climate record, covering the Holocene and the Glacial Termination, but rather to the study of the NEGIS itself, including its causes and
30 temporal changes. The high-resolution chemical impurity records from the EGRIP ice core provide not only a climate record, but allow the detailed study of the interactions between impurities, micro-structure and ice rheology in this unique glaciological setting (Stoll et al., 2021, 2022).

In this paper we present data from the Bern wet-chemistry CFA, covering Ca^{2+} (calcium), NH_4^+ (ammonium) and NO_3^- (nitrate) concentrations, as well as electrolytic meltwater conductivity. Furthermore we provide data from the online ICP-
35 TOFMS measurements of Ca and Na (sodium) concentrations. Along with the data sets, we describe the changes made to the Bern CFA system for the EGRIP CFA campaigns since the last major measurement campaigns (Kaufmann et al., 2008; Erhardt et al., 2022a). These consist of the addition of an inductively coupled plasma time-of-flight mass spectrometer (ICP-TOFMS) to the system (Erhardt et al., 2019) as well as updated melting and calibration procedures for the wet-chemistry CFA system. We also provide a discussion of the uncertainties related to these changes covering analytical precision, depth-assignment and
40 resolution. This paper accompanies the publication of the revision of the Greenland ice core chronology (GICC21) presented in [Sinnl et al. \(2022\)](#), that makes extensive use of the data sets shown and discussed here.

2 Coring location

The EGRIP drill camp is located 359 km NNE of the Greenland Summit at 75.63° N, 36.00° W and an altitude of 2708 m a.s.l. as shown in Figure 1. Current [ice-equivalent](#) accumulation rates at EGRIP are around 11 cm a^{-1} (Vallelonga et al., 2014).
45 Drilling at the site commenced in 2016 and has reached a depth of 2121 m in 2019 when the drilling was interrupted due to technical difficulties and had to be postponed in the following seasons due to the COVID-19 pandemic. [Bedrock at the drill site](#)

[was reached in July 2023](#). The site is located within NEGIS, 150 km down-stream of the ice divide. Horizontal surface velocity at the coring location is around 55 m a^{-1} as determined by GPS measurements between 2015–2019 (Hvidberg et al., 2020). For all the data from the EGRIP ice core, including the data presented here, the coring location in the ice stream means, that going back further in time, the snow that formed the archive was deposited further upstream, at higher elevation and closer to the ice divide. [At An inversion of a two-dimensional flow model along different flow-lines to EGRIP using radar-stratigraphic horizons shows, that at](#) 3.8 ka, the ice likely originated approximately 111 km further upstream at an elevation about 160 m higher than the coring location and closer to the ice divide (Gerber et al., 2021). This area is characterized by a slightly higher [ice-equivalent](#) accumulation rate of around 15 cm a^{-1} . This upstream accumulation rate increase balances out the down-core thinning of the annual layers, leading to almost constant annual layer thickness over large parts of the Holocene (Gerber et al., 2021; Mojtabavi et al., 2020).




Figure 1. Coring location of the EGRIP ice core in bold with other major ice core drilling sites in Greenland

3 Sample description

For the continuous flow analysis measurements, samples with a cross section of 36 mm by 36 mm and a length of 0.55m were melted on a gold-plated melthead and only the innermost 26 mm by 26 mm area was used to supply the clean meltwater stream (Bigler et al., 2011). The measurements of the data down to 479 m depth presented here were performed in two melting campaigns in 2018 and 2019 covering the depth range of 13.75–350.35 m and 350.35–904.75 m respectively. The samples for the measurements down to 350.35 m were entirely cut in the field and 0.55 m pieces were then shipped via Copenhagen to

Bern for analysis. Below that, the chemistry CFA sticks and the neighboring gas CFA sticks were not split in the field but were shipped as 36 mm by 72 mm by 0.55 m slabs to Copenhagen, where they were split prior to the shipment to Bern. This was done to guarantee higher sample quality for the CFA measurements in the brittle-ice section of the core, starting below the depth section presented here.

4 EGRIP continuous flow analysis

All measurements were performed at the continuous flow analysis lab at the University of Bern, instead of in the field as had been the case for the NGRIP and NEEM deep-drilling campaigns in Greenland. This change was necessary due to the addition of an ICP-TOFMS to the setup that is not easily field-deployable. In the CFA lab, the melthead is located in a -20°C cold cell that is also used to prepare the samples for the measurements. The meltwater is pumped to the warm part of the lab for subsequent analysis.

For the EGRIP measurement campaigns, the analytical methods of the Bern CFA system were used basically unchanged from the NEEM (North Greenland Eemian Ice drilling, Dahl-Jensen et al. (2002)) system (Kaufmann et al., 2008; Erhardt et al., 2022a) with the exception of some changes to the melting procedures and the addition of a directly coupled ICP-TOFMS (icpTOF R, TOFWERK, Thun, Switzerland) (Erhardt et al., 2019). The addition of an ICP-TOFMS constitutes the largest change to the previously described system and will be outlined separately in Section 4.2.




Figure 2. Water flow distribution from the central part of the melthead between all channels in the EGRIP CFA setup used for the top 350 m of the core in the 2018 melt campaign. Not shown is the meltwater stream from the outside of the core sample that was collected for cosmogenic radionuclide measurements. All arrows denote peristaltic pump channels with pump rates given in ml min^{-1} . Dashed connections denote mixed air/water streams, solid lines pure water streams. Bold labels indicate channels used for the data presented here. Bag mean FC denotes fraction collection (FC) at 0.55 m resolution.

In addition to the measurements described here, the CFA system supplied clean meltwater streams to additional analysis channels and sampling efforts. The complete distribution of the meltwater streams between the different analysis channels of the Bern system and the measurements of sampling efforts of the partner institutes for the 2018 melting campaign are shown

in Figure 2. These were setup and run in Bern by the international partners in the EGRIP project and include the continuous analysis of dissolved iron (Fe) and FeII (Burgay et al. (2018), run by the University of Venice, Italy), Black Carbon using an SP2 instrument (Mori et al. (2016), run by the National Institute for Polar Research, Japan) and an absorption spectroscopic measurement of the meltwater acidity (Kjær et al. (2016), run by the Center for Ice and Climate, Copenhagen, Denmark). The acidity measurements were integrated into the Bern CFA system in 2019 (for data starting from 350.35 m) using a purpose-built spectrometer similar to the once used for all other channels in the Bern system. Furthermore, samples at ~~5~~2.5 cm and 2.5 cm depth resolution for ion chromatography (run by the Alfred Wegener Institute, Bremerhaven, Germany) as well as multiple ~~aliquotes~~aliquots at 55-cm resolution were collected for various offline measurements run by the international partners. Finally, excess meltwater from the outer part of the ice stick was collected in 50 ml aliquotes for high-resolution measurements of cosmogenic ^{10}Be and ^{36}Cl (Paleari et al., 2022a, b).

4.1 Wet chemistry measurements

In the following section we provide the main references that describe the system and refer the reader to Erhardt et al. (2022a) for a more detailed discussion of the system and its development over the years. The wet-chemistry (or main Bern CFA) system uses a range of analyte specific absorption and fluorescence spectroscopy methods to measure the low concentration of dissolved Ca^{2+} (Tsien et al., 1982), Na^+ (Quiles et al., 1993), NH_4^+ (Genfa and Dasgupta, 1989), NO_3^- (McCormack et al., 1994) and H_2O_2 (Dasgupta and Hwang, 1985) (hydrogen peroxide) in the ice. All of these methods were specifically adapted for the CFA ice-core measurements (Sigg et al., 1994; Röthlisberger et al., 2000; Kaufmann et al., 2008) and employ custom-built spectrometers as described in detail in (Kaufmann et al., 2008). Furthermore, the system employs a conductivity cell (Amber Science) to determine the electrolytic meltwater conductivity (Cond.) and a laser attenuation particle counter and sizer (Abalus, Klotz GmbH) for insoluble particle measurements. In 2019 (starting from 350.35 m) the absorption spectroscopic Na^+ measurement was dropped in favor of the more reliable and higher resolution measurement by the ICP-TOFMS introduced prior to the 2018 melting and discussed below (Erhardt et al., 2019). Furthermore, H_2O_2 measurements were halted after the 2018 season because of the loss of a useful signal due to excessive diffusion. Here we only provide data of Ca^{2+} , NH_4^+ and NO_3^- concentrations, electrolytic meltwater conductivity ~~and ICP-TOFMS measurements of and concentrations~~from the wet-chemistry CFA.

4.2 Continuous ICP-TOFMS

The addition of an inductively coupled plasma time-of-flight mass spectrometry (ICP-TOFMS) is the biggest analytical change of the Bern CFA system in comparison to previous melting campaigns. Coupling and integration of the ICP-TOFMS to the Bern CFA setup and the instrument parameters have been described in detail in Erhardt et al. (2019) and will only briefly be outlined in the following. The main feature of the icpTOF is the fact that the time-of-flight mass spectrometer can measure the complete mass range of m/z 23 to 275 at a sampling rate of 33 kHz allowing the temporal resolution of short transient signals and coverage of almost the entire table of elements. For the CFA analysis, the instrument is run in two different data acquisition modes: either at an integration time of 250 ms for normal data acquisition of total elemental concentrations and

at 2 ms integration time to also resolve the ionization events of individual particles entering the plasma in the single-particle
115 mode.

The sample stream for the ICP-TOFMS is diverted close to the melthead from the main meltwater stream. The air-water mixture coming from the melthead is used to efficiently transport particulates to the instrument and to limit signal dispersion. ~~Note, that the~~ The air is released from the ice during melting and thus does not pose a contamination risk as would other external gas sources. The air is only removed very close to the instrument using a custom build Teflon micro-volume debubbler
120 with a volume of approximately 100 μl . The bubble-free sample stream is then slightly acidified to 1 % using high-purity nitric acid (Optima Grade, Fisher Scientific) close to the sample introduction of the instrument. In this way washout times from the nebulizer and spray chamber can be minimized without risking the dissolution of particles in the ice which are the focus of the single-particle measurements. During the acidification, Rh (rhodium) is added to the sample as internal standard to monitor system stability and correct for sensitivity drifts. The sample is introduced into the plasma via a quartz cyclonic spray chamber
125 at a flow rate of 400 $\mu\text{l min}^{-1}$ using a glass concentric nebulizer (Micro Mist, Glass Expansion) for the 2018 melting campaign (to 350 m depth) and a Teflon nebulizer (MicroFlow PFA-ST, Elemental Scientific) thereafter. Detailed information on plasma conditions used for both cases can be found in (Erhardt et al., 2019).

Before and after each measurement run the instrument is calibrated with a range of multi-element standards. All standards are prepared by gravimetrically mixing multi-element standards (Sigma Aldrich, Inorganic Ventures) for a stock solution which
130 is then further diluted into working standards that span the expected range of concentrations in the ice. Due to the wide mass range that is covered by the TOFMS and the fact that now mass-scanning is necessary to acquire different masses, the range of analytes that can be targeted is basically only limited by their calibration and their possible interference. Even though the data presented here only contains Na and Ca concentrations, the instrument is routinely calibrated for a wide range of analytes including halogens, rare earth elements as well as heavy metals and platinum group elements. These data are still
135 under evaluation and are not provided here. For the calibration measurements a SC- μDX auto sampler with FAST injection valve (Elemental Scientific) was used to allow for repeatable and time efficient calibrations. Measurements were performed by injection of a 400 μl loop into a dilute nitric acid carrier stream which was run continuously between measurement runs.

For the data presented in this study, both normal (4 Hz) and single-particle (500 Hz) acquisition data were averaged to the nominal 1-mm resolution used for all of the data series from the main CFA system. The final data set is then smoothed with
140 a 10 s cutoff frequency Gaussian filter to remove high-frequency measurement noise from the data set. The depth assignment of the ICP-TOFMS data was achieved by aligning the Ca data to the raw fluorimetric Ca^{2+} measurements of the main CFA system and thus to its depth-scale.

Figure 3 shows an overview of all data sets provided here at a 10 yr resolution on the GICC21 ~~(?)~~ (Sinnl et al., 2022) age scale, both from the wet-chemistry CFA as well as from the ICP-TOFMS. Note, that even though the final data set of Ca^{2+}
145 exhibits large gaps, the depth assignment of the ICP-TOFMS Ca data set is done using the raw fluorescence intensity data, as elaborated above. Thus the depth assignment of the Ca data is well constrained even when calibration issues affects the Ca^{2+} data as described in Section 5.1.




Figure 3. Overview of the complete data set at 10 yr resolution on the GICC21 (Sinnl et al., 2022) agescale. From top to bottom the data series are electrolytic meltwater conductivity (Cond.), Ca^{2+} measured by fluorescence spectrometry, Ca and Na measured by ICP-TOFMS, NH_4^+ measured by fluorescence spectrometry and NO_3^- , measured by absorption spectrometry. Note, that the Ca^{2+} record partly suffers from analytical issues leading to a large number of gaps in the final data set. However, the Ca data from the ICP-TOFMS covers ~~not~~ most of these gaps and provides an almost complete data set.

4.3 Updated melting and measurement procedure

Previously, the Bern CFA system was run with 1.10 m to 1.65 m of ice per run at most (Erhardt et al., 2022a), i.e. 2–3 times 150 0.55 m per continuous measurement run. This maximum length is dictated by the length of the frames used to guide the ice during melting and by the vertical space available for the melting unit. Between each of the measurement runs, all connected instruments are run on an ultra pure water stream. Depending on the method it can take a significant time to reach a stable baseline signal between the runs, which is needed to catch and correct for drift in the analysis units. In a typical campaign, 2 to 3 runs were performed before a calibration run approximately every two to three hours which means that the waiting time 155 between each run adds up to a large fraction of each measurement day.

To capitalize on the stability of the well matured Bern CFA system and the environmentally more stable lab setting as compared to the field, the melting procedure for the EGRIP ice core was changed to allow the uninterrupted measurement of multiple frames worth of ice-core samples by exchanging the frames mid-run. This is achieved by adding an additional fixed sample guide shown in Figure 4 to the Teflon melthead cover similar to what is also done in other CFA labs. During a




Figure 4. Picture of the melthead and the additional transparent sample guide used for the stacking of core segments during melting.

160 measurement run, the frames can then be exchanged as soon as there is only so much ice left on the melthead (~ 10 cm) that it reaches the top of the new sample guide. To do so, the weight connected to the depth encoder [of the CFA system](#) is lifted from the remaining ice, the empty frame is removed from the holder and a new one is put in its place. The new piece of ice is then lowered carefully down to the remainder on the melthead using a Teflon wedge. Finally the encoder weight is then placed on top of the new sample. After some training, the whole procedure can be done within a few seconds during which the melting

165 of the small remainder of ice continues. In this time a little less meltwater is produced, however the amount is still enough to avoid contamination by inflow from meltwater from the outside of the sample. During the procedure no depth recording is possible, however due to the short amount of time needed to re-position the encoder on the new piece of ice only very little ice melts during this time, as discussed below. The total amount of ice that can be measured in each measurement run is limited by the system stability and the frequency of calibration runs between extended ice measurements.

170 For the data presented here, usually a total of 3.30 m of ice were melted within each run, amounting to approximately two hours of uninterrupted measurement time per run at a meltspeed of 2.7 cm min^{-1} . During each measurement day, a total of four runs with bracketing calibrations were performed. Depending on sample quality, this amounts to a total daily production of approximately 12–14 m per ~ 16 hour workday including the daily startup and shutdown of all instruments.

5 Data quality

175 The discussion of the uncertainties is split into three major parts: The analytical precision of the measurements, the depth assignment and the resulting depth and temporal resolution. To complete the discussion we added a few notes on the effect of the meltspeed stability on the data resolution as well as a brief comparison of the to calcium data sets.

5.1 Analytical precision

Because the spectroscopic methods used here were essentially not modified since the last measurement campaign (i.e. NEEM), the detection limits and uncertainties are assumed to have remained largely unchanged. The relevant numbers for each of the detection channels are listed in Table 1. As discussed in Gfeller et al. (2014), the uncertainty in the wet-chemistry is dominated by the uncertainty of the standard preparation by use of the water dispenser and microliter-pipette. For the concentration ranges discussed here, these uncertainties typically amount to less than 10%. For the wet-chemistry CFA data, the limit of detection (LOD) was determined as three times the standard deviation of the baseline/blank signal from ultra pure water (mQ, Merck/Millipore). In the case of the Ca^{2+} data, an issue with the fluorescence detector caused a low sensitivity in the top 322.3 m of the data set leading to a much higher limit of detection in this section of the core as indicated by the two different values in Table 1. The lower sensitivity and the resulting higher limit of detection also affects the Ca^{2+} concentration range for the upper part of the core as values below the LOD were removed from the data.

Table 1. Analytical figures of merit for the wet-chemistry data. The sample concentration ranges are given by the 5 th and 95 th percentile and the median value in parentheses. The limit of detection is calculated as three times the standard deviation of the blank signal. The temporal resolution of the detection channels is determined as the long-term average 10–90% rise time which is converted to an equivalent depth resolution assuming a melt speed of 2.7 cm min^{-1} . Note that this resolution does not account for signal dispersion during melting and in the debubbler.

	Range (ppb)	LOD (ppb)	t_{10-90} (s)	Res. (cm)
Ca^{2+} (<322.3 m)	2.3–23.3 (6.8)	2.2	22.9	1.06
Ca^{2+} (>322.3 m)	1.5–20.1 (5.4)	0.2	22.9	1.06
NH_4^+	0.8–28.0 (3.7)	0.3	22.1	1.00
NO_3^-	61.8–162.4 (99.1)	1.2	26.9	1.21
Cond.	0.88–1.82 (1.16)	—	20.6	0.93

For the ICP-TOFMS measurements of Ca and Na, the relevant performance numbers are given in Table 2. Here the values include the relative uncertainties from the standard errors of the calibrations. In the case of the ICP-TOFMS data, LODs were determined using three times the standard error of the regression for the calibrations. These regressions include the blank measurements and allow for a non-zero intercept. This approach was chosen as it yielded more reliable results and allows for the determination of the LODs even in the occasional case if no reliable blank measurement could be performed.

5.2 Depth assignment

Previous investigations (Röthlisberger et al., 2000; Hiscock et al., 2013) have reported the uncertainty of the depth assignment of the 110–165 cm measurements runs as $\pm 1 \text{ cm}$. The total precision of the depth assignment is likely limited by the precision of the length measurements during sample preparation due to possible influence of parallax effects and the limited precision of rulers (Erhardt et al., 2022a). However, due to the aforementioned change in the melt protocol in comparison to previous

Table 2. Analytical figures of merit for the ICP-TOFMS data. The sample concentration ranges are given by the 5 th and 95 th percentile and the median value in parentheses. The relative uncertainty (Rel. Uncert.) values are given for the respective concentrations in the range column. The limit of detection is determined from three times the standard error of the regression line at its intercept. The temporal resolution of the detection channels is determined as the long-term average 10–90 % rise time which is converted to an equivalent depth resolution assuming a melt speed of 2.7 cm min⁻¹. Note that this resolution does not account for signal dispersion during melting and in the debubbler.

	Range (ppb)	Rel. Uncert. (%)	LOD (ppb)	t_{10-90} (s)	Res. (cm)
Ca	1.8–23.9 (5.8)	6.6–3.6 (3.8)	0.28	20.5	0.92
Na	0.4–22.3 (3.9)	14.1–1.3 (1.9)	0.17	17.9	0.81

studies, the quality of the depth assignment merits some additional discussion below. Overall, the uncertainty of the absolute
 200 depth assignment of all records to the ice-core depth is about ± 2 cm, while the offsets between the individual CFA records is only on the order of 1 mm.

As mentioned above, during the stacking, the weight that is connected to the depth encoder of the CFA system is lifted from the ice. During that time, the remaining ~ 10 cm of ice ~~slowly continues~~ continue to melt, albeit slower than usual, due to the missing pressure from the encoder weight. Due to the encoder being removed during that time, usually for around 5–15 s, the
 205 progress of the melting is unrecorded. The resulting gaps in the depth assignment are filled during the data processing using linear interpolation. ~~Additionally~~ Furthermore, because the amount of ice is greater per run as previously, the total amount of ice contains more breaks ~~, each of which carries the risk of~~ adding to the depth-assignment uncertainty ~~due to~~. This is due to the fact that each break surface will need to be cleaned and squared, ~~leading to possible mis-measurement of the~~ each of which carries the risk of mis-measurement of the total length of ice.

210 To ~~check whether the changed procedure introduces an additional~~ investigate the influence of the stacking procedure on the depth-assignment ~~error~~ uncertainty, we look at the ~~total~~ discrepancy between the length of ice as measured by the encoder ~~to and~~ the length of ice measured ~~using sample preparation. The relative deviation of these values provides a measure of the accumulated uncertainty in the depth assignment at the end of the run. This relative deviation is shown in~~ during sample preparation, shown as relative deviation in Figure 5G ~~alongside other variables that describe the melting process. Note, however, that these deviations include the uncertainty of the length measurements during sample preparation and only show their discrepancy to the length measured by the encoder. Thus, they only provide a measure of the combined uncertainty.~~

Comparing the deviations to the length-statistics shown in Panels A-C of Figure 5, most notable, the relative deviation between melted and measured sample lengths for the runs does not show any correlation with the total amount of ice melted or the number of pieces that make up the total length of the ice. ~~Overall, we judge the relative uncertainty of the depth assignment not to be affected by the core-stacking process beyond the approximately 1~~ previously reported for solid ice (Erhardt et al., 2022a) ~~. However, in~~ This indicates, that the stacking procedure does not introduce excessive additional uncertainty.

220 In the shallow part of the ice core, ~~above the firn/ice transition down to approximately 100 m depth~~ where the melt speed is ~~significantly~~ higher than 2.7 cm min⁻¹, the melted length of ice is systematically lower than the amount measured before the

melting. We suggest ~~, that~~ this can be attributed to a slight compaction of the fragile firn during sample handling and the upright
225 melting due to the overburden of the sample and the encoder weight. ~~The change in melting procedure has not increased the
uncertainty in the depth assignment~~

The deviation between the two length measurements can also be used to estimate the overall uncertainty of the depth
assignment. On average, if the measurements are not biased and their errors are not correlated, the difference between them
should be zero and its variance is the sum of the variances of the two measurements, providing the overall squared uncertainty
230 of the depth measurement. To provide an average uncertainty we report half the overall relative standard deviation of the
difference of the known encoder calibration (cm^{-1}) to the observed number of encoder counts per cm of ice melted as upper
estimate of the depth assignment. In this way the provided uncertainty is independent from the exact amount of melted ice.
Furthermore, the beginning and the end of the run are used as reference points to fix the length of ice measured on the overall
depth-scale of the ice core. That means that the error of the depth assignment relative to the ice-core depth scale is by definition
235 zero at these points, further reducing the overall uncertainty. In the shallow part of the data, as mentioned above, one of the
measurements is biased and the aforementioned assumption not valid. Excluding measurements above 100 m depth, the relative
uncertainty of the depth assignment is 0.54 % amounting to 1.8 cm for a 3.3 m run. Over the complete depth range of the data
here the same calculation yields 0.64 % or 2.1 cm for a 3.3 m run.

Overall, we judge the relative uncertainty of the depth assignment not to be affected by the core-stacking process beyond
240 the approximately 1 ~~in solid ice that were previously reported~~ % of the amount of ice melted per measurement as previously
reported for solid ice (Erhardt et al., 2022a). As discussed in Erhardt et al. (2022a) this uncertainty mainly arises from the
sample preparation, and only to a small part from the melting ~~itself-procedure itself~~. We note that the overall uncertainty can
further be reduced by incorporating the 55-cm bag marker into the depth assignment procedure.

It is worth noting however, that these sources of uncertainty for the absolute depth assignment do not contribute to the phasing
245 error between the different analytes. The relative alignment between the parameters of the main CFA system is performed as
described previously using short injections of a multi-component standard to determine the delays to each of the wet-chemistry
channels. As discussed in (Erhardt et al., 2022a), the uncertainty of this relative phasing is on the order of 1–2 s, translating
to less than a millimeter at the melt speeds employed here. The relative alignment of the ICP-TOFMS data is performed by
alignment of the Ca to the Ca^{2+} , transferring the relative phasing uncertainty of the wet-chemistry CFA to the ICP-TOFMS
250 measurements (Erhardt et al., 2019). In summary, while the uncertainty in absolute depth of all records may be about ± 2 cm
the relative depth offsets between individual CFA records is only on the order of 1 mm.

5.3 Depth and temporal resolution

The resolution for both the wet-chemistry CFA data sets as well as the ICP-TOFMS was determined by the 10–90 % rise
time of standard measurements, treated in the same way as the sample measurements with respect to temporal resolution and
255 smoothing of the raw signals. Overall, the resolution of the data is on the order of approximately 1 cm as listed in Tables 1 and 2.
The NO_3^- data, due to the long tubing and the reaction column involved, exhibits a slightly lower resolution at 1.2 cm. Both the
electrolytic meltwater conductivity and the ICP-TOFMS measurements have slightly better resolution, below 1 cm. Note, that

these values do not include the additional dispersion that happens during melting and in the debubbling volumes. However, as argued before (Erhardt et al., 2022a; Sigg et al., 1994), the response time of the detection usually dominates the overall signal dispersion and the numbers reported can serve as a good guidance for the overall resolution of the data. Generally, the resolution of the CFA-ICP-TOFMS data is slightly better than that of the corresponding wet-chemistry measurement including the dispersion during the melting and in the respective debubbler volumes as a direct comparison of the high-frequency content of a section of the data has shown (Erhardt et al., 2019).

Figure 7 shows a 10-m section of the data at the full nominal 1-mm resolution. This example shows the very clear resolution of the seasonal variability in all of the parameters as exploited in the revision of the GICC time scale (Sinnl et al., 2022). The peak dominating the electrolytic meltwater conductivity record in Figure 7 is a typical signal from aerosol deposition after a volcanic eruption. This specific eruption signal has previously been attributed to the 79 CE Vesuvius eruption which has recently been ruled out by tephra evidence pointing to a source in the Aleutian ark (Plunkett et al., 2022). In correspondence with other dating efforts (e.g. Sigl et al., 2015), the eruption is now dated to 87/88 CE, corresponding to 1910 on the GICC21 age scale (Sinnl et al., 2022).

For the depth resolution, the melt-speed and its stability plays an important role in determining the overall resolution as well as the stability of this resolution. Especially in the shallow parts of the core, above the depth when the firn reaches the density of ice at 917 kg m^{-3} , this is an important factor. For technical reasons, i.e., to supply enough meltwater for the system, the melt speed above the firn-ice transition needs to be increased to match the melt speed in ice-equivalent depth. This is clearly visible in Figure 5E, with the exponential decrease of the melt speed for top $\sim 100 \text{ m}$ of the core. Through the adjustment of the melt speed, the data which is measured at constant time intervals is measured at a constant resolution in terms of the ice-equivalent depth. Assuming an approximate constant annual layer thickness, this translates to an approximate constant temporal resolution as well. Panel F in Figure 5 shows the variability of the melt speed within each of the measurement runs. This is a measure of how constant the resolution of the data is: high melt speed variability will translate directly into variability of the measurement resolution in terms of depth along the core. Overall, this variability given by the relative standard deviation is significantly lower than 10 %, and mostly around 5 %. Runs with much higher variability are usually caused by sample quality issues which resulted in interrupted melting. In some cases such variability can be caused by too-thin ice sections that were intentionally melted faster to avoid contamination by intrusion of meltwater from the outside of the melthead to the clean part inside. However, because of the mixing in the system and because the response time of detection method on the order of $\sim 20 \text{ s}$, any short-term variability likely plays a minor role in terms of jitter or the determination of annual layer thicknesses from the data.

5.4 Ca records

Because the Ca and Ca^{2+} concentrations were measured using two different and entirely independently calibrated methods, they allow for direct comparison between the fluorimetric detection and ICP-TOFMS measurements at high resolution. Figure 6 shows a direct comparison of the records at 10 cm resolution alongside a linear regression between the two, their correlation, root mean square (rms) difference and average ratio. Generally both data sets agree very well with each other despite the

differences in measurement techniques as indicated by the low rms difference, high correlation and close to unity average ratio. The linear regression between the two indicates a small constant offset between the two. This is likely caused by the aforementioned Ca^{2+} sensitivity and LOD issues in the top part of the data set as the offset vanishes when looking only at data
295 below 322.3 m. However, measurements below the baseline for Ca^{2+} were removed from the data set and the apparent offset is small in comparison to the concentrations and their uncertainties.

At higher resolution, as illustrated in Figure 7, the agreement between the two measurement is also very good. Focusing on 1-m sections of the full resolution 1-mm data, the average correlation between the two measurements is 0.95 with an rms difference of 2.44 ppb.

300 In summary, the agreement both at low and high resolution between the two data sets illustrates their compatibility as well as their high quality. However, it is worth noting that it is not given a priori that the two should agree as the fundamentally measure two different things. Ca^{2+} in the meltwater steam is the result of water-soluble calcium species that are dissolved during the melting and transport of the sample. These are only a subset of the total amount of Ca that is present in the meltwater which is what the ICP-TOFMS measures. For example, in case of the presence of a high load of poorly soluble Ca-bearing mineral
305 dust particles in the ice the ICP-TOFMS will detect significantly more Ca than the fluorimetric Ca^{2+} measurement. That being said, the close agreement between the two suggests that most of the total Ca present in the ice is soluble in the meltwater stream. A detailed comparison of the differences between the two and how they arise is planned.

For the drilling of the EGRIP ice core the same drill liquid combination of COASOAL and ESTISOL 240 (Popp et al., 2014) was used as for the NEEM ice core. The drill liquid acts as an absorber of bivalent cations in the CFA system if it is introduced
310 by contaminated ice. This can have a detrimental effect on the measurement of Ca^{2+} and likely all other bivalent cations by producing strong absorption/desorption signals during measurement runs (Erhardt et al., 2022a). As almost all of the Calcium detected by the ICP-TOFMS is also dissolved Ca^{2+} , the same is true for the total Ca measurements. To counteract the risk of drill liquid entering the melt system, special care was taken during sample preparation to remove all possibly contaminated ice
315 either by scraping off or cutting out possibly affected ice. This is especially important for small, refrozen and often hard to see cracks in the sample. For the data presented here, this effort, possibly in combination with the low Ca concentrations in the Holocene ice, has proven to be very fruitful. The data was found to be free of the previously described absorption/desorption effects and no correction was necessary.

5.5 Possible upstream effects

6 Possible upstream effects

320 As mentioned above, the EGRIP ice core is located inside of the Northeast Greenland Ice Stream and at the coring location surface velocities are 55 m a^{-1} (Hvidberg et al., 2020) and ice down-core originates from a location closer to the ice divide, likely characterized by higher accumulation rate (Gerber et al., 2021). This change in location and especially in the accumulation rate of about $\sim 30\%$ likely also has an influence on the aerosol data presented here both due to the change in deposition characteristics as well as due to the change in distance from possible source areas (e.g. Fischer et al., 1998a, b). Furthermore,

325 short-term variability in the data sets is likely affected by the varying influence of the local topography changes upstream (e.g. Winski et al., 2019), leading for example to accumulation variability independent of precipitation on length scales of meters (sastrugies) to kilometers (dunes) due to wind reworking. That means, that both long-term trends as well as short-term variability in the data will need to be interpreted with great care and in the light of the influence of the upstream effects.

7 Conclusions

330 Here ~~were~~we present a multi-proxy record of impurity concentrations from the EGRIP ice core covering the past 3.8 ka. The data was measured using the state-of-the-art continuous flow analysis system at the University of Bern and includes measurements of dissolved calcium, ammonium and nitrate from spectrophotometric measurements, electrolytic meltwater conductivity as well as total calcium and sodium concentrations measured by online ICP-TOFMS. The high-resolution records presented have sub-annual resolution and have formed part of the data base for the revised Greenland ice-core chronology, the
335 GICC21, over the past 3.8 ka (~~?)~~([Sinnl et al., 2022](#)).

Summarizing from the previous section we stress the following points that should be considered when using this data:

1. The data sets have undergone extensive quality control both manually and automated. However, we can not guarantee the complete absence of any spurious signals, especially at the full resolution.
2. Similar to previous CFA data sets, the depth assignment to a given down-core depth is likely accurate to better than
340 ± 2 cm. The relative depth assignment within the CFA data set is much more accurate and uncertainties are on the order of 1 mm.
3. Analytical uncertainties for the wet-chemistry data (Ca^{2+} , NH_4^+ , NO_3^-) is typically better than 10 %, for the ICP-TOFMS data (Ca, Na) better than 5 %.
4. The fluorescence spectrometric Ca^{2+} data above 322.3 m depth suffers from decreased sensitivity of the measurement
345 setup and has higher-than-usual limits of detection. Values below the baseline were removed from the data set.
5. All the aerosol concentrations in the EGRIP ice core, including the ones presented here, are likely affected by upstream effects due to changes in the local deposition regime and distance from the aerosol source regions further upstream. These effects are subject to ongoing investigation and they need to be taken into account when interpreting the data at any temporal scale.
- 350 6. Last but not least, we stress that any detailed interpretation, especially of short-term signals in the data sets should always be confirmed at least within the multi-parameter data series, if not with additional data or measurements.

We invite any future user of these data sets to actively involve CFA or ice-core specialists into their investigation for an expert assessment of the data quality/limitations in which may be specific to the application intended by the data user in order to avoid any shortcomings and pitfalls of the complex data sets.

355 **8 Data availability**

Both the full 1-mm-resolution as well as the decadal averaged data are available on PANGAEA under CC BY license (<https://doi.pangaea.de/10.1594/PANGAEA.945293> Erhardt et al., 2022b):

- Full resolution: <https://doi.org/10.1594/PANGAEA.945290>
- Decadal averages on the GICC21 age scale: <https://doi.org/10.1594/PANGAEA.945291>

360 The GICC21 age scale can be found in the supplement of Sinnl et al. (2022).

Author contributions. The CFA campaigns at the University of Bern were organized and hosted by TE and CMJ under the lead of TE. FA supported the wet-chemistry CFA measurements. The wet-chemistry CFA data was processed and quality controlled by CMJ, the ICP-TOFMS data by TE. TE wrote the initial draft of the manuscript and prepared the figures. The manuscript and figures were finalized with input from HF, CMJ, HAK, MB, DS, AS, NM, RM, FB, FA and IC. All authors took part in the measurement campaigns or contributed
365 to their success. We thank the two reviewers for their thoughtful comments and suggestions that improved the manuscript. We also thank Xingchen Wang for handling the review process as topic editor.

Competing interests. The authors declare no competing interests.

Acknowledgements. Purchase of the icpTOF has been made possible through the Swiss National Science Foundation (SNF) R'Equip proposal iceCP-TOF (grant no. 206021_170739) and specific institutional funds made available by the University of Bern. The long-term
370 financial support of ice core research at the University of Bern by the Swiss National Science Foundation (grant no. 200020_172506 (iCEP) and 20FI21_164190 (EGRIP)) is gratefully acknowledged.

EGRIP is directed and organized by the Centre for Ice and Climate at the Niels Bohr Institute, University of Copenhagen. It is supported by funding agencies and institutions in Denmark (A. P. Møller Foundation, University of Copenhagen), USA (US National Science Foundation, Office of Polar Programs), Germany (Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research), Japan (National Institute
375 of Polar Research and Arctic Challenge for Sustainability), Norway (University of Bergen and Trond Mohn Foundation), Switzerland (Swiss National Science Foundation), France (French Polar Institute Paul-Emile Victor, Institute for Geosciences and Environmental research), Canada (University of Manitoba) and China (Chinese Academy of Sciences and Beijing Normal University).

MH, KF and OJ acknowledge financial support by JSPS KAKENHI (grant number JP18H04140).

The authors gratefully acknowledge the contributions of the countless people that facilitated and took part in the field campaigns, the
380 ice-core drilling and processing as well helped during the CFA melting campaigns.

Furthermore the authors are especially grateful for the help of Priska Lehmann, Sam Black, Patrick Zens, Miriam Läderach, and Michelle Shu-Ting Lee during the campaigns.

References

- Bigler, M., Svensson, A., Kettner, E., Vallelonga, P., Nielsen, M. E., and Steffensen, J. P.: Optimization of High-Resolution
385 Continuous Flow Analysis for Transient Climate Signals in Ice Cores, *Environmental Science & Technology*, 45, 4483–4489,
<https://doi.org/10.1021/es200118j>, 2011.
- Burgay, F., Erhardt, T., Lunga, D. D., Jensen, C. M., Spolaor, A., Vallelonga, P., Fischer, H., and Barbante, C.: Fe²⁺ in
ice cores as a new potential proxy to detect past volcanic eruptions, *Science of The Total Environment*, 654, 1110–1117,
<https://doi.org/10.1016/J.SCITOTENV.2018.11.075>, 2018.
- 390 Dahl-Jensen, D., Gundestrup, N. S., Miller, H., Watanabe, O., Johnsen, S. J., Steffensen, J. P., Clausen, H. B., Svensson, A., and Larsen,
L. B.: The NorthGRIP deep drilling programme, *Annals of Glaciology*, 35, 1–4, <https://doi.org/10.3189/172756402781817275>, 2002.
- Dasgupta, P. K. and Hwang, H.: Application of a nested loop system for the flow injection analysis of trace aqueous peroxides, *Analytical
Chemistry*, 57, 1009–1012, <https://doi.org/10.1021/ac00283a010>, 1985.
- Erhardt, T., Jensen, C. M., Borovinskaya, O., and Fischer, H.: Single particle characterization and total elemental concentration measurements
395 in polar ice using CFA-icpTOF, *Environmental Science & Technology*, 53, 13 275–13 283, <https://doi.org/10.1021/acs.est.9b03886>, 2019.
- Erhardt, T., Bigler, M., Federer, U., Gfeller, G., Leuenberger, D., Stowasser, O., Röthlisberger, R., Schüpbach, S., Ruth, U., Twarloh, B.,
Wegner, A., Goto-Azuma, K., Kuramoto, T., Kjær, H. A., Vallelonga, P. T., Siggaard-Andersen, M.-L., Hansson, M. E., Benton, A. K.,
Fleet, L. G., Mulvaney, R., Thomas, E. R., Abram, N., Stocker, T. F., and Fischer, H.: High-resolution aerosol concentration data from the
Greenland NorthGRIP and NEEM deep ice cores, *Earth System Science Data*, 14, 1215–1231, <https://doi.org/10.5194/essd-14-1215-2022>,
400 2022a.
- Erhardt, T., Jensen, C., and Fischer, H.: High resolution aerosol records over the past 3.8ka from the EastGRIP ice core, PANGAEA, 2022b.
- Fischer, H., Wagenbach, D., and Kipfstuhl, J.: Sulfate and Nitrate Firm Concentrations on the Greenland Ice Sheet. 1. Large-Scale Geograph-
ical Deposition Changes, *Journal of Geophysical Research*, 103, 1–8, <https://doi.org/10.1029/98JD01885>, 1998a.
- Fischer, H., Werner, M., Wagenbach, D., Schwager, M., Thorsteinsson, T., Wilhelms, F., Kipfstuhl, J., and Sommer, S.: Little Ice Age clearly
405 recorded in northern Greenland ice cores, *Geophysical Research Letters*, 25, 1749–1752, <https://doi.org/10.1029/98GL01177>, 1998b.
- Genfa, Z. and Dasgupta, P. K.: Fluorometric measurement of aqueous ammonium ion in a flow injection system, *Analytical Chemistry*, 61,
408–412, <https://doi.org/10.1021/ac00180a006>, 1989.
- Gerber, T. A., Hvidberg, C. S., Rasmussen, S. O., Franke, S., Sinnl, G., Grinsted, A., Jansen, D., and Dahl-Jensen, D.: Upstream flow effects
revealed in the EastGRIP ice core using Monte Carlo inversion of a two-dimensional ice-flow model, *The Cryosphere*, 15, 3655–3679,
410 <https://doi.org/10.5194/tc-15-3655-2021>, 2021.
- Gfeller, G., Fischer, H., Bigler, M., Schüpbach, S., Leuenberger, D., and Mini, O.: Representativeness and seasonality of major ion records
derived from NEEM firn cores, *The Cryosphere*, 8, 1855–1870, <https://doi.org/10.5194/tc-8-1855-2014>, 2014.
- Hiscock, W. T., Fischer, H., Bigler, M., Gfeller, G., Leuenberger, D., and Mini, O.: Continuous Flow Analysis of Labile Iron in Ice-Cores,
Environmental Science & Technology, 47, 4416–4425, <https://doi.org/10.1021/es3047087>, 2013.
- 415 Hvidberg, C. S., Grinsted, A., Dahl-Jensen, D., Khan, S. A., Kusk, A., Andersen, J. K., Neckel, N., Solgaard, A., Karlsson, N. B., and Kjær,
H. A.: Surface velocity of the Northeast Greenland Ice Stream (NEGIS): assessment of interior velocities derived from satellite data by
GPS, *The Cryosphere*, 14, 3487–3502, <https://doi.org/10.5194/tc-14-3487-2020>, 2020.

- Kaufmann, P. R., Federer, U., Hutterli, M. A., Bigler, M., Schüpbach, S., Ruth, U., Schmitt, J., and Stocker, T. F.: An Improved Continuous Flow Analysis System for High-Resolution Field Measurements on Ice Cores, *Environmental Science & Technology*, 42, 8044–8050, <https://doi.org/10.1021/es8007722>, 2008.
- 420 Kjør, H. A., Vallelonga, P., Svensson, A., Elleskov L. Kristensen, M., Tibuleac, C., Winstrup, M., and Kipfstuhl, J.: An Optical Dye Method for Continuous Determination of Acidity in Ice Cores, *Environmental Science & Technology*, 50, 10485–10493, <https://doi.org/10.1021/acs.est.6b00026>, 2016.
- McCormack, T., David, A. R., Worsfold, P. J., and Howland, R.: Flow injection determination of nitrate in estuarine and coastal waters, *Analytical Proceedings including Analytical Communications*, 31, 81–83, <https://doi.org/10.1039/A19943100081>, 1994.
- 425 Mojtabavi, S., Wilhelms, F., Cook, E., Davies, S. M., Sinnl, G., Skov Jensen, M., Dahl-Jensen, D., Svensson, A., Vinther, B. M., Kipfstuhl, S., Jones, G., Karlsson, N. B., Faria, S. H., Gkinis, V., Kjør, H. A., Erhardt, T., Berben, S. M. P., Nisancioglu, K. H., Koldtoft, I., and Rasmussen, S. O.: A first chronology for the East Greenland Ice-core Project (EGRIP) over the Holocene and last glacial termination, *Climate of the Past*, 16, 2359–2380, <https://doi.org/10.5194/cp-16-2359-2020>, 2020.
- 430 Mori, T., Moteki, N., Ohata, S., Koike, M., Goto-Azuma, K., Miyazaki, Y., and Kondo, Y.: Improved technique for measuring the size distribution of black carbon particles in liquid water, *Aerosol Science and Technology*, 50, 242–254, <https://doi.org/10.1080/02786826.2016.1147644>, 2016.
- Paleari, C., Mekhaldi, F., Adolphi, F., Christl, M., Vockenhuber, C., Gautschi, P., Beer, J., Brehm, N., Erhardt, T., Synal, H., Wacker, L., Wilhelms, F., and Muscheler, R.: Cosmogenic radionuclides reveal an extreme solar particle storm near a solar minimum 9125 years BP, *Nature Communications*, 13, 214, <https://doi.org/10.1038/s41467-021-27891-4>, 2022a.
- 435 Paleari, C. I., Mekhaldi, F., Erhardt, T., Zheng, M., Christl, M., Adolphi, F., Hörhold, M., and Muscheler, R.: Evaluating the 11-year solar cycle and short-term ^{10}Be deposition events with novel excess water samples from the EGRIP project, *Climate of the Past Discussions*, in review, <https://doi.org/10.5194/cp-2022-94>, 2022b.
- Plunkett, G., Sigl, M., Schwaiger, H. F., Tomlinson, E. L., Toohey, M., McConnell, J. R., Pilcher, J. R., Hasegawa, T., and Siebe, C.: No evidence for tephra in Greenland from the historic eruption of Vesuvius in 79 CE: implications for geochronology and paleoclimatology, *Climate of the Past*, 18, 45–65, <https://doi.org/10.5194/cp-18-45-2022>, 2022.
- 440 Popp, T. J., Hansen, S. B., Sheldon, S. G., and Panton, C.: Deep ice-core drilling performance and experience at NEEM, Greenland, *Annals of Glaciology*, 55, 53–64, <https://doi.org/10.3189/2014AoG68A042>, 2014.
- Quiles, R., Fernández-Romero, J., Fernández, E., Luque de Castro, M., and Valcárcel, M.: Automated enzymatic determination of sodium in serum, *Clinical Chemistry*, 39, 500–503, <https://doi.org/10.1093/clinchem/39.3.500>, 1993.
- 445 Röthlisberger, R., Bigler, M., Hutterli, M. A., Sommer, S., Stauffer, B., Junghans, H. G., Wagenbach, D., Stauffer, B., Junghans, H. G., and Wagenbach, D.: Technique for continuous high-resolution analysis of trace substances in firn and ice cores, *Environmental Science & Technology*, 34, 338–342, <https://doi.org/10.1021/es9907055>, 2000.
- Sigg, A., Fuhrer, K., Anklin, M., Staffelbach, T., and Zurmühle, D.: A continuous analysis technique for trace species in ice cores, *Environmental Science & Technology*, 28, 204–209, <https://doi.org/10.1021/es00051a004>, 1994.
- 450 Sigl, M., Winstrup, M., McConnell, J. R., Welten, K. C., Plunkett, G., Ludlow, F., Büntgen, U., Caffee, M. W., Chellman, N. J., Dahl-Jensen, D., Fischer, H., Kipfstuhl, J., Kostick, C., Maselli, O. J., Mekhaldi, F., Mulvaney, R., Muscheler, R., Pasteris, D. R., Pilcher, J. R., Salzer, M., Schüpbach, S., Steffensen, J. P., Vinther, B. M., and Woodruff, T. E.: Timing and climate forcing of volcanic eruptions for the past 2,500 years, *Nature*, 523, 543–549, <https://doi.org/10.1038/nature14565>, 2015.

- 455 Sinnl, G., Winstrup, M., Erhardt, T., Cook, E., Jensen, C. M., Svensson, A., Vinther, B. M., Muscheler, R., and Rasmussen, S. O.: A multi-ice-core, annual-layer-counted Greenland ice-core chronology for the last 3800 years: GICC21, *Climate of the Past*, 18, 1125–1150, 2022.
- Stoll, N., Eichler, J., Hörhold, M., Erhardt, T., Jensen, C., and Weikusat, I.: Microstructure, micro-inclusions, and mineralogy along the EGRIP ice core – Part 1: Localisation of inclusions and deformation patterns, *The Cryosphere*, 15, 5717–5737, <https://doi.org/10.5194/tc-15-5717-2021>, 2021.
- 460 Stoll, N., Hörhold, M., Erhardt, T., Eichler, J., Jensen, C., and Weikusat, I.: Microstructure, micro-inclusions, and mineralogy along the EGRIP (East Greenland Ice Core Project) ice core – Part 2: Implications for palaeo-mineralogy, *The Cryosphere*, 16, 667–688, <https://doi.org/10.5194/tc-16-667-2022>, 2022.
- Tsien, R., Pozzan, T., and Rink, T.: Calcium homeostasis in intact lymphocytes: cytoplasmic free calcium monitored with a new, intracellularly trapped fluorescent indicator., *The Journal of cell biology*, 94, 325–334, <https://doi.org/10.1083/jcb.94.2.325>, 1982.
- Vallelonga, P., Christianson, K., Alley, R. B., Anandakrishnan, S., Christian, J. E. M., Dahl-Jensen, D., Gkinis, V., Holme, C., Jacobel, R. W., Karlsson, N. B., Keisling, B. A., Kipfstuhl, J., Kjær, H. A., Kristensen, M. E. L., Muto, A., Peters, L. E., Popp, T., Riverman, K. L., Svensson, A., Tibuleac, C., Vinther, B. M., Weng, Y., and Winstrup, M.: Initial results from geophysical surveys and shallow coring of the Northeast Greenland Ice Stream (NEGIS), *The Cryosphere*, 8, 1275–1287, <https://doi.org/10.5194/tc-8-1275-2014>, 2014.
- 470 Winski, D. A., Fudge, T. J., Ferris, D. G., Osterberg, E. C., Fegyveresi, J. M., Cole-Dai, J., Thundercloud, Z., Cox, T. S., Kreutz, K. J., and Ortman, N.: The SP19 chronology for the South Pole Ice Core–Part 1: volcanic matching and annual layer counting, *Climate of the Past*, 15, 1793–1808, <https://doi.org/10.5194/cp-15-1793-2019>, 2019.




Figure 5. Depth assignment statistics. Panels A-D show information on the amount of ice melted during each run (A) and the number of pieces of ice (B), their average length and the amount of ice that is missing as indicators for the sample quality. Panels E-G show the melt speed (E) alongside the target meltspeed of 2.7 cm min^{-1} in ice equivalent, the interrune variability of the melt speed (F) as an indicator for the meltspeed stability and the encoder deviation (G) as an indicator of the depth-assignment precision. In panel F, the horizontal lines mark 5 and 10 % variability, in Panel G 0 and ± 1 % deviation.




Figure 6. Cross plot of ICP-TOFMS Ca and fluorimetric Ca^{2+} concentrations at 10 cm resolution, considering only sections with less than 50 % missing data. Regression, Pearson correlation (r), root mean square error (rms) and average ratio were calculated on the concentrations (i.e. not log-transformed). The solid line is the result of the regression, the dashed line marks the 1:1 line, the dotted lines indicate a $\pm 10\%$ deviation from that line.




Figure 7. Example of high-resolution data over a 10-m section of the core. The parameters shown are the same as in Figure 3, however, data is shown on the depth scale, with the secondary x-axis at the top indicating the age at annual resolution. The large peak in conductivity at 244.3 m is a typical signal of a volcanic eruption, and has been used to synchronize the age scales between different ice cores.