

A 30 m annual cropland dataset of China from 1986 to 2021

Ying Tu¹, Shengbiao Wu², Bin Chen^{2,3,4}, Qihao Weng⁵, Peng Gong^{3,6}, Yuqi Bai¹, Jun Yang¹, Le Yu¹, Bing Xu^{1,7,*}

¹Ministry of Education Ecological Field Station for East Asian Migratory Birds, Department of Earth System Science, Institute for Global Change Studies, Tsinghua University, Beijing 100084, China

²Future Urbanity & Sustainable Environment (FUSE) Lab, Division of Landscape Architecture, Faculty of Architecture, The University of Hong Kong, Hong Kong SAR, China

³Urban Systems Institute, The University of Hong Kong, Hong Kong SAR, China

⁴HKU Musketeers Foundation Institute of Data Science, The University of Hong Kong, Hong Kong SAR, China

⁵Department of Land Surveying and Geo-Informatics, The Hong Kong Polytechnic University, Kowloon, Hong Kong SAR, China

⁶Department of Geography, and Department of Earth Science, The University of Hong Kong, Hong Kong SAR, China

⁷International Research Center of Big Data for Sustainable Development Goals, Beijing, 100094, China

15 *Correspondence to:* Bing Xu (bingxu@tsinghua.edu.cn)

Abstract. Accurate, detailed, and up-to-date information on cropland extent is crucial for provisioning food security and environmental sustainability. However, because of the complexity of agricultural landscapes and lack of sufficient training samples, it remains challenging to monitor cropland dynamics at high spatial and temporal resolutions across large geographical extents, especially for places where agricultural land use is changing dramatically. Here we developed a cost-effective annual cropland mapping framework that integrated time-series Landsat imagery, automated training sample generation, and machine learning and change detection techniques. We implemented the proposed scheme to a cloud computing platform of Google Earth Engine and generated China's annual cropland dataset (CACD) at a 30 m spatial resolution for the first time. Results demonstrated that our approach was capable of tracking dynamic cropland changes in different agricultural zones. The pixel-wise F1 scores for annual maps and change maps of CACD were 0.79 ± 0.02 and 0.81, respectively. A further cross-product comparison in terms of accuracy assessment, correlations with statistics, and spatial details indicated the precision and robustness of CACD than other datasets. **According to our estimation, from 1986 to 2021, China's total cropland area expanded by 30,300 km² (1.79%), which underwent an increase before 2002 but a general decline between 2002-2015 and a slight recovery afterward.** Cropland expansion was concentrated in the northwest while the eastern coastal region experienced substantial cropland loss. In addition, we observed 419,342 km² (17.57%) of croplands that were abandoned at least once during the study period. The consistent, high-resolution data of CACD can support progress toward sustainable agricultural use and food production in various research applications. The full archive of CACD is freely available at <https://doi.org/10.5281/zenodo.7936885> (Tu et al., 2023a).

Keywords. Cropland change detection; Landsat; Training sample; Class probability; LandTrendr; Google Earth Engine; China

1 Introduction

35 The global population has boosted during the past century: from less than 2 billion in 1920 to 7.8 billion in 2020 and is expected to reach 9.7 billion by 2050 (Roberts, 2011). Feeding this growing population, as well as coping with increasing food demand from shifting diets, has become a major challenge for global food security and environmental sustainability (Godfray et al., 2010; Searchinger et al., 2008; Tilman et al., 2011). The United Nations has specified the need for realizing “no poverty and zero hunger” and “balancing increasing agricultural production with the maintenance of ecosystem services” in its sustainable
40 development goals (SDGs) (United Nations, 2015). Knowledge of where farmlands locate and how they change over time not only facilitates people’s understanding of crop area and yield, but also supports early warnings and adaptation initiatives of the agriculture system (See et al., 2015). In this context, detailed, timely, and accurate mapping of cropland extent appears as a critical prerequisite for food market stability, land and water resources management, and the assessment of environmental impacts of agriculture (Waldner et al., 2015b).

45 As the most populous country on the planet, China has a long history of farming dating back more than 8000 years (Bryan et al., 2018). Today, China feeds about 20% of the world’s population with only 7% of the total farmland (Cui and Shoemaker, 2018). This has inevitably placed huge pressure on the country’s food security and agricultural sustainability. During the past few decades, extensive croplands were lost in China because of rapid urbanization (Liu et al., 2019; Tu et al., 2021; Tu et al., 2023b). To compensate for the loss, substantial cropland expansion and intensification took place in the meantime (Zuo et al.,
50 2018), which seriously threatens ecosystem functioning and biodiversity (Tang et al., 2021; Zabel et al., 2019). It is therefore of great importance to track spatial-temporal dynamics of croplands in China to propose more adaptive policies and strategies in the face of dramatically changing agricultural land use.

Earth observations provide a reliable and cost-effective way of long-term, large-scale, and up-to-date cropland information gathering. The Moderate Resolution Imaging Spectroradiometer (MODIS) imagery, for example, provides a unique capability
55 to map cropland extent at 250-500 m spatial resolutions (Gumma et al., 2014; Wardlow and Egbert, 2008; Xiong et al., 2017; Zhang et al., 2022a; Zhang et al., 2015). However, such kinds of data may fail to detect small field patches (<2.56 ha) (Fritz et al., 2015), especially for heterogeneous agricultural landscapes. In recent years, higher-resolution data like Landsat and Sentinel-2 have been well used in global cropland mapping initiatives at tens-of-meter levels, including Finer Resolution Observation and Monitoring-Global Cropland (FROM-GC) (Yu et al., 2013), Global Food Security-Support Analysis Data
60 (GFSAD30) (Thenkabail et al., 2021), and Global Land Analysis & Discovery Cropland Data (GLAD) (Potapov et al., 2022). These are often one-phase or multi-year cropland data products. In some developed countries and regions, more detailed cropland maps, such as the United States National Agricultural Statistics Service Cropland Data Layer (NASS-CDL) (Boryan et al., 2011) and the European Union 10 m crop type map (D’andrimont et al., 2021), have been produced. To date, no fine-resolution annual cropland dataset of China exists yet.

65 Monitoring cropland dynamics is demanding given the high inter- and intra-annual spectral variabilities of cropland (Prishchepov et al., 2012; Yin et al., 2014). From the perspective of mapping schemes, current approaches undergo a shift

from multi-year cropland classification with traditional machine learning to annual cropland mapping that combines change detection techniques. In the former strategy, cropland is usually considered as a general land use category, of which classifications are performed individually for each year using *in situ* data and supervised classifiers such as decision tree
70 (Boryan et al., 2011; Pittman et al., 2010; Potapov et al., 2022), random forest (Li et al., 2022; Yang and Huang, 2021; Yin et al., 2020), and support vector machine (Lambert et al., 2016; Shao and Lunetta, 2012; Traoré et al., 2019). However, the accuracy of classification maps is highly dependent on data quality, and errors present in each of the maps are compounded in the final change map, which may impair its ability in capturing real cropland use transitions (Zhu, 2017). To overcome this issue, post-classification processing methods, such as consistency checking (Li et al., 2015) or spatial-temporal filtering (Liu and Cai, 2012), have been employed to increase mapping accuracy. Instead, the latter strategy leverages all time-series information of remote sensing imagery into a change detection model and has become increasingly popular. Normally, temporal features of spectral bands or indices are used to fit temporal trajectories to identify both abrupt changes and continuous trends (Zhu, 2017). Representative change detection algorithms include Landsat-based detection of Trends in Disturbance and Recovery (LandTrendr) (Kennedy et al., 2010), Breaks for Additive Season and Trend (BFAST) (Verbesselt et al., 2010), and Continuous Change Detection and Classification (CCDC) (Zhu and Woodcock, 2014), of which trials have been given to characterize cropland patterns (Schneibel et al., 2017; Zhu et al., 2019). Nevertheless, since most of these algorithms are not originally developed for cropland change detection, the result can be highly variable and is often mixed with other vegetation types such as forest and grassland (Pasquarella et al., 2022).

One possible solution to deal with the shortcomings mentioned above is to take the advantages of both strategies, that is, to
85 first perform supervised classifications to generate multi-epoch land cover maps or cropland probabilities and then apply this intermediate information as inputs for change detections. A few efforts have been made in this manner so far, but challenges remain, including (1) limited exploration of the time-series sequence. Change detection analyses are simply used as a final step for updating the year of change (Dara et al., 2018; Xu et al., 2020; Xu et al., 2018), while the rich information stored in fitted trajectories of temporal segmentation is not well-incorporated in cropland distinguishing. (2) Unreasonable settings of land
90 use conversion rules. Xu et al. (2020), for instance, assumed there was a low probability of frequent land use changes and therefore only detected one single change during each 5-year interval. However, this is often not the case with croplands. (3) Most studies are conducted for specific type of cropland use change (e.g., cropland abandonment) and tested in small areas (Dara et al., 2018; Yin et al., 2018; Xu et al., 2021). It remains unclear whether the mapping scheme can be applied to monitor large-scale cropland dynamics.

95 Training samples are another essential issue in effective cropland classification. Existing studies rely extensively on *in situ* data or human interpretation of spectral signatures, making the classification process resource-intensive, time-consuming, and difficult to repeat over space and time (Zhong et al., 2014). To cope with the constrain, recent initiatives have been made to extract labeling or training data from existing land cover information (so called “baseline maps”) (Huang et al., 2020; Radoux et al., 2014; Waldner et al., 2015a; Xie et al., 2019; Zhang and Roy, 2017). Zhang and Roy (2017) derived training samples of
100 classification from previous 500 m MODIS land cover products. Huang et al. (2020) proposed an automatic training sample

migration method by measuring the spectral similarity and spectral distance between the reference spectra and image spectra. These achievements pave a new way for mapping annual croplands cost-effectively.

The aim of this study is to propose an incorporated paradigm for large-scale fine-resolution cropland dynamics monitoring. We first generated training samples automatically based on prerequisite knowledge, existing land cover baseline maps, and a 105 time-weighted dynamic time warping method. We then input multi-year phenological features derived from time-series Landsat imagery into a random forest classifier to obtain per-pixel annual cropland probabilities. By further applying the LandTrendr change detection algorithm, we classified the cropland probability time series into several segments, in which breakpoints and corresponding years of changes were recorded. Here we inspected cropland changes as a continuous dynamic process that consisted of several trajectories including no change, cropland gain, or cropland loss. We established a set of rules 110 to discriminate annual cropland types from the LandTrendr segmentation results. Based on the proposed framework, we produced the first annual cropland dataset of China (CACD) for 1986-2021 at a 30 m spatial resolution. We examined the accuracy of CACD using independent validation sample sets and compared the results with existing cropland maps. Finally, we assessed spatial and temporal changes of croplands in China since 1980s with CACD.

2 Methodology

115 As the framework delivered in Fig.1, we have proposed a trajectory-based approach that combines machine learning and change detection techniques for mapping annual cropland dynamics. Annual cropland in this study is defined as a piece of land of 0.25 ha in minimum (minimum width of 30 m) that is sowed/planted and harvestable at least once within the 12 months after the sowing or planting date. This definition aligns with the criteria established by the Joint Experiment of Crop Assessment and Monitoring (JECAM) network (Defourny et al., 2014) and adopts a shared scope of cropland that meets FAO's 120 Land Cover Meta Language (Di Gregorio, 2005). One crucial criterion for discerning annual cropland in this study is that its vegetation signals in remote sensing imagery must demonstrate noticeable variations over a 12-month period, reflecting the planting and harvesting activities (Fig. S1a). Consequently, certain exceptions are excluded in the definition of annual cropland: (1) Perennial crops like sugarcane and cassava, which have longer vegetation cycles and are not planted annually (Fig. S1b-c). However, if they are planted and harvested within a 12-month timeframe, we would consider them as croplands for that specific 125 year. (2) Fruit, tea, and coffee plantations, as their vegetation signals more closely resemble those of trees (Fig. S1d-f). (3) Greenhouse crops, as they exhibit distinct remote sensing characteristics compared to other cropland types (Fig. S1g). (4) Small plots such as legumes that do not meet the minimum size criteria of cropland.

China, a traditional agricultural nation in east Asia, is chosen as the study area for testing and verifying the proposed approach under large scales and dynamic climates. The country shares a total area of 9,600,000 km², including mainland China, Hong 130 Kong, Macau, and Taiwan. According to the national comprehensive agricultural zoning promulgated by the Chinese government, our study area consists of nine agricultural zones in general (Fig. S2). Their geographical and climatic conditions, growing seasons, cropping patterns, and major crops are provided in Table S1. Considering local differences and computation

capacity, we further divide the study area into around 1700 subregions with a size of $0.8^\circ \times 0.8^\circ$ (Fig. S2). Within each subregion, we perform annual cropland classifications individually on Google Earth Engine (GEE, <https://earthengine.google.com/>) based on the proposed methods in Fig. 1, of which details are described in the following Sections 2.1-2.7.

2.1 Training data generation

Training samples emerged as a key component in supervised classification. Here we developed an effective approach for automated training sample generation using existing land cover maps and the time-weighted dynamic time warping (TWDTW) method. First, we built a Cropland Inspector Tool on GEE that depicted time-series NDVI profiles and corresponding Landsat

140 imagery for a given point of interest. An illustration of the interpretation process of annual cropland types was given in the supplementary materials Fig. S3. Based on the tool, we manually interpreted 100 stable cropland points and 100 non-cropland (including forests, water ponds, and impervious areas) points within each agricultural zone as the referencing set. Second, we overlapped two annual land cover datasets in China, CLCD (Yang and Huang, 2021) and CLUD (Xu et al., 2020), to extract 145 cropland and non-cropland masks for each year respectively. For each subregion, we produced thousands of potential cropland and non-cropland points randomly and compared their monthly NDVI sequences with those of the referencing set using TWDTW, an algorithm that had been developed for distinguishing between different types of land use and land cover (Maus et al., 2016). TWDTW works by comparing the similarity between two temporal sequences and finds their optimal alignment, resulting in a dissimilarity measure (Belgiu and Csillik, 2018). Based on the discrimination results, we retained 20% invariant 150 samples that had the lowest dissimilarity value compared to the referencing set. The threshold value was set following recommendations by Ghorbanian et al. (2020). To determine how many training samples were least required for a robust classification, we conducted multiple experiments with different training sizes. As shown in Fig. S4, the F1 score of cropland classification increased as the number of training samples grew in the initial stage but became rather stable after reaching 800. Consequently, we generated ~800 training samples within each subregion.

155 In practice, we employed the widely adopted areal-proportional sampling strategy (Huang et al., 2002; Jin et al., 2014) for allocating both cropland and non-cropland samples. However, one limitation of this approach is that it can result in extremely small sample sizes for rare land-cover types in homogeneous landscapes. When these types are underrepresented in the samples, it may lead to subpar classification performance. To address this concern, Zhu et al. (2016) recommended a minimum sample size of 600 for rare land-cover types. A recent study also applied this parameter for sample balancing in global annual land 160 cover mapping within each $5^\circ \times 5^\circ$ geographical tile (Zhang et al., 2023). Given that the subregion size in our research ($0.8^\circ \times 0.8^\circ$) is significantly smaller than that in Zhang's study ($5^\circ \times 5^\circ$), we established the minimum sample size as 100. This means that we initially generated 800 samples for each $0.8^\circ \times 0.8^\circ$ subregion, encompassing both cropland and non-cropland categories. If either category fell below the specified threshold, we increased it to a minimum of 100 samples.

2.2 Feature space construction

We used all available Landsat TM/ETM+/OLI Tier 1 surface reflectance imagery from 1986 to 2021 covering our study area 165 as of March 2022. The data had been atmospherically corrected by USGS using the LEDAPS (Masek et al., 2006) and LaSRC

(Vermote et al., 2016) algorithms. We further masked out all the clouds, cloud shadows, and snow/ice contained in the Landsat

imagery using the FMASK (Zhu and Woodcock, 2012). Considering the differences between sensors, we applied the

coefficients suggested by Roy et al. (2016) to normalize OLI reflectance to that of TM and ETM+. This enabled multi-temporal

images to be consistent and comparable across years. We then calculated spectral indices of NDVI, NDSI, NDBI, NBR, and

170 EVI based on the formulas provided in Table S2. To improve classification performance, we finally carried out the tasseled

cap transformation to include Brightness, Greenness, and Wetness indices using the coefficients provided by Crist (1985).

Multi-temporal metrics have been widely used for capturing the spatial-temporal features of different land covers, such as forests, croplands, and urban areas (Potapov et al., 2012; Yin et al., 2020). Due to the nature of crop growth, spectral signals

175 of the croplands exhibit phenological patterns periodically. An example is shown in Fig. S5, where NDVI is high during the

growing season and low during sowing. Accordingly, based on the processed Landsat data, we extracted a series of multi-

temporal metrics each year as input features for annual cropland classification. These included the 10th, 25th, 50th, 75th, and 90th

percent quantiles of both growing and non-growing seasons for the spectral bands and indices listed in Table S2. In addition,

we augmented elevation information into feature space using the Shuttle Radar Topography Mission (SRTM) digital elevation

dataset (Jarvis et al., 2008). This resulted in a total of 141 input features for each year.

180 2.3 Cropland probability estimation

We estimated pixel-wise cropland probabilities for each year between 1986-2021 using the random forest algorithm. Random 185 forest is a bagging ensemble learning approach that constructs a multitude of decision trees at the training time (Ho, 1995).

The final class (classification) or prediction (regression) is assigned based on the majority “vote” of all trees (Breiman, 2001).

The algorithm has been widely used in land cover and land use classifications, given its accuracy and efficiency in dealing

190 with high-dimensional data (Gong et al., 2019; Tu et al., 2020). In this study, the parameter for the number of trees

(*numberOfTrees*) was set to 500, the number of variables per split (*variablesPerSplit*) was set to the square root of *numberOfTrees*, and other parameters were set as default as those on GEE.

2.4 Temporal segmentation

We employed the trajectory-based spectral-temporal segmentation algorithm, LandTrendr, to decompose the pixel-wise annual 195 cropland probability time series into different cropland use segments. The overarching goal of LandTrendr is to characterize a

temporal trajectory of data values using a sequence of connected linear segments bounded by breakpoints known as “vertex”

(Kennedy et al., 2018). By fitting a series of linear segments, LandTrendr reduces inter-annual signal noise while capturing

both long-term gradual and short-term drastic changes (Kennedy et al., 2010). The fitting result of LandTrendr is a series of

195 segments composed of breakpoints that separate periods of durable change or stability in spectral trajectory (Fig. 2). In this way, the magnitude of changes as well as years when changes occur can be recorded. LandTrendr was originally developed for detecting forest disturbances, but recent applications also demonstrated its potential in monitoring cropland dynamics (Dara et al., 2018; Yin et al., 2018).

196 The implementation of LandTrendr requires a series of control parameters and filtering steps designed to reduce overfitting while still capturing the desired features of trajectories. Nevertheless, when leveraging LandTrendr to monitor cropland
200 dynamics, we found few experiences with parameter settings in the literature. In addition, differences in climatic, geographic, and cropping conditions across regions would also affect algorithm performance. Accounting for these issues, we defined one test region for each agricultural zone with a size of 100 km×100 km (Fig. S2). Within each test region, we interpreted 100 cropland and non-cropland samples likewise, and compared classification results under different LandTrendr arguments using the developed approach (Table S3). Practically, we assigned LandTrendr parameter settings with the highest classification
205 accuracy of F1 score as the input for annual cropland mapping in the corresponding agricultural zone (Table S4). Details are given in the supplementary materials.

2.5 Annual cropland mapping

210 We have established a set of rules that automatically discriminate cropland transition types based on the LandTrendr segmentation results of annual cropland probability time series. Suppose each segment represents a specific cropland use status or change, the key is to determine its transition type characterized by the fitting breakpoints in the beginning and end, and then to reverse cropland types for each year within the segment. Take the pixel in Fig. 2 as an example, LandTrendr has divided its time series of cropland probabilities into three end-to-end segments (s) and four breakpoints (p), where t and v represent the year in which cropland use changes occur and the fitted cropland probability. First, land use conditions for all breakpoints $p_{(t,v)}$ are identified based on the fitted value: cropland if $v > 0.5$ and non-cropland if vice versa. Since each segment s_i is delineated by a starting breakpoint p_{is} and an ending breakpoint p_{ie} , this results in four scenarios for the transition type of s_i (Table 1). If both p_{is} and p_{ie} are cropland (non-cropland), land use types of the pixel during $t_{is}-t_{ie}$ remain unchanged and should always be cropland (non-cropland). If $v_{is}>0.5$ and $v_{ie}\leq0.5$, it means cropland loss takes place during the period. In this case, the pixel will be recognized as cropland in t_{is} and non-cropland from t_{is+1} to t_{ie} . Conversely, under the scenario where p_{is} is non-cropland and p_{ie} is cropland, land use types of the pixel are assigned as cropland in all years of s_i except for t_{is} (which is non-cropland). **More**
215 **examples on discerning dynamic cropland changes with our proposed annual cropland framework can be found in**
220 **supplementary materials Figs. S6-9.**

2.6 Post-classification processing

225 In post-classification processing, we first applied a 2-D Gaussian low pass filter to the classified cropland maps. The Gaussian filter is one of the most widely used and effective window-based filtering methods for reducing noise and blurring regions of an image (Canny, 1986). Weights for each pixel in the filtering window are determined by a Gaussian function:

$$G_{x,y} = \frac{1}{2\pi\sigma^2} e^{-\frac{(x^2+y^2)}{2\sigma^2}}, \quad (1)$$

where x and y are the coordinates of the pixels in the filtering window. The coordinate of the center pixel of the window is (0,0). σ is the standard deviation of the Gaussian filter that controls the degree of blurring spectral information. The filtering window is also controlled by a kernel, which determines the neighborhood extent that can be computed on the central pixel

230 (usually odd numbers). In this study, a 3×3 Gaussian filter was performed using the OpenCV package in Python.

Furthermore, we implemented a spatial-temporal consistency check approach, modified based on the method proposed by Li et al. (2015), to refine the annual cropland maps. Specifically, for each pixel i in year t , we calculated its spatial-temporal consistency probability $Prob_{i,t}$ within the surrounding $3\times 3\times 3$ window:

$$Prob_{i,t} = \frac{1}{N} \sum_{t'=t-1}^{t+1} \sum_{x=m-1}^{m+1} \sum_{y=n-1}^{n+1} Con(L_{i,t} = L_j), \quad (2)$$

235 where $L_{i,t}$ denotes the label of the target pixel i in year t , and L_j denotes the label of pixels in the neighborhood window. N signifies the total number of pixels (i.e., $N=27$), and x and y indicate the coordinates of pixel i within the window. The crux of this approach lies in the consistency check function $Con()$, which equals 1 if $L_{i,t} = L_j$, and 0 otherwise. Here we followed the threshold of 0.5, as suggested by Li et al. (2015), to discern the label transition between cropland (1) and non-cropland (0): if $Prob_{i,t}$ is less than 0.5, then the label of pixel i in year t is altered to the opposite category.

240 Lastly, we used the JRC global surface water (Pekel et al., 2016) and global artificial impervious area (GAIA) (Gong et al., 2020) datasets to mask out permanent water and impervious areas in each corresponding year respectively.

2.7 Accuracy assessment and inter-comparison

Map confidence is a key reflection of classification performance and data quality. In this study, three independent sample sets were used to evaluate the accuracy of CACD comprehensively. The first one was the annual validation sample set generated

245 by the research team. Specifically, we divided the study area into nearly one thousand hexagons and randomly created five points within each hexagon, following the work by Gong et al. (2013) on sample design. For each random point, land cover status and changes were interpreted manually. This was realized based on the Cropland Inspector Tool and historical high-resolution images from Google Earth. Six well-trained researchers were involved in the interpretation task. Each validation point would be identified by at least two researchers, of which yearly land cover types between 1986-2021 (cropland or non-
250 cropland) as well as confidence level (low or high) were filled in the result sheet. If there was a disagreement between the interpretation results, a final judgment would be made through discussions among the research team. We discarded points with low confidence and retrieved a total of 4972 validation samples by the end (Fig. 3).

Moreover, we leveraged two third-party sample sets (namely GLCVSS and GeoWiki) for the accuracy assessment of CACD.

The global land cover validation sample set (GLCVSS) was the first all-season validation sample set generated for global land
255 cover mapping (Li et al., 2017). It contained ~35000 validation samples interpreted on Landsat 8 with records of the date of image acquisition (interpreted from 2013 to 2015), spectral reflectance for each season, and level of interpretation uncertainty and sample sizes (3×3 , 9×9 , 17×17 , 33×33 , 1 unit =30 m). The classification system of GLCVSS was the same as FROM-

GLC. We selected all GLCVSS points in China with high confidence and a sample size of 30 m×30 m (Fig. S10). The other sample set Geo-Wiki was a global cropland reference dataset collected through a crowdsourcing campaign, which enveloped 260 80 participants from around the world to review almost 36,000 sample units during September 2016 (Laso Bayas et al., 2017). Each record in the Geo-Wiki dataset represents a cropland site identified by different participants. Likewise, we extracted all the cropland points located in China from Geo-Wiki (Fig. S11). In summary, there were 2018 GLCVSS and 1674 Geo-Wiki samples included in the research.

A group of accuracy metrics was calculated based on the confusion matrix, including F1 score, overall accuracy (OA), 265 producer's accuracy (PA), user's accuracy (UA), and Kappa coefficient. The F1 score was used as a major indicator of accuracy as it conveyed the balance between UA and PA, which reached its best value at 1 and worst value at 0 (Chen et al., 2021). Accuracy metrics of both yearly maps and change maps of CACD were calculated based on the annual validation sample set while GLCVSS was used for assessing the accuracy of CACD in 2015. In terms of Geo-Wiki, because it only contained 270 cropland samples, we calculated the agreement level between Geo-Wiki and CACD in 2016, that is, how many points of GeoWiki were classified as cropland in CACD.

In addition, we compared CACD with other publicly available land cover or cropland datasets (i.e., CLCD, CLUD, GLAD, and GFSAD). Both CLCD (Yang and Huang, 2021) and CLUD (Xu et al., 2020) were 30 m annual land cover datasets in 275 China with a long time span. For each year in CLCD and CLUD, we extracted cropland extent (1) and grouped other land cover types as 'non-cropland' (0). GLAD represented a globally consistent cropland extent time series at 30 m spatial resolution and was mapped in four-year intervals (2000-2003, 2004-2007, 2008-2011, 2012-2015, and 2016-2019) (Potapov et al., 2022). GFSAD was a 30 m global cropland dataset for 2015 (Thenkabail et al., 2021). We calculated accuracies of CLCD, CLUD, 280 GLAD, and GFSAD using the three validation sample sets. Besides, we estimated the total area of croplands at the provincial scale and compared it with statistics.

3 Results

280 3.1 Accuracy assessment of CACD

We performed pixel-wise accuracy assessment of CACD based on the annual validation sample set. The F1 score, OA, UA, PA, and Kappa coefficient of annual maps on average were 0.79 ± 0.02 , 0.93 ± 0.01 , 0.79 ± 0.01 , 0.79 ± 0.02 , and 0.75 ± 0.02 , respectively. Temporally, mapping accuracy after 2000 was generally higher than that before (Table S5), probably due to the improvement of data quantity and quality of Landsat images in the later period. Besides, we estimated mapping accuracy of 285 CACD at hexagon and province levels using the annual validation samples set. Spatially, except for parts of the western and southeastern coastal areas, the mean F1 score for most of the study area was above 0.8, with omission and commission errors less than 0.25 (Fig. 4a-c). When it comes to provincial performance, the mean F1 score for Northeast Plain, North China Plain, Jiangsu, Zhejiang, Jiangxi, and Yunnan reached up to 0.8, whereas a lower value was found in inland provinces such as Inner Mongolia and Xinjiang (Fig. 4d). The relatively high omission error in Qinghai and Fujian (Fig. 4e) demonstrated an

290 underestimation of cropland in these regions. In contrast, Guangdong had the highest commission error on average (Fig. 4f), indicating that croplands were somewhat overestimated in the province.

In terms of change accuracy, we reclassified the annual validation sample set as changed if any cropland use conversion was identified between 1986-2021 and vice versa. Since the number of non-changed samples (98%) was substantially greater than that of changed samples (2%), we randomly subset 100 non-changed samples and combined them with changed samples for 295 further evaluation. Results demonstrated that the F1 score, OA, and Kappa coefficient of CACD change maps could be as high as 0.81, 0.84, and 0.68 (Table 2). Accuracy for the year of change ranged between 0.76-0.87 as the tolerance years increased from 0 to ± 5 years (Table S6).

Furthermore, using two third-part sample sets, the F1 score of CACD in 2015 was verified as 0.82 based on the GLCVSS sample set (Table 3), while 86% of the sample points in Geo-Wiki were correctly classified as cropland in CACD (Table 4).

300 Evaluation results of third-party sample sets complemented the accuracy of CACD.

3.2 Comparison with other products

We compared CACD with four existing land cover (cropland) datasets (CLCD, CLUD, GLAD, and GFSAD) at 30 spatial resolutions from multiple aspects. CACD was superior to other products in all comparison years in terms of mapping accuracy (Fig. 5). In 2015, for example, the F1 score of CACD was 0.85, followed by GFSAD (0.77), GLAD (0.76), CLCD (0.76), and 305 CLUD (0.69). Similar results were observed in the accuracy assessment results of GLCVSS and Geo-Wiki sample sets (Tables 3-4). Spatially, the five products were consistent in delineating cropland distributions in major production areas including the Northeast Plain, North China Plain, Sichuan Basin, and Xinjiang (Fig. 6). However, differences occurred in Loess Plateau, Yunan-Guinan Plateau, and the vast hilly regions along the southeast coast. Overall, according to the frequency that how often pixels were identified as cropland, 65.62% of the study area were labeled as croplands at least three times out of the five 310 products (Fig. 6, matched level ≥ 3).

Fig. 7 compares spatial details of cropland distribution of the five products in six selected regions. In the Northeast Plain (region A), CACD, GLAD, and GFSAD depicted the extent of cropland well, while CLCD slightly overestimated and CLUD severely underestimated croplands. In Xinjiang and North China Plain (region B and C), high spatial consistency was observed between the five products, with CACD and GLAD displaying more details (like the road network). Classification differences 315 mainly appeared in the southern hilly areas (region D, E, and F). In general, CACD, CLCD, and CLUD were more consistent and closer to the actual cropland distribution. For instance, region D was a terracing land in Sichuan Province, southwestern China. The region was mainly used for rice cultivation, yet those croplands were massively underestimated in GLAD and GFSAD. **Further illustrations on the multi-year cropland distributions among different products can be found in the supplementary materials Figs. S12-16.**

320 Using the five products, we counted cropland area at the provincial scale separately and compared it with statistical data in 2015. As shown in Fig. 8, CACD showed the best agreement with statistics ($R^2=0.93$), followed by CLUD ($R^2=0.91$), CLCD ($R^2=0.89$), GFSAD ($R^2=0.87$), and GLAD ($R^2=0.74$). The scattered points between CLCD and statistics were basically above

the 1:1 diagonal, indicating an overestimation of cropland area. On the contrary, GLAD had an underestimated cropland area as more scattered points were located below the 1:1 diagonal. This was mainly caused by GLAD's misclassification in southern China. Taking together, the comparison results between different products demonstrated the accuracy of our annual cropland datasets.

4 Discussion

4.1 Cropland dynamics in China

Based on the produced CACD, we calculated provincial net changes in cropland area between 1986-2021 and plotted annual cropland area dynamics at the national scale. We also employed the annual validation samples to an error-adjusted stratified estimation approach following Olofsson et al. (2014) to estimate sample-based cropland area and its 95% confidence interval. As shown in Fig. 9a, cropland gain mainly occurred in northern China, including Xinjiang, Inner Mongolia, and Heilongjiang. In Xinjiang, for example, 44635 km² of the land area (~3%) was developed as cropland during the past 36 years, a figure that was equivalent to 1.3 times the area of Hainan Province. On the contrary, most provinces in eastern, central, and southern China experienced massive cropland loss. The top five provinces with the largest area of cropland loss were Sichuan (12739 km²), Jiangsu (10865 km²), Shaanxi (10700 km²), Hebei (8566 km²), and Shandong (8407 km²). In the capital city of Beijing, more than 2000 km² croplands were occupied during the study period, which accounted for 12% of the city's total area. Additionally, cropland areas in some inland provinces (such as Guizhou) remained rather stable. Overall, according to sample-based estimation, China's cropland area increased from 1,694,900 ± 193,700 km² in 1986 to 1,725,200 ± 212,400 km² in 2021, a net increase of 30,300 km² (1.79%). Temporally, the country's cropland area increased before 2002 but gradually decreased until 2015, after which a recovery phase began (Fig. 9b). These findings are generally consistent with those reported in previous empirical studies (Liu et al., 2014a; Gao et al., 2019).

Rapid urbanization process in China since the 1980s has caused significant losses of cropland, particularly in large cities and the eastern coastal region (Qiu et al., 2020; Tu et al., 2021; Tu et al., 2023b). In response to the increasing demand for food, the Chinese government has implemented a series of farmland protection policies, including the Requisition–Compensation Balance of Farmland (RCBF) policy, which mandates that any lost cropland must be offset by the expansion of cropland elsewhere (Liang et al., 2015; Liu et al., 2014b). Despite these efforts, croplands in China have undergone dramatic and unbalanced changes over the past few decades. While the net change in cropland area has been relatively small, the spatial-temporal variance is noteworthy (Gao et al., 2019; Liu et al., 2014a; Yu et al., 2018). Fig. 10 showcases four typical types of cropland dynamic change in China with CACD. In the Ar Horqin Banner of Chifeng city, Inner Mongolia, large-scale croplands were developed for the reclamation and cultivation of crops during the past decades (Fig. 10a). Similarly, as shown in Fig. 10b, vast agricultural land parcels sprang up in Aksu, Xinjiang for cotton cultivation (Li et al., 2021; Liu, 2022). In Shanghai, China's most significant metropolitan, remarkable socioeconomic development since the reform and opening up in the 1980s had led to extensive urban sprawl, encroaching substantial areas of pre-existing croplands (Fig. 10c). In the

355 mountainous area of Tongren, Guizhou, some croplands were abandoned in recent years as a result of urban-rural migration and declining marginal benefits of arable land (Fig. 10d). In summary, these specific examples demonstrated that CACD was capable of capturing fine-resolution cropland dynamics across different landscapes.

360 Cropland abandonment is a widespread and rapidly increasing phenomenon worldwide (Queiroz et al., 2014), as a result of land marginalization caused by unsatisfactory land suitability and reduced economic viability (Macdonald et al., 2000). It has wide-ranging effects on the environment and social economy, such as biodiversity conservation (Isbell et al., 2019), carbon storage (Vuichard et al., 2008), and land economics (Ito et al., 2016). However, knowledge of the extent and timing of cropland abandonment in China is lacking. According to FAO, cropland abandonment refers to land that has not been cultivated for at least five consecutive years (Fao, 2016). Following this definition, we identified a pixel of CACD as abandoned if it was classified as cropland in the initial year but changed to non-cropland for the following five years. Also, we were conservative and included only those croplands as abandoned that converted to natural cover, not those to built-up lands. To do so, we masked out cropland conversions to impervious areas based on the GAIA data. When abandonment occurred, we labeled its timing as the first year in which cropland was no longer active (starting from 1990 to 2015).

365 Our results revealed that 419,342 km² (17.57%) of the croplands in China were abandoned at least once during the past decades. Most abandoned croplands were distributed in the central and the west (from 100° to 125°E and from 20° to 50°N, Fig. 11a), 370 particularly in Inner Mongolia, Gansu, Shaanxi, and Yunan provinces (Fig. S17). This spatial pattern was consistent with Li et al. (2018), who estimated the extent of cropland abandonment in China's mountainous regions using multiple regression models. The average altitude of abandoned cropland was 860 m, 81% of which were in regions below 1500 m (Fig. 11b). Temporally, the annual cropland abandonment area in China showed an upward trend from 1990 to 2015, which increased from 7516 km² to 14823 km² (Fig. 11c). Rural-urban migration, reductions in agricultural labor forces, and rural population 375 ageing have been invoked as key drivers of cropland abandonment in China (Zhang et al., 2014; Yan et al., 2016; Ren et al., 2023).

4.2 Strengths and potential implications

380 This research introduces an integrated framework for monitoring annual cropland dynamics at the 30 m spatial resolution. Our methods offer several contributions to the existing literature. First, we leverage baseline land cover maps and the TWDTW discrimination algorithm to realize automated training sample generation. This eliminates the time-consuming and labor-costive process of traditional training sample collection, enabling cost-effective cropland mapping at large scales. Second, we 385 adopt the random forest classifier for annual cropland probabilities estimation and then integrate time series of these probabilities as spectral metrics into LandTrendr for cropland trajectory modeling. The incorporation of machine learning and change detection techniques not only increases accuracy but also improves the spatial-temporal consistency of classification results. Third, we establish a set of transition rules to convert the LandTrendr fitting results to pixel-wise annual cropland types, which can capture various cropland use changes such as abandonment or fallow. This cropland transition discrimination strategy is distinct from existing initiatives (Potapov et al., 2022; Dara et al., 2018; Xu et al., 2018; Yin et al., 2020). Our

results highlight the potential of change detection algorithms like LandTrendr to complement traditional classification processes in identifying dynamic land cover changes effectively.

390 In practice, we partitioned the study area into nine agricultural zones and performed localized annual cropland mapping within each $0.8^\circ \times 0.8^\circ$ subregion. We also evaluated the impacts of training sample size and LandTrendr parameter settings on classification accuracy. Localized classifications may greatly improve the accuracy in heterogeneous regions such as southern China. Our experiments provide valuable insights for future land cover and land use mapping endeavors. Moreover, we took advantage of the powerful data storage, computing, and analysis capabilities of GEE to build an end-to-end framework, which
395 enables fast annual cropland mapping in any given area of interest worldwide. Theoretically, the proposed framework is highly adaptable and can be extended to map other land use types.

Based on the aforementioned mapping scheme, we produced the first 30-m annual cropland maps of China for the period 1986-2021. The accuracy of the dataset was validated through three independent validation sample sets and multi-perspective comparisons with other products. The CACD dataset is a valuable contribution to the field of agriculture and land use
400 management in China. Its accuracy and high-resolution information provide insights into the changing dynamics of cropland use over time, helping inform policymakers and stakeholders in developing sustainable land use practices. Our subsequent analyses with CACD revealed a dramatic but heterogeneous change in cropland dynamics in China, with over a third of croplands experiencing at least one change of land use during the study period. Furthermore, we identified a total of 419,342 km² (17.57%) of abandoned croplands from 1990 to 2015, mostly located in the central and western mountainous areas. With
405 the ability to detect cropland abandonment patterns, targeted interventions can be implemented to bring abandoned land back into use, promoting food security and protecting arable land resources. Further efforts can also be directed towards scrutinizing the underlying driving forces of cropland abandonment as well as assessing its impacts on ecosystem service and biodiversity. Additionally, CACD can be utilized to conduct agriculture-related studies, such as analyzing crop yield and productivity, assessing the impact of climate change on crop growth, and monitoring land use changes over time. Overall, CACD represents
410 a powerful tool for promoting sustainable agriculture practices and ensuring the long-term availability of arable land resources in China.

4.3 Limitations and prospects

A few limitations regarding methodology and data are included in the study. First, we track cropland dynamics on an annual basis but do not account for intra-annual variations in crops, such as different crop calendars and cropping intensity.
415 Consequently, our results may not be effective in identifying perennial crops. Second, we depict the general extent of croplands but do not differentiate specific types. This becomes essential when evaluating yields or climate responses of different crops. Future work needs to explore advanced methods for fine-resolution crop types mapping. Third, despite the high accuracy, CACD is subject to several levels of uncertainty. Temporally, CACD has a relatively low accuracy before 2000, given the uneven coverage of Landsat 5 data. Spatially, mapping accuracy varied across different regions, where the western and
420 southeastern coastal areas have a comparatively high error compared to others. Accurately classifying cropland in these regions

has consistently been challenging, with recent research highlighting significant discrepancies between products (Lu et al., 2016; Zhang et al., 2022b; Xue et al., 2023). This challenge stems from characteristics of local topography and landscapes, including factors like elevation, slope, field size, and farmland fragmentation (Lu et al., 2016; Zhang et al., 2022b; Xue et al., 2023). Besides, the low quality of Landsat data (e.g., unmasked clouds), as well as the sensitivity of LandTrendr parameters, further 425 constitute the influence on mapping outcomes. Looking ahead, it is crucial to embark on comprehensive investigations and formulate targeted adaptation strategies to enhance the accuracy of cropland classification in these specific areas.

5 Data availability

The atlas of CACD generated in this study can be accessed at <https://doi.org/10.5281/zenodo.7936885> (Tu et al., 2023a). All maps are at the 30 m spatial resolution under the EPSG:4326 (WGS84) spatial reference system.

430 6 Conclusions

Timely and dependable cropland distribution maps appear as key variables for crop yield estimation, agricultural sustainability assessment, and climate modeling. This research put forward a cost-effective fine-resolution cropland dynamics monitoring scheme by synthesizing automated training sample generation, random forest supervised classification, and the LandTrendr temporal segmentation algorithm. We showed how labor on training sample collection can be substantially reduced by making 435 full use of existing land cover datasets. We also demonstrated how state-of-art machine learning and change detection approaches can be synthesized to characterize cropland dynamics. Leveraging the full archive of Landsat imagery and the GEE cloud computing platform, we mapped the first annual cropland distributions in China at a 30 m resolution from 1986 to 2021. The generated dataset reached a promising accuracy of F1 score of 0.79 ± 0.02 , which was superior to other products including CLCD, CLUD, GLAD, and GFSAD. Furthermore, validations of third-party sample sets, regressions with provincial statistics, 440 as well as comparisons of spatial details between multiple products, indicated that CACD was reasonable in delineating spatial distributions and temporal trends of cropland dynamics. The total cropland area of China in 2021 was $1,725,200 \pm 212,400$ km², which increased by 30,300 km² (1.79%) compared to that in 1986. Cropland expansion mainly took place in northwestern China while the eastern coastal region withstood substantial cropland loss due to rapid urbanization. The annual cropland abandonment area was 16,128 km² on average and showed an increasing trend from 1990 to 2015. The fine-resolution annual 445 cropland data of this study is expected to serve as the basis for a wide spectrum of applications and decision makings in the future, for example, facilitating crop condition monitoring and early warning, promoting progress towards sustainable food production, and assessing the environmental impacts of agricultural expansion and intensification.

Author contributions

Ying Tu and Bing Xu conceived the research idea. Ying Tu designed the study, created the dataset, carried out the analysis, 450 and wrote the first draft of the manuscript, under the supervision of other authors. All authors participated in the review and editing of the manuscript.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

455 The authors would like to express their great gratitude to the following organizations for their contributions to this research: USGS for providing free access to Landsat imagery and GFSAD data, Wuhan University for providing CLCD data, Tsinghua University for providing CLUD and GLCVSS data, the University of Maryland for providing GLAD data, and Geo-Wiki for sharing the third-party validation samples. We thank Prof. Jian Xu and students from Jiangxi Normal University for their efforts in sample collection. We also thank Prof. Jinwei Dong and Prof. Li Wang from the Chinese Academy of Sciences for 460 their valuable comments on the manuscript.

Financial support

This study was supported by the Open Research Program of the International Research Center of Big Data for Sustainable Development Goals (CBAS2022ORP02), National Key Research and Development Program of China (2022YFB3903703, 2022YFE0209300), Science and Technology Commission of Shanghai Municipality (22dz1209602)

465 References

Belgiu, M. and Csillik, O.: Sentinel-2 cropland mapping using pixel-based and object-based time-weighted dynamic time warping analysis, *Remote Sensing of Environment*, 204, 509-523, <https://doi.org/10.1016/j.rse.2017.10.005>, 2018.

Boryan, C., Yang, Z., Mueller, R., and Craig, M.: Monitoring US agriculture: the US Department of Agriculture, National Agricultural Statistics Service, Cropland Data Layer Program, *Geocarto International*, 26, 341-358, 10.1080/10106049.2011.562309, 2011.

Breiman, L.: Random forests, *Machine learning*, 45, 5-32, 2001.

Bryan, B. A., Gao, L., Ye, Y., Sun, X., Connor, J. D., Crossman, N. D., Stafford-Smith, M., Wu, J., He, C., Yu, D., Liu, Z., Li, A., Huang, Q., Ren, H., Deng, X., Zheng, H., Niu, J., Han, G., and Hou, X.: China's response to a national land-system sustainability emergency, *Nature*, 559, 193-204, 10.1038/s41586-018-0280-2, 2018.

Canny, J.: A Computational Approach to Edge Detection, *IEEE Transactions on Pattern Analysis and Machine Intelligence, PAMI-8*, 679-698, 10.1109/TPAMI.1986.4767851, 1986.

Chen, B., Tu, Y., Song, Y., Theobald, D. M., Zhang, T., Ren, Z., Li, X., Yang, J., Wang, J., Wang, X., Gong, P., Bai, Y., and Xu, B.: Mapping essential urban land use categories with open big data: Results for five metropolitan areas in the United States of America, *ISPRS Journal of Photogrammetry and Remote Sensing*, 178, 203-218, <https://doi.org/10.1016/j.isprsjprs.2021.06.010>, 2021.

480 Crist, E. P.: A TM Tasseled Cap equivalent transformation for reflectance factor data, *Remote Sensing of Environment*, 17, 301-306, [https://doi.org/10.1016/0034-4257\(85\)90102-6](https://doi.org/10.1016/0034-4257(85)90102-6), 1985.

Cui, K. and Shoemaker, S. P.: A look at food security in China, *npj Science of Food*, 2, 4, 10.1038/s41538-018-0012-x, 2018.

485 d'Andrimont, R., Verhegghen, A., Lemoine, G., Kempeneers, P., Meroni, M., and van der Velde, M.: From parcel to continental scale – A first European crop type map based on Sentinel-1 and LUCAS Copernicus in-situ observations, *Remote Sensing of Environment*, 266, 112708, <https://doi.org/10.1016/j.rse.2021.112708>, 2021.

Dara, A., Baumann, M., Kuemmerle, T., Pflugmacher, D., Rabe, A., Griffiths, P., Hözel, N., Kamp, J., Freitag, M., and Hostert, P.: Mapping the timing of cropland abandonment and recultivation in northern Kazakhstan using annual Landsat time series, *Remote Sensing of Environment*, 213, 49-60, <https://doi.org/10.1016/j.rse.2018.05.005>, 2018.

490 Defourny, P., Jarvis, I., and Blaes, X.: JECAM Guidelines for cropland and crop type definition and field data collection, *JECAM*, 2014.

Di Gregorio, A.: Land cover classification system: classification concepts and user manual: LCCS, Food & Agriculture Org. 2005.

FAO: FAOSTAT, Methods & Standards, 2016.

495 Fritz, S., See, L., McCallum, I., You, L., Bun, A., Moltchanova, E., Duerauer, M., Albrecht, F., Schill, C., Perger, C., Havlik, P., Mosnier, A., Thornton, P., Wood-Sichra, U., Herrero, M., Becker-Reshef, I., Justice, C., Hansen, M., Gong, P., Abdel Aziz, S., Cipriani, A., Cumani, R., Cecchi, G., Conchedda, G., Ferreira, S., Gomez, A., Haffani, M., Kayitakire, F., Malanding, J., Mueller, R., Newby, T., Nonguierma, A., Olusegun, A., Ortner, S., Rajak, D. R., Rocha, J., Schepaschenko, D., Schepaschenko, M., Terekhov, A., Tiangwa, A., Vancutsem, C., Vintrou, E., Wenbin, W., van der Velde, M., Dunwoody, A., Kraxner, F., and Obersteiner, M.: Mapping global cropland and field size, *Global Change Biology*, 21, 1980-1992, <https://doi.org/10.1111/gcb.12838>, 2015.

500 Gao, X., Cheng, W., Wang, N., Liu, Q., Ma, T., Chen, Y., and Zhou, C.: Spatio-temporal distribution and transformation of cropland in geomorphologic regions of China during 1990–2015, *Journal of Geographical Sciences*, 29, 180-196, 10.1007/s11442-019-1591-4, 2019.

505 Ghorbanian, A., Kakooei, M., Amani, M., Mahdavi, S., Mohammadzadeh, A., and Hasanolou, M.: Improved land cover map of Iran using Sentinel imagery within Google Earth Engine and a novel automatic workflow for land cover classification using migrated training samples, *ISPRS Journal of Photogrammetry and Remote Sensing*, 167, 276-288, <https://doi.org/10.1016/j.isprsjprs.2020.07.013>, 2020.

510 Godfray, H. C. J., Beddington, J. R., Crute, I. R., Haddad, L., Lawrence, D., Muir, J. F., Pretty, J., Robinson, S., Thomas, S. M., and Toulmin, C.: Food Security: The Challenge of Feeding 9 Billion People, *Science*, 327, 812-818, 10.1126/science.1185383, 2010.

Gong, P., Li, X., Wang, J., Bai, Y., Chen, B., Hu, T., Liu, X., Xu, B., Yang, J., Zhang, W., and Zhou, Y.: Annual maps of global artificial impervious area (GAIA) between 1985 and 2018, *Remote Sensing of Environment*, 236, 111510, <https://doi.org/10.1016/j.rse.2019.111510>, 2020.

515 Gong, P., Liu, H., Zhang, M., Li, C., Wang, J., Huang, H., Clinton, N., Ji, L., Li, W., Bai, Y., Chen, B., Xu, B., Zhu, Z., Yuan, C., Ping Suen, H., Guo, J., Xu, N., Li, W., Zhao, Y., Yang, J., Yu, C., Wang, X., Fu, H., Yu, L., Dronova, I., Hui, F., Cheng, X., Shi, X., Xiao, F., Liu, Q., and Song, L.: Stable classification with limited sample: transferring a 30-m resolution sample set collected in 2015 to mapping 10-m resolution global land cover in 2017, *Science Bulletin*, 64, 370-373, <https://doi.org/10.1016/j.scib.2019.03.002>, 2019.

520 Gong, P., Wang, J., Yu, L., Zhao, Y., Zhao, Y., Liang, L., Niu, Z., Huang, X., Fu, H., Liu, S., Li, C., Li, X., Fu, W., Liu, C., Xu, Y., Wang, X., Cheng, Q., Hu, L., Yao, W., Zhang, H., Zhu, P., Zhao, Z., Zhang, H., Zheng, Y., Ji, L., Zhang, Y., Chen, H., Yan, A., Guo, J., Yu, L., Wang, L., Liu, X., Shi, T., Zhu, M., Chen, Y., Yang, G., Tang, P., Xu, B., Giri, C., Clinton, N., Zhu, Z., Chen, J., and Chen, J.: Finer resolution observation and monitoring of global land cover: first mapping results with Landsat TM and ETM+ data, *International Journal of Remote Sensing*, 34, 2607-2654, 10.1080/01431161.2012.748992, 2013.

525

530 Gumma, M. K., Thenkabail, P. S., Maunahan, A., Islam, S., and Nelson, A.: Mapping seasonal rice cropland extent and area in the high cropping intensity environment of Bangladesh using MODIS 500m data for the year 2010, *ISPRS Journal of Photogrammetry and Remote Sensing*, 91, 98-113, <https://doi.org/10.1016/j.isprsjprs.2014.02.007>, 2014.

Ho, T. K.: Random decision forests, *Proceedings of 3rd international conference on document analysis and recognition*, 278-282,

Huang, C., Davis, L. S., and Townshend, J. R. G.: An assessment of support vector machines for land cover classification, *International Journal of Remote Sensing*, 23, 725-749, 10.1080/01431160110040323, 2002.

Huang, H., Wang, J., Liu, C., Liang, L., Li, C., and Gong, P.: The migration of training samples towards dynamic global land cover mapping, *ISPRS Journal of Photogrammetry and Remote Sensing*, 161, 27-36, <https://doi.org/10.1016/j.isprsjprs.2020.01.010>, 2020.

535 Isbell, F., Tilman, D., Reich, P. B., and Clark, A. T.: Deficits of biodiversity and productivity linger a century after agricultural abandonment, *Nature Ecology & Evolution*, 3, 1533-1538, 10.1038/s41559-019-1012-1, 2019.

Ito, J., Nishikori, M., Toyoshi, M., and Feuer, H. N.: The contribution of land exchange institutions and markets in countering farmland abandonment in Japan, *Land Use Policy*, 57, 582-593, <https://doi.org/10.1016/j.landusepol.2016.06.020>, 2016.

540 Jarvis, A., Reuter, H. I., Nelson, A., and Guevara, E.: Hole-filled SRTM for the globe Version 4, available from the CGIAR-CSI SRTM 90m Database ([http://srtm.cgiar.org](http://srtm.csi.cgiar.org)), 15, 5, 2008.

Jin, H., Stehman, S. V., and Mountrakis, G.: Assessing the impact of training sample selection on accuracy of an urban classification: a case study in Denver, Colorado, *International Journal of Remote Sensing*, 35, 2067-2081, 10.1080/01431161.2014.885152, 2014.

545 Kennedy, R. E., Yang, Z., and Cohen, W. B.: Detecting trends in forest disturbance and recovery using yearly Landsat time series: 1. LandTrendr — Temporal segmentation algorithms, *Remote Sensing of Environment*, 114, 2897-2910, <https://doi.org/10.1016/j.rse.2010.07.008>, 2010.

Kennedy, R. E., Yang, Z., Gorelick, N., Braaten, J., Cavalcante, L., Cohen, W. B., and Healey, S.: Implementation of the LandTrendr Algorithm on Google Earth Engine, 10.3390/rs10050691, 2018.

550 Lambert, M.-J., Waldner, F., and Defourny, P.: Cropland Mapping over Sahelian and Sudanian Agrosystems: A Knowledge-Based Approach Using PROBA-V Time Series at 100-m, 10.3390/rs8030232, 2016.

Laso Bayas, J. C., Lesiv, M., Waldner, F., Schucknecht, A., Duerauer, M., See, L., Fritz, S., Fraisl, D., Moorthy, I., McCallum, I., Perger, C., Danylo, O., Defourny, P., Gallego, J., Gilliams, S., Akhtar, I. u. H., Baishya, S. J., Baruah, M., Bungnamei, K., Campos, A., Changkakati, T., Cipriani, A., Das, K., Das, K., Das, I., Davis, K. F., Hazarika, P., Johnson, B. A., Malek, Z., Molinari, M. E., Panging, K., Pawe, C. K., Pérez-Hoyos, A., Sahariah, P. K., Sahariah, D., Saikia, A., Saikia, M., Schlesinger, P., Seidacaru, E., Singha, K., and Wilson, J. W.: A global reference database of crowdsourced cropland data collected using the Geo-Wiki platform, *Scientific Data*, 4, 170136, 10.1038/sdata.2017.136, 2017.

555 Li, C., Gong, P., Wang, J., Zhu, Z., Biging, G. S., Yuan, C., Hu, T., Zhang, H., Wang, Q., Li, X., Liu, X., Xu, Y., Guo, J., Liu, C., Hackman, K. O., Zhang, M., Cheng, Y., Yu, L., Yang, J., Huang, H., and Clinton, N.: The first all-season sample set for mapping global land cover with Landsat-8 data, *Science Bulletin*, 62, 508-515, <https://doi.org/10.1016/j.scib.2017.03.011>, 2017.

Li, Q., Liu, G., and Chen, W.: Toward a Simple and Generic Approach for Identifying Multi-Year Cotton Cropping Patterns Using Landsat and Sentinel-2 Time Series, *Remote Sensing*, 13, 10.3390/rs13245183, 2021.

560 Li, S., Li, X., Sun, L., Cao, G., Fischer, G., and Tramberend, S.: An estimation of the extent of cropland abandonment in mountainous regions of China, *Land Degradation & Development*, 29, 1327-1342, <https://doi.org/10.1002/ldr.2924>, 2018.

Li, X.-Y., Li, X., Fan, Z., Mi, L., Kandakji, T., Song, Z., Li, D., and Song, X.-P.: Civil war hinders crop production and threatens food security in Syria, *Nature Food*, 3, 38-46, 10.1038/s43016-021-00432-4, 2022.

565 Li, X., Gong, P., and Liang, L.: A 30-year (1984–2013) record of annual urban dynamics of Beijing City derived from Landsat data, *Remote Sensing of Environment*, 166, 78-90, <https://doi.org/10.1016/j.rse.2015.06.007>, 2015.

Liang, C., Penghui, J., wei, C., Manchun, L., Liyan, W., Yuan, G., Yuzhe, P., Nan, X., Yuewei, D., and Qiuhan, H.: Farmland protection policies and rapid urbanization in China: A case study for Changzhou City, *Land Use Policy*, 48, 552-566, <https://doi.org/10.1016/j.landusepol.2015.06.014>, 2015.

570 Liu, D. and Cai, S.: A Spatial-Temporal Modeling Approach to Reconstructing Land-Cover Change Trajectories from Multi-temporal Satellite Imagery, *Annals of the Association of American Geographers*, 102, 1329-1347, 10.1080/00045608.2011.596357, 2012.

Liu, F., Zhang, Z., Zhao, X., Wang, X., Zuo, L., Wen, Q., Yi, L., Xu, J., Hu, S., and Liu, B.: Chinese cropland losses due to urban expansion in the past four decades, *Science of The Total Environment*, 650, 847-857, <https://doi.org/10.1016/j.scitotenv.2018.09.091>, 2019.

580 Liu, G.: Understanding cotton cultivation dynamics in Aksu Oases (NW China) by reconstructing change trajectories using multi-temporal Landsat and Sentinel-2 data, *Geocarto International*, 37, 4406-4424, 10.1080/10106049.2021.1886337, 2022.

Liu, J., Kuang, W., Zhang, Z., Xu, X., Qin, Y., Ning, J., Zhou, W., Zhang, S., Li, R., Yan, C., Wu, S., Shi, X., Jiang, N., Yu, D., Pan, X., and Chi, W.: Spatiotemporal characteristics, patterns, and causes of land-use changes in China since the late 1980s, *Journal of Geographical Sciences*, 24, 195-210, 10.1007/s11442-014-1082-6, 2014a.

585 Liu, Y., Fang, F., and Li, Y.: Key issues of land use in China and implications for policy making, *Land Use Policy*, 40, 6-12, <https://doi.org/10.1016/j.landusepol.2013.03.013>, 2014b.

Lu, M., Wu, W., Zhang, L., Liao, A., Peng, S., and Tang, H.: A comparative analysis of five global cropland datasets in China, *Science China Earth Sciences*, 59, 2307-2317, 10.1007/s11430-016-5327-3, 2016.

590 MacDonald, D., Crabtree, J. R., Wiesinger, G., Dax, T., Stamou, N., Fleury, P., Gutierrez Lazpita, J., and Gibon, A.: Agricultural abandonment in mountain areas of Europe: Environmental consequences and policy response, *Journal of Environmental Management*, 59, 47-69, <https://doi.org/10.1006/jema.1999.0335>, 2000.

Masek, J. G., Vermote, E. F., Saleous, N. E., Wolfe, R., Hall, F. G., Huemmrich, K. F., Feng, G., Kutler, J., and Teng-Kui, L.: A Landsat surface reflectance dataset for North America, 1990-2000, *IEEE Geoscience and Remote Sensing Letters*, 3, 68-72, 10.1109/LGRS.2005.857030, 2006.

595 Maus, V., Câmara, G., Cartaxo, R., Sanchez, A., Ramos, F. M., and Queiroz, G. R. d.: A Time-Weighted Dynamic Time Warping Method for Land-Use and Land-Cover Mapping, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 9, 3729-3739, 10.1109/JSTARS.2016.2517118, 2016.

Olofsson, P., Foody, G. M., Herold, M., Stehman, S. V., Woodcock, C. E., and Wulder, M. A.: Good practices for estimating area and assessing accuracy of land change, *Remote Sensing of Environment*, 148, 42-57, <https://doi.org/10.1016/j.rse.2014.02.015>, 2014.

600 Pasquarella, V. J., Arévalo, P., Bratley, K. H., Bullock, E. L., Gorelick, N., Yang, Z., and Kennedy, R. E.: Demystifying LandTrendr and CCDC temporal segmentation, *International Journal of Applied Earth Observation and Geoinformation*, 110, 102806, <https://doi.org/10.1016/j.jag.2022.102806>, 2022.

Pekel, J.-F., Cottam, A., Gorelick, N., and Belward, A. S.: High-resolution mapping of global surface water and its long-term changes, *Nature*, 540, 418-422, 10.1038/nature20584, 2016.

605 Pittman, K., Hansen, M. C., Becker-Reshef, I., Potapov, P. V., and Justice, C. O.: Estimating Global Cropland Extent with Multi-year MODIS Data, 10.3390/rs2071844, 2010.

Potapov, P., Turubanova, S., Hansen, M. C., Tyukavina, A., Zalas, V., Khan, A., Song, X.-P., Pickens, A., Shen, Q., and Cortez, J.: Global maps of cropland extent and change show accelerated cropland expansion in the twenty-first century, *Nature Food*, 3, 19-28, 2022.

610 Potapov, P. V., Turubanova, S. A., Hansen, M. C., Adusei, B., Broich, M., Altstatt, A., Mane, L., and Justice, C. O.: Quantifying forest cover loss in Democratic Republic of the Congo, 2000–2010, with Landsat ETM+ data, *Remote Sensing of Environment*, 122, 106-116, <https://doi.org/10.1016/j.rse.2011.08.027>, 2012.

Prishchepov, A. V., Radeloff, V. C., Dubinin, M., and Alcantara, C.: The effect of Landsat ETM/ETM+ image acquisition dates on the detection of agricultural land abandonment in Eastern Europe, *Remote Sensing of Environment*, 126, 195-209, <https://doi.org/10.1016/j.rse.2012.08.017>, 2012.

Qiu, B., Li, H., Tang, Z., Chen, C., and Berry, J.: How cropland losses shaped by unbalanced urbanization process?, *Land Use Policy*, 96, 104715, 2020.

620 Queiroz, C., Beilin, R., Folke, C., and Lindborg, R.: Farmland abandonment: threat or opportunity for biodiversity conservation? A global review, *Frontiers in Ecology and the Environment*, 12, 288-296, <https://doi.org/10.1890/120348>, 2014.

Radoux, J., Lamarche, C., Van Bogaert, E., Bontemps, S., Brockmann, C., and Defourny, P.: Automated Training Sample Extraction for Global Land Cover Mapping, 10.3390/rs6053965, 2014.

Ren, C., Zhou, X., Wang, C., Guo, Y., Diao, Y., Shen, S., Reis, S., Li, W., Xu, J., and Gu, B.: Ageing threatens sustainability of smallholder farming in China, *Nature*, 10.1038/s41586-023-05738-w, 2023.

625 Roberts, L.: 9 Billion?, *Science*, 333, 540-543, 10.1126/science.333.6042.540, 2011.

Roy, D. P., Kovalskyy, V., Zhang, H. K., Vermote, E. F., Yan, L., Kumar, S. S., and Egorov, A.: Characterization of Landsat-7 to Landsat-8 reflective wavelength and normalized difference vegetation index continuity, *Remote Sensing of Environment*, 185, 57-70, <https://doi.org/10.1016/j.rse.2015.12.024>, 2016.

Schneibel, A., Stellmes, M., Röder, A., Frantz, D., Kowalski, B., Haß, E., and Hill, J.: Assessment of spatio-temporal changes of smallholder cultivation patterns in the Angolan Miombo belt using segmentation of Landsat time series, *Remote Sensing of Environment*, 195, 118-129, <https://doi.org/10.1016/j.rse.2017.04.012>, 2017.

Searchinger, T., Heimlich, R., Houghton, R. A., Dong, F., Elobeid, A., Fabiosa, J., Tokgoz, S., Hayes, D., and Yu, T.-H.: Use of U.S. Croplands for Biofuels Increases Greenhouse Gases Through Emissions from Land-Use Change, *Science*, 319, 1238-1240, 10.1126/science.1151861, 2008.

See, L., Fritz, S., You, L., Ramankutty, N., Herrero, M., Justice, C., Becker-Reshef, I., Thornton, P., Erb, K., Gong, P., Tang, H., van der Velde, M., Erickson, P., McCallum, I., Kraxner, F., and Obersteiner, M.: Improved global cropland data as an essential ingredient for food security, *Global Food Security*, 4, 37-45, <https://doi.org/10.1016/j.gfs.2014.10.004>, 2015.

Shao, Y. and Lunetta, R. S.: Comparison of support vector machine, neural network, and CART algorithms for the land-cover classification using limited training data points, *ISPRS Journal of Photogrammetry and Remote Sensing*, 70, 78-87, <https://doi.org/10.1016/j.isprsjprs.2012.04.001>, 2012.

Tang, L., Ke, X., Chen, Y., Wang, L., Zhou, Q., Zheng, W., and Xiao, B.: Which impacts more seriously on natural habitat loss and degradation? Cropland expansion or urban expansion?, *Land Degradation & Development*, 32, 946-964, <https://doi.org/10.1002/ldr.3768>, 2021.

Thenkabail, P. S., Teluguntla, P. G., Xiong, J., Oliphant, A., Congalton, R. G., Ozdogan, M., Gumma, M. K., Tilton, J. C., Giri, C., Milesi, C., Phalke, A., Massey, R., Yadav, K., Sankey, T., Zhong, Y., Aneece, I., and Foley, D.: Global cropland-extent product at 30-m resolution (GCEP30) derived from Landsat satellite time-series data for the year 2015 using multiple machine-learning algorithms on Google Earth Engine cloud, Reston, VA, Report 1868, 63, 10.3133/pp1868, 2021.

Tilman, D., Balzer, C., Hill, J., and Befort, B. L.: Global food demand and the sustainable intensification of agriculture, *Proceedings of the National Academy of Sciences*, 108, 20260-20264, 10.1073/pnas.1116437108, 2011.

Traoré, F., Bonkoungou, J., Compaoré, J., Kouadio, L., Wellens, J., Hallot, E., and Tychon, B.: Using Multi-Temporal Landsat Images and Support Vector Machine to Assess the Changes in Agricultural Irrigated Areas in the Mogtedo Region, Burkina Faso, 10.3390/rs11121442, 2019.

Tu, Y., Chen, B., Zhang, T., and Xu, B.: Regional Mapping of Essential Urban Land Use Categories in China: A Segmentation-Based Approach, 10.3390/rs12071058, 2020.

Tu, Y., Chen, B., Yu, L., Xin, Q., Gong, P., and Xu, B.: How does urban expansion interact with cropland loss? A comparison of 14 Chinese cities from 1980 to 2015, *Landscape Ecology*, 36, 243-263, 10.1007/s10980-020-01137-y, 2021.

Tu, Y., Wu, S., Chen, B., Weng, Q., Gong, P., Bai, Y., Yang, J., Yu, L., and Xu, B.: A 30 m annual cropland dataset of China from 1986 to 2021 [dataset], 10.5281/zenodo.7936884, 2023a.

Tu, Y., Chen, B., Yu, L., Song, Y., Wu, S., Li, M., Wei, H., Chen, T., Lang, W., Gong, P., and Xu, B.: Raveling the nexus between urban expansion and cropland loss in China, *Landscape Ecology*, 10.1007/s10980-023-01653-7, 2023b.

United Nations: United Nations Transforming Our World: The 2030 Agenda for Sustainable Development, Division for Sustainable Development Goals: New York, NY, USA, 2015.

Verbesselt, J., Hyndman, R., Newnham, G., and Culvenor, D.: Detecting trend and seasonal changes in satellite image time series, *Remote Sensing of Environment*, 114, 106-115, <https://doi.org/10.1016/j.rse.2009.08.014>, 2010.

Vermote, E., Justice, C., Claverie, M., and Franch, B.: Preliminary analysis of the performance of the Landsat 8/OLI land surface reflectance product, *Remote Sensing of Environment*, 185, 46-56, <https://doi.org/10.1016/j.rse.2016.04.008>, 2016.

Vuichard, N., Ciais, P., Belelli, L., Smith, P., and Valentini, R.: Carbon sequestration due to the abandonment of agriculture in the former USSR since 1990, *Global Biogeochemical Cycles*, 22, <https://doi.org/10.1029/2008GB003212>, 2008.

Waldner, F., Canto, G. S., and Defourny, P.: Automated annual cropland mapping using knowledge-based temporal features, *ISPRS Journal of Photogrammetry and Remote Sensing*, 110, 1-13, <https://doi.org/10.1016/j.isprsjprs.2015.09.013>, 2015a.

Waldner, F., Fritz, S., Di Gregorio, A., and Defourny, P.: Mapping Priorities to Focus Cropland Mapping Activities: Fitness Assessment of Existing Global, Regional and National Cropland Maps, 10.3390/rs70607959, 2015b.

675 Wardlow, B. D. and Egbert, S. L.: Large-area crop mapping using time-series MODIS 250 m NDVI data: An assessment for the U.S. Central Great Plains, *Remote Sensing of Environment*, 112, 1096-1116, <https://doi.org/10.1016/j.rse.2007.07.019>, 2008.

680 Xie, Y., Lark, T. J., Brown, J. F., and Gibbs, H. K.: Mapping irrigated cropland extent across the conterminous United States at 30 m resolution using a semi-automatic training approach on Google Earth Engine, *ISPRS Journal of Photogrammetry and Remote Sensing*, 155, 136-149, <https://doi.org/10.1016/j.isprsjprs.2019.07.005>, 2019.

685 Xiong, J., Thenkabail, P. S., Gumma, M. K., Teluguntla, P., Poehnelt, J., Congalton, R. G., Yadav, K., and Thau, D.: Automated cropland mapping of continental Africa using Google Earth Engine cloud computing, *ISPRS Journal of Photogrammetry and Remote Sensing*, 126, 225-244, <https://doi.org/10.1016/j.isprsjprs.2017.01.019>, 2017.

690 Xu, H., Qi, S., Li, X., Gao, C., Wei, Y., and Liu, C.: Monitoring three-decade dynamics of citrus planting in Southeastern China using dense Landsat records, *International Journal of Applied Earth Observation and Geoinformation*, 103, 102518, <https://doi.org/10.1016/j.jag.2021.102518>, 2021.

695 Xu, Y., Yu, L., Zhao, F. R., Cai, X., Zhao, J., Lu, H., and Gong, P.: Tracking annual cropland changes from 1984 to 2016 using time-series Landsat images with a change-detection and post-classification approach: Experiments from three sites in Africa, *Remote Sensing of Environment*, 218, 13-31, <https://doi.org/10.1016/j.rse.2018.09.008>, 2018.

700 Xu, Y., Yu, L., Peng, D., Zhao, J., Cheng, Y., Liu, X., Li, W., Meng, R., Xu, X., and Gong, P.: Annual 30-m land use/land cover maps of China for 1980–2015 from the integration of AVHRR, MODIS and Landsat data using the BFAST algorithm, *Science China Earth Sciences*, 63, 1390-1407, 10.1007/s11430-019-9606-4, 2020.

705 **Xue, J., Zhang, X.-l., Chen, S.-c., Hu, B.-f., Wang, N., and Shi, Z.: Quantifying the agreement and accuracy characteristics of four satellite-based LULC products for cropland classification in China, *Journal of Integrative Agriculture*, <https://doi.org/10.1016/j.jia.2023.06.005>, 2023.**

710 Yan, J., Yang, Z., Li, Z., Li, X., Xin, L., and Sun, L.: Drivers of cropland abandonment in mountainous areas: A household decision model on farming scale in Southwest China, *Land Use Policy*, 57, 459-469, <https://doi.org/10.1016/j.landusepol.2016.06.014>, 2016.

715 Yang, J. and Huang, X.: The 30 m annual land cover dataset and its dynamics in China from 1990 to 2019, *Earth System Science Data*, 13, 3907-3925, 10.5194/essd-13-3907-2021, 2021.

720 Yin, H., Pflugmacher, D., Kennedy, R. E., Sulla-Menashe, D., and Hostert, P.: Mapping Annual Land Use and Land Cover Changes Using MODIS Time Series, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 7, 3421-3427, 10.1109/JSTARS.2014.2348411, 2014.

725 Yin, H., Prishchepov, A. V., Kuemmerle, T., Bleyhl, B., Buchner, J., and Radeloff, V. C.: Mapping agricultural land abandonment from spatial and temporal segmentation of Landsat time series, *Remote Sensing of Environment*, 210, 12-24, <https://doi.org/10.1016/j.rse.2018.02.050>, 2018.

730 Yin, H., Brandão, A., Buchner, J., Helmers, D., Iuliano, B. G., Kimambo, N. E., Lewińska, K. E., Razenkova, E., Rizayeva, A., Rogova, N., Spahn, S. A., Xie, Y., and Radeloff, V. C.: Monitoring cropland abandonment with Landsat time series, *Remote Sensing of Environment*, 246, 111873, <https://doi.org/10.1016/j.rse.2020.111873>, 2020.

735 Yu, L., Wang, J., Clinton, N., Xin, Q., Zhong, L., Chen, Y., and Gong, P.: FROM-GC: 30 m global cropland extent derived through multisource data integration, *International Journal of Digital Earth*, 6, 521-533, 2013.

740 Yu, Q., Hu, Q., van Vliet, J., Verburg, P. H., and Wu, W.: GlobeLand30 shows little cropland area loss but greater fragmentation in China, *International Journal of Applied Earth Observation and Geoinformation*, 66, 37-45, <https://doi.org/10.1016/j.jag.2017.11.002>, 2018.

745 Zabel, F., Delzeit, R., Schneider, J. M., Seppelt, R., Mauser, W., and Václavík, T.: Global impacts of future cropland expansion and intensification on agricultural markets and biodiversity, *Nature Communications*, 10, 2844, 10.1038/s41467-019-10775-z, 2019.

750 Zhang, C., Dong, J., and Ge, Q.: Mapping 20 years of irrigated croplands in China using MODIS and statistics and existing irrigation products, *Scientific Data*, 9, 407, 10.1038/s41597-022-01522-z, 2022a.

755 Zhang, C., Dong, J., and Ge, Q.: Quantifying the accuracies of six 30-m cropland datasets over China: A comparison and evaluation analysis, *Computers and Electronics in Agriculture*, 197, 106946, <https://doi.org/10.1016/j.compag.2022.106946>, 2022b.

760 Zhang, G., Xiao, X., Dong, J., Kou, W., Jin, C., Qin, Y., Zhou, Y., Wang, J., Menarguez, M. A., and Biradar, C.: Mapping paddy rice planting areas through time series analysis of MODIS land surface temperature and vegetation index data,

Zhang, H. K. and Roy, D. P.: Using the 500m MODIS land cover product to derive a consistent continental scale 30m Landsat
725 land cover classification, *Remote Sensing of Environment*, 197, 15-34, <https://doi.org/10.1016/j.rse.2017.05.024>, 2017.

Zhang, X., Zhao, T., Xu, H., Liu, W., Wang, J., Chen, X., and Liu, L.: **GLC_FCS30D**: The first global 30-m land-cover
730 dynamic monitoring product with a fine classification system from 1985 to 2022 using dense time-series Landsat imagery
and continuous change-detection method, *Earth Syst. Sci. Data Discuss.*, 2023, 1-32, 10.5194/essd-2023-320, 2023.

Zhang, Y., Li, X., and Song, W.: Determinants of cropland abandonment at the parcel, household and village levels in mountain
735 areas of China: A multi-level analysis, *Land Use Policy*, 41, 186-192, <https://doi.org/10.1016/j.landusepol.2014.05.011>,
2014.

Zhong, L., Gong, P., and Biging, G. S.: Efficient corn and soybean mapping with temporal extendability: A multi-year
735 experiment using Landsat imagery, *Remote Sensing of Environment*, 140, 1-13, <https://doi.org/10.1016/j.rse.2013.08.023>,
2014.

Zhu, L., Liu, X., Wu, L., Tang, Y., and Meng, Y.: Long-Term Monitoring of Cropland Change near Dongting Lake, China,
Using the LandTrendr Algorithm with Landsat Imagery, 10.3390/rs11101234, 2019.

Zhu, Z.: Change detection using landsat time series: A review of frequencies, preprocessing, algorithms, and applications,
740 *ISPRS Journal of Photogrammetry and Remote Sensing*, 130, 370-384, <https://doi.org/10.1016/j.isprsjprs.2017.06.013>,
2017.

Zhu, Z. and Woodcock, C. E.: Object-based cloud and cloud shadow detection in Landsat imagery, *Remote Sensing of
745 Environment*, 118, 83-94, <https://doi.org/10.1016/j.rse.2011.10.028>, 2012.

Zhu, Z. and Woodcock, C. E.: Continuous change detection and classification of land cover using all available Landsat data,
745 *Remote Sensing of Environment*, 144, 152-171, <https://doi.org/10.1016/j.rse.2014.01.011>, 2014.

Zhu, Z., Gallant, A. L., Woodcock, C. E., Pengra, B., Olofsson, P., Loveland, T. R., Jin, S., Dahal, D., Yang, L., and Auch, R.
F.: Optimizing selection of training and auxiliary data for operational land cover classification for the LCMAP initiative,
750 *ISPRS Journal of Photogrammetry and Remote Sensing*, 122, 206-221, <https://doi.org/10.1016/j.isprsjprs.2016.11.004>,
2016.

Zuo, L., Zhang, Z., Carlson, K. M., MacDonald, G. K., Brauman, K. A., Liu, Y., Zhang, W., Zhang, H., Wu, W., Zhao, X.,
Wang, X., Liu, B., Yi, L., Wen, Q., Liu, F., Xu, J., Hu, S., Sun, F., Gerber, J. S., and West, P. C.: Progress towards
750 sustainable intensification in China challenged by land-use change, *Nature Sustainability*, 1, 304-313, 10.1038/s41893-
018-0076-2, 2018.





Figure 1: Flowchart of this study for mapping annual cropland dynamics.



760

Figure 2: Schematic diagram of the segmentation results of annual cropland probability time series based on LandTrendr.






Figure 3: Spatial distribution of validation samples collected in this study (taking the year 1986 as an example).



765

Figure 4: Spatial variation of mean accuracy of CACD for 1986-2020. (a-c) Hexagon level. Noted only hexagons containing both cropland and non-cropland samples were calculated. (d-f) Province-level. Noted Xizang, Hong Kong, and Macau were excluded from the analysis due to small numbers of cropland samples.



770 **Figure 5: Comparisons of the pixel-wise accuracy of F1 score between the five products based on the annual validation samples.**




Figure 6: Spatial agreement of cropland extent of the five products (CACD, CLCD, CLUD, GLAD, and GFSAD) in 2015. Matched level indicates how often a pixel is identified as cropland. For example, '4' means four out of the five products recognize it as cropland. The bar chart in the lower-left corner denotes the area proportion of different matched levels (1-5). Locations A-F respectively correspond to zoomed areas in Figure 7.





Figure 7: Regional comparisons of the five products in 2015. Referencing Landsat imagery in the first column is composed by SWIR1, NIR, and Red bands in RGB channels. Columns 2-6 display the spatial distribution of the five products, with cropland shown in white and non-cropland shown in black. All maps have a scale of 1:100000. Locations of A-F are shown in Figure 6.



780

Figure 8: Scatterplots of the provincial cropland area in 2015 between the five products and the statistical data. The red dotted line represents the 1:1 diagonal.




Figure 9: Cropland area changes in China between 1986-2021. (a) Provincial cropland change area derived from the CACD data.

785

(b) Annual dynamics of the map-based and the sample-based cropland area at the national scale.




Figure 10: Examples of cropland expansion and loss between 1986-2021 in four selected regions in China. (a) Chifeng, Inner Mongolia, where large-scale cropland expansion occurred after 2006 for pasture cultivation. (b) Aksu, Xinjiang. The region was mainly dominated by cotton cultivation. (c) Shanghai, where rapid urbanization induced extensive cropland loss. (d) Tongren, Guizhou, where some croplands in the mountainous area had been abandoned due to soil infertile and rural out-migration. All the Landsat images are freely provided by USGS. Both base maps in (c) and (d) are open web maps provided by Esri.




Figure 11: Cropland abandonment in China between 1990-2015. (a) Spatial distribution of abandonment time. (b) Histogram distribution within different elevation ranges. (c) Annual trends of abandonment area. The elevation information is derived from the SRTM digital elevation dataset (Jarvis et al., 2008).

795

Tables

Table. 1. Rules for annual cropland transition discrimination based on the LandTrendr segmentation results. $p_{is}(t_{is}, v_{is})$ and $p_{ie}(t_{ie}, v_{ie})$ are the starting and ending breakpoints of the i -th segment s_i , respectively, where t is the year and v is the fitted cropland probability.

Fitted cropland probabilities of p_{is} and p_{ie}	Transition type of s_i	Land use type for each year in s_i
$v_{is} > 0.5, v_{ie} > 0.5$	Stable cropland	Cropland from t_{is} to t_{ie}
$v_{is} \leq 0.5, v_{ie} \leq 0.5$	Stable non-cropland	Non-cropland from t_{is} to t_{ie}
$v_{is} > 0.5, v_{ie} \leq 0.5$	Cropland loss	Cropland in t_{is} and non-cropland from t_{is+1} to t_{ie}
$v_{is} \leq 0.5, v_{ie} > 0.5$	Cropland gain	Non-cropland in t_{is} and cropland from t_{is+1} to t_{ie}

800

Table. 2. Change accuracy of CACD based on the annual validation samples. Change is defined as any cropland use conversion identified between 1986-2021. UA: user's accuracy. PA: producer's accuracy.

		Map			
		No-change	Change	Total	PA
Reference	No-change	95	5	100	0.95
	Change	24	62	86	0.72
	Total	119	67	186	
	UA	0.80	0.93		
F1 score: 0.81; Overall accuracy: 0.84; Kappa coefficient: 0.68					

Table. 3. Comparison of mapping accuracy of CACD, CLUD, CLCD, GLAD, and GFSAD in 2015 based on GLCVSS. F1: F1 score. OA: overall accuracy. Kappa: Kappa coefficient. UA: user's accuracy. PA: producer's accuracy.

Product	F1	OA	UA	PA	Kappa
CACD	0.82	0.94	0.83	0.81	0.78
CLCD	0.77	0.92	0.82	0.73	0.73
CLUD	0.70	0.90	0.72	0.67	0.64
GLAD	0.72	0.92	0.64	0.83	0.68
GFSAD	0.76	0.93	0.77	0.76	0.72

Table. 4. Comparison of agreement level of CACD, CLUD, CLCD, GLAD, and GFSAD with Geo-Wiki sample set. Noted CACD, CLUD, and CLCD in 2016 are used while GLAD and GFSAD are from 2015.

Product	Agreement level
CACD	0.86
CLCD	0.83
CLUD	0.65
GLAD	0.60
GFSAD	0.73