

The patterns of soil nitrogen stocks and C:N stoichiometry under impervious surfaces in China

Qian Ding¹, Hua Shao², Chi Zhang^{2, 1, 3, *}, Xia Fang⁴

5

¹Shandong Provincial Key Laboratory of Water and Soil Conservation and Environmental Protection, College of Resources and Environment, Linyi University, Linyi, 270600, China.

²State Key Laboratory of Desert and Oasis Ecology, Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences, Urumqi, 830011, China.

10 ³Research Center for Ecology and Environment of Central Asia, Chinese Academy of Sciences, Urumqi, 830011, China.

⁴Xinjiang Institute of Engineering, Urumqi, 830091, China.

Correspondence to: Chi Zhang (zc@ms.xjb.ac.cn)

Abstract. Accurate assessment of soil nitrogen (N) storage and carbon (C):N stoichiometry under impervious
15 surface areas (ISAs) is key to understanding the impact of urbanization on soil health and the N cycle. Based
on 888 soil profiles from 148 sampling sites in 41 cities across China, we estimated the country's N stock
(100 cm depth) in the ISA soil to be 98.74 ± 59.13 Tg N with a mean N density (N_{ISA}) of 0.59 ± 0.35 kg m⁻²,
which was significantly lower (at all depths) than the soil N density ($N_{PSA} = 0.83 \pm 0.46$ kg m⁻²) under the
reference permeable surface areas (PSAs). The N_{ISA} was also only about 53–69% of the reported national
20 mean soil N density, indicating ISA expansion caused soil N loss. The C:N ratio of ISA (10.33 ± 2.62) was
26–34% higher than that of natural ecosystems (forests, grasslands, etc.), but close to the C:N of PSA.
Moreover, there was a significant C–N correlation in ISA soil, showing no signs of C–N decoupling as
suggested by the previous studies. The ISA had smaller variances in the C:N ratio than did the PSA at regional
scale, indicating convergence of soil C:N stoichiometry due to ISA conversion. The East subregion of China
25 had the highest N_{ISA} , although its natural soil N density was the among lowest in the country. Unlike the
vertical pattern in natural permeable soils, whose N density declined faster in the upper soil layers than in the
lower layers, N_{ISA} decreased linearly with depth. Similar to natural soil N, N_{ISA} was negatively correlated
with temperature; but unlike natural soil C:N which was positively correlated with temperature, the C: N_{ISA}
was negatively correlated with temperature. N_{ISA} was not correlated with net primary productivity, but
30 significantly correlated with the soil N density of adjacent PSA and the urbanization rate. These findings
indicate the ISA soil had unique N distribution pattern, possibly as the result of intensive disturbances during
land conversion.

1 Introduction

Nitrogen (N) is an essential nutrient that regulates ecosystem structure and function and maintains nutrient
35 cycling (Fowler et al., 2013). It affects ecosystem carbon sequestration in various ways (Vitousek and
Howarth, 1991), and the C:N stoichiometry conveys a rough measure of the mineralization and humification

of soil organic matter (SOM) (Chapin et al., 2011). Currently, global ecosystem structure and functions are intensively disrupted by urbanization, especially the rapid expansion of impervious surface area (ISA). The global ISA area in 2018 was 1.5 times larger than in 1990, at approximately $7.97 \times 10^5 \text{ km}^2$ (Gong et al., 2020).

40 It has been suggested that the expansion of ISA blocks soil–atmosphere carbon/water exchanges, alters the physicochemical properties of soil, and seriously disrupts soil biogeochemical processes (Wei et al., 2014a; Yu et al., 2019). N loss from disturbed urban soils may cause groundwater N contamination (Li et al., 2022). Thus, there is an urgent need to study N_{ISA} and SOC: N_{ISA} patterns to provide a solid basis for assessing the potential risk of N loss and other negative impacts on urban ecosystems (Pereira et al., 2021).

45 Due to the difficulties in sampling beneath impervious surfaces, our knowledge about the biogeochemical processes in sealed soils is still very limited. Although the knowledge gap has gained more attention recently, most of the studies in ISAs have focused on the soil organic carbon (SOC) pool but have generally overlooked the soil N pool (Yan et al., 2015; Bae and Ryu, 2020; Cambou et al., 2018; Short et al., 1986). These studies showed that soil sealing not only causes a large amount of SOC loss but also alters the structure of the SOC

50 pools, indicating profound changes in soil carbon (C) processes (Wei et al., 2013; Raciti et al., 2012; Ding et al., 2022). To date, only a few isolated studies from seven cities (three in China, three in Europe and one in the USA) have reported the soil N content/density under ISA (N_{ISA}) (Table 1). All of these studies indicated that N_{ISA} was lower than the N content/density (N_{PSA}) in pervious surface area (PSA). Two of them found extremely high C:N ratios in ISA (164 vs. 19 in PSA soil, 12.44 vs. 6.99 in PSA, respectively), suggesting

55 decoupling of C and N processes as the result of soil sealing (Raciti et al., 2012; O'riordan et al., 2021). A study from Nanjing city, China, however, found that ISAs had a lower C:N ratio than PSAs (Wei et al., 2014a).

Considering the high heterogeneity of urban soils, the available observations from 7 cities around the world are far from enough to provide useful information about the storage and characteristic distribution of N_{ISA} at

60 large scale. Previous studies focused on individual cities, but regional scale surveys are required to investigate the influences of climatic, ecological, geographic, and socioeconomic factors on N_{ISA} distribution. Such information is not only necessary to evaluate global N_{ISA} pool size, but also helpful in revealing the environmental-control mechanisms over the soil biogeochemical processes in ISA (Ding et al., 2022). For example, the urban ecosystem convergence theory suggests that cities from different regions tend to have

65 similar soil properties (e.g., SOC density) as a result of intensive human disturbances, even if their native soil properties differ significantly (Pouyat et al., 2003). Regional soil surveys from multiple cities are required to evaluate this theory with soil nutrient data. In addition, more observational data are required to evaluate whether ISA soil has extremely high C:N ratio, which might indicate decoupling of soil C and N processes (Raciti et al., 2012; O'riordan et al., 2021).

70 Investigations on the vertical distribution pattern of soil N are also important, because the nutrient distribution patterns through soil profiles are influenced by both natural and human factors. In natural ecosystems, vertical nutrient distributions are dominated by plant cycling relative to leaching, weathering dissolution, and

atmospheric deposition, leading to nutrient concentrating in topsoil (Jobbágy and Jackson, 2001). Previous studies in urban areas, however, showed that the removal of plants and topsoil in the ISA may alter the vertical pattern of SOC, resulting in a more homogeneous SOC distribution through the soil profile (Yan et al., 2015; Ding et al., 2022). Based on the observed SOC pattern, previous studies suggested that the changes in soil biogeochemistry in ISA was mainly caused by plant and topsoil removals and initial disturbance as opposed to postconstruction processes (Jobbágy and Jackson, 2001). Investigations on the vertical distribution patterns of N_{ISA} can help us to evaluate this mechanism. However, most previous studies only sampled the topsoil or upper soil layers (Table 1) and thus could not obtain a complete picture of the vertical distribution pattern of the N_{ISA} .

To address these issues, we investigated the patterns of China's N_{ISA} pool and C: N_{ISA} (C:N ratio of the ISA) based on 148 observations from 41 major cities across China (sampled at 100 cm depth and 20 cm intervals). The objectives of this study were to (a) compare N_{ISA} with N_{PSA} , (b) reveal the spatial pattern of N_{ISA} and C: N_{ISA} , and (c) identify the environmental factors correlated with N_{ISA} and discuss the underlying mechanism. We chose China as the study area because its urbanization rate is twice the global average, and approximately 2/3 of its urban area is occupied by ISA, which is also higher than the global average (Kuang, 2019). There are also relatively more previous N_{ISA} studies in Chinese cities than in other countries (Table 1), which makes it easier to evaluate our results.

Table 1: Compilation of soil N_{ISA} studies


City, country	Previous studies				This study		
	N density (kg m ⁻²)	N content (g kg ⁻¹)	Depth (cm)	References	N density (kg m ⁻²)*	N content (g kg ⁻¹)*	Depth (cm)
Beijing, China	NA	0.61	0–10	(Zhao et al., 2012)	0.08±0.02	0.34±0.06	0–20
	NA	0.54	10–20		0.08±0.02	0.34±0.06	0–20
	NA	0.42	20–30		0.09±0.02	0.4±0.11	20–40
	NA	0.26	30–40		0.09±0.02	0.4±0.11	20–40
	NA	0.37	0–15	(Hu et al., 2018)	0.08±0.02	0.34±0.06	0–20
Nanjing, China	NA	0.49	0–20	(Wei et al., 2014b)	0.38±0.05	0.13±0.15	0–20
Yixing, China	0.25	NA	0–20	(Wei et al., 2014a)	0.15±0.01	NA	0–20
New York, USA	0.014	NA	0–15	(Raciti et al., 2012)	0.10±0.06	NA	0–15
Lancaster, UK	NA	2.08	0–10	(Pereira et al., 2021)	0.07±0.04	NA	0–10
Greater Manchester, UK	0.081	NA	0–10	(O'riordan et al., 2021)	0.07±0.04	NA	0–10
Toruń, Poland	0.027	0.17	15–25 or 10–20	(Piotrowska-Długosz and Charzyński, 2015)	0.12±0.08	NA	0–20

*±1SD


2 Materials and methods

2.1 Soil sampling

The soil samples were collected from 148 sample sites in 41 cities that were evenly distributed across 95 mainland China except for the Tibetan Plateau during 2013–2014 (Figure 1). Depending on the city size, multiple sample sites were identified in each city. Each site belonged to a separate city district, i.e., the soil samples were taken from 148 different city districts across China. The sample sites included various ISA types (roads, elevated piers, buildings, etc.) and PSA types (trees, shrubs, herbs, bare ground, vegetable plots, etc.). Detailed descriptions of the cities and sample sites can be found in (Ding et al., 2023).



100



(b)

Figure 1: Study area. (a) Spatial distribution of the sampled cities. The numbers in the map are the IDs of the studied cities, which can be used to retrieve detailed information of the sample sites from the online dataset of this study (Ding et al., 2023). The background land-use type shows the regional dominant land-use/land-cover type where the cities locate. To facilitate spatial analysis, we divided the country into six subregions – E: eastern China, S: southern China, N: northern China, NE: northeastern China, NW: northwestern China, SW: southwestern China. (b) Example photos for different sampling sites.

At each sample site, 3 representative ISA sampling plots, more than 10 m apart from each other, were randomly selected. In addition, three paired sampling plots in adjacent PSAs were randomly selected for comparison. In each plot, a 100 cm depth profile pit was dug, and the soil profile was sampled at 20 cm intervals to the 100 cm depth with a 100 cm³ ring knife. Our study across China found that most of the Ekranic (sealed) Technosol profiles have a clear boundary between the building material layer and the soil. Where the boundary is unclear, we treated the topsoil with a high amount of hard building materials, where artifacts >0.15 mm accounted for over half of the soil volume, as the building material layer. We only took samples in the soil below the building material layer. Samples with notable additions of anthropogenic artifacts, e.g., coal fly ash, mixed in the soil were discarded. Following the protocol of China's National Soil Surveys, the visible non-soil artifacts in the remaining soil samples, such as fragmentations of bricks, glasses, stones, roots, etc., were picked out and discarded (Shi et al., 2004). A total of 4356 soil samples were eventually collected from 888 soil profiles. These samples (ID# XJBIZC0001–XJBIZC4356) are currently stored in the herbarium of the Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences.

To facilitate spatial analysis, we divided the country into six subregions – the northeast, north, northwest, east, south, and southwest, according to geography, climate, and socioeconomics following Ding et al. (2022). To estimate the N_{ISA} storage in each subregion, we multiple the mean N_{ISA} density in the region with the region's ISA land area, which was derived from the 30 m resolution ISA map of mainland China (Zhang et al., 2020). Then, the N_{ISA} stock of all subregions were added up to estimate the national N_{ISA} storage.

2.2 Soil physical and chemical analyses

In this study, soil bulk density (BD) and N content were measured for each soil sample. Soil samples inside the ring knife were dried at 105 °C, and soil bulk weight (BD) (g cm⁻³) was measured, while the rest of the samples were air dried and passed through a 0.15 mm sieve, and the N content (g kg⁻¹) was measured by Kjeldahl digestion (Bremner and Mulvaney, 1982). The N density (kg m⁻²) per 20 cm soil layer was calculated according to Eq. 1, and the N density for the entire 100 cm profile was obtained by summing the N density per 20 cm soil layer (Eq. 2).

$$N_i = \frac{NC_i \times BD_i \times 20}{100}, \quad (1)$$

$$N_{100cm} = \sum_{i=1}^n N_i, \quad (2)$$

where N represents N density (kg m^{-2}), $i \in [1,5]$ represents soil layer (each 20 cm in thickness), NC is N content (g kg^{-1}), BD is soil bulk density (g cm^{-3}).

2.3 Comparing N_{ISA} and N_{PSA} , $C:N_{\text{ISA}}$ and $C:N_{\text{PSA}}$

140 A paired T test (2 tailed) was used to determine the difference between N_{ISA} and N_{PSA} and the difference between $C:N_{\text{ISA}}$ and $C:N_{\text{PSA}}$ ($C:N$ ratio of the PSA). The $C:N$ stoichiometry, i.e., the $C:N$ ratio, shows the connection between the C process and N process. An extremely high $C:N_{\text{ISA}}$ in comparison with the reference $C:N_{\text{PSA}}$ indicates C–N decoupling due to soil sealing (Raciti et al., 2012).

145 The SOC density of the samples was reported in a previous study (Ding et al., 2022). We noticed that some research (Hu et al., 2018; Pereira et al., 2021; O'riordan et al., 2021) used the ratio between total C and total N to investigate the $C:N$ stoichiometry in ISA soil. However, the content of soil inorganic C under impervious surfaces is likely altered by anthropogenic C from construction materials, and black C (Zhao et al., 2017; O'riordan et al., 2021). In this study, we used the ratio between SOC and N to investigate the soil $C:N$ stoichiometry, just like most soil studies in both ISA (Wei et al., 2014a; Raciti et al., 2012; Piotrowska-Długosz and Charzyński, 2015) and PSA (Lu et al., 2023; Schroeder et al., 2022; Yang et al., 2021)..

150 We further investigate whether soil sealing may influence the variations of $C:N$ stoichiometry at the national scale. According to the urban ecosystem convergence theory, intensive human disturbances (e.g., soil sealing) could reduce variations in soil property at large scale (i.e., among different cities) even if the intensively disturbed areas may have similar or higher variations in soil properties at city scale compared to the less disturbed areas (e.g., PSA) (Pouyat et al., 2003). To evaluate this theory, we compared the mean inter-city
155 $C:N$ stoichiometry dissimilarity and the mean intra-city $C:N$ stoichiometry dissimilarity between the ISA and PSA. The inter-city dissimilarity (or regional scale dissimilarity) measured the Euclidean distance (Eq. 3) in $C:N$ between each pair of different cities, while the intra-city dissimilarity (or local scale dissimilarity) measured the Euclidean distance in $C:N$ between each pair of sampling sites within the same city, all combinations included. If the urban ecosystem convergence theory was correct, we expect to see ISA having
160 lower inter-city $C:N$ dissimilarity than PSA, but higher or similar intra-city $C:N$ dissimilarity than/to PSA.

$$\text{Euclidean distance} = \sqrt{(C:N_i - C:N_j)^2}, \quad (3)$$

where $C:N_i$ and $C:N_j$ are the soil $C:N$ ratios of site i and site j , respectively, when measuring the intra-city dissimilarity, or the city-averaged $C:N$ ratios of city i and city j , respectively, when measuring the inter-city dissimilarity.

165 2.4 Investigating the vertical pattern of N_{ISA}

Unlike other studies that focused on topsoil, our multiple-layer soil sampling data made it possible to study the vertical pattern of N_{ISA} to a 100 cm depth. The proportions of N stored in the 0–20 cm depth, 0–40 cm depth, 0–60 cm depth, and 0–80 cm depth to the total (100 cm depth) N stock in each sample profile were

calculated and plotted against the soil depth to reveal the vertical distribution pattern of N_{ISA} and N_{PSA} . Based on these data, we could model how N storage changed with soil depth. According to Yang et al. (2007), 46% of the N stock (in 1 m depth) of natural soil is stored in the top 0–30 cm soil, and 68% of the N stock is stored in the top 0–50 cm, translating into a power function fitting model (Figure 7):

$$N_{\text{Natural}}\%_d = -0.0074d^2 + 1.7378d = (1.7378 - 0.0074d) \times d, \quad (4)$$

where $N_{\text{Natural}}\%_d$ is the proportion of total N stock (in 100 cm depth) stored to depth d cm in natural soil in China. The equation shows that the $N_{\text{Natural}}\%_d$ does not increase linearly with soil depth, its increasing rate (i.e., $1.7378 - 0.0074d$) reduces with soil depth d . This pattern indicates the natural soil N does not have homogeneous N density through the soil profile, it decreases with depth.

2.5 Correlation analysis between N_{ISA} and potential environmental factors

Our large scale soil survey made it possible, for the first time, to investigate the correlations between soil N and various environmental factors so as to identify the climatic, ecological, geographical and socio-economic factors that may control or influence the N and C:N dynamics in sealed soil. In natural ecosystems, the distribution of N pools is significantly influenced by climate factors (Zhang et al., 2021). Temperature and precipitation are key drivers of soil biogeochemical processes (Wiesmeier et al., 2019). A previous study indicated that the ISA soil may also be affected indirectly by adjacent PSA (Yan et al., 2015), because many ISAs were converted from urban PSA during urban infilling (Delgado-Baquerizo et al., 2021; Kuang, 2019; Kuang et al., 2021). The soil organic matter input is influenced by ecosystem net primary productivity (NPP) (Chan, 2001). The N_{ISA} could also be correlated with the intensity of urbanization or human disturbances, which are influenced by population size, GDP, and the spatio-temporal patterns of built-up areas in a city (Bloom et al., 2008). Moreover, elevation and terrain may influence both the soil biogeochemical processes and ISA expansion (Zhu et al., 2022; Pan et al., 2023). Therefore, we selected 15 indicators to investigate the factors associated with N_{ISA} , including mean temperature, annual precipitation, background NPP (averaged natural ecosystem NPP in a 5 km buffer outside the city), C: N_{PSA} and N_{PSA} , longitude, latitude, elevation, population density, built-up area in a city, urbanization rate as indicated by the fraction of the built-up area that expanded after 2002, ISA coverage in built-up areas, greenspace coverage in built-up areas, per capita greenspace, city GDP, and per capita GDP. We also investigated the correlation between soil BD and the N_{ISA} content.

Gridded datasets of environmental factors, including mean annual temperature (Figure 1a), annual precipitation (Figure 1b), and elevation (Figure 1d) at 1 km resolution, were obtained from the Data Center for Resource and Environmental Sciences, Chinese Academy of Sciences (<http://www.resdc.cn/>). The national NPP (1985–2015) estimates at 1 km resolution was obtained from the Digital Journal of Global Change Data Repository (<https://www.geodoi.ac.cn/>) (Figure 1c). Statistical datasets include the Ministry of Housing and Urban–Rural Development of China (www.mohurd.gov.cn/) urban built-up area, population density, built-up area green space rate, per capita built-up area green space, the National Bureau of Statistics of China (data.stats.gov.cn/) total urban GDP, and per capita GDP. We used Pearson’s correlation (2-tailed)




Figure 2: A subset of the spatial datasets of climatic, ecological and terrain factors whose correlations with N_{ISA} were investigated in this study, including (a) annual precipitation normal (1981–2010), (b) air temperature normal (1981–2010), (c) mean annual NPP (1985–2015), and (d) digital elevation model (DEM). E: eastern China, S: southern China, N: northern China, NE: northeastern China, NW: northwestern China, SW: southwestern China.

3 Results

3.1 N densities and storage under ISA in China

The national mean N_{ISA} density in the 100 cm soil profile was $0.59 \pm 0.35 \text{ kg m}^{-2}$ (mean \pm 1 SD), ranging from 0.08–1.88 kg m^{-2} with a median value of 0.48 kg m^{-2} . Paired t tests showed that the N_{ISA} was significantly (approximately 30%) lower ($P < 0.01$) than the reference N_{PSA} (Figure 3a). Moreover, the N_{ISA} was lower than the reference N_{PSA} at all soil depths and in all subregions of China (Figure 4a). The national total N_{ISA} stock was about $98.74 \pm 59.13 \text{ Tg N}$.

$C:N_{ISA}$ (10.33 ± 2.62) was significantly lower than $C:N_{PSA}$ (10.93 ± 3.19) (Figure 3b). Moreover, N_{PSA} and SOC_{PSA} were significantly correlated ($R = 0.893$, $P < 0.01$), and N_{ISA} and SOC_{ISA} were also significantly correlated ($R = 0.926$, $P < 0.01$). There were no signs of C–N decoupling according to our data.





Figure 3: Comparing soil N density (a) and C:N (b) in the ISA and the reference PSA. The square box shows median and quad values, the inner small rectangle is the mean value, and the letters indicate the significance of the difference among the groups.

3.2 Spatial variation and spatial trend analysis

225 To facilitate spatial analysis, we divided the country into six subregions – the Northeast, North, Northwest, East, South, and Southwest, according to geography, climate, and socioeconomics (Ding et al., 2022) (Figure 1a). The highest N_{ISA} density of $0.84 \pm 0.45 \text{ kg m}^{-2}$ was found in the East, while the lowest N_{ISA} density of $0.51 \pm 0.20 \text{ kg m}^{-2}$ was found in the South (Figure 4a). Notably, the urban soil N densities, both in ISA and PSA, in the East were significantly higher than that of the other subregions except for the Southwest where the variations of soil N densities were extremely high (Figure 4a). As the result, the East subregion accounted

230 the variations of soil N densities were extremely high (Figure 4a). As the result, the East subregion accounted for the largest share (34%) of the N_{ISA} stock in China (Figure 4b).



235 **Figure 4: The spatial patterns of N_{ISA} in the six subregions of China. (a) the regional mean N_{ISA} density and N_{PSA} density, (b) the regional N_{ISA} storage. The letters indicate the significance of the difference among the subregions. * and ** indicate significant differences between ISA and PSA, $p < 0.01$ and $p < 0.05$, respectively.**

Figure 5 shows the ISA soil samples had lower inter-city C:N dissimilarity (1.86 ± 1.40 vs. 2.43 ± 2.15) than PSA, but similar intra-city C:N dissimilarity to PSA. This pattern indicates that although the ISA soil and the PSA soil had similar variations in C:N stoichiometry at the local scale (within a city), the C:N variations at national scale (among the cities) were reduced for the ISA soil, possibly due to the intensive human disturbances on ISA soil as predicted by the urban ecosystem convergence theory (Pouyat et al., 2003).




Figure 5: Comparing the inter-city variation and the intra-city variation of C:N ratios between the ISA and PSA. The variations were measured by the dissimilarity of (or the Euclidean distance between) paired observations. For intra-city variation, the soil C:N dissimilarity between each pair of different sampling sites within the same city were calculated and averaged; for inter-city variation, the soil C:N dissimilarity between each pair of different cities under investigation were calculated and averaged. The letters indicate the significance of the difference among the groups.

The spatial trend analysis of the N_{ISA} showed a slow decline followed by a rapid increase in the north–south direction and a rapid decline followed by a rapid increase in the east–west direction (Figure 6a); the spatial trend analyses of the N_{PSA} produced similar concave lines in both the north–south and the east–west directions but with a more drastic initial decline in the north–south direction and a flatter trend in the east–west direction (Figure 6b). The spatial trend analysis of $C:N_{ISA}$ showed a rapid increase in the north–south direction and a rapid decrease followed by a slow increase in the east–west direction (Figure 6c); the spatial trend analysis of $C:N_{PSA}$ showed a slow increase in the north–south direction and a slow decrease in the east–west direction (Figure 6d). According to the spatial trend analyses, the change rate of $C:N_{ISA}$ was significantly higher than that of $C:N_{PSA}$, which was consistent with the results of inter–city C:N ratio dissimilarity analysis (Figure 5).




Figure 6: The variation trend of N and C:N under the two surfaces.

3.4 Vertical distribution pattern of N_{ISA}

In this study, the vertical profiles of soils under ISAs were systematically sampled and analysed at 20 cm intervals to a 100 cm depth, and the storage of N_{ISA} increased linearly with soil depth (Figure 7). This linear distribution pattern was evident at the national scale ($R^2 = 1$, $P < 0.001$), and the vertical distribution pattern of N_{ISA} in China can be described by a linear model (Eq. 5):

$$N_{ISA} \%_d = 1.0324d, \quad (5)$$

where $N_{ISA} \%_d$ (%) is the percentage of total N storage (of 100 cm depth) stored in the top d (cm) depth of the soil. The unit of N_{ISA} is kg m^{-2} .




Figure 7: Comparing the vertical distribution pattern of N between the sealed soil (N_{ISA}) and the natural soil ($N_{Natural}$) in China (refer to Section 2.4 Equation 4).

3.5 The natural and socioeconomic factors correlated with N_{ISA} and $C:N_{ISA}$

270 The impacts of climate and geographic factors were confirmed by correlation analyses, which showed N_{ISA} to be negatively correlated with temperature ($R=-0.471$) but positively correlated with latitude ($R=0.386$) (Table 2). In addition, the N_{ISA} had a positive correlation with the N of urban PSA ($R=0.715$) and a negative correlation ($R=-0.34$) with the urbanization rate as indicated by the fraction of the newly expanded ISA since 2002 (i.e., $f_{new_ISA} = \frac{ISA_{2015}-ISA_{2002}}{ISA_{2015}}$). Surprisingly, we did not find significant correlations between N_{ISA} and

275 common environmental drivers like precipitation and NPP at 95% significant level, although the N_{ISA} had a weak negative correlation with precipitation ($R=-0.268$) at 90% significant level.

The $C:N_{ISA}$ was negatively correlated with both precipitation ($R=-0.539$) and temperature ($R=-0.496$) as well as longitude ($R=-0.316$), but positively correlated with latitude ($R=0.482$) (Table 2). In addition, the $C:N_{ISA}$ had a positive correlation with the N of urban PSA ($R=0.575$) and a negative correlation ($R=-0.485$) with the

280 NPP.

Table 2: Correlations between N_{ISA} , $C:N_{ISA}$ and potential environmental drivers

Factors	N density (kg m^{-2})		$C:N_{ISA}$	
	Correlation Coefficient	Sig. (2 tailed)	Correlation Coefficient	Sig. (2 tailed)
Longitude	0.196	0.22	-0.316*	0.04
Latitude	0.386*	0.01	0.482**	0.00
DEM (m)	0.141	0.38	0.378*	0.01

Annual precipitation (mm)	-0.268	0.09	-0.539**	0.00
Mean Temperature (°C)	-0.471**	0.00	-0.496**	0.00
NPP (g m ⁻²)	-0.096	0.55	-0.485**	0.00
ISA coverage in built-up area (%)	-0.126	0.43	-0.240	0.13
Built-up area (km ²)	-0.072	0.65	0.049	0.76
Greenspace coverage in built-up area (%)	-0.229	0.15	-0.001	0.99
Population density (person/km ²)	-0.032	0.84	-0.051	0.75
Per capita GDP (person/10 ⁴ yuan)	-0.012	0.94	0.000	1.00
City GDP (billion yuan)	-0.015	0.93	-0.028	0.86
Per capita greenspace (m ²)	0.098	0.54	0.098	0.54
The fraction of the newly expanded ISA since 2002 (%)	-0.340*	0.03	-0.090	0.58
N _{PSA} density (kg m ⁻²)	0.715**	0.00	NA	NA
C:N _{PSA}	NA	NA	0.575**	0.00
BD	-0.104	0.52	NA	NA

* p < 0.05;

** p < 0.01.

4 Data availability

285 The dataset "Observations of soil nitrogen and soil organic carbon to soil nitrogen stoichiometry under the
 impervious surfaces areas (ISA) of China" includes N density, N content, BD, C:N, and other related data
 under ISA and PSA. It also contains geographical coordinates of sampling locations, as well as spatial pattern
 layer files of N_{ISA} density and C:N_{ISA} in China. This dataset is available from the National Cryosphere Desert
 Data Center (www.ncdc.ac.cn/portal/metadata/review/04cee3f5-64bb-4b22-9368-ee1c55f9c2bb?lang=en)
 290 (Ding et al., 2023).

5 Discussion

5.1 Comparing the N density and C:N stoichiometry in ISA soil with those in natural soils

Our results were comparable to or moderately lower than the previously reported topsoil (0–20 cm) N_{ISA}
 contents/densities in Chinese cities, including Beijing (0.34±0.06 g kg⁻¹ vs. 0.37–0.61 g kg⁻¹) (Hu et al., 2018;
 295 Zhao et al., 2012), Nanjing city (0.13±0.15 g kg⁻¹ vs. 0.49 g kg⁻¹) (Wei et al., 2014b) and Yixing city
 (0.15±0.01 kg m⁻² vs. 0.25 kg m⁻²) (Wei et al., 2014a) (Table 1). Our observed N_{ISA} content (0.4 g kg⁻¹) in
 the 20–40 cm soil layer in Beijing was also comparable to the reports by Zhao et al. (2012) (0.26–0.42 g kg⁻¹)
 1). Outside China, the reported topsoil (0–10 cm) N_{ISA} density in Greater Manchester, UK (0.081 kg m⁻²),
 was comparable to our estimated mean N_{ISA} density in 0–10 cm (0.07±0.04 kg m⁻²) in China (O'riordan et

300 al., 2021), but the reported N_{ISA} in New York, USA and Toruń, Poland, was much lower than our results and the reports from other Chinese city studies (Table 1) (Raciti et al., 2012; Piotrowska-Długosz and Charzyński, 2015).

Compared with previous assessments of China's soil N stock, N_{ISA} accounted for 0.96–1.47% of the total soil N pool in China (Zhang et al., 2021; Xu et al., 2020). The N_{ISA} pool size (98.74 ± 59.13 Tg) exceeded the
305 vegetation N of scrubland (8.1–50 Tg) (Xu et al., 2020) and grassland (48.8 Tg) (Zhang et al., 2021) in China. The N_{ISA} density (0.58 ± 0.12 kg m⁻²) was lower than that of natural soil and equivalent to 53–69% of the national average (Yang et al., 2007; Xu et al., 2020). The N densities in ISA soil (0.59 ± 0.35 kg m⁻²) were lower than those in other ecosystems, such as forest (1.29 kg m⁻²), farmland (1.13 kg m⁻²), and grassland (1.11 kg m⁻²) (Xu et al., 2020), indicating that ISA construction resulted in N loss. Previous studies ignored
310 the impacts of ISAs (Tian et al., 2006; Yang et al., 2007) and thus might have overestimated China's soil N pool size.

Our estimated C: N_{ISA} (10.33 ± 2.62) in China matched the previously observed C: N_{ISA} (10.8) in Yixing city, China (Wei et al., 2014a). It has been suggested that different terrestrial ecosystems may have similar C:N ratios (Yang et al., 2021). This study showed that the C: N_{ISA} (10.33 ± 2.62) was only slightly lower than the
315 C: N_{PSA} (10.93 ± 3.19) in urban ecosystems but much higher than the C:N ratios of natural ecosystem soils such as forests (8.21), croplands (8.18), and grasslands (7.7) (Xu et al., 2020; Tang et al., 2018). Therefore, it is possible that the C:N stoichiometry could remain relatively stable within the same land–use type (natural ecosystems, urban areas, etc.) but might differ significantly among different natural or human–disturbed land–use types.

320 **5.2. The C:N stoichiometry analysis showing no sign of C–N decoupling in the ISA soil**

The above comparison indicates that ISA soil has a higher C:N than natural soils. A study in New York City reported that the N density in the ISA was 95% lower than that in the PSA, leading to an extremely high soil total C:total N ratio (Raciti et al., 2012; Majidzadeh et al., 2017). Therefore, Raciti et al. (2012) suggested that paving decouples the C and N cycles. Our observations, however, showed that the soil N of ISA was
325 only 30% lower than that of PSA, and the C: N_{ISA} was lower than the C: N_{PSA} in China. Furthermore, there was a significant positive correlation ($R=0.926$, $P<0.01$) between N and SOC in ISA soil. Similarly, Wei et al. (2014a) found that C: N_{ISA} was lower than C: N_{PSA} in Yixing city, China, and O'Riordan et al. (2021) found a significant positive correlation between N and C in ISA soil in Greater Manchester, UK, even though they also observed an increased total C:total N ratio in ISA soil compared to PSA soil. There were no signs of C–
330 N decoupling according to our data and others (O'Riordan et al., 2021; Wei et al., 2014a). It is possible that the extremely high C:N ratio observed in previous studies might be merely caused by anthropogenic C inputs that partially compensated for SOC loss during land conversion (O'Riordan et al., 2021). Because the construction materials could add large amounts of inorganic C into soil, it is preferable to investigate the C:N stoichiometry under ISAs with the C:N ratio rather than the total C:total N ratio. This study highlights the

335 important role of N in urban biogeochemical research, which helps to prevent us from being confused/misled
by the complex C dynamics in urban soil due to anthropogenic C inputs.

5.3 Potential driving factors of the N_{ISA} and $C:N_{ISA}$

The spatial distribution pattern of soil N was significantly correlated with climate factors such as temperature and precipitation in natural ecosystems (Yang et al., 2007). In general, the soil N in China's temperate and
340 subtropical ecosystems were negatively correlated with temperature (Lu et al., 2017). Similarly, our study found a negative correlation between N_{ISA} and temperature, and a positive correlation between N_{ISA} and latitude. There was no significant correlation between precipitation and the soil N in natural ecosystem, except for dryland where a positive correlation has been found (Lu et al., 2017). We didn't find significant
345 correlation between precipitation and N_{ISA} at the 95% confidence level, although there was a weak negative correlation at the 90% confidence level. Previous study showed the SOC_{ISA} was also negatively correlated with precipitation, and it was suggested that the observed soil biogeochemistry (SOC, nutrient content etc.) under impervious surface was mainly determined by the losses (esp. in topsoil) during land conversion (Majidzadeh et al., 2018; Cambou et al., 2018; Edmondson et al., 2012). Higher precipitation leads to higher soil nutrient loss during land conversion (Ding et al., 2022). The relatively weak correlation between N_{ISA}
350 and precipitation (compared with the correlation between SOC_{ISA} and precipitation) as well as the negative correlation between $C:N_{ISA}$ and precipitation might indicate that the N loss during land conversion was not as significant as the loss in SOC. It is also possible that the high N deposition in urban ISA might somehow replenish the N_{ISA} pool.

N_{ISA} was not correlated with background NPP but positively correlated with the soil N in the adjacent urban
355 PSA. This pattern agrees with the previous report that the SOC_{ISA} was mainly influenced by the SOC in the adjacent urban PSA rather than the background SOC and NPP (Ding et al., 2022). However, there was a negative correlation between $C:N_{ISA}$ and background NPP.

The soil C:N ratio could be a more stable parameter (Yang et al., 2021). Tian et al. (2010) found the soil C:N ratio was relatively stable among climate zones in rural ecosystems in China. It has been observed that the
360 soil stoichiometric characteristics in China are influenced by geographical parameters such as altitude and latitude (Sheng et al., 2022). Lu et al. (2023) found a lower C:N ratio at higher latitudes in China, suggesting a positive correlation between C:N and temperature in natural ecosystem soils. Our study, however, found that the $C:N_{ISA}$ ratio increased with latitude and that there was a significant negative correlation between the $C:N_{ISA}$ ratio and temperature. The soil C:N ratio of natural ecosystems is influenced by plant litter input and
365 N uptake. Ecosystems in warmer regions have higher NPP, resulting in higher inputs of litter with a high C:N ratio (compared with the soil C:N ratio) and higher N uptake by roots, thus reducing soil inorganic N. Therefore, the C:N ratio is positively correlated with temperature in natural ecosystems. However, the C:N ratio under the impervious surface is solely determined by the relative mineralization rate of C and N. It seems that soil ecosystems have a higher retention capacity for N than for C (C fixation is unlikely to be
370 found in sealed soil). Therefore, while both the soil N_{ISA} pool and the SOC_{ISA} pool decrease when the

temperature increases, the net N mineralization rate is lower than the C mineralization rate, leading to a negative correlation between the C:N_{ISA} ratio and temperature.

5.4 N_{ISA} had different regional distribution pattern and vertical distribution pattern from rural soil N

Our study and Tian et al. (2006)'s study on China's soil N had same subregion zone design. However, we
375 found the urban soil (both the ISA and PSA) in the East zone had the highest N density while Tian et al.
(2006) found the rural soil N density in the East zone was among the lowest in the country. The relatively
high precipitation and temperature in the East China may lead to high SOM decomposition rate and nutrient
leaching rate, which explains its low rural soil N density (Tian et al., 2010). However, the East region was
380 also the most developed region in China for the last several centuries. Its cities had high population density
and long urbanization history. The long-term intensive human activities might leave profoundly footprint in
the soil biogeochemical processes, significantly elevated its N content. This finding, together with the
relatively low inter-city C:N variations/dissimilarities in the ISA (see section 3.2), indicate intensive human
disturbances might override the nature environmental effects in shaping regional distribution pattern of soil
N processes, further confirmed the urban ecosystem convergence theory (Pouyat et al., 2003).

385 The vertical N distribution patterns are also different between the ISA soil and rural soil. Unlike the vertical
distribution pattern of natural soil N, which decreased with depth and should be modelled with a second-
degree polynomial fitting function (Eq 4), the vertical distribution of N_{ISA} was relatively homogeneous
through the soil profile and could be modelled with a linear function (Eq 5; Figure 7). The SOC under the
impervious surface had similar vertical distribution pattern (Ding et al., 2022). The unique vertical pattern
390 reflects the effect of human disturbance during land conversion, which led to topsoil soil organic matter and
nutrient loss and reduced the SOC and N gradient through the soil profile (Majidzadeh et al., 2018; Cambou
et al., 2018; Edmondson et al., 2012).

5.5 Potential applications of the data

N plays an important and complex role in natural ecosystems. Soil microbial C use efficiency is negatively
395 correlated with C:N (Schroeder et al., 2022). Incorporating N into Earth system models can improve the
accuracy of C cycle estimates (Fleischer et al., 2019), and a good description of N can help understand and
predict the patterns and mechanisms of global C dynamics (Zhang et al., 2021) and provide a reliable basis
for exploring how geochemical cycles are coupled.

For a long time, knowledge of biogeochemical cycles under impervious surfaces has been a major gap in
400 urban biogeochemical research. Until recently, the size and pattern of N_{ISA} pools and their contributions to
the global N cycle have remained unclear. Our research, which is the first national-scale study on N_{ISA} and
C:N_{ISA}, helps to fill this gap by improving our understanding of the special pattern of soil N under impervious
surfaces. Such information is necessary when assessing urbanization impacts on global C and N cycles
(Lorenz and Lal, 2009).

405 **Author contributions.** Conceptualization: CZ. Data curation: CZ. Formal analysis: QD. Funding acquisition: ZC and HS. Investigation: HS. and XF. Methodology: QD and CZ. Project administration: CZ and HS. Resources: HS. and XF. Software: QD. Supervision: CZ. Validation: QD. Visualization: QD. Writing – original draft preparation: QD. Writing – review & editing: QD., HS. and CZ.

Competing interests. The authors declare that they have no conflicts of interest.

410 **Acknowledgements.** We want to thank the reviewers for their constructive comments which are helpful for improving our article. This project was funded by the Strategic Priority Research Program of the Chinese Academy of Sciences (Grant No. XDA2006030201), the Natural Science Foundation of Xinjiang Uygur Autonomous Region (2022D01D02) and the National Natural Science Foundation of China (Grant 31770515). Chi Zhang is supported by the Taishan Scholars Program of Shandong, China (Grant
415 ts201712071).

References

- Bae, J. and Ryu, Y.: High soil organic carbon stocks under impervious surfaces contributed by urban deep cultural layers, *Landscape and Urban Planning*, 204, <https://doi.org/10.1016/j.landurbplan.2020.103953>, 2020.
- 420 Bloom, D. E., Canning, D., and Fink, G.: Urbanization and the wealth of nations, *Science*, 319, 772-775, <https://doi.org/10.1126/science.1153057>, 2008.
- Bremner, J. M. and Mulvaney, C. S.: *Methods of Soil Analysis. Part 2. Chemical and Microbiological Properties, Nitrogen-Total*, American Society of Agronomy, Soil Science Society of America, 595–624 pp.1982.
- 425 Cambou, A., Shaw, R. K., Huot, H., Vidal-Beaudet, L., Hunault, G., Cannavo, P., Nold, F., and Schwartz, C.: Estimation of soil organic carbon stocks of two cities, New York City and Paris, *Sci. Total Environ.*, 644, 452-464, <https://doi.org/10.1016/j.scitotenv.2018.06.322>, 2018.
- Chan, K. Y.: Soil particulate organic carbon under different land use and management, *Soil Use and Management*, 17, 217-221, <https://doi.org/10.1079/sum200180>, 2001.
- 430 Chapin, F. S., Matson, P. A., and Vitousek, P. M.: *Principles of Terrestrial Ecosystem Ecology*, Springer, New York, 2011.
- Delgado-Baquerizo, M., Eldridge, D. J., Liu, Y. R., Sokoya, B., Wang, J. T., Hu, H. W., He, J. Z., Bastida, F., Moreno, J. L., Bamigboye, A. R., Blanco-Pastor, J. L., Cano-Diaz, C., Illan, J. G., Makhalyane, T. P., Siebe, C., Trivedi, P., Zaady, E., Verma, J. P., Wang, L., Wang, J., Grebenc, T., Penaloza-Bojaca, G. F.,
435 Nahberger, T. U., Teixido, A. L., Zhou, X., Berdugo, M., Duran, J., Rodriguez, A., Zhou, X., Alfaro, F., Abades, S., Plaza, C., Rey, A., Singh, B. K., Tedersoo, L., and Fierer, N.: Global homogenization of the structure and function in the soil microbiome of urban greenspaces, *Science Advances*, 7, <https://doi.org/10.1126/sciadv.abg5809>, 2021.
- Ding, Q., Shao, H., Chen, X., and Zhang, C.: Urban Land Conversion Reduces Soil Organic Carbon Density Under Impervious Surfaces, *Global Biogeochemical Cycles*, 36, e2021GB007293, <https://doi.org/10.1029/2021GB007293>, 2022.
- Ding, Q., Shao, H., Zhang, C., and Fang, X.: Observations of soil nitrogen and soil organic carbon to soil nitrogen stoichiometry under the impervious surfaces areas (ISA) of China, *National Cryosphere Desert Data Center*, <https://doi.org/10.12072/ncdc.socn.db2851.2023>, 2023.
- 445 Edmondson, J. L., Davies, Z. G., McHugh, N., Gaston, K. J., and Leake, J. R.: Organic carbon hidden in urban ecosystems, *Sci Rep*, 2, <https://doi.org/10.1038/srep00963>, 2012.
- Fleischer, K., Rammig, A., De Kauwe, M. G., Walker, A. P., Domingues, T. F., Fuchslueger, L., Garcia, S., Goll, D. S., Grandis, A., Jiang, M., Haverd, V., Hofhansl, F., Holm, J. A., Kruijt, B., Leung, F., Medlyn, B. E., Mercado, L. M., Norby, R. J., Pak, B., von Randow, C., Quesada, C. A., Schaap, K., Valverde-Barrantes,

- 450 O. J., Wang, Y. P., Yang, X., Zaehle, S., Zhu, Q., and Lapola, D. M.: Amazon forest response to CO₂ fertilization dependent on plant phosphorus acquisition, *Nature Geoscience*, 12, 736-741, <https://doi.org/10.1038/s41561-019-0404-9>, 2019.
- Fowler, D., Coyle, M., Skiba, U., Sutton, M. A., Cape, J. N., Reis, S., Sheppard, L. J., Jenkins, A., Grizzetti, B., Galloway, J. N., Vitousek, P., Leach, A., Bouwman, A. F., Butterbach-Bahl, K., Dentener, F., Stevenson, D., Amann, M., and Voss, M.: The global nitrogen cycle in the twenty-first century, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368, 20130164, <https://doi.org/10.1098/rstb.2013.0164>, 2013.
- 455 Gong, P., Li, X., Wang, J., Bai, Y., Chen, B., Hu, T., Liu, X., Xu, B., Yang, J., Zhang, W., and Zhou, Y.: Annual maps of global artificial impervious area (GAIA) between 1985 and 2018, *Remote Sensing of Environment*, 236, 111510, <https://doi.org/10.1016/j.rse.2019.111510>, 2020.
- 460 Hu, Y., Dou, X., Li, J., and Li, F.: Impervious Surfaces Alter Soil Bacterial Communities in Urban Areas: A Case Study in Beijing, China, *Frontiers in Microbiology*, 9, <https://doi.org/10.3389/fmicb.2018.00226>, 2018.
- Jobbágy, E. G. and Jackson, R. B.: The distribution of soil nutrients with depth: Global patterns and the imprint of plants, *Biogeochemistry*, 53, 51-77, <https://doi.org/10.1023/A:1010760720215>, 2001.
- 465 Kuang, W.: Mapping global impervious surface area and green space within urban environments, *Science China-Earth Sciences*, 62, 1591-1606, <https://doi.org/10.1007/s11430-018-9342-3>, 2019.
- Kuang, W., Liu, J., Tian, H., Shi, H., Dong, J., Song, C., Li, X., Du, G., Hou, Y., Lu, D., Chi, W., Pan, T., Zhang, S., Hamdi, R., Yin, Z., Yan, H., Yan, C., Wu, S., Li, R., Yang, J., Dou, Y., Wu, W., Liang, L., Xiang, B., and Yang, S.: Cropland redistribution to marginal lands undermines environmental sustainability, *Natl. Sci. Rev.*, 9, nwab091-nwab091, <https://doi.org/10.1093/nsr/nwab091>, 2021.
- 470 Li, S., Liu, X., Yue, F., Yan, Z., Wang, T., Li, S., and Liu, C.: Nitrogen dynamics in the Critical Zones of China, *Progress in Physical Geography: Earth and Environment*, 46, 869-888, <https://doi.org/10.1177/03091333221114732>, 2022.
- Lorenz, K. and Lal, R.: Biogeochemical C and N cycles in urban soils, *Environment International*, 35, 1-8, <https://doi.org/10.1016/j.envint.2008.05.006>, 2009.
- 475 Lu, M., Zeng, F., Lv, S., Zhang, H., Zeng, Z., Peng, W., Song, T., Wang, K., and Du, H.: Soil C:N:P stoichiometry and its influencing factors in forest ecosystems in southern China, *Frontiers in Forests and Global Change*, 6, <https://doi.org/10.3389/ffgc.2023.1142933>, 2023.
- Lu, T., Zhang, W., Niu, J., Lin, Y., and Wu, M.: Study on Spatial Variability and Driving Factors of Stoichiometry of Nitrogen and Phosphorus in Soils of Typical Natural Zones of China, *Acta Pedologica Sinica*, 54, 682-692, 2017.
- Majidzadeh, H., Lockaby, B. G., and Governo, R.: Effect of home construction on soil carbon storage-A chronosequence case study, *Environmental Pollution*, 226, 317-323, <https://doi.org/10.1016/j.envpol.2017.04.005>, 2017.
- 485 Majidzadeh, H., Graeme Lockaby, B., Price, R., and Governo, R.: Soil carbon and nitrogen dynamics beneath impervious surfaces, *Soil Science Society of America Journal*, 82, 663-670, <https://doi.org/10.2136/sssaj2017.11.0381>, 2018.
- O'Riordan, R., Davies, J., Stevens, C., and Quinton, J. N.: The effects of sealing on urban soil carbon and nutrients, *SOIL*, 7, 661-675, <https://doi.org/10.5194/soil-7-661-2021>, 2021.
- 490 Pan, T., Kuang, W., Shao, H., Zhang, C., Wang, X., and Wang, X.: Urban expansion and intra-urban land evolution as well as their natural environmental constraints in arid/semiarid regions of China from 2000–2018, *Journal of Geographical Sciences*, 33, 1419-1441, <https://doi.org/10.1007/s11442-023-2136-4>, 2023.
- Pereira, M. C., O'Riordan, R., and Stevens, C.: Urban soil microbial community and microbial-related carbon storage are severely limited by sealing, *J. Soils Sediments*, 21, 1455-1465, <https://doi.org/10.1007/s11368-021-02881-7>, 2021.
- 495 Piotrowska-Długosz, A. and Charzyński, P.: The impact of the soil sealing degree on microbial biomass, enzymatic activity, and physicochemical properties in the Ekranic Technosols of Toruń (Poland), *J. Soils Sediments*, 15, 47-59, <https://doi.org/10.1007/s11368-014-0963-8>, 2015.
- Pouyat, R. V., Russell-Anelli, J., Yesilonis, I. D., and Groffman, P. M.: Soil carbon in urban forest ecosystems, in: *Potential of U.S. Forest Soils to Sequester Carbon and Mitigate the Greenhouse Effect*, CRC Press, 347-362, 2003.
- 500 Raciti, S. M., Hutyra, L. R., and Finzi, A. C.: Depleted soil carbon and nitrogen pools beneath impervious surfaces, *Environmental Pollution*, 164, 248-251, <https://doi.org/10.1016/j.envpol.2012.01.046>, 2012.

- Schroeder, J., Peplau, T., Pennekamp, F., Gregorich, E., Tebbe, C. C., and Poeplau, C.: Deforestation for agriculture increases microbial carbon use efficiency in subarctic soils, *Biology and Fertility of Soils*, <https://doi.org/10.1007/s00374-022-01669-2>, 2022.
- Sheng, H., Yin, Z., Zhou, P., and Thompson, M. L.: Soil C:N:P ratio in subtropical paddy fields: variation and correlation with environmental controls, *J. Soils Sediments*, 22, 21-31, <https://doi.org/10.1007/s11368-021-03046-2>, 2022.
- Shi, X., Yu, D., Warner, E., Pan, X., Petersen, G., Gong, Z., and Weindorf, D.: Soil database of 1: 1,000,000 digital soil survey and reference system of the Chinese genetic soil classification system, *Soil Survey Horizons*, 45, 129-136, <https://doi.org/10.2136/sh2004.4.0129>, 2004.
- Short, J. R., Fanning, D. S., Foss, J. E., and Patterson, J. C.: SOILS OF THE MALL IN WASHINGTON, DC .2. GENESIS, CLASSIFICATION, AND MAPPING, *Soil Science Society of America Journal*, 50, 705-710, <https://doi.org/10.2136/sssaj1986.03615995005000030031x>, 1986.
- Tang, X., Zhao, X., Bai, Y., Tang, Z., Wang, W., Zhao, Y., Wan, H., Xie, Z., Shi, X., Wu, B., Wang, G., Yan, J., Ma, K., Du, S., Li, S., Han, S., Ma, Y., Hu, H., He, N., Yang, Y., Han, W., He, H., Yu, G., Fang, J., and Zhou, G.: Carbon pools in China's terrestrial ecosystems: New estimates based on an intensive field survey, *Proceedings of the National Academy of Sciences*, 115, 4021-4026, <https://doi.org/10.1073/pnas.1700291115>, 2018.
- Tian, H., Chen, G., Zhang, C., Melillo, J. M., and Hall, C. A. S.: Pattern and variation of C:N:P ratios in China's soils: a synthesis of observational data, *Biogeochemistry*, 98, 139-151, <https://doi.org/10.1007/s10533-009-9382-0>, 2010.
- Tian, H., Wang, S., Liu, J., Pan, S., Chen, H., Zhang, C., and Shi, X.: Patterns of soil nitrogen storage in China, *Global Biogeochemical Cycles*, 20, <https://doi.org/10.1029/2005GB002464>, 2006.
- Vitousek, P. M. and Howarth, R. W.: Nitrogen limitation on land and in the sea: How can it occur?, *Biogeochemistry*, 13, 87-115, <https://doi.org/10.1007/BF00002772>, 1991.
- Wei, Z., Wu, S., Yan, X., and Zhou, S.: Density and Stability of Soil Organic Carbon beneath Impervious Surfaces in Urban Areas, *Plos One*, 9, <https://doi.org/10.1371/journal.pone.0109380>, 2014a.
- Wei, Z., Wu, S., Zhou, S., and Lin, C.: Installation of impervious surface in urban areas affects microbial biomass, activity (potential C mineralisation), and functional diversity of the fine earth, *Soil Research*, 51, 59-67, <https://doi.org/10.1071/sr12089>, 2013.
- Wei, Z., Wu, S., Zhou, S., Li, J., and Zhao, Q.: Soil Organic Carbon Transformation and Related Properties in Urban Soil Under Impervious Surfaces, *Pedosphere*, 24, 56-64, [https://doi.org/10.1016/s1002-0160\(13\)60080-6](https://doi.org/10.1016/s1002-0160(13)60080-6), 2014b.
- Wiesmeier, M., Urbanski, L., Hobbey, E., Lang, B., von Luetzow, M., Marin-Spiotta, E., van Wesemael, B., Rabot, E., Liess, M., Garcia-Franco, N., Wollschlaeger, U., Vogel, H.-J., and Koegel-Knabner, I.: Soil organic carbon storage as a key function of soils - A review of drivers and indicators at various scales, *Geoderma*, 333, 149-162, <https://doi.org/10.1016/j.geoderma.2018.07.026>, 2019.
- Xu, L., He, N., and Yu, G.: Nitrogen storage in China's terrestrial ecosystems, *Sci. Total Environ.*, 709, 136201, <https://doi.org/10.1016/j.scitotenv.2019.136201>, 2020.
- Yan, Y., Kuang, W., Zhang, C., and Chen, C.: Impacts of impervious surface expansion on soil organic carbon - a spatially explicit study, *Sci Rep*, 5, <https://doi.org/10.1038/srep17905>, 2015.
- Yang, J., Yuan, D., Zhao, Y., He, Y., and Zhang, G.: Stoichiometric relations of C, N, and P in urban top soils in Nanjing, China, and their biogeochemical implications, *J. Soils Sediments*, 21, 2154-2164, <https://doi.org/10.1007/s11368-020-02826-6>, 2021.
- Yang, Y., Ma, W., Mohammat, A., and Fang, J.: Storage, Patterns and Controls of Soil Nitrogen in China, *Pedosphere*, 17, 776-785, [https://doi.org/10.1016/S1002-0160\(07\)60093-9](https://doi.org/10.1016/S1002-0160(07)60093-9), 2007.
- Yu, W. W., Hu, Y. H., Cui, B. W., Chen, Y. Y., and Wang, X. K.: The Effects of Pavement Types on Soil Bacterial Communities across Different Depths, *Int. J. Environ. Res. Public Health*, 16, 11, <https://doi.org/10.3390/ijerph16101805>, 2019.
- Zhang, X., Liu, L., Wu, C., Chen, X., Gao, Y., Xie, S., and Zhang, B.: Development of a global 30-m impervious surface map using multisource and multitemporal remote sensing datasets with the Google Earth Engine platform, *Earth Syst. Sci. Data*, 12, 1625-1648, <https://doi.org/10.5194/essd-12-1625-2020>, 2020.
- Zhang, Y. W., Guo, Y., Tang, Z., Feng, Y., Zhu, X., Xu, W., Bai, Y., Zhou, G., Xie, Z., and Fang, J.: Patterns of nitrogen and phosphorus pools in terrestrial ecosystems in China, *Earth Syst. Sci. Data*, 13, 5337-5351, <https://doi.org/10.5194/essd-13-5337-2021>, 2021.

- Zhao, D., Li, F., Wang, R., Yang, Q., and Ni, H.: Effect of soil sealing on the microbial biomass, N transformation and related enzyme activities at various depths of soils in urban area of Beijing, China, *J. Soils Sediments*, 12, 519-530, <https://doi.org/10.1007/s11368-012-0472-6>, 2012.
- 560 Zhao, H., Wu, S., Xu, X., Zhou, S., and Li, X.: Spatial Distribution of Soil Inorganic Carbon in Urban Soil and Its Relationship with Urbanization History of the City, *Acta Pedologica Sinica*, 54, 1540-1546, <https://doi.org/10.11766/trxb201703300075>, 2017.
- 565 Zhu, G., Zhou, L., He, X., Wei, P., Lin, D., Qian, S., Zhao, L., Luo, M., Yin, X., Zeng, L., Long, Y., Hu, S., Ouyang, X., and Yang, Y.: Effects of Elevation Gradient on Soil Carbon and Nitrogen in a Typical Karst Region of Chongqing, Southwest China, *Journal of Geophysical Research: Biogeosciences*, 127, e2021JG006742, <https://doi.org/10.1029/2021JG006742>, 2022.