



MODIS Daily Cloud-gap-filled Fractional Snow Cover Dataset of the Asian Water Tower Region (2000-2022)

Fangbo Pan¹, Lingmei Jiang^{1, *}, Gongxue Wang², Jinmei Pan³, Jinyu Huang¹, Cheng Zhang¹, Huizhen Cui¹, Jianwei Yang¹, Zhaojun Zheng⁴, Shengli Wu⁴, Jiancheng Shi⁵

5 ¹State Key Laboratory of Remote Sensing Science, Jointly Sponsored by Beijing Normal University and Aerospace Information Research Institute of Chinese Academy of Sciences, Faculty of Geographical Science, Beijing Normal University, Beijing 100875, China

² College of Geomatics, Xi'an University of Science and Technology, Xi'an 710054, China

10 ³ State Key Laboratory of Remote Sensing Science, Aerospace Information Research Institute, Chinese Academy of Sciences, Beijing 100101, China.

⁴ Satellite Meteorological Institute, National Satellite Meteorological Center, China Meteorological Administration, Beijing 100081, China

⁵ National Space Science Center, Chinese Academy of Sciences, Beijing 100190, China

Correspondence to: Lingmei Jiang (jiang@bnu.edu.cn)

15 **Abstract.** Accurate long-term daily Cloud-gap-filled fractional snow cover products are essential for climate change and snow hydrological studies in the Asia Water Tower (AWT) region, but existing Moderate Resolution Imaging Spectroradiometer (MODIS) snow cover products are not sufficient. In this study, the multiple endmember spectral mixture analysis algorithm based on automatic endmember extraction (MESMA-AGE) and the multistep spatiotemporal interpolation algorithm (MSTI) are used to produce the MODIS daily cloud-gap-filled fractional snow cover product over the AWT region
20 (AWT MODIS FSC). The AWT MODIS FSC product have a spatial resolution of 0.005°, and spans from 2000 to 2022. The 2745 scenes of Landsat-8 images are used for the areal scale accuracy assessment. The fractional snow cover accuracy metrics, including coefficient of determination (R^2), root mean squared error (RMSE), and mean absolute error (MAE) are 0.80, 0.16 and 0.10, respectively. The binarized identification accuracy metrics, including overall accuracy (OA), producer's accuracy (PA), and user's accuracy (UA), are 95.17%, 97.34% and 97.59%, respectively. Snow depth data observed at 175
25 meteorological stations are used to evaluate accuracy at point scale, yielding the following accuracy metrics: an OA of 93.26%, a PA of 84.41%, a UA of 82.14%, and a Cohen's kappa (CK) value of 0.79. Snow depth observations from meteorological stations are also used to assess the fractional snow cover resulting from different weather conditions, with an OA of 95.36% (88.96%), a PA of 87.75% (82.26%), a UA of 86.86% (78.86%) and a CK of 0.84 (0.72) under the MODIS clear sky observations (spatiotemporal reconstruction based on the MSTI algorithm). The AWT MODIS FSC product can provide
30 quantitative spatial distribution information of snowpack for mountain hydrological models, land surface models, and numerical weather prediction in the Asia Water Tower region. This dataset is freely available from the National Tibetan Plateau Data Centre at <https://doi.org/10.11888/Cryos.tpd.c.272503>(Jiang et al., 2022) or from the Zenodo platform at <https://zenodo.org/doi/10.5281/zenodo.10005826>.



1 Introduction

35 Snow cover has the characteristics of high albedo, low emissivity and strong water-holding energy (Yang et al., 2014; Wang et al., 2022; Pan et al., 2023; Wang et al., 2023). The extent and variability of snow cover have profound implications for global and regional water and energy cycles (Elguindi et al., 2005; Senan et al., 2016) and climate change (Barnett et al., 2005; Li et al., 2018). The Asian Water Tower region centered on the Tibetan Plateau is the region with the largest snow accumulation outside the North Pole and South Pole (Immerzeel et al., 2020). In recent years, this region has been in a state of imbalance, 40 which is mainly reflected in the massive conversion of snow (one of the two forms of solid water) into liquid water (Yao et al., 2022). Furthermore, since 2008, the snow cover in the Asian Water Tower region has surpassed the tipping point and has become unstable (Liu et al., 2023), which had a strong and stable relationship with the changes in the Amazon rainforest ecosystem. Therefore, it is important to produce long-term series and high spatiotemporal resolution Cloud-gap-filled fractional snow cover datasets in the Asian Water Tower region.

45 Remote sensing technology has become an essential tool for monitoring snow cover globally. Polar orbit satellites such as NOAA/AVHRR, Terra/Aqua MODIS, Landsat, and Sentinel-2, which are often used to monitor snow cover, have spatial resolutions ranging from meters to kilometers. Toward the requirement of daily, large-scale, and long-time series of fractional snow cover monitoring, only moderate to coarse resolution sensors, such as AVHRR and MODIS are currently available. However, multispectral images at moderate and coarse spatial resolution have mixed pixels near the snow line, the edge zone 50 of snow patches and the forest area covered by snow (Painter et al., 2009; Pan et al., 2022; Wang et al., 2022). The classification of snow and non-snow alone will lead to significant overestimation or underestimation. Classification errors will be further transferred the subsequent applications in various fields (Wang et al., 2013; Niittynen et al., 2020; Notarnicola, 2020). The existing optical remote sensing snow cover mapping methods mainly include the reflectivity linear interpolation method (Metsämäki et al., 2012; Metsamaki et al., 2005; Wang et al., 2017), snow index empirical relationship method (Hall et al., 1995; Salomonson and Appel, 2004; Wang et al., 2021; Salomonson and Appel, 2006; Wang et al., 2020), machine 55 learning method (Dobrevá and Klein, 2011; Czyzowska-Wisniewski et al., 2015; Kuter, 2021; Xiao et al., 2022) and spectral mixture analysis method (Painter et al., 2003, 2009; Bair et al., 2021). The accuracy of the first three methods depends on training data, and the methods need to be retrained when used in different regions and on different dates. The MEAMA-AGE algorithm is a kind of automatic extraction of pure snow and non-snow endmembers based on the single-band reflectance of 60 MODIS multispectral images and the normalized differential snow index (NDSI), normalized differential vegetation index (NDVI) and normalized differential water index (NDWI), and then fractional snow cover is retrieved by the MESMA-AGE algorithm (Shi, 2012; Zhu and Shi, 2018). This algorithm can ensure the representativeness of the endmember, improve the computational efficiency and effectively adapt to the characteristics of strong topographic heterogeneity and thin and broken snow, with better accuracy and robustness than other algorithms (Hao et al., 2019; Pan et al., 2022).

65 Terra and Aqua MODIS provide two daily daytime observations, but the MODIS annual average cloud cover in the Asian Water Tower region is proximately 50% (Wang et al., 2019; Huang et al., 2022). Snow cover observations can be obscured by



clouds, resulting in many data gaps in daily snow cover products, which greatly limits the application of daily snow cover products. To improve the spatiotemporal continuity of snow cover products, researchers have proposed various spatiotemporal reconstruction algorithms, such as temporal methods (Dozier et al., 2008; Tang et al., 2017; Tran et al., 2019), spatial methods (López-Burgos et al., 2013; Shea et al., 2013; Hou et al., 2019), spatiotemporal methods (Li et al., 2017; Huang et al., 2018; Li et al., 2020; Xing et al., 2022), and multisource data fusion methods (Yang et al., 2014; Yu et al., 2016; Dai et al., 2017). Most existing algorithms were developed for binary snow cover products, and although they have good accuracy, they are difficult to apply to continuous values such as fractional snow cover. The multistep grouping algorithm used in this study is an improved spatio-temporal method that combines spatial and temporal methods through multiple step implementations (Parajka and Blöschl, 2008; Gafurov and Bárdossy, 2009; López-Burgos et al., 2013). These simple multistep combinations have been shown to be effective and efficient in cloud removal, and agree very well with *in situ* observations (Paudel and Andersen, 2011).

Currently, there are various snow cover datasets for the Asian Water Tower region, such as the Interactive Multi-sensor Snow and Ice Mapping System (IMS) (Mazari et al., 2013), MODIS/Terra Snow Cover Daily L3 Global 500 m SIN Grid product (MOD10A1) (Hall and Riggs, 2016), MODIS Snow-Covered Area and Grain size product (MODSCAG) (Painter et al., 2009), Japan Aerospace Exploration Agency (JAXA) long-term snow cover extent dataset (JASMES) (Hori et al., 2017), Northwest Institute of Eco-Environment and Resources (NIEER), Chinese Academy of Sciences AVHRR/MODIS snow cover extent product (NIEER AVHRR/MODIS SCE) (Hao et al., 2021, 2022), Tibetan Plateau long-term daily gap-free snow cover product based on the Hidden Markov Random Field model (HMRFS-TP) (Huang et al., 2022), and the European Space Agency (ESA) Snow Climate Change Initiative (Snow CCI: MODIS (Nagler et al., 2022) and AVHRR (Naegeli et al., 2022)). The IMS, JASMES, and NIEER AVHRR SCE products are binary products and have relatively coarse resolutions. The NIEER AVHRR/MODIS SCE and HMRFS-TP products are binary, and their range cannot fully cover the Asian Water Tower area. The MODSCAG product has better accuracy, but it cannot freely available. The MOD10A1 product is the most widely used, but it has data gaps, and the linear relationship between the NDSI and FSC is not always valid. Compared to MODSCAG and MOD10A1, the AWT MODIS FSC product has overall better accuracy in the AWT region (Hao et al., 2019). Although Snow CCI is a fractional snow cover product, the key parameter for retrieving canopy transmittance is calculated using static forest data from early the 2000s, which makes it difficult to capture the dynamic changes in snow cover in forest areas, and it has data gaps. Therefore, there is an urgent need for a high precision, high spatiotemporal resolution, and long-term series cloud-gap-filled fractional snow cover dataset to meet the growing demand for snow monitoring in the Asian Water Tower region.


This study used the MESMA-AGE algorithm and the MSTI algorithm to produce a MODIS long-term series daily fractional snow cover dataset for the Asian Water Tower region from 2000 to 2022. This work is organized as follows: first, the study area and datasets are presented. Then, the MESMA-AGE algorithm framework, the MSTI algorithm framework, and the data processing process are introduced. The two algorithms are used to produce a daily cloud-gap-filled fractional snow cover dataset for the Asian Water Tower region. Finally, the accuracy of this product is evaluated using high spatial resolution Landsat-8 images and meteorological station snow depth data from the China Meteorological Administration (CMA).



2 Study area and data

2.1 Study area

The Asian Water Tower region consists mainly of the Pamir Plateau, Xinjiang, and the Qinghai-Tibet Plateau in China (Fig.1). The latitude and longitude ranges are 24° - 54° N and 60° - 106° E respectively, with an average elevation of over 4000 meters. The Asian Water Tower region is the birthplace of more than 10 major rivers in Asia, sustaining nearly 2 billion people in its vicinity (Immerzeel et al., 2020; Li et al., 2022). In the Asian Water Tower region, the monthly average snowmelt runoff ratio is greater than 30% in more than half of the months, far exceeding the surrounding area (Yang et al., 2022). Over the past 50 years, the temperature in Asia Water Tower region has increased by an average of 0.3 to 0.4 °C per 10 years, which is twice the global average rate (Barnett et al., 2005; Kraaijenbrink et al., 2017). As one of the most important climate response factors (Liu and Chen, 2000; Immerzeel et al., 2010), the distribution and change in snow cover is of great importance for the study of climatic and ecological changes across the region. Meanwhile, fractional snow cover data are important input for the Snowmelt Runoff Model (SRM) (Martinec, 1975) and can also be used for snow water equivalent reconstruction and optimization (Rittger et al., 2016). Therefore, a set of high-precision fractional snow cover products is necessary for hydrological simulation and hydrological applications in the Asian Water Tower region.



115

Figure 1: DEM (a) and Land cover (b) maps of the AWT with the positions of the MODIS tiles, Landsat scenes and CMA stations used in the validation



2.2 MODIS surface reflectance data

This study used MODIS surface reflectance products MOD09GA and MYD09GA in Collection 6 spanning from 2000 to 2022. These surface reflectance products have two data layers. The 500 m reflectance data layer provides reflectance, quality assessment level, observation area, observation number, and 250 m scan information for bands 1-7. The 1 km geographic information data layer provides additional information, such as observation times, quality assessment levels, sensor azimuth zenith angles, solar azimuth altitude angles, and orbit pointers. In addition, it also includes metadata information of the file (production information, geographical scope, etc.). The characteristics of MODIS solar reflective bands are shown in table 1. The cloud information used in this study was obtained from the 'state_1km' layer, which includes 'cloud state' is not a clear and 'cirrus detected' is a high. Moreover, 12 MODIS tiles ('h23v03', 'h23v04', 'h23v05', 'h24v04', 'h24v05', 'h24v06', 'h25v04', 'h25v05', 'h25v06', 'h26v05', 'h26v06', and 'h27v06') with sinusoidal projection were used in this study, as shown in Figure 1 (a).

Table 1: MODIS spectral characteristics

Band Name	Spectral Range(μm)	Central Wavelength(μm)	Spatial Resolution(m)
band1	0.62-0.67	0.645	500
band2	0.841-0.876	0.858	500
band3	0.459-0.479	0.469	500
band4	0.545-0.565	0.555	500
band5	1.23-1.25	1.24	500
band6	1.628-1.652	1.64	500
band7	2.105-2.155	2.13	500

2.3 Landsat-8 images

This study used the Google Earth Engine (GEE) cloud platform to select a total of 2745 scenes of Landsat-8 images from 2013 to 2021 that met the cloud coverage ratio of less than 10% and snow coverage ratio of more than 30% as "ground truth" to validate our fractional snow cover product. Landsat-5 Thematic Mapper (TM) has obvious attenuation since 2000, and the Landsat-7 Enhanced Thematic Mapper Plus (ETM+) sensor has been affected by striping in 25% of the image area due to scanner failure since June 2003. Therefore, this study mainly focused on Landsat-8 images to evaluate the accuracy of the AWT MODIS FSC dataset. To better evaluate the MESMA-AGE algorithm and the AWT MODIS FSC product, this study also applied the MESMA-AGE algorithm to retrieve Landsat-8 fractional snow cover, which has been demonstrated to have good accuracy on Landsat-8 using higher resolution Gaofen-2 imagery with an OA of 94.46% and RMSE of 0.094 (Hao et al., 2019). The Landsat-8 fractional snow cover results at 30 m were resampled to the resolution of the AWT MODIS FSC product (0.005°) through aggregation and averaging. Subsequently, the Landsat fractional snow cover results at 0.005° resolution were used to assess the accuracy of the MODIS clear sky retrieval results in the AWT MODIS FSC product.



2.4 Ground snow-depth measurements

As Landsat images can only assess the accuracy under clear sky, this study chose to use snow depth data from meteorological stations to support the validation of the accuracy of the spatio-temporal reconstruction results under cloud cover. This study used a total of 175 *in situ* stations provided by the China Meteorological Administration in the Asian Water Tower area from 26 February 2000 to 30 April 2019, as shown in Figure 1b. Figure 1b shows that the *in situ* stations are more evenly distributed in the southeast of the Qinghai Tibet Plateau, Tianshan Mountains and Altay Mountains, where seasonal snow is prevalent. Snow depth data are measured in an open field at 8:00 am using a professional meter ruler. If the fractional snow cover is greater than 50% and the snow depth is greater than 1 cm, it is considered snow and recorded. The geographical coordinates, time of observation, and snow pressure are also recorded. In this study, a threshold of 3 cm was used to classify snow using *in-situ* snow depth data (Huang et al., 2022), where a snow depth less than 3 cm was reclassified as no snow; otherwise, it was recognized as snow. To further illustrate the accuracy of snow identification, this study excluded stations with snow depths greater than 1 cm but snow cover days less than 20 (Zhang et al., 2020; Hao et al., 2021).

2.5 Auxiliary data

To better evaluate the accuracy of the MESMA-AGE algorithm and the AWT MODIS FSC product, auxiliary information, such as elevation and the land cover type of the Asian Water Tower, were used. The GEE cloud platform provided the MCD12Q1 V6.1 annual International Geosphere-Biosphere Programme (IGBP) classification data (Sulla-Menashe et al., 2019). The surface types were further divided into four categories: bare land, grassland, forest, and plateau mountain. The GEE cloud platform was utilized to obtain Shuttle Radar Topography Mission (STRM) digital elevation model (DEM) data. The DEM data were then resampled from 90 m to the 0.005° resolution of the AWT MODIS FSC product (Reuter et al., 2007).

3 Methodology

Figure 2 shows the flowchart of the AWT MODIS FSC production. According to the accuracy evaluation of the MOD10A1, MODSCAG, and MODAGE fractional snow cover products in the Qinghai Tibet Plateau region, the MODAGE product had the highest accuracy (Hao et al., 2019). Therefore, the MODAGE fractional snow cover retrieval algorithm (MESMA-AGE algorithm) was selected for the fractional snow cover retrieval of Terra and Aqua MODIS surface reflectance version 6 data in the Asian Water Tower region. Second, based on the Terra/MODIS fractional snow cover retrieval results, the Aqua/MODIS fractional snow cover retrieval results were used to fill in data gaps due to clouds and missing observations (Li et al., 2014). Third, the Geospatial Data Abstraction Library (GDAL) was used to reproject and mosaic the fractional snow cover retrieval results of 12 MODIS tiles (GDAL Development Team, 2022). Fourth, the MSTI algorithm was developed for performing spatiotemporal interpolation on pixels with cloud cover or missing data, enabling the generation of a daily cloud-gap-filled fractional snow cover product. Finally, accuracy evaluation and algorithm optimization of the MESMA-AGE algorithm and the AWT MODIS FSC product were performed using snow depth data from meteorological stations and Landsat-8 imagery.




Figure 2: Overall flowchart of the AWT MODIS FSC product

175 3.1 MESMA-AGE algorithm

When a pixel contains information from multiple surface types, it is called a mixed pixel, whereas a pixel containing only one type of ground object can be called an endmember of that surface type. The algorithm for unmixing mixed pixels is mainly based on the linear combination of the spectral information of the endmember (Roberts et al., 1998). To analyze the spectral information combination of a pixel, a linear spectral mixing analysis model can be used, which assumes that different endmember energies only undergo single scattering mixing and that there is no nonlinear mixing process (Painter et al., 2003).
 180 The linear spectral mixing analysis expression and constraints can be expressed as Equations 1-3:

$$R_{\lambda} = \sum_{i=1}^N F_i R_{i,\lambda} + \varepsilon_{\lambda} \quad (1)$$

$$\sum_{i=1}^N F_i = 1 \quad (2)$$

$$F_i \geq 0 \quad (3)$$



Painter et al. (2003) established an endmember spectral library by collecting spectra of various types of vegetation, rocks, soils, and lake ice from the field and lab and optimized the endmember metadata of snow cover of different grain sizes using radiative transfer models. They then used this spectral library with MODIS images to produce the MODIS Snow Covered Area and Grain Size algorithm (MODSCAG)(Painter et al., 2009). Due to the phenomenon of “the same object with different spectra”, spectra from limited observation conditions in the field and lab have difficulty representing the actual complex surface. Meanwhile, the spectrum simulated by the Mie/DISORT model can represent the reflection characteristics of snow under different snow properties and observation conditions, but it is also susceptible to the simulation errors of the model itself. In this study, the MEAMA-AGE algorithm was used to retrieve the fractional snow cover in the Asian Water Tower region, which combines an image-based automatic endmember extraction algorithm(Shi, 2012) with a spectral library optimization method (Xu et al., 2015). The MESMA-AGE algorithm can improve the computational efficiency while ensuring the representativeness of the endmembers(Hao et al., 2019). The rules for extracting snow and non-snow endmembers are shown in Table 2.

Table 2. Endmember extraction rule of the MESMA-AGE algorithm

End-Member	Rule for MODIS Surface Reflectance data
Snow	$NDSI > 0.75$ & $NDVI < -0.035$ & $R_{0.55} > 0.7$
Vegetation	$NDSI < -0.4$ & $NDVI > 0.7$
Soil/rock	$NDSI < -0.4$ & $0 < NDVI < 0.15$
Waterbody	$NDWI > 0.2$ & $R_{0.86} < 0.2$




Figure 3: The NDSI and NDVI pattern of vegetation, soil/rock, and snow endmembers from MODIS images



For this study, 8893 samples were selected from the MODIS data in January, November, and December 2020. The sample types include snow (3136 samples), vegetation (1116 samples), and soil (4641 samples). The samples were used to create the ground object feature map (Fig. 3). The x-axis in Fig. 3 is the NDSI value, and the y-axis is the NDVI. From Figure 3, snow, vegetation, and bare ground pixels can be easily located in the NDSI-NDVI feature space. The endmember extraction rules in Table 2 can effectively achieve endmember extraction for different land types.

3.2 Multistep spatiotemporal interpolation algorithm

This study developed a multistep cloud removal algorithm that combines temporal and spatial information. The MSTI algorithm prioritizes the use of nearby spatiotemporal information based on the characteristics of snow cover and achieves complete cloud removal for still cloudy pixels by further expanding their spatiotemporal range. This algorithm is mainly divided into four steps: temporal filtering with a 3×3 temporal window, 4×4 spatial interpolation, piecewise cubic hermite interpolating polynomial (PCHIP) for the 19-day period, and further spatial interpolation using a 10×10 window. The process is shown in the MSTI algorithm flowchart in Figure 2.

1) The temporal filtering algorithm assumes that FSC does not change during a short period. (Hou et al., 2019). In previous studies, the size window of the adjacent time filter ranged from 1 d to 8 d. Due to the unique climate conditions and terrain conditions of the Asian Water Tower (high wind speeds can easily redistribute snow, and thin layers of snow can melt and sublime quickly), the snow cover changes rapidly. Therefore, choosing a longer time window may introduce errors. In this study, the time window of the adjacent time filtering algorithm was set to 3 d (the day of cloud cover, and the day before and after cloud cover). If a given pixel is covered by clouds and there are no clouds before and after two days, the FSC value of the cloud pixel can be calculated using the following formula:

$$FSC_{predict_{cloud}}^T(x, y) = (FSC_{observed_{cloud-free}}^{T-1}(x, y) + FSC_{observed_{cloud-free}}^{T+1}(x, y))/2 \quad (4)$$

where $FSC_{predict_{cloud}}^T(x, y)$ is the predicted FSC value of cloud pixels (x, y) at time T, $FSC_{observed_{cloud-free}}^{T-1}(x, y)$ is the observed FSC value of cloud-free pixel (x, y) at time T-1, and $FSC_{observed_{cloud-free}}^{T+1}(x, y)$ is the observed FSC value of cloud-free pixel (x, y) at time T+1.

2) Based on the continuity of snow cover in spatial continuity, snow cover can be interpolated based on information from non-cloud pixels around a cloud pixel (Gafurov and Bárdossy, 2009; Paudel and Andersen, 2011; Lindsay et al., 2015). Considering the situation where there are at least 3 identical pixels in the 4 pixels adjacent to a cloud pixel, the cloud pixel discrimination rule is as follows: if at least 3 of the 4 pixels above, below, left, and right of a cloud pixel are covered with snow, the central cloud pixel is assigned the mean FSC of the snow pixels in the adjacent 8 pixels. If at least 3 out of the 4 pixels above, below, left, and right of a cloud pixel are land, the central cloud pixel is assigned as land. In other cases, the pixels retain their cloud pixel values.

3) For the remaining cloud pixels after the previous two steps, the PCHIP algorithm is used to interpolate the time series of the missing data. Compared with the spline curve used in previous studies (Dozier et al., 2008; Tang et al., 2013), the PCHIP



algorithm ensures the monotonicity of the interpolation result, which allows it to achieve spatiotemporally continuous fractional snow cover while suppressing the effects of noise. Through statistical analysis of cloud cover in the Asian Water Tower region over the past approximately 20 years, 78.92% of the regions has less than 16 days of cloud cover, and 93.14% has less than 19 days of cloud cover. Therefore, the time window selected for this study was set to 9 days before and after the presence of cloud cover pixels.

The PCHIP algorithm assumes that the known function $f(x)$ satisfies $f(x_i) = f_i$ and $f'(x_i) = f'_i$ ($i = 0, 1, 2, \dots, n$) at $n+1$ distinct nodes x_i ($i = 0, \dots, n$) on the interpolation interval $[a, b]$. A segmented cubic Hermite interpolation function $G(x)$ can be constructed to satisfy Equations (5), (6) and (7).

The polynomial degree of $G(x)$ between each cell is 3 (5)

$$G(x) \in C^1[a, b] \quad (6)$$

$$G(x_i) = f(x_i), G'(x_i) = f'(x_i), i = (0, 1, \dots, n) \quad (7)$$

The expression of $G(x)$ between cells $[x_k, x_{k+1}]$ can be directly obtained from the above conditions:

$$G(x) = \left(1 + 2 \frac{x - x_k}{x_{k+1} - x_k}\right) \left(\frac{x - x_{k+1}}{x_k - x_{k+1}}\right)^2 y_k + \left(1 + 2 \frac{x - x_{k+1}}{x_k - x_{k+1}}\right) \left(\frac{x - x_k}{x_{k+1} - x_k}\right)^2 y_{k+1} + (x - x_k) \left(\frac{x - x_{k+1}}{x_k - x_{k+1}}\right)^2 y'_k + (x - x_{k+1}) \left(\frac{x - x_k}{x_{k+1} - x_k}\right)^2 y'_{k+1} \quad (8)$$

where x_k and x_{k+1} are the positions of two adjacent time points to be interpolated, y_k and y_{k+1} are the FSC corresponding to the two observations before and after the corresponding interpolation point, and y'_k and y'_{k+1} are the corresponding derivatives.

4) After the first three steps of spatiotemporal interpolation, there are still a few cloud pixels left. In this study, the observation information from the 11*11 interpolation window centred on the cloud pixel was used based on the inverse distance weight (IDW) interpolation algorithm, which considers elevation information for spatial interpolation. The IDW interpolation algorithm is an important application of the first law of geography, which uses the distance between the interpolation point and the sample point as the weight for weighted averaging (Zhao et al., 2022). The closer the interpolation point is, the greater the weight assigned to the sample point. According to existing studies, elevation is important for the distribution of fractional snow cover (Li et al., 2017), but traditional IDW algorithms only consider spatial distance. Therefore, this study incorporated the influence of elevation on fractional snow cover on this basis. The SNOWL method is a commonly used algorithm for spatiotemporal interpolation of snow cover, and scholars often use 100 m as the interval (Huang et al., 2016; Li et al., 2017). Therefore, this study mainly used clear sky pixel information within the range of elevation differences less than 100 m around the pixels. The process of the IDW interpolation algorithm considering elevation information is described in Eq. (9), Eq. (10) and Eq. (11):

$$d_i = \sqrt{(x - x_i)^2 + (y - y_i)^2} \quad (9)$$



$$w_i = \frac{\Delta E_i/d_i}{\sum_1^n \Delta E_i/d_i}, \Delta E_i = \begin{cases} 1 - \frac{|\Delta Elev_i|}{100}, & |\Delta Elev_i| \leq 100 \text{ m} \\ 0, & |\Delta Elev_i| > 100 \text{ m} \end{cases} \quad (10)$$

$$FSC(x, y) = \sum_{i=1}^n w_i * FSC(x_i, y_i) \quad (11)$$

where (x, y) is the position of the cloud pixel, (x_i, y_i) is the observing pixel positions for the surrounding clear sky pixels, d_i is the distance between the sample point and the position to be interpolated, ΔE_i is the weight of the i -th sample point obtained based on elevation, $\Delta Elev_i$ is the elevation difference, w_i is the weight of the i -th sample point, $FSC(x_i, y_i)$ is the fractional snow cover for clear sky pixels, and $FSC(x, y)$ is the interpolated fractional snow cover for the cloud pixel.

3.3 Evaluation Metrics

Selected metrics for validation of the AWT MODIS FSC product included the OA, PA, UA, CK, R^2 , RMSE, and MAE, which are defined below:

$$OA = \frac{TP + TN}{TP + TN + FP + FN} \quad (12)$$

$$PA = \frac{TP}{TP + FP} \quad (13)$$

$$UA = \frac{TP}{TP + FN} \quad (14)$$

$$CK = \frac{OA - P}{1 - P}, P = \frac{(TP+FN)(TP+FP)+(TN+FN)(TN+FP)}{(TP+TN+FP+FN)^2} \quad (15)$$

$$R^2 = \frac{[\sum(f_{est} - \overline{f_{est}})(f_{ref} - \overline{f_{ref}})]^2}{\sum(f_{est} - \overline{f_{est}})^2 \cdot \sum(f_{ref} - \overline{f_{ref}})^2} \quad (16)$$

$$MAE = \frac{1}{N} \sum_N |f_{est} - f_{ref}| \quad (17)$$

$$RMSE = \sqrt{\frac{1}{N} \sum_N (f_{est} - f_{ref})^2} \quad (18)$$

where TP indicates true positive, TN indicates true negative, FP indicates false positive, FN indicates false negative, f_{est} indicates fractional snow cover estimation derived from MODIS, and similarly, f_{ref} indicates reference fractional snow cover derived from Landsat-8. To calculate the overall accuracy, binary snow cover was labeled for pixels with fractional snow cover $\geq 15\%$ (Rittger et al., 2013; Wang et al., 2019).



4 Results

265 In this study, a comprehensive evaluation of the accuracy of the AWT MODIS FSC product in two dimensions, namely
binary and fractional snow cover, was conducted. The binary and fractional snow cover accuracies of the AWT MODIS FSC
product were quantitatively assessed under three different conditions: overall, different surface types, and altitudes. This
evaluation was performed using 2745 Landsat-8 images. Additionally, the binary accuracies of the AWT MODIS FSC product
were specifically evaluated in three different conditions, i.e., overall (including clear and cloud conditions), clear sky, and
270 cloud cover, utilizing snow depth data from 175 meteorological stations.

4.1 Validation with Landsat-8 images

4.1.1 Overall results

In this study, the 2745 Landsat-8 scenes were used as "ground truth" to quantitatively evaluate the AWT MODIS FSC
product obtained from clear sky in two dimensions: binary (OA, PA, and UA) and fractional snow cover (R^2 , MAE, and
275 RMSE). Figure 4 shows violin charts of the accuracy evaluation metrics, and the violin charts of each accuracy metric are
composed of two parts: the outer violin chart and the inner box plot. The left side of the outer violin chart is the kernel density
map, the larger the area of a certain range is; the greater the probability of the distribution near a certain value, and the
horizontal line in the left area is where the median is located. To the right of the violin plot is a histogram of the frequency of
a value, i.e., the longer the line is, the more points there are for that value. The internal box plot contains a gray rectangle
280 consisting of the upper and lower quartiles, with the mean position represented by the white point. From Figure 4, the minimum
OA value of the AWT MODIS FSC product is 80.38%, and the average value is 95.17%. The minimum PA value is 58.71%,
and the average value is 97.34%, i.e., the average omission error of this product is 2.66%. The minimum UA value is 67.02%,
and the average value is 97.59%, i.e., the average commission error is 2.41%. The R^2 value distribution range is 0.40-0.97, and
the average value is 0.80. The MAE ranges from 0.01 to 0.23, with an average of 0.10. The RMSE ranges from 0.02 to 0.26,
285 with an average of 0.16. The above results provide a good illustration of the consistency of the AWT MODIS FSC product
with the "ground truth" data.




Figure 4: Violin charts for the accuracy evaluation metrics of the AWT MODIS FSC product validated by Landsat images

To better explore the interannual differences in each accuracy metric, this study divided the 2745 Landsat-8 scenes by year. Figure 5 shows the interannual distribution of each accuracy metric, and Table 3 shows the number of Landsat images and the interannual average of each accuracy metric. Figure 5 and Table 3 show that the interannual means of the OA range from 92.41% to 96.22%, PA from 94.77% to 97.88%, UA from 95.35% to 98.55%, R^2 from 0.76 to 0.81, MAE from 0.09 to 0.11, and RMSE from 0.15 to 0.17. The accuracy metrics perform better except for 2013, where the poor accuracy indicators are mainly due to the overall low number of validation data and the fact that a significantly larger proportion of the validation data are located at high altitudes in mountainous areas than in other years. The results of this part of the retrieval are strongly influenced by topographical effects.




Figure 5: Violin chart for the interannual accuracy evaluation metrics



Table 3. The number of Landsat images and the interannual average of each accuracy metric

Year	Image Number	OA	PA	UA	R ²	MAE	RMSE
2013	123	92.41%	94.77%	95.35%	0.81	0.11	0.17
2014	328	95.60%	97.67%	98.38%	0.79	0.11	0.16
2015	287	95.08%	97.26%	97.58%	0.79	0.10	0.16
2016	309	96.22%	97.88%	98.55%	0.81	0.09	0.15
2017	315	95.85%	97.25%	98.27%	0.79	0.09	0.15
2018	389	95.16%	97.30%	97.95%	0.76	0.11	0.17
2019	360	95.23%	97.32%	97.37%	0.79	0.10	0.16
2020	320	95.01%	97.21%	97.53%	0.81	0.10	0.16
2021	314	94.20%	97.33%	95.07%	0.81	0.10	0.17

300 4.1.2 Evaluation for different land surface types

Spectra of snow-covered surface also encounters the impacts of land cover type. In particular, in forested areas, snow below the forest canopy is difficult to observe with spaceborne sensors because the forest blocks the visible, near-infrared, and shortwave infrared bands (Wang et al., 2021, 2023). Therefore, in this study, 2745 scenes of Landsat-8 images were divided into four categories according to land surface types, namely, grassland area (1500 scenes), bare land area (264 scenes), forest area (410 scenes), and mountain area of the Himalayas and Pamir Plateau (571 scenes). The accuracy evaluation metrics under the two dimensions of binary value and fractional snow cover of the four land surface types are shown in Figure 6. Regarding the accuracy metrics of binarization, the accuracy of the grassland area is better than that of other land surface types, and its average OA is 95.99%, while the bare land area is the worst but also reaches 93.69%. This is because the bare land area is mainly the hinterland of the Asian Water Tower region, which is mainly desert/Gobi with little vegetation growth, and there are also many large and small lakes distributed in this area. The bright land and the winter water surface lead to deviations in the retrieval algorithm. In addition, the snow in this region is relatively broken, and the observation scales of MODIS and Landsat are quite different, making it difficult for MODIS to capture the broken snow information as effectively as Landsat. These two reasons lead to many errors in the results. The UA also illustrates this problem well. Figure 6 shows that the average UA of bare land is only 94.06%, approximately 4% lower than that of the other surface types. From the perspective of fractional snow cover accuracy metrics, grassland and forest are slightly worse, mainly because it is difficult to observe the snow signal shielded by the vegetation canopy at the MODIS scale.




Figure 6: Violin charts for accuracy evaluation metrics of the AWT MODIS FSC product validated by Landsat images under different surface types

320 4.1.3 Evaluation for different altitudes

Topographic effects challenge accurate snow cover mapping with optical imagery as well. The snow cover products such as MOD10A1 have been reported uncertainties related with altitudes (Zhang et al., 2020; Wu et al., 2021; Huang et al., 2022). Therefore, the 2745 Landsat-8 images are divided into four sections according to altitude, namely < 3 km (1603 scenes), 3-4 km (395 scenes), 4-5 km (355 scenes) and > 5 km (392 scenes). The results of the accuracy evaluation metrics according to different heights are shown in Figure 7. As shown in Figure 7, both binary and fractional snow cover accuracy metrics show a decreasing trend with increasing altitude. The areas smaller than 3 km are mostly distributed in northern Xinjiang, China, i.e., the area north of 40°N, where the snow distribution is relatively concentrated, and the surface type is mostly grassland with a small amount of forest, so the accuracy is highest. The three elevation regions greater than 3 km are mainly distributed in the Tianshan Mountains, the Pamir Plateau and the Tibetan Plateau. Snow fragmentation and topographic heterogeneity in these regions increase with altitude. This results in a slight reduction in the accuracy of the AWT MODIS FSC product.




Figure 7: Violin chart for accuracy evaluation metrics of the AWT MODIS FSC product validated by Landsat under different altitudes

4.2 Validation with *in situ* snow depth measurements

Landsat images can only be used to evaluate the accuracy of the fractional snow cover retrieval algorithm and FSC product under clear sky conditions, and the fractional snow cover information reconstructed by the MSTI algorithm needs to be verified by snow depth observations at meteorological stations. Therefore, this study used a total of nearly 1 million observations collected from 175 *in situ* stations during the period from 26 February 2000 to 30 April 2019 to evaluate the accuracy under different weather conditions. The numbers of meteorological stations and SD observations obtained each year are shown in Figure 8. The number of stations fluctuates to some extent each year, and the number of snow depth observations available after 2014 has nearly tripled compared with before 2014.




Figure 8: The number of *in situ* stations and observed data volume per year (3.12, 2000-4.30, 2019)

4.2.1 Overall results

To better assess the accuracy of snow identification, only *in situ* stations with more than 20 days of snow observations per year were selected for evaluation (Zhang et al., 2020; Hao et al., 2021). This reduced the number of *in situ* stations and the total observation data by approximately half. In this study, snow depth observations from 175 *in situ* stations were used to perform a binary evaluation of the AWT MODIS FSC product. Table 4 shows the results of the overall accuracy evaluation. It can be seen from the results that the OA of the product reaches 93.26%, PA can reach 84.41%, and UA can reach 82.14%, i.e., the omission error is 15.59%, and the commission error is 17.86%. In addition, CK reach 0.79. The results of the above accuracy metrics exclude the stations without snow observations, which indicates that the AWT MODIS FSC product has good accuracy and good consistency with the snow depth observation data of meteorological stations.

Table 4. Confusion matrix and accuracy results of the AWT MODIS FSC product based on snow depth measurements from CMA. OA, PA, UA and CK

		AWT MODIS FSC	
Class		Snow	Non-Snow
<i>In situ</i> snow depth measurements	Snow	102617	18946
	Non-Snow	22316	468105
	OA	93.26%	
	PA	84.41%	
	UA	82.14%	
	CK	0.79	

To verify the stability of the product accuracy over time, this study performed a binary accuracy assessment of the snow depth observations at each station by year. The overall results of each accuracy metric over the last 20 years are shown in Figure 9. Each accuracy metric is relatively stable before and after 2014, but there is a large fluctuation in 2014. The OA metric exhibits the most significant temporal variation. Before 2014, the fluctuation range of OA is 88.69%-92.96%, and after 2014,



the fluctuation range of OA is 95.05-97.54%. Meanwhile, CK and PA increase significantly after 2014. This also indicates that the consistency between the AWT MODIS FSC product and the snow depth observations from meteorological stations has improved significantly since 2014. The fluctuation in the above accuracy indicators is mainly due to the significant increase in the number of meteorological station observations used in this study after 2014 and the improvement in the accuracy of snow identification, which ultimately leads to a significant increase in OA. The percentage of cloud cover in different years is also shown in Figure 9 below. Combined with Figures 8 and 9, the increase in the amount of station data and observations acquired in this study, coupled with the decrease in cloud cover in the study area, are two factors that lead to better results in the station-based accuracy assessment after 2014 than before.




Figure 9: Accuracy fluctuations of the AWT MODIS FSC product based on *in situ* snow-depth measurements and the percentage of cloud cover in the past 20 years

4.2.2 Accuracy metrics at each *in situ* station

Figure 10 shows the detailed results of the accuracy metrics of the AWT MODIS FSC product verified by the snow depth data of the stations. As shown in Figure 10, the OA of most *in situ* stations is above 90%, with only one *in situ* station below 70%. However, the figure shows that, unlike the OA, the accuracy of the entire Asia Water Tower region is relatively consistent. PA, UA and CK are severely affected by the region. The PA and UA metrics at stations in northern Xinjiang, China, are generally greater than 90%, and CK is also greater than 0.8. This is mainly due to the stable snow cover in the region. The spatiotemporal reconstruction algorithm of snow cover developed in this research can well grasp the spatiotemporal variation characteristics of snow cover in this region, so that high-precision spatiotemporal reconstruction of snow cover information can be achieved. However, the snow cover in the eastern part of the Asia Water Tower region and the northwestern edge of the Tarim Basin is relatively broken, and the MODIS resolution is coarse. These areas are seriously affected by clouds, so the PA, UA and CK metrics in these areas are generally not high.




Figure 10: Point-based accuracy results of the AWT MODIS FSC product: (a) OA; (b) PA; (c) UA; (d) CK.

4.2.3 Performance of spatiotemporal reconstruction algorithm

The above two sections presented the overall accuracy of the AWT MODIS FSC product using snow depth data from meteorological stations. The AWT MODIS FSC product is derived from the composition of two parts: the real MODIS observation under clear sky and the spatiotemporal reconstruction with the MSTI algorithm for cloudy conditions. To further explore the accuracy of the fractional snow cover results of these two parts, the snow depth observation data of the meteorological stations are divided into two categories based on MODIS clear sky and cloudy conditions. First, the stability of the accuracy evaluation metric under clear sky and cloudy conditions is evaluated, respectively and the results are shown in Figure 11. Comparison of Figure 10 shows that there is an increase in accuracy in years with lower cloud cover. Figure 11 (a) presents the interannual variation results of the accuracy metrics of fractional snow cover obtained from MODIS clear sky observations using snow depth observations at the selected meteorological stations. OA, PA, UA and CK all exhibit good stability, and the range of variation in OA is 92.80-99.01%. The range of UA is 82.39-90.26%. After 2014, the two indices of PA and CK improved, with the maximum of PA reaching 95.46% and CK reaching 0.91. Figure 11 (b) shows the interannual variation results of the accuracy metrics of the fractional snow cover result obtained by the spatiotemporal reconstruction of the MSTI algorithm using the snow depth observations at the selected meteorological stations. All accuracy metrics decrease



to some extent compared to the clear sky condition. Influenced by the amount of *in situ* station data used, the spatiotemporal reconstruction results show a relatively obvious jump in 2014. The variation range of OA before 2014 is 84.51-90.00%, but the variation range of OA after 2014 is 92.76-95.60%. PA and CK have interannual variations in the years before 2014, but the value of the years after 2014 has a large increase, with the maximum PA reaching 89.61% and CK reaching 0.83. UA has only a large interannual variation, and there is no significant jump in 2014. The results show that the accuracy of the fractional snow cover based on clear sky observations is significantly better than that of the spatiotemporal reconstruction. This is mainly due to the presence of clouds over a long period of time and over a large area in most of the Asian Water Tower region, the interpolation of this part of the area relies heavily on the last two steps of the MSTI algorithm. The last two steps of the MSTI algorithm require a larger space-time window to complete the interpolation. However, a larger space-time window introduces more error, especially for snow cover, which has strong spatial heterogeneity and changes rapidly over time.





Figure 11: Point-based accuracy results of the AWT MODIS FSC product: (a) clear sky observations; (b) spatiotemporal reconstruction.

This study further analyzed the accuracy of the AWT MODIS FSC product obtained from MODIS clear sky and cloud cover observations at each station, and the results of the binarization accuracy metrics are shown in Figure 12. It can be seen from the figure that the accuracy of fractional snow cover obtained by the clear sky retrievals is significantly better than that obtained by the MSTI algorithm. According to the OA in Figure 12 (a) and (e), the accuracy of the verification results of fractional snow cover based on MODIS clear sky observations is good at all stations, and only a few stations are less than 90%. However, the OA of the fractional snow cover reconstructed by the MSTI algorithm shows some regional differences. When comparing the accuracy metrics (PA, UA, and CK) of fractional snow cover between the real MODIS observations and the spatiotemporal reconstruction results achieved through the MSTI algorithm, notable regional variations are observed in all three metrics. In other words, the accuracy of the stable snow cover area in northern Xinjiang, China is obviously better than that in the central and eastern parts of the Asia Water Tower region and the northwestern edge of the Tarim Basin, where the snow cover is relatively fragmented and rapidly changing.



420

Figure 12: Point-based accuracy results of the AWT MODIS FSC product under clear sky conditions ((a)OA, (b)PA, (c)UA, and (d)CK) and spatiotemporal reconstruction ((e)OA, (f)PA, (g)UA, and (h)CK)



5 Discussion

5.1 Comparing AWT MODIS FSC with MOD10A1 and HMRFS-TP product

425 In order to evaluate the accuracy of AWT MODIS FSC products more objectively, MOD10A1(Hall and Riggs, 2016) and
HMRFS-TP(Huang et al., 2022) products are selected as benchmarks for this study using Landsat-8 imagery. MOD10A1 is
the most widely used MODIS snow product, which has a long time series (since 2000) and highly spatial and temporal
resolution (i.e., 500 m and daily). This study is based on the GEE platform to obtain the MOD10A1 (Collection 61) data from
2013 to 2022 for the Asian water tower region, and uses this data product to evaluate the accuracy difference with the AWT
430 MODIS FSC product under the clear sky scenario. The HMRFS-TP product is a continuous spatio-temporal binary snow
product based on the MOD10A1 product and the HMRFS spatio-temporal interpolation algorithm covering the Tibetan Plateau
region within China, and is used in this study to compare the accuracy of the two sets of continuous spatio-temporal products.

The accuracy of the MOD10A1 product depends on the threshold value of NDSI, and the commonly used thresholds are 0.1
(Zhang et al., 2019), 0.29 (Zhang et al., 2021; Tang et al., 2022) and 0.4 (Riggs et al., 2017). Since no spatio-temporal
435 interpolation is performed for the MOD10A1 product, in order to make a comprehensive and objective comparative assessment,
this study uses 1805 Landsat images to compare the clear sky pixels of the two products, and the results are shown in Figure
13 below. The OA, PA and UA of the AWT MODIS FSC product are 97.69%, 98.73% and 98.83% respectively. The NDSI
threshold is 0.1, the OA, PA and UA of the MOD10A1 products are 94.51%, 95.70% and 99.04%. The NDSI threshold is 0.29,
the OA, PA and UA of the MOD10A1 products are 94.75%, 98.49% and 96.52%. The NDSI threshold is 0.4, the OA, PA and
440 UA of the MOD10A1 products are 91.86%, 98.96% and 90.82%. As the NDSI threshold increases, Figure 13 shows that the
PA of the MOD10A1 product gradually increases and the UA gradually decreases. This also means that the percentage of
omission error is decreasing and the percentage of commission error is increasing. This is because the larger the NDSI, the
higher the probability that the image pixel is snow, and the probability of correctly judging snow increases accordingly, which
also leads to a lower probability of correctly judging non-snow, resulting in a decrease in overall accuracy. From the results in
445 Figure 13, the MOD10A1 product has the best accuracy when the NDSI is 0.29, but its accuracy is still lower than that of the
AWT MODIS FSC product.




Figure 13: Violin charts for the accuracy evaluation metrics of the AWT MODIS FSC and MOD10A1 product validated by Landsat images



450

The NDSI of the HMRFS-TP product is 0.4 as the threshold for snow identification of MOD10A1, and the spatio-temporal continuous product is obtained by the HMRFS spatio-temporal interpolation algorithm. In this study, 372 Landsat images were used to quantitatively evaluate and compare two sets of spatio-temporal continuous snow products, and the results are shown in Figure 14 below. As shown in Figure 14, the OA, PA and UA of the AWT MODIS FSC product are 89.71%, 94.29% and 86.21%. The OA, PA and UA of the HMRFS-TP product are 79.45%, 99.20% and 66.82%, respectively. Comparing the accuracy indices of the two sets of products, AWT MODIS FSC products are significantly better than HMRFS-TP products, and various accuracy evaluation indices are around 90%. The poor accuracy of the HMRFS-TP products is mainly due to the value of NDSI. Combined with Figure 13, the threshold of 0.4 will lead to serious misclassification of products and is not applicable to the Tibetan Plateau region.

455

460




Figure 14: Violin charts for the accuracy evaluation metrics of the AWT MODIS FSC and HMRFS-TP product validated by Landsat images

5.2 Area differences in fractional snow cover and binary snow cover


Long-term series and high precision fractional snow cover products are of great importance for snow hydrology research in the Asia Water Tower region. However, most of the existing snow cover products are binary products or space-time gap fill fractional snow cover products. In this study, the clear sky fractional snow cover was retrieved by the MEASMA-AGE algorithm based on MODIS observations. The missing fractional snow cover information caused by cloud cover was reconstructed by the MSTI algorithm, and finally, the spatiotemporally continuous long-term series AWT MODIS FSC product was obtained. The actual snow distribution, which is difficult to capture, was identified with binary values, especially in scenarios with mixed pixels at medium and coarse resolutions, and the subsequent direct application of binary products will introduce large errors. Therefore, the AWT MODIS FSC product produced in this study was used to quantitatively analyze the actual difference between the binary snow cover product and the fractional snow cover product, and the results are shown in Figure 15. In this study, pixels with $FSC > 15\%$ were identified as the binary snow product (Rittger et al., 2013; Wang et al., 2019). Figure 15 (a) shows the difference between the total snow cover area obtained by the binary snow product and that obtained by the fractional snow cover product in the Asian Water Tower region. There is a significant overestimation of the

470

475



480 binary snow product, with an average difference of 39400 km² and a maximum difference of 102,000 km², which is very large for the Asia Water Tower region with a total area of only 623,000 km². Figure 15 (b) shows the proportion of the difference in the total snow area obtained by the two snow cover products in the total snow area obtained by the binary snow product. The average difference is 34.53%, and the maximum difference is 59.52%. Comparing Figure 15 (a) and (b), the smaller the total snow cover area is, the greater the difference between the two sets of products, indicating that greater errors in the binary snow cover product occur for more broken and smaller areas.



485 **Figure 15:** Difference between the binary snow cover product and the fractional snow cover product in the Asia Water Tower region: (a) the difference in total snow cover area between the binary snow cover product and the fractional snow cover product, (b) proportion of total snow area difference between the two snow products compared with the total snow area obtained by the binary snow product

5.3 Limitations of the AWT MODIS FSC product

490 Clouds in most of the Asia Water Tower region have the characteristics of wide coverage and long duration. If clouds exist for a long-term, relying only on MODIS data will lead to a serious data gap, and the accuracy of snow cover monitoring will be reduced, regardless of any spatiotemporal reconstruction algorithm. With the launch of a new generation of geostationary satellites (FY-4A/B, GOES-17/18, Himawari-8/9, and MSG/MTG), their sensor performance can be comparable to that of MODIS sensors, and at the same time, the observation can be realized once every 5-15 minutes. Combined with geostationary sensors, these platforms are expected to provide the highest precision fractional snow cover monitoring. At the same time, the MODIS cloud product is overestimated, and the observation of multitemporal geostationary satellites can help improve the MODIS cloud product, thus improving the accuracy of snow cover monitoring. In this study, the monthly average cloud cover data MODIS and FY4A during the period 2018.04-2022.03 were collected, and the results are shown in Figure 16. Figure 16 (a) shows the average monthly cloud cover statistics from MODIS. It can be seen from the figure that the average monthly cloud cover in areas with more snow cover, such as the Pamir Plateau, Tianshan Mountain and Altai Mountain, is generally



more than 15 days, and some areas of the Hengduan Mountain range can reach more than 25 days. As shown in Figure 16 (b), the average monthly cloud cover in most regions based on the FY4A data is generally less than 10 days, and the average monthly cloud cover in the Hengduan Mountain region is also less than 19 days, which will provide strong data support for the subsequent high-precision spatiotemporal reconstruction of fractional snow cover information. However, the large amount of multitemporal observation data from geostationary satellites, the lack of necessary preprocessing steps, such as atmospheric correction, angle correction and geometric registration, and the serious misjudgment of cloud snow in existing cloud products will limit its further application.




Figure 16: Monthly average cloud cover over the Asia Water Tower region (2018.04-2022.03) from (a) MODIS and (b) FY4A

6 Data and Code availability

The AWT MODIS FSC product is the daily cloud-gap-filled snow cover data for the Asia Water Tower region. It has a spatial resolution of 0.005° and a daily temporal resolution. This dataset is freely available from the National Tibetan Plateau Data Center at <https://doi.org/10.11888/Cryos.tpd.c.272503> (Jiang et al., 2022) or from the Zenodo platform at <https://zenodo.org/doi/10.5281/zenodo.10005826>. It contains 8347 daily data files from 26 February 2000 to 31 December 2022 in NetCDF format. The filename rule is 'AWT_MODIS_FSC_yyyymmdd.NC', where AWT_MODIS_FSC represents the daily cloud-gap-filled MODIS fractional snow cover product over the Asian Water Tower region, and yyyymmdd indicates the year, month, and day of the data. The dataset contains two layer, 'fSCA' layer: fractional snow cover (non-snow (0), snow (1-100), water (237), cloud (250), and filling value (255)), 'QA' layer: cloud mask (0: clear sky, 1: cloud mask and 2: invalid value).

The Landsat-8 fractional snow cover dataset for verification is available on the Zenodo platform: <https://doi.org/10.5281/zenodo.10008227>. The binary value (snow/no-snow) snow depth dataset based on ground stations is available on the Github platform: https://github.com/FangboPan/AWT_Site_SD. The code is available on the Github platform: https://github.com/FangboPan/AWT_MODIS_DailyFSC_Product_code_v1.



7 Conclusions

In this study, based on the MESMA-AGE algorithm and the MSTI spatiotemporal reconstruction algorithm, the daily AWT MODIS FSC product was produced with long-term series, high precision, and spatiotemporal continuity in the Asian Water Tower region. The spatial resolution of the product is 0.005° from 2000 to 2022. The new AWT MODIS FSC product was quantitatively evaluated in two dimensions: binary value and fractional snow cover using snow depth observations from meteorological stations and high spatial resolution Landsat-8 images. Based on the results of the Landsat-8 image accuracy evaluation, the binarized identification accuracy metrics OA, PA and UA are 95.17%, 97.34% and 97.59%, respectively. The total fractional snow cover accuracy metrics R^2 , RMSE and MAE are 0.80, 0.16 and 0.10, respectively, compared with 2745 Landsat-8 images. All these results indicate that the AWT MODIS FSC product has good consistency with the high spatial resolution Landsat-8 images and has high accuracy. Based on the accuracy evaluation results after excluding the stations that cannot observe snow at all, the OA, PA, UA and CK of the AWT MODIS FSC product can reach 93.26%, 84.41%, 82.14% and 0.79, respectively. The AWT MODIS FSC product consists of two parts: the retrieval results of MODIS clear sky observations and the spatiotemporal reconstruction results based on the MSTI algorithm. Snow depth observations from meteorological stations are also used to evaluate these two parts. The binary precision metrics of fractional snow cover based on MODIS clear-sky observations are as follows: OA (95.36%), PA (87.75%), UA (86.86%) and CK (0.84). The binarization accuracy metrics of the fractional snow cover based on the spatiotemporal reconstruction of the MSTI algorithm are as follows: OA (88.96%), PA (82.26%), UA (78.86%), CK (0.72). Therefore, it can be shown that both the binarized identification and fractional snow cover metrics are excellent at both the point scale and the areal scale, which further indicates that this AWT MODIS FSC product has relatively high precision (Wu et al., 2021; Huang et al., 2022; Hao et al., 2022). The AWT MODIS FSC product is expected to offer robust and highly accurate data support for future snow hydrology studies.

Author contribution

PFB performed the study and wrote the manuscript. JLM conceived the study and supervised the manuscript construction and revision. WGX, PJM, HJY, ZC, CHZ, and YJW helped in algorithm development and data processing. ZZJ provided the meteorological station data. WSL and SJC provided many meaningful suggestions. All authors have read and agreed to the published version of the manuscript.

Competing interests

The authors declare that they have no conflicts of interest.



Acknowledgement

The authors would like to thank the National Meteorological Information Center (NMIC) for the weather station data. We also
550 acknowledge that the Google Earth Engine dramatically facilitated the work on image reprocessing.

Financial support

This research was funded by the Second Tibetan Plateau Scientific Expedition and Research Program (STEP) under Grant No. 2019QZKK0206, the Strategic Priority Research Program of Chinese Academy of Sciences (XDA20100300) and the National Natural Science Foundation of China (42171317, 42090014).

555 References

- Bair, E. H., Stilling, T., and Dozier, J.: Snow Property Inversion From Remote Sensing (SPIReS): A Generalized Multispectral Unmixing Approach With Examples From MODIS and Landsat 8 OLI, *IEEE Trans. Geosci. Remote Sens.*, 59, 7270–7284, <https://doi.org/10.1109/TGRS.2020.3040328>, 2021.
- Barnett, T. P., Adam, J. C., and Lettenmaier, D. P.: Potential impacts of a warming climate on water availability in snow-dominated regions, *Nature*, 438, 303–309, <https://doi.org/10.1038/nature04141>, 2005.
- 560 Czychowska-Wisniewski, E. H., van Leeuwen, W. J. D., Hirschboeck, K. K., Marsh, S. E., and Wisniewski, W. T.: Fractional snow cover estimation in complex alpine-forested environments using an artificial neural network, *Remote Sens. Environ.*, 156, 403–417, <https://doi.org/10.1016/j.rse.2014.09.026>, 2015.
- Dai, L., Che, T., Ding, Y., and Hao, X.: Evaluation of snow cover and snow depth on the Qinghai–Tibetan Plateau derived from passive microwave remote sensing, *The Cryosphere*, 11, 1933–1948, <https://doi.org/10.5194/tc-11-1933-2017>, 2017.
- 565 Dobreva, I. D. and Klein, A. G.: Fractional snow cover mapping through artificial neural network analysis of MODIS surface reflectance, *Remote Sens. Environ.*, 115, 3355–3366, <https://doi.org/10.1016/j.rse.2011.07.018>, 2011.
- Dozier, J., Painter, T. H., Rittger, K., and Frew, J. E.: Time–space continuity of daily maps of fractional snow cover and albedo from MODIS, *Adv. Water Resour.*, 31, 1515–1526, <https://doi.org/10.1016/j.advwatres.2008.08.011>, 2008.
- 570 Elguindi, N., Hanson, B., and Leathers, D.: The Effects of Snow Cover on Midlatitude Cyclones in the Great Plains, *J. Hydrometeorol.*, 6, 263–279, <https://doi.org/10.1175/JHM415.1>, 2005.
- Gafurov, A. and Bárdossy, A.: Cloud removal methodology from MODIS snow cover product, *Hydrol. Earth Syst. Sci.*, 13, 1361–1373, <https://doi.org/10.5194/hess-13-1361-2009>, 2009.
- GDAL Development Team: GDAL -- Geospatial Data Abstraction Library, 2022.
- 575 Hall, D. K. and Riggs, G. A.: MODIS/Terra Snow Cover Daily L3 Global 500m SIN Grid, Version 6, <https://doi.org/10.5067/MODIS/MOD10A1.006>, 2016.



- Hall, D. K., Riggs, G. A., and Salomonson, V. V.: Development of Methods for Mapping Global Snow Cover Using Moderate Resolution Imaging Spectroradiometer Data, *Remote Sens. Environ.*, 54, 127–140, [https://doi.org/10.1016/0034-4257\(95\)00137-P](https://doi.org/10.1016/0034-4257(95)00137-P), 1995.
- 580 Hao, S., Jiang, L., Shi, J., Wang, G., and Liu, X.: Assessment of MODIS-Based Fractional Snow Cover Products Over the Tibetan Plateau, *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.*, 12, 533–548, <https://doi.org/10.1109/JSTARS.2018.2879666>, 2019.
- Hao, X., Huang, G., Che, T., Ji, W., Sun, X., Zhao, Q., Zhao, H., Wang, J., Li, H., and Yang, Q.: The NIEER AVHRR snow cover extent product over China – A long-term daily snow record for regional climate research, *Cryosphere – Snow and Sea Ice*, <https://doi.org/10.5194/essd-2021-189>, 2021.
- 585 Hao, X., Huang, G., Zheng, Z., Sun, X., Ji, W., Zhao, H., Wang, J., Li, H., and Wang, X.: Development and validation of a new MODIS snow-cover-extent product over China, *Hydrol. Earth Syst. Sci.*, 26, 1937–1952, <https://doi.org/10.5194/hess-26-1937-2022>, 2022.
- Hori, M., Sugiura, K., Kobayashi, K., Aoki, T., Tanikawa, T., Kuchiki, K., Niwano, M., and Enomoto, H.: A 38-year (1978–2015) Northern Hemisphere daily snow cover extent product derived using consistent objective criteria from satellite-borne optical sensors, *Remote Sens. Environ.*, 191, 402–418, <https://doi.org/10.1016/j.rse.2017.01.023>, 2017.
- Hou, J., Huang, C., Zhang, Y., Guo, J., and Gu, J.: Gap-Filling of MODIS Fractional Snow Cover Products via Non-Local Spatio-Temporal Filtering Based on Machine Learning Techniques, *Remote Sens.*, 11, 90, <https://doi.org/10.3390/rs11010090>, 2019.
- 595 Huang, X., Deng, J., Ma, X., Wang, Y., Feng, Q., Hao, X., and Liang, T.: Spatiotemporal dynamics of snow cover based on multi-source remote sensing data in China, *The Cryosphere*, 10, 2453–2463, <https://doi.org/10.5194/tc-10-2453-2016>, 2016.
- Huang, Y., Liu, H., Yu, B., Wu, J., Kang, E. L., Xu, M., Wang, S., Klein, A., and Chen, Y.: Improving MODIS snow products with a HMRF-based spatio-temporal modeling technique in the Upper Rio Grande Basin, *Remote Sens. Environ.*, 204, 568–582, <https://doi.org/10.1016/j.rse.2017.10.001>, 2018.
- 600 Huang, Y., Xu, J., Xu, J., Zhao, Y., Yu, B., Liu, H., Wang, S., Xu, W., Wu, J., and Zheng, Z.: HMRFs-TP: long-term daily gap-free snow cover products over the Tibetan Plateau from 2002 to 2021 based on Hidden Markov Random Field model, *Snow and Sea Ice*, <https://doi.org/10.5194/essd-2022-134>, 2022.
- Immerzeel, W. W., van Beek, L. P. H., and Bierkens, M. F. P.: Climate Change Will Affect the Asian Water Towers, *Science*, 328, 1382–1385, <https://doi.org/10.1126/science.1183188>, 2010.
- 605 Immerzeel, W. W., Lutz, A. F., Andrade, M., Bahl, A., Biemans, H., Bolch, T., Hyde, S., Brumby, S., Davies, B. J., Elmore, A. C., Emmer, A., Feng, M., Fernández, A., Haritashya, U., Kargel, J. S., Koppes, M., Kraaijenbrink, P. D. A., Kulkarni, A. V., Mayewski, P. A., Nepal, S., Pacheco, P., Painter, T. H., Pellicciotti, F., Rajaram, H., Rupper, S., Sinisalo, A., Shrestha, A. B., Viviroli, D., Wada, Y., Xiao, C., Yao, T., and Baillie, J. E. M.: Importance and vulnerability of the world’s water towers, *Nature*, 577, 364–369, <https://doi.org/10.1038/s41586-019-1822-y>, 2020.
- 610 Jiang, L., Pan, F., Wang, G., Pan, J., Shi, J., and Zhang, C.: MODIS daily cloud-free fractional snow cover data set for Asian water tower area (2000–2022), <https://doi.org/10.11888/Cryos.tpdc.272503>. <https://cstr.cn/18406.11.Cryos.tpdc.272503>, 2022.
- Kraaijenbrink, P. D. A., Bierkens, M. F. P., Lutz, A. F., and Immerzeel, W. W.: Impact of a global temperature rise of 1.5 degrees Celsius on Asia’s glaciers, *Nature*, 549, 257–260, <https://doi.org/10.1038/nature23878>, 2017.



- 615 Kuter, S.: Completing the machine learning saga in fractional snow cover estimation from MODIS Terra reflectance data: Random forests versus support vector regression, *Remote Sens. Environ.*, 255, https://doi.org/ARTN_112294 10.1016/j.rse.2021.112294, 2021.
- Li, M., Zhu, X., Li, N., and Pan, Y.: Gap-Filling of a MODIS Normalized Difference Snow Index Product Based on the Similar Pixel Selecting Algorithm: A Case Study on the Qinghai–Tibetan Plateau, *Remote Sens.*, 12, <https://doi.org/10.3390/rs12071077>, 2020.
- 620 Li, W., Guo, W., Qiu, B., Xue, Y., Hsu, P.-C., and Wei, J.: Influence of Tibetan Plateau snow cover on East Asian atmospheric circulation at medium-range time scales, *Nat. Commun.*, 9, 4243, <https://doi.org/10.1038/s41467-018-06762-5>, 2018.
- Li, X., Shen, H., Zhang, L., Zhang, H., and Yuan, Q.: Dead Pixel Completion of Aqua MODIS Band 6 Using a Robust M-Estimator Multiregression, *IEEE Geosci. Remote Sens. Lett.*, 11, 768–772, <https://doi.org/10.1109/LGRS.2013.2278626>, 2014.
- 625 Li, X., Fu, W., Shen, H., Huang, C., and Zhang, L.: Monitoring snow cover variability (2000–2014) in the Hengduan Mountains based on cloud-removed MODIS products with an adaptive spatio-temporal weighted method, *J. Hydrol.*, 551, 314–327, <https://doi.org/10.1016/j.jhydrol.2017.05.049>, 2017.
- Li, X., Long, D., Scanlon, B. R., Mann, M. E., Li, X., Tian, F., Sun, Z., and Wang, G.: Climate change threatens terrestrial water storage over the Tibetan Plateau, *Nat. Clim. Change*, 12, 801–807, <https://doi.org/10.1038/s41558-022-01443-0>, 2022.
- 630 Lindsay, C., Zhu, J., Miller, A., Kirchner, P., and Wilson, T.: Deriving snow cover metrics for Alaska from MODIS, *Remote Sens.*, 7, 12961–12985, <https://doi.org/10.3390/rs71012961>, 2015.
- Liu, T., Chen, D., Yang, L., Meng, J., Wang, Z., Ludescher, J., Fan, J., Yang, S., Chen, D., Kurths, J., Chen, X., Havlin, S., and Schellnhuber, H. J.: Teleconnections among tipping elements in the Earth system, *Nat. Clim. Change*, 13, 67–74, <https://doi.org/10.1038/s41558-022-01558-4>, 2023.
- Liu, X. and Chen, B.: Climatic warming in the Tibetan Plateau during recent decades, *Int. J. Climatol.*, 20, 1729–1742, [https://doi.org/10.1002/1097-0088\(20001130\)20:14<1729::AID-JOC556>3.0.CO;2-Y](https://doi.org/10.1002/1097-0088(20001130)20:14<1729::AID-JOC556>3.0.CO;2-Y), 2000.
- López-Burgos, V., Gupta, H. V., and Clark, M.: Reducing cloud obscuration of MODIS snow cover area products by combining spatio-temporal techniques with a probability of snow approach, *Hydrol. Earth Syst. Sci.*, 17, 1809–1823, <https://doi.org/10.5194/hess-17-1809-2013> 10.5194/hessd-9-13693-2012, 2013.
- 640 Martinec, J.: Snowmelt - runoff model for stream flow forecasts, *Hydrol. Res.*, 6, 145–154, <https://doi.org/10.2166/nh.1975.0010>, 1975.
- Mazari, N., Tekeli, A. E., Xie, H., Sharif, H. I., and Hassan, A. A. E.: Assessment of ice mapping system and moderate resolution imaging spectroradiometer snow cover maps over Colorado Plateau, *J. Appl. Remote Sens.*, 7, 073540, <https://doi.org/10.1117/1.JRS.7.073540>, 2013.
- 645 Metsämäki, S., Mattila, O.-P., Pulliainen, J., Niemi, K., Luojus, K., and Böttcher, K.: An optical reflectance model-based method for fractional snow cover mapping applicable to continental scale, *Remote Sens. Environ.*, 123, 508–521, <https://doi.org/10.1016/j.rse.2012.04.010>, 2012.
- Metsamaki, S. J., Anttila, S. T., Markus, H. J., and Vepsäläinen, J. M.: A feasible method for fractional snow cover mapping in boreal zone based on a reflectance model, *Remote Sens. Environ.*, 95, 77–95, <https://doi.org/10.1016/j.rse.2004.11.013>, 2005.



- 650 Naegeli, K., Neuhaus, C., Salberg, A.-B., Schwaizer, G., Weber, H., Wiesmann, A., Wunderle, S., and Nagler, T.: ESA Snow Climate Change Initiative (Snow_cci): Daily global Snow Cover Fraction - snow on ground (SCFG) from AVHRR (1982 - 2018), version 2.0., <https://dx.doi.org/10.5285/3f034f4a08854eb59d58e1fa92d207b6>, 2022.
- Nagler, T., Schwaizer, G., Molg, N., Keuris, L., Hetzenecker, M., and Metsämäki, S.: ESA Snow Climate Change Initiative (Snow_cci): Daily global Snow Cover Fraction - snow on ground (SCFG) from MODIS (2000-2020), version 2.0,
655 <https://dx.doi.org/10.5285/8847a05eeda646a29da58b42bdf2a87c>, 2022.
- Niittynen, P., Heikkinen, R. K., and Luoto, M.: Decreasing snow cover alters functional composition and diversity of Arctic tundra, *Proc. Natl. Acad. Sci.*, 117, 21480–21487, <https://doi.org/10.1073/pnas.2001254117>, 2020.
- Notarnicola, C.: Hotspots of snow cover changes in global mountain regions over 2000–2018, *Remote Sens. Environ.*, 243, 111781, <https://doi.org/10.1016/j.rse.2020.111781>, 2020.
- 660 Painter, T. H., Dozier, J., Roberts, D. A., Davis, R. E., and Green, R. O.: Retrieval of subpixel snow-covered area and grain size from imaging spectrometer data, *Remote Sens. Environ.*, 85, 64–77, [https://doi.org/10.1016/S0034-4257\(02\)00187-6](https://doi.org/10.1016/S0034-4257(02)00187-6), 2003.
- Painter, T. H., Rittger, K., McKenzie, C., Slaughter, P., Davis, R. E., and Dozier, J.: Retrieval of subpixel snow covered area, grain size, and albedo from MODIS, *Remote Sens. Environ.*, 113, 868–879, <https://doi.org/10.1016/j.rse.2009.01.001>, 2009.
- 665 Pan, F., Jiang, L., Zheng, Z., Wang, G., Cui, H., Zhou, X., and Huang, J.: Retrieval of Fractional Snow Cover over High Mountain Asia Using 1 km and 5 km AVHRR/2 with Simulated Mid-Infrared Reflective Band, *Remote Sens.*, 14, 3303, <https://doi.org/10.3390/rs14143303>, 2022.
- Pan, J., Yang, J., Jiang, L., Xiong, C., Pan, F., Gao, X., Shi, J., and Chang, S.: Combination of Snow Process Model Priors and Site Representativeness Evaluation to Improve the Global Snow Depth Retrieval Based on Passive Microwaves, *IEEE Trans. Geosci. Remote Sens.*, 61, 1–20, <https://doi.org/10.1109/TGRS.2023.3276651>, 2023.
- 670 Parajka, J. and Blöschl, G.: Spatio-temporal combination of MODIS images - potential for snow cover mapping, *Water Resour. Res.*, 44, W03406, <https://doi.org/10.1029/2007WR006204>, 2008.
- Paudel, K. P. and Andersen, P.: Monitoring snow cover variability in an agropastoral area in the Trans Himalayan region of Nepal using MODIS data with improved cloud removal methodology, *Remote Sens. Environ.*, 115, 1234–1246,
675 <https://doi.org/10.1016/j.rse.2011.01.006>, 2011.
- Reuter, H. I., Nelson, A., and Jarvis, A.: An evaluation of void-filling interpolation methods for SRTM data, *Int. J. Geogr. Inf. Sci.*, 21, 983–1008, <https://doi.org/10.1080/13658810601169899>, 2007.
- Riggs, G. A., Hall, D. K., and Román, M. O.: Overview of NASA’s MODIS and Visible Infrared Imaging Radiometer Suite (VIIRS) snow-cover Earth System Data Records, *Earth Syst. Sci. Data*, 9, 765–777, <https://doi.org/10.5194/essd-9-765-2017>,
680 2017.
- Rittger, K., Painter, T. H., and Dozier, J.: Assessment of methods for mapping snow cover from MODIS, *Adv. Water Resour.*, 51, 367–380, <https://doi.org/10.1016/j.advwatres.2012.03.002>, 2013.
- Rittger, K., Bair, E. H., Kahl, A., and Dozier, J.: Spatial estimates of snow water equivalent from reconstruction, *Adv. Water Resour.*, 94, 345–363, <https://doi.org/10.1016/j.advwatres.2016.05.015>, 2016.



- 685 Roberts, D. A., Gardner, M., Church, R., Ustin, S., Scheer, G., and Green, R. O.: Mapping Chaparral in the Santa Monica Mountains Using Multiple Endmember Spectral Mixture Models, *Remote Sens. Environ.*, 65, 267–279, [https://doi.org/10.1016/S0034-4257\(98\)00037-6](https://doi.org/10.1016/S0034-4257(98)00037-6), 1998.
- Salomonson, V. V. and Appel, I.: Estimating fractional snow cover from MODIS using the normalized difference snow index, *Remote Sens. Environ.*, 89, 351–360, <https://doi.org/10.1016/j.rse.2003.10.016>, 2004.
- 690 Salomonson, V. V. and Appel, I.: Development of the Aqua MODIS NDSI fractional snow cover algorithm and validation results, *Ieee Trans. Geosci. Remote Sens.*, 44, 1747–1756, <https://doi.org/10.1109/Tgrs.2006.876029>, 2006.
- Senan, R., Orsolini, Y. J., Weisheimer, A., Vitart, F., Balsamo, G., Stockdale, T. N., Dutra, E., Doblas-Reyes, F. J., and Basang, D.: Impact of springtime Himalayan–Tibetan Plateau snowpack on the onset of the Indian summer monsoon in coupled seasonal forecasts, *Clim. Dyn.*, 47, 2709–2725, <https://doi.org/10.1007/s00382-016-2993-y>, 2016.
- 695 Shea, J. M., Menounos, B., Moore, R. D., and Tennant, C.: An approach to derive regional snow lines and glacier mass change from MODIS imagery, western North America, *The Cryosphere*, 7, 667–680, <https://doi.org/10.5194/tc-7-667-2013>, 2013.
- Shi, J.: An Automatic Algorithm on Estimating Sub-Pixel Snow Cover from MODIS, *Quat. Sci.*, 32, 6–15, 2012.
- Sulla-Menashe, D., Gray, J. M., Abercrombie, S. P., and Friedl, M. A.: Hierarchical mapping of annual global land cover 2001 to present: The MODIS Collection 6 Land Cover product, *Remote Sens. Environ.*, 222, 183–194, <https://doi.org/10.1016/j.rse.2018.12.013>, 2019.
- 700 Tang, Z., Wang, J., Li, H., and Yan, L.: Spatiotemporal changes of snow cover over the Tibetan plateau based on cloud-removed moderate resolution imaging spectroradiometer fractional snow cover product from 2001 to 2011, *J. Appl. Remote Sens.*, 7, 073582, <https://doi.org/10.1117/1.JRS.7.073582>, 2013.
- Tang, Z., Wang, X., Wang, J., Wang, X., Li, H., and Jiang, Z.: Spatiotemporal Variation of Snow Cover in Tianshan Mountains, Central Asia, Based on Cloud-Free MODIS Fractional Snow Cover Product, 2001–2015, *Remote Sens.*, 9, 1045, <https://doi.org/10.3390/rs9101045>, 2017.
- Tang, Z., Deng, G., Hu, G., Zhang, H., Pan, H., and Sang, G.: Satellite observed spatiotemporal variability of snow cover and snow phenology over high mountain Asia from 2002 to 2021, *J. Hydrol.*, 613, 128438, <https://doi.org/10.1016/j.jhydrol.2022.128438>, 2022.
- 710 Tran, H., Nguyen, P., Ombadi, M., Hsu, K., Sorooshian, S., and Qing, X.: A cloud-free MODIS snow cover dataset for the contiguous United States from 2000 to 2017, *Sci. Data*, 6, 180300, <https://doi.org/10.1038/sdata.2018.300>, 2019.
- Wang, G., Jiang, L., Wu, S., Shi, J., Hao, S., and Liu, X.: Fractional Snow Cover Mapping from FY-2 VISSR Imagery of China, *Remote Sens.*, 9, 983, <https://doi.org/10.3390/rs9100983>, 2017.
- 715 Wang, G., Jiang, L., Shi, J., Liu, X., Yang, J., and Cui, H.: Snow-Covered Area Retrieval from Himawari–8 AHI Imagery of the Tibetan Plateau, *Remote Sens.*, 11, <https://doi.org/10.3390/rs11202391>, 2019.
- Wang, G., Jiang, L., Shi, J., and Su, X.: A Universal Ratio Snow Index for Fractional Snow Cover Estimation, *IEEE Geosci. Remote Sens. Lett.*, 18, 721–725, <https://doi.org/10.1109/LGRS.2020.2982053>, 2021.
- Wang, G., Jiang, L., Xiong, C., and Zhang, Y.: Characterization of NDSI Variation: Implications for Snow Cover Mapping, *IEEE Trans. Geosci. Remote Sens.*, 60, 1–18, <https://doi.org/10.1109/TGRS.2022.3165986>, 2022.



- 720 Wang, G., Jiang, L., Pan, F., Weng, H., and Zhang, Y.: Sensitivity of Snow NDSI to Simulated Snow Grain Shape Characteristics, *IEEE Geosci. Remote Sens. Lett.*, 20, 1–5, <https://doi.org/10.1109/LGRS.2022.3233379>, 2023.
- Wang, T., Peng, S., Lin, X., and Chang, J.: Declining snow cover may affect spring phenological trend on the Tibetan Plateau, *Proc. Natl. Acad. Sci.*, 110, <https://doi.org/10.1073/pnas.1306157110>, 2013.
- 725 Wang, X., Chen, S., and Wang, J.: An Adaptive Snow Identification Algorithm in the Forests of Northeast China, *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.*, 13, 5211–5222, <https://doi.org/10.1109/JSTARS.2020.3020168>, 2020.
- Wu, X. D., Naegeli, K., Premier, V., Marin, C., Ma, D. J., Wang, J. P., and Wunderle, S.: Evaluation of snow extent time series derived from Advanced Very High Resolution Radiometer global area coverage data (1982–2018) in the Hindu Kush Himalayas, *Cryosphere*, 15, 4261–4279, <https://doi.org/10.5194/tc-15-4261-2021>, 2021.
- 730 Xiao, X., He, T., Liang, S., Liu, X., Ma, Y., Liang, S., and Chen, X.: Estimating fractional snow cover in vegetated environments using MODIS surface reflectance data, *Int. J. Appl. Earth Obs. Geoinformation*, 114, 103030, <https://doi.org/10.1016/j.jag.2022.103030>, 2022.
- Xing, D., Hou, J., Huang, C., and Zhang, W.: Spatiotemporal Reconstruction of MODIS Normalized Difference Snow Index Products Using U-Net with Partial Convolutions, *Remote Sens.*, 14, 1795, <https://doi.org/10.3390/rs14081795>, 2022.
- 735 Xu, Y., Shi, J., and Du, J.: An Improved Endmember Selection Method Based on Vector Length for MODIS Reflectance Channels, *Remote Sens.*, 7, 6280–6295, <https://doi.org/10.3390/rs70506280>, 2015.
- Yang, J., Jiang, L., Shi, J., Wu, S., Sun, R., and Yang, H.: Monitoring snow cover using Chinese meteorological satellite data over China, *Remote Sens. Environ.*, 143, 192–203, <https://doi.org/10.1016/j.rse.2013.12.022>, 2014.
- Yang, Y., Chen, R., Liu, G., Liu, Z., and Wang, X.: Trends and variability in snowmelt in China under climate change, *Hydrol. Earth Syst. Sci.*, 26, 305–329, <https://doi.org/10.5194/hess-26-305-2022>, 2022.
- 740 Yao, T., Bolch, T., Chen, D., Gao, J., Immerzeel, W., Piao, S., Su, F., Thompson, L., Wada, Y., Wang, L., Wang, T., Wu, G., Xu, B., Yang, W., Zhang, G., and Zhao, P.: The imbalance of the Asian water tower, *Nat. Rev. Earth Environ.*, 3, 618–632, <https://doi.org/10.1038/s43017-022-00299-4>, 2022.
- 745 Yu, J., Zhang, G., Yao, T., Xie, H., Zhang, H., Ke, C., and Yao, R.: Developing Daily Cloud-Free Snow Composite Products From MODIS Terra–Aqua and IMS for the Tibetan Plateau, *IEEE Trans. Geosci. Remote Sens.*, 54, 2171–2180, <https://doi.org/10.1109/TGRS.2015.2496950>, 2016.
- Zhang, H., Zhang, F., Zhang, G., Che, T., Yan, W., Ye, M., and Ma, N.: Ground-based evaluation of MODIS snow cover product V6 across China: Implications for the selection of NDSI threshold, *Sci. Total Environ.*, 651, 2712–2726, <https://doi.org/10.1016/j.scitotenv.2018.10.128>, 2019.
- 750 Zhang, H., Zhang, F., Che, T., and Wang, S.: Comparative evaluation of VIIRS daily snow cover product with MODIS for snow detection in China based on ground observations, *Sci. Total Environ.*, 724, 138156, <https://doi.org/10.1016/j.scitotenv.2020.138156>, 2020.
- Zhang, H., Zhang, F., Zhang, G., Yan, W., and Li, S.: Enhanced scaling effects significantly lower the ability of MODIS normalized difference snow index to estimate fractional and binary snow cover on the Tibetan Plateau, *J. Hydrol.*, 592, 125795, <https://doi.org/10.1016/j.jhydrol.2020.125795>, 2021.



- 755 Zhao, K., Peng, D., Gu, Y., Luo, X., Pang, B., and Zhu, Z.: Temperature lapse rate estimation and snowmelt runoff simulation in a high-altitude basin, *Sci. Rep.*, 12, 13638, <https://doi.org/10.1038/s41598-022-18047-5>, 2022.
- Zhu, J. and Shi, J.: An Algorithm for Subpixel Snow Mapping: Extraction of a Fractional Snow-Covered Area Based on Ten-Day Compositing AVHRR V2 Data of the Qinghai-Tibet Plateau, *IEEE Geosci. Remote Sens. Mag.*, 6, 86–98, <https://doi.org/10.1109/mgrs.2018.2850963>, 2018.

760