# **Seeing the Wood for the Trees: Active human-**

2

# environmental interactions in arid northwest China

3	Hui Shen <sup>1,2</sup> , Robert N. Spengler <sup>3,4</sup> , Xinying Zhou <sup>1,2</sup> , Alison Betts <sup>5</sup> , Peter Weiming
4	Jia <sup>5</sup> , Keliang Zhao <sup>1,2</sup> , Xiaoqiang Li <sup>1,2</sup>
5	<sup>1</sup> Key Laboratory of Vertebrate Evolution and Human Origins, Institute of Vertebrate Paleontology and
6	Paleoanthropology, Chinese Academy of Sciences, Beijing, 100044, China
7	<sup>2</sup> University of Chinese Academy of Sciences, Beijing, 100049, China
8	<sup>3</sup> Domestication and Anthropogenic Evolution Research Group, Max Planck Institute of
9	Geoanthropology, Jena, 07745, Germany
10	<sup>4</sup> Department of Archaeology, Max Planck Institute of Geoanthropology, Jena, 07745, Germany
11	<sup>5</sup> Department of Archaeology, University of Sydney, Sydney, NSW 2006, Australia
12	
13	Corresponding author: Xiaoqiang Li
14	email: lixiaoqiang@ivpp.ac.cn
15	
16	
17	
18	
19	

**Abstract:** Due largely to demographic growth, agricultural populations during the Holocene became increasingly more impactful ecosystem engineers. Multidisciplinary research has revealed a deep history of human-environmental dynamics; however, these pre-modern anthropogenic ecosystem transformations and cultural adaptions are still poorly understood. Here, we synthesis anthracological data to explore the complex array of human-environmental interactions in the regions of the prehistoric Silk Road. Our results suggest that these ancient humans were not passively impacted by environmental change, but rather they culturally adapted to, and in turn altered, arid ecosystems. Underpinned by the establishment of complex agricultural systems on the western Loess Plateau, people may have started to manage chestnut trees, likely through conservation of economically significant species, as early as 4600 BP. Since ca. 3500 BP, with the appearance of high-yielding wheat/barley farming in Xinjiang and the Hexi Corridor, people appear to have been cultivating *Prunus* and *Morus* trees. We also argue that people were transporting the preferred coniferous woods over long distances to meet the need for fuel and timber. After 2500 BP, people in our study area were making conscious selections between wood types for craft production, and were also clearly cultivating a wide range of long-generation perennials, showing a remarkable traditional knowledge tied into the arid environment. At the same time, the data suggest that there was significant deforestation throughout the chronology of occupation, including a rapid decline of slow-growing spruce forests and riparian woodlands across the northwest China. The wood charcoal dataset is publicly available at https://doi.org/10.5281/zenodo.8158277 (Shen et al., 2023).

- 43 **Keywords:** Human-environmental interaction, human adaption, fruit management,
- 44 deforestation, northwest China

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

# 1 Introduction

45

68

46	The extent of prehistoric anthropogenic environmental change long-term interaction
47	between our humans and their environment, especially relating to the ways early
48	agricultural practices groups reshaped and adapted to terrestrial ecosystems, has been
49	the subject of ongoing debate (Ruddiman, 2003, 2008; Zong et al., 2007; Zalasiewicz
50	et al., 2017; ArchaeoGLOBE Project, 2019; Renn, 2020; Asouti and Kabukcu, 2014;
51	Asouti et al., 2015; Dong et al., 2020a, 2022a; Cowie et al., 2022). Over the past
52	decade, scholars have adopted big data approaches to understanding long-term-
53	anthropogenic changes to the Earth's surface (Zalasiewicz et al., 2017;
54	ArchaeoGLOBE Project, 2019; Renn, 2020; Cowie et al., 2022). While humans have
55	undoubtedly been reshaping interactinged with their environments since before the
56	Holocene, the magnitude and complexity of these impacts this interaction following
57	the adoption of agricultural economies increased immensely. During this process,
58	people shifted their subsistence system from hunting-gathering to cereal cultivation
59	and animal husbandry, and increasingly gained the ability to alter and adapt their
60	ecological surroundings (Bellwood, 2005; Zeder, 2008; Zohary et al., 2012). During
61	the fifth millennium BP, agricultural populations across Europe and Asia first came
62	into contact via diffusion of crops, contributing to food globalization in prehistory
63	(Sherratt, 2006; Jones et al., 2011; Dong et al., 2017, 2022b; Boivin et al., 2016; Liu
64	et al., 2019; Zhou et al., 2020). The intermingling of millets, adapted for arid and
65	short-season grasslands in northern China, with cereals, adapted for rainy season
66	growth in arid southwest Asia, eventually facilitated a greater intensification of
67	farming systems (Spengler 2019; Miller et al. 2016).

Mounting evidence shows that the development of <u>intensive</u> farming systems

69 was accompanied by a series of ecological and social changes, including 70 deforestation, wild species loss, and demographic expansion (Bellwood, 2005; 71 Weisdorf, 2005; Atahan et al., 2008; Kaplan et al., 2009; Bocquet-Appel, 2011; Fuller 72 et al., 2011a; Asouti et al., 2015; Ruddiman, 2013). For instance, the dispersal and expansion of agriculture largely altered the natural geographic distributions of 73 74 anthropophilic plants (crops and weeds) and directly influenced vegetation 75 communities worldwide (Vigne et al., 2012; Fuller et al., 2011b; Crowther et al., 2016; Boivin et al., 2017; Spengler et al., 2021). Forest clearing, either to increase the 76 77 surface area of arable land or to acquire wood for construction or fuel, has caused large-scale deforestation and created a more open landscape (Zong et al., 2007; 78 Atahan et al., 2008; Kaplan et al., 2009; Innes et al., 2013; Zheng et al., 2021). 79 80 Meanwhile, human-mediated management of local woodlands encouraged to 81 encourage the growth of fruit- and nut-bearing trees, shifting land-use strategies from 82 an emphasis on short-term returns of annual cereals to long-term investment with 83 delayed return crops, was widely recognized (Fall et al., 2002; Janick, 2005; Miller 84 and Gross, 2011; Miller, 2013; Asouti and Kabukcu, 2014; Asouti et al., 2015). Today, essentially all ecosystems on the planet are anthropogenic constructs, 85 recognized through the increasingly prominent use of the term Anthropocene 86 87 (Crutzen, 2002; Ruddiman, 2003, 2013; Monastersky, 2015). Northwest China, the focus region of this paper, is of particular interest, because 88 89 it is located at the core of the ancient trade routes that are colloquially referred to as 90 the Silk Road and farmers in the region were the first to experiment with agricultural crops from both West and East Asia (Wang et al., 2017; Dong et al., 2017, 2018, 91 92 2022b; Zhou et al., 2020; Li, 2021). Archaeobotanical data have pinpointed the broad-93 region and time period when humans first started to cultivated millets in East Asia.

Specifically, evidence from the Dadiwan site has revealed that broomcorn millet cultivation began as early as the eighth millennium BP (Liu et al., 2004; Li, 2018), and the gradual diffusion of broomcorn millet reached famers in the mountains of Central Asia by 4500 BP (Spengler et al. 2014; Yatoo et al. 2020). The remains of barley (Hordeum vulgare var. nudum) and wheat (Triticum aestivum) found at the Tongtian Cave site, have been dated to around 5200 BP, representing the earliest known southwest Asian cereals found in East Asia (Zhou et al., 2020). In addition to long-distance exchange diffusion of cereals and knowledge of their cultivation, this area also fostered the trans-continental dispersals of sheep, goat, bronze-smelting technology, mudbrick-manufacturing techniques, and a variety of other cultural attributes (Mei and Shell, 1991; Dodson et al., 2009; Li et al., 2011; Yang et al., 2017; Dong et al., 2017; Chen et al., 2018; Ren et al., 2022). Additionally, most of this region is characterized by a hyper-arid desert and fragile oasis ecosystem, which are especially vulnerable to human activity, making it a prime zone for studying the interaction between early agricultural societies and the environment.

Archaeologists and geologists working in this region have mainly focused their attention on the relationship between climate change and Neolithic cultural development, as well as anthropogenic impacts on regional ecosystems. These scholars have argued that enhanced precipitation during the Late Yangshao (5500-5000 BP), Majiayao type (5300-4800 BP), and Qijia (4200-3800 BP) periods played an important role in the expansion of these early farmers (An et al., 2004; 2005, 2006; Hou et al. 2009; Liu et al., 2010; Dong et al., 2012, 2013, 2016, 2020a). A reduction in the number of archaeological sites during the gap between early and middle Majiayao (4800-4400 BP), and the decline of the Qijia culture are thought to be a response to increasingly aridity (Dong et al., 2012, 2013). Concurrent with these

changes, people were actively engaged in reshaping the landscape. For instance, a wood charcoal study from the Hexi Corridor has suggested that prehistoric wood collection led to a rapid reduction in local woodlands and a decline in woody plant diversity (Shen et al., 2018). In a different study, an increase in large-scale fire-frequency was proposed based on micro carbon records from Tian'e Lake, which was further correlated with high Cu content, suggesting the consequence of large scale bronze smelting activities (Dong et al., 2020b). However, relatively less attention has been paid to how agriculture influenced the cultural responses and adaption strategies employed by early famers in these arid environments. Meanwhile, scientific records are geographically uneven, with regions, such as the Hexi Corridor, attracting considerable attention, while few studies have targeted the vast area of Xinjiang, leading to an incomplete picture of prehistoric human-environmental interactions along the ancient Silk Road.

In this study, we present a comprehensive synthesis of wood charcoal records from northwest China. As the result of incomplete burning, wood charcoal fragments from archaeological sites shed light on the practices of local woody plant use (Asoutiand Austin, 2005; Marguerie and Hunot, 2007; Théry Parisot et al., 2010). Since the first charcoal analyse, beginning in the 1940s (Salysbury and Jane, 1940), the application of reflected light microscopy has allowed for the rapid identification of charcoal, making it widely used in: 1) the reconstruction of firewood collection strategies (Asouti and Asutin, 2005; Marguerie and Hunot, 2007; Li et al., 2016; Shen et al., 2018; Kabukcu, 2017; Mas et al., 2021); 2) elucidating the impacts that wood cutting had on local forests (Li et al., 2011; Asouti et al., 2015; Knapp et al., 2015; Shen et al., 2018); 3) identifying compositions of woody communities (Wang et al., 2014; Asouti et al., 2015; Allué and Zaidner, 2022; Mas et al., 2022); and 4)

determining fruit and/or nut tree management (Miller, 2013; Asouti and Kabukcu, 2014; Shen and Li, 2021). Here, we seek to identify patterns with in wood charcoal assemblages recovered from seven archaeological sites in Xinjiang, which we contrast with more than 30 other published regional records. We aim to explore multiple perspectives on the complexities of human-environmental interactions within the agricultural background, including the influence of farming and wood cutting on woody vegetation change, as well as the strategies applied in response to climatic aridification.

#### 2 Study area

#### 2.1 Regional setting

Our study focuses on the provinces of Xinjiang and Gansu, because of the important roles people in this region played in exchange along the ancient Silk Road. This region is characterized by montane ecoclines, including those of the Tianshan, Altai, Altun, and Qilian mountains (Figure 1). Due to glacial snowmelt, alluvial plains are widely distributed across the low-land basins, and fine-grained nutrients and water brought by the runoff nourish a network of oases, especially within the Hexi Corridor and Tarim Basin (Zheng et al., 2015). Climatically, mean annual precipitation (MAP) is geographically uneven, due to differences in prevailing air masses. For the West Loess Plateau, which is under the control of the Asian monsoons, MAP usually exceeds 400 mm (https://data.cma.cn/). Water vapour carried by the westerlies mainly concentrates in the Ili or Irtysh valleys and Junggar Basin, and the MAP sometimes can reach more than 500 mm (Xiao et al., 2006; Zheng et al., 2015). In the Tarim Basin and the Hexi Corridor, the MAP is usually less than 200 mm

(<a href="https://data.cma.cn/">https://data.cma.cn/</a>). Temperatures are also spatially and seasonally unevenly distributed; likewise, the mean annual temperature in the Kunlun, Tianshan, and Altai mountains is below zero, while that of the Turpan Basin is around 14°C (Chen, 2010).




Figure 1. The location of archaeological sites mentioned in this study. 1 Xintala; 2 Wupaer; 3

Xiakalangguer; 4 Shirenzigou; 5 Sidaogou; 6 Xicaozi; 7 Qiongkeke; 8 Tongtian Cave; 9

Ji'rzankal; 10 Yanghai; 11 Jiayi; 12 Shengjindian; 13 Yuergou; 14 Xiaohe; 15 Gumugou; 16 South

Aisikexiaer Cemetery; 17 Wupu; 18 Xihetan; 19 Zhaojiashuimo; 20 Huoshaogou 21

Huoshiliang; 22 Ganggangwa 23 Lifuzhai; 24 Xichengyi; 25 Sanjiao; 26 Mozuizi; 27

Donghuishan; 28 Jingbaoer; 29 Yingwoshu; 30 Sanjiaocheng; 31 Majiayao; 32 Xishanping; 33

Dadiwan; 34 Shannashuzha; 35 Daping; 36 Gaozhuang; 37 Jiangjiazui; 38 Laohuzui; 39 Qiaocun,

the base map was obtained at https://www.ncei.noaa.gov/maps/grid-extract/.

Due to the arid climate, vegetation types here are characterized by expansive deserts (Xinjiang Integrated Expedition Team and Institute of Botany, 1978). Along the rivers in the low-land basins, riparian woodlands are mainly composed of *Populus*, *Elaeagnus*, *Ulmus*, and *Salix* (Chen, 2010). Within the montane belt, vegetation usually changes from grassland (dominated by *Stipa*), coniferous forest (mainly *Picea* and *Larix*), subalpine steppe (mainly *Stipa*), alpine meadows (including *Stipa*, *Carex*, and *Artemisia*), and alpine cushion vegetation (represented by

Androsace, Stellaria media, and Geranium wilfordii), in banded ecoclines from 186 lowest to highest elevation (Chen, 2010; Zheng et al., 2015; Xinjiang Integrated 188 Expedition Team and Institute of Botany, 1978). Wild fruit and nut woodlands are distributed throughout the Tianshan Mountains, especially in the Ili valley, and the main wild fruit trees include Malus sp., Juglans regia, and Prunus spp. (Chen, 2009; 190 191 Abudureheman et al., 2016).

#### 2.2 Prehistoric cultures and agriculture

187

189

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

As an important cultural bridge connecting East and West Asia, northwest China has fostered a variety of cultural communities. The early Neolithic cultures included the Dadiwan and Yangshao, mainly distributed in southern Gansu (Institute of Cultural Relics and Archaeology of Gansu, 2006). Later, people with material culture ascribed to the Majiayao expanded quickly into the Hexi Corridor around 4800 BP (Xie, 2002; Dong et al., 2020b). From 4000-3000 BP, the main archaeological cultures in Gansu consisted of the Xichengyi, Qijia, Siba, and Dongjiatai (Li et al., 2010), and the Shanma and Shajing cultures gradually developed after 3000 BP (Li, 2009; Gansu Provincial Institute of Cultural Relics and Archaeology et al., 2015). In Xinjiang, the prehistoric peoples before 4000 BP were represented by material culture categorized as the Afanasievo and Chemurchek (Shao, 2018). From 4000-3500 BP, the Andronovo Culture expanded into western Xinjiang, and the Tianshanbeilu and Xiaohe cultures occupied the eastern Tianshan and Tarim Basin, respectively (Mei and Shell, 1999; Ruan, 2014; Jia et al., 2017; Shao and Zhang, 2019; Xinjiang Institute of Cultural Relics and Archaeology, 2004, 2014). Since 3500 BP, cultural communities have continually diversified, with more localized groups forming, like the Subeixi Culture in the Turpan Basin (Chen, 2002).

210 Archaeobotanical evidence shows that millet cultivation was already practiced by ca. 7800-7350 BP (Liu et al., 2004; Li, 2018). By at least 5500 years ago, people 211 212 were engaging in an intensive intermixed crop-livestock system by integrating pig 213 maintenance and with millet cultivation (Yang et al., 2022). From 5000-4000 BP, 214 both East Asian millets diffused into the Hexi Corridor, while agricultural practices in Xinjiang were restricted to limited microenvironmental pockets (Zhou et al., 2016; 215 216 Dong et al., 2017, 2018, 2020b; Li, 2021). Since 4000 BP, mixed agricultural systems 217 composed of both East and southwest Asian crops became more prominent; although, 218 barley and wheat had reached northwest China about a millennium prior (Flad et al., 2010; Zhao et al., 2013; Yang et al., 2014; Zhang et al., 2017; Zhou et al., 2016, 2020; 219 220 Jiang et al., 2017a, 2017b; Tian et al., 2021). Stable carbon isotope data also suggest 221 that the consumption of both C<sub>3</sub> and C<sub>4</sub> plants was widely practiced after 4000 BP (Liu et al., 2014; Zhang et al., 2015; An et al., 2017; Wang et al., 2016, 2017; Ma et 222 al., 2016; Qu et al., 2018). Around 3700-3300 BP, wheat and barley gradually 223 224 replaced the millets, becoming the dominant crops within the Hexi Corridor (Zhou et 225 al., 2016). From 3300-2200 BP, agriculture in Xinjiang gradually developed into something more complex and spread to larger areas and more diverse ecozones, as 226 227 evidenced by the diversification of crops, and the appearance of irrigation technology 228 and various types of farming tools (Li, 2021). Meanwhile, secondary crops, such as 229 Vitis vinifera and Ziziphus jujuba, appeared more widely after ca. 2500 BP, indicating a strong concept of land tenure associated with the development of agriculture (Jiang 230 231 et al., 2009, 2013; Li, 2021)

## 3 Archaeobotanical Data and Chronology

232

233

## 3.1 Chronology of the archaeological sites

234 In this study, we present data from seven archaeological sites and have developed a chronology based on AMS <sup>14</sup>C dating through the Beta Analytic Testing Laboratory 235 and Australian Nuclear Science and Technology Organisation. For dating, we focused 236 237 on wheat seeds and wood charcoal, and the calibrated ages were generated using Oxcal 4.4 with IntCal20 (Table 1 and Figure 2) (Reimer et al., 2020). The dating 238 results show that the seven archaeological sites cover a time span between 3900 and 239 240 2000 BP, and the oldest dates come from Xintala, at ca. 3900-3500 BP. The 241 Xiakalangguer, Sidagou, Xicaozi, and Qiongkeke sites fall in-to the period of 3500-242 3000 BP. The chronology for Shirenzigou covers roughly 2700-2000 BP. At Wupaer, we collected wood charcoal samples from two sections, S1 and S3, and the date of the 243 244 S3 section is about 2900-2800 BP. The S1 section shows two different timespans of 245 occupation, specifically ca. 3400-3300 BP and 2500-2300 BP.

Table 1. Dates for the seven archaeological sites in this study.

Site	Latitude	Longitude	Culture	Lab no.	Material	Date (BP)	Calibrated date (2δ, BP)	References
				OZM448	charcoal	3395±30	3815-3561	
				OZM449	charcoal	3515±30	3877-3696	
Xintala	42.22	86.39	Xintala type	OZM450	charcoal	3335±30	3680-3469	Zhao et al., 2013
				OZM451	wheat	3460±35	3835-3593	
				OZL437	wheat	3515±50	3960-3642	
Qiongkeke	43.83	82.75	Andronovo	Beta-642945	charcoal	3220±30	3482-3375	
Qioligheke	45.65	62.73	Andronovo	Beta-642946	charcoal	3320±30	3591-3458	this study
Xiakalangguer	46.74	83.03	Andronovo	Beta-642943	charcoal	3140±30	3447-3327	tills study
Alakalaligguei	40.74	83.03	Andronovo	Beta-642944	charcoal	3070±30	3365-3209	
Sidaogou	43.79	90.19	Nonvyon tymo	OZK664	wheat	3030±50	3362-3075	Dodson et al.,
Sidaogou	ogou 43.79 90.19 Nanwan type	Nanwan type	OZK665	wheat	3080±60	3445-3080	2013	
Xicaozi	44.00	89.68	Unknown	OZM674	wheat	2975±45	3331-2997	2013
				Beta-642939	charcoal	3160±30	3451-3339	
Wupaer	39.28	75.52	Wupaer	Beta-642940	charcoal	2450±30	2544-2361	
wupaei	39.26	13.32	w upaer	Beta-642941	charcoal	2420±30	2515-2351	
				Beta-642942	charcoal	2800±30	2967-2844	this study
				Beta-642947	charcoal	2350±30	2466-2329	•
Shirenzigou	42.56	94.09	94.09 Shirenzigou type	Beta-642948	charcoal	2180±30	2313-2099	
				Beta-642949	charcoal	2150±30	2178-2041	
				-				




Figure 2. The chronology of seven archaeologic sites in this study.

## 3.2 Wood charcoal assemblages

The identification of wood charcoal was accomplished via scanning electron microsemicroscopyope, with 2,960 fragments of charcoal analysed and reported here (Appendix A). Three of the sites are located in oases and wood charcoal assemblages show clear similarities, with a dominance of *Tamarix* wood (Figure 3). In sediments from Xintala, we identified 878 wood charcoal fragments, with *Tamarix* accounting for 74-95%. *Elaeagnus angustifolia* increased across the chronology and reached its highest level (13%) in the latest layer. There were limited occurrences of *Populus*, *Salix*, and cf. *Nitraria*. Wood charcoal from Wupaer also shows an abundance of

Tamarix (ca. 80%), followed by fragments of Populus, Salix, and Chenopodioideae. Fruit tree remains include *Prunus*, usually less than 3% in abundance. At the Xiakalangguer site, Salix and Tamarix account for 44 and 28% of the assemblage respectively, followed by Chenopodioideae (17%). A small number of fragments of Betula and Prunus were also identified.

258

259

260

261

262

266

267

268




Figure 3. Wood charcoal assemblages from seven archaeological sites in northwest China.

In the eastern Tianshan, wood charcoal from three sites revealed an abundance of coniferous wood fragments. At Shirenzigou, wood charcoal fragments from cultural strata included Picea, Juniperus, Tamarix, Populus, Salix, and Rhamnus, with

conifers accounting for over 90% of the fragments. However, 14 wood samples taken from coffins suggest that they are all made from coniferous woods, including *Picea* (11) and *Juniperus* (3). At Sidaogou, wood charcoal from five samples was dominated by *Picea* and *Populus*, followed by *Salix* and *Tamarix*. Progressively over time, *Picea* fragments decreased from 52% to less than 20%, while *Populus* increased quickly from 37% to over 70%. Similarly, *Picea* and *Populus* also constituted a dominant percentage of the Xicaozi assemblage and the other taxa only cover a small percentage, represented by *Prunus*, *Juniperus*, *Salix*, and *Betula*. The Qiongkeke site is located in the Ili Valley, with five taxa identified among 229 wood charcoal fragments. *Prunus* and *Rhamnus* account for 30% each. The proportion of *Picea* is around 20%, followed by *Tamarix* and Maloideae.

In addition, we compiled wood charcoal data from published studies. In the Altai Mountains, wood charcoal from Tongtian Cave indicates that people widely collected *Larix, Picea, Betula, Populus, Salix,* Maloideae, and *Prunus* (Zhou et al., 2020). On the Pamir Plateau, the data we have assembled from the Ji'rzankal Cemetery show that *Populus* was used for making fire tools, *Betula* for wooden plates, *Salix* for wooden sticks, *Juniperus* for fire altars, and *Lonicera* for arrow shafts (Shen et al., 2015). Similarly, in the Turpan Basin, *Populus* was also selected for making fire tools at the Yanghai Cemetery, and there was selective use of a variety of other woods, including *Picea, Spiraea, Tamarix, Betula, Morus, Salix, Clematis,* and *Vitis vinifera* (Jiang, 2022). *Lonicera* was also used for arrow shafts and composite bows at the Jiayi and Shengjindian cemeteries (Nong et al., 2023). *Picea* was widely used at Yuergou for coffin manufacture and firewood (Jiang et al., 2013). While in the Tarim and Hami basins, *Populus* and *Tamarix* were largely used for coffins and wooden utensils, as revealed by studies at the Xiaohe, Gumugou, South Aisikexiaer, and

Wupu cemeteries (Institute of Cultural Relics and Archaeology of Xinjiang, 2007, Zhang et al., 2017, 2019; Wang et al., 2021).

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

In the Hexi Corridor, *Picea* and/or *Juniperus* constituted the dominant portion of wood charcoal fragments in sites located near the Oilian Mountains, such as Xihetan and Zhaojiashuimo (Shen et al., 2018). While wood charcoal from oasis sites, like Huoshaogou, Huoshiliang, and Ganggangwa, also record the abundance of *Tamarix*, and woody Polygonaceae and Salix disappear from later phases of Huoshiliang, presumably due to over harvesting for fuel (Shen et al., 2018, Li et al., 2011). The other sites in this area are characterized by abundant broadleaved taxa, with a small percentage of coniferous wood fragments, such as at the Lifuzhai, Xichengyi, and Sanjiao sites (Wang et al., 2014; Shen et al., 2018; Liu et al., 2019). Meanwhile, wood charcoal assemblages from the Mozuizi and Donghuishan sites suggest a rapid decline of local wood sources, including those of *Picea*, Maloideae, and *Betula* (Shen et al., 2018). Additionally, an abundance of *Prunus* wood fragments was found in these two sites, and people might have transported *Picea* wood over long distances to burn at Donghuishan (Shen et al., 2018). The long-distance transport of *Picea* and *Pinus* was also recognized in the assemblage from the Jingbaoer jade mine (Liu et al., 2021). At the Yingwoshu and Sanjiaocheng sites, abundant Morus wood fragments were identified, possibly indicating the early cultivation of mulberry (Shen et al., 2018).

As with the Hexi Corridor, wood taxa recovered from the western Loess Plateau also suggest a quick decline in the abundance of *Picea*, notably from 37% to less than 4% at Majiayao (Shen et al., 2021). In the assemblage from Xishanping, *Picea*, *Betula*, *Acer*, and *Quercus* decreased markedly after 4600 BP, and *Picea* declined from a peak value of 28% to less than 5%, while Bambusoideae increased sharply (Li

et al., 2012). The sudden spike on abundance of bamboo is thought to be due to rapid successional colonization after significant deforestation or clearing of woody competitive species. Meanwhile, fruit trees, including *Castanea*, *Prunus* (what the wood specialists in this study called *Cerasus* and *Padus*), and *Diospyros* expressed a considerable increase in abundance (Li et al., 2012). The use of fruit tree wood was also recognized in the Dadiwan, Shannashuzha, Daping, and Gaozhuang sites, with the abundance of *Prunus* (these researchers subdivided this group into *Prunus* and *Padus*, which we have clumped together in this study for consistency, since that the wide conception of *Prunus*, that is, *Prunus sensu lato* (s.l.) includes *Prunus sensu stricto* (s.s.), *Amygdalus*, *Cerasus*, *Padus*, as well as *Armeniaca*), Maloideae, and *Ziziphus* (Sun et al., 2013; An et al., 2014; Li et al., 2017).

#### **4 Discussions and Conclusion**

#### 4.1 Wood collection strategies and the transport of conifers

As the result of wood <u>fuel useburning</u>, wood charcoal provides insights into the decision-making process regarding the collection <u>of fuelstrategies</u>. In this study, we found that wood charcoal assemblages from all oasis sites were dominated by *Tamarix*. Most species from the *Tamarix* genus are deciduous shrubs, generally 2-5 meters high, with slender and soft branches (Yang and Gaskin, 2012). The twigs are often browsed by sheep, camel, and donkey, and the branches can serve as a rapidly-regenerating fuel (Editorial Board of Flora of China, CAS, 1990). Therefore, this widely-distributed, arid-tolerant, and rapid-growing shrubby *Tamarix*, <u>might-would have</u> constituted the best fuel for ancient oases groups. For the archaeological sites located in mountainous areas, wood fragments from coniferous trees are more

prevalent. For example, abundant *Picea* and *Juniperus* wood fragments were found at Shirenzigou in the eastern Tianshan. Similarly, *Picea/Juniperus* constitutes the dominant portion of the fragments from sites near the Qilian Mountains (Shen et al., 2018). All of the assemblages show that people were largely opportunistic in their choices and the availability of wood sources played a key role in the wood collection strategies.

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

Additionally, as wood resources in arid northwest China are relatively limited, coping with localized wood shortages would have been an issue that people inevitably dealt with. Among these wood charcoal assemblages, we found that there are some fragments of coniferous woods that likely represent people traveling over long distances on collection trips. The earliest known evidence might come from Donghuishan (3700-3400 BP), in which *Picea* charcoal experienced a sharp decrease and then suddenly increased to its highest level (Shen et al., 2018). Given that spruce forests are very slow to regenerate, the sudden increase of spruce fragments was likely the result of long-distance collection from the Oilian Mountains (Shen et al., 2018). Generally, spruce wood has preferential properties, as its timber is straight and tall, and easily worked, presumably contributing to the selection and transportation of this specific species. Since 2500 BP, the long-distance collection of coniferous woods seems to have been a more regular activity, as evidenced at the Jingbaoer jade mine, where Picea and Pinus wood fragments are recovered well outside their natural ecological distribution (Liu et al., 2021). In the Turpan Basin, *Picea* wood fragments were found in sediments from a series of Subeixi sites, which may have been collected from the Tianshan Mountains (Jiang et al., 2013; Jiang, 2022).

In addition to noting the likely long-distance collection of coniferous woods, the

abundance of conifers in most of our study sites hints to the likelihood that people might also have a preference for this specific wood type. At Sidaogou, spruce wood fragments comprise more than 60% of the total fragment assemblage. Similarly, charcoal from Majiayao recorded spruce fragments as the most used taxon right from the onset of when people settled down at the location (Shen et al., 2021). Meanwhile, the exclusive use of coniferous wood for coffin construction is also recognizable in this study. At Shirenzigou, the analysis of 14 wooden coffins show that they were all made of coniferous woods. However, in sediments from the site, we found a variety of carbonized wood types, including *Tamarix*, *Populus*, *Rhamnus*, *Salix*, etc. Historically, a preference towards coniferous woods is widely noted in ancient China (Ding, 2022), and archaeological wood studies in Central Asia have also noted similar patterns (Spengler and Willcox 2013). Many ethnographic and historical references to ritual juniper twig burning as incense are noted from across Inner Asia. The fact that the wooden coffins at Shirenzigou are all constructed from conifers, suggests that the ritual significance of the resinous trees may stretch much further back in time. Ultimately, we conclude that aAn awareness of the properties and special meaning of these woods probably plays a key role in their wide use.

# 4.2 Collection and cultivation of fruit trees

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

In addition to the prehistoric expansion of agricultural systems, the significant amounts of fruit wood fragments in our study may imply that the anthropogenic processes were increasing the density of fruit trees near human settlements. Presently, scholars continue to grapple with the question of what evidence is necessary to differentiate between wild foraging, conservation of economically significant trees and low-investment cultivation of wild populations (Dal Martello et al., 2023). In our

study, fruit wood fragments before 4600 BP were usually found in low percentages, indicating limited collection of seasonally available wild fruits (Sun et al., 2013; Li et al., 2017; Shen et al., 2021). Roughly between 4600-4300 BP, *Castanea*, *Prunus*, and *Diospyros* charcoal shows a rapid increase in abundance at Xishanping on the western Loess Plateau (Li et al., 2012). Pollen data at this time also demonstrates that *Castanea* became the dominant broadleaved taxon, which is quite different from the reconstructed natural vegetation, likely indicating the management of wild chestnut forests or at least that humans were choosing not to cut these trees down, increasing their populations (Li et al., 2007). Also, archaeobotanical records at this site illustrate that a complex agricultural system based on a variety of crops, including millets\_ (*Setaria italica* and *Panicum miliaceum*), rice (*Oryza sativa*), oats (*Avena* sp.), soybean (*Glycine soja*), and buckwheat (*Fagopyrum* sp.), appeared synchronously with the management of chestnut. This cooccurrence probably suggests that the exploitation of secondary crops was closely related to and underpinned by the well-organized agricultural system.

During the period from 4300 to 3500 years ago, there is an increase in the abundance of fruit wood remains in Xinjiang and the Hexi Corridor. For example, *Elaeagnus angustifolia* charcoal was found throughout the whole section and shows a gradually increasing trend at Xintala. In the Hexi Corridor, *Prunus* wood fragments were found discovered in great abundance at Mozuizi and Donghuishan, far higher than its percentage is believed to have been in the natural vegetation, possibly showing an intensive collection of *Prunus* (Shen et al., 2019). However, there is no clear sign of fruit management during this period, given that a wide range of wild fruit types, such as *Nitraria* and *Cotoneaster* were also widely exploited (Zhou et al., 2016; Shen et al., 2019). Meanwhile, previous studies show that, although a mixed

agricultural system consisting of both millets, wheat, and barley existed in Xinjiang and the Hexi Corridor after 4000 BP, people still relied heavily on animal herding and/or feeding (Dong et al., 2020b; Li, 2021).

From 3500-2500 BP, the cultivation or maintenance of *Prunus* and *Morus* trees was probably adopted into the agricultural system. As in Wupaer, located in the Kashgar oasis, the presence of *Prunus* charcoal remains is-recovered beyond its the natural distribution of the tree and the climatic conditions around the site are not suitable for the growth of *Prunus*, likely resulted from anthropogenic planting. On the other hand, considering that the distribution of wild *Prunus* trees had largely shrunk or even disappeared presumably due to long-term human activity, we should still be cautious about this conclusion. Almost at the same time, people in the Hexi Corridor probably also started engaging in horticultural practices, supported by the abundant discovery of *Morus* charcoal (Shen et al., 2019). Synchronously, a high-yield wheat and barley farming system was developed in the Hexi Corridor (Zhou et al., 2012), and a more intensified agricultural system developed in Xinjiang (Li, 2021), likely providing a fundamental basis for the exploration of delayed-return perennial crops.

After 2500 BP, the cultivation of fruit trees was probably a-widely practiced in northwest China. For instance, evidence from the Turpan Basin shows the presence of *Morus* woods and *Vitis vinifera* stems at the Yanghai cemetery (Jiang, 2022; Jiang et al., 2009), *Vitis vinifera* seeds in the Shengjindian cemetery (Jiang et al., 2015), and *Ziziphus jujuba* stones in the Yuergou site (Jiang et al., 2013). At the Sampula cemetery, fruit, nut and seed types were more abundant, including *Prunus-persica*, *P. armeniaca*, *Juglans regia*, *Coix lacryma-jobi*, etc. (Jiang et al., 2008). The appearance of such a rich and diverse array of fruit crops indicates that people in northwest China

had developed a complex indigenous knowledge to for survivale in this hyper arid environment and conducted more and more frequent exchange across the Eurasian continent.

## 4.3 Indigenous knowledge of plant resources

Due to the extreme arid climate, wooden objects found in our study area are usually well-preserved and the data suggest that people might have also captured the knowledge of deliberately selecting certain types of woods when making various utensils. For example, within the Subeixi groups in the Turpan Basin, *Lonicera* was harvested from wild stands for making arrow shafts at Jiayi and Shengjingdian (Nong et al., 2023). At the Yanghai cemetery, *Betula* was selected for making dippers or ladles, for its rigidity; flammable *Populus* and *Picea* were used for fire tool manufacture (Jiang et al., 2018, 2021). People at this time also used *Lithospermum officinale* seeds for decoration (Jiang et al., 2007a), *Nitraria tangutorum* for making necklaces (Jiang, 2022), and *Cannabis* for ritualized consumption and/or medical purposes, as revealed in both the Turpan Basin (Jiang et al., 2006, 2007b, 2016) and the Pamir Plateau (Ren et al., 2019).

Similarly, on the Pamir Plateau, *Betula*, which has high rigidity and density, and homogeneous texture, was selected for making wooden plates (Shen et al., 2015).

Additionally, the study of other wooden objects it appears suggests that people specifically chose flammable *Populus* wood to make fire tools; *Salix*, with long and straight branches, was used for fashioning wooden sticks; sweetresinous-scented *Juniperus* was the preferred choice for making fire altars, and *Lonicera* was selected for arrow shaft manufacture (Shen et al., 2015). Such conscious utilization of different wood properties illustrates the ingenuity of these ancient people. Although the current

archaeobotanical research data related to wooden utensils is are still limited, studies from the Turpan Basin and the Pamir Plateau clearly suggest that the conscious selection of wood types for specific properties was a particularly pronounced practice after 2500 BP, especially among cultural contexts of a well-established agriculture based on with millets, wheat, and barley. Meanwhile, the appearance of horticulture based on a variety of secondary crops at the time indicated a more settled lifestyle, which might provide opportunities for prehistoric people to fully explore and make the best use of the indigenous plant resources.

#### 4.4 Anthropogenic deforestation

Presumably-Largely due to via slash and burn agriculture, people have largely altered terrestrial ecosystems across the globe (Zong et al., 2007; Schlütz et al., 2009; Li et al., 2009; Neumann et al., 2012; Innes et al., 2013; Ma et al., 2020; Zheng et al., 2021). For northwest China, wood charcoal data in this study show that, apart from diversified cultural adaptions, human-induced landscape alteration also occurred widely, not only throughout the whole history of agricultural activity, but also across different vegetation contexts. Along the Tianshan mountains, For example, pollen records from the Bosten and Balikun lakes suggest a relatively stable climate during 3900-3500 BP, and a long-term increase of humidity after 3800 BP (Chen et al., 2006; Huang et al., 2009; An et al., 2012). However, wood charcoal data from Sidaogou (3400-3000 BP) in the eastern Tianshan recorded a significant decrease in abundance of spruce wood fragments (Figure 4). Meanwhile, *Tamarix* and *Salix* nearly disappeared in the later stage, showing that wood cutting caused thea sharp attenuation of spruce forests and broadleaved woodlands was caused by intensive wood cutting rather than climate change. Similarly, *Tamarix* charcoal from the

Xintala (3900-3500 BP)section \_\_in the Yanqi Oasis firstly increased and then decreased to its lowest level in the upper layer, suggesting that continuous woodeutting resulted in the decline of *Tamarix* shrubs. At the same time, *Populus* and *Salix* charcoal disappeared in the middle layer, implying that local riparian woodlands were largely deforested.




Figure 4. The wood charcoal and pollen records show synchronous deforestation of spruce forests across all of northwest China. (a) the change of *Picea* wood charcoal (bar) and pollen (curve) from archaeological sites including Sidaogou, Donghuishan (Zhou et al., 2012; Shen et al., 2018), Majiayao (Zhou, 2009; Shen et al., 2021), and Xishanping (Li et al., 2007, 2012).

The column chart shows the stratum layer. (b) the comparison of spruce forests between prehistoric times and now, the squares represent archaeological sites with *Picea* charcoal

# remains and the red areas show the current distribution of spruce forests in northwest China (after Hou, 2019).

The Neolithic deforestation and reduction in range of spruce forests have also
been widely recognized across the western Loess Plateau and the Hexi Corridor. At-
On the western Loess Plateau, high-resolutionved (ca. ~5 year incrementss) stalagmite
$\delta^{18}O$ data recorded no abrupt climate changes at around 5300-5100 BP and 4600 BP
(Tan et al., 2020). While, the Majiayao site, wood charcoal recorded record from the
Majiayao site showed the a rapid decline of Picea from its highest level of nearly 40%
to the lowest of less than 4% during the early stages of the site's occupation at ca.
5300-5100 BP, implying that anthropogenic exploration exerted a significant impact
on local spruce forests (Figure 4a) (Shen et al., 2021). Not far from Majiayao, wood
charcoal from the Xishanping section revealed a similar pattern, with Picea, Betula,
Acer, Ulmus, and Quercus, illustrating a marked decrease after 4600 BP, while
Bambusoideae quickly colonized after the clearing of the original forest (Li et al.,
2012). In the Hexi Corridor, <u>studies of</u> wood charcoal <u>fragments</u> <u>assemblagesdata</u> _
from the Mozuizi and Donghuishan sites <u>also</u> show a quick decline in plant diversity
concurrent with human settlement, and the percentage of Picea from Donghuishan
recorded experienced a sharp decrease (Figure 4) (Shen et al, 2018). Similarly, wood
charcoal fragments from Huoshiliang show that Salix and Polygonaceae almost
disappear, likely due to the large demand for fuel used in bronze smelting activities
(Li et al., 2011). Collectively, we interpret the broader trend throughout all of these
wood charcoal assemblages as revealing a rather rapid process of deforestation across
northwest China, especially shown in the large-scale reduction in spruce forests. Our
results are also supported by evidence from pollen records, especially <i>Picea</i> pollen
from Majiayao (Zhou, 2009), Xishanping (Li et al., 2007), Donghuishan (Zhou et al.,

2012), and other sections from the Loess Plateau (Zhou and Li, 2011). All of these records document considerable reduction in spruce forests (Figure 4a). Today, the distribution of spruce forests has shrunk down to a few constrained small forest patches (Figure 4b).

#### 5 Data availability

The datasets of archaeobotanical wood charcoal records in northwest China including taxa types, absolute counts of wood charcoal fragments, and the locations and AMS <sup>14</sup>C dates of each archaeological site are available at the open-access repository Zenodo (Shen et al., 2023; <a href="https://doi.org/10.5281/zenodo.8158277">https://doi.org/10.5281/zenodo.8158277</a>).

#### **6 Summary**

The synthesis of wood charcoal data from nearly 40 archaeological sites shows that prehistoric human-environmental interactions in northwest China were closely related to the development of agriculture and considerably more complicated than previously thought (Figure 5). Although anthropogenic deforestation occurred throughout the whole period, most evidently relating to the decline of spruce forests, people also actively applied a range of adaptive strategies to survive in this harsh environment. As early as 4600 BP, people on the western Loess Plateau might have started managing or at least conserving chestnut trees, likely underpinned by the development of a complex agricultural system. Since ca. 3500 BP, with the appearance of high-yielding agriculture based on wheat and barley in Xinjiang and the Hexi Corridor, people appear to have been planting perennial tree crops, such as *Prunus* and *Morus*.

Additionally, they likely engaged in long-distance transportation of preferred woods, specifically coniferous trees. After 2500 BP, people successfully mastered a wide

range of adaption strategies along the ancient Silk Road, as they began manufacturing wooden utensils with conscious selection of wood properties. Moreover, the consumption of a further diversity of fruit types, including grapes, signalled more intensive horticultural practices and complex social structure.





Figure 5. A summary of prehistory human-environmental interactions in northwest China.

**Appendix A**. The selected scanning electron microscopic images of wood charcoal in Xinjiang. (a-c) *Picea*. (d-f) *Prunus*. (g-I) *Populus*. (j-l) *Tamarix*.



**Author contributions.** HS and XL designed the archaeobotanical dataset; HS was responsible for construction of the database; HS performed numerical analyses and organized the manuscript, and XZ, RS, PJ and AB revised the draft of the paper. All authors discussed the results and contributed to the final paper.

**Competing interests.** The contact author has declared that none of the authors has any competing interests.

5/3	<b>Acknowledgements.</b> We sincerely thank Ming Ji and Hongbin Zhang for their help in
574	the wood charcoal sample collection, and Nan Sun for her assistance with data
575	collection.
576	
577	Financial support. This research has been supported by the National Natural Science
578	Foundation of China (grant no. 42002202), the Youth Innovation Promotion
579	Association of Chinese Academy of Sciences (grant no. 2022071), and the National
580	Key Research and Development Program of China (grant no. 2022YFF0801502).
581	
582	References
583 584 585	Abudureheman, B., Liu, H-L., Liu, N., Zhang, D-Y., and Guan, K.: Spermatophyte floras in the wild fruit forest in Tianshan Mountains, Arid Land Geogr., 39(4), 828–834, 2016 (in Chinese with English abstract).
586 587 588 589 590 591 592 593	Allué, E. and Zaidner, Y.: The charcoal assemblage from Nesher Ramla, Israel: A contribution to the paleo-environmental dataset from Marine Isotope Stage (MIS) 5 in the Levant, Quat. Int., 624, 117-127, https://doi.org/10.1016/j.quaint.2021.04.025, 2022. Mas, B., Oms, F. X., and Allué, E.: Anthropogenic impacts on vegetation landscapes and environmental implications during the Middle-Late Holocene in the Iberian Central Pre-Pyrenees: An anthracological approach, Rev. Palaeobot. Palynol., 300, 104624, https://doi.org/10.1016/j.revpalbo.2022.104624, 2022.
594 595 596	An, C., Feng, Z., and Tang, L.: Environment change and cultural response between 8000 and 4000 cal. yr BP in the western Loess Plateau, NW China, J. Quat. Sci., 19, 529-535, https://doi.org/10.1002/jqs.849, 2004.
597 598 599	An, C., Tang, L., Barton, L., and Chen, F.: Climatic Change and Cultural Response around 4,000 cal. yr B.P. in the western part of the Chinese Loess Plateau, Quat. Res., 63(3), 347-352, https://doi.org/10.1016/j.yqres.2005.02.004, 2005.
600 601 602	An, C., Wang, L., Ji, D., Chen, F., and Wang, P.: The temporal and spatial changes of Neolithic cultures in Gansu-Qinghai region and possible environmental forcing, Quat. Sci. 26(6), 923-927, 2006 (in Chinese with English abstract).
603 604	An, C., Wang, W., Duan, F., Huang, W., and Chen, F.: Environmental changes and cultural exchange between East and West along the Silk Road in arid Central

605 606	Asia, Acta Geogr. Sin., 72(5), 875–891, http://doi.org/ 10.11821/dlxb201705009, 2017 (in Chinese with English abstract).
607	An, C-B., Li, H., Dong, W., Chen, Y., Zhao, Y. and Shi, C.: How prehistoric humans
608	use plant resources to adapt to environmental change: A case study in the
609	western Chinese Loess Plateau during Qijia Period, Holocene, 24(4), 512–517,
610	https://doi.org/10.1177/0959683613520259, 2014.
611	An, C-B., Lu, Y., Zhao, J., Tao, S., Dong, W., Li, H., Jin, M., and Wang, Z.: A high-
612	resolution record of Holocene environmental and climatic changes from Lake
613	Balikun (Xinjiang, China): Implications for central Asia, Holocene, 22(1), 43-52,
614	https://doi.org/10.1177/0959683611405244, 2012.
615	ArchaeoGLOBE Project: Archaeological assessment reveals Earth's early
616	transformation through land use, Science, 365, 897–902,
617	https://doi.org/10.1126/science.aax1192, 2019.
618	Asouti, E. and Austin, P.: Reconstructing woodland vegetation and its exploitation by
619	past societies, based on the analysis and interpretation of archaeological wood
620	charcoal macro-remains, Environ. Archaeol., 10(1), 1–18,
621	https://doi.org/10.1179/env.2005.10.1.1, 2005.
622	Asouti, E. and Kabukcu, C.: Holocene semi-arid oak woodlands in the Irano-
623	Anatolian region of Southwest Asia: natural or anthropogenic? Quat. Sci. Rev.,
624	90, 158–182, https://doi.org/10.1016/j.quascirev.2014.03.001, 2014.
625	Asouti, E., Kabukcu, C., White, C. E., Kuijt, I., Finlayson, B., and Makarewicz, C.:
626	Early Holocene woodland vegetation and human impacts in the arid zone of the
627	southern Levant, Holocene, 25(10): 1565–1580,
628	https://doi.org/10.1177/0959683615580199, 2015.
629	Atahan, P., Itzstein-Davey, F., Taylor, D., Dodson, J., Qin, J., Zheng, H., and Brooks,
630	A.: Holocene-aged sedimentary records of environmental changes and early
631	agriculture in the lower Yangtze, China, Quat. Sci. Rev., 27(5), 556-570,
632	https://doi.org/10.1016/j.quascirev.2007.11.003, 2008.
633	Bellwood, P. (Eds.): The first farmers: the origins of agricultural societies, Blackwell
634	Pub, London, ISBN 0-631-20565-9, 2005.
635	Bocquet-Appel, J. P.: When the world's population took off: The springboard of the
636	Neolithic Demographic Transition, Science, 333(6042), 560–561,
637	10.1126/science.1208880, 2011.
638	Boivin, N. L., Zeder, M. A., Fuller, D. Q., Crowther, A., Larson, G., Erlandson, J. M.,
639	Denham, T., and Petraglia, M. D.: Ecological consequences of human niche
640	construction: Examining long-term anthropogenic shaping of global species
641	distributions, Proc. Natl. Acad. Sci. USA, 113(23), 6388–6396,

642	https://doi.org/10.1073/pnas.1525200113, 2016.
643 644	Boivin, N., Crassard, R., and Petraglia, M. (Eds.): Human Dispersal and Species Movement: From Prehistory to the Present, Cambridge University Press,
645	Cambridge, ISBN 9781316686942, 2017.
646	Chen, F. H., Huang, X. Z., Yang, M. L., Yang, X. L., Fan, Y. X., and Zhao, H.:
647 648	Westerly dominated Holocene climate model in arid Central Asia—case study on Bosten Lake, Xinjiang, China, Quat. Sci., 26(6), 881-887, 2006 (in Chinese with
649	English abstract).
650	Chen, G.: The origin of Subeixi Culture and its relationship with other cultures, West
651	Reg. Studies, 2, 11–18, 2002 (in Chinese).
652	Chen, K., Mei, J., and Qian, W.: The Silk Road and the exchange of early bronze and
653 654	iron technology, West. Reg. Studies, 2, 127-150, 2018 (in Chinese with English abstract).
654	abstract).
655	Chen, X. (Eds.): Physical geography of arid regions of China, Science Press, Beijing,
656	ISBN 9787030275837, 2010.
657	Chen, X.: Construction of Xinjiang wild fruit resources database system, Master
658	thesis, Xinjiang Agricultural University, Xinjiang, 2009.
659	Cowie, R. H., Bouchet, P., and Fontaine, B.: The sixth mass extinction: fact, fiction or
660	speculation? Biol. Rev., 97(2), 640–663, https://doi.org/10.1111/brv.12816,
661	2022.
662	Crowther, A., Lucas, L., Helm, R, Horton, M., Shipton, C., Wright, H. T., Walshaw,
663 664	S., Pawlowicz, M., Radimilahy, C., Douka, K., Picornell-Gelabertm, L., Fuller, D. Q., and Boivin, N. L.: Ancient crops provide first archaeological signature of
665	the westward Austronesian expansion, Proc. Natl. Acad. Sci. USA, 113(24),
666	6635–6640, https://doi.org/10.1073/pnas.1522714113, 2016.
667	Crutzen, P.: Geology of mankind, Nature, 415, 23, https://doi.org/10.1038/415023a,
668	2002.
669	Dal Martello, R., von Baeyer, M., Hudson, M., Bjorn, R. G., Leipe, C., Zach, B., Mir-
670	Makhamad, B., Billings, T. N., Muñoz Fernández, I. M., Huber, B., Boxleitner,
671	K., Lu, J-C., Chi, K-A., Liu, H. L., Kistler, L., and Spengler, R.N.: The
672	Domestication and Dispersal of Large-Fruiting Prunus spp.: A Metadata
673 674	Analysis of Archaeobotanical Material, Agronomy, 13, 1027, https://doi.org/10.3390/agronomy13041027, 2023.
014	1111ps.//doi.01g/10.5570/ag10110111y15041021, 2025.
675	Ding, J.: On the custom about planting pines and cypresses in cemeteries in ancient
676	China and its cultural connotation of funeral, J. Beijing Forestry University (Soc
677	Sci.), 21(2), 39–46, http://doi.org/10.13931/j.cnki.bjfuss.2021066, 2022.

- Dodson, J. R., Li, X., Zhou, X., Zhao, K., Sun, N., and Atahan, P.: Origin and spread
- of wheat in China, Quat. Sci. Rev., 72(2): 108-111,
- 680 https://doi.org/10.1016/j.quascirev.2013.04.021, 2013.
- Dodson, J., Li, X., Ji, M., Zhao, K., Zhou, X., and Levchenko, V.: Early bronze in two
- Holocene archaeological sites in Gansu, NW China, Quat. Res., 72(3), 309–314,
- 683 https://doi.org/10.1016/j.yqres.2009.07.004, 2009.
- Dong, G. H., Zhang, S. J., Yang, Y. S., Chen, J. H., and Chen, F.H.: Agricultural
- intensification and its impact on environment during Neolithic Age in northern
- China, Chin. Sci. Bull., 61, 2913–2925, https://doi.org/10.1360/N972016-00547,
- 687 2016 (in Chinese with English abstract).
- Dong, G., Du, L., Yang, L., Lu, M., Qiu, M., Li, H., Ma, M., and Chen, F.: Dispersal
- of crop-livestock and geographical-temporal variation of subsistence along the
- Steppe and Silk Roads across Eurasia in prehistory, Sci. China Earth Sci., 65(7),
- 691 1187–1210, https://doi.org/10.1007/s11430-021-9929-x, 2022b.
- 692 Dong, G., Li, R., Lu, M., Zhang, D., and James, N.: Evolution of human-
- 693 environmental interactions in China from the Late Paleolithic to the Bronze Age,
- 694 Prog. Phys. Geogr.: Earth Environ., 44(2), 233–250,
- 695 https://doi.org/10.1177/0309133319876802, 2020a.
- 696 Dong, G., Lu, Y., Zhang, S., Huang, X., and Ma, M.: Spatiotemporal variation in
- human settlements and their interaction with living environments in Neolithic
- and Bronze Age China, Prog. Phys. Geogr.: Earth Environ., 46(6), 949–967,
- 699 https://doi.org/10.1177/03091333221087992, 2022a.
- 700 Dong, G., Xin, J., An, C., Chen, F., Zhao, Y., Tao, S., and Ma, M.: Mid-Holocene
- 701 climate change and its effect on prehistoric cultural evolution in eastern Qinghai
- 702 Province, China, Quat. Res., 77(1), 23-30,
- 703 https://doi.org/10.1016/j.yqres.2011.10.004, 2012.
- 704 Dong, G., Yang, Y., Han, J., Wang, H., and Chen, F.: Exploring the history of cultural
- exchange in prehistoric Eurasia from the perspectives of Crop diffusion and
- 706 consumption, Sci. China Earth Sci., 47, 530–543,
- 707 https://doi.org/10.1007/s11430-016-9037-x, 2017.
- 708 Dong, G., Yang, Y., Liu, X., Li, H., Cui, Y., Wang, H., Chen, G., Dodson, J., and
- 709 Chen, F.: Prehistoric trans-continental cultural exchange in the Hexi Corridor,
- 710 northwest China, Holocene, 28, 621–628,
- 711 https://doi.org/10.1177/0959683617735585, 2018.
- 712 Dong, G., Yang, Y., Ren, L., and Ma, M. (Eds.): Subsistence economy and human-
- environment interaction in prehistoric Hexi Corridor, Science Press, Beijing,
- 714 ISBN 978-7-03-048932-6, 2020b.

- Dong, G.: Neolithic cultural evolution and its environmental driving force in the
- Gansu-Qinghai region: problems and perspectives, Mar. Geol. Quat. Geol.,
- 717 33(4), 67-75, 2013 (in Chinese with English abstract).
- 718 Editorial Board of Flora of China, Chinese Academy of Sciences.: Flora of China,
- 719 Volume 13, Science Press, Beijing & St. Louis, 146–165, 1990.
- 720 Fall, P. L., Falconer, S. E., and Lines, L.: Agricultural intensification and the
- secondary products revolution along the Jordan Rift, Hum. Ecol., 30(4), 445–
- 722 482, https://doi.org/10.1023/A:1021193922860, 2002.
- Flad, R., Li, S., Wu, X., and Zhao, Z.: Early wheat in China: results from new studies
- at Donghuishan in the Hexi Corridor. Holocene, 20(6), 955-965,
- 725 https://doi.org/10.1177/0959683609358914, 2010.
- Fuller, D. Q., van Etten, J., Manning, K., Castillo, C., Kingwell-Banham, E.,
- Weisskopf, A., Qin, L., Sato, Y-I., and Hijmans, R. J.: The contribution of rice
- agriculture and livestock pastoralism to prehistoric methane levels: An
- archaeological assessment, Holocene, 21(5), 743–759,
- 730 https://doi.org/10.1177/0959683611398052, 2011a.
- Fuller, D., Boivin, N., Hoogervorst, T., and Allaby, R.: Across the Indian Ocean: The
- prehistoric movement of plants and animals, Antiquity, 85(328), 544-558.
- 733 https://doi.org/10.1017/S0003598X00067934, 2011b.
- Gansu Provincial Institute of Cultural Relics and Archaeology, Institute of Historical
- 735 Metallurgy and Materials, USTB, Institute of Archaeology, CASS, and
- Northwest University: The Xichengyi site in Zhangye city, Gansu, Archaeology,
- 737 15, 14–25, 2015.
- Hou, G., Liu, F., Liu, C., and Fang, X.: Prehistorical Cultural Transition Forced by
- 739 Environmental Change in Mid-Holocene in Gansu-Qinghai Region. Acta Geogr.
- 740 Sin., 64(1), 53-58, 2009 (in Chinese with English abstract).
- Hou, X.: 1:1 million vegetation map of China. National Tibetan Plateau/Third Pole
- Environment Data Center, Beijing, 2019.
- Huang, X. Z., Chen, F. H., Fan, Y. X., and Yang, M. L.: Dry late-glacial and early
- Holocene climate in arid Central Asia indicated by lithological and palynological
- evidence from Bosten Lake, China, Quat. Int., 194, 19-27,
- 746 <u>https://doi.org/10.1016/j.quaint.2007.10.002, 2009.</u>
- Innes, J. B., Blackford, J. J., and Rowley-Conwy, P. A.: Late Mesolithic and early
- Neolithic forest disturbance: A high resolution palaeoecological test of human
- impact hypotheses, Quat. Sci. Rev., 77, 80–100,
- 750 https://doi.org/10.1016/j.quascirev.2013.07.012, 2013.

- 751 Institute of Cultural Relics and Archaeology of Gansu. (Eds.): The excavation report
- of Dadiwan site, Qin'an, Cultural Relics Publishing House, Beijing, ISBN
- 753 9787501018024, 2006.
- 754 Institute of Cultural Relics and Archaeology of Xinjiang.: A brief excavation report
- on Xiaohe grave yard in Luobupo, Xinjiang Autonomous Region, Cult. Relics,
- 756 10, 4–42, 2007.
- Janick, J.: The origins of fruit, fruit growing, and fruit breeding, Plant Breed Rev., 25,
- 758 255–320, https://doi.org/10.1002/9780470650301.ch8, 2005.
- Jia, P., Betts, A., Cong, D., Jia, X., and Dupuy, P.: Adunqiaolu: New evidence for the
- Andronovo in Xinjiang, China, Antiquity, 91(357), 621–639,
- 761 https://doi.org/10.15184/aqy.2017.67, 2017.
- Jiang, H. E. (Eds.): Agricultural activities and plant utilization of the ancient Yanghai
- people, Turpan of Xinjiang, China. Science Press, Beijing, ISBN
- 764 9787030720054, 2022.
- Jiang, H. E., Li, X., Ferguson, D. K., Wang Y-F., Liu, C-J., and Li, C-S.: The
- discovery of Capparis spinosa L. (Capparidaceae) in the Yanghai Tombs (2800
- years B.P.), NW China, and its medicinal implications, J. Ethnopharmacol., 113,
- 768 409–420, https://doi.org/10.1016/j.jep.2007.06.020, 2007b.
- Jiang, H., Feng, G., Liu, X., Cao, H., Wang, S., Ma, L., and Ferguson, D. K.: Drilling
- wood for fire: discoveries and studies of the fire-making tools in the Yanghai
- cemetery of ancient Turpan, China, Veget. Hist. Archaeobot., 27, 197–206,
- 772 https://doi.org/10.1007/s00334-017-0611-5, 2018.
- Jiang, H., Li, C-S., Cao, H., Shading, P., and Cheng, Y-M.: Wood utilization during
- the Late Bronze to Early Iron Age in the Turpan Basin of Xinjiang, China, with
- special emphasis on Betula (Betulaceae), SAGE Open, 11(3),
- 776 https://doi.org/10.1177/21582440211046950, 2021.
- Jiang, H., Wang, L., Merlin, M. D., Clarke, R.C., Pan, Y., Zhang, Y., Xiao, G., and
- 778 Ding, X.: Ancient Cannabis burial shroud in a Central Eurasian Cemetery, Econ.
- 779 Bot., 70(3), 213–221, https://doi.org/10.1007/s12231-016-9351-1, 2016.
- Jiang, H., Wu, Y., Wang, H., Ferguson, D. K., and Li, C-S.: Ancient plant use at the
- site of Yuergou, Xinjiang, China: implications from desiccated and charred plant
- remains, Veg. Hist. Archaeobot., 22, 129–140, https://doi.org/10.1007/s00334-
- 783 012-0365-z, 2013.
- Jiang, H., Zhang, Y., Lü, E., and Wang, C.: Archaeobotanical evidence of plant
- 785 utilization in the ancient Turpan of Xinjiang, China: a case study at the
- Shengjindian cemetery, Veget. Hist. Archaeobot., 24, 165–177,
- 787 https://doi.org/10.1007/s00334-014-0495-6, 2015.

- Jiang, H-E., Li X., Liu, C-J., Wang, Y-F., and Li, C-S.: Fruits of Lithospermum
- officinaleL. (Boraginaceae) used as an early plant decoration (2500 years BP) in
- 790 Xinjiang, China, J. Archaeol. Sci., 34, 167–170,
- 791 https://doi.org/10.1016/j.jas.2006.04.003, 2007a.
- Jiang, H-E., Li, X., Zhao, Y-X., Ferguson, D. K., Hueber, F., Bera, S., Wang, Y-F.,
- 793 Zhao, L-C., Liu, C-J., and Li, C-S.: A new insight into Cannabis sativa
- 794 (Cannabaceae) utilization from 2500-year-old Yanghai Tombs, Xinjiang, China,
- 795 J. Ethnopharmacol, 108, 414–422, https://doi.org/10.1016/j.jep.2006.05.034,
- 796 2006.
- Jiang, H-E., Wang, B., Li, X., Lü, E-G., and Li, C-S.: A consideration of the involucre
- remains of Coix lacryma-jobi L. (Poaceae) in the Sampula Cemetery (2000 years
- 799 BP), Xinjiang, China, J. Archaeol. Sci., 35, 1311–1316,
- 800 https://doi.org/10.1016/j.jas.2007.09.006, 2008.
- Jiang, H-E., Zhang, Y-B., Li, X., Yao, Y-F., Ferguson, D. K., Lü, E-G., and Li, C-S.:
- 802 Evidence for early viticulture in China: proof of a grapevine (Vitis vinifera, L.
- Vitaceae) in the Yanghai Tombs, Xinjiang, J. Archaeol. Sci., 36(7), 1458–1465,
- 804 https://doi.org/10.1016/j.jas.2009.02.010, 2009.
- Jiang, Y. C., Chen, G. K., and Li, S. C.: Analysis of flotation results of the Xichengyi
- site in the Zhangye city, Gansu, Huaxia Archaeol., 1, 62–68, 2017a (in Chinese
- with English abstract).
- Jiang, Y. C., Wang, H. and Li, S. C.: Flotation result of the Donghuishan site in the
- Minle city, Gansu. Archaeol. Cult. Relics, 1, 62–68, 2017b (in Chinese with
- 810 English abstract).
- Jones, M. K., Hunt, H., Lightfoot, E., Lister, D., Liu, X. Y., and Matuzeviciute, G.
- M.: Food globalization in prehistory, World Archaeol., 43, 665–675,
- 813 https://doi.org/10.1080/00438243.2011.624764, 2011.
- 814 Kabukcu, C.: Identification of woodland management practices and tree growth
- conditions in archaeological fuel waste remains: A case study from the site of
- Catalhöyük in central Anatolia, Turkey, Quat. Int., 486, 282-297,
- 817 https://doi.org/10.1016/j.quaint.2017.03.017, 2017.
- 818 Kaplan, J. O., Krumhardt, K. M., and Zimmermann, N.: The prehistoric and
- preindustrial deforestation of Europe, Quat. Sci. Rev., 28(27), 3016–3034,
- 820 https://doi.org/10.1016/j.quascirev.2009.09.028, 2009.
- 821 Knapp, H., Nelle, O., and Kirleis, W.: Charcoal usage in medieval and modern times
- in the Harz Mountains Area, Central Germany: Wood selection and fast
- overexploitation of the woodlands, Quat. Int. 366(1), 51-69,
- 824 https://doi.org/10.1016/j.quaint.2015.01.053, 2015.

- 825 Li, H. M.: Study on the utilization strategies of major crops from Neolithic to
- historical periods in the western Loess Plateau, Ph.D. thesis, Lanzhou University,
- 827 Lanzhou, 2018.
- 828 Li, H., An, C. B., Dong, W., Wang, W., Hu, Z., Wang S., Zhao, X., and Yang, Y.:
- Woodland vegetation composition and prehistoric human fuel collection strategy
- at the Shannashuzha site, Gansu Province, northwest China, during the middle
- Holocene, Veg. Hist. Archaeobot., 26(1-2), 1-9, https://doi.org/10.1007/s00334-
- 832 016-0585-8, 2016.
- 833 Li, H., An, C-B., Dong, W., Wang, W., Hu, Z., Wang, S., Zhao, X., and Yang, Y.:
- Woodland vegetation composition and prehistoric human fuel collection strategy
- at the Shannashuzha site, Gansu Province, northwest China, during the middle
- Holocene, Veget. Hist. Archaeobot., 26, 213–221,
- 837 https://doi.org/10.1007/s00334-016-0585-8, 2017.
- Li, S. C. (Eds.): Prehistoric culture evolution in Northwest China. Cultural Relics Press, Beijing, ISBN 7-5010-2655-5, 2009.
- 840 Li, S. C., Shui, T., and Wang, H.: Report on prehistoric archaeological survey in Hexi
- Corridor, Acta Archaeol. Sin., 2, 229–274, 2010 (in Chinese with English
- abstract).
- 843 <u>Li, X., Zhou, X., Zhou, J., Dodson, J., Zhang H., and Shang X.: The earliest</u>
- archaeobiological evidence of the broadening agriculture in China recorded at
- Xishanping site in Gansu Province, Sci. China Ser. D-Earth Sci. 50, 1707—1714,
- 846 <u>https://doi.org/10.1007/s11430-007-0066-0, 2007.</u>
- Li, X., Shang, X., Dodson, J., and Zhou, X.: Holocene agriculture in the Guanzhong
- Basin in NW China indicated by pollen and charcoal evidence, Holocene, 19(8),
- 849 1213–1220, https://doi.org/10.1177/0959683609345083, 2009.
- Li, X., Sun, N., Dodson, J., and Zhou, X.: Human activity and its impact on the
- landscape at the Xishanping site in the western Loess Plateau during 4800–4300
- cal yr BP based on the fossil charcoal record, J. Archaeol. Sci., 39(10), 3141–
- 853 3147, https://doi.org/10.1016/j.jas.2012.04.052, 2012.
- Li, X., Sun, N., Dodson, J., Ji, M., Zhao, K., and Zhou, X.: The impact of early
- smelting on the environment of Huoshiliang in Hexi Corridor, NW China, as
- recorded by fossil charcoal and chemical elements, Paleogeogr. Paleoclimatol.
- Paleoecol., 305(1–4), 329–336, https://doi.org/10.1016/j.palaeo.2011.03.015,
- 858 2011.
- Li, Y.: Agriculture and palaeoeconomy in prehistoric Xinjiang, China (3000–200
- 860 BC), Veg. Hist. Archaeobot., 30, 287–303, https://doi.org/10.1007/s00334-020-
- 861 00774-2, 2021.

- Liu, C. J., Kong, Z. C., and Lang, S. D.: Plant remains found in the Dadiwan site and
- its implication on past environment, Cult. Relics Cent. China, 4, 26–30, 2004 (in
- Chinese with English abstract).
- 865 Liu, F., Li, H., Cui, Y., Yang, Y., Lee, H. F., Ding, D., Hou, Y., and Dong, G.:
- Chronology and Plant Utilization from the Earliest Walled Settlement in the
- Hexi Corridor, Northwestern China, Radiocarbon, 61(4), 971–989,
- 868 https://doi.org/10.1017/RDC.2019.57, 2019.
- Liu, F., Yang, Y., Chen, G., Zhang, S., and Zhang, H.: Changes in Wood Utilization
- Due to Iron Age Jade Mining in the Western Hexi Corridor: Wood Charcoal
- 871 Investigations, Front Earth Sci., 9, 636534,
- https://doi.org/10.3389/feart.2021.636534, 2021.
- Liu, F., Zhang, Y., Feng, Z., Hou, G., Zhou, Q., and Zhang, H.: The impacts of
- climate change on the Neolithic cultures of Gansu-Qinghai region during the late
- Holocene Megathermal, J. Geogr. Sci., 20, 417–430,
- 876 https://doi.org/10.1007/s11442-010-0417-1, 2010.
- Liu, X., Jones, P. J., Matuzeviciute, G. M., Hunt, H. V., Lister, D. L., An, T.,
- Przelomska, N., Kneale, C. J., Zhao, Z., and Jones, M. K.: From ecological
- opportunism to multi-cropping: Mapping food globalisation in prehistory, Quat.
- 880 Sci. Rev., 206, 21–28, https://doi.org/10.1016/j.quascirev.2018.12.017, 2019.
- Liu, X., Lightfoot, E., O'Connell, T. C., Wang, H., Li, S., Zhou, L., and Hu, Y.,
- Motuzaite-Matuzeviciute, G., Jones, M.K.: From necessity to choice: Dietary
- revolutions in west China in the second millennium BC, World Archaeol., 46,
- 884 661–680, https://doi.org/10.1080/00438243.2014.953706, 2014.
- Ma, M., Dong, G., Jia, X., Wang, H., Cui, Y., and Chen, F.: Dietary shift after 3600
- cal yr BP and its influencing factors in northwestern China: evidence from stable
- isotopes, Quat. Sci. Rev., 145, 57–70,
- https://doi.org/10.1016/j.quascirev.2016.05.041, 2016.
- 889 Ma, T., Rolett, B. V., Zheng, Z., and Zong, Y.: Holocene coastal evolution preceded
- the expansion of paddy field rice farming, Proc. Natl. Acad. Sci. USA, 117,
- 891 24138–24143, https://doi.org/10.1073/pnas.1919217117, 2020.
- Marguerie, D. and Hunot, J.Y.: Charcoal analysis and dendrology: data from
- archaeological sites in north-western France, J. Archaeol. Sci., 34(9), 1417–
- 894 1433, https://doi.org/10.1016/j.jas.2006.10.032, 2007.
- Mas, B., Allué, E., Alonso, E. S., and Vaquero, M.: From forest to settlement:
- Magdalenian hunter-gatherer interactions with the wood vegetation environment
- based on anthracology and intra-site spatial distribution, Archaeol. Anthropol.
- 898 Sci., 13, 12, https://doi.org/10.1007/s12520-020-01264-2, 2021.

- 899 Mei, J. and Shell, C.: The existence of Andronovo cultural influence in Xinjiang
- 900 during the second millennium BC, Antiquity, 73, 570–78,
- 901 https://doi.org/10.1017/S0003598X00065121, 1999.
- 902 Miller, A. and Gross, B.L.: From forest to field: perennial fruit crop domestication.
- 903 Am. J. Bot. 98, 1389–1414, https://doi.org/10.3732/ajb.1000522, 2011.
- 904 Miller, N. F., Spengler, R. N., and Frachetti, M. D.: Millet cultivation across Eurasia:
- Origins, spread, and the influence of seasonal climate, Holocene, 26(10), 1566–
- 906 1575, https://doi.org/10.1177/0959683616641742, 2016.
- 907 Miller, N. F.: Agropastoralism and archaeobiology: Connecting plants, animals and
- people in West and Central Asia, Environ. Archaeol. 18(3), 247–256,
- 909 https://doi.org/10.1179/1749631413Y.0000000003, 2013.
- 910 Monastersky, R.: Anthropocene: The human age, Nature, 519, 144–147,
- 911 https://doi.org/10.1038/519144a, 2015.
- 912 Neumann, K., Bostoen, K., Höhn, A., Kahlheber, S., Ngomanda, A., and Tchiengué,
- 913 B.: First farmers in the Central African rainforest: A view from southern
- 914 Cameroon, Quat. Int., 249, 53–62, https://doi.org/10.1016/j.quaint.2011.03.024,
- 915 2012.
- Nong, K., Zhang, G., Wang, L., Cheng, Y., and Jiang H.: Prehistoric wooden bows
- and arrows in the Turpan Basin, Northwest China: Wood selection and
- 918 utilization in a mosaic landscape. Archaeometry, 65(4), 881–896,
- 919 https://doi.org/10.1111/arcm.12846, 2023.
- 920 Qu, Y., Hu, Y., Rao, H., Abuduresule, I., Li, W., Hu, X., Jiang, H., Wang, C., and
- Yang, Y.: Diverse lifestyles and populations in the Xiaohe culture of the Lop
- 922 Nur region, Xinjiang, China, Archaeol. Anthropol. Sci., 10, 2005–2014,
- 923 https://doi.org/10.1007/s12520-017-0520-7, 2018.
- 924 Reimer, P., Austin, W., Bard, E., Bayliss, A., Blackwell, P., Bronk Ramsey, C.,
- Butzin, M., Cheng, H., Edwards, R. L., Friedrich, M., Grootes, P. M.,
- Guilderson, T. P., Hajdas, I., Heaton, T. G., Hogg, A. G., Hughen, K. A.,
- 927 Kromer, B., Manning, S. W., Muscheler, R., Palmer, J. G., Pearson, C., van der
- 928 Plicht, J., Reimer, R. W., Richards, D. A., Scott, E. M., Southon, J. R., Turney,
- 929 C. S. M., Wacker, L., Adolphi, F., Büntgen, U., Capano, M., Fahrni, S. M.,
- 930 Fogtmann-Schulz, A., Friedrich, R., Köhler, P., Kudsk, S., Miyake, F., Olsen, J.,
- Reinig, F., Sakamoto, M., Sookdeo, A., and Talamo, S.: The IntCal20 Northern
- 932 Hemisphere Radiocarbon Age Calibration Curve (0–55 cal kBP), Radiocarbon,
- 933 62(4), 725-757, https://doi.org/10.1017/RDC.2020.41, 2020.
- Ren, L., Yang, Y., Qiu, M., Brunson, K., Chen, G., and Dong, G.: Direct dating of the
- earliest domesticated cattle and caprines in northwestern China reveals the
- history of pastoralism in the Gansu-Qinghai region, J. Archaeol. Sci., 144,

937	105627, https://doi.org/10.1016/j.jas.2022.105627, 2022.
938 939 940 941	Ren, M., Tang, Z., Wu, X., Spengler, R., Jiang, H., Yang, Y., and Boivin, N.: The origins of cannabis smoking: Chemical residue evidence from the first millennium BCE in the Pamirs, Sci. Adv., 5(6), eaaw1391, https://doi.org/10.1126/sciadv.aaw1391, 2019.
942 943 944	Renn, J. (Eds.): The Evolution of Knowledge: Rethinking Science for the Anthropocene, Princeton University Press, Princeton, ISBN 9780691171982, 2020.
945 946	Ruan, Q. R.: Research on the Andronovo remains discovered in Xinjiang, Xibu Kaogu, 125–154, 2014 (in Chinese with English abstract).
947 948	Ruddiman, W. F. (Eds.): Earth's climate: past and future, second edition, WH Freeman & Co Ltd, New York, ISBN 10978-0-7167-8490-6, 2008.
949 950	Ruddiman, W. F.: The Anthropocene, Annu. Rev. Earth Planet Sci., 4, 45–68, https://doi.org/10.1146/annurev-earth-050212-123944, 2013.
951 952 953	Ruddiman, W. F.: The anthropogenic greenhouse era began thousands of years ago, Clim. Change, 61(3), 261–293, https://doi.org/10.1023/B:CLIM.0000004577.17928.fa, 2018.
954 955 956	Salysbury, K. J. and Jane, F. W.: Charcoal from maiden Castle and their significance in relation to the vegetation and climatic conditions in Prehistoric times, J. Ecology, 28, 310–325, 1940.
957 958 959 960	Schlütz, F. and Lehmkuhl, F.: Holocene climatic change and the nomadic Anthropocene in eastern Tibet: Palynological and geomorphological results from the Nianbaoyeze Mountains, Quat. Sci. Rev., 28(15), 1449–1471, https://doi.org/10.1016/j.quascirev.2009.01.009, 2009.
961 962 963	Shao, H. Q. (Eds.): The evolution of prehistorical cultural patterns in Xinjiang and the relationship with the its neighbours, Science Press, Beijing, ISBN 978-7-03-058710-7, 2018.
964 965	Shao, H. Q. and Zhang, W. S.: A review of Andronovo culture in Xinjiang, West Reg. Studies, 2, 113–121, 2019 (in Chinese).
966 967 968	Shen, H. and Li, X.: From extensive collection to intensive cultivation, the role of fruits and nuts in subsistence economy on Chinese Loess Plateau, Archaeol. Anthropol. Sci., https://doi.org/10.1007/s12520-020-01255-3, 2021.
969 970 971	Shen, H., Li, X., Spengler, R., Zhou, X., and Zhao, K.: Forest cover and composition on the Loess Plateau during the Middle to Late-Holocene: Integrating wood charcoal analyses, Holocene, 31(1), 38–49,

972	https://doi.org/10.1177/0959683620961486, 2021.
973 974 975	Shen, H., Wu, X., Tang, Z., Zhou, X., Sun, N., and Li, X.: Wood usage and fire veneration in the Pamir, Xinjiang, 2500 yr BP, Plos One, 10(8), e0134847, https://doi.org/10.1371/journal.pone.0134847, 2015.
976	Shen, H., Zhou, X., Betts, A., Jia, P, W., Zhao, K., and Li, X.: Fruit collection and
977	early evidence for horticulture in the Hexi Corridor, NW China, based on
978	charcoal evidence, Veget. Hist. Archaeobot., 28, 187-197,
979	https://doi.org/10.1007/s00334-018-0691-x, 2019.
980	Shen, H., Zhou, X., Zhao, K., Betts, A., Jia, P.W., and Li, X.: Wood types and human
981	impact between 4300 and 2400 yr BP in the Hexi Corridor, NW China, inferred
982	from charcoal records, Holocene, 28(4), 629–639,
983	https://doi.org/10.1177/0959683617735586, 2018.
984	Sherratt, A.: The Trans-Eurasian exchange: the prehistory of Chinese relations with
985	the West, in: Contact and Exchange in the Ancient World, edited by: Mair, V.,
986	Hawaii University Press, Honolulu, 32–53,
987	https://doi.org/10.1515/9780824841676-005, 2006.
988	Spengler, R. N. (Eds.): Fruit from the Sands: The Silk Road Origins of the Food We
989	Eat, University of California Press, Oakland, ISBN 9780520972780, 2019.
990	Spengler, R. N. and Wilcox, G.: Archaeobotanical results from Sarazm, Tajikistan, an
991	early Bronze Age village on the edge: agriculture and exchange, J. Environ.
992	Archaeol., 18(3), 211–221, https://doi.org/10.1179/1749631413Y.0000000008,
993	2013.
994	Spengler, R. N., Frachetti, M. D., Doumani, P. N., Rouse, L. M., Cerasetti, B.,
995	Bullion, E., and Mar'yashev, A.N.: Early agriculture and crop transmission
996	among Bronze Age mobile pastoralists of Central Eurasia, Proc. R Soc. B, 281,
997	20133382, http://dx.doi.org/10.1098/rspb.2013.3382, 2014.
998	Spengler, R. N., Stark, S., Zhou, X., Fuks, D., Tang, L., Mir-Makhamad, B., Bjørn,
999	R., Jiang, H., Olivieri, L.M., Begmatov, A., and Boivin, N.: A journey to the
1000	west: the ancient dispersal of rice out of East Asia, Rice, 14, 83,
1001	https://doi.org/10.1186/s12284-021-00518-4, 2021.
1002	Sun, N., Li, X., Dodson, J., Zhou, X., Zhao, K., and Yang, Q.: Plant diversity of the
1003	Tianshui Basin in the western Loess Plateau during the mid-Holocene – Charcoal
1004	records from archaeological sites, Quat. Int., 308-309, 27–35,
1005	https://doi.org/10.1016/j.quaint.2013.05.014, 2013.
1006	Tan, L., Li, Y., Wang, X., Cai, Y., Lin, F., Cheng, H., Ma, L., Sinha, A., and Edwards
1007	L.: Holocene monsoon change and abrupt events on the western Chinese Loess
1008	Plateau as revealed by accurately dated stalagmites, Geophys. Res. Lett., 46,

1009	e2020GL090273, https://doi.org/10.1029/2020GL090273, 2020.
1010 1011 1012	Théry Parisot, I., Chabal, L., and Chrzavzez, J.: Anthracology and taphonomy, from- wood gathering to charcoal analysis. A review of the taphonomic processes— modifying charcoal assemblages, in archaeological contexts, Palaeogeogr.
1013	Palaeoclimatol. Palaeoecol., 291(1), 142–153,
1014	https://doi.org/10.1016/j.palaeo.2009.09.016, 2010.
 1015	Tian, D., Festa, M., Cong, D., Zhao, Z., Jia, P. W., and Betts, A.: New evidence for
1016	supplementary crop production, foddering and fuel use by Bronze Age
1017	transhumant pastoralists in the Tianshan Mountains, Sci. Rep., 11, 13718,
1018	https://doi.org/10.1038/s41598-021-93090-2, 2021.
1019	Vigne, J. D., Briois, F., Zazzo, A., Willcox, G., Cucchi, T., Thiébault, S., Carrère, I.,
1020	Franel, Y., Touquet, R., Martin, C., Moreau, C., Comby, C., and Guilaine, J.:
1021	First wave of cultivators spread to Cyprus at least 10,600 y ago, Proc. Natl.
1022	Acad. Sci. USA, 109(22), 8445–8449, https://doi.org/10.1073/pnas.1201693109,
1023	2012.
1024	Wang, L., Wang, Y., Li, W., Spate, M., Reheman, K., Sun, Q., Wang, B., Xu, H.,
1025	Zhang, Z., Zhang, G., and Jiang, H.: Inner Asian agro-pastoralism as optimal
1026	adaptation strategy of Wupu inhabitants (3000-2400 cal BP) in Xinjiang, China,
1027	Holocene, 31(2), 203–216, https://doi.org/10.1177/0959683620941139, 2021.
1028	Wang, S., Li, H., Zhang, L., Chen, G., Wang, P., and Zhao, Z.: Tree exploration and
1029	palaeo-environment at Heishuiguo Xichengyi site, Zhangye city, Gansu
1030	province—revealed with excavated charcoal analysis, Quat. Sci., 34(1), 43–50,
1031	https://doi.org/10.3969/j.issn.1001-7410.2014.06 shu, 2014 (in Chinses with
1032	English abstract).
1033	Wang, T. T., Fuller, B. T., Wei, D., Chang, X. E., and Hu, Y. W.: Investigating
1034	dietary patterns with stable isotope ratios of collagen and starch grain analysis of
1035	dental calculus at the Iron Age cemetery site of Heigouliang, Xinjiang, China,
1036	Int. J. Osteoarchaeol., 26, 693–704, https://doi.org/10.1002/oa.2467, 2016.
1037	Wang, T., Wei, D., Chang, X., Yu, Z., Zhang, X., Wang, C., Hu, Y., and Fuller, B. T.:
1038	Tianshanbeilu and the Isotopic Millet Road: reviewing the late Neolithic/Bronze
1039	Age radiation of human millet consumption from north China to Europe, Nat.
1040	Sci. Rev., 6(5), 1024–1039, https://doi.org/10.1093/nsr/nwx015, 2017.
1041	Weisdorf, J. L.: From foraging to farming: explaining the Neolithic Revolution, J.
1042	Econ. Surv., 19(4), 561–586, https://doi.org/10.1111/j.0950-0804.2005.00259.x,
1043	2005.
1044	Xiao, D-S., Zhao, Y-F., and An, Z-Q.: Analysis on the characteristics and change
1045	trend of precipitation in Ili region, J. Arid Land Resour. Environ., 20(4), 47–50,
1046	2006 (in Chinese with English abstract).

1047 Xie, D. J. (Eds.): Prehistoric archaeology of Gansu-Qinghai region. Cultural Relics Publishing House, Beijing, ISBN 9787501014170, 2002. 1048 Xinjiang Institute of Cultural Relics and Archaeology: 2013 excavation report of 1049 Sayituer cemetery in Hami, J. Nat. Mus. China, 9, 24–38, 2014 (in Chinese with 1050 English abstract). 1051 Xinjiang Institute of Cultural Relics and Archaeology: The excavation report of 1052 Xiaohe cemetery, Res. China's Front. Archaeol., 3, 338–398, 2004 (in Chinese 1053 with English abstract). 1054 1055 Xinjiang integrated Expedition Team and Institute of Botany, CAS. (Eds.): Vegetation and its utilization in Xinjiang, Science Press, Beijing, ISBN 1056 1057 13031836, 1978. Yang, J., Zhang, D., Yang, X., Wang, W., Perry, L., Fuller, D. Q., Li, H., Wang, J., 1058 Ren, L., Xia, H., Shen, X., Wang, H., Yang, Y., Yao, J., Gao, Y., and Chen, F.: 1059 Sustainable intensification of millet-pig agriculture in Neolithic North China, 1060 Nat. Sustain., 5, 780–786, https://doi.org/10.1038/s41893-022-00905-9, 2022. 1061 Yang, O. and Gaskin, J.: Flora of China, Volume 13, Science Press & Missouri 1062 Botanical Garden Press, Beijing & St. Louis, 59-65, 2007. 1063 1064 Yang, R., Yang, Y., Li, W., Abuduresule, Y., Hu, X., Wang, C., and Jiang, H.: 1065 Investigation of cereal remains at the Xiaohe Cemetery in Xinjiang, China, J. 1066 Archaeol. Sci., 49, 42–47, https://doi.org/10.1016/j.jas.2014.04.020, 2014. 1067 Yang, Y., Dong, G., Zhang, S., Cui, Y., Li, H., Chen, G., Dodson, J., and Chen, F.: Copper content in anthropogenic sediments as a tracer for detecting smelting 1068 activities and its impact on environment during prehistoric period in Hexi 1069 1070 Corridor, Northwest China, Holocene, 27(2), 282–291, https://doi.org/10.1177/0959683616658531, 2017. 1071 Yatoo, M. A., Spate, M., Betts, A., Pokharia, A. K., and Shah, M. A.: New evidence 1072 from the Kashmir Valley indicates the adoption of East and West Asian crops in 1073 the western Himalayas by 4400 years ago, Quat. Sci. Adv., 2, 100011, 1074 https://doi.org/10.1016/j.qsa.2020.100011, 2020. 1075 1076 Zalasiewicz, J., Waters, C., and Head, M.: Anthropocene: its stratigraphic basis, 1077 Nature, 541, 289, https://doi.org/10.1038/541289b, 2017. Zeder, M. A.: Domestication and early agriculture in the Mediterranean Basin: 1078 Origins, diffusion, and impact, Proc. Natl. Acad. Sci. USA, 105, 11597–11604, 1079 https://doi.org/10.1073/pnas.0801317105, 2008. 1080 Zhang, G., Wang, S., Ferguson, D. K., Yang, Y., Liu, X., and Jiang, H.: Ancient plant 1081

- China: Implication from desiccated plant remains, Archaeol. Anthropol. Sci., 9, 1083 145–152, https://doi.org/10.1007/s12520-015-0246-3, 2017. 1084 Zhang, G., Wang, Y., Spate, M., Wang, S., and Jiang, H.: Investigation of the diverse 1085 plant uses at the South Aisikexiaer cemetery (~ 2700~2400 years BP) in the 1086 Hami basin of Xinjiang, Northwest China, Archaeol. Anthropol. Sci., 11(2), 1087 699–711, https://doi.org/10.1007/s12520-017-0563-9, 2019. 1088 Zhang, X. L., Zhang, J., Li, Z. P., Zhang, L. R., Chen, G. K., Wang, P., and Wang, H.: 1089 Analysis of subsistence economy of Xichengyi people in the Zhangye city, 1090 Gansu, Archaeology, 7, 110–120, 2015. 1091 Zhao, K., Li, X., Zhou, X., Dodson, J., and Ji, M.: Impact of agriculture on an oasis 1092 landscape during the late Holocene: Palynological evidence from the Xintala site 1093 1094 in Xinjiang, NW China, Quat. Int., 311, 81–86, 1095 https://doi.org/10.1016/j.quaint.2013.06.035, 2013. Zheng, D., Yang, Q. Y., and Wu, S. H. (Eds.): Physical geography of China. Science 1096 Press, Beijing, ISBN 978-7-03-045397-6, 2015. 1097 Zheng, Z., Ma, T., Roberts, P., Li, Z., Yue, Y., Peng, H., Huang, K., Han, Z., Wan, 1098 Q., Zhang, Y., Zhang, X., Zheng, Y., and Satio, Y.: Anthropogenic impacts on 1099 Late Holocene land-cover change and floristic biodiversity loss in tropical 1100 southeastern Asia, Proc. Natl. Acad. Sci. USA, 118(40), e2022210118, 1101 https://doi.org/10.1073/pnas.2022210118, 2021. 1102 1103 Zhou, X. and Li, X.: Variation in spruce (Picea sp.) distribution in the Chinese Loess Plateau and surrounding areas during the Holocene, Holocene, 22(6), 687-696, 1104 https://doi.org/10.1177/0959683611400195, 2012. 1105 1106 Zhou, X., Li, X., Dodson, J., and Zhao, K.: Rapid agricultural transformation in the 1107 prehistoric Hexi corridor, China, Quat. Int., 426, 33-41, https://doi.org/10.1016/j.quaint.2016.04.021, 2016. 1108 Zhou, X., Li, X., Dodson, J., Zhao, K., Atahan, P., Sun, N., and Yang, Q.: Land 1109 degradation during the Bronze Age in Hexi Corridor (Gansu, China), Quat. Int., 1110 254(1), 42-48, https://doi.org/10.1016/j.quaint.2011.06.046, 2012. 1111
- 1112 Zhou, X., Yu, J., Spengler, R. N., Shen, H., Zhao, K., Ge, J., Bao, Y., Liu, J., Yang,
- Q., Chen, G., Jia, P. W., and Li, X.: 5,200-year-old cereal grains from the eastern
- Altai Mountains redate the trans-Eurasian crop exchange, Nat. Plants, 6, 78–87,
- 1115 https://doi.org/10.1038/s41477-019-0581-y, 2020.
- 2116 Zhou, X.: Mid-Holocene environmental changes, agricultural activity and human
- adaption in Gansu-Qinghai region, Ph.D. thesis, Institute of Earth Environment,
- 1118 Chinese Academy of Sciences, Xi'an, 2009.

1119	Zohary, D., Hopf, M., and Weiss, E. (Eds.): Domestication of plants in the Old
1120	World: The origin and spread of cultivated plants in West Asia, Europe and Nile
1121	Valley, fourth edition, Oxford University Press, Oxford, ISBN 9780191810046,
1122	2012.
1123	Zong, Y., Chen, Z., Innes, J. B., Chen, C., Wang, Z., and Wang, H.: Fire and flood
1124	management of coastal swamp enabled first rice paddy cultivation in east China,
1125	Nature, 449(7161), 459–462, https://doi.org/10.1038/nature06135, 2007.