



A 10 m resolution land cover map of the Tibetan Plateau with detailed vegetation types

Xingyi Huang^{1,#}, Yuwei Yin^{1,#}, Luwei Feng¹, Xiaoye Tong², Xiaoxin Zhang², Jiangrong Li³, and Feng Tian^{1,4,*}

¹Hubei Key Laboratory of Quantitative Remote Sensing of Land and Atmosphere, School of Remote Sensing and Information Engineering, Wuhan University, Wuhan, China

²Department of Geosciences and Natural Resource Management, University of Copenhagen, Copenhagen, Denmark

³Institute of Tibet Plateau Ecology, Tibet Agricultural and Animal Husbandry University, Linzhi, China

⁴Perception and Effectiveness Assessment for Carbon-neutrality Efforts, Engineering Research Center of Ministry of Education, Wuhan, China

These authors contributed equally to this work.

* Corresponding author

Correspondence: Feng Tian (tian.feng@whu.edu.cn)

Abstract. The Tibetan Plateau (TP) hosts a variety of vegetation types ranging from broadleaved and needle-leaved forests at the lower altitudes and mesic areas to alpine grassland at the higher altitudes and xeric areas. Accurate and detailed mapping of the vegetation distribution on TP is essential for an improved understanding of climate change effects on terrestrial ecosystems. Yet, existing land cover datasets of TP are either provided at a low spatial resolution or have insufficient vegetation types to characterize certain unique TP ecosystems, such as the alpine scree. Here, we produced a 10 m resolution TP land cover map with 12 vegetation classes and 3 non-vegetation classes for the year 2022 (referred as TP_LC10-2022) by leveraging state-of-the-art remote sensing approaches including the Sentinel-1 and Sentinel-2 imagery, environmental and topographic datasets, and 4 machine learning models using Google Earth Engine platform. Our dataset TP_LC10-2022 achieved an overall classification accuracy of 86.5% with a Kappa coefficient of 0.854. By comparing with 4 existing global land cover products, TP_LC10-2022 showed significant improvements in terms of reflecting local-scale vertical variations in the southeast TP region. Moreover, we found that alpine scree occupied 13.99% of the TP region which was ignored in existing land cover datasets, and that shrublands occupied 4.63% of the TP region characterized by distinct forms of deciduous shrublands and evergreen shrublands largely determined by topography and missed in existing land cover datasets. Our dataset provides a solid foundation for further analyses which need accurate delineation of these unique vegetation types in TP. The TP_LC10-2022 and the sample dataset are freely available at <https://doi.org/10.5281/zenodo.8228112> (Huang et al., 2023b) and <https://doi.org/10.5281/zenodo.8227942> (Huang et al., 2023a) respectively. Additionally, the classification map can be viewed through <https://cold-classifier.users.earthengine.app/view/tplc10-2022>.



1 Introduction

The Earth's surface is physically covered by various types of land cover, including forests, grasslands, croplands, lakes, wet-
20 lands, etc. Accurate mapping and classification of land cover are fundamental components for Earth observations. By under-
standing the distribution and characteristics of different land cover types, land cover mapping supports the assessment of carbon
stocks, vegetation dynamics, and land-atmosphere interactions, contributing to the implementation of effective climate change
mitigation measures (Wang et al., 2022; Liu et al., 2022; Li et al., 2018).

The advent of remote sensing technology has enabled the generation of global-scale land cover products at various res-
25 olutions. For instance, products like MCD12Q1, produced using MODIS data (Friedl et al., 2010, 2002), and the ESA CCI
product, derived from sensors like MERIS (Agency, 2014), have significantly contributed to the understanding of global ecosys-
tem responses to climate change. However, their spatial resolutions are at hundreds of meters, unable to provide an accurate
representation of the land surface conditions (Tian et al., 2021), particularly in spatially heterogenous regions, such as the
mountainous southeast Tibetan Plateau (TP) (Yang et al., 2017; Grekousis et al., 2015). In response to this limitation, several
30 medium- to high-resolution land cover products have been created using satellite images from Landsat and Sentinel-2. Notable
examples include GlobeLand30 (Chen et al., 2021, 2015), FROM_GLC30 (Gong et al., 2013), GLC_FCS30 (Zhang et al.,
2021) based on Landsat, and FROM_GLC10 (Chen et al., 2019), Dynamic World (Brown et al., 2022), Esri Land Cover (Karra
et al., 2021), and ESA WorldCover (Zanaga et al., 2022) based on Sentinel-2. However, these products use different classifica-
tion systems, resulting in large divergence in certain regions (Shi et al., 2023; Hua et al., 2018), and are often inadequate to
35 reflect the diverse and unique land cover types for important ecosystems (Liu et al., 2023), such as those in the TP.

Renowned as the "Third Pole" of the world (Shukla and Sen, 2021), TP holds a dual significance as a sensitive area and
an indicator zone for global climate change (Hua et al., 2021; Li et al., 2022; Trew and Maclean, 2021; Pepin et al., 2022).
It hosts a variety of vegetation types, ranging from broadleaved and needle-leaved forests at the lower altitudes and mesic
areas to alpine grassland at the higher altitudes and xeric areas. However, many of the unique vegetation types in TP are not
40 well represented in existing land cover datasets. For example, the alpine scree ecosystem in the transitional zone from alpine
grasslands to bare rocks at very high altitudes and the shrubland ecosystem in the transitional zone from forests to grasslands
(Li et al., 2014). Furthermore, shrublands in TP can have either evergreen leaves or deciduous leaves depending on the local
environments they grow, yet are largely ignored in existing 10 m resolution land cover datasets (Venter et al., 2022). These
unique ecosystems in TP are of high significance to monitor, given that TP has experienced dramatic warming (Fu et al.,
45 2021), increased humidity (Yang et al., 2014), rapid glacier retreat (Zhao et al., 2022), permafrost thawing (Gao et al., 2021),
expansion of lakes (Zhang et al., 2020), and vegetation changes (Wang et al., 2020; Duan et al., 2021; Gao et al., 2014) in the
last decades. Thus, a detailed and accurate mapping of the diverse vegetation types in TP is required for understanding climate
change effects on the terrestrial ecosystem, yet is challenging to accomplish given that shrublands are often confused with
forests or alpine meadows and alpine grasslands are commonly misclassified as bare land in most products (Liu et al., 2021;
50 Cai et al., 2022; Yu et al., 2014). Moreover, the extremely rough terrain in TP results in large mountain shadows and variations



in slope aspects which complicates the accurate detection of vegetation types from satellite imagery (Pizarro et al., 2022; Wang et al., 2023b).

To address the aforementioned challenges, we developed a specific vegetation remote sensing fine classification system tailored for the TP, consisting of 12 vegetation classes and 3 non-vegetation classes. We then created a comprehensive training and validation dataset consisting of 10,242 samples through manual interpretation and field trips, based on which we performed land cover classification of the TP by integrating multiple data sources on the Google Earth Engine (GEE) platform, including satellite imagery of Sentinel-1 and Sentinel-2, topography, temperature, and precipitation. We investigated the performance of 4 different classification models provided in GEE and selected the highest-accuracy one to generate a 10 m resolution land cover product for the TP in 2022, referred as TP_LC10-2022.

60 2 Methodology

2.1 Study area

The TP spans from the northern foot of the West Kunlun Mountains and Qilian Mountains to the southern foot of the Himalayas and other mountain ranges, extending from the western edge of the Kunlun Mountains and Pamir Plateau to the eastern edge of the Hengduan Mountains (Fig. 1). It lies between latitudes $25^{\circ}59'30''\text{N}$ to $40^{\circ}1'0''\text{N}$ and longitudes $67^{\circ}40'37''\text{E}$ to $104^{\circ}40'57''\text{E}$, covering a total area of 3.083 million km^2 . Its average elevation is approximately 4,320 m (Zhang et al., 2022).

Due to the combined influence of climate, topography, and human activities over time, the vegetation cover types vary significantly at different altitudes in the TP. The northwestern and central regions are characterized by extensive bare lands, alpine screes, and persistent snow cover. In the southern and eastern areas, there is a distribution of evergreen forests and mixed forests consisting of needle-leaved and broadleaved trees. The transitional zone between these regions is characterized by shrublands, alpine grasslands, and alpine meadows. We investigated the vegetation cover in a field trip carried out along the national road No. 318 and 109 in July 2023 (Fig. 1), covering all the vegetation types in TP.

2.2 Data

2.2.1 Satellite imagery

We used both the optical imagery from Copernicus Sentinel-2 and radar imagery from Copernicus Sentinel-1 for the classification. Sentinel-2 comprises two high-resolution multispectral imaging satellites, each equipped with a multispectral imager. It consists of 13 bands, with spatial resolutions of 10 m for 4 bands, 20 m for 6 bands, and 60 m for 3 bands. The study utilized Level-2A products from the year 2022, which had undergone processing via the Sen2Cor tool at the Copernicus Scientific Data Hub (Doxani et al., 2018). Sentinel-1 comprises two polar-orbiting satellites positioned in the same orbital plane. For this research, the Ground Range Detected (GRD) data obtained in wide swath (IW) mode was chosen. The GRD data consists of dual polarization (VV, VH) interferometric wave modes, offering a 10 m resolution (Prats-Iraola et al., 2015). It enables the provision of radar images suitable for land and maritime services, regardless of weather conditions and time of day.




Figure 1. Overview of the study area colored by elevation. The black lines are the field trip routes along national roads. The photos show examples of the landscape views of typical vegetation types in the Tibetan Plateau.

2.2.2 Topography data

Shuttle Radar Topography Mission (SRTM) (Farr et al., 2000) was designed to generate high-quality digital elevation models (DEMs) globally using synthetic aperture radar technology. The data collected by SRTM was used to create a global elevation model with a horizontal accuracy of 16 m and vertical accuracy of 6 m, at a spatial resolution of 30 m (Yang et al., 2011).

2.2.3 Precipitation data

The Climate Hazards Group InfraRed Precipitation with Station data (CHIRPS) (Funk et al., 2015) is a comprehensive dataset documenting global precipitation from 1981 to the present. CHIRPS integrates satellite imagery with in-situ station data, providing a resolution of 0.05° to generate gridded rainfall time-series suitable for trend analysis and seasonal drought monitoring.



90 2.2.4 Temperature data

The ERA5-Land dataset (Muñoz-Sabater et al., 2021) offers a comprehensive reanalysis of land variables, presenting a consistent perspective on their evolution over multiple decades at a higher resolution than ERA5. As the land component of the ECMWF ERA5 climate reanalysis, ERA5-Land combines model data and global observations to create a coherent dataset utilizing the principles of physics. Nineteen extra bands were incorporated by GEE, with each corresponding to an accumulation
95 band, and the hourly values were calculated as the difference between 2 successive forecast steps (Muñoz-Sabater, 2019). For this study, hourly temperature data from 2022 were used.

2.3 Land cover classification

The advancement of cloud computing technology in remote sensing has revolutionized the rapid analysis and application of Earth system science on a large scale, even globally. GEE stands out among these technologies, offering online visualization,
100 computation, and analysis capabilities for extensive Earth science data (Gorelick et al., 2017; Kumar and Mutanga, 2018). Consequently, we opted to utilize GEE for data processing, and analysis. Importantly, the satellite data and auxiliary data relevant to this study can be readily accessed through GEE. Fig. 2 presents our comprehensive classification system, which comprises 4 main steps: 1) sampling strategy, 2) data preprocessing and feature construction, 3) classification model comparison, and 4) accuracy assessment and inter-comparison.

105 2.3.1 Classification system

The TP harbors the world's highest and one of the most distinctive alpine vegetation communities, which pose challenges to their inclusion in both global and Chinese land cover classification systems. To address this issue, we have developed an adapted classification system specifically tailored to the alpine vegetation types found in the TP. The advantages of our classification system are as follows:

110 (1) Comprehensive vegetation functional types: We have categorized the vegetation in the TP based on plant growth form (trees, shrubs, and herbs), leaf phenology (evergreen and deciduous), leaf type (broadleaved and needle-leaved), and ecosystem type. This classification system results in 12 vegetation types, including 5 types of tree cover including evergreen needle-leaved forest (ENF), deciduous needle-leaved forest (DNF), evergreen broadleaved forest (EBF), deciduous broadleaved forest (DBF), and mixed forest (MF); 2 types of shrub cover including evergreen shrubland (ES) and deciduous shrubland (DS); 2 types of
115 herb cover including alpine grassland (AG) and alpine meadow (AM); 3 special vegetation cover types including alpine scree (AS), wetland (WL), and cultivated vegetation (CV); and 3 non-vegetation land cover types, including bare land (BL), water body (WB), and permanent ice and snow (PIS).

(2) Discriminability in remote sensing imagery: The utilization of high-resolution remote sensing imagery, such as Google Earth imagery (with a spatial resolution of up to 0.3 m), enhances the distinguishability of land cover types within this clas-
120 sification system. This ensures the feasibility of visually interpreting large-scale samples from remote sensing imagery and obtaining reliable and up-to-date information.




Figure 2. Flowchart of the land cover classification carried out in this study.

2.3.2 Sampling strategy

Supervised classification models heavily depend on a substantial number of labeled samples for effective training and validation (Foody and Mathur, 2004). While extracting samples directly from existing land cover products can save manpower, it introduces several issues: (1) Extracted training samples may inherit errors from previous land cover products (Xi et al., 2022); (2) Utilizing low-resolution products to extract training samples for high-resolution land cover mapping can lead to information loss and boundary effects between adjacent land parcels (Zhang et al., 2021; Zhang and Roy, 2017); (3) Reconciling classification systems of different products is difficult, and global land cover products may not include specific land cover types for certain regions. Therefore, collecting samples through visual interpretation emerges as a more feasible approach (Schepaschenko et al., 2019).




Figure 3. Examples of auxiliary data used for explaining visual interpretation, including Google Earth imagery and the Landsat monthly mean Normalized Difference Vegetation Index (NDVI) time-series for 2013-2022. The x-axis represents month of the year and the y-axis represents NDVI value.



Google Earth integrates high-resolution imagery from sources like QuickBird and GeoEye, providing reliable remote sensing data sources for visual interpretation. Selecting samples in areas without Google Earth image coverage in 2022 poses a challenge. Normalized Difference Vegetation Index (NDVI) time-series have thus been used as auxiliary data for land cover sample selection (Yang and Huang, 2021; Feng et al., 2016). To ensure the selection of stable samples, this study examines the stability of land features by reviewing the Landsat NDVI time-series from 2013 to 2022. To eliminate the interference of clouds and snow in the NDVI time-series, the following operations were performed on Landsat images: 1) Filtering out pixels with cloud coverage greater than 50%; 2) When selecting forest and shrub samples, applying a Normalized Difference Snow Index (NDSI) mask to filter out pixels with NDSI greater than -0.4. To obtain a more continuous NDVI time-series, the harmonic analysis of time series (HANTS) model was used for data interpolation and smoothing to remove noise and reconstruct missing data (Zhou et al., 2015). Additionally, the monthly mean value of the NDVI time-series for 2013-2022 was calculated to determine the phenological characteristics of each sample point (Chu et al., 2021).

For instance, in Fig. 3, different color characteristics are observed for evergreen shrubs and deciduous shrubs in Google Earth imagery. Evergreen shrubs maintain their green color even during winter, while deciduous shrubs appear yellow-brown. However, during spring or summer, direct differentiation between the two from imagery is not possible. Therefore, phenological characteristics are extracted from their mean NDVI time-series. Evergreen shrubs exhibit relatively stable NDVI values, whereas deciduous shrubs show a decrease in NDVI due to seasonal leaf shedding.

Table 1. Number of training and validation samples for the 15 land cover types.

Land cover type	Number of training samples	Number of validation samples	Total
Bare land	883	234	1117
Alpine scree	583	129	712
Alpine grassland	607	149	756
Alpine meadow	949	226	1175
Evergreen needle-leaved forest	654	169	823
Deciduous needle-leaved forest	432	106	538
Evergreen broadleaved forest	550	130	680
Deciduous broadleaved forest	459	132	591
Mixed forest	215	78	293
Evergreen shrubland	566	153	719
Deciduous shrubland	663	157	820
Water body	504	114	618
Wetland	280	72	352
Cultivated vegetation	441	91	532
Permanent ice and snow	414	102	516
Total	8200	2042	10242




Figure 4. Spatial distribution of the 10, 242 samples for land cover classification in the Tibetan Plateau.

Google Earth imagery does not accurately determine the presence of herbaceous plant growth. Nevertheless, grasslands display a significant periodic increase in NDVI during the growing season, while bare land exhibit a relatively flat NDVI time-series. This characteristic is utilized for identifying bare land. Regarding alpine grasslands and alpine meadows, judgments are based on area size, vegetation composition, moisture condition, and terrain. Meadows typically have a smaller area compared to grasslands, better moisture conditions and are often accompanied by trees or shrubs in the vicinity. Grasslands have a flatter distribution area compared to meadows, as depicted in Fig. 3. Consequently, effective differentiation between alpine grasslands and alpine meadows is achieved.

Topography data (elevation, slope, aspect) (Farr et al., 2000), the 1:1 million Chinese vegetation map (Yang and Huang, 2021), and high-quality Google Maps photos were selected for auxiliary judgment. Ultimately, a total of 10,242 samples were collected, as illustrated in Fig. 4. Subsequently, the 10,242 samples were mixed, and the samples for each category were randomly divided into training and validation sets in an approximate 4:1 ratio, as presented in Table 1.

2.3.3 Data preprocessing and feature construction

Interannual remote sensing images have proven to accurately capture phenological changes in specific vegetation cover and have been successfully utilized in various large-scale land cover classification studies (Verde et al., 2020). Hence, in this study, the Sentinel-2 remote sensing images from the entire year of 2022 were selected for band feature extraction.



In this study, the initial step involved retaining the images with a cloud cover of less than 10%. Subsequently, the quality assessment information (QA band) was utilized to exclude pixels with inadequate quality through cloud masking. The selected input bands for Sentinel-2 consisted of B2-B8, B8A, B9, B11, and B12, while for Sentinel-1 images, VV and VH were chosen as input features. The median compositing method in GEE (Souza Jr et al., 2020; Phan et al., 2020) was applied to process all bands of Sentinel-1 and Sentinel-2.

To better discern the characteristics of various land features, we calculated several indices using Sentinel-2 imagery. These included the NDVI, NDSI, Normalized Difference Water Index (NDWI), and Optimized Soil-Adjusted Vegetation Index (OS-AVI). NDVI is highly sensitive to vegetation growth and is commonly used to distinguish between vegetated and non-vegetated areas (Rouse et al., 1974). NDSI effectively detects snow by utilizing the reflective properties of snow in the short infrared

Table 2. Features used for land cover classification.

Data source	Feature	Description
Sentinel-1	VV	Single Polarization Radar Reflectivity
	VH	Dual-Band Cross-Polarization Radar Reflectivity
Sentinel-2	B2	Blue Band Reflectance (<i>Blue</i>)
	B3	Green Band Reflectance (<i>Green</i>)
	B4	Red Band Reflectance (<i>Red</i>)
	B5	Vegetation Red-Edge 1 Band Reflectance
	B6	Vegetation Red-Edge 2 Band Reflectance
	B7	Vegetation Red-Edge 3 Band Reflectance
	B8	Near-Infrared Band Reflectance (<i>NIR</i>)
	B8A	Narrow Near-Infrared Band Reflectance
	B9	Water Vapor Band Reflectance
	B11	Shortwave Infrared 1 Band Reflectance (<i>SWIR1</i>)
	B12	Shortwave Infrared 2 Band Reflectance (<i>SWIR2</i>)
	NDVI	$NDVI = (NIR - Red) / (NIR + Red)$
	NDSI	$NDSI = (Green - SWIR1) / (Green + SWIR1)$
	NDWI	$NDWI = (Green - NIR) / (Green + NIR)$
OSAVI	$OSAVI = (NIR - Red) / (NIR + Red + 0.16)$	
SRTM	Elevation	
	Slope	
	Aspect	
CHIRPS	Annual precipitation	
ERA5-Land	Annual mean temperature	Temperature of air at 2m above the surface of land or in-land waters.



band, making it advantageous for studying ice and snow coverage in high mountain regions (Dozier, 1989). NDWI effectively distinguishes between water and non-water features (Xu, 2006). OSAVI improves the sensitivity and stability of vegetation indices by considering the influence of soil reflectance, providing a more accurate reflection of vegetation coverage and growth conditions, particularly in cases of bare soil or sparse vegetation (Rondeaux et al., 1996).

175 Topography significantly influence the vertical distribution of vegetation in high mountain areas (Zou et al., 2023). Therefore, in this study, we included elevation, slope, and aspect as input features for classification. Additionally, we incorporated annual precipitation and mean annual temperature as classification feature indicators (Wang et al., 2023a; Shen et al., 2015). For bands with a spatial resolution different from 10 m, we employed bicubic interpolation to resample them to 10 m resolution for mapping (Liu et al., 2020). All the features and their detailed descriptions are presented in Table 2.

180 **2.3.4 Classification models comparison**

Machine learning is a typically employed technique in remote sensing image classification. To identify the most appropriate classification model, we compared 4 widely-used machine learning models in GEE, including Random Forest (RF) (Breiman, 2001), Gradient Tree Boosting (GTB) (Friedman, 2001), Support Vector Machine (SVM) (Hearst et al., 1998), and Minimum Distance (MD) (Wacker and Landgrebe, 1972). We fine-tuned the parameters of all the classification models to achieve optimal
185 results (Table A1). The classification model with the highest overall performance was chosen to generate the land cover map and calculate the area proportion of each land cover type.

2.3.5 Accuracy assessment and inter-comparison

The accuracy of remote sensing image classification is commonly assessed using a confusion matrix, which provides 4 quantitative indicators: Producer's Accuracy (P.A.) for measuring omission errors, User's Accuracy (U.A.) for measuring commission
190 errors, Overall Accuracy (O.A.), and Kappa coefficient.

To compare with existing 4 global land cover datasets, namely ESA WorldCover2021, FROM_GLC10-2017, FROM_GLC30-2015, and GLC_FCS30-2020, we merged pixels belonging to the same class (Table A2) and employed randomly sampled validation samples. Additionally, we selected three $0.1^\circ \times 0.1^\circ$ grids within the TP to compare the visual classification results of TP_LC10-2022 with the existing 4 land cover products.

195 **3 Results and discussion**

3.1 Comparison of classification models

Table 3 presents the evaluation results of different classification models applied in the study area using GEE. The results demonstrate that the RF model achieved the highest accuracy, with an Overall Accuracy (O.A.) of 86.5% and a Kappa coefficient of 0.854. The Gradient Tree Boosting (GTB) model closely followed with an O.A. of 85.6% and a Kappa coefficient of



Table 3. Comparison of classification results for Random Forest (RF), Gradient Tree Boosting (GTB), Minimum Distance (MD), and Support Vector Machine (SVM) at their best performance. Bold font denotes highest U.A. and P.A. within each land cover type, as well as the highest O.A. and Kappa among the 4 models.

		BL	AS	AG	AM	ENF	DNF	EBF	DBF	MF	ES	DS	WB	WL	CV	PIS	O.A.	Kappa
RF	P.A.	0.974	0.884	0.953	0.894	0.805	0.830	0.915	0.750	0.462	0.856	0.847	1.000	0.611	0.923	0.941	0.865	0.854
	U.A.	0.942	0.851	0.953	0.831	0.764	0.871	0.815	0.786	0.800	0.851	0.821	0.983	0.898	0.884	0.941		
GTB	P.A.	0.979	0.868	0.919	0.894	0.769	0.774	0.892	0.788	0.513	0.843	0.834	0.991	0.639	0.923	0.902	0.856	0.844
	U.A.	0.942	0.855	0.951	0.835	0.756	0.872	0.835	0.759	0.741	0.838	0.775	0.983	0.885	0.894	0.902		
SVM	P.A.	0.688	0.791	0.584	0.681	0.396	0.717	0.646	0.636	0.269	0.621	0.643	0.833	0.597	0.780	0.784	0.647	0.618
	U.A.	0.703	0.729	0.561	0.661	0.429	0.623	0.587	0.592	0.447	0.674	0.574	0.969	0.642	0.664	0.930		
MD	P.A.	0.885	0.814	0.711	0.863	0.876	0.660	0.777	0.780	0.397	0.745	0.771	1.000	0.625	0.758	0.971	0.797	0.781
	U.A.	0.885	0.766	0.914	0.739	0.643	0.946	0.863	0.665	0.674	0.891	0.742	0.950	0.750	0.852	0.846		

BL: bare land; AS: alpine scree; AG: alpine grassland; AM: alpine meadow; ENF: evergreen needle-leaved forest; DNF: deciduous needle-leaved forest;
 EBF: evergreen broadleaved forest; DBF: deciduous broadleaved forest; MF: mixed forest; ES: evergreen shrubland; DS: deciduous shrubland;
 WB: water body; WL: wetland; CV: cultivated vegetation; PIS: permanent ice and snow

200 0.844. The Minimum Distance (MD) model yielded an accuracy of O.A. 79.7% and Kappa 0.781, while the Support Vector Machine (SVM) exhibited significantly lower classification results, with an O.A. of 64.7% and Kappa of 0.618.

The high accuracy achieved by RF and GTB models can be attributed to their ensemble learning algorithms based on decision trees. These algorithms combine multiple decision trees to enhance model performance and generalization capabilities (Salditt et al., 2022). In contrast to the findings of Abdi (2020), where RF and SVM exhibited similar O.A., our SVM showed a decline
 205 of 21.8% compared to RF (Tu et al., 2020). This discrepancy may be attributed to RF's ability to mitigate the correlation between samples and features through random sampling and feature selection, resulting in improved classification performance and robustness. Moreover, RF can effectively handle high-dimensional data and capture nonlinear relationships by integrating multiple decision trees (Tu et al., 2020; Gislason et al., 2006).

The three classification models, excluding SVM, effectively distinguished water bodies from other land cover types, achiev-
 210 ing a P.A. exceeding 0.99. However, all classification models performed fairly in differentiating mixed forests. For instance, SVM achieved a low P.A. of only 0.269 for mixed forest classification. Despite the integration of various machine learning models within GEE, including algorithms like RF, a distinct absence of direct support for deep learning persists. This is notable even in light of the well-established and showcased capabilities of deep learning in the fine-grained classification of land cover (Wang et al., 2023c). This limitation, to a certain extent, poses a hindrance to the extensive application of large-scale land cover
 215 mapping.

The utilization of multi-source remote sensing data can offer a more comprehensive understanding of land cover (Xu et al., 2022; Chen et al., 2017). Given to the importance of features, all features contributed to the mapping and elevation contributed slightly more to the accuracy of the classification (Fig. A1). This is attributed to the impact of the TP's rugged terrain on the hydrothermal conditions in distinct regions, leading to notable variations in vegetation phenology (Hwang et al., 2011).




Figure 5. The overall Comparison for TP_LC10-2022 and other land cover maps. (a) TP_LC10-2022 and the proportion of 15 land cover types. (b) An overview of 4 land cover products in the Tibetan Plateau, including ESA WorldCover2021, FROM_GLC10-2017, FROM_GLC30-2015 and GLC_FCS30-2020. Legend fusion rules for WorldCover2021 and FROM_GLC10-2017 are provided in Table A2, and for FROM_GLC30-2015 and GLC_FCS30-2020, refer to Table A3.




Figure 6. The confusion proportions for each of the land cover types in TP_LC10-2022. BL: bare land; AS: alpine scree; AG: alpine grassland; AM: alpine meadow; ENF: evergreen needle-leaved forest; DNF: deciduous needle-leaved forest; EBF: evergreen broadleaved forest; DBF: deciduous broadleaved forest; MF: mixed forest; ES: evergreen shrubland; DS: deciduous shrubland; WB: water body; WL: wetland; CV: cultivated vegetation; PIS: permanent ice and snow.

220 3.2 Land cover classification map

Fig. 5a provides an overview of the TP_LC10-2022 product and 4 global land cover products, along with the proportion of each land cover type in TP_LC10-2022. Alpine meadow and alpine grassland account for the proportions at 23.76% and 16.48%, respectively. Alpine scree surprisingly ranks fourth, with a proportion of 13.99%, after alpine meadow, bare land, and alpine grassland. Evergreen needle-leaved forest has the largest area among the forest types, and deciduous shrubland has a larger area than evergreen shrubland, reaching 3.57%, surpassing other forest types except for evergreen needle-leaved forest.

According to Fig. 5b, ESA WorldCover2021, FROM_GLC10-2017, and FROM_GLC30-2015 products overestimate the area of bare land in the TP, similar to the issues observed in FROM_GLC-agg and ESA CCI land cover products (Liu et al., 2021; Yu et al., 2014). This may be due to the misclassification of alpine grassland as bare land because these products captured less spectral information during the growing season of alpine grasslands. GLC_FCS30-2020 classified more grasslands but failed to distinguish between grasslands and meadows.

Table 4 illustrates the confusion matrix of TP_LC10-2022, with an overall accuracy of 86.5% and a Kappa coefficient of 0.854. Water body achieved a P.A. of 100%, while mixed forest only reached 46.2%. Fig. 6 shows that most mixed forests are challenging to differentiate from other forest types, with over 25% of mixed forests misclassified as evergreen needle-leaved forests and 11.5% misclassified as evergreen broadleaved forests or deciduous broadleaved forests. The classification accuracy of wetlands is also unsatisfactory, with a P.A. of only 61.1%. Over 16% of wetlands were classified as bare land, and over 15%



Table 4. Confusion matrix of TP_LC10-2022 product extracted using Random Forest (RF) classification model. Bold font denotes correctly classified sample points.

	BA	AS	AG	AM	ENF	DNF	EBF	DBF	MF	ES	DS	WB	WL	CV	PIS	Total	P.A.	
BA	228	2	2	0	0	0	0	0	0	0	0	1	0	1	0	234	0.974	
AS	1	114	0	5	0	1	0	0	0	0	4	0	0	0	4	129	0.884	
AG	1	1	141	4	0	0	0	0	0	0	0	0	1	1	0	149	0.953	
AM	0	3	4	202	0	0	0	2	0	2	6	0	2	4	1	226	0.894	
ENF	0	0	0	2	136	5	9	1	3	12	1	0	0	0	0	169	0.805	
DNF	0	0	0	2	1	88	0	7	0	3	5	0	0	0	0	106	0.830	
EBF	0	0	0	0	11	0	119	0	0	0	0	0	0	0	0	130	0.915	
DBF	0	0	0	3	5	1	8	99	6	2	3	0	0	5	0	132	0.750	
MF	0	0	0	0	20	1	9	9	36	2	1	0	0	0	0	78	0.462	
ES	0	2	0	2	5	1	1	2	0	131	8	0	0	0	1	153	0.856	
DS	0	5	0	8	0	4	0	5	0	2	133	0	0	0	0	157	0.847	
WB	0	0	0	0	0	0	0	0	0	0	0	114	0	0	0	114	1.000	
WL	12	0	1	11	0	0	0	1	0	0	1	1	44	1	0	72	0.611	
CV	0	1	0	4	0	0	0	0	0	0	0	0	2	84	0	91	0.923	
PIS	0	6	0	0	0	0	0	0	0	0	0	0	0	0	96	102	0.941	
Total	242	134	148	243	178	101	146	126	45	154	162	116	49	96	102	2042		
U.A.	0.942	0.851	0.953	0.831	0.764	0.871	0.815	0.786	0.800	0.851	0.821	0.983	0.898	0.884	0.941			
O.A.																	0.865	
Kappa																	0.854	

BL: bare land; AS: alpine scree; AG: alpine grassland; AM: alpine meadow; ENF: evergreen needle-leaved forest; DNF: deciduous needle-leaved forest; EBF: evergreen broadleaved forest; DBF: deciduous broadleaved forest; MF: mixed forest; ES: evergreen shrubland; DS: deciduous shrubland; WB: water body; WL: wetland; CV: cultivated vegetation; PIS: permanent ice and snow

were incorrectly classified as alpine meadows. The U.A. for the water body reached 98.3%, while evergreen needle-leaved forests had the lowest U.A. at 76.4%.

240 Although we employed the Sentinel-2 median composition method in this study, we acknowledge the potential enhancement that time-series analysis could bring to our research. In comparison to median composition, time-series analysis has the potential to more comprehensively capture phenological information of vegetation, thereby yielding more accurate land cover classification results (Xie et al., 2019; Nguyen et al., 2020). However, time-series methods also have their limitations, such as the requirement for a greater number of valid observations (Hemmerling et al., 2021). This constraint proves inapplicable to regions in southeastern TP due to the inefficacy of conventional cloud removal techniques in these densely cloud-covered areas, precluding the attainment of desired outcomes (Chu et al., 2021).

245 3.3 Inter-comparison with other products

We validated the aggregation of samples into eight categories and assessed the performance of TP_LC10-2022 and 4 additional land cover products in the Tibetan Plateau region, as depicted in Table 5.




Figure 7. Comparison of TP_LC10-2022 with 4 other global land cover products in details. (a), (b) and (c) represent the $0.1^\circ \times 0.1^\circ$ grids for detailed comparisons. Legend fusion rules for WorldCover2021 and FROM_GLC10-2017 are provided in Table A2, and for FROM_GLC30-2015 and GLC_FCS30-2020, refer to Table A3.



Table 5. Comparison of mapping accuracy based on validation samples merged into 8 land cover types. Legend for the fusion rules of the 5 land cover products is provided in Table A2. Bold font denotes highest U.A. and P.A. within each land cover type, as well as the highest O.A. and Kappa among the 5 products.

		BL	GL	FST	SHR	WB	WL	CV	PIS	O.A	Kappa
GLC_FCS30-2020	P.A.	0.726	0.757	0.956	0.026	0.965	0.167	0.516	1.000	0.691	0.604
	U.A.	0.769	0.538	0.709	0.216	0.965	1.000	0.797	0.903		
FROM_GLC30-2015	P.A.	0.902	0.518	0.936	0.063	0.981	0.048	0.440	0.931	0.663	0.578
	U.A.	0.531	0.524	0.758	0.593	0.737	0.375	0.688	0.979		
FROM_GLC10-2017	P.A.	0.961	0.493	0.953	0.058	0.982	0.085	0.824	0.971	0.683	0.604
	U.A.	0.520	0.474	0.820	0.563	0.974	0.188	0.773	1.000		
WorldCover2021	P.A.	0.936	0.604	0.935	0.000	0.991	0.521	0.802	1.000	0.706	0.631
	U.A.	0.569	0.509	0.787	0.000	0.863	0.974	1.000	0.990		
TP_LC10-2022	P.A.	0.970	0.928	0.935	0.881	1.000	0.556	0.923	0.961	0.919	0.900
	U.A.	0.912	0.872	0.962	0.889	0.974	0.889	0.857	0.980		

BL: bare land; GL: grassland; FST: forest; SHR: shrubland; WB: water body; WL: wetland; CV: cultivated vegetation;
 PIS: Permanent ice and snow

For shrubland, the classification performance of the 4 global land cover products is remarkably low. Notably, ESA World-Cover2021 achieves a P.A. and U.A. of 0 for shrubland classification. Among these land cover products, FROM_GLC30-2015 exhibits the highest U.A. for shrubland classification, albeit at a mere 59.3%. This suggests substantial shortcomings in the precise classification of shrubland in the TP region by the current land cover products.

A simultaneous visual comparison was conducted among the 5 products. In Fig. 7a, TP_LC10-2022 and FROM_GLC30-2015 exhibited superior performance, revealing more intricate forest details compared to other products. Notably, other products largely disregarded vast areas of high-elevation alpine shrublands above the timberline, while TP_LC10-2022 delineated them (shown in brown) and exhibited distinct vertical zonation. In Fig. 7b, the other 4 products tended to misclassify shrublands as forests, particularly FROM_GLC30-2015 and GLC_FCS30-2020, whereas TP_LC10-2022 accurately differentiated between forests and shrubs. In Fig. 7c, both FROM_GLC10-2017 and FROM_GLC30-2015 depicted scattered shrublands but lacked continuity. These 4 products overestimated grasslands and underestimated the extent of shrubland areas. This discrepancy may stem from the similar phenological characteristics between deciduous shrublands and meadows, posing difficulties in their sole distinction based on spectral features (Li et al., 2021). However, TP_LC10-2022 integrates topographic and climatic factors as classification features, facilitating precise differentiation between shrublands and grasslands.

Alpine screes are extensively distributed across the TP, yet they are frequently disregarded from other products. Our product presents the initial description of alpine scree vegetation locations, which will contribute to environmental monitoring and biodiversity research in the periglacial zone of the TP (Li et al., 2014). Shrubbylands play a vital role as carbon sinks in ecosystems and hold substantial implications for global carbon cycling (Ma et al., 2021). TP_LC10-2022 precisely portrays the distribution



of shrublands, which holds significant importance in forecasting the impact of future changes in the carbon cycle on global-scale ecosystems.

4 Conclusions

We present a detailed land cover map including 12 vegetation types and 3 non-vegetation types at 10 m spatial resolution of the year 2022 for the Tibet Plateau (TP_LC10-2022) by integrating multi-source data including Sentinel-1, Sentinel-2, SRTM, CHIRPS, and ERA5-Land and comparing 4 classification models via GEE. The TP_LC10-2022 achieved an overall accuracy of 86.5% and a Kappa coefficient of 0.854% using the RF model, which outperforms other classification models, including GTB, MD and SVM. The comparisons between TP_LC10-2022 and 4 widely used land cover products (GLC_FCS30-2020, FROM_GLC30-2015, FROM_GLC10-2017, and WorldCover2021) demonstrated that TP_LC10-2022 has higher overall accuracy and reflects the local-scale variations of vegetation types along latitudes. In particular, TP_LC10-2022 incorporated unique land cover types like alpine scree, alpine grassland, and alpine meadow, which accounts for 54.23% of the total coverage. Moreover, it accurately depicted the distribution of shrubland that occupied 4.63% of the TP and was underestimated in the other products. The proposed vegetation classification system for the TP can serve as a foundation for land cover mapping in this region and a reference approach for mapping shrubland globally. The developed TP_LC10-2022 product can facilitate monitoring vegetation changes and studying the response to climate change in the TP.

Data availability. The TP_LC-2022 product generated in this paper is available at <https://doi.org/10.5281/zenodo.8228112> (Huang et al., 2023b). Across the entire Tibetan Plateau, the TP_LC-2022 product is grouped by $54\ 3^{\circ} \times 3^{\circ}$ tiles in the GeoTIFF format (EPSG: 4326), which are named “TP_LC10-2022_E**N**.tif”, where “E**N**” explains the longitude and latitude information of the upper left corner of each regional land cover map. The multi-source data used in this study, including Sentinel-2, can be directly accessed from Google Earth Engine.

The corresponding sample dataset, produced by manual interpretation and field trips, is available at <https://doi.org/10.5281/zenodo.8227942> (Huang et al., 2023a). The classification map can be viewed through <https://cold-classifier.users.earthengine.app/view/tplc10-2022>.



Appendix A

Table A1. Optimal parameters for Random Forest (RF), Gradient Tree Boosting (GTB), Minimum Distance (MD), and Support Vector Machine (SVM) in this study.

Model	Optical parameters
RF	numberOfTrees: 100
GTB	numberOfTrees: 75
MD	metric: 'mahalanobis' kNearest: 1
SVM	decisionProcedure: 'Voting' kernelType: 'RBF' gamma: 0.000005 cost: 2000



Table A2. Cross-walking table between different land cover products.

Target type	TP_LC2022	WorldCover2021	FROM_GLC10-2017	FROM_GLC30-2015	GLC_FCS30-2020
Bare land	Bare land	Bare / sparse vegetation Built-up	Bare land Impervious area	Bareland Impervious surface	Bare areas Impervious surfaces Consolidated bare areas Unconsolidated bare areas
Grassland	Alpine grassland Alpine meadow	Grassland	Grassland	Natural grassland Grassland, leaf-off	Grassland Sparse vegetation Sparse herbaceous Herbaceous cover
Forest	Evergreen broadleaved forest Deciduous broadleaved forest Evergreen needle-leaved forest Deciduous needle-leaved forest Mixed forest	Tree cover	Forest	Broadleaf, leaf-on Broadleaf, leaf-off Needleleaf, leaf-on Needleleaf, leaf-off Mixed leaf, leaf-on	Open evergreen broadleaved forest Closed evergreen broadleaved forest Open deciduous broadleaved forest Closed deciduous broadleaved forest Open evergreen needle-leaved forest Closed evergreen needle-leaved forest Open deciduous needle-leaved forest Closed deciduous needle-leaved forest
Shrubland	Evergreen Shrubland Deciduous Shrubland	Shrubland	Shrubland	Shrubland, leaf-on Shrubland, leaf-off	Shrubland Evergreen Shrubland Deciduous Shrubland
Water body	Water body	Permanent water bodies	Water body	Water	Water body
Wetland	Wetland	Herbaceous wetland	Wetland	Marshland Mudflat Marshland, leaf-off	Wetlands
Cultivated vegetation	Cultivated vegetation	Cropland	Cropland	Rice paddy Greenhouse Orchard Bare farmland	Rainfed cropland Tree or shrub cover (Orchard) Irrigated cropland
Permanent ice and snow	Permanent ice and snow	Snow and ice	Snow and ice	Snow Ice	Permanent ice and snow
Excluded	Alpine scree	Moss and lichen	Tundra	Herbaceous tundra	Lichens and mosses

• This table includes only land cover types present within the study area.



Table A3. Cross-walking table between FROM_GLC30-2015 and GLC_FCS30-2020.

Target type	FROM_GLC30-2015	GLC_FCS30-2020
Bare land	Bareland	Bare areas
	Impervious surface	Impervious surfaces
		Consolidated bare areas
		Unconsolidated bare areas
Grassland	Natural grassland	Grassland
	Grassland, leaf-off	Sparse vegetation
		Sparse herbaceous
		Herbaceous cover
Evergreen broadleaved forest	Broadleaf, leaf-on	Open evergreen broadleaved forest
		Closed evergreen broadleaved forest
Deciduous broadleaved forest	Broadleaf, leaf-off	Open deciduous broadleaved forest
		Closed deciduous broadleaved forest
Evergreen needle-leaved forest	Needleleaf, leaf-on	Open evergreen needle-leaved forest
		Closed evergreen needle-leaved forest
Deciduous needle-leaved forest	Needleleaf, leaf-off	Open deciduous needle-leaved forest
		Closed deciduous needle-leaved forest
Mixed forest	Mixed leaf, leaf-on	
Shrubland	Shrubland, leaf-on	Evergreen Shrubland
	Shrubland, leaf-off	Deciduous Shrubland
Water body	Water	Water body
Wetland	Marshland	Wetlands
	Mudflat	
	Marshland, leaf-off	
Cultivated vegetation	Rice paddy	Rainfed cropland
	Greenhouse	Tree or shrub cover (Orchard)
	Orchard	Irrigated cropland
	Bare farmland	
Permanent ice and snow	Snow	Permanent ice and snow
	Ice	
Tundra / Lichens and mosses	Herbaceous tundra	Lichens and mosses

- This table includes only land cover types present within the study area.
- The 'shrubland' class in the GLC_FCS30-2020 product has been omitted from the table due to its small area.




Figure A1. Statistical chart of the importance of different features for the Random Forest classification model. AP: annual precipitation; AMT: annual mean temperature.

Author contributions. FT conceived the study. XH and YY collected the samples and carried out the analysis while LF provided the technical support. XH and YY wrote the original draft. FT and JL carried out the field trip. All authors helped revise the draft.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This work is funded by the National Key Research and Development Program of China (2020YFA0608704), the National Natural Science Foundation of China (Grant No. 42001299), and the Seed Fund Program for Sino-Foreign Joint Scientific Research Platform of Wuhan University (No. WHUZZJJ202205).



References

- Abdi, A. M.: Land cover and land use classification performance of machine learning algorithms in a boreal landscape using Sentinel-2 data, *GISci. Remote Sens.*, 57, 1–20, <https://doi.org/10.1080/15481603.2019.1650447>, 2020.
- Agency, E. S.: Land Cover CCI Product user guide version 2, https://www.esa-landcover-cci.org/?q=webfm_send/84, last access: 9 Aug, 2023, 2014.
- Breiman, L.: Random forests, *Mach. Learn.*, 45, 5–32, https://doi.org/10.1007/978-3-030-56485-8_3, 2001.
- Brown, C. F., Brumby, S. P., Guzder-Williams, B., Birch, T., Hyde, S. B., Mazzariello, J., Czerwinski, W., Pasquarella, V. J., Haertel, R., Ilyushchenko, S., et al.: Dynamic World, Near real-time global 10 m land use land cover mapping, *Sci. Data*, 9, 251, <https://doi.org/10.1038/s41597-022-01307-4>, 2022.
- Cai, L., Wang, S., Jia, L., Wang, Y., Wang, H., Fan, D., and Zhao, L.: Consistency Assessments of the land cover products on the Tibetan Plateau, *IEEE J. Sel. Top. Appl. Earth Observ. Remote Sens.*, 15, 5652–5661, <https://doi.org/10.1109/JSTARS.2022.3188650>, 2022.
- Chen, B., Huang, B., and Xu, B.: Multi-source remotely sensed data fusion for improving land cover classification, *ISPRS-J. Photogramm. Remote Sens.*, 124, 27–39, <https://doi.org/10.1016/j.isprsjprs.2016.12.008>, 2017.
- Chen, B., Xu, B., Zhu, Z., Yuan, C., Suen, H. P., Guo, J., Xu, N., Li, W., Zhao, Y., Yang, J., et al.: Stable classification with limited sample: Transferring a 30-m resolution sample set collected in 2015 to mapping 10-m resolution global land cover in 2017, *Sci. Bull.*, 64, 3, <https://doi.org/10.1016/j.scib.2019.03.002>, 2019.
- Chen, J., Chen, J., Liao, A., Cao, X., Chen, L., Chen, X., He, C., Han, G., Peng, S., Lu, M., et al.: Global land cover mapping at 30 m resolution: A POK-based operational approach, *ISPRS-J. Photogramm. Remote Sens.*, 103, 7–27, <https://doi.org/10.1016/j.isprsjprs.2014.09.002>, 2015.
- Chen, J., Chen, L., Chen, F., Ban, Y., Li, S., Han, G., Tong, X., Liu, C., Stamenova, V., and Stamenov, S.: Collaborative validation of GlobeLand30: Methodology and practices, *Geo-Spat. Inf. Sci.*, 24, 134–144, <https://doi.org/10.1080/10095020.2021.1894906>, 2021.
- Chu, D., Shen, H., Guan, X., Chen, J. M., Li, X., Li, J., and Zhang, L.: Long time-series NDVI reconstruction in cloud-prone regions via spatio-temporal tensor completion, *Remote Sens. Environ.*, 264, 112 632, <https://doi.org/10.1016/j.rse.2021.112632>, 2021.
- Doxani, G., Vermote, E., Roger, J.-C., Gascon, F., Adriaensen, S., Frantz, D., Hagolle, O., Hollstein, A., Kirches, G., Li, F., et al.: Atmospheric correction inter-comparison exercise, *Remote Sensing*, 10, 352, <https://doi.org/10.3390/rs10020352>, 2018.
- Dozier, J.: Spectral signature of alpine snow cover from the Landsat Thematic Mapper, *Remote Sens. Environ.*, 28, 9–22, [https://doi.org/10.1016/0034-4257\(89\)90101-6](https://doi.org/10.1016/0034-4257(89)90101-6), 1989.
- Duan, H., Xue, X., Wang, T., Kang, W., Liao, J., and Liu, S.: Spatial and temporal differences in alpine meadow, alpine steppe and all vegetation of the Qinghai-Tibetan Plateau and their responses to climate change, *Remote Sens.*, 13, 669, <https://doi.org/10.3390/rs13040669>, 2021.
- Farr, T. G., Hensley, S., Rodriguez, E., Martin, J., and Kobrick, M.: The shuttle radar topography mission, in: SAR workshop: CEOS Committee on Earth Observation Satellites, vol. 450, p. 361, <https://doi.org/10.1029/2005RG000183>, 2000.
- Feng, M., Sexton, J. O., Huang, C., Anand, A., Channan, S., Song, X.-P., Song, D.-X., Kim, D.-H., Noojipady, P., and Townshend, J. R.: Earth science data records of global forest cover and change: Assessment of accuracy in 1990, 2000, and 2005 epochs, *Remote Sens. Environ.*, 184, 73–85, <https://doi.org/10.1016/j.rse.2016.06.012>, 2016.
- Foody, G. M. and Mathur, A.: Toward intelligent training of supervised image classifications: directing training data acquisition for SVM classification, *Remote Sens. Environ.*, 93, 107–117, <https://doi.org/10.1016/j.rse.2004.06.017>, 2004.



- Friedl, M. A., McIver, D. K., Hodges, J. C., Zhang, X. Y., Muchoney, D., Strahler, A. H., Woodcock, C. E., Gopal, S., Schneider, A., Cooper, A., et al.: Global land cover mapping from MODIS: algorithms and early results, *Remote Sens. Environ.*, 83, 287–302, [https://doi.org/10.1016/S0034-4257\(02\)00078-0](https://doi.org/10.1016/S0034-4257(02)00078-0), 2002.
- Friedl, M. A., Sulla-Menashe, D., Tan, B., Schneider, A., Ramankutty, N., Sibley, A., and Huang, X.: MODIS Collection 5 global land cover: Algorithm refinements and characterization of new datasets, *Remote Sens. Environ.*, 114, 168–182, <https://doi.org/10.1016/j.rse.2009.08.016>, 2010.
- Friedman, J. H.: Greedy function approximation: a gradient boosting machine, *Ann. Stat.*, pp. 1189–1232, <https://doi.org/10.1214/aos/1013203451>, 2001.
- Fu, Y.-H., Gao, X.-J., Zhu, Y.-M., and Guo, D.: Climate change projection over the Tibetan Plateau based on a set of RCM simulations, *Adv. Clim. Chang. Res.*, 12, 313–321, <https://doi.org/10.1016/j.accre.2021.01.004>, 2021.
- Funk, C., Peterson, P., Landsfeld, M., Pedreros, D., Verdin, J., Shukla, S., Husak, G., Rowland, J., Harrison, L., Hoell, A., et al.: The climate hazards infrared precipitation with stations—a new environmental record for monitoring extremes, *Sci. Data*, 2, 1–21, <https://doi.org/10.1038/sdata.2015.66>, 2015.
- Gao, Q.-z., Li, Y., Xu, H.-m., Wan, Y.-f., and Jiangcun, W.-z.: Adaptation strategies of climate variability impacts on alpine grassland ecosystems in Tibetan Plateau, *Mitig. Adapt. Strateg. Glob. Chang.*, 19, 199–209, <https://doi.org/10.1007/s11027-012-9434-y>, 2014.
- Gao, T., Zhang, Y., Kang, S., Abbott, B. W., Wang, X., Zhang, T., Yi, S., and Gustafsson, Ö.: Accelerating permafrost collapse on the eastern Tibetan Plateau, *Environ. Res. Lett.*, 16, 054 023, <https://doi.org/10.1088/1748-9326/abf7f0>, 2021.
- Gislason, P. O., Benediktsson, J. A., and Sveinsson, J. R.: Random forests for land cover classification, *Pattern Recognit. Lett.*, 27, 294–300, <https://doi.org/10.1016/j.patrec.2005.08.011>, 2006.
- Gong, P., Wang, J., Yu, L., Zhao, Y., Zhao, Y., Liang, L., Niu, Z., Huang, X., Fu, H., Liu, S., et al.: Finer resolution observation and monitoring of global land cover: First mapping results with Landsat TM and ETM+ data, *Int. J. Remote Sens.*, 34, 2607–2654, <https://doi.org/10.1080/01431161.2012.748992>, 2013.
- Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., and Moore, R.: Google Earth Engine: Planetary-scale geospatial analysis for everyone, *Remote Sens. Environ.*, 202, 18–27, <https://doi.org/10.1016/j.rse.2017.06.031>, 2017.
- Grekousis, G., Mountrakis, G., and Kavouras, M.: An overview of 21 global and 43 regional land-cover mapping products, *Int. J. Remote Sens.*, 36, 5309–5335, <https://doi.org/10.1080/01431161.2015.1093195>, 2015.
- Hearst, M. A., Dumais, S. T., Osuna, E., Platt, J., and Scholkopf, B.: Support vector machines, *IEEE Intell. Syst. Appl.*, 13, 18–28, <https://doi.org/10.1109/5254.708428>, 1998.
- Hemmerling, J., Pflugmacher, D., and Hostert, P.: Mapping temperate forest tree species using dense Sentinel-2 time series, *Remote Sens. Environ.*, 267, 112 743, <https://doi.org/10.1016/j.rse.2021.112743>, 2021.
- Hua, T., Zhao, W., Liu, Y., Wang, S., and Yang, S.: Spatial consistency assessments for global land-cover datasets: A comparison among GLC2000, CCI LC, MCD12, GLOBCOVER and GLCNMO, *Remote Sens.*, 10, 1846, <https://doi.org/10.3390/rs10111846>, 2018.
- Hua, T., Zhao, W., Cherubini, F., Hu, X., and Pereira, P.: Sensitivity and future exposure of ecosystem services to climate change on the Tibetan Plateau of China, *Landsc. Ecol.*, 36, 3451–3471, <https://doi.org/10.1007/s10980-021-01320-9>, 2021.
- Huang, X., Yin, Y., Feng, L., Tong, X., Zhang, X., Li, J., and Tian, F.: A Dataset of Land Cover Samples over the Tibetan Plateau, Zenodo [data set], <https://doi.org/10.5281/zenodo.8227942>, 2023a.
- Huang, X., Yin, Y., Feng, L., Tong, X., Zhang, X., Li, J., and Tian, F.: A 10 m resolution land cover map of the Tibetan Plateau with detailed vegetation types, Zenodo [data set], <https://doi.org/10.5281/zenodo.8228112>, 2023b.



- Hwang, T., Song, C., Vose, J. M., and Band, L. E.: Topography-mediated controls on local vegetation phenology estimated from MODIS vegetation index, *Landsc. Ecol.*, 26, 541–556, <https://doi.org/10.1007/s10980-011-9580-8>, 2011.
- Karra, K., Kontgis, C., Statman-Weil, Z., Mazzariello, J. C., Mathis, M., and Brumby, S. P.: Global land use/land cover with Sentinel 2 and deep learning, in: 2021 IEEE international geoscience and remote sensing symposium IGARSS, pp. 4704–4707, Brussels, Belgium, 11–16 Jul 2021, <https://doi.org/10.1109/IGARSS47720.2021.9553499>, 2021.
- 375 Kumar, L. and Mutanga, O.: Google Earth Engine applications since inception: Usage, trends, and potential, *Remote Sens.*, 10, 1509, <https://doi.org/10.3390/rs10101509>, 2018.
- Li, J., Chen, F., Zhang, G., Barlage, M., Gan, Y., Xin, Y., and Wang, C.: Impacts of land cover and soil texture uncertainty on land model simulations over the central Tibetan Plateau, *J. Adv. Model. Earth Syst.*, 10, 2121–2146, <https://doi.org/10.1029/2018MS001377>, 2018.
- 380 Li, X., Zhu, X., Niu, Y., and Sun, H.: Phylogenetic clustering and overdispersion for alpine plants along elevational gradient in the Hengduan Mountains Region, southwest China, *J. Syst. Evol.*, 52, 280–288, <https://doi.org/10.1111/jse.12027>, 2014.
- Li, X., Zhu, W., Xie, Z., Zhan, P., Huang, X., Sun, L., and Duan, Z.: Assessing the effects of time interpolation of NDVI composites on phenology trend estimation, *Remote Sens.*, 13, 5018, <https://doi.org/10.3390/rs13245018>, 2021.
- Li, X., Long, D., Scanlon, B. R., Mann, M. E., Li, X., Tian, F., Sun, Z., and Wang, G.: Climate change threatens terrestrial water storage over the Tibetan Plateau, *Nat. Clim. Chang.*, 12, 801–807, <https://doi.org/10.1038/s41558-022-01443-0>, 2022.
- 385 Liu, C., Zhang, Q., Tao, S., Qi, J., Ding, M., Guan, Q., Wu, B., Zhang, M., Nabil, M., Tian, F., et al.: A new framework to map fine resolution cropping intensity across the globe: Algorithm, validation, and implication, *Remote Sens. Environ.*, 251, 112095, <https://doi.org/10.1016/j.rse.2020.112095>, 2020.
- Liu, C., Xu, X., Feng, X., Cheng, X., Liu, C., and Huang, H.: CALC-2020: a new baseline land cover map at 10 m resolution for the circumpolar Arctic, *Earth System Science Data*, 15, 133–153, <https://doi.org/10.5194/essd-15-133-2023>, 2023.
- 390 Liu, Q., Wang, X., Zhang, Y., and Li, S.: Complex ecosystem impact of rapid expansion of industrial and mining land on the Tibetan Plateau, *Remote Sens.*, 14, 872, <https://doi.org/10.3390/rs14040872>, 2022.
- Liu, S., Liu, X., Yu, L., Wang, Y., Zhang, G. J., Gong, P., Huang, W., Wang, B., Yang, M., and Cheng, Y.: Climate response to introduction of the ESA CCI land cover data to the NCAR CESM, *Clim. Dyn.*, 56, 4109–4127, <https://doi.org/10.1007/s00382-021-05690-3>, 2021.
- 395 Ma, H., Mo, L., Crowther, T. W., Maynard, D. S., van den Hoogen, J., Stocker, B. D., Terrer, C., and Zohner, C. M.: The global distribution and environmental drivers of aboveground versus belowground plant biomass, *Nat. Ecol. Evol.*, 5, 1110–1122, <https://doi.org/10.1038/s41559-021-01485-1>, 2021.
- Muñoz-Sabater, J.: ERA5-Land hourly data from 1981 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [data set], <https://doi.org/10.24381/cds.e2161bac>, 2019.
- 400 Muñoz-Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo, G., Boussetta, S., Choulga, M., Harrigan, S., Hersbach, H., et al.: ERA5-Land: A state-of-the-art global reanalysis dataset for land applications, *Earth Syst. Sci. Data*, 13, 4349–4383, <https://doi.org/10.5194/essd-13-4349-2021>, 2021.
- Nguyen, L. H., Joshi, D. R., Clay, D. E., and Henebry, G. M.: Characterizing land cover/land use from multiple years of Landsat and MODIS time series: A novel approach using land surface phenology modeling and random forest classifier, *Remote Sens. Environ.*, 238, 111017, <https://doi.org/10.1016/j.rse.2018.12.016>, 2020.
- 405 Pepin, N., Arnone, E., Gobiet, A., Haslinger, K., Kotlarski, S., Notarnicola, C., Palazzi, E., Seibert, P., Serafin, S., Schöner, W., et al.: Climate changes and their elevational patterns in the mountains of the world, *Rev. Geophys.*, 60, e2020RG000730, <https://doi.org/10.1029/2020RG000730>, 2022.



- Phan, T. N., Kuch, V., and Lehnert, L. W.: Land cover classification using Google Earth Engine and random forest classifier—The role of
410 image composition, *Remote Sens.*, 12, 2411, <https://doi.org/10.3390/rs12152411>, 2020.
- Pizarro, S. E., Pricope, N. G., Vargas-Machuca, D., Huanca, O., and Ñaupari, J.: Mapping land cover types for highland Andean ecosystems
in Peru using google earth engine, *Remote Sens.*, 14, 1562, <https://doi.org/10.3390/rs14071562>, 2022.
- Prats-Iraola, P., Nannini, M., Scheiber, R., De Zan, F., Wollstadt, S., Minati, F., Vecchioli, F., Costantini, M., Borgstrom, S., De Martino,
P., et al.: Sentinel-1 assessment of the interferometric wide-swath mode, in: 2015 IEEE international geoscience and remote sensing
415 symposium (IGARSS), pp. 5247–5251, Milan, Italy, 26–31 Jul 2015, <https://doi.org/10.1109/IGARSS.2015.7327018>, 2015.
- Rondeaux, G., Steven, M., and Baret, F.: Optimization of soil-adjusted vegetation indices, *Remote Sens. Environ.*, 55, 95–107,
[https://doi.org/10.1016/0034-4257\(95\)00186-7](https://doi.org/10.1016/0034-4257(95)00186-7), 1996.
- Rouse, J. W., Haas, R. H., Schell, J. A., Deering, D. W., et al.: Monitoring vegetation systems in the Great Plains with ERTS, *NASA Spec.*
Publ., 351, 309, 1974.
- 420 Salditt, M., Humberg, S., and Nestler, S.: Gradient Tree Boosting for Hierarchical Data, *Multivariate Behav. Res.*, pp. 1–27,
<https://doi.org/10.1080/00273171.2022.2146638>, 2022.
- Schepaschenko, D., See, L., Lesiv, M., Bastin, J.-F., Mollicone, D., Tsendbazar, N.-E., Bastin, L., McCallum, I., Laso Bayas, J. C., Baklanov,
A., et al.: Recent advances in forest observation with visual interpretation of very high-resolution imagery, *Surv. Geophys.*, 40, 839–862,
<https://doi.org/10.1007/s10712-019-09533-z>, 2019.
- 425 Shen, M., Piao, S., Cong, N., Zhang, G., and Jassens, I. A.: Precipitation impacts on vegetation spring phenology on the Tibetan Plateau,
Glob. Change Biol., 21, 3647–3656, <https://doi.org/10.1111/gcb.12961>, 2015.
- Shi, W., Zhao, X., Zhao, J., Zhao, S., Guo, Y., Liu, N., Sun, N., Du, X., and Sun, M.: Reliability and consistency assessment of land cover
products at macro and local scales in typical cities, *Int. J. Digit. Earth*, 16, 486–508, <https://doi.org/10.1080/17538947.2023.2181992>,
2023.
- 430 Shukla, T. and Sen, I. S.: Preparing for floods on the Third Pole, *Science*, 372, 232–234, <https://doi.org/10.1126/science.abh3558>, 2021.
- Souza Jr, C. M., Z. Shimbo, J., Rosa, M. R., Parente, L. L., A. Alencar, A., Rudorff, B. F., Hasenack, H., Matsumoto, M., G. Ferreira, L.,
Souza-Filho, P. W., et al.: Reconstructing three decades of land use and land cover changes in brazilian biomes with landsat archive and
earth engine, *Remote Sens.*, 12, 2735, <https://doi.org/10.3390/rs12172735>, 2020.
- Tian, F., Cai, Z., Jin, H., Hufkens, K., Scheifinger, H., Tagesson, T., Smets, B., Van Hoolst, R., Bonte, K., Ivits, E., et al.: Calibrating vegetation
435 phenology from Sentinel-2 using eddy covariance, PhenoCam, and PEP725 networks across Europe, *Remote Sensing of Environment*,
260, 112456, <https://doi.org/10.1016/j.rse.2021.112456>, 2021.
- Trew, B. T. and Maclean, I. M.: Vulnerability of global biodiversity hotspots to climate change, *Glob. Ecol. Biogeogr.*, 30, 768–783,
<https://doi.org/10.1111/geb.13272>, 2021.
- Tu, Y., Lang, W., Yu, L., Li, Y., Jiang, J., Qin, Y., Wu, J., Chen, T., and Xu, B.: Improved mapping results of 10 m resolution land cover
440 classification in Guangdong, China using multisource remote sensing data with Google Earth Engine, *IEEE J. Sel. Top. Appl. Earth*
Observ. Remote Sens., 13, 5384–5397, <https://doi.org/10.1109/JSTARS.2020.3022210>, 2020.
- Venter, Z. S., Barton, D. N., Chakraborty, T., Simensen, T., and Singh, G.: Global 10 m Land Use Land Cover Datasets: A Comparison of
Dynamic World, World Cover and Esri Land Cover, *Remote Sens.*, 14, 4101, <https://doi.org/10.3390/rs14164101>, 2022.
- Verde, N., Kokkoris, I. P., Georgiadis, C., Kaimaris, D., Dimopoulos, P., Mitsopoulos, I., and Mallinis, G.: National scale land cover classifi-
445 cation for ecosystem services mapping and assessment, using multitemporal copernicus EO data and google earth engine, *Remote Sens.*,
12, 3303, <https://doi.org/10.3390/rs12203303>, 2020.



- Wacker, A. and Landgrebe, D.: Minimum distance classification in remote sensing, LARS Technical Reports, p. 25, 1972.
- Wang, F., Ma, Y., Darvishzadeh, R., and Han, C.: Annual and Seasonal Trends of Vegetation Responses and Feedback to Temperature on the Tibetan Plateau since the 1980s, *Remote Sens.*, 15, 2475, <https://doi.org/10.3390/rs15092475>, 2023a.
- 450 Wang, X., Zhou, G., Lv, X., Zhou, L., Hu, M., He, X., and Tian, Z.: Comparison of Lake Extraction and Classification Methods for the Tibetan Plateau Based on Topographic-Spectral Information, *Remote Sens.*, 15, 267, <https://doi.org/10.3390/rs15010267>, 2023b.
- Wang, Y., Feng, L., Zhang, Z., and Tian, F.: An unsupervised domain adaptation deep learning method for spatial and temporal transferable crop type mapping using Sentinel-2 imagery, *ISPRS Journal of Photogrammetry and Remote Sensing*, 199, 102–117, <https://doi.org/10.1016/j.isprsjprs.2023.04.002>, 2023c.
- 455 Wang, Z., Wu, J., Niu, B., He, Y., Zu, J., Li, M., and Zhang, X.: Vegetation expansion on the Tibetan Plateau and its relationship with climate change, *Remote Sens.*, 12, 4150, <https://doi.org/10.3390/rs12244150>, 2020.
- Wang, Z., Song, W., and Yin, L.: Responses in ecosystem services to projected land cover changes on the Tibetan Plateau, *Ecol. Indic.*, 142, 109 228, <https://doi.org/10.1016/j.ecolind.2022.109228>, 2022.
- Xi, X., Liu, Z., Sun, L., Xie, S., and Wang, Z.: High-Confidence Sample Generation Technology and Application for Global Land-Cover Classification, *IEEE J. Sel. Top. Appl. Earth Observ. Remote Sens.*, 16, 3248–3263, <https://doi.org/10.1109/JSTARS.2022.3227911>, 2022.
- 460 Xie, S., Liu, L., Zhang, X., Yang, J., Chen, X., and Gao, Y.: Automatic land-cover mapping using landsat time-series data based on google earth engine, *Remote Sens.*, 11, 3023, <https://doi.org/10.3390/rs11243023>, 2019.
- Xu, H.: Modification of normalised difference water index (NDWI) to enhance open water features in remotely sensed imagery, *Int. J. Remote Sens.*, 27, 3025–3033, <https://doi.org/10.1080/01431160600589179>, 2006.
- 465 Xu, P., Tsendbazar, N.-E., Herold, M., Clevers, J. G., and Li, L.: Improving the characterization of global aquatic land cover types using multi-source earth observation data, *Remote Sens. Environ.*, 278, 113 103, <https://doi.org/10.1016/j.rse.2022.113103>, 2022.
- Yang, J. and Huang, X.: The 30 m annual land cover dataset and its dynamics in China from 1990 to 2019, *Earth Syst. Sci. Data*, 13, 3907–3925, <https://doi.org/10.5194/essd-13-3907-2021>, 2021.
- Yang, K., Wu, H., Qin, J., Lin, C., Tang, W., and Chen, Y.: Recent climate changes over the Tibetan Plateau and their impacts on energy and 470 water cycle: A review, *Glob. Planet. Change*, 112, 79–91, <https://doi.org/10.1016/j.gloplacha.2013.12.001>, 2014.
- Yang, L., Meng, X., and Zhang, X.: SRTM DEM and its application advances, *International Journal of Remote Sensing*, 32, 3875–3896, <https://doi.org/10.1080/01431161003786016>, 2011.
- Yang, Y., Xiao, P., Feng, X., and Li, H.: Accuracy assessment of seven global land cover datasets over China, *ISPRS-J. Photogramm. Remote Sens.*, 125, 156–173, <https://doi.org/10.1016/j.isprsjprs.2017.01.016>, 2017.
- 475 Yu, L., Wang, J., Li, X., Li, C., Zhao, Y., and Gong, P.: A multi-resolution global land cover dataset through multisource data aggregation, *Science China Earth Sciences*, 57, 2317–2329, <https://doi.org/10.1007/s11430-014-4919-z>, 2014.
- Zanaga, D., Van De Kerchove, R., Daems, D., De Keersmaecker, W., Brockmann, C., Kirches, G., Wevers, J., Cartus, O., Santoro, M., Fritz, S., Lesiv, M., Herold, M., Tsendbazar, N.-E., Xu, P., Ramoino, F., and Arino, O.: ESA WorldCover 10 m 2021 v200, Zenodo [data set], <https://doi.org/10.5281/zenodo.5571936>, 2022.
- 480 Zhang, G., Yao, T., Xie, H., Yang, K., Zhu, L., Shum, C., Bolch, T., Yi, S., Allen, S., Jiang, L., et al.: Response of Tibetan Plateau lakes to climate change: Trends, patterns, and mechanisms, *Earth-Sci. Rev.*, 208, 103 269, <https://doi.org/10.1016/j.earscirev.2020.103269>, 2020.
- Zhang, H. K. and Roy, D. P.: Using the 500 m MODIS land cover product to derive a consistent continental scale 30 m Landsat land cover classification, *Remote Sens. Environ.*, 197, 15–34, <https://doi.org/10.1016/j.rse.2017.05.024>, 2017.



- 485 Zhang, X., Liu, L., Chen, X., Gao, Y., Xie, S., and Mi, J.: GLC_FCS30: Global land-cover product with fine classification system at 30 m using time-series Landsat imagery, *Earth Syst. Sci. Data*, 13, 2753–2776, <https://doi.org/10.5194/essd-13-2753-2021>, 2021.
- Zhang, Y., Li, B., and Zheng, D.: Integration dataset of Tibet Plateau boundary, <https://data.tpdc.ac.cn/en/data/61701a2b-31e5-41bf-b0a3-607c2a9bd3b3/>, last access: 9 Aug, 2023, 2022.
- Zhao, F., Long, D., Li, X., Huang, Q., and Han, P.: Rapid glacier mass loss in the Southeastern Tibetan Plateau since the year 2000 from satellite observations, *Remote Sens. Environ.*, 270, 112 853, <https://doi.org/10.1016/j.rse.2021.112853>, 2022.
- 490 Zhou, J., Jia, L., and Menenti, M.: Reconstruction of global MODIS NDVI time series: Performance of Harmonic ANalysis of Time Series (HANTS), *Remote Sens. Environ.*, 163, 217–228, <https://doi.org/10.1016/j.rse.2015.03.018>, 2015.
- Zou, L., Tian, F., Liang, T., Eklundh, L., Tong, X., Tagesson, T., Dou, Y., He, T., Liang, S., and Fensholt, R.: Assessing the upper elevational limits of vegetation growth in global high-mountains, *Remote Sens. Environ.*, 286, 113 423, <https://doi.org/10.1016/j.rse.2022.113423>, 2023.