

CEDAR-GPP: spatiotemporally upscaled estimates of gross primary productivity incorporating CO₂ fertilization

Yanghui Kang^{1,2,3}, Maoya Bassiouni^{1,2}, Max Gaber^{1,4}, Xinchun Lu^{1,2}, Trevor F. Keenan^{1,2}

¹ Department of Environmental Science, Policy, and Management, University of California, Berkeley, Berkeley, CA 94720, USA

² Climate and Ecosystem Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

³ Department of Biological Systems Engineering, Virginia Tech, Blacksburg, VA 24061, USA

⁴ Department of Geosciences and Natural Resource Management, University of Copenhagen, Copenhagen, 1350, Denmark.

Correspondence: Yanghui Kang (yanghuikang@berkeley.edu, vt.edu)

Trevor Keenan (trevorkeen@berkeley.edu)

Abstract: Gross primary productivity (GPP) is the largest carbon flux in the Earth system, playing a crucial role in removing atmospheric carbon dioxide and providing carbohydrates needed for ecosystem metabolism. Despite the importance of GPP, however, existing estimates present significant uncertainties and discrepancies. A key issue is the underrepresentation of the CO₂ fertilization effect, a major factor contributing to the increased terrestrial carbon sink over recent decades. This omission could potentially bias our understanding of ecosystem responses to climate change.

Here, we introduce CEDAR-GPP, the first global machine-learning-upscaled GPP product that incorporates the direct CO₂ fertilization effect on photosynthesis. Our product is comprised of monthly GPP estimates and their uncertainty at 0.05° resolution from 1982 to 2020, generated using a comprehensive set of eddy covariance measurements, multi-source satellite observations, climate variables, and machine learning models. Importantly, we used both theoretical and data-driven approaches to incorporate the direct CO₂ effects. Our machine learning models effectively predicted monthly GPP ($R^2 \sim 0.72$), the mean seasonal cycles ($R^2 \sim 0.77$), and spatial variabilities ($R^2 \sim 0.63$) based on cross-validation at flux sites. After incorporating the direct CO₂ effects, predicted long-term

GPP trend across global flux towers substantially increases~~ed~~ from $3.1 \text{ gC}_m^{-2}\text{year}^{-1}$ to $4.5 - 5.4 \text{ gC}_m^{-2}\text{year}^{-1}$, which aligns more closely with the $7.7 \text{ gC}_m^{-2}\text{year}^{-1}$ trend detected from eddy covariance data. While the global patterns of annual mean GPP, seasonality, and interannual variability generally align~~ed~~ with existing satellite-based products, CEDAR-GPP demonstrates~~ed~~ higher long-term trends globally after incorporating CO_2 fertilization and reflected a strong temperature control on direct CO_2 effects. The estimated global GPP trend ~~was-is~~ $0.57 - 0.76 \text{ PgC per-year}^{-1}$ from 2001 to 2018 and $0.32 - 0.34 \text{ PgC per-year}^{-1}$ from 1982 to 2020. Estimating and validating GPP trends in data-scarce regions, such as the tropics, remains challenging, underscoring the importance of ongoing ground-based monitoring and advancements in modeling techniques. ~~Nevertheless,~~ CEDAR-GPP offers a comprehensive representation of GPP temporal and spatial dynamics, providing valuable insights into ecosystem-climate interactions. The CEDAR-GPP product is available at <https://zenodo.org/doi/10.5281/zenodo.8212706> (Kang et al., 2024).

1. Introduction

Terrestrial ecosystem photosynthesis, known as Gross Primary Productivity (GPP), is the primary source of food and energy for the Earth system and human society (Keenan and Williams, 2018). Through photosynthesis, terrestrial ecosystems also mitigate climate change, by removing thirty percent of anthropogenic carbon emissions from the atmosphere each year (Friedlingstein et al., 2023). However, due to the lack of direct measurements at the global scale, our understanding of photosynthesis and its spatiotemporal dynamics is limited, leading to considerable disagreements among various GPP estimates (Anav et al., 2015; O’Sullivan et al., 2020; Smith et al., 2016; Yang et al., 2022). Addressing these uncertainties is crucial for improving the predictability of ecosystem dynamics under climate change (Friedlingstein et al., 2014).

Over the past three decades, global networks of eddy covariance flux towers collected *in situ* carbon flux measurements that allow for accurate estimates of GPP, providing valuable insights into photosynthesis dynamics under various environmental conditions (Baldocchi, 2020; Beer et al., 2010). To quantify and understand GPP at scales and locations beyond the $\sim 1\text{km}^2$ flux tower footprints, machine learning has been employed with gridded satellite and climate datasets to upscale site-based measurements and produce wall-to-wall GPP maps (Dannenberg et al., 2023; Joiner and Yoshida, 2020; Jung et al., 2011; Tramontana et al., 2016; Xiao et al., 2008; Yang et al., 2007; Zeng et al., 2020). This “upscaling” approach provides data-driven and observation-based quantifications without prescribed functional relations between GPP and its climatic or environmental drivers. It offers unique empirical constraints of ecosystem carbon dynamics, complementing those derived from process-based and semi-process-based approaches such as terrestrial biosphere models or the Light Use Efficiency (LUE) models (Beer et al., 2010; Gampe et al., 2021; Jung et al., 2017; Schwalm et al., 2017). In recent years, the growth of global and regional flux networks, coupled with increasing efforts in data standardization, has offered new opportunities for the advancement of upscaling frameworks, enabling comprehensive quantifications of terrestrial photosynthesis (Joiner and Yoshida, 2020; Nelson et al., 2024; Pastorello et al., 2020).

Effective machine learning upscaling depends on a complete set of input predictors that fully explain GPP dynamics. Upscaled datasets have primarily relied on satellite-observed greenness indicators, such as vegetation indices, Leaf Area Index (LAI), the fraction of absorbed photosynthetically active radiation (fAPAR), which effectively capture canopy-level GPP dynamics related to leaf area changes (Joiner and Yoshida, 2020; Ryu et al., 2019; Tramontana et al., 2016).

However, important aspects of leaf-level physiology, such as those controlled by climate factors, are often omitted in major upscaled datasets, preventing accurate characterization of GPP responses to climate change (Bloomfield et al., 2023; Stocker et al., 2019). In particular, none of the previous upscaled datasets have considered the direct effect of atmospheric CO₂ on leaf-level photosynthesis, which is a key factor contributing to at least half of the enhanced land carbon sink observed over the past decades (Keenan et al., 2016, 2023; Ruehr et al., 2023; Walker et al., 2021). This omission can lead to incorrect inferences regarding long-term trends in various components of the terrestrial carbon cycle (De Kauwe et al., 2016).

Multiple independent lines of evidence from the atmospheric inversion (Wenzel et al., 2016), atmospheric ¹³C/¹²C measurements (Keeling et al., 2017), ice core records of carbonyl sulfide (Campbell et al., 2017), glucose isotopomers (Ehlers et al., 2015), as well as free-air CO₂ enrichment experiments (FACE) (Walker et al., 2021), suggest a widespread positive effect of elevated atmospheric CO₂ on GPP from site to global scales. Increasing atmospheric CO₂ *directly* stimulates the biochemical rate or the light use efficiency (LUE) of leaf-level photosynthesis, known as the direct CO₂ fertilization effect (CFE). Enhanced photosynthesis could lead to greater net carbon assimilation, contributing to an increase in total leaf area. This expansion, contributing to a higher light interception, further enhances canopy-level photosynthesis (i.e. GPP), which is referred to as the indirect CFE. The direct CFE has been found to dominate GPP responses to CO₂ compared to the indirect effect, from both theoretical and observational analyses (Chen et al., 2022; Haverd et al., 2020).

Satellite-based estimates have shown an increasing global GPP trend in the past few decades largely attributable to CO₂-induced increases in LAI (Chen et al., 2019; De Kauwe et al., 2016; Piao et al., 2020; Zhu et al., 2016). However, previous upscaled GPP datasets, as well as most LUE models such as the MODIS GPP product, have failed to consider the direct CO₂ effects on leaf-level biochemical processes (Jung et al., 2020; Zheng et al., 2020). Consequently, these products likely underestimated the long-term trend of global GPP, leading to large discrepancies when compared to process-based models, which typically consider both direct and indirect CO₂ effects (Anav et al., 2015; De Kauwe et al., 2016; Keenan et al., 2023; O’Sullivan et al., 2020). Notably, recent improvements in LUE models have included the CO₂ response and show improved long-term changes in GPP globally (Zheng et al., 2020), yet, this important mechanism is still missing in GPP products upscaled from *in situ* eddy covariance flux measurements based on machine learning models.

To improve the quantification of GPP spatial and temporal dynamics and provide a robust representation of long-term dynamics in global photosynthesis, we developed the CEDAR-GPP¹ data product. CEDAR-GPP was upscaled from global eddy covariance carbon flux measurements using machine learning along with a broad range of multi-source satellite observations and climate variables. In addition to incorporating direct CO₂ fertilization effects on photosynthesis, we also account for indirect effects via greenness indicators and include novel satellite datasets such as solar-induced fluorescence (SIF), Land Surface Temperature (LST) and soil moisture to explain variability under environmental stresses. We provide monthly GPP estimations and associated uncertainties at 0.05° resolution derived from ten model setups. These setups differ by the temporal range depending on satellite data availability, the method for incorporating the direct CO₂ fertilization effects, and the partitioning approach used to derive GPP from eddy covariance measurements. Short-term model setups ~~were~~are primarily based on data derived from MODIS satellites generating GPP estimates from 2001 to 2020, while long-term estimates span~~ned~~ 1982 to 2020 using combined Advanced Very High Resolution Radiometer (AVHRR) and MODIS data. We used two approaches to incorporate the direct CO₂ fertilization effects, including direct prescription with eco-evolutionary theory and machine learning inference from the eddy-covariance data. Additionally, we ~~provided~~ a baseline configuration that did not incorporate the direct CO₂ effects. Uncertainties in GPP estimation were quantified using bootstrapped model ensembles. We evaluated the machine learning models' skills in predicting monthly GPP, seasonality, interannual variability, and trend against eddy covariance measurements, and compared the CEDAR-GPP spatial and temporal variability to existing satellite-based GPP estimates.

2. Data and Methods

2.1 Eddy covariance data

We obtained monthly eddy covariance GPP measurements from 2001 to 2020 from the FLUXNET2015 (Pastorello et al., 2020), AmeriFlux FLUXNET (<https://ameriflux.lbl.gov/data/flux-data-products/>), and ICOS Warm Winter 2020 (Warm Winter 2020 Team, 2022) datasets. All data were processed with the ONEFLUX pipeline (Pastorello et al., 2020). Following previous upscaling efforts (Tramontana et al., 2016), we selected monthly GPP data

¹ CEDAR stands for upsCaling Ecosystem Dynamics with ARtificial intelliGence

with at least 80% of high-quality hourly or half-hourly data for temporal aggregation. High-quality data refers to GPP derived from measured or high-quality gap-filled Net Ecosystem Exchange (NEE) data. We further excluded large negative GPP values, setting a cutoff of $-1 \text{ gC}_m^{-2}\text{day}^{-1}$. We utilized GPP estimates from both the night-time (GPP_REF_NT_VUT) and day-time (GPP_REF_DT_VUT) partitioning approaches. We classified flux tower sites according to the C3 and C4 plant categories reported in metadata and related publications when available and used a C4 plant percentage map (Still et al., 2003) otherwise. This classification information is included in Supplementary Text S1. Our analysis encompassed 233 sites, predominantly located in North America, Western Europe, and Australia (Figure 1). A list of the sites is provided in Appendix A. Despite their uneven geographical distribution, these sites effectively cover a diverse range of climatic conditions and are representative of global biomes (Figure 1c, 1d). In total, our dataset included over 18000 site-months. Note that we did not include eddy covariance data before 2001, since it was limited to only a few sites with only four sites containing data before 1996. This scarcity might introduce biases in the machine learning models, particularly in the relationship between GPP and CO_2 , leading to unreliable extrapolations across space and time in the long-term predictions.




Figure 1. (a) Spatial distribution of eddy covariance sites used to generate the CEDAR-GPP product. (b) Annual site counts. (c) Site counts by biomes. ENF: evergreen needleleaf forests, EBF: evergreen broadleaf forests, DBF: deciduous broadleaf forests, MF: mixed forests, WSA: woody savannas, SAV: savannas, OSH: open shrublands, CSH: closed shrublands, GRA: grasslands, CRO: croplands, WET: wetlands. (d) Sites distributions in the annual temperature and precipitation space. Whittaker biome classification is shown as a reference of natural vegetation based on long-term climatic conditions. It does not directly indicate the actual biome associated with each site. The base map in (a) was obtained from the NASA Earth Observatory map by Joshua Stevens using data from NASA’s MODIS Land Cover, the Shuttle Radar Topography Mission (SRTM), the General Bathymetric Chart of the Oceans (GEBCO), and Natural Earth boundaries. Whittaker biomes were plotted using the “plotbiomes” R package (Stefan and Levin, 2018).

2.2 Global input datasets

We compiled an extensive set of covariates from gridded climate reanalysis data, multi-source satellite datasets including optical, thermal, and microwave observations, as well as categorical information on land cover, climate zone, and C3/C4 classification. The datasets that we compiled offer comprehensive information about GPP dynamics and its responses to climatic variabilities and stresses. Table 1 lists the datasets and associated variables used to generate CEDAR-GPP.

Table 1. Datasets used in different model setups to generate the CEDAR GPP product. Refer to Table S1 for a list of specific variables from each dataset.

Category	Dataset	Temporal coverage	Spatial resolution	Temporal resolution	Usage in model setups		Reference
					Short-term	Long-term	
Climate	ERA5-Land	1950 – present	0.1°	Monthly	✓	✓	(Sabater, 2019)
	ESRA Global Monitoring Laboratory Atmospheric Carbon Dioxide	1976 – present	-	Monthly	✓	✓	(Thoning et al., 2021)
Satellite-based datasets	MODIS Nadir BRDF-adjusted reflectance (MCD43C4v006)	2000 – present	0.05°	Daily	✓		(Schaaf and Wang, 2015)
	MODIS Terra and Aqua LAI/fPAR (MCD15A3H, MOD15A2H, v006)	2000 – present	500m	4-day, 8-day	✓		(Myneni et al., 2015a, b)
	MODIS Terra and Aqua LST (MYD11A1, MOD11A1, v006)	2000 – present	1 km	Daily	✓		(Wan et al., 2015b, a)
	BESS_Rad	2000 – 2020	0.05°	Daily	✓		(Ryu et al., 2018)
	Continuous-SIF (from OCO-2 and MODIS)	2000 – 2020	0.05°	4-day	✓		(Zhang, 2021)
	ESA CCI Soil Moisture Combined Passive and Active v06.1	1979 – 2021	0.25°	Daily	✓		(Gruber et al., 2019)
	GIMMS LAI4g	1982 – 2021	0.0833°	Half-month		✓	(Cao et al., 2023)
	GIMMS NDVI4g	1982 – 2021	0.0833 °	Half-month		✓	(Li et al., 2023b)
Static categorical datasets	MODIS Land Cover (MCD12Q1v006)	Average status used between 2001 and 2020	500m	-	✓	✓	(Friedl and Sulla-Menashe, 2019)
	Koppen-Geiger Climate Classification	present	1 km	-	✓	✓	(Beck et al., 2018)
	C4 percentage map	present	1°	-	✓	✓	(Still et al., 2003, 2009)

168

169

2.2.1 Climate variables

We obtained air temperature, vapor pressure deficit, precipitation, potential evapotranspiration, and skin temperature from the EAR5-Land reanalysis dataset (Sabater, 2019) (Table 1; Table S1). We applied a three-month lag to precipitation, to reflect the memory of soil moisture and represent the root zone water availability. Averaged monthly atmospheric CO₂ concentrations were calculated as an average of records from the Mauna Loa Observatory and South Pole Observation stations, retrieved from NOAA’s Earth System Research Laboratory (Thoning et al., 2021).

2.2.2 Satellite datasets

We assembled a broad collection of satellite-based observations of vegetation greenness and structure, LST, solar radiation, solar-induced fluorescence (SIF), and soil moisture (Table 1, Table S1).

We used three MODIS version 6 products: surface reflectance, LAI/fAPAR, and LST. Surface reflectance from optical to infrared bands (band 1 to 7) was sourced from the MODIS Nadir BRDF-adjusted reflectance (NBAR) daily dataset (MCD43C4) (Schaaf and Wang, 2015). From these data, we derived vegetation indices, including NIRv (Badgley et al., 2019), kNDVI (Camps-Valls et al., 2021), NDVI, Enhanced Vegetation Index (EVI), Normalized Difference Water Index (NDWI) (Gao, 1996), and the green chlorophyll index (CIgreen) (Gitelson, 2003). We also used snow percentages from the NBAR dataset. We used the 4-day LAI and fPAR composite derived from Terra and Aqua satellites (MCD15A3H) (Myneni et al., 2015a; Yan et al., 2016a, b) from July 2002 onwards and the MODIS 8-day LAI and fPAR dataset from Terra only (MOD15A2H) prior to July 2002 (Myneni et al., 2015b). We used day-time and night-time LST from the Aqua satellite (MYD11A1) (Wan et al., 2015b), with the Terra-based LST product (MOD11A1) used after July 2002 (Wan et al., 2015a). Terra LST was bias-corrected with the differences in the mean seasonal cycles between Aqua and Terra following Walther et al. (2022).

We used the PKU GIMMS NDVI4g dataset (Li et al., 2023b) and PKU GIMMS LAI4g (Cao et al., 2023) datasets available from 1982 to 2020. PKU GIMMS NDVI4g is a harmonized time series that includes AVHRR-based NDVI from 1982 to 2003 (with biases and corrections mitigated through inter-calibration with Landsat surface reflectance images) and MODIS NDVI from 2004 onward. PKU GIMMS LAI4g consisted of consolidated AVHRR-based LAI from 1982 to 2003 (generated using machine learning models trained with Landsat-based LAI data and NDVI4g) and reprocessed MODIS LAI (Yuan et al., 2011) from 2004 onwards.

We utilized photosynthetically active radiation (PAR), diffusive PAR, and shortwave downwelling radiation from the BESS_Rad dataset (Ryu et al., 2018). We obtained the continuous-SIF (CSIF) dataset (Zhang, 2021; Zhang et al., 2018) produced by a machine learning algorithm trained using OCO-2 SIF observations and MODIS surface reflectance. We used surface soil moisture from the ESA CCI soil moisture combined passive and active product (version 6.1) (Dorigo et al., 2017; Gruber et al., 2019).

2.2.3 Other categorical datasets

We used plant functional type (PFT) information derived from the MODIS Land Cover product (MCD12Q1) (Friedl and Sulla-Menashe, 2019). We followed the International Geosphere-Biosphere Program classification scheme but merged several similar categories to maximize the amount of eddy covariance sites/observations available for each category. Closed shrublands and open shrublands are combined into a shrubland category. Woody savannas and savannas are combined into savannas. We generated a static PFT map by taking the mode of the MODIS land cover time series between 2001 – 2020 at each pixel to mitigate uncertainties from misclassification in the MODIS dataset. Nevertheless, changes in vegetation structure induced by land use and land cover change are reflected in the dynamics surface reflectance and LAI/fAPAR datasets we used. We used the Koppen-Geiger main climate groups (tropical, arid, temperate, cold, and polar) (Beck et al., 2018). We also utilized a C4 plant percentage map to account for different photosynthetic pathways when incorporating CO₂ fertilization (Still et al., 2003, 2009). The C4 percentage dataset was constant over time.

2.2.4 Data preprocessing

We implemented a three-step preprocessing strategy for the satellite datasets: 1) quality control, 2) gap-filling, and 3) spatial and temporal aggregation. Firstly, we selected high-quality data based on the quality control flags of the satellite products when available. For the MODIS NBAR dataset (MCD43C3), we used data with 75% or more high-resolution NBAR pixels retrieved with full inversions for each band. For MODIS LST, we selected the best quality data from the quality control bitmask as well as data where retrieved values had an average emissivity error of no more than 0.02. For MODIS LAI/fAPAR, we used retrievals from the main algorithm with or without saturation. We used all available data in ESA-CCI soil moisture due to the presence of substantial data gaps. In the gap-filling step, missing values in satellite datasets were temporally filled at the native temporal resolution, following a two-step protocol adapted from Walther et al (2021). Short temporal gaps were first filled with medians from a moving window, and the remaining gaps were filled with the mean

seasonal cycle. For datasets with a high temporal resolution, including MODIS NBAR (daily), LAI/fPAR (4-day), BESS (4-day), CSIF (4-day), ESA-CCI (daily), temporal gaps no longer than 5 days (8 days for 4-day resolution products) were filled with medians of 15-day moving windows in the first step. An exception is MODIS LST (daily), for which we used a shorter moving window of 9 days due to rapid changes in surface temperature. GIMMS LAI4g and NDVI4g data were only filled with mean seasonal cycle due to their low temporal resolution (half-month). This is because vegetation structure could experience significant changes at half-month intervals, and gap-filling using temporal medians within moving windows could introduce considerable uncertainties and potentially over-smooth the time series.

Finally, all the datasets were aggregated to a monthly time step and 0.05-degree spatial resolution. We employed the conservative resampling approach using the xESMF python package (Zhuang et al., 2023). To generate the machine learning model training data, we extracted values from the nearest 0.05 degree pixel relative to the site locations within the gridded dataset.




Figure 2. Schematic overview of the CEDAR-GPP model setups.

2.3 Machine learning upscaling

2.3.1 CEDAR-GPP model setups

We trained machine learning models with eddy covariance GPP measurements as targets and climate/satellite variables as input features. We created ten model setups to produce different global monthly GPP estimates (Figure 2; Table 2). The model setups were characterized by the temporal range depending on input data availability, the configuration of CO₂ fertilization effects, and the partitioning approach used to derive the GPP from eddy covariance measurements.

The short-term (ST) model configuration produced GPP from 2001 to 2020, and the long-term (LT) configuration spanned 1982 to 2020. Each temporal configuration uses a different set of input variables depending on their availability. Inputs for the short-term configuration included MODIS,

CSIF, BESS PAR, ESA-CCI soil moisture, ERA5-Land, as well as PFT and Koppen Climate zone as categorical variables with one-hot encoding. The long-term used GIMMS NDVI4g and LAI4g data, ERA5-land, PFT and Koppen climate. ESA CCI soil moisture datasets were excluded from the long-term model setups due to concerns about the product quality in the early years when the number and quality of microwave satellite data were limited (Dorigo et al., 2015). A detailed list of input features for each setup is provided in Table S1.

Regarding the direct CO₂ fertilization effects (CFE), we established a “Baseline” configuration that did not incorporate these effects, a “CFE-Hybrid” configuration that incorporated the effects via eco-evolutionary theory, and a “CFE-ML” configuration that inferred the direct effects from eddy covariance data using machine learning. Detailed information about these approaches is provided in Sec. 2.3.2. Furthermore, separate models were trained for GPP target variables from the night-time (NT) and daytime (DT) partitioning approaches.

Table 2 lists the characteristics of ten model setups. Due to the limited availability of eddy covariance observations before 2001, we did not apply the CFE-ML approach to the long-term setups. The CFE-ML model, when trained on data from 2001 to 2020 with atmospheric CO₂ ranging from 370 to 412 ppm, would not accurately predict GPP response to CO₂ for the period 1982 – 2000 when the CO₂ levels were markedly lower (340 – 369 ppm). This is because machine learning models, especially tree-based models, could not extrapolate beyond the range of the training data.

Table 2. Specifications of the CEDAR-GPP model setups.

Model Setup Name	Temporal range	Direct CO ₂ Fertilization Effects		GPP Partitioning Method
		Configuration	Method	
ST_Baseline_NT	Short-term (ST) 2001 – 2020	Baseline	Not incorporated	Night-time (NT)
ST_Baseline_DT				Day-time (DT)
ST_CFE-Hybrid_NT		CFE-Hybrid	Theoretical	NT
ST_CFE-Hybrid_DT				DT
ST_CFE-ML_NT		CFE-ML	Data-driven	NT
ST_CFE-ML_DT				DT
LT_Baseline_NT	Long-term (LT) 1982 – 2020	Baseline	Not incorporated	NT
LT_Baseline_DT				DT
LT_CFE-Hybrid_NT		CFE-Hybrid	Theoretical	NT
LT_CFE-Hybrid_DT				DT

2.3.2 CO₂ fertilization effect

We established three configurations regarding the direct CO₂ fertilization effects on photosynthesis. In the baseline configuration, we trained machine learning models with eddy

covariance GPP measurements, input climate and satellite features, but excluding CO₂ concentration. As such, the models only include indirect CO₂ effects from the satellite-based proxies of vegetation greenness or structure representing changes in canopy light interception, and they do not consider the direct effect of CO₂ on leaf-level photosynthetic rates (or light use efficiency, LUE). Our baseline model is therefore directly comparable to other satellite-derived GPP products that only account for indirect CO₂ effects (Joiner and Yoshida, 2020; Jung et al., 2020).

In the CFE-ML configuration, we added monthly CO₂ concentration into the feature set in addition to those incorporated in the baseline models. Models inferred the functional relationship between GPP and CO₂ from the eddy covariance data. They thus encompass both CO₂ fertilization pathways – direct effects on LUE and indirect effects from the satellite-based proxies of vegetation greenness and structure.

In the CFE-Hybrid configuration, we applied biophysical theory to estimate the response of LUE to elevated CO₂, i.e. the direct CFE (Appendix A). First, we estimated a reference GPP, where LUE was not affected by any increase in atmospheric CO₂, by applying the CFE-ML model with a constant atmospheric CO₂ concentration equal to the 2001 level while keeping all other variables temporally dynamic. Then, the impacts of CO₂ on LUE were prescribed onto the reference GPP estimates using a theoretical CO₂ sensitivity function of LUE according to the optimal coordination theory (Appendix A). The theoretical CO₂ sensitivity function represents a CO₂ sensitivity that is equivalent to that of the electron-transport-limited (light-limited) photosynthetic rate. When light is limited, elevated CO₂ suppresses photorespiration leading to increased photosynthesis at a lower rate than when photosynthesis is limited by CO₂ (Lloyd and Farquhar, 1996; Smith and Keenan, 2020). Thus, the CFE-Hybrid scenario provides a conservative estimation of the direct CO₂ effects on LUE. Note that the theoretical sensitivity function describes the fractional change in LUE due to direct CO₂ effects relative to a reference period (i.e. 2001). Therefore, we used the CFE-ML model to establish this reference GPP by fixing the CO₂ effects to the 2001 level, rather than simply using the GPP from the Baseline model in which the direct CO₂ effects were not represented. Long-term trends from the reference and the Baseline models are consistent.

For both CFE-ML and CFE-Hybrid scenarios, we made another conservative assumption that C4 plants do not benefit from elevated CO₂, despite potential increases in photosynthesis during water-limited conditions due to enhanced water use efficiency (Walker et al., 2021). Data from flux tower sites dominated by C4 plants were removed from our training set, so the machine learning models inferred CO₂ fertilization only from flux tower sites dominated by C3 plants. When applying

models globally, we assumed the reference GPP values (with constant atmospheric CO₂ concentration equal to the 2001 level) to represent C4 plants, and GPP estimates from CFE-ML or CFE-Hybrid models were applied in proportion to the percentage of C3 plants in a grid cell.

2.3.3 Machine learning model training and validation

We employed the state-of-the-art XGBoost machine learning model, known for its high accuracy in regression problems across various domains, including environmental and ecological predictions (Berdugo et al., 2022; Chen and Guestrin, 2016; Kang et al., 2020). XGBoost is a scalable and parallelized implementation of the gradient boosting technique that iteratively trains an ensemble of decision trees, with each iteration targeting to minimize the residuals from the last iteration. A notable merit of XGBoost is its ability to make predictions in the presence of missing values, a common issue in remote sensing datasets. Without relying on prior assumptions about the functional forms or statistical distributions, the model is also robust to multi-collinearity between the predictors in our dataset, particularly for the variables derived from MODIS data.

We used five-fold cross-validation for model evaluation. Training data was randomly split into five groups (folds), with each fold held out for testing while the rest four folds were used for model training. We imposed two restrictions on fold splitting: each flux site was entirely assigned to a fold to test model performance over unseen locations; the random sampling was stratified based on PFT to ensure coverage of the full range of PFTs in both training and testing. We also used a nested-cross-validation strategy, during which we performed a randomized search of hyperparameters using three-fold cross-validation within the training set. The nested-cross-validation was aimed to reduce the risk of overfitting and improve the robustness of the evaluation.

We assessed the models' ability to capture the temporal and spatial characteristics of GPP, including monthly GPP, mean seasonal cycles, monthly anomalies, and cross-site variability. Model performance was assessed separately for each model setup (Table 2) and summarized by PFT and Koppen climate zone. Mean seasonal cycles were calculated as the mean monthly GPP over the site observation period, and monthly anomalies were the residuals of monthly GPP after subtracting mean seasonal cycles. Monthly GPP averaged over years for each site was used to assess cross-site variability. Goodness-of-fit metrics include RMSE, bias, and coefficient of determination (R^2).

To evaluate the models' ability to capture long-term GPP trends, we aggregated the monthly GPP to annual values following Chen et al. (2022), which detected the CO₂ fertilization effect across global eddy covariance sites. For sites with at least five years of observations, GPP anomalies were

computed by subtracting the multi-year mean GPP from the annual GPP for each site. Anomalies were aggregated across sites to achieve a single multi-site GPP anomaly per year. We excluded a site-year if less than 11 months of data was available and used linear interpolation to fill the remaining temporal gaps. This resulted in 81 sites used in the GPP trend evaluation. We used the Sen slope and Mann-Kendall test to examine the GPP trends from 2002 to 2019, excluding 2001 and 2020, due to the limited number of available sites with more than five years of data. We further assessed the aggregated annual trend by grouping the sites based on plant functional types and the Koppen climate zones. Categories with less than six long-term sites available were excluded from the analysis, which includes EBF and Tropics.

To further analyze GPP responses to CO₂ in the CFE-ML models, we leveraged two explainable machine learning approaches: ALE (Accumulated Local Effects) (Apley and Zhu, 2020; Baniecki et al., 2021) and SHAP (SHapley Additive exPlanations) (Lundberg and Lee, 2017). SHAP is a model interpretation method derived from game theory, providing a value for each feature's contribution to a prediction, elucidating how each feature impacts the model's output in a specific instance. Conversely, ALE quantifies the average effect of a feature across the data, isolating its impact by aggregating local effects and avoiding the biases associated with correlated features.

2.3.4 Product generation and uncertainty quantification

In the CEDAR-GPP product, we generated GPP estimates from each of the ten model setups, by applying the model to global gridded datasets within the corresponding temporal range (Table 2). GPP estimates were named after the corresponding model setups. We used bootstrapping to quantify estimate prediction uncertainties. For each model setup, we generated 30 bootstrapped sample sets of eddy covariance data, which were then used to train an ensemble of 30 XGBoost models. The bootstrapping was performed at the site level, and each bootstrapped sample set contained around 140 to 150 unique sites, 17000 to 19000 site months covering all PFTs. The relative PFT composition in the bootstrapped sample sites was consistent with the full dataset. Hyperparameters of the XGBoost models used in the final product generation were described in Supplementary Text S2. The 30 models trained with bootstrapped samples generated an ensemble of 30 GPP values. We provided the ensemble GPP mean and used standard deviation to indicate uncertainties, for each of the ten model setups.

2.4 Product inter-comparison

We compared the global spatial and temporal patterns of CEDAR-GPP with other major satellite-based GPP products, including three machine learning upscaled and two LUE-based datasets. We obtained two FLUXCOM products (Jung et al., 2020), the latest version of FLUXCOM-RS (FLUXCOM-RSv006) available from 2001 to 2020 based on remote sensing (MODIS collection 6) datasets only, as well as the FLUXCOM-RS+METEO ensemble available between 1979 to 2018 and based on the climatology of remote sensing observations and ERA5 forcings (hereafter FLUXCOM-ERA5). We used FluxSat (Joiner and Yoshida, 2020), available from 2001 to 2019, which is an upscaled dataset based on MODIS NBAR surface reflectance and PAR from Modern-Era Retrospective analysis for Research and Applications 2 (MERRA-2). Importantly, FluxSat does not incorporate climate forcings. We used the MODIS GPP product (MOD17) available since 2001, which was generated based on MODIS fAPAR and LUE as a function of air temperature and vapor pressure deficit but not atmospheric CO₂ concentration (Running et al., 2015). We also used the rEC-LUE products, available from 1982 to 2018 and based on a revised LUE model that incorporated the effect of atmospheric CO₂ concentration and the fraction of diffuse PAR on LUE (Zheng et al., 2020). Additionally, to evaluate GPP trends, we compared our product against three process-based models forced by remote sensing data – BEPS (Leng et al., 2024), BESSv2 (Li et al., 2023a), and PML V2 (Zhang et al., 2019). These products estimate GPP by scaling leaf-level biochemical photosynthesis models to the canopy level, using satellite-derived vegetation structural variables such as LAI. All three products incorporate the direct CO₂ effects within their biochemical photosynthesis models.

All datasets were resampled to 0.1 ° spatial resolution, and a common mask for the vegetated land area was applied. We evaluated global mean annual GPP, mean seasonal cycle, interannual variability, and trend among different datasets, comparing them over a common time period determined by their data availability. Global total GPP was computed by scaling the global area-weighted average GPP flux with the global land area (122.4 million km²) following Jung et al. (2020). Mean seasonal cycle was defined as above (Sec. 2.3.3). We used the standard deviation of annual GPP to indicate the magnitude of interannual variability, the Sen slope to indicate the GPP annual trend, and the Mann-Kendall test for the statistical significance of trends.

3. Results

3.1 Evaluation of model performance

3.1.1 Overall performance

The short-term and long-term models explained approximately 72% and 67%, respectively, of the variation in monthly GPP across global eddy covariance sites (Figure 3a). The long-term models consistently yielded lower performance than the short-term models, likely due to differences in the satellite remote sensing datasets used, as the short-term models benefited from richer information from surface reflectance of individual bands, LST, CSIF, as well as soil moisture, while the long-term model only exploited NDVI and LAI. The models with different CFE configurations and target GPP variables (i.e. partitioning approaches) have similar performance in predicting monthly GPP (Figure 3b, Table S2). All models exhibited minimal bias of less than 0.1.

Model performance in terms of the different temporal and spatial characteristics of monthly GPP ~~was-is~~ variable (Figure 3c-h). The models ~~were-are~~ most successful at predicting mean seasonal cycles, with the short-term and long-term models explaining around 77% and 72% of the variability, respectively (Figure 3c-d). The short-term and long-term models captured 63% and 54% , respectively, of the spatial variabilities in multi-year mean GPP across global sites (i.e., cross-site variability) (Figure 3g-h). However, all models underestimated monthly anomalies across the sites, with R^2 values below 0.12 (Figure 3e-f). Patterns from the DT setups do not significantly differ from those of the NT setups (Figure S1, Table S2). Model performance also varies across sites, and models ~~were-are~~ more advantageous in explaining mean seasonal cycles than monthly anomalies in most sites (Figure S2).




Figure 3. Machine learning model performance in predicting monthly GPP and its spatial and temporal variability. Only NT models are shown and DT results ~~is~~are provided in Supplementary Figure S1. Scatter plots illustrated relationships between model predictions and observations for monthly GPP (a), mean seasonal cycles (MSC) (c), monthly anomaly (e), and cross-site variability (g) for ST_CFE-Hybrid_NT (left, blue) and LT_CFE-Hybrid_NT (right, green) models. Corresponding bar plots show the R^2 values for five NT model setups in predicting monthly GPP (b), MSC (d), monthly anomaly (f), and cross-site variability (h).

3.1.2 Performance by biome and climate zone

The predictive ability of our models varied across different PFTs and Koppen climate zones (Figure 4). Here we present results from the CFE-Hybrid LT and ST models based on NT partitioning and note that patterns for the other CFE configurations and the DT GPP were similar (Figure S3, S4, S5).

Model performance in terms of monthly GPP was the highest for deciduous broadleaf forests, mixed forests, and evergreen needleleaf forests, with R^2 values above 0.76. Model accuracies were also high for savannas and grasslands, followed by croplands and wetlands, with R^2 values between 0.48 and 0.76. Model accuracies were lowest in evergreen broadleaf forests and shrublands, with R^2 values as low as 0.13. Across climate zones, models achieved the highest accuracy in predicting monthly GPP in cold climates with R^2 around 0.73 – 0.78, followed by tropics and temperate zones ($R^2 \sim 0.47 - 0.65$). The short-term models had the lowest performance in polar regions with an R^2 value of around 0.37, and the long-term models had the lowest performance in arid regions with an R^2 value of 0.28. Interestingly, short-term and long-term models exhibited substantial differences in arid regions and shrublands marked by strong seasonality and interannual variabilities.

Model performance in terms of mean seasonal cycles across PFTs and climate zones followed patterns for monthly GPP, while disparities emerged for performance in terms of GPP anomaly and cross-site variability (Figure 4, Figure S3, S4, S5). The short-term model showed the highest predictive power in explaining monthly anomalies in arid regions with an R^2 value of 0.48, where savanna and shrublands sites are primarily located. Model performance in all other climate zones was significantly lower. The short-term model also demonstrated good performance in capturing anomalies in deciduous broadleaf forests. The long-term model's relative performance between PFTs and climate zones was mostly consistent with that of the short-term model, with lower accuracy in shrublands when compared to the short-term model.

Models demonstrated the highest accuracy in predicting cross-site variability in savannas, grasslands, evergreen needleleaf forests, and evergreen broadleaf forests ($R^2 > 0.36$) and the lowest accuracy in deciduous broadleaf forests, mixed forests, and croplands ($R^2 < 0.1$). The short-term model additionally showed good performance in shrublands and wetlands ($R^2 > 0.36$), whereas the long-term model failed to capture any variability for shrublands. In terms of climate zones, models were most successful at explaining the variabilities within tropical and cold climate zones ($R^2 > 0.50$), the short-term model had moderate performance in temperate and polar regions ($R^2 \sim 0.22$),

and the long-term model has low performance for both temperate and arid regions with R^2 values below 0.16.




Figure 4. Performance of the ST_CFE-Hybrid_NT (blue) and LT_CFE-Hybrid_NT (green) models on GPP spatiotemporal estimation by plant functional types (a) and climate zones (b). The cross-site panels included the number of sites within each category. Color indicates short-term (ST) or long-term (LT) models. ENF: evergreen needleleaf forest, EBF: evergreen broadleaf forest, DBF: deciduous broadleaf forest, MF: mixed forest, SH: shrubland, SA: savanna, GRA: grassland, CRO: cropland, WET: wetland. Tr: tropical, Ar: arid, Tp: temperate, Cd: cold, Pl: polar. The performance of DT models is displayed in Supplementary Figure S3.

3.1.3 Prediction of long-term trends

Eddy covariance derived GPP presents a substantial increasing trend across flux sites between 2002 and 2019 (Figure 5a, Figure S6a). The eddy covariance GPP from the night-time partitioning approach indicates an overall trend of $7.7 \text{ gC}_m^{-2}\text{year}^{-2}$. In contrast, the ST_Baseline_NT model predicts a more modest overall trend of $3.1 \text{ gC}_m^{-2}\text{year}^{-2}$ across the flux sites, primarily reflecting the indirect CO_2 effect manifested through the growth of LAI. Both the ST_CFE-ML_NT and ST_CFE-hybrid_NT models predicted much higher trends of 5.4 and $4.5 \text{ gC}_m^{-2}\text{year}^{-2}$ respectively, representing an improvement from the Baseline model by 74% and 45%, aligning more closely to eddy covariance observations. Similarly, the LT_CFE-Hybrid_NT model shows an improved trend estimation than the LT_Baseline_NT model. All trends were statistically significant ($p < 0.05$).

Aggregated eddy covariance GPP experiences increasing trends of varied magnitudes across different climate zones and plant functional types (Figure 5b,c; Figure S6b,c). While the machine learning models generally do not fully capture the enhancement in GPP for most categories, the CFE-ML and/or CFE-hybrid models consistently outperformed the Baseline models in both ST and LT setups. The CFE-ML setup predicts a higher trend than CFE-hybrid in most cases, suggesting that the data-driven approach captures more dynamics not represented in the theoretical model, which was based on conservative assumptions regarding the CO_2 sensitivity of photosynthesis (see Sect. 2.3.2 and Appendix A). The choice of remote sensing data (ST vs. LT configurations) did not lead to substantial differences in the predicted GPP trend. Most long-term flux sites (at least 10 years of records) with a significant trend experienced an increase in GPP, and the CFE-ML and/or CFE-hybrid models aligned closer to eddy covariance data than the Baseline models (Figure S7). Additionally, we found a considerably higher trend in eddy covariance GPP measurements derived from the day-time versus night-time partitioning approach, potentially associated with uncertainties in GPP partitioning methods (Figure S6). Yet, machine learning model predicted trends were not strongly affected by GPP partitioning methods.




Figure 5. Comparison of observed and predicted GPP (from NT models only) trends across eddy covariance flux towers. (a) Aggregated annual GPP anomaly from 2002 to 2019 and trend lines from eddy covariance (EC) data, and three CFE model setups (short-term, night-time partitioning) for ST (left) and LT (right) models. The size of grey circle markers is proportional to the number of sites. (b) Comparison of annual GPP trends from eddy covariance measurements and the short-term (ST) CEDAR-GPP model setups by plant functional types and climate zones. (c) Comparison of annual GPP trends from eddy covariance measurements and the long-term (LT) CEDAR-GPP model setups by plant functional types and climate zones. In (b) and (c), Categories with less than 6 sites, including Tropics and EBF, were not shown. While dots on the bars indicate statistically significant trends with p -value < 0.1 . Results for the DT models are shown in Supplementary Figure S3.

The differences in estimated GPP trends between the Baseline and CFE models underscore the significant long-term GPP changes driven by the direct CO₂ effect. Using explainable machine learning

approaches – ALE and SHAP, we further assessed the CFE-ML models for quantifying the direct CO₂ effect. Both approaches revealed a consistently positive influence of CO₂ on GPP, aligning with biophysical theories (Figure S8). Compared to the effects from light (PAR) and vegetation structures (e.g. NIRv), the impacts of CO₂ are considerably smaller, which explains the minimal differences in overall model accuracy between the Baseline and CFE models.

Finally, we evaluated CEDAR-GPP using independent eddy covariance data (11 sites, Table S3) that was not involved in model training and obtained from the OzFlux FluxNet dataset (Ozflux, 2024). Among these sites, only two - AU-Cpr (Tropical) and AU-Stp (Arid) - with more than five years of records exhibit a GPP trend with p-value less than 0.3. CEDAR-GPP shows strong consistency with the observed trend (Figure S9). Additionally, CEDAR-GPP achieves reasonable accuracy in predicting monthly GPP ($R^2 \sim 0.73 - 0.75$), mean seasonal cycle ($R^2 \sim 0.74 - 0.78$), and monthly anomalies ($R^2 \sim 0.26 - 0.50$) (Table S4, Figure S10), closely aligning with the cross-validation results.

3.2 Evaluation of GPP spatial and temporal dynamics

We compared CEDAR-GPP estimates with other upscaled or LUE-based datasets regarding the mean annual GPP (Sect. 3.2.1), GPP seasonality (Sect. 3.2.2), interannual variability (Sect. 3.2.3), and annual trends (Sect. 3.2.4). CEDAR-GPP model setups generally showed similar patterns in mean annual GPP, seasonality, and interannual variability, therefore, in corresponding sections, we present the CFE-Hybrid model setups as representative examples for comparisons with other datasets, unless otherwise stated. Supplementary figures include comparisons involving CEDAR-GPP estimates from all model setups.

3.2.1 Mean annual GPP

Global patterns of mean annual GPP ~~were~~are generally consistent among CEDAR-GPP model setups, FLUXCOM, FLUXSAT, MODIS, and rEC-LUE, with few noticeable regional differences (Figure 6, Figure S11~~8~~). Differences among CEDAR-GPP model setups ~~were~~are minimal and only evident between the NT and DT setups in the tropics (Figure 6b-c, Figure S8~~S~~11). CEDAR-GPP short-term datasets showed highest consistency with FLUXSAT in terms of mean annual GPP magnitudes (2001 – 2018) and latitudinal variations, although FLUXSAT presents slightly higher GPP values in the tropics compared to CEDAR-GPP (Figure 6b). Mean annual GPP magnitudes for FLUXCOM-RS006 and MODIS ~~was~~are lower globally than CEDAR-GPP and FLUXSAT, with the most pronounced differences observed in the tropical areas. Among the long-term datasets (CEDAR-

GPP LT, FLUXCOM-ERA5, and rEC-LUE), mean annual GPP (1982 – 2018) exhibits greater disparities in the northern mid-latitudes than in the tropics and southern hemisphere (Figure 6c). CEDAR-GPP aligned more closely with FLUXCOM-ERA5 than with rEC-LUE, with the latter showing lower annual mean GPP globally, particularly between 20°N to 50° N.




Figure 6. Global distributions of mean annual GPP from CEDAR-GPP and other machine learning upscaled and LUE-based reference datasets. (a) Global patterns of mean annual GPP from two short-term datasets including ST_CFE-Hybrid_NT, and FLUXCOM-RS006, and two long-term datasets including LT_CFE-Hybrid_NT, and FLUXCOM-ERA5. (b) Latitudinal distributions of mean annual GPP from short-term

552 datasets (ST_CFE-Hybrid_NT, FLUXSAT, FLUXCOM-RS006, and MODIS). (c)
553 Latitudinal distributions of mean annual GPP from long-term datasets (LT_CFE-
554 Hybrid_NT, FLUXCOM-ERA5, and rEC-LUE). Mean annual GPP was computed
555 between 2001 and 2018 for short-term datasets and between 1982 and 2018 for long-
556 term datasets.

557 3.2.2 Seasonal variability

558 CEDAR-GPP agrees~~d~~ with other GPP datasets on seasonal variabilities (average between 2001
559 and 2018) at the global scale, characterized by a peak in GPP in July and a nadir between December
560 and January (Figure 7, Figure S129). At the global scale, CEDAR-GPP ~~was~~is most closely aligned
561 with FLUXSAT in GPP seasonal magnitude and amplitude, while both FLUXCOM and MODIS
562 display~~d~~ a relatively less pronounced magnitude.

563 In boreal and temperate regions of the Northern Hemisphere-, all datasets agree~~d~~ on seasonal
564 GPP variation, with only minor variances in the magnitude of peak GPP. In Southern Hemisphere
565 temperate regions, datasets demonstrate~~d~~ similar seasonality, though with greater variability in peak
566 amplitudes compared to the Northern Hemisphere. The largest disparities ~~were~~are found in the South
567 American tropical areas, where seasonal variation is less prominent. Here, FLUXSAT show~~s~~ed a
568 distinct bi-modal pattern with peaks in March-April and September-October. CEDAR-GPP and
569 FLUXCOM-ERA5 align~~s~~ed with the second peak, but exhibit~~d~~ a less pronounced first peak.
570 Interestingly, the DT setups of CEDAR-GPP show~~d~~ slightly higher peaks in March-April in this
571 region (Figure S130). MODIS, in contrast, indicat~~e~~s an inverse seasonal pattern with a small peak
572 from June to August. Across all regions, CEDAR-GPP's seasonality align~~s~~ed more closely with
573 FLUXSAT and FLUXCOM-ERA5 than with other datasets. Differences among the ten C~~E~~D~~A~~R-
574 GPP model setups ~~were~~are minimal, except for small variations in GPP magnitude in some tropical
575 areas between NT and DT setups (Figure S130).




Figure 7. Comparison of global and regional GPP mean seasonal cycle between different datasets on a global scale. Monthly means were averaged from 2001 to 2018 for all datasets. Geographic boundaries of [these the 11 TransCom land](#) regions were obtained from the CarbonTracker (CT2022) dataset [and shown in -Figure S19.](#)

3.2.3 Interannual variability

We found distinct spatial patterns in GPP interannual variability between upscaled and LUE-based datasets and a high level of agreement within each category, with the exception of FLUXCOM-ERA5, which showed minimal interannual variability globally (Figure 8, Figure S144). All datasets agreed on the presence of GPP interannual variability hotspots in eastern and southern South America, central North America, southern Africa, and western Australia. These hotspots primarily corresponded to arid and semi-arid areas characterized by grasslands, shrubs, and croplands (Figure 9). CEDAR-GPP ~~was~~^{is} highly consistent with FLUXSAT, and both datasets also displayed relatively high interannual variability in the dry subhumid areas of Europe, predominantly covered by croplands. FLUXCOM-RS006 mirrored the relative spatial patterns of CEDAR-GPP and FLUXSAT, albeit at lower magnitudes. The LUE-based datasets (MODIS and rEC-LUE) predicted a much higher interannual variability than the upscaled datasets in the tropical areas, particularly in evergreen broadleaf forests and woody savannas (Figure 8, Figure 9). These datasets also depicted slightly higher interannual variability for other types of forests, including evergreen needleleaf forests and deciduous

broadleaf forests, compared to the upscaled datasets. The lack of interannual variability in FLUXCOM-ERA5 is attributable to the use of mean seasonal cycles of remotely sensed vegetation greenness indicators rather than their dynamic time series. Ten CEDAR-GPP model setups presented consistent patterns in interannual variability, and differences were minimal (Figure S144).




Figure 8. Spatial patterns of GPP interannual variability extracted over 2001 to 2018 for CEDAR-GPP (ST_CFE-Hybrid_NT), FLUXSAT, FLUXCOM-RS006, MODIS, FLUXCOM-ERA5, and rEC-LUE.




Figure 9. Comparison of GPP interannual variability (IAV) across global datasets by PFT. Colored dots represent the median IAV, thicker gray bars indicate the 25% to 75% percentiles of IAV distributions, and thinner grey bars show the 10% to 90% percentiles.

3.2.4 Trends

Differences in annual GPP trends among CEDAR-GPP model setups and other upscaled and LUE-based datasets mainly reflected the variability in the representation of CO₂ fertilization effects (Figure 10, Figure S152). From 2001 to 2018, the CEDAR-GPP Baseline model setups showed spatial variations in GPP trends consistent with the other upscaled datasets without direct CO₂ fertilization effects, including FLUXSAT and FLUXCOM-RSv006. In these datasets, substantial increases were seen in southeastern China and India, western Europe, and part of North and South America. These increases were largely associated with rising LAI due to land use changes and indirect CO₂ fertilization effects, as identified by previous studies (Chen et al., 2019; Zhu et al., 2016). Although MODIS, which also does not include a direct CO₂ fertilization effect, generally agreed with these increasing trends, it also showed a declining GPP in the tropical Amazon and a stronger positive trend in central South America. After incorporating the direct CO₂ fertilization effects, both the CFE-Hybrid and CFE-ML setups predicted positive trends in tropical forests, an observation absent in all other datasets. Furthermore, the CFE-Hybrid and CFE-ML models also revealed increasing GPP in

temperate and boreal forests of North America and Eurasia. Notably, all datasets agreed on a pronounced GPP decrease in eastern Brazil.

From 2001 to 2018, a positive trend in global annual GPP ~~was-is~~ uniformly detected by all datasets, albeit with varying magnitudes (Figure 11a-b). The ST_Baseline_NT model predicted a GPP growth rate of 0.35 (± 0.02) Pg C per year, aligning with FLUXCOM-RS, but lower than FLUXSAT (0.51 Pg C yr⁻²) and MODIS (0.39 Pg C yr⁻²) (Figure 11b). The CFE-hybrid models estimated a notably faster GPP growth at 0.58 (± 0.03) Pg C yr⁻². The CFE-ML models predicted the highest trends, up to 0.76 (± 0.15) Pg C yr⁻² from the ST_CFE-ML_NT model and 0.59 (± 0.13) Pg C yr⁻² from the ST_CFE-ML_DT model. Also, a higher variance ~~was-is~~ observed among ensemble members in the ST_CFE-ML setups compared to the ST_Baseline and ST_CFE-Hybrid models.

The LT_Baseline_NT model identified increasing GPP trends in large areas of Europe, East and South Asia, as well as the Northern Amazon from 1982 to 2020 (Figure 10b). The pattern from the LT_CFE-Hybrid_NT model aligned closely with the LT_Baseline_NT model but exhibited a stronger positive trend in global tropical areas as well as Eurasian boreal forests. In contrast, FLUXCOM-ERA5 showed overall negative trends in the tropics, with a small magnitude. Lastly, rEC-LUE agreed with positive GPP trends identified in CEDAR-GPP in the extratropical areas, but predicted a pronounced negative trend in the tropics. At the global scale, all the CEDAR-GPP long-term models predicted a positive global GPP trend (Figure 11d). The LT_Baseline_NT and LT_Baseline_DT models showed a trend of 0.13 (± 0.02) ~~to-and~~ 0.15 (± 0.02) Pg C yr⁻² ~~respectively~~, while the LT_CFE-Hybrid_NT and LT_CFE-Hybrid_DT models ~~setups-doubled these~~ rates with 0.33 (± 0.02) and 0.31 (± 0.03) Pg C yr⁻² respectively. rEC-LUE showed a two-phased pattern with a strong increase in GPP from 1982 to 2000 (0.54 Pg C yr⁻²), followed by a decreasing trend after 2001 (-0.20 Pg C yr⁻²) (Figure S16). This resulted in an overall positive change at a rate comparable to that of the Baseline model. FLUXCOM-ERA5 exhibited a small negative trend.




Figure 10. Annual GPP trend over 2001 – 2018 for short-term CEDAR-GPP, FLUXCOM-RS006, FLUXSAT, and MODIS datasets (a) and over 1982 – 2018 for long-term CEDAR-GPP, FLUXCOM-ERA5 and rEC-LUE datasets (b). Hatched

areas indicate the GPP trend that is statistically significant at $p < 0.05$ level under the Mann-Kendall test.




Figure 11. Global annual GPP variations (a) and trends (b) from 2001 to 2018 for short-term CEDAR-GPP, FLUXCOM-RS006, FLUXSAT, and MODIS datasets. Global annual GPP variations (c) and trends (d) over 1982 to 2018 for long-term for long-term CEDAR-GPP, FLUXCOM-ERA5, and rEC-LUE datasets. Error bars in (b) and (d) represent the 25% to 75% percentile from the model ensembles of CEDAR-GPP. Dots in (b) and (d) indicate the minimum and maximum from the model ensembles of CEDAR-GPP.

Among the process-based models forced by remote sensing data, CEDAR-GPP aligns closely with BESS v2 and BEPS, showing widespread positive GPP trends in the boreal and temperate regions

of the Northern Hemisphere, as well as across tropical areas (Figure S17, S18). In contrast, PML V2 presents minimal GPP changes in tropics and substantial reduction in Africa. The CEDAR-GPP CFE-ML model exhibits higher trends in tropical and northern hemisphere temperate regions, while BEPS shows more pronounced trends in boreal regions. None of the products indicates significant GPP trends in Australia. Globally, trends from BEPS and BESS V2 are around 0.55 PgC year⁻¹, consistent with CEDAR's ST_CFE-Hybrid_NT model at 0.58 PgC year⁻¹, while PML V2 displays a neutral trend of 0.08 PgC year⁻¹.

3.3 GPP estimation uncertainties

We analyzed the spread between the 30 model ensemble members in CEDAR-GPP as an indicator of uncertainties in GPP estimations. The spatial pattern of uncertainty in estimating annual mean GPP largely resembles that of the mean map (Figure 12, Figure 6a). The largest model spread ~~was-is~~ found in highly productive tropical forests, and this uncertainty decreased in temperate and cold areas (Figure 12a). Tropical ecosystems, with a mean annual GPP between 1000 to 3500 PgC year⁻¹, only exhibited a 2% and 6% variation within the model ensemble (Figure 12b). Ecosystems in the temperate and cold climates ~~had-have~~ a smaller annual GPP and proportionally small uncertainties of up to 6%. However, ecosystems in Arid and Polar climates, despite their similarly low GPP, showed higher model uncertainty, reaching 10% to 40% of the ensemble mean.

The estimation uncertainty of GPP trends ~~was-is~~ generally below 15% to 20% in the CEDAR-GPP datasets under the ST_Baseline and ST_CFE-Hybrid setups (Figure 12c). However, in the ST_CFE-ML setup, the estimation increases substantially, with model spread reaching up to 40% in tropical areas. Figure S20 further illustrates the trend uncertainties with the ensemble mean error range based on one standard deviation. Both the CFE-ML models show large discrepancies between the upper and lower uncertainty ranges particularly within the tropics. Additionally, Notably, the long-term models also showed a higher uncertainty compared to the short-term models.




Figure 12. CEDAR-GPP estimation uncertainty derived from ensemble spread (standard deviation of 30 model predictions). (a) Spatial patterns of the absolute standard deviation from ensemble members in estimating the mean annual GPP from 2001 to 2018, using data from the ST_CFE-Hybrid_NT setup. (b) Relationships between ensemble standard deviation and ensemble mean in mean annual GPP. Colored contours denote clusters of Koppen climate zones. Dashed lines indicate the ratio between the ensemble standard deviation and the ensemble mean with values shown in percentage. (c) Spatial patterns of model uncertainty in GPP long-term trend estimation. Only areas where 90% of the ensemble members showed a statistically significant trend ($p < 0.05$) are shown in the maps. The trend for the short-term datasets

(left column) was computed between 2001 to 2018. The trend for the long-term datasets (right column) was computed between 1982 to 2018.

4. Discussion

4.1 Reducing uncertainties in GPP upscaling

Here we examine the three predominant sources of uncertainties in machine learning upscaling of GPP: eddy covariance measurements, input datasets, and the machine learning model. We discuss strategies used in CEDAR-GPP to reduce the impacts of these uncertainties and highlight potential future research directions.

4.1.1 Eddy covariance data

Uncertainties associated with eddy covariance measurement and data processing can propagate through the upscaling process. CEDAR-GPP was produced using monthly aggregated eddy covariance data, where the impact of random errors in half-hourly measurements was minimized due to the temporal aggregation (Jung et al., 2020). Our stringent quality screening further reduced data processing uncertainties such as those associated with gap-filling. Yet, the discrepancy in GPP patterns between the CEDAR-GPP NT and DT setups is indicative of systematic biases linked to the partitioning approaches used to derive GPP from the NEE measurements (Keenan et al., 2019; Pastorello et al., 2020). Interestingly, the mean annual GPP from the DT setup ~~was~~is slightly higher than that from the NT setup (Figure 6), and the DT setup also predicts ~~a~~ed a higher GPP trend in the long-term dataset (Figure 11). While these discrepancies ~~were~~are relatively small compared to the predominant spatiotemporal patterns, the separate DT and NT setups in CEDAR-GPP offered ~~an~~ interesting quantification of the GPP partitioning uncertainties over space and time, providing insights for future methodology improvements.

The unbalanced spatial representativeness of the eddy covariance data constitutes a more significant source of uncertainty, as highlighted by previous studies (Jung et al., 2020; Tramontana et al., 2015). Effective generalization of machine learning models requires a substantial volume of training data that adequately represents and balances varied conditions. In CEDAR-GPP, this issue was mitigated with a large set of eddy covariance data (~18000 site-months) integrating FLUXNET2015 and two regional networks. However, data availability remains limited in critical carbon exchange hotspots such as tropics, subtropics, drylands, and boreal regions, as well as in

mountainous areas (Figure 1). Contrary to widespread perception that sparse training data leads to high upscaling uncertainties, our findings from the bootstrapped model spread indicate modest uncertainties in tropical areas relative to their high GPP magnitude (Figure 12). This observation aligns with findings from the FLUXCOM product, revealing low extrapolation uncertainty in humid tropical regions (Jung et al., 2020). Nevertheless, to fully understand the upscaling uncertainty, it is essential to evaluate the generalization or extrapolation errors within the predictor space and consider the potential limitations of model structures (van der Horst et al., 2019; Villarreal and Vargas, 2021). Additionally, data limitations in mountainous areas and the absence of topology information in the predictor space in our models suggest potential uncertainties related to topographical effects on GPP (Hao et al., 2022; Xie et al., 2023).

Furthermore, our analysis suggests that the estimated global GPP magnitudes were related to the specific eddy covariance GPP data used in upscaling. Notably, global GPP magnitudes derived from CEDAR-GPP closely aligned with those from FLUXSAT, while the estimates from FLUXCOM were considerably lower (Figure 6, Figure 11). FLUXSAT used eddy covariance data from FLUXNET2015, which largely overlapped with that included in CEDAR-GPP (Joiner and Yoshida, 2020). FLUXCOM utilized data from FLUXNET La Thuile set and CarboAfrica network, which consist of a distinct set of sites (Tramontana et al., 2016). The influence from the predictor datasets was minimal since all three datasets relied on MODIS-derived products. For a more in-depth evaluation of the impacts of flux site representativeness on upscaling, future research directions could include conducting synthetic experiments with simulations of ensembles of terrestrial biosphere models.

4.1.2 Input predictors and controlling factors

Upscaled GPP inherent uncertainties from the input predictors, including satellite and climate datasets. First, satellite remote sensing data contains noises resulting from sun-earth geometry, atmospheric conditions, soil background, and geolocation inaccuracies. The models or algorithms used for retrieving LAI, fAPAR, LST, and soil moisture, also contain random errors and systematic biases specific to certain regions, biome types, or climatic conditions (Fang et al., 2019; Ma et al., 2019; Yan et al., 2016b). Moreover, satellite observations frequently contain missing values due to clouds, aerosols, snow, and algorithm failure, leading to both systematic and random uncertainties. In producing CEDAR-GPP, we mitigated these uncertainties through comprehensive preprocessing procedures. Our temporal gap-filling strategy exploits both the temporal dependency of vegetation

status and long-term climatology, to reduce biases from missing values. Temporal and spatial aggregation further reduced the remaining data gaps and random noises. Nevertheless, considerable uncertainties likely remained in satellite datasets impacting the upscaled estimations.

A potentially more impactful source of uncertainty is the mismatch between the footprint of the eddy covariance measurements and the coarse resolution of satellite observations. While flux towers typically have a footprint of around $\sim 1 \text{ km}^2$ (Chu et al., 2021), satellite observations employed in CEDAR-GPP and most other upscaled datasets ~~were~~are at 5 km or lower resolution. Systematic and random errors could be introduced due to this mismatch, particularly in heterogenous biomes and areas with a mixture of vegetation and non-vegetated land covers. One mitigation strategy is to generate upscaled datasets at a higher spatial resolution (e.g. 500m). Alternatively, models could be trained at a high resolution and applied to the coarse resolution to reduce computation and storage requirements (Dannenberg et al., 2023; Gaber et al., 2024). However, this approach does not address inherent scaling errors in coarse-resolution satellite images (Dong et al., 2023; Yan et al., 2016a).

Besides the quality of predictors, successful machine learning upscaling also requires a comprehensive set of features representing all controlling factors. For example, the lack of GPP interannual variabilities in FLUXCOM-ERA5 manifests the importance of incorporating dynamic vegetation signals from remote sensing in the upscaling framework. CEDAR-GPP used satellite observations from optical, thermal, and microwave systems as well as climate variables thoroughly representing GPP dynamics. Particularly, the inclusion of LST and soil moisture data provides important information about resource limitations and stress factors, which are crucial for certain biomes and/or under specific conditions (Green et al., 2022; Stocker et al., 2018, 2019). Dannenberg et al. (2023) showed that incorporating LST from MODIS and soil moisture from the SMAP satellite datasets substantially improved the machine learning estimation accuracy of GPP in North American drylands. Nevertheless, accurately capturing interannual anomalies remains challenging for certain biomes, such as evergreen needleleaf forest, cropland, and wetland (Figure 4), as acknowledged by previous studies (Tramontana et al., 2016; Jung et al., 2020). High prediction uncertainties (Figure 12) in drylands also suggest the machine learning models did not sufficiently represent the mechanisms of water stress and drought responses. Potential improvement may be achieved by incorporating datasets related to agricultural management practices (crop type, cultivar, irrigation, fertilization) (Xie et al., 2021), plant hydraulic and physiological properties (Liu et al., 2021), dynamic C4 plant distributions (Luo et al., 2024), root and soil characteristics (Stocker et al., 2023), as well as topography (Xie et al., 2023).

4.1.3 Machine learning models and uncertainty quantification

The choice of machine learning models and their parameterization has been found to have a relatively minor impact on GPP upscaling uncertainties (Tramontana et al., 2015). CEDAR used the state-of-the-art boosting algorithm, XGBoost, which provided high performance given the current data availability. Further reduction of model uncertainty will likely rely on additional information, such as increasing the number of eddy covariance sites or incorporating more high-quality predictors. Additionally, temporal dependency of carbon fluxes responses to atmospheric controls may also be exploited with specialized deep neural networks such as recurrent neural networks or transformers (Besnard et al., 2019; Ma and Liang, 2022).

A key challenge, however, is the quantification of uncertainties in machine learning upscaling (Reichstein et al., 2019). The limited availability of eddy covariance data hinders a comprehensive assessment of the extrapolation errors; consequently, metrics of predictive performance from cross-validation are inherently biased. CEDAR derived estimation uncertainty for each GPP prediction using bootstrapping model ensemble, which naturally mimics the sampling bias associated with flux tower locations. Notably, the choice of input climate reanalysis datasets could also induce systematic differences in GPP spatial and temporal patterns (Tramontana et al., 2015). As a result, the FLUXCOM product generates model ensembles based on different reanalysis datasets to capture these uncertainties. Additionally, different satellite datasets of vegetation structural proxies, such as LAI, also exhibited significant discrepancies (Jiang et al., 2017). Thus, an ensemble approach combining site-level bootstrapping with multiple sources of input predictors could potentially provide a more comprehensive quantification of uncertainties. Furthermore, tree-based models do not generalize well to unseen conditions, and the uncertainty estimates derived from bootstrapping of XGBoost models may underrepresent actual biases stemming from limitations in training data representation. Future work may explore Bayesian neural networks, which provide uncertainty along with predictions and, at the same time, present high predictive power comparable to ensemble tree-based algorithms (Ma et al., 2021).

4.2 Long-term GPP changes and CO₂ fertilization effect

CEDAR-GPP was constructed using a comprehensive set of climate variables and multi-source satellite observations, thus encapsulating long-term GPP dynamics from both direct and indirect effects of climate controls. Particularly, CEDAR-GPP included the direct CO₂ fertilization effect,

which has been shown to dominate the increasing trend of global photosynthesis (Chen et al., 2022). Incorporating these effects substantially improved long-term trends of GPP from site to global scales (Figure 5, 10, 11). CEDAR's CFE-Hybrid setup offered a conservative estimation of the direct CO₂ effects by simulating the CO₂ sensitivity of light-limited LUE for C3 plants (Walker et al., 2021). However, the model ~~did~~ does not account for the impacts of nutrient availability, which could potentially constrain CO₂ fertilization (Peñuelas et al., 2017; Reich et al., 2014; Terrer et al., 2019). Robust modeling of LUE responses to rising CO₂ under various environmental conditions remains challenging (Wang et al., 2017). Future work is needed to better understand how these factors affect the quantification of GPP and its long-term temporal variations.

The CFE-ML model adopted a data-driven approach to infer CO₂ effects directly from eddy covariance data. This strategy allowed the model to potentially capture multiple physiological pathways of the CO₂ impact evidenced in the eddy covariance measurements, including the increases of the biochemical rates and enhancements in the water use efficiency (Keenan et al., 2013). The model detected a strong positive effect of CO₂ on eddy covariance measured GPP, consistent with previous studies based on process-based and statistical models (Chen et al., 2022; Fernández-Martínez et al., 2017; Ueyama et al., 2020). Moreover, spatial patterns of GPP trends derived from the CFE-ML model reflected a strong temperature dependency, aligning with the anticipated temperature sensitivity of photosynthetic biochemical processes (Keenan et al., 2023). Yet, the considerable ensemble spread in the CO₂ trends from the CFE-ML model and discrepancies between the CFE setups (Figure 11, Figure 13) underscored a high level of uncertainty in the machine learning quantified CO₂ effects.

Several limitations should be noted regarding GPP trend estimation and validation. First, the CFE-ML model may not fully capture the intricate mechanisms of plant physiological responses to CO₂. For example, eddy covariance towers, especially long-term sites, are typically located in homogeneous and undisturbed ecosystems, not representative of the full diversity of ecosystems globally. Thus, interactions between CO₂ and natural or human-induced disturbance, as well as many other stresses, are likely underrepresented in the models. Ultimately, the model's capacity to robustly quantify CO₂ fertilization is constrained by the scope and diversity of the eddy covariance data. Additionally, the use of spatially invariant CO₂ data may not fully represent the actual CO₂ variations that plants experience across different environments.

Secondly, CO₂ effects inferred by the CFE-ML models may be confounded by other factors that correlate with CO₂ over time. Industrialization-induced nitrogen deposition could synergistically boost GPP alongside CO₂ (O'Sullivan et al., 2019). Technological and management improvements in

agriculture that contribute to a global enhancement of crop photosynthesis (Zeng et al., 2014), might also be indirectly reflected in the model estimates. Moreover, interactions with the other input features that exhibit long-term trends, such as those induced by non-biological factors (e.g. sensor orbital drifts), also affect the CO₂ effects inference. Additionally, other factors that could lead to long-term GPP trends (e.g. forest aging, disturbances) might also be underrepresented in our models.

Finally, direct validation of GPP trends is limited, particularly in tropical regions, constrained by the availability of long-term records. Detecting and evaluating trends is challenging and typically requires long monitoring records (e.g. over 10 to 15 years), since long-term changes, such as those induced by CO₂, are very small relative to large interannual variations. Evaluating aggregated GPP trends across multiple sites presents an alternative approach; however, there were still insufficient sites in tropical and evergreen broadleaf forest areas to robustly validate our estimates for those ecosystems (Figure 5). Partly due to data limitations, uncertainties in GPP estimated from bootstrapped samples are very high in tropical areas (Figure 12). Thus, trend estimates in these areas should be interpreted in the context of associated uncertainties and limitations.

Our results also suggested that variations in the estimated GPP long-term trends from different products ~~were~~are largely related to the representation of CO₂ fertilization. Products that ~~did~~do not consider the direct CO₂ effect, including our Baseline models, FLUXSAT, FLUXCOM, and MODIS, showed minimal long-term changes in tropical GPP, while the CEDAR CFE-ML and CFE-Hybrid models demonstrated significant GPP increases aligning with predictions from the terrestrial biosphere models (Anav et al., 2015). FLUXCOM-ERA5, not accounting for dynamics changes in vegetation structures and CO₂, ~~did~~does not capture either the direct or indirect CO₂ fertilization resulting in a slight negative GPP trend attributable to shifted climate patterns. Notably, rEC-LUE exhibited contrasting trends before and after circa 2000, primarily attributed to changes in vapor pressure deficit, PAR, and LAI, while the direct CO₂ fertilization effect remained consistent (Zheng et al., 2020). CEDAR CFE-ML and CFE-Hybrid models align well with two process-based models forced with remote sensing data which consider direct CO₂ effects (BESS and BEPS). Nevertheless, considerable differences ~~between~~between CEDAR-GPP ~~and~~and other remote sensing products that include direct CO₂ effects (rEC-LUE₇ and PML V2) ~~as well as between our CFE-ML and CFE-Hybrid products,~~ warrant more in-depth investigations into long-term GPP responses to changes in atmospheric CO₂ and climate patterns.

Lastly, quantifications of GPP trends and their causes remain highly uncertain from site to global scales. Trend detection is often complicated by data noises and interannual variabilities, thus

requiring long-term records which are limited in certain areas, biomes, and environmental conditions, such as tropics, polar regions, wetlands, as well as ecosystems with regular or anthropogenic disturbances (Baldocchi et al., 2018; Zhan et al., 2022). Moreover, isolating the effect of CO₂ is challenging, as it is confounded by other factors, such as forest regrowth, land cover change, and disturbances, which also significantly impacts long-term GPP variations. To this end, continued efforts in expanding ecosystem flux measurements and standardizing data processing present new opportunities to assess ecosystem productivity responses to changing climate conditions (Delwiche et al., 2024; Pastorello et al., 2020). Future research could also leverage novel machine learning techniques, such as knowledge-guided machine learning (Liu et al., 2024) and hybrid modeling that combines process-based and machine learning approaches (Kraft et al., 2022; Reichstein et al., 2019).

5. Data availability and usage note

The CEDAR-GPP product, comprising ten GPP datasets, can be accessed at <https://zenodo.org/doi/10.5281/zenodo.8212706> (Kang et al., 2024). These datasets were generated at a spatial resolution of 0.05° and monthly time steps. Each dataset includes an ensemble mean GPP (“GPP_mean”) and an ensemble standard deviation (“GPP_std”). Data is formatted in netCDF with the following naming convention: “CEDAR-GPP_<version>_<model setup>_<YYYYMM>.nc”.

The CEDAR GPP product offers GPP estimates derived from ten different models. Models are characterized by 1) temporal coverage, 2) configuration of CO₂ fertilization, and 3) GPP partitioning approach (Table 2). We provide a structured approach to selecting the most appropriate dataset for research or applications.

1) Study period considerations: the Short-Term (ST) setup is ideal for studies focusing on periods after 2000. These models are constructed using a broader range of explanatory predictors, offering higher precision and smaller random errors. The Long-Term (LT) datasets shall be used for research assessing GPP dynamics over a longer time period (before 2001). It is important to note that trends from the ST and LT datasets are not directly comparable, as they were derived from different satellite remote sensing data.

2) CO₂ Fertilization Effect (CFE) configurations: the CFE-Hybrid and CFE-ML setups are preferable when assessing temporal GPP dynamics, especially long-term trends. The CFE-Hybrid setup includes a hypothetical trend from the direct CO₂ effect, while CFE-ML is purely data-driven and does not make any specific assumption about the sensitivity of photosynthesis to CO₂. Averaging

the CFE-Hybrid and CFE-ML estimates is acceptable, with the difference between them reflecting the uncertainty surrounding the direct CO₂ effect. Note that the Baseline setup should not be used to study long-term GPP dynamics, especially those induced by elevated CO₂. The Baseline setup may be useful to compare with other remote sensing-derived GPP datasets that do not consider the direct CO₂ effect. Differences between these setups regarding mean GPP spatial patterns, seasonal and interannual variations are considered to be minor.

3) GPP partitioning methods: We recommend using the mean value derived from both the “NT” (Nighttime) and “DT” (Daytime). The difference between these two provides insight into the uncertainties arising from the partitioning approaches used in GPP estimation from eddy covariance measurements.

Finally, like other upscaled or remote sensing-based GPP datasets, CEDAR-GPP should not be regarded as “observations” but rather as model estimates informed by remote sensing and ground-based data. The extent of assumptions or structural constraints varies across such datasets. CEDAR-GPP, particularly in its CFE-Baseline and CFE-ML configurations, is entirely data-driven and incorporates no explicit assumptions regarding the biological and environmental processes underlying photosynthesis, apart from the generic assumptions inherent in machine learning models. Consequently, the usage and interpretation of this dataset should be carefully framed within the context of the input eddy covariance and environmental data as well as their limitations.

6. Code availability

The code for upscaling and generating global GPP datasets can be accessed at <https://doi.org/10.5281/zenodo.8400968>.

7. Conclusions

We present the CEDAR-GPP product generated by upscaling global eddy covariance measurements with machine learning and a broad range of satellite and climate variables. CEDAR-GPP comprises four long-term datasets from 1982 to 2020 and six short-term datasets from 2001 to 2020. These datasets encompass three configurations regarding the incorporation of direct CO₂ fertilization effects and two partitioning approaches to derive GPP from eddy covariance data. The machine learning models of CEDAR-GPP demonstrated high capability in predicting monthly GPP,

its seasonal cycles, and spatial variability within the global eddy covariance sites, with cross-validated R^2 between 0.56 to 0.79. Short-term model setups consistently outperformed long-term models due to considerably more and higher-quality information from multi-source satellite observations.

CEDAR-GPP advances satellite-based GPP estimations, as the first upscaled dataset that considered the direct biochemical effects of elevated atmospheric CO_2 on photosynthesis, which is responsible for an increasing land carbon sink over the past decades. We showed that incorporating this effect in our CFE-ML and CFE-Hybrid models substantially improved the estimation of GPP trends at eddy covariance sites. Global patterns of long-term GPP trends in the CFE-ML setups showed a strong temperature dependency consistent with biophysical theories. However, trend estimation and validation remain particularly challenging in data-scarce regions, such as the tropics, emphasizing the need for enhanced data availability and methodological advancements. Beyond trends, global spatial and temporal GPP patterns from CEDAR generally aligned with other satellite-based GPP datasets.

In conclusion, CEDAR-GPP, informed by global eddy covariance measurements and a broad range of multi-source remote sensing observations and climatic variables, offered a comprehensive representation of global GPP spatial and temporal dynamics over the past four decades. The different CO_2 fertilization ~~configures-configurations~~ integrated in CEDAR-GPP offer new opportunities for understanding global ecosystem photosynthesis's response to increases in atmospheric CO_2 along different pathways over space and time. CEDAR-GPP is expected to serve as a valuable tool for benchmarking process-based modeling and constraining the global carbon cycle.

Site ID	IGBP	Data Range	Citation
AR-SLu	MF	2010 - 2011	(Garcia et al., 2016)
AR-Vir	ENF	2010 - 2012	(Posse et al., 2016)
AT-Neu	GRA	2002 - 2012	(Wohlfahrt et al., 2016)
AU-Ade	SAV	2010 - 2014	(Beringer and Hutley, 2016a)
AU-ASM	WSA	2007 - 2009	(Cleverly et al., 2016)
AU-Cpr	SAV	2010 - 2014	(Meyer et al., 2016)
AU-Cum	EBF	2012 - 2014	(Pendall et al., 2016)
AU-DaP	GRA	2007 - 2013	(Beringer and Hutley, 2016b)
AU-DaS	SAV	2008 - 2014	(Beringer and Hutley, 2016g)
AU-Dry	SAV	2008 - 2014	(Beringer and Hutley, 2016c)
AU-Emr	GRA	2011 - 2013	(Schroder et al., 2016)
AU-Fog	WET	2006 - 2008	(Beringer and Hutley, 2016d)
AU-Gin	WSA	2011 - 2014	(Macfarlane et al., 2016)
AU-How	WSA	2001 - 2014	(Beringer and Hutley, 2016e)
AU-RDF	WSA	2011 - 2013	(Beringer and Hutley, 2016f)
AU-Rig	GRA	2011 - 2014	(Beringer et al., 2016a)
AU-Tum	EBF	2001 - 2014	(Woodgate et al., 2016)
AU-Wac	EBF	2005 - 2008	(Beringer et al., 2016b)
AU-Whr	EBF	2011 - 2014	(Beringer et al., 2016c)
AU-Wom	EBF	2010 - 2014	(Arndt et al., 2016)
AU-Ync	GRA	2012 - 2014	(Beringer and Walker, 2016)
BE-Bra	MF	2001 - 2020	(Warm Winter 2020 Team, 2022)
BE-Dor	GRA	2011 - 2020	(Warm Winter 2020 Team, 2022)
BE-Lon	CRO	2004 - 2020	(Warm Winter 2020 Team, 2022)
BE-Maa	CSH	2016 - 2020	(Warm Winter 2020 Team, 2022)
BE-Vie	MF	2001 - 2020	(Warm Winter 2020 Team, 2022)
BR-Sa1	EBF	2002 - 2011	(Saleska, 2016)
BR-Sa3	EBF	2001 - 2004	(Goulden, 2016a)
CA-Ca1	ENF	2001 - 2002	(Black, 2023a)
CA-Ca2	ENF	2001 - 2010	(Black, 2023b)
CA-Ca3	ENF	2001 - 2010	(Black, 2018)
CA-Cbo	DBF	2001 - 2003	(Staebler, 2022)
CA-Gro	MF	2003 - 2014	(McCaughey, 2022)
CA-Man	ENF	2001 - 2008	(Amiro, 2016a)
CA-NS1	ENF	2002 - 2005	(Goulden, 2022a)
CA-NS2	ENF	2001 - 2005	(Goulden, 2022b)
CA-NS3	ENF	2001 - 2005	(Goulden, 2022c)
CA-NS4	ENF	2002 - 2005	(Goulden, 2016b)
CA-NS5	ENF	2001 - 2005	(Goulden, 2022d)
CA-NS6	OSH	2001 - 2005	(Goulden, 2022e)
CA-NS7	OSH	2002 - 2005	(Goulden, 2016c)
CA-Oas	DBF	2001 - 2010	(Black, 2016a)
CA-Obs	ENF	2001 - 2010	(Black, 2016b)
CA-Qc2	MF	2008 - 2010	(Margolis, 2018)
CA-Qfo	ENF	2003 - 2010	(Margolis, 2023)
CA-SF1	ENF	2003 - 2006	(Amiro, 2016b)
CA-SF2	ENF	2003 - 2005	(Amiro, 2023)
CA-SF3	OSH	2003 - 2006	(Amiro, 2016c)

CA-SJ2	ENF	2003 - 2007	(Barr and Black, 2018)
CA-TP1	ENF	2003 - 2014	(Arain, 2016b)
CA-TP2	ENF	2003 - 2007	(Arain, 2016c)
CA-TP3	ENF	2003 - 2014	(Arain, 2016d)
CA-TP4	ENF	2003 - 2017	(Arain, 2016a)
CA-TPD	DBF	2012 - 2014	(Arain, 2016e)
CA-WP1	WET	2003 - 2009	(Flanagan, 2018a)
CA-WP2	WET	2004 - 2006	(Flanagan, 2018b)
CA-WP3	WET	2004 - 2006	(Flanagan, 2018c)
CG-Tch	SAV	2006 - 2009	(Nouvellon, 2016)
CH-Aws	GRA	2006 - 2020	(Warm Winter 2020 Team, 2022)
CH-Cha	GRA	2005 - 2020	(Warm Winter 2020 Team, 2022)
CH-Dav	ENF	2001 - 2020	(Warm Winter 2020 Team, 2022)
CH-Fru	GRA	2005 - 2020	(Warm Winter 2020 Team, 2022)
CH-Lae	MF	2004 - 2020	(Warm Winter 2020 Team, 2022)
CH-Oe1	GRA	2002 - 2008	(Ammann, 2016)
CH-Oe2	CRO	2004 - 2020	(Warm Winter 2020 Team, 2022)
CN-Cha	MF	2003 - 2005	(Zhang and Han, 2016)
CN-Cng	GRA	2007 - 2010	(Dong, 2016)
CN-Din	EBF	2003 - 2005	(Zhou and Yan, 2016)
CN-Du2	GRA	2007 - 2008	(Chen, 2016c)
CN-Ha2	WET	2003 - 2005	(Li, 2016)
CN-HaM	GRA	2002 - 2004	(Tang et al., 2016)
CN-Qia	ENF	2003 - 2005	(Wang and Fu, 2016)
CN-Sw2	GRA	2011 - 2012	(Shao, 2016)
CZ-BK1	ENF	2004 - 2020	(Warm Winter 2020 Team, 2022)
CZ-BK2	GRA	2006 - 2012	(Sigut et al., 2016)
CZ-KrP	CRO	2014 - 2020	(Warm Winter 2020 Team, 2022)
CZ-Lnz	DBF	2015 - 2020	(Warm Winter 2020 Team, 2022)
CZ-RAJ	ENF	2012 - 2020	(Warm Winter 2020 Team, 2022)
CZ-Stn	DBF	2010 - 2020	(Warm Winter 2020 Team, 2022)
CZ-wet	WET	2006 - 2020	(Warm Winter 2020 Team, 2022)
DE-Akm	WET	2009 - 2020	(Warm Winter 2020 Team, 2022)
DE-Geb	CRO	2001 - 2020	(Warm Winter 2020 Team, 2022)
DE-Gri	GRA	2004 - 2020	(Warm Winter 2020 Team, 2022)
DE-Hai	DBF	2001 - 2020	(Warm Winter 2020 Team, 2022)
DE-HoH	DBF	2015 - 2020	(Warm Winter 2020 Team, 2022)
DE-Hte	WET	2009 - 2018	(Drought 2018 Team, 2020)
DE-Hzd	DBF	2010 - 2020	(Warm Winter 2020 Team, 2022)
DE-Kli	CRO	2004 - 2020	(Warm Winter 2020 Team, 2022)
DE-Lkb	ENF	2009 - 2013	(Lindauer et al., 2016)
DE-Lnf	DBF	2002 - 2012	(Knobl et al., 2016)
DE-Obe	ENF	2008 - 2020	(Warm Winter 2020 Team, 2022)
DE-RuR	GRA	2011 - 2020	(Warm Winter 2020 Team, 2022)
DE-RuS	CRO	2011 - 2020	(Warm Winter 2020 Team, 2022)
DE-RuW	ENF	2012 - 2020	(Warm Winter 2020 Team, 2022)
DE-Seh	CRO	2007 - 2010	(Schneider and Schmidt, 2016)
DE-SfN	WET	2012 - 2014	(Klatt et al., 2016)
DE-Spw	WET	2010 - 2014	(Bernhofer et al., 2016)
DE-Tha	ENF	2001 - 2020	(Warm Winter 2020 Team, 2022)
DK-Eng	GRA	2005 - 2007	(Pilegaard and Ibrom, 2016)
DK-Sor	DBF	2001 - 2020	(Warm Winter 2020 Team, 2022)

ES-Abr	WSA	2015 - 2020	(Warm Winter 2020 Team, 2022)
ES-Agu	OSH	2006 - 2019	(Warm Winter 2020 Team, 2022)
ES-Amo	OSH	2007 - 2012	(Poveda et al., 2016)
ES-LgS	OSH	2005 - 2020	(Reverter et al., 2016)
ES-LJu	WSA	2014 - 2020	(Warm Winter 2020 Team, 2022)
ES-LM1	WSA	2014 - 2020	(Warm Winter 2020 Team, 2022)
ES-LM2	OSH	2007 - 2009	(Warm Winter 2020 Team, 2022)
FI-Hyy	ENF	2001 - 2020	(Warm Winter 2020 Team, 2022)
FI-Jok	CRO	2001 - 2003	(Lohila et al., 2016)
FI-Ken	ENF	2018 - 2020	(Warm Winter 2020 Team, 2022)
FI-Let	ENF	2009 - 2020	(Warm Winter 2020 Team, 2022)
FI-Lom	WET	2007 - 2009	(Aurela et al., 2016a)
FI-Qvd	CRO	2018 - 2020	(Warm Winter 2020 Team, 2022)
FI-Sii	GRA	2016 - 2020	(Warm Winter 2020 Team, 2022)
FI-Sod	ENF	2001 - 2014	(Aurela et al., 2016b)
FI-Var	ENF	2016 - 2020	(Warm Winter 2020 Team, 2022)
FR-Aur	CRO	2005 - 2020	(Warm Winter 2020 Team, 2022)
FR-Bil	ENF	2014 - 2020	(Warm Winter 2020 Team, 2022)
FR-FBn	MF	2008 - 2020	(Warm Winter 2020 Team, 2022)
FR-Fon	DBF	2005 - 2020	(Warm Winter 2020 Team, 2022)
FR-Gri	CRO	2004 - 2020	(Warm Winter 2020 Team, 2022)
FR-Hes	DBF	2014 - 2020	(Warm Winter 2020 Team, 2022)
FR-Lam	ENF	2001 - 2008	(Warm Winter 2020 Team, 2022)
FR-LBr	WET	2017 - 2020	(Berbigier et al., 2016)
FR-LGt	CRO	2005 - 2020	(Warm Winter 2020 Team, 2022)
FR-Pue	EBF	2001 - 2014	(Ourcival et al., 2016)
FR-Tou	GRA	2018 - 2020	(Warm Winter 2020 Team, 2022)
GF-Guy	EBF	2015 - 2015	(Warm Winter 2020 Team, 2022)
GH-Ank	EBF	2011 - 2014	(Valentini et al., 2016a)
GL-NuF	WET	2008 - 2014	(Hansen, 2016)
GL-ZaF	WET	2009 - 2011	(Lund et al., 2016a)
GL-ZaH	GRA	2001 - 2014	(Lund et al., 2016b)
IL-Yat	ENF	2001 - 2020	(Warm Winter 2020 Team, 2022)
IT-CA1	DBF	2011 - 2014	(Sabbatini et al., 2016a)
IT-CA2	CRO	2011 - 2014	(Sabbatini et al., 2016b)
IT-CA3	DBF	2011 - 2014	(Sabbatini et al., 2016c)
IT-Col	DBF	2001 - 2014	(Matteucci, 2016)
IT-Cp2	EBF	2012 - 2020	(Warm Winter 2020 Team, 2022)
IT-Cpz	EBF	2001 - 2008	(Valentini et al., 2016b)
IT-La2	ENF	2001 - 2002	(Cescatti et al., 2016)
IT-Lav	ENF	2003 - 2020	(Warm Winter 2020 Team, 2022)
IT-Lsn	OSH	2016 - 2020	(Warm Winter 2020 Team, 2022)
IT-MBo	GRA	2003 - 2020	(Warm Winter 2020 Team, 2022)
IT-Noe	CSH	2004 - 2014	(Spano et al., 2016)
IT-PT1	DBF	2002 - 2004	(Manca and Goded, 2016)
IT-Ren	ENF	2001 - 2020	(Warm Winter 2020 Team, 2022)
IT-Ro1	DBF	2001 - 2008	(Valentini et al., 2016c)
IT-Ro2	DBF	2002 - 2012	(Papale et al., 2016)
IT-SR2	ENF	2013 - 2020	(Warm Winter 2020 Team, 2022)
IT-SRo	ENF	2001 - 2012	(Gruening et al., 2016)
IT-Tor	GRA	2008 - 2020	(Warm Winter 2020 Team, 2022)
JP-MBF	DBF	2004 - 2005	(Kotani, 2016a)

JP-SMF	MF	2002 - 2006	(Kotani, 2016b)
MY-PSO	EBF	2003 - 2009	(Kosugi and Takanashi, 2016)
NL-Hor	GRA	2004 - 2011	(Dolman et al., 2016a)
NL-Loo	ENF	2001 - 2018	(Drought 2018 Team, 2020)
PA-SPn	DBF	2007 - 2009	(Wolf et al., 2016)
RU-Che	WET	2002 - 2005	(Merbold et al., 2016)
RU-Cok	OSH	2003 - 2013	(Dolman et al., 2016b)
RU-Fy2	ENF	2015 - 2020	(Warm Winter 2020 Team, 2022)
RU-Fyo	ENF	2001 - 2020	(Warm Winter 2020 Team, 2022)
RU-Ha1	GRA	2002 - 2004	(Belelli et al., 2016)
SD-Dem	SAV	2007 - 2009	(Ardö et al., 2016)
SE-Deg	WET	2001 - 2020	(Warm Winter 2020 Team, 2022)
SE-Htm	ENF	2015 - 2020	(Warm Winter 2020 Team, 2022)
SE-Lnn	CRO	2014 - 2018	(Drought 2018 Team, 2020)
SE-Nor	ENF	2014 - 2020	(Warm Winter 2020 Team, 2022)
SE-Ros	ENF	2014 - 2020	(Warm Winter 2020 Team, 2022)
SE-Svb	ENF	2014 - 2020	(Warm Winter 2020 Team, 2022)
SJ-Adv	WET	2013 - 2014	(Christensen, 2016)
SN-Dhr	SAV	2010 - 2013	(Tageson et al., 2016)
US-ARM	CRO	2004 - 2018	(Biraud et al., 2022)
US-Atq	WET	2003 - 2008	(Zona and Oechel, 2016a)
US-Bar	DBF	2005 - 2017	(Richardson and Hollinger, 2023)
US-Blo	ENF	2001 - 2007	(Goldstein, 2016)
US-Cop	CRO	2011 - 2013	(Bowling, 2016)
US-CRT	GRA	2001 - 2007	(Chen and Chu, 2023)
US-Dk1	GRA	2004 - 2008	(Oishi et al., 2016a)
US-Dk2	DBF	2004 - 2008	(Oishi et al., 2016b)
US-Dk3	ENF	2004 - 2008	(Oishi et al., 2016c)
US-Fmf	WSA	2005 - 2008	(Dore and Kolb, 2023a)
US-FR2	ENF	2005 - 2010	(Litvak, 2016)
US-Fuf	ENF	2005 - 2010	(Dore and Kolb, 2023b)
US-GBT	ENF	2001 - 2003	(Massman, 2016a)
US-GLE	ENF	2005 - 2014	(Massman, 2016b)
US-Goo	GRA	2002 - 2006	(Meyers, 2016)
US-Ha1	DBF	2001 - 2012	(Munger, 2016)
US-Ho1	ENF	2012 - 2018	(Hollinger, 2016)
US-Ivo	WET	2004 - 2007	(Zona and Oechel, 2016b)
US-KFS	GRA	2009 - 2017	(Brunsell, 2022)
US-KS2	CSH	2003 - 2006	(Drake and Hinkle, 2016)
US-Los	WET	2001 - 2014	(Desai, 2016a)
US-Me2	DBF	2001 - 2017	(Law, 2022)
US-Me3	ENF	2003 - 2017	(Law, 2016a)
US-Me5	ENF	2004 - 2009	(Law, 2016b)
US-Me6	ENF	2001 - 2002	(Law, 2016c)
US-MMS	ENF	2010 - 2014	(Novick and Phillips, 2022)
US-Mpj	OSH	2008 - 2017	(Litvak, 2021)
US-Myb	WET	2011 - 2014	(Sturtevant et al., 2016)
US-Ne1	ENF	2001 - 2014	(Suyker, 2016a)
US-Ne2	CRO	2001 - 2013	(Suyker, 2016b)
US-Ne3	CRO	2001 - 2013	(Suyker, 2016c)
US-NR1	CRO	2001 - 2013	(Blanken et al., 2016)
US-Oho	DBF	2004 - 2013	(Chen et al., 2023)

US-PFa	MF	2001 - 2014	(Desai, 2016b)
US-Prr	ENF	2010 - 2016	(Iwahana et al., 2016)
US-Rls	CSH	2014 - 2017	(Flerchinger, 2023)
US-Rms	CSH	2014 - 2017	(Flerchinger, 2022a)
US-Ro1	CRO	2004 - 2016	(Baker et al., 2022)
US-Rws	OSH	2014 - 2017	(Flerchinger, 2022b)
US-Seg	MF	2008 - 2014	(Litvak, 2023a)
US-Ses	WSA	2004 - 2014	(Litvak, 2023b)
US-SRC	GRA	2007 - 2017	(Kurc, 2016)
US-SRM	OSH	2007 - 2017	(Scott, 2016a)
US-Sta	OSH	2005 - 2009	(Ewers and Pendall, 2016)
US-Syv	MF	2001 - 2014	(Desai, 2016c)
US-Ton	WSA	2001 - 2014	(Baldocchi and Ma, 2016)
US-Tw1	WET	2011 - 2017	(Valach et al., 2021)
US-Tw4	WET	2014 - 2017	(Eichelmann et al., 2023)
US-Twt	CRO	2009 - 2014	(Baldocchi, 2016)
US-Uaf	DBF	2007 - 2017	(Ueyama et al., 2018)
US-UMB	DBF	2008 - 2017	(Gough et al., 2023)
US-UMd	ENF	2003 - 2017	(Gough et al., 2022)
US-Var	GRA	2001 - 2014	(Baldocchi et al., 2016)
US-Vcm	ENF	2008 - 2017	(Litvak, 2023c)
US-Vcp	ENF	2007 - 2017	(Litvak, 2023d)
US-WCr	DBF	2001 - 2014	(Desai, 2016d)
US-Whs	WET	2011 - 2013	(Scott, 2016b)
US-Wi3	OSH	2007 - 2014	(Chen, 2016a)
US-Wi4	DBF	2002 - 2004	(Chen, 2016b)
US-Wjs	ENF	2002 - 2005	(Litvak, 2022)
US-WPT	SAV	2007 - 2017	(Chen and Chu, 2016)
ZM-Mon	DBF	2007 - 2009	(Kutsch et al., 2016)

963

964

Appendix B: CO₂ sensitivity function of Light Use Efficiency

In the CFE-Hybrid model, the direct CO₂ fertilization effect was prescribed onto machine learning estimated GPP at a reference CO₂ level using a theoretical CO₂ sensitivity function of LUE. The sensitivity function, which describes the fractional change in LUE due to CO₂ relative to the reference period, is described below.

The Light Use Efficiency (LUE) model (Monteith, 1972) of GPP states that,

$$GPP = APAR \times LUE = PAR \times fAPAR \times LUE \quad (A1)$$

where PAR is the photosynthetic active radiation, $fAPAR$ is the fraction of PAR that plant canopy has absorbed, and $APAR$ is the absorbed PAR . Eco-evolutionary theory, specifically the optimal coordination hypothesis, predicts that the electron-transport-limited (light-limited) (A_j) and Rubisco-limited (A_c) rates of photosynthesis converge on the time scale of physiological acclimation, which is in the order of a few weeks (Harrison et al., 2021; Haxeltine and Prentice, 1996; Wang et al., 2017). Thus, at a monthly time scale, we assume that

$$A = A_c = A_j \quad (A2)$$

where A is the gross photosynthetic rate, here equivalent to GPP.

In the following, we derive our sensitivity function based on A_j , which has a smaller response to CO₂ than A_c , thus providing conservative estimates of the direct CO₂ fertilization effect (Walker et al., 2021). According to the Farquhar, von Caemmerer and Berry (FvCB) model (Farquhar et al., 1980),

$$A_j = \varphi_0 I \frac{c_i - \Gamma^*}{c_i + 2\Gamma^*} \quad (A3)$$

where φ_0 is the intrinsic quantum efficiency of photosynthesis, I is the absorbed PAR ($I = APAR$), c_i is the leaf-internal partial pressure of CO₂, and Γ^* is the photorespiratory compensation point that depends on temperature:

$$\Gamma^* = r_{25} e^{\frac{\Delta H(T-298.15)}{298.15RT}} \quad (A4)$$

where $r_{25} = 4.22 \text{ Pa}$ is the photorespiratory point at 25 °C, ΔH is the activation energy ($37.83 \cdot 10^3 \text{ J mol}^{-1}$), T is the air temperature in Kelvin, and R is the molar gas constant ($8.314 \text{ J mol}^{-1} \text{ K}^{-1}$). We denote atmospheric CO₂ concentration as c_a , and χ is the ratio of leaf internal and external CO₂, so

$$c_i = \chi c_a \quad (A5)$$

Combining (A1), (A3), (A5), and assuming (A2), LUE can be written as,

$$LUE = \varphi_0 \frac{c_i - \Gamma^*}{c_i + 2\Gamma^*} = \varphi_0 \frac{\chi c_a - \Gamma^*}{\chi c_a + 2\Gamma^*} \quad (A6)$$

We can therefore show that under constant absorbed light (I or $APAR$), the sensitivity of GPP to CO_2 is proportional to that of LUE,

$$\frac{\partial GPP}{\partial c_a} = \frac{\partial \varphi_0 I \frac{\chi c_a - \Gamma^*}{\chi c_a + 2\Gamma^*}}{\partial c_a} = I \frac{\partial LUE}{\partial c_a} \quad (A7)$$

Thus from (A7), we can express the actual GPP at the time t and a CO_2 level c_a^t as the product of a reference GPP with a CO_2 level c_a^0 and the ratio between actual and reference LUE (A8-9). We denote the actual GPP as time t as $GPP_{c_a=c_a^t}^t$, and the reference GPP at time t as $GPP_{c_a=c_a^0}^t$.

$$\frac{GPP_{c_a=c_a^t}^t}{GPP_{c_a=c_a^0}^t} = \frac{LUE_{c_a=c_a^t}^t}{LUE_{c_a=c_a^0}^t} = \frac{\frac{\chi c_a^t - \Gamma^*}{\chi c_a^t + 2\Gamma^*}}{\frac{\chi c_a^0 - \Gamma^*}{\chi c_a^0 + 2\Gamma^*}} = \frac{\phi_{CO2}^t}{\phi_{CO2}^0} \quad (A8)$$

$$GPP_{c_a=c_a^t}^t = GPP_{c_a=c_a^0}^t \times \frac{\phi_{CO2}^t}{\phi_{CO2}^0} \quad (A9)$$

The reference GPP represents the GPP value at time t if the CO_2 were at the level of a reference level, while all other factors, such as PAR , $fAPAR$, temperature, and other environmental controls remain unchanged. Here the CO_2 impacts on LUE depend on atmospheric CO_2 (c_a), χ , and air temperature. We fixed χ to the global long-term average value 0.7 typical to C3 plants (Prentice et al., 2014; Wang et al., 2017). We further tested a dynamic model that quantified χ as a function of air temperature and vapor pressure deficit following an eco-evolutionary theory across global flux sites (Keenan et al., 2023). The estimated χ had a mean and median of 0.7 and a standard deviation of 0.04 (Figure S2144a). Differences in the direct CO_2 effect between the dynamic and fixed χ approaches were minimal, with an R^2 of 0.99 and a slope of 0.99 from a least squares linear regression line (Figure S2144b). GPP trends across flux towers were also highly consistent between the two approaches, with a difference less than 0.1 $gC\ m^{-2}\ year^{-2}$ (Figure S2144b, c). Since these results indicated that χ is relatively stable, we used the fixed χ approach to produce the CEDAR-GPP dataset.

In the CFE-Hybrid model, we estimated the reference GPP by fixing the CO_2 at the level of the year 2001 while keeping all other variables dynamic in the CFE-ML model. Then the actual GPP can be estimated following (A9). Fixing CO_2 values to the 2001 level, the start year of eddy covariance data used in model training, essentially removed the effects of CO_2 inferred by the CFE-ML model.

Supplement

The supplement related to this article is available online.

Author contributions

T. K. and Y. K. conceptualized the study. Y. K. performed the formal analysis and generated the final product. Y. K., T. K., M. B., and M. G. contributed to the development and investigation of the research. Y. K., M. G., and X. L. contributed to data curation and processing. Y. K. prepared the manuscript with contributions from all co-authors. T. K. supervised the project.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgments

We are grateful to Dr. Youngryel Ryu for providing the BESS_Rad dataset and Dr. Martin Jung for sharing the FLUXCOM-RS006 dataset. We also thank Dr. Muyi Li, Dr. Zaichun Zhu, and Dr. Sen Cao for sharing early versions of the PKU GIMMS NDVI4g and LAI4g datasets with us. We extend our gratitude to four reviewers for their constructive feedback for improving this paper.

Financial support

This research was supported by the U.S. Department of Energy Office of Science Early Career Research Program award #DE-SC0021023 and a NASA Award 80NSSC21K1705. YK acknowledges additional support from a NASA award 80NSSC24K1562. TFK acknowledges ~~additional-further~~ support from the LEMONTREE (Land Ecosystem Models based On New Theory, obseRvations and ExperiMents) project, funded through the generosity of Eric and Wendy Schmidt by recommendation of the Schmidt Futures programme, support from the RUBISCO SFA, which is sponsored by the Regional and Global Model Analysis (RGMA) Program in the Climate and Environmental Sciences Division (CESD) of the Office of Biological and Environmental Research (BER) in the U.S. Department of Energy Office of Science, and NASA awards ~~#~~80NSSC20K1801 and 80NSSC25K7327. MB acknowledges additional support from the U.S. Department of Agriculture NIFA award #2023-67012-40086.

References

- Amiro, B.: FLUXNET2015 CA-Man Manitoba - Northern Old Black Spruce (former BOREAS Northern Study Area), <https://doi.org/10.18140/FLX/1440035>, 2016a.
- Amiro, B.: FLUXNET2015 CA-SF1 Saskatchewan - Western Boreal, forest burned in 1977, <https://doi.org/10.18140/FLX/1440046>, 2016b.
- Amiro, B.: FLUXNET2015 CA-SF3 Saskatchewan - Western Boreal, forest burned in 1998, <https://doi.org/10.18140/FLX/1440048>, 2016c.
- Amiro, B.: AmeriFlux FLUXNET-1F CA-SF2 Saskatchewan - Western Boreal, forest burned in 1989, <https://doi.org/10.17190/AMF/2006961>, 2023.
- Ammann, C.: FLUXNET2015 CH-Oe1 Oensingen grassland, <https://doi.org/10.18140/FLX/1440135>, 2016.
- Anav, A., Friedlingstein, P., Beer, C., Ciais, P., Harper, A., Jones, C., Murray-Tortarolo, G., Papale, D., Parazoo, N. C., Peylin, P., Piao, S., Sitch, S., Viovy, N., Wiltshire, A., and Zhao, M.: Spatiotemporal patterns of terrestrial gross primary production: A review, *Reviews of Geophysics*, 1–34, <https://doi.org/10.1002/2015RG000483>, 2015.
- Apley, D. W. and Zhu, J.: Visualizing the Effects of Predictor Variables in Black Box Supervised Learning Models, *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 82, 1059–1086, <https://doi.org/10.1111/rssb.12377>, 2020.
- Arain, M. A.: AmeriFlux AmeriFlux CA-TP4 Ontario - Turkey Point 1939 Plantation White Pine, <https://doi.org/10.17190/AMF/1246012>, 2016a.
- Arain, M. A.: FLUXNET2015 CA-TP1 Ontario - Turkey Point 2002 Plantation White Pine, <https://doi.org/10.18140/FLX/1440050>, 2016b.
- Arain, M. A.: FLUXNET2015 CA-TP2 Ontario - Turkey Point 1989 Plantation White Pine, <https://doi.org/10.18140/FLX/1440051>, 2016c.
- Arain, M. A.: FLUXNET2015 CA-TP3 Ontario - Turkey Point 1974 Plantation White Pine, <https://doi.org/10.18140/FLX/1440052>, 2016d.
- Arain, M. A.: FLUXNET2015 CA-TPD Ontario - Turkey Point Mature Deciduous, <https://doi.org/10.18140/FLX/1440112>, 2016e.
- Ardö, J., El Tahir, B. A., and ElKhidir, H. A. M.: FLUXNET2015 SD-Dem Demokeya, <https://doi.org/10.18140/FLX/1440186>, 2016.
- Arndt, S., Hinko-Najera, N., Griebel, A., Beringer, J., and Livesley, S. J.: FLUXNET2015 AU-Wom Wombat, <https://doi.org/10.18140/FLX/1440207>, 2016.
- Aurela, M., Lohila, A., Tuovinen, J.-P., Hatakka, J., Rainne, J., Mäkelä, T., and Lauria, T.: FLUXNET2015 FI-Lom Lompolojankka, <https://doi.org/10.18140/FLX/1440228>, 2016a.

1079 Aurela, M., Tuovinen, J.-P., Hatakka, J., Lohila, A., Mäkelä, T., Rainne, J., and Lauria, T.:
 1080 FLUXNET2015 FI-Sod Sodankyla, <https://doi.org/10.18140/FLX/1440160>, 2016b.

1081 Badgley, G., Anderegg, L. D. L., Berry, J. A., and Field, C. B.: Terrestrial gross primary production:
 1082 Using NIRV to scale from site to globe, *Global Change Biology*, 25, 3731–3740,
 1083 <https://doi.org/10.1111/gcb.14729>, 2019.

1084 Baker, J., Griffis, T., and Griffis, T.: AmeriFlux FLUXNET-1F US-Ro1 Rosemount- G21,
 1085 <https://doi.org/10.17190/AMF/1881588>, 2022.

1086 Baldocchi, D.: FLUXNET2015 US-Twt Twitchell Island, <https://doi.org/10.18140/FLX/1440106>,
 1087 2016.

1088 Baldocchi, D. and Ma, S.: FLUXNET2015 US-Ton Tonzi Ranch,
 1089 <https://doi.org/10.18140/FLX/1440092>, 2016.

1090 Baldocchi, D., Ma, S., and Xu, L.: FLUXNET2015 US-Var Vaira Ranch- Ione,
 1091 <https://doi.org/10.18140/FLX/1440094>, 2016.

1092 Baldocchi, D., Chu, H., and Reichstein, M.: Inter-annual variability of net and gross ecosystem carbon
 1093 fluxes: A review, *Agricultural and Forest Meteorology*, 249, 520–533,
 1094 <https://doi.org/10.1016/j.agrformet.2017.05.015>, 2018.

1095 Baldocchi, D. D.: How eddy covariance flux measurements have contributed to our understanding of
 1096 Global Change Biology, *Global Change Biology*, 26, 242–260, <https://doi.org/10.1111/gcb.14807>,
 1097 2020.

1098 Baniecki, H., Kretowicz, W., Piątysek, P., Wiśniewski, J., and Biecek, P.: dalex: Responsible Machine
 1099 Learning with Interactive Explainability and Fairness in Python, *Journal of Machine Learning*
 1100 *Research*, 22, 1–7, 2021.

1101 Barr, A. and Black, A. T.: AmeriFlux AmeriFlux CA-SJ2 Saskatchewan - Western Boreal, Jack Pine
 1102 forest harvested in 2002, <https://doi.org/10.17190/AMF/1436321>, 2018.

1103 Beck, H. E., Zimmermann, N. E., McVicar, T. R., Vergopolan, N., Berg, A., and Wood, E. F.: Present
 1104 and future köppen-geiger climate classification maps at 1-km resolution, *Scientific Data*, 5, 1–12,
 1105 <https://doi.org/10.1038/sdata.2018.214>, 2018.

1106 Beer, C., Reichstein, M., Tomelleri, E., Ciais, P., Jung, M., Carvalhais, N., Rödenbeck, C., Arain, M.
 1107 A., Baldocchi, D., Bonan, G. B., Bondeau, A., Cescatti, A., Lasslop, G., Lindroth, A., Lomas, M.,
 1108 Luyssaert, S., Margolis, H., Oleson, K. W., Rouspard, O., Veenendaal, E., Viovy, N., Williams, C.,
 1109 Woodward, F. I., and Papale, D.: Terrestrial gross carbon dioxide uptake: Global distribution and
 1110 covariation with climate, *Science*, 329, 834–838, <https://doi.org/10.1126/science.1184984>, 2010.

1111 Belelli, L., Papale, D., and Valentini, R.: FLUXNET2015 RU-Ha1 Hakasia steppe,
 1112 <https://doi.org/10.18140/FLX/1440184>, 2016.

1113 Berbigier, P., Loustau, D., Bonnefond, J. M., Bosc, A., and Trichet, P.: FLUXNET2015 FR-LBr Le
 1114 Bray, <https://doi.org/10.18140/FLX/1440163>, 2016.

1115 Berdugo, M., Gaitán, J. J., Delgado-Baquerizo, M., Crowther, T. W., and Dakos, V.: Prevalence and
 1116 drivers of abrupt vegetation shifts in global drylands, *Proceedings of the National Academy of*
 1117 *Sciences*, 119, e2123393119, <https://doi.org/10.1073/pnas.2123393119>, 2022.

1118 Beringer, J. and Hutley, L.: FLUXNET2015 AU-Ade Adelaide River,
 1119 <https://doi.org/10.18140/FLX/1440193>, 2016a.

1120 Beringer, J. and Hutley, L.: FLUXNET2015 AU-DaP Daly River Savanna,
 1121 <https://doi.org/10.18140/FLX/1440123>, 2016b.

1122 Beringer, J. and Hutley, L.: FLUXNET2015 AU-Dry Dry River,
 1123 <https://doi.org/10.18140/FLX/1440197>, 2016c.

1124 Beringer, J. and Hutley, L.: FLUXNET2015 AU-Fog Fogg Dam,
 1125 <https://doi.org/10.18140/FLX/1440124>, 2016d.

1126 Beringer, J. and Hutley, L.: FLUXNET2015 AU-How Howard Springs,
 1127 <https://doi.org/10.18140/FLX/1440125>, 2016e.

1128 Beringer, J. and Hutley, L.: FLUXNET2015 AU-RDF Red Dirt Melon Farm, Northern Territory,
 1129 <https://doi.org/10.18140/FLX/1440201>, 2016f.

1130 Beringer, J. and Hutley, P. L.: FLUXNET2015 AU-DaS Daly River Cleared,
 1131 <https://doi.org/10.18140/FLX/1440122>, 2016g.

1132 Beringer, J. and Walker, J.: FLUXNET2015 AU-Ync Jaxa, <https://doi.org/10.18140/FLX/1440208>,
 1133 2016.

1134 Beringer, J., Cunningham, S., Baker, P., Cavagnaro, T., MacNally, R., Thompson, R., and McHugh, I.:
 1135 FLUXNET2015 AU-Rig Riggs Creek, <https://doi.org/10.18140/FLX/1440202>, 2016a.

1136 Beringer, J., Hutley, L., McGuire, D., U, P., and McHugh, I.: FLUXNET2015 AU-Wac Wallaby Creek,
 1137 <https://doi.org/10.18140/FLX/1440127>, 2016b.

1138 Beringer, J., Cunningham, S., Baker, P., Cavagnaro, T., MacNally, R., Thompson, R., and McHugh, I.:
 1139 FLUXNET2015 AU-Whr Whroo, <https://doi.org/10.18140/FLX/1440206>, 2016c.

1140 Bernhofer, C., Grünwald, T., Moderow, U., Hehn, M., Eichelmann, U., Prasse, H., and Postel, U.:
 1141 FLUXNET2015 DE-Spw Spreewald, <https://doi.org/10.18140/FLX/1440220>, 2016.

1142 Besnard, S., Carvalhais, N., Altaf Arain, M., Black, A., Brede, B., Buchmann, N., Chen, J., Clevers, J.
 1143 G. P. W., Dutrieux, L. P., Gans, F., Herold, M., Jung, M., Kosugi, Y., Knohl, A., Law, B. E., Paul-
 1144 Limoges, E., Lohila, A., Merbold, L., Rouspard, O., Valentini, R., Wolf, S., Zhang, X., and Reichstein,
 1145 M.: Memory effects of climate and vegetation affecting net ecosystem CO₂ fluxes in global forests,
 1146 *PLoS ONE*, 14, 1–22, <https://doi.org/10.1371/journal.pone.0211510>, 2019.

1147 Biraud, S., Fischer, M., Chan, S., and Torn, M.: AmeriFlux FLUXNET-1F US-ARM ARM Southern
 1148 Great Plains site- Lamont, <https://doi.org/10.17190/AMF/1854366>, 2022.

1149 Black, T. A.: FLUXNET2015 CA-Oas Saskatchewan - Western Boreal, Mature Aspen,
1150 <https://doi.org/10.18140/FLX/1440043>, 2016a.

1151 Black, T. A.: FLUXNET2015 CA-Obs Saskatchewan - Western Boreal, Mature Black Spruce,
1152 <https://doi.org/10.18140/FLX/1440044>, 2016b.

1153 Black, T. A.: AmeriFlux AmeriFlux CA-Ca3 British Columbia - Pole sapling Douglas-fir stand,
1154 <https://doi.org/10.17190/AMF/1480302>, 2018.

1155 Black, T. A.: AmeriFlux FLUXNET-1F CA-Ca1 British Columbia - 1949 Douglas-fir stand,
1156 <https://doi.org/10.17190/AMF/2007163>, 2023a.

1157 Black, T. A.: AmeriFlux FLUXNET-1F CA-Ca2 British Columbia - Clearcut Douglas-fir stand
1158 (harvested winter 1999/2000), <https://doi.org/10.17190/AMF/2007164>, 2023b.

1159 Blanken, P. D., Monson, R. K., Burns, S. P., Bowling, D. R., and Turnipseed, A. A.: FLUXNET2015
1160 US-NR1 Niwot Ridge Forest (LTER NWT1), <https://doi.org/10.18140/FLX/1440087>, 2016.

1161 Bloomfield, K. J., Stocker, B. D., Keenan, T. F., and Prentice, I. C.: Environmental controls on the
1162 light use efficiency of terrestrial gross primary production, *Global Change Biology*, 29, 1037–1053,
1163 <https://doi.org/10.1111/gcb.16511>, 2023.

1164 Bowling, D.: FLUXNET2015 US-Cop Corral Pocket, <https://doi.org/10.18140/FLX/1440100>,
1165 2016.

1166 Brunsell, N.: AmeriFlux FLUXNET-1F US-KFS Kansas Field Station,
1167 <https://doi.org/10.17190/AMF/1881585>, 2022.

1168 Campbell, J. E., Berry, J. A., Seibt, U., Smith, S. J., Montzka, S. A., Launois, T., Belviso, S., Bopp, L.,
1169 and Laine, M.: Large historical growth in global terrestrial gross primary production, *Nature*, 544, 84–
1170 87, <https://doi.org/10.1038/nature22030>, 2017.

1171 Camps-Valls, G., Campos-Taberner, M., Moreno-Martínez, Á., Walther, S., Duveiller, G., Cescatti, A.,
1172 Mahecha, M. D., Muñoz-Marí, J., García-Haro, F. J., Guanter, L., Jung, M., Gamon, J. A., Reichstein,
1173 M., and Running, S. W.: A unified vegetation index for quantifying the terrestrial biosphere, *Science*
1174 *Advances*, 7, eabc7447, <https://doi.org/10.1126/sciadv.abc7447>, 2021.

1175 Cao, S., Li, M., Zhu, Z., Zha, J., Zhao, W., Duanmu, Z., Chen, J., Zheng, Y., and Chen, Y.:
1176 Spatiotemporally consistent global dataset of the GIMMS Leaf Area Index (GIMMS LAI4g) from
1177 1982 to 2020, *Earth System Science Data Discussions*, 1–31, <https://doi.org/10.5194/essd-2023-68>,
1178 2023.

1179 Cescatti, A., Marcolla, B., Zorer, R., and Gianelle, D.: FLUXNET2015 IT-La2 Lavarone2,
1180 <https://doi.org/10.18140/FLX/1440235>, 2016.

1181 Chen, C., Park, T., Wang, X., Piao, S., Xu, B., Chaturvedi, R. K., Fuchs, R., Brovkin, V., Ciais, P.,
1182 Fensholt, R., Tømmervik, H., Bala, G., Zhu, Z., Nemani, R. R., and Myneni, R. B.: China and India
1183 lead in greening of the world through land-use management, *Nature Sustainability*, 2, 122–129,
1184 <https://doi.org/10.1038/s41893-019-0220-7>, 2019.

1185 Chen, C., Riley, W. J., Prentice, I. C., and Keenan, T. F.: CO₂ fertilization of terrestrial photosynthesis
 1186 inferred from site to global scales, *Proceedings of the National Academy of Sciences*, 119, 1–8,
 1187 <https://doi.org/10.1073/pnas.2115627119/>, 2022.

1188 Chen, J.: FLUXNET2015 US-Wi3 Mature hardwood (MHW),
 1189 <https://doi.org/10.18140/FLX/1440057>, 2016a.

1190 Chen, J.: FLUXNET2015 US-Wi4 Mature red pine (MRP), <https://doi.org/10.18140/FLX/1440058>,
 1191 2016b.

1192 Chen, J. and Chu, H.: FLUXNET2015 US-WPT Winous Point North Marsh,
 1193 <https://doi.org/10.18140/FLX/1440116>, 2016.

1194 Chen, J. and Chu, H.: AmeriFlux FLUXNET-1F US-CRT Curtice Walter-Berger cropland,
 1195 <https://doi.org/10.17190/AMF/2006974>, 2023.

1196 Chen, J., Chu, H., and Noormets, A.: AmeriFlux FLUXNET-1F US-Oho Oak Openings,
 1197 <https://doi.org/10.17190/AMF/2229385>, 2023.

1198 Chen, S.: FLUXNET2015 CN-Du2 Duolun_grassland (D01),
 1199 <https://doi.org/10.18140/FLX/1440140>, 2016c.

1200 Chen, T. and Guestrin, C.: XGBoost: a scalable tree boosting system, in: *Proceedings of the 22nd*
 1201 *ACM SIGKDD International Conference on Knowledge Discovery and Data Mining - KDD '16*,
 1202 785–794, <https://doi.org/10.1145/2939672.2939785>, 2016.

1203 Christensen, T.: FLUXNET2015 SJ-Adv Adventdalen, <https://doi.org/10.18140/FLX/1440241>,
 1204 2016.

1205 Chu, H., Luo, X., Ouyang, Z., Chan, W. S., Dengel, S., Biraud, S. C., Torn, M. S., Metzger, S., Kumar,
 1206 J., Arain, M. A., Arkebauer, T. J., Baldocchi, D., Bernacchi, C., Billesbach, D., Black, T. A., Blanken,
 1207 P. D., Bohrer, G., Bracho, R., Brown, S., Brunsell, N. A., Chen, J., Chen, X., Clark, K., Desai, A. R.,
 1208 Duman, T., Durden, D., Fares, S., Forbrich, I., Gamon, J. A., Gough, C. M., Griffis, T., Helbig, M.,
 1209 Hollinger, D., Humphreys, E., Ikawa, H., Iwata, H., Ju, Y., Knowles, J. F., Knox, S. H., Kobayashi,
 1210 H., Kolb, T., Law, B., Lee, X., Litvak, M., Liu, H., Munger, J. W., Noormets, A., Novick, K.,
 1211 Oberbauer, S. F., Oechel, W., Oikawa, P., Papuga, S. A., Pendall, E., Prajapati, P., Prueger, J., Quinton,
 1212 W. L., Richardson, A. D., Russell, E. S., Scott, R. L., Starr, G., Staebler, R., Stoy, P. C., Stuart-Haëntjens,
 1213 E., Sonnentag, O., Sullivan, R. C., Suyker, A., Ueyama, M., Vargas, R., Wood, J. D., and Zona, D.:
 1214 Representativeness of Eddy-Covariance flux footprints for areas surrounding AmeriFlux sites,
 1215 *Agricultural and Forest Meteorology*, 301–302, <https://doi.org/10.1016/j.agrformet.2021.108350>,
 1216 2021.

1217 Cleverly, J., Eamus, D., and Isaac, P.: FLUXNET2015 AU-ASM Alice Springs,
 1218 <https://doi.org/10.18140/FLX/1440194>, 2016.

1219 Dannenberg, M. P., Barnes, M. L., Smith, W. K., Johnston, M. R., Meerdink, S. K., Wang, X., Scott,
 1220 R. L., and Biederman, J. A.: Upscaling dryland carbon and water fluxes with artificial neural networks
 1221 of optical, thermal, and microwave satellite remote sensing, *Biogeosciences*, 20, 383–404,
 1222 <https://doi.org/10.5194/bg-20-383-2023>, 2023.

1223 De Kauwe, M. G., Keenan, T. F., Medlyn, B. E., Prentice, I. C., and Terrer, C.: Satellite based estimates
 1224 underestimate the effect of CO₂ fertilization on net primary productivity, *Nature Climate Change*, 6,
 1225 892–893, <https://doi.org/10.1038/nclimate3105>, 2016.

1226 Delwiche, K. B., Nelson, J., Kowalska, N., Moore, C. E., Shirkey, G., Tarin, T., Cleverly, J. R., and
 1227 Keenan, T. F.: Charting the Future of the FLUXNET Network, *Bulletin of the American*
 1228 *Meteorological Society*, 105, E466–E473, <https://doi.org/10.1175/BAMS-D-23-0316.1>, 2024.

1229 Desai, A.: FLUXNET2015 US-Los Lost Creek, <https://doi.org/10.18140/FLX/1440076>, 2016a.

1230 Desai, A.: FLUXNET2015 US-PFa Park Falls/WLEF, <https://doi.org/10.18140/FLX/1440089>,
 1231 2016b.

1232 Desai, A.: FLUXNET2015 US-Syv Sylvania Wilderness Area,
 1233 <https://doi.org/10.18140/FLX/1440091>, 2016c.

1234 Desai, A.: FLUXNET2015 US-WCr Willow Creek, <https://doi.org/10.18140/FLX/1440095>, 2016d.

1235 Dolman, H., Hendriks, D., Parmentier, F.-J., Marchesini, L. B., Dean, J., and van Huissteden, K.:
 1236 FLUXNET2015 NL-Hor Horstermeer, <https://doi.org/10.18140/FLX/1440177>, 2016a.

1237 Dolman, H., van der Molen, M., Parmentier, F.-J., Marchesini, L. B., Dean, J., van Huissteden, K., and
 1238 Maximov, T.: FLUXNET2015 RU-Cok Chokurdakh, <https://doi.org/10.18140/FLX/1440182>,
 1239 2016b.

1240 Dong, G.: FLUXNET2015 CN-Cng Changling, <https://doi.org/10.18140/FLX/1440209>, 2016.

1241 Dong, Y., Li, J., Jiao, Z., Liu, Q., Zhao, J., Xu, B., Zhang, H., Zhang, Z., Liu, C., Knyazikhin, Y., and
 1242 Myneni, R. B.: A Method for Retrieving Coarse-Resolution Leaf Area Index for Mixed Biomes Using
 1243 a Mixed-Pixel Correction Factor, *IEEE Transactions on Geoscience and Remote Sensing*, 61, 1–17,
 1244 <https://doi.org/10.1109/TGRS.2023.3235949>, 2023.

1245 Dore, S. and Kolb, T.: AmeriFlux FLUXNET-1F US-Fmf Flagstaff - Managed Forest,
 1246 <https://doi.org/10.17190/AMF/2007173>, 2023a.

1247 Dore, S. and Kolb, T.: AmeriFlux FLUXNET-1F US-Fuf Flagstaff - Unmanaged Forest,
 1248 <https://doi.org/10.17190/AMF/2007174>, 2023b.

1249 Dorigo, W., Wagner, W., Albergel, C., Albrecht, F., Balsamo, G., Brocca, L., Chung, D., Ertl, M.,
 1250 Forkel, M., Gruber, A., Haas, E., Hamer, P. D., Hirschi, M., Ikonen, J., de Jeu, R., Kidd, R., Lahoz,
 1251 W., Liu, Y. Y., Miralles, D., Mistelbauer, T., Nicolai-Shaw, N., Parinussa, R., Pratola, C., Reimer, C.,
 1252 van der Schalie, R., Seneviratne, S. I., Smolander, T., and Lecomte, P.: ESA CCI Soil Moisture for
 1253 improved Earth system understanding: State-of-the art and future directions, *Remote Sensing of*
 1254 *Environment*, 203, 185–215, <https://doi.org/10.1016/j.rse.2017.07.001>, 2017.

1255 Dorigo, W. A., Gruber, A., De Jeu, R. A. M., Wagner, W., Stacke, T., Loew, A., Albergel, C., Brocca,
 1256 L., Chung, D., Parinussa, R. M., and Kidd, R.: Evaluation of the ESA CCI soil moisture product using
 1257 ground-based observations, *Remote Sensing of Environment*, 162,
 1258 <https://doi.org/10.1016/j.rse.2014.07.023>, 2015.

1259 Drake, B. and Hinkle, R.: FLUXNET2015 US-KS2 Kennedy Space Center (scrub oak),
1260 <https://doi.org/10.18140/FLX/1440075>, 2016.

1261 Drought 2018 Team: Drought-2018 ecosystem eddy covariance flux product for 52 stations in
1262 FLUXNET-Archive format, <https://doi.org/10.18160/YVR0-4898>, 2020.

1263 Ehlers, I., Augusti, A., Betson, T. R., Nilsson, M. B., Marshall, J. D., and Schleucher, J.: Detecting
1264 long-term metabolic shifts using isotopomers: CO₂-driven suppression of photorespiration in C₃
1265 plants over the 20th century, *Proceedings of the National Academy of Sciences*, 112, 15585–15590,
1266 <https://doi.org/10.1073/pnas.1504493112>, 2015.

1267 Eichelmann, E., Shortt, R., Knox, S., Sanchez, C. R., Valach, A., Sturtevant, C., Szutu, D., Verfaillie,
1268 J., and Baldocchi, D.: AmeriFlux FLUXNET-1F US-Tw4 Twitchell East End Wetland,
1269 <https://doi.org/10.17190/AMF/2204881>, 2023.

1270 Ewers, B. and Pendall, E.: FLUXNET2015 US-Sta Saratoga,
1271 <https://doi.org/10.18140/FLX/1440115>, 2016.

1272 Fang, H., Baret, F., Plummer, S., and Schaepman-Strub, G.: An overview of global leaf area index
1273 (LAI): Methods, products, validation, and applications, *Reviews of Geophysics*, 2018RG000608,
1274 <https://doi.org/10.1029/2018RG000608>, 2019.

1275 Farquhar, G. D., von Caemmerer, S., and Berry, J. A.: A biochemical model of photosynthetic CO₂
1276 assimilation in leaves of C₃ species, *Planta*, 149, 78–90, <https://doi.org/10.1007/BF00386231>, 1980.

1277 Fernández-Martínez, M., Vicca, S., Janssens, I. A., Ciais, P., Obersteiner, M., Bartrons, M., Sardans, J.,
1278 Verger, A., Canadell, J. G., Chevallier, F., Wang, X., Bernhofer, C., Curtis, P. S., Gianelle, D.,
1279 Grünwald, T., Heinesch, B., Ibrom, A., Knohl, A., Laurila, T., Law, B. E., Limousin, J. M., Longdoz,
1280 B., Loustau, D., Mammarella, I., Matteucci, G., Monson, R. K., Montagnani, L., Moors, E. J., Munger,
1281 J. W., Papale, D., Piao, S. L., and Peñuelas, J.: Atmospheric deposition, CO₂, and change in the land
1282 carbon sink, *Sci Rep*, 7, 9632, <https://doi.org/10.1038/s41598-017-08755-8>, 2017.

1283 Flanagan, L. B.: AmeriFlux AmeriFlux CA-WP1 Alberta - Western Peatland - LaBiche River, Black
1284 Spruce/Larch Fen, <https://doi.org/10.17190/AMF/1436323>, 2018a.

1285 Flanagan, L. B.: AmeriFlux AmeriFlux CA-WP2 Alberta - Western Peatland - Poor Fen (Sphagnum
1286 moss), <https://doi.org/10.17190/AMF/1436324>, 2018b.

1287 Flanagan, L. B.: AmeriFlux AmeriFlux CA-WP3 Alberta - Western Peatland - Rich Fen (Carex),
1288 <https://doi.org/10.17190/AMF/1436325>, 2018c.

1289 Flerchinger, G.: AmeriFlux FLUXNET-1F US-Rms RCEW Mountain Big Sagebrush,
1290 <https://doi.org/10.17190/AMF/1881587>, 2022a.

1291 Flerchinger, G.: AmeriFlux FLUXNET-1F US-Rws Reynolds Creek Wyoming big sagebrush,
1292 <https://doi.org/10.17190/AMF/1881592>, 2022b.

1293 Flerchinger, G.: AmeriFlux FLUXNET-1F US-Rls RCEW Low Sagebrush,
1294 <https://doi.org/10.17190/AMF/2229387>, 2023.

1295 Friedl, M. and Sulla-Menashe, D.: MCD12Q1 MODIS/Terra+Aqua Land Cover Type Yearly L3
 1296 Global 500m SIN Grid V006 [Data set], NASA EOSDIS Land Processes DAAC,
 1297 <https://doi.org/10.5067/MODIS/MCD12Q1.006>, 2019.

1298 Friedlingstein, P., Meinshausen, M., Arora, V. K., Jones, C. D., Anav, A., Liddicoat, S. K., and Knutti,
 1299 R.: Uncertainties in CMIP5 climate projections due to carbon cycle feedbacks, *Journal of Climate*, 27,
 1300 511–526, <https://doi.org/10.1175/JCLI-D-12-00579.1>, 2014.

1301 Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Gregor, L., Hauck, J., Le Quéré, C.,
 1302 Luijkx, I. T., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J.
 1303 G., Ciais, P., Jackson, R. B., Alin, S. R., Alkama, R., Arneeth, A., Arora, V. K., Bates, N. R., Becker, M.,
 1304 Bellouin, N., Bittig, H. C., Bopp, L., Chevallier, F., Chini, L. P., Cronin, M., Evans, W., Falk, S., Feely,
 1305 R. A., Gasser, T., Gehlen, M., Gkritzalis, T., Gloege, L., Grassi, G., Gruber, N., Gürses, Ö., Harris, I.,
 1306 Hefner, M., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Jain, A. K., Jersild, A., Kadono, K.,
 1307 Kato, E., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Landschützer, P., Lefèvre,
 1308 N., Lindsay, K., Liu, J., Liu, Z., Marland, G., Mayot, N., McGrath, M. J., Metzl, N., Monacci, N. M.,
 1309 Munro, D. R., Nakaoka, S.-I., Niwa, Y., O’Brien, K., Ono, T., Palmer, P. I., Pan, N., Pierrot, D.,
 1310 Pocock, K., Poulter, B., Resplandy, L., Robertson, E., Rödenbeck, C., Rodriguez, C., Rosan, T. M.,
 1311 Schwinger, J., Séférian, R., Shutler, J. D., Skjelvan, I., Steinhoff, T., Sun, Q., Sutton, A. J., Sweeney, C.,
 1312 Takao, S., Tanhua, T., Tans, P. P., Tian, X., Tian, H., Tilbrook, B., Tsujino, H., Tubiello, F., van der
 1313 Werf, G. R., Walker, A. P., Wanninkhof, R., Whitehead, C., Willstrand Wranne, A., et al.: Global
 1314 Carbon Budget 2022, *Earth System Science Data*, 14, 4811–4900, 2023.

1315 Gaber, M., Kang, Y., Schurgers, G., and Keenan, T.: Using automated machine learning for the
 1316 upscaling of gross primary productivity, *Biogeosciences*, 21, 2447–2472, [https://doi.org/10.5194/bg-](https://doi.org/10.5194/bg-21-2447-2024)
 1317 21-2447-2024, 2024.

1318 Gampe, D., Zscheischler, J., Reichstein, M., Sullivan, M. O., Smith, W. K., Sitch, S., and Buermann,
 1319 W.: Increasing impact of warm droughts on northern ecosystem productivity over recent decades,
 1320 *Nature Climate Change*, <https://doi.org/10.1038/s41558-021-01112-8>, 2021.

1321 Gao, B. C.: NDWI - A normalized difference water index for remote sensing of vegetation liquid
 1322 water from space, *Remote Sensing of Environment*, 58, 257–266, [https://doi.org/10.1016/S0034-](https://doi.org/10.1016/S0034-4257(96)00067-3)
 1323 4257(96)00067-3, 1996.

1324 Garcia, A., Di Bella, C., Houspanossian, J., Magliano, P., Jobbágy, E., Posse, G., Fernández, R., and
 1325 Noretto, M.: FLUXNET2015 AR-SLu San Luis, 2016.

1326 Gitelson, A. A.: Remote estimation of leaf area index and green leaf biomass in maize canopies,
 1327 *Geophysical Research Letters*, 30, 1248, <https://doi.org/10.1029/2002GL016450>, 2003.

1328 Goldstein, A.: FLUXNET2015 US-Blo Blodgett Forest, <https://doi.org/10.18140/FLX/1440068>,
 1329 2016.

1330 Gough, C., Bohrer, G., and Curtis, P.: AmeriFlux FLUXNET-1F US-UMd UMBS Disturbance,
 1331 <https://doi.org/10.17190/AMF/1881597>, 2022.

1332 Gough, C., Bohrer, G., and Curtis, P.: AmeriFlux FLUXNET-1F US-UMB Univ. of Mich. Biological
 1333 Station, <https://doi.org/10.17190/AMF/2204882>, 2023.

1334 Goulden, M.: FLUXNET2015 BR-Sa3 Santarem-Km83-Logged Forest,
1335 <https://doi.org/10.18140/FLX/1440033>, 2016a.

1336 Goulden, M.: FLUXNET2015 CA-NS4 UCI-1964 burn site wet,
1337 <https://doi.org/10.18140/FLX/1440039>, 2016b.

1338 Goulden, M.: FLUXNET2015 CA-NS7 UCI-1998 burn site,
1339 <https://doi.org/10.18140/FLX/1440042>, 2016c.

1340 Goulden, M.: AmeriFlux FLUXNET-1F CA-NS1 UCI-1850 burn site,
1341 <https://doi.org/10.17190/AMF/1902824>, 2022a.

1342 Goulden, M.: AmeriFlux FLUXNET-1F CA-NS2 UCI-1930 burn site,
1343 <https://doi.org/10.17190/AMF/1902825>, 2022b.

1344 Goulden, M.: AmeriFlux FLUXNET-1F CA-NS3 UCI-1964 burn site,
1345 <https://doi.org/10.17190/AMF/1902826>, 2022c.

1346 Goulden, M.: AmeriFlux FLUXNET-1F CA-NS5 UCI-1981 burn site,
1347 <https://doi.org/10.17190/AMF/1902828>, 2022d.

1348 Goulden, M.: AmeriFlux FLUXNET-1F CA-NS6 UCI-1989 burn site,
1349 <https://doi.org/10.17190/AMF/1902829>, 2022e.

1350 Green, J. K., Ballantyne, A., Abramoff, R., Gentine, P., Makowski, D., and Ciais, P.: Surface
1351 temperatures reveal the patterns of vegetation water stress and their environmental drivers across the
1352 tropical Americas, *Global Change Biology*, 28, 2940–2955, <https://doi.org/10.1111/gcb.16139>, 2022.

1353 Gruber, A., Scanlon, T., Van Der Schalie, R., Wagner, W., and Dorigo, W.: Evolution of the ESA CCI
1354 Soil Moisture climate data records and their underlying merging methodology, *Earth System Science*
1355 *Data*, 11, 717–739, <https://doi.org/10.5194/essd-11-717-2019>, 2019.

1356 Gruening, C., Goded, I., Cescatti, A., Manca, G., and Seufert, G.: FLUXNET2015 IT-SRo San
1357 Rossore, <https://doi.org/10.18140/FLX/1440176>, 2016.

1358 Hansen, B. U.: FLUXNET2015 GL-NuF Nuuk Fen, <https://doi.org/10.18140/FLX/1440222>, 2016.

1359 Hao, D., Bisht, G., Huang, M., Ma, P.-L., Tesfa, T., Lee, W.-L., Gu, Y., and Leung, L. R.: Impacts of
1360 Sub-Grid Topographic Representations on Surface Energy Balance and Boundary Conditions in the
1361 E3SM Land Model: A Case Study in Sierra Nevada, *Journal of Advances in Modeling Earth Systems*,
1362 14, e2021MS002862, <https://doi.org/10.1029/2021MS002862>, 2022.

1363 Harrison, S. P., Cramer, W., Franklin, O., Prentice, I. C., Wang, H., Brännström, Å., de Boer, H.,
1364 Dieckmann, U., Joshi, J., Keenan, T. F., Lavergne, A., Manzoni, S., Mengoli, G., Morfopoulos, C.,
1365 Peñuelas, J., Pietsch, S., Rebel, K. T., Ryu, Y., Smith, N. G., Stocker, B. D., and Wright, I. J.: Eco-
1366 evolutionary optimality as a means to improve vegetation and land-surface models, *New Phytologist*,
1367 <https://doi.org/10.1111/nph.17558>, 2021.

1368 Haverd, V., Smith, B., Canadell, J. G., Cuntz, M., Mikaloff-Fletcher, S., Farquhar, G., Woodgate, W.,
1369 Briggs, P. R., and Trudinger, C. M.: Higher than expected CO₂ fertilization inferred from leaf to global
1370 observations, *Global Change Biology*, 26, 2390–2402, <https://doi.org/10.1111/gcb.14950>, 2020.

1371 Haxeltine, A. and Prentice, I. C.: A General Model for the Light-Use Efficiency of Primary Production,
1372 *Functional Ecology*, 10, 551–561, <https://doi.org/10.2307/2390165>, 1996.

1373 Hollinger, D.: AmeriFlux AmeriFlux US-Ho1 Howland Forest (main tower),
1374 <https://doi.org/10.17190/AMF/1246061>, 2016.

1375 van der Horst, S. V. J., Pitman, A. J., De Kauwe, M. G., Ukkola, A., Abramowitz, G., and Isaac, P.:
1376 How representative are FLUXNET measurements of surface fluxes during temperature extremes?,
1377 *Biogeosciences*, 16, 1829–1844, <https://doi.org/10.5194/bg-16-1829-2019>, 2019.

1378 Iwahana, G., Kobayashi, H., Ikawa, H., and Suzuki, R.: AmeriFlux AmeriFlux US-Prr Poker Flat
1379 Research Range Black Spruce Forest, <https://doi.org/10.17190/AMF/1246153>, 2016.

1380 Jiang, C., Ryu, Y., Fang, H., Myneni, R., Claverie, M., and Zhu, Z.: Inconsistencies of interannual
1381 variability and trends in long-term satellite leaf area index products, *Global Change Biology*, 23, 4133–
1382 4146, <https://doi.org/10.1111/gcb.13787>, 2017.

1383 Joiner, J. and Yoshida, Y.: Satellite-based reflectances capture large fraction of variability in global
1384 gross primary production (GPP) at weekly time scales, *Agricultural and Forest Meteorology*, 291,
1385 108092, <https://doi.org/10.1016/j.agrformet.2020.108092>, 2020.

1386 Jung, M., Reichstein, M., Margolis, H. A., Cescatti, A., Richardson, A. D., Arain, M. A., Arneth, A.,
1387 Bernhofer, C., Bonal, D., Chen, J., Gianelle, D., Gobron, N., Kiely, G., Kutsch, W., Lasslop, G., Law,
1388 B. E., Lindroth, A., Merbold, L., Montagnani, L., Moors, E. J., Papale, D., Sottocornola, M., Vaccari,
1389 F., and Williams, C.: Global patterns of land-atmosphere fluxes of carbon dioxide , latent heat , and
1390 sensible heat derived from eddy covariance , satellite , and meteorological observations, *Journal of*
1391 *Geophysical Research: Biogeosciences*, 116, 1–16, <https://doi.org/10.1029/2010JG001566>, 2011.

1392 Jung, M., Reichstein, M., Schwalm, C. R., Huntingford, C., Sitch, S., Ahlström, A., Arneth, A., Camps-
1393 Valls, G., Ciais, P., Friedlingstein, P., Gans, F., Ichii, K., Jain, A. K., Kato, E., Papale, D., Poulter, B.,
1394 Raduly, B., Rödenbeck, C., Tramontana, G., Viovy, N., Wang, Y. P., Weber, U., Zaehle, S., and Zeng,
1395 N.: Compensatory water effects link yearly global land CO₂ sink changes to temperature, *Nature*, 541,
1396 516–520, <https://doi.org/10.1038/nature20780>, 2017.

1397 Jung, M., Schwalm, C., Migliavacca, M., Walther, S., Camps-Valls, G., Koirala, S., Anthoni, P., Besnard,
1398 S., Bodesheim, P., Carvalhais, N., Chevallier, F., Gans, F., S Goll, D., Haverd, V., Köhler, P., Ichii, K.,
1399 K Jain, A., Liu, J., Lombardozzi, D., E M S Nabel, J., A Nelson, J., O’Sullivan, M., Pallandt, M., Papale,
1400 D., Peters, W., Pongratz, J., Rödenbeck, C., Sitch, S., Tramontana, G., Walker, A., Weber, U., and
1401 Reichstein, M.: Scaling carbon fluxes from eddy covariance sites to globe: Synthesis and evaluation of
1402 the FLUXCOM approach, *Biogeosciences*, 17, 1343–1365, [https://doi.org/10.5194/bg-17-1343-](https://doi.org/10.5194/bg-17-1343-2020)
1403 2020, 2020.

1404 Kang, Y., Ozdogan, M., Zhu, X., Ye, Z., Hain, C., and Anderson, M.: Comparative assessment of
1405 environmental variables and machine learning algorithms for maize yield prediction in the US Midwest,
1406 *Environmental Research Letters*, 15, <https://doi.org/10.1088/1748-9326/ab7df9>, 2020.

1407 Kang, Y., Bassiouni, M., Gaber, M., Lu, X., and Keenan, T.: CEDAR-GPP: A Spatiotemporally
1408 Upscaled Dataset of Gross Primary Productivity Incorporating CO₂ Fertilization (v1.0),
1409 <https://doi.org/10.5281/zenodo.8212706>, 2024.

1410 Keeling, R. F., Graven, H. D., Welp, L. R., Resplandy, L., Bi, J., Piper, S. C., Sun, Y., Bollenbacher,
1411 A., and Meijer, H. A. J.: Atmospheric evidence for a global secular increase in carbon isotopic
1412 discrimination of land photosynthesis, *Proceedings of the National Academy of Sciences of the United*
1413 *States of America*, 114, 10361–10366, <https://doi.org/10.1073/pnas.1619240114>, 2017.

1414 Keenan, T. F., Hollinger, D. Y., Bohrer, G., Dragoni, D., Munger, J. W., Schmid, H. P., and
1415 Richardson, A. D.: Increase in forest water-use efficiency as atmospheric carbon dioxide
1416 concentrations rise, *Nature*, 499, 324–327, <https://doi.org/10.1038/nature12291>, 2013.

1417 Keenan, T. F., Prentice, I. C., Canadell, J. G., Williams, C. A., Wang, H., Raupach, M., and Collatz, G.
1418 J.: Recent pause in the growth rate of atmospheric CO₂ due to enhanced terrestrial carbon uptake,
1419 *Nature Communications*, 7, 1–9, <https://doi.org/10.1038/ncomms13428>, 2016.

1420 Keenan, T. F., Migliavacca, M., Papale, D., Baldocchi, D., Reichstein, M., Torn, M., and Wutzler, T.:
1421 Widespread inhibition of daytime ecosystem respiration, *Nature Ecology and Evolution*, 3, 407–415,
1422 <https://doi.org/10.1038/s41559-019-0809-2>, 2019.

1423 Keenan, T. F., Luo, X., Stocker, B. D., De Kauwe, M. G., Medlyn, B. E., Prentice, I. C., Smith, N. G.,
1424 Terrer, C., Wang, H., Zhang, Y., and Zhou, S.: A constraint on historic growth in global
1425 photosynthesis due to rising CO₂, *Nat. Clim. Chang.*, 1–6, [https://doi.org/10.1038/s41558-023-](https://doi.org/10.1038/s41558-023-01867-2)
1426 [01867-2](https://doi.org/10.1038/s41558-023-01867-2), 2023.

1427 Klatt, J., Schmid, H. P., Mauder, M., and Steinbrecher, R.: FLUXNET2015 DE-SfN Schechenfilz
1428 Nord, <https://doi.org/10.18140/FLX/1440219>, 2016.

1429 Knohl, A., Tiedemann, F., Kolle, O., Schulze, E.-D., Anthoni, P., Kutsch, W., Herbst, M., and Siebicke,
1430 L.: FLUXNET2015 DE-Lnf Leinefelde, <https://doi.org/10.18140/FLX/1440150>, 2016.

1431 Kosugi, Y. and Takanashi, S.: FLUXNET2015 MY-PSO Pasoh Forest Reserve (PSO),
1432 <https://doi.org/10.18140/FLX/1440240>, 2016.

1433 Kotani, A.: FLUXNET2015 JP-MBF Moshiri Birch Forest Site,
1434 <https://doi.org/10.18140/FLX/1440238>, 2016a.

1435 Kotani, A.: FLUXNET2015 JP-SMF Seto Mixed Forest Site,
1436 <https://doi.org/10.18140/FLX/1440239>, 2016b.

1437 Kraft, B., Jung, M., Körner, M., Koirala, S., and Reichstein, M.: Towards hybrid modeling of the global
1438 hydrological cycle, *Hydrology and Earth System Sciences*, 26, 1579–1614,
1439 <https://doi.org/10.5194/hess-26-1579-2022>, 2022.

1440 Kurc, S.: FLUXNET2015 US-SRC Santa Rita Creosote, <https://doi.org/10.18140/FLX/1440098>,
1441 2016.

1442 Kutsch, W. L., Merbold, L., and Kolle, O.: FLUXNET2015 ZM-Mon Mongu,
1443 <https://doi.org/10.18140/FLX/1440189>, 2016.

1444 Law, B.: FLUXNET2015 US-Me3 Metolius-second young aged pine,
1445 <https://doi.org/10.18140/FLX/1440080>, 2016a.

1446 Law, B.: FLUXNET2015 US-Me5 Metolius-first young aged pine,
1447 <https://doi.org/10.18140/FLX/1440082>, 2016b.

1448 Law, B.: FLUXNET2015 US-Me6 Metolius Young Pine Burn,
1449 <https://doi.org/10.18140/FLX/1440099>, 2016c.

1450 Law, B.: AmeriFlux FLUXNET-1F US-Me2 Metolius mature ponderosa pine,
1451 <https://doi.org/10.17190/AMF/1854368>, 2022.

1452 Leng, J., Chen, J. M., Li, W., Luo, X., Xu, M., Liu, J., Wang, R., Rogers, C., Li, B., and Yan, Y.: Global
1453 datasets of hourly carbon and water fluxes simulated using a satellite-based process model with
1454 dynamic parameterizations, *Earth System Science Data*, 16, 1283–1300,
1455 <https://doi.org/10.5194/essd-16-1283-2024>, 2024.

1456 Li, B., Ryu, Y., Jiang, C., Dechant, B., Liu, J., Yan, Y., and Li, X.: BESSv2.0: A satellite-based and
1457 coupled-process model for quantifying long-term global land–atmosphere fluxes, *Remote Sensing of*
1458 *Environment*, 295, 113696, <https://doi.org/10.1016/j.rse.2023.113696>, 2023a.

1459 Li, M., Cao, S., and Zhu, Z.: Spatiotemporally consistent global dataset of the GIMMS Normalized
1460 Difference Vegetation Index (PKU GIMMS NDVI) from 1982 to 2020, *Earth System Science Data*
1461 *Discussions*, 1–31, <https://doi.org/10.5194/essd-2023-1>, 2023b.

1462 Li, Y.: FLUXNET2015 CN-Ha2 Haibei Shrubland, <https://doi.org/10.18140/FLX/1440211>, 2016.

1463 Lindauer, M., Steinbrecher, R., Wolpert, B., Mauder, M., and Schmid, H. P.: FLUXNET2015 DE-
1464 Lkb Lackenberg, <https://doi.org/10.18140/FLX/1440214>, 2016.

1465 Litvak, M.: AmeriFlux AmeriFlux US-FR2 Freeman Ranch- Mesquite Juniper,
1466 <https://doi.org/10.17190/AMF/1246054>, 2016.

1467 Litvak, M.: AmeriFlux FLUXNET-1F US-Mpj Mountainair Pinyon-Juniper Woodland,
1468 <https://doi.org/10.17190/AMF/1832161>, 2021.

1469 Litvak, M.: AmeriFlux FLUXNET-1F US-Wjs Willard Juniper Savannah,
1470 <https://doi.org/10.17190/AMF/1871146>, 2022.

1471 Litvak, M.: AmeriFlux FLUXNET-1F US-Seg Sevilleta grassland,
1472 <https://doi.org/10.17190/AMF/1984572>, 2023a.

1473 Litvak, M.: AmeriFlux FLUXNET-1F US-Ses Sevilleta shrubland,
1474 <https://doi.org/10.17190/AMF/1984573>, 2023b.

1475 Litvak, M.: AmeriFlux FLUXNET-1F US-Vcm Valles Caldera Mixed Conifer,
1476 <https://doi.org/10.17190/AMF/2229391>, 2023c.

1477 Litvak, M.: AmeriFlux FLUXNET-1F US-Vcp Valles Caldera Ponderosa Pine,
1478 <https://doi.org/10.17190/AMF/2229392>, 2023d.

1479 Liu, L., Zhou, W., Guan, K., Peng, B., Xu, S., Tang, J., Zhu, Q., Till, J., Jia, X., Jiang, C., Wang, S.,
1480 Qin, Z., Kong, H., Grant, R., Mezbahuddin, S., Kumar, V., and Jin, Z.: Knowledge-guided machine
1481 learning can improve carbon cycle quantification in agroecosystems, *Nat Commun*, 15, 357,
1482 <https://doi.org/10.1038/s41467-023-43860-5>, 2024.

1483 Liu, Y., Holtzman, N. M., and Konings, A. G.: Global ecosystem-scale plant hydraulic traits retrieved
1484 using model–data fusion, *Hydrology and Earth System Sciences*, 25, 2399–2417,
1485 <https://doi.org/10.5194/hess-25-2399-2021>, 2021.

1486 Lohila, A., Aurela, M., Tuovinen, J.-P., Hatakka, J., and Laurila, T.: FLUXNET2015 FI-Jok Jokioinen,
1487 <https://doi.org/10.18140/FLX/1440159>, 2016.

1488 Lund, M., Jackowicz-Korczyński, M., and Abermann, J.: FLUXNET2015 GL-ZaF Zackenberg Fen,
1489 <https://doi.org/10.18140/FLX/1440223>, 2016a.

1490 Lund, M., Jackowicz-Korczyński, M., and Abermann, J.: FLUXNET2015 GL-ZaH Zackenberg Heath,
1491 <https://doi.org/10.18140/FLX/1440224>, 2016b.

1492 Lundberg, S. M. and Lee, S.-I.: A Unified Approach to Interpreting Model Predictions, in: *Advances*
1493 *in Neural Information Processing Systems*, 2017.

1494 Luo, X., Zhou, H., Satriawan, T. W., Tian, J., Zhao, R., Keenan, T. F., Griffith, D. M., Sitch, S., Smith,
1495 N. G., and Still, C. J.: Mapping the global distribution of C4 vegetation using observations and
1496 optimality theory, *Nat Commun*, 15, 1219, <https://doi.org/10.1038/s41467-024-45606-3>, 2024.

1497 Ma, H. and Liang, S.: Development of the GLASS 250-m leaf area index product (version 6) from
1498 MODIS data using the bidirectional LSTM deep learning model, *Remote Sensing of Environment*,
1499 273, 112985, <https://doi.org/10.1016/j.rse.2022.112985>, 2022.

1500 Ma, H., Zeng, J., Chen, N., Zhang, X., Cosh, M. H., and Wang, W.: Satellite surface soil moisture from
1501 SMAP, SMOS, AMSR2 and ESA CCI: A comprehensive assessment using global ground-based
1502 observations, *Remote Sensing of Environment*, 231, 111215,
1503 <https://doi.org/10.1016/j.rse.2019.111215>, 2019.

1504 Ma, Y., Zhang, Z., Kang, Y., and Özdoğan, M.: Corn yield prediction and uncertainty analysis based
1505 on remotely sensed variables using a Bayesian neural network approach, *Remote Sensing of*
1506 *Environment*, 259, 112408, <https://doi.org/10.1016/j.rse.2021.112408>, 2021.

1507 Macfarlane, C., Lambert, P., Byrne, J., Johnstone, C., and Smart, N.: FLUXNET2015 AU-Gin Gingin,
1508 <https://doi.org/10.18140/FLX/1440199>, 2016.

1509 Manca, G. and Goded, I.: FLUXNET2015 IT-PT1 Parco Ticino forest,
1510 <https://doi.org/10.18140/FLX/1440172>, 2016.

1511 Margolis, H.: AmeriFlux CA-Qc2 Quebec - 1975 Harvested Black Spruce (HBS75),
1512 <https://doi.org/10.17190/AMF/1419514>, 2018.

1513 Margolis, H. A.: AmeriFlux FLUXNET-1F CA-Qfo Quebec - Eastern Boreal, Mature Black Spruce,
1514 <https://doi.org/10.17190/AMF/2006960>, 2023.

1515 Massman, B.: FLUXNET2015 US-GBT GLEES Brooklyn Tower,
1516 <https://doi.org/10.18140/FLX/1440118>, 2016a.

1517 Massman, B.: FLUXNET2015 US-GLE GLEES, <https://doi.org/10.18140/FLX/1440069>, 2016b.

1518 Matteucci, G.: FLUXNET2015 IT-Col Collelongo, <https://doi.org/10.18140/FLX/1440167>, 2016.

1519 McCaughey, H.: AmeriFlux FLUXNET-1F CA-Gro Ontario - Groundhog River, Boreal Mixedwood
1520 Forest, <https://doi.org/10.17190/AMF/1902823>, 2022.

1521 Merbold, L., Rebmann, C., and Corradi, C.: FLUXNET2015 RU-Che Cherski,
1522 <https://doi.org/10.18140/FLX/1440181>, 2016.

1523 Meyer, W., Cale, P., Koerber, G., Ewenz, C., and Sun, Q.: FLUXNET2015 AU-Cpr Calperum,
1524 <https://doi.org/10.18140/FLX/1440195>, 2016.

1525 Meyers, T.: FLUXNET2015 US-Goo Goodwin Creek, <https://doi.org/10.18140/FLX/1440070>,
1526 2016.

1527 Munger, J. W.: FLUXNET2015 US-Ha1 Harvard Forest EMS Tower (HFR1),
1528 <https://doi.org/10.18140/FLX/1440071>, 2016.

1529 Myneni, R., Knyazikhin, Y., and Park, T.: MCD15A3H MODIS/Terra+Aqua Leaf Area Index/FPAR
1530 4-day L4 Global 500m SIN Grid V006 [Data set], NASA EOSDIS Land Processes DAAC.,
1531 <https://doi.org/10.5067/MODIS/MCD15A3H.006>, 2015a.

1532 Myneni, R., Knyazikhin, Y., and Park, T.: MOD15A2H MODIS/Terra Leaf Area Index/FPAR 8-
1533 Day L4 Global 500m SIN Grid V006 [Data set], NASA EOSDIS Land Processes DAAC,
1534 <https://doi.org/10.5067/MODIS/MOD15A2H.006>, 2015b.

1535 Nelson, J. A., Walther, S., Gans, F., Kraft, B., Weber, U., Novick, K., Buchmann, N., Migliavacca, M.,
1536 Wohlfahrt, G., Šigut, L., Ibrom, A., Papale, D., Göckede, M., Duveiller, G., Knohl, A., Hörtnagl, L.,
1537 Scott, R. L., Zhang, W., Hamdi, Z. M., Reichstein, M., Aranda-Barranco, S., Ardö, J., Op de Beeck,
1538 M., Billesbach, D., Bowling, D., Bracho, R., Brümmer, C., Camps-Valls, G., Chen, S., Cleverly, J. R.,
1539 Desai, A., Dong, G., El-Madany, T. S., Euskirchen, E. S., Feigenwinter, I., Galvagno, M., Gerosa, G.,
1540 Gielen, B., Godec, I., Goslee, S., Gough, C. M., Heinesch, B., Ichii, K., Jackowicz-Korczynski, M. A.,
1541 Klosterhalfen, A., Knox, S., Kobayashi, H., Kohonen, K.-M., Korkiakoski, M., Mammarella, I., Mana,
1542 G., Marzuoli, R., Matamala, R., Metzger, S., Montagnani, L., Nicolini, G., O'Halloran, T., Ourcival, J.-
1543 M., Peichl, M., Pendall, E., Ruiz Reverter, B., Roland, M., Sabbatini, S., Sachs, T., Schmidt, M.,
1544 Schwalm, C. R., Shekhar, A., Silberstein, R., Silveira, M. L., Spano, D., Tagesson, T., Tramontana, G.,
1545 Trotta, C., Turco, F., Vesala, T., Vincke, C., Vitale, D., Vivoni, E. R., Wang, Y., Woodgate, W., Yezpe,
1546 E. A., Zhang, J., Zona, D., and Jung, M.: X-BASE: the first terrestrial carbon and water flux products

1547 from an extended data-driven scaling framework, FLUXCOM-X, EGU sphere, 1–51,
1548 <https://doi.org/10.5194/egusphere-2024-165>, 2024.

1549 Nouvellon, Y.: FLUXNET2015 CG-Tch Tchizalamou, <https://doi.org/10.18140/FLX/1440142>,
1550 2016.

1551 Novick, K. and Phillips, R.: AmeriFlux FLUXNET-1F US-MMS Morgan Monroe State Forest,
1552 <https://doi.org/10.17190/AMF/1854369>, 2022.

1553 Oishi, C., Novick, K., and Stoy, P.: AmeriFlux AmeriFlux US-Dk1 Duke Forest-open field,
1554 <https://doi.org/10.17190/AMF/1246046>, 2016a.

1555 Oishi, C., Novick, K., and Stoy, P.: AmeriFlux AmeriFlux US-Dk2 Duke Forest-hardwoods,
1556 <https://doi.org/10.17190/AMF/1246047>, 2016b.

1557 Oishi, C., Novick, K., and Stoy, P.: AmeriFlux AmeriFlux US-Dk3 Duke Forest - loblolly pine,
1558 <https://doi.org/10.17190/AMF/1246048>, 2016c.

1559 O’Sullivan, M., Spracklen, D. V., Batterman, S. A., Arnold, S. R., Gloor, M., and Buermann, W.: Have
1560 Synergies Between Nitrogen Deposition and Atmospheric CO₂ Driven the Recent Enhancement of
1561 the Terrestrial Carbon Sink?, *Global Biogeochemical Cycles*, 33, 163–180,
1562 <https://doi.org/10.1029/2018GB005922>, 2019.

1563 O’Sullivan, M., Smith, W. K., Sitch, S., Friedlingstein, P., Arora, V. K., Haverd, V., Jain, A. K., Kato,
1564 E., Kautz, M., Lombardozzi, D., Nabel, J. E. M. S., Tian, H., Vuichard, N., Wiltshire, A., Zhu, D., and
1565 Buermann, W.: Climate-Driven Variability and Trends in Plant Productivity Over Recent Decades
1566 Based on Three Global Products, *Global Biogeochemical Cycles*, 34,
1567 <https://doi.org/10.1029/2020GB006613>, 2020.

1568 Ourcival, J.-M., Piquemal, K., Joffre, R., and Jean-Marc, L.: FLUXNET2015 FR-Pue Puechabon,
1569 <https://doi.org/10.18140/FLX/1440164>, 2016.

1570 Ozflux, P.: FluxNet Data OzFlux: Australian and New Zealand Flux Research and Monitoring,
1571 [https://data.ozflux.org.au/portal/pub/viewColDetails.jsp?collection.id=1882723&collection.owner](https://data.ozflux.org.au/portal/pub/viewColDetails.jsp?collection.id=1882723&collection.owner.id=450&viewType=anonymous)
1572 [.id=450&viewType=anonymous](https://data.ozflux.org.au/portal/pub/viewColDetails.jsp?collection.id=1882723&collection.owner.id=450&viewType=anonymous), 2024.

1573 Papale, D., Tirone, G., Valentini, R., Arriga, N., Beilelli, L., Consalvo, C., Dore, S., Manca, G.,
1574 Mazzenga, F., Sabbatini, S., Stefani, P., Boschi, A., and Tomassucci, M.: FLUXNET2015 IT-Ro2
1575 Roccarespampani 2, <https://doi.org/10.18140/FLX/1440175>, 2016.

1576 Pastorello, G., Trotta, C., Canfora, E., Chu, H., Christianson, D., Cheah, Y. W., Poindexter, C., Chen,
1577 J., Elbashandy, A., Humphrey, M., Isaac, P., Polidori, D., Ribeca, A., van Ingen, C., Zhang, L., Amiro,
1578 B., Ammann, C., Arain, M. A., Ardö, J., Arkebauer, T., Arndt, S. K., Arriga, N., Aubinet, M., Aurela,
1579 M., Baldocchi, D., Barr, A., Beamesderfer, E., Marchesini, L. B., Bergeron, O., Beringer, J., Bernhofer,
1580 C., Berveiller, D., Billesbach, D., Black, T. A., Blanken, P. D., Bohrer, G., Boike, J., Bolstad, P. V.,
1581 Bonal, D., Bonnefond, J. M., Bowling, D. R., Bracho, R., Brodeur, J., Brümmer, C., Buchmann, N.,
1582 Burban, B., Burns, S. P., Buysse, P., Cale, P., Cavagna, M., Cellier, P., Chen, S., Chini, I., Christensen,
1583 T. R., Cleverly, J., Collalti, A., Consalvo, C., Cook, B. D., Cook, D., Coursolle, C., Cremonese, E.,
1584 Curtis, P. S., D’Andrea, E., da Rocha, H., Dai, X., Davis, K. J., De Cinti, B., de Grandcourt, A., De

1585 Ligne, A., De Oliveira, R. C., Delpierre, N., Desai, A. R., Di Bella, C. M., di Tommasi, P., Dolman,
1586 H., Domingo, F., Dong, G., Dore, S., Duce, P., Dufrêne, E., Dunn, A., Dušek, J., Eamus, D.,
1587 Eichelmann, U., ElKhidir, H. A. M., Eugster, W., Ewenz, C. M., Ewers, B., Famulari, D., Fares, S.,
1588 Feigenwinter, I., Feitz, A., Fensholt, R., Filippa, G., Fischer, M., Frank, J., Galvagno, M., Gharun, M.,
1589 Gianelle, D., et al.: The FLUXNET2015 dataset and the ONEFlux processing pipeline for eddy
1590 covariance data, *Scientific data*, 7, 225, <https://doi.org/10.1038/s41597-020-0534-3>, 2020.

1591 Pendall, E., Griebel, A., Barton, C., and Metzen, D.: FLUXNET2015 AU-Cum Cumberland Plains,
1592 <https://doi.org/10.18140/FLX/1440196>, 2016.

1593 Peñuelas, J., Ciais, P., Canadell, J. G., Janssens, I. A., Fernández-Martínez, M., Carnicer, J., Obersteiner,
1594 M., Piao, S., Vautard, R., and Sardans, J.: Shifting from a fertilization-dominated to a warming-
1595 dominated period, *Nature Ecology and Evolution*, 1, 1438–1445, [https://doi.org/10.1038/s41559-](https://doi.org/10.1038/s41559-017-0274-8)
1596 017-0274-8, 2017.

1597 Piao, S., Wang, X., Park, T., Chen, C., Lian, X., He, Y., Bjerke, J. W., Chen, A., Ciais, P., Tømmervik,
1598 H., Nemani, R. R., and Myneni, R. B.: Characteristics, drivers and feedbacks of global greening, *Nature*
1599 *Reviews Earth and Environment*, 1, 14–27, <https://doi.org/10.1038/s43017-019-0001-x>, 2020.

1600 Pilegaard, K. and Ibrom, A.: FLUXNET2015 DK-Eng Enghave,
1601 <https://doi.org/10.18140/FLX/1440153>, 2016.

1602 Posse, G., Lewczuk, N., Richter, K., and Cristiano, P.: FLUXNET2015 AR-Vir Virasoro, FluxNet;
1603 Instituto Nacional de Tecnología Agropecuaria, <https://doi.org/10.18140/FLX/1440192>, 2016.

1604 Poveda, F. D., Ballesteros, A. L., Cañete, E. P. S., Ortiz, P. S., Jiménez, M. R. M., Priego, O. P., and
1605 Kowalski, A. S.: FLUXNET2015 ES-Amo Amoladeras, <https://doi.org/10.18140/FLX/1440156>,
1606 2016.

1607 Prentice, I. C., Dong, N., Gleason, S. M., Maire, V., and Wright, I. J.: Balancing the costs of carbon
1608 gain and water transport: testing a new theoretical framework for plant functional ecology, *Ecology*
1609 *Letters*, 17, 82–91, <https://doi.org/10.1111/ele.12211>, 2014.

1610 Reich, P. B., Hobbie, S. E., and Lee, T. D.: Plant growth enhancement by elevated CO₂ eliminated by
1611 joint water and nitrogen limitation, *Nature Geoscience*, 7, 920–924,
1612 <https://doi.org/10.1038/ngeo2284>, 2014.

1613 Reichstein, M., Camps-Valls, G., Stevens, B., Jung, M., Denzler, J., Carvalhais, N., and Prabhat: Deep
1614 learning and process understanding for data-driven Earth system science, *Nature*, 566, 195–204,
1615 <https://doi.org/10.1038/s41586-019-0912-1>, 2019.

1616 Reverter, B. R., Perez-Cañete, E. S., and Kowalski, A. S.: FLUXNET2015 ES-LgS Laguna Seca,
1617 <https://doi.org/10.18140/FLX/1440225>, 2016.

1618 Richardson, A. and Hollinger, D.: AmeriFlux FLUXNET-1F US-Bar Bartlett Experimental Forest,
1619 <https://doi.org/10.17190/AMF/2006969>, 2023.

1620 Ruehr, S., Keenan, T. F., Williams, C., Zhou, Y., Lu, X., Bastos, A., Canadell, J. G., Prentice, I. C.,
 1621 Sitch, S., and Terrer, C.: Evidence and attribution of the enhanced land carbon sink, *Nat Rev Earth*
 1622 *Environ*, 1–17, <https://doi.org/10.1038/s43017-023-00456-3>, 2023.

1623 Running, S., Mu, Q., and Zhao, M.: MOD17A2H MODIS/Terra Gross Primary Productivity 8-Day
 1624 L4 Global 500m SIN Grid V006, <https://doi.org/10.5067/MODIS/MOD17A2H.006>, 2015.

1625 Ryu, Y., Jiang, C., Kobayashi, H., and Detto, M.: MODIS-derived global land products of shortwave
 1626 radiation and diffuse and total photosynthetically active radiation at 5 km resolution from 2000,
 1627 *Remote Sensing of Environment*, 204, 812–825, <https://doi.org/10.1016/j.rse.2017.09.021>, 2018.

1628 Ryu, Y., Berry, J. A., and Baldocchi, D. D.: What is global photosynthesis? History, uncertainties and
 1629 opportunities, *Remote Sensing of Environment*, 223, 95–114,
 1630 <https://doi.org/10.1016/j.rse.2019.01.016>, 2019.

1631 Sabater, J. M.: ERA5-Land monthly averaged data from 1981 to present., Copernicus Climate Change
 1632 Service (C3S) Climate Data Store (CDS), <https://doi.org/doi:10.24381/cds.68d2bb30>, 2019.

1633 Sabbatini, S., Arriga, N., Papale, D., Boschi, A., and Tomassucci, M.: FLUXNET2015 IT-CA1 Castel
 1634 d’Asso1, <https://doi.org/10.18140/FLX/1440230>, 2016a.

1635 Sabbatini, S., Arriga, N., Gioli, B., Papale, D., Boschi, A., and Tomassucci, M.: FLUXNET2015 IT-
 1636 CA2 Castel d’Asso2, <https://doi.org/10.18140/FLX/1440231>, 2016b.

1637 Sabbatini, S., Arriga, N., Matteucci, G., Papale, D., Boschi, A., and Tomassucci, M.: FLUXNET2015
 1638 IT-CA3 Castel d’Asso 3, <https://doi.org/10.18140/FLX/1440232>, 2016c.

1639 Saleska, S.: FLUXNET2015 BR-Sa1 Santarem-Km67-Primary Forest,
 1640 <https://doi.org/10.18140/FLX/1440032>, 2016.

1641 Schaaf, C. and Wang, Z.: MCD43C4 MODIS/Terra+Aqua BRDF/Albedo Nadir BRDF-Adjusted
 1642 Ref Daily L3 Global 0.05Deg CMG V006 [Data set], NASA EOSDIS Land Processes DAAC.,
 1643 <https://doi.org/10.5067/MODIS/MCD43C4.006>, 2015.

1644 Schneider, K. and Schmidt, M.: FLUXNET2015 DE-Seh Selhausen,
 1645 <https://doi.org/10.18140/FLX/1440217>, 2016.

1646 Schroder, I., Zegelin, S., Palu, T., and Feitz, A.: FLUXNET2015 AU-Emr Emerald,
 1647 <https://doi.org/10.18140/FLX/1440198>, 2016.

1648 Schwalm, C. R., Anderegg, W. R. L., Michalak, A. M., Fisher, J. B., Biondi, F., Koch, G., Litvak, M.,
 1649 Ogle, K., Shaw, J. D., Wolf, A., Huntzinger, D. N., Schaefer, K., Cook, R., Wei, Y., Fang, Y., Hayes,
 1650 D., Huang, M., Jain, A., and Tian, H.: Global patterns of drought recovery, *Nature*, 548, 202–205,
 1651 <https://doi.org/10.1038/nature23021>, 2017.

1652 Scott, R.: FLUXNET2015 US-SRM Santa Rita Mesquite, <https://doi.org/10.18140/FLX/1440090>,
 1653 2016a.

1654 Scott, R.: FLUXNET2015 US-Whs Walnut Gulch Lucky Hills Shrub,
1655 <https://doi.org/10.18140/FLX/1440097>, 2016b.

1656 Shao, C.: FLUXNET2015 CN-Sw2 Siziwang Grazed (SZWG),
1657 <https://doi.org/10.18140/FLX/1440212>, 2016.

1658 Sigut, L., Havrankova, K., Jocher, G., Pavelka, M., Janouš, D., Czerny, R., Stanik, K., and Trusina, J.:
1659 FLUXNET2015 CZ-BK2 Bily Kriz grassland, <https://doi.org/10.18140/FLX/1440144>, 2016.

1660 Smith, W. K., Reed, S. C., Cleveland, C. C., Ballantyne, A. P., Anderegg, W. R. L., Wieder, W. R., Liu,
1661 Y. Y., and Running, S. W.: Large divergence of satellite and Earth system model estimates of global
1662 terrestrial CO₂ fertilization, *Nature Climate Change*, 6, 306–310,
1663 <https://doi.org/10.1038/nclimate2879>, 2016.

1664 Spano, D., Duce, P., Marras, S., Sirca, C., Arca, A., Zara, P., Ventura, A., Mereu, S., and Sanna, L.:
1665 FLUXNET2015 IT-Noe Arca di Noe - Le Prigionette, <https://doi.org/10.18140/FLX/1440171>,
1666 2016.

1667 Staebler, R.: AmeriFlux FLUXNET-1F CA-Cbo Ontario - Mixed Deciduous, Borden Forest Site,
1668 <https://doi.org/10.17190/AMF/1854365>, 2022.

1669 Ştefan, V. and Levin, S.: plotbiomes: R package for plotting Whittaker biomes with ggplot2, ,
1670 <https://doi.org/10.5281/zenodo.7145245>, 2018.

1671 Still, C. J., Berry, J. A., Collatz, G. J., and DeFries, R. S.: Global distribution of C₃ and C₄ vegetation:
1672 Carbon cycle implications, *Global Biogeochemical Cycles*, 17,
1673 <https://doi.org/10.1029/2001gb001807>, 2003.

1674 Still, C. J., Berry, J. A., Collatz, G. J., and DeFries, R. S.: ISLSCP II C₄ Vegetation Percentage, in: Hall,
1675 Forrest G., G. Collatz, B. Meeson, S. Los, E. Brown de Colstoun, and D. Landis (eds.). ISLSCP
1676 Initiative II Collection. Data set., <http://dx.doi.org/10.3334/ORNLDAAAC/932>, 2009.

1677 Stocker, B. D., Zscheischler, J., Keenan, T. F., Prentice, I. C., Peñuelas, J., and Seneviratne, S. I.:
1678 Quantifying soil moisture impacts on light use efficiency across biomes, *New Phytologist*, 218, 1430–
1679 1449, <https://doi.org/10.1111/nph.15123>, 2018.

1680 Stocker, B. D., Zscheischler, J., Keenan, T. F., Prentice, I. C., Seneviratne, S. I., and Peñuelas, J.:
1681 Drought impacts on terrestrial primary production underestimated by satellite monitoring, *Nature*
1682 *Geoscience*, 12, 264–270, <https://doi.org/10.1038/s41561-019-0318-6>, 2019.

1683 Stocker, B. D., Tumber-Dávila, S. J., Konings, A. G., Anderson, M. C., Hain, C., and Jackson, R. B.:
1684 Global patterns of water storage in the rooting zones of vegetation, *Nat. Geosci.*, 16, 250–256,
1685 <https://doi.org/10.1038/s41561-023-01125-2>, 2023.

1686 Sturtevant, C., Szutu, D., Baldocchi, D., Matthes, J. H., Oikawa, P., and Chamberlain, S. D.:
1687 FLUXNET2015 US-Myb Mayberry Wetland, <https://doi.org/10.18140/FLX/1440105>, 2016.

1688 Suyker, A.: FLUXNET2015 US-Ne1 Mead - irrigated continuous maize site,
1689 <https://doi.org/10.18140/FLX/1440084>, 2016a.

- 1690 Suyker, A.: FLUXNET2015 US-Ne2 Mead - irrigated maize-soybean rotation site,
1691 <https://doi.org/10.18140/FLX/1440085>, 2016b.
- 1692 Suyker, A.: FLUXNET2015 US-Ne3 Mead - rainfed maize-soybean rotation site,
1693 <https://doi.org/10.18140/FLX/1440086>, 2016c.
- 1694 Tagesson, T., Ardö, J., and Fensholt, R.: FLUXNET2015 SN-Dhr Dahra,
1695 <https://doi.org/10.18140/FLX/1440246>, 2016.
- 1696 Tang, Y., Kato, T., and Du, M.: FLUXNET2015 CN-HaM Haibei Alpine Tibet site,
1697 <https://doi.org/10.18140/FLX/1440190>, 2016.
- 1698 Terrer, C., Jackson, R. B., Prentice, I. C., Keenan, T. F., Kaiser, C., Vicca, S., Fisher, J. B., Reich, P.
1699 B., Stocker, B. D., Hungate, B. A., Peñuelas, J., McCallum, I., Soudzilovskaia, N. A., Cernusak, L. A.,
1700 Talhelm, A. F., Van Sundert, K., Piao, S., Newton, P. C. D., Hovenden, M. J., Blumenthal, D. M., Liu,
1701 Y. Y., Müller, C., Winter, K., Field, C. B., Viechtbauer, W., Van Lissa, C. J., Hoosbeek, M. R.,
1702 Watanabe, M., Koike, T., Leshyk, V. O., Polley, H. W., and Franklin, O.: Nitrogen and phosphorus
1703 constrain the CO₂ fertilization of global plant biomass, *Nature Climate Change*, 9, 684–689,
1704 <https://doi.org/10.1038/s41558-019-0545-2>, 2019.
- 1705 Thoning, K. W., Crotwell, A. M., and Mund, J. W.: Atmospheric Carbon Dioxide Dry Air Mole
1706 Fractions from continuous measurements at Mauna Loa, Hawaii, Barrow, Alaska, American Samoa
1707 and South Pole. 1973–2020, Version 2021-08-09, National Oceanic and Atmospheric Administration
1708 (NOAA), Global Monitoring Laboratory (GML), Boulder, Colorado, USA,
1709 <https://doi.org/10.15138/yaf1-bk21>, 2021.
- 1710 Tramontana, G., Ichii, K., Camps-Valls, G., Tomelleri, E., and Papale, D.: Uncertainty analysis of
1711 gross primary production upscaling using Random Forests, remote sensing and eddy covariance data,
1712 *Remote Sensing of Environment*, 168, 360–373, <https://doi.org/10.1016/j.rse.2015.07.015>, 2015.
- 1713 Tramontana, G., Jung, M., Schwalm, C. R., Ichii, K., Camps-Valls, G., Ráduly, B., Reichstein, M.,
1714 Arain, M. A., Cescatti, A., Kiely, G., Merbold, L., Serrano-Ortiz, P., Sickert, S., Wolf, S., and Papale,
1715 D.: Predicting carbon dioxide and energy fluxes across global FLUXNET sites with regression
1716 algorithms, *Biogeosciences*, 13, 4291–4313, <https://doi.org/10.5194/bg-13-4291-2016>, 2016.
- 1717 Ueyama, M., Iwata, H., and Harazono, Y.: AmeriFlux AmeriFlux US-Uaf University of Alaska,
1718 Fairbanks, <https://doi.org/10.17190/AMF/1480322>, 2018.
- 1719 Ueyama, M., Ichii, K., Kobayashi, H., Kumagai, T., Beringer, J., Merbold, L., Euskirchen, E. S., Hirano,
1720 T., Marchesini, L. B., Baldocchi, D., Saitoh, T. M., Mizoguchi, Y., Ono, K., Kim, J., Varlagin, A., Kang,
1721 M., Shimizu, T., Kosugi, Y., Bret-Harte, M. S., Machimura, T., Matsuura, Y., Ohta, T., Takagi, K.,
1722 Takanashi, S., and Yasuda, Y.: Inferring CO₂ fertilization effect based on global monitoring land-
1723 atmosphere exchange with a theoretical model, *Environmental Research Letters*, 15, 84009,
1724 <https://doi.org/10.1088/1748-9326/ab79e5>, 2020.
- 1725 Valach, A., Shortt, R., Szutu, D., Eichelmann, E., Knox, S., Hemes, K., Verfaillie, J., and Baldocchi,
1726 D.: AmeriFlux FLUXNET-1F US-Tw1 Twitchell Wetland West Pond,
1727 <https://doi.org/10.17190/AMF/1832165>, 2021.

1728 Valentini, R., Nicolini, G., Stefani, P., de Grandcourt, A., and Stivanello, S.: FLUXNET2015 GH-Ank
1729 Ankasa, <https://doi.org/10.18140/FLX/1440229>, 2016a.

1730 Valentini, R., Dore, S., Mazzenga, F., Sabbatini, S., Stefani, P., Tirone, G., and Papale, D.:
1731 FLUXNET2015 IT-Cpz Castelporziano, <https://doi.org/10.18140/FLX/1440168>, 2016b.

1732 Valentini, R., Tirone, G., Vitale, D., Papale, D., Arriga, N., Beilelli, L., Dore, S., Manca, G., Mazzenga,
1733 F., Pegoraro, E., Sabbatini, S., Stefani, P., Boschi, A., and Tomassucci, M.: FLUXNET2015 IT-Ro1
1734 Roccarespampani 1, <https://doi.org/10.18140/FLX/1440174>, 2016c.

1735 Villarreal, S. and Vargas, R.: Representativeness of FLUXNET Sites Across Latin America, *Journal of*
1736 *Geophysical Research: Biogeosciences*, 126, e2020JG006090,
1737 <https://doi.org/10.1029/2020JG006090>, 2021.

1738 Walker, A. P., De Kauwe, M. G., Bastos, A., Belmecheri, S., Georgiou, K., Keeling, R. F., McMahon,
1739 S. M., Medlyn, B. E., Moore, D. J. P., Norby, R. J., Zachle, S., Anderson-Teixeira, K. J., Battipaglia,
1740 G., Brien, R. J. W., Cabugao, K. G., Cailleret, M., Campbell, E., Canadell, J. G., Ciais, P., Craig, M.
1741 E., Ellsworth, D. S., Farquhar, G. D., Fatichi, S., Fisher, J. B., Frank, D. C., Graven, H., Gu, L., Haverd,
1742 V., Heilmann, K., Heimann, M., Hungate, B. A., Iversen, C. M., Joos, F., Jiang, M., Keenan, T. F.,
1743 Knauer, J., Körner, C., Leshyk, V. O., Leuzinger, S., Liu, Y., MacBean, N., Malhi, Y., McVicar, T. R.,
1744 Penuelas, J., Pongratz, J., Powell, A. S., Riutta, T., Sabot, M. E. B., Schleucher, J., Sitch, S., Smith, W.
1745 K., Sulman, B., Taylor, B., Terrer, C., Torn, M. S., Treseder, K. K., Trugman, A. T., Trumbore, S. E.,
1746 van Mantgem, P. J., Voelker, S. L., Whelan, M. E., and Zuidema, P. A.: Integrating the evidence for a
1747 terrestrial carbon sink caused by increasing atmospheric CO₂, *New Phytologist*, 229, 2413–2445,
1748 <https://doi.org/10.1111/nph.16866>, 2021.

1749 Walther, S., Besnard, S., Nelson, J. A., El-Madany, T. S., Migliavacca, M., Weber, U., Carvalhais, N.,
1750 Ermida, S. L., Brümmer, C., Schrader, F., Prokushkin, A. S., Panov, A. V., and Jung, M.: Technical
1751 note: A view from space on global flux towers by MODIS and Landsat: the FluxnetEO data set,
1752 *Biogeosciences*, 19, 2805–2840, <https://doi.org/10.5194/bg-19-2805-2022>, 2022.

1753 Wan, Z., Hook, S., and Hulley, G.: MOD11A1 MODIS/Terra Land Surface Temperature/Emissivity
1754 Daily L3 Global 1km SIN Grid V006 [Data set], NASA EOSDIS Land Processes DAAC,
1755 <https://doi.org/10.5067/MODIS/MOD11A1.006>, 2015a.

1756 Wan, Z., Hook, S., and Hulley, G.: MYD11A1 MODIS/Aqua Land Surface Temperature/Emissivity
1757 Daily L3 Global 1km SIN Grid V006 [Data set], NASA EOSDIS Land Processes DAAC,
1758 <https://doi.org/10.5067/MODIS/MYD11A1.006>, 2015b.

1759 Wang, H. and Fu, X.: FLUXNET2015 CN-Qia Qianyanzhou,
1760 <https://doi.org/10.18140/FLX/1440141>, 2016.

1761 Wang, H., Prentice, I. C., Keenan, T. F., Davis, T. W., Wright, I. J., Cornwell, W. K., Evans, B. J., and
1762 Peng, C.: Towards a universal model for carbon dioxide uptake by plants, *Nature Plants*, 3, 734–741,
1763 <https://doi.org/10.1038/s41477-017-0006-8>, 2017.

1764 Warm Winter 2020 Team: Warm Winter 2020 ecosystem eddy covariance flux product for 73 stations
1765 in FLUXNET-Archive format—release 2022-1 (Version 1.0), <https://doi.org/10.18160/2G60-ZHAK>, 2022.

1767 Wenzel, S., Cox, P. M., Eyring, V., and Friedlingstein, P.: Projected land photosynthesis constrained
1768 by changes in the seasonal cycle of atmospheric CO₂, *Nature*, 538, 499–501,
1769 <https://doi.org/10.1038/nature19772>, 2016.

1770 Wohlfahrt, G., Hammerle, A., and Hörtnagl, L.: FLUXNET2015 AT-Neu Neustift,
1771 <https://doi.org/10.18140/FLX/1440121>, 2016.

1772 Wolf, S., Eugster, W., and Buchmann, N.: FLUXNET2015 PA-SPn Sardinilla Plantation,
1773 <https://doi.org/10.18140/FLX/1440180>, 2016.

1774 Woodgate, W., van Gorsel, E., Leuning, R., Hughes, D., Kitchen, M., and Zegelin, S.: FLUXNET2015
1775 AU-Tum Tumburumba, <https://doi.org/10.18140/FLX/1440126>, 2016.

1776 Xiao, J., Zhuang, Q., Baldocchi, D. D., Law, B. E., Richardson, A. D., Chen, J., Oren, R., Starr, G.,
1777 Noormets, A., Ma, S., Verma, S. B., Wharton, S., Wofsy, S. C., Bolstad, P. V., Burns, S. P., Cook, D.
1778 R., Curtis, P. S., Drake, B. G., Falk, M., Fischer, M. L., Foster, D. R., Gu, L., Hadley, J. L., Hollinger,
1779 D. Y., Katul, G. G., Litvak, M., Martin, T. A., Matamala, R., McNulty, S., Meyers, T. P., Monson, R.
1780 K., Munger, J. W., Oechel, W. C., Paw U, K. T., Schmid, H. P., Scott, R. L., Sun, G., Suyker, A. E.,
1781 and Torn, M. S.: Estimation of net ecosystem carbon exchange for the conterminous United States
1782 by combining MODIS and AmeriFlux data, *Agricultural and Forest Meteorology*, 148, 1827–1847,
1783 <https://doi.org/10.1016/j.agrformet.2008.06.015>, 2008.

1784 Xie, X., Chen, J. M., Yuan, W., Guan, X., Jin, H., and Leng, J.: A Practical Algorithm for Correcting
1785 Topographical Effects on Global GPP Products, *Journal of Geophysical Research: Biogeosciences*,
1786 128, e2023JG007553, <https://doi.org/10.1029/2023JG007553>, 2023.

1787 Xie, Y., Gibbs, H. K., and Lark, T. J.: Landsat-based Irrigation Dataset (LANID): 30m resolution
1788 maps of irrigation distribution, frequency, and change for the US, 1997–2017, *Earth System Science*
1789 *Data*, 13, 5689–5710, <https://doi.org/10.5194/essd-13-5689-2021>, 2021.

1790 Yan, K., Park, T., Yan, G., Chen, C., Yang, B., Liu, Z., Nemani, R. R., Knyazikhin, Y., and Myneni, R.
1791 B.: Evaluation of MODIS LAI/FPAR product collection 6. Part 1: Consistency and improvements,
1792 *Remote Sensing*, 8, 1–16, <https://doi.org/10.3390/rs8050359>, 2016a.

1793 Yan, K., Park, T., Yan, G., Liu, Z., Yang, B., Chen, C., Nemani, R. R., Knyazikhin, Y., and Myneni, R.
1794 B.: Evaluation of MODIS LAI/FPAR product collection 6. Part 2: Validation and intercomparison,
1795 *Remote Sensing*, 8, 460, <https://doi.org/10.3390/rs8060460>, 2016b.

1796 Yang, F., Ichii, K., White, M. A., Hashimoto, H., Michaelis, A. R., Votava, P., Zhu, A. X., Huete, A.,
1797 Running, S. W., and Nemani, R. R.: Developing a continental-scale measure of gross primary
1798 production by combining MODIS and AmeriFlux data through Support Vector Machine approach,
1799 *Remote Sensing of Environment*, 110, 109–122, <https://doi.org/10.1016/j.rse.2007.02.016>, 2007.

1800 Yang, R., Wang, J., Zeng, N., Sitch, S., Tang, W., McGrath, M. J., Cai, Q., Liu, D., Lombardozzi, D.,
1801 Tian, H., Jain, A. K., and Han, P.: Divergent historical GPP trends among state-of-the-art multi-model
1802 simulations and satellite-based products, *Earth System Dynamics*, 13, 833–849,
1803 <https://doi.org/10.5194/esd-13-833-2022>, 2022.

1804 Yuan, H., Dai, Y., Xiao, Z., Ji, D., and Shangguan, W.: Reprocessing the MODIS Leaf Area Index
 1805 products for land surface and climate modelling, *Remote Sensing of Environment*, 115, 1171–1187,
 1806 <https://doi.org/10.1016/j.rse.2011.01.001>, 2011.

1807 Zeng, J., Matsunaga, T., Tan, Z.-H., Saigusa, N., Shirai, T., Tang, Y., Peng, S., and Fukuda, Y.: Global
 1808 terrestrial carbon fluxes of 1999–2019 estimated by upscaling eddy covariance data with a random
 1809 forest, *Sci Data*, 7, 313, <https://doi.org/10.1038/s41597-020-00653-5>, 2020.

1810 Zeng, N., Zhao, F., Collatz, G. J., Kalnay, E., Salawitch, R. J., West, T. O., and Guanter, L.: Agricultural
 1811 Green Revolution as a driver of increasing atmospheric CO₂ seasonal amplitude, *Nature*, 515, 394–
 1812 397, <https://doi.org/10.1038/nature13893>, 2014.

1813 Zhan, C., Orth, R., Migliavacca, M., Zaehle, S., Reichstein, M., Engel, J., Rammig, A., and Winkler, A.
 1814 J.: Emergence of the physiological effects of elevated CO₂ on land–atmosphere exchange of carbon
 1815 and water, *Global Change Biology*, 28, 7313–7326, <https://doi.org/10.1111/gcb.16397>, 2022.

1816 Zhang, J. and Han, S.: FLUXNET2015 CN-Cha Changbaishan,
 1817 <https://doi.org/10.18140/FLX/1440137>, 2016.

1818 Zhang, Y.: A global spatially contiguous solar-induced fluorescence (CSIF) dataset using neural
 1819 networks (2000–2020), National Tibetan Plateau Data Center,
 1820 <https://doi.org/10.11888/Ecolo.tpd.c.271751>, 2021.

1821 Zhang, Y., Joiner, J., Hamed Alemohammad, S., Zhou, S., and Gentine, P.: A global spatially
 1822 contiguous solar-induced fluorescence (CSIF) dataset using neural networks, *Biogeosciences*, 15,
 1823 5779–5800, <https://doi.org/10.5194/bg-15-5779-2018>, 2018.

1824 Zhang, Y., Kong, D., Gan, R., Chiew, F. H. S., McVicar, T. R., Zhang, Q., and Yang, Y.: Coupled
 1825 estimation of 500 m and 8-day resolution global evapotranspiration and gross primary production in
 1826 2002–2017, *Remote Sensing of Environment*, 222, 165–182,
 1827 <https://doi.org/10.1016/j.rse.2018.12.031>, 2019.

1828 Zheng, Y., Shen, R., Wang, Y., Li, X., Liu, S., Liang, S., Chen, J. M., Ju, W., Zhang, L., and Yuan, W.:
 1829 Improved estimate of global gross primary production for reproducing its long-Term variation, 1982–
 1830 2017, *Earth System Science Data*, 12, 2725–2746, <https://doi.org/10.5194/essd-12-2725-2020>, 2020.

1831 Zhou, G. and Yan, J.: FLUXNET2015 CN-Din Dinghushan,
 1832 <https://doi.org/10.18140/FLX/1440139>, 2016.

1833 Zhu, Z., Piao, S., Myneni, R. B., Huang, M., Zeng, Z., Canadell, J. G., Ciais, P., Sitch, S., Friedlingstein,
 1834 P., Arneth, A., Cao, C., Cheng, L., Kato, E., Koven, C., Li, Y., Lian, X., Liu, Y., Liu, R., Mao, J., Pan,
 1835 Y., Peng, S., Peuelas, J., Poulter, B., Pugh, T. A. M., Stocker, B. D., Viovy, N., Wang, X., Wang, Y.,
 1836 Xiao, Z., Yang, H., Zaehle, S., and Zeng, N.: Greening of the Earth and its drivers, *Nature Climate*
 1837 *Change*, 6, 791–795, <https://doi.org/10.1038/nclimate3004>, 2016.

1838 Zhuang, J., dussin, raphael, Huard, D., Bourgault, P., Banihirwe, A., Raynaud, S., Malevich, B.,
 1839 Schupfner, M., Filipe, Levang, S., Gauthier, C., Jüling, A., Almansi, M., RichardScottOZ, RondeauG,
 1840 Rasp, S., Smith, T. J., Stachelek, J., Plough, M., Pierre, Bell, R., Caneill, R., and Li, X.: xESMF: v0.8.2, ,
 1841 <https://doi.org/10.5281/zenodo.8356796>, 2023.

1842 Zona, D. and Oechel, W.: FLUXNET2015 US-Atq Atqasuk,
1843 <https://doi.org/10.18140/FLX/1440067>, 2016a.

1844 Zona, D. and Oechel, W.: FLUXNET2015 US-Ivo Ivotuk, <https://doi.org/10.18140/FLX/1440073>,
1845 2016b.

1846