

1 **A global database of dissolved organic matter (DOM) concentration**
2 **measurements in coastal waters (CoastDOM v1)**

3 Christian Lønborg^{1*}, Cátia Carreira², Gwenaël Abril³, Susana Agustí⁴, Valentina Amaral⁵,
4 Agneta Andersson⁶, Javier Arístegui⁷, Punyasloke Bhadury⁸, Mariana B. Bif⁹, Alberto V.
5 Borges¹⁰, Steven Bouillon¹¹, Maria Ll. Calleja^{4,12}, Luiz C. Cotovicz Jr^{13,49}, Stefano Cozzi¹⁴,
6 Maryló Doval¹⁵, Carlos M. Duarte⁴, Bradley Eyre¹⁶, Cédric G. Fichot¹⁷, E. Elena García-
7 Martín¹⁸, Alexandra Garzon-Garcia¹⁹, Michele Giani^{20,21}, Rafael Gonçalves-Araujo²²,
8 Renee Gruber²³, Dennis A. Hansell²⁴, Fuminori Hashihama²⁵, Ding He²⁶, Johnna M.
9 Holding²⁷, William R. Hunter²⁸, J. Severino P. Ibánhez²⁹, Valeria Ibello³⁰, Shan Jiang³¹,
10 Guebuem Kim³², Katja Klun³³, Piotr Kowalczyk³⁴, Atsushi Kubo³⁵, Choon Weng Lee³⁶,
11 Cláudia B. Lopes³⁷, Federica Maggioni³⁸, Paolo Magni³⁹, Celia Marrase⁴⁰, Patrick
12 Martin⁴¹, S. Leigh McCallister⁴², Roisin McCallum⁴³, Patricia M. Medeiros⁴⁴, Xosé Anxelu
13 G. Morán^{4,45}, Frank E. Muller-Karger⁴⁶, Allison Myers-Pigg⁴⁷, Marit Norli⁴⁸, Joanne M.
14 Oakes¹⁵, Helena Osterholz⁴⁹, Hyekyung Park³², Maria Lund Paulsen⁵⁰, Judith A.
15 Rosentreter^{16,51}, Jeff D. Ross⁵², Digna Rueda-Roa⁴⁶, Chiara Santinelli⁵³, Yuan Shen⁵⁴,
16 Eva Teira⁵⁵, Tinkara Tinta³³, Guenther Uher⁵⁶, Masahide Wakita⁵⁷, Nicholas Ward⁴⁷,
17 Kenta Watanabe⁵⁸, Yu Xin⁵⁹, Youhei Yamashita⁶⁰, Liyang Yang⁶¹, Jacob Yeo¹⁶, Huamao
18 Yuan⁶², Qiang Zheng^{54,63}, Xosé Antón Álvarez-Salgado²⁹

19 ¹Section for Marine Diversity and Experimental Ecology, Department of Ecoscience,
20 Aarhus University, 4000 Roskilde, Denmark.

21 ²Department of Environmental Science, Aarhus University, 4000 Roskilde, Denmark.

22 ³Laboratoire de Biologie des Organismes et Ecosystèmes Aquatiques (BOREA), CNRS,
23 Muséum National d'Histoire Naturelle, 61 rue Buffon, 75005, Paris, France.

24 ⁴King Abdullah University of Science and Technology, Thuwal 23955-6900, Kingdom of
25 Saudi Arabia.

26 ⁵Departamento Interdisciplinario de Sistemas Costero Marinos, Centro Universitario
27 Regional Este, Universidad de la República, Ruta 9 y 15, CP 27000, Rocha, Uruguay.
28 ⁶Umeå Marine Sciences Centre, Umeå University, Sweden.
29 ⁷Instituto de Oceanografía y Cambio Global (IOCAG), Universidad de Las Palmas de
30 Gran Canaria, Las Palmas, Spain.
31 ⁸Centre for Climate and Environmental Studies, Indian Institute of Science Education
32 and Research Kolkata, Mohanpur, Nadia, West Bengal, India.
33 ⁹Monterey Bay Aquarium Research Institute, Moss Landing, California, United States.
34 ¹⁰University of Liège, Chemical Oceanography Unit, Liège, Belgium.
35 ¹¹KU Leuven, Department of Earth and Environmental Sciences, Leuven, Belgium.
36 ¹²Marine Ecology and Systematics (MarES), Department of Biology, Universitat de les
37 Illes Balears, 07122 Palma de Mallorca, Spain.
38 ¹³Departamento de Geoquímica, Universidade Federal Fluminense, Outeiro São João
39 Batista s/n, 24020015 Niterói, RJ, Brazil.
40 ¹⁴Consiglio Nazionale delle Ricerche, Istituto di Scienze Marine (CNR-ISMAR), Strada
41 Statale 14, km 163.5, 34149 Trieste, Italy.
42 ¹⁵Instituto Tecnológico para o Control do Medio Mariño de Galicia, 36611 Vilagarcía de
43 Arousa, Spain.
44 ¹⁶Centre for Coastal Biogeochemistry, Faculty of Science and Engineering, Southern
45 Cross University, Lismore 2480, NSW, Australia.
46 ¹⁷Department of Earth and Environment, Boston University, Boston, MA, United States.
47 ¹⁸National Oceanography Centre, European Way, Southampton, SO14 3ZH, United
48 Kingdom.
49 ¹⁹Department of Environment and Science, PO Box 5078, Brisbane, Queensland 4001,
50 Australia.
51 ²⁰National Institute of Oceanography and Applied Geophysics (OGS), Trieste, Italy.

52 ²¹Istituto Centrale per la Ricerca scientifica e tecnologia Applicata al MAre, Chioggia,
53 Italy

54 ²²National Institute of Aquatic Resources, Technical University of Denmark, Kgs.
55 Lyngby, Denmark.

56 ²³Australian Institute of Marine Science, PMB 3, Townsville QLD 4810, Australia.

57 ²⁴Department of Ocean Sciences, Rosenstiel School of Marine and Atmospheric
58 Science, University of Miami, Miami, FL, United States.

59 ²⁵Tokyo University of Marine Science and Technology, Japan.

60 ²⁶Department of Ocean Science and Center for Ocean Research in Hong Kong and
61 Macau, The Hong Kong University of Science and Technology, Clear Water Bay, Hong
62 Kong, China.

63 ²⁷Department of Ecoscience, Aarhus University, 8000 Aarhus, Denmark.

64 ²⁸Fisheries and Aquatic Ecosystems Branch, Agri-Food and Biosciences Institute,
65 Belfast, Northern Ireland, United Kingdom.

66 ²⁹CSIC, Instituto de Investigaci3n Mariñas, Eduardo Cabello 6, 36208 Vigo, Spain.

67 ³⁰Institute of Marine Sciences, Middle East Technical University, 33731 Erdemli-Mersin,
68 Turkey.

69 ³¹State Key Laboratory of Estuarine and Coastal Research, East China Normal
70 University, 200241, Shanghai, China.

71 ³²School of Earth and Environmental Sciences, Seoul National University, Seoul 08826,
72 Korea.

73 ³³Marine Biology Station, National Institute of Biology, Fornače 41, 6330 Piran,
74 Slovenia.

75 ³⁴Remote Sensing Laboratory, Institute of Oceanology, Polish Academy of Sciences,
76 Sopot, Poland.

77 ³⁵Department of Geosciences, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka,
78 422-8529, Japan.

79 ³⁶Laboratory of Microbial Ecology, Institute of Biological Sciences, Institute of Ocean
80 and Earth Sciences, Universiti Malaya, 50603 Kuala Lumpur, Malaysia.

81 ³⁷CICECO – Aveiro Institute of Materials, Department of Chemistry, University of Aveiro,
82 Campus de Santiago, 3810-193 Aveiro, Portugal.

83 ³⁸University of New Caledonia and Institute de recherche pour le developement (IRD),
84 New Caledonia.

85 ³⁹Consiglio Nazionale delle Ricerche, Istituto per lo studio degli impatti Antropici e
86 Sostenibilità in ambiente marino (CNR-IAS), Loc. Sa Mardini, Torregrande, 09170,
87 Oristano, Italy.

88 ⁴⁰Institut de Ciències del Mar (CSIC), Passeig Marítim de la Barceloneta, 37
89 08003 Barcelona, Spain.

90 ⁴¹Asian School of the Environment, Nanyang Technological University, 639798,
91 Singapore.

92 ⁴²Virginia Commonwealth University, Department of Biology, Richmond, Virginia, United
93 States.

94 ⁴³Centre for Marine Ecosystems Research, School of Science, Edith Cowan University,
95 270 Joondalup Drive, Joondalup 6027 WA, Australia.

96 ⁴⁴Department of Marine Sciences, University of Georgia, Athens, Georgia, United
97 States.

98 ⁴⁵Centro Oceanográfico de Gijón/Xixón, Instituto Español de Oceanografía, Gijón/Xixón,
99 Spain.

100 ⁴⁶College of Marine Science, University of South Florida, Saint Petersburg, Florida
101 33701, United States.

102 ⁴⁷Pacific Northwest National Laboratory, Marine and Coastal Research Laboratory,
103 Sequim, Washington, United States.

104 ⁴⁸Norwegian Institute for Water Research, Oslo, Norway.

105 ⁴⁹Leibniz Institute for Baltic Sea Research Warnemünde, D-18119 Rostock-
106 Warnemünde, Germany.

107 ⁵⁰Marine microbiology, University of Bergen, Norway.

108 ⁵¹Yale School of the Environment, Yale University, New Haven, Connecticut, United
109 States.

110 ⁵²Institute for Marine and Antarctic Studies, University of Tasmania, Private Bag 49,
111 Hobart, Tasmania 7001, Australia

112 ⁵³Biophysics Institute, CNR, Pisa, Italy.

113 ⁵⁴State Key Laboratory of Marine Environmental Science (MEL) & College of Ocean
114 and Earth Sciences, Xiamen University, China.

115 ⁵⁵Departamento de Ecología y Biología Animal, Universidade de Vigo, Centro de
116 Investigación Mariña da Universidade de Vigo (CIM-UVigo), Vigo, Spain.

117 ⁵⁶School of Natural and Environmental Science, Newcastle University, Newcastle upon
118 Tyne, United Kingdom.

119 ⁵⁷Mutsu Institute for Oceanography, Research Institute for Global Change, Japan
120 Agency for Marine-Earth Science and Technology, 690 Kitasekine, Sekine, Mutsu,
121 Aomori, Japan.

122 ⁵⁸Coastal and Estuarine Environment Research Group, Port and Airport Research
123 Institute, Yokosuka 239-0826, Japan.

124 ⁵⁹Key Laboratory of Marine Chemistry Theory and Technology, Ministry of Education,
125 Institute for Advanced Ocean Study, Ocean University of China, Qingdao, Shandong,
126 China.

127 ⁶⁰Faculty of Environmental Earth Science, Hokkaido University, Hokkaido 060-0810,
128 Japan.

129 ⁶¹College of Environment and Safety Engineering, Fuzhou University, China.

130 ⁶²Key Laboratory of Marine Ecology and Environmental Sciences, Institute of
131 Oceanology, Chinese Academy of Sciences, Qingdao, China.

132 ⁶³Fujian Key Laboratory of Marine Carbon Sequestration, Xiamen University, Xiamen
133 361102, China.

134

135 *Corresponding author:

136 E-mail: c.lonborg@ecos.au.dk or clonborg@gmail.com

137

138 **ORCID nr.:**

139 Christian Lønborg: 0000-0001-8380-0238

140 Cátia Carreira: 0000-0002-1520-9320

141 Gwenaël Abril: 0000-0002-4914-086X

142 Susana Agustí: 0000-0003-0536-7293

143 Valentina Amaral: 0000-0002-1088-1484

144 Agneta Andersson: 0000-0001-7819-9038

145 Javier Arístegui: 0000-0002-7526-7741

146 Punyasloke Bhadury: 0000-0001-8714-7475

147 Mariana Bernardi Bif: 0000-0002-2148-4556

148 Alberto V. Borges: 0000-0002-5434-2247

149 Steven Bouillon: 0000-0001-7669-2929

150 Maria Ll. Calleja: 0000-0002-5992-2013

151 Luiz C. Cotovicz Jr: 0000-0002-3914-8155

152 Stefano Cozzi: 0000-0003-0116-742X

153 Maryló Doval: 0000-0002-8565-8703

154 Carlos M. Duarte: 0000-0002-1213-1361

155 Bradley Eyre: 0000-0001-5502-0680

156 Cédric G. Fichot: 0000-0002-1099-5764

157 E. Elena García-Martín: 0000-0003-4807-3287

158 Alexandra Garzon-Garcia: 0000-0002-6804-8890

159 Michele Giani: 0000-0002-3306-7725

160 Rafael Gonçalves-Araujo: 0000-0001-8344-8326

161 Renee Gruber: 0000-0002-8788-6910

162 Dennis A. Hansell: 0000-0001-9275-3445

163 Fuminori Hashihama: 0000-0003-3835-7681

164 Ding He: 0000-0001-9620-6115

165 Johnna M. Holding: 0000-0002-7364-0055

166 William R. Hunter: 0000-0001-8801-7947

167 J. Severino P. Ibánhez: 0000-0001-6093-3054
168 Valeria Ibello: 0000-0002-1067-0425
169 Shan Jiang: 0000-0002-1121-6080
170 Guebuem Kim: 0000-0002-5119-0241
171 Katja Klun: 0000-0001-6111-1650
172 Piotr Kowalczyk: 0000-0001-6016-0610
173 Atsushi Kubo: 0000-0002-6457-5386
174 Choon Weng Lee: 0000-0001-9805-9980
175 Cláudia Lopes: 0000-0001-7378-8677
176 Federica Maggioni: 0000-0002-7109-4257
177 Paolo Magni: 0000-0001-5955-6829
178 Celia Marrase: 0000-0002-5097-4829
179 Patrick Martin: 0000-0001-8008-5558
180 S. Leigh McCallister: 0000-0002-9041
181 Roisin McCallum: 0000-0002-0358-2371
182 Patricia M. Medeiros: 0000-0001-6818-2603
183 Xosé Anxelu G. Morán: 0000-002-9823-5339
184 Frank Muller-Karger: 0000-0003-3159-5011
185 Allison Myers-Pigg: 0000-0002-6905-6841
186 Marit Norli: 0000-0001-7472-1562
187 Joanne M. Oakes: 0000-0002-9287-2652
188 Helena Osterholz: 0000-0002-2858-9799
189 Hyekyung Park: 0000-0002-4743-5883
190 Maria Lund Paulsen: 0000-0002-1474-7258
191 Jeff D. Ross: 0000-0002-8659-3833
192 Judith A. Rosentreter: 0000-0001-5787-5682
193 Digna Rueda-Roa: 000-0003-4621-009X
194 Chiara Santinelli: 0000-0002-8921-275X
195 Yuan Shen: 0000-0001-6618-4226
196 Eva Teira: 0000-0002-4333-0101
197 Tinkara Tinta: 0000-0001-6740-8973
198 Guenther Uher: 0000-0001-5105-4445
199 Masahide Wakita: 0000-0002-3333-0546
200 Nicholas Ward: 0000-0001-6174-5581
201 Kenta Watanabe: 0000-0002-0106-3623
202 Yu Xin: 0000-0002-5328-7717
203 Youhei Yamashita: 0000-0002-9415-8743
204 Liyang Yang: 0000-0001-8767-8698
205 Jacob Yeo: 0000-0003-2443-5378
206 Huamao Yuan: 0000-0003-2014-619X
207 Qiang Zheng: 0000-0002-6836-2310
208 Xosé Antón Álvarez-Salgado: 0000-0002-2387-9201

209 **Abstract**

210 Measurements of dissolved organic carbon (DOC), nitrogen (DON), and phosphorus
211 (DOP) concentrations are used to characterize the dissolved organic matter (DOM) pool
212 and are important components of biogeochemical cycling in the coastal ocean. Here, we
213 present the first edition of a global database (CoastDOM v1; available at
214 <https://doi.pangaea.de/10.1594/PANGAEA.964012>) compiling previously published and
215 unpublished measurements of DOC, DON, and DOP in coastal waters. These data are
216 complemented by hydrographic data such as temperature and salinity and, to the extent
217 possible, other biogeochemical variables (e.g., Chlorophyll-a, inorganic nutrients) and the
218 inorganic carbon system (e.g., dissolved inorganic carbon and total alkalinity). Overall,
219 CoastDOM v1 includes observations of concentrations from all continents. However,
220 most data were collected in the Northern Hemisphere, with a clear gap in coastal water
221 DOM measurements from the Southern Hemisphere. The data included were collected
222 from 1978 to 2022 and consist of 62339 data points for DOC, 20360 for DON, and 13440
223 for DOP. The number of measurements decreases progressively in the sequence DOC
224 > DON > DOP, reflecting both differences in the maturity of the analytical methods and
225 the greater focus on carbon cycling by the aquatic science community. The global
226 database shows that the average DOC concentration in coastal waters (average \pm
227 standard deviation (SD): $182 \pm 314 \mu\text{mol C L}^{-1}$; median: $103 \mu\text{mol C L}^{-1}$) is 13-fold higher
228 than the average coastal DON concentration (average \pm SD: $13.6 \pm 30.4 \mu\text{mol N L}^{-1}$;
229 median: $8.0 \mu\text{mol N L}^{-1}$), which is itself 39-fold higher than the average coastal DOP
230 concentration (average \pm SD: $0.34 \pm 1.11 \mu\text{mol P L}^{-1}$; median: $0.18 \mu\text{mol P L}^{-1}$). This
231 dataset will be useful for identifying global spatial and temporal patterns in DOM and help
232 facilitate the reuse of DOC, DON, and DOP data in studies aimed at better characterizing
233 local biogeochemical processes, closing nutrient budgets, estimating carbon, nitrogen,

234 and phosphorous pools, as well as establishing a baseline for modelling future changes
235 in coastal waters.

236

237 **Keywords:** Dissolved organic matter, Dissolved organic carbon, Dissolved organic
238 nitrogen, Dissolved organic phosphorus, Coastal waters, Global database.

239 1. Introduction

240 Coastal waters are the most biogeochemical dynamic areas of the ocean, exhibiting
241 the highest standing stocks, process rates and transport fluxes of carbon (C), nitrogen
242 (N), and phosphorus (P) per unit area (Bauer et al., 2013; Mackenzie et al., 2011). In
243 these areas, organic matter plays a critical role in numerous biogeochemical processes,
244 serving as both a C, N, and P reservoir and substrate (Carreira et al., 2021).

245 Organic material found in the marine environment is commonly distinguished by its
246 size; material retained on a filter with a pore size typically between 0.2 and 0.7 μm is
247 classified as particulate organic matter (POM), whereas organic matter that passes
248 through the filter is referred to as dissolved organic matter (DOM). This partitioning is
249 operational but has implications for biogeochemical cycling: POM can be suspended in
250 the water column or sink to the sediments controlled by its size, shape and density
251 (Laurenceau-Cornec et al., 2015), whereas DOM is a solute that mostly remains in the
252 water column. In most coastal waters, DOM concentrations are higher than POM, with
253 POM having a larger proportion of known biochemical classes (e.g., carbohydrates,
254 proteins) than the dissolved fraction, suggesting that generally, DOM is more reworked
255 and recalcitrant (Boudreau and Ruddick, 1991; Lønborg et al., 2018; Benner and Amon,
256 2015).

257 The DOM pool consists mainly of C (DOC), N (DON), and P (DOP) but it also includes
258 other elements such as oxygen, sulphur and trace elements (Lønborg et al., 2020). In
259 coastal waters, DOM originates from multiple sources. Internal, or autochthonous,
260 sources include planktonic organisms (Lønborg et al., 2009; Carlson and Hansell, 2015),
261 benthic microalgae, macrophytes, and sediment porewater (Burdige and Komada, 2014;
262 Wada et al., 2008). On the other hand, DOM from external, or allochthonous, sources,
263 has mainly terrestrial origins, including wetlands, river and surface runoff, groundwater
264 discharges, and atmospheric deposition (Lavorivska et al., 2016; Raymond and Spencer,

265 2015; Taniguchi et al., 2019; Santos et al., 2021). The main sinks for DOM from the water
266 column in coastal waters are: 1) bubble coagulation and abiotic flocculation (Kerner et al.,
267 2003) or sorption to particles (Chin et al., 1998); 2) sunlight-mediated photodegradation
268 (Mopper et al., 2015); and 3) microbial degradation by mainly heterotrophic prokaryotes
269 (Lønborg and Álvarez-Salgado, 2012).

270 Given the importance of DOM as a source of nutrients and for coastal biogeochemical
271 cycling in general, numerous studies have measured the C, N and P content of the DOM
272 pool over the last few decades (e.g.,(García-Martín et al., 2021; Cauwet, 2002; Osterholz
273 et al., 2021). Most data, however, are often unavailable or stored in an inaccessible
274 manner, making it difficult to e.g., analyse global spatial and temporal patterns effectively.
275 Global open ocean DOM data compilation for DOC total dissolved nitrogen (TDN)
276 (Hansell et al., 2021) and DOP (Liang et al., 2022; Karl and Björkman, 2015) already exist
277 and contains few coastal samples (< 200m) (Hansell et al., 2021), but there are no
278 compilation specifically focused on coastal waters. Hence, there is a clear need for a
279 comprehensive global and integrated database of DOC, DON and DOP measurements
280 for coastal waters. To address this need, we have prepared the first edition of a coastal
281 DOM database (named CoastDOM v1), by compiling both previously reported as well as
282 unpublished data. These data have been obtained from authors of the original studies or
283 extracted directly from the original studies. In order to allow the DOM measurements to
284 be interpreted across larger scales, and to better understand their relationship with local
285 environmental conditions, we have included concurrently collected ancillary data (such
286 as physical and/or chemical seawater properties) whenever available. The objective of
287 this database is multifaceted. Firstly, we aimed to compile all available coastal DOM data
288 into a single repository. Secondly, our intention was to make these data easily accessible
289 to the research community and thirdly, we sought to achieve long-term consistency of the

290 measurements, to enable data intercomparison and establish a robust baseline for
291 assessing, for example, the impacts of climate change and land use changes.

292 **2. Methods**

293 **2.1. Data compilation**

294 The measurements included in CoastDOM v1 were obtained either directly from
295 authors of previously published studies, online databases, or scientific papers. An
296 extensive search of published reports, Ph.D. theses, and peer-reviewed literature was
297 performed to identify studies dealing with DOM in coastal waters. First, a formal search
298 was performed using Google Scholar in January 2022 using the search terms “dissolved
299 organic carbon”, “dissolved organic nitrogen”, and “dissolved organic phosphorus” in
300 connection with “marine” or “ocean”, which yielded a total of 897 articles (after filtering
301 the query by searching content in the title and abstract and excluding non-coastal
302 articles). When data could not be obtained directly from the corresponding authors,
303 relevant data were extracted. Further searches for relevant datasets were conducted
304 using the reference lists of the identified scientific papers as well as databases and
305 repositories to capture as many datasets as possible. Additionally, research groups that
306 were invited to participate in this effort were also encouraged to submit unpublished data
307 to CoastDOM v1.

308

309 **2.2. Dissolved organic matter analysis**

310 The DOC concentrations included in CoastDOM v1 were commonly measured using a
311 total organic carbon (TOC) hightemperature catalytic oxidation (HTCO) analyser (81% of
312 samples (Sharp et al., 1993). Some were measured by a combined wet chemical
313 oxidation (WCO) step and/or UV digestion, after which the carbon dioxide generated was
314 quantified (19% of samples). Similarly, concentrations of total dissolved nitrogen (TDN;

315 Sipler and Bronk, 2015) were determined using either a nitric oxide chemiluminescence
316 detector connected in series with the HTOCO analyser used for DOC analyses (31% of the
317 samples), or by employing a UV and/or chemical oxidation step (69%). In the latter
318 approach, both organic and inorganic N compounds were oxidised to nitrate, which was
319 subsequently quantified through a colorimetric method to determine the concentration of
320 inorganic N (Valderrama, 1981; Álvarez-Salgado et al., 2023; Halewood et al., 2022;
321 Foreman et al., 2019). Another method used for DON determination is oxidizing the
322 sample and measuring the resulting total nitrate by the nitric oxide chemiluminescence
323 method (Knapp et al., 2005). However, none of the concentration measurements included
324 in CoastDOM v1 applied this method. The reported DON concentrations were calculated
325 as the difference between TDN and dissolved inorganic nitrogen (DIN: sum of ammonium
326 (NH_4^+) and nitrate/nitrite ($\text{NO}_3^- + \text{NO}_2^-$); $\text{DON} = \text{TDN} - \text{DIN}$) (Álvarez-Salgado et al., 2023).
327 Analyses of total dissolved phosphorus (TDP) were determined by UV (4%) or wet
328 chemical oxidation (66%), or a combination of these (30%), and subsequently were
329 analysed for inorganic phosphorus by a colorimetric method (Álvarez-Salgado et al.,
330 2023). Another method also previously used for TDP analysis is the ash/hydrolysis
331 method (Solorzano and Sharp, 1980), even though none of the data included in
332 CoastDOM v1 used this method. The DOP concentrations were calculated as the
333 difference between TDP and soluble reactive phosphorus (SRP: HPO_4^{2-}) ($\text{DOP} = \text{TDP} -$
334 SRP) (Álvarez-Salgado et al., 2023).

335

336 **3. Description of the dataset**

337 The data compiled in CoastDOM v1 were collected, analysed and processed by different
338 laboratories, however, all data included have undergone quality control measures, either
339 by using reference samples or internal quality assurance procedures. While many of the
340 included DOC and TDN data have been systematically compared against consensus

341 reference material (CRM) mainly provided by the University of Miami's CRM program
342 (Hansell, 2005), there is a limitation in CoastDOM v1 regarding the intercalibration across
343 different measurement systems used for both DOP and DON determination. While the
344 CRM could be used for DOC, DON and DOP measurements, this has not yet been
345 attempted for DOP and measurement uncertainties increase in the sequence DOC >
346 DON > DOP. Although some of the reported measurements have quantified the DOP
347 recovery based on commercially available DOP compounds such as Adenosine
348 triphosphate (ATP), it is not known if these were conducted systematically in all cases.
349 Therefore, we strongly recommend undertaking further intercalibration across
350 laboratories for future measurements of TDP, as has been done for DOC and TDN
351 measurements (e.g.,(Sharp et al., 2002). Since additional quality control is not possible
352 in retrospect, we assessed the quality of CoastDOM v1 based on its internal consistency.

353 In CoastDOM v1, we defined "coastal water" as encompassing estuaries (salinity >
354 0.1) to the continental shelf break (water depth < 200 m). However, some locations, such
355 as deep fjords which are close to the coast cannot be classed as coastal due to
356 bathymetry (deeper than > 200 m). Therefore, we evaluated the inclusion of some
357 datasets on a case-by-case basis. For inclusion in the database, each DOM
358 measurement needed at a minimum to contain the following information (if reported in the
359 original publication or otherwise available):

360

- 361 - Country where samples were collected
- 362 - Latitude of measurement (in decimal units)
- 363 - Longitude of measurement (in decimal units)
- 364 - Year of sampling
- 365 - Month of sampling
- 366 - Sampling day (when available)

- 367 - Depth (m) at which the discrete samples were collected
- 368 - Temperature (°C) of the sample
- 369 - Salinity of the sample
- 370 - Dissolved organic carbon (DOC) concentration ($\mu\text{mol L}^{-1}$)
- 371 - Method used to measure DOC concentration
- 372 - DOC - QA flag: Quality flag for DOC measurement
- 373 - Dissolved organic nitrogen (DON) concentration ($\mu\text{mol L}^{-1}$)
- 374 - Total dissolved nitrogen (TDN) concentration ($\mu\text{mol L}^{-1}$)
- 375 - Method used to measure TDN concentration
- 376 - TDN - QA flag: Quality flag for TDN measurement
- 377 - Dissolved organic phosphorus (DOP) concentration ($\mu\text{mol L}^{-1}$)
- 378 - Total dissolved phosphorus (TDP) concentration ($\mu\text{mol L}^{-1}$)
- 379 - Method used to measure TDP concentration
- 380 - TDP - QA flag: Quality flag for TDP measurement
- 381 - Responsible person
- 382 - Originator institution
- 383 - Contact of data originator
- 384 It should be noted that in all entries, at least DOC, DON or DOP should have been
- 385 measured. In addition, we also included other relevant data, when available, in the
- 386 CoastDOM v1 dataset:
- 387
- 388 - Depth at the station where the sample was collected (Bottom depth, m).
- 389 - Total suspended solids (TSS) concentration (mg L^{-1})
- 390 - Chlorophyll-a (Chl a) concentration ($\mu\text{g L}^{-1}$)
- 391 - Chl a - QA flag: Quality flag for chlorophyll-a measurement
- 392 - Sum of nitrate and nitrite ($\text{NO}_3^- + \text{NO}_2^-$) concentration ($\mu\text{mol L}^{-1}$)

- 393 - $\text{NO}_3^- + \text{NO}_2^-$ - QA flag: Quality flag for $\text{NO}_3^- + \text{NO}_2^-$ measurement
- 394 - Ammonium (NH_4^+) concentration ($\mu\text{mol L}^{-1}$)
- 395 - NH_4^+ - QA flag: Quality flag for NH_4^+ measurement
- 396 - Soluble reactive phosphorus (HPO_4^{2-}) concentration ($\mu\text{mol L}^{-1}$)
- 397 - HPO_4^{2-} - QA flag: Quality flag for HPO_4^{2-} measurement
- 398 - Particulate organic carbon (POC) concentration ($\mu\text{mol L}^{-1}$)
- 399 - Method used to measure POC concentration
- 400 - POC - QA flag: Quality flag for POC measurement
- 401 - Particulate nitrogen (PN) concentration ($\mu\text{mol L}^{-1}$)
- 402 - Method used to measure PN concentration
- 403 - PN - QA flag: Quality flag for PN measurement
- 404 - Particulate phosphorus (PP) concentration ($\mu\text{mol L}^{-1}$)
- 405 - Method used to measure PP concentration
- 406 - PP - QA flag: Quality flag for PP measurement
- 407 - Dissolved inorganic carbon (DIC) concentration ($\mu\text{mol kg}^{-1}$)
- 408 - DIC - QA flag: Quality flag for DIC measurement
- 409 - Total alkalinity (TA) concentration ($\mu\text{mol kg}^{-1}$)
- 410 - TA - QA flag: Quality flag for TA measurement

411

412 Quality control of large datasets is crucial to ensure their reliability and usefulness.
413 Thus, we have not included data that were deemed compromised, such as records that
414 had not gone through quality control by the data originators. We also accepted a certain
415 degree of measurement error since multiple groups have been involved in the collection,
416 analysis, and/or compilation of the information. Some of these errors were corrected (e.g.,
417 when a value was placed in a wrong column, or clearly inaccurate locations were
418 reallocated for consistency with the place of study), while others could not be rectified

419 (e.g., values showing clear signs of contamination) and were consequently excluded from
420 CoastDOM v1 (Fig. 1). It should also be noted that differences in analytical capabilities
421 between laboratories and individual measurement campaigns likely caused additional
422 uncertainty. Outliers, arising for example from contamination, were removed from the
423 dataset. The data were moreover screened for zero values (i.e., concentrations below the
424 detection limit or absence of data). In cases where concentrations were below the
425 detection limit, the zero values were replaced with half the value of the limit-of-detection.
426 Commonly reported detection limits are $4 \mu\text{mol L}^{-1}$ for DOC, $0.3 \mu\text{mol L}^{-1}$ for DON and
427 are $0.03 \mu\text{mol L}^{-1}$ for DOP.

428 To ensure the inclusion of only high-quality data, we only accepted entries with specific
429 World Ocean Circulation Experiment (WOCE) quality codes: “2- Acceptable
430 measurement” and “6- Mean of replicate measurements”. In our quality control
431 assessments, we carefully avoided overly strict criteria, known as “data grooming”, which
432 could potentially overlook genuine patterns and changes in the dataset that may be
433 significant over longer temporal and/or wider spatial scales. Coastal waters are known to
434 exhibit a wide range of environmental concentrations, influenced by factors such as
435 seasonality and local anthropogenic activities. Consequently, these data points may
436 encompass a wide concentration range. However, obtaining consistent long-term
437 datasets is important to enable data intercomparison and establish a robust baseline.
438 Such long-term consistency can be achieved by using the CRM standards provided by
439 the Hansell laboratory for DOC and TDN. Another helpful approach is comparing the
440 DOM concentrations obtained by different laboratories in the same study area and time
441 of year.

442

443 **3.1 Summary of dissolved organic carbon (DOC) concentration observations**

444 Measurements of DOC concentrations were conducted between 1978 to 2022, with a
445 total of 62339 individual data points (Table 1). The DOC concentrations ranged from 17
446 to 30327 $\mu\text{mol C L}^{-1}$ (average \pm Standard Deviation (SD): $182 \pm 314 \mu\text{mol C L}^{-1}$; median:
447 $103 \mu\text{mol C L}^{-1}$; Table 1). The majority (53%) of the concentrations fell within the range of
448 60 to 120 $\mu\text{mol C L}^{-1}$ (Fig. 2). A large number of DOC concentration observations (17%)
449 ranged between 300 and 600 $\mu\text{mol C L}^{-1}$, which were predominantly collected in eutrophic
450 and river-influenced coastal waters of the Northern Hemisphere, such as the Baltic Sea
451 (Fig. 2). It was observed that 75% of the DOC concentrations were higher than 77 μmol
452 C L^{-1} , while 25% of the measurements surpassed 228 $\mu\text{mol C L}^{-1}$ (Table 1).

453 Coastal environments that experience minimal continental runoff, such as Palmer
454 Station in Antarctica, typically exhibit low DOC concentrations. On the other hand, coastal
455 waters heavily influenced by humic-rich terrigenous inputs, such as the Sarawak region
456 in Malaysia, tended to have high DOC concentrations. In addition, some extremely high
457 DOC concentrations were measured in the Derwent River in Australia which is impacted
458 by paper mill effluents. There has been a large increase in the number of DOC
459 concentration observations after 1992 (Fig. 3), and those measurements were from a
460 wide range of locations. However, these concentration observations were not evenly
461 distributed around the globe, with the Southern Hemisphere being under-sampled (10%
462 of observations), especially in the African, South American and Antarctic continents (Fig.
463 3, 4).

464

465 **3.2. Summary of dissolved organic nitrogen (DON) concentration observations**

466 The DON concentration measurements were collected between 1990 and 2021, with
467 a total of 20357 data points (Table 1). Concentrations of DON ranged from < 0.1 to 2095.3
468 $\mu\text{mol N L}^{-1}$ (average \pm SD: $13.6 \pm 30.4 \mu\text{mol N L}^{-1}$; median: $8.0 \mu\text{mol N L}^{-1}$; Table 1), with
469 the most common range (42%) for DON concentrations between 4 to 8 $\mu\text{mol N L}^{-1}$ (Fig.

470 2). Overall, 75% of DON concentrations were above $5.5 \mu\text{mol N L}^{-1}$, while 25% were
471 above $15.8 \mu\text{mol N L}^{-1}$ (Table 1).

472 The lowest DON concentrations were recorded in Young Sound, Greenland, which
473 receives direct run-off from the Greenland Ice Sheet, whereas the highest concentrations
474 were detected during a flood event in the Richmond River Estuary, Australia. Since 1995,
475 there has been a large increase in the number of DON measurements conducted in
476 coastal waters globally (Fig. 3); however, the majority of those measurements have been
477 in the Northern Hemisphere (79% of observations), mostly in Europe and the United
478 States (Figs. 3, 4).

479

480 **3.3. Summary of dissolved organic phosphorus (DOP) concentration**

481 **observations**

482 CoastDOM v1 includes a total of 13534 DOP measurements, collected between 1990
483 and 2021 (Table 1). Overall, DOP concentrations ranged from < 0.10 to $84.27 \mu\text{mol P L}^{-1}$
484 ¹ (average \pm SD: $0.34 \pm 1.11 \mu\text{mol P L}^{-1}$; median: $0.18 \mu\text{mol P L}^{-1}$; Table 2). The majority
485 (74%) of DOP concentrations were below $0.30 \mu\text{mol P L}^{-1}$ (Fig. 2). Analysis of the DOP
486 dataset revealed that 75% of the concentrations were above $0.11 \mu\text{mol P L}^{-1}$, while 25%
487 were above $0.30 \mu\text{mol P L}^{-1}$ (Table 1).

488 The lowest DOP concentrations were measured off the Kimberley Coast in Australia,
489 while the highest concentrations were found in the Vasse-Wonnerup Estuary in the South
490 west region of Australia. Similar to DOC and DON, most of the DOP measurements have
491 been conducted from the 1990s onwards, with a predominant focus in the Northern
492 Hemisphere (70% of observations), particularly in Europe and the United States (Figs. 3,
493 4).

494

495 **3.4. Summary of dissolved organic matter (DOM) concentration observations**

496 In CoastDOM v1 the number of measurements decreases progressively in the
497 sequence DOC > DON > DOP (62339, 20357, and 13534, respectively), reflecting both
498 differences in the maturity of the analytical methods and the greater focus on carbon
499 cycling by the aquatic science community. In addition, the average DOC concentration in
500 coastal waters (182 ± 314 $\mu\text{mol C L}^{-1}$), was 13-fold higher than the average coastal DON
501 concentrations (13.6 ± 30.4 $\mu\text{mol N L}^{-1}$), which was itself 39-fold higher than the average
502 coastal DOP concentrations (0.34 ± 1.11 $\mu\text{mol P L}^{-1}$) (Table 1). Interestingly the coefficient
503 of variation (C.V.- dispersion of the data around the mean) increased from DOC (173%)
504 to DON (224%) and DOP (326%), which is related to the fact that the % contribution of
505 refractory organic material decreases in the same sequence (Table 1). It should be noted
506 that CoastDOM v1 only contains 7058 paired measurements of DOC, DON, and DOP,
507 and therefore only a subset of observations reported all three element pools. The average
508 C: N: P stoichiometry for these paired DOM measurements was 1171 (± 4248): 100 (\pm
509 580): 1 (Table 1), which was very N- and P- depleted compared to the Redfield Ratio
510 (Redfield et al., 1963). However, the large variations in C:N, C:P and N:P ratios reveals
511 large variations in the composition of the DOM pool in coastal waters.

512

513 **3.5. Potential use of the dataset**

514 The use of the CoastDOM v1 dataset should be accompanied by the citation of this
515 paper and the inclusion of the correct doi-reference. CoastDOM v1 is available in full open
516 access on the PANGEA homepage as a *.csv file. The dataset includes a brief description
517 of the metadata and methods employed, with emphasis on measurement techniques and
518 data units. We chose the terminology most familiar to the ocean science community. It is
519 important to note that all data included in CoastDOM v1, as well as this manuscript, are
520 considered public domain; as such, a subset of this global dataset is also available in
521 previous data compilations (e.g.,(Hansell et al., 2021). The list of citations and links

522 referenced in CoastDOM v1 also provide users with information on how these data have
523 been previously used in publications or databases.

524

525 **3.6. Recommendations and conclusions**

526 In CoastDOM v1, we have compiled available coastal DOM data in a single repository,
527 making it openly and freely available to the research community. This compilation has
528 established a consistent global dataset, serving as a valuable information source to
529 investigate a variety of environmental questions and to explore spatial and temporal
530 trends. We suggest a set of recommendations for the future expansion of this global
531 dataset. First, our analysis highlights a spatial bias, with a concentration of sampling
532 efforts and/or data availability predominantly concentrated in the Northern Hemisphere.
533 The data gap in coastal DOM measurements in the Southern Hemisphere needs to be
534 addressed to provide a more representative global understanding of the role of DOM in
535 coastal water biogeochemistry. Additionally, increased sampling efforts especially around
536 Africa and South America, and island nations are warranted due to the vulnerability of
537 many coastal areas to climate change and intensifying human activities, which will
538 undoubtedly impact DOM biogeochemistry. Furthermore, there are comparatively few
539 data from coastal waters affected by river discharge into the tropics, e.g., the Amazon,
540 and Indian and Indonesian rivers that together dominate freshwater inputs to the coastal
541 ocean. Second, there is a need for more comprehensive temporal and spatial datasets to
542 capture the variability of DOM concentrations in highly dynamic and productive coastal
543 systems. Focused efforts should be made to resolve these temporal and spatial changes.
544 Third, only a fraction of data entries report paired DOC, DON and DOP measurements,
545 we encourage that these be measured and reported together in order to better determine
546 changes in stoichiometry and composition. Fourth, collecting and reporting ancillary data,
547 such as temperature, salinity, nutrient measurements, and particulate components, is

548 important to provide context and better understand the underlying processes driving the
549 observed DOM concentrations. Fifth, studies need to collect a minimum of metadata and
550 report it in standardized manner. Lastly, we recommend regular inter-calibration
551 exercises to establish standardised and interoperable methods and data, particularly for
552 DON and DOP measurements. This will ensure the comparability and reliability of data
553 across different studies and enhance our understanding of DON and DOP dynamics in
554 coastal waters.

555 In light of ongoing global environmental changes, the mobilisation and open sharing of
556 existing data for important biogeochemical variables, such as the DOM pool, are crucial
557 for establishing baselines and determining global trends and changes in coastal waters.
558 The aim is to publish an updated version of the database periodically to determine global
559 trends of DOM levels in coastal waters, and we therefore encourage researchers to
560 submit new data to the corresponding author. The CoastDOM v1 dataset was developed
561 according to the FAIR principles regarding Findability, Accessibility, Interoperability and
562 Reusability of data. Thus, CoastDOM v1 will serve as a reliable open-source information
563 resource, enabling in-depth analyses and providing quality-controlled input data for large-
564 scale ecosystem models.

565

566 **4. Data availability**

567 The dataset is available at the PANGAEA database
568 (<https://doi.pangaea.de/10.1594/PANGAEA.964012>; (Lønborg et al., 2023)). The file can
569 be downloaded as a *.csv merged file and is available in full open access.

570

571 **Competing interests**

572 The authors declare no competing interests.

573 **Author Contribution**

574 C.L., C.C., and X.A.A-S started the initiative and finalised the data compilation. All co-
575 authors contributed data. C.L. wrote the manuscript with input from all co-authors.

576 **Acknowledgement**

577 Qi Chen is acknowledged for his very skilful help with Figure 4.

578 **Funding**

579 During the drafting of the manuscript C.L. received funding from the Independent
580 Research Fund Denmark Grant No. 1127-00033B. The monitoring data obtained from
581 Bermuda received funding from the Bermuda Government Department of Environment
582 and Natural Resources. A subset of the data obtained from UK coastal estuaries received
583 funding from the Natural Environment Research Council (Grant NE/N018087/1). Data
584 retrieved from the Palmer LTER data were collected with support from the Office of Polar
585 Programs, US National Science Foundation. Data obtained from the Great Barrier Reef
586 Marine Monitoring Program for Inshore Water Quality, which is a partnership between the
587 Great Barrier Reef Marine Park Authority, the Australian Institute of Marine Science,
588 James Cook University, and the Cape York Water Partnership. The contribution by P.K.
589 was supported by DiSeDOM project contract no. UMO-2019/33/B/ST10/01232 funded by
590 the NCN - National Science Centre, Poland. N.W. and A.M.-P. participated in this
591 synthesis effort with funding provided by the U.S. Department of Energy funded
592 COMPASS-FME project; the provided data was collected with funding from the PREMIS
593 Initiative, conducted under the Laboratory Directed Research and Development Program
594 at Pacific Northwest National Laboratory. The data obtained from the Levantine Sea (Med
595 Sea) received funding from the Scientific and Technological Research Council of Türkiye
596 (TÜBİTAK, 1001 program, Grant 115Y629). The data obtained from Gulf of Trieste

597 (Slovenian waters) were financed by Research Program No. P1-0237 (Slovenian
598 Research and Innovation Agency). Data obtained from the northern Baltic Sea were
599 financed by the research program EcoChange (Swedish research council FORMAS). The
600 contribution by D.H. was supported by National Natural Science Foundation of China
601 (42222061), and funding support from the Center for Ocean Research in Hong Kong and
602 Macau (CORE). CORE is a joint research center for ocean research between Laoshan
603 Laboratory and HKUST. The contribution by Y.S. was supported by National Natural
604 Science Foundation of China (42106040), the Fundamental Research Funds for the
605 Central Universities of China (20720210076), and Fujian Provincial Central Guided Local
606 Science and Technology Development Special Project (Grant No. 2022L3078). The data
607 provided by LCCJr were supported by the Brazilian National Council of Research and
608 Development (CNPq-Pve No. 401.726/2012-6) and by the Carlos Chagas Foundation for
609 Research Support of the State of Rio de Janeiro (FAPERJ; No. E-26202.785/2016). Data
610 provided by SC were collected in the framework of Italian (PRISMA 1 and 2, ANOCSIA
611 and VECTOR) and European (OCEANCERTAIN) research projects. Data provided by
612 MG were collected within the Project "Mucilages in the Adriatic and Tyrrhenian Seas
613 (MAT) ", coordinated by the Istituto Centrale per la Ricerca Scientifica e Tecnologica
614 Applicata al Mare and financially supported by the Italian Ministry of the Environment.
615 Data obtained from the Georgia coast (USA) were supported by the National Science
616 Foundation through grants OCE-1832178 (GCE-LTER Program) and OCE-1902131.
617 Observations in the southern Caribbean Sea including the Cariaco Basin were collected
618 as part of the CARIACO Ocean Time Series program (supported by the Consejo Nacional
619 de Ciencia y Tecnología of Venezuela, the Ley de Ciencia, Tecnología e Innovación de
620 Venezuela, the Estación de Investigaciones Marinas de Venezuela; the National Science
621 Foundation grants OCE-0752139, OCE-9216626, OCE-9729284, OCE-491 9401537,
622 OCE-9729697, OCE-9415790, OCE-9711318, OCE-0326268, OCE-0963028, OCE-

623 0326313, and OCE-0326268, NASA grants NAG5-6448, NAS5-97128, and
624 NNX14AP62A; the Inter-American Institute for Global Change Research grant IAI-
625 CRN3094), and the Marine Biodiversity Observation Network/MBON of the Group on
626 Earth Observations Biodiversity Observation Network). Data provided by D.J.R. and
627 B.D.E. was supported by ARC Linkage (LP0770222), with Norske-Skog Boyer and the
628 Derwent Estuary Program providing financial and in-kind assistance.

629

630

631 **References**

- 632 Álvarez-Salgado, X. A., Nieto-Cid, M., and Rossel, P. E.: Dissolved Organic Matter, in:
633 Marine Analytical Chemistry, edited by: Blasco, J., and Tovar-Sánchez, A., Springer
634 International Publishing, Cham, 39-102, 10.1007/978-3-031-14486-8_2, 2023.
- 635 Bauer, J. E., Cai, W. J., Raymond, P. A., Bianchi, T. S., Hopkinson, C. S., and Regnier,
636 P. A.: The changing carbon cycle of the coastal ocean, *Nature*, 504, 61-70,
637 10.1038/nature12857, 2013.
- 638 Benner, R. and Amon, R. M.: The size-reactivity continuum of major bioelements in the
639 ocean, *Annual review of marine science*, 7, 185-205, 10.1146/annurev-marine-
640 010213-135126, 2015.
- 641 Boudreau, B. P. and Ruddick, B. R.: On a reactive continuum representation of organic
642 matter diagenesis, *American Journal of Science*, 291, 507-538, 1991.
- 643 Burdige, D. J. and Komada, T.: Sediment pore waters, in: *Biogeochemistry of marine*
644 *dissolved organic matter*, edited by: Hansen, D. A., and Carlson, C. A., Elsevier, 535-
645 577, 2014.
- 646 Carlson, C. A. and Hansell, D. A.: DOM Sources, Sinks, Reactivity, and Budgets, in:
647 *Biogeochemistry of Marine Dissolved Organic Matter*, edited by: Carlson, C. A., and
648 Hansell, D. A., Elsevier Science & Technology, 65-126, 10.1016/b978-0-12-405940-
649 5.00003-0, 2015.
- 650 Carreira, C., Talbot, S., and Lønborg, C.: Bacterial consumption of total and dissolved
651 organic carbon in the Great Barrier Reef, *Biogeochemistry*, 10.1007/s10533-021-
652 00802-x, 2021.
- 653 Cauwet, G.: DOM in coastal areas, in: *Biogeochemistry of Dissolved organic matter*,
654 edited by: Hansell, D., and Carlson, C. A., Academic Press, London, 579-609, 2002.
- 655 Chin, W. C., Orellana, M. V., and Verdugo, P.: Spontaneous assembly of marine
656 dissolved organic matter into polymer gels, *Nature*, 391, 568-572, 1998.
- 657 Foreman, R. K., Björkman, K. M., Carlson, C. A., Opalk, K., and Karl, D. M.: Improved
658 ultraviolet photo-oxidation system yields estimates for deep-sea dissolved organic
659 nitrogen and phosphorus, *Limnology and Oceanography: Methods*, 17, 277-291,
660 <https://doi.org/10.1002/lom3.10312>, 2019.
- 661 García-Martín, E. E., Sanders, R., Evans, C. D., Kitidis, V., Lapworth, D. J., Rees, A. P.,
662 Spears, B. M., Tye, A., Williamson, J. L., Balfour, C., Best, M., Bowes, M., Breimann,
663 S., Brown, I. J., Burden, A., Callaghan, N., Felgate, S. L., Fishwick, J., Fraser, M.,
664 Gibb, S. W., Gilbert, P. J., Godsell, N., Gomez-Castillo, A. P., Hargreaves, G., Jones,
665 O., Kennedy, P., Lichtschlag, A., Martin, A., May, R., Mawji, E., Mounteney, I.,
666 Nightingale, P. D., Olszewska, J. P., Painter, S. C., Pearce, C. R., Pereira, M. G.,

667 Peel, K., Pickard, A., Stephens, J. A., Stinchcombe, M., Williams, P., Woodward, E.
668 M. S., Yarrow, D., and Mayor, D. J.: Contrasting Estuarine Processing of Dissolved
669 Organic Matter Derived From Natural and Human-Impacted Landscapes, *Global*
670 *Biogeochemical Cycles*, 35, e2021GB007023,
671 <https://doi.org/10.1029/2021GB007023>, 2021.

672 Halewood, E., Opalk, K., Custals, L., Carey, M., Hansell, D. A., and Carlson, C. A.:
673 Determination of dissolved organic carbon and total dissolved nitrogen in seawater
674 using High Temperature Combustion Analysis, *Frontiers in Marine Science*, 9,
675 10.3389/fmars.2022.1061646, 2022.

676 Hansell, D. A.: Dissolved Organic Carbon Reference Material Program, *Eos*,
677 *Transactions American Geophysical Union*, 86, 318-318,
678 <https://doi.org/10.1029/2005EO350003>, 2005.

679 Hansell, D. A., Carlson, C. A., Amon, R. M. W., Álvarez-Salgado, X. A., Yamashita, Y.,
680 Romera-Castillo, C., and Bif, M. B.: Compilation of dissolved organic matter (DOM)
681 data obtained from global ocean observations from 1994 to 2021. Version 2. (NCEI
682 Accession 0227166) [dataset], doi.org/10.25921/s4f4-ye35, 2021.

683 Iavorivska, L., Boyer, E. W., and DeWalle, D. R.: Atmospheric deposition of organic
684 carbon via precipitation, *Atmospheric Environment*, 146, 153-163,
685 <https://doi.org/10.1016/j.atmosenv.2016.06.006>, 2016.

686 Karl, D. M. and Björkman, K. M.: Dynamics of Dissolved Organic Phosphorus, in:
687 *Biogeochemistry of marine dissolved organic matter*, edited by: Hansell, D. A., and
688 Carlson, C. A., 233-334, 10.1016/b978-0-12-405940-5.00005-4, 2015.

689 Kerner, M., Hohenberg, H., Ertl, S., Reckermann, M., and Spitzzy, A.: Self-organization
690 of dissolved organic matter tomicelle-like microparticles in river water, *Nature*, 422,
691 150-154, 2003.

692 Knapp, A. N., Sigman, D. M., and Lipschultz, F.: N isotopic composition of dissolved
693 organic nitrogen and nitrate at the Bermuda Atlantic Time-series Study site, *Global*
694 *Biogeochemical Cycles*, 19, <https://doi.org/10.1029/2004GB002320>, 2005.

695 Laruelle, G. G., Dürr, H. H., Lauerwald, R., Hartmann, J., Slomp, C. P., Goossens, N.,
696 and Regnier, P. A. G.: Global multi-scale segmentation of continental and coastal
697 waters from the watersheds to the continental margins, *Hydrol. Earth Syst. Sci.*, 17,
698 2029-2051, 10.5194/hess-17-2029-2013, 2013.

699 Laurenceau-Cornec, E. C., Trull, T. W., Davies, D. M., De La Rocha, C. L., and Blain,
700 S.: Phytoplankton morphology controls on marine snow sinking velocity, *Marine*
701 *Ecology Progress Series*, 520, 35-56, 2015.

702 Liang, Z., McCabe, K., Fawcett, S. E., Forrer, H. J., Hashihama, F., Jeandel, C.,
703 Marconi, D., Planquette, H., Saito, M. A., Sohm, J. A., Thomas, R. K., Letscher, R. T.,
704 and Knapp, A. N.: A global ocean dissolved organic phosphorus concentration
705 database (DOPv2021), *Scientific Data*, 9, 772, 10.1038/s41597-022-01873-7, 2022.

706 Lønborg, C. and Álvarez-Salgado, X. A.: Recycling versus export of bioavailable
707 dissolved organic matter in the coastal ocean and efficiency of the continental shelf
708 pump, *Global biogeochemical cycles* 26, GB3018, 10.1029/2012GB004353, 2012.

709 Lønborg, C., Álvarez-Salgado, X. A., Davidson, K., and Miller, A. E. J.: Production of
710 bioavailable and refractory dissolved organic matter by coastal heterotrophic
711 microbial populations, *Estuarine, Coastal and Shelf science*, 82, 682–688,
712 10.1016/j.ecss.2009.02.026, 2009.

713 Lønborg, C., Álvarez-Salgado, X. A., Duggan, S., and Carreira, C.: Organic matter
714 bioavailability in tropical coastal waters: The Great Barrier Reef, *Limnology and*
715 *Oceanography*, 63, 1015-1035, 10.1002/lno.10717, 2018.

716 Lønborg, C., Carreira, C., Jickells, T., and Álvarez-Salgado, X. A.: Impacts of Global
717 Change on Ocean Dissolved Organic Carbon (DOC) Cycling, *Frontiers in Marine*
718 *Science*, 7, 466, 10.3389/fmars.2020.00466, 2020.

719 Lønborg, C., Carreira, C., Abril, G., Agustí, S., Amaral, V., Andersson, A., Arístegui, J.,
720 Bhadury, P., Bif, M. B., Borges, A. V., Bouillon, S., Calleja, M. L., Cotovicz Jr, L. C.,
721 Cozzi, S., Doval, M. D., Duarte, C. M., Eyre, B. D., Fichot, C. G., García-Martín, E.
722 E., Garzon-Garcia, A., Giani, M., Gonçalves-Araujo, R., Gruber, R., Hansell, D. A.,
723 Hashihama, F., He, D., Holding, J. M., Hunter, W. R., Ibánhez, J. S., Ibello, V., Jiang,
724 S., Kim, G., Klun, K., Kowalczyk, P., Kubo, A., Weng Lee, C., Lopes, C., Maggioni,
725 F., Magni, P., Marrase, C., Martin, P., McCallister, S. L., McCallum, R., Medeiros, P.
726 M., Morán, X. A. G., Muller-Karger, F., Myers-Pigg, A., Norli, M., Oakes, J. M.,
727 Osterholz, H., Park, H., Paulsen, M. L., Rosentreter, J. A., Rueda-Roa, D., Santinelli,
728 C., Shen, Y., Teira, E., Tinta, T., Uher, G., Wakita, M., Ward, N., Watanabe, K., Xin,
729 Y., Yamashita, Y., Yang, L., Yeo, J., Yuan, H., Zheng, Q., and Álvarez-Salgado, X.
730 A.: A global database of dissolved organic matter (DOM) concentration
731 measurements in coastal waters (CoastDOM v1) [dataset], 2023.

732 Mackenzie, F. T., De Carlo, E. H., and Lerman, A.: 5.10 - Coupled C, N, P, and O
733 Biogeochemical Cycling at the Land–Ocean Interface, in: *Treatise on Estuarine and*
734 *Coastal Science*, edited by: Wolanski, E., and McLusky, D., Academic Press,
735 Waltham, 317-342, <https://doi.org/10.1016/B978-0-12-374711-2.00512-X>, 2011.

736 Mopper, K., Kieber, D. J., and Stubbins, A.: Marine Photochemistry of Organic Matter,
737 in: *Biogeochemistry of Marine Dissolved Organic Matter*, edited by: Carlson, C. A.,
738 and Hansell, D. A., 389-450, 10.1016/b978-0-12-405940-5.00008-x, 2015.

739 Osterholz, H., Burmeister, C., Busch, S., Dierken, M., Frazão, H. C., Hansen, R.,
740 Jeschek, J., Kremp, A., Kreuzer, L., and Sadkowiak, B.: Nearshore dissolved and
741 particulate organic matter dynamics in the southwestern Baltic Sea: environmental
742 drivers and time series analysis (2010–2020), *Frontiers in Marine Science*, 8,
743 795028, 2021.

744 Raymond, P. A. and Spencer, R. G. M.: Riverine DOM, in: *Biogeochemistry of Marine*
745 *Dissolved Organic Matter*, edited by: Hansell, D. A., and Carlson, C. A., Elsevier,
746 Amsterdam, 509-533, 10.1016/b978-0-12-405940-5.00011-x, 2015.

747 Redfield, A. C., Ketchum, B. K., and Richards, F. A.: The influence of organisms on the
748 composition of sea-water, in: *The sea*, vol. 2, *The composition of sea water:*
749 *Comparative and descriptive oceanography*, edited by: Hill, M. N., Wiley-Interscience,
750 26- 77, 1963.

751 Santos, I. R., Burdige, D. J., Jennerjahn, T. C., Bouillon, S., Cabral, A., Serrano, O.,
752 Wernberg, T., Filbee-Dexter, K., Guimond, J. A., and Tamborski, J. J.: The
753 renaissance of Odum's outwelling hypothesis in 'Blue Carbon' science, *Estuarine,*
754 *Coastal and Shelf Science*, 255, 10.1016/j.ecss.2021.107361, 2021.

755 Sharp, J. H., Benner, R., Bennett, L., Carlson, C. A., Dow, R., and Fitzwater, S. E.: Re-
756 evaluation of high temperature combustion and chemical oxidation measurements of
757 dissolved organic carbon in seawater, *Limnology and Oceanography*, 38, 1774-1782,
758 <https://doi.org/10.4319/lo.1993.38.8.1774>, 1993.

759 Sharp, J. H., Rinker, K. R., Savidge, K. B., Abell, J., Benaim, J. Y., Bronk, D., Burdige,
760 D. J., Cauwet, G., Chen, W., Doval, M. D., Hansell, D., Hopkinson, C., Kattner, G.,
761 Kaumeyer, N., McGlathery, K. J., Merriam, J., Morley, N., Nagel, K., Ogawa, H.,
762 Pollard, C., Pujo-Pay, M., Raimbault, P., Sambrotto, R., Seitzinger, S., Spyres, G.,
763 Tirendi, F., Walsh, T. W., and Wong, C. S.: A preliminary methods comparison for
764 measurement of dissolved organic nitrogen in seawater, *Marine Chemistry*, 78, 171-
765 184, 2002.

766 Sipler, R. E. and Bronk, D. A.: Dynamics of Dissolved Organic Nitrogen, in:
767 Biogeochemistry of Marine Dissolved Organic Matter, edited by: Hansell, D. A., and
768 Carlson, C. A., 127-232, 10.1016/b978-0-12-405940-5.00004-2, 2015.

769 Solorzano, L. and Sharp, J. H.: Determination of total dissolved phosphorus and
770 particulate phosphorus in natural waters, *Limnology and Oceanography*, 25, 754-
771 758, 1980.

772 Taniguchi, M., Dulai, H., Burnett, K. M., Santos, I. R., Sugimoto, R., Stieglitz, T., Kim,
773 G., Moosdorf, N., and Burnett, W. C.: Submarine Groundwater Discharge: Updates
774 on Its Measurement Techniques, Geophysical Drivers, Magnitudes, and Effects,
775 *Frontiers in Environmental Science*, 7, 10.3389/fenvs.2019.00141, 2019.

776 Valderrama, J. C.: The simultaneous analysis of total nitrogen and total phosphorus in
777 natural waters, *Marine Chemistry*, 10, 109-122, 1981.

778 Wada, S., Aoki, M. N., Mikami, A., Komatsu, T., Tsuchiya, Y., Sato, T., Shinagawa, H.,
779 and Hama, T.: Bioavailability of macroalgal dissolved organic matter in seawater,
780 *Marine Ecology Progress Series*, 370, 33-44, 2008

781

782 **Figure legends**

783 **Figure 1.** Flow diagram of data collation, quality control and inclusion into CoastDOM v1
784 database.

785 **Figure 2.** Histograms showing the distribution of observations for **a)** dissolved organic
786 carbon (DOC), **b)** nitrogen (DON) and **c)** phosphorus (DOP), within defined
787 concentration ranges in the coastal ocean. Note that the concentration ranges are not
788 uniform in all cases due to the large difference in concentrations.


789 **Figure 3. a)** Cumulative number of concentration observations for dissolved organic
790 carbon (DOC), nitrogen (DON), and phosphorus (DOP). Number of concentration
791 observations shown as a function of **b)** sampling month (“N.S” are samples for which
792 the sampling month is not specified), **c)** latitude, and **d)** longitude, grouped into bins of
793 10° latitude or longitude.

794 **Figure 4** Global distribution of concentration observations included in CoastDOM v1 for
795 **a)** dissolved organic carbon (DOC), **b)** nitrogen (DON), and **c)** phosphorus (DOP). The
796 black dots on the map represent the reported data that are included in the CoastDOM
797 v1 database. Histograms show the distribution of observations for DOC, DON and DOP
798 within defined concentration ranges in the continents where measurements are
799 available. Maps were created using the GIS shape file obtained from Laurelle et al.
800 (Laruelle et al., 2013)

801 **Table 1.** Descriptive statistics for the dissolved organic carbon (DOC), dissolved organic
802 nitrogen (DON), and dissolved organic phosphorus (DOP) concentration observations
803 included in the CoastDOM v1 dataset. The DOC:DON, DOC:DOP and DON:DOP ratios
804 are also reported. The minimum (Min), maximum (Max), average values (Avg.) and
805 standard deviation (SD), coefficient of variation (CV %), median, 25th and 75th
806 percentiles (perc.) and number of samples (N) for each variable are shown.


	DOC $\mu\text{mol L}^{-1}$	DON $\mu\text{mol L}^{-1}$	DOP $\mu\text{mol L}^{-1}$	DOC:DON	DOC:DOP	DON:DOP
Min	17	< 0.1	< 0.01	1	18	0.14
Max	30327	2095.3	84.27	3046	248024	8894
Avg. \pm SD	182 \pm 314	13.6 \pm 30.4	0.34 \pm 1.11	18 \pm 43	1171 \pm 4248	100 \pm 580
Median	103	8.0	0.18	14	583	47
CV	173	224	324	244	363	578
25%iles	77	5.5	0.11	11	401	30
75%iles	228	15.8	0.30	18	1034	78
N	62339	20357	13534	12632	7415	12954

807




808

809 **Figure 1.**




810

811 **Figure 2.**



812

813 **Figure 3.**



814

815 **Figure 4.**

1 **A global database of dissolved organic matter (DOM) concentration**

2 **measurements in coastal waters (CoastDOM v1)**

3 Christian Lønborg¹, Cátia Carreira², Gwenaël Abril³, Susana Agustí⁴, Valentina Amaral⁵,
4 Agneta Andersson⁶, Javier Arístegui⁷, Punyasloke Bhadury⁸, Mariana B. Bif⁹, Alberto V.
5 Borges¹⁰, Steven Bouillon¹¹, Maria LI. Calleja^{4,12}, Luiz C. Cotovicz Jr^{13,49}, Stefano Cozzi¹⁴,
6 Maryló Doval¹⁵, Carlos M. Duarte⁴, Bradley Eyre¹⁶, Cédric G. Fichot¹⁷, E. Elena García-
7 Martín¹⁸, Alexandra Garzon-Garcia¹⁹, Michele Giani^{20,21}, Rafael Gonçalves-Araujo²²,
8 Renee Gruber²³, Dennis A. Hansell²⁴, Fuminori Hashihama²⁵, Ding He²⁶, Johnna M.
9 Holding²⁷, William R. Hunter²⁸, J. Severino P. Ibánhez²⁹, Valeria Ibello³⁰, Shan Jiang³¹,
10 Guebuem Kim³², Katja Klun³³, Piotr Kowalczyk³⁴, Atsushi Kubo³⁵, Choon Weng Lee³⁶,
11 Cláudia B. Lopes³⁷, Federica Maggioni³⁸, Paolo Magni³⁹, Celia Marrase⁴⁰, Patrick
12 Martin⁴¹, S. Leigh McCallister⁴², Roisin McCallum⁴³, Patricia M. Medeiros⁴⁴, Xosé Anxelu
13 G. Morán^{4,45}, Frank E. Muller-Karger⁴⁶, Allison Myers-Pigg⁴⁷, Marit Norli⁴⁸, Joanne M.
14 Oakes¹⁵, Helena Osterholz⁴⁹, Hyekyung Park³², Maria Lund Paulsen⁵⁰, Judith A.
15 Rosentreter^{16,51}, Jeff D. Ross⁵², Digna Rueda-Roa⁴⁶, Chiara Santinelli⁵³, Yuan Shen⁵⁴,
16 Eva Teira⁵⁵, Tinkara Tinta³³, Guenther Uher⁵⁶, Masahide Wakita⁵⁷, Nicholas Ward⁴⁷,
17 Kenta Watanabe⁵⁸, Yu Xin⁵⁹, Youhei Yamashita⁶⁰, Liyang Yang⁶¹, Jacob Yeo¹⁶, Huamao
18 Yuan⁶², Qiang Zheng^{54,63}, Xosé Antón Álvarez-Salgado²⁹

19 ¹Section for Marine Diversity and Experimental Ecology, Department of Ecoscience,
20 Aarhus University, 4000 Roskilde, Denmark.

21 ²Department of Environmental Science, Aarhus University, 4000 Roskilde, Denmark.

22 ³Laboratoire de Biologie des Organismes et Ecosystèmes Aquatiques (BOREA), CNRS,
23 Muséum National d'Histoire Naturelle, 61 rue Buffon, 75005, Paris, France.

24 ⁴King Abdullah University of Science and Technology, Thuwal 23955-6900, Kingdom of
25 Saudi Arabia.

26 ⁵Departamento Interdisciplinario de Sistemas Costero Marinos, Centro Universitario
27 Regional Este, Universidad de la República, Ruta 9 y 15, CP 27000, Rocha, Uruguay.
28 ⁶Umeå Marine Sciences Centre, Umeå University, Sweden.
29 ⁷Instituto de Oceanografía y Cambio Global (IOGAG), Universidad de Las Palmas de
30 Gran Canaria, Las Palmas, Spain.
31 ⁸Centre for Climate and Environmental Studies, Indian Institute of Science Education
32 and Research Kolkata, Mohanpur, Nadia, West Bengal, India.
33 ⁹Monterey Bay Aquarium Research Institute, Moss Landing, California, United States.
34 ¹⁰University of Liège, Chemical Oceanography Unit, Liège, Belgium.
35 ¹¹KU Leuven, Department of Earth and Environmental Sciences, Leuven, Belgium.
36 ¹²Marine Ecology and Systematics (MarES), Department of Biology, Universitat de les
37 Illes Balears, 07122 Palma de Mallorca, Spain.
38 ¹³Departamento de Geoquímica, Universidade Federal Fluminense, Outeiro São João
39 Batista s/n, 24020015 Niterói, RJ, Brazil.
40 ¹⁴Consiglio Nazionale delle Ricerche, Istituto di Scienze Marine (CNR-ISMAR), Strada
41 Statale 14, km 163.5, 34149 Trieste, Italy.
42 ¹⁵Instituto Tecnológico para o Control do Medio Mariño de Galicia, 36611 Vilagarcía de
43 Arousa, Spain.
44 ¹⁶Centre for Coastal Biogeochemistry, Faculty of Science and Engineering, Southern
45 Cross University, Lismore 2480, NSW, Australia.
46 ¹⁷Department of Earth and Environment, Boston University, Boston, MA, United States.
47 ¹⁸National Oceanography Centre, European Way, Southampton, SO14 3ZH, United
48 Kingdom.
49 ¹⁹Department of Environment and Science, PO Box 5078, Brisbane, Queensland 4001,
50 Australia.
51 ²⁰National Institute of Oceanography and Applied Geophysics (OGS), Trieste, Italy.

52 ²¹Istituto Centrale per la Ricerca scientifica e tecnologia Applicata al MAre, Chioggia,
53 Italy

54 ²²National Institute of Aquatic Resources, Technical University of Denmark, Kgs.
55 Lyngby, Denmark.

56 ²³Australian Institute of Marine Science, PMB 3, Townsville QLD 4810, Australia.
57 ²⁴Department of Ocean Sciences, Rosenstiel School of Marine and Atmospheric
58 Science, University of Miami, Miami, FL, United States.

59 ²⁵Tokyo University of Marine Science and Technology, Japan.

60 ²⁶Department of Ocean Science and Center for Ocean Research in Hong Kong and
61 Macau, The Hong Kong University of Science and Technology, Clear Water Bay, Hong
62 Kong, China.

63 ²⁷Department of Ecoscience, Aarhus University, 8000 Aarhus, Denmark.

64 ²⁸Fisheries and Aquatic Ecosystems Branch, Agri-Food and Biosciences Institute,
65 Belfast, Northern Ireland, United Kingdom.

66 ²⁹CSIC, Instituto de Investigaci3n Mariñas, Eduardo Cabello 6, 36208 Vigo, Spain.

67 ³⁰Institute of Marine Sciences, Middle East Technical University, 33731 Erdemli-Mersin,
68 Turkey.

69 ³¹State Key Laboratory of Estuarine and Coastal Research, East China Normal
70 University, 200241, Shanghai, China.

71 ³²School of Earth and Environmental Sciences, Seoul National University, Seoul 08826,
72 Korea.

73 ³³Marine Biology Station, National Institute of Biology, Fornače 41, 6330 Piran,
74 Slovenia.

75 ³⁴Remote Sensing Laboratory, Institute of Oceanology, Polish Academy of Sciences,
76 Sopot, Poland.

77 ³⁵Department of Geosciences, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka,
78 422-8529, Japan.

79 ³⁶Laboratory of Microbial Ecology, Institute of Biological Sciences, Institute of Ocean
80 and Earth Sciences, Universiti Malaya, 50603 Kuala Lumpur, Malaysia.

81 ³⁷CICECO – Aveiro Institute of Materials, Department of Chemistry, University of Aveiro,
82 Campus de Santiago, 3810-193 Aveiro, Portugal.

83 ³⁸University of New Caledonia and Institute de recherche pour le development (IRD),
84 New Caledonia.

85 ³⁹Consiglio Nazionale delle Ricerche, Istituto per lo studio degli impatti Antropici e
86 Sostenibilità in ambiente marino (CNR-IAS), Loc. Sa Mardini, Torregrande, 09170,
87 Oristano, Italy.

88 ⁴⁰Institut de Ciències del Mar (CSIC), Passeig Marítim de la Barceloneta, 37
89 08003 Barcelona, Spain.

90 ⁴¹Asian School of the Environment, Nanyang Technological University, 639798,
91 Singapore.

92 ⁴²Virginia Commonwealth University, Department of Biology, Richmond, Virginia, United
93 States.

94 ⁴³Centre for Marine Ecosystems Research, School of Science, Edith Cowan University,
95 270 Joondalup Drive, Joondalup 6027 WA, Australia.

96 ⁴⁴Department of Marine Sciences, University of Georgia, Athens, Georgia, United
97 States.

98 ⁴⁵Centro Oceanográfico de Gijón/Xixón, Instituto Español de Oceanografía, Gijón/Xixón,
99 Spain.

100 ⁴⁶College of Marine Science, University of South Florida, Saint Petersburg, Florida
101 33701, United States.

102 ⁴⁷Pacific Northwest National Laboratory, Marine and Coastal Research Laboratory,
103 Sequim, Washington, United States.

104 ⁴⁸Norwegian Institute for Water Research, Oslo, Norway.

105 ⁴⁹Leibniz Institute for Baltic Sea Research Warnemünde, D-18119 Rostock-
106 Warnemünde, Germany.

107 ⁵⁰Marine microbiology, University of Bergen, Norway.

108 ⁵¹Yale School of the Environment, Yale University, New Haven, Connecticut, United
109 States.

110 ⁵²Institute for Marine and Antarctic Studies, University of Tasmania, Private Bag 49,
111 Hobart, Tasmania 7001, Australia

112 ⁵³Biophysics Institute, CNR, Pisa, Italy.

113 ⁵⁴State Key Laboratory of Marine Environmental Science (MEL) & College of Ocean
114 and Earth Sciences, Xiamen University, China.

115 ⁵⁵Departamento de Ecología y Biología Animal, Universidade de Vigo, Centro de
116 Investigación Mariña da Universidade de Vigo (CIM-UVigo), Vigo, Spain.

117 ⁵⁶School of Natural and Environmental Science, Newcastle University, Newcastle upon
118 Tyne, United Kingdom.

119 ⁵⁷Mutsu Institute for Oceanography, Research Institute for Global Change, Japan
120 Agency for Marine-Earth Science and Technology, 690 Kitasekine, Sekine, Mutsu,
121 Aomori, Japan.

122 ⁵⁸Coastal and Estuarine Environment Research Group, Port and Airport Research
123 Institute, Yokosuka 239-0826, Japan.

124 ⁵⁹Key Laboratory of Marine Chemistry Theory and Technology, Ministry of Education,
125 Institute for Advanced Ocean Study, Ocean University of China, Qingdao, Shandong,
126 China.

127 ⁶⁰Faculty of Environmental Earth Science, Hokkaido University, Hokkaido 060-0810,
128 Japan.

129 ⁶¹College of Environment and Safety Engineering, Fuzhou University, China.

130 ⁶²Key Laboratory of Marine Ecology and Environmental Sciences, Institute of
131 Oceanology, Chinese Academy of Sciences, Qingdao, China.

132 ⁶³Fujian Key Laboratory of Marine Carbon Sequestration, Xiamen University, Xiamen
133 361102, China.

134

135 *Corresponding author:

136 E-mail: c.lonborg@ecos.au.dk or clonborg@gmail.com

137

138 **ORCID nr.:**

139 Christian Lønborg: 0000-0001-8380-0238

140 Cátia Carreira: 0000-0002-1520-9320

141 Gwenaël Abril: 0000-0002-4914-086X

142 Susana Agustí: 0000-0003-0536-7293

143 Valentina Amaral: 0000-0002-1088-1484

144 Agneta Andersson: 0000-0001-7819-9038

145 Javier Arístegui: 0000-0002-7526-7741

146 Punyasloke Bhadury: 0000-0001-8714-7475

147 Mariana Bernardi Bif: 0000-0002-2148-4556

148 Alberto V. Borges: 0000-0002-5434-2247

149 Steven Bouillon: 0000-0001-7669-2929

150 Maria LI. Calleja: 0000-0002-5992-2013

151 Luiz C. Cotovicz Jr: 0000-0002-3914-8155

152 Stefano Cozzi: 0000-0003-0116-742X

153 Maryló Doval: 0000-0002-8565-8703

154 Carlos M. Duarte: 0000-0002-1213-1361

155 Bradley Eyre: 0000-0001-5502-0680

156 Cédric G. Fichot: 0000-0002-1099-5764

157 E. Elena García-Martín: 0000-0003-4807-3287

158 Alexandra Garzon-Garcia: 0000-0002-6804-8890

159 Michele Giani: 0000-0002-3306-7725

160 Rafael Gonçalves-Araujo: 0000-0001-8344-8326

161 Renee Gruber: 0000-0002-8788-6910

162 Dennis A. Hansell: 0000-0001-9275-3445

163 Fuminori Hashihama: 0000-0003-3835-7681

164 Ding He: 0000-0001-9620-6115

165 Johnna M. Holding: 0000-0002-7364-0055

166 William R. Hunter: 0000-0001-8801-7947

167 J. Severino P. Ibánhez: 0000-0001-6093-3054
168 Valeria Ibello: 0000-0002-1067-0425
169 Shan Jiang: 0000-0002-1121-6080
170 Guebuem Kim: 0000-0002-5119-0241
171 Katja Klun: 0000-0001-6111-1650
172 Piotr Kowalczyk: 0000-0001-6016-0610
173 Atsushi Kubo: 0000-0002-6457-5386
174 Choon Weng Lee: 0000-0001-9805-9980
175 Cláudia Lopes: 0000-0001-7378-8677
176 Federica Maggioni: 0000-0002-7109-4257
177 Paolo Magni: 0000-0001-5955-6829
178 Celia Marrase: 0000-0002-5097-4829
179 Patrick Martin: 0000-0001-8008-5558
180 S. Leigh McCallister: 0000-0002-9041
181 Roisin McCallum: 0000-0002-0358-2371
182 Patricia M. Medeiros: 0000-0001-6818-2603
183 Xosé Anxelu G. Morán: 0000-002-9823-5339
184 Frank Muller-Karger: 0000-0003-3159-5011
185 Allison Myers-Pigg: 0000-0002-6905-6841
186 Marit Norli: 0000-0001-7472-1562
187 Joanne M. Oakes: 0000-0002-9287-2652
188 Helena Osterholz: 0000-0002-2858-9799
189 Hyekyung Park: 0000-0002-4743-5883
190 Maria Lund Paulsen: 0000-0002-1474-7258
191 Jeff D. Ross: 0000-0002-8659-3833
192 Judith A. Rosentreter: 0000-0001-5787-5682
193 Digna Rueda-Roa: 000-0003-4621-009X
194 Chiara Santinelli: 0000-0002-8921-275X
195 Yuan Shen: 0000-0001-6618-4226
196 Eva Teira: 0000-0002-4333-0101
197 Tinkara Tinta: 0000-0001-6740-8973
198 Guenther Uher: 0000-0001-5105-4445
199 Masahide Wakita: 0000-0002-3333-0546
200 Nicholas Ward: 0000-0001-6174-5581
201 Kenta Watanabe: 0000-0002-0106-3623
202 Yu Xin: 0000-0002-5328-7717
203 Youhei Yamashita: 0000-0002-9415-8743
204 Liyang Yang: 0000-0001-8767-8698
205 Jacob Yeo: 0000-0003-2443-5378
206 Huamao Yuan: 0000-0003-2014-619X
207 Qiang Zheng: 0000-0002-6836-2310
208 Xosé Antón Álvarez-Salgado: 0000-0002-2387-9201

209 **Abstract**

210 ~~M~~The measurements of dissolved organic carbon (DOC), nitrogen (DON), and
211 phosphorus (DOP) concentrations are used to characterize the dissolved organic matter
212 (DOM) pool and are important components of biogeochemical cycling in the coastal
213 ocean. Here, we present the first edition of a global database (CoastDOM v1; available
214 at <https://doi.pangaea.de/10.1594/PANGAEA.964012>) compiling previously published
215 and unpublished measurements of DOC, DON, and DOP ~~collected~~ in coastal waters.
216 These data are complemented by hydrographic data such as temperature and salinity
217 and, to the extent possible, other biogeochemical variables (e.g., Chlorophyll-a, inorganic
218 nutrients) and the inorganic carbon system (e.g., dissolved inorganic carbon and total
219 alkalinity). Overall, CoastDOM v1 includes observations of concentrations from all
220 continents. ~~H~~However, most data were collected in the Northern Hemisphere, with a clear
221 gap in coastal water DOM measurements from the Southern Hemisphere. The data
222 included were collected from 1978 to 2022 and consist of 62339 data points for DOC,
223 20360 for DON, and 13440 for DOP. The number of measurements decreases
224 progressively in the sequence DOC > DON > DOP, reflecting both differences in the
225 maturity of the analytical methods and the greater focus on carbon cycling by the aquatic
226 science community. The global database shows that the average DOC concentration in
227 coastal waters (average ~~±~~ (standard deviation; ~~-~~ (SD): 182 ~~±~~ (314) $\mu\text{mol C L}^{-1}$; median:
228 103 $\mu\text{mol C L}^{-1}$); is 13-fold ~~greathigher~~ than the average coastal DON concentrations
229 (average ~~±~~ (SD): 13.6 ~~±~~ (30.4) $\mu\text{mol N L}^{-1}$; median: 8.0 $\mu\text{mol N L}^{-1}$), which ~~was-is~~ itself 39-
230 fold ~~highgreater~~ than the average coastal DOP concentrations (average ~~±~~ (SD): 0.34 \pm
231 1.11 $\mu\text{mol P L}^{-1}$; median: 0.18 $\mu\text{mol P L}^{-1}$). This dataset will be useful ~~to-for~~ identifying
232 global spatial and temporal patterns in DOM and ~~help to-facilitate~~ ing thee reuse of DOC,
233 DON, and DOP data in studies aimed at better ~~characteris~~ characterizing ~~inge~~ local
234 biogeochemical processes, closing ~~e~~ nutrient budgets, estimating ~~e~~ carbon, nitrogen, and

235 phosphorous pools, as well as ~~establishing identifying~~ a baseline for modelling future
236 changes in coastal waters.

237

238 **Keywords:** Dissolved organic matter, Dissolved organic carbon, Dissolved organic
239 nitrogen, Dissolved organic phosphorus, Coastal waters, Global database.

240 **1. Introduction**

241 Coastal waters are the most biogeochemical dynamic areas of the ocean, exhibiting
242 the highest standing stocks, process rates and transport fluxes of carbon (C), nitrogen
243 (N), and phosphorus (P) per unit area (Bauer et al., 2013; Mackenzie et al., 2011). In
244 these areas, organic matter plays a critical role in numerous biogeochemical processes,
245 serving as both a C, N, and P reservoir and substrate (Carreira et al., 2021).

246 Organic material found in the marine environment is commonly distinguished by its
247 size; material retained on a filter with a pore size typically between 0.2 and 0.7 μm is
248 classified as particulate organic matter (POM), whereas organic matter that passes
249 through the filter is referred to as dissolved organic matter (DOM). This partitioning is
250 operational but has implications for biogeochemical cycling: POM can be suspended in
251 the water column or sink to the sediments controlled by its size, shape and density
252 (Laurenceau-Cornec et al., 2015), whereas DOM is a solute that mostly remains in the
253 water column. In most coastal waters, DOM concentrations are higher than POM, with
254 POM having a larger proportion of known biochemical classes (e.g., carbohydrates,
255 proteins) than the dissolved fraction, suggesting that generally, DOM is more reworked
256 and recalcitrant ~~In most coastal waters, the DOM concentrations are greater than POM,~~
257 ~~with the POM fraction being less degraded and more bioavailable~~ (Boudreau and
258 Ruddick, 1991; Lønborg et al., 2018; Benner and Amon, 2015).

259 The DOM pool consists mainly of C (DOC), N (DON), and P (DOP) but it also includes
260 other elements such as oxygen, sulphur and trace elements (Lønborg et al., 2020). In
261 coastal waters, DOM originates from multiple sources. Internal, or autochthonous,
262 sources include planktonic organisms (Lønborg et al., 2009; Carlson and Hansell, 2015),
263 benthic microalgae, macrophytes, and sediment porewater (Burdige and Komada, 2014;
264 Wada et al., 2008). On the other hand, DOM from external, or allochthonous, sources,
265 has mainly terrestrial origins, including wetlands, river and surface runoff, groundwater

266 discharges, and atmospheric deposition (Lavorivska et al., 2016; Raymond and Spencer,
267 2015; Taniguchi et al., 2019; Santos et al., 2021). The main sinks for DOM from the water
268 column in coastal waters are: 1) bubble coagulation and abiotic flocculation (Kerner et al.,
269 2003) or sorption to particles (Chin et al., 1998); 2) sunlight-mediated photodegradation
270 (Mopper et al., 2015); and 3) microbial degradation by mainly heterotrophic prokaryotes
271 (Lønborg and Álvarez-Salgado, 2012).

272 Given the importance of DOM as a source of nutrients and for coastal biogeochemical
273 cycling in general, numerous studies have measured the C, N and P content of the DOM
274 pool over the last few decades (e.g., García-Martín et al., 2021; Cauwet, 2002; Osterholz
275 et al., 2021). Most data, however, are often unavailable or stored in an inaccessible
276 manner, making it difficult to e.g., analyse global spatial and temporal patterns effectively.

277 ~~A~~ global open ocean DOM data compilation for DOC, total dissolved nitrogen (TDN),
278 DON (Hansell et al., 2021) and DOP (Liang et al., 2022; Karl and Björkman, 2015) already
279 exists, and contains few coastal samples (< 200m) (Hansell et al., 2021), but there are
280 no compilation specifically -focused on coastal waters. Hence, there is a clear need for a
281 comprehensive global and integrated database of DOC, DON and DOP measurements
282 for coastal waters. To address this need, we have prepared the first edition of a coastal
283 DOM database (named CoastDOM v1), by compiling both previously reported as well as
284 unpublished data. These data have been obtained from authors of the original studies or
285 extracted directly from the original studies. In order to allow the DOM measurements to
286 be interpreted across larger scales, and to better understand their relationship with local
287 environmental conditions, we have included concurrently collected ancillary data (such
288 as physical and/or chemical seawater properties) whenever available. The objective of
289 this database is multifaceted. Firstly, we aimed to compile all available coastal DOM data
290 into a single repository. Secondly, our intention was to make these data easily accessible
291 to the research community and thirdly, we sought to achieve long-term consistency of the

292 measurements, to enable data intercomparison, and establish a robust baseline for
293 assessing, for example, the impacts of climate change and land use changes.

294 **2. Methods**

295 **2.1. Data compilation**

296 The measurements included in CoastDOM v1 were obtained either directly from
297 authors of previously published studies, online databases, or scientific papers. An
298 extensive search of published reports, Ph.D. theses, and peer-reviewed literature was
299 performed to identify studies dealing with DOM in coastal waters. First, a formal search
300 was performed using Google Scholar in January 2022 using the search terms “dissolved
301 organic carbon”, “dissolved organic nitrogen”, and “dissolved organic phosphorus” in
302 connection with “marine” or “ocean”, which yielded a total of 897 articles (after filtering
303 the query by searching content in the title and abstract and excluding non-coastal
304 articles). When data could not be obtained directly from the corresponding authors,
305 relevant data were extracted. Further searches for relevant datasets were conducted
306 using the reference lists of the identified scientific papers as well as databases and
307 repositories to capture as many datasets as possible. Additionally, research groups that
308 were invited to participate in this effort were also encouraged to submit unpublished data
309 to CoastDOM v1.

310

311 **2.2. Dissolved organic matter analysis**

312 The DOC concentrations included in CoastDOM v1 were commonly measured using a
313 total organic carbon (TOC) high-temperature catalytic oxidation (HTCO) analyser (81%
314 of samples; (Sharp et al., 1993). Some were measured by a combined wet chemical
315 oxidation (WCO) step and/or UV digestion, after which the carbon dioxide generated was
316 quantified (19% of samples). Similarly, concentrations of total dissolved nitrogen (TDN;

317 ~~(~~Sipler and Bronk, 2015) were determined using either a nitric oxide chemiluminescence
318 detector connected in series with the HTOCO analyser used for DOC analyses (31% of the
319 samples), or by employing a UV and/or chemical oxidation step (69%). In the latter
320 approach, both organic and inorganic N compounds were oxidised to nitrate, which was
321 subsequently quantified through a colorimetric method to determine the concentration of
322 inorganic N (Valderrama, 1981; Álvarez-Salgado et al., 2023; Halewood et al., 2022;
323 Foreman et al., 2019). Another method used for DON determination is oxidizing the
324 sample and measuring the resulting total nitrate by the nitric oxide chemiluminescence
325 method (Knapp et al., 2005). However, none of the concentration measurements included
326 in CoastDOM v1 applied this method. The reported DON concentrations were calculated
327 as the difference between TDN and dissolved inorganic nitrogen (DIN; sum of ammonium
328 (NH_4^+) and nitrate/nitrite ($\text{NO}_3^- + \text{NO}_2^-$); $\text{DON} = \text{TDN} - \text{DIN}$) (Álvarez-Salgado et al., 2023).
329 Analyses of total dissolved phosphorus (TDP) were determined by UV (4%) or wet
330 chemical oxidation (66%), or a combination of these (30%), and subsequently were
331 analysed for inorganic phosphorus by a colorimetric method (Álvarez-Salgado et al.,
332 2023). Another method also previously used for TDP analysis is the ash/hydrolysis
333 method (Solorzano and Sharp, 1980), even though none of the data included in
334 CoastDOM v1 used this method. The DOP concentrations were calculated as the
335 difference between TDP and soluble reactive phosphorus (SRP: HPO_4^{2-}) ($\text{DOP} = \text{TDP} -$
336 SRP) (Álvarez-Salgado et al., 2023).

Formatted: Font: English (United States)

337

338 3. Description of the dataset

339 The data compiled in CoastDOM v1 were collected, analysed and processed by different
340 laboratories, however, all data included have undergone quality control measures, either
341 by using reference samples or internal quality assurance procedures. While many of the
342 included DOC and TDN data have been systematically compared against consensus

343 reference material (CRM) mainly provided by the University of Miami's CRM program
344 (Hansell, 2005), there is a limitation in CoastDOM v1 regarding the intercalibration across
345 different measurement systems used for both DOP and DON determination. While the
346 CRM could be used for DOC, DON and DOP measurements, this has not yet been
347 attempted for DOP and measurement uncertainties increase in the sequence DOC >
348 DON > DOP. Although some of the reported measurements have quantified the DOP
349 recovery based on commercially available DOP compounds such as Adenosine
350 triphosphate (ATP), it is not known if these were conducted systematically in all cases.

351 Therefore, we strongly recommend undertaking further intercalibrations across
352 laboratories for future measurements of TDP, as has been done for DOC and TDN
353 measurements (e.g., Sharp et al., 2002). Since additional quality control is not possible
354 in retrospect, we assessed the quality of CoastDOM v1 based on its internal consistency.

355 In CoastDOM v1, we defined "coastal water" as encompassing estuaries (salinity >
356 0.1) to the continental shelf break (water depth < 200 m). However, some locations, such
357 as deep fjords which are close to the coast cannot be classed as coastal due to
358 bathymetry (deeper than > 200 m). Therefore, we evaluated the inclusion of some
359 datasets on a case-by-case basis. For inclusion in the database, each DOM
360 measurement needed at a minimum to contain the following information (if reported in the
361 original publication or otherwise available):

362

- 363 - Country where samples were collected
- 364 - Latitude of measurement (in decimal units)
- 365 - Longitude of measurement (in decimal units)
- 366 - Year of sampling
- 367 - Month of sampling
- 368 - Sampling day (when available)

- 369 - Depth (m) at which the discrete samples were collected
- 370 - Temperature (°C) of the sample
- 371 - Salinity of the sample
- 372 - Dissolved organic carbon (DOC) concentration ($\mu\text{mol L}^{-1}$)
- 373 - Method used to measure DOC concentration
- 374 - DOC - QA flag: Quality flag for DOC measurement
- 375 - Dissolved organic nitrogen (DON) concentration ($\mu\text{mol L}^{-1}$)
- 376 - Total dissolved nitrogen (TDN) concentration ($\mu\text{mol L}^{-1}$)
- 377 - Method used to measure TDN concentration
- 378 - TDN - QA flag: Quality flag for TDN measurement
- 379 - Dissolved organic phosphorus (DOP) concentration ($\mu\text{mol L}^{-1}$)
- 380 - Total dissolved phosphorus (TDP) concentration ($\mu\text{mol L}^{-1}$)
- 381 - Method used to measure TDP concentration
- 382 - TDP - QA flag: Quality flag for TDP measurement
- 383 - Responsible person
- 384 - Originator institution
- 385 - Contact of data originator

386

387 It should be noted that in all entries, at least DOC, DON or DOP should have been

388 measured. In addition, we also included other relevant data, when available, in the

389 CoastDOM v1 dataset:

- 390
- 391 - Depth at the station where the sample was collected (Bottom depth, m).
- 392 - Total suspended solids (TSS) concentration (mg L^{-1})
- 393 - Chlorophyll-a (Chl a) concentration ($\mu\text{g L}^{-1}$)
- 394 - Chl a - QA flag: Quality flag for chlorophyll-a measurement

- 395 - Sum of nitrate and nitrite ($\text{NO}_3^- + \text{NO}_2^-$) concentration ($\mu\text{mol L}^{-1}$)
- 396 - $\text{NO}_3^- + \text{NO}_2^-$ - QA flag: Quality flag for $\text{NO}_3^- + \text{NO}_2^-$ measurement
- 397 - Ammonium (NH_4^+) concentration ($\mu\text{mol L}^{-1}$)
- 398 - NH_4^+ - QA flag: Quality flag for NH_4^+ measurement
- 399 - Soluble reactive phosphorus (HPO_4^{2-}) concentration ($\mu\text{mol L}^{-1}$)
- 400 - HPO_4^{2-} - QA flag: Quality flag for HPO_4^{2-} measurement
- 401 - Particulate organic carbon (POC) concentration ($\mu\text{mol L}^{-1}$)
- 402 - Method used to measure POC concentration
- 403 - POC - QA flag: Quality flag for POC measurement
- 404 - Particulate nitrogen (PN) concentration ($\mu\text{mol L}^{-1}$)
- 405 - Method used to measure PN concentration
- 406 - PN - QA flag: Quality flag for PN measurement
- 407 - Particulate phosphorus (PP) concentration ($\mu\text{mol L}^{-1}$)
- 408 - Method used to measure PP concentration
- 409 - PP - QA flag: Quality flag for PP measurement
- 410 - Dissolved inorganic carbon (DIC) concentration ($\mu\text{mol kg}^{-1}$)
- 411 - DIC - QA flag: Quality flag for DIC measurement
- 412 - Total alkalinity (TA) concentration ($\mu\text{mol kg}^{-1}$)
- 413 - TA - QA flag: Quality flag for TA measurement

414

415 Quality control of large datasets is crucial to ensure their reliability and usefulness.

416 Thus, we have not included data that were deemed compromised, such as records that
 417 had not gone through quality control by the data originators. We also accepted a certain
 418 degree of measurement error since multiple groups have been involved in the collection,
 419 analysis, and/or compilation of the information. Some of these errors were corrected (e.g.,
 420 when a value was placed in a wrong column, or clearly inaccurate locations were

421 reallocated for consistency with the place of study), while others could not be rectified
422 (e.g., values showing clear signs of contamination) and were consequently excluded from
423 CoastDOM v1 (Fig. 1). It should also be noted that differences in analytical capabilities
424 between laboratories and individual measurement campaigns likely caused additional
425 uncertainty. Outliers, arising for example from contamination, were removed from the
426 dataset. The data were moreover screened for zero values (i.e., concentrations below the
427 detection limit or absence of data). In cases where concentrations were below the
428 detection limit, the zero values were replaced with half the value of the limit-of-detection.

429 Commonly reported detection limits are reach $-4 \mu\text{mol L}^{-1}$ for DOC, $\rightarrow 0.3 \mu\text{mol L}^{-1}$ for
430 DON and are $\sim 0.03 \mu\text{mol L}^{-1}$ for DOP.

431 To ensure the inclusion of only high-quality data, we only accepted entries with specific
432 World Ocean Circulation Experiment (WOCE) quality codes: “2- Acceptable
433 measurement” and “6- Mean of replicate measurements”. In our quality control
434 assessments, we carefully avoided overly strict criteria, known as “data grooming”, which
435 could potentially overlook genuine patterns and changes in the dataset that may be
436 significant over longer temporal and/or wider spatial scales. Coastal waters are known to
437 exhibit a wide range of environmental concentrations, influenced by factors such as
438 seasonality and local anthropogenic activities. Consequently, these data points may
439 encompass a wide concentration range. However, obtaining consistent long-term
440 datasets is important to enable data intercomparison, and establish a robust baseline.
441 Such long-term consistency can be achieved by using the CRM standards provided by
442 the Hansell laboratory for DOC and TDN. Another helpful approach is comparing the
443 DOM concentrations obtained by different laboratories in the same study area and time
444 of year.

445
446 **3.1 Summary of dissolved organic carbon (DOC) concentration observations**

447 Measurements of DOC concentrations were conducted between 1978 to 2022, with a
448 total of 62339 individual data points (Table 1). The DOC concentrations ranged from 17
449 to 30327 $\mu\text{mol C L}^{-1}$ (average ~~\pm~~ (Standard Deviation; ~~\pm~~ (SD): 182 ~~\pm~~ (314) $\mu\text{mol C L}^{-1}$;
450 median: 103 $\mu\text{mol C L}^{-1}$; Table 1). The majority (53%) of the concentrations fell within the
451 range of 60 to 120 $\mu\text{mol C L}^{-1}$ (Fig. 24). A large number of DOC concentration
452 observations (17%) ranged between 300 and 600 $\mu\text{mol C L}^{-1}$, which were predominantly
453 collected in eutrophic and river-influenced coastal waters of the Northern Hemisphere,
454 such as the Baltic Sea (Fig. 24). It was observed that 75% of the DOC concentrations
455 were higher than 77 $\mu\text{mol C L}^{-1}$, while 25% of the measurements surpassed 228 $\mu\text{mol C}$
456 L^{-1} (Table 1).

457 Coastal environments that experience minimal continental runoff, such as Palmer
458 Station in Antarctica, typically exhibit low DOC concentrations. On the other hand, coastal
459 waters heavily influenced by humic-rich terrigenous inputs, such as the Sarawak region
460 in Malaysia, tended to have high DOC concentrations. In addition, some extremely high
461 DOC concentrations were measured in the Derwent River in Australia which is impacted
462 by paper mill effluents. There has been a large increase in the number of DOC
463 concentration observations after 1992 (Fig. 32), and those measurements were from a
464 wide range of locations. However, these concentration observations were not evenly
465 distributed around the globe, with the Southern Hemisphere being ~~relatively~~ under-
466 sampled (10% of observations), especially in the African, South American and Antarctic
467 continents (Fig. 32, 43).

468

469 3.2. Summary of dissolved organic nitrogen (DON) concentration observations

470 The DON concentration measurements were collected between 1990 and 2021, with
471 a total of 20357 data points (Table 1). Concentrations of DON ranged from < 0.1 to 2095.3
472 $\mu\text{mol N L}^{-1}$ (average ~~\pm~~ (SD): 13.6 ~~\pm~~ (30.4) $\mu\text{mol N L}^{-1}$; median: 8.0 $\mu\text{mol N L}^{-1}$; Table 1),

473 with the most common range (42%) for DON concentrations between 4 to 8 $\mu\text{mol N L}^{-1}$
474 (Fig. 42). Overall, 75% of DON concentrations were above 5.5 $\mu\text{mol N L}^{-1}$, while 25%
475 were above 15.8 $\mu\text{mol N L}^{-1}$ (Table 1).

476 The lowest DON concentrations were recorded in Young Sound, Greenland, which
477 receives direct run-off from the Greenland Ice Sheet, whereas the highest concentrations
478 were detected during a flood event in the Richmond River Estuary, Australia. Since 1995,
479 there has been a large increase in the number of DON measurements conducted in
480 coastal waters globally (Fig. 23); however, the majority of those measurements have been
481 in the Northern Hemisphere (79% of observations), mostly in Europe and the United
482 States (Figs. 23, 43).

484 3.3. Summary of dissolved organic phosphorus (DOP) concentration 485 observations

486 CoastDOM v1 includes a total of 13534 DOP measurements, collected between 1990
487 and 2021 (Table 1). Overall, DOP concentrations ranged from < 0.10 to $84.27 \mu\text{mol P L}^{-1}$
488 (average \pm (SD): $0.34 \pm 1.11 \mu\text{mol P L}^{-1}$; median: $0.18 \mu\text{mol P L}^{-1}$; Table 2). The
489 majority (74%) of DOP concentrations were below $0.30 \mu\text{mol P L}^{-1}$ (Fig. 42). Analysis of
490 the DOP dataset revealed that 75% of the concentrations were above $0.11 \mu\text{mol P L}^{-1}$,
491 while 25% were above $0.30 \mu\text{mol P L}^{-1}$ (Table 1).

492 The lowest DOP concentrations were measured off the Kimberley Coast in Australia,
493 while the highest concentrations were found in the Vasse-Wonnerup Estuary in the South
494 west region of Australia. Similarly to DOC and DON, most of the DOP measurements
495 have been conducted from the 1990s onwards, with a predominant focus in the Northern
496 Hemisphere (70% of observations), particularly in Europe and the United States (Figs.
497 32, 43).

3.4. Summary of dissolved organic matter (DOM) concentration observations

In CoastDOM v1 the number of measurements decreases progressively in the sequence DOC > DON > DOP (62339, 20357, and 13534, respectively), reflecting both differences in the maturity of the analytical methods and the greater focus on carbon cycling by the aquatic science community. In addition, the average DOC concentration in coastal waters (182 ± 314 $\mu\text{mol C L}^{-1}$), was 13-fold ~~great~~higher than the average coastal DON concentrations 13.6 ± 30.4 $\mu\text{mol N L}^{-1}$), which was itself 39-fold higher than the average coastal DOP concentrations (0.34 ± 1.11) $\mu\text{mol P L}^{-1}$) (Table 1). Interestingly the coefficient of variation (C.V. - dispersion of the data around the mean) increased from DOC (173%) to DON (224%) and DOP (326%), which is related to the fact that the % contribution of refractory organic material decreases in the same sequence (Table 1). It should be noted that CoastDOM v1 only contains 7058 paired measurements of DOC, DON, and DOP, and therefore only a subset of observations reported all three element pools. The average C: N: P stoichiometry for these paired DOM measurements was 1171 (± 4248): 100 (± 580): 1 (Table 1), which was very N- and P- depleted compared to the Redfield Ratio (Redfield et al., 1963). -However, the large variations in C:N, C:P and N:P ratios reveals large variations in the composition of the DOM pool in coastal waters.

3.5. Potential use of the dataset

The use of the CoastDOM v1 dataset should be accompanied by the citation of this paper and the inclusion of the correct doi-reference. CoastDOM v1 is available in full open access on the PANGEA homepage ~~after acceptance of the manuscript, where it will be available~~ as a *.csv file. The dataset includes a brief description of the metadata and methods employed, with emphasis on measurement techniques and data units. We chose the terminology most familiar to the ocean science community. It is important to note that all data included in CoastDOM v1, as well as this manuscript, are considered

public domain; as such, a subset of this global dataset ~~is may~~ also ~~available be present~~ in previous data compilations (e.g., Hansell et al., 2021). The list of citations and links referenced in CoastDOM v1 also provide users with information ~~as to on~~ how these data ~~have~~ been previously used in publications or databases.

3.6. Recommendations and conclusions

In CoastDOM v1, we have compiled available coastal DOM data in a single repository, making it openly and freely available to the research community. This compilation has established a consistent global dataset, serving as a valuable information source to investigate a variety of environmental questions and to explore spatial and temporal trends. We suggest a set of recommendations for the future expansion of this global dataset. ~~Firstly,~~ our analysis highlights a spatial bias, with a concentration of sampling efforts and/or data availability predominantly concentrated in the Northern Hemisphere. The data gap in coastal DOM measurements in the Southern Hemisphere needs to be addressed to provide a more representative global understanding of the role of DOM in coastal water biogeochemistry. Additionally, increased sampling efforts especially around ~~in the African and South American,~~ and island nations ~~-continents~~ are warranted due to the vulnerability of many coastal areas to climate change and intensifying human activities, which will undoubtedly impact DOM biogeochemistry. Further more, ~~it is also worth noting that~~ there are is comparatively few data from coastal waters affected by river discharge into the tropics, e.g., the Amazon, and Indian and Indonesian rivers that together dominate freshwater inputs to the coastal ocean. ~~Secondly,~~ there is a need for more comprehensive temporal and spatial datasets to capture the variability of DOM ~~concentrations levels~~ in highly dynamic and productive coastal systems. Focused efforts should be made to resolve these temporal and spatial changes. Third, only a fraction of data entries report paired DOC, DON and DOP measurements, we encourage that these

551 be measured and reported together in order to better determine changes in stoichiometry
552 and composition. Fourth~~Thirdly, it is also important to collect~~ing and reporting ancillary
553 data, such as temperature, salinity, nutrient measurements, and particulate components,
554 is important to provide context and better understand the underlying processes driving
555 the observed DOM concentrationslevels. Fifth, studies need to collect a minimum of
556 metadata and report it in standardized manner. Lastly, we ~~strongly~~ recommend ~~that the~~
557 ~~DOM research community conducts~~ regular inter-calibration exercises to establish
558 standardised and interoperable methods and data, particularly for DON and DOP
559 measurements. This will ensure the comparability and reliability of data across different
560 studies and enhance our understanding of DON and DOP dynamics in coastal waters.

561 In light of ongoing global environmental changes, the mobilisation and open sharing of
562 existing data ~~of~~for important biogeochemical variables, such as the DOM pool, are crucial
563 for establishing baselines and determining global trends and changes in coastal waters.
564 The aim is to publish an updated version of the database periodically to determine global
565 trends of DOM levels in coastal waters, and we therefore encourage researchers to
566 submit new data to the corresponding author. The CoastDOM v1 dataset was developed
567 according to the FAIR principles regarding Findability, Accessibility, Interoperability and
568 Reusability of data. Thus, CoastDOM v1 will serve as a reliable open-source information
569 resource, enabling in-depth analyses and providing quality-controlled input data for large
570 -scale ecosystem models.

572 4. Data availability

573 ~~The dataset is available for the review process at Figshare~~
574 ~~<https://figshare.com/s/512289eb43c4f8e8eae0f>.~~The dataset is available at furthermore
575 submitted to the PANGEA database
576 (<https://doi.pangaea.de/10.1594/PANGAEA.964012>; and is currently waiting to be

577 ~~assigned a Doi number~~ (Lønborg et al., 2023). The file ~~is will be~~ available as a *.csv
578 merged file and ~~is will be~~ available in full open access ~~in the PANGAEA database after~~
579 ~~acceptance of the manuscript~~.

Commented [CL1]: update

581 **Competing interests**

582 The authors declare no competing interests.

583 **Author Contribution**

584 C.L., C.C., and X.A.A-S started the initiative and finalised the data compilation. All co-
585 authors contributed data. C.L. wrote the manuscript with input from all co-authors.

586 **Acknowledgement**

587 Qi Chen is acknowledged for his very skilful help with Figure ~~43~~.

588 **Funding**

589 During the drafting of the manuscript C.L. received funding from the Independent
590 Research Fund Denmark Grant No. 1127-00033B. The monitoring data obtained from
591 Bermuda received funding from the Bermuda Government Department of Environment
592 and Natural Resources. A subset of the data obtained from UK coastal estuaries received
593 funding from the Natural Environment Research Council (Grant NE/N018087/1). Data
594 retrieved from the Palmer LTER data were collected with support from the Office of Polar
595 Programs, US National Science Foundation. Data obtained from the Great Barrier Reef
596 Marine Monitoring Program for Inshore Water Quality, which is a partnership between the
597 Great Barrier Reef Marine Park Authority, the Australian Institute of Marine Science,
598 James Cook University, and the Cape York Water Partnership. The contribution by P.K.
599 was supported by DiSeDOM project contract no. UMO-2019/33/B/ST10/01232 funded by
600 the NCN - National Science Centre, Poland. N.W. and A.M.-P. participated in this

601 synthesis effort with funding provided by the U.S. Department of Energy funded
602 COMPASS-FME project; the provided data was collected with funding from the PREMIS
603 Initiative, conducted under the Laboratory Directed Research and Development Program
604 at Pacific Northwest National Laboratory. The data obtained from the Levantine Sea (Med
605 Sea) received funding from the Scientific and Technological Research Council of Türkiye
606 (TÜBİTAK, 1001 program, Grant 115Y629). The data obtained from Gulf of Trieste
607 (Slovenian waters) were financed by Research Program No. P1-0237 (Slovenian
608 Research and Innovation Agency). Data obtained from the northern Baltic Sea were
609 financed by the research program EcoChange (Swedish research council FORMAS). The
610 contribution by D.H. was supported by National Natural Science Foundation of China
611 (42222061), and funding support from the Center for Ocean Research in Hong Kong and
612 Macau (CORE). CORE is a joint research center for ocean research between Laoshan
613 Laboratory and HKUST. The contribution by Y.S. was supported by National Natural
614 Science Foundation of China (42106040), the Fundamental Research Funds for the
615 Central Universities of China (20720210076), and Fujian Provincial Central Guided Local
616 Science and Technology Development Special Project (Grant No. 2022L3078). The data
617 provided by LCCJr were supported by the Brazilian National Council of Research and
618 Development (CNPq-Pve No. 401.726/2012-6) and by the Carlos Chagas Foundation for
619 Research Support of the State of Rio de Janeiro (FAPERJ; No. E-26202.785/2016). Data
620 provided by SC were collected in the framework of Italian (PRISMA 1 and 2, ANOCSIA
621 and VECTOR) and European (OCEANCERTAIN) research projects. Data provided by
622 MG were collected within the Project “Mucilages in the Adriatic and Tyrrhenian Seas
623 (MAT)”, coordinated by the Istituto Centrale per la Ricerca Scientifica e Tecnologica
624 Applicata al Mare and financially supported by the Italian Ministry of the Environment.
625 Data obtained from the Georgia coast (USA) were supported by the National Science
626 Foundation through grants OCE-1832178 (GCE-LTER Program) and OCE-1902131.

627 [Observations in the southern Caribbean Sea including the Cariaco Basin were collected](#)
628 [as part of the CARIACO Ocean Time Series program \(supported by the Consejo Nacional](#)
629 [de Ciencia y Tecnología of Venezuela, the Ley de Ciencia, Tecnología e Innovación de](#)
630 [Venezuela, the Estación de Investigaciones Marinas de Venezuela; the National Science](#)
631 [Foundation grants OCE-0752139, OCE-9216626, OCE-9729284, OCE-491 9401537,](#)
632 [OCE-9729697, OCE-9415790, OCE-9711318, OCE-0326268, OCE-0963028, OCE-](#)
633 [0326313, and OCE-0326268, NASA grants NAG5-6448, NAS5-97128, and](#)
634 [NNX14AP62A; the Inter-American Institute for Global Change Research grant IAI-](#)
635 [CRN3094\), and the Marine Biodiversity Observation Network/MBON of the Group on](#)
636 [Earth Observations Biodiversity Observation Network\). Data provided by D.J.R. and](#)
637 [B.D.E. was supported by ARC Linkage \(LP0770222\), with Norske-Skog Boyer and the](#)
638 [Derwent Estuary Program providing financial and in-kind assistance.](#)

640 **References**

- 641 Álvarez-Salgado, X. A., Nieto-Cid, M., and Rossel, P. E.: Dissolved Organic Matter, in:
642 Marine Analytical Chemistry, edited by: Blasco, J., and Tovar-Sánchez, A., Springer
643 International Publishing, Cham, 39-102, 10.1007/978-3-031-14486-8_2, 2023.
- 644 Bauer, J. E., Cai, W. J., Raymond, P. A., Bianchi, T. S., Hopkinson, C. S., and Regnier,
645 P. A.: The changing carbon cycle of the coastal ocean, Nature, 504, 61-70,
646 10.1038/nature12857, 2013.
- 647 Benner, R. and Amon, R. M.: The size-reactivity continuum of major bioelements in the
648 ocean, Annual review of marine science, 7, 185-205, 10.1146/annurev-marine-
649 010213-135126, 2015.
- 650 Boudreau, B. P. and Ruddick, B. R.: On a reactive continuum representation of organic
651 matter diagenesis, American Journal of Science, 291, 507-538, 1991.
- 652 Burdige, D. J. and Komada, T.: Sediment pore waters, in: Biogeochemistry of marine
653 dissolved organic matter, edited by: Hansen, D. A., and Carlson, C. A., Elsevier, 535-
654 577, 2014.
- 655 Carlson, C. A. and Hansell, D. A.: DOM Sources, Sinks, Reactivity, and Budgets, in:
656 Biogeochemistry of Marine Dissolved Organic Matter, edited by: Carlson, C. A., and
657 Hansell, D. A., Elsevier Science & Technology, 65-126, 10.1016/b978-0-12-405940-
658 5.00003-0, 2015.
- 659 Carreira, C., Talbot, S., and Lønborg, C.: Bacterial consumption of total and dissolved
660 organic carbon in the Great Barrier Reef, Biogeochemistry, 10.1007/s10533-021-
661 00802-x, 2021.
- 662 Cauwet, G.: DOM in coastal areas, in: Biogeochemistry of Dissolved organic matter,
663 edited by: Hansell, D., and Carlson, C. A., Academic Press, London, 579-609, 2002.

665 Chin, W. C., Orellana, M. V., and Verdugo, P.: Spontaneous assembly of marine
666 dissolved organic matter into polymer gels, *Nature*, 391, 568-572, 1998.

667 Foreman, R. K., Björkman, K. M., Carlson, C. A., Opalk, K., and Karl, D. M.: Improved
668 ultraviolet photo-oxidation system yields estimates for deep-sea dissolved organic
669 nitrogen and phosphorus, *Limnology and Oceanography: Methods*, 17, 277-291,
670 <https://doi.org/10.1002/lom3.10312>, 2019.

671 García-Martín, E. E., Sanders, R., Evans, C. D., Kitidis, V., Lapworth, D. J., Rees, A. P.,
672 Spears, B. M., Tye, A., Williamson, J. L., Balfour, C., Best, M., Bowes, M., Breimann,
673 S., Brown, I. J., Burden, A., Callaghan, N., Felgate, S. L., Fishwick, J., Fraser, M.,
674 Gibb, S. W., Gilbert, P. J., Godsell, N., Gomez-Castillo, A. P., Hargreaves, G., Jones,
675 O., Kennedy, P., Lichtschlag, A., Martin, A., May, R., Mawji, E., Mounteney, I.,
676 Nightingale, P. D., Olszewska, J. P., Painter, S. C., Pearce, C. R., Pereira, M. G.,
677 Peel, K., Pickard, A., Stephens, J. A., Stinchcombe, M., Williams, P., Woodward, E.
678 M. S., Yarrow, D., and Mayor, D. J.: Contrasting Estuarine Processing of Dissolved
679 Organic Matter Derived From Natural and Human-Impacted Landscapes, *Global
680 Biogeochemical Cycles*, 35, e2021GB007023,
681 <https://doi.org/10.1029/2021GB007023>, 2021.

682 Halewood, E., Opalk, K., Custals, L., Carey, M., Hansell, D. A., and Carlson, C. A.:
683 Determination of dissolved organic carbon and total dissolved nitrogen in seawater
684 using High Temperature Combustion Analysis, *Frontiers in Marine Science*, 9,
685 10.3389/fmars.2022.1061646, 2022.

686 Hansell, D. A.: Dissolved Organic Carbon Reference Material Program, *Eos,
687 Transactions American Geophysical Union*, 86, 318-318,
688 <https://doi.org/10.1029/2005EO350003>, 2005.

689 Hansell, D. A., Carlson, C. A., Amon, R. M. W., Álvarez-Salgado, X. A., Yamashita, Y.,
690 Romera-Castillo, C., and Bif, M. B.: Compilation of dissolved organic matter (DOM)
691 data obtained from global ocean observations from 1994 to 2021. Version 2. (NCEI
692 Accession 0227166) [dataset], doi.org/10.25921/s4f4-ye35, 2021.

693 Iavorivska, L., Boyer, E. W., and DeWalle, D. R.: Atmospheric deposition of organic
694 carbon via precipitation, *Atmospheric Environment*, 146, 153-163,
695 <https://doi.org/10.1016/j.atmosenv.2016.06.006>, 2016.

696 Karl, D. M. and Björkman, K. M.: Dynamics of Dissolved Organic Phosphorus, in:
697 *Biogeochemistry of marine dissolved organic matter*, edited by: Hansell, D. A., and
698 Carlson, C. A., 233-334, 10.1016/b978-0-12-405940-5.00005-4, 2015.

699 Kerner, M., Hohenberg, H., Ertl, S., Reckermann, M., and Spitzky, A.: Self-organization
700 of dissolved organic matter to micelle-like microparticles in river water, *Nature*, 422,
701 150-154, 2003.

702 Knapp, A. N., Sigman, D. M., and Lipschultz, F.: N isotopic composition of dissolved
703 organic nitrogen and nitrate at the Bermuda Atlantic Time-series Study site, *Global
704 Biogeochemical Cycles*, 19, <https://doi.org/10.1029/2004GB002320>, 2005.

705 Laruelle, G. G., Dürr, H. H., Lauerwald, R., Hartmann, J., Slomp, C. P., Goossens, N.,
706 and Regnier, P. A. G.: Global multi-scale segmentation of continental and coastal
707 waters from the watersheds to the continental margins, *Hydrol. Earth Syst. Sci.*, 17,
708 2029-2051, 10.5194/hess-17-2029-2013, 2013.

709 Laurenceau-Cornec, E. C., Trull, T. W., Davies, D. M., De La Rocha, C. L., and Blain,
710 S.: Phytoplankton morphology controls on marine snow sinking velocity, *Marine
711 Ecology Progress Series*, 520, 35-56, 2015.

712 Liang, Z., McCabe, K., Fawcett, S. E., Forrer, H. J., Hashihama, F., Jeandel, C.,
713 Marconi, D., Planquette, H., Saito, M. A., Sohm, J. A., Thomas, R. K., Letscher, R. T.,
714 and Knapp, A. N.: A global ocean dissolved organic phosphorus concentration
715 database (DOPv2021), *Scientific Data*, 9, 772, 10.1038/s41597-022-01873-7, 2022.

716 Lønborg, C. and Álvarez-Salgado, X. A.: Recycling versus export of bioavailable
717 dissolved organic matter in the coastal ocean and efficiency of the continental shelf
718 pump, *Global biogeochemical cycles* 26, GB3018, 10.1029/2012GB004353, 2012.

719 Lønborg, C., Álvarez-Salgado, X. A., Davidson, K., and Miller, A. E. J.: Production of
720 bioavailable and refractory dissolved organic matter by coastal heterotrophic
721 microbial populations, *Estuarine, Coastal and Shelf science*, 82, 682–688,
722 10.1016/j.ecss.2009.02.026, 2009.

723 Lønborg, C., Álvarez-Salgado, X. A., Duggan, S., and Carreira, C.: Organic matter
724 bioavailability in tropical coastal waters: The Great Barrier Reef, *Limnology and
725 Oceanography*, 63, 1015-1035, 10.1002/lno.10717, 2018.

726 Lønborg, C., Carreira, C., Jickells, T., and Álvarez-Salgado, X. A.: Impacts of Global
727 Change on Ocean Dissolved Organic Carbon (DOC) Cycling, *Frontiers in Marine
728 Science*, 7, 466, 10.3389/fmars.2020.00466, 2020.

729 Lønborg, C., Carreira, C., Abril, G., Agustí, S., Amaral, V., Andersson, A., Arístegui, J.,
730 Bhadury, P., Bif, M. B., Borges, A. V., Bouillon, S., Calleja, M. L., Cotovicz Jr, L. C.,
731 Cozzi, S., Doval, M. D., Duarte, C. M., Eyre, B. D., Fichot, C. G., García-Martín, E.
732 E., Garzon-Garcia, A., Giani, M., Gonçalves-Araujo, R., Gruber, R., Hansell, D. A.,
733 Hashihama, F., He, D., Holding, J. M., Hunter, W. R., Ibánhez, J. S., Ibello, V., Jiang,
734 S., Kim, G., Klun, K., Kowalczyk, P., Kubo, A., Weng Lee, C., Lopes, C., Maggioni,
735 F., Magni, P., Marrase, C., Martin, P., McCallister, S. L., McCallum, R., Medeiros, P.
736 M., Morán, X. A. G., Muller-Karger, F., Myers-Pigg, A., Norli, M., Oakes, J. M.,
737 Osterholz, H., Park, H., Paulsen, M. L., Rosentreter, J. A., Rueda-Roa, D., Santinelli,
738 C., Shen, Y., Teira, E., Tinta, T., Uher, G., Wakita, M., Ward, N., Watanabe, K., Xin,
739 Y., Yamashita, Y., Yang, L., Yeo, J., Yuan, H., Zheng, Q., and Álvarez-Salgado, X.
740 A.: A global database of dissolved organic matter (DOM) [concentration](#)
741 measurements in coastal waters (CoastDOM v1) [dataset], 2023.

742 Mackenzie, F. T., De Carlo, E. H., and Lerman, A.: 5.10 - Coupled C, N, P, and O
743 Biogeochemical Cycling at the Land–Ocean Interface, in: *Treatise on Estuarine and
744 Coastal Science*, edited by: Wolanski, E., and McLusky, D., Academic Press,
745 Waltham, 317-342, <https://doi.org/10.1016/B978-0-12-374711-2.00512-X>, 2011.

746 Mopper, K., Kieber, D. J., and Stubbins, A.: Marine Photochemistry of Organic Matter,
747 in: *Biogeochemistry of Marine Dissolved Organic Matter*, edited by: Carlson, C. A.,
748 and Hansell, D. A., 389-450, 10.1016/b978-0-12-405940-5.00008-x, 2015.

749 Osterholz, H., Burmeister, C., Busch, S., Dierken, M., Frazão, H. C., Hansen, R.,
750 Jeschek, J., Kremp, A., Kreuzer, L., and Sadkowiak, B.: Nearshore dissolved and
751 particulate organic matter dynamics in the southwestern Baltic Sea: environmental
752 drivers and time series analysis (2010–2020), *Frontiers in Marine Science*, 8,
753 795028, 2021.

754 Raymond, P. A. and Spencer, R. G. M.: Riverine DOM, in: *Biogeochemistry of Marine
755 Dissolved Organic Matter*, edited by: Hansell, D. A., and Carlson, C. A., Elsevier,
756 Amsterdam, 509-533, 10.1016/b978-0-12-405940-5.00011-x, 2015.

757 Redfield, A. C., Ketchum, B. K., and Richards, F. A.: The influence of organisms on the
758 composition of sea-water, in: *The sea*, vol. 2, The composition of sea water:
759 Comparative and descriptive oceanography, edited by: Hill, M. N., Wiley-Interscience,
760 26- 77, 1963.

761 Santos, I. R., Burdige, D. J., Jennerjahn, T. C., Bouillon, S., Cabral, A., Serrano, O.,
762 Wernberg, T., Filbee-Dexter, K., Guimond, J. A., and Tamborski, J. J.: The
763 renaissance of Odum's outwelling hypothesis in 'Blue Carbon' science, *Estuarine,
764 Coastal and Shelf Science*, 255, 10.1016/j.ecss.2021.107361, 2021.

765 Sharp, J. H., Benner, R., Bennett, L., Carlson, C. A., Dow, R., and Fitzwater, S. E.: Re-
766 evaluation of high temperature combustion and chemical oxidation measurements of

767 dissolved organic carbon in seawater, *Limnology and Oceanography*, 38, 1774-1782,
768 <https://doi.org/10.4319/lo.1993.38.8.1774>, 1993.
769 Sharp, J. H., Rinker, K. R., Savidge, K. B., Abell, J., Benaim, J. Y., Bronk, D., Burdige,
770 D. J., Cauwet, G., Chen, W., Doval, M. D., Hansell, D., Hopkinson, C., Kattner, G.,
771 Kaumeyer, N., McGlathery, K. J., Merriam, J., Morley, N., Nagel, K., Ogawa, H.,
772 Pollard, C., Pujo-Pay, M., Raimbault, P., Sambrotto, R., Seitzinger, S., Spyres, G.,
773 Tirendi, F., Walsh, T. W., and Wong, C. S.: A preliminary methods comparison for
774 measurement of dissolved organic nitrogen in seawater, *Marine Chemistry*, 78, 171-
775 184, 2002.
776 Sipler, R. E. and Bronk, D. A.: Dynamics of Dissolved Organic Nitrogen, in:
777 *Biogeochemistry of Marine Dissolved Organic Matter*, edited by: Hansell, D. A., and
778 Carlson, C. A., 127-232, 10.1016/b978-0-12-405940-5.00004-2, 2015.
779 Solorzano, L. and Sharp, J. H.: Determination of total dissolved phosphorus and
780 particulate phosphorus in natural waters, *Limnology and Oceanography*, 25, 754-
781 758, 1980.
782 Taniguchi, M., Dulai, H., Burnett, K. M., Santos, I. R., Sugimoto, R., Stieglitz, T., Kim,
783 G., Moosdorf, N., and Burnett, W. C.: Submarine Groundwater Discharge: Updates
784 on Its Measurement Techniques, Geophysical Drivers, Magnitudes, and Effects,
785 *Frontiers in Environmental Science*, 7, 10.3389/fenvs.2019.00141, 2019.
786 Valderrama, J. C.: The simultaneous analysis of total nitrogen and total phosphorus in
787 natural waters, *Marine Chemistry*, 10, 109-122, 1981.
788 Wada, S., Aoki, M. N., Mikami, A., Komatsu, T., Tsuchiya, Y., Sato, T., Shinagawa, H.,
789 and Hama, T.: Bioavailability of macroalgal dissolved organic matter in seawater,
790 *Marine Ecology Progress Series*, 370, 33-44, 2008

791

792 **Figure legends**

793 **Figure 1.** Flow diagram of data collation, quality control and inclusion into CoastDOM v1
794 database.

795 **Figure 24.** Histograms showing the distribution of observations for **a)** dissolved organic
796 carbon (DOC), **b)** nitrogen (DON) and **c)** phosphorus (DOP), within defined
797 concentration ranges in the coastal ocean. Note that the concentration ranges are not
798 uniform in all cases due to the large difference in concentrations_s levels.

799 **Figure 23.** **a)** Cumulative number of concentration observations for dissolved organic
800 carbon (DOC), nitrogen (DON), and phosphorus (DOP). Number of concentration
801 observations shown as a function of **b) sampling month** (“N.S” are samples for which
802 the sampling month is not specified), **cb)** latitude, and **de)** longitude, grouped into bins
803 of 10° latitude or longitude.


804 **Figure 34.** Global distribution of concentration observations included in CoastDOM v1 for
805 **a)** dissolved organic carbon (DOC), **b)** nitrogen (DON), and **c)** phosphorus (DOP). The
806 black dots on the map represent the reported data that are included in the CoastDOM
807 v1 database. Histograms show the distribution of observations for DOC, DON and DOP
808 within defined concentration ranges in the continents where measurements are
809 available. Maps were created using the GIS shape file obtained from Laurelle et al.
810 (Laruelle et al., 2013)

811 **Table 1.** Descriptive statistics for the dissolved organic carbon (DOC), dissolved organic
 812 nitrogen (DON), and dissolved organic phosphorus (DOP) concentration observations
 813 measurements included in the CoastDOM v1 dataset. The DOC:DON, DOC:DOP and
 814 DON:DOP ratios are also reported. The minimum (Min), maximum (Max), average values
 815 (Avg.) and standard deviation (SD), coefficient of variation (CV %), median, 25th and 75th
 816 percentiles (perc.) and number of samples (N) for each variable are shown.

	DOC	DON	DOP			
	$\mu\text{mol C L}^{-1}$	$\mu\text{mol N L}^{-1}$	$\mu\text{mol P L}^{-1}$			
Min	17	<0.1	<0.01			
Max	30327	2095.3	84.27			
Avg. (SD)	182 (314)	13.6 (30.4)	0.34 (1.11)			
Median	103	8.0	0.18			
CV %	173	224	326			
25th perc.	77	5.5	0.11			
75th perc.	228	15.8	0.30			
N	62339	20357	13534			

	DOC	DON	DOP	DOC:DON	DOC:DOP	DON:DOP
	$\mu\text{mol L}^{-1}$	$\mu\text{mol L}^{-1}$	$\mu\text{mol L}^{-1}$			
Min	17	<0.1	<0.01	1	18	0.14
Max	30327	2095.3	84.27	3046	248024	8894
Avg. \pm SD	182 \pm 314	13.6 \pm 30.4	0.34 \pm 1.11	18 \pm 43	1171 \pm 4248	100 \pm 580
Median	103	8.0	0.18	14	583	47
CV	173	224	324	244	363	578
25%iles	77	5.5	0.11	11	401	30
75%iles	228	15.8	0.30	18	1034	78
N	62339	20357	13534	12632	7415	12954


817




§18

§19




Figure 1.






821

822 **Figure 24.**






823 -
824




825

826 **Figure 23.**

■ Africa
 ■ Antarctica
 ■ Asia
 ■ Europe
 ■ Oceania
 ■ South America
 ■ North America



827
828



829

830 Figure 43.