

1 Reconciliation-Comparison of observation- and inventory- based methane emissions for
2 eight large global emitters

3
4 Ana Maria Roxana Petrescu¹, Glen P. Peters², Richard Engelen³, Sander Houweling¹, Dominik Brunner⁴, Aki
5 Tsuruta⁵, Bradley Matthews⁶, Prabir K. Patra^{7,8,9}, Dmitry Belikov⁹, Rona L. Thompson¹⁰, Lena Höglund-
6 Isaksson¹¹, Wenxin Zhang¹², Arjo J. Segers¹³, Giuseppe Etiope^{14,15}, Giancarlo Ciotoli^{16,14}, Philippe Peylin¹⁷,
7 Frédéric Chevallier¹⁷, Tuula Aalto⁵, Robbie M. Andrew², David Bastviken¹⁸, Antoine Berchet¹⁷, Grégoire
8 Broquet¹⁷, Giulia Conchedda¹⁹, Johannes Gütschow²⁰, Jean-Matthieu Haussaire⁴, Ronny Lauerwald²¹, Tiina
9 Markkanen⁵, Jacob C. A. van Peet¹, Isabelle Pison¹⁷, Pierre Regnier²², Espen Solum¹⁰, Marko Scholze¹², Maria
10 Tenkanen⁵, Francesco N. Tubiello¹⁹, Guido R. van der Werf²³, John R. Worden²⁴

11
12 ¹Department of Earth Sciences, Vrije Universiteit Amsterdam, 1081HV, Amsterdam, the Netherlands

13 ²CICERO Center for International Climate Research, Oslo, Norway

14 ³European Centre for Medium-Range Weather Forecasts (ECMWF), Reading, RG2 9AX, UK

15 ⁴Empa, Swiss Federal Laboratories for Materials Science and Technology, 8600 Dübendorf, Switzerland

16 ⁵Finnish Meteorological Institute, P. O. Box 503, FI-00101 Helsinki, Finland

17 ⁶Umweltbundesamt GmbH, Climate change mitigation & emission inventories, 1090, Vienna, Austria

18 ⁷Research Institute for Humanity and Nature, Kyoto 6038047, Japan

19 ⁸Research Institute for Global Change, JAMSTEC, Yokohama 2360001, Japan

20 ⁹Chiba University, 1-33 Yayoicho, Inage Ward, Chiba, 263-8522, Japan

21 ¹⁰NILU - Norsk Institutt for Luftforskning, Kjeller, Norway

22 ¹¹International Institute for Applied Systems Analysis (IIASA), 2361 Laxenburg, Austria

23 ¹²Department of Physical Geography and Ecosystem Science, Lund University, SE-223 62 Lund, Sweden

24 ¹³Department of Climate, Air and Sustainability, TNO, Princetonlaan 6, 3584 CB Utrecht, the Netherlands

25 ¹⁴Istituto Nazionale di Geofisica e Vulcanologia, Sezione Roma 2, via V. Murata 605, Roma, Italy

26 ¹⁵Faculty of Environmental Science and Engineering, Babes-Bolyai University, Cluj-Napoca, Romania

27 ¹⁶Consiglio Nazionale delle Ricerche, Istituto di Geologia Ambientale e Geoingegneria, Via Salaria km 29300,
28 00015 Monterotondo, Rome, Italy

29 ¹⁷Laboratoire des Sciences du Climat et de l'Environnement, 91190 Gif-sur-Yvette, France

30 ¹⁸Department of Thematic Studies – Environmental Change, Linköping University, Sweden

31 ¹⁹Food and Agriculture Organization of the United Nations, Statistics Division. 00153 Rome, Italy

32 ²⁰Climate Resource, Northcote, Australia

33 ²¹Université Paris-Saclay, INRAE, AgroParisTech, UMR ECOSYS, Palaiseau, France

34 ²²Biogeochemistry and Modeling of the Earth System, Université Libre de Bruxelles, 1050 Bruxelles, Belgium

35 ²³Meteorology and Air Quality Groep, Wageningen University and Research, Wageningen, the Netherlands

36 ²⁴Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA

37
38 *Correspondence to:* A.M. Roxana Petrescu (a.m.r.petrescu@vu.nl)
39

40 **Abstract**

41

42 Monitoring the spatial distribution and trends in surface greenhouse gas (GHG) fluxes, as well as flux
43 attribution to natural and anthropogenic processes, is essential to track progress under the Paris Agreement and
44 to inform its Global Stocktake. This study updates earlier syntheses (Petrescu et al., 2020, 2021, 2023^a) and
45 provides a consolidated synthesis of CH₄ emissions using bottom-up (BU) and top-down (TD) approaches for
46 the European Union (EU) and is expanded to include seven additional countries with large anthropogenic and/or
47 natural emissions (USA, Brazil, China, India, Indonesia, Russia, and the Democratic Republic of Congo (DR
48 Congo)). Our aim is to demonstrate the use of different emission estimates to help improve national GHG
49 emission inventories for a diverse geographical range of stakeholders.

50 We use updated National GHG Inventories (NGHGIs) reported by Annex I Parties under the United
51 Nations Framework Convention on Climate Change (UNFCCC) in 2023 and the latest available Biennial Update
52 Reports (BURs) reported by non-Annex I Parties. Comparing NGHGIs with other approaches highlights that
53 different system boundaries are a key source of divergence. A key system boundary difference is whether both
54 anthropogenic and natural fluxes are included and, if they are, how fluxes belonging to these two sources are
55 partitioned.

56 Over the studied period, the total CH₄ emission estimates in the EU, USA, and Russia show a steady
57 decreasing trend since 1990, while for the non-Annex I emitters analyzed in this study, Brazil, China, India,
58 Indonesia, and DR Congo, CH₄ emissions have generally increased. Quantitatively, in the EU the mean of 2015-
59 2020 anthropogenic UNFCCC NGHGIs (15 ± 1.8 Tg CH₄ yr⁻¹) and the mean of the BU CH₄ emissions (17.8 (16-
60 19) Tg CH₄ yr⁻¹) generally agree on the magnitude, while inversions show higher emission estimates (medians
61 of 21 (19-22) Tg CH₄ yr⁻¹ and 24 (22-25) Tg CH₄ yr⁻¹ for the three regional and six global inversions,
62 respectively), as they include natural emissions, which for the EU were quantified at 6.6 Tg CH₄ yr⁻¹ (Petrescu
63 et al., 2023^a). Similarly, for the other Annex I Parties in this study (**USA and Russia**), the gap between the BU
64 anthropogenic and total TD emissions is partly explained by the natural emissions.

65 For the **non-Annex I Parties**, anthropogenic CH₄ estimates from UNFCCC BURs show large
66 differences with the other global inventory-based estimates and even more with atmospheric-based ones. This
67 poses an important potential challenge to monitoring the progress of the global CH₄ pledge and the Global
68 Stocktake. Our analysis provides a useful baseline to prepare for the influx of inventories from non-Annex I
69 Parties as regular reporting starts under the Enhanced Transparency Framework of the Paris Agreement.

70 By systematically comparing the BU with TD methods, this study provides recommendations for more
71 robust comparisons of available data sources and hopes to steadily engage more Parties in using observational
72 methods to complement their UNFCCC inventories, as well as considering their natural emissions. With
73 anticipated improvements in atmospheric modeling and observations, as well as modeling of natural fluxes,
74 future development needs to resolve knowledge gaps in both BU and TD approaches and to better quantify

75 remaining uncertainty. TD methods may emerge as a powerful tool to help improve NGHGs of CH₄ emissions,
76 but further confidence is needed in the comparability and robustness of the estimates.

77 The referenced datasets related to figures are available at <https://doi.org/10.5281/zenodo.10276087>
78 [12582667](https://doi.org/10.5281/zenodo.12582667) (Petrescu et al., 2023b).

79 1. Introduction

80

81 In 2021, the NOAA Global Monitoring Laboratory (GML) reported the largest annual increase in
82 atmospheric CH₄ mixing ratios since records began in 1983, at 17 parts per billion (ppb) (NOAA
83 (https://gml.noaa.gov/ccgg/trends_ch4/). In 2022, atmospheric CH₄ concentrations averaged 1912 ppb yr⁻¹, 162
84 % higher than pre-industrial levels. A similar, abnormally large growth rate of 14.8 ppb yr⁻¹ was detected from
85 total column mixing ratio measurements (XCH₄) by the Greenhouse Gases Observing Satellite (GOSAT) (Peng
86 et al., 2022). ~~The drivers of the recent growth are most likely driven primarily by biogenic emissions (Basu et al., 2022; Lan, et al., 2021a; Lan et al., 2021b; Lan et al., 2022; Nisbet et al., 2016, 2019), with smaller~~
87 ~~contributions from increased fossil fuel emissions and a reduced atmospheric sink (Nisbet et al., 2023).~~ ~~These~~
88 ~~processes drove the near record increase in atmospheric CH₄ growth in 2020, despite and furthermore outweighed~~
89 ~~the slight the observed decrease in anthropogenic CH₄ emissions accumulated from March–June 2020 as impact~~
90 ~~of the COVID-19 slowdown (e.g. China) which might be small relative to the long-term positive trend in~~
91 ~~emissions. (McNorton et al., 2022, Peng et al., 2022, Qu et al. 2022).~~

93 CH₄ in the atmosphere has many different sources, of both natural and anthropogenic origin. The natural
94 sources of CH₄ are dominated by wetlands, while anthropogenic emissions principally come from agricultural
95 activities (livestock and rice farming), waste management (landfills and water treatment plants) and the
96 production, transportation, and use of fossil fuels. Most of the agricultural sources are distributed sources, while
97 the energy-related industrial sources of CH₄ are a mix of large point sources, of which some are detectable by
98 satellite ([Jacob et al., 2022](#)) and smaller point and distributed sources of fugitive emissions (e.g., leaks in
99 pipelines and compression stations), which are more challenging to identify (Rutherford et al., 2021; Omara et
100 al., 2022).

101 While anthropogenic CH₄ emissions from fossil fuels, agriculture, and waste can be reduced by
102 mitigation actions, increased natural emissions lead to different challenges. It has been suggested that the drivers
103 of the recent growth are most likely driven primarily by biogenic emissions (Basu et al., 2022; Lan, et al., 2021a;
104 Lan et al., 2021b; Lan et al., 2022; Nisbet et al., 2016, 2019), with smaller contributions from increased fossil
105 fuel emissions and a reduced atmospheric sink (Nisbet et al., 2023). Fluctuations in natural sources - dominated
106 by wetlands and open water bodies - were the main reasons for some of the atmospheric CH₄ anomalies observed
107 during the last decades (Rocher-Ros et al., 2023; Zhang et al., 2023; Nisbet et al., 2023; Lunt et al., 2019). Nisbet
108 et al., 2023 review recent studies, including those which quantified the observed methane growth in the last years.
109 Using a global inverse analysis of GOSAT satellite observations, ~~it has been shown that increases in the range~~
110 ~~of 22-32 Tg CH₄ yr⁻¹ were detected~~ between 2019 ~~and 2020~~ ~~were in the range of 22–32 Tg CH₄ yr⁻¹~~ and were
111 attributed to biogenic sources, half of which took place in East Africa (~ 15 Tg yr⁻¹), and some were observed in
112 Canada and Alaska (4.8 Tg yr⁻¹) (Qu et al., 2022 and Basu et al., 2022).

113 Chandra et al., 2021 identified a few main sectors that triggered increases and decreases in the
114 anthropogenic CH₄ emissions of different countries. The first is energy, with its fugitive emissions from the oil
115 and gas industry whose decline in emissions helped stabilize CH₄ concentration in the 1990s, before they
116 contributed to the renewed CH₄ growth since the late 2000s (increased emissions). The other major sectors that
117 drove changes in the CH₄ growth rate were agriculture (increase in emissions from enteric fermentation and
118 manure management) and waste. The increase in emissions from enteric fermentation and manure management
119 was caused primarily by increased animal numbers, and in addition by the greater intensity of ruminant farming
120 as estimated by the FAO and the emission inventories (e.g. EDGAR) which might take into account productivity
121 increases (Crippa et al., 2020; Wolf et al., 2017; FAOSTAT, 2018) while inventory emissions from waste can
122 account for up to 43 % of the linear increase in emissions for the rest of the world.

123 The contribution of CH₄ to global warming has been estimated to be about 0.5°C relative to the period
124 1850–1900 (IPCC, 2021) (Stavert et al., 2022). Methane has a relatively short perturbation lifetime (averaging
125 12.4 years, Balcombe et al., 2018) and a high global warming potential (86 and 34 for 20- and 100-years times
126 horizons respectively, compared to that of CO₂ emissions, IPCC, 2021, Table 7.15). Given the short lifetime, a
127 decline in CH₄ emissions will rapidly reduce the global warming contribution from CH₄ and help mitigate the
128 impact of climate change at decadal time scale (Cain et al., 2021). However, efforts to reduce CH₄ emissions
129 require a thorough understanding of the dominant CH₄ sources and sinks and their temporal and regional
130 distribution and trends (Stavert et al., 2022).

131 The Paris Agreement, a milestone of the UNFCCC to combat climate change and adapt to its effects,
132 entered into force on November 4, 2016. It asks each signatory to define and communicate its planned climate
133 actions, known as Nationally Determined Contributions (NDCs), and to report their progress towards their
134 targets. Next to commitments adopted by countries at COP26, the Global Methane Pledge (GMP) was launched.
135 The goal of the GMP is to cut anthropogenic CH₄ emissions by at least 30 % by 2030 with respect to 2020 levels
136 and is seen as the fastest way to reduce near-term warming and is necessary to keep a 1.5°C temperature limit
137 within reach. Achieving this goal will drive significant energy security, food security, health, and development
138 gains, through specific energy and agriculture defined pathways including innovative actions, national targeted
139 policies, and green climate funds to help smallholder farmers ([https://www.state.gov/global-methane-pledge-](https://www.state.gov/global-methane-pledge-from-moment-to-momentum/)
140 from-moment-to-momentum/). About 150 countries joined this pledge and about fifty already have already
141 developed national CH₄ action plans or are doing so. As agriculture and waste are the main anthropogenic
142 sources for CH₄ emissions, a GMP Food and agriculture pathway and a GMP waste pathway were launched at
143 COP27, foreseeing actions that increase agricultural productivity, while reducing emissions from dairy, food loss
144 and waste by supporting small farmers and innovation ([https://www.state.gov/global-methane-pledge-from-](https://www.state.gov/global-methane-pledge-from-moment-to-momentum/)
145 moment-to-momentum/).

146 Starting in 2024, non-Annex I Parties to the UNFCCC must - given they have sufficient capacities -
147 report formal inventories under the Paris Agreement's Enhanced Transparency Framework following the same
148 guidelines and rules as the Annex I countries (Perugini et al., 2021). Furthermore, they will undergo more
149 stringent reviews than those that previously looked at the Biennial Update Reports (BURs) and NDCs. This will
150 also allow strengthening the robustness of such comparison exercises when using independent atmospheric
151 observations in estimating trends and patterns for regional and national CH₄ emissions (IPCC, 2006). The influx

152 of new inventories will place additional demands on independent emission estimates to help improve and inform
153 National Greenhouse Gas Inventories (NGHGs), particularly in countries with low capacity.

154 With increased focus on CH₄ in climate policy, there is a demand to ensure that planned emission
155 reductions are realized. Further, as non-Annex I countries begin regular reporting of emission estimates under
156 the Enhanced Transparency Framework, there is a need to help countries improve their GHG emission estimates.
157 This has created an active field of research comparing NGHGI and independent estimates. Our analysis builds
158 on the three-year EU funded project CoCO₂, which had as main objective the building of prototype systems for
159 a European Monitoring and Verification Support capacity for anthropogenic CO₂ (and CH₄) emissions
160 (CO₂MVS). In this context, one of the results of the CoCO₂ project was the production of a Blueprint for a
161 decision support system to be used in an eventual CO₂MVS, aiming at informing and attracting attention of
162 diverse climate stakeholders on the use of the results needed beyond research. Therefore, the objectives of this
163 study reflect those of the Blueprint and focus on user engagement. It builds on dialogues with a broad community
164 of users (e.g. scientists, inventory agencies, policy makers), considering their opinions and needs when it comes
165 to comparisons between independent approaches. Furthermore, this study expands beyond the EU to include
166 seven countries that have large anthropogenic and/or natural CH₄ emissions (USA, Brazil, China, India,
167 Indonesia, Russia and the Democratic Rep. of Congo). It examines both Annex I (EU, USA and Russia) and non-
168 Annex I estimates from observation-based BU process-based models and inversions-based TD approaches (using
169 satellite observations) by identifying and explaining differences with official inventory reports submitted by
170 parties to the UNFCCC. The seven countries were chosen based on location and the importance / magnitude of
171 their anthropogenic and natural emissions. By using multiple methodologies, uncertainties can be estimated by
172 looking at the range in both emissions and trends.

173

174 **2. Methods and data**

175 In this work we focus on comparing BU and TD emission estimates. The ‘reconciliation process’
176 described in this work is the action of making one dataset comparable with another to assess their consistency.
177 In this respect, we attempt to obtain consistent results from both BU and TD estimates, through harmonization
178 of the results, concepts and definitions. After the reconciliation process, the estimates do not necessarily agree,
179 representing uncertainties in the different methods and datasets. We now describe the key data and methods used
180 in our analysis.

181 **2.1. Verification practices in official UNFCCC NGHGs**

182

183 Quality assurance/quality control (QA/QC) is a key component of NGHGs development. Verification
184 is an additional step and refers specifically to methods that are external to the inventory and apply independent
185 data. There are two main methods of verification: 1) independent inventory-based estimates, 2) observation-
186 based emission estimates.

187 A challenge with comparisons against *independent inventory-based estimates* is that none of them is
188 truly independent as they may rely on, for example, the same activity data reported by a country (Andrew 2020).
189 Experience has shown that performing detailed comparisons (Petrescu et al., 2021, 2023a, Lauerwald et al., 2024)

190 can help clarify differences in system boundaries or even identify errors (Andrew 2020). Improving independent
191 emission inventories also has value, as these are often used in global studies where common methods across all
192 countries are desired.

193 *Observation-based estimates* use observations of atmospheric concentrations and prior fluxes that are
194 then coupled to a transport model. These methods are more complex and computationally expensive and can
195 make use of both direct observations and emission inventories.

196 ~~Since most developed countries have reported UNFCCC inventories for decades and these have been~~
197 ~~continually reviewed and refined, the focus of this work is on observation-based estimates. As an increasing~~
198 ~~number of developing countries begin more detailed and frequent reporting, comparisons with independent TD~~
199 ~~approaches will be an important method of verification for those countries.~~

200 The 2019 refinement of the 2006 IPCC guidelines highlighted notable advances in the application of
201 inverse models of atmospheric transport for estimating emissions at the national scale. Building on this progress,
202 they extend the guidance on the use of atmospheric measurements for verification (IPCC, 2019). There are
203 several countries that currently use atmospheric measurements for verification of parts of their inventories.
204 Australia (Luhar et al 2020, AUS NIR, 2023) and New Zealand (Geddes et al., 2021) have estimated regional
205 CH₄ emissions to help better understand the methods and their potential. Germany performs various cross-
206 validation checks with available data (German NIR, 2023), some of which are based on observations. The UK
207 and Switzerland (Annex 6 CHE NIR, 2023) have developed more comprehensive methods based on inversion
208 modeling, covering several GHGs in addition to CH₄. Building on modeling experience, the country reporting
209 confirms that most potential lies in using observations to verify fluorinated gases (Annex 6 UK NIR, 2023), but
210 the large uncertainty in CH₄ emissions gives the potential for verification if a sufficient observation network is
211 used in inversion modeling (Bergamaschi et al., 2018, Thompson et al., 2014).

212 While inversions of CH₄ fluxes are associated with significant uncertainty, so are NGHGI estimates of
213 anthropogenic CH₄ emissions. Furthermore, inversions can provide information on subannual and subnational
214 variations in time and space that may indicate differences in source sector emission estimates. In geographic
215 areas with sufficiently dense ground-based observation networks, the inversions will have more value.

216

217 **2.2. Anthropogenic CH₄ emissions from the NGHGs**

218

219 Annex I countries report their annual GHG emissions to the UNFCCC in the so-called Common
220 Reporting Format (CRFs) data tables and National Inventory Reports (NIRs). Here, anthropogenic CH₄
221 emissions from the five UNFCCC sectors, incl. Land Use, Land Use Change and Forestry (LULUCF) are
222 grouped together. As part of the LULUCF sector, we also have the CH₄ emissions from wetlands, which
223 according to the IPCC guidelines are defined as managed "where the water table is artificially changed (i.e.
224 lowered or raised) or those created through human activity (e.g. damming a river) and that do not fall into Forest
225 Land, Cropland, or Grassland categories (IPCC, 2014)". Reporting CH₄ emissions from managed wetlands is not
226 mandatory, but if done, parties are encouraged to make use of the 2013 IPCC Wetlands supplement (IPCC,
227 2014). In the EU, if Member States report these emissions, they report not only restored (rewetted) wetlands but
228 also emissions from drained organic and mineral soils (e.g. peatlands, ditches, etc.). These are not large by
229 magnitude but are large by area in the Nordic countries. According to NGHGI data, in 2021, managed wetlands

230 in the EU, for which emissions were reported under the LULUCF (CRF Table 4(II) and Summary 1.As2
231 accessible for each EU country), summed up to 0.21 Tg CH₄ yr⁻¹, in comparison to total emissions of ~15 Tg
232 CH₄ yr⁻¹. Furthermore, the NGHGs do not include any lateral fluxes from inland waters but do include biomass
233 burning anthropogenic emissions reported under the LULUCF sector.

234 The presented uncertainties in the CH₄ emission levels of the individual countries and the EU are not
235 always reported in a complete and harmonized format, and therefore were calculated applying gap-filling and
236 harmonization procedures that are used to compile the EU GHG inventory reported under UNFCCC (EU NIR,
237 2023) (see SI and Appendix A1.1 in Petrescu et al., 2023a). The EU uncertainty analysis reported in the bloc's
238 National Inventory Report (NIR) is based on country-level, Approach 1 uncertainty estimates (IPPC, 2006, Vol.
239 1, Chap. 3) that are reported by EU Member States, previously under Article 7(1)(p) of Regulation (EU) 525/2013
240 and since 2023 under Article 26(3) and Annex V(Part 1)(m) of the Governance Regulation (EU) 2018/1999.

241 Non-Annex I countries report their updated NGHGs to the UNFCCC, including a national inventory
242 report and information on mitigation actions, needs and support received in Biennial Update Reports (BURs). In
243 this study, Brazil, China, Indonesia, India and the Democratic Rep. of Congo (DR Congo) were investigated. For
244 Brazil, information from its fourth BUR (Brazil, 2020) was used, giving both total and sectoral split emission
245 values for years 1994, 2000, 2010, 2012, 2015 and 2016. For China, information from its second BUR Tables 2-
246 10, 2-13, 2-14, 2-15, and 2-16 was used (China, 2019). The information was available for both total and sectoral
247 split emission values for 1994, 2005, 2010 and 2014. Uncertainties for 2014 are available in Table 2-12.
248 Indonesia submitted its third BUR in 2021 (Indonesia, 2021). Indonesian total sectoral CH₄ emissions time series
249 as reported by the 2nd ~~UNFCCC~~-BUR (2001-2016) were revised in the 3rd BUR (2000 and 2019, Table 2). For
250 2017 and 2018, only the agricultural CH₄ emissions were detailed by the 3rd BUR (Fig. 2-24), but the total
251 emissions for these two years were not reported. Uncertainty for 2019 activity data and emission factors (EFs)
252 are the same as reported in the 2nd BUR (2018). The ~~result of the uncertainty analysis showed that the~~ overall
253 uncertainty of Indonesia's National GHG inventory with agriculture and AFOLU-LULUCF (including peat fires)
254 for 2000 and 2019 were approximately 20.0% and 19.9%, respectively. A much smaller uncertainty, 10.4 % for
255 2000 and 13.8 % for 2019, occurred when the FOLUforestry and land use sector (including forest fires), was
256 excluded from the analysis, pointing to the high uncertainty of, ~~This shows that Indonesian emission inventories~~
257 ~~are highly uncertain when emissions from~~ forest fires in Indonesia are included in the analysis. The DR Congo
258 submitted its first BUR in 2022, and we used timeseries reported for 2000-2018 (Table 12 Congo, 2022). India
259 has submitted three BURs and information on sectoral CH₄ emissions are in each of them only for one year. We
260 compiled information for 2010 from the first BUR (India, 2016), for 2014 from the second BUR (India, 2018)
261 and for 2016 from the third and latest BUR (India, 2021).

262 **2.3. Other CH₄ data sources and estimation approaches**

263

264 The CH₄ emissions in the EU and non-Annex I countries used in the atmospheric inversions and
265 anthropogenic and natural emissions estimates from various BU approaches and inventories (i.e., UNFCCC
266 CRFs and BURs) covering specific products, sectors and activities are summarized in Table 1. The data and the
267 detailed description of most products (Tables S1 and S2, Supplementary Information) span the period from 1990
268 to 2021, with some of the data only available for shorter time periods. The estimates are available both from peer-

269 reviewed literature and from unpublished research results from the VERIFY and CoCO₂ projects (Supplementary
 270 Information, SI) and in this work they are compared with NGHGs reported in 2023 (time series for all (Annex
 271 I) or some years (non-Annex I) of the 1990-2021 period). The BU anthropogenic sources are from UNFCCC
 272 NGHGs and three global inventory datasets/models: EDGARv7.0, FAOSTAT/[PRIMAP-hist 2.4](#) and GAINS.
 273 In this synthesis, ~~data from~~ FAOSTAT (Tubiello et al., 2022; FAO, 2023) data includes estimates for all
 274 economic sectors: Energy, [Industrial Processes and Products Use \(IPPU\)](#), Waste and Other, which are sourced
 275 from the PRIMAP-hist v2.4 dataset (Gütschow et al., 2022) to build emissions indicators on agrifood systems
 276 and on the entire economy. Emission totals from the agrifood domain are computed following the Tier 1 methods
 277 of the Intergovernmental Panel on Climate Change (IPCC) Guidelines for NGHGs. Agrifood systems emissions
 278 in FAOSTAT are largely based on FAO crop, livestock and land--use statistics (Tubiello et al., 2022; FAO,
 279 2023). They are complemented with activity data from the UN Statistics~~at~~ Division (UNSD), the International
 280 Energy Agency (IEA) and with geospatial information on drained organic soils and biomass fires (Conchedda
 281 and Tubiello, 2020; Prospero et al., 2020).

282 The analysis focuses on both total and sectoral or partitioned information from both BU and TD
 283 estimates. As detailed in Table 1, not all inversions distinguish between sources but in the following sections we
 284 discuss comparability between BU and TD for both total and partitioned results.

285 *Table 1: Sectors included in this study and data sources providing estimates for these sectors. CAMS stands for*
 286 *Copernicus Atmosphere Monitoring Service. References to data products are found in Table 2 Petrescu et al.,*
 287 *2023a and Table S1 and S2, SI.*

Anthropogenic (BU) ¹ CH ₄	Natural (BU) ² CH ₄	Regional TD CH ₄	Global TD CH ₄
1. Energy: UNFCCC NGHGI (CRFs and BURs), GAINS, EDGAR v7.0, FAOSTAT/ PRIMAP-hist 2.4	Wetlands EU: JSBACH-HIMMELI Global: LPJ-GUESS	No partitions – total emissions FLEXkF_v2023	Totals and partitioned emissions:
2. Industrial Products and Products in Use (IPPU): UNFCCC NGHGI (CRFs and BURs), EDGAR v7.0, FAOSTAT/ PRIMAP-hist 2.4	Peatlands, mineral soils: EU: JSBACH-HIMMELI Global: LPJ-GUESS	CIF-FLEXPARTv10.4 CIF-CHIMERE	MIROC4-ACTM (control and OH varying runs) CAMSv21r1 (NOAA and NOAA_GOSAT runs)
3. Agriculture: UNFCCC NGHGI (CRFs and BURs), GAINS, EDGAR v7.0, FAOSTAT	Inland waters fluxes EU: lakes, rivers and reservoirs (RECCAP2)		TM5-4DVAR (TROPOMI)
4. LULUCF: UNFCCC NGHGI (CRFs and BURs) and FAOSTAT	Global: lakes and reservoirs ORNL DAAC		CTE-CH ₄ (GCP2021)

¹ For consistency with the NGHGI, here we refer to the five reporting sectors as defined by the UNFCCC and the Paris Agreement decision (18/CMP.1), the IPCC Guidelines (IPCC, 2006), and their Refinement (IPCC, 2019a), with the only exception that the latest IPCC Refinement groups together Agriculture and LULUCF sectors in one sector (Agriculture, Forestry and Other land Use - AFOLU).

² The term **natural** refers here to unmanaged natural CH₄ emissions (peatlands, mineral soils, geological, inland waters and biomass burning) not reported under the anthropogenic UNFCCC LULUCF sector.

5. Waste: UNFCCC NGHGI (CRFs and BURs), GAINS, EDGAR v7.0, FAOSTAT/PRIMAP-hist 2.4	Geological fluxes updated activity (see SI) Biomass burning (GFEDv4.1s)		CEOS (GOSAT) GEOS-Chem CTM (TROPOMI) for USA only
--	--	--	--

288 note: Not all models have a version id. Those that have, are used in previous syntheses (Petrescu et al., 2021 and 2023a).

289 We define natural sources as all sources which do not belong to the anthropogenic partition: wetlands,
290 geological, termites, ocean, inland waters, soils and biomass burning-(Table 2). The BU natural components for
291 the EU were computed as the sum of the VERIFY products (biomass burning, inland waters and undisturbed
292 peatlands plus mineral soils (as described in Petrescu et al., 2021 and 2023) and geological emissions (Etiope et
293 al. 2019) updated for the VERIFY project). For the seven non-EU emitters, the BU natural fluxes are the sum of
294 wetland emissions (LPJ-GUESS), lake and reservoir emissions (ORNL DAAC), biomass burning emissions
295 (GFED4.1s) and geological emissions (updated activity, SI). The TD natural global estimates were calculated as
296 the sum of all natural partitions reported by the inversions. Adjustments were made to have a consistent
297 comparison between partitions, adding the missing ones from the BU estimates (Table 4). The error bar on the
298 TD natural represents the range of the min/max between inversion estimates.

299 The total regional TD estimates (for EU) and their uncertainties were calculated as the mean and
300 min/max range between FLEXkF_v2023, CIF-FLEXPART and CIF-CHIMERE inversions (see Priors table in
301 Petrescu et al., 2023b). For the USA, we considered the optimized emissions from the GEOS-Chem CTM (based
302 on TROPOMI data for 2019) from Nesser et al., (2023), with the range from the eight members of the inversion
303 ensemble shown as uncertainty (Table 2 in Nesser et al., 2023).

304 For all countries, the total global TD inversion estimates (time series) and uncertainties were calculated
305 over the period 2015-2021 using the mean and min/max between CTE-GCP2021, MIROC4-ACTM both runs,
306 CAMS v21r1 (both runs), and TM5-4DVAR (TROPOMI based). CEOS (GOSAT) provided an estimate only
307 for 2019.

308 The units used in this paper are metric tons (t) [1kt = 10⁹ g; 1Mt (Tg) = 10¹² g] of CH₄. The referenced
309 data for replicability purposes are available for download at <https://doi.org/10.5281/zenodo.1258266740276087>
310 (Petrescu et al., 2023b). Upon request, the computer code for plotting figures in the same style and layout can be
311 provided. Throughout the paper and mostly for the complex figures, the following ISO3 country codes are used:
312 USA (United States of America), BRA (Brazil), CHN (China), IDN (Indonesia), RUS (Russia), COD (DR
313 Congo) and IND (India). Next to these we also refer to CHE (Switzerland) and AUS (Australia). The European
314 Union consists of 27 Member States, excludes the United Kingdom (UK) and is further abbreviated as EU. All
315 abbreviations are summarized in the SI, Table S5.

316 **3. Results**

318 **3.1. NGHGI official reported estimates (UNFCCC)**


319

320 Figure 1 presents anthropogenic CH₄ emissions reported to the UNFCCC in 2023 from the NGHGI
321 CRFs (EU, USA and Russia) and BURs (Brazil (4th in 2021), China (2nd in 2019), Indonesia (3rd in 2021), DR
322 Congo (1st in 2022) and India (all three BURs). The following section provides additional details for all the
323 countries.

324 For the EU, the total anthropogenic CH₄ emissions in 2021 amount to ~~tofer~~ 14.8 ± 1.8 Tg CH₄ yr⁻¹ and
325 represent 12.8 % of the total EU greenhouse gas emissions (in CO₂ equivalents, GWP 100 years, IPCC AR5³).
326 CH₄ emissions are predominantly from agriculture (Figure 1, brown), which accounted for 8.3 Tg CH₄ yr⁻¹ ± 0.8
327 Tg CH₄ yr⁻¹ or 56 % of the total EU CH₄ emissions in 2021 (incl. LULUCF). Anthropogenic CH₄ emissions from
328 the LULUCF sector are very small for the EU: -0.5 Tg CH₄ yr⁻¹ or 3 % in 2021, including emissions from biomass
329 burning. The EU data from Figure 1 shows steadily decreasing trends for all sectors with respect to the 1990
330 levels. The reduction in total CH₄ emissions in 2021 with respect to 1990 is 8.9 Tg CH₄ yr⁻¹ (37 %) at an average
331 yearly rate of -1%.

332 In 2021, the USA reported anthropogenic CH₄ emissions of 28.3 Tg and, compared to 1990, the reported
333 USA CH₄ emissions show a small decrease of 4.6 Tg CH₄ yr⁻¹, more pronounced for the last two years (2020-
334 2021), with an average reduction rate of -0.5 % per year (Fig. 1 black dotted line). In the USA, the largest share
335 of emissions comes from the energy sector (38%), and next to IPPU and waste, had the highest reductions since
336 1990 (42%, 34% and 26%, respectively). Emissions from agriculture (35%, the second largest sector) and
337 LULUCF increased 16 % and 23-%, respectively. CH₄ emissions have been slowly declining since 1990 but had
338 a notable decrease of 1.5 Tg CH₄ yr⁻¹ in 2016 compared to 2015, before increasing again and had a second
339 decreasing trend in 2020 and 2021, possibly due to the COVID pandemic. Overall, reported data indicates that
340 reductions in the USA CH₄ emissions have declined more slowly than that in the EU. The EU also has much
341 lower CH₄ emissions than the US on a per capita basis (Figure 2).


³ IPCC AR4 GWP 100 values are still used by the Member States in their NGHGI reporting to the UNFCCC.



342

343 *Figure 1: Total and sectoral CH₄ emissions (incl. LULUCF) from the UNFCCC NGHGI (2023) CRFs (EU, USA*
 344 *and Russia) and BURs (Brazil (4th in 2021), China (2nd in 2019), Indonesia (3rd in 2021), DR Congo (1st in 2022)*
 345 *and India (all three BURs: 2016, 2018 and 2021). The relative error on the UNFCCC value represents the*
 346 *NGHGI (2023) reported uncertainties computed with the error propagation method (95% confidence interval)*
 347 *and gap-filled to provide respective estimates for each year. Information on Indonesian sectoral CH₄ emissions*
 348 *in 2017 and 2018 are only available for Agriculture. The overall uncertainty of Indonesia's National GHG*
 349 *inventory with AFOLU (including peat fires) for 2000 and 2019 were approximately 20.0% and 19.9%,*
 350 *respectively. In 2014, China reported uncertainty as well (min 5.2 % and max 5.3 %).*

351 The trend in total anthropogenic CH₄ emissions in *Brazil* is strongly increasing, with 32.5 % more
 352 emissions in 2016 compared to 1994. Given that the Brazilian BUR inventory does not include data between
 353 2001 and 2019, it is difficult to discuss the yearly growth rates. We can only note that maximum +22% in 2010
 354 compared to 2000, minimum increase (+1% 2016 compared to 2015. The agricultural sector (76 % of the total)
 355 was the main driver of the growth, followed by the waste sector (16 % of the total). There are only small CH₄
 356 emissions from the Energy sector (some oil and gas activities). The Brazilian agricultural CH₄ emissions are the
 357 highest of the eight compared to all other countries on a per capita basis (see Figure 2).



358

359 *Figure 2: Per capita emissions (kg) intensity per sector based (IPCC, 2006) on reported emissions and*
 360 *population data from the last reported NIRs as follows: 2021 for the EU, USA and Russia, 2016 for Brazil and*
 361 *India, 2014 for China, 2019 for Indonesia and 2018 for DR Congo.*

362 *China's total anthropogenic CH₄ emissions are much larger than the emissions reported by many*
 363 *developed countries or the entire EU (see Figure 1), but on a per capita basis it is only fifth of the eight countries*
 364 *considered (Figure 2). China's CH₄ emissions have grown 114% from 1995 to 2014, when they reached 32 Tg*
 365 *CH₄. The highest contributions to China's CH₄ emissions are from energy (40%) and agriculture (36%). The*
 366 *rapid growth of China's coal demand has important implications for CH₄ emissions from coal mining or coal*
 367 *mine methane emissions (Gao et al., 2020). The energy and agriculture sectors have respectively increased by*
 368 *214 % and 54 % in 2014 compared to 1994.*

369 *Indonesia's 3rd BUR data (2000 and 2019) show increasing trends in total anthropogenic CH₄ emissions.*
 370 *The time series 2001-2006 derives from the 2nd BUR submitted in 2018. In 2019, Indonesian CH₄ emissions had*
 371 *increased by +44 % compared to 2000, corresponding to 2.6 Tg CH₄ yr⁻¹, an average yearly increase of 3 %, and*
 372 *the sector which contributes the most to this increase is the waste sector, which nearly doubled its emissions in*
 373 *2019 compared to 2000. According to Qonitan et al., 2021, the major solid waste source in Indonesia is the*
 374 *household sector, which contributed 44-75% to total waste generated. The composition of municipal waste*
 375 *consists of 44% food waste, 16% paper, and 14% plastics. CH₄ emissions from the other sectors remained nearly*
 376 *constant. For 2017 and 2018 the Indonesian 3rd BUR does not report total emissions other than agricultural*
 377 *emissions, which were taken from the report Figure 2-24. The last data point (2019) shows lower total emissions*
 378 *because it belongs to the revised versions of the (3rd) BUR while the previous data points 2000-2016 belong to*
 379 *the 2nd BUR.*

380 *Russia's anthropogenic CH₄ emissions have decreased by -25 % from 1990 to 2021, but most of this*
 381 *decrease happened during the dissolution of the Soviet Union. Since 2000, CH₄ emissions have increased*
 382 *slightly, but remain lower than pre-2000 levels. The decline seen between 1990 and 2000 is primarily due to the*
 383 *agricultural sector (-52 %) and energy (-27 %). At the same time, the waste sector started to increase its emissions*

384 (6 %). Between 2001 and 2021, the CH₄ emissions from the agriculture and energy sectors continue to decrease
385 (by 17 % and 11 %, respectively), while the emissions from the waste sector register an additional 76 % increase.
386 IPPU emissions increased by 85 % but remain negligible compared to other sectors. Since the 2000s, LULUCF
387 emissions have also increased, by 53 %.


388 For its first BUR, *DR Congo* submitted emissions from energy, AFOLU (agriculture plus LULUCF)
389 and waste for 2000-2018. Since 2000, the DR Congo total anthropogenic CH₄ emissions have increased by a
390 factor of four. Most of the CH₄ emissions are reported for the waste sector, and account for 90 % of the total
391 emissions. The high percentage of waste emissions in DR Congo is also seen in the per capita emissions (Figure
392 2). Assè-Wassa Sama and Berenger (2023) confirm that between 2000 and 2021, CH₄ emissions, which in 2021
393 represent in DR Congo ~97% of total waste generated emissions, grew at a rate of 4 % yr⁻¹, compared with 2.7
394 % yr⁻¹ for total emissions. This increase was driven by the increase in emissions caused by solid waste disposal
395 (+6.2 %). The CH₄ waste emissions come mainly from the treatment and discharge of wastewater (69 % in 2021,
396 compared with 80 % in 2000), followed by the elimination of solid waste (31 % in 2021, compared with 20 %
397 in 2000). The weight of emissions caused by the elimination of solid waste in the sector's total emissions has
398 nevertheless increased by 11 percentage points between 2000 and 2021 (Assè-Wassa Sama and Berenger, 2023).

399 Each of *India*'s BURs provide detailed information on sectoral CH₄ emissions only for one year. Most
400 of the emissions in India belong to the agriculture sector, amounting to almost 15 Tg CH₄ yr⁻¹ (in 2016),
401 representing 74 % of the total anthropogenic emissions. However, with only three years of reported data
402 available, no clear or notable trend is observed.

403 **3.2. NGHGI compared to other bottom-up estimates**

404

405 Figure 3 shows UNFCCC (CRFs and BURs) estimates from EU and seven non-EU countries compared
406 to global bottom-up inventories. The EU and USA show decreasing trends in emissions from all data sets (except
407 for GAINS in the USA), while all the other countries show increasing trends in all datasets. The match between
408 UNFCCC reported emissions and all other data sources is satisfactory, with a few notable exceptions.



409

410 *Figure 3: Total anthropogenic CH₄ emissions (excl. LULUCF) from bottom-up (BU) inventories as: UNFCCC*
 411 *NGHGIs (2023) of CRFs (EU, USA and Russia) and BURs (Brazil (4th in 2021), China (2nd in 2019), Indonesia*
 412 *(3rd in 2021), DR Congo (1st in 2022), India (all three BURs:2016, 2018 and 2021) and three other global*
 413 *datasets: EDGARv7.0, GAINS (no IPPU) and FAOSTAT (PRIMAP based, except for AFOLU). The relative error*
 414 *on the UNFCCC value represents the NGHGI (2023) reported uncertainties computed with the error*
 415 *propagation method (95% confidence interval) and gap-filled to provide respective estimates for each year.*
 416 *China and Indonesia report uncertainties, for 2014 and 2000 and 2019 respectively (BUR). Total COD*
 417 *UNFCCC BUR emissions do not include IPPU. The EDGARv7.0 uncertainty is only for 2015 and was*
 418 *calculated according to Solazzo et al., 2021 for EDGARv5.0. The mean of overlapping time series was calculated*
 419 *for 1990-last available year as following: 2021 for UNFCCC NGHGI (2023), EDGARv7.0 and FAOSTAT and*
 420 *2020 for GAINS.*

421 For the EU, the difference between the UNFCCC NGHGI 1990-2020 average and the other three data
 422 sets is less than 5 %. As previously discussed, the inventory-based data sources are consistent with each other
 423 for capturing recent CH₄ emission reductions, but they are not independent because they use similar methodology
 424 with different versions of the same [activity data \(AD\)](#) (Petrescu et al., 2020, Figure 4).

425 For the USA, GAINS reports high emissions after 2010, with strong growth. This divergence is largely
 426 found in the Energy sector, resulting from the EFs used for conventional gas production as well as for
 427 unconventional shale gas extraction, which has increased rapidly since 2006 due to the development of hydraulic
 428 fracturing technology (Supplementary Figure S6-1 in Höglund-Isaksson et al., 2020). The high share of
 429 emissions from unconventional shale gas can be explained by the GAINS EFs which, in the absence of published
 430 factors, are derived from the residual emissions after having subtracted estimated emissions for oil production
 431 and conventional gas production from the total upstream emission estimated by Alvarez et al., (2018, Table 1)
 432 As Alvarez et al. 2018 do not specify [emission factors EFs](#) by type of gas produced, GAINSv4 splits it based on
 433 activity data from other references, International Energy Agency-World Energy Outlook (IEA-WEO, 2018) and
 434 [Energy Information Administration \(EIA, 2019\)](#). On the other hand, the NGHGI EF seems to be too low, and

435 this is reflected by the low oil and gas emissions reported by the [Unites States Environmental Protection Agency](#)
436 [\(USEPA 2017\)](#) for 2015, compared to Alvarez et al., 2018 (Supplementary Table S6-3, Höglund-Isaksson et al.,
437 2020). For the USA, total gas production increased by 47 % between 2006 and 2017. Revisions for the
438 agricultural livestock emissions concern updates of AD and reported EFs to statistics from FAOSTAT (2018)
439 and CRFs [\(UNFCCC \(2016; 2018\)\)](#), and a review of available technical abatement options for CH₄.

440 For *Brazil*, UNFCCC and GAINS report emissions of similar magnitudes and trends. The EDGARv7.0
441 and FAOSTAT report on average around 23 % more emissions for the 1990-2021 period, but closely follow the
442 NGHGI trends. The similarity between trends could be explained by the use of the same EFs following Tier-1
443 IPCC 2006 Guidelines and UNFCCC NIRs (Janssens-Maenhout et al., 2019), while the higher emissions could
444 appear when using different AD information.

445 For *China* the inventory estimates agree [reasonably](#) with the BUR reported data, with EDGARv7.0
446 showing the highest estimates. According to both GAINS and EDGARv7.0, the primary drivers for growth in
447 Chinese CH₄ emissions are due to a mix of sources, mainly from the IPCC 2006 sector 1.B.1, fugitive emissions
448 from solid fuels activity linked to increased coal mining.

449 In *Indonesia* the three global datasets agree well up until 2010. From 2010, the third-party [datasets show](#)
450 a continued increase in emissions, while the UNFCCC BUR emissions suggest a decline. EDGARv7.0 reports a
451 large increase in emissions from fugitive emissions from solid fuels (coal mining) (IPCC 2006, sector 1.B.1.) at
452 an increased average rate of 19 % per year and has increased by a factor of 152 until 2021 compared to 1990
453 (Figure 3).

454 For *Russia*, GAINS emissions are much higher than NGHGI and the other two data sets due to the
455 revisions of the assumptions on the average composition of the associated gas generated from oil production
456 based on information provided in Huang et al. (2015). The higher emissions in GAINsv4 might be caused by a
457 greater source from venting of associated gas instead of flaring. GAINsv4 estimates a decline in global CH₄
458 emissions in the first half of the 1990s, primarily a consequence of the ~~collapse~~ [dissolution](#) of the Soviet Union
459 and the associated general decline in production levels in agriculture and fossil fuels (see regional emission
460 illustrations in figures S2–1 of the SI). In addition, as described by Evans and Roshchanka (2014) and assumed
461 in Höglund-Isaksson (2017), venting of associated petroleum gas declined significantly in Russia due to an
462 increase in flaring. It is unclear why this happened, but a possible explanation could be that the privatization of
463 oil production in this period meant that the new private owners were less willing to take the security risks of
464 venting and invested in flaring devices to avoid potential production disruptions. This hypothesis is however yet
465 to be confirmed (Höglund-Isaksson et al., 2020). FAOSTAT data for the Russian Federation starts in 1992, [but](#)
466 [the former USSR statistics were used prior to 1992 without adjustments and this is the cause of the 1990 and](#)
467 1991 outliers in time series. The slightly increasing trend observed in EDGARv7.0 and FAOSTAT are set by
468 emissions from the Energy sector.

469 For *DR Congo* estimates from GAINS are not available because they only report aggregated emissions
470 from a few African regions. Both FAOSTAT (PRIMAP based) and EDGARv7.0 estimates show similar slowly
471 increasing trends, potentially indicating the use of similar prior statistics (EFs). For non-AFOLU sectors the
472 PRIMAP-hist third party data priority scenario used in FAOSTAT also uses EDGAR data as an input data source

473 explaining similarities in these sectors. On the other hand, UNFCCC BUR data reports a strong increase in
474 emissions, which is due to a rapid growth of CH₄ emissions from the Waste sector, by a factor of four until 2018
475 compared to 2000. This increase happened at an average yearly rate of +8 %, with an initial sharp increase of
476 +30 % between 2000 and 2001. As previously discussed, (section 3.1.) we believe that DR Congo BUR reported
477 waste emissions are improbable and further investigation is needed.

478 For *India*, all bottom-up global inventories show similar trends and magnitudes of anthropogenic CH₄
479 emissions. The emissions of CH₄ averaged across EDGARv7.0, GAINS and FAOSTAT are 67% (2010), 68 %
480 (2014) and 65 % (2016) higher than the Indian BURs. All three BU inventories show an averaged steady increase
481 of 1 % yr⁻¹ between 1990-2020.


482 3.3. NGHGs compared to TD atmospheric-based CH₄ estimates

483

484 In Figure 4 we compare the reported TD anthropogenic estimates, after removing natural emissions,
485 with the UNFCCC official reported emissions for the EU and the seven non-EU emitters. The mean column on
486 the right of each chart represents the mean of the overlapping time series (2009-last available year, except for
487 TROPOMI, which was available only for 2018-2020). For the EU, the three regional inversions report total
488 emissions because they do not separate anthropogenic from natural emissions. Therefore, we subtracted from the
489 total the natural emissions as calculated in Petrescu et al., 2023a which amount to 6.6 Tg CH₄ yr⁻¹ and are the
490 sum of biomass burning, wetlands, geological and inland water CH₄ emissions. For the global inversions, the
491 anthropogenic estimates were calculated by subtracting from the total fluxes the reported natural partitions as
492 follows: for the two CAMS inversions and TM5-4DVAR (TROPOMI based) the sum of biomass burning and
493 wetlands, for MIROC4-ACTM runs the natural is the represented by the sum of the biomass burning, geologic,
494 ocean, termites, soils and wetlands, for CTE-GCP2021 the sum of the biologic (wetlands + soils) and other
495 (ocean, termites, geological). Because not all inversions report the same partitions, we consider this a coarse
496 comparison, and we detail the harmonization of the natural emissions in the next section (Table 4 and Figure 7).
497 For China, the last BUR is available for 2014, and therefore we used that value. ~~The inversions show total CH₄~~
498 ~~emissions, including both anthropogenic and natural sources. We present here the total TD estimate against the~~
499 ~~anthropogenic NGHGs, emphasizing that the difference between BU and TD estimates might be due to the~~
500 ~~natural emissions.~~

501

502



503

504 *Figure 4: Anthropogenic CH₄ emissions from UNFCCC NGHGI (2023) CRFs (EU, USA and Russia) and BURs*
 505 *(Brazil (4th in 2021), China (2nd in 2019), Indonesia (3rd in 2021), DR Congo (1st in 2022), India (all three BURs:*
 506 *2016, 2018 and 2021) and ~~total~~ TD estimates as following: for EU anthropogenic regional inversions*
 507 *(FLEXkF_v2023, CIF-FLEXPART and CIF-CHIMERE) and for global inversions anthropogenic estimates*
 508 *calculated as total TD minus natural TD reported partitions (TM5-4DVAR, CAMSv21r1_NOAA,*
 509 *CAMSv21r1_NOAA_GOSAT, CTE-GCP2021 and MIROC4-ACTM both runs) products. The relative error on the*
 510 *UNFCCC value represents the NGHGI (2023) reported uncertainties computed with the error propagation*
 511 *method (95% confidence interval) and gap-filled to provide respective estimates for each year. China reports*
 512 *uncertainties for 2014 (min 5.2 %, max 5.3 %) and Indonesia reports for 2000 and ~~2019,2019~~, 20 % and 19.9*
 513 *% respectively. Total COD UNFCCC BUR emissions do not include IPPU. The last available years are CIF-*
 514 *CHIMERE (2022), TM5-4DVAR, CIF-FLEXPART and CTE-GCP2021 (2020) and FLEXkF_v2023, MIROC4-*
 515 *ACTM both runs, UNFCCC CRFs, and CAMSv21r1 both runs (2021). The mean of overlapping time series was*
 516 *calculated for 2009-2021, except for TM5-4DVAR (2018-2020).*

517

518 In the EU, the average anthropogenic CH₄ emissions from global inversions for 2009-2021 were 19 Tg
 519 CH₄ yr⁻¹ with a min-max range of 15-23 Tg CH₄ yr⁻¹, in line with previous estimates published in Petrescu et al.
 520 (2021, 2023a) and the recent RECCAP2 European GHG budgets study of Lauerwald et al., 2024. This is
 521 consistent with the UNFCCC NGHGI (2023) which report for the same period anthropogenic emissions of (15.8
 522 ± 1.8 Tg CH₄ yr⁻¹), noting the uncertainty ranges of both estimates overlap. There is good agreement in trends,
 523 but with inversions showing a larger year to year variability. The regional inversions, for the same period, report
 524 averaged emissions of 14 Tg CH₄ yr⁻¹ with a min-max range of 13-20 Tg CH₄ yr⁻¹. We note that the regional
 525 inversions tend to report slightly lower emissions than the at-of-global inversions, closer to the UNFCCC
 526 estimates. One reason could be that regional inversions use better-constrained regional observations (e.g. ICOS,

527 not just NOAA), have higher spatial resolution, and may thus better resolve the transport. However, they may
528 still have problems with the regional boundary conditions.

529 For the *USA*, averaged over the period 2009-2021, inversions indicate anthropogenic CH₄ emissions of
530 30 Tg CH₄ yr⁻¹ with min-max range of 26-35 Tg CH₄ yr⁻¹, well in line with the UNFCCC NGHGs (2023) which
531 for the same period report anthropogenic total emissions of 29 Tg CH₄ yr⁻¹. The trends observed in TD products
532 are slightly increasing after 2010, except for CAMS which shows no trend (Figure 4). The striking discrepancy
533 between the trends from CAMS and those from MIROC4-ACTM and CTE-GCP2021 are most likely caused by
534 the increasing oil and gas emissions from the Eastern USA (Permian Basin). The same increasing trend is also
535 captured by GAINS (Figure 3). In their runs, both MIROC4-ACTM and CTE-GCP2021 use oil and gas priors
536 from GAINS, while CAMS uses priors from EDGAR (Figure 3). We discuss further differences in having CTE-
537 GCP2021 run with both EDGAR and GAINS oil & gas prior estimates in the SI.

538 For *Brazil*, inversions yield an average (range) of anthropogenic CH₄ emissions of 23 (17-27) Tg CH₄
539 yr⁻¹, slightly higher than the UNFCCC estimate of 16.6 Tg CH₄ yr⁻¹. The two CAMS inversions have trends
540 which match the trend of the UNFCCC reports estimates.

541 For *China*, approximately 80 % of the CH₄ emission increase (21.5 Tg yr⁻¹) during 2000 – 2015 was
542 from fugitive emissions from coal (mines), consistent with what GAINS and EDGAR reports (Figure 3). The
543 TD estimates mostly agree with the BURs, except for CAMS inversions which show 10 to 20 Tg CH₄ yr⁻¹ higher
544 emission than the other inversions. Both MIROC4-ACTM runs (control and OH inter-annual variability (IAV)
545 varying run; Patra et al., 2021) are in line with the BURs. Trend wise, all inversions agree on increased emissions
546 after 2019, with a slight decrease- after 2013 which picks up again after 2018 seen in all inversion trends.

547 For *Indonesia*, most TD results agree on the trend and show a slight increase in emissions. A similar
548 trend is also seen by the BURs. However, the CAMS inversions show linear increased trends while the other
549 inversions have a more variable trend. Regarding the East Asian estimates, MIROC4_ACTM inversion simulates
550 higher fluxes compared to the other inversions. Only recently they found that annual total East Asian emissions
551 have lowered more significantly than in Patra et al. (2016) or Chandra et al. (2021), therefore new runs with
552 updated input set-ups are currently being investigated.

553 For *Russia*, the estimates from MIROC4-ACTM runs and CTE-GCP2021 are both in the same range as
554 the BU GAINS estimate (see Figure 2) from 2000 onwards (between 30-40 Tg CH₄ yr⁻¹) but does not show such
555 a strong decrease as GAINS from 1990 to 2000), while CAMS runs report about 10 Tg CH₄ yr⁻¹ lower emissions
556 than the other two inversions, matching the UNFCCC estimates. The reason for higher estimates reported by
557 CTE-GCP2021 and MIROC4-ACTM is most likely the use of oil and gas priors from GAINS.

558 For *DR Congo*, inversions show the same slightly increasing trend, similar to that of UNFCCC BURs,
559 without the abrupt increase after 2010. The inversions appear to confirm the overreported growth in emissions
560 from waste.

561 For *India*, all the TD anthropogenic estimates agree well on increased trends and magnitudes. In
562 contrast, UNFCCC reporting does not show any trend, but given the insufficient data from BURs, a plausible
563 conclusion cannot be drawn.

564

565 **3.4. Sectoral attribution of CH₄ emissions in TD products**

566

567 In some cases, inversions can be used to partition emissions to different sources. Table 2 shows the
568 partitions as originally reported by some of the inversions, which we name here “unharmonized partitions”. A
569 straightforward, direct comparison of the fluxes is not possible because of the different ways each inversion
570 allocates and groups the natural/anthropogenic fluxes. For example, not all inversions report soil fluxes as done
571 by MIROC4-ACTM and CTE-GCP2021 (together with wetlands); or report the biomass burning fluxes
572 separately from anthropogenic emissions (MIROC4-ACTM and TM5-4DVAR). Rice is also sometimes
573 allocated to natural emissions. Termites, oceans and geological fluxes are sometimes reported separately
574 (MIROC4-ACTM) or grouped in “Other” (CTE-GCP2021, TM5-4DVAR). Regarding the anthropogenic
575 emissions, TM5-4DVAR reports them as other, providing a separate partition for rice. Figure 5 shows the
576 UNFCCC NGHGI anthropogenic total reported estimate (diamond) next to all TD estimates. All global
577 inversions report total and disaggregated partitions, while the regional inversions report only the total emissions
578 (green column).

579
580 *Table 2: Unharmonized partitions originally reported by inverse products:*

Inversion	Anthropogenic	Rice	Soils	Wetlands	Ocean	Termites	Geological	Biomass burning	Other
CAMsv21r1 (both runs)	Yes (in-as Other)	Yes	No	Yes	Yes (in Other)	Yes (in Other)	No	Yes	Yes**
MIROC4-ACTM (control and OH var)	Yes ((Agr, Waste, Oil/Gas, Biofuel, coal)	Yes (in Agr.)	Yes	Yes	Yes	Yes	Yes	Yes	Yes (separated)
CTE-GCP2021*	Yes (Agr, waste, fossil fuel, biofuel, biomass burning)	Yes (in Agr.)	Yes (BIO)		Yes (in as Other)	Yes (in-as Other)	Yes (in-as Other)	In-as anthr.	Yes (Ocean, Termites, Geological)
CEOS (GOSAT)	Yes (Livestock, rice, waste, coal, oil, fire)	In-as anthr.	No	Yes	No	No	Yes (seeps)	In-as anthr.(but separate)	only seeps
TM5-4DVAR (TROPOMI)	Yes (in-as Other)	Yes	No	Yes	Yes (in as other)	Yes (in-as other)	Yes (in-as Other)	Yes	Yes**
GEOS-Chem CTM (TROPOMI for USA)	Yes (Livestock, Oil Gas, Landfills, Wastewater, Other anthro (rice)	In-as other anthr.	No	Yes	Yes (in as Other)	Yes (in-as Other)	Yes (in-as Other)	Yes (In as Other)	Yes***


581 *CTE-GCP2021 partitions refer to anthropogenic, bio and other.

582 ** In TM5-4DVAR (similar to the CAMsv20 set-up and CAMsv21r1), the "Other" partition includes anthropogenic sources
583 except for the rice paddies. It also includes the small fluxes from termites, oceans, soil sink, geological etc.). More details on
584 priors are found in Petrescu et al., 2023b, Priors table.

585 ***Named Other biogenic

586
587 Figure 5 shows the UNFCCC NGHGI anthropogenic total reported estimate (diamond) next to all TD
588 estimates. All global inversions report total and disaggregated partitions, while the regional inversions report
589 only the total emissions (green column).

UNFCCC anthropogenic and unharmonized CH_4 emissions from partitions reported by TD estimates (average 2015-last available year)



5

592 *Figure 5: Total (green) and disaggregated anthropogenic and natural CH_4 emissions from TD estimates*
 593 *compared to UNFCCC NGHGI (2023) anthropogenic emissions (incl. LULUCF) (diamond) for the EU and*
 594 *seven global emitters outside the EU (USA, Brazil, China, Indonesia, Russia, DR Congo and India). The*
 595 *UNFCCC anthropogenic value represents the sum of all five IPCC sectors (Energy, IPPU, Agriculture,*
 596 *LULUCF and Waste). The partitions reported by the TD global inversions are detailed in Table 2. The relative*
 597 *error on the UNFCCC CRF value represents the NGHGI (2023) reported uncertainties computed with the error*
 598 *propagation method (95% confidence interval) and gap-filled to provide respective estimates for each year (see*
 599 *Petrescu et al., 2023a, Appendix). China value and uncertainties (min 5.2 %, max 5.3 %) are for 2014 only and*
 600 *Indonesia uncertainties for 2019, 19.9 %. For the USA CEOS (GOSAT) we used the Nessar et al., 2023 total*
 601 *uncertainty of min 1.1 and max 1 $Tg\ yr^{-1}$. CTE-GCP2021 provides uncertainties for each partition, but here the*
 602 *uncertainty of the total flux is shown. FLEXkF_v2023 reports the relative uncertainty (%) of the posterior*
 603 *emissions. The plotted data represents the average between 2015 and last available year as follows: CIF-*
 604 *CHIMERE (2022), TM5-4DVAR, CIF-FLEXPART and CTE-GCP2021 (2020) and FLEXkF_v2023, MIROC4-*
 605 *ACTM both runs, UNFCCC CRFs, and CAMSv21r1 both runs (2021). GEOS-Chem CTM (TROPOMI) USA*
 606 *reports only for 2019 (Nessar et al., 2023).*

607 Since the different models define sectors differently, also whether they are natural or anthropogenic,
608 harmonization is required to make them comparable. CTE-GCP2021 reports the net natural land-biosphere flux
609 “BIO flux” (soil+wetlands), while other inversions report wetlands and soil separately. Rice emissions are
610 sometimes a part of the agriculture component (anthropogenic partition) (MIROC4-ACTM, CTE-GCP2021)
611 while CEOS (GOSAT) and GEOS-Chem CTM (USA TROPOMI) report separate partitions for rice in
612 anthropogenic emissions, while CAMS reports rice separate from anthropogenic and natural. Same for the
613 biomass burning - CTE-GCP2021 and CEO report it as part of anthropogenic emissions, while GEOS-Chem
614 CTM as part of Others. The rest of the inversions report it separately; this different allocation makes comparisons
615 for these two sources challenging. To facilitate comparisons between all TD products, we aggregated and
616 harmonized the partitions in three main categories, as summarized in Table 3 and Figure 6. The dark green
617 columns in Figure 6 show the total flux for regional EU inversions which did not report partitions.

618


619 *Table 3: Harmonized partitions from inverse products:*

Inversions	Anthropogenic + Rice + Biomass burning			Soils + Wetlands		Other (Ocean + Termites + Geological)		
	Anthropogenic	Rice	Biomass burning	Soils	Wetlands	Ocean	Termites	Geological
CAMsv21r1 (both runs)	= Other	Yes	Yes	No	Yes	Yes	Yes	Yes
MIROC4-ACTM (control and OH var)	Yes ((Agr (livestock + rice), Waste, Oil/Gas, Biofuel, coal)	In Agr.	Yes, summed to anthr.	Yes	Yes	Yes	Yes	Yes
CTE-GCP2021*	Yes (Agr (rice is in), waste, fossil fuel, biofuel, biomass burning)	in Agr.	In anthr.	Yes (BIO)		Yes (Other)		
CEOS (GOSAT)	Yes (Livestock, rice, waste, coal, oil, fire)	In anthr.	In anthr.	No	Yes	No	No	Yes
TM5-4DVAR (TROPOMI)	Others + Rice+ BB	In anthr.	Yes, summed to anthr.	In Other	Yes	Yes	Yes	Yes
GEOS-Chem CTM (TROPOMI) USA	Yes	In anthr.	In other biogenic	No	Yes	Yes	Yes	Yes

620 *CTE-GCP2021 partitions refer to Anthropogenic, Bio and Other. Other fluxes are imposed

621

UNFCCC anthropogenic and harmonized CH_4 emissions from partitions reported by TD estimates (average 2015-last available year)



622

623 *Figure 6: Total (green) and disaggregated anthropogenic and natural CH_4 emissions from TD estimates*
 624 *compared to UNFCCC NGHGI (2023) anthropogenic emissions (incl. LULUCF) for the EU and seven global*
 625 *emitters (USA, Brazil, China, Indonesia, Russia and DR Congo). The UNFCCC anthropogenic value represents*
 626 *the sum of all five IPCC sectors (Energy, IPPU, Agriculture, LULUCF and Waste). The partitions reported by*
 627 *the TD global inversions are harmonized and detailed in Table 3. The relative error on the UNFCCC CRF value*
 628 *represents the NGHGI (2023) reported uncertainties computed with the error propagation method (95%*
 629 *confidence interval) and gap-filled to provide respective estimates for each year (see Petrescu et al., 2023a,*
 630 *Appendix). In 2014, China UNFCCC value and reported uncertainties (min 5.2 % and max 5.3 %) are for 2014*
 631 *while Indonesia reported uncertainties for 2019, 19.9 %. India UNFCCC value is for 2016. CTE-GCP2021*
 632 *provides uncertainties for each partition, but here we plotted the uncertainty of the total flux. FLEKf_y2023*
 633 *reports the relative uncertainty (%) of the posterior emissions. The plotted data represents the average between*
 634 *2015 and last available reported year as follows: CIF-CHIMERE (2022), UNFCCC CRFs, TM5-4DVAR, CIF-*
 635 *FLEXPART and CTE-GCP2021 (2020) and FLEKf_y2023, MIROC4-ACTM both runs, and CAMSv21r1 both*
 636 *runs (2021). GEOS-Chem CTM (TROPOMI) USA reports only for 2019 (Nesser et al., 2023).*

637 **3.5. 3.4.2. Reconciliation-Comparison of BU and TD CH_4 estimates**

638

639 Figure 7 summarizes the total CH_4 fluxes for the EU and the seven global emitters as following: BU
 640 anthropogenic sources disaggregated per sector, BU natural emissions, TD natural emissions from regional and

641 global inversions, and total emissions from global TD estimates (see 2.3 and SI for description of all data
 642 products). This figure brings all the estimates together to demonstrate the reconciliation process.

643 Inversions currently report in a way that makes comparison between BU natural and TD natural sources
 644 difficult. TD products differ in the sources they report (Table 2) or they allocate them to different categories. We
 645 consider natural the following sources: biomass burning, soils, oceans and termites (often reported by inversions
 646 under category “Other”), wetlands, geological and lakes & reservoirs (or freshwaters). Due to lack of
 647 information, biomass burning emissions were considered among the natural sources, recognizing that in regions
 648 like tropical forests, some of these events are influenced by human intervention. To make the products from
 649 Figure 7 comparable, we added the missing BU information from TD, and vice-versa, presented in hatched
 650 pattern. In this way, comparison between BU and TD natural emission estimates is consistent regarding the
 651 “apples to apples” comparison, but became “apples of different flavors” (see Table 4):

652

653 *Table 4: BU and TD natural partitions as presented in Figure 7:*

Product name	TD natural partitions			
	Reported	Missing* <u>(not in priors)</u>	<u>Not reported**</u>	<u>Missing was added from:</u>
CAMsv21r1_NOAA	BB, wetlands, “Others” include anthropogenic and was not used	termites, oceans, soils, lakes and reservoirs, geological	<u>termites, oceans, soil sink</u>	MIROC4-ACTM (termites, oceans and soils), DAAC lakes and reservoirs, geological, updated for <u>in</u> this study (see SI)
CAMsv21r1_ _NOAA_GOSAT	BB, wetlands, “Others” include anthropogenic and was not used	termites, oceans, soils, lakes and reservoirs, geological	<u>termites, oceans, soil sink</u>	MIROC4-ACTM (termites, oceans and soils), DAAC lakes and reservoirs, geological, updated for <u>in</u> this study (SI)
MIROC4-ACTM control	BB, wetlands, oceans, termites, soils, geological	lakes and reservoirs		DAAC lakes and reservoirs
MIROC4-ACTM_OHvar	BB, wetlands, oceans, termites, soils, geological	lakes and reservoirs		DAAC lakes and reservoirs
CTE-GCP2021	soils + wetlands (BIO), termites and oceans	<u>lakes and reservoirs</u>	<u>BB, geologic</u>	<u>DAAC lakes and reservoirs</u> BB from GFEDv4.1s
CEOS (GOSAT)	Fires (BB), Seeps and wetlands	termites, oceans, soils, lakes and reservoirs		MIROC4-ACTM (termites, oceans and soils), DAAC lakes and reservoirs
TM5-4DVAR (TROPOMI)	BB and wetlands	oceans, termites, soils, lakes and reservoirs, <u>geological,</u>	<u>termites, oceans, soil sink</u>	MIROC4-ACTM (termites, oceans and soils), DAAC lakes and reservoirs, geological, <u>updated in this study (see SI)</u>
Product name	BU natural partitions			
		available <u>Reported</u>	Not reported**	added -Added from

<u>Biomass burning</u> <u>Lakes and reservoirs</u> <u>Wetlands</u> Geological	GFEDv4.1s DAAC LPJ-GUESS Geological <u>emissions</u> updated in this study (SI)	soils termites oceans	MIROC4-ACTM
--	--	-----------------------------	-------------

654 note: in TD products termites, oceans emissions are imposed from existing literature

655 * missing = not in the priors, presented as hatched pattern in the figure “\\”

656 **Not reported** = data not available, presented as hatched pattern in the figure “//”

657

658 For an easier visual comparison and reconciliation between BU and TD estimates, we added the mean
659 of the BU anthropogenic estimates (off-white), underneath the BU and TD natural estimates. To note that for
660 some countries (e.g. Russia, DR Congo) this area might look like subtracted from the BU natural estimates, but
661 this is due to the sign convention used in this study (sink = negative and source = positive). In most cases, the
662 missing soil sink emissions are represented as a downward area.


663 We note that for most countries, the sum of the anthropogenic and natural components matches those
664 of the TD global total estimates. This gives confidence that, to a certain extent and albeit with inconsistencies
665 between products, BU anthropogenic emission estimates are accurate and consistent with the observation-based
666 estimates and can be used to reconcile with the atmospheric-based estimates. We note from Figure 7 that in all
667 Annex I countries (EU, USA, Russia) and China, TD and BU natural emissions are consistent with each other,
668 after including the missing sources, as detailed in Table 4. For Brazil and DR Congo, the gap between the two
669 natural components is highly significant, while less for Indonesia and India. We hypothesize that mapping of the
670 wetland extent might cause these inconsistencies.

671

672

673

Total anthropogenic and natural CH₄ emissions from UNFCCC, other BU and TD estimates (average 2015-last available year)



67.

675 Figure 7: Total anthropogenic and natural CH₄ emissions from BU and TD estimates presented as average of
 676 2015-last available year for EU and seven global emitters (USA, Brazil, China, Indonesia, Russia, DR Congo
 677 and India). The BU anthropogenic estimates belong to: UNFCCC NGHGI (2023) CRFs and BURs (incl.
 678 LULUCF) as totals and sectoral shares, EDGARv7.0, GAINS and FAOSTAT/PRIMAP-hist. The relative error
 679 on the UNFCCC CRF value represents the NGHGI (2023) reported uncertainties computed with the error
 680 propagation method (95% confidence interval) and gap-filled to provide respective estimates for each year (see
 681 Petrescu et al., 2023a, Appendix). In 2014, China reported an uncertainty of min 5.2% - max 5.3%. The BU
 682 Natural emissions for the EU are the sum of the VERIFY products (biomass burning, inland waters, geological
 683 and peatlands plus mineral soils as described in Petrescu et al., 2021 and 2023a, Appendix A2.1). For the seven
 684 non-EU emitters, the BU Natural fluxes are the sum of wetland emissions (LPJ-GUESS), lakes and reservoirs
 685 fluxes (ORNL DAAC, Johnson et al., 2022), geological (updated activity in SI) and biomass burning emissions
 686 (GFED4.1s). The TD natural global estimates are presented in Table 1. The uncertainty on the TD natural
 687 emissions is the min/max of all estimates. To both BU and TD estimates, missing (as not reported or not included
 688 information in the priors) was added (see Table 4). The natural emissions have been plotted starting at the mean
 689 of the BU anthropogenic estimates, to retain comparability across the natural emission estimates, but also
 690 compare with the total TD estimates. The total regional TD estimates (for EU) belong to the mean and min/max
 691 of FELXkF_v2023, CIF-FLEXPART and CIF-CHIMERE and for USA GEOS-Chem CTM (TROPOMI) for the
 692 year 2019 (Nesser et al., 2023). The total global TD inversions represent the average of the 2015-last available

693 year of the mean and min/max of CTE-GCP2021, MIROC4-ACTM both runs, CAMS v21r both runs and TM5-
694 4DVAR. The last available years are 2022 for CIF-CHIMERE, 2021 for EDGARv7.0, FAOSTAT, MIROC4-
695 ACTM both runs, UNFCCC CRFs, and CAMSv21r1 both runs, and 2020 for CIF-FLEXPART and CTE-
696 GCP2021. TM5-4DVAR partitioned data is only available between 2018 and 2020.

697 However, [Figure 7](#) should be interpreted with caution because in Europe, natural emission priors come
698 from regional ecosystem model simulations, where drained peatland, drainage ditches areas, and pristine areas
699 are lumped together. Therefore, if both LULUCF sector and natural BU emissions are included in the total budget
700 estimation, there is some overlap and possible double counting. Especially, ecosystem model estimates of ‘soil
701 sink’ or ‘inundated soil emissions’ may overlap with NGHGI managed peatland forest soil category (or
702 agricultural soils). The separation of emissions into different categories requires further clarification together
703 with inventory makers. Furthermore, it should be assessed which emissions should be called natural and which
704 anthropogenic (e.g., LULUCF, Agriculture) by inversions.

705

706 **4. Discussion and recommendations on reconciliation procedures** **Challenges comparing** 707 **bottom-up and top-down estimates** 708

709 An off-the-shelf comparison of BU and TD estimates is not possible, with a variety of adjustments
710 needed for comparability, often without the necessary data. Broadly speaking, inversions have not necessarily
711 been designed to compare directly to NGHGIs. A valid comparison should have consistent system boundaries
712 and perform a full uncertainty analysis to determine whether differences between estimates are statistically
713 significant given the constraining observational data.

714 The two most common issues limiting comparability are geographic scope and system boundaries
715 (Petrescu et al., 2021, 2023; McGrath et al., 2023, Andrew 2020; Grassi et al., 2018). The **geographical scope**
716 of inverse modelling versus inventory estimates should be controllable, but it can be challenging for small
717 countries or coarse inversions. Inversions are generally performed on a spatial grid and require aggregation, in
718 line with how official NGHGIs are reported (EEA, 2013). Inconsistent system boundaries have implications in
719 comparing the inventory- with inversions-based estimates for **source attribution**, e.g., anthropogenic vs. natural.
720 Most emission inventories aim at estimating anthropogenic emissions, while most inversions estimate both
721 anthropogenic and natural emissions. This is a particularly important issue for CH₄ where, globally, natural
722 emissions are of similar magnitude as anthropogenic emissions, with larger variations at regional scales, mainly
723 due to seasonality (i.e. wetlands). Thus, methods are needed to separate the anthropogenic flux from the total
724 flux (Deng et al. 2022, and above section 3.4). Similar issues arise with fossil CO₂ (Andrew 2020) as different
725 datasets can report different emission sources. Standardization procedures, such as The Community Inversion
726 Framework (CIF; Berchet et al., 2021), may help resolve some of these issues.

727 When comparing inventory- and inversion-based emissions, there are difficulties in analyzing **trends**
728 due to different time scale **variability**. Inventory-based approaches report emissions at the annual level, but often
729 do not consider interannual variations. Further, the Paris Agreement is set around five-yearly Global Stocktakes,
730 which indicates a desire to average trends, prioritizing the multi-annual trend over IAV, canceling out extremes
731 from both weather and socio-economic fluctuations. Inversion models, on the other hand, include variations over

732 a wide range of timescales, but in particular for IAV (e.g. OH and weather) that remains challenging to assess.
733 For an effective comparison, inversion-based estimates need to have IAVs statistically removed to make
734 comparisons with NGHGIs easier (e.g., 5-year or 10-year averages or trend analysis). Additionally, averages of
735 ensembles of inversions may mask underlying differences and trends in individual inversions. Many research
736 projects make use of multi-model ensembles (Saunois et al., 2020; Deng et al. 2022, Lauerwald et al., 2024,
737 Zhang et al., 2024). From a scientific perspective, the model ensemble is often considered a more robust estimate
738 of the mean and uncertainty, as often individual estimates make errors due to some limitations and biases, while
739 in an ensemble, these errors are averaged out. From an inventory perspective, individual model comparisons may
740 be more efficient, as various input variables or processes can be compared directly to the inventory. Currently,
741 most inventory comparisons in UNFCCC NGHGI (e.g. UK, Switzerland) use single-model comparisons.

742 A strength of inversions is that they provide high temporal and spatial resolutions, which are not directly
743 capitalized when comparing with inventories. CH₄ from the fossil-fuel industry can contribute to large releases
744 to the atmosphere over a short period of time, given the large number of uncontrolled emission point sources in
745 oil and gas (O&G) and coal production areas worldwide (Jackson et al., 2020). Such processes include leakage
746 from landfills, spontaneous events from oil and gas production activities, so-called uncontrolled gas well blasts
747 etc. (Jacob et al., 2016, 2022). These uncontrolled events are difficult to include in the national inventories
748 leading to a potential underestimate of emissions (Massackers et al., 2016, 2022). Recently, under the CoCO₂
749 project (<https://coco2-project.eu/>) a hot-spot satellite detection interactive map (Published studies on hot spot
750 detection (CO₂, CH₄) - uMap (openstreetmap.fr) was released as a user-centric interface featuring published
751 studies on hot-spot detection between 2010 and 2021. It allows for advanced filtering by year, gas, activity,
752 geographical zone, and country.

753 A key challenge when comparing inversions with NGHGIs is ensuring independence from the assumed
754 prior emissions. A more valid comparison between inversions is made when all inversions use the same **priors**.
755 In this context, we define as priors input data in the form of atmospheric observations (e.g. satellite retrievals,
756 ground-based observation networks (ICOS)) and/or bottom-up emissions datasets (e.g. EDGAR, GAINS) used
757 as input parameters to the inverse models. A key issue is the prior emission estimate. Theoretically, a constant
758 emission prior could be used, but this would require a dense observational network. Because of sparse
759 observations, inversion modelers assess how far observations have shifted the prior emissions to the posterior
760 emissions, preferably incorporating a full uncertainty analysis. The posterior emissions depend to a varying
761 extent on the prior that was used; the extent of this dependency is determined by the number of observations
762 used in the inversion, by how the observations relate to the emissions (governed by atmospheric transport) and
763 by the uncertainties assigned on the prior emissions and the observations. Thus, better quantified uncertainties
764 for the prior emissions would lead to more robust ~~inversions would be more robust with better quantified~~
765 ~~uncertainties for the prior emissions~~. Whereas the comparison of an inversion with NGHGIs or other inversions
766 would be made more robust by having more information on how dependent the posterior estimate is on the prior.
767 This stresses the need for more systematic in-situ data measurements of fluxes necessary to produce adequate
768 prior data (Bastviken et al., 2022) and synthesized atmospheric observations with their uncertainties to robustly
769 constrain the inversions.

770 It is not generally clear how inventory uncertainties can be compared to inversion uncertainties; however, it is
771 important that both methods provide comprehensive **uncertainty estimates**. The prior emissions used as input
772 into an inversion model should have robust uncertainty estimates, particularly with correlations in space and
773 time. This allows a full inversion system to better characterize how observations reduce uncertainty when
774 estimating the posterior estimate. Very few inversions routinely report this information. The inventory-based
775 emission estimate will additionally have uncertainty estimates, though these statistics may not be sufficiently
776 robust for verification purposes (National Academies of Sciences, Engineering, and Medicine 2022). There are
777 often offsets in inversion models, because of systematic inconsistencies between observations and chemistry-
778 transport models, which may make trends more robust than instantaneous estimates. Though, estimating
779 uncertainty in trends also requires understanding the correlation structure in time.

780 A key challenge for comparisons between NGHGI and independent estimates is to understand the
781 reasons for differences. In the case of BU comparisons, obtaining sufficient activity data and emission factors
782 should enable an accurate reconciliation of different estimates. However, in practice, it is often not possible to
783 obtain the necessary data. For inversions it is more complex. Often a close collaboration may be needed between
784 the inversion modeler and NGHGI team (e.g. UK NIR). If an inversion indicated a different trend in agricultural
785 CH₄ emissions, it is necessary to track down if this is a real difference or artifact of the inversion system. After
786 this, the spatial and temporal data in the inversion could be useful to the NGHGI team to locate what is causing
787 the difference. Many of the comparisons we show in this article ultimately remain comparisons, with detailed
788 reconciliations likely requiring intensive country-level case studies.

789 **5. Data availability**

790

791 Data files reported in this work which were used for calculations and figures are available for public
792 download at <https://doi.org/10.5281/zenodo.1258266740276087> (Petrescu et al., 2023b). The data are reachable
793 with one click (without the need for entering login and password), with a second click to download the data,
794 consistent with the two click access principle for data published in ESSD (Carlson and Oda, 2018). The data and
795 the DOI number are subject to future updates and only refer to this version of the paper. The raw gridded data is
796 available upon request, directly from the data providers, as detailed in the Supplementary Information, Table S2.

797 **6. Conclusions**

798

799 We analyzed data from both anthropogenic and natural CH₄ fluxes, from both BU and TD observation-
800 based estimates (Table 1). BU estimates show that the largest sectors depend on the country: Agriculture (EU,
801 Brazil, India), Energy (US, China, Russia), and Waste (Indonesia, DR Congo). The inversions attribute most of
802 the fluxes to the anthropogenic emissions, with tropical countries having a larger share of natural emissions
803 (wetlands). The EU and the seven other large emitters analyzed here contribute an anthropogenic emission of
804 173 Tg CH₄ yr⁻¹ (sum of last UNFCCC reported year, Figure 1,2), representing roughly half of the total global
805 anthropogenic emissions (386 Tg CH₄ yr⁻¹) reported by EDGARv7.0 in 2021. For comparison, the average of
806 the anthropogenic component from the atmospheric global inversions (MIROC booth runs, CTE-GCP2021,
807 CEOS and CAMS booth runs) is 181 Tg CH₄ yr⁻¹ (Figure 5).

808 We performed comparisons to the UNFCCC NGHGs using the BU and TD data. Comparisons between
809 UNFCCC and BU products (Figure 3) reveal some deviations, particularly related to assumptions on gas/oil
810 emissions (e.g., GAINS for Russia and the USA) and waste (e.g., Indonesia, DR Congo). It is more challenging
811 to compare BU and TD estimates, due to different attribution to source activities (Table 2 and Figure 5) and
812 different priors used in the simulations (Petrescu et al., 2023b, Priors Table). The comparison between UNFCCC
813 and the TD estimates (Figure 4) agrees largely with the findings of Deng et al. (2022) who applied different
814 methodologies to calculate natural emissions. In most cases, the gap between the anthropogenic BU fluxes from
815 inventories and total TD fluxes can be largely explained by the natural fluxes (Figure 7). It is difficult to draw
816 definitive conclusions on emissions trends seen by inversions, as the adjustments for natural emissions and IAV
817 and seasonal variability might strongly influence trends. Despite this, given that, in most cases, the UNFCCC
818 BURs reports are incomplete for the non-Annex I parties (China, Indonesia, DR Congo) it is important to
819 acknowledge that the TD estimates might become a useful way to complement inventories and play a role in the
820 validation of the BU estimates.

821 There is still a pressing need for reporting of uncertainties in both prior and posterior emissions, even if
822 some TD inversions do report it as the standard deviation of ensemble members (CTE-GCP2021 and
823 FLEXkF_v2023, Figure 4). The use of a variety of priors across different inversion systems can also inhibit
824 comparability with inventories and between inversions. Generally, inversions are still ill-constrained by
825 observations (only 60 sites globally plus satellites) and the prior flux uncertainty for each of the 54 regions is
826 large. Therefore, the monthly results could be more ill-constrained than the annual totals. Even if comparisons
827 between CH₄ inversion estimates and NGHGs are currently uncertain because of the spread in the inversion
828 results, TD inversions inferred from atmospheric observations represent partly independent data against which
829 inventory totals and trends can be compared, considering the recommendations most encountered issues
830 discussed -listed at the end of in section 4.

831

832 **7. Appendix**

833

834 All the information regarding models/methods descriptions is available in the Supplementary Information (SI)
835 file. Appendices A1 and A2 in Petrescu et al., 2023a contain detailed information about Table 1 products. Further
836 information on new products together with references and contact details are found in Tables S1 and S2 in SI.

837

838 The tables with priors used by all the products and the matrix highlighting the comparability issues identified in
839 section 4 are found in the Zenodo data repository, Petrescu et al., 2023b.

840

841 **Supplementary Information (link)**

842

843

844 **Author contributions**

845

846 AMRP designed research and led the discussions; AMRP wrote the initial draft of the paper and edited all the
847 following versions; GPP drafted the initial version of section 4, edited the final version of this manuscript,

848 contributed to the revised version and advised on the context; PP processed all the original EU data submitted to
849 the VERIFY portal; RLT, SH, BM, DaB, RL, PKP, AT, RMA, LHI, FNT, GC and JG edited and gave consistent
850 comments and suggestions to the initial manuscript; [DaB, RL and RMA provided input to the final revised](#)
851 [version](#); all co-authors are data providers and contributed to subsequent versions of the manuscript by providing
852 specific comments and information related to their data in the main text, providing as well product descriptions
853 for the Supplementary Information file.

854 **Competing interests**

855 At least one of the (co-)authors is a member of the editorial board of Earth System Science Data.

856 **Acknowledgements**

857 The lead author would like to thank former colleagues, Dr. Chunjing Qiu and Dr. Matthew McGrath for previous
858 work done for the EU funded VERIFY project. FAOSTAT statistics are produced and disseminated with the
859 support of its member countries to the FAO regular budget. The views expressed in this publication are those of
860 the author(s) and do not necessarily reflect the views or policies of FAO. We acknowledge the work of current
861 and former members of the EDGAR group (Marilena Muntean, Diego Guizzardi, Monica Crippa, Edwin Schaaf,
862 Efisio Solazzo, Gabriel David Orreggioni and Jos Olivier).

864 **Financial support**

865 This research has been supported by the European Commission, Horizon 2020 Framework Programme (CoCO2,
866 grant no. 958927).

867 [GPP was supported by the European Union's Horizon Europe Research and Innovation program under grant](#)
868 [agreement no. 101081395 \(EYE-CLIMA\)](#). Development of MIROC4-ACTM is supported by the Environment
869 Research and Technology Development Fund (grant no. JP-MEERF21S20800) and the Arctic Challenge for
870 Sustainability phase II (ArCS-II; grant no. JP- MXD1420318865) project. DB was supported by FORMAS
871 (Grant No. 2018-01794), the European Union (H2020 Grant No. 101015825; TRIAGE) and the Swedish
872 Research Council VR (Grant No. 2022-03841). PR acknowledges funding from the European Union's Horizon
873 2020 research and innovation program under Grant Agreement No. 101003536 (ESM2025-Earth System Models
874 for the Future) and from the FRS-FRNS PDR project T.0191.23 CH4-lakes. Annual, gap-filled and harmonized
875 NGHGI uncertainty estimates for the EU were provided by the EU GHG inventory team (European Environment
876 Agency and its European Topic Centre on Climate change mitigation). RL acknowledges funding from French
877 state aid, managed by ANR under the "Investissements d'avenir" programme (ANR-16-CONV-0003). WZ was
878 supported by grants from the Swedish Research Council VR (2020-05338) and Swedish National Space Agency
879 (209/19). [Part of JRW's work was carried out at the Jet Propulsion Laboratory, California Institute of Technology,](#)
880 [under a contract with the National Aeronautics and Space Administration \(NASA\). Part of PP's work was](#)
881 [supported by the European Commission H2020 Framework Programme under grant agreement no. 776810](#)
882 [\(VERIFY\)](#).

883

884 **8. References**

- 885 Alvarez, R. A., Zavala-Araiza, D., Lyon, D. R., Allen, D. T., Barkley, Z. R., Brandt, A. R., Davis, K. J., Herndon,
886 S. C., Jacob, D. J., Karion, A., Kort, E. A., Lamb, B. K., Lauvaux, T., Maasakkers, J. D., Marchese, A. J.,
887 Omara, M., Pacala, S. W., Peischl, J., Robinson, A. L., Shepson, P. B., Sweeney, C., Townsend-Small,
888 A., Wofsy, S. C., and Hamburg, S. P.: Assessment of methane emissions from the U.S. oil and gas supply
889 chain, *Science*, 361, 186–188, <https://doi.org/10.1126%2Fscience.aar7204>, 2018.
- 890 Andrew, R. M.: A comparison of estimates of global carbon dioxide emissions from fossil carbon sources, *Earth*
891 *Syst. Sci. Data*, 12, 1437–1465, <https://doi.org/10.5194/essd-12-1437-2020>, 2020.
- 892 Balcombe, P., Speirs, J. F., Brandon, N. P. and Hawkes, A. D.: Methane emissions: Choosing the right climate
893 metric and time horizon, *Environ. Sci.: Processes Impacts* 20, 1323 (2018).
- 894 Bastviken, D., Wilk, J., Duc, N. T., Gålfalk, M., Karlson, M., Neset, T.-S., Neset, Opach, T., Enrich-Prast, A.,
895 & Sundgren, I.: Critical method needs in measuring greenhouse gas fluxes. *Environmental Research*
896 *Letters*. <https://doi:10.1088/1748-9326/ac8fa9>, 2022.
- 897 Bastviken, D., Wilk, J., Duc, N.T., Gålfalk, M., Karlson, M., Neset, T.-S., Opach, T., Enrich-Prast, A., and
898 Sundgren, I.: Critical method needs in measuring greenhouse gas fluxes. *Environmental Research Letters*
899 17, 104009. 10.1088/1748-9326/ac8fa9, 2022.
- 900 Basu, S., Lan, X., Dlugokencky, E., Michel, S., Schwietzke, S., Miller, J. B., Bruhwiler, L., Oh, Y., Tans, P. P.,
901 Apadula, F., Gatti, L. V., Jordan, A., Necki, J., Sasakawa, M., Morimoto, S., Di Iorio, T., Lee, H., Arduini,
902 J., and Manca, G.: Estimating emissions of methane consistent with atmospheric measurements of
903 methane and $\delta^{13}\text{C}$ of methane, *Atmos. Chem. Phys.*, 22, 15351–15377, [https://doi.org/10.5194/acp-22-](https://doi.org/10.5194/acp-22-15351-2022)
904 15351-2022, 2022.
- 905 Berchet, A., Sollum, E., Thompson, R. L., Pison, I., Thanwerdas, J., Broquet, G., Chevallier, F., Aalto, T.,
906 Berchet, A., Bergamaschi, P., Brunner, D., Engelen, R., Fortems-Cheiney, A., Gerbig, C., Groot
907 Zwaafink, C. D., Haussaire, J.-M., Henne, S., Houweling, S., Karstens, U., Kutsch, W. L., Lujikx, I. T.,
908 Monteil, G., Palmer, P. I., van Peet, J. C. A., Peters, W., Peylin, P., Potier, E., Rödenbeck, C., Saunio,
909 M., Scholze, M., Tsuruta, A., and Zhao, Y.: The Community Inversion Framework v1.0: a unified system
910 for atmospheric inversion studies, *Geosci. Model Dev.*, 14, 5331–5354, [https://doi.org/10.5194/gmd-14-](https://doi.org/10.5194/gmd-14-5331-2021)
911 5331-2021, 2021.
- 912 Bergamaschi, P., Karstens, U., Manning, A. J., Saunio, M., Tsuruta, A., Berchet, A., Vermeulen, A. T., Arnold,
913 T., Janssens-Maenhout, G., Hammer, S., Levin, I., Schmidt, M., Ramonet, M., Lopez, M., Lavric, J.,
914 Aalto, T., Chen, H., Feist, D. G., Gerbig, C., Haszpra, L., Hermansen, O., Manca, G., Moncrieff, J.,
915 Meinhardt, F., Necki, J., Galkowski, M., O'Doherty, S., Paramonova, N., Scheeren, H. A., Steinbacher,
916 M., and Dlugokencky, E.: Inverse modelling of European CH₄ emissions during 2006–2012 using
917 different inverse models and reassessed atmospheric observations, *Atmos. Chem. Phys.*, 18, 901–920,
918 <https://doi.org/10.5194/acp-18-901-2018>, 2018.

- 919 Cain, M., Jenkins, S., Allen, M.R., Lynch, J., Frame, D.J., Macey, A.H. and Peters, G.P.: Methane and the
 920 Paris Agreement temperature goals, *Phil. Trans. R. Soc. A.*, **380**2020045620200456,
 921 <http://doi.org/10.1098/rsta.2020.0456>, 2022.
- 922 Carlson, D. and Oda, T.: Editorial: Data publication – ESSD goals, practices and recommendations, *Earth Syst.*
 923 *Sci. Data*, 10, 2275–2278, <https://doi.org/10.5194/essd-10-2275-2018>, 2018.
- 924 Chandra, N., Patra, P. K., Bisht, J. S. H., Ito, A., Umezawa, T., Saigusa, N., Morimoto, S., Aoki, S., Janssens-
 925 Maenhout, G., Fujita, R., Takigawa, M., Watanabe, S., Saitoh, N., and Canadell, J. G.: Emissions from
 926 the Oil and Gas Sectors, Coal Mining and Ruminant Farming Drive Methane Growth over the Past Three
 927 Decades, *J. Meteorol. Soc. Japan. Ser. II*, 99, 2021–015, <https://doi.org/10.2151/jmsj.2021-015>, 2021.
- 928 ~~Chandra, Naveen., Patra, P., Bisht, J., Ito, A., Umezawa, T., Saigusa, N., Morimoto, S., Aoki, S., Janssens-~~
 929 ~~Maenhout, G., Fujita, R., Takigawa, M., Watanabe, S., Saitoh, N. and Canadell, J.G., Emissions from the~~
 930 ~~oil and gas sectors, coal mining and ruminant farming drive methane growth over the past three decades,~~
 931 ~~JOURNAL OF THE METEOROLOGICAL SOCIETY OF JAPAN, ISSN 0026-1165, 99 (2), p. 309-337,~~
 932 ~~JRC123075, 2021.~~
- 933 CoCO₂ project: <https://coco2-project.eu/>, last access October, 2023.
- 934 Crippa, M., Solazzo, E., Huang, G., Guizzardi, D., Koffi, E., Muntean, M., Schieberle, C., Friedrich, R., and
 935 Janssens-Maenhout, G.: High resolution temporal profiles in the Emissions Database for Global
 936 Atmospheric Research, *Sci. Data*, 17, 121, <https://doi.org/10.1038/s41597-020-0462-2>, 2020.
- 937 Deng, Z., Ciais, P., Tzompa-Sosa, Z. A., Saunio, M., Qiu, C., Tan, C., Sun, T., Ke, P., Cui, Y., Tanaka, K., Lin,
 938 X., Thompson, R. L., Tian, H., Yao, Y., Huang, Y., Lauerwald, R., Jain, A. K., Xu, X., Bastos, A., Sitch,
 939 S., Palmer, P. I., Lauvaux, T., d'Aspremont, A., Giron, C., Benoit, A., Poulter, B., Chang, J., Petrescu, A.
 940 M. R., Davis, S. J., Liu, Z., Grassi, G., Albergel, C., Tubiello, F. N., Perugini, L., Peters, W., and
 941 Chevallier, F.: Comparing national greenhouse gas budgets reported in UNFCCC inventories against
 942 atmospheric inversions, *Earth Syst. Sci. Data*, 14, 1639–1675, <https://doi.org/10.5194/essd-14-1639-2022>,
 943 2022.
- 944 EEA Technical report No 20/2013 European Union CO₂ emissions: different accounting perspectives, ISSN
 945 1725-2237, [https://www.google.com/url?q=https://www.eea.europa.eu/publications/european-union-](https://www.google.com/url?q=https://www.eea.europa.eu/publications/european-union-co2-emissions-accounting/download&sa=D&source=docs&ust=1700125746315585&usg=AOvVaw1scG2zBViuEOMu3YxdnJGZ)
 946 [co2-emissions-](https://www.google.com/url?q=https://www.eea.europa.eu/publications/european-union-co2-emissions-accounting/download&sa=D&source=docs&ust=1700125746315585&usg=AOvVaw1scG2zBViuEOMu3YxdnJGZ)
 947 [accounting/download&sa=D&source=docs&ust=1700125746315585&usg=AOvVaw1scG2zBViuEOM](https://www.google.com/url?q=https://www.eea.europa.eu/publications/european-union-co2-emissions-accounting/download&sa=D&source=docs&ust=1700125746315585&usg=AOvVaw1scG2zBViuEOMu3YxdnJGZ)
 948 [u3YxdnJGZ](https://www.google.com/url?q=https://www.eea.europa.eu/publications/european-union-co2-emissions-accounting/download&sa=D&source=docs&ust=1700125746315585&usg=AOvVaw1scG2zBViuEOMu3YxdnJGZ), last access October 2023, 2013.
- 949 Etiope, G. and Schwietzke, S.: Global geological methane emissions: an update of top-down and bottom-up
 950 estimates, *Elem. Sci. Anth.*, 7, 47, <https://doi.org/10.1525/elementa.383>, 2019.
- 951 Evans A and Steven D 2009 An institutional architecture for climate change—a concept paper Report
 952 commissioned by the Department for International Development and produced by Center on International

- 953 Cooperation (New York: New York University) (
- 954 http://envirosecurity.org/gpc/publications/Institutional_architecture_climate_change.pdf)
- 955 FAO: FAOSTAT, Statistics Division of the Food and Agricultural Organisation of the UN, Emissions statistics
- 956 till 2021: <https://www.fao.org/faostat/en/#data/GT> (last access November 2023), 2023.
- 957 FAO: FAOSTAT, Statistics Division of the Food and Agricultural Organisation of the UN—,(
- 958 <http://fao.org/faostat/en/#home>), 2018.
- 959 Gao, J., Guan, C., Zhang, B. :China's CH₄ emissions from coal mining: a review of current bottom-up
- 960 inventories, *Sci. Total Environ.*,
- 961 <https://www.sciencedirect.com/science/article/abs/pii/S0048969720318088>, 2020.
- 962 Geddes, A., Mikaloff-Fletcher, S., Schaefer, H., Smale, D., and Brailsford, G., Law, R., Ausseil, A-G.:
- 963 Atmospheric Methane Modelling: Evaluating Regional Methane Emission Using Inverse Modelling, MPI
- 964 Technical Paper No: 2021/02, <https://www.mpi.govt.nz/dmsdocument/46222/direct>, 2021, last access
- 965 November 2023.
- 966 Grassi, G., House, J., Kurz, W. A., Cescatti, A., Houghton, R. A., Peters, G. P., Sanz, M. J., Vinas, R. A., Alkama,
- 967 R., Arneth, A., Bondeau, A., Dentener, F., Fader, M., Federici, S., Friedlingstein, P., Jain, A. K., Kato, E.,
- 968 Koven, C. D., Lee, D., Nabel, J. E. M. S., Nassikas, A. A., Perugini, L., Rossi, S., Sitch, S., Viovy, N.,
- 969 Wiltshire, A., and Zaehle, S.: Reconciling global-model estimates and country reporting of anthropogenic
- 970 forest CO₂ sinks, *Nat. Clim. Change*, 8, 914–920, <https://doi.org/10.1038/s41558-018-0283-x>, 2018.
- 971 Gütschow, J. and Pflüger, M.: The PRIMAP-hist national historical emissions time series (1750-2021) v2.4.
- 972 Zenodo: <https://doi.org/10.5281/zenodo7179775>, 2022.
- 973 Höglund-Isaksson L., Gómez-Sanabria, A., Klimont, Z., Rafaj, P., and Schöpp, W.: Technical potentials and
- 974 costs for reducing global anthropogenic methane emissions in the 2050 timeframe – results from the
- 975 GAINS model, *Environ. Res. Commun.*, 2, 025004, <https://doi.org/10.1088/2515-7620/ab7457>, 2020.
- 976 Höglund-Isaksson L.: Bottom-up simulations of methane and ethane from global oil and gas systems *Environ.*
- 977 *Res. Lett.* 12 024007, 2017.
- 978 Huang K, Fu J. S., Prikhodko V. Y., Storey J. M., Romanov A., Hodson E. L., Cresko J., Morozova I., Ignatieva
- 979 Y. and Cabaniss J.:Russian anthropogenic black carbon: Emission reconstruction and Arctic black carbon
- 980 simulation *Journal of Geophysical Research: Atmospheres*, 2015.
- 981 IPCC Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth
- 982 Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, R.K.
- 983 Pachauri and L.A. Meyer (eds.)]. IPCC, Geneva, Switzerland, 151 pp., 2014.
- 984 IPCC, 2023: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth
- 985 Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and

- 986 J. Romero (eds.)]. IPCC, Geneva, Switzerland, pp. 35-115, doi: 10.59327/IPCC/AR6-9789291691647,
987 2023.
- 988 IPCC: 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories, available at:
989 [https://www.ipcc.ch/report/2019-refinement-to-the-2006-ipcc-guidelines-for-national-greenhouse-gas-](https://www.ipcc.ch/report/2019-refinement-to-the-2006-ipcc-guidelines-for-national-greenhouse-gas-inventories)
990 [inventories](https://www.ipcc.ch/report/2019-refinement-to-the-2006-ipcc-guidelines-for-national-greenhouse-gas-inventories) (last access: January 2020), 2019.
- 991 IPCC: Guidelines for National Greenhouse Gas Inventories (NGHGI), available at: [https://www.ipcc-](https://www.ipcc-nggip.iges.or.jp/public/2006gl/)
992 [nggip.iges.or.jp/public/2006gl/](https://www.ipcc-nggip.iges.or.jp/public/2006gl/) (last access: January 2020), 2006.
- 993 IPCC: Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands, edited by:
994 Hiraishi, T., Krug, T., Tanabe, K., Srivastava, N., Baasansuren, J., Fukuda, M., and Troxler, T. G., IPCC,
995 Switzerland, 2014.
- 996 IPCC: vol. 1, chap. 3, [https://www.ipcc-](https://www.ipcc-nggip.iges.or.jp/public/2006gl/pdf/1_Volume1/V1_3_Ch3_Uncertainties.pdf)
997 [nggip.iges.or.jp/public/2006gl/pdf/1_Volume1/V1_3_Ch3_Uncertainties.pdf](https://www.ipcc-nggip.iges.or.jp/public/2006gl/pdf/1_Volume1/V1_3_Ch3_Uncertainties.pdf) (last access: June 2022),
998 2006.
- 999 Jackson, R.B., Saunio, M., Bousquet, P., Canadell, J.G., Pulter, B., Stavert, A.R., Bergamaschi, P., Niwa, Y.,
1000 Segers, A. and Tsutra, A.: Increasing anthropogenic methane emissions arise equally from agricultural
1001 and fossil fuel sources, *Environ. Res. Lett.* 15, 071002, 2020.
- 1002 Jacob, D. J., Turner, A. J., Maasakkers, J. D., Sheng, J., Sun, K., Liu, X., Chance, K., Aben, I., McKeever, J.,
1003 and Frankenberg, C.: Satellite observations of atmospheric methane and their value for quantifying
1004 methane emissions, *Atmos. Chem. Phys.*, 16, 14371–14396, <https://doi.org/10.5194/acp-16-14371-2016>,
1005 2016.
- 1006 [Jacob, D.J., D.J. Varon, D.H. Cusworth, P.E. Dennison, C. Frankenberg, R. Gautam, L. Guanter, J.](https://doi.org/10.5194/acp-22-9617-2022)
1007 [Kelley, J. McKeever, L.E. Ott, B. Poulter, Z. Qu, A.K. Thorpe, J.R. Worden, and R.M. Duren,](https://doi.org/10.5194/acp-22-9617-2022)
1008 [Quantifying methane emissions from the global scale down to point sources using satellite](https://doi.org/10.5194/acp-22-9617-2022)
1009 [observations of atmospheric methane, *Atmos. Chem. Phys.*, 22, 9617–9646,](https://doi.org/10.5194/acp-22-9617-2022)
1010 [https://doi.org/10.5194/acp-22-9617-2022, 2022.](https://doi.org/10.5194/acp-22-9617-2022)
- 1011 Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Dentener, F., Bergamaschi, P.,
1012 Pagliari, V., Olivier, J. G. J., Peters, J. A. H. W., van Aardenne, J. A., Monni, S., Doering, U., Petrescu,
1013 A. M. R., Solazzo, E., and Oreggioni, G. D.: EDGAR v4.3.2 Global Atlas of the three major greenhouse
1014 gas emissions for the period 1970–2012, *Earth Syst. Sci. Data*, 11, 959–1002,
1015 <https://doi.org/10.5194/essd-11-959-2019>, 2019.
- 1016 Johnson, M.S., Matthews, E., Du, J., Genovese, V., and Bastviken, D.: Methane Emission From Global Lakes:
1017 New Spatiotemporal Data and Observation-Driven Modeling of Methane Dynamics Indicates Lower
1018 Emissions. *Journal of Geophysical Research: Biogeosciences* 127, e2022JG006793.
1019 <https://doi.org/10.1029/2022JG006793>, 2022.

- 1020 Lan, X., Basu, S., Schwietzke, S., Bruhwiler, L. M. P., Dlugokencky, E. J., Michel, S. E., Sherwood, O. A., Tans,
 1021 P. P., Thoning, K., Etiope, G., Zhuang, Q., Liu, L., Oh, Y., Miller, J. B., Pétron, G., Vaughn, B. H., and
 1022 Crippa, M.: Improved Constraints on Global Methane Emissions and Sinks Using $\delta^{13}\text{C-CH}_4$, *Global*
 1023 *Biogeochem. Cy.*, 35, e2021GB007000, <https://doi.org/10.1029/2021GB007000>, 2021a.
- 1024 Lan, X., Nisbet, E.G., Dlugokencky, E. J., and Michel, S. E.: What do we know about the global methane
 1025 budget? Results from four decades of atmospheric CH_4 observations and the way forward *Phil. Trans.*
 1026 *R. Soc. A*. **379**2020044020200440, <http://doi.org/10.1098/rsta.2020.0440>, 2021. b
- 1027 Lan, X., Dlugokencky, E., Englund Michel, S., Basu, S., Schuldt, K., Mund, J., Aoki, S., di Sarra, A., Vermeulen,
 1028 A., Andrews, A., Jordan, A., Baier, B., Labuschagne, C., Lund Myhre, C., Sweeney, C., Kubistin, D.,
 1029 Smale, D., Worthy, D., Cuevas, E., Apadula, F., Brailsford, G., Lee, H., Moosen, H., Schaefer, H., Jui,
 1030 H., Necki, J., Arduini, J., Miller, J., Moncrieff, J., Hatakka, J., Uhse, K., McKain, K., Haszpra, L., Gatti,
 1031 L., Ries, L., Steinbacher, M., Schmidt, M., Ramonet, M., Arshinov, M., Sasakawa, M., Paramonova, N.,
 1032 Bergamaschi, P., Langenfelds, R., Kim, S.-Y., Morimoto, S., Takatsuji, S., Nichol, S., Umezawa, T., Di
 1033 Iorio, T., and Kawasaki, T.: Database of methane (CH_4) abundance and its stable carbon isotope ($\delta^{13}\text{CCH}_4$)
 1034 composition from atmospheric measurements, NOAA Global Monitoring Laboratory [data set],
 1035 <https://doi.org/10.15138/64w0-0g71>, 2022.
- 1036
 1037 [Lauerwald, R., A. Bastos, M.J. McGrath, A.M.R. Petrescu, F. Ritter, R.M. Andrew, A. Berchet, G. Broquet, D.](https://doi.org/10.22541/essoar.171320253.37867733/v1)
 1038 [Brunner, F. Chevallier, A. Cescatti, S. Filipek, A. Fortems-Cheiney, G. Forzieri, P. Friedlingstein, R.](https://doi.org/10.22541/essoar.171320253.37867733/v1)
 1039 [Fuchs, C. Gerbig, S. Houweling, P. Ke, B.J.W. Lerink, W. Li, W. Li, X. Li, I.T. Luijkx, G. Monteil, S.](https://doi.org/10.22541/essoar.171320253.37867733/v1)
 1040 [Munassar, G.-J. Nabuurs, P.K. Patra, P. Peylin, J. Pongratz, P. Regnier, M. Saunois, M.-J. Schelhaas, M.](https://doi.org/10.22541/essoar.171320253.37867733/v1)
 1041 [Scholze, S. Sitch, R.L. Thompson, H. Tian, A. Tsuruta, C. Wilson, J.-P. Wigneron, Y. Yao, S. Zaehle, P.](https://doi.org/10.22541/essoar.171320253.37867733/v1)
 1042 [Ciais, 2024. Carbon and greenhouse gas budgets of Europe: trends, interannual and spatial variability, and](https://doi.org/10.22541/essoar.171320253.37867733/v1)
 1043 [their drivers. *ESS Open Archive* . April 15, 2024.](https://doi.org/10.22541/essoar.171320253.37867733/v1)
 1044 <https://doi.org/10.22541/essoar.171320253.37867733/v1>.
- 1045
 1046 Luhar, A., Etheridge, D. M., Loh, Z.M., Noonan, J., Spencer, D., Smith, L., and Ong, C.: Quantifying methane
 1047 emissions from Queensland's coal seam gas producing Surat Basin using inventory data and a regional
 1048 Bayesian inversion, *Journal of Atmos. Chem. Phys.*, 20, 15487–15511, 2020
 1049 <https://acp.copernicus.org/articles/20/15487/2020/>, 2020.
- 1050 Lunt, M. F., Palmer, P. I., Feng, L., Taylor, C. M., Boesch, H., and Parker, R. J.: An increase in methane
 1051 emissions from tropical Africa between 2010 and 2016 inferred from satellite data, *Atmos. Chem. Phys.*,
 1052 19, 14721–14740, <https://doi.org/10.5194/acp-19-14721-2019>, 2019.
- 1053 Maasackers, J. D., Varon, D. J., Elfarsdóttir, A., McKeever, J., Jarvis, D., Mahapatra, G., Pandey, S., Lorente,
 1054 A., Borsdorff, T., Foorthuis, L. R., Schuit, B. J., Tol, P., van Kempen, T. A., van Hees, R., and Aben, I.:
 1055 Using satellites to uncover large methane emissions from landfills, *Sci. Adv.*, 8, eabn9683,
 1056 <https://doi.org/10.1126/sciadv.abn9683>, 2022.

- 1057 Maasackers, J. D., Jacob, D. J., Sulprizio, M. P., Turner, A. J., Weitz, M., Wirth, T., Hight, C., DeFigueiredo,
1058 M., Desai, M., and Schmeltz, R.: Gridded national inventory of US methane emissions, *Environ. Sci.*
1059 *Technol.*, 50, 13123–13133, 2016.
- 1060 McGrath, M. J., Petrescu, A. M. R., Peylin, P., Andrew, R. M., Matthews, B., Dentener, F., Balkovič, J.,
1061 Bastrikov, V., Becker, M., Broquet, G., Ciais, P., Fortems-Cheiney, A., Ganzenmüller, R., Grassi, G.,
1062 Harris, I., Jones, M., Knauer, J., Kuhnert, M., Monteil, G., Munassar, S., Palmer, P. I., Peters, G. P., Qiu,
1063 C., Schelhaas, M.-J., Tarasova, O., Vizzarri, M., Winkler, K., Balsamo, G., Berchet, A., Briggs, P.,
1064 Brockmann, P., Chevallier, F., Conchedda, G., Crippa, M., Dellaert, S. N. C., Denier van der Gon, H. A.
1065 C., Filipek, S., Friedlingstein, P., Fuchs, R., Gauss, M., Gerbig, C., Guizzardi, D., Günther, D., Houghton,
1066 R. A., Janssens-Maenhout, G., Lauerwald, R., Lerink, B., Luijkx, I. T., Moulas, G., Muntean, M., Nabuurs,
1067 G.-J., Paquirissamy, A., Perugini, L., Peters, W., Pilli, R., Pongratz, J., Regnier, P., Scholze, M., Serengil,
1068 Y., Smith, P., Solazzo, E., Thompson, R. L., Tubiello, F. N., Vesala, T., and Walther, S.: The consolidated
1069 European synthesis of CO₂ emissions and removals for the European Union and United Kingdom: 1990–
1070 2020, *Earth Syst. Sci. Data*, 15, 4295–4370, <https://doi.org/10.5194/essd-15-4295-2023>, 2023.
- 1071 McNorton, J., Bousserez, N., Agustí-Panareda, A., Balsamo, G., Cantarello, L., Engelen, R., Huijnen, V., Inness,
1072 A., Kipling, Z., Parrington, M., and Ribas, R.: Quantification of methane emissions from hotspots and
1073 during COVID-19 using a global atmospheric inversion, *Atmos. Chem. Phys.*, 22, 5961–5981,
1074 <https://doi.org/10.5194/acp-22-5961-2022>, 2022.
- 1075 National Academies of Sciences, Engineering, and Medicine, *Greenhouse Gas Emissions Information for*
1076 *Decision Making: A Framework Going Forward*. Washington, DC: The National Academies Press.
1077 <https://doi.org/10.17226/26641>, 2022.
- 1078 Nesser, H., Jacob, D. J., Maasackers, J. D., Lorente, A., Chen, Z., Lu, X., Shen, L., Qu, Z., Sulprizio, M. P.,
1079 Winter, M., Ma, S., Bloom, A. A., Worden, J. R., Stavins, R. N., and Randles, C. A.: High-resolution U.S.
1080 methane emissions inferred from an inversion of 2019 TROPOMI satellite data: contributions from
1081 individual states, urban areas, and landfills, *EGUsphere* [preprint], [https://doi.org/10.5194/egusphere-](https://doi.org/10.5194/egusphere-2023-946)
1082 [2023-946](https://doi.org/10.5194/egusphere-2023-946), 2023.
- 1083 Nisbet, E. G., Dlugokencky, E. J., Manning, M. R., Lowry, D., Fisher, R. E., France, J. L., Michel, S. E., Miller,
1084 J. B., White, J. W. C., Vaughn, B., Bousquet, P., Pyle, J. A., Warwick, N. J., Cain, M., Brownlow, R.,
1085 Zazzeri, G., Lanoisellé, M., Manning, A. C., Gloor, E., Worthy, D. E. J., Brunke, E.-G., Labuschagne, C.,
1086 Wolff, E. W., and Ganesan, A. L.: Rising Atmospheric Methane: 2007–2014 Growth and Isotopic Shift,
1087 *Global Biogeochem. Cycles*, 30, 1356–1370, <https://doi.org/10.1002/2016GB005406>, 2016.
- 1088 Nisbet, E. G., Manning, M. R., Dlugokencky, E. J., Fisher, R. E., Lowry, D., Michel, S. E., Myhre, C. L., Platt,
1089 S. M., Allen, G., Bousquet, P., Brownlow, R., Cain, M., France, J. L., Hermansen, O., Hossaini, R., Jones,
1090 A. E., Levin, I., Manning, A. C., Myhre, G., Pyle, J. A., Vaughn, B. H., Warwick, N. J., and White, J. W.
1091 C.: Very Strong Atmospheric Methane Growth in the 4 Years 2014–2017: Implications for the Paris

1092 Agreement, Global Biogeochem. Cycles, 33, 318–342,
1093 <https://doi.org/10.1029/2018GB006009>, 2019.

1094 Nisbet, E. G. (2023). Climate change feedback on methane emissions from global wetlands. *Nature Climate*
1095 *Change*, 13(5), 421–422. <https://doi.org/10.1038/s41558-023-01634-3>

1096 NOAA: Atmospheric data, https://www.esrl.noaa.gov/gmd/ccgg/trends_ch4/, last access: June 2023.

1097 Omara, M., Zavala-Araiza, D., Lyon, D.R., Hmiel, B., Roberts, K.A., and Hamburg, S.P.: Methane emissions
1098 from US low production oil and natural gas well sites. *Nature Communications* 13, 2085. 10.1038/s41467-
1099 022-29709-3, 2022.

1100 Patra, P. K., Saeki, T., Dlugokencky, E. J., Ishijima, K., Umezawa, T., Ito, A., Aoki, S., Morimoto, S., Kort, E.
1101 A., Crotwell, A., Ravikumar, K., and Nakazawa, T.: Regional methane emission estimation based on
1102 observed atmospheric concentrations (2002–2012), *J. Meteorol. Soc. Jpn.*, 94, 91–113, 2016.

1103
1104 Patra, P. K., Takigawa, M., Watanabe, S., Chandra, N., Ishijima, K., and Yamashita, Y.: Improved Chemical
1105 Tracer Simulation by MIROC4.0-based Atmospheric Chemistry-Transport Model (MIROC4-ACTM),
1106 *SOLA*, 14, 91–96, 2018.

1107 Patra, P.K., Krol, M.C., Prinn, R.G., Takigawa, M., Muhle, J., Montzka, S., Lal, S., Yamashita, Y., Naus, S.,
1108 Chandra, N., Weiss, R.F., Krummerl, P.B., Fraser, P.J., O’Doherty, S., Elkins, J.W.: Methyl Chloroform
1109 Continues to Constrain the Hydroxyl (OH) Variability in the Troposphere. *Journal of Geophysical*
1110 *Research: Atmospheres*, 126(4), e2020JD033862, doi: 10.1029/2020jd033862, 2021.

1111 Peng, S., Lin, X., Thompson, R.L., Xi, Y., Liu, G., Hauglustaine, D., Lan, X., Poulter, B., Ramonet, M., Sauniois,
1112 M., Yin, Y., Zhang, Z., Zheng, B., & Ciais, P.: Wetland emission and atmospheric sink changes explain
1113 methane growth in 2020, *Nature* 612, 477–482, <https://doi.org/10.1038/s41586-022-05447-w>, 2022.

1114 Petrescu, A. M. R., Peters, G. P., Janssens-Maenhout, G., Ciais, P., Tubiello, F. N., Grassi, G., Nabuurs, G.-J.,
1115 Leip, A., Carmona-Garcia, G., Winiwarter, W., Höglund-Isaksson, L., Günther, D., Solazzo, E., Kiesow,
1116 A., Bastos, A., Pongratz, J., Nabel, J. E. M. S., Conchedda, G., Pilli, R., Andrew, R. M., Schelhaas, M.-
1117 J., and Dolman, A. J.: European anthropogenic AFOLU greenhouse gas emissions: a review and
1118 benchmark data, *Earth Syst. Sci. Data*, 12, 961–1001, <https://doi.org/10.5194/essd-12-961-2020>, 2020.

1119 Petrescu, A. M. R., Qiu, C., Ciais, P., Thompson, R. L., Peylin, P., McGrath, M. J., Solazzo, E., Janssens-
1120 Maenhout, G., Tubiello, F. N., Bergamaschi, P., Brunner, D., Peters, G. P., Höglund-Isaksson, L., Regnier,
1121 P., Lauerwald, R., Bastviken, D., Tsuruta, A., Winiwarter, W., Patra, P. K., Kuhnert, M., Oreggioni, G.
1122 D., Crippa, M., Sauniois, M., Perugini, L., Markkanen, T., Aalto, T., Groot Zwaafink, C. D., Tian, H.,
1123 Yao, Y., Wilson, C., Conchedda, G., Günther, D., Leip, A., Smith, P., Haussaire, J.-M., Leppänen, A.,
1124 Manning, A. J., McNorton, J., Brockmann, P., and Dolman, A. J.: The consolidated European synthesis
1125 of CH₄ and N₂O emissions for the European Union and United Kingdom: 1990–2017, *Earth Syst. Sci.*
1126 *Data*, 13, 2307–2362, <https://doi.org/10.5194/essd-13-2307-2021>, 2021.

- 1127 Petrescu, A. M. R., Qiu, C., McGrath, M. J., Peylin, P., Peters, G. P., Ciais, P., Thompson, R. L., Tsuruta, A.,
 1128 Brunner, D., Kuhnert, M., Matthews, B., Palmer, P. I., Tarasova, O., Regnier, P., Lauerwald, R.,
 1129 Bastviken, D., Höglund-Isaksson, L., Winiwarter, W., Etiope, G., Aalto, T., Balsamo, G., Bastrikov, V.,
 1130 Berchet, A., Brockmann, P., Ciotoli, G., Conchedda, G., Crippa, M., Dentener, F., Groot Zwaaftink, C.
 1131 D., Guizzardi, D., Günther, D., Haussaire, J.-M., Houweling, S., Janssens-Maenhout, G., Kouyate, M.,
 1132 Leip, A., Leppänen, A., Lugato, E., Maisonnier, M., Manning, A. J., Markkanen, T., McNorton, J.,
 1133 Muntean, M., Oreggioni, G. D., Patra, P. K., Perugini, L., Pison, I., Raivonen, M. T., Saunois, M., Segers,
 1134 A. J., Smith, P., Solazzo, E., Tian, H., Tubiello, F. N., Vesala, T., van der Werf, G. R., Wilson, C., and
 1135 Zaehle, S.: The consolidated European synthesis of CH₄ and N₂O emissions for the European Union and
 1136 United Kingdom: 1990–2019, *Earth Syst. Sci. Data*, 15, 1197–1268, [https://doi.org/10.5194/essd-15-](https://doi.org/10.5194/essd-15-1197-2023)
 1137 1197-2023, 2023a.
- 1138 Petrescu, A.M.R., Peters, G.P., Engelen, R., Houweling, S., Brunner, D., Tsuruta, A., Matthews, B., Patra, P.K.,
 1139 Belikov, D., Thompson, R.L., Höglund-Isaksson, L., Zhang, W., Segers, A.J., Etiope, G., Ciotoli, G.,
 1140 Peylin, P., Chevallier, F., Aalto, T., Andrew, R.M., Bastviken, D., Berchet, A., Broquet, G., Conchedda,
 1141 G., Gütschow, J., Haussaire, J.-M., Lauerwald, R., Markkanen, T., van Peet, J.C.A., Pison, I., Regnier, P.,
 1142 Solum, E., Scholze, M., Tenkanen, M., Tubiello, F.N., van der Werf, G.R., Worden, J.R.: Reconciliation
 1143 of observation- and inventory- based CH₄ emissions for eight large global emitters, version 1, Zenodo
 1144 [data set], <https://doi.org/10.5281/zenodo.10276087>, 2023b.
- 1145 Prosperi, P., Bloise, M., Tubiello, F. N., Conchedda, G., Rossi, S., Boschetti, L., Salvatore, M., and Bernoux,
 1146 M.: New estimates of greenhouse gas emissions from biomass burning and peat fires using MODIS
 1147 Collection 6 burned areas, *Clim. Change*, 161, 415–432, <https://doi.org/10.1007/s10584-020-02654-0>,
 1148 2020.
- 1149 [Perugini, L., Pellis, G., Grassi, G., Ciais, P., Dolman, H., House, J. I., Peters, G. P., Smith, P.,](#)
 1150 [Günther, D., and Peylin, P.: Emerging reporting and verification needs under the](#)
 1151 [Paris Agreement: How can the research community effectively contribute?, *Environ.*](#)
 1152 [Sci. Policy, 122, 116–126, <https://doi.org/10.1016/j.envsci.2021.04.012>, 2021.](#)
- 1153 Qonitan, F.D., Wayan, I., Suryawan, K., and Rahman, A.: Overview of Municipal Solid Waste Generation and
 1154 Energy Utilization Potential in Major Cities of Indonesia, *J. Phys. Conf. Ser.* 1858,
 1155 <https://doi.org/10.1088/1742-6596/1858/1/012064>, 2021.
- 1156 Qu, Z., Jacob, D. J., Zhang, Y., Shen, L., Varon, D. J., Lu, X., et al. (2022). Attribution of the 2020 surge in
 1157 atmospheric methane by inverse analysis of GOSAT observations. *Environmental Research Letters*,
 1158 17(9), 094003. <https://doi.org/10.1088/1748-9326/ac8754>
- 1159 Regulation (EU) 2018/1999 (European Climate Law): Amended proposal for a regulation of the European
 1160 parliament and of the council on establishing the framework for achieving climate neutrality,

1161 https://ec.europa.eu/clima/sites/clima/files/eu-climate-action/docs/prop_reg_ecl_en.pdf, last access:
1162 October 2023.

1163 Regulation (EU) 525/2013 of the European Parliament and of the Council: [https://eur-lex.europa.eu/legal-](https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32013R0525&from=EN)
1164 [content/EN/TXT/PDF/?uri=CELEX:32013R0525&from=EN](https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32013R0525&from=EN), last access: November 2022.

1165 Rocher-Ros, G., Stanley, E.H., Loken, L.C., Casson, N.J., Raymond, P.A., Liu, S., Amatulli, G. & Sponseller,
1166 A.R.: Global methane emissions from rivers and streams, *Nature* 621, 530–535,
1167 <https://doi.org/10.1038/s41586-023-06344-6>, 2023.

1168 Rutherford, J.S., Sherwin, E.D., Ravikumar, A.P., Heath, G.A., Englander, J., Cooley, D., Lyon, D., Omara, M.,
1169 Langfitt, Q., and Brandt, A.R.: Closing the methane gap in US oil and natural gas production emissions
1170 inventories. *Nature Communications* 12, 4715. [10.1038/s41467-021-25017-4](https://doi.org/10.1038/s41467-021-25017-4), 2021.

1171

1172 Assè-Wassa Sama, M. and Berenger, V.: In Africa, Greenhouse gas emissions from the waste sector increase,
1173 despite efforts from local actors, Observatory on Climate Action in Africa report, [https://www.climate-](https://www.climate-chance.org/wp-content/uploads/2023/06/obsaf_emag6_note-economie-circulaire-et-dechets_eng-2.pdf)
1174 [chance.org/wp-content/uploads/2023/06/obsaf_emag6_note-economie-circulaire-et-dechets_eng-2.pdf](https://www.climate-chance.org/wp-content/uploads/2023/06/obsaf_emag6_note-economie-circulaire-et-dechets_eng-2.pdf),
1175 last access November 2023, 2023.

1176 Saunio, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A.,
1177 Dlugokencky, E. J., Houweling, S., Patra, P. K., Ciais, P., Arora, V. K., Bastviken, D., Bergamaschi, P.,
1178 Blake, D. R., Brailsford, G., Bruhwiler, L., Carlson, K. M., Carrol, M., Castaldi, S., Chandra, N.,
1179 Crevoisier, C., Crill, P. M., Covey, K., Curry, C. L., Etiope, G., Frankenberg, C., Gedney, N., Hegglin,
1180 M. I., Höglund-Isaksson, L., Hugelius, G., Ishizawa, M., Ito, A., Janssens-Maenhout, G., Jensen, K. M.,
1181 Joos, F., Kleinen, T., Krummel, P. B., Langenfelds, R. L., Laruelle, G. G., Liu, L., Machida, T.,
1182 Maksyutov, S., McDonald, K. C., McNorton, J., Miller, P. A., Melton, J. R., Morino, I., Müller, J.,
1183 Murguia-Flores, F., Naik, V., Niwa, Y., Noce, S., O'Doherty, S., Parker, R. J., Peng, C., Peng, S., Peters,
1184 G. P., Prigent, C., Prinn, R., Ramonet, M., Regnier, P., Riley, W. J., Rosentreter, J. A., Segers, A.,
1185 Simpson, I. J., Shi, H., Smith, S. J., Steele, L. P., Thornton, B. F., Tian, H., Tohjima, Y., Tubiello, F. N.,
1186 Tsuruta, A., Viovy, N., Voulgarakis, A., Weber, T. S., van Weele, M., van der Werf, G. R., Weiss, R. F.,
1187 Worthy, D., Wunch, D., Yin, Y., Yoshida, Y., Zhang, W., Zhang, Z., Zhao, Y., Zheng, B., Zhu, Q., Zhu,
1188 Q., and Zhuang, Q.: The Global Methane Budget 2000–2017, *Earth Syst. Sci. Data*, 12, 1561–1623,
1189 <https://doi.org/10.5194/essd-12-1561-2020>, 2020.

1190 Stavert, A. R., Saunio, M., Canadell, J. G., Poulter, B., Jackson, R. B., Regnier, P., Lauerwald, R., Raymond,
1191 P. A., Allen, G. H., Patra, P. K., Bergamaschi, P., Bousquet, P., Chandra, N., Ciais, P., Gustafson, A.,
1192 Ishizawa, M., Ito, A., Kleinen, T., Maksyutov, S., Joe McNorton, J., Melton, J.R., Müller, J., Niwa, J.,
1193 Peng, S., Riley, W.J., Segers, A., Tian, H., Tsuruta, A., Yin, Y., Zhang, Z., Zheng, B., and Zhuang, Q.:
1194 Regional trends and drivers of the global methane budget, *Global Change Biol.*, 28, 182–200,
1195 <https://doi.org/10.1111/gcb.15901>, 2022.

- 1196 The Global methane Pledge, <https://www.state.gov/global-methane-pledge-from-moment-to-momentum/>, last
1197 access September 2023.
- 1198 Thompson, R. L. and Stohl, A.: FLEXINVERT: an atmospheric Bayesian inversion framework for determining
1199 surface fluxes of trace species using an optimized grid, *Geosci. Model Dev.*, 7, 2223–2242,
1200 <https://doi.org/10.5194/gmd-7-2223-2014>, 2014.
- 1201 Tubiello, F. N., Karl, K., Flammini, A., Gütschow, J., Obli-Laryea, G., Conchedda, G., Pan, X., Qi, S. Y.,
1202 Halldórudóttir Heiðarsdóttir, H., Wanner, N., Quadrelli, R., Rocha Souza, L., Benoit, P., Hayek, M.,
1203 Sandalow, D., Mencos Contreras, E., Rosenzweig, C., Rosero Moncayo, J., Conforti, P., and Torero, M.:
1204 Pre- and post-production processes increasingly dominate greenhouse gas emissions from agri-food
1205 systems, *Earth Syst. Sci. Data*, 14, 1795–1809, <https://doi.org/10.5194/essd-14-1795-2022>, 2022.
- 1206 UNFCCC 1st BUR DR Congo: <https://unfccc.int/documents/624762>, last access may 2023
- 1207 UNFCCC 1st BUR India: <https://unfccc.int/documents/180646>, 2016, last access May 2023
- 1208 UNFCCC Common Reporting Formats (CRFs) of the National Inventory Reports Unpublished to the UNFCCC
1209 by various Annex-1 countries (Bonn: United Nations Framework Convention on Climate Change) (
1210 [https://unfccc.int/process-and-meetings/transparency-and-reporting/reporting-and-review-under-the-](https://unfccc.int/process-and-meetings/transparency-and-reporting/reporting-and-review-under-the-convention/greenhouse-gas-inventories-annex-i-parties/national-inventory-submissions-2019)
1211 [convention/greenhouse-gas-inventories-annex-i-parties/national-inventory-submissions-2019](https://unfccc.int/process-and-meetings/transparency-and-reporting/reporting-and-review-under-the-convention/greenhouse-gas-inventories-annex-i-parties/national-inventory-submissions-2019)), 2016
- 1212 UNFCCC Common Reporting Formats (CRFs) of the National Inventory Reports Unpublished to the UNFCCC
1213 by various Annex-1 countries (Bonn: United Nations Framework Convention on Climate Change) (
1214 [https://unfccc.int/process-and-meetings/transparency-and-reporting/reporting-and-review-under-the-](https://unfccc.int/process-and-meetings/transparency-and-reporting/reporting-and-review-under-the-convention/greenhouse-gas-inventories-annex-i-parties/national-inventory-submissions-2019)
1215 [convention/greenhouse-gas-inventories-annex-i-parties/national-inventory-submissions-2019](https://unfccc.int/process-and-meetings/transparency-and-reporting/reporting-and-review-under-the-convention/greenhouse-gas-inventories-annex-i-parties/national-inventory-submissions-2019)), 2018.
- 1216 UNFCCC Australia NIR: <https://unfccc.int/documents/627765>, last access November 2023.
- 1217 UNFCCC 2nd BUR China: [https://unfccc.int/documents/197666?gclid=CjwKCAiA3aeqBhBzEiwAxFiOBpb-](https://unfccc.int/documents/197666?gclid=CjwKCAiA3aeqBhBzEiwAxFiOBpb-OsdbvprVkIXVi5QeaNHBj22J8z7q6yznj0y8T3qqS1SzL4nWtBoCZ4oQAvD_BwE)
1218 [OsdbvprVkIXVi5QeaNHBj22J8z7q6yznj0y8T3qqS1SzL4nWtBoCZ4oQAvD_BwE](https://unfccc.int/documents/197666?gclid=CjwKCAiA3aeqBhBzEiwAxFiOBpb-OsdbvprVkIXVi5QeaNHBj22J8z7q6yznj0y8T3qqS1SzL4nWtBoCZ4oQAvD_BwE), 2019, last access
1219 April 2023
- 1220 UNFCCC 2nd BUR India: <https://unfccc.int/documents/192316>, 2018, last access May 2023.
- 1221 UNFCCC 3rd BUR India:
1222 [https://unfccc.int/documents/268470?gclid=CjwKCAiA3aeqBhBzEiwAxFiOBjSI1whAzuMQOqDCtDg](https://unfccc.int/documents/268470?gclid=CjwKCAiA3aeqBhBzEiwAxFiOBjSI1whAzuMQOqDCtDgXewzSFSQiimB4kj9fzQOc0NKODxLwrESB8hoCOHQQA_VD_BwE)
1223 [XewzSFSQiimB4kj9fzQOc0NKODxLwrESB8hoCOHQQA_VD_BwE](https://unfccc.int/documents/268470?gclid=CjwKCAiA3aeqBhBzEiwAxFiOBjSI1whAzuMQOqDCtDgXewzSFSQiimB4kj9fzQOc0NKODxLwrESB8hoCOHQQA_VD_BwE), 2021, last access May 2023
- 1224 UNFCCC 3rd BUR Indonesia: [https://unfccc.int/documents/403577?gclid=CjwKCAiA3aeqBhBzEiwAxFiOBt-](https://unfccc.int/documents/403577?gclid=CjwKCAiA3aeqBhBzEiwAxFiOBt-IUluMufU7_wGQWfi968cP_4c59-5_9Sq9MR4pf80FbjXAYU1jBoC-gAQAvD_BwE)
1225 [I_UluMufU7_wGQWfi968cP_4c59-5_9Sq9MR4pf80FbjXAYU1jBoC-gAQAvD_BwE](https://unfccc.int/documents/403577?gclid=CjwKCAiA3aeqBhBzEiwAxFiOBt-IUluMufU7_wGQWfi968cP_4c59-5_9Sq9MR4pf80FbjXAYU1jBoC-gAQAvD_BwE) , 2021 last
1226 access May 2023
- 1227 UNFCCC 4th BUR Brazil: <https://unfccc.int/documents/267661>, 2020 last access May 2023

- 1228 UNFCCC EU NIR: <https://unfccc.int/documents/627851>, last access: April 2023.
- 1229 UNFCCC Germany NIR: <https://unfccc.int/documents/627785>, last access: April 2023.
- 1230 UNFCCC Switzerland NIR, Annex 6: <https://unfccc.int/documents/627731>, last access: April 2023.
- 1231 UNFCCC UK NIR, Annex 6: <https://unfccc.int/documents/627789>, last access: April 2023.
- 1232 VERIFY H2020 Project: <https://verify.lsce.ipsl.fr/>, last access May, 2023
- 1233 Wolf, J., Asrar, G.R. & West, T.O. Revised methane emissions factors and spatially distributed annual carbon
1234 fluxes for global livestock. Carbon Balance Manage 12, 16, <https://doi.org/10.1186/s13021-017-0084-y>,
1235 2017.
- 1236 Zhang, X., Waugh, D.W. and Orbe, C.:Dependence of Northern Hemisphere tropospheric transport on the
1237 midlatitude jet under abrupt CO₂ increase. J. Geophys. Res. Atmos., 128, no. 13, e2022JD038454,
1238 doi:10.1029/2022JD038454, 2023.
- 1239 [Zhang, Z., Poulter, B., Melton, J. R., Riley, W. J., Allen, G. H., Beerling, D. J., Bousquet, P.,](https://doi.org/10.5194/egusphere-2024-1584)
1240 [Canadell, J. G., Fluet-Chouinard, E., Ciais, P., Gedney, N., Hopcroft, P. O., Ito, A.,](https://doi.org/10.5194/egusphere-2024-1584)
1241 [Jackson, R. B., Jain, A. K., Jensen, K., Joos, F., Kleinen, T., Knox, S., Li, T., Li, X., Liu, X.,](https://doi.org/10.5194/egusphere-2024-1584)
1242 [McDonald, K., McNicol, G., Miller, P. A., Müller, J., Patra, P. K., Peng, C., Peng, S., Qin,](https://doi.org/10.5194/egusphere-2024-1584)
1243 [Z., Riggs, R. M., Saunois, M., Sun, Q., Tian, H., Xu, X., Yao, Y., Yi, X., Zhang, W., Zhu,](https://doi.org/10.5194/egusphere-2024-1584)
1244 [Q., Zhu, Q., and Zhuang, Q.: Ensemble estimates of global wetland methane](https://doi.org/10.5194/egusphere-2024-1584)
1245 [emissions over 2000–2020, EGUsphere \[preprint\],](https://doi.org/10.5194/egusphere-2024-1584)
1246 [https://doi.org/10.5194/egusphere-2024-1584, 2024.](https://doi.org/10.5194/egusphere-2024-1584)