

Decadal Growth in Emission Load of Major Air Pollutants in Delhi

Saroj Kumar Sahu¹, Poonam Mangaraj^{1,2}, Gufran Beig²

¹Environmental Science, Department of Botany, Utkal University, Bhubaneswar, India

²National Institute of Advanced Studies, Indian Institute of Science, Campus, Bangalore, India

Correspondence to: Saroj Kumar Sahu (saroj.bot@utkaluniversity.ac.in); Poonam Mangaraj (poonammangaraj92@gmail.com); Gufran Beig (beig@nias.res.in)

Abstract

Indian capital megacity Delhi is reeling under deteriorating air quality and control measures are not yielding any significant changes mainly due to a poor understanding of sources of emissions, hence priority option in mitigation planning is lacking. In this paper, we have made an attempt to develop a spatially resolved technological high-resolution gridded (~0.4km × 0.4km) emission inventory for eight major pollutants of the Delhi region where high-resolution activity data of all possible major and unattended minor sources are generated by organizing a mega campaign involving 100s of volunteers. It is for the first time that we are able to estimate the decadal growth in emissions of various pollutants by comparing newly developed 2020 emissions with SAFAR emissions of 2010 using the identical methodology and quantum of activity data. The estimated annual emission for PM_{2.5}, PM₁₀, CO, NO_x, VOC, SO₂, BC and OC over Delhi-NCR are estimated to be 123.8 Gg/yr, 243.6 Gg/yr, 799.0 Gg/yr, 488.9 Gg/yr, 730.0 Gg/yr, 425.8 Gg/yr, 33.6 Gg/yr, and 20.3 Gg/yr respectively for the year 2020. The decadal growth (2010-2020) in PM_{2.5} and PM₁₀ are found to be marginal 31% and 3% respectively. The maximum growth is found to be in the transport sector followed by the industrial and other sectors. Maximum decadal growth found for pollutants BC, OC and NO_x is 57%, 34% and 91% respectively. The decadal shift of sectorial emissions with changing policies is examined. The complete dataset is available on Zenodo at <https://doi.org/10.5281/zenodo.7715595> (Sahu et al., 2023).

Keywords: Megacity, Emission Inventory, Hotspots, Air quality, Anthropogenic Emission, Major/Minor Sources, Mitigation Strategies

32 **1. Introduction:**

33 Clean air is a basic need for a healthy life but air pollution has emerged as a global emergency
34 where cities are more vulnerable due to high population density. Asian mega-cities are even
35 more polluted than before and have drawn all global attention (IPCC, 2000; Molina & Molina,
36 2004; Permadi et al., 2018). Air quality in Indian megacity Delhi makes headlines across the
37 print and media with the onset of the winter months (Beig et al., 2021; India Today, 2022).
38 Worldwide, air pollution is a widespread problem and a major contemporary public health threat.
39 Air pollutants are treated like a modern-day curse due to their association with premature
40 mortality and disease burden has a significant impact on low-income developing countries,
41 especially India. Air pollution emerged as the fourth leading risk factor contributing to disease
42 burden and early death worldwide (HEI 2019, 2020). The Global Burden of Disease (GBD)
43 reported that ~4.9 million premature deaths across the globe occur because of air pollution
44 (Stanaway et al., 2018; Manisalidis et al., 2020). People from any geographical region could
45 suffer from its adverse impacts irrespective of the place of origin (Akimoto 2003). Certainly,
46 Indian urban have emerged as one of the most adversely affected polluting places as well as
47 global health risks (Down to Earth, 2015; GBD, 2018). 22 cities of the world's 30 most polluted
48 cities are in India from which Delhi, the capital of India tops the ranking for consecutive years
49 with its annual particulate matter (PM_{2.5}) level nearly ten times the WHO permissible limits and
50 is intricately caught in the toxic web of air quality and health-based standards (UNEP, 2019;
51 World Air Quality Report, 2019, 2020). This led to alarming levels of Air Quality Index (AQI)
52 in National Capital mega-city Delhi that has dragged first-ever such a large-scale media and
53 political attention in recent years. No doubt the mega-cities have emerged as a better place to live
54 but at the same time, are prone to degrading air quality due to elevated concentration of
55 particulate matter (PM) (Molina et al., 2004; Beig et al, 2020, Sahu et al, 2011, 2021).
56 Combating mega-city air pollution has become utmost challenging due to a poor understanding
57 of the complexity of air pollution sources and its dynamic mixture of both natural and
58 anthropogenic.

59 Numerous studies have constantly manifested higher rates of respiratory and
60 cardiovascular diseases in megacities due to alarming pollution levels where the school-going
61 students and old generation are the largely affected (Sahu et al, 2011, Mangaraj et al, 2022).

62 Delhi's worse air quality during winter months is also linked with stubble burning in Punjab and
63 Haryana (Beig et al, 2019, 2020). The government introduced Odd & Even vehicle ply on roads
64 to reduce the impact of emission load (Transport Department, Govt. of Delhi, 2019). However,
65 the impact was not significant. The blame game keeps on running from one agency (or) state to
66 another where each one has its independent opinion to combat the rising level of pollutants in
67 Delhi. Despite many initiatives from stakeholders, Delhi air has shown no sign of improvement
68 and has drawn the attention of global researchers. Understanding the complexity of pollution
69 sources and their magnitude in a megacity is essential for air quality study as well as regional
70 atmospheric chemistry and climate point of view (Li et al, 2017). However, identification of the
71 unattended sources and their precise quantification becomes challenging due to the diversity of
72 contributing major/minor sources along with the complicity of technology being used during
73 combustion activities. The problem becomes even more complex due to the heterogeneity of
74 pollution sources and their temporal variation. A comprehensive high-resolution emission
75 inventory (EI) may solve the purpose because EIs are critical research and regulatory tools to
76 address the air pollution issues in many cities. Moreover, the surface emission is the most
77 sensitive input data chemical transport model to understand the impact of emission on
78 atmospheric chemistry on different scales urban to regional, national to global scale (Sahu et al.,
79 2011; Mangaraj et al., 2022).


80 There are few limited comprehensive detail studies, that focus on Delhi emission
81 estimation but each study has some or the other limitations. So far, many attempts from various
82 attempts like that of NEERI, 2010; Guttikunda and Calori, 2013 and TERI & ARAI, 2018 have
83 failed to get a concrete alternative to get rid of this air quality issue/problem. In order to frame
84 appropriate mitigation strategies to curve air pollution load in megacity Delhi, we have identified
85 the new emerging sources and have estimated the pollutant load from all possible major/minor
86 sectors responsible for the emission of various pollutants directly or indirectly. Unlike the
87 previous studies, the present study is unique of its kind by targeting 17 organized as well as
88 unorganized sectors responsible directly or indirectly for changing air quality in Delhi-NCR
89 regions. The present findings provide a comprehensive assessment of sources of air pollutants
90 and their magnitude, which has shifted with changing policies in the last one decade. One of the
91 main objective behind developing this reliable high-resolution (~0.4km × 0.4km) gridded
92 emission inventory of eight major pollutants over a domain of 70km×65km covering Delhi and

93 its adjacent NCR region for the base year 2020 (i.e. April 2019 to March 2020) is not only to
94 frame desired mitigation strategy to combat air pollutant issue but also to understand the decadal
95 growth of emission over same region under the flagship of SAFAR program of MoES. It will be
96 also an integral input to air quality forecasting based modeling study to understand the regional
97 atmospheric chemistry.

98

99 **1.1. Source of Emission, Activity Data & Emission Factors:**

100 Megacity Delhi (Figure 1), the capital of India, which is designated as the National Capital
101 Territory region (NCT), is located towards the Northern part of the country straddling the
102 Yamuna River. This megacity is stretched over an area of 1484 km² and shares border with Uttar
103 Pradesh in East and Haryana to the rest directions. It is situated at an elevation of ~216 m above
104 the sea level at 28.7041° N, 77.1025° E. The NCT of Delhi is divided into nine districts. The
105 estimated population of megacity Delhi is 28.5 million making it the largest metropolitan in
106 India. The overgrown population density of Delhi, has led to the expansion of city and increase
107 in use of energy and fossil fuels associated with alarming levels of air pollution and health risks.



108

109

Fig 1: Domain of interest

110 In order to suffice the objective of developing an authentic emission inventory, the
111 collection of primary activity data is of great significance. In the present attempt, for the first
112 time 17 minor/major sectors responsible for direct/indirect emission of pollutants have been
113 taken into account in the emission estimation process. To understand the emission practices, the
114 primary activity data were generated through a three-month-long extensive emission campaign
115 (SAFAR-Delhi, 2018) carried out over Delhi and surrounding National Capital Regions (NCR)
116 in 2018 (Figure 2). This initiative was undertaken by the Indian Institute of Tropical
117 Meteorology (IITM, Pune) in collaboration with the School of Planning and Architecture (SPA-
118 Delhi) and Utkal University under the Ministry of Earth Sciences (MoES)'s project "System of
119 Air Quality and Weather Forecasting And Research (SAFAR)". In order to serve the purpose of
120 understanding the complex source of pollutants, primary activity data is of great role in building
121 a high-resolution gridded emission inventory, which has to be generated through a
122 comprehensive field campaign only. This is a unique attempt to collect micro-level primary
123 activity database like the type of fuel used, the quantity of fuel being in various technology in
124 various sectors like a slum, residential cooking, brick industry, construction sites, street vendors,
125 large hotels, vehicle load around tourist places/railway stations/shopping malls/large
126 hospitals/large school/colleges and traffic junctions, airport, biomass/crop residue burning,
127 crematorium, use of cow dung as an alternative fuel for cooking, road dust, construction, open
128 waste burning, diesel generators in commercial purpose and mobile towers. Apart from
129 traditionally dominating sectors like transport, wind-blown road dust, industry, thermal power
130 plants, and residential, there are several unattended minor sectors, which collectively have a
131 relatively significant contribution to air pollution issue in Delhi. Apart from this, the most
132 important objective is to check the authenticity and accuracy of the existing secondary data
133 collected from various government agencies and reports as well as to fill the data gap. For the
134 same, meticulously ~150 students from various universities and colleges put an extensive
135 painstaking approx. 40,000 hrs effort to compile a comprehensive and robust activity database
136 under the supervision of a group of scientists/experts. This will not only help to understand all
137 possible major/minor sources better but also the prevailing changing trend in megacity Delhi and
138 its surrounding regions. The generated data will play an instrumental role in understanding the
139 changing trend of the source of pollutants in the last decade.



140
141 **Figure 2: Snapshot of Delhi Survey**


142 We have elaborated all the 17 sources into 5 major sectors further which are:- (a) Transport, (b)
143 Windblown Road Dust, (c) Industry, (d) Residential (includes sub sectors: household, slum,
144 street vendor, crop residue burning, cow dung, and diesel generators), (e) Others (includes sub
145 sectors: municipal solid waste burning, construction, incense sticks/mosquito coils/cigarettes,
146 and crematory).

147 **(a) Transport:**

148 Delhi has been witnessing a consistent increase in number of motor vehicles in recent years. It is
149 a home to approximately 13.3 million registered vehicles as of 2020 March (MoRTH, 2020) that
150 has grown to two and four folds in last one and two decade respectively (Sahu et al, 2011,
151 SAFAR-Delhi, 2010). In transport sector, vehicles have been classified broadly into eight
152 categories Two-wheelers (2W), three-wheelers (3W), Buses, Personal cars, Commercial Cars,
153 Light commercial vehicles (LCV), Heavy commercial vehicles (HCV) and Miscellaneous
154 (MSLV). Overall, the relative contribution of each category showed a higher contribution of 2W
155 with ~56%, personal cars with ~23%, followed by commercial cars with ~17%, 3W and buses
156 with ~3% and the remaining 2% by rest vehicle categories. The Supreme Court of India in 1998
157 sanctioned a rule for all the transport system of Delhi to be run by compressed natural gas (CNG)

158 in order to deal with the increase in vehicular emission. Delhi as of now has ~1 million vehicles
159 running on CNG that constitutes ~26% of CNG-3W, 67% CNG-cars and 7% CNG-buses. The
160 government has been concerned for the air pollution crisis in Delhi since long and therefore BS-
161 IV emission norms were implemented in Delhi in 2010 before any was implemented for rest of
162 the nation in 2017. BS-IV has been implemented in Delhi since 2018 but has been proposed to
163 implement in other cities by the month April 2020. The National Automobile Scrapping policy
164 was introduced in India lately in 13th August 2021 to reduce India's vehicular air pollution with
165 effect from 25th September 2021 (MoRTH, 2021). The transport department of Delhi has lately
166 passed an order for diesel vehicles more than 10 years old would be deregistered automatically
167 from January 2022. At the same time, the calculations tally that a fraction of the fleet registered
168 during 2000-2010 might still be active on the roads of Delhi in 2020 despite the phasing out
169 process. The present area of interest has road network of ~2450 km of major roads and ~31000
170 km of minor roads. The manual vehicle counts were computed over 87 survey locations (Figure
171 3 a) in Delhi and its surrounding NCR region to identify the density of vehicle (Figure 3 b) and
172 its composition according to vehicle age was also estimated(Figure 3 c).

173 The enumerating task was carried out for both in weekdays and weekends with the help
174 of digital click counters. The counting task was carried out for continuously for around 14 - 16
175 hrs. per day. Vehicle density was recorded to be as high as around 110000 - 160000 during
176 weekdays in many major roads as shown in Figure 3 d. However, it was observed that vehicle
177 number increased during weekend over the couple of roads like India Gate Circle, Chandni
178 Chowk, and Lajpat Nagar etc. High vehicular density of more than 100000 per day were
179 observed on roads like Delhi Meerut Expressway, Dhaula Kuan, Peeragarhi, Ashram Road,
180 South Extension Airport Road etc. Delhi is surrounded by other populous states like Uttar
181 Pradesh, Haryana, and Punjab, which are directly/indirectly linked with various activities over
182 Delhi-NCR region. Therefore, the other state cars contribute as high as nearly 40% in majority of
183 well-known busy roads in Delhi. An approx. of 2600 samples was collected for the random
184 survey with several real time diverse data like fuel consumption pattern, hours of usage, vehicle
185 density, Vehicle Kilometers Travelled (VKT) per day, type of fuel used, etc. The real time VKT
186 generated during random survey is depicted in Table 1.



187
 188 **Figure 3: (a) Survey locations for primary activity data for transport sector in Delhi-NCR;**
 189 **(b) Category-wise vehicle density in various roads across Delhi-NCR; (c) Age-wise vehicle**
 190 **category; (d) Comparison of vehicle density on weekdays and weekends on major roads of**
 191 **Delhi**

192

Vehicle Category	Fuel	VKT (km/day)
Two Wheeler (2W)	Gasoline	75
Three Wheeler 2S/4S (3W)	CNG/Gasoline	120
Bus	Diesel	210
Personal Car (P Car)	Gasoline	60
Commercial Car (C Car)	CNG/Diesel	200
Heavy Commercial Vehicle (HCV)	Diesel	75
Light Commercial Vehicle (LCV)	Diesel	150
Miscellaneous (MSLV)	Diesel	50

193 **Table 1: Vehicle Category specific VKT collected during field survey**

194

195 **(b) Windblown:**

196 Delhi has a huge and dense road network but all the roads are not certainly maintained. Road
197 condition of Delhi was observed keenly. The roads joining towards the outskirts of the city were
198 found to be worn out and lead to huge dust load. At the same time, random survey in different
199 roads was undertaken in order to assess the driving cycle/pattern of different vehicle categories.
200 The other state car contributes as high as nearly 40-50% in majority of well-known busy roads in
201 Delhi. Due to rise in the number of vehicles, the average speed of vehicles is found to be
202 decreasing in trend (i.e.18-25 km/hr in most of the major roads and 35-55 km/hr on airport roads
203 and few more important roads). The average weight-age of vehicles in Delhi was determined
204 based on vehicle category and its composition, which was estimated to be 1.23 tons. The number
205 of precipitation days in Delhi is hardly 50 days with an annual rainfall of just 547 mm (Rainfall
206 Statistics of India, 2019); therefore, the soil moisture content over study area was determined to
207 be considered just6%. The paved and unpaved road ratio was closely monitored and ~40% of
208 roads were found to be unpaved with broken road shoulders, poor infrastructure and the rest
209 ~60% were considered to be paved. The silt load on these roads was estimated to be 10% for
210 paved roads and 12% unpaved roads which comparatively better than any other cities in India.
211 The resuspension of dust load increases with increasing weight of vehicle and speed. The
212 average vehicle weight and mean vehicle speed derived from fleet composition running on the
213 road were determined from field survey across many roads along with the number of
214 precipitation days and moisture content to arrive at total dust load over study area.

215 **(c) Industry:**

216 In case of industries, it is one of the most diverse sectors with more than 3182 industries
217 scattered over Delhi-NCR where the large fraction is much unorganized with no/limited fuel
218 activity data followed by small fraction of organized sector. The Central Pollution Control Board
219 (CPCB) and Delhi Pollution Control Committee (DPCC) have classified the polluting industries
220 of Delhi into three categories as Red (highly polluting), Orange (moderately polluting) and
221 Green (non-polluting). The red category industries are strictly banned within the Delhi city
222 however; orange category industries are allowed for operation. There is no comprehensive
223 database for all industries with their technological details. As per the primary survey, large
224 numbers of unorganized small industries were found to be confined over Eastern, Southeastern,

225 and Southwestern part of Delhi region. Central Delhi has relatively very low number of
226 industries in comparison to others part of city. The spatial distribution of diverse range of small,
227 medium and large industries is depicted in Figure 4. Major industries include – Engineering
228 industries, which carry a frequency of 546, Machine and tools industries of 169, Electricals 175,
229 Iron and Steel industries 114, etc. Most of the detailed information on industrial areas, fuel
230 consumption pattern, production capacity etc. has been collected from DPCC. Fuels used in these
231 industries include Low Sulphur heavy Stock (LSHS), Light Diesel Oil (LDO), High Speed
232 Diesel (HSD), Liquid Petroleum Gas (LPG), Natural Gas (NG) and coal.

233 **(d) Residential:**

234 Delhi's estimated population was ~22.7 million which within a decade increased to a total
235 population of ~30.2 million (2020) and is known as the first most populous city of India and
236 second largest populated city of the world. According to the
237 Ministry of Housing & Urban Poverty Alleviation, Govt. of India, around 13-14% of Delhi's
238 population lives in slums. The Delhi Urban Shelter Improvement Board, 2019 reported ~675
239 clusters of slum in Delhi. The actual slum population data is very uncertain. During the field
240 survey, nearly 187 locations were covered to collect over 3000 samples over slum clusters
241 confined over the Central, Eastern and South Eastern part of Delhi. The total population is
242 estimated to be distributed among 4 million households with an average household size of five.
243 The cooking fuel activity data collected confirm the changing trend in cooking fuel used in Delhi
244 slum in last one decade. Unlike traditional like wood, dung, bio-fuel, LPG is being widely used
245 as main fuel which accounts around 95%, followed by wood 3% and coal 2%. In winters, the
246 relative contribution of wood as fuel increases (for heating of water). This indicates that there is
247 excellent penetration of government awareness and promotion of LPG connection in slum
248 pockets. Apart from this, it came into notice that people residing in the outskirts of Delhi are
249 using cow dung as fuel for heating and cooking purpose, especially during winters. The mixture
250 of generated agricultural residue with cow-dung and raw materials like biomass and coal dust are
251 still being used for domestic cooking in the peripherals of Delhi and its adjoining districts. The
252 mixture is dried and molded into circular shapes with a curvature staked to the walls and left for
253 sunbathing called as 'Uplah' in local language. Later, they are piled up into mounds to be
254 preserved for months and are used as an alternative for domestic fuels. As per the survey, a

255 single household size of 5-6 members use approx. 30kg of cow-dung per month as a source of
256 fuel for cooking and heating of water in winter.

257 With changing lifestyle with eating habits, vending in megacity Delhi holds up ~5,00,000
258 street vendors which are well scattered all across the city. Nearly 1653 samples on cooking fuel
259 activities were collected by interacting with people working in various hotels, restaurants and
260 street vendors to know the exact situation prevailing in Delhi-NCR regions where the coal and
261 wood are combusted using traditional approach as well as traditional stove. A large proportion of
262 these street vendors were certified under the regional Municipal Corporations and were situated
263 at permanent vending zones and many were found to be unauthorized ones who kept shifting
264 from one place to another. During the field survey, it was observed that LPG is being
265 predominantly used as a source of fuel by the street vendors (i.e. 83%) followed by coal (15%)
266 and wood (2%). Few street vending zones were found to be predominantly using coal for
267 'tandoor' food making activities especially near tourist places like India Gate, Jam Masjid, Lal
268 Qila etc. Kerosene is found not to be in use primarily as a source of fuel for cooking activities.
269 However, crop residue burning is not prominent in core urban region of Delhi but the peri-urban
270 areas towards the northeastern fringes hold less cultivated cropland. Hence, crop residue burning
271 in the urban region is of little significance. The activity data with respect to cultivated area and
272 amount are accounted from government portals like ICAR (Indian Council of Agricultural
273 Research), MoSPI (Ministry of Statistics and Programme Implementation), Ministry of
274 Agriculture & Farmers' Welfare and paid sites Indiastat. The approach used for estimating the
275 total crop residue generated and the fraction burnt is adopted from Sahu et al., 2021.

276 Due to load shedding, diesel generator (DG) sets as a source for power backup are
277 increasingly frequent in commercial establishments and apartments. In most parts of Delhi 1-2
278 hrs. of power failure is quiet common in summer then. Besides that, DG sets are also used in
279 base transceiver stations (BTS). According to the Department of Telecommunication (DoT),
280 2019, Delhi has more than 26,000 telecom towers, which have ~1 lakh BTSs that run with DG
281 sets for a constant or substitute source of power. A common BTS is equipped with a 12-25kWh
282 DG set, which on an average consumes ~9000-12000 liters of diesel annually (Sahu et al., 2015).
283 For estimating the number of diesel generator sets in commercial premises, ratio of gensets and
284 population was taken and total number of commercial establishments in Delhi with their spatial

285 locations was assembled from paid sources. Total emission from DG sector was based on the
286 number of diesel generator sets and power failure hours.

287 **(e) Other:**

288 The Indira Gandhi International Airport is the primary international airport spread over an area
289 of 2066 ha situated at 9.9 miles from city centre of New Delhi. It is the busiest airport and sixth
290 busiest airport in Asia in terms of passenger traffic. According to the bulletin of Indira Gandhi
291 International Airport, the calendar year of 2019-20, it witnessed ~67 million passenger traffic
292 and 450,012 aircraft movements. The Landing/Take-off (LTO) cycle, which happens below the
293 altitude of ~ 1000 m (3000 feet) basically, contributes to the air pollution. The activity data of
294 aircraft movement and passenger traffic are collected from government reports of the Directorate
295 General of Civil Aviation (DGCA), 2020 and the Ministry of Civil Aviation, 2020.

296 According to the Delhi Pollution Control Committee, Delhi generates ~11,144 tons of
297 Municipal Solid Waste (MSW) per day on average, which are dumped over three uncontrolled
298 and unlined landfill sites of Delhi i.e. Ghazipur, Bhalaswa, and Okhla dump yard (DPCC, 2020).
299 Ghazipur landfill is the largest dump yard located towards the Eastern perimeters of Delhi
300 covering an area of ~70 acres, receives around 2500-3500 metric tons of solid waste every day.
301 Many print/news reports also stated the unexpected overflowing of wastes at the Ghazipur
302 landfill site. The Bhalsawa landfill is situated to the North-west of megacity covers an area of
303 ~36 acres where every day ~2000-3000 tons of waste are dumped. The dumping ground of
304 Okhla is yet another landfill site with area of ~46 acres is great concern which receives at least
305 ~1500-2000 tons of waste dumped every day despite the site was declared exhausted in 2010.
306 The zones covered in waste collection for Okhla includes South, Central, Najafgarh, and Delhi
307 Cantonment Board (DCB).

308 Delhi has three operational Waste to Energy (WTE) Plants of total waste intake capacity
309 of ~5250-5750 tons per day (TPD) at three locations in Delhi namely Ghazipur, Bawana, and
310 Okhla. The waste-to-energy plant installed in Ghazipur has an installed capacity of 12 MW
311 processes ~1300-1350 TPD. The Bawana Integrated MSW plant processes ~2000-2300 TPD of
312 solid waste with an installed capacity of 24 MW. The Okhla WTE plant has an installed capacity
313 of 16 MW and processes ~1950 tons of municipal solid waste per day. Another WTE plant of

314 processing capacity ~2000 TPD is proposed at Tehkhand and another at Ghonda Gujran. After
315 the commission of these two proposed WTE plants, the total capacity would likely be increased
316 from 5750 TPD to at least 8450 TPD in upcoming future. These WTE plants potentially process
317 the waste for energy generation to some extent reduces the volume of landfills while providing a
318 renewable source of energy. Limited evidence has put forward that well-planned and well-
319 operated WTE plants might seem significant to reduce adverse health impacts, due to lesser
320 perilous emissions when compared to burning of waste at landfills, whereas, poorly fed WTE
321 plants potentially emit particulate matter and strenuous toxins with severe health risks (Cole-
322 Hunter et al., 2020). As a recent study reported that so far in India, only 23% of total generated
323 municipal solid waste is treated by various processing and approximately 43% of waste is
324 dumped some. Remaining 34% is allowed to burn openly at the landfill site itself in order to
325 prevent spilling over (Sharma et al., 2019). As there are three WTE plants installed within
326 megacity Delhi and quite evidently it processes ~22% of total MSW generated annually,
327 therefore it is estimated that only 48% of the total MSW is dumped and the left over 30% is
328 burnt right away on the dumping site which contributes to the air pollution issues in Delhi.


329 Along the same line, construction activities in Delhi are also one of the significant
330 contributors to particulate matter emission in Delhi. Construction activities include demolition,
331 site preparation and removal of debris. During the survey, at least 20 construction sites were
332 observed from which some of the major sites at DDA-Housing Sector 19-B, GH-project Sector
333 10, Megamall- sector 14, DDA-Housing, sector 16-B, Bhagwati C.G.H.S- Sector 22-Dwarka,
334 SaritaVihar- Metro enclave, Maharani Bagh flyover and Naraina flyover. HCVs and multi-utility
335 vehicles like bulldozers, tractors, scrapers, compactors involved in loading and unloading of
336 construction materials, preparation of site, demolition and disposal of debris which in a certain
337 way contribute to the dust load. Additional information on area and duration of construction
338 activities were procured from Public Works Department (PWD), 2020 and Delhi Development
339 Authority (DDA), 2020. In case of brick kilns industry, which is very much confined across the
340 outskirts of Delhi areas like Jhajjar, Faridabad and Ghaziabad region where there is a cluster of
341 kiln industries (like approximately 300 brick kilns in Jhajjar region). Operation of these brick
342 kilns is very seasonal in nature as their peak business month between December to June month.
343 It is also noticed that approximately 10 tons of coal or 13 tons of tudi/ rubber is being used to
344 produce one lakhs of bricks using semi-zig-zag technology. The sector is widely scattered in a

345 much unorganized manner where it is observed that coal (~70%) is being used as primary fuel
346 followed by tudi (i.e. mustard husk) (~25%) and rubber/other biomass/waste/etc. (~5%) as an
347 alternative fuel.

348 The practice of using Incense Sticks/Mosquito Coils/Cigarettes (IMC) has remained as an
349 unattended sector, which is of vital significance source to indoor and also moderately contributes
350 to outdoor air pollution. Use of incense sticks in festivals and holy places is common in India.
351 Besides that, during the field survey it was observed that maximum street vendors (this includes
352 both food zones and non-food zones) of Delhi as well as small scale dhabas using incense sticks
353 during business hours. The composition of incense sticks is responsible for continuous
354 smoldering. It comprises of resin, charcoal and wood dust mixed altogether and wrapped to thin
355 sticks made from coconut leaves or bamboos. Generally, incense sticks comprises of 45%
356 biomass, 25% wooden chips/bakhoor 15% coal and 15% resin/jigit (Cohen et al., 2013; Kumar et
357 al., 2014) and are responsible for emission of hazardous mixture of pollutants causing indoor air
358 pollution as well. It is very astonishing that most street vendors as well as dhaba/hotel lights
359 incense sticks/cake during business hours. Also, mosquito coils have been widely used by the
360 low/middle income grade households (Kumar et al., 2014) especially in slum zones which were
361 quite fascinating to observe during the field survey. The smouldering of the contents of coils:-
362 biomass, wood dust, and charcoal releases deadly pollutants responsible for acute respiratory
363 infections. Similarly, smoking of cigarettes/tobacco has caused over 10 million fatalities every
364 year in India. In fact, India has been declared home to at least 120 million smokers by World
365 Health Organization (WHO). The estimated emission for these sectors was based upon the
366 activity data of household population and street vendors with their per capita consumption.

367 Open-air funeral pyre the traditional system of cremating human bodies is a wide custom
368 in South Asian countries especially in India and Nepal (Chakrabarty et al., 2013) as the
369 population of Hindu religion is a majority. During the field campaign, around 62 crematoriums
370 were surveyed where it was found that only 6 crematoriums were observed to be using modern
371 electrical burning method as compared to 56 crematoriums with traditional method of burning of
372 wood. The pyre is built by using roughly ~450-550 kg of wood along with assorted materials,
373 such as shells of coconut, cow-dung, camphor, and pure ghee/clarified-butter. The dead body is
374 basically placed on top of the pyre and flaming process is carried out which takes around 4 to 6

375 hours. As stated by the vital statistics of Municipal Corporation of Delhi (MCD), the crude death
376 rate for Delhi has been reported 6.51 per 1000 people in 2020. No authentic data was accessible
377 regarding the number of dead bodies cremated everyday/annually in each crematorium except in
378 a few crematoria. So, the emission estimation was based on the population statistics on religion
379 data of Census from crematoriums, annual death rate, number of deaths, and quantity of wood
380 used. Later, the emission was spatially allocated to the respective crematorium grids. The Land
381 use and Land Cover pattern with activity data incorporated are highlighted in Figure 4.



382

383 Figure 4: Land Use and Land Cover with Spatial Surrogates over Megacity Delhi and NCR

384 **2. Methodology:**

385 Emission factors (EFs) are the most critical and sensitive components to build a reliable emission
386 inventory and the selection of an appropriate regional sector-specific technological emission
387 factor is the most crucial and challenging task and should be validated through scientific
388 judgments and acceptability. A dynamic EF can epitomize a better scenario of transport
389 emission, especially in a developing country like India where the usage of vehicle is much longer
390 as compared to developed countries. Based on our best judgment, in some cases, the EFs for
391 aging vehicle type are derived by averaging out the EFs given for 10 yr & 15 yr old vehicle
392 category. There are few EFs which are adopted from other countries due to lack of indigenous
393 EFs. Although many uncertainties prevail due to the sensitivity of EFs for development of
394 emission inventory, but the present effort is towards the best possible estimate by including EFs
395 already adopted in several authenticated reports and experiments conducted by certified agencies
396 of government as well as government authorized non-government and autonomous agencies
397 which provide best estimates for EFs, that are being referred in our latest studies Mangaraj et al.,
398 (2022 a, b) and taken into account in this present study.

399 **2.1. Calculation:**

400 The total emission i.e. the sum of emission from all individual sectors is expressed by Equation-1
401 with respect to particular pollutant. Most of sector's emission is estimated using IPCC Tire-2
402 approach. In the absence of activity data, the Tire-1 approach is adopted for few sectors. In case
403 of transport sector, the EFs are defined on the basic of kilometer travel, which is highly sensitive
404 to technology, and age of vehicle. In the presence of country specific technological EFs for
405 transport sector developed by ARAI, it is highly useful to prove the estimation. The emission
406 from transport sector has been calculated as per the Equation -2. In case of road dust emission,
407 the method is adopted from widely used AP-42, USEPA (Equation 3 and Equation 4) where the
408 country specific parameter like silt load, moisture content, no. of precipitation days and average
409 vehicular weight.

410 **Equations Used:**

411 $TE = \sum_r \sum_s FU_{r,s} [\sum_t Ef_{r,s,t} A_{r,s,t}]$ -----**(Equation 1)**

412 Where,
 413 r, s, t = sector, fuel type, technology, TE= Total emission, FU= Sector and fuel specific amount
 414 Ef= Technology specific EFs, A = fraction of fuel for a sector with particular technology, where
 415 $\sum A = 1$ for each fuel and sector.

416 $E_t = \sum(Vh_l \times D_l) \times Ef_{l,km}$ -----**(Equation 2)**

417 Where,
 418 E_t = Total Emission of compound, Vh_l =Number of Vehicle per type, D_l =Distance travelled in a
 419 year per different vehicle type, Ef_l, km = emission of compound, vehicle type per driven
 420 kilometre

421 **For Paved Road Dust:**

422
 423 $E_p = [k (st/2)^{0.91} (wt)^{1.02}] (1 - \frac{pt}{4N})$ -----**(Equation 3)**

424
 425 Where,
 426 E_p = particulate emission factor (having units matching the units of k),k = particle size multiplier
 427 for particle size range and units of interest, st = road surface silt loading (grams per square meter)
 428 (g/m²),
 429 wt = average weight (tons) of the vehicles travelling on the road, pt = number of “wet” days with
 430 at least 0.254 mm (0.01 in) of precipitation during the averaging period, N = number of days in
 431 the averaging period (e.g., 365 for annual, 91 for seasonal, 30 for monthly)

432
 433 **For Unpaved Road Dust:**

434 $E_{up} = \left\{ \left[k \left(\frac{st}{12} \right)^a \left(\frac{VS}{30} \right)^d / \left(\frac{m}{0.5} \right)^c - C \right] * [(365 - pt)/365] \right\}$ -----**(Equation 4)**

435
 436 where,
 437 E_{up} = size-specific emission factor (lb/VMT), st = surface material silt content (%), m = surface
 438 material moisture content (%), VS = mean vehicle speed (mph), C = emission factor for 1980's

439 vehicle fleet exhaust, brake wear and tire wear, pt = number of days in a year with at least 0.254
440 mm (0.01 in) of precipitation; k , a , c and d are empirical constants

441

442 **2.2. Spatial allocation of emission:**

443 The Geographical Information System (GIS) organizes the geographic data from various sources
444 followed by being a key aspect that allows these tools to transform large spatially uniformed
445 emission dataset to systematic thematic layers used for developing gridded emission inventory.
446 A high-resolution Land Use Land Cover (LULC) digital database over the megacity is used to
447 improve the spatial distribution of emission from various sectors. Before input of calculated
448 emission into the GIS environment, several preliminary tasks like geo-referencing, digitization
449 and building of attribute activity database are undertaken. A GIS based statistical approach is
450 developed to spatially distribute the emissions across the Delhi-NCR. Different layers of spatial
451 proxies have been taken into account to grid the emission values to required resolution
452 ($\sim 0.4 \times \sim 0.4$ km) for each sector, which can be used as tool for further analysis. The basic spatial
453 features are points, lines and polygons; layers of road networks- national and state highways,
454 major and minor roads; population density of village/district-level, the urban spread of the grid;
455 database on the economic activity of hospitals, market complexes, industrial estates, hotels,
456 residential blocks etc. These spatial features are used as proxies to determine the emission both
457 spatially and temporally where grid level emissions are allocated by overlaying the facility
458 location layer with the grid cell layer and aggregating the facility points in each cell covering
459 Delhi-NCR.

460 **3. Result & Discussion:**

461 The developed emission inventory for major air pollutants like $PM_{2.5}$, PM_{10} , CO, NO_x , SO_2 ,
462 VOC, BC and OC covering Delhi-NCR in 2020 are calculated to be 123.8 Gg/yr, 243.6 Gg/yr,
463 799.0 Gg/yr, 488.9 Gg/yr, 730.0 Gg/yr, 425.8 Gg/yr, 33.6 Gg/yr, and 20.3 Gg/yr respectively.
464 The sector-wise total emission of pollutants across Delhi-NCR is provided in Table-2. Also a
465 dataset has been provided at <https://doi.org/10.5281/zenodo.7715595> (Sahu et al., 2023) for
466 gridded pollutant wise sectorial spatial distribution. Keeping the space constraint in mind,
467 comprehensive analysis of the spatial distribution of PM_{10} and CO is elaborated further.

Sector	PM _{2.5}	PM ₁₀	CO	NO _x	VOC	SO ₂	BC	OC
Windblown-Road Dust	10.867	99.975	-	-	-	-	-	-
Transport	41.369	42.330	540.100	342.650	709.380	77.230	23.640	-
Industry	20.370	37.076	10.218	85.091	-	338.096	4.327	-
Household	0.311	1.310	1.038	0.867	0.005	0.227	0.065	0.113
Slum	0.216	0.550	1.443	0.463	0.010	0.086	0.018	0.107
Street Vendor	0.687	1.175	1.440	0.286	0.011	0.743	0.092	0.242
Crop Residue Burning	11.094	13.820	113.086	6.131	17.107	1.276	1.432	4.969
Cow-Dung	2.519	3.149	21.345	0.408	0.175	0.099	0.273	1.643
Diesel Generators	3.620	4.590	2.070	9.590	-	0.640	1.880	-
Aviation	-	-	21.007	36.068	3.297	2.871	0.021	0.036
MSW Burning	11.915	12.831	61.407	3.428	-	0.458	0.917	11.915
WTE Plants	10.217	10.441	0.786	2.021	0.022	1.853	-	-
Construction	5.956	9.926	-	-	-	-	-	-
Brick Kiln	2.727	4.017	12.807	0.913	0.041	2.106	0.896	0.773
IMC	1.161	1.379	4.143	0.060	0.005	0.105	0.031	0.021
Crematory	0.863	1.078	8.134	0.987	0.042	0.014	0.078	0.550
TOTAL	123.891	243.649	799.023	488.963	730.093	425.804	33.669	20.370

468 **Table 2: Pollutant-wise and sector-specific total emission across Delhi-NCR**

469 **3.1. Anthropogenic PM₁₀ Emission in Delhi-NCR:**

470 The total PM₁₀ emission across is estimated to be 243.6 Gg/yr, where windblown dust is
471 emerged as largest sources (99.9 Gg/yr) followed by traditionally dominating sector like
472 transport sector (42.3 Gg/yr) and industry (37.0 Gg/yr). It is also noticed that crop residue
473 burning (13.8 Gg/yr) and municipal solid waste burning (12.8 Gg/yr) in open area along with
474 waste-to-energy plants (10.4 Gg/yr) are emerging as larges source of particulate matters across
475 the city.

476 A high emission in the order of 1000-6000 tons/grid/yr and 120-1000 tons/grid/yr is
477 found over Central, Eastern, Northern, some parts towards the South and South-eastern fringes of
478 Delhi confined over national highways, many major and busy roads as shown in Figure 5.
479 Moderate emission in the order just 30-120 tons/grid/yr is well scattered across the study regions.
480 It has been noted that Central and Eastern Delhi region are one of the highly polluted regions.
481 Recent rising trend of vehicle numbers along with vehicle from surrounding states in Delhi road
482 in last ten year has put tremendous pressure on road network expansion, leading heavy traffic
483 congestion. All major traffic junctions are experiencing high emission load. However, the

484 highest emitting grids in the order of ~1300-6000 tons/grid/yr are also found in small patches
485 driven by sources like WTE plants and industrial practices followed by municipal solid waste
486 burning as well. It has been found that the Okhla region is one of the highly polluted hotspots
487 where WTE plants, municipal solid waste burning followed by windblown road dust are the
488 dominating sectors responsible for elevated PM₁₀ emissions. The next dominating hotspots
489 identified in Bawana and Ghazipur regions are dominated by large point sources like the WTE
490 plant with ~2566 tons/grid/yr and 1704 tons/grid/yr respectively. Furthermore, Anand Parbat
491 (~1300-1700 tons/grid/yr), Badli Industrial Area (~648 tons/grid/yr), Wazirpur Industrial Area
492 (~508 tons/grid/yr), Mayapuri Industrial Area (~400-500 tons/grid/yr), Rohini Industrial Area
493 (~481 tons/grid/yr) are some of the industrial dominating hotspots. It is noticed that coal is
494 predominantly used in both organized and unorganized industrial sector followed by diesel as
495 fuel. Dense major road networks across these regions led to slow-moving traffic congestion,
496 moreover these roads are concurrent to the major junctions of industrial area and they tend to
497 witness the large movement of heavy weighted HCVs and LCVs for the supply of raw materials
498 and goods. The continuous movement of these vehicles undoubtedly is responsible for the
499 broken and worn-out roads. Besides that, this vehicle-induced turbulence and poor road
500 condition are the leading factors accountable for road dust resuspension in an order of ~150-750
501 tons/grid/yr making it the second dominating sector overall. The gross weight of the HCVs and
502 LCVs also affects their speed while carrying the goods, which intensifies the vehicular exhaust
503 emission too, which is why the transport sector is the third dominating sector with ~70-300
504 tons/grid/yr. High vehicular density over many busy roads is the main cause of high particulate
505 emission due to moderate vehicular speed of ~25 km/hr. This speed increases towards the
506 outskirts of the city. The load of windblown road dust depends on vehicle speed, therefore the
507 traffic congestion leading to a decrease in average vehicle speed in Delhi is regarded as one of
508 the important factors that lead to suppressing the windblown dust but at the same time, it
509 increases the transport emission due to traffic congestion. Apart from this, heavy commercial
510 vehicles loaded beyond their carrying capacity cause resuspension of road dust, which results in
511 severe particulate pollution. Moreover, a significant amount of vehicle fleet plying over megacity
512 Delhi belongs to other states where the share of the personal and commercial car (taxi) can go as
513 high as ~30-40% on various road types.


514

515 **3.2. Anthropogenic CO Emission in Delhi-NCR:**

516 The estimated total CO emission from all the sources is found to be around 799.02 Gg/yr. The
517 relative contributions of CO from transport, industrial, residential and other sector are estimated
518 to be 67.5% (540 Gg/yr), 1.2% (10.21 Gg/yr), 17.5% (140.42 Gg/yr) and 13.5% (108.28 Gg/yr)
519 respectively. The spatial pattern as shown in Figure 5 depicts that CO emission hotspots in the
520 order of 750-6500 tons/grid/yr are found to be over the large region of Central, Eastern and
521 South-eastern Delhi regions along with few more over surrounding NCR regions like Noida,
522 Gurgaon, Gaziabad and Faridabad etc. Transport sector is the dominating source in the above
523 discussed regions due to high population and dense road network driving to high vehicular
524 activities. The estimated emission from transport is found to be around 540.10 Gg/yr, where the
525 petrol driven vehicles emits more CO as compared to diesel and CNG vehicle. The petrol
526 vehicles are mostly the personal vehicle in India whereas the vehicle numbers have gone up
527 nearly two folds in Delhi during last 10 years, contribute more than 80% of total CO emission.
528 Commercial vehicle growth contributes less to CO emission. Most of the major traffic junctions
529 in down town are highly polluted by transports related CO emission (~1200-1800 ton/yr). Most
530 of CO emitting Industrial zones in Delhi is more confined to Central & Eastern Delhi and few
531 more specific regions outskirts of Delhi.

532 The second most dominant source is residential sector where major slum clusters
533 contribute significantly. The regions are more confined to the Central, Eastern, South-Easter part
534 of Delhi and few surrounding regions. It is also found that highly dense population with middle
535 and lower income group is lying over above discuss areas too and associated slum cooking,
536 residential cooking, street vendors and commercial cooking etc. Low technological cum soil fuel
537 based cooking practices in slum areas drive to high CO emission. Moreover, the slum population
538 located in the Eastern and Central Delhi is dense aggravates CO emission further. A relatively
539 low emission of the order of 25-150 ton/yr is found to be in the outskirts of Delhi and adjacent
540 districts like Rotak, Jhajjar and Guttam Budhanagar etc. Low population density along with
541 agricultural lands cover are the main reasons for low emission of CO. Collectively, the street
542 vendor cooking and commercial cooking contribute a significant amount of CO emission in
543 densely populated regions and are well uniformly scattered over large area. Similar hotspots are

544 also identified over the Noida, Gurgaon, Faridabad regions surrounding the Delhi where an
 545 emission of the order 1000-1500 tons/yr is found.




546
 547 **Figure5: Spatial distribution of PM₁₀ and CO load across Delhi-NCR region**

548 **3.3. Decadal Change in Emission (2010-2020):**

549 Shifting of emission sources and its trend over the years is vital to access the impact of air
 550 pollution especially in megacities. The present estimated PM₁₀ emission is compared with our
 551 own previous estimation for the base year 2010 (SAFAR-Delhi, 2010) for same domain, it is
 552 clearly concluded that the effective net increase of PM₁₀ emission over the last decade is just
 553 ~3%. This is small growth could be due to various new policy being adopted by government
 554 which is directly or indirectly influence the emission. At the same time, there is couple of shift in
 555 sectorial emission load as well as addition of new unorganized sectors in 2020 emission
 556 estimation. If you look at the sector specific change then there are significant shift in emission
 557 pattern and required attention. It can be observed that there has been an increase by 39% in

558 emission load from transport sector as compared to another 36% in industrial sector during same
559 period. In case of windblown road dust emission, there is a decrease of 23% as shown in Figure
560 7. Due to penetration of LPG in slums, the cooking related emission is improved significantly as
561 well as in residential sector. The rise in number of vehicles with increase in spread of road
562 networks turned out to be the major cause along with the overburdening of four wheeler cars,
563 where the contribution of other state cars is significant. However, there is an increase in traffic
564 congestion but better paved road condition and road shoulder maintenance has resulted in a
565 decrease in emission load from windblown road dust in last one decade. The discontinuation
566 (permanent closure) of the thermal power plants in Delhi has resulted in exclusion of thermal
567 power plant as a sector contributing to total emission load.

568 As far as the residential sector is concerned, there is a rapid reduction in relative
569 contribution. The decrease in number of slums in Delhi when compared to 2010 period has
570 resulted in a reduction in consumption of cooking fuels, which shows a significant decline in
571 residential PM₁₀ load by 31%. Primarily the awareness among the people led to penetration of
572 LPG in slum areas, street vendors, household etc., which reduced the emissions to great extent.
573 However, emissions from other sectors have significant contribution to the present PM₁₀ load.
574 The new emerging sectors like WTE plant, MSW burning, crematory, use of incense
575 sticks/mosquito coil/cigarettes and construction, were not considered in the previous report in
576 2010 so the relative contribution has increased significantly. This decadal change in emission is
577 also observed in the case of CO in similar trend except residential where there is a substantial
578 decreasing trend as shown in Figure 6. A summary of the growth trend for all the pollutants is
579 shown in Table 3.



580

581 **Figure 6: Decadal change of emission with sectorial relative contribution**

582

Base Year	2010*						2020**					
	PM _{2.5}	PM ₁₀	CO	NO _x	BC	OC	PM _{2.5}	PM ₁₀	CO	NO _x	BC	OC
Transport	30.25	30.29	427.55	162.28	9.77	-	41.37	42.33	540.10	342.65	23.64	-
Windblown Road Dust	26.20	131.27	-	-	-	-	10.87	99.98	-	-	-	-
Industry	16.29	27.20	10.92	79.84	8.67	12.60	20.37	37.08	10.22	85.09	4.33	-
Power	2.87	11.02	0.29	6.90	0.04	-	-	-	-	-	-	-
Residential	18.65	36.07	264.41	6.40	2.96	2.60	18.45	24.60	140.42	17.75	3.76	7.07
Others	-	-	-	-	-	-	32.84	39.67	108.28	43.48	1.94	13.29
Total	94.26	235.85	703.17	255.42	21.44	15.20	123.89	243.65	799.02	488.96	33.67	20.37

*SAFAR Delhi 2010; **Present Study All Emission in Gg/yr

583 **Table 3: Comparison of sectorial emission during 2010 and 2020**

584 **3.4. Uncertainty in emissions and limitations:**

585 Emission inventories may have errors due to activity data and EFs gaps. Therefore, the collection
 586 of data and the evaluation of uncertainty are unambiguously linked. We have made an attempt to
 587 estimate the uncertainty in the sectorial emissions for which, error propagation was calculated by

588 following the Monte Carlo methodology. The factors included for uncertainty estimation include
589 the (a) emission factors used, (b) activity data collection, (c) proxy data used, (d) data gaps
590 leading to approximation, and (e) efficiency of emission control. Uncertainty estimation for the
591 transport sector seems very complex as it involves fuel-specific technological vehicle categories
592 that have diversity in emission factors according to the age of vehicles. In the case of transport,
593 the disparity in activity data and VKT is not much as a robust ground survey was performed.
594 Therefore, the contribution of vehicular emission to gross uncertainty is the least with a
595 maximum uncertainty for ranging $\pm 23\%$. Emission from windblown road dust has heterogeneous
596 factors like the speed of the vehicle along with its weight, soil moisture content and silt load.
597 These modulating factors are responsible for defining the emission load and their combined
598 uncertainty ranges $\pm 33\%$. The residential/domestic emission source comprises of per capita fuel-
599 induced activity data and corresponding emission factors so the combined uncertainty in this
600 sector is $\pm 28\%$. The industrial sector has the highest disparity in secondary activity data and its
601 availability of relevant technological emission factors is the key factor to a higher uncertainty
602 level of $\pm 41\%$. The sources belonging to other sectors comprise several minor unorganized
603 sources, which have comparatively less contribution to total emission and have high uncertainty
604 ranging $\pm 47\%$. The gross uncertainty in the inventory is estimated to be around $\pm 29\%$, which is
605 found to be in an acceptable range. As of date, no comprehensive study has been done to
606 determine the uncertainty for the emission inventory of Delhi. This is the first approach to do the
607 same and in accordance with our best scientific judgment, it can be said that the present surface
608 emission dataset both in terms of quality and quantity has the least errors. The emission
609 inventory's limitation lies in various steps like limited access to industrial information like the
610 one fuel quantity used in various techniques used. Similarly, the exact number of other state
611 vehicle plying in the megacity is very uncertain and need a better approach to improve the
612 estimation. There are many unorganized sectors like street vendors; small-scale waste burning
613 across the local level, silt load on various roads, driving conditions varies with road type and its
614 condition etc. Still, we believe the kind of micro-level activity data used is better than any other
615 earlier inventories developed over the study region.

616

617 **3.5. Inter-comparison among studies:**


618 In this section, a comparative analysis of the present study with the past studies is taken into
619 account and has been elaborated. As mentioned earlier, Delhi has been in the spotlight when air
620 quality issues are concern. Here, the present study is compared with previous eight studies done
621 over Delhi. NEERI in 2010 presented sector-wise emission inventory at 2 km resolution covering
622 the metropolitan area of Delhi for the base year 2007, targeting only four pollutants (PM₁₀, SO₂,
623 NO_x, CO). The calculated emissions were found to be 147 tons/day, 268 tons/day, 460 tons/day,
624 and 374.1 tons/day respectively. Guttikunda and Calori (2013) worked on the National Capital
625 Territory (NCT) region that includes Delhi and its suburbs (Gurgaon, Noida, Faridabad, and
626 Ghaziabad) over an area of 6400 km² at ~1 km resolution. This was done for the base year 2010
627 for PM, SO₂, NO_x, CO, and VOCs. It includes sectors of re-suspended road dust, construction,
628 vehicular exhaust, domestic cooking, power plants, industries, brick kilns, diesel gen-sets, and
629 waste burning. About 35% of the total PM₁₀ emission is contributed by the transport sector and
630 road dust and around 37% are contributed by the major point sources (brick kilns, industries, and
631 power plants). It has been highlighted that brick kilns located outside the city affect the city air to
632 some extent but the origin of certain sources like diesel gen-sets, waste burning, and construction
633 remains unclear whether they have been influenced by the surrounding areas or not. In addition,
634 the domain of interest considered is around 69% of the total area of Delhi, which is huge, and
635 therefore it doesn't give a clear representation of the exact emissions prevailing in Delhi.
636 Sindhvani et al., (2015) estimated PM₁₀, CO, NO_x, and SO₂ emissions in the NCR-Delhi region
637 that comprises the neighboring states of Haryana and Uttar Pradesh. This study was done in the
638 year 2010 at a 2 km×2 km resolution. The estimated total emissions for PM₁₀, CO, NO_x, and SO₂
639 were 107.47 Gg/yr, 1290.13 Gg/yr, 342.30 Gg/yr, and 83.16 Gg/yr respectively. The contribution
640 of sectors like road transport, road-dust and domestic sources altogether is ~47% of total PM₁₀
641 emissions. A quantitative assessment of only three pollutants i.e., PM, NO_x, and CO was carried
642 out for Delhi Urban Area for the base year 2010 by Mishra and Goyal, (2015). The major
643 contributors included vehicles, industries, power plants, and domestic and dust. The CO and NO_x
644 emissions from the transport sector (210.83 kt and 92 kt respectively), were found to be the
645 largest contributor followed by the domestic sector. Road dust (25.50 kt) has a significant
646 contribution to PM while vehicular, industries and power plants are approximately having equal
647 contributions. Similarly, Jaiprakash et al., (2016) reported an experimental-based study focusing
648 on specifically vehicular emissions (CO, CO₂, and NO_x) in Delhi for the base year 2012. The

649 study estimates an on-road tailpipe measurement of 14 passenger cars of different types of fuel
650 and vintage and reported that the share of diesel, gasoline, and CNG to total CO, CO₂, and NO_x
651 emissions were in order of 7:84:9, 50:48:2 and 58:41:1 respectively. These studies majorly lack
652 in accounting for the impactful active sources like commercial cooking (street vendors),
653 crematoria, WTE plants, crop residue burning, and many more, which makes this inventory
654 insignificant for further use.

655 Sharma and Dikshit, (2016) attempted a comprehensive study on PM₁₀, PM_{2.5}, NO_x, SO₂,
656 and CO in Delhi city focusing on ~14 sources for the base year 2014 (November 2013 – June
657 2014) at 2 km resolution. The results showed that road dust (56%), concrete batching (10%),
658 industrial sources (10%) and vehicular (9%) are the major contributors to PM₁₀ emission.
659 Though the study involved site sampling for a few of the sectors it also lacks an absolute
660 sampling number (limitation) and most of the activity data were collected from secondary
661 sources. Singh et al., (2018) attempted the estimation of emissions from the road transport sector
662 of NCT-Delhi for the base year 2010. The study stated that major roads contribute to more than
663 50% of total PM emissions. When specifically focusing on limited pollutants, which most
664 importantly include PM, this study has certain limitations in terms of non-exhaust emission
665 (vehicular dust resuspension) from road transport, which is a significant contributor to the city's
666 PM₁₀ load. Thereafter, TERI & ARAI (2018) initiated a source apportionment study for
667 identifying sources responsible for PM_{2.5} and PM₁₀ in Delhi-NCR and developed an coarse
668 resolution (4 km×4 km) based emission inventory of a few pollutants (PM, NO_x, SO₂, CO,
669 NMVOC) for 2016. The results stated that in the case of PM₁₀, road dust and construction dust
670 contributed significantly, where the contribution of dust from surrounding regions was
671 comparatively higher in summers, which reduced the proportion of major sectors in the PM₁₀.

672 Taken as a whole, a large disparity is found between the reported past studies and present
673 emission estimations as shown in Figure 7. The basic reasons for these variations point towards
674 the differences in sectors being focused on or the activity data being considered for the past
675 works in conjunction with the use of technological emission factors used are also an additional
676 reason of concern. The base years as well as domain considered differ significantly from each
677 other. As the sources of emission tend to change with time and the evolution of a region hence,
678 upgrading an emission inventory is the most fundamental segment to be taken care of. As a

679 consequence, this present study has premeditated all such important factors in the most potent
 680 ways to build up this gridded surface-emission dataset. In addition to this, unlike the previous
 681 works, this study is the first-ever ultra-high-resolution-gridded (~400 mts) emission data set
 682 targeting eight major pollutants for the latest base year 2020. This new dataset could be a
 683 valuable element in air quality management (mitigation strategies) and air quality modelling a
 684 study, which is why it is believed to be more reliable data.




685
 686 **Figure 7: Inter-comparison among studies over the domain of interest**
 687

688 **3.6. Mitigation strategy using developed emission:**

689 As emission inventory acts as a fundamental tool by both policymakers and scientific
 690 communities for mitigation strategies in combating air pollution in any cities. In the present

691 study, a developed sensitive piece of surface-gridded emission database is unique in many ways
 692 and will pave a path to understanding the air quality issues in megacity Delhi. For the same, a
 693 thorough analysis has been made to identify the contribution of major sectors to the high
 694 emitting polluted zone across the Delhi. Following that, a number of hotspot regions were
 695 identified as shown in Figure 8 (a), from which the top ten hotspots are being identified along
 696 with first three dominating sectors affecting the air over the hotspots significantly as shown in
 697 Figure 8 (b). Since, PM₁₀ is considered to be one of the dominating pollutants in modulating
 698 urban air quality. In one of the applications to the developed emission inventories, sector-specific
 699 control strategies are recommended based on the input of available activity data and emission
 700 factors, which would possibly benefit the policymakers and help in the improvement of megacity
 701 air quality. The ten most dominating hotspots are identified with the relative contribution of three
 702 major sectors in descending order as identified in the table to follow. Each area mentioned
 703 against each megacity below is accompanied by several color codes which denote a specific
 704 sector associated with the pollution where; TRN- Transport, WB- Wind-blown road dust, IND- Industry,
 705 TPP- Thermal Power plant, SLM- Slum, MSW- Municipal Solid Waste burning, DG- Diesel Generator, WTE-
 706 Waste-to-energy plant, RES- Residential, CON- Construction.



707
 708 **Figure 8: (a) Hotspots across the Delhi-NCR domain, with (b) First three dominating**
 709 **sectors affecting the air over the hotspots**

711 Based on the analysis of each hotspot as delineated in Figure 8, the mitigation strategy can be
712 framed accordingly to control the emission at source through various approaches on the ground.
713 Apart from this, a few sector-specific, generalized recommendations have been listed below for
714 all the megacities:-

- 715 a) The discard of ageing vehicles (more than 10 yr) especially commercial cars and heavy
716 commercial vehicles category from the system followed by fast traffic movement along with
717 enhanced penetration of electric vehicles can reduce the transport-related emission
718 significantly. The heavy and light commercial (diesel) vehicles together contribute ~40-50%
719 of road transport emissions where strict implementation of BS-VI norms needs to be applied.
- 720 b) Vehicles from surrounding states/regions play a significant role, where the average low traffic
721 speed is major roads cause of elevated emission of pollutants across the megacity, so a similar
722 stringent vehicular policy has to be implemented in surrounding states of Delhi too.
- 723 c) Major identified roads in megacity need road diversions in order to reduce the vehicle density,
724 which will ultimately increase the speed of vehicles by reducing emission load from tailpipes.
- 725 d) Flexible office hours and work from home culture could be an alternative approach to reduce
726 traffic congestion and at the same time, will increase average speed of vehicles and associated
727 reduction in emissions.
- 728 e) In order to reduce the impact of silt load, Road shoulders must be repaired in regular intervals
729 to avoid impaired and fractured ways. Similar approach should be adopted around outskirts of
730 Delhi too. They should be cleaned periodically.
- 731 f) Implementation of more stringent standards for both large and small-scale industries along
732 with better solid/fossil fuels utilization.
- 733 g) Open burning at Municipal solid waste dumping sites should be replaced with other substitute
734 approaches like vermi-composting, natural decomposition, or mulching and encourage WTE
735 plants.
- 736 h) Slum clusters with better penetration of LPG-based cooking fuel usage to discourage solid
737 fuels like fuel wood, cow dung, and coal.
- 738 i) Construction sites should be properly handle materials while loading and unloading
739 procedures.
- 740 j) Discouraging usage of DG-set usage in unorganized industries and commercial and private
741 zone could potentially help reduce the emission further.

742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771

4. Data availability:

The emission dataset can be accessed through open access data repository <https://doi.org/10.5281/zenodo.7715595> (Sahu et al., 2023). The dataset is presented in .shp file format covering Delhi-NCR region having domain size of 70km×65 km.

5. Conclusion:

Present megacities are facing pressing air quality challenges in South Asia due to variety of individual regional sources and changing policy, therefore, the present study is attempt to decode the understanding of present air quality over megacity Delhi through ultra-fine Emission Inventory for 2020 proclaims to be an essential component not only to address the mitigation plan towards improving megacity air quality but also understand the decadal change (2010-2020) in emission patter in megacity Delhi and surrounding NCR. The decadal change with changing government policy and action plan has modulated the emission from various unattended sources. However, only a single strategy cannot tackle the elevated air pollution issues in Delhi-NCR. A mixture of policy measures well adapted for domain's hotspot-specific, source-specific strategies is imperative to improve air quality. The developed surface emission dataset provides every such detail which can be comprehended as robust in all terms.

Author contributions:

Saroj Kumar Sahu (SKS) conceived the present idea and Poonam Mangaraj (PM) wrote the whole paper and analyzed the data. Gufran Beig (GB) provided useful discussion and suggested a conclusion.

Acknowledgements:

Authors acknowledge Ministry of Earth Science, Govt. of India, for partial support through QUISARC project (Grant No. MoES/Indo-Nor/PS-10/2015). This full research work did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors

772 **References:**

- 773 Akimoto, H.: Global air quality and pollution, *Science.*, 302, 1716-1719,
774 <https://doi.org/10.1126/science.1092666>, 2003.
- 775 Beig, G., Sahu, S. K., Singh, V., Tikle, S., Sobhana S. B., Gargeva, P., Ramakrishna, K., Rathod,
776 A., Murthy, B. S.: Objective evaluation of stubble emission of North India and
777 quantifying its impact on air quality of Delhi, *Science of The Total Environment.*, 709,
778 136126, <https://doi.org/10.1016/j.scitotenv.2019.136126>, 2020.
- 779 Beig, G., Sahu, S.K., Anand, V., Bano, S., Maji, S., Rathod, A., Korhale, N., Sobhana, S.B.,
780 Parkhi, N., Mangaraj, P., Srinivas, R., Peshin, S.N., Singh, S., Latha, R., Shinde, R.,
781 Trimbake, K.: India's Maiden air quality forecasting framework for megacities of
782 divergent environments: The SAFAR-project, *Environmental Modelling and Software*,
783 145, 105204, <https://doi.org/10.1016/j.envsoft.2021.105204>, 2021.
- 784 Beig, G., Srinivas, R., Parkhi, N. S., Carmichael, G. R., Singh, S., Sahu, S. K., Rathod, A., Maji,
785 S.: Anatomy of the winter 2017 air quality emergency in Delhi, *Science of The Total*
786 *Environment*, 681, 305-311, <https://doi.org/10.1016/j.scitotenv.2019.04.347>, 2019.
- 787 Chakrabarty, R.K., Pervez, S., Chow, J.C., Watson, J.G., Dewangan, S., Robles, J. and Tian, G.:
788 Funeral pyres in South Asia: Brown carbon aerosol emissions and climate impacts,
789 *Environmental Science & Technology Letters*, 1, 44-48,
790 <https://doi.org/10.1021/ez4000669>, 2013.
- 791 Cohen, R., Sexton, K.G., Yeatts, K.B.: Hazard assessment of United Arab Emirates (UAE)
792 incense smoke, *Sci Total Environ.*, 458, 176-86,
793 <https://doi.org/10.1016/j.scitotenv.2013.03.101>, 2013.
- 794 Cole-Hunter, T., Johnston, F.H., Marks, G.B., Morawska, L., Morgan, G.G., Overs, M., Porta-
795 Cubas, A., and Cowie, C.T.: The health impacts of waste-to-energy emissions: a
796 systematic review of the literature, *Environ. Res. Lett.*, 15, 123006,
797 <https://doi.org/10.1088/1748-9326/abae9f>, 2020.

798 DDA (Delhi Development Authority): Annual Report 2019-
799 20,https://dda.gov.in/sites/default/files/Annual_Report_2019-20_English.pdf, (last
800 access: 12Jun, 2021), 2020.

801 Delhi Urban Shelter Improvement Board: List of JJ Bastis,[https://delhishelterboard.in/main/wp-](https://delhishelterboard.in/main/wp-content/uploads/2019/10/JJBastisList675.pdf)
802 [content/uploads/2019/10/JJBastisList675.pdf](https://delhishelterboard.in/main/wp-content/uploads/2019/10/JJBastisList675.pdf), (last access: 15Apr, 2020), 2019.

803 DoT (Department of Telecommunications): Telecom Statistics of India. Economics Research
804 Unit Ministry of Communications. Government of India, New Delhi.,
805 <https://dot.gov.in/reportsstatistics/telecom-statistics-india-2019>, (last access: 5Sep,
806 2021), 2019.

807 DGCA (Directorate General of Civil Aviation): Handbook on Civil Aviation Statistics 2019-
808 20,<https://www.dgca.gov.in/digigov-portal/?page=4252/4205/sericename>, (last access:
809 12 Jun, 2020), 2020.

810 Down to Earth: [https://www.downtoearth.org.in/coverage/when-will-india-be-able-to-control-](https://www.downtoearth.org.in/coverage/when-will-india-be-able-to-control-pollution--17950)
811 [pollution--17950](https://www.downtoearth.org.in/coverage/when-will-india-be-able-to-control-pollution--17950), (last access: 8 June 2022), 2015.

812 DPCC (Delhi Pollution Control Committee): Status of Solid Waste Management in Delhi,
813 <https://www.dpcc.delhigovt.nic.in/solidwastemanagement>, (last access: 15 March
814 2020), 2020

815 GBD (Global Burden of Disease): Burden of disease from the joint effects of household and
816 ambient air pollution for 2016. Version 2. Summary of results. World Health
817 Organization, Geneva Switzerland,[https://cdn.who.int/media/docs/default-source/air-](https://cdn.who.int/media/docs/default-source/air-quality-database/aqd-2018/ap_joint_effect_bod_results_may2018.pdf)
818 [quality-database/aqd-2018/ap_joint_effect_bod_results_may2018.pdf](https://cdn.who.int/media/docs/default-source/air-quality-database/aqd-2018/ap_joint_effect_bod_results_may2018.pdf), (last access: 9
819 June, 2020), 2018.

820 Guttikunda, S., and Calori, G.: A GIS based emissions inventory at 1km x 1km spatial resolution
821 for air pollution analysis in Delhi, India, Atmospheric Environment, 67, 101-
822 111,<https://doi.org/10.1016/j.atmosenv.2012.10.040>, 2013.

823 HEI (Health Effects Institute): State of Global Air/2019. A special report on global exposure to
824 air pollution and its disease burden, ISSN 2578-6873,

825 https://www.stateofglobalair.org/sites/default/files/soga_2019_report.pdf, (last access:
826 10 Sep, 2020), 2019.

827 HEI (Health Effects Institute): State of Global Air/2020. A special report on global exposure to
828 air pollution and its health impacts, ISSN 2578-6873, <https://www.stateofglobalair.org/>
829 (last access: 14 Sep, 2020), 2020.

830 India Today: Air quality worsens in Delhi with onset of winter, 'severe' in Noida,
831 [https://www.indiatoday.in/india/story/air-quality-worsens-delhi-onset-of-winter-severe-](https://www.indiatoday.in/india/story/air-quality-worsens-delhi-onset-of-winter-severe-category-noida-2303441-2022-11-30)
832 [category-noida-2303441-2022-11-30](https://www.indiatoday.in/india/story/air-quality-worsens-delhi-onset-of-winter-severe-category-noida-2303441-2022-11-30) (last access: 30 Nov, 2022), 2022.

833 IPCC (Intergovernmental Panel on Climate Change): Special Report on Emissions Scenarios.
834 ISBN 0-521-80081. <https://www.ipcc.ch/report/emissions-scenarios/>(last access: 27Apr,
835 2021), 2000.

836 Jaiprakash, Habib, G., Kumar, A., Sharma, A., Haider, M.: On-road emissions of CO, CO₂, and
837 NO_x from four wheeler and emission estimates for Delhi. Journal of Environmental
838 Sciences, 53, 39 – 47, <https://doi.org/10.1016/j.jes.2016.01.034>, 2016.

839 Kumar, R., Gupta, N., Kumar, D., Mavi, A.K., Singh, K., Kumar, M.: Monitoring of indoor
840 particulate matter during burning of mosquito coil, incense sticks and dhoop. Indian
841 Journal of Allergy, Asthma and Immunology, 28, 68-73, [https://doi.org/10.4103/0972-](https://doi.org/10.4103/0972-6691.140770)
842 [6691.140770](https://doi.org/10.4103/0972-6691.140770), 2014.

843 Li, L., Maher, K., Navarre-Sitchler, A., Druhan, J., Meile, C., Lawrence, C., Moore, J., Perdrial,
844 J., Sullivan, P., Thompson, A., Jin, L., Bolton, E.W., Brantley, S.L., Dietrich, W.E.,
845 Mayer, K.U., Steefel, C.I., Valocchi, A., Zachara, J., Kocar, B., Mcintosh, J., Tutolo,
846 B.M., Kumar, M., Sonnenthal, E., Bao, C., Beisman, J.: Expanding the role of reactive
847 transport models in critical zone processes, Earth-Science Reviews, 165, 280-301,
848 <https://doi.org/10.1016/j.earscirev.2016.09.001>, 2017.

849 Mangaraj, P., Sahu, S.K., Beig, G., Samal, B.: Development and assessment of inventory of air
850 pollutants that deteriorate air quality in Indian Megacity Bengaluru, Journal of Cleaner
851 Production, 360, 132209, <https://doi.org/10.1016/j.jclepro.2022.132209>, 2022b.

852 Mangaraj, P., Sahu, S.K., Beig, G., Yadav, R.: A comprehensive high-resolution gridded
853 emission inventory of anthropogenic sources of air pollutants in Indian megacity
854 Kolkata, SN Appl. Sci., 4, 117.<https://doi.org/10.1007/s42452-022-05001-3>, 2022a.

855 Manisalidis, I., Stavropoulou, E., Stavropoulos, A. and Bezirtzoglou, E.: Environmental and
856 Health Impacts of Air Pollution: A Review, Front. Public Health., 8, 14,
857 <https://doi.org/10.3389/fpubh.2020.00014>, 2020.

858 Ministry of Civil Aviation: Annual Report 2019-
859 20,<https://www.civilaviation.gov.in/sites/default/files/annual-report-2019-2020.pdf> (last
860 access: 12Jun, 2020), 2020.

861 Mishra, D., and Goyal, P.: Quantitative Assessment of the Emitted Criteria Pollutant in Delhi
862 Urban Area, Aerosol and Air Quality Research, 15, 1601–1612,
863 <https://doi.org/10.4209/aaqr.2014.05.0104>, 2015.

864 Molina, L.T., Molina, M.J., Slott, R.S., Kolb, C.E., Gbor, P.K., Meng, F., Singh, R.B., Galvez,
865 O., Sloan, J.J., Anderson, W.P., Tang, X., Hu, M., Xie, S., Shao, M., Zhu, T., Zhang,
866 Y.H., Gurjar, B.R., Artaxo, P.E., Oyola, P., Gramsch, E., Hidalgo, D., Gertler, A.W.:
867 Air Quality in Selected Megacities, Journal of the Air & Waste Management
868 Association, 54, 1-73, <https://doi.org/10.1080/10473289.2004.10471015>, 2004.

869 Molina, M.J. and Molina, L.T.: Megacities and Atmospheric Pollution, Journal of the Air &
870 Waste Management Association, ISSN 1047-3289, J. Air & Waste Manage. Assoc.,
871 54,644–680.<https://doi.org/10.1080/10473289.2004.10470936>, 2004.

872 MoRTH (Ministry of Road Transport and Highways): Annual Report 2019-20,
873 <https://www.morth.nic.in/>, (last access: 11 Sep, 2020), 2020.

874 MoRTH (Ministry of Road Transport and Highways, Government of India): Notification. G.S.R.
875 653(E), THE GAZETTE OF INDIA: EXTRAORDINARY [PART II—SEC. 3(i)].
876 Ministry of Road Transport and Highways, Government of India, New Delhi,
877 India,<https://morth.nic.in/Circulars-Notifications-related-to-Vehicle-Scrapping-Policy>
878 (last access: 21 Dec, 2021), 2021.

879 NEERI (National Environmental Engineering Research Institute): Air Quality Monitoring,
880 Emission Inventory & Source Apportionment Studies for Delhi,
881 <https://cpcb.nic.in/displaypdf.php?id=RGVsaGkucGRm>, (last access: 21 Feb, 2020),
882 2010.

883 Permadi, D.A., Kim Oanh, N.T., Vautard, R.: Integrated emission inventory and modeling to
884 assess distribution of particulate matter mass and black carbon composition in Southeast
885 Asia, *Atmos. Chem. Phys.*, 18, 2725–2747, <https://doi.org/10.5194/acp-18-2725-2018>,
886 2018.

887 PWD (Public Works Department): Road and Building Construction Works,
888 [https://www.pwddelhi.gov.in/Home/ProjectProgress?pwd=Tg3814v+U9IMV5dBcmZue](https://www.pwddelhi.gov.in/Home/ProjectProgress?pwd=Tg3814v+U9IMV5dBcmZueR/JLRTSNPoIqdU52zWDuaG5M+Um4MKP/xsb5s6W0V9t)
889 [R/JLRTSNPoIqdU52zWDuaG5M+Um4MKP/xsb5s6W0V9t](https://www.pwddelhi.gov.in/Home/ProjectProgress?pwd=Tg3814v+U9IMV5dBcmZueR/JLRTSNPoIqdU52zWDuaG5M+Um4MKP/xsb5s6W0V9t) (last access: 12 Jun,
890 2020), 2020.

891 Rainfall Statistics of India: Hydromet Division, Indian Meteorological Department, Lodi Road,
892 New Delhi, Report No. MoES/IMD/HS/RAINFALL REPORT/30, (last access: 15Aug,
893 2020),
894 [http://hydro.imd.gov.in/hydrometweb/\(S\(k1llxc451oipskqly0iivw55\)\)/PRODUCTS/Pub](http://hydro.imd.gov.in/hydrometweb/(S(k1llxc451oipskqly0iivw55))/PRODUCTS/Publications/Rainfall%20Statistics%20of%20India%20-%202019/Rainfall%20Statistics%20of%20India%20-%202019.pdf)
895 [lications/Rainfall%20Statistics%20of%20India%20-](http://hydro.imd.gov.in/hydrometweb/(S(k1llxc451oipskqly0iivw55))/PRODUCTS/Publications/Rainfall%20Statistics%20of%20India%20-%202019/Rainfall%20Statistics%20of%20India%20-%202019.pdf)
896 [%202019/Rainfall%20Statistics%20of%20India%20-%202019.pdf](http://hydro.imd.gov.in/hydrometweb/(S(k1llxc451oipskqly0iivw55))/PRODUCTS/Publications/Rainfall%20Statistics%20of%20India%20-%202019/Rainfall%20Statistics%20of%20India%20-%202019.pdf), 2019.

897 SAFAR-Delhi-2010: Development of High Resolution Emission Inventory for National Capital
898 Region Delhi, Ministry of Earth Sciences, Govt. of India, (Special Scientific Report;
899 SAFAR-2010-A), <http://koha.moes.gov.in:8083/jspui/handle/123456789/18001>, (last
900 access: 1 Sept, 2022), 2010

901 SAFAR-Delhi-2018: SAFAR-High Resolution Emission Inventory of Megacity Delhi 2018.
902 Ministry of Earth Sciences, Govt. of India, (Special Scientific Report; SAFAR-Delhi-
903 2018-A, ISSN: 0252-1075), <http://safar.tropmet.res.in/source.pdf> (last access: 1 Sept,
904 2022), 2018.

905 Sahu, S.K., Beig, G., Parkhi, N.: High Resolution Emission Inventory of NO_x and CO for Mega
906 City Delhi, India, *Aerosol and Air Quality Research*. 15, 1137–1144,
907 <https://doi.org/10.4209/aaqr.2014.07.0132>, 2015.

908 Sahu, S.K., Beig, G., Parkhi, N.S.: Emissions inventory of anthropogenic PM_{2.5} and PM₁₀ in
909 Delhi during Commonwealth Games 2010, *Atmos. Environ.*, 45, 6180-6190.
910 <https://doi.org/10.1016/j.atmosenv.2011.08.014>, 2011.

911 Sahu, S.K., Mangaraj, P., Beig, G.: Gridded Surface Emission Data for Delhi-NCR (Version V2)
912 [Data set], Zenodo, <https://doi.org/10.5281/zenodo.7715595>, 2023.

913 Sahu, S.K., Mangaraj, P., Beig, G., Samal, A., Pradhan, C., Dash, S., Tyagi, B.: Quantifying the
914 high resolution seasonal emission of air pollutants from crop residue burning in India.
915 *Environmental Pollution*, 286, 117165, <https://doi.org/10.1016/j.envpol.2021.117165>,
916 2021.

917 Sharma, G., Sinha, B., Pallavi, Hakkim, H., Chandra, B.P., Kumar, A., Sinha, V.: Gridded
918 Emissions of CO, NO_x, SO₂, CO₂, NH₃, HCl, CH₄, PM_{2.5}, PM₁₀, BC, and NMVOC
919 from Open Municipal Waste Burning in India, *Environ. Sci. Technol.*, 53, 4765–4774,
920 <https://doi.org/10.1021/acs.est.8b07076>, 2019.

921 Sharma, M., and Dikshit, O.: Comprehensive Study on Air Pollution and Green House Gases
922 (GHGs) in Delhi. <https://cerca.iitd.ac.in/uploads/Reports/1576211826iitk.pdf>(last
923 access: 5 Sep, 2022), 2016.

924 Sindhwani, R., Goyal, P., Kumar, S., Kumar, A.: Anthropogenic Emission Inventory of Criteria
925 Air Pollutants of an Urban Agglomeration - National Capital Region (NCR), Delhi,
926 *Aerosol and Air Quality Research*, 15, 1681–1697,
927 <https://doi.org/10.4209/aaqr.2014.11.0271>, 2015.

928 Singh, V., Sahu, S. K., Kesarkar, A. P., Biswal, A.: Estimation of high resolution emissions from
929 road transport sector in a megacity Delhi, *Urban Climate*, 26, 109–120,
930 <https://doi.org/10.1016/j.uclim.2018.08.011>, 2018.

931 Stanaway, J.D. et al.: Global, regional, and national comparative risk assessment of 84
932 behavioural, environmental and occupational, and metabolic risks or clusters of risks for
933 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of
934 Disease Study 2017. The Lancet, 392, 1923-1994, [https://doi.org/10.1016/S0140-6736\(18\)32225-6](https://doi.org/10.1016/S0140-6736(18)32225-6), 2018.

936 TERI (The Energy and Resources Institute) and ARAI (The Automotive Research Association of
937 India): Source Apportionment of PM2.5 & PM10 of Delhi NCR for Identification of
938 Major Sources, https://www.teriin.org/sites/default/files/2018-08/Report_SA_AQM-Delhi-NCR_0.pdf, (last access: 17 Feb, 2022) 2018.

940 Transport Department, Govt. of Delhi, <https://transport.delhi.gov.in/home/transport-department>
941 (last access: 15Feb, 2023), 2019.

942 UNEP (United Nations Environment Programme): Air Pollution in Asia and the Pacific:
943 Science-based solutions, ISBN: 978-92-807-3725-7,
944 <https://wedocs.unep.org/20.500.11822/26861>, (last access: 8 Feb, 2022), 2019.

945 World Air Quality Report: Region & City PM2.5 Ranking, <https://www.iqair.com/world-most-polluted-cities/world-air-quality-report-2019-en.pdf>, (last access: 8 Feb, 2022), 2019.

947 World Air Quality Report: Region & City PM2.5 Ranking, <https://www.iqair.com/world-most-polluted-cities/world-air-quality-report-2020-en.pdf>, (last access: 10 Feb, 2022), 2020.