

Providing quality-assessed and standardised soil data to support global mapping and modelling (WoSIS snapshot 2023)

Niels H. Batjes¹, Luis Calisto¹, Luis Moreira de Sousa¹

¹ISRIC – World Soil Information, Droevendaalsesteeg 3, 6708 PB Wageningen, The Netherlands

5 *Correspondence:* Niels H. Batjes (niels.batjes@isric.org)

Abstract. Snapshots derived from the World Soil Information Service (WoSIS) are served freely to the international community. These static datasets provide quality-assessed and standardised soil profile data that can be used to support digital soil mapping and environmental applications at broad scale levels. Since the release of the preceding snapshot in 2019, refactored ETL (Extract, Load, Transform) procedures for screening, ingesting and standardising disparate source data have been developed. In conjunction with this, the WoSIS data model was overhauled making it compatible with the ISO 28258 and Observations and Measurements (O&M) domain models. Additional procedures for querying, serving, and downloading the publicly available standardised data have been implemented using open software (e.g. GraphQL API). Following up on a short discussion of these methodological developments we discuss the structure and content of the “WoSIS 2023-snapshot”. A range of new soil datasets was shared with us, registered in the ISRIC World Data Centre for Soils (WDC-Soils) data repository, and subsequently processed in accordance with the licences specified by the data providers. An important effort has been the processing of forest soil data collated in the framework of the EU-HoliSoils project. We paid special attention to the standardisation of soil property definitions, description of the soil analytical procedures, and standardisation of the units of measurement. The “2023 snapshot” considers the following soil chemical properties (total carbon, organic carbon, inorganic carbon (total carbonate equivalent), total nitrogen, phosphorus (extractable-P, total-P, and P-retention), soil pH, cation exchange capacity, and electrical conductivity) and physical properties (soil texture (sand, silt, and clay), bulk density, coarse fragments, and water retention), grouped according to analytical procedures that are operationally comparable. Method options are defined for each analytical procedure (e.g. pH measured in water, KCl or CaCl₂ solution, molarity of the solution, and soil/solution ratio). For each profile we also provide the original soil classification (i.e. FAO, WRB and USDA system with their version) and pedological horizon designations as far as these have been specified in the source databases. Three measures for “fitness-for-intended-use” are provided to facilitate informed data use: a) positional uncertainty of the profile’s site location, b) possible uncertainty associated with the operationally defined analytical procedures, and c) date of sampling. The most recent (i.e. *dynamic*) dataset, called *wosis_latest*, is freely accessible via various webservices. To permit consistent referencing and citation we also provide a *static* snapshot (in casu, December 2023). This snapshot comprises quality-assessed and standardised data for 228k geo-referenced profiles. The data come from 174 countries and represent more than 900k soil layers (or horizons) and over 6 million records. The number of measurements for each soil

property varies (greatly) between profiles and with depth, this generally depending on the objectives of the initial soil sampling programmes. In the coming years, we aim to gradually fill gaps in the geographic distribution of the profiles, as well as in the soil observations themselves, this subject to the sharing of a wider selection of “public” soil data by prospective data contributors; possible solutions for this are discussed. The WoSIS 2023-snapshot is archived and freely available at
5 <https://doi.org/10.17027/isric-wdcsoils-20231130> (Calisto et al., 2023).

1 Introduction

The World Soil Information service (WoSIS) draws on a large complement of soil profile data that have been shared by numerous data providers. Nonetheless, a large proportion of the 800k “so-called” freely available soil profiles (see Arrouays et al., 2017),
10 in practice, still remain “inaccessible” due to various licence constraints (e.g. Cornu et al., 2023). Soil data submitted for consideration in WoSIS come from a wide range of legacy holdings (e.g. traditional soil surveys), and increasingly include data derived from proximal sensing (e.g. Shepherd et al., 2022; Viscarra Rossel et al., 2016). The source data come in various formats and were determined according to a range of field sampling and soil analytical procedures, requiring standardisation and harmonisation during their ingestion/processing into WoSIS.

15 Prior to discussing the “2023 snapshot”, we provide a short retrospective of activities that lead to the development of WoSIS. In the early days of desktop computers, ISRIC with its partners compiled a range of project-specific databases such as ISIS (van de Ven and Tempel, 1994), created to manage data for the ISRIC World Soil Reference Collection, several national and continental scale Soil and Terrain (SOTER) databases (e.g. FAO and ISRIC, 2003; FAO et al., 2007; FAO et al., 1998), the WISE database (Batjes, 1997; Batjes and Bridges, 1994), and the Africa Soil Profile (AfSP) database (Leenaars et al., 2014). While
20 these different databases were structured along the general principles and criteria of the FAO Guidelines for Soil Description (FAO, 1977; 2006) and USDA Soil Survey Manual (Soil Survey Division Staff, 1993), the ISIS, SOTER, WISE, and AfSP databases each had their own data models and conventions. Further, out of necessity at the time, the databases were developed and implemented on stand-alone computers using a range of commercial software packages. In 2009, ISRIC management decided to bring the above stand-alone products together in a centralised enterprise database, known as WoSIS (World Soil Information
25 Service), developed using PostgreSQL with the PostGIS extension for handling spatial data. After the initial ingest and standardisation of the above “ISRIC holdings” the service was to be expanded with datasets shared by a diverse range of soil data providers.

The original aim of WoSIS was to accommodate any type of soil data (profile, vector and grid) (Ribeiro et al., 2015; Tempel et al., 2013). However, from 2015 onwards, in view of technical considerations and institutional developments, the scope of
30 WoSIS was changed to “safeguarding, processing, standardizing and serving geo-referenced soil profile (point) data for the

world” (Ribeiro et al., 2020). Alternatively, vector and grid maps derived from traditional soil mapping (e.g., Batjes, 2016; Dijkshoorn et al., 2005; FAO et al., 2012; van Engelen et al., 2006) and digital soil mapping (e.g., Hengl et al., 2017; Poggio et al., 2021; Turek et al., 2022) would be managed and served through other components of our spatial data infrastructure, such as the ISRIC data hub (<https://data.isric.org>) and the SoilGrids/WoSIS portal (<https://soilgrids.org>; last access: 24 April 2024). All these web services were developed using free-and-open-software (FOSS).

The ultimate goal of WoSIS, like for related global data compilation activities (Baritz et al., 2017; de Sousa et al., 2019), is full data harmonisation (Batjes et al., 2020; Ribeiro et al., 2015; Ribeiro et al., 2020). According to the Global Soil Partnership (GSP, Baritz et al., 2014), harmonisation involves “providing mechanisms for the collation, analysis and exchange of consistent and comparable global soil data and information” and considers the following domains: a) soil description, classification and mapping, b) soil analyses, c) exchange of digital soil data, and d) interpretations. In view of the breadth and magnitude of the task, as well as the limited availability of comparative “multiple analytical procedures” data sets as required for full harmonisation (Batjes, 2023; Bispo et al., 2021; van Leeuwen et al., 2022) we have limited ourselves to the standardisation of soil property definitions, soil analytical procedures descriptions, plausibility checks for soil observation values and the standardisation of measurement units for commonly required soil properties (see Appendix A). Importantly, users should always keep in mind that the source datasets themselves (e.g., Armas et al., 2023; NPDB, 2023; USDA-NCSS, 2021) will provide more detailed information than WoSIS albeit not in a consistent, globally standardised format.

This paper discusses methodological changes to the WoSIS workflow and new data additions since the release of the previous snapshot (Batjes et al., 2020). First, we describe the new data model, the refactored data screening/ingestion process and indicate how the “shared” data are being served to the user community upon their standardisation. Thereafter, we describe the actual data screening, quality control and standardisation process. Subsequently, we describe the spatial distribution of soil profile sites and list the number of soil observations represented in the “WoSIS 2023-snapshot” (hereafter referred to as 2023-snapshot). In conjunction with this, we provide three measures for “fitness-for-intended-use” of the standardised data and discuss possible limitations of the snapshot. Finally, following up on a discussion concerning the scope for “full data harmonisation” in WoSIS, future developments and possible constraints arising are outlined.

The naming conventions and standard units of measurement are listed in Appendix A, while the structure of the snapshot files is described in Appendix B. In Appendix C we list the number of sites by country and continent (Table C1) as well as their distribution by world terrestrial ecosystems (Table C2) and biomes (Table C3).

Soils are important providers of ecosystem services (FAO and ITPS, 2015). WoSIS-served data have been used for a range of applications, such as predictive soil property mapping (Guevara et al., 2018; Moulatlet et al., 2017; Nenkam et al., 2022; Poggio et al., 2021; Turek et al., 2023), space and time modelling of soil organic carbon stock change (Heuvelink et al., 2021), and a diverse range of environmental assessments (e.g., Hassani et al., 2024; Huang et al., 2024; Luo et al., 2021; Lutz et al., 2019; Maire et al., 2015; Sanderman et al., 2017; Sothe et al., 2022). For example, based on the “2016 snapshot” and “2019

snapshot” respectively, Ivushkin et al. (2019) mapped global soil salinity change, while Wang et al. (2024) analysed responses of soil organic carbon under warming across global biomes. Ultimately, such information can help to inform the global conventions such as the UNCCD (United Nations Convention to Combat Desertification) and UNFCCC (United Nations Framework Convention on Climate Change), so that policymakers and business leaders can make informed decisions about the environment, biodiversity, and human well-being at an appropriate scale level.

2 WoSIS data model and workflow

2.1 Workflow

The data model and workflow for acquiring, ingesting, processing and serving data as described in Ribeiro *et al.* (2020) was overhauled. This proved necessary as this procedure was essentially designed as a series of dataset-specific python and SQL scripts, which was adequate as long as WoSIS was still relatively small. However, in view of the rapidly growing population of “shared” soil data and overall complexity of the data model itself it proved necessary to implement a new, state-of-the-art ISO domain model (de Sousa, 2023; de Sousa et al., 2023), with re-factored ETL (extract, transform and load) procedures, to ultimately better serve our diverse user community in our capacity as World Data Centre for Soils (WDC-Soils).

The main stages of the new workflow are visualised in Fig. 1: a) Data providers share their data with ISRIC WDC-Soils, b) the submitted data sets with associated metadata are screened for “completeness of information provided” (e.g. the licence defining access rights and description of terms and units) and, once considered adequate, subsequently stored “as is” in the WDC-Soils data repository (see “ISRIC Admin” in Fig. 1); and c) the source datasets are imported into the new WoSIS PostgreSQL relational database (see Sect. 2.2), using re-factored ETL procedures (see Sect. 2.3). Step c) includes: c1) basic data quality assessment and control, c2) standardising descriptions for the soil analytical procedures and units of measurement, and c3) automated checks against plausibility limits for each soil observation, see Sect. 3 for details. Subsequently, d) distribution of the quality-assessed and standardised data via various services such as dashboards and WFS (OpenGIS web feature service) as well as a metadata catalogue service.




Figure 1. Schematic WoSIS workflow for ingesting, standardising and distributing soil profile data.

5 2.2 Data model

As indicated earlier, a new data model for WoSIS was developed, aligned where possible with the ISO 28258 domain model (de Sousa et al., 2023) and the GloSIS web ontology (Palma et al., 2024), both stemming from O&M (Observations and Measurement, Cox and David, 2011), all the while preserving legacy data. Main features of interest are the dataset (describes source of data), site (geo-spatial location where a soil investigation took place) and profile (sequence of pedo-genetic horizons along the depth of the profile). The key modification vis à vis the previous data model (Ribeiro et al., 2020) is the conditioning of analytical methods to the observation (see <https://git.wur.nl/isric/databases/wosis-docs>, last access: 26 April 2024). Changes made to the database schema and data over time are tracked using a migration tool (<https://github.com/graphile/migrate>, last access 24 April 2024). It maintains a record of the history, state, and dependencies of the database, including the conversion to the new data model.

Special attention was paid to the succinct description of the analytical procedures (see c2 above) using seven database tables, as summarised below:

- `thes_method_value`: Thesaurus of values that match the keys used to define an analytical method. For example, "natural clod" for the "sample type" key for method "bulk density".
 - `thes_method_key`: Thesaurus of keys used to define an analytical method. For example, "reported pH", "exchange solution" and "index cation".
- 5
- `thes_method_option`: Encodes the possible combinations of key-value pairs for each numerical observation on layers. Note that only a small sub-set of observations can be associated with particular method options.
 - `method_source`: Analytical methods descriptions as defined in the respective source databases (i.e. prior to standardisation). This table was imported "as is" from the old data model (Ribeiro et al., 2020) with the addition of a synthetic primary key. The records in this table remain essential to identify the method referred by each result.
- 10
- `method_standard`: Distinguishes each source method by the particular observation to which it applies. It can be regarded as a standardised description of the source method. Each record corresponds to a collection of key-value pairs in the `method_option` table for a single observation. Results for numerical observations reference this table to identify the corresponding analytical method.
 - `method_option_standard`: Defines a many-to-many relationship between the `method_standard` and `method_option` tables. It determines the exact collection of key-value option pairs that constitute a standard method. The standard method is a specialisation of the source method for a specific observation.
- 15

2.3 ETL procedure

Extract, Transform and Load (ETL) is a standardised, semi-automatic process that guides the data processor during the ingestion of new datasets. Re-factored ETL procedures were developed to align with the structure of the ISO data model. During the initial phase, newly shared datasets are submitted to a quick consistency check (i.e. format, data model, metadata and licence after which they are uploaded "as is" to a staging area in the WoSIS system. Subsequently, during the transform stage the uploaded datasets are parsed by the system. During this process validation and standardisation occurs (see Sect. 3.3 for details). In case of (possible) unconformities the system will generate descriptive messages that guide the data processor towards possible actions that may be needed to resolve the flagged unconformities. The data processor then needs to correct these issues in conformity with the requirements of the WoSIS procedure manual in steps guided by the system; in some cases the original data providers may need to be consulted. At the end of this phase, the cleaned and standardised data remain in the staging area for final verification by a

soil expert. After this verification, the final stage of the ETL process, “Load”, can start. This is a fully automated process during which the cleaned and standardised data are copied into the WoSIS database and subsequently removed from the staging area (note: the source data themselves are permanently preserved in the ISRIC data repository). The newly ingested data can now be used to create a range of WoSIS-derived products (e.g., *wosis_latest*, *wosis_internal*, and dashboards, see Fig.11) in accord with the licences and possible restrictions specified by the data providers.

2.4 Operational definitions

Soil characteristics, such as texture, bulk density and organic carbon content, are collated according to a wide range of procedures in different countries. For such incongruent data to be interpreted correctly during the ETL process, the procedures for their collection, analysis, and reporting need to be well documented and understood. Results can differ when different analytical procedures are used even though these procedures may carry the same name (e.g. clay, silt and sand size fraction) or concept (see Soil Survey Staff, 2011). This makes the inter-comparison of different datasets difficult if it is not known how these data were collected/analysed. Therefore we use “operational definitions”, as defined by USDA Soil Survey Staff (2011), for soil properties that are linked to specific analytical procedures. To properly characterise the “pH of a soil”, for example, we need information on sample pre-treatment, soil/solution ratio, and description of solution (e.g. H₂O, 1 M KCl, 0.02 M CaCl₂, or 1 M NaF). Soil pH measured in Sodium Fluoride (pH NaF), for example, provides a measure for the phosphorus (P) retention of a soil whereas pH measured in water (pH H₂O) is an indicator for soil nutrient status. Consequently, in WoSIS soil properties are named according to and defined by the analytical procedures and corresponding “method options”, based on common practice in soil science (e.g. BDFIOD for “bulk density (BD), fine earth fraction (FI), oven dry (OD)”). The current list of soil properties standardised in WoSIS is described in Sect. 3.3.

2.5 Data provisioning

Upon completion of the semi-automated ETL process, the quality-assessed and standardised data are distributed freely through various channels (see Fig. 1) in accordance with the license agreements (see Sect. 2.6):

- As *wosis_latest* (dynamic) via WFS; the respective endpoints are catalogued at the ISRIC Data Hub. (https://data.isric.org/geonetwork/srv/eng/catalog.search#/search?any=wosis_latest, last access: 26 April 2024)
- As “fixed” snapshots (in TSV format) with a unique digital object identifier (DOI) to permit consistent citation (https://data.isric.org/geonetwork/srv/eng/catalog.search#/search?any=wosis_snapshot, last access:26 April 2024).
- The contents of *wosis_latest* can also be visualised using a dashboard with some querying and zooming facilities (https://dashboards.isric.org/superset/dashboard/wosis_latest, last access: 26 April 2024).

- Profile data from *wosis_latest* can also be queried through the “SoilGrids web platform” (<https://soilgrids.org/>, last access: 26 April 2024), which also provides access to a range of soil property maps derived from the WoSIS-served profile data and a set of environmental covariates using digital soil mapping (Poggio et al., 2021; Turek et al., 2023).
- The *wosis_latest* holdings can also be queried using a GraphQL interface (<https://graphql.isric.org/>, last access: 26 April 2024) that facilitates exploration of the data (e.g. select data for organic carbon, bulk density, proportion of coarse fragments per layer (horizon) for profiles located in a given geography). Results of such tailor-made queries can then be exported as input in scripting languages such as Python or R (R Core Team, 2021), for example to calculate regional carbon stocks.

2.6 Licence agreements

10 It is not a simple task to find potential providers of “open” soil data (Arrouays et al., 2017; Batjes, 2009; Cornu et al., 2023). This may be due to technical issues, access arrangements, reasons for sharing (e.g., “Why share the data and for what purpose? What is in it for us?”), as well as legal requirements (Bispo et al., 2021; Robinson et al., 2019). All data sets that are shared with our centre are first registered in the ISRIC Data Repository together with their metadata; data sharing agreements should align with the ISRIC Data Policy (ISRIC, 2016). During the subsequent WoSIS standardisation workflow, we are faced with three different
15 types of datasets. First, those with a non-restrictive Creative Commons (CC-BY) licence, defined here as at least a CC-BY (Attribution) or CC-BY-NC (Attribution Non-Commercial) licence (these are later served as *wosis_latest*). Second, datasets with a more “restrictive” licence in the sense that they can exclusively be used for “visualisations”, such as SoilGrids™ (i.e. *wosis_internal*, see Fig. 1), by ISRIC itself. The latter, generally because the coordinates cannot be disclosed as stipulated by certain data providers (for details see <https://www.isric.org/explore/wosis/wosis-contributing-institutions-and-experts>, last
20 access: 26 April 2024). Finally, several data sets have licences that stipulate that they should only be safeguarded in the ISRIC repository and cannot be used for any data processing (i.e. permanent embargo).

The number of profiles in WoSIS per licence category, i.e. “public” respectively “restricted”, can be viewed and filtered using a dashboard (https://dashboards.isric.org/superset/dashboard/wosis_licenses/, last access: 26 April 2024). As shown in Table 1, the number of “public access” profiles served from WoSIS as snapshots increased from 96k in 2016 to 228k in 2023. Conversely,
25 it should be noted here that a large proportion of the forest soil data “shared” in the framework of the EU-Holisoiils project, for instance, could not be included in the “2023 snapshot” due to licence restrictions specified by the data providers. As a result, only 34k out of the total of 107k profiles “shared” with ISRIC between 2019 and 2023 could actually be included in the 2023-snapshot (resp. *wosis_latest*).

Table 1. Number of soil profiles and properties served in successive WoSIS snapshots.

Snapshot	No. of profiles	No. of properties ^a
2016-07	96k	22
2019-09	196k	45
2023-12	228k	45

^aProperty names are based on “operational definitions”, i.e. a combination of a property and procedure in the terminology of the WoSIS data model (see Sect. 2.4 and Sect. 3.3).

5 3 Data screening, quality control and standardisation

3.1 Consistency checks

Soil profile data shared for possible consideration in WoSIS were sampled and analysed according to various national or international standards and presented in various formats (from paper to digital). They are of varying degree of completeness as discussed below. To be considered in the WoSIS standardisation workflow (Fig. 1), each soil profile must meet several criteria as described earlier in Batjes et al (2020, p. 301). Summarising, they must be associated to a site correctly geo-referenced, have consistently defined upper and lower depths for each layer (or pedogenetic horizon), and have observations for at least some of the soil properties that are being served (e.g. sand, silt, clay and pH) as well as a succinct description of the analytical procedures and units of measurement. A soil (taxonomic) classification is considered desirable though not mandatory. Profiles associated to a valid site, for which only the classification is specified in the source data can still be useful for mapping of soil taxonomic classes.

Consistency in layer depth (i.e. sequential increase of the upper and lower depth reported for each layer down the profile) is checked using automated procedures (see Sect. 3.2). In line with current internationally accepted conventions, such depth increments are given as “measured from the soil surface, including organic layers and mineral covers” (FAO, 2006; IUSS Working Group WRB, 2022; Schoeneberger et al., 2012; Soil Survey Staff, 2022b). Until 1993, however, the begin (zero datum) of the profile was set at the top of the mineral surface (the *solum* proper), except for “thick” organic layers as defined for peat soils (FAO, 1977; 1990). Organic horizons were recorded as above and mineral horizons recorded as below, relative to the mineral

surface (Schoeneberger et al., 2012, p. 2-6). As far as possible, such “organic_surface” layers are flagged in the snapshot (see Appendix B) so that they may be filtered-out during auxiliary computations of soil organic carbon stocks, for example.

3.2 Screening for duplicate profiles

5 In the early stage of WoSIS, many source databases were compilations of shared soil profile data necessitating intricate procedures for identifying and flagging possibly repeated profiles (see Batjes et al., 2017; Ribeiro et al., 2020). Soil profiles located within 100 m of each other are flagged as possible duplicates, provided the year of sampling is identical (this criterion allows for reporting results of soil monitoring campaigns at the same *site*). Upon additional automated checks concerning the thickness of the first three soil layers (i.e. upper and lower depth), sand, silt and clay content, the duplicate profiles with the least-comprehensive component of observations are flagged and excluded from further processing (i.e. distribution). When still in doubt after these rigorous tests a final visual “similarity check” is made with respect to other commonly reported soil properties such as pH_{water} and organic carbon content, possibly leading to the flagging (exclusion) of some additional profiles.

3.3 Standardisation of property names, analytical procedure descriptions and units of measurement

15 A crucial step during data ingestion is the standardisation of the, regularly non-English, soil property names used in the source databases to the WoSIS conventions, as well as the standardisation of the soil analytical procedures according to consistent “operational definitions” (see Appendix A). Subsequently, the units of measurement are standardised, and the reported measurement values assessed according to soil observation-specific plausibility ranges for the respective soil properties (i.e. likely minimum and maximum). Some of these plausibility limits may change when more data become available for so far under-
20 represented soil observations, similar to ICP Forests (2020, p. 25), and appropriate PostgreSQL “trigger mechanisms” have been implemented for this. Data that do not meet these conditions are flagged and not processed further in the ETL workflow (see above), unless the observed “inconsistencies” can easily be solved (e.g. blatant typos in pH values). Alternatively, the data provider(s) may be contacted to resolve the observed errors.

Similar to the 2019-snapshot, the following soil properties are considered in the 2023-snapshot:

- 25
- Chemical: total carbon (i.e. organic plus inorganic carbon), organic carbon, inorganic carbon (i.e. total carbonate equivalent), total nitrogen, soil pH, cation exchange capacity, electrical conductivity, and phosphorus (extractable-P, total-P, and P-retention),
 - Physical: Soil texture (clay, silt, sand), coarse fragments, bulk density, and water retention.

All measurement values are served as recorded in the source data, after the above consistency checks and standardisation of the units of measurement to the target units (see Appendix A). As such, we *do not* apply any “gap filling” procedures during ETL nor do we apply any pedotransfer functions (PTF) to derive missing bulk density data or soil hydrological properties, respectively harmonise particle class size limits to a common standard, for example. This follow up stage of data processing is seen as the task of the data users (modellers) themselves. In practice, the required PTFs or ways for depth-aggregating the layer data will be determined by the projected use(s) of the standardised data (see Finke, 2006; Heuvelink et al., 2021; Poggio et al., 2021; Turek et al., 2023; van Leeuwen et al., 2024; Van Looy et al., 2017). It should be noted, however, that inadvertently some PTF-derived values (e.g. for bulk density) could have slipped through the above consistency checks in situations where procedures were mis-coded in the metadata of a source data set; critical modellers should exclude such values during their analyses.

10

3.4 Providing measures for fitness-for-intended-use

As indicated earlier, data served from WoSIS are used for a wide range of environmental applications(e.g., Guevara et al., 2018; Heuvelink et al., 2021; Luo et al., 2021; Maire et al., 2015 ; Moulatlet et al., 2017; Poggio et al., 2021; Sanderman et al., 2017; Sothe et al., 2022; Turek et al., 2023), but many of these assessments do not explicitly consider the uncertainties that are associated with the data. However, it is well known that “soil observations used for calibration and interpolation are themselves not error-free” (e.g., Baroni et al., 2017; Cressie and Kornak, 2003; Folberth et al., 2016; Grimm and Behrens, 2010; Guevara et al., 2018; Heuvelink, 2014; van Leeuwen et al., 2022). Therefore, since 2019, we provide three measures for “fitness-for-intended-use” in *wosis_latest* namely: a) positional uncertainty of the profiles (i.e. site location), b) inferred accuracy of the laboratory measurements, and c) date of sampling. These three measures, although approximative, should be duly considered in digital soil mapping and subsequent earth system modelling as they can affect the prediction uncertainty and “area-of-applicability” of the resulting derived products (Dai et al., 2019; Meyer and Pebesma, 2021; Shi et al., 2023). For example, large areas of the globe are still poorly represented in WoSIS (basically the yellow areas in Fig. 3). As indicated earlier, this issue can only be remedied when a larger selection of datasets is shared by the international soil community for consideration in WoSIS.

15

20

Importantly, prospective data users should also realise that the point/profile data shared for consideration in WoSIS are largely based on purposive sampling. During such “traditional” surveys, soil surveyors identify sample locations based on their knowledge of the survey area, desired level of detail (scale) and objective of the survey, for example detailed or exploratory surveys (FAO, 2006; IUSS Working Group WRB, 2022; Soil Survey Staff, 2017). Hence, such “legacy” data are not based on a probabilistic sampling scheme as recommended for digital soil mapping (Brus et al., 2011; Brus, 2022; Cramer et al., 2019; Heuvelink et al., 2007).

30

3.4.1 Positional uncertainty

Profiles in WoSIS are georeferenced through the *site* in which they were sampled in accord with ISO 28258 standards (de Sousa et al., 2023). The coordinates themselves are presented according to the World Geodetic System datum ensemble (i.e. WGS84, EPSG code 4326) upon their conversion from a diverse range of national projections. For most profiles (86 %, see Table 2) the approximate positional uncertainty of the profile locations, as inferred from the coordinates given in the source datasets, is ~100 m. Typically, geo-referencing before the advent of GPS (Global Positioning Systems) in the 1970s is less accurate; often we just do not know the “true” accuracy. Nonetheless, digital soil mappers should be aware of this issue (Grimm and Behrens, 2010), because the soil observations and environmental covariates may not actually overlap (Cressie and Kornak, 2003), this both in space and time.

Table 2. Positional uncertainty of profile site locations.

Positional uncertainty	Number of profiles	
	n	%
~ 100 m	195,554	86
100 m - 1 km	21,653	9
1 km – 10 km	3,846	2
Over 10 km	7,037	3

3.4.2 Measurement uncertainty

Soil data managed in WoSIS have been analysed according to a diverse range of analytical procedures in multiple laboratories. A measure for measurement uncertainty is thus desired. Soil laboratory-specific Quality Management Systems and laboratory proficiency-testing (PT) can provide this type of information (GLOSOLAN, 2023; Magnusson and Örnemark, 2014; Munzert et al., 2007; NATP, 2015; WEPAL, 2019). Calculation of laboratory-specific measurement uncertainty for a single procedure, respectively multiple analytical procedures, will require several measurement rounds (years of observation) and solid statistical

analyses (van Leeuwen et al., 2022). Generally, however, this type of information is not provided with the source data sets submitted to the ISRIC data repository. Therefore, pragmatically, we have distilled the required information from the PT-literature (Al-Shammary et al., 2018; ICP Forests, 2021a; Kalra and Maynard, 1991; Rayment and Lyons, 2011; Rossel and McBratney, 1998; van Reeuwijk, 1983; WEPAL, 2019), as far as technically feasible. In the case of organic carbon content, for example, the mean variability was 17 % (with a range of 12 to 42 %) and for “CEC buffered at pH 7” of 18 % (range 13 to 25%) when multiple laboratories analyse a standard set of reference materials using similar operational procedures (WEPAL, 2019).

The figures for measurement accuracy presented in Appendix A represent first approximations. They are derived from the inter-laboratory comparison of analyses on well-homogenised, reference samples for a still relatively small range of soil types. These indicatory figures should be refined, for example using probability distribution functions (Heuvelink et al., 2007; van Leeuwen et al., 2022), once sufficient laboratory and procedure-related accuracy (i.e. systematic and random error) information is provided with the shared soil data. (Magnusson and Örnemark, 2014) Alternatively, this type of information may be collated in the context of international laboratory PT-networks such as GLOSOLAN and WEPAL, and in the framework of the ongoing LUCAS topsoil monitoring round (Bispo et al., 2021; Cornu et al., 2023). Meanwhile, the present first estimates can already be considered when calculating the uncertainty of predictive digital soil maps and of any interpretations derived from them (e.g. studies of soil organic carbon stock change).

Realistically, full harmonisation of analytical data derived from disparate sources, the ultimate ambition in WoSIS, will first become feasible once results of a representative set of multi-procedure, inter-laboratory comparison data sets become (freely) available, as discussed by Baritz *et al.* (2014), Bispo *et al.* (2021) and Batjes (2023), and a common set of reference Standard Operating Procedures (SOPs) has been accepted as a global standard.

20

3.4.3 Year of sampling

For each profile site, the date of sampling has been recorded as far as documented in the source data. This information is important to consider when superimposing the profile data with environmental co-variates, such as land cover, for example in the context of space and time analyses (Giller et al., 2006; Heuvelink et al., 2021). Most (54%) profiles represented in the snapshot were described/sampled between 1980 and 2020 (Table 3), and less than 4% before 1960. Alternatively, the date of site description and sampling is not known for almost 27% of the profiles as the information was not provided in the source materials.

Table 3. Period of sampling/analysis.

Period	N of profiles	Percentage
< 1920	37	0
1920-1940	253	0.1
1940-1960	8,632	3.8
1960-1980	35,358	15.5
1980-2000	75,686	33.2
2000-2020	47,768	20.9
Not specified	60,356	26.5

4 Spatial distribution of soil profiles and number of observations

4.1 Spatial distribution

5 The 2023-snapshot includes standardised data for 228k profiles, sampled at 217k different sites (Fig. 2). The greatest number of profiles come from north America (35 %) followed by Oceania (19%) and Europe (17%), while there are still few profiles for Asia (3%) and Antarctica (Table 4). The profiles come from sites in 174 countries. The average density of observations varies greatly both between countries (Table C1) and within each country.

10 Changes in the spatial distribution and density of profiles (per 1000 km²) in the successive WoSIS snapshots (Fig. 3) reflect the degree to which our data acquisition efforts were successful, as further discussed in Sect. 6. Overall, the density of soil observations is still low for Central Asia, Southeast Asia, Central and Eastern Europe, Russia, and the northern circumpolar region in the 2023-snapshot.




Figure 2. Distribution of sites represented in the 2023 snapshot of WoSIS (Good homolosine equal-area projection).

The number of profiles by biome (Olson et al., 2001b) and broad climatic region (Sayre et al., 2014) respectively, as derived
 5 from GIS overlays, are listed in Table C2 and C3.

Table 4. Number of soil profiles per continent.

Continent	Number of profiles		
	2023-snapshot	2019-snapshot	2016-snapshot
Africa	32,198	27,688	17,153
Antarctica	35	9	0
Asia	7,763	6,704	3,089
Europe	39,728	35,311	1,908
North America	78,996	73,604	63,066

Oceania	43,013	42,918	235
South America	26,457	10,218	8,790




Figure 3. Density and spatial distribution of profiles served with the 2017, 2019 and 2023 WoSIS snapshots.

4.2 Number and depth of observations

In total, the profiles considered in the 2023-snapshot are described by 0.9 million soil layers (or horizons). This corresponds with over 6.1 million records that include both numeric (e.g. silt content, soil pH, and cation exchange capacity) as well as class (e.g. WRB soil classification and horizon designation) properties. There are more observations for the chemical properties than the physical properties (see Table A1). Further, the number of observations generally decreases with depth, this largely depending on the objectives of the original soil surveys. The interquartile range (Q1-Q3) for maximum depth of soil sampled in the field is 33-150 cm, with a median (Q2) of 100 cm (mean= 107 cm). It should be noted here that specific purpose surveys only consider the topsoil (e.g. soil fertility surveys), while others systematically sample soil layers up to depths exceeding 20 m (with a maximum of 32 m). When data from such “specific purpose surveys” (defined here as < 30 cm and >300 cm) are excluded, the figures for maximum depth sampled become: Q1= 90 cm, Q2= 122 cm, Q3= 155 cm with a mean of 126 cm.

Table 5 provides an overview of the maximum depth of soil sampled during the various surveys that underpin WoSIS, by continent. Unfortunately, we are not able to show the “depth to bedrock” as this information is seldom made explicit in the source databases.

15

Table 5. Maximum depth of soil sampled per continent.

Continent	Maximum depth sampled (cm)		
	<=30	30-60	>60
Africa	5307	1779	22458
Antarctica	6	7	17
Asia	635	505	4310
Europe	9411	2848	23195
Northern America	6190	6728	60698
Oceania	9216	3792	27839
South America	1730	810	8477

5 Distributing the standardised data

The standardised data are distributed through ISRIC’s Spatial Data Infrastructure (SDI). The SDI is based on open source technologies and open web-services (WFS, WMS, WCS, CSW) following Open Geospatial Consortium (OGC) standards and aimed specifically at handling soil data. Our metadata are organised following standards of the International Organization for Standardization (ISO-19139, 2019) using GeoNetwork (see <https://data.isric.org>). The WoSIS database is hosted in a PostgreSQL database, with the spatial extension PostGIS. The PostgreSQL database itself is connected to MapServer to permit data download from GeoNetwork. These processes are aimed at facilitating global data interoperability and citation in compliance with FAIR principles. The data should be “findable, accessible, interoperable, and reusable” (Wilkinson et al., 2016).

Static snapshots are given a unique DOI (digital object identifier) to permit consistent citation. The 2023-snapshot is distributed in tab-separated values format (see Appendix B for file structure) and as GeoPackage (<https://doi.org/10.17027/isric-wdcsoils-20231130>). An online Readme file, that includes links to two short tutorials, provides additional technical information (https://www.isric.org/sites/default/files/Readme_WoSIS_202312_v2.pdf). Alternatively, the evolving *dynamic* version of the standardised data (i.e. *wosis_latest*) can be accessed/queried through the ISRIC Data Hub (<https://data.isric.org>) and the SoilGrids platform (<https://soilgrids.org>). Tutorials describing how to access *wosis_latest* from QGIS using WFS and with GraphQL (Calisto, 2023) can be found on the ISRIC website (see <https://www.isric.org/explore/wosis/faq-wosis>, last access: 26 April 2024).

By its nature, the *dynamic* version will grow when new profile data are shared and processed, additional soil properties are considered in the WoSIS workflow, and/or when possible corrections are required. Potential errors can be reported via a “Google group” (<https://groups.google.com/forum/#!forum/isric-world-soil-information>, last access: 26 April 2024) so that these may be addressed in the *dynamic* version.

6 Discussion

We describe new procedures for handling and standardising disparate world soil profile data in WoSIS. The data model was fully harmonised to ISO 25828 and O&M requirements, with minor adjustments, and refactored ETL procedures were implemented. Alternatively, it should be stressed that the ultimate, desired full harmonisation of observations to an agreed reference analytical procedure Y, for example “pH H₂O, 1:2.5 soil/water solution” for say all “pH 1:x H₂O” measurements, will first become feasible once the target procedure (Y) for analysing each property has been defined, and subsequently accepted as “global standard” by the international soil community. A next step would be to collate/develop “comparative” data sets for each soil property (i.e. sets with samples analysed according to a given reference procedure (Y_i) and the corresponding national procedures (X_j)) for pedotransfer function development. These relationships, however, will often be soil type and region specific (GlobalSoilMap,

2015) and difficult to develop (i.e. calibrate and validate) when datasets for the comparisons do not yet exist or are simply not freely shared/available (Batjes, 2023; Bispo et al., 2021; Cornu et al., 2023; van Leeuwen et al., 2024). Hence the importance of regional laboratory inter-comparison programmes, such as those undertaken in the framework of for example ANSIS (2023), GLOSOLAN (2023), ICP Forests (2021a) and LUCAS (Bispo et al., 2021), that aim to develop consistent, context-specific (e.g. by country or land use/soil type) pedotransfer functions towards an agreed set of SOPs. However, it should be noted that the standard SOPs specified by these various programmes need not be comparable. In his context, Suvannang *et al.* (2018) observed that “comparable and useful soil information (at the global level) will only be attainable once laboratories agree to follow common standards and norms”. Over the years, however, many organisations/countries have implemented analytical procedures and quality assurance systems that are well suited for their specific purposes (e.g., ANSIS, 2023; Cornu et al., 2023; Orgiazzi et al., 2018; Soil Survey Staff, 2022a). Consequently, they may not be inclined to harmonise their data to a (still to be decided) set of global “reference” SOPs. However, agreed upon procedures for such a full scale harmonisation will be required when developing a globally federated, and ultimately interoperable, spatial soil data infrastructure (GLOSIS, de Sousa et al., 2021) through which (pre-harmonised) source data are served and updated by the respective data providers, and made queryable according to a common standard (de Sousa et al., 2023; OGC, 2019).

It is our intention to gradually fill gaps in the geographic distribution (Fig. 3) and range of soil properties (Appendix A) in the coming years. This work is part of ISRIC's remit as a regular member of the World Data System (<https://worlddatasystem.org>, last access: 26 April 2024). The degree to which this will be feasible, however, will largely depend on the willingness and ability of data providers to share (some of) their data for consideration in WoSIS. For the northern Boreal and Arctic region, for example, ISRIC can draw on new profiles collated by the International Soil Carbon Network (ISCN, see Malhotra et al., 2019). Alternatively, it should be reiterated that several datasets in our repository (e.g. ICP Forests, 2021a) can *only* be standardised and used for SoilGrids™ applications due to existing licence restrictions. Conversely, some countries such as New Zealand distribute their national soil profile dataset with a CC-BY-ND 4.0 licence, which implicitly precludes making any derivatives and hence their standardisation in WoSIS (see <https://viewer-nsdr.landcareresearch.co.nz/datasets/downloads/1042-2>, last access 10 June 2024).

Concerning the actual scope for expanding *wosis_latest* in the coming years, we noted that getting positive responses to our requests for sharing soil data is becoming increasingly cumbersome; the overall success rate during the “2019-2023” acquisition effort was around 25%. However, many of these datasets are being “shared” with ISRIC with the provision that the profile coordinates themselves may not be shown; hence, the corresponding soil data cannot be “openly” served to our user community through *wosis_latest*. Further, the site and profile coordinates are then regularly shared as “theoretical coordinates” only (e.g. ICP Forests, 2021b; Poeplau et al., 2020), highlighting the need for considering positional uncertainty in digital soil mapping and other applications. Another source of concern is that major soil monitoring programmes, such as LUCAS (e.g. Ballabio et al., 2016; Orgiazzi et al., 2018), only consider the top 20 or 30 cm of the soil. That is, they do not consider the actual soil profile depth as required for more comprehensive soil assessments such as computing changes in global carbon stocks or mapping plant-

available water holding capacity in the root zone (e.g. Batlle-Bayer et al., 2010; Leenaars et al., 2018; von Haden et al., 2020; Wang et al., 2022).

7 Data availability

The 2023-snapshot is archived for long-term storage at ISRIC – World Soil Information, the World Data Centre for Soils (WDC-
5 Soils) of the ISC (International Council for Science) World Data System (WDS). It is freely accessible at <https://doi.org/10.17027/isric-wdcsoils-20231130> (Calisto et al., 2023). The zip file (446 Gb) includes a copy of the “readme” file, the data in TSV format (see Appendix B) respectively in OGC GeoPackage format.

8 Conclusions

10 Bringing disparate soil profile data from different sources under a common global standard poses many and diverse challenges. A major improvement has been the harmonisation of the WoSIS data model to ISO 28258 and O&M domain specifications. In conjunction with this re-factored ETL procedures greatly improved the data ingestion and standardisation process, and new ways for visualising, querying and serving the data were developed to better serve our user community.

There are still numerous gaps in terms of geographic distribution as well as range of soil taxonomic units and/or soil properties
15 represented. We aspire to address such gaps in future updates of ‘*wosis_latest*’. However, as World Data Centre, we are largely dependent on the ability of soil data owners to share some of their data freely for the greater benefit of the international community. To facilitate and stimulate this process, we are developing a web-based facility (front-end) to permit data providers to directly upload their soil data to WoSIS in a consistent format based on the refactored ETL procedures. As an incentive, upon their standardisation, we aim to provide each data provider with a tailor-made dashboard for viewing and querying the datasets
20 they shared, possibly with a DOI to facilitate citation.

Various sources of uncertainty are associated with the data. Therefore, we provide three measures for “fitness-for-intended-use” of the standardised data. This information, although coarse, should be duly considered by prospective users of the snapshot.

Unfortunately, numerous soil datasets worldwide are not freely accessible for various reasons. Standardised procedures, mechanisms, policies and incentives aimed at encouraging soil data sharing by different categories of data owners/providers are
25 needed (e.g., Fantappie et al., 2021; Gobezie and Biswas, 2023; Padarian and McBratney, 2020; Robinson et al., 2019). At a transnational level, these pressing and complex issues are being addressed by the Global Soil Partnership, hosted by UN-FAO, in the context of the evolving federated Global Soil Information System.

Appendix A: Coding conventions

Table A1. Coding conventions for observations (i.e. a combination of property, procedure and unit of measurement), number of profiles and layers provided in the WoSIS 2023-snapshot, and inferred accuracy of measurements (Codes are listed in alphabetical order).

5

Code	Property	Procedure ^a	Unit	Profiles	Layers	Accuracy (± %) ^b
BDFI33	Bulk density fine earth ^c	Bulk density of a soil sample that has been desorbed to 33 kPa (1/3 bar)	kg/dm ³	14886	78007	25.0
BDFIAD	Bulk density fine earth	Bulk density of a soil sample that has been air dried	kg/dm ³	4238	14485	25.0
BDFIFM	Bulk density fine earth	Bulk density of a soil sample at field-soil water content at time of sampling	kg/dm ³	5265	14075	25.0
BDFIOD	Bulk density fine earth	Bulk density of a soil sample that has been dried in an oven at 110 °C	kg/dm ³	26064	131623	25.0
BDWSAD	Bulk density whole soil ^c	Bulk density of a soil sample that has been air dried	kg/dm ³	0	0	25.0
BDWSOD	Bulk density whole soil	Bulk density of a soil sample that has been dried in an oven at 110 °C	kg/dm ³	14596	75397	25.0
CECPH7	Cation exchange capacity	CEC estimated by buffering the soil at "pH7" (e.g., NH ₄ Oac)	cmol(c)/kg	60339	320532	20.0
CECPH8	Cation exchange capacity	CEC estimated by buffering the soil at "pH7" (e.g., NH ₄ Oac)	cmol(c)/kg	6838	25100	20.0
CFGR	Coarse fragments	Gravimetric content of soil material larger than 2 mm ^c	g/100g	39481	202414	20.0
CFVO	Coarse fragments	Volumetric content of soil material larger than 2 mm ^c	cm ³ /100cm ³	48891	246580	30.0
CLAY	Clay ^d	Determination of total gravimetric content of clay-size fraction (for class-size limits and analytical methods see 'method_options')	g/100g	153319	652347	15.0
ECEC	Cation exchange capacity	Effective CEC conventionally approximated by summation of exchangeable bases (Ca ²⁺ , Mg ²⁺ , K ⁺ , and Na ⁺) plus 1 M KCl exchangeable acidity (Al ³⁺ and H ⁺) in acidic soils	cmol(c)/kg	35123	143693	25.0
ELCO20	Electrical conductivity	Electrical conductivity assessed on a 1:2 soil water extract. Used for saline soils.	dS/m	7971	44350	10.0
ELCO25	Electrical conductivity	Electrical conductivity assessed on a 1:2.5 soil water extract. Used for saline soils.	dS/m	4395	17825	10.0
ELCO50	Electrical conductivity	Electrical conductivity assessed on a 1:5 soil water extract. Used for saline soils.	dS/m	23121	90959	10.0

ELCOSP	Electrical conductivity	Electrical conductivity assessed on water saturated soil paste. Used for saline soils.	dS/m	22052	85020	10.0
NITKJD	Total nitrogen (N)	Kjeldahl wet-oxidation digestion procedure	g/kg	72905	240433	10.0
ORGC	Organic carbon (C)	Amount of organic carbon determined according to method specified under 'method_options'	g/kg	135655	526953	15.0
ORGM	Organic matter	Determination of organic compounds that accompany soil particles through a 2-mm sieve using loss-on-ignition (LOI) at about 400 degrees Celsius.	g/kg	3871	16282	15.0
PHAQ	pH	A measure of the acidity or alkalinity in soils, defined as the negative logarithm (base 10) of the activity of hydronium ions (H ⁺) in water ^a .	unitless	140326	655336	0.3
PHCA	pH	A measure of the acidity or alkalinity in soils, defined as the negative logarithm (base 10) of the activity of hydronium ions (H ⁺), in the specified CaCl ₂ solution.	unitless	69437	325153	0.3
PHKC	pH	A measure of the acidity or alkalinity in soils, defined as the negative logarithm (base 10) of the activity of hydronium ions (H ⁺), in the specified KCl solution.	unitless	38022	173464	0.3
PHNF	pH	A measure of the acidity or alkalinity in soils, defined as the negative logarithm (base 10) of the activity of hydronium ions (H ⁺), in the specified NaF solution.	unitless	4965	25409	0.3
PHETB1	Phosphorus (P)	Phosphorus determined according to the Bray-I method, a combination of HCl and NH ₄ -F to remove easily acid soluble P forms, largely Al- and Fe-phosphates (mainly applicable for acid soils)	mg/kg	10719	40379	40.0
PHETM3	Phosphorus (P)	Determined according to Mehlich-3 method, a weak acid soil extraction procedure that is considered suitable for removing P and other elements in acid and neutral soil. The extract is composed of 0.2 M glacial acetic acid, 0.25 M ammonium nitrate, 0.015 M ammonium fluoride, 0.013 M nitric acid, and 0.001 M ethylene diamine tetraacetic acid (EDTA).	mg/kg	1444	7230	25.0
PHETOL	Phosphorus (P)	Phosphorus determined according to the Olsen method (0.5 M sodium bicarbonate (NaHCO ₃) solution at a pH of 8.5); used extract P from calcareous, alkaline, and neutral soils.	mg/kg	4266	12291	25.0
PHPRTN	Phosphorus (P)	Phosphorus retention measured according to the New Zealand method (Blakemore, 1981).	g/100g	5599	26569	20.0
PHPTOT	Phosphorus (P)	Phosphorus determined with a "harsh" digest procedure to liberate and measure all forms of element.	mg/kg	7561	19310	15.0
PHPWSL	Phosphorus (P)	Phosphorus soluble in soluble in water	mg/kg	282	1241	15.0

SAND	Sand	Determination of total gravimetric content of sand-size fraction (for class-size limits and analytical methods see 'method_options').	g/100g	119127	542463	15.0
SILT	Silt ^f	Determination of total gravimetric content of silt-size fraction (for class-size limits and analytical methods see 'method_options').	g/100g	145906	620790	15.0
TCEQ	Calcium carbonate equivalent (TCEQ)	Determination of the gravimetric loss of carbonates as carbon dioxide in the presence of excess hydrochloric acid. The quantity of carbonate (CO ₃) in the soil is expressed as CaCO ₃ and as a weight percentage of the less than 2 mm size fraction.	g/kg	59294	247368	10.0
TOTC	Total carbon (C)	Total C is quantified by two basic methods: wet or dry combustion (see 'method_options'). In total C determinations, all forms of C in a soil are converted to CO ₂ followed by a quantification of the evolved CO ₂ . Total C can be used to estimate the organic C content of a soil. The difference between total and inorganic C is an estimate of the organic C.	g/kg	33527	112787	10.0
WG0006	Water retention gravimetric	Water retention assessed at tension 6 kPa (see method options')	g/100g	827	3828	20.0
WG0010	Water retention gravimetric	Water retention assessed at tension 10 kPa (see 'method_options').	g/100g	2970	12517	20.0
WG0033	Water retention gravimetric	Water retention assessed at tension 33 kPa (see 'method_options').	g/100g	20994	94707	20.0
WG0100	Water retention gravimetric	Water retention assessed at tension 100 kPa (see 'method_options').	g/100g	687	3360	20.0
WG0200	Water retention gravimetric	Water retention assessed at tension 200 kPa (see 'method_options').	g/100g	4391	27773	20.0
WG0500	Water retention gravimetric	Water retention assessed at tension 500 kPa (see 'method_options').	g/100g	326	1414	20.0
WG1500	Water retention gravimetric	Water retention assessed at tension 1500 kPa (see 'method_options').	g/100g	33782	181999	20.0
WV0010	Water retention volumetric	Water retention assessed at tension 10 kPa (see 'method_options').	cm ³ /100cm ³	1914	6883	20.0
WV0033	Water retention volumetric	Water retention assessed at tension 33 kPa (see 'method_options').	cm ³ /100cm ³	7444	22291	20.0
WV0100	Water retention volumetric	Water retention assessed at tension 100 kPa (see 'method_options').	cm ³ /100cm ³	747	2553	20.0
WV0500	Water retention volumetric	Water retention assessed at tension 500 kPa (see 'method_options').	cm ³ /100cm ³	702	1758	20.0
WV1500	Water retention volumetric	Water retention assessed at tension 1500 kPa (see 'method_options').	cm ³ /100cm ³	7904	23331	20.0

- ^a Method options for each analytical procedure are described in Batjes and van Oostrum (2023), and provided in file *Wosis_202312_xxxx.tsv*, see Appendix C.
- ^b Inferred accuracy (or uncertainty), rounded to the nearest 5%, unless otherwise indicated (i.e. units for soil pH) as derived from the following sources (Al-Shammary et al., 2018; Kalra and Maynard, 1991; Rayment and Lyons, 2011; Rossel and McBratney, 1998; van Reeuwijk, 1983; WEPAL, 2019).
5 These figures are first approximations that should be fine-tuned once more specific results of laboratory proficiency tests, resp. national Soil Quality Management systems, become freely available (e.g. from the GLOSOLAN laboratory proficiency programme).
- ^c Generally, the fine earth fraction is defined as being < 2 mm. Alternatively, an upper limit of 1 mm was used in the former Soviet Union and its satellite states (Katchynsky scheme). The actual size limits are specified under “method options” (see Appendix C).
- ^d Provided only when the sum of clay, silt and sand fraction is ≥ 90 and ≤ 100 percent (Note that users should normalise the totals to 100 percent
10 before using them for mapping or modelling purposes; further, more stringent limits (e.g. ≥ 98 and ≤ 102) may be considered).
- ^e No data are being served for this property because the associated licences are flagged as ‘restricted’ by the data providers.
- ^f The lower and upper limits for the ‘silt’ size fraction can vary markedly between countries, hence these limits have been specified explicitly in WoSIS under “method options” (see Appendix B). Development and application of conversion procedures to one common “silt” fraction (e.g. 0.002-0.05 mm) is beyond the remit of the WoSIS project itself. The necessary pedotransfer functions should be developed (and tested) prior to generating particle size class related soil
15 property maps for a given geography. Research in this direction is being undertaken by the SoilGrids team, based on the “best available” comparative datasets for calibration.

Table A2. Coding conventions and brief descriptions for soil classification, horizon designations and number of occurrences in the WoSIS 2023-snapshot.

Code	Description	Count
CSTX	Classification of the soil profile according to specified edition (year) of USDA Soil Taxonomy, at least at soil order level	31400
CWRB	Classification of the soil profile according to specified edition (year) of the World Reference Base for Soil Resources (WRB), at least at reference soil group level	39649
CFAO	Classification of the soil profile according to specified edition (year) of the FAO-Unesco Legend, at least at major group level	38792
HODS ^a	Horizon designations as provided in the source databases	80849 / 396522 ^b

5 ^a Where available, the “cleaned” (original) layer/horizon designation is provided for general information; these codes have not been standardised as they vary widely between different classification systems (Bridges, 1993; Gerasimova et al., 2013). When no horizon designations are provided in the source data bases, we have flagged all layers with an upper depth given as being negative (e.g. -10 to 0 cm that is using pre-1993 conventions (see Sect. 3.1) in the source databases as likely being a shallow “organic surface” layer above a mineral soil layer.

^b Number of profiles with horizon descriptions respectively total number of layers with horizon designations.

10

Appendix B: Structure of WoSIS 2023-snapshot

This Appendix describes the structure of the data files served with the WoSIS 2023-snapshot, namely
 15 *wosis_202312_observations.tsv*, *wosis_20312_site.tsv*, *wosis_202312_profiles.tsv*, *wosis_202312_layers.tsv*, and
wosis_202312_xxxx.tsv (where “xxxx” is the name of the observation). The data files are also distributed as OGC
 GeoPackage, which stores the files within an SQLite database. Technical details are provided in a Readme file
 (https://www.isric.org/sites/default/files/Readme_WoSIS_202312_v2.pdf).

20

- *wosis_202312_observations.tsv*: This file lists the four to six letter codes for each observation, whether the observation is for a site/profile or layer (horizon), the unit of measurement and the number of profiles respectively layers represented in the snapshot. It also provides the inferred accuracy for the laboratory measurements (see Appendix A).

	code	Code for the observation
5	property	Description of soil property
	procedure	Description of analytical procedure
	unit	Standard unit of measurement
	profiles	Number of profiles that have at least one measurement for the observation
	layers	Number of layers that have measurements for the observation
10	accuracy	Inferred accuracy of the laboratory measurements (First approximation, see Sect. 3.4.2)

- *Wosis_202312_site.tsv*: This file characterises the site location where profiles were sampled. The following field names are used:

15	site_id	Primary key
	latitude	Latitude in degrees (WGS84)
	longitude	Longitude in degrees (WGS84)
	positional_uncertainty	Positional uncertainty of the profile's site location, expressed in four classes (see Table 2)
20	country_name	Name of country where site is located
	region	Region in which site is located
	continent	Continent in which site is located

- 25 • *wosis_202312_profiles.tsv*: Presents the unique profile ID (i.e. primary key), site_id, source of the data, country ISO code and name, positional uncertainty, latitude and longitude (WGS 1984), maximum depth of soil described and sampled, as well as information on the soil classification system and edition. Depending on the soil classification system used, the number of fields will vary. For example, for the World Soil Reference Base (WRB) system the

options are publication year (i.e. version), reference_soil_group_code, reference_soil_group_name, and the name(s) of the prefix (primary) qualifier(s) respectively suffix (supplementary) qualifier(s). The terms principal qualifier and supplementary qualifier are used since 2015 (IUSS Working Group WRB, 2015; 2022); earlier WRB versions used prefix and suffix for this (e.g. IUSS Working Group WRB, 2006). Alternatively, for USDA Soil Taxonomy, the version (year), order, suborder, great group, and subgroup can be accommodated (Soil Survey Staff, 2014). The following field names are used:

	profile_id	Primary key
	profile_code	Code for the profile
	dataset_code	Identifier for source data set
5	site_id	Identifier for site where profile is located
	positional_uncertainty	Positional uncertainty of the profile's site location, expressed in four classes (see Table 2).
	country_name	Name of country where site is located.
	latitude	Latitude in degrees (WGS84)
15	longitude	Longitude in degrees (WGS84)
	wrb_reference_soil_group_code	Code for WRB group (in given version of WRB)
	wrb_reference_soil_group	Full name for reference soil group
	wrb_prefix_qualifiers	Name for prefix (i.e. for WRB1988)
	wrb_suffix_qualifiers	Name for suffix (i.e. for WRB1988)
20	wrb_principal_qualifiers	Name for principal qualifiers (i.e. for WRB 2015 and WRB 2022)
	wrb_supplementary_qualifiers	Name for supplementary qualifiers (i.e. for WRB 2015 and WRB 2022)
	wrb_publication_year	Version of World Reference Base for Soil Resources
	fao_major_group_code	Code for major group (in given version of the Legend),

	fao_major_group	Name of major group
	fao_soil_unit_code	Code for soil unit
	fao_soil_unit	Name of soil unit
	fao_publication_year	Version of FAO Legend (e.g. 1974 or 1988)
5	usda_order_name	Name of USDA Soil Taxonomy order
	usda_suborder	Name of USDA Soil Taxonomy suborder
	usda_great_group	Name of USDA Soil Taxonomy greatgroup
	usda_subgroup	Name of USDA Soil Taxonomy subgroup
	usda_publication_year	Version of USDA Soil Taxonomy
10		
	<ul style="list-style-type: none"> • <i>Wosis_202312_layers.tsv</i>. This file characterises the layers (or horizons) per profile: 	
	profile_id	Primary key
	layer_id	Sequential number for the layer (or horizon)
	profile_code	Code for the profile
15	site_id	Identifier for site where profile is located
	layer_name	Name of pedogenetic horizon (“as is”)
	upper_depth	Upper depth of layer
	lower_depth	Lower depth of layer
	layer_number	Sequential number for the layer (or horizon)
20	organic_surface	Flag for the presence of an organic layer above the mineral soil

	dataset_id	Abbreviation for source data set (e.g. WD-ISCN)
	licence	Licence for observation as indicated by the data provider (e.g. CC BY)
	<ul style="list-style-type: none"> • <i>Wosis_202312_xxxx.tsv</i>. For each observation (e.g. “xxxx” = “BDFIOD”), as defined under “code” in file <i>wosis_202312_observation.tsv</i>, the following are listed: 	
5	profile_id	Primary key
	layer_id	Primary key (number, sequential from top to bottom)
	profile_code	Code for given profile
	layer_name	Name of pedogenetic horizon (“as is”)
	upper_depth	Upper depth of layer
10	lower_depth	Lower depth of layer
	organic_surface	Indicates if there is an organic layer above the mineral surface
	value	Array listing all measurement values for observation “xxxx” for the given layer. In some cases, more than one observation is reported for a given horizon (layer) in the source, for example four values for TOTC: [1:5.4, 2:8.2, 3:6.3, 4:7.7] (see value_avg below)
15	method_options	Array listing the method options for each analytical procedure as distilled from the source data. The content of this array varies with the soil observation under consideration as described in the method option table for each analytical procedure. For example, in the case of electrical conductivity (ELCO), the method options include sample pretreatment (e.g. sieved over 2 mm size, solution (e.g. water), ratio (e.g. 1:5), and ratio base (e.g. weight /volume). For details see Batjes and van Oostrum (2023).
20	value_avg	Average, for above (it is recommended to use this value for “routine” modelling)

dataset_id	Abbreviation for source data set (e.g. WD-ISCN)
country_name	Name of country where site is located.
latitude	Latitude in degrees (WGS84)
longitude	Longitude in degrees (WGS84)
5 positional_uncertainty	Positional uncertainty of the profile's site location (see Table 2).
region	Region in which site is located
continent	Continent where the profile's site is located
date	Date the profile was described/sampled
licence	Licence for given data, as indicated by the data provider (i.e. CC BY or CC BY-NC)

10

Format: All fields in the above files are tab-delimited, with double quotation marks as text delimiters. File coding is according to the UTF-8 unicode transformation format.

Using the data: Tutorials for downloading and querying the data, using various platforms, are provided on the WoSIS FAQ webpage (<https://www.isric.org/explore/wosis/faq-wosis>, last access: 24 April 2024).

15

Appendix C: Distribution of sites

Table C1. Number of sites per continent and country.

Continent	Country	Country code	No. of sites	Area (km ²)	Site density (per 1000 km ²)	
Africa	Abyei		4	0	9943	0
	Algeria	DZ		10	2308647	0.004
	Angola	AO		1168	1246690	0.937
	Benin	BJ		743	115247	6.447
	Botswana	BW		994	578247	1.719
	British Indian Ocean Territory	IO		0	49	0
	Burkina Faso	BF		2023	273281	7.403
	Burundi	BI		36	26857	1.34
	Cameroon	CM		1417	465363	3.045
	Cape Verde	CV		0	4056	0
	Central African Republic	CF		88	619591	0.142
	Chad	TD		7	1265392	0.006
	Comoros	KM		0	1652	0
	Congo	CG		70	340599	0.206
	Côte d'Ivoire	CI		255	321762	0.793
	Democratic Republic of the Congo	CD		378	2329162	0.162
	Djibouti	DJ		0	21670	0
	Egypt	EG		26	982161	0.026
	Equatorial Guinea	GQ		0	27000	0
	Eritrea	ER		0	120763	0
	Ethiopia	ET		1712	1129314	1.516
	Gabon	GA		47	264022	0.178
	Gambia	GM		0	11203	0
	Ghana	GH		432	238842	1.809
	Guinea	GN		128	243023	0.527
	Guinea-Bissau	GW		15	30740	0.488
	Hala'ib triangle		10	0	17684	0
	Ilemi triangle		13	0	3179	0

Kenya	KE	1603	582342	2.753	
Lesotho	LS	33	30453	1.084	
Liberia	LR	50	96103	0.52	
Libya	LY	14	1620583	0.009	
Madagascar	MG	130	588834	0.221	
Malawi	MW	3050	118715	25.692	
Mali	ML	885	1251471	0.707	
Ma'tan al-Sarra		11	0	1993	0
Mauritania	MR	13	1038527	0.013	
Mauritius	MU	0	2014	0	
Mayotte	YT	0	378	0	
Morocco	MA	113	414030	0.273	
Mozambique	MZ	565	787305	0.718	
Namibia	NA	1569	823989	1.904	
Niger	NE	520	1182602	0.44	
Nigeria	NG	1402	908978	1.542	
Réunion	RE	0	2504	0	
Rwanda	RW	1016	25388	40.018	
Saint Helena, Ascension and Tristan da Cunha	SH	0	399	0	
Sao Tome and Principe	ST	0	991	0	
Senegal	SN	312	196200	1.59	
Seychelles	SC	0	499	0	
Sierra Leone	SL	12	72281	0.166	
Somalia	SO	245	632562	0.387	
South Africa	ZA	879	1220127	0.72	
South Sudan	SS	82	629821	0.13	
Sudan	SD	130	1843196	0.071	
Swaziland	SZ	14	17290	0.81	
Togo	TG	9	56767	0.159	
Tunisia	TN	60	155148	0.387	
Uganda	UG	84	241495	0.348	
United Republic of Tanzania	TZ	1910	939588	2.033	
Western Sahara	EH	0	268617	0	

	Zambia	ZM	603	751063	0.803	
	Zimbabwe	ZW	413	390648	1.057	
Antarctica	Antarctica	AQ	30	12537967	0.002	
	Bouvet Island	BV	0	45	0	
	French Southern and Antarctic Territories	TF	0	7738	0	
	Heard Island and McDonald Islands	HM	0	412	0	
	South Georgia and the South Sandwich Islands	GS	0	3870	0	
Asia	Afghanistan	AF	19	641827	0.03	
	Aksai Chin		1	0	30666	0
	Armenia	AM	509	29624	17.182	
	Arunachal Pradesh		2	2	67965	0.029
	Azerbaijan	AZ	28	164780	0.17	
	Bahrain	BH	2	673	2.97	
	Bangladesh	BD	207	139825	1.48	
	Bhutan	BT	85	37674	2.256	
	Brunei Darussalam	BN	0	5899	0	
	Cambodia	KH	424	181424	2.337	
	China	CN	1644	9345214	0.176	
	China/India		3	0	3526	0
	Christmas Island	CX	0	136	0	
	Cocos (Keeling) Islands	CC	0	16	0	
	Cyprus	CY	12	9249	1.297	
	Democratic People's Republic of Korea	KP	0	122465	0	
	Georgia	GE	18	69785	0.258	
	Hong Kong	HK	2	1081	1.851	
	India	IN	199	2961118	0.067	
	Indonesia	ID	179	1888620	0.095	
	Iran (Islamic Republic of)	IR	2010	1677319	1.198	
	Iraq	IQ	14	435864	0.032	

Israel	IL		17	20720	0.82
Jammu and Kashmir		12	4	186035	0.022
Japan	JP		197	373651	0.527
Jordan	JO		47	89063	0.528
Kazakhstan	KZ		52	2841103	0.018
Kuril islands		5	0	4996	0
Kuwait	KW		1	17392	0.057
Kyrgyzstan	KG		1	199188	0.005
Lao People's Democratic Republic	LA		20	230380	0.087
Lebanon	LB		10	10136	0.987
Macau	MO		0	17	0
Malaysia	MY		155	329775	0.47
Maldives	MV		0	223	0
Mongolia	MN		9	1564529	0.006
Myanmar	MM		0	667085	0
Nepal	NP		142	147437	0.963
Occupied Palestinian Territory	PS		18	6225	2.892
Oman	OM		11	308335	0.036
Pakistan	PK		45	788439	0.057
Paracel Islands		6	0	8	0
Philippines	PH		78	296031	0.263
Qatar	QA		0	11549	0
Republic of Korea	KR		23	99124	0.232
Saudi Arabia	SA		7	1925621	0.004
Scarborough Reef		7	0	44	0
Senkaku Islands		8	0	5	0
Singapore	SG		1	594	1.683
Spratly Islands		9	0	1	0
Sri Lanka	LK		73	66173	1.103
Syrian Arab Republic	SY		69	188128	0.367
Taiwan	TW		35	36127	0.969
Tajikistan	TJ		5	142004	0.035
Thailand	TH		479	515417	0.929
Timor-Leste	TL		0	14892	0

Turkey	TR	69	781229	0.088
Turkmenistan	TM	0	555052	0
United Arab Emirates	AE	12	71079	0.169
Uzbekistan	UZ	9	449620	0.02
Viet Nam	VN	29	327575	0.089
Yemen	YE	284	453596	0.626

Europe

Albania	AL	97	28682	3.382
Andorra	AD	0	475	0
Austria	AT	128	83964	1.524
Belarus	BY	96	207581	0.462
Belgium	BE	7013	30669	228.667
Bosnia and Herzegovina	BA	32	51145	0.626
Bulgaria	BG	134	111300	1.204
Croatia	HR	78	56589	1.378
Czech Republic	CZ	666	78845	8.447
Denmark	DK	72	44458	1.619
Estonia	EE	241	45441	5.304
Faroe Islands	FO	0	1400	0
Finland	FI	442	336892	1.312
France	FR	3183	548785	5.8
Germany	DE	4362	357227	12.211
Gibraltar	GI	0	6	0
Greece	GR	374	132549	2.822
Guernsey	GG	0	79	0
Holy See	VA	0	0	0
Hungary	HU	1421	93119	15.26
Iceland	IS	17	102566	0.166
Ireland	IE	124	69809	1.776
Isle of Man	IM	0	573	0
Italy	IT	576	301651	1.909
Jersey	JE	0	120	0
Latvia	LV	102	64563	1.58
Liechtenstein	LI	0	151	0

Lithuania	LT	127	64943	1.956
Luxembourg	LU	142	2621	54.184
Malta	MT	0	316	0
Monaco	MC	0	8	0
Montenegro	ME	12	13776	0.871
Netherlands	NL	958	35203	27.214
Norway	NO	507	324257	1.564
Poland	PL	796	311961	2.552
Portugal	PT	455	91876	4.952
Republic of Moldova	MD	35	33798	1.036
Romania	RO	113	238118	0.475
Russian Federation	RU	1464	16998830	0.086
San Marino	SM	0	60	0
Serbia	RS	69	88478	0.78
Slovakia	SK	161	49072	3.281
Slovenia	SI	67	20320	3.297
Spain	ES	907	505752	1.793
Svalbard and Jan Mayen Islands	SJ	4	63464	0.063
Sweden	SE	594	449212	1.322
Switzerland	CH	10928	41257	264.874
The Republic of North Macedonia	MK	20	25424	0.787
Ukraine	UA	462	600526	0.769
United Kingdom	GB	1727	244308	7.069

Northern America	Anguilla	AI	0	79	0
	Antigua and Barbuda	AG	0	452	0
	Aruba	AW	0	180	0
	Bahamas	BS	0	11904	0
	Barbados	BB	3	433	6.928
	Belize	BZ	26	21764	1.195
	Bermuda	BM	0	63	0
	British Virgin Islands	VG	0	154	0
	Canada	CA	8778	9875646	0.889
	Cayman Islands	KY	0	269	0

	Clipperton Island	CP	0	9	0
	Costa Rica	CR	560	51042	10.971
	Cuba	CU	53	110863	0.478
	Dominica	DM	0	751	0
	Dominican Republic	DO	10	48099	0.208
	El Salvador	SV	38	20732	1.833
	Greenland	GL	2	2165159	0.001
	Grenada	GD	0	318	0
	Guadeloupe	GP	5	1697	2.947
	Guatemala	GT	28	109062	0.257
	Haiti	HT	0	27022	0
	Honduras	HN	38	112124	0.339
	Jamaica	JM	74	10965	6.749
	Martinique	MQ	0	1104	0
	Mexico	MX	12599	1949527	6.463
	Montserrat	MS	0	101	0
	Netherlands Antilles	AN	4	790	5.066
	Nicaragua	NI	21	128376	0.164
	Panama	PA	50	74850	0.668
	Puerto Rico	PR	280	8937	31.329
	Saint Kitts and Nevis	KN	0	262	0
	Saint Lucia	LC	0	603	0
	Saint Pierre and Miquelon	PM	0	233	0
	Saint Vincent and the Grenadines	VC	0	427	0
	Trinidad and Tobago	TT	2	5144	0.389
	Turks and Caicos Islands	TC	0	530	0
	United States Minor Outlying Islands	UM	0	348	0
	United States of America	US	56322	9315946	6.046
	United States Virgin Islands	VI	46	352	130.555
Oceania	American Samoa	AS	0	200	0
	Australia	AU	42767	7687634	5.563
	Cook Islands	CK	0	241	0

	Fiji	FJ	6	18293	0.328
	French Polynesia	PF	0	3967	0
	Guam	GU	15	544	27.579
	Kiribati	KI	0	1020	0
	Marshall Islands	MH	0	268	0
	Micronesia (Federated States of)	FM	75	740	101.343
	Nauru	NR	0	22	0
	New Caledonia	NC	2	18574	0.108
	New Zealand	NZ	52	270415	0.192
	Niue	NU	0	263	0
	Norfolk Island	NF	0	38	0
	Northern Mariana Islands	MP	0	476	0
	Palau	PW	18	451	39.924
	Papua New Guinea	PG	24	462230	0.052
	Pitcairn Islands	PN	0	49	0
	Samoa	WS	18	2835	6.349
	Solomon Islands	SB	1	28264	0.035
	Tokelau	TK	0	15	0
	Tonga	TO	0	700	0
	Tuvalu	TV	0	48	0
	Vanuatu	VU	1	12236	0.082
	Wake Island	WK	0	7	0
	Wallis and Futuna Islands	WF	0	142	0
South America	Argentina	AR	253	2780175	0.091
	Bolivia (Plurinational State of)	BO	87	1084491	0.08
	Brazil	BR	9262	8485946	1.091
	Chile	CL	13662	753355	18.135
	Colombia	CO	236	1137939	0.207
	Ecuador	EC	94	256249	0.367
	Falkland Islands (Malvinas)	FK	0	12084	0
	French Guiana	GF	30	83295	0.36
	Guyana	GY	43	211722	0.203
	Paraguay	PY	2	399349	0.005
	Peru	PE	158	1290640	0.122

Suriname	SR	31	145100	0.214
Uruguay	UY	136	177811	0.765
Venezuela (Bolivarian Republic of)	VE	204	912025	0.224

^a Disputed territory. Country names and areas are based on the Global Administrative Layers (GAUL) database, see: <https://data.apps.fao.org/map/catalogsrv/eng/catalog.search?id=12691#/metadata/9c35ba10-5649-41c8-bdfc-eb78e9e65654> (last access: 26 April 2024).

Table C2. Number of sites by world terrestrial ecosystems (WTE)^a.

Temperature zone	Moisture zone	No. of sites	Percent (%)
Polar	Dry	224	0.1
Polar	Moist	532	0.2
Boreal	Dry	1789	0.8
Boreal	Moist	3398	1.6
Cool Temperate	Desert	25	0
Cool Temperate	Dry	10968	5
Cool Temperate	Moist	53245	24.5
Warm Temperate	Desert	238	0.1
Warm Temperate	Dry	29209	13.4
Warm Temperate	Moist	46533	21.4
Sub Tropical	Desert	296	0.1
Sub Tropical	Dry	25748	11.8
Sub Tropical	Moist	17906	8.2
Tropical	Desert	178	0.1
Tropical	Dry	11315	5.2
Tropical	Moist	11095	5.1
No data	-	4674	2.2

^a World Terrestrial Ecosystems (WTE) as defined by Sayre (2022). Total may differ from 100% due to rounding.

Table C3. Number of sites by WWF biome^b.

WWF biome	No. of sites	Percent (%)
Boreal Forests/Taiga	5519	2.5
Deserts and Xeric Shrublands	13410	6.2
Flooded Grasslands and Savannas	792	0.4
Lakes	85	0
Mangroves	765	0.4
Mediterranean Forests, Woodlands, and Scrub	24459	11.3
Montane Grasslands and Shrublands	2796	1.3
Rock and Ice	20	0
Temperate Broadleaf and Mixed Forests	74068	34.1
Temperate Coniferous Forests	14436	6.6
Temperate Grasslands, Savannas, and Shrublands	23890	11
Tropical and Subtropical Coniferous Forests	2363	1.1
Tropical and Subtropical Dry Broadleaf Forests	4120	1.9
Tropical and subtropical grasslands, savannas, and shrublands	31376	14.4
Tropical and Subtropical Moist Broadleaf Forests	16478	7.6
Tundra	2072	1
No data	724	0.3

^a Biomes defined according to “Terrestrial Ecoregions of the World” (WWF) (Olson et al., 2001a). Total may differ from 100% due to rounding.

Author contributions. NB is scientific lead of the WoSIS project and wrote the first draft. LC developed the ETL and GraphQL procedures while LdS developed the new data model. All authors performed quality checks, data analyses and contributed to the writing/editing of the final manuscript.

Competing interests. The authors declare that they have no conflict of interest.

- 5 **Disclaimer.** ISRIC – World Soil Information remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Acknowledgements

The development of WoSIS has been made possible thanks to the contributions and shared knowledge of a steadily growing number of data providers, including soil survey organisations, research institutes and individual experts, for which we are grateful.

- 10 Regrettably, we can impossibly acknowledge all contributors (e.g., field surveyors, laboratory personnel, soil experts and database experts) individually. Therefore, we do this largely in a generic way (see <https://www.isric.org/explore/wosis/wosis-contributing-institutions-and-experts>; last access: 26 April 2024).

- Our special thanks go to Eloi Ribeiro, former WoSIS database management expert at ISRIC, for his sustained support and advice during the refactoring of the ETL procedure. We also thank our colleague Laura Poggio for useful methodological
15 discussions concerning methodological linkages between WoSIS and SoilGrids.

We gratefully acknowledge Dr. Alessandro Samuel Rosa, the second anonymous reviewer, and the topic editor Dr. Sybille Haßler for their comments which greatly improved the scope of the initial submission.

- The ETL procedures and new data model were co-developed in the framework of the European Union’s Horizon 2020
20 HoliSoils project (Grant agreement 101000289) and ISRIC’s “global soil information and standards” workstream.

ISRIC – World Soil Information, legally registered as International Soil Reference and Information Centre, receives core funding from the Dutch Government.

References

- Al-Shammary, A. A. G., Kouzani, A. Z., Kaynak, A., Khoo, S. Y., Norton, M., and Gates, W.: Soil Bulk Density Estimation Methods: A Review, *Pedosphere*, 28, 581-596, [https://doi.org/10.1016/S1002-0160\(18\)60034-7](https://doi.org/10.1016/S1002-0160(18)60034-7), 2018.
- 5 ANSIS: Australian National Soil Information System, Australian National Soil Information System, Canberra (AU), <https://ansis.net/>, 2023.
- Armas, D., Guevara, M., Bezares, F., Vargas, R., Durante, P., Osorio, V., Jiménez, W., and Oyonarte, C.: Harmonized Soil Database of Ecuador (HESD): data from 2009 to 2015, *Earth Syst. Sci. Data*, 15, 431-445, <https://doi.org/10.5194/essd-15-431-2023>, 2023.
- 10 Arrouays, D., Leenaars, J. G. B., Richer-de-Forges, A. C., Adhikari, K., Ballabio, C., Greve, M., Grundy, M., Guerrero, E., Hempel, J., Hengl, T., Heuvelink, G., Batjes, N., Carvalho, E., Hartemink, A., Hewitt, A., Hong, S.-Y., Krasilnikov, P., Lagacherie, P., Lelyk, G., Libohova, Z., Lilly, A., McBratney, A., McKenzie, N., Vasquez, G. M., Leatitia Mulder, V., Minasny, B., Luca, M., Odeh, I., Padarian, J., Poggio, L., Roudier, P., Saby, N., Savin, I., Searle, R., Solbovoy, V., Thompson, J., Smith, S., Sulaeman, Y., Vintila, R., Rossel, R. V., Wilson, P., Zhang, G.-L., Swerts, M., Oorts, K., Karklins, A., Feng, L.,
- 15 Ibelles Navarro, A. R., Levin, A., Laktionova, T., Dell'Acqua, M., Suvannang, N., Ruam, W., Prasad, J., Patil, N., Husnjak, S., Pasztor, L., Okx, J., Hallet, S., Keay, C., Farewell, T., Lilja, H., Juilleret, J., Marx, S., Takata, Y., Kazuyuki, Y., Mansuy, N., Panagos, P., Van Liedekerke, M., Skalsky, R., Sobocka, J., Kobza, J., Eftekhari, K., Kacem Alavipanah, S., Moussadek, R., Badraoui, M., Da Silva, M., Paterson, G., da Conceicao Gonsalves, M., Theocharopoulos, S., Yemefack, M., Tedou, S., Vrscaj, B., Grob, U., Kozak, J., Boruvka, L., Dobos, E., Taboada, M., Moretti, L., and Rodriguez, D.: Soil legacy data rescue via
- 20 GlobalSoilMap and other international and national initiatives, *GeoResJ*, 14, 1-19, <https://doi.org/10.1016/j.grj.2017.06.001>, 2017.
- Ballabio, C., Panagos, P., and Monatanarella, L.: Mapping topsoil physical properties at European scale using the LUCAS database, *Geoderma*, 261, 110-123, <https://doi.org/10.1016/j.geoderma.2015.07.006>, 2016.
- Baritz, R., Erdogan, H., Fujii, K., Takata, Y., Nocita, M., Bussian, B., Batjes, N. H., Hempel, J., Wilson, P., and Vargas, R.:
- 25 Harmonization of methods, measurements and indicators for the sustainable management and protection of soil resources (Providing mechanisms for the collation, analysis and exchange of consistent and comparable global soil data and information), Global Soil Partnership, FAO, 44 pp., <http://www.fao.org/3/a-az922e.pdf>, 2014.
- Baritz, R., Erdogan, H., Ahmadov, H., Ghanma, I., Lalljee, V. B., Wongmaneroj, A., Collins, A., Monger, C., Ribeiro, J. L., Bertsch, F., Lalljee, V. B., with Montanarella, L., Comerma, J., Khan, A., VandenBygaart, B., Gaistardo, C. C., Constantini,
- 30 E., Galbraith, J. M., Schad, P., Lame, F., Suvannang, N., Hartmann, C., Medyckyj-Scott, D., Batjes, N. H., van Liedekerke, M., and Ziadat, F.: Implementation Plan for Pillar Five of the Global Soil Partnership: Providing mechanisms for the collation,

- analysis and exchange of consistent and comparable global soil data and information [Submitted to ITPS 05/2017], Rome, 48 pp., <http://www.fao.org/3/a-bs756e.pdf>, 2017.
- Baroni, G., Zink, M., Kumar, R., Samaniego, L., and Attinger, S.: Effects of uncertainty in soil properties on simulated hydrological states and fluxes at different spatio-temporal scales, *Hydrology and Earth System Sciences*, 21, 2301-2320, 5 <https://doi.org/10.5194/hess-21-2301-2017>, 2017.
- Batjes, N. H., and Bridges, E. M.: Potential emissions of radiatively active gases from soil to atmosphere with special reference to methane: development of a global database (WISE), *Journal of Geophysical Research*, 99(D8), 16479-16489, <http://dx.doi.org/10.1029/93JD03278>, 1994.
- Batjes, N. H.: A world dataset of derived soil properties by FAO–UNESCO soil unit for global modelling, *Soil Use and* 10 *Management*, 13, 9-16, <https://doi.org/10.1111/j.1475-2743.1997.tb00550.x>, 1997.
- Batjes, N. H.: Harmonized soil profile data for applications at global and continental scales: updates to the WISE database, *Soil Use and Management*, 25, 124-127, <https://doi.org/10.1111/j.1475-2743.2009.00202.x>, 2009.
- Batjes, N. H.: Harmonised soil property values for broad-scale modelling (WISE30sec) with estimates of global soil carbon stocks, *Geoderma*, 269, 61-68, <http://dx.doi.org/10.1016/j.geoderma.2016.01.034> 2016.
- 15 Batjes, N. H., Ribeiro, E., van Oostrum, A., Leenaars, J., Hengl, T., and Mendes de Jesus, J.: WoSIS: providing standardised soil profile data for the world, *Earth Syst. Sci. Data*, 9, 1-14, <http://dx.doi.org/10.5194/essd-9-1-2017>, 2017.
- Batjes, N. H., Ribeiro, E., and van Oostrum, A.: Standardised soil profile data to support global mapping and modelling (WoSIS snapshot 2019), *Earth Syst. Sci. Data*, 12, 299-320, <https://doi.org/10.5194/essd-12-299-2020>, 2020.
- Batjes, N. H.: Options for harmonising soil data obtained from different sources ISRIC - World Soil Information, Wageningen, 20 21 pp., <https://dx.doi.org/10.17027/isric-wdc-6ztd-eb19> 2023.
- Batjes, N. H., and van Oostrum, A. J. M.: WoSIS Procedures for standardizing soil analytical method descriptions, ISRIC - World Soil Information, Wageningen, 46 pp., <https://doi.org/10.17027/isric-1dq0-1m83>, 2023.
- Battle-Bayer, L., Batjes, N. H., and Bindraban, P. S.: Changes in organic carbon stocks upon land use conversion in the Brazilian Cerrado: A review, *Agriculture, Ecosystems & Environment*, 137, 47-58, 25 <http://dx.doi.org/10.1016/j.agee.2010.02.003>, 2010.
- Bispo, A., Arrouays, D., Saby, N., Boulonne, L., and Fantappiè, M.: Proposal of methodological development for the LUCAS programme in accordance with national monitoring programmes. Towards climate-smart sustainable management of agricultural soils (EU H2020-SFS-2018-2020 / H2020-SFS-2019) *EJP Soil*, 135 pp., https://ejpsoil.eu/fileadmin/projects/ejpsoil/WP6/EJP_SOIL_Deliverable_6.3_Dec_2021_final.pdf, 2021.
- 30 Bridges, E. M.: Soil horizon designations: past use and future prospects, *CATENA*, 20, 363-373, [https://doi.org/10.1016/S0341-8162\(05\)80002-5](https://doi.org/10.1016/S0341-8162(05)80002-5), 1993.

- Brus, D. J., Kempen, B., and Heuvelink, G. B. M.: Sampling for validation of digital soil maps, *European Journal of Soil Science*, 62, 394-407, <https://bsssjournals.onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-2389.2011.01364.x>, 2011.
- Brus, J.: *Spatial sampling with R*, Chapman and Hall R/C, New York, 2022.
- Calisto, L.: ISRIC GraphQL web services for WoSIS and ISIS data access, ISRIC - World Soil Information, Wageningen, 5 <https://graphql.isric.org/>, 2023.
- Calisto, L., de Souza, L. M., and Batjes, N. H.: Standardised soil profile data for the world (WoSIS, December snapshot) [Dataset, <https://doi.org/10.17027/isric-wdcsoids-20231130>]. ISRIC - World Soil Information, Wageningen, 2023.
- Cornu, S., Keesstra, S., Bispo, A., Fantappie, M., van Egmond, F., Smreczak, B., Wawer, R., Pavlů, L., Sobocká, J., Bakacsi, Z., Farkas-Iványi, K., Molnár, S., Møller, A. B., Madenoglu, S., Feiziene, D., Oorts, K., Schneider, F., Gonçalves, M. d. C., 10 Mano, R., Garland, G., Skalský, R., O'Sullivan, L., Kasparinskis, R., and Chenu, C.: National soil data in EU countries, where do we stand?, *European Journal of Soil Science*, e13398, <https://doi.org/10.1111/ejss.13398>, 2023.
- Cox, S., and David, J.: ISO 19156:2011 Geographic information – Observations and measurements International Organization for Standardization., <https://www.iso.org/standard/32574.html>, 2011.
- Cramer, M. D., Wootton, L. M., van Mazijk, R., and Verboom, G. A.: New regionally modelled soil layers improve prediction 15 of vegetation type relative to that based on global soil models, *Diversity and Distributions*, 25, 1736-1750, <https://onlinelibrary.wiley.com/doi/abs/10.1111/ddi.12973>, 2019.
- Cressie, N., and Kornak, J.: Spatial statistics in the presence of location error with an application to remote sensing of the environment, *Statistical Science*, 18, 436-456, <https://doi.org/10.1214/ss/1081443228>, 2003.
- Dai, Y., Shangguan, W., Wang, D., Wei, N., Xin, Q., Yuan, H., Zhang, S., Liu, S., and Yan, F.: A review on the global soil 20 datasets for earth system models, *SOIL*, 5, 137-158, <https://doi.org/10.5194/soil-5-137-2019>, 2019.
- de Sousa, L., Kempen, B., Mendes de Jesus, J., Yigini, Y., Viatkin, K., Medyckyj-Scott, D., Richie, D. A., Wilson, P., van Egmond, F., and Baritz, R.: Conceptual design of the Global Soil Information System infrastructure, Rome, FAO and ISRIC, Wageningen, Netherlands, 30 pp., <http://www.fao.org/3/cb4355en/cb4355en.pdf>, 2021.
- de Sousa, L. M., Kempen, B., Mendes de Jesus, J., Yigini, Y., Viatkin, K., Medyckyj-Scott, D., Richie, A., Wilson, P., van 25 Egmond, F., and Baritz, R.: Conceptual desing of the Global Soil Information System infrastructure, ISRIC, FAO, Manaaki Whenua (Landcare Research), CSIRO, Wageningen UR, European Environment Agency, 30 pp., <http://www.fao.org/3/cb4355en/cb4355en.pdf>, 2019.
- de Sousa, L. M.: WoSIS data model 2023. Procedures Manual - Technical documentation, ISRIC - World Soil Information, Wageningen, <https://git.wur.nl/isric/databases/wosis-docs>, 2023.
- 30 de Sousa, L. M., Calisto, L., van Genuchten, P., Turdukulov, U., and Kempen, B.: Data model for the ISO 28258 domain model, ISRIC - World Soil Information, <https://iso28258.isric.org/>, 2023.

- Dijkshoorn, J. A., Huting, J. R. M., and Tempel, P.: Update of the 1:5 million Soil and Terrain Database for Latin America and the Caribbean (SOTERLAC, ver. 2.0), ISRIC - World Soil Information, Wageningen, Report 2005/01, <https://www.isric.org/documents/document-type/isric-report-200501-update-15-million-soil-and-terrain-database-latin>, 2005.
- Fantappie, M., Peruginelli, G., Conti, S., Rennes, S., van Egmond, F. M., and Le Bas, C.: Towards climate-smart sustainable management of agricultural soils: Deliverable 6.2 Report on the national and EU regulations on agricultural soil data sharing and national monitoring activities, 202 pp., <https://edepot.wur.nl/642353>, 2021.
- FAO: Guidelines for the description of soils (2nd ed.), FAO, Rome, 66 pp., 1977.
- FAO: Guidelines for soil description (3rd rev. ed.), FAO, Rome, 45 pp., <https://edepot.wur.nl/570291>, 1990.
- FAO, ISRIC, UNEP, and CIP: Soil and terrain digital database for Latin America and the Caribbean at 1:5 million scale, Food and Agriculture Organization of the United Nations, Rome, Land and Water Digital Media Series No. 51998.
- FAO, and ISRIC: Soil and Terrain database for Southern Africa (1:2 million scale), ISRIC and FAO, Rome, FAO Land and Water Digital Media Series 252003.
- FAO: Guidelines for soil description (4th ed.), FAO, Rome, 97 pp., <http://www.fao.org/docrep/019/a0541e/a0541e.pdf>, 2006.
- FAO, ISRIC, and UG: Soil and terrain database for central Africa (Burundi and Rwanda 1:1 million scale; Democratic Republic of the Congo 1:2 million scale), Food and Agricultural Organization of the United Nations, ISRIC - World Soil Information and Universiteit Gent, Rome, Land and Water Digital Media Series 33, https://www.isric.org/sites/default/files/isric_report_2006_07.pdf, 2007.
- FAO, IIASA, ISRIC, ISSCAS, and JRC: Harmonized World Soil Database (version 1.2), Prepared by Nachtergaele FO, van Velthuisen H, Verelst L, Wiberg D, Batjes NH, Dijkshoorn JA, van Engelen VWP, Fischer G, Jones A, Montanarella L., Petri M, Prieler S, Teixeira E and Xuezheng Shi. Food and Agriculture Organization of the United Nations (FAO), International Institute for Applied Systems Analysis (IIASA), ISRIC - World Soil Information, Institute of Soil Science - Chinese Academy of Sciences (ISSCAS), Joint Research Centre of the European Commission (JRC), Laxenburg, Austria, http://webarchive.iiasa.ac.at/Research/LUC/External-World-soil-database/HWSD_Documentation.pdf, 2012.
- FAO, and ITPS: Status of the world's soil resources (SWSR) - Main report, Food and Agriculture Organization of the United Nations and Intergovernmental Technical Panel on Soils, Rome, 650 pp., <http://www.fao.org/3/a-i5199e.pdf>, 2015.
- Finke, P.: Quality assessment of digital soil maps: producers and users perspectives, in: Digital soil mapping: An introductory perspective, edited by: Lagacherie, P., McBratney, A., and Voltz, M., Elsevier, Amsterdam, 523-541, 2006.
- Folberth, C., Skalsky, R., Moltchanova, E., Balkovic, J., Azevedo, L. B., Obersteiner, M., and van der Velde, M.: Uncertainty in soil data can outweigh climate impact signals in global crop yield simulations, Nature Communications, 7, <https://doi.org/10.1038/ncomms11872>, 2016.
- Gerasimova, M. I., Lebedeva, I. I., and Khitrov, N. B.: Soil horizon designation: State of the art, problems, and proposals, Eurasian Soil Science, 46, 599-609, <https://dx.doi.org/10.1134/S1064229313050037>, 2013.

- Giller, K. E., Rowe, E. C., de Ridder, N., and van Keulen, H.: Resource use dynamics and interactions in the tropics: Scaling up in space and time, *Agricultural Systems*, 88, 8-27, <https://doi.org/10.1016/j.agsy.2005.06.016>, 2006.
- GlobalSoilMap: Specifications Tiered GlobalSoilMap products (Release 2.4), 52 pp., <https://www.isric.org/documents/document-type/globalsoilmap-specifications-v24-07122015>, 2015.
- 5 GLOSOLAN: GLOSOLAN best practice manual (on-line), FAO, GSP, Rome, <https://www.fao.org/global-soil-partnership/glosolan-old/soil-analysis/standard-operating-procedures/en/#c763834>, 2023.
- Gobezie, T. B., and Biswas, A.: Break barriers in soil data stewardship by rewarding data generators, *Nature Reviews Earth & Environment*, 4, 353-354, <https://doi.org/10.1038/s43017-023-00439-4>, 2023.
- Grimm, R., and Behrens, T.: Uncertainty analysis of sample locations within digital soil mapping approaches, *GEODERMA*, 10 155, 154-163, <https://doi.org/10.1016/j.geoderma.2009.05.006>, 2010.
- Guevara, M., Olmedo, G. F., Stell, E., Yigini, Y., Aguilar Duarte, Y., Arellano Hernández, C., Arévalo, G. E., Arroyo-Cruz, C. E., Bolivar, A., Bunning, S., Bustamante Cañas, N., Cruz-Gaistardo, C. O., Davila, F., Dell Acqua, M., Encina, A., Figueredo Tacona, H., Fontes, F., Hernández Herrera, J. A., Ibelle Navarro, A. R., Loayza, V., Manueles, A. M., Mendoza Jara, F., Olivera, C., Osorio Hermosilla, R., Pereira, G., Prieto, P., Alexis Ramos, I., Rey Brina, J. C., Rivera, R., Rodríguez-Rodríguez, 15 J., Roopnarine, R., Rosales Ibarra, A., Rosales Riveiro, K. A., Schulz, G. A., Spence, A., Vasques, G. M., Vargas, R. R., and Vargas, R.: No Silver Bullet for Digital Soil Mapping: Country-specific Soil Organic Carbon Estimates across Latin America, *SOIL*, 2018, 173-193, <https://doi.org/10.5194/soil-4-173-2018>, 2018.
- Hassani, A., Smith, P., and Shokri, N.: Negative correlation between soil salinity and soil organic carbon variability, *Proceedings of the National Academy of Sciences*, 121, e2317332121, 20 24 <https://www.pnas.org/doi/abs/10.1073/pnas.2317332121>, 2024.
- Hengl, T., de Jesus, J. M., Heuvelink, G. B. M., Gonzalez, M. R., Kilibarda, M., Blagotic, A., Shangguan, W., Wright, M. N., Geng, X. Y., Bauer-Marschallinger, B., Guevara, M. A., Vargas, R., MacMillan, R. A., Batjes, N. H., Leenaars, J. G. B., Ribeiro, E., Wheeler, I., Mantel, S., and Kempen, B.: SoilGrids250m: Global gridded soil information based on machine learning, *PLoS ONE*, 12, <https://doi.org/10.1371/journal.pone.0169748>, 2017.
- 25 Heuvelink, G. B. M., Brown, J. D., and van Loon, E. E.: A probabilistic framework for representing and simulating uncertain environmental variables, *International Journal of Geographical Information Science*, 21, 497-513, <https://doi.org/10.1080/13658810601063951>, 2007.
- Heuvelink, G. B. M.: Uncertainty quantification of GlobalSoilMap products in: *GlobalSoilMap. Basis of the Global Spatial Soil Information System*, edited by: Arrouays, D., McKenzie, N., Hempel, J., Forges, A. R. d., and McBratney, A., Taylor & Francis 30 Group, London, UK, 335-240, 2014.

- Heuvelink, G. B. M., Angelini, M. E., Poggio, L., Bai, Z. G., Batjes, N. H., van den Bosch, R., Bossio, D., Estella, S., Lehmann, J., Olmedo, G. F., and Sanderman, J.: Machine learning in space and time for modelling soil organic carbon change, *European Journal of Soil Science*, 72, 1607-1623, <https://doi.org/10.1111/ejss.12998>, 2021.
- Huang, Y., Song, X., Wang, Y.-P., Canadell, J. G., Luo, Y., Ciais, P., Chen, A., Hong, S., Wang, Y., Tao, F., Li, W., Xu, Y., Mirzaeitalarposhti, R., Elbasiouny, H., Savin, I., Shechepashchenko, D., Rossel, R. A. V., Goll, D. S., Chang, J., Houlton, B. Z., Wu, H., Yang, F., Feng, X., Chen, Y., Liu, Y., Niu, S., and Zhang, G.-L.: Size, distribution, and vulnerability of the global soil inorganic carbon, *Science*, 384, 233-239, <https://www.science.org/doi/abs/10.1126/science.adi7918>, 2024.
- ICP Forests: ICP Forests monitoring Manual. Part XVI: Quality assurance and control in laboratories (ver 2020-1), Eberswalde, Germany, 46 pp., https://www.icp-forests.org/pdf/manual/2020/ICP_Manual_part16_2020_QAOC_Labs_version_2020-1.pdf, 2020.
- ICP Forests: ICP Forests monitoring Manual. Part X: Sampling and analysis of soil, Eberswalde, Germany, <https://storage.ning.com/topology/rest/1.0/file/get/9995584862?profile=original>, 2021a.
- ICP Forests: ICP Forests monitoring Manual Eberswalde (Germany), <http://icp-forests.net/page/icp-forests-manual>, 2021b.
- ISO-19139: Geographic information XML schema implementation Part 1: Encoding rules, <https://www.iso.org/standard/67253.html>, 2019.
- ISRIC: Data and Software Policy, ISRIC - World Soil Information (WDC - Soils) Wageningen, 6 pp., https://www.isric.org/sites/default/files/user/ISRIC_Data_Policy_2016jun21doi.pdf, 2016.
- IUSS Working Group WRB: World Reference Base for Soil Resources (2nd ed.), FAO, Rome, World Soil Resources Report 103, 145 pp., <http://www.fao.org/ag/agl/agll/wrb/doc/wrb2006final.pdf>, 2006.
- IUSS Working Group WRB: World Reference Base for soil resources 2014 - International soil classification system for naming soils and creating legends for soil maps (update 2015), Global Soil Partnership, International Union of Soil Sciences, and Food and Agriculture Organization of the United Nations, Rome, World Soil Resources Reports 106, 182 pp., <http://www.fao.org/3/i3794en/I3794en.pdf>, 2015.
- IUSS Working Group WRB: World Reference Base for soil resources 2022 - International soil classification system for naming soils and creating legends for soil maps, International Union of Soil Sciences, Vienna (Austria), 284 pp., https://www.isric.org/sites/default/files/WRB_fourth_edition_2022-12-18.pdf, 2022.
- Ivushkin, K., Bartholomeus, H., Bregt, A. K., Pulatov, A., Kempen, B., and de Sousa, L.: Global mapping of soil salinity change, *Remote Sensing of Environment*, 231, <https://doi.org/10.1016/j.rse.2019.111260>, 2019.
- Kalra, Y. P., and Maynard, D. G.: Methods manual for forest soil and plant analysis, Forestry Canada, Edmonton (Alberta), 116 pp., <https://cfs.nrcan.gc.ca/publications/download-pdf/11845>, 1991.
- Leenaars, J. G. B., van Oostrum, A. J. M., and Ruiperez Gonzalez, M.: Africa Soil Profiles Database: A compilation of georeferenced and standardised legacy soil profile data for Sub Saharan Africa (version 1.2), Africa Soil Information Service

- (AFSIS) and ISRIC - World Soil Information, Wageningen, Report 2014/01, 160 pp., http://www.isric.org/sites/default/files/isric_report_2014_01.pdf, 2014.
- Leenaars, J. G. B., Claessens, L., Heuvelink, G. B. M., Hengl, T., Ruiperez González, M., van Bussel, L. G. J., Guilpart, N., Yang, H., and Cassman, K. G.: Mapping rootable depth and root zone plant-available water holding capacity of the soil of sub-Saharan Africa, *Geoderma*, 324, 18-36, <https://doi.org/10.1016/j.geoderma.2018.02.046>, 2018.
- Luo, Z., Viscarra-Rossel, R. A., and Qian, T.: Similar importance of edaphic and climatic factors for controlling soil organic carbon stocks of the world, *Biogeosciences*, 18, 2063-2073, <https://bg.copernicus.org/articles/18/2063/2021/>, 2021.
- Lutz, F., Stoorvogel, J. J., and Müller, C.: Options to model the effects of tillage on N₂O emissions at the global scale, *Ecological Modelling*, 392, 212-225, <https://www.sciencedirect.com/science/article/pii/S0304380018304034>, 2019.
- 10 Magnusson, B., and Örnemark, U.: The Fitness for Purpose of Analytical Methods – A Laboratory Guide to Method Validation and Related Topics (2nd ed.), *Eurachem*, https://www.eurachem.org/images/stories/Guides/pdf/MV_guide_2nd_ed_EN.pdf, 2014.
- Maire, V., Wright, I. J., Prentice, I. C., Batjes, N. H., Bhaskar, R., van Bodegom, P. M., Cornwell, W. K., Ellsworth, D., Niinemets, U., Ordonez, A., Reich, P. B., and Santiago, L. S.: Global effects of soil and climate on leaf photosynthetic traits and rates, *Global Ecology and Biogeography*, 24, 706-717, <https://doi.org/10.1111/geb.12296>, 2015.
- 15 Malhotra, A., Todd-Brown, K., Nave, L. E., Batjes, N. H., Holmquist, J. R., Hoyt, A. M., Iversen, C. M., Jackson, R. B., Lajtha, K., Lawrence, C., Vinduskova, O., Wieder, W., Williams, M., Hugelius, G., and Harden, J.: The landscape of soil carbon data: emerging questions, synergies and databases, *Progress in Physical Geography-Earth and Environment*, 43, 707-719, <https://doi.org/10.1177/0309133319873309>, 2019.
- 20 Meyer, H., and Pebesma, E.: Predicting into unknown space? Estimating the area of applicability of spatial prediction models, *Methods in Ecology and Evolution*, 12, 1620-1633, <https://besjournals.onlinelibrary.wiley.com/doi/abs/10.1111/2041-210X.13650>, 2021.
- Moulatlet, G. M., Zuquim, G., Figueiredo, F. O. G., Lehtonen, S., Emilio, T., Ruokolainen, K., and Tuomisto, H.: Using digital soil maps to infer edaphic affinities of plant species in Amazonia: Problems and prospects, *Ecol Evol*, 7, 8463-8477, <https://doi.org/10.1002/ece3.3242>, 2017.
- 25 Munzert, M., Kießling, G., Übelhör, W., Nätscher, L., and Neubert, K.-H.: Expanded measurement uncertainty of soil parameters derived from proficiency-testing data, *Journal of Plant Nutrition and Soil Science*, 170, 722-728, <https://onlinelibrary.wiley.com/doi/abs/10.1002/jpln.200620701>, 2007.
- NATP: North American Proficiency Testing (NAPT) Program, <http://www.naptprogram.org/>, 2015.
- 30 Nenkam, A. M., Wadoux, A. M. J. C., Minasny, B., McBratney, A. B., Traore, P. C. S., Falconier, G. N., and Whitbread, A. M.: Using homosols for quantitative extrapolation of soil mapping models, *European Journal of Soil Science*, n/a, e13285, <https://doi.org/10.1111/ejss.13285>, 2022.

- NPDB: National Pedon Database Canada, Agriculture and Agri-food Canada, <https://sis.agr.gc.ca/cansis/nsdb/npdb/index.html>, 2023.
- OGC: Soil Data IE (Interoperability Experiment), Open Geospatial Consortium (OGC), <https://www.opengeospatial.org/projects/initiatives/soildataie>, 2019.
- 5 Olson, D. M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell, G. V. N., Underwood, E. C., D'amico, J. A., Itoua, I., Strand, H. E., Morrison, J. C., Loucks, C. J., Allnutt, T. F., Ricketts, T. H., Kura, Y., Lamoreux, J. F., Wettengel, W. W., Hedao, P., and Kassem, K. R.: Terrestrial Ecoregions of the World: A New Map of Life on Earth: A new global map of terrestrial ecoregions provides an innovative tool for conserving biodiversity, *BioScience*, 51, 933-938, [https://doi.org/10.1641/0006-3568\(2001\)051\[0933:TEOTWA\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2001)051[0933:TEOTWA]2.0.CO;2), 2001a.
- 10 Olson, R. J., Johnson, K. R., Zheng, D. L., and Scurlock, J. M. O.: Global and regional ecosystem modelling: databases of model drivers and validation measurements, Oak Ridge National Laboratory, Oak Ridge, ORNL/TM-2001/196, 95 pp., http://www-eosdis.ornl.gov/npp/GPPDI/comp/NPP_TM196.pdf, 2001b.
- Orgiazzi, A., Ballabio, C., Panagos, P., Jones, A., and Fernandez-Ugalde, O.: LUCAS Soil, the largest expandable soil dataset for Europe: a review, *European Journal of Soil Science*, 69, 140-153, <https://doi.org/10.1111/ejss.12499>, 2018.
- 15 Padarian, J., and McBratney, A. B.: A new model for intra- and inter-institutional soil data sharing, *SOIL*, 6, 89-94, <https://soil.copernicus.org/articles/6/89/2020/>, 2020.
- Palma, R., Janiak, B., Sousa, L. M. d., Schleidt, K., Tomáš Rezník, Egmond, F. v., Leenaars, J., Moshou, D., Mouazen, A., Peter Wilson, Medyckyj-Scott, D., Ritchie, A., Yigini, Y., and Vargas, R.: GloSIS: The Global Soil Information System Web Ontology, arXiv, :2403.16778, <https://doi.org/10.48550/arXiv.2403.16778>, 2024.
- 20 Poeplau, C., Don, A., Flessa, H., Heidkamp, A., Jacobs, A., and Prietz, R.: Erste Bodenzustandserhebung Landwirtschaft – Kerndatensatz. . Thünen-Institut, I. f. A. (Ed.), Göttingen, 2020.
- Poggio, L., de Sousa, L., Batjes, N. H., Heuvelink, G. B. M., Kempen, B., Riberio, E., and Rossiter, D.: SoilGrids 2.0: producing soil information for the globe with quantified spatial uncertainty, *SOIL*, 7, 217–240, <https://doi.org/10.5194/soil-7-217-2021>, 2021.
- 25 R Core Team: R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, 2021.
- Rayment, E. R., and Lyons, D. J.: Soil chemical methods - Australasia, CSIRO Publishing, 495 pp., 2011.
- Ribeiro, E., Batjes, N. H., Leenaars, J. G. B., Van Oostrum, A. J. M., and Mendes de Jesus, J.: Towards the standardization and harmonization of world soil data: Procedures Manual ISRIC World Soil Information Service (WoSIS version 2.0) ISRIC -
- 30 World Soil Information, Wageningen, Report 2015/03, 110 pp., http://www.isric.org/sites/default/files/isric_report_2015_03.pdf, 2015.

- Ribeiro, E., Batjes, N. H., and Van Oostrum, A. J. M.: World Soil Information Service (WoSIS) - Towards the standardization and harmonization of world soil data. Procedures Manual 2020, ISRIC - World Soil Information, Wageningen, ISRIC Report 2020/01, 153 pp., <http://dx.doi.org/10.17027/isric-wdc-2020-01>, 2020.
- Robinson, N. J., Dahlhaus, P. G., Wong, M., MacLeod, A., Jones, D., and Nicholson, C.: Testing the public-private soil data and information sharing model for sustainable soil management outcomes, *Soil Use and Management*, 35, 94-104, <https://bsssjournals.onlinelibrary.wiley.com/doi/abs/10.1111/sum.12472>, 2019.
- Rossel, R. A. V., and McBratney, A. B.: Soil chemical analytical accuracy and costs: implications from precision agriculture, *Australian Journal of Experimental Agriculture*, 38, 765-775, 1998.
- Sanderman, J., Hengl, T., and Fiske, G. J.: Soil carbon debt of 12,000 years of human land use, *P Natl Acad Sci USA*, 114, 9575-9580, <https://doi.org/10.1073/pnas.1706103114>, 2017.
- Sayre, R., Dangermond, J., Frye, C., Vaughan, R., Aniello, P., Breyer, S., Cribbs, D., Hopkins, D., Nauman, R., Derrenbacher, W., Burton, D., Grosse, A., True, D., Metzger, M., Hartmann, J., Moosdorf, N., Dürr, H., Paganini, M., DeFourmy, P., Arino, O., and Maynard, S.: A New Map of Global Ecological Land Units — An Ecophysiological Stratification Approach, Association of American Geographers, Washington DC, 46 pp., https://www.aag.org/wp-content/uploads/2021/12/AAG_Global_Ecosyst_bklt72.pdf, 2014.
- Sayre, R.: World Terrestrial Ecosystems (WTE) 2020 [Dataset], <https://doi.org/10.5066/P9DO61LP>, 2022.
- Schoeneberger, P. J., Wysocki, D. A., E.C. Benham, and Soil Survey Staff: Field book for describing and sampling soils (ver. 3.0, Reprint 2021), National Soil Survey Center Natural Resources Conservation Service, U.S. Department of Agriculture, Lincoln (NE), 2012.
- Shepherd, K. D., Ferguson, R., Hoover, D., van Egmond, F., Sanderman, J., and Ge, Y.: A global soil spectral calibration library and estimation service, *Soil Security*, 7, 100061, <https://doi.org/10.1016/j.soisec.2022.100061>, 2022.
- Shi, G., Shangguan, W., Zhang, Y., Li, Q., Wang, C., and Li, L.-J.: Reducing Location Error of Legacy Soil Profiles Leads to Significant Improvement in Digital Soil Mapping, SSRN, <https://ssrn.com/abstract=4643055> or <http://dx.doi.org/10.2139/ssrn.4643055>, 2023.
- Soil Survey Division Staff: Soil survey manual, Soil Conservation Service, U.S. Department of Agriculture, Washington, 503 pp., 1993.
- Soil Survey Staff: Soil Survey Laboratory Information Manual (Ver. 2.0), National Soil Survey Center, Soil Survey Laboratory, USDA-NRCS, Lincoln (NE), Soil Survey Investigation Report No. 45, 506 pp., http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_052226.pdf, 2011.
- Soil Survey Staff: Keys to Soil Taxonomy, 12th ed., USDA-Natural Resources Conservation Service, Washington, DC., 2014.
- Soil Survey Staff: Soil Survey Manual (rev. ed.), edited by: Ditzler, C., Scheffe, K., and Monger, H. C., United States Agriculture Handbook 18, USDA, Washington, 2017.

- Soil Survey Staff: Soil Survey Laboratory Methods Manual (Version 6.0., Part1: Curren methods), U.S. Department of Agriculture, Natural Resources Conservation Service, Lincoln (Nebraska), 1001 pp., 2022a.
- Soil Survey Staff: Keys to Soil Taxonomy, 13th ed., USDA-Natural Resources Conservation Service, Washington, DC., 2022b.
- 5 Sothe, C., Gonsamo, A., Arabian, J., and Snider, J.: Large scale mapping of soil organic carbon concentration with 3D machine learning and satellite observations, *Geoderma*, 405, 115402, <https://www.sciencedirect.com/science/article/pii/S0016706121004821>, 2022.
- Suvannang, N., Hartmann, C., Yakimenko, O., Solokha, M., Bertsch, F., and Moody, P.: Evaluation of the First Global Soil Laboratory Network (GLOSOLAN) online survey for assessing soil laboratory capacities, Global Soil Partnership (GSP) / Food and Agriculture Organization of the United Nations (FAO), Rome, GLOSOLAN/18/Survey Report, 54 pp., <http://www.fao.org/3/CA2852EN/ca2852en.pdf>, 2018.
- 10 Tempel, P., van Kraalingen, D., Mendes de Jesus, J., and Reuter, H. I.: Towards an ISRIC World Soil Information Service (WOSIS ver. 1.0), ISRIC - World Soil Information, Wageningen, ISRIC Report 2013/02, 188 pp., https://www.isric.org/sites/default/files/isric_report_2013_02.pdf, 2013.
- Turek, M. E., Poggio, L., Batjes, N. H., Armindo, R. A., de Jong van Lier, Q., de Sousa, L., and Heuvelink, G. B. M.: Global mapping of volumetric water retention at 100, 330 and 15 000 cm suction using the WoSIS database, *International Soil and Water Conservation Research*, <https://www.sciencedirect.com/science/article/pii/S2095633922000636>, 2022.
- 15 Turek, M. E., Poggio, L., Batjes, N. H., Armindo, R. A., de Jong van Lier, Q., de Sousa, L., and Heuvelink, G. B. M.: Global mapping of volumetric water retention at 100, 330 and 15 000 cm suction using the WoSIS database, *International Soil and Water Conservation Research*, 11, 225-239, <https://www.sciencedirect.com/science/article/pii/S2095633922000636>, 2023.
- 20 USDA-NCSS: National Cooperative Soil Survey (NCSS) Soil Characterization Database, United States Department of Agriculture, Natural Resources Conservation Service, Lincoln, https://ncsslabdatamart.sc.egov.usda.gov/database_download.aspx, 2021.
- van de Ven, T., and Tempel, P.: ISIS 4.0 - ISRIC Soil Information System: User Manual, International Soil Reference and Information Centre, Wageningen, Technical Paper 15 (rev. ed.), https://www.isric.org/sites/default/files/ISRIC_TechPap15b.pdf, 1994.
- 25 van Engelen, V. W. P., Verdoodt, A., Dijkshoorn, K., and van Ranst, E.: SOTER database for Central Africa -- DR Congo, Burundi and Rwanda (SOTERCAF; ver. 1.0), Laboratory of Soil Science (University of Ghent), FAO and ISRIC - World Soil Information, Wageningen (http://www.isric.org/Isric/Webdocs/Docs/ISRIC_Report_2006_07.pdf; accessed 15 August 2007), ISRIC REport 2006/07, 28 pp.2006.
- 30 van Leeuwen, C., Mulder, V. L., Batjes, N. H., and Heuvelink, G. B. M.: Statistical modelling of measurement error in wet chemistry soil data, *European Journal of Soil Science*, 73, 13137, <https://doi.org/10.1111/ejss.13137>, 2022.

- van Leeuwen, C. C. E., Mulder, V. L., Batjes, N. H., and Heuvelink, G. B. M.: Effect of measurement error in wet chemistry soil data on the calibration and model performance of pedotransfer functions, *Geoderma*, 442, 116762, <https://www.sciencedirect.com/science/article/pii/S0016706123004391>, 2024.
- Van Looy, K., Bouma, J., Herbst, M., Koestel, J., Minasny, B., Mishra, U., Montzka, C., Nemes, A., Pachepsky, Y., Padarian, J., Schaap, M., Tóth, B., Verhoef, A., Vanderborght, J., van der Ploeg, M., Weihermüller, L., Zacharias, S., Zhang, Y., and Vereecken, H. C. R. G.: Pedotransfer functions in Earth system science: challenges and perspectives, *Reviews of Geophysics*, 55, 1199-1256, <http://dx.doi.org/10.1002/2017RG000581>, 2017.
- van Reeuwijk, L. P.: On the way to improve international soil classification and correlation: the variability of soil analytical data, ISRIC, Wageningen, Annual Report 1983, 7-13 pp., https://www.isric.org/sites/default/files/isric_annual_report_1983.pdf, 1983.
- Viscarra Rossel, R. A., Behrens, T., Ben-Dor, E., Brown, D. J., Demattê, J. A. M., Shepherd, K. D., Shi, Z., Stenberg, B., Stevens, A., Adamchuk, V., Aichi, H., Barthès, B. G., Bartholomeus, H. M., Bayer, A. D., Bernoux, M., Böttcher, K., Brodský, L., Du, C. W., Chappell, A., Fouad, Y., Genot, V., Gomez, C., Grunwald, S., Gubler, A., Guerrero, C., Hedley, C. B., Knadel, M., Morrás, H. J. M., Nocita, M., Ramirez-Lopez, L., Roudier, P., Campos, E. M. R., Sanborn, P., Sellitto, V. M., Sudduth, K. A., Rawlins, B. G., Walter, C., Winowiecki, L. A., Hong, S. Y., and Ji, W.: A global spectral library to characterize the world's soil, *Earth-Science Reviews*, 155, 198-230, <http://dx.doi.org/10.1016/j.earscirev.2016.01.012>, 2016.
- von Haden, A. C., Yang, W. H., and DeLucia, E. H.: Soils' dirty little secret: Depth-based comparisons can be inadequate for quantifying changes in soil organic carbon and other mineral soil properties, *Global Change Biology*, n/a, <https://doi.org/10.1111/gcb.15124>, 2020.
- Wang, M., Guo, X., Zhang, S., Xiao, L., Mishra, U., Yang, Y., Zhu, B., Wang, G., Mao, X., Qian, T., Jiang, T., Shi, Z., and Luo, Z.: Global soil profiles indicate depth-dependent soil carbon losses under a warmer climate, *Nature Communications*, 13, 5514, <https://doi.org/10.1038/s41467-022-33278-w>, 2022.
- Wang, M., Zhang, S., Guo, X., Xiao, L., Yang, Y., Luo, Y., Mishra, U., and Luo, Z.: Responses of soil organic carbon to climate extremes under warming across global biomes, *Nature Climate Change*, 14, 98-105, <https://doi.org/10.1038/s41558-023-01874-3>, 2024.
- WEPAL: ISE Reference Material - A list with all available ISE reference material samples, WEPAL (Wageningen Evaluating Programmes for Analytical Laboratories), Wageningen, 110 pp., <http://www.wepal.nl/website/products/RefMatISE.htm>, 2019.
- Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-W., da Silva Santos, L. B., Bourne, P. E., Bouwman, J., Brookes, A. J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C. T., Finkers, R., Gonzalez-Beltran, A., Gray, A. J. G., Groth, P., Goble, C., Grethe, J. S., Heringa, J., 't Hoen, P. A. C., Hooft, R., Kuhn, T., Kok, R., Kok, J., Lusher, S. J., Martone, M. E., Mons, A., Packer, A. L., Persson, B., Rocca-Serra, P., Roos, M., van Schaik, R., Sansone, S.-A., Schultes, E., Sengstag, T., Slater, T., Strawn, G., Swertz, M. A., Thompson, M., van der Lei, J., van Mulligen, E., Velterop, J., Waagmeester, A., Wittenburg, P., Wolstencroft, K., Zhao, J., and Mons, B.: The FAIR Guiding

