



Providing quality-assessed and standardised soil data to support global mapping and modelling (WoSIS snapshot 2023)

Niels H. Batjes¹, Luis Calisto¹, Luis Moreira de Sousa¹

¹ISRIC – World Soil Information, Droevendaalsesteeg 3, 6708 PB Wageningen, The Netherlands

5 *Correspondence:* Niels H. Batjes (niels.batjes@isric.org)

Abstract. Snapshots derived from the World Soil Information Service (WoSIS) are served freely to the international community. These static datasets provide quality-assessed and standardised soil profile data that can be used to support digital soil mapping and environmental applications at broad scale levels. Since the release of the preceding snapshot in 2019, new ETL (Extract, Load, Transform) procedures for screening, ingesting and standardising disparate source data have been developed. In conjunction with this, the WoSIS data model was overhauled making it compatible with the ISO 28258 and Observations and Measurements (O&M) domain models. Additional procedures for querying, serving, and downloading the publicly available standardised data have been implemented using open software (e.g. GraphQL API). Following up on a short discussion of these methodological developments we discuss the structure and content of the “WoSIS 2023-snapshot”. A range of new soil datasets was shared with us, registered in the ISRIC World Data Centre for Soils (WDC-Soils) data repository, and subsequently processed in accordance with the licences specified by the data providers. An important effort has been the processing of forest soil data collated in the framework of the EU-HoliSoils project. We paid special attention to the standardisation of soil property definitions, description of the soil analytical procedures, and standardisation of the units of measurement. The “2023 snapshot” considers the following soil chemical properties (total carbon, organic carbon, inorganic carbon (total carbonate equivalent), total nitrogen, phosphorus (extractable-P, total-P, and P-retention), soil pH, cation exchange capacity, and electrical conductivity) and physical properties (soil texture (sand, silt, and clay), bulk density, coarse fragments, and water retention), grouped according to analytical procedures that are operationally comparable. Method options are defined for each analytical procedure (e.g. pH measured in water, KCl or CaCl₂ solution, molarity of the solution, and soil/solution ratio). For each profile we also provide the original soil classification (i.e. FAO, WRB and USDA system with their version) and pedological horizon designations as far as these have been specified in the source databases. Three measures for “fitness-for-intended-use” are provided to facilitate informed data use: a) positional uncertainty of the profile’s site location, b) possible uncertainty associated with the operationally defined analytical procedures, and c) date of sampling. The most recent (i.e. *dynamic*) dataset, called *wosis_latest*, is freely accessible via various webservices. To permit consistent referencing and citation we also provide a *static* snapshot (in casu, December 2023). This snapshot comprises quality-assessed and standardised data for 228k geo-referenced profiles. The data come from 174 countries



and represent more than 900k soil layers (or horizons) and over 6 million records. The number of measurements for each soil property vary (greatly) between profiles and with depth, this generally depending on the objectives of the initial soil sampling programmes. In the coming years, we aim to gradually fill gaps in the geographic distribution of the profiles, as well as in the soil observations themselves, this subject to the sharing of a wider selection of “public” soil data by prospective data contributors.

5 The WoSIS 2023-snapshot is archived and freely available at <https://doi.org/10.17027/isric-wdcssoils-20231130> (Calisto et al., 2023).

1 Introduction

The World Soil Information service (WoSIS) draws on a large complement of soil profile data that have been shared by numerous data providers (ISRIC, 2023a). Nonetheless, a large proportion of the 800k “so-called” freely available soil profiles (see Arrouays et al., 2017), in practice, still remain “inaccessible” due to various licence constraints (e.g. Cornu et al., 2023). Soil data submitted for consideration in WoSIS come from a wide range of legacy holdings (e.g. traditional soil surveys), and increasingly include data derived from proximal sensing (e.g. Shepherd et al., 2022; Viscarra Rossel et al., 2016). The source data come in various formats and were determined according to a range of field sampling and soil analytical procedures, requiring standardisation and harmonisation during their ingestion/processing into WoSIS.

10
15

As discussed earlier (Batjes et al., 2020; Ribeiro et al., 2020), we follow the definition of harmonisation as defined by the Global Soil Partnership (GSP, Baritz et al., 2014). Harmonisation involves “providing mechanisms for the collation, analysis and exchange of consistent and comparable global soil data and information” and considers the following domains according to GSPs definition: a) soil description, classification and mapping, b) soil analyses, c) exchange of digital soil data, and d) interpretations.

20 In view of the breadth and magnitude of the task, as well as the limited availability of comparative “multiple analytical procedures” data sets as required for full harmonisation (Batjes, 2023; Bispo et al., 2021; van Leeuwen et al., 2022) we have limited ourselves to the standardisation of soil property definitions, soil analytical procedures descriptions, plausibility checks for soil observation values and the standardisation of measurement units for commonly required soil properties (see Appendix A). Importantly, users should always keep in mind that the source datasets themselves (e.g., Armas et al., 2023; NPDB, 2023; USDA-NCSS, 2021) will provide more detailed information than WoSIS albeit not in a consistent, globally standardised format.

25

Soil characteristics, such as texture, bulk density and organic carbon content, are collated according to a wide range of procedures in different countries. For such incongruent data to be interpreted correctly, the procedures for their collection, analysis, and reporting need to be well documented and understood. Results can differ when different analytical procedures are used even though these procedures may carry the same name (e.g. clay, silt and sand size fraction) or concept (see Soil Survey Staff, 2011). This makes the inter-comparison of different datasets difficult if it is not known how these data were

30



collected/analysed. Therefore we use “operational definitions”, as defined by USDA Soil Survey Staff (2011), for soil properties that are linked to specific analytical procedures. To properly characterise the “pH of a soil”, for example, we need information on sample pre-treatment, soil/solution ratio, and description of solution (e.g. H₂O, 1 M KCl, 0.02 M CaCl₂, or 1 M NaF). Soil pH measured in Sodium Fluoride (pH NaF), for example, provides a measure for the phosphorus (P) retention of a soil whereas pH measured in water (pH H₂O) is an indicator for soil nutrient status. Consequently, in WoSIS soil properties are named according to and defined by the analytical procedures and corresponding “method options”, based on common practice in soil science (e.g. BDFIOD for “bulk density (BD), fine earth fraction (FI), oven dry (OD)”).

This paper discusses methodological changes to the WoSIS workflow and new data additions since the release of the 2019-snapshot (Batjes et al., 2020). First, we describe the new data ingestion/screening process and indicate how the “shared” standardised data are being served to the user community. Thereafter, we describe the actual data screening, quality control and standardisation. In conjunction with this we provide three measures for “fitness-for-intended-use” of the standardised data. Subsequently, we describe the spatial distribution of soil profile sites and list the number of soil observations represented in the “WoSIS 2023-snapshot” (hereafter referred to as 2023-snapshot) Following up on a discussion concerning the scope for “full data harmonisation” in WoSIS, future developments are outlined.

The naming conventions and standard units of measurement are described in Appendix A and B, the file structure in Appendix C, the number of sites by country and continent in Appendix D, and their distribution by world terrestrial ecosystems respectively biomes in Appendix E.

Soils are important providers of ecosystem services. WoSIS-served data have been used for a range of applications, such as predictive soil property mapping (Guevara et al., 2018; Moulatlet et al., 2017; Nenkam et al., 2022; Poggio et al., 2021; Turek et al., 2023), space and time modelling of soil organic carbon stock change (Heuvelink et al., 2021), and a diverse range of environmental assessments (e.g., Luo et al., 2021; Lutz et al., 2019; Maire et al., 2015; Sanderman et al., 2017; Sothe et al., 2022). For example, based on the “2016 snapshot” and “2019 snapshot” respectively, Ivushkin et al. (2019) mapped global soil salinity change, while Wang et al. (2024) analysed responses of soil organic carbon under warming across global biomes. Ultimately, such information can help to inform the global conventions such as the UNCCD (United Nations Convention to Combat Desertification) and UNFCCC (United Nations Framework Convention on Climate Change), so that policymakers and business leaders can make informed decisions about the environment, biodiversity, and human well-being at an appropriate scale level.

2 WoSIS data model and workflow

2.1 Workflow



The original data model and workflow for acquiring, ingesting, processing and serving data, as described in Batjes *et al.* (2020), was overhauled. The main stages of the new workflow are visualised in Fig. 1: a) Data providers share their data with ISRIC WDC-Soils, b) the submitted data sets with associated metadata are screened for “completeness of information provided” (e.g. the licence defining access rights and description of terms and units) and, once considered adequate, subsequently stored “as is” in the WDC-Soils data repository (see “ISRIC Admin” in Fig. 1); and c) the source datasets are imported into the WoSIS PostgreSQL relational database, using newly developed ETL procedures. Step c) includes: c1) basic data quality assessment and control, c2) standardising descriptions for the soil analytical procedures and units of measurement, and c3) automated checks against plausibility limits for each soil observation, see Sect. 3 for details. Subsequently, d) distribution of the quality-assessed and standardised data via various services such as dashboards and WFS (OpenGIS web feature service) as well as a metadata catalogue service.




Figure 1. Schematic WoSIS workflow for ingesting, standardising and distributing soil profile data.

15 2.2 Data model

A new data model for WoSIS was developed, aligned where possible with the ISO 28258 domain model (de Sousa *et al.*, 2023) and the GloSIS web ontology (Palma *et al.*, 2023), both stemming from O&M (Observations and Measurement, Cox and David,



2011), all the while preserving legacy data (see <https://git.wur.nl/isric/databases/wosis-docs>, last access: 11 December 2023). Special attention was paid to the succinct description of the analytical procedures (see c2 above) using seven database tables, as summarised below:

- `thes_method_value`: Thesaurus of values that match the keys used to define an analytical method. For example, "natural
5 clod" for the "sample type" key for method "bulk density".
- `thes_method_key`: Thesaurus of keys used to define an analytical method. For example, "reported pH", "exchange
solution" and "index cation".
- `thes_method_option`: Encodes the possible combinations of key-value pairs for each numerical observation on layers.
Note that only a small sub-set of observations can be associated with particular method options.
- 10 • `method_source`: Analytical methods descriptions as defined in the respective source databases (i.e. prior to standardisation).
This table was imported "as is" from the old data model (Ribeiro et al., 2020) with the addition of a synthetic primary key. The
records in this table remain essential to identify the method referred by each result.
- `method_standard`: Distinguishes each source method by the particular observation to which it applies. It can be regarded
as a standardised description of the source method. Each record corresponds to a collection of key-value pairs in the
15 `method_option` table for a single observation. Results for numerical observations reference this table to identify the
corresponding analytical method.
- `method_option_standard`: Defines a many-to-many relationship between the `method_standard` and
`method_option` tables. It determines the exact collection of key-value option pairs that constitute a standard method. The
standard method is a specialisation of the source method for a specific observation.

20

2.3 ETL procedure

Extract, Transform and Load (ETL) is a standardised, semi-automatic process that guides the data processor during the ingestion
of new datasets. During the initial phase, newly shared datasets are submitted to a quick consistency check (i.e. format, data
model, metadata and licence) after which they are uploaded "as is" to a staging area in the WoSIS system. Subsequently, during
25 the transform stage the uploaded datasets are parsed by the system. During this process validation and standardisation occurs. In



case of (possible) unconformities the system will generate descriptive messages that guide the data processor towards possible actions that may be needed to resolve the flagged unconformities. The data processor then needs to correct these issues in conformity with the requirements of the WoSIS procedure manual in steps guided by the system; in some cases the original data providers may need to be consulted. At the end of this phase, the cleaned and standardised data remain in the staging area for
5 final verification by a soil expert. After this verification, the final stage of the ETL process, “Load”, can start. This is a fully automated process during which the cleaned and standardised data are copied into the WoSIS database and subsequently removed from the staging area. The newly ingested data can now be used to create a range of WoSIS-derived products (e.g., *wosis_latest*, *wosis_internal*, and dashboards, see Figure 1) in accord with the licences and possible restrictions specified by the data providers.

10 2.4 Soil properties standardised

We currently standardise the following soil properties (with associated analytical procedure descriptions): Bulk density (fine earth respectively for whole soil); Calcium carbonate equivalent (TCEQ); Cation exchange capacity; Coarse fragments; Electrical conductivity; Organic carbon (C); Organic matter; pH; Phosphorus (P); Soil texture (i.e. Sand, Silt and Clay size fraction); Total carbon (C); Total nitrogen (N); Water retention (gravimetric and volumetric, at defined tensions). Further details are provided in
15 Appendix A.

2.5 Data provisioning

Upon completion of the semi-automated ETL process, the quality-assessed and standardised data are distributed freely through
20 various channels (see Figure 1):

- As *wosis_latest* (dynamic) via WFS; the respective end-points are catalogued at the ISRIC Data Hub (ISRIC, 2023b).
- As “fixed” snapshots (in TSV format) with a unique digital object identifier (DOI) to permit consistent citation (e.g., Batjes et al., 2020).
- The contents of *wosis_latest* can also be visualised using a dashboard with some querying and zooming facilities (ISRIC, 25 2023c).
- Profile data from *wosis_latest* can also be queried through the “SoilGrids web platform” (ISRIC, 2020), which also provides access to a range of soil property maps derived from the WoSIS-served profile data and a set of environmental covariates using digital soil mapping (Poggio et al., 2021; Turek et al., 2023).



- The *wosis_latest* holdings can also be queried using a GraphQL interface (Calisto, 2023) that facilitates exploration of the data (e.g. select data for organic carbon, bulk density, proportion of coarse fragments per layer (horizon) for profiles located in a given geography). Results of such tailor-made queries can then be exported as input in scripting languages such as Python or R (R Core Team, 2021), for example to calculate regional carbon stocks.

5

2.6 Licence agreements

It is not a simple task to find potential providers of “open” soil data, (Arrouays et al., 2017; Batjes, 2009; Cornu et al., 2023). This may be due to technical issues, reasons for sharing (e.g., “Why share the data and for what purpose? What is in it for us?”) as well as legal requirements, as discussed by Bispo et al. (2021). All data sets that are shared with our centre are first registered
10 in the ISRIC Data Repository together with their metadata; data sharing agreements should align with the ISRIC Data Policy (ISRIC, 2016). During the subsequent WoSIS standardisation workflow, we are faced with three different types of datasets. First, those with a non-restrictive Creative Commons (CC-BY) licence, defined here as at least a CC-BY (Attribution) or CC-BY-NC (Attribution Non-Commercial) licence (these are later served as *wosis_latest*). Second, datasets with a “restrictive” licence in the
15 sense that they can exclusively be used for “visualisations”, such as SoilGridsTM (i.e. *wosis_internal*, see Figure 1), by ISRIC itself. The latter, generally because the coordinates cannot be disclosed based on our letter of agreement with some data providers. Hence, for third parties to get access to such “restricted” datasets they would have to approach the source data provider and make their own licencing arrangements. Third, several data sets have licences that stipulate that they should only be safeguarded in the ISRIC repository and cannot be used for any data processing (i.e. permanent embargo).

The number of profiles in WoSIS per licence category can be viewed using a [dashboard](#), and the number of profiles by
20 continent and country can be queried using a number of filters (ISRIC, 2023d). The majority of soil datasets from Europe represented in WoSIS, for example, have a “restricted” licence.

2.7 Number of profiles

The number of “public access” profiles served from WoSIS as snapshots increased from 96k in 2016 to 228k in 2023 (Table 1).
25 It should be noted here, however, that a large proportion of the forest soil data contributed in the framework of the EU-Holisoils project, for instance, could not be included in the “2023 snapshot” due to licence restrictions specified by the data providers. As a result, only 34k out of the total of 107k profiles “shared” with ISRIC between 2019 and 2023 could actually be included in the 2023-snapshot (resp. *wosis_latest*).



Table 1. Number of soil profiles and properties served in successive WoSIS snapshots.

Snapshot	No. of profiles	No. of properties ^a
2016-07	96k	22
2019-09	196k	45
2023-10	228k	45

^a Properties based on “operational definitions”, i.e. a combination of a property and procedure in the terminology of the WoSIS data model.

5

3 Data screening, quality control and standardisation

3.1 Consistency checks

Soil profile data shared for possible consideration in WoSIS were sampled and analysed according to various national or international standards and presented in various formats (from paper to digital). They are of varying degree of completeness as discussed below. To be considered in the WoSIS standardisation workflow (Fig. 1), each soil profile must meet several criteria as described earlier in Batjes et al (2020, p. 301). Summarising, they must be associated to a site correctly geo-referenced, have consistently defined upper and lower depths for each layer (or pedogenetic horizon), and have observations for at least some of the soil properties that are being served (e.g. sand, silt, clay and pH) as well as a succinct description of the analytical procedures and units of measurement. A soil (taxonomic) classification is considered desirable though not mandatory. Profiles associated to a valid site, for which only the classification is specified in the source data can still be useful for mapping of soil taxonomic classes. Similarly, profiles not associated to a valid site may still prove useful to develop pedotransfer functions; however, they cannot be served through WFS (because there is no geometry). The remaining cases are automatically excluded from the WoSIS ETL workflow.

Consistency in layer depth (i.e. sequential increase of the upper and lower depth reported for each layer down the profile) is checked using automated procedures (see Sect. 3.2). In line with current internationally accepted conventions, such depth



increments are given as “measured from the soil surface, including organic layers and mineral covers” (FAO, 2006; IUSS Working Group WRB, 2022; Schoeneberger et al., 2012; Soil Survey Staff, 2022a). Until 1993, however, the begin (zero datum) of the profile was set at the top of the mineral surface (the *solum* proper), except for “thick” organic layers as defined for peat soils (FAO-ISRIC, 1986; FAO, 1977). Organic horizons were recorded as above and mineral horizons recorded as below, relative to
5 the mineral surface (Schoeneberger et al., 2012, p. 2-6). As far as possible, such “organic_surface” layers are flagged in the snapshot (see C) so that they may be filtered-out during auxiliary computations of soil organic carbon stocks, for example.

3.2 Screening for duplicate profiles

In the early stage of WoSIS, many source databases were compilations of shared soil profile data necessitating intricate procedures
10 for identifying and flagging possibly repeated profiles (see Batjes et al., 2017; Ribeiro et al., 2020). Soil profiles located within 100 m of each other are flagged as possible duplicates, provided the year of sampling is identical (this criterion allows for reporting results of soil monitoring campaigns at the same *site*). Upon additional automated checks concerning the thickness of the first three soil layers (i.e. upper and lower depth), sand, silt and clay content, the duplicate profiles with the least-comprehensive component of observations are flagged and excluded from further processing (i.e. distribution). When still in doubt after these
15 rigorous tests a final visual “similarity check” is made with respect to other commonly reported soil properties such as pH_{water} and organic carbon content, possibly leading to the flagging (exclusion) of some additional profiles.

3.3 Standardisation of property names, analytical procedure descriptions and units of measurement

A crucial step during data ingestion is the standardisation of the, regularly non-English, soil property names used in the source
20 databases to the WoSIS conventions, as well as the standardisation of the soil analytical procedures according to consistent “operational definitions” (see Appendix A). Subsequently, the units of measurement are standardised, and the reported measurement values assessed according to soil observation-specific plausibility ranges for the respective soil properties (i.e. likely minimum and maximum). Data that do not meet these conditions are flagged and not processed further in the ETL workflow (see above), unless the observed “inconsistencies” can easily be solved (e.g. blatant typos in pH values). Alternatively, the data
25 provider(s) may be contacted to resolve the observed errors.

The following set of soil properties are considered in the 2023-snapshot:



- Chemical: total carbon (i.e. organic plus inorganic carbon), organic carbon, inorganic carbon (i.e. total carbonate equivalent), total nitrogen, soil pH, cation exchange capacity, electrical conductivity, and phosphorus (extractable-P, total-P, and P-retention),
- Physical: Soil texture (clay, silt, sand), coarse fragments, bulk density, and water retention.

5

All measurement values are served as recorded in the source data, after the above consistency checks and standardisation of the units of measurement to the target units (see Appendix A). As such, we *do not* apply any “gap filling” procedures during ETL nor do we apply any pedotransfer functions (PTF) to derive missing bulk density or soil hydrological properties, for example. This follow up stage of data processing is seen as the task of the data users (modellers) themselves. In practice, the required
10 functions or ways for depth-aggregating the layer data will be determined by the projected use(s) of the standardised data (see Finke, 2006; Heuvelink et al., 2021; Turek et al., 2023; van Leeuwen et al., 2024; Van Looy et al., 2017). It should be noted, however, that inadvertently some PTF-derived values (e.g. for bulk density) could have slipped through the above consistency checks in situations where procedures were mis-coded in the metadata of a source data set; critical modellers should exclude such values during their analyses.

15

3.4 Providing measures for fitness-for-intended-use

As indicated earlier, data served from WoSIS are used for a wide range of environmental applications (e.g., Guevara et al., 2018; Heuvelink et al., 2021; Luo et al., 2021; Maire et al., 2015 ; Moulatlet et al., 2017; Poggio et al., 2021; Sanderman et al., 2017; Sothe et al., 2022; Turek et al., 2023), but many of these assessments do not explicitly consider the uncertainties that are associated
20 with the data. However, it is well known that “soil observations used for calibration and interpolation are themselves not error-free” (e.g., Baroni et al., 2017; Cressie and Kornak, 2003; Folberth et al., 2016; Grimm and Behrens, 2010; Guevara et al., 2018; Hengl et al., 2017; Heuvelink, 2014; van Leeuwen et al., 2022). Therefore, we provide three measures for “fitness-for-intended-use” in *wosis_latest* namely: a) positional uncertainty of the profiles (i.e. site location), b) inferred accuracy of the laboratory measurements, and c) date of sampling. These three measures, although approximative, should be duly considered in digital soil
25 mapping and subsequent earth system modelling as they can affect the prediction uncertainty and “area-of-applicability” of the resulting derived products (Dai et al., 2019; Meyer and Pebesma, 2021; Shi et al., 2023).



3.4.1 Positional uncertainty

Profiles in WoSIS are georeferenced through the *site* in which they were sampled in accord with ISO 28258 standards (de Sousa et al., 2023). The coordinates themselves are presented according to the World Geodetic System datum ensemble (i.e. WGS84, EPSG code 4326) upon their conversion from a diverse range of national projections. For most profiles (86 %, see Table 2) the approximate positional uncertainty of the profile locations, as inferred from the coordinates given in the source datasets, is ~100 m. Typically, geo-referencing before the advent of GPS (Global Positioning Systems) in the 1970s is less accurate; often we just do not know the “true” accuracy. Nonetheless, digital soil mappers should be aware of this issue (Grimm and Behrens, 2010), because the soil observations and environmental covariates may not actually overlap (Cressie and Kornak, 2003), this both in space and time.

10

Table 2. Positional uncertainty of profile site locations.

Positional uncertainty	Number of profiles	
	n	%
~ 100 m	195,554	86
100 m - 1 km	21,653	9
1 km – 10 km	3,846	2
Over 10 km	7,037	3

3.4.2 Measurement uncertainty

Soil data managed in WoSIS have been analysed according to a diverse range of analytical procedures in multiple laboratories. A measure for measurement uncertainty is thus desired. Soil laboratory-specific Quality Management Systems and laboratory proficiency-testing (PT) can provide this type of information (GLOSOLAN, 2023; Magnusson and Örnemark, 2014; Munzert et al., 2007; NATP, 2015; WEPAL, 2019). Calculation of laboratory-specific measurement uncertainty for a single procedure, respectively multiple analytical procedures, will require several measurement rounds (years of observation) and solid statistical analyses (van Leeuwen et al., 2022). Generally, however, this type of information is not provided with the source data sets submitted to the ISRIC data repository. Therefore, pragmatically, we have distilled the required information from the PT-literature

20



(Al-Shammary et al., 2018; ICP Forests, 2021; Kalra and Maynard, 1991; Rayment and Lyons, 2011; Rossel and McBratney, 1998; van Reeuwijk, 1983; WEPAL, 2019), as far as technically feasible. In the case of organic carbon content, for example, the mean variability was 17 % (with a range of 12 to 42 %) and for “CEC buffered at pH 7” of 18 % (range 13 to 25%) when multiple laboratories analyse a standard set of reference materials using similar operational procedures (WEPAL, 2019).

5 The figures for measurement accuracy presented in Appendix A represent first approximations. They are derived from the inter-laboratory comparison of analyses on well-homogenised, reference samples for a still relatively small range of soil types. These indicative figures should be refined once specific, laboratory and procedure-related accuracy (i.e. systematic and random error) information is provided with the shared soil data, for instance using the procedures described by Eurachem (Magnusson and Örnemark, 2014). Alternatively, this type of information may be collated in the context of international laboratory PT-
10 networks such as GLOSOLAN and WEPAL, and in the framework of the ongoing LUCAS topsoil monitoring round (Bispo et al., 2021; Cornu et al., 2023). Meanwhile, the present first estimates can already be considered when calculating the uncertainty of predictive digital soil maps and of any interpretations derived from them (e.g. studies of soil organic carbon stock change).

Full harmonisation of analytical data derived from disparate sources, the ultimate ambition in WoSIS, will first become feasible once results of a representative set of multi-procedure, inter-laboratory comparison data sets become (freely) available,
15 as discussed by Baritz *et al.* (2014), Bispo *et al.* (2021) and Batjes (2023), and a common set of reference Standard Operating Procedures (SOPs) has been accepted as a global standard.

3.4.3 Year of sampling

For each profile site, the date of sampling has been recorded as far as documented in the source data. This information is important
20 to consider when superimposing the profile data with environmental co-variates, such as land cover, for example in the context of space and time analyses (Giller et al., 2006; Heuvelink et al., 2021). Most (54%) profiles represented in the snapshot were described/sampled between 1980 and 2020 (Table 3), and less than 4% before 1960. Alternatively, the date of site description and sampling is unknown for almost 26% of the profiles as the information was not provided in the source materials.

25



Table 3. Period of sampling/analysis.

Period	N of profiles	Percentage
< 1920	37	0
1920-1940	253	0.1
1940-1960	8,632	3.8
1960-1980	35,358	15.5
1980-2000	75,686	33.2
2000-2020	47,768	20.9
Not specified	60,356	26.5

4 Spatial distribution of soil profiles and number of observations

The 2023-snapshot of WoSIS includes standardised data for 228k profiles, characterising 217k different sites (Fig. 2). These are represented by 0.9 million soil layers (or horizons). In total, this corresponds with over 6.1 million records that include both numeric (e.g. silt content, soil pH, and cation exchange capacity) as well as class (e.g. WRB soil classification and horizon designation) properties. The naming conventions and standard units of measurement are described in Appendix A, and the file structure in Appendix C.




Figure 2. Distribution of sites represented in the 2023 snapshot of WoSIS (Good homolosine equal-area projection).

The greatest number of profiles come from north America (35 %) followed by Oceania (19%) and Europe (17%), while there are still few profiles for Asia (3%) and Antarctica (Table 4). The profiles come from sites in 174 countries. The average density of observations per 1000 km² varies greatly both between countries (Appendix D) and within each country. The highest densities of “shared” profiles are reported for Belgium (229 profiles per 1000 km²) and Switzerland (265 profiles per 1000 km²). There are still few profiles for Central Asia, Southeast Asia, Central and Eastern Europe, Russia, and the northern circumpolar region in the 2023-snapshot. Appendix E lists the number of profiles by biome (Olson et al., 2001b) respectively broad climatic region (Sayre et al., 2014), as derived from GIS overlays.

Table 4. Number of soil profiles per continent.

Continent	Number of profiles		
	2023-snapshot	2019-snapshot	2016-snapshot
Africa	32,198	27,688	17,153



Antarctica	35	9	0
Asia	7,763	6,704	3,089
Europe	39,728	35,311	1,908
North America	78,996	73,604	63,066
Oceania	43,013	42,918	235
South America	26,457	10,218	8,790

5 There are more observations for the chemical properties than the physical properties (see Appendix A) and the number of observations generally decreases with depth, this largely depending on the objectives of the original soil surveys. The interquartile range (Q1-Q3) for maximum depth of soil sampled in the field is 33-150 cm, with a median (Q2) of 100 cm (mean= 107 cm). It should be noted here that specific purpose surveys only consider the topsoil (e.g. soil fertility surveys), while others systematically sample soil layers up to depths exceeding 20 m (with a maximum of 32 m). When data from such “specific purpose surveys” (defined here as < 30 cm and >300 cm) are excluded, the figures for maximum depth sampled become: Q1= 90 cm, Q2= 122 cm, Q3= 155 cm with a mean of 126 cm. Unfortunately, we are not able to show the “depth to bedrock” as this information is seldom made explicit in the source databases.

10 It is our intention to fill present gaps in the geographic distribution (Appendix D and E) and range of soil properties (Appendix A) in the coming years. The degree to which this will be feasible, however, will mainly depend on the willingness or ability of data providers to share (some of) their data for consideration in WoSIS. For the northern Boreal and Arctic region, in principle, ISRIC can draw on new profiles collated by the International Soil Carbon Network (ISCN, see Malhotra et al., 2019). Alternatively, it should be reiterated that some datasets (e.g. EU LUCAS topsoil database, see Fernandez-Ugalde et al., 2022; ICP Forests, 2021) can *only* be standardised and used for SoilGrids™ applications due to current licence restrictions. Consequently, the corresponding profiles (~116k, see [dashboard](#)) are *not* shown in Figure 2, nor are they considered in Appendix A and B.



5 Distributing the standardised data

The standardised data are distributed through ISRIC's Spatial Data Infrastructure (SDI). The SDI is based on open source technologies and open web-services (WFS, WMS, WCS, CSW) following Open Geospatial Consortium (OGC) standards and aimed specifically at handling soil data. Our metadata are organised following standards of the International Organization for Standardization (ISO-19139, 2019) using GeoNetwork (see <https://data.isric.org>). The WoSIS database is hosted in a PostgreSQL database, with the spatial extension PostGIS. The PostgreSQL database itself is connected to MapServer to permit data download from GeoNetwork. These processes are aimed at facilitating global data interoperability and citation in compliance with FAIR principles. The data should be “findable, accessible, interoperable, and reusable” (Wilkinson et al., 2016).

Static snapshots are given a unique DOI (digital object identifier) to permit consistent citation. The 2023-snapshot is distributed in tab-separated values format and GeoPackage (<https://doi.org/10.17027/isric-wdcsoils-20231130>). Alternatively, the evolving *dynamic* version of the standardised data (i.e. *wosis_latest*) can be accessed/queried through the ISRIC Data Hub (<https://data.isric.org>) and the SoilGrids platform (<https://soilgrids.org>). Tutorials describing how to access *wosis_latest* from QGIS using WFS and with GraphQL (Calisto, 2023) can be found on the ISRIC website (see <https://www.isric.org/explore/wosis/faq-wosis>).

By its nature, the *dynamic* version will grow when new profile data are shared and processed, additional soil properties are considered in the WoSIS workflow, and/or when possible corrections are required. Potential errors can be reported via a “Google group” (<https://groups.google.com/forum/#!forum/isric-world-soil-information>, accessed 8/01/2024) so that these may be addressed in the *dynamic* version.

6 Discussion

We describe new procedures for handling and standardising disparate world soil profile data in WoSIS. The data model was fully harmonised to ISO 25828 and O&M requirements, with minor adjustments, and new ETL procedures were implemented. Alternatively, it should be stressed that the ultimate, desired full harmonisation of observations to an agreed reference analytical procedure Y, for example “pH H₂O, 1:2.5 soil/water solution” for say all “pH 1:x H₂O” measurements, will first become feasible once the target procedure (Y) for analysing each property has been defined, and subsequently accepted as “global standard” by the international soil community. A next step would be to collate/develop “comparative” data sets for each soil property (i.e. sets with samples analysed according to a given reference procedure (Y_i) and the corresponding national procedures (X_j)) for pedotransfer function development. These relationships, however, will often be soil type and region specific (GlobalSoilMap, 2015) and difficult to develop (i.e. calibrate and validate) when datasets for the comparisons do not yet exist or are simply not



freely shared/available (Batjes, 2023; Bispo et al., 2021; Cornu et al., 2023; van Leeuwen et al., 2024). Hence the importance of regional laboratory inter-comparison programmes, such as those undertaken in the framework of for example ANSIS (2023), GLOSOLAN (2023), ICP Forests (2021) and LUCAS (Bispo et al., 2021), that aim to develop consistent, context-specific (e.g. by country or land use/soil type) pedotransfer functions towards an agreed set of SOPs. However, it should be noted that the standard SOPs specified by these various programmes need not be comparable. In his context, Suvannang *et al.* (2018) observed that “comparable and useful soil information (at the global level) will only be attainable once laboratories agree to follow common standards and norms”. Over the years, however, many organisations/countries have implemented analytical procedures and quality assurance systems that are well suited for their specific purposes (e.g., Cornu et al., 2023; Orgiazzi et al., 2018; Soil Survey Staff, 2022b). Consequently, they may not be inclined to harmonise their data to a (still to be decided) set of global “reference” SOPs. However, procedures for such a full scale harmonisation will be required when developing a globally federated, and ultimately interoperable, spatial soil data infrastructure (GLOSIS, de Sousa et al., 2021) through which (harmonised) source data are served and updated by the respective data providers, and made queryable according to a common standard (de Sousa et al., 2023; OGC, 2019).

7 Data availability

The 2023-snapshot is archived for long-term storage at ISRIC – World Soil Information, the World Data Centre for Soils (WDC-Soils) of the ISC (International Council for Science) World Data System (WDS). It is freely accessible at <https://dx.doi.org/10.17027/isric-wdcsoils.20231130> (Calisto et al., 2023). The zip file (446 Gb) includes a “readme” file, the data in TSV format (see Appendix C) respectively in OGC GeoPackage format.

20

8 Conclusions

Bringing disparate soil profile data from different sources under a common global standard poses many and diverse challenges. A major improvement has been the harmonisation of the WoSIS data model to ISO 28258 and O&M domain specifications. In conjunction with this newly developed ETL procedures greatly improved the data ingestion and standardisation process, and new ways for visualising, querying and serving the data were developed to better serve our user community.

25

There are still numerous gaps in terms of geographic distribution as well as range of soil taxonomic units and/or soil properties represented. We aspire to address such gaps in future updates of ‘*wosis_latest*’. However, as World Data Centre, we are largely



dependent on the ability of soil data owners to share some of their data freely for the greater benefit of the international community. To facilitate this process, we are developing a web-based facility (front-end) to permit data providers to directly upload their soil data to WoSIS in a consistent format based on the newly developed ETL procedures. In return, upon their standardisation, we aim to provide each data provider with a tailor-made dashboard for viewing and querying the datasets they
5 shared, possibly with a DOI to facilitate citation.

Various sources of uncertainty are associated with the data. Therefore, we provide three measures for “fitness-for-intended-use” of the standardised data. This information, although coarse, should be duly considered by prospective users of the snapshot.

Unfortunately, numerous soil datasets worldwide are not yet freely accessible for various reasons. Standardised procedures, mechanisms and policies aimed at encouraging soil data sharing, by different categories of data owners/providers, need to be
10 developed, for example in the framework of the Global Soil Partnership.



8 Appendices

Appendix A: Coding conventions for observations (i.e. a combination of property, procedure and unit of measurement), number of profiles and layers provided in the WoSIS 2023-snapshot, and inferred accuracy of measurements (Codes are listed in alphabetical order).

5

Code	Property	Procedure ^a	Unit	Profiles	Layers	Accuracy (± %) ^b
BDFI33	Bulk density fine earth ^c	Bulk density of a soil sample that has been desorbed to 33 kPa (1/3 bar)	kg/dm ³	14886	78007	25.0
BDFIAD	Bulk density fine earth	Bulk density of a soil sample that has been air dried	kg/dm ³	4238	14485	25.0
BDFIFM	Bulk density fine earth	Bulk density of a soil sample at field-soil water content at time of sampling	kg/dm ³	5265	14075	25.0
BDFIOD	Bulk density fine earth	Bulk density of a soil sample that has been dried in an oven at 110 °C	kg/dm ³	26064	131623	25.0
BDWSAD	Bulk density whole soil ^c	Bulk density of a soil sample that has been air dried	kg/dm ³	0	0	25.0
BDWSOD	Bulk density whole soil	Bulk density of a soil sample that has been dried in an oven at 110 °C	kg/dm ³	14596	75397	25.0
CECPH7	Cation exchange capacity	CEC estimated by buffering the soil at "pH7" (e.g., NH ₄ Oac)	cmol(c)/kg	60339	320532	20.0
CECPH8	Cation exchange capacity	CEC estimated by buffering the soil at "pH7" (e.g., NH ₄ Oac)	cmol(c)/kg	6838	25100	20.0
CFGR	Coarse fragments	Gravimetric content of soil material larger than 2 mm ^c	g/100g	39481	202414	20.0
CFVO	Coarse fragments	Volumetric content of soil material larger than 2 mm ^c	cm ³ /100cm ³	48891	246580	30.0
CLAY	Clay ^d	Determination of total gravimetric content of clay-size fraction (for class-size limits and analytical methods see 'method_options')	g/100g	153319	652347	15.0
ECEC	Cation exchange capacity	Effective CEC conventionally approximated by summation of exchangeable bases (Ca ²⁺ , Mg ²⁺ , K ⁺ , and Na ⁺) plus 1 M KCl exchangeable acidity (Al ³⁺ and H ⁺) in acidic soils	cmol(c)/kg	35123	143693	25.0
ELCO20	Electrical conductivity	Electrical conductivity assessed on a 1:2 soil water extract. Used for saline soils.	dS/m	7971	44350	10.0
ELCO25	Electrical conductivity	Electrical conductivity assessed on a 1:2.5 soil water extract. Used for saline soils.	dS/m	4395	17825	10.0



ELCO50	Electrical conductivity	Electrical conductivity assessed on a 1:5 soil water extract. Used for saline soils.	dS/m	23121	90959	10.0
ELCOSP	Electrical conductivity	Electrical conductivity assessed on water saturated soil paste. Used for saline soils.	dS/m	22052	85020	10.0
NITKJD	Total nitrogen (N)	Kjeldahl wet-oxidation digestion procedure	g/kg	72905	240433	10.0
ORGC	Organic carbon (C)	Amount of organic carbon determined according to method specified under 'method_options'	g/kg	135655	526953	15.0
ORGM	Organic matter	Determination of organic compounds that accompany soil particles through a 2-mm sieve using loss-on-ignition (LOI) at about 400 degrees Celsius.	g/kg	3871	16282	15.0
PHAQ	pH	A measure of the acidity or alkalinity in soils, defined as the negative logarithm (base 10) of the activity of hydronium ions (H ⁺) in water ^a .	unitless	140326	655336	0.3
PHCA	pH	A measure of the acidity or alkalinity in soils, defined as the negative logarithm (base 10) of the activity of hydronium ions (H ⁺), in the specified CaCl ₂ solution.	unitless	69437	325153	0.3
PHKC	pH	A measure of the acidity or alkalinity in soils, defined as the negative logarithm (base 10) of the activity of hydronium ions (H ⁺), in the specified KCl solution.	unitless	38022	173464	0.3
PHNF	pH	A measure of the acidity or alkalinity in soils, defined as the negative logarithm (base 10) of the activity of hydronium ions (H ⁺), in the specified NaF solution.	unitless	4965	25409	0.3
PHETB1	Phosphorus (P)	Phosphorus determined according to the Bray-I method, a combination of HCl and NH ₄ -F to remove easily acid soluble P forms, largely Al- and Fe-phosphates (mainly applicable for acid soils)	mg/kg	10719	40379	40.0
PHETM3	Phosphorus (P)	Determined according to Mehlich-3 method, a weak acid soil extraction procedure that is considered suitable for removing P and other elements in acid and neutral soil. The extract is composed of 0.2 M glacial acetic acid, 0.25 M ammonium nitrate, 0.015 M ammonium fluoride, 0.013 M nitric acid, and 0.001 M ethylene diamine tetraacetic acid (EDTA).	mg/kg	1444	7230	25.0
PHETOL	Phosphorus (P)	Phosphorus determined according to the Olsen method (0.5 M sodium bicarbonate (NaHCO ₃) solution at a pH of 8.5); used extract P from calcareous, alkaline, and neutral soils.	mg/kg	4266	12291	25.0
PHPRTN	Phosphorus (P)	Phosphorus retention measured according to the New Zealand method (Blakemore, 1981).	g/100g	5599	26569	20.0
PHPTOT	Phosphorus (P)	Phosphorus determined with a "harsh" digest procedure to liberate and measure all forms of element.	mg/kg	7561	19310	15.0



PHPWSL	Phosphorus (P)	Phosphorus soluble in water	mg/kg	282	1241	15.0
SAND	Sand	Determination of total gravimetric content of sand-size fraction (for class-size limits and analytical methods see 'method_options').	g/100g	119127	542463	15.0
SILT	Silt	Determination of total gravimetric content of silt-size fraction (for class-size limits and analytical methods see 'method_options').	g/100g	145906	620790	15.0
TCEQ	Calcium carbonate equivalent (TCEQ)	Determination of the gravimetric loss of carbonates as carbon dioxide in the presence of excess hydrochloric acid. The quantity of carbonate (CO ₃) in the soil is expressed as CaCO ₃ and as a weight percentage of the less than 2 mm size fraction.	g/kg	59294	247368	10.0
TOTC	Total carbon (C)	Total C is quantified by two basic methods: wet or dry combustion (see 'method_options'). In total C determinations, all forms of C in a soil are converted to CO ₂ followed by a quantification of the evolved CO ₂ . Total C can be used to estimate the organic C content of a soil. The difference between total and inorganic C is an estimate of the organic C.	g/kg	33527	112787	10.0
WG0006	Water retention gravimetric	Water retention assessed at tension 6 kPa (see method options')	g/100g	827	3828	20.0
WG0010	Water retention gravimetric	Water retention assessed at tension 10 kPa (see 'method_options').	g/100g	2970	12517	20.0
WG0033	Water retention gravimetric	Water retention assessed at tension 33 kPa (see 'method_options').	g/100g	20994	94707	20.0
WG0100	Water retention gravimetric	Water retention assessed at tension 100 kPa (see 'method_options').	g/100g	687	3360	20.0
WG0200	Water retention gravimetric	Water retention assessed at tension 200 kPa (see 'method_options').	g/100g	4391	27773	20.0
WG0500	Water retention gravimetric	Water retention assessed at tension 500 kPa (see 'method_options').	g/100g	326	1414	20.0
WG1500	Water retention gravimetric	Water retention assessed at tension 1500 kPa (see 'method_options').	g/100g	33782	181999	20.0
WV0010	Water retention volumetric	Water retention assessed at tension 10 kPa (see 'method_options').	cm ³ /100cm ³	1914	6883	20.0
WV0033	Water retention volumetric	Water retention assessed at tension 33 kPa (see 'method_options').	cm ³ /100cm ³	7444	22291	20.0
WV0100	Water retention volumetric	Water retention assessed at tension 100 kPa (see 'method_options').	cm ³ /100cm ³	747	2553	20.0
WV0500	Water retention volumetric	Water retention assessed at tension 500 kPa (see 'method_options').	cm ³ /100cm ³	702	1758	20.0



WV1500	Water retention volumetric	Water retention assessed at tension 1500 kPa (see 'method_options').	cm ³ /100cm ³	7904	23331	20.0
--------	-------------------------------	-------------------------------------------------------------------------	-------------------------------------	------	-------	------

^a Method options for each analytical procedure are described in Batjes and van Oostrum (2023), and provided in file *Wosis_202312_xxx.tsv*, see Appendix C.

^b Inferred accuracy (or uncertainty), rounded to the nearest 5%, unless otherwise indicated (i.e. units for soil pH) as derived from the following sources (Al-Shammary et al., 2018; Kalra and Maynard, 1991; Rayment and Lyons, 2011; Rossel and McBratney, 1998; van Reeuwijk, 1983; WEPAL, 2019).

5 These figures are first approximations that should be fine-tuned once more specific results of laboratory proficiency tests, resp. national Soil Quality Management systems, become freely available (e.g. from the GLOSOLAN laboratory proficiency programme).

^c Generally, the fine earth fraction is defined as being < 2 mm. Alternatively, an upper limit of 1 mm was used in the former Soviet Union and its satellite states (Katchynsky scheme). The actual size limits are specified under “method options” (see Appendix C).

10 ^d Provided only when the sum of clay, silt and sand fraction is ≥ 90 and ≤ 100 percent (Note that users should normalise the totals to 100 percent before using them for mapping or modelling purposes).

^e No data are being served for this property because the associated licences are flagged as ‘restricted’ by the data providers.



Appendix B: Coding conventions and brief descriptions for soil classification, horizon designations and number of occurrences in the WoSIS 2023-snapshot.

Code	Description	Count
CSTX	Classification of the soil profile according to specified edition (year) of USDA Soil Taxonomy, at least at soil order level	31400
CWRB	Classification of the soil profile according to specified edition (year) of the World Reference Base for Soil Resources (WRB), at least at reference soil group level	39649
CFAO	Classification of the soil profile according to specified edition (year) of the FAO-Unesco Legend, at least at major group level	38792
HODS ^a	Horizon designations as provided in the source databases	80849 / 396522 ^b

5 ^a Where available, the “cleaned” (original) layer/horizon designation is provided for general information; these codes have not been standardised as they vary widely between different classification systems (Bridges, 1993; Gerasimova et al., 2013). When no horizon designations are provided in the source data bases, we have flagged all layers with an upper depth given as being negative (e.g. -10 to 0 cm that is using pre-1993 conventions (see Sect. 3.1) in the source databases as likely being a shallow “organic surface” layer above a mineral soil layer.

^b Number of profiles with horizon descriptions respectively total number of layers with horizon designations.

10

Appendix C: Structure of WoSIS 2023-snapshot

This Appendix describes the structure of the data files served with the WoSIS 2023-snapshot, namely
15 *README_wosis202312.pdf*, *wosis_202312_observations.tsv*, *wosis_20312_site.tsv*, *wosis_202312_profiles.tsv*,
wosis_202312_layers.tsv, and *wosis_202312_xxxx.tsv* (where “xxxx” is the name of the observation). The files are also distributed as OGC GeoPackage, which stores the files within an SQLite database.



- *Readme_wosis202312.pdf*. This file gives a short description of the contents of the snapshot with links to the corresponding documentation.

- *wosis_202312_observations.tsv*: This file lists the four to six letter codes for each observation, whether the observation is for a site/profile or layer (horizon), the unit of measurement and the number of profiles respectively layers represented in the snapshot. It also provides the inferred accuracy for the laboratory measurements (see Appendix A).

code	Code for the observation
property	Description of soil property
procedure	Description of analytical procedure
unit	Standard unit of measurement
profiles	Number of profiles that have at least one measurement for the observation
layers	Number of layers that have measurements for the observation
accuracy	Inferred accuracy of the laboratory measurements (First approximation, see Sect. 3.4.2)

15

- *Wosis_202312_site.tsv*: This file characterises the site location where profiles were sampled. The following field names are used:

site_id	Primary key
latitude	Latitude in degrees (WGS84)
longitude	Longitude in degrees (WGS84)
positional_uncertainty	Positional uncertainty of the profile's site location, expressed in four classes (see Table 2)
country_name	Name of country where site is located
region	Region in which site is located
continent	Continent in which site is located

25



- 5
- *wosis_202312_profiles.tsv*: Presents the unique profile ID (i.e. primary key), `site_id`, source of the data, country ISO code and name, positional uncertainty, latitude and longitude (WGS 1984), maximum depth of soil described and sampled, as well as information on the soil classification system and edition. Depending on the soil classification system used, the number of fields will vary. For example, for the World Soil Reference Base (WRB) system the options are: publication year (i.e. version), `reference_soil_group_code`, `reference_soil_group_name`, and the name(s) of the prefix (primary) qualifier(s) respectively suffix (supplementary) qualifier(s). The terms principal qualifier and supplementary qualifier are used since 2015 (IUSS Working Group WRB, 2015; 2022); earlier WRB versions used prefix and suffix for this (e.g. IUSS Working Group WRB, 2006). Alternatively, for USDA Soil Taxonomy, the version (year), order, suborder, great group, and subgroup can be accommodated (Soil Survey Staff, 2014). The following field names are used:
- 10

	<code>profile_id</code>	Primary key
	<code>profile_code</code>	Code for the profile
	<code>dataset_code</code>	Identifier for source data set
	<code>site_id</code>	Identifier for site where profile is located
15	<code>positional_uncertainty</code>	Positional uncertainty of the profile's site location, expressed in four classes (see Table 2).
	<code>country_name</code>	Name of country where site is located.
	<code>latitude</code>	Latitude in degrees (WGS84)
	<code>longitude</code>	Longitude in degrees (WGS84)
20	<code>wrb_reference_soil_group_code</code>	Code for WRB group (in given version of WRB)
	<code>wrb_reference_soil_group</code>	Full name for reference soil group
	<code>wrb_prefix_qualifiers</code>	Name for prefix (i.e. for WRB1988)
	<code>wrb_suffix_qualifiers</code>	Name for suffix (i.e. for WRB1988)
	<code>wrb_principal_qualifiers</code>	Name for principal qualifiers (i.e. for WRB 2015 and WRB 2022)



	wrb_supplementary_qualifiers	Name for supplementary qualifiers (i.e. for WRB 2015 and WRB 2022)
	wrb_publication_year	Version of World Reference Base for Soil Resources
	fao_major_group_code	Code for major group (in given version of the Legend),
	fao_major_group	Name of major group
5	fao_soil_unit_code	Code for soil unit
	fao_soil_unit	Name of soil unit
	fao_publication_year	Version of FAO Legend (e.g. 1974 or 1988)
	usda_order_name	Name of USDA Soil Taxonomy order
	usda_suborder	Name of USDA Soil Taxonomy suborder
10	usda_great_group	Name of USDA Soil Taxonomy greatgroup
	usda_subgroup	Name of USDA Soil Taxonomy subgroup
	usda_publication_year	Version of USDA Soil Taxonomy
	• <i>Wosis_202312_layers.tsv</i> . This file characterises the layers (or horizons) per profile:	
15	profile_id	Primary key
	layer_id	Sequential number for the layer (or horizon)
	profile_code	Code for the profile
	site_id	Identifier for site where profile is located
	layer_name	Name of pedogenetic horizon (“as is”)
20	upper_depth	Upper depth of layer



	lower_depth	Lower depth of layer
	layer_number	Sequential number for the layer (or horizon)
	organic_surface	Flag for the presence of an organic layer above the mineral soil
	dataset_id	Abbreviation for source data set (e.g. WD-ISCN)
5	licence	Licence for observation as indicated by the data provider (e.g. CC BY)
	• <i>Wosis_202312_xxxx.tsv</i> . For each observation (e.g. “xxxx” = “BDFIOD”), as defined under “code” in file <i>wosis_202312_observation.tsv</i> , the following are listed:	
	profile_id	Primary key
	layer_id	Primary key (number, sequential from top to bottom)
10	profile_code	Code for given profile
	layer_name	Name of pedogenetic horizon (“as is”)
	upper_depth	Upper depth of layer
	lower_depth	Lower depth of layer
	organic_surface	Indicates if there is an organic layer above the mineral surface
15	value	Array listing all measurement values for observation “xxxx” for the given layer. In some cases, more than one observation is reported for a given horizon (layer) in the source, for example four values for TOTC: [1:5.4, 2:8.2, 3:6.3, 4:7.7] (see <i>value_avg</i> below)
20	method_options	Array listing the method options for each analytical procedure as distilled from the source data. The content of this array varies with the soil observation under consideration as described in the method option table for each analytical procedure. For example, in the case of electrical conductivity (ELCO), the method options



include sample pretreatment (e.g. sieved over 2 mm size, solution (e.g. water), ratio (e.g. 1:5), and ratio base (e.g. weight /volume). For details see Batjes and van Oostrum (2023).

	value_avg	Average, for above (it is recommended to use this value for “routine” modelling)
5	dataset_id	Abbreviation for source data set (e.g. WD-ISCN)
	country_name	Name of country where site is located.
	latitude	Latitude in degrees (WGS84)
	longitude	Longitude in degrees (WGS84)
	positional_uncertainty	Positional uncertainty of the profile’s site location (see Table 2).
10	region	Region in which site is located
	continent	Continent where the profile’s site is located
	date	Date the profile was described/sampled
	licence	Licence for given data, as indicated by the data provider (i.e. CC BY or CC BY-NC)

15 *Format:* All fields in the above files are tab-delimited, with double quotation marks as text delimiters. File coding is according to the UTF-8 unicode transformation format.

Using the data: Tutorials for downloading and querying the data, using various platforms, are provided on the WoSIS FAQ webpage (<https://www.isric.org/explore/wosis/faq-wosis>, last access: 15 December 2023).



Appendix D: Number of sites by country and continent.

Continent	Country	Country code	No. of sites	Area (km ²)	Site density (per 1000 km ²)	
Africa	Abyei		4	0	9943	0
	Algeria	DZ	10	2308647	0.004	
	Angola	AO	1168	1246690	0.937	
	Benin	BJ	743	115247	6.447	
	Botswana	BW	994	578247	1.719	
	British Indian Ocean Territory	IO	0	49	0	
	Burkina Faso	BF	2023	273281	7.403	
	Burundi	BI	36	26857	1.34	
	Cameroon	CM	1417	465363	3.045	
	Cape Verde	CV	0	4056	0	
	Central African Republic	CF	88	619591	0.142	
	Chad	TD	7	1265392	0.006	
	Comoros	KM	0	1652	0	
	Congo	CG	70	340599	0.206	
	Côte d'Ivoire	CI	255	321762	0.793	
	Democratic Republic of the Congo	CD	378	2329162	0.162	
	Djibouti	DJ	0	21670	0	
	Egypt	EG	26	982161	0.026	
	Equatorial Guinea	GQ	0	27000	0	
	Eritrea	ER	0	120763	0	
	Ethiopia	ET	1712	1129314	1.516	
	Gabon	GA	47	264022	0.178	
	Gambia	GM	0	11203	0	
	Ghana	GH	432	238842	1.809	
	Guinea	GN	128	243023	0.527	
	Guinea-Bissau	GW	15	30740	0.488	
	Hala'ib triangle		10	0	17684	0
	Ilemi triangle		13	0	3179	0
	Kenya	KE	1603	582342	2.753	



Lesotho	LS	33	30453	1.084	
Liberia	LR	50	96103	0.52	
Libya	LY	14	1620583	0.009	
Madagascar	MG	130	588834	0.221	
Malawi	MW	3050	118715	25.692	
Mali	ML	885	1251471	0.707	
Ma'tan al-Sarra		11	0	1993	0
Mauritania	MR	13	1038527	0.013	
Mauritius	MU	0	2014	0	
Mayotte	YT	0	378	0	
Morocco	MA	113	414030	0.273	
Mozambique	MZ	565	787305	0.718	
Namibia	NA	1569	823989	1.904	
Niger	NE	520	1182602	0.44	
Nigeria	NG	1402	908978	1.542	
Réunion	RE	0	2504	0	
Rwanda	RW	1016	25388	40.018	
Saint Helena, Ascension and Tristan da Cunha	SH	0	399	0	
Sao Tome and Principe	ST	0	991	0	
Senegal	SN	312	196200	1.59	
Seychelles	SC	0	499	0	
Sierra Leone	SL	12	72281	0.166	
Somalia	SO	245	632562	0.387	
South Africa	ZA	879	1220127	0.72	
South Sudan	SS	82	629821	0.13	
Sudan	SD	130	1843196	0.071	
Swaziland	SZ	14	17290	0.81	
Togo	TG	9	56767	0.159	
Tunisia	TN	60	155148	0.387	
Uganda	UG	84	241495	0.348	
United Republic of Tanzania	TZ	1910	939588	2.033	
Western Sahara	EH	0	268617	0	



	Zambia	ZM	603	751063	0.803	
	Zimbabwe	ZW	413	390648	1.057	
Antarctica	Antarctica	AQ	30	12537967	0.002	
	Bouvet Island	BV	0	45	0	
	French Southern and Antarctic Territories	TF	0	7738	0	
	Heard Island and McDonald Islands	HM	0	412	0	
	South Georgia and the South Sandwich Islands	GS	0	3870	0	
Asia	Afghanistan	AF	19	641827	0.03	
	Aksai Chin		1	0	30666	0
	Armenia	AM	509	29624	17.182	
	Arunachal Pradesh		2	2	67965	0.029
	Azerbaijan	AZ	28	164780	0.17	
	Bahrain	BH	2	673	2.97	
	Bangladesh	BD	207	139825	1.48	
	Bhutan	BT	85	37674	2.256	
	Brunei Darussalam	BN	0	5899	0	
	Cambodia	KH	424	181424	2.337	
	China	CN	1644	9345214	0.176	
	China/India		3	0	3526	0
	Christmas Island	CX	0	136	0	
	Cocos (Keeling) Islands	CC	0	16	0	
	Cyprus	CY	12	9249	1.297	
	Democratic People's Republic of Korea	KP	0	122465	0	
	Georgia	GE	18	69785	0.258	
	Hong Kong	HK	2	1081	1.851	
	India	IN	199	2961118	0.067	
	Indonesia	ID	179	1888620	0.095	
	Iran (Islamic Republic of)	IR	2010	1677319	1.198	
	Iraq	IQ	14	435864	0.032	
	Israel	IL	17	20720	0.82	



Jammu and Kashmir		12	4	186035	0.022
Japan	JP		197	373651	0.527
Jordan	JO		47	89063	0.528
Kazakhstan	KZ		52	2841103	0.018
Kuril islands		5	0	4996	0
Kuwait	KW		1	17392	0.057
Kyrgyzstan	KG		1	199188	0.005
Lao People's Democratic Republic	LA		20	230380	0.087
Lebanon	LB		10	10136	0.987
Macau	MO		0	17	0
Malaysia	MY		155	329775	0.47
Maldives	MV		0	223	0
Mongolia	MN		9	1564529	0.006
Myanmar	MM		0	667085	0
Nepal	NP		142	147437	0.963
Occupied Palestinian Territory	PS		18	6225	2.892
Oman	OM		11	308335	0.036
Pakistan	PK		45	788439	0.057
Paracel Islands		6	0	8	0
Philippines	PH		78	296031	0.263
Qatar	QA		0	11549	0
Republic of Korea	KR		23	99124	0.232
Saudi Arabia	SA		7	1925621	0.004
Scarborough Reef		7	0	44	0
Senkaku Islands		8	0	5	0
Singapore	SG		1	594	1.683
Spratly Islands		9	0	1	0
Sri Lanka	LK		73	66173	1.103
Syrian Arab Republic	SY		69	188128	0.367
Taiwan	TW		35	36127	0.969
Tajikistan	TJ		5	142004	0.035
Thailand	TH		479	515417	0.929
Timor-Leste	TL		0	14892	0



	Turkey	TR	69	781229	0.088
	Turkmenistan	TM	0	555052	0
	United Arab Emirates	AE	12	71079	0.169
	Uzbekistan	UZ	9	449620	0.02
	Viet Nam	VN	29	327575	0.089
	Yemen	YE	284	453596	0.626
Europe	Albania	AL	97	28682	3.382
	Andorra	AD	0	475	0
	Austria	AT	128	83964	1.524
	Belarus	BY	96	207581	0.462
	Belgium	BE	7013	30669	228.667
	Bosnia and Herzegovina	BA	32	51145	0.626
	Bulgaria	BG	134	111300	1.204
	Croatia	HR	78	56589	1.378
	Czech Republic	CZ	666	78845	8.447
	Denmark	DK	72	44458	1.619
	Estonia	EE	241	45441	5.304
	Faroe Islands	FO	0	1400	0
	Finland	FI	442	336892	1.312
	France	FR	3183	548785	5.8
	Germany	DE	4362	357227	12.211
	Gibraltar	GI	0	6	0
	Greece	GR	374	132549	2.822
	Guernsey	GG	0	79	0
	Holy See	VA	0	0	0
	Hungary	HU	1421	93119	15.26
	Iceland	IS	17	102566	0.166
	Ireland	IE	124	69809	1.776
	Isle of Man	IM	0	573	0
	Italy	IT	576	301651	1.909
	Jersey	JE	0	120	0
	Latvia	LV	102	64563	1.58
	Liechtenstein	LI	0	151	0



	Lithuania	LT	127	64943	1.956
	Luxembourg	LU	142	2621	54.184
	Malta	MT	0	316	0
	Monaco	MC	0	8	0
	Montenegro	ME	12	13776	0.871
	Netherlands	NL	958	35203	27.214
	Norway	NO	507	324257	1.564
	Poland	PL	796	311961	2.552
	Portugal	PT	455	91876	4.952
	Republic of Moldova	MD	35	33798	1.036
	Romania	RO	113	238118	0.475
	Russian Federation	RU	1464	16998830	0.086
	San Marino	SM	0	60	0
	Serbia	RS	69	88478	0.78
	Slovakia	SK	161	49072	3.281
	Slovenia	SI	67	20320	3.297
	Spain	ES	907	505752	1.793
	Svalbard and Jan Mayen Islands	SJ	4	63464	0.063
	Sweden	SE	594	449212	1.322
	Switzerland	CH	10928	41257	264.874
	The Republic of North Macedonia	MK	20	25424	0.787
	Ukraine	UA	462	600526	0.769
	United Kingdom	GB	1727	244308	7.069
Northern America	Anguilla	AI	0	79	0
	Antigua and Barbuda	AG	0	452	0
	Aruba	AW	0	180	0
	Bahamas	BS	0	11904	0
	Barbados	BB	3	433	6.928
	Belize	BZ	26	21764	1.195
	Bermuda	BM	0	63	0
	British Virgin Islands	VG	0	154	0
	Canada	CA	8778	9875646	0.889
	Cayman Islands	KY	0	269	0



Clipperton Island	CP	0	9	0	
Costa Rica	CR	560	51042	10.971	
Cuba	CU	53	110863	0.478	
Dominica	DM	0	751	0	
Dominican Republic	DO	10	48099	0.208	
El Salvador	SV	38	20732	1.833	
Greenland	GL	2	2165159	0.001	
Grenada	GD	0	318	0	
Guadeloupe	GP	5	1697	2.947	
Guatemala	GT	28	109062	0.257	
Haiti	HT	0	27022	0	
Honduras	HN	38	112124	0.339	
Jamaica	JM	74	10965	6.749	
Martinique	MQ	0	1104	0	
Mexico	MX	12599	1949527	6.463	
Montserrat	MS	0	101	0	
Netherlands Antilles	AN	4	790	5.066	
Nicaragua	NI	21	128376	0.164	
Panama	PA	50	74850	0.668	
Puerto Rico	PR	280	8937	31.329	
Saint Kitts and Nevis	KN	0	262	0	
Saint Lucia	LC	0	603	0	
Saint Pierre and Miquelon	PM	0	233	0	
Saint Vincent and the Grenadines	VC	0	427	0	
Trinidad and Tobago	TT	2	5144	0.389	
Turks and Caicos Islands	TC	0	530	0	
United States Minor Outlying Islands	UM	0	348	0	
United States of America	US	56322	9315946	6.046	
United States Virgin Islands	VI	46	352	130.555	
Oceania	American Samoa	AS	0	200	0
Australia	AU	42767	7687634	5.563	
Cook Islands	CK	0	241	0	



	Fiji	FJ	6	18293	0.328
	French Polynesia	PF	0	3967	0
	Guam	GU	15	544	27.579
	Kiribati	KI	0	1020	0
	Marshall Islands	MH	0	268	0
	Micronesia (Federated States of)	FM	75	740	101.343
	Nauru	NR	0	22	0
	New Caledonia	NC	2	18574	0.108
	New Zealand	NZ	52	270415	0.192
	Niue	NU	0	263	0
	Norfolk Island	NF	0	38	0
	Northern Mariana Islands	MP	0	476	0
	Palau	PW	18	451	39.924
	Papua New Guinea	PG	24	462230	0.052
	Pitcairn Islands	PN	0	49	0
	Samoa	WS	18	2835	6.349
	Solomon Islands	SB	1	28264	0.035
	Tokelau	TK	0	15	0
	Tonga	TO	0	700	0
	Tuvalu	TV	0	48	0
	Vanuatu	VU	1	12236	0.082
	Wake Island	WK	0	7	0
	Wallis and Futuna Islands	WF	0	142	0
South America	Argentina	AR	253	2780175	0.091
	Bolivia (Plurinational State of)	BO	87	1084491	0.08
	Brazil	BR	9262	8485946	1.091
	Chile	CL	13662	753355	18.135
	Colombia	CO	236	1137939	0.207
	Ecuador	EC	94	256249	0.367
	Falkland Islands (Malvinas)	FK	0	12084	0
	French Guiana	GF	30	83295	0.36
	Guyana	GY	43	211722	0.203
	Paraguay	PY	2	399349	0.005



Peru	PE	158	1290640	0.122
Suriname	SR	31	145100	0.214
Uruguay	UY	136	177811	0.765
Venezuela (Bolivarian Republic of)	VE	204	912025	0.224

^a Disputed territory. Country names and areas are based on the Global Administrative Layers (GAUL) database, see: <https://data.apps.fao.org/map/catalogsrv/eng/catalog.search?id=12691#/metadata/9c35ba10-5649-41c8-bdfc-eb78e9e65654>; last access: 5 December 2023.



Appendix E. Distribution of sites by world terrestrial ecosystems and by WWF biomes

A) Number of sites by world terrestrial ecosystems (WTE)^a

Temperature zone	Moisture zone	No. of sites	Percent (%)
Polar	Dry	224	0.1
Polar	Moist	532	0.2
Boreal	Dry	1789	0.8
Boreal	Moist	3398	1.6
Cool Temperate	Desert	25	0
Cool Temperate	Dry	10968	5
Cool Temperate	Moist	53245	24.5
Warm Temperate	Desert	238	0.1
Warm Temperate	Dry	29209	13.4
Warm Temperate	Moist	46533	21.4
Sub Tropical	Desert	296	0.1
Sub Tropical	Dry	25748	11.8
Sub Tropical	Moist	17906	8.2
Tropical	Desert	178	0.1
Tropical	Dry	11315	5.2
Tropical	Moist	11095	5.1
No data	-	4674	2.2

5 ^a World Terrestrial Ecosystems (WTE) as defined by Sayre (2022). Total may differ from 100% due to rounding.



B) Number of sites by WWF biome^b

WWF biome	No. of sites	Percent (%)
Boreal Forests/Taiga	5519	2.5
Deserts and Xeric Shrublands	13410	6.2
Flooded Grasslands and Savannas	792	0.4
Lakes	85	0
Mangroves	765	0.4
Mediterranean Forests, Woodlands, and Scrub	24459	11.3
Montane Grasslands and Shrublands	2796	1.3
Rock and Ice	20	0
Temperate Broadleaf and Mixed Forests	74068	34.1
Temperate Coniferous Forests	14436	6.6
Temperate Grasslands, Savannas, and Shrublands	23890	11
Tropical and Subtropical Coniferous Forests	2363	1.1
Tropical and Subtropical Dry Broadleaf Forests	4120	1.9
Tropical and subtropical grasslands, savannas, and shrublands	31376	14.4
Tropical and Subtropical Moist Broadleaf Forests	16478	7.6
Tundra	2072	1
No data	724	0.3

^a Biomes defined according to “Terrestrial Ecoregions of the World” (WWF) (Olson et al., 2001a). Total may differ from 100% due to rounding.



Author contributions. NB is scientific lead of the WoSIS project and wrote the first draft. LC developed the ETL and GraphQL procedures while LdS developed the new data model. All authors performed quality checks, data analyses and contributed to the writing/editing of the final manuscript.

Competing interests. The authors declare that they have no conflict of interest.

5 **Disclaimer.** ISRIC – World Soil Information remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Acknowledgements

The development of WoSIS has been made possible thanks to the contributions and shared knowledge of a steadily growing number of data providers, including soil survey organisations, research institutes and individual experts, for which we are grateful.

10 Regrettably, we can impossibly acknowledge all contributors (e.g., field surveyors, laboratory personnel, soil experts and database experts) individually. Therefore, we do this largely in a generic way (see <https://www.isric.org/explore/wosis/wosis-contributing-institutions-and-experts>; last access: 15 December 2023).

Our special thanks go to Eloi Ribeiro, former WoSIS database management expert at ISRIC, for his sustained support and advice during the transition from the former data ingestion and standardisation scripts to the new ETL procedures. We also thank
15 our colleague Laura Poggio for useful methodological discussions concerning methodological linkages between WoSIS and SoilGrids.

The ETL procedures and new data model were co-developed in the framework of the European Union’s Horizon 2020 HoliSoils project (Grant agreement 101000289) and ISRIC’s “global soil information & standards” workstream.

ISRIC – World Soil Information, legally registered as International Soil Reference and Information Centre, receives core
20 funding from the Dutch Government.



References

- Al-Shammary, A. A. G., Kouzani, A. Z., Kaynak, A., Khoo, S. Y., Norton, M., and Gates, W.: Soil Bulk Density Estimation Methods: A Review, *Pedosphere*, 28, 581-596, [https://doi.org/10.1016/S1002-0160\(18\)60034-7](https://doi.org/10.1016/S1002-0160(18)60034-7), 2018.
- 5 ANSIS: Australian National Soil Information System, Australian National Soil Information System, Canberra (AU), <https://ansis.net/>, 2023.
- Armas, D., Guevara, M., Bezares, F., Vargas, R., Durante, P., Osorio, V., Jiménez, W., and Oyonarte, C.: Harmonized Soil Database of Ecuador (HESD): data from 2009 to 2015, *Earth Syst. Sci. Data*, 15, 431-445, <https://doi.org/10.5194/essd-15-431-2023>, 2023.
- 10 Arrouays, D., Leenaars, J. G. B., Richer-de-Forges, A. C., Adhikari, K., Ballabio, C., Greve, M., Grundy, M., Guerrero, E., Hempel, J., Hengl, T., Heuvelink, G., Batjes, N., Carvalho, E., Hartemink, A., Hewitt, A., Hong, S.-Y., Krasilnikov, P., Lagacherie, P., Lelyk, G., Libohova, Z., Lilly, A., McBratney, A., McKenzie, N., Vasquez, G. M., Leatitia Mulder, V., Minasny, B., Luca, M., Odeh, I., Padarian, J., Poggio, L., Roudier, P., Saby, N., Savin, I., Searle, R., Solbovoy, V., Thompson, J., Smith, S., Sulaeman, Y., Vintila, R., Rossel, R. V., Wilson, P., Zhang, G.-L., Swerts, M., Oorts, K., Karklins, A., Feng, L., Ibelles Navarro, A. R., Levin, A., Laktionova, T., Dell'Acqua, M., Suvannang, N., Ruam, W., Prasad, J., Patil, N., Husnjak, S., Pasztor, L., Okx, J., Hallet, S., Keay, C., Farewell, T., Lilja, H., Juilleret, J., Marx, S., Takata, Y., Kazuyuki, Y., Mansuy, N., Panagos, P., Van Liedekerke, M., Skalsky, R., Sobocka, J., Kobza, J., Eftekhari, K., Kacem Alavipanah, S., Moussadek, R., Badraoui, M., Da Silva, M., Paterson, G., da Conceicao Gonsalves, M., Theocharopoulos, S., Yemefack, M., Tedou, S., Vrscaj, B., Grob, U., Kozak, J., Boruvka, L., Dobos, E., Taboada, M., Moretti, L., and Rodriguez, D.: Soil legacy data rescue via GlobalSoilMap and other international and national initiatives, *GeoResJ*, 14, 1-19, <https://doi.org/10.1016/j.grj.2017.06.001>, 2017.
- 20 Baritz, R., Erdogan, H., Fujii, K., Takata, Y., Nocita, M., Bussian, B., Batjes, N. H., Hempel, J., Wilson, P., and Vargas, R.: Harmonization of methods, measurements and indicators for the sustainable management and protection of soil resources (Providing mechanisms for the collation, analysis and exchange of consistent and comparable global soil data and information), *Global Soil Partnership, FAO*, 44 pp., <http://www.fao.org/3/a-az922e.pdf>, 2014.
- 25 Baroni, G., Zink, M., Kumar, R., Samaniego, L., and Attinger, S.: Effects of uncertainty in soil properties on simulated hydrological states and fluxes at different spatio-temporal scales, *Hydrology and Earth System Sciences*, 21, 2301-2320, <https://doi.org/10.5194/hess-21-2301-2017>, 2017.
- Batjes, N. H.: Harmonized soil profile data for applications at global and continental scales: updates to the WISE database, *Soil Use and Management*, 25, 124-127, 2009.
- 30 Batjes, N. H., Ribeiro, E., van Oostrum, A., Leenaars, J., Hengl, T., and Mendes de Jesus, J.: WoSIS: providing standardised soil profile data for the world, *Earth Syst. Sci. Data*, 9, 1-14, <http://dx.doi.org/10.5194/essd-9-1-2017>, 2017.
- Batjes, N. H., Ribeiro, E., and van Oostrum, A.: Standardised soil profile data to support global mapping and modelling (WoSIS snapshot 2019), *Earth Syst. Sci. Data*, 12, 299-320, <https://doi.org/10.5194/essd-12-299-2020>, 2020.
- 35 Batjes, N. H.: Options for harmonising soil data obtained from different sources [HoliSoils project - H2020 Grant Agreement No 101000289], *ISRIC - World Soil Information, Wageningen*, 21 pp., <https://dx.doi.org/10.17027/isric-wdc-6ztd-eb19> 2023.
- Batjes, N. H., and van Oostrum, A. J. M.: WoSIS Procedures for standardizing soil analytical method descriptions, *ISRIC - World Soil Information, Wageningen*, 46 pp., <https://doi.org/10.17027/isric-1dq0-1m83>, 2023.
- 40 Bispo, A., Arrouays, D., Saby, N., Boulonne, L., and Fantappiè, M.: Proposal of methodological development for the LUCAS programme in accordance with national monitoring programmes. Towards climate-smart sustainable management of agricultural soils (EU H2020-SFS-2018-2020 / H2020-SFS-2019) *EJP Soil*, 135 pp., https://ejpsoil.eu/fileadmin/projects/ejpsoil/WP6/EJP_SOIL_Deliverable_6.3_Dec_2021_final.pdf, 2021.
- Bridges, E. M.: Soil horizon designations: past use and future prospects, *CATENA*, 20, 363-373, [https://doi.org/10.1016/S0341-8162\(05\)80002-5](https://doi.org/10.1016/S0341-8162(05)80002-5), 1993.



- Calisto, L.: ISRIC GraphQL web services for WoSIS and ISIS data access, ISRIC - World Soil Information, Wageningen, <https://graphql.isric.org/>, 2023.
- Calisto, L., de Souza, L. M., and Batjes, N. H.: Standardised soil profile data for the world (WoSIS, December snapshot) [Dataset], ISRIC - World Soil Information, Wageningen, <https://doi.org/10.17027/isric-wdcsoils-20231130>, 2023.
- 5 Cornu, S., Keesstra, S., Bispo, A., Fantappie, M., van Egmond, F., Smreczak, B., Wawer, R., Pavlů, L., Sobocká, J., Bakacsi, Z., Farkas-Iványi, K., Molnár, S., Møller, A. B., Madenoglu, S., Feiziene, D., Oorts, K., Schneider, F., Gonçalves, M. d. C., Mano, R., Garland, G., Skalský, R., O'Sullivan, L., Kasparinskis, R., and Chenu, C.: National soil data in EU countries, where do we stand?, *European Journal of Soil Science*, e13398, <https://doi.org/10.1111/ejss.13398>, 2023.
- Cox, S., and David, J.: ISO 19156:2011 Geographic information – Observations and measurements International Organization for Standardization., <https://www.iso.org/standard/32574.html>, 2011.
- 10 Cressie, N., and Kornak, J.: Spatial statistics in the presence of location error with an application to remote sensing of the environment, *Statistical Science*, 18, 436-456, <https://doi.org/10.1214/ss/1081443228>, 2003.
- Dai, Y., Shangguan, W., Wang, D., Wei, N., Xin, Q., Yuan, H., Zhang, S., Liu, S., and Yan, F.: A review on the global soil datasets for earth system models, *SOIL*, 5, 137-158, <https://doi.org/10.5194/soil-5-137-2019>, 2019.
- 15 de Sousa, L., Kempen, B., Mendes de Jesus, J., Yigini, Y., Viatkin, K., Medyckyj-Scott, D., Richie, D. A., Wilson, P., van Egmond, F., and Baritz, R.: Conceptual design of the Global Soil Information System infrastructure, Rome, FAO and ISRIC, Wageningen, Netherlands, 30 pp., <http://www.fao.org/3/cb4355en/cb4355en.pdf>, 2021.
- de Sousa, L. M., Calisto, L., van Genuchten, P., Turdukulov, U., and Kempen, B.: Data model for the ISO 28258 domain model, ISRIC - World Soil Information, <https://iso28258.isric.org/>, 2023.
- 20 FAO-ISRIC: Guidelines for soil description (3rd Edition, Rev.), FAO, Rome, 70 pp., 1986.
- FAO: Guidelines for the description of soils, FAO, Rome, 66 pp., 1977.
- FAO: Guidelines for soil description (Fourth ed.), FAO, Rome, 97 pp., <http://www.fao.org/docrep/019/a0541e/a0541e.pdf>, 2006.
- Fernandez-Ugalde, O., Scarpa, S., Orgiazzi, A., Panagos, P., Van Liedekerke, M., A., M., and Jones, A.: LUCAS 2018 Soil Module. Presentation of dataset and results, Publications Office of the European Union, Luxembourg, 128 pp., <https://dx.doi.org/10.2760/215013>, 2022.
- 25 Finke, P.: Quality assessment of digital soil maps: producers and users perspectives, in: *Digital soil mapping: An introductory perspective*, edited by: Lagacherie, P., McBratney, A., and Voltz, M., Elsevier, Amsterdam, 523-541, 2006.
- Folberth, C., Skalsky, R., Moltchanova, E., Balkovic, J., Azevedo, L. B., Obersteiner, M., and van der Velde, M.: Uncertainty in soil data can outweigh climate impact signals in global crop yield simulations, *Nature Communications*, 7, <https://doi.org/10.1038/ncomms11872>, 2016.
- 30 Gerasimova, M. I., Lebedeva, I. I., and Khitrov, N. B.: Soil horizon designation: State of the art, problems, and proposals, *Eurasian Soil Science*, 46, 599-609, <https://dx.doi.org/10.1134/S1064229313050037>, 2013.
- Giller, K. E., Rowe, E. C., de Ridder, N., and van Keulen, H.: Resource use dynamics and interactions in the tropics: Scaling up in space and time, *Agricultural Systems*, 88, 8-27, <https://doi.org/10.1016/j.agsy.2005.06.016>, 2006.
- 35 GlobalSoilMap: Specifications Tiered GlobalSoilMap products (Release 2.4), 52 pp., <https://www.isric.org/documents/document-type/globalsoilmap-specifications-v24-07122015>, 2015.
- GLOSOLAN: GLOSOLAN best practice manual (on-line), FAO, GSP, Rome, <https://www.fao.org/global-soil-partnership/glosolan-old/soil-analysis/standard-operating-procedures/en/#c763834>, 2023.
- 40 Grimm, R., and Behrens, T.: Uncertainty analysis of sample locations within digital soil mapping approaches, *GEODERMA*, 155, 154-163, <https://doi.org/10.1016/j.geoderma.2009.05.006>, 2010.
- Guevara, M., Olmedo, G. F., Stell, E., Yigini, Y., Aguilar Duarte, Y., Arellano Hernández, C., Arévalo, G. E., Arroyo-Cruz, C. E., Bolivar, A., Bunning, S., Bustamante Cañas, N., Cruz-Gaistardo, C. O., Davila, F., Dell Acqua, M., Encina, A., Figueredo Tacona, H., Fontes, F., Hernández Herrera, J. A., Ibelle Navarro, A. R., Loayza, V., Manueles, A. M., Mendoza Jara, F., Olivera, C., Osorio Hermosilla, R., Pereira, G., Prieto, P., Alexis Ramos, I., Rey Brina, J. C., Rivera, R., Rodríguez-
- 45



- Rodríguez, J., Roopnarine, R., Rosales Ibarra, A., Rosales Riveiro, K. A., Schulz, G. A., Spence, A., Vasques, G. M., Vargas, R. R., and Vargas, R.: No Silver Bullet for Digital Soil Mapping: Country-specific Soil Organic Carbon Estimates across Latin America, *SOIL*, 2018, 173-193, <https://doi.org/10.5194/soil-4-173-2018>, 2018.
- 5 Hengl, T., Mendes de Jesus, J., Heuvelink, G. B. M., Ruiperez Gonzalez, M., Kilibarda, M., Blagotić, A., Shangguan, W., Wright, M. N., Geng, X., Bauer-Marschallinger, B., Guevara, M. A., Vargas, R., MacMillan, R. A., Batjes, N. H., Leenaars, J. G. B., Ribeiro, E., Wheeler, I., Mantel, S., and Kempen, B.: SoilGrids250m: Global gridded soil information based on machine learning, *PLoS ONE*, 12, e0169748, <http://dx.doi.org/10.1371/journal.pone.0169748>, 2017.
- 10 Heuvelink, G. B. M.: Uncertainty quantification of GlobalSoilMap products in: *GlobalSoilMap. Basis of the Global Spatial Soil Information System*, edited by: Arrouays, D., McKenzie, N., Hempel, J., Forges, A. R. d., and McBratney, A., Taylor & Francis Group, London, UK, 335-240, 2014.
- Heuvelink, G. B. M., Angelini, M. E., Poggio, L., Bai, Z. G., Batjes, N. H., van den Bosch, R., Bossio, D., Estella, S., Lehmann, J., Olmedo, G. F., and Sanderman, J.: Machine learning in space and time for modelling soil organic carbon change, *European Journal of Soil Science*, 72, 1607-1623, <https://doi.org/10.1111/ejss.12998>, 2021.
- 15 ICP Forests: ICP Forests monitoring Manual. Part X: Sampling and analysis of soil, Eberswalde, Germany, <https://storage.ning.com/topology/rest/1.0/file/get/9995584862?profile=original>, 2021.
- ISO-19139: Geographic information XML schema implementation Part 1: Encoding rules, <https://www.iso.org/standard/67253.html>, 2019.
- ISRIC: Data and Software Policy, ISRIC - World Soil Information (WDC - Soils) Wageningen, 6 pp., https://www.isric.org/sites/default/files/user/ISRIC_Data_Policy_2016jun21doi.pdf, 2016.
- 20 ISRIC: SoilGrids and WoSIS web platform (ver. 2.0), ISRIC - World Soil Information, Wageningen, <https://soilgrids.org/>, 2020.
- ISRIC: WoSIS - Contributing institutions and experts, ISRIC - World Soil Information, Wageningen, <https://www.isric.org/explore/wosis/wosis-contributing-institutions-and-experts>, 2023a.
- ISRIC: ISRIC Data Hub, ISRIC - World Soil Information, Wageningen, <https://data.isric.org/>, 2023b.
- 25 ISRIC: WoSIS-latest dashboard, ISRIC - World Soil Information, Wageningen, https://dashboards.isric.org/superset/dashboard/wosis_latest, 2023c.
- ISRIC: WoSIS soil profile database (License categories), ISRIC - World Soil Information, Wageningen, https://dashboards.isric.org/superset/dashboard/wosis_licenses, 2023d.
- 30 IUSS Working Group WRB: World Reference Base for Soil Resources (2nd ed.), FAO, Rome, World Soil Resources Report 103, 145 pp., <http://www.fao.org/ag/agl/agll/wrb/doc/wrb2006final.pdf>, 2006.
- IUSS Working Group WRB: World Reference Base for soil resources 2014 - International soil classification system for naming soils and creating legends for soil maps (update 2015), Global Soil Partnership, International Union of Soil Sciences, and Food and Agriculture Organization of the United Nations, Rome, World Soil Resources Reports 106, 182 pp., <http://www.fao.org/3/i3794en/i3794en.pdf>, 2015.
- 35 IUSS Working Group WRB: World Reference Base for soil resources 2022 - International soil classification system for naming soils and creating legends for soil maps, International Union of Soil Sciences, Vienna (Austria), 284 pp., https://www.isric.org/sites/default/files/WRB_fourth_edition_2022-12-18.pdf, 2022.
- Ivushkin, K., Bartholomeus, H., Bregt, A. K., Pulatov, A., Kempen, B., and de Sousa, L.: Global mapping of soil salinity change, *Remote Sensing of Environment*, 231, <https://doi.org/10.1016/j.rse.2019.111260>, 2019.
- 40 Kalra, Y. P., and Maynard, D. G.: Methods manual for forest soil and plant analysis, Forestry Canada, Edmonton (Alberta), 116 pp., <https://cfs.nrcan.gc.ca/publications/download-pdf/11845>, 1991.
- Luo, Z., Viscarra-Rossel, R. A., and Qian, T.: Similar importance of edaphic and climatic factors for controlling soil organic carbon stocks of the world, *Biogeosciences*, 18, 2063-2073, <https://bg.copernicus.org/articles/18/2063/2021/>, 2021.
- 45 Lutz, F., Stoorvogel, J. J., and Müller, C.: Options to model the effects of tillage on N₂O emissions at the global scale, *Ecological Modelling*, 392, 212-225, <https://www.sciencedirect.com/science/article/pii/S0304380018304034>, 2019.



- Magnusson, B., and Örnemark, U.: The Fitness for Purpose of Analytical Methods – A Laboratory Guide to Method Validation and Related Topics (2nd ed.), Eurachem, https://www.eurachem.org/images/stories/Guides/pdf/MV_guide_2nd_ed_EN.pdf, 2014.
- 5 Maire, V., Wright, I. J., Prentice, I. C., Batjes, N. H., Bhaskar, R., van Bodegom, P. M., Cornwell, W. K., Ellsworth, D., Niinemets, U., Ordonez, A., Reich, P. B., and Santiago, L. S.: Global effects of soil and climate on leaf photosynthetic traits and rates, *Global Ecology and Biogeography*, 24, 706-717, <https://doi.org/10.1111/geb.12296>, 2015.
- Malhotra, A., Todd-Brown, K., Nave, L. E., Batjes, N. H., Holmquist, J. R., Hoyt, A. M., Iversen, C. M., Jackson, R. B., Lajtha, K., Lawrence, C., Vinduskova, O., Wieder, W., Williams, M., Hugelius, G., and Harden, J.: The landscape of soil carbon data: emerging questions, synergies and databases, *Progress in Physical Geography-Earth and Environment*, 43, 707-719, <https://doi.org/10.1177/0309133319873309>, 2019.
- 10 Meyer, H., and Pebesma, E.: Predicting into unknown space? Estimating the area of applicability of spatial prediction models, *Methods in Ecology and Evolution*, 12, 1620-1633, <https://besjournals.onlinelibrary.wiley.com/doi/abs/10.1111/2041-210X.13650>, 2021.
- 15 Moulatlet, G. M., Zuquim, G., Figueiredo, F. O. G., Lehtonen, S., Emilio, T., Ruokolainen, K., and Tuomisto, H.: Using digital soil maps to infer edaphic affinities of plant species in Amazonia: Problems and prospects, *Ecol Evol*, 7, 8463-8477, <https://doi.org/10.1002/ece3.3242>, 2017.
- Munzert, M., Kießling, G., Übelhör, W., Nätscher, L., and Neubert, K.-H.: Expanded measurement uncertainty of soil parameters derived from proficiency-testing data, *Journal of Plant Nutrition and Soil Science*, 170, 722-728, <https://onlinelibrary.wiley.com/doi/abs/10.1002/jpln.200620701>, 2007.
- 20 NATP: North American Proficiency Testing (NAPT) Program, <http://www.naptprogram.org/>, 2015.
- Nenkam, A. M., Wadoux, A. M. J. C., Minasny, B., McBratney, A. B., Traore, P. C. S., Falconier, G. N., and Whitbread, A. M.: Using homosols for quantitative extrapolation of soil mapping models, *European Journal of Soil Science*, n/a, e13285, <https://doi.org/10.1111/ejss.13285>, 2022.
- 25 NPDB: National Pedon Database Canada, Agriculture and Agri-food Canada, <https://sis.agr.gc.ca/cansis/nsdb/npdb/index.html>, 2023.
- OGC: Soil Data IE (Interoperability Experiment), Open Geospatial Consortium (OGC), <https://www.opengeospatial.org/projects/initiatives/soildataie>, 2019.
- 30 Olson, D. M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell, G. V. N., Underwood, E. C., D'amico, J. A., Itoua, I., Strand, H. E., Morrison, J. C., Loucks, C. J., Allnutt, T. F., Ricketts, T. H., Kura, Y., Lamoreux, J. F., Wettengel, W. W., Hedao, P., and Kassem, K. R.: Terrestrial Ecoregions of the World: A New Map of Life on Earth: A new global map of terrestrial ecoregions provides an innovative tool for conserving biodiversity, *BioScience*, 51, 933-938, [https://doi.org/10.1641/0006-3568\(2001\)051\[0933:TEOTWA\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2001)051[0933:TEOTWA]2.0.CO;2), 2001a.
- Olson, R. J., Johnson, K. R., Zheng, D. L., and Scurlock, J. M. O.: Global and regional ecosystem modelling: databases of model drivers and validation measurements, Oak Ridge National Laboratory, Oak Ridge, ORNL/TM-2001/196, 95 pp., http://www-eosdis.ornl.gov/npp/GPPDI/comp/NPP_TM196.pdf, 2001b.
- 35 Orgiazzi, A., Ballabio, C., Panagos, P., Jones, A., and Fernandez-Ugalde, O.: LUCAS Soil, the largest expandable soil dataset for Europe: a review, *European Journal of Soil Science*, 69, 140-153, <https://doi.org/10.1111/ejss.12499>, 2018.
- Palma, R., Janiak, B., Sousa, L. M. d., Schleidt, K., Tomáš Rezník, Egmond, F. v., Leenaars, J., Moshou, D., Mouazen, A., Peter Wilson, Medyckyj-Scott, D., Ritchie, A., Yigini, Y., and Vargas, R.: GloSIS: The Global Soil Information System Web Ontology, *Semantic Web Journal* (in review), <https://w3id.org/gloSIS/model>, 2023.
- 40 Poggio, L., de Sousa, L., Batjes, N. H., Heuvelink, G. B. M., Kempen, B., Riberio, E., and Rossiter, D.: SoilGrids 2.0: producing soil information for the globe with quantified spatial uncertainty, *SOIL*, 7, 217-240, <https://doi.org/10.5194/soil-7-217-2021>, 2021.
- 45 R Core Team: R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, 2021.
- Rayment, E. R., and Lyons, D. J.: Soil chemical methods - Australasia, CSIRO Publishing, 495 pp., 2011.



- Ribeiro, E., Batjes, N. H., and Van Oostrum, A. J. M.: World Soil Information Service (WoSIS) - Towards the standardization and harmonization of world soil data. Procedures Manual 2020, ISRIC - World Soil Information, Wageningen, ISRIC Report 2020/01, 153 pp., <http://dx.doi.org/10.17027/isric-wdc-2020-01>, 2020.
- 5 Rossel, R. A. V., and McBratney, A. B.: Soil chemical analytical accuracy and costs: implications from precision agriculture, *Australian Journal of Experimental Agriculture*, 38, 765-775, 1998.
- Sanderman, J., Hengl, T., and Fiske, G. J.: Soil carbon debt of 12,000 years of human land use, *P Natl Acad Sci USA*, 114, 9575-9580, <https://doi.org/10.1073/pnas.1706103114>, 2017.
- 10 Sayre, R., Dangermond, J., Frye, C., Vaughan, R., Aniello, P., Breyer, S., Cribbs, D., Hopkins, D., Nauman, R., Derrenbacher, W., Burton, D., Grosse, A., True, D., Metzger, M., Hartmann, J., Moosdorf, N., Dürr, H., Paganini, M., DeFourny, P., Arino, O., and Maynard, S.: A New Map of Global Ecological Land Units — An Ecophysiological Stratification Approach, *Association of American Geographers*, Washington DC, 46 pp., https://www.aag.org/wp-content/uploads/2021/12/AAG_Global_Ecosyst_bklt72.pdf, 2014.
- Sayre, R.: World Terrestrial Ecosystems (WTE) 2020 [Dataset], <https://doi.org/10.5066/P9DO61LP>, 2022.
- 15 Schoeneberger, P. J., Wysocki, D. A., E.C. Benham, and Soil Survey Staff: Field book for describing and sampling soils (ver. 3.0, Reprint 2021), National Soil Survey Center Natural Resources Conservation Service, U.S. Department of Agriculture, Lincoln (NE), 2012.
- Shepherd, K. D., Ferguson, R., Hoover, D., van Egmond, F., Sanderman, J., and Ge, Y.: A global soil spectral calibration library and estimation service, *Soil Security*, 7, 100061, <https://www.sciencedirect.com/science/article/pii/S2667006222000284>, 2022.
- 20 Shi, G., Shangguan, W., Zhang, Y., Li, Q., Wang, C., and Li, L.-J.: Reducing Location Error of Legacy Soil Profiles Leads to Significant Improvement in Digital Soil Mapping, SSRN, <https://ssrn.com/abstract=4643055> or <http://dx.doi.org/10.2139/ssrn.4643055>, 2023.
- Soil Survey Staff: Soil Survey Laboratory Information Manual (Ver. 2.0), National Soil Survey Center, Soil Survey Laboratory, USDA-NRCS, Lincoln (NE), Soil Survey Investigation Report No. 45, 506 pp., http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_052226.pdf, 2011.
- 25 Soil Survey Staff: Keys to Soil Taxonomy, 12th ed., USDA-Natural Resources Conservation Service, Washington, DC., 2014.
- Soil Survey Staff: Keys to Soil Taxonomy, 13th ed., USDA-Natural Resources Conservation Service, Washington, DC., 2022a.
- Soil Survey Staff: Soil Survey Laboratory Methods Manual (Version 6.0., Part1: Current methods), U.S. Department of Agriculture, Natural Resources Conservation Service, Lincoln (Nebraska), 1001 pp., 2022b.
- 30 Sothe, C., Gonsamo, A., Arabian, J., and Snider, J.: Large scale mapping of soil organic carbon concentration with 3D machine learning and satellite observations, *Geoderma*, 405, 115402, <https://www.sciencedirect.com/science/article/pii/S0016706121004821>, 2022.
- Suvannang, N., Hartmann, C., Yakimenko, O., Solokha, M., Bertsch, F., and Moody, P.: Evaluation of the First Global Soil Laboratory Network (GLOSOLAN) online survey for assessing soil laboratory capacities, *Global Soil Partnership (GSP) / Food and Agriculture Organization of the United Nations (FAO)*, Rome, GLOSOLAN/18/Survey Report, 54 pp., <http://www.fao.org/3/CA2852EN/ca2852en.pdf>, 2018.
- 35 Turek, M. E., Poggio, L., Batjes, N. H., Armindo, R. A., de Jong van Lier, Q., de Sousa, L., and Heuvelink, G. B. M.: Global mapping of volumetric water retention at 100, 330 and 15 000 cm suction using the WoSIS database, *International Soil and Water Conservation Research*, 11, 225-239, <https://www.sciencedirect.com/science/article/pii/S2095633922000636>, 2023.
- 40 USDA-NCSS: National Cooperative Soil Survey (NCSS) Soil Characterization Database, United States Department of Agriculture, Natural Resources Conservation Service, Lincoln, https://ncsslabsdatamart.sc.egov.usda.gov/database_download.aspx, 2021.
- van Leeuwen, C., Mulder, V. L., Batjes, N. H., and Heuvelink, G. B. M.: Statistical modelling of measurement error in wet chemistry soil data, *European Journal of Soil Science*, 73, 13137, <https://doi.org/10.1111/ejss.13137>, 2022.



- van Leeuwen, C. C. E., Mulder, V. L., Batjes, N. H., and Heuvelink, G. B. M.: Effect of measurement error in wet chemistry soil data on the calibration and model performance of pedotransfer functions, *Geoderma*, 442, 116762, <https://www.sciencedirect.com/science/article/pii/S0016706123004391>, 2024.
- 5 Van Looy, K., Bouma, J., Herbst, M., Koestel, J., Minasny, B., Mishra, U., Montzka, C., Nemes, A., Pachepsky, Y., Padarian, J., Schaap, M., Tóth, B., Verhoef, A., Vanderborght, J., van der Ploeg, M., Weihermüller, L., Zacharias, S., Zhang, Y., and Vereecken, H. C. R. G.: Pedotransfer functions in Earth system science: challenges and perspectives, *Reviews of Geophysics*, 55, 1199-1256, <http://dx.doi.org/10.1002/2017RG000581>, 2017.
- 10 van Reeuwijk, L. P.: On the way to improve international soil classification and correlation: the variability of soil analytical data, ISRIC, Wageningen, Annual Report 1983, 7-13 pp., https://www.isric.org/sites/default/files/isric_annual_report_1983.pdf, 1983.
- 15 Viscarra Rossel, R. A., Behrens, T., Ben-Dor, E., Brown, D. J., Demattê, J. A. M., Shepherd, K. D., Shi, Z., Stenberg, B., Stevens, A., Adamchuk, V., Aichi, H., Barthès, B. G., Bartholomeus, H. M., Bayer, A. D., Bernoux, M., Böttcher, K., Brodský, L., Du, C. W., Chappell, A., Fouad, Y., Genot, V., Gomez, C., Grunwald, S., Gubler, A., Guerrero, C., Hedley, C. B., Knadel, M., Morrás, H. J. M., Nocita, M., Ramirez-Lopez, L., Roudier, P., Campos, E. M. R., Sanborn, P., Sellitto, V. M., Sudduth, K. A., Rawlins, B. G., Walter, C., Winowiecki, L. A., Hong, S. Y., and Ji, W.: A global spectral library to characterize the world's soil, *Earth-Science Reviews*, 155, 198-230, <http://dx.doi.org/10.1016/j.earscirev.2016.01.012>, 2016.
- 20 Wang, M., Zhang, S., Guo, X., Xiao, L., Yang, Y., Luo, Y., Mishra, U., and Luo, Z.: Responses of soil organic carbon to climate extremes under warming across global biomes, *Nature Climate Change*, 14, 98-105, <https://doi.org/10.1038/s41558-023-01874-3>, 2024.
- 25 WEPAL: ISE Reference Material - A list with all available ISE reference material samples, WEPAL (Wageningen Evaluating Programmes for Analytical Laboratories), Wageningen, 110 pp., <http://www.wepal.nl/website/products/RefMatISE.htm>, 2019.
- 30 Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-W., da Silva Santos, L. B., Bourne, P. E., Bouwman, J., Brookes, A. J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C. T., Finkers, R., Gonzalez-Beltran, A., Gray, A. J. G., Groth, P., Goble, C., Grethe, J. S., Heringa, J., 't Hoen, P. A. C., Hooft, R., Kuhn, T., Kok, R., Kok, J., Lusher, S. J., Martone, M. E., Mons, A., Packer, A. L., Persson, B., Rocca-Serra, P., Roos, M., van Schaik, R., Sansone, S.-A., Schultes, E., Sengstag, T., Slater, T., Strawn, G., Swertz, M. A., Thompson, M., van der Lei, J., van Mulligen, E., Velterop, J., Waagmeester, A., Wittenburg, P., Wolstencroft, K., Zhao, J., and Mons, B.: The FAIR Guiding Principles for scientific data management and stewardship, *Scientific Data*, 3, 160018, <http://dx.doi.org/10.1038/sdata.2016.18>, 2016.