

# A global monthly 3D-field of seawater pH over 3 decades: a machine learning approach

Guorong Zhong<sup>1,2,3,4</sup>, Xuegang Li<sup>1,2,3,4\*</sup>, Jinming Song<sup>1,2,3,4\*</sup>, Baoxiao Qu<sup>1,2,3,4</sup>, Fan Wang<sup>1,2,3,4</sup>, Yanjun Wang<sup>1,4</sup>, Bin Zhang<sup>1,4</sup>, Lijing Cheng<sup>4,5</sup>, Jun Ma<sup>1,2,3,4</sup>, Huamao Yuan<sup>1,2,3,4</sup>, Liqin Duan<sup>1,2,3,4</sup>, Ning Li<sup>1,2,3,4</sup>, Qidong Wang<sup>1,2,3,4</sup>, Jianwei Xing<sup>1,2,3,4</sup>, Jiajia Dai<sup>1,2,3,4</sup>

<sup>1</sup>Institute of Oceanology, Chinese Academy of Sciences, Qingdao 266071, China

<sup>2</sup>Laboratory for Marine Ecology and Environmental Science, Qingdao National Laboratory for Marine Science and Technology, Qingdao, China

<sup>3</sup>University of Chinese Academy of Sciences, Beijing 101407, China

<sup>4</sup>Center for Ocean Mega-Science, Chinese Academy of Sciences, Qingdao 266071, China

<sup>5</sup>Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China

Correspondence to: Xuegang Li (lixuegang@qdio.ac.cn) and Jinming Song (jmsong@qdio.ac.cn)

**Abstract.** The continuous uptake of anthropogenic CO<sub>2</sub> by the ocean leads to ocean acidification, which is an ongoing threat to the marine ecosystem. The ocean acidification rate was globally documented in the surface ocean but limited below the surface. Here, we present a monthly four-dimensional 1°×1° gridded product of global seawater pH at total scale and in-situ temperature, derived from a machine learning algorithm trained on pH observations at total scale and in-situ temperature from the Global Ocean Data Analysis Project (GLODAP). The proposedeconstructed pH product covers the years 1992-2020 and depths from the surface to 2 km on 41 levels. A three-step machine learning-based algorithm was used to construct the pH product, incorporating region division by the self-organizing map neural network, predictor selection by the stepwise regression algorithm that adds and removes variables from network inputs based on their contribution to reducing reconstruction errors, and non-linear relationship regression by feed-forward neural networks (FFNN). ~~Three types of machine learning algorithms were used in the pH product construction, including self-organizing map neural networks for region dividing, a stepwise algorithm for predictor selection, and feed forward neural networks (FFNN) for non-linear relationship regression.~~ The performance of the machine learning algorithm was validated using real observations by a cross validation method, where four repeating iterations were carried out with 25% varied observations for each evaluation and 75% for training. The proposedeconstructed pH product is evaluated through comparisons to time series observations and the GLODAP pH climatology. The overall root mean square error between the FFNN reconstructed pH and the GLODAP measurements is 0.028, ranging from 0.044 in the surface to 0.013 at 2000 m. The pH product is distributed through the data repository of the Marine Science Data Center of the Chinese Academy of Sciences at <http://dx.doi.org/10.12157/IOCAS.20230720.001> (Zhong et al., 2023).

## 1 Introduction

Since the Industrial Revolution, the oceans have absorbed approximately one-quarter of the carbon dioxide emitted by human activities (Le Quéré et al., 2010; Friedlingstein et al., 2023). The continuous absorption of carbon dioxide from the atmosphere results in a decline in carbonate saturation states and surface seawater pH, which is a phenomenon of great concern: ocean acidification (Caldeira et al., 2003; Feely et al., 2004; Orr et al., 2005; Feely et al., 2009). As one of the primary environmental challenges the ocean faces today, ocean acidification will have extensive impacts on marine organisms and the ecological environment, resulting in notable changes to the marine ecosystem. Therefore, the assessment of ocean acidification is crucial for researching the response of marine organisms to changes in seawater pH and understanding the potential future changes in the capacity of the global ocean to uptake CO<sub>2</sub> (Sabine et al., 2010; Guallart et al., 2015).

40 However, acidification research is greatly limited in terms of temporal and spatial coverage due to the lack of long-term, global coverage, and continuous seawater pH measurements. Accurate seawater pH measurements are only available from select ship surveys and a limited number of time series stations in recent decades (Fay et al., 2013; Takahashi et al., 2014). Recent research using discrete ship survey measurements revealed rapid surface ocean acidification in the Arctic Ocean, with some areas showing an average decreasing pH trend of  $-0.0086 \text{ yr}^{-1}$  (Luo et al., 2016; Terhaar et al., 2020; Qi et al., 2022).  
45 Both seawater pH measurements from time series stations and discrete ship surveys suggest notable regional differences in surface ocean acidification rates (Bates et al., 2014; Lauvset et al., 2015). In the Japan/East Sea, the acidification rate in the deep ocean may be faster than previously considered and even faster than in the surface ocean (Chen et al., 2017; Li et al., 2022). Meanwhile, relatively slow acidification was found in the deep Atlantic Ocean below 2000 m (Guallart et al., 2015), and rising pH in deep waters around 1000 m was also reported in the North Pacific Ocean (Ishizu et al., 2021). With limited  
50 reports about acidification below the surface, there remains a need to enhance our understanding of global ocean acidification rates across varying depths-at different depths still needs to be improved.

The lack of long-term, global coverage, and continuous seawater pH measurements makes it difficult to expand the understanding of global deep ocean acidification using classic regression methods. Recent applications of machine learning methods in the global reconstructions mapping of marine carbonate system variables have facilitated global-scale research on  
55 the acidification and carbon cycle, such as including the single/ensemble-based FFNN method and the SOM-FFNN method for reconstruction of surface ocean partial pressure of  $\text{CO}_2$  ( $p\text{CO}_2$ , Landschützer et al., 2014; Chau et al., 2022; Zhong et al., 2022; Chau et al., 2024), dissolved inorganic carbon (DIC-, Broullón et al., 2020; Keppler et al., 2020; Gregor and Gruber, 2021; Chau et al., 2024), and alkalinity (Broullón et al., 2019; Gregor and Gruber, 2021; Chau et al., 2024), indicated a feasible way to construct a global gridded seawater pH dataset. These methods have inspired our methodology for constructing the  
60 global gridded seawater pH dataset. Until now, there are only surface ocean gridded pH products are available in acidification research, including the 1° JMA product (Iida et al., 2021), the 1° OceanSODA-ETHZ product (- Gregor and Gruber, 2021), the 0.25° remote-sensing-based product (Jiang et al., 2022), and the 0.25° CMEMS-LSCE product (Chau et al., 2024), which were derived from reconstructing  $p\text{CO}_2$ , DIC, or alkalinity-other carbonate system variables using machine learning algorithms and subsequently calculating pH with the CO2SYS program (Lewis and Wallace, 1998). In this paper, we present a monthly  
65 gridded global ocean pH product covering depths of 0-2000 m from January 1992 to December 2020, using a machine learning method trained on pH measurements from the Global Ocean Data Analysis Project (GLODAP) dataset (Lauvset et al., 20222023). The construct-proposed pH product provides regional and global insight into ocean acidification on timescales ranging from a few years to multiple decades.

## 2 Methods

### 70 2.1 Data sources and processing

The pH measurements at total scale and in-situ temperature and pressure from the Global Ocean Data Analysis Project (GLODAP) dataset 2023 version were used for neural network training (Lauvset et al., 20222023). The reconstructed pH product is also at total scale and in-situ temperature based on a gridded global seawater temperature product (Cheng et al., 2017). We have collected gridded products of different variables as potential pH predictors (Table 1), and the selection of these  
75 products was based on two reasons. The first reason was their potential association with physical, chemical, and biological ocean process which may affect the seawater pH. Another reason was the sufficient availability in time and spatial coverage and their potential association with the unavailable interannual variability of some climatological products used. Specifically, the mixed layer depth, bathymetry, and ocean currents were related to the physical mixing of seawater and spatial distribution of pH. Sea level pressure, surface pressure, wind speed, sea surface height, surface ocean  $p\text{CO}_2$ , and dry air mixing ratio of  
80 atmospheric  $\text{CO}_2$  were related to the  $\text{CO}_2$  exchange across the interface. The Multivariate ENSO ~~oceanic~~ El Niño index, Arctic Oscillation index, and Southern Oscillation index may be related to pH variability over years or decades in particular regions.

The total alkalinity and DIC reflect the ocean carbonate system and were generally used to calculate seawater pH indirectly. However, 3D field products with sufficient time and spatial coverage are currently not available for these two variables, so monthly climatological 3D products were used for better pH spatial distribution. The remote sensing products may be useful to provide extra interannual variability information from currently used climatological products, and they were also used to test the correlation with seawater pH and its potential drivers. Temporal and spatial sample information, including latitude, longitude, depth and sample time, was also used as supplementary variables. Latitude and longitude were normalized to radians using sine and cosine transformations, to present connected sample position information. The spatial sample position and time information of GLODAP measurements were input in the training of FFNNs, and the spatial position and time of defined 1° and monthly product grids were input into FFNNs during the interpolation process to output a gridded product. Most predictor products were obtained with a monthly and 1°×1° resolution, which can be directly used without any treatments. Differently, products with higher resolutions were integrated into the same monthly and 1°×1° resolution by averaging, before they can be used in the relationship fitting. For instance, the mixed layer depth product, originally obtained with a resolution of 0.25°×0.25°, was converted to a 1°×1° resolution by averaging 16 0.25° grids into one 1° grid. Similarly, such as the xCO<sub>2</sub> product, predictor products obtained with ~~daily or~~ weekly resolutions were converted to the monthly resolution by directly averaging all values within the same month, ~~such as the ocean currents product.~~ Products used for variables listed in Table 1 was chose due to their sufficient temporal and spatial coverage and the application in previous research on reconstruct of carbonate system variables. For example, the ECCO2 MLD product has been used in reconstructions of the CMEMS-LSCE surface ocean carbonate system variables product (Chau, et al., 2024) and the MPI-SOM-FFN pCO<sub>2</sub> product (Landschützer et al., 2014).

100

**Table 1. Data products used as pH predictors.**

Predictor	Abbreviation	Data product <u>and reference</u>	Resolution	<u>Related process to affect pH</u>
Sine of $(\text{latitude} \cdot \pi/180^\circ)$ <u>latitude</u>	sin(Lat)	-	-	<u>Sample position and time of GLODAP pH measurements</u>
Sine of $(\text{longitude} \cdot \pi/180^\circ)$ <u>longitude</u>	sin(Lon)	-	-	
Cosine of $(\text{longitude} \cdot \pi/180^\circ)$ <u>longitude</u>	cos(Lon)	-	-	
Number of months since January 1992	N <sub>mon</sub>	-	-	
Year	Year	-	-	
Month	Month	-	-	
Depth	Depth	-	-	
Temperature and monthly anomaly	Temp, Temp <sub>anom</sub>	IAP global ocean temperature gridded product <u>(Cheng et al., 2016; 2017)</u>	1°, <u>monthly</u> since 1940, <u>0-2000 m with 41 levels</u>	<u>State of carbonate system</u>
Salinity and monthly anomaly	Sal, Sal <sub>anom</sub>	IAP global ocean salinity gridded product <u>(Cheng et al., 2020)</u>	1°, <u>monthly</u> since 1940, <u>0-2000 m with 41 levels</u>	
Climatological total alkalinity	Alk	AT_NNGv2_climatology <u>(Broullón et al., 2019)</u>	1°, <u>monthly climatological, 0-5500 m with 102 levels</u>	
Climatological dissolved inorganic carbon	DIC	TCO2_NNGv2LDEO_climatology <u>(Broullón et al., 2020)</u>	1°, <u>monthly climatological, 0-5500 m with 102 levels</u>	
Climatological dissolved oxygen	DO	WOA18 <u>(Garcia et al., 2020a)</u>	1°, <u>monthly climatological, 0-5500 m with 102 levels</u>	<u>Biological production and drawdown of organic matter</u>
Climatological nitrate	Nitrate	WOA18 <u>(Garcia et al., 2020b)</u>	1°, <u>monthly climatological, 0-5500 m with 102 levels</u>	
Climatological phosphate	Phosphate	<del>WOA18</del>		
Climatological silicate	Silicate	<del>WOA18</del>		
Mixed layer depth and monthly anomaly	MLD, MLD <sub>anom</sub>	ECCO2 cube92 <u>(Menemenlis et al., 2008)</u>	0.25°, <u>monthly</u> since 1992	<u>Physical mixing of seawater and stratification</u>
Sea surface height and monthly anomaly	SSH, SSH <sub>anom</sub>	<del>ECCO2 cube92</del> <del>ECCO2 cube92</del>		<u>ocean wave, tides, current, and sea-level rise</u>
W velocity of ocean currents at 5 m, 65m, 105m, 195m, and in-situ depth	W <sub>vel(5m)</sub> –W <sub>vel(i n-situ)</sub>			<u>Ocean current and upwelling</u>
Sea level pressure	SLP	ERA5 <u>(Hersbach et al., 2020)</u>	1°, <u>monthly</u> since 1940	<u>CO<sub>2</sub> exchange between surface seawater and atmosphere</u>
Surface pressure	P <sub>surf</sub>	<del>ERA5</del>		
dry air mixing ratio of atmospheric CO <sub>2</sub> and monthly anomaly	xCO <sub>2</sub> , xCO <sub>2 anom</sub>	NOAA Greenhouse Gas Marine Boundary Layer Reference <u>(Lan et al., 2023)</u>	0.25°, <u>weekly</u> since 1979	
<u>Multivariate ENSO Index</u> <u>Oceanic Niño Index</u>	MEI	bi-monthly Multivariate El Niño/Southern Oscillation index <u>(Wolter et al., 2011)</u>	<u>monthly</u> since 1979	<u>El Niño and Southern Oscillation</u>

Arctic Oscillation index	AOI	Climate Prediction Center Daily Arctic Oscillation Index ( <a href="#">CPC, 2002</a> )	<a href="#">monthly</a> since 1950	<a href="#">Arctic Oscillation</a>
Southern Oscillation Index	SOI	Climate Prediction Center Southern Oscillation Index ( <a href="#">CPC, 2005</a> )	<a href="#">monthly</a> since 1951	<a href="#">Southern Oscillation</a>
Bathymetry	Bathy	GEBCO_2022 Grid ( <a href="#">GEBCO, 2022</a> )	15 arc-second	<a href="#">Vertical volume of seawater</a>
10 m Wind speed and monthly anomaly	Wind, Wind <sub>anom</sub>	ERA5 ( <a href="#">Hersbach et al., 2020</a> )	1°, <a href="#">monthly</a> since 1940	<a href="#">CO<sub>2</sub> exchange between surface seawater and atmosphere</a>
Surface ocean pCO <sub>2</sub>	pCO <sub>2</sub>	Stepwise FFNN ( <a href="#">Zhong et al., 2022</a> )	1°, <a href="#">monthly</a> since 1992	
Climatology of Surface Ocean pCO <sub>2</sub>	pCO <sub>2 clim</sub>	MPI-ULB-SOM_FFNN_clim ( <a href="#">Landschützer et al., 2020</a> )	0.25°, <a href="#">monthly climatological</a>	
Chlorophyll and monthly anomaly*	Chl, Chl <sub>anom</sub>	MODIS-Aqua Chlorophyll Data ( <a href="#">NASA, 2022a</a> )	9km, <a href="#">monthly</a> since 2002	<a href="#">Biological production of organic matter</a>
Photosynthetically Available Radiation	PAR	<a href="#">MODIS-Aqua Photosynthetically Available Radiation Data (NASA, 2022b)</a>		
Diffuse attenuation coefficient at 490 nm	KD490	<a href="#">MODIS-Aqua Downwelling Diffuse Attenuation Coefficient Data (NASA, 2022c)</a>		<a href="#">Supplementary for lacking interannual variability of other variables, or potential correlation with unclear process affecting pH</a>
Remote sensing reflectance at 412-678 nm**	RRS412–RRS678	<a href="#">MODIS-Aqua Remote-Sensing Reflectance Data (NASA, 2022d)</a>		
Total absorption at 412-678 nm	Ta412–Ta678	<a href="#">MODIS-Aqua Inherent Optical Properties Data (NASA, 2022e)</a>		
Total backscattering at 412-678 nm	Tb412–Tb678	<a href="#">MODIS-Aqua Inherent Optical Properties Data (NASA, 2022e)</a>		

(\*): products from Chlorophyll to Total backscattering are satellite remote sensing products;


(\*\*): Remote sensing reflectance, total absorption, and total backscattering both include 10 wavelengths: 412nm, 443nm, 469nm, 488nm, 531nm, 547nm, 555nm, 645nm, 667nm, and 678nm, with each wave length regard as one individual parameter.)

On the other hand, the discrete GLODAP measurements did not match the monthly 1°×1° resolution of pH predictor products. To be consistent in the temporal and spatial resolution, the discrete GLODAP measurements were also merged into a monthly and 1°×1° resolution by averaging. The vertical layer of the temperature and salinity gridded product were used as reference standards for adjusting other collected products and constructing the pH product (Cheng et al., 2016; Cheng et al., 2017; Cheng et al., 2020). These layers covered a depth range of 0-2000 m depth, with a total of 41 layers, including 0 m, 5 m, 10-100 m at 10 m intervals, 120-200 m at 20 m intervals, 250-900 m at 50 m intervals, and 1000-2000 m at 100 m intervals. Subsequently, the in-situ seawater measurements of pH, temperature, salinity, latitude, longitude, and depth from the GLODAP dataset were averaged monthly within the same 1°×1° grid (first grid centered at 89.5°S, 0.5°E) and within the same vertical layer to match the resolution of the predictor products. Since a direct average was used instead of a weighted average, the average latitude, longitude, and depth values from the initial measurements within the same 1°×1° grid were then used as new sample position for the derived monthly measurements, instead of being located at the center point of grids. The pH measurements obtained after the 1°×1° grid and monthly averaging were employed to establish a neural network model and fit a non-linear relationship with the pH predictors.


## 2.2 Biogeochemical province

To identify predictors that are most relevant to pH drivers in different regions, we divide the global ocean into distinct biogeochemical provinces using self-organizing map neural networks (SOM). This was achieved by inputting climatological surface seawater temperature, salinity, mixed layer depth, chlorophyll concentration, dissolved oxygen, nitrate, phosphate,

silicate, and pH (Lauvset et al., 2016) into a 4×4 SOM network, resulting in the partitioning of the global ocean into preliminary 16 provinces. Subsequently, the small “island” provinces with fewer than ten connected grids or covered by fewer than 100 GLODAP pH measurements were merged with the nearest neighboring provinces, as the pH reconstruction errors tend to be notably higher due to the extremely few training samples in the non-linear relationship fitting by networks. Subsequently, the province with fewer than ten connected grids or less than 100 GLODAP pH measurements was merged with the nearest neighboring provinces. In addition, the province separated by continents was manually subdivided into distinct provinces, such as the province spanning the North Pacific and the North Atlantic. As a result, the global ocean was divided into 14 biogeochemical provinces, as shown in Figure 1. The boundary of SOM provinces was treated with a cross-boundary method to relieve the discontinuity of spatial distribution near the SOM boundaries (Zhong et al., 2022). Due to much more dynamic variation in coastal seawater pH, the global coastal areas have higher reconstruction errors than the open oceans. In this study, we removed all coastal areas shallower than 200 m bathymetry. Furthermore, because the drivers of seawater pH near the surface is different with deeper waters, the ocean area was divided into two layers: the mixed layer (ranging from 0 m to mixed layer depth) and the intermediate layer (ranging from mixed layer depth to 2000 m). Consequently, the gridded product construction in each province was carried out separately for the two layers. Application of SOM method can effectively reduce regional reconstruction errors, but it also generates discontinuity problems near the boundary. Therefore, a cross-boundary method was used to improve the FFNN performance near the SOM and vertical boundary (Zhong et al., 2022). The spatial scale of training samples in each SOM province was expanded out of the boundary for 10 grids, and out of the vertical boundary for 2 layers (Figure 2). By increasing additional training sample outside the SOM province and vertical layer boundary, the cross-boundary method can effectively reduce the appearance of dysconnectivity near boundaries (Figures S1 and S2).



145 **Figure 1: Map of the biogeochemical province.**



**Figure 2: Cross-boundary method for better connectivity near the SOM boundary and vertical boundary.**



### 2.3 pH product construction

The forward feedback neural network (FFNN) with a single hidden layer was applied to fit the non-linear relationship between seawater pH and its predictors to perform spatial interpolation and construct the gridded product:

$$\text{pH} = f(\text{Predictors}_1, \text{Predictors}_2, \dots, \text{Predictors}_N) \quad (1)$$

where  $f$  was a non-linear function built by FFNN, and predictors related to chemical, physical, and biological properties were selected from products in Table 1. Considering the regional difference in pH variability and its drivers, identifying the combination of most relevant predictors in each region was a critical precondition. Thus, the entire product construction method includes two steps (Figure 3):


(1) Selection of seawater pH predictors in each province using the Stepwise FFNN algorithm (referred as (1) Stepwise FFNN in Figure 3). All the collected products were input into the Stepwise FFNN algorithm to identify the predictors that yield the lowest reconstructionpredicting errors for seawater pH (Zhong et al., 2022). The variation in standard deviation (MAE) calculated by the K-fold cross validation method will feed back to update the input products. The input variables are selected as pH predictors one by one in the way MAE decreases the fastest. Specifically, by comparing reconstruction errors of using each collected environmental variable in Table 1 as the only predictor input to the FFNN, the variable with the lowest error is selected as the first pH predictor and moved out from the environmental variables list used in the subsequent steps. Subsequently, while keeping the first predictor unchanged, compare reconstruction errors when using each remaining environmental variable as the second input for the FFNN. The variable with the lowest error is determined to be the second pH predictor. In the same way, new predictors are sequentially determined. This selection process continued through multiple iterations until no further reduction in MAE was observed, regardless of whether a variable was added or removed. The variables identified in previous iterations were then output as the optimal pH predictors. The co-correlation among pH predictors was not considered when choosing the variables in Table 1, but was considered in the design of the Stepwise FFNN algorithm. Since both overfitting caused by co-correlation and underfitting caused by an insufficient number of predictors result in both lead to a significant increases in pH reconstructionpredicting errors, the lowest reconstructionpredicting error is considered to occurs between these two states. In order to eliminate potential co-correlation and prevent overfitting, whenever after a new predictor is identified, the algorithm also tests whether the reconstruction error will decrease when sequentially removing each determined predictor. The algorithm individually removes each previously identified predictors immediately after adding one variable as a predictor. If the error decreases after removing a previously determined predictor, this predictor is highly correlated with other identified predictors. ~~The Stepwise FFNN algorithm tests all products based on changes in predicting errors, to reach a balance avoiding overfitting and underfitting. If a certain predictor is highly correlated with existing predictors, this predictor tends to fail to compete with other variables in the adding of predictors and is generally removed in the following removal step to reduce reconstruction errors.~~ Therefore, most of the co-correlation among the selected predictors has been removed in this Stepwise FFNN selection procedure. If products with co-correlations are still selected, some products may provide important additional information in specific regions, leading to a greater reduction in reconstructionpredicting errors compared to the increase caused by overfitting. In each province, pH predictors were selected separately for the mixed layer (Table 2) and intermediate layer (Table 3). In certain polar areas and prior to August 2002 when satellite remote sensing products (products from  $Z_{eu}$  to  $Tb678$  in Table 1) were not available, the additional selection of predictors was carried out without the use of satellite remote sensing products (Table S1). These satellite products were not used in the intermediate layer due to low correlation with seawater pH, with no need for additional selection.

(2) Fitting the non-linear relationship between seawater pH and selected predictors (referred as (2) FFNN in Figure 3). In each province, a group of FFNNs ~~was~~ were trained separately for the mixed layer and intermediate layer to fit the non-linear relationship, based on the predictors selected in the first step and GLODAP pH measurements. To mitigate the influence of the FFNN's initial state on reconstructedpredicting values, multiple networks with the same structure but different initial states

190 were trained and their results were averaged (Standard deviation showing in Figure S5). ~~Then~~ Subsequently, the seawater pH  
was calculated ~~using the trained FFNNs, along with by inputting~~ the product of pH predictors ~~into the trained FFNNs~~. ~~Since~~  
~~the satellite remote sensing products used in this work lack data during the period before August 2002 and in certain polar~~  
~~areas during winter, the FFNN generated missing values in these grids when remote sensing products were used as predictors.~~  
~~If satellite remote sensing products were used~~ To address these missing values, we selected additional groups of predictors after  
195 ~~removing remote sensing products (Table S1), and then trained~~ additional FFNNs ~~to predict pH in grids with missing values.~~  
~~This procedure was the same as the reconstruction process in the intermediate layer, in which the remote sensing products~~  
~~were also not used, were trained with predictors in Table S1, for pH product construction in some polar areas and before August~~  
~~2002 when satellite products were unavailable.~~ Finally, the seawater pH ~~predicting~~ values from all FFNNs were combined to  
construct the global ocean 0-2000 m seawater pH gridded product from January 1992 to December 2020, with a  $1^{\circ} \times 1^{\circ}$  spatial  
200 resolution. ~~The pH data earlier than 1992 is unavailable because the predictors used from ECCO2 cube92 product (Menemenlis~~  
~~et al., 2008) also start from 1992. Data after 2020 is limited by the coverage of used surface ocean  $p\text{CO}_2$  product and will be~~  
~~updated in future works.~~

All FFNNs used in these two steps have the same structure with a single hidden layer, as using deeper structures tends to  
cause overfitting and increase pH reconstruction errors. The number of neurons was determined by comparing reconstruction  
205 errors of FFNNs with different neurons based on the same training samples, testing samples, and pH predictors, and then  
adopting the number with the lowest reconstruction error. Specifically, for the stepwise FFNN regression step, the number of  
neurons in FFNNs was determined using provisional predictors from preliminary experiments with the number of neurons set  
to 25.





**Figure 3: The procedure of pH product construction.** (1) Stepwise FFNN: the algorithm for selecting predictors (Zhong et al., 2022); (2) FFNN: fitting the non-linear relationship between seawater pH and its predictors. *Collected Environmental variables: collected products listed in Table 1. pH predictors: the selected most informative variables listed in Tables 2 and 3. Remote sensing products: variables from Chlorophyll to Total backscattering in Table 1. Mixed layer: from 0 m to mixed layer depth; intermediate layer: from mixed layer depth to*

215 ~~2000 m. Available gridded product: products listed in table 1, from which pH predictors were selected; Remote sensing products were not available in some polar areas and during the period before August 2002. Mixed layer depth: using ECCO2 cube92 mixed layer depth product (Menemenlis et al., 2008).~~

**Table 2. Predictors selected by the stepwise FFNN algorithm in the Mixed layer.**

Province	FFNN neurons	pH Predictor
P1 Arctic Ocean	10	$p\text{CO}_2$ , sin(Lat), Depth, Sal, $W_{\text{vel}}(105\text{m})$
P2 Subpolar North Atlantic	10	Phosphate, DO, $N_{\text{mon}}$ , DIC, Sal, Bathy
P3 Seasonally stratified North Atlantic	75	$p\text{CO}_2_{\text{clim}}$ , Depth, Temp, Silicate, $p\text{CO}_2$ , DIC
P4 Permanently stratified North Atlantic	20	$p\text{CO}_2$ , Phosphate, sin(Lat), Depth, SSH <sub>anom</sub> , Sal <sub>anom</sub> , $W_{\text{vel}}(195\text{m})$ , Temp, $W_{\text{vel}}(\text{in-situ})$ , $p\text{CO}_2_{\text{clim}}$ , DO
P5 Equatorial Atlantic	50	sin(Lat), Tb469, Temp, Tb555, Tb547, Nitrate, Tb667, Tb678, Tb488, Tb645, Tb531, Sal
P6 Subpolar North Pacific	10	DIC, sin(Lat), sin(Lon), Depth, Salinity, Temp, $p\text{CO}_2$ , $W_{\text{vel}}(\text{in-situ})$
P7 Subtropical North Pacific	50	Temp, sin(Lon), sin(Lat), $p\text{CO}_2$ , Phosphate, Sal, $p\text{CO}_2_{\text{clim}}$ , Depth, cos(Lon), Nitrate, Sal <sub>anom</sub> , Alk
P8 Equatorial Pacific	20	$p\text{CO}_2$ , Silicate, Depth, Sal, Temp, Wind, Alk, RRS645, Ta555, Ta547
P9 Equatorial Indian Ocean	10	DO, Temp <sub>anom</sub> , $p\text{CO}_2$ , Depth, $W_{\text{vel}}(\text{in-situ})$ , $W_{\text{vel}}(195\text{m})$ , $W_{\text{vel}}(65\text{m})$
P10 Subtropical South Atlantic	10	$p\text{CO}_2$ , DIC, Silicate, RRS645, $W_{\text{vel}}(\text{in-situ})$ , Ta547, Temp, Ta667, Sal, Phosphate, Tb412, Ta412, Tb443, DO, $x\text{CO}_2$
P11 Subtropical South Pacific	10	Silicate, $p\text{CO}_2$ , Tb412, Phosphate, Depth, Ta488, Temp <sub>anom</sub> , Ta531
P12 Subtropical South Indian Ocean	10	$p\text{CO}_2$ , Silicate, Phosphate, Nitrate, Depth, Wind
P13 Subpolar Southern Ocean	20	Phosphate, Depth, $p\text{CO}_2$ , $p\text{CO}_2_{\text{clim}}$ , Sal, DIC, Nitrate
P14 Southern Ocean ice	20	Phosphate, Temp, $p\text{CO}_2$ , Depth, Sal, Alk, SSH

220 (The predictors are arranged in order of relative importance, with the variables listed at the front of each province being more effective in reducing reconstructionpredicting errors when used as pH predictors.)

**Table 3. Predictors selected by the stepwise FFNN algorithm in the intermediate layer.**

Province	FFNN neurons	pH Predictor
P1 Arctic Ocean	50	Phosphate, Nitrate, Sal, Depth, sin(Lat), SSH
P2 Subpolar North Atlantic	20	Phosphate, DO, Depth, Year, Sal, Temp, Nitrate, sin(Lat), Alk, $W_{\text{vel}}(195\text{m})$
P3 Seasonally stratified North Atlantic	10	DIC, Nitrate, Temp, Depth, sin(Lon), Year
P4 Permanently stratified North Atlantic	20	Phosphate, Temp, Depth, sin(Lat), $N_{\text{mon}}$ , sin(Lon), Sal, Sal <sub>anom</sub> , Nitrate, $W_{\text{vel}}(\text{in-situ})$
P5 Equatorial Atlantic	25	Depth, DIC, Sal, sin(Lat), Temp, Phosphate, SSH, cos(Lon), Nitrate, Silicate
P6 Subpolar North Pacific	25	Phosphate, Sal, Depth, Temp, sin(Lat), Silicate, $x\text{CO}_2_{\text{anom}}$ , Alk, Nitrate
P7 Subtropical North Pacific	50	Phosphate, Sal, Temp, Silicate, $N_{\text{mon}}$ , sin(Lat), sin(Lon), Depth, Alk, DIC, Nitrate
P8 Equatorial Pacific	25	Phosphate, Depth, Temp, sin(Lat), Sal, Silicate, $x\text{CO}_2$ , Nitrate, $W_{\text{vel}}(105\text{m})$
P9 Equatorial Indian Ocean	10	Phosphate, Depth, $p\text{CO}_2$ , $W_{\text{vel}}(\text{in situ})$
P10 Subtropical South Atlantic	10	Temp, DIC, Sal, Depth, Nitrate, $W_{\text{vel}}(65\text{m})$ , $p\text{CO}_2$ , $p\text{CO}_2_{\text{clim}}$ , DO, $W_{\text{vel}}(195\text{m})$
P11 Subtropical South Pacific	25	Phosphate, Depth, Temp, $x\text{CO}_2$ , sin(Lat), Silicate, Sal, Alk
P12 Subtropical South Indian Ocean	25	Phosphate, $p\text{CO}_2$ , Depth, Temp, Sal, $p\text{CO}_2_{\text{clim}}$ , Silicate, DO
P13 Subpolar Southern Ocean	50	DIC, Temp, Depth, $N_{\text{mon}}$ , Sal, Alk, DO, Silicate, $P_{\text{surf}}$ , Temp <sub>anom</sub>
P14 Southern Ocean ice	25	cos(Lon), sin(Lat), Depth, DIC, Temp, Sal

(The predictors are arranged in order of relative importance, with the variables listed at the front of each province being more effective in reducing reconstructionpredicting errors when used as pH predictors.)

## 225 2.4 Validation and uncertainty

The reconstructed pH product was validated based on pH measurements from GLODAP and time series stations. First, the root mean square error (RMSE) between the FFNN predicting-pH and GLODAP pH measurements was calculated using the K-fold cross validation method. The GLODAP pH measurements were divided by years, and the K value was 4 to keep

230 aside 25% independent measurements for testing in each one of the total 4 iterations. Thus, within every set of four consecutive years, pH measurements from three years were utilized for training the FFNN model, while the measurements from the remaining year were employed for testing. This approach ensured the independence between the training and testing groups (Gregor et al., 2019; Zhong et al., 2022). Subsequently, the pH measurements in the testing group were compared against the FFNN-predicted pH values based on the training group. A total of 4 iterations were carried out with each iteration designating different years as the testing groups, ensuring that measurements from all years have been set as the test group once and  
235 matched with a FFNN-predicted value. By comparing all FFNN-predicted pH values with GLODAP pH measurements, the RMSE of pH and the molar hydrogen ion concentration ( $[H^+]$ ) was calculated to evaluate the performance of the FFNN model. The reconstruction-predicting of the testing group from the training group is similar to the interpolation process, wherein the FFNN is trained with existing measurements to reconstruct-predicting pH in unknown areas.

240 Second, the reconstructed seawater pH product was compared with independent pH measurements from the Hawaii Ocean Time-series (HOT, 22° 45' N, 158° 00' W, since October 1988) (Dore et al., 2009), Bermuda Atlantic Time-series Study (BAT, 31°50' N, 64°10' W, since October 1988) (Bates et al., 2007; Bates et al., 2020), and The European Station for Time Series in the Ocean Canary Islands (ESTOC, 29°10' N, 15°30' W, from 1995 to 2009) (González-Dávila et al., 2010). The long-term trend was further compared with data from the Irminger Sea station (64.3°N, 28.0°W, from 1983 to 2019, Ólafsson, 2016; Ólafsdóttir et al., 2020a), the Iceland Sea station (68.0°N, 12.7°W, from 1985 to 2019, Ólafsson, 2012; Ólafsdóttir et al., 2020b), and the DYFAMED station (42.3°N, 7.5° E, from 1991 to 2017, Coppola et al., 2024). For better evaluating the performance of FFNN below the surface, the constructed pH product was also compared to independent pH data from the Integrated Marine Observing System Biogeochemical ARGO profiles core data collection (IMOS BGC ARGO, <https://imos.org.au/facilities/argofloats/bgcargo>). Validation based on these independent measurements from time series stations and ARGO profiles provides additional evidence of data accuracy.

250 A comparison between the method of training FFNN with pH and the method of training FFNN with  $[H^+]$  then converting to pH was carried out, to validate which way has a lower pH reconstruction-predicting error (Figure S3). In addition, to identify the difference in pH variability uncertainty hidden by logarithm among regions with the same pH RMSE but different pH level, the uncertainty of reconstructed pH values was converted from  $[H^+]$  RMSE instead of directly using pH RMSE. The pH obtained from the FFNN was first converted to  $[H^+]$  to estimate RMSE. Subsequently, the pH values were shown as  $pH_0 \pm \sigma$  at  
255 each given  $pH_0$  value, and the uncertainty  $\sigma$  stem from FFNN reconstruction-predicting errors was calculated as the following:

$$\sigma = -\log_{10}(10^{-pH_0} - RMSE_{[H^+]}) - pH_0 \quad (2)$$

where  $RMSE_{[H^+]}$  was the RMSE of  $[H^+]$  converted from FFNN pH in each layer of all 14 biogeochemical provinces,  $pH_0$  was the specific predicted-FFNN pH value. The uncertainty  $\sigma$  calculated by this method is simultaneously related to the pH reconstruction-predicting error and pH level, which better distinguishes the differences in uncertainty across different regions.  
260 The uncertainty of products used as pH predictors is one ineluctable source for pH reconstruction-predicting errors of the FFNN model. However, the direct estimation of pH uncertainty from summing the uncertainty of each used product is not feasible, as the relationships between pH and predictors are not established as specific formulas but as matrix manipulations. Therefore, the uncertainty of our pH product was directly estimated from the FFNN pH reconstruction-predicting errors following formula (2), instead of synthesizing the inherent uncertainty of each used predictor product. The inherent uncertainty and construction  
265 method of predictor products are described in the Supplementary text.


### 3 Results and discussion

#### 3.1 Validation of algorithm

##### 3.1.1 Validation based on GLODAP and time series measurements

270 Compared with the GLODAP dataset, most reconstructed predicting values of Stepwise FFNN are close to the GLODAP pH measurements, concentrated around the  $y=x$  line (Figure 4). Only a few samples notably differ between the pH measurements and the reconstructed predicting-values, with the RMSE of 0.028 in the global ocean of 0-2000 m. A better performance of the FFNN was found in the intermediate layer, with testing samples more concentrated on the  $y=x$  line. The RMSE in the mixed layer is 0.034, higher than 0.026 in the intermediate layer. The minor difference between the reconstructed predicting value and the pH measurements and the  $R^2$  of 0.97 in the intermediate layer may be caused by less pH variability at depth and better model fit with broader pH value range suggested a sufficient fitting ability of FFNN.

275



280 **Figure 4: Comparison between FFNN predicting-pH and GLODAP pH measurements.** a): mixed layer from the surface to mixed layer depth; b): intermediate layer from mixed layer depth to 2000 m. Black lines: the  $y=x$  line; red lines: the linear regression between GLODAP pH and Stepwise FFNN pH (Lauvset et al., 2023); Slope: slope of the linear regression.

The RMSE between the FFNN predicting-pH and GLODAP pH measurements at most grids were lower than 0.03 (Figure 5a). The performance of FFNN was relatively better in the temperate oceans, with the RMSE lower than 0.02 at some temperate grids. However, relatively higher RMSE was found in the equatorial and polar oceans, especially in the eastern equatorial Pacific, the near-polar North Pacific, and the northwest Indian Ocean. The RMSE was relatively lower in regions with concentrated GLODAP measurements, such as the near-polar North Atlantic, south Atlantic, and south Indian Ocean.





285

Due to the higher seasonal and interannual variability of seawater pH near the surface ocean, the RMSE decreases with depth in all basins (Figure 5b). At the surface ocean, the RMSE between the FFNN predicting-pH and the GLODAP pH measurements was 0.044. The RMSE fluctuates between 0.032 and 0.048 at the subsurface 0-200 m. The RMSE between the FFNN predicting-pH and the GLODAP pH measurements decreased rapidly from the 200 m depth. In the global ocean 1500-2000 m depth, the global RMSE was lower than 0.015. While at 2000 m depth, the global ocean RMSE at 2000m was only





290




0.013, with the higher RMSE in the Arctic Ocean and the lower in the Southern Ocean. The vertical distribution of the RMSE and statistical distribution of pH difference in different basins all suggested a relatively higher **reconstruction**~~predicting~~ error in the mixed layer than in the intermediate layer (Figure 5d). The vertical difference of RMSE between the mixed layer and intermediate layer was most notable in the Arctic and Indian Ocean, where the RMSE at different depths was also higher than the other basins. The RMSE in the surface Arctic Ocean was higher than 0.10 and decreased rapidly to 0.025 by 450 m depth. On the contrary, the RMSE of the surface Indian Ocean was ~~low, only~~ 0.018, but increased to 0.053 by 80 m depth and then decreased continuously with depth. The high RMSE of subsurface oceans is because there are almost no GLODAP pH measurements in the entire Indian Ocean at 50-150 m depth. The RMSE in different years also suggested the notable influence of pH measurement amount on the FFNN **reconstruction**~~predicting~~ errors. The RMSE in the early years was relatively higher than in recent years, while the number of GLODAP measurements increased with the years (Figure 5c).







305 **Figure 5: Distribution of RMSE between FFNN predicted-pH values and GLODAP pH measurements.** a): global spatial distribution of RMSE between FFNN predicted-pH and GLODAP pH measurements at 0-2000 m (Lauvset et al., 2023); b): basin average RMSE at different depth; c): temporal distribution of global RMSE; d): Statistical distribution of pH difference between predicted-reconstructed pH values and GLODAP pH measurements in each basin.



310 **Figure 6: Comparison between FFNN predicted-pH and time-series measurements.** a-b): pH value, pH difference and its distribution, pH seasonal variability of FFNN result and time series measurements at the BAT station; c-d): the ESTOC station; e-f): the HOT station.

The Stepwise FFNN pH product showed variability of seawater pH close to the independent time series observations in the surface ocean from HOT, ESTOC, and BAT stations (Figure 6). At the BAT station, the reconstructed pH product and time series observations was only 0.013. The surface seawater pH of our Stepwise FFNN product decreased by  $0.0017 \pm 0.0007 \text{ yr}^{-1}$  on average during the past three decades at the BAT station, close to the  $-0.0018 \pm 0.0001 \text{ yr}^{-1}$  of BAT time series observations in the same period (Bates et al., 2020). At the ESTOC station, the Stepwise FFNN product and time series observations were also well consistent, with the RMSE of only 0.009 and a similar long-term trend (González-Dávila et al., 2010; Chau et al., 2022). The RMSE between the Stepwise FFNN product and HOT time series observations was also only 0.010, and the long-term trends of the Stepwise FFNN pH product was  $0.0018 \pm 0.0004 \text{ yr}^{-1}$ , consistent with the HOT time series observations. Although at the BAT station, the Stepwise FFNN product suggested a smaller seasonal change scale than the time series observations, the seasonal patterns of surface seawater pH were consistent between the Stepwise FFNN product and time series observations at all three stations. The extreme values not reconstructed by the FFNN are mainly observed at the BAT station near 2010 and at the HOT station near 2000 during La Niña events, and at the HOT station before 2000 during El Niño events. Differently, the extreme values not reconstructed by the FFNN are less observed at the ESTOC station, where the surface pH did not notably fluctuate during El Niño/La Niña events. It can be inferred that the extreme values not reconstructed by the FFNN may be due to its underestimating of the impact of El Niño/La Niña events on pH of certain temperate areas. Compared to previous surface ocean seawater pH product, which were derived from reconstructed DIC, TA,

or  $p\text{CO}_2$  products, the Stepwise FFNN product had lowest surface RMSE at the ESTOC station, and similar RMSE values to other products at the BAT and HOT stations was consistent with the pH trend from the majority of time series stations (Table 4). The long-term pH trend of our product at the ESTOC station differed from was slower than other gridded products at the ESTOC station, but the result is still close to the  $-0.0016 \pm 0.0001 \text{ yr}^{-1}$  of real observations. In the Irminger Sea station, the FFNN pH trend was notably faster than the result of time series observations. However, differences in pH trend among pH products were most remarkable in this station. On the global scale, the pH trend of our FFNN product is  $-0.0015 \pm 0.0002$  over the period from 1992 to 2020. There is no significant difference between our FFNN product, the CMEMS product, and the Copernicus product under the current uncertainty.

**Table 4: Comparison of surface acidification rate with previous product in different time series stations and on a global scale.**

Stations	Period	Time series observation	Stepwise FFNN (This study)	JMA (Iida et al., 2021)	CMEMS (Chau et al., 2024)	OS-ETHZ (Gregor et al., 2021)	Copernicus (Copernicus Marine Service, 2020)
BAT	1992~2020	$-0.0018 \pm 0.0001$	$-0.0017 \pm 0.0007$	$-0.0018 \pm 0.0002$	$-0.0018 \pm 0.0002$	$-0.0018 \pm 0.0002$	-
ESTOC	1995~2010	$-0.0016 \pm 0.0001$	$-0.0014 \pm 0.0005$	$-0.0022 \pm 0.0003$	$-0.0020 \pm 0.0002$	$-0.0017 \pm 0.0003$	-
HOT	1992~2020	$-0.0018 \pm 0.0001$	$-0.0018 \pm 0.0004$	$-0.0020 \pm 0.0001$	$-0.0021 \pm 0.0001$	$-0.0019 \pm 0.0001$	-
Iceland Sea	1992~2019	$-0.0020 \pm 0.0004$	$-0.0028 \pm 0.0002$	$-0.0030 \pm 0.0003$	$-0.0015 \pm 0.0002$	$-0.0020 \pm 0.0002$	-
Irminger Sea	1992~2019	$-0.0025 \pm 0.0004$	$-0.0022 \pm 0.0002$	$-0.0027 \pm 0.0002$	$-0.0017 \pm 0.0003$	$-0.0016 \pm 0.0003$	-
DYFAMED	1998~2017	$-0.0010 \pm 0.0008$	$-0.0005 \pm 0.0003$	-	$-0.0017 \pm 0.0003$	$-0.0023 \pm 0.0004$	-
Global	1992~2020	-	$-0.0015 \pm 0.0002$	$-0.0018 \pm 0.0000$	$-0.0017 \pm 0.0004$	$-0.0018 \pm 0.0000$	$-0.0017 \pm 0.0002$

(RMSE: root mean square error between pH products and time series observations; the trend from different products for comparison were recalculated based on data during same period noted in the second column;


Stepwise FFNN product: reconstructed from pH measurements with  $1^\circ \times 1^\circ$  and monthly resolution from 1992 to 2020, covering global open ocean 0-2000 m;

JMA product: reconstructed from DIC and Alk with a  $1^\circ \times 4^\circ$  and monthly resolutions from 1990 to 2021+2022, covering global surface ocean except a portion of the Arctic;

CMEMS product: reconstructed from  $p\text{CO}_2$  and Alk with a  $1^\circ$  or  $0.25^\circ$  resolutions  $\times 4^\circ$  and monthly resolution from 1985 to 2021, covering global surface ocean except a portion of the Arctic;

OS-ETHZ product: reconstructed from  $p\text{CO}_2$  and Alk with a  $1^\circ \times 4^\circ$  and monthly resolutions from 1982 to 2022, covering global surface ocean except the Arctic; Copernicus product: mean sea water pH time series and trend from Multi-Observations Reprocessing, from 1985 to 2021)

Compared with the time series data below the surface, the FFNN pH was close to the pH observations at upper few hundred meters in the BAT and HOT station (Figure 7). However, higher RMSE and larger ranges of pH difference were observed at 500-1500 m in the BAT station and below 300 m in the HOT station. This may be due to the sparser GLODAP observations used to train the FFNN model in these areas. Additionally, as depth is used as a pH predictor, in the validation based on the GLODAP dataset the FFNN pH values used for validation were outputted at the same depth of the GLODAP observations. When comparing FFNN pH with independent time series observations, differences in depth between the pH product and the observations can amplify the calculated pH difference and RMSE. For example, the FFNN pH product was reconstructed at depths of 1800 m and 2000 m in the bottom. If the time series observation is at 1910 m depth, it will be compared with the FFNN pH value at 2000 m in the independent validation. This depth difference significantly increases the pH error in validation based on independent data. Despite higher RMSE at certain depths, the RMSE at most depths in the deep areas of the BAT station and DYFAMED station is below 0.03, indicating that the notable deviations may only occur at local scale.




365 **Figure 7. RMSE and pH difference between FFNN pH and time series observations at different depths. a) BAT station at 31°50' N, 64°10' W based on data from 1992 to 2020; b) HOT station at 22° 45' N, 158° 00' W based on data from 1992 to 2020; c) DYFMED station at 42.3°N, 7.5° E based on data from 1998 to 2017.**

### 3.1.2 Comparison of performance between FFNNs training based on pH and $[H^+]$

370 Due to the logarithmic relationship between pH value and  $[H^+]$  concentration, results obtained from training FFNN with pH and from training FFNN with  $[H^+]$  then converting outputs into pH may differ. A comparison of predicting errors was conducted between these two training methods. The results show a nearly consistent pH RMSE between the FFNN training with pH and with  $[H^+]$  (Figure 7). As the pH measurements of all GLODAP samples are closer to a normal distribution than the  $[H^+]$ , the predicting error was slightly lower in most regions when the FFNN was trained with pH, but the difference in predicting errors was extremely small. In addition, the FFNN trained using  $[H^+]$  occasionally produced negative  $[H^+]$  in regions with extremely low  $[H^+]$ . Therefore, it is better to train FFNN using pH rather than using  $[H^+]$  in the reconstruction process of the pH product.


375 The distribution patterns of regional pH RMSE and  $[H^+]$  RMSE are inconsistent whenever the FFNN was trained using pH or  $[H^+]$ . In fact, the pH RMSE of the intermediate layer in regions such as the subarctic North Pacific and the equatorial Pacific is significantly lower than that in the intermediate layer of the Arctic Ocean, but their  $[H^+]$  RMSE is higher than that of the intermediate layer in the Arctic Ocean (Figure 7a and 7b). This is caused by the effect of the logarithmic relationship. If the pH values are different for the same pH RMSE, the corresponding  $[H^+]$  RMSE will be different. Therefore, the uncertainty of the pH product is calculated based on the  $[H^+]$  RMSE and pH value, rather than solely based on the pH RMSE.



385 **Figure 7: Comparison of pH RMSE and  $[H^+]$  RMSE from training FFNN using pH and using  $[H^+]$ . a) pH RMSE of FFNN trained using pH and  $[H^+]$  in each biogeochemical province, the predicted  $[H^+]$  from FFNN trained using  $[H^+]$  was converted to pH for estimating pH RMSE. b)  $[H^+]$  RMSE of FFNN trained using pH and  $[H^+]$  in each biogeochemical province; c)  $[H^+]$  RMSE of FFNN trained using pH and  $[H^+]$  in each vertical layer; the predicted pH from FFNN trained using pH was converted to  $[H^+]$  for estimating  $[H^+]$  RMSE. The numbers shown in the X axis represent the SOM province in Figure 1.**

390 **3.1.2 Validation based on biogeochemical ARGO float pH measurements**

Comparison with time series observations in deeper oceans suggested that the distribution of pH reconstruction errors with depth varies notably across different stations. To better assess the performance of FFNN in the reconstruction of pH at different depths, the FFNN reconstructed pH was further evaluated by comparing with independent pH data at various depths from the IMOS BGC ARGO dataset (IMOS 2002-2020, 2021). Different from the validation results based on the GLODAP dataset, the RMSE between FFNN pH and BGC ARGO pH data in the intermediate layer is 0.055, higher than the 0.033 in the mixed layer (Figures 8a and 8b). In both the mixed layer and intermediate layer, most samples were evenly distributed around the  $y=x$  line. However, in the intermediate layer, some samples were slightly offset and distributed below the  $y=x$  line, which may be the main reason for the notably higher RMSE between FFNN pH and BGC ARGO pH data in the intermediate layer. Overall, there is a good linear correlation between FFNN reconstructed pH and independent BGC ARGO pH data, with  $R^2$  values of 0.78 and 0.77 in the mixed layer and intermediate layer, respectively.



405 **Figure 8. Difference between FFNN pH and BGC ARGO floats pH.** a) comparison between FFNN pH and BGC ARGO floats pH in the mixed layer; b) comparison between FFNN pH and BGC ARGO floats pH in the intermediate layer; c) Statistical distribution of pH difference (FFNN pH minus BGC ARGO floats pH) at different depth levels. FFNN pH: pH data reconstructed in this work; BGC ARGO floats pH: pH data from IMOS Biogeochemical ARGO floats core data collection (IMOS 2002-2020, 2021).

The distribution of pH differences between FFNN pH and BGC ARGO pH data at different depths reveals a relatively smaller biases above 500 m (Figure 8c). However, below 500 m, the bias between FFNN pH and BGC ARGO pH data increases with depth and was the most remarkable at 2000 m. Comparing the pH bias calculated based on BGC ARGO dataset and GLODAP dataset, it is evident that only the bias between FFNN pH and BGC ARGO pH data tends to increase with depth in most basins except the Pacific Ocean. In contrast, greater biases between FFNN pH and GLODAP pH occur mainly in the surface layer, with the most in the surface Indian ocean. This disparity in distribution patterns between biases based on BGC ARGO dataset and GLODAP dataset is most remarkable in the Southern Ocean, where the bias between FFNN pH and GLODAP pH is nearly zero below 1000 m, compared to biases between FFNN pH and BGC ARGO pH data ranging from 0.053 to 0.076. These differences between FFNN pH and BGC ARGO pH data are primarily attributed to the discrepancies between GLODAP dataset and the BGC ARGO dataset in the deep ocean, as our product was based on the GLODAP dataset and small biases with GLODAP pH were observed in the deep ocean.

**Table 5. pH bias by area and depth computed with BGC ARGO and GLODAP dataset.**

Area		0-50 m	50-200 m	200-500 m	500-1000 m	1000-1500 m	1500-2000 m
Arctic	BGC ARGO	-0.006	0.021	0.014	0.026	0.037	0.029
	GLODAP	-0.005	-0.003	0.001	0.000	0.002	0.003
Pacific	BGC ARGO	0.012	0.011	-0.008	-0.015	0.008	0.000




	<u>GLODAP</u>	<u>-0.001</u>	<u>-0.001</u>	<u>0.000</u>	<u>0.000</u>	<u>0.000</u>	<u>-0.001</u>
<u>Atlantic</u>	<u>BGC ARGO</u>	<u>0.016</u>	<u>0.017</u>	<u>0.010</u>	<u>-0.024</u>	<u>0.029</u>	<u>0.068</u>
	<u>GLODAP</u>	<u>0.000</u>	<u>0.000</u>	<u>-0.001</u>	<u>-0.001</u>	<u>0.000</u>	<u>0.000</u>
<u>Indian</u>	<u>BGC ARGO</u>	<u>0.024</u>	<u>0.026</u>	<u>0.014</u>	<u>-0.056</u>	<u>0.002</u>	<u>0.059</u>
	<u>GLODAP</u>	<u>-0.006</u>	<u>-0.001</u>	<u>-0.003</u>	<u>-0.004</u>	<u>-0.004</u>	<u>-0.001</u>
<u>Southern</u>	<u>BGC ARGO</u>	<u>0.011</u>	<u>0.002</u>	<u>0.002</u>	<u>0.027</u>	<u>0.053</u>	<u>0.076</u>
	<u>GLODAP</u>	<u>0.004</u>	<u>0.001</u>	<u>0.001</u>	<u>0.000</u>	<u>0.000</u>	<u>0.000</u>
<u>Global</u>	<u>BGC ARGO</u>	<u>0.011</u>	<u>0.005</u>	<u>0.001</u>	<u>0.021</u>	<u>0.046</u>	<u>0.066</u>
	<u>GLODAP</u>	<u>-0.001</u>	<u>0.000</u>	<u>0.000</u>	<u>-0.001</u>	<u>-0.001</u>	<u>0.000</u>

## 420 3.2 Gridded pH product

### 3.2.1 Spatial pH distribution

The spatial distribution of long-term average seawater pH in the ~~constructed~~ Stepwise FFNN product suggests the lowest surface seawater pH in the equatorial Pacific with an average value near 8.00 (Figure [8a9a](#)), which is in good agreement with the surface seawater pH range of 7.91-8.12 observed in the equatorial Pacific in recent decades (Sutton et al., 2014). The upwelling transporting the deep water with high dissolved inorganic carbon and low pH to the surface was the main driver. The equatorial Indian Ocean and the equatorial Atlantic also show a low surface pH of about 8.05, consistent with the distribution patterns of the GLODAP pH climatology (Lauvset et al., 2016). The highest surface pH is found in the Atlantic sector of the Arctic Ocean, where the average surface pH was around 8.15 during the past three decades. Besides, the average surface pH in temperate oceans is relatively higher, such as the south Indian and south Atlantic Oceans. In the ~~temperate~~ Pacific Ocean, ~~differences in surface pH levels were observed between the west and east in both our product and GLODAP pH climatology, which may be caused by the spread of eastern equatorial seawater with extremely low pH the high surface pH is mainly concentrated in the west, while the average surface pH in the East temperate Pacific was about 8.05.~~ At the deeper depth of 1000 m, the spatial distribution pattern of FFNN pH product is generally consistent with the GLODAP climatology, despite still some disturbance of bad FFNN performance along the SOM province boundary ~~and the higher FFNN pH in the~~ [Southern Ocean](#).






**Figure 89: Average pH distribution from FFNN pH product and GLODAP climatology normalized to the year 2002.** The GLODAP climatology data is from Lauvset et al., 2016.

440 The vertical distribution of average pH in the ~~proposed~~~~constructed~~ product showed a notable pH decrease with increasing depth in the upper 500 m of different basins (Figure 910). The seawater pH was the lowest at nearly 500 m and rose with increasing depth at 500-2000 m in the Pacific and Atlantic oceans. The distribution pattern of seawater pH in the Indian Ocean was similar to that in the South Pacific, with the lowest seawater pH appearing near 1000 m. The subsurface seawater with low pH in the Atlantic Ocean and Indian Ocean was mainly concentrated in the equatorial region. In contrast, subsurface seawater with low pH in the Pacific Ocean appeared in subpolar and equatorial regions. The overall distribution pattern of the ~~reconstructed~~ pH ~~product~~ is in good agreement with previous research (Lauvset et al., 2016; Lauvset et al., 2020). It can be concluded that the FFNN fitted the relationship between GLODAP seawater pH and its predictors well, and the ~~proposed~~ ~~constructed~~ pH product has good accuracy.

450 Based on the pH predictors selected by the Stepwise FFNN algorithm, differences in processes driving pH variability were identified between the mixed layer and intermediate layer in most provinces. In the mixed layer, surface ocean  $p\text{CO}_2$  was identified as the most informative predictor in many provinces, followed by temperature and nutrient concentration. This suggests that the  $\text{CO}_2$  exchange between surface ocean and atmosphere is the primary driver of pH variability, followed by biological  $\text{CO}_2$  utilization and seasonal changes in seawater temperature. In contrast, phosphate was identified as the most informative predictor in the intermediate layer, followed by temperature and depth. This suggests that the primary process driving pH variability is the remineralization of organic matter, converting organic carbon into inorganic forms and also releasing nitrogen and phosphorus. Given the notably smaller seasonal temperature changes in the intermediate layer compared to the mixed layer, the selection of temperature as an important pH predictor may indicate a notable influence of ocean warming on seawater pH variability. Additionally, depth was also selected as an important predictor in the intermediate layer. The observed pattern of seawater pH decreasing with increasing depth in most provinces, as suggested by the constructed pH product, may be the main reason.

460



**Figure 910:** Climatological vertical distribution of zonal average FFNN pH in main basins. The pH values shown at each latitude were averaged from pH values across all longitudes within each major basin.


### 3.2.2 Uncertainty

465 As described in the method section, the FFNN ~~predicted~~ pH was converted to  $[H^+]$  to calculate the regional RMSE of  $[H^+]$  between FFNN results and GLODAP measurements, and then the RMSE of  $[H^+]$  in each SOM province was used to calculate the pH product uncertainty caused by the construction algorithm (Equation 2). Due to higher ~~reconstruction~~~~predicting~~ errors, the pH product uncertainty is relatively higher near the surface (Figure 4011). The uncertainty is generally lower than 0.02 at depths from 500 m to 2000 m, except for some regions near the SOM province and vertical boundary. Although we have used

470 a cross-boundary method to improve the FFNN performance near the SOM and vertical boundary, there are still some discontinuity problems and relatively higher uncertainty. This is because the pH values on two sides of the SOM boundary were reconstructed from two different FFNN models, which were trained with different samples and used different predictors. If one of the FFNN models experienced a worse performance due to insufficient training samples or predictors, the pH values on two sides of the SOM boundary will still differ notably, that is, discontinuity along the boundary. Therefore, the analyze on

475 a regional scale based on pH values near SOM boundaries should be more cautious when using our product. In addition, the equatorial and polar regions show an uncertainty higher than 0.04. This is because the FFNN performance tends to be worse in regions with the highest and lowest pH levels than in regions where pH values are near the average level. Especially in the Arctic Ocean, the pH measurements are much sparser leading to the highest ~~reconstruction~~~~predicting~~ error and pH uncertainty. Therefore, the ~~proposed constructed~~ pH product should be cautiously used in regional analysis near the boundaries or equatorial

480 and polar regions.



**Figure 1011:** Uncertainty of FFNN pH product in main basins.

#### 4 Data availability

The materials used in this research including the gridded seawater pH product (NetCDF files for all individual years), MATLAB code for reconstruction and validation, and other materials (available as .m or .mat files) are available from the Marine Science Data Center of the Chinese Academy of Sciences at <http://dx.doi.org/10.12157/IOCAS.20230720.001> (Zhong et al., 2023). The used pH measurements are available from GLODAP (<https://glodap.info/index.php/merged-and-adjusted-data-product-v2-2023/>, Lauvset et al., 20222023), data products used for predictors are available from references listed in Table 1.

#### 490 5 Conclusions

Quantifying the global seawater pH variability is important for understanding the future responses of oceans on the uptake of anthropogenic CO<sub>2</sub>. A four-dimensional global seawater pH product covering depths from the surface to 2000 m and years from 1992 to 2020 was reconstructed in this work. This product serves as a reference for guiding acidification surveys by providing a general understanding of acidification process at different depths on a basin scale and indicating areas with potential fast or slow acidification rates. Additionally, the pH product brings insights into acidification research and can be used to analyze the influence of specific ocean processes on acidification rates and the broader impacts of acidification on a large scale when direct observations are unavailable, but still should be used carefully in regional analyses. However, caution should be exercised when using this product for regional analyses at a small spatial scale. The analysis of pH RMSE and

uncertainty suggested that the ~~proposed~~ reconstructed pH product remains limited in equatorial and polar regions and along the SOM boundary lines. This limitation was caused by sparse measurements and method disadvantages, which can be mitigated in future improvement works. Potential improvement may be achieved by increasing more predictor products to capture the pH drivers, testing more machine learning algorithms, and accumulating more seawater pH observations. Besides, the method used to reconstruct the pH product can be applied in ~~constructing~~ reconstruction of global fields of other ocean chemical variables, such as nutrients, particulate organic carbon, and dissolved inorganic carbon. The global field of these variables may further improve the pH product accuracy, as climatological products of these variables were used as pH predictors and lacked interannual variability information. Overall, decreasing seawater pH will influence the metabolism of marine organisms and result in notable changes to the marine ecosystem. The discrete observations may be insufficient to support research on large scales. With the machine learning method in this work, the discrete pH measurements were mapped to global gridded fields to fill the unsampled areas. Our product can be used for analysis of seasonal to decadal and regional to global pH variability, to break through the limitation of discrete observations.

### Acknowledgments

We thank the data support of the Marine Science Data Center and Public Technical Service Center, Institute of Oceanology, Chinese Academy of Sciences. We thank GLODAP for sharing the pH observation data. We thank the support by the National Natural Science Foundation of China (grant nos.42176200); [National Key Research and Development Program \(2022YFC3104305\)](#); Laboratory for Marine Ecology and Environmental Science, Qingdao National Laboratory for Marine Science and Technology (LSKJ202204001, LSKJ202205001); ~~National Key Research and Development Program (2022YFC3104305)~~; Shandong Province and Yantai City Talent Programs; and Science Fund for Creative Research Groups of the National Natural Science Foundation of China (42221005).

### Author contributions:

Data product collection: Baoxiao Qu, Yanjun Wang, Bin Zhang; Data product synthesize: Jun Ma, Qidong Wang, Jianwei Xing; Methodology: Guorong Zhong, Jiajia Dai, Liqin Duan, Ning Li; Model improve: Xuegang Li, Jinming Song, Huamao Yuan; Writing – original draft: Guorong Zhong, Xuegang Li; Writing – review & editing: Jinming Song, Fan Wang, Lijing Cheng.

**Competing interests:** Authors declare that they have no competing interests.

### References

- Bates, N. R. Interannual variability of the oceanic CO<sub>2</sub> sink in the subtropical gyre of the North Atlantic Ocean over the last 2 decades. *Journal of Geophysical Research: Oceans*, 112, C9, <https://doi.org/10.1029/2006JC003759>, 2007.
- Bates, N. R., Astor, Y. M., Church, M. J., Currie, K., Dore, J. E., González-Dávila, M., ... & Santana-Casiano, J. M. A time-series view of changing surface ocean chemistry due to ocean uptake of anthropogenic CO<sub>2</sub> and ocean acidification. *Oceanography*, 27, 126-141, <https://doi.org/10.5670/oceanog.2014.16>, 2014.
- Bates, N. R., & Johnson, R. J. Acceleration of ocean warming, salinification, deoxygenation and acidification in the surface subtropical North Atlantic Ocean. *Communications Earth & Environment*, 1, 33, <https://doi.org/10.1038/s43247-020-00030-5>, 2020.
- Broullón, D., Pérez, F. F., Velo, A., Hoppema, M., Olsen, A., Takahashi, T., ... & van Heuven, S. M. A global monthly climatology of total alkalinity: a neural network approach. *Earth System Science Data*, 11, 1109-1127, <https://doi.org/10.5194/essd-11-1109-2019>, 2019.
- Broullón, D., Pérez, F. F., Velo, A., Hoppema, M., Olsen, A., Takahashi, T., ... & Kozyr, A. A global monthly climatology of oceanic total dissolved inorganic carbon: a neural network approach. *Earth System Science Data*, 12, 1725-1743, <https://doi.org/10.5194/essd-12-1725-2020>, 2020.



- 540 Caldeira, K., & Wickett, M. E. Anthropogenic carbon and ocean pH. *Nature*, 425, 365-365, <https://doi.org/10.1038/425365a>, 2003.
- Chau T T T, Gehlen M, Chevallier F. A seamless ensemble-based reconstruction of surface ocean  $p\text{CO}_2$  and air-sea  $\text{CO}_2$  fluxes over the global coastal and open oceans. *Biogeosciences*, 19, 1087-1109, <https://doi.org/10.5194/bg-19-1087-2022>, 2022.
- 545 [Chau, T.-T.-T., Gehlen, M., Metzl, N., and Chevallier, F.: CMEMS-LSCE: a global, 0.25°, monthly reconstruction of the surface ocean carbonate system, \*Earth Syst. Sci. Data\*, 16, 121–160, <https://doi.org/10.5194/essd-16-121-2024>, 2024.](https://doi.org/10.5194/essd-16-121-2024)
- Chen, C. T. A., Lui, H. K., Hsieh, C. H., Yanagi, T., Kosugi, N., Ishii, M., & Gong, G. C. Deep oceans may acidify faster than anticipated due to global warming. *Nature Climate Change*, 7, 890-894, <https://doi.org/10.1038/s41558-017-0003-y>, 2017.
- Cheng, L., & Zhu, J. Benefits of CMIP5 multimodel ensemble in reconstructing historical ocean subsurface temperature variations. *Journal of Climate*, 29, 5393-5416, <https://doi.org/10.1175/JCLI-D-15-0730.1>, 2016.
- 550 Cheng, L., Trenberth, K. E., Fasullo, J., Boyer, T., Abraham, J., & Zhu, J. Improved estimates of ocean heat content from 1960 to 2015. *Science Advances*, 3, e1601545, <https://doi.org/10.1126/sciadv.1601545>, 2017.
- Cheng, L., Trenberth, K. E., Gruber, N., Abraham, J. P., Fasullo, J. T., Li, G., ... & Zhu, J. Improved estimates of changes in upper ocean salinity and the hydrological cycle. *Journal of Climate*, 33, 10357-10381, <https://doi.org/10.1175/JCLI-D-20-0366.1>, 2020.
- 555 Climate Prediction Center. Daily Arctic Oscillation Index. [Accessed on 2023/08/20]. [https://www.cpc.ncep.noaa.gov/products/precip/CWlink/daily\\_ao\\_index/ao\\_index.html](https://www.cpc.ncep.noaa.gov/products/precip/CWlink/daily_ao_index/ao_index.html). (2002).
- Climate Prediction Center. Southern Oscillation Index. [Accessed on 2023/08/20]. [https://www.cpc.ncep.noaa.gov/products/analysis\\_monitoring/ensocycle/soi.shtml](https://www.cpc.ncep.noaa.gov/products/analysis_monitoring/ensocycle/soi.shtml). (2005).
- 560 [Copernicus Marine Service. Global Ocean acidification - mean sea water pH time series and trend from Multi-Observations Reprocessing \[Dataset\]. Mercator Ocean International. <https://doi.org/10.48670/MOI-00224>, 2020.](https://doi.org/10.48670/MOI-00224)
- [Coppola L., Diamond R. E., Carval T., Irisson J., and Desnos C.: Dyfamed observatory data, SEANOE \[data set\], <https://doi.org/10.17882/43749>, 2024.](https://doi.org/10.17882/43749)
- Dore, J. E., Lukas, R., Sadler, D. W., Church, M. J., & Karl, D. M. Physical and biogeochemical modulation of ocean acidification in the central North Pacific. *Proceedings of the National Academy of Sciences*, 106, 12235-12240, <https://doi.org/10.1073/pnas.0906044106>, 2009.
- 565 Fay, A. R., & McKinley, G. A. Global trends in surface ocean  $p\text{CO}_2$  from in situ data. *Global Biogeochemical Cycles*, 27, 541-557, <https://doi.org/10.1002/gbc.20051>, 2013.
- Feely, R. A., Sabine, C. L., Lee, K., Berelson, W., Kleypas, J., Fabry, V. J., & Millero, F. J. Impact of anthropogenic  $\text{CO}_2$  on the  $\text{CaCO}_3$  system in the oceans. *Science*, 305, 362-366, <https://doi.org/10.1126/science.1097329>, 2004.
- 570 Feely, R. A., Doney, S. C., & Cooley, S. R. Ocean acidification: Present conditions and future changes in a high- $\text{CO}_2$  world. *Oceanography*, 22, 36-47, <https://doi.org/10.5670/oceanog.2009.95>, 2009.
- Friedlingstein, P., O'sullivan, M., Jones, M. W., Andrew, R. M., Bakker, D. C., Hauck, J., ... & Zheng, B. Global carbon budget 2023. *Earth System Science Data*, 15(12), 5301-5369, <https://doi.org/10.5194/essd-15-5301-2023>, 2023.
- Garcia He, Weathers Kw, Paver Cr, Smolyar I, Boyer Tp, Locarnini Mm, Zweng Mm, Mishonov Av, Baranova Ok, Seidov
- 575 D, Reagan Jr. World Ocean Atlas 2018, Volume 3: Dissolved Oxygen, Apparent Oxygen Utilization, and Dissolved Oxygen Saturation. NOAA Atlas NESDIS 83, 38pp, <https://archimer.ifremer.fr/doc/00651/76337/>, 2019a.
- Garcia He, Weathers Kw, Paver Cr, Smolyar I, Boyer Tp, Locarnini Mm, Zweng Mm, Mishonov Av, Baranova Ok, Seidov D, Reagan Jr. World Ocean Atlas 2018. Vol. 4: Dissolved Inorganic Nutrients (phosphate, nitrate and nitrate+nitrite, silicate). NOAA Atlas NESDIS 84, 35pp, <https://archimer.ifremer.fr/doc/00651/76336/>, 2019b.
- 580 GEBCO Compilation Group. GEBCO\_2022 Grid. [Accessed on 2022/12/01]. [https://www.gebcoset.net/data\\_and\\_products/gridded\\_bathymetry\\_data/#global](https://www.gebcoset.net/data_and_products/gridded_bathymetry_data/#global), 2022.

- González-Dávila, M., Santana-Casiano, J. M., Rueda, M. J., & Llinás, O. The water column distribution of carbonate system variables at the ESTOC site from 1995 to 2004. *Biogeosciences*, 7, 3067–3081, <https://doi.org/10.5194/bg-7-3067-2010>, 2010.
- Guallart, E. F., Fajar, N. M., Padín, X. A., Vázquez-Rodríguez, M., Calvo, E., Ríos, A. F., ... & Pérez, F. F. Ocean acidification along the 24.5° N section in the subtropical North Atlantic. *Geophysical Research Letters*, 42, 450-458, <https://doi.org/10.1002/2014gl062971>, 2015.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., ... & Thépaut, J. N. The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*, 146(730), 1999-2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 590 Iida, Y., Takatani, Y., Kojima, A., & Ishii, M. Global trends of ocean CO<sub>2</sub> sink and ocean acidification: an observation-based reconstruction of surface ocean inorganic carbon variables. *Journal of Oceanography*, 77, 323-358, <https://doi.org/10.1007/s10872-020-00571-5>, 2021.
- [IMOS 2002-2020, Argo Profiles - biogeochemical data. https://imos.org.au/facilities/argofloats/bgcargo](https://imos.org.au/facilities/argofloats/bgcargo), accessed 2024-07-30, 2021.
- 595 Ishizu, M., Miyazawa, Y., & Guo, X. Long-term variations in ocean acidification indices in the Northwest Pacific from 1993 to 2018. *Climatic Change*, 168, 1-20, <https://doi.org/10.1007/s10584-021-03239-1>, 2021.
- [Jiang, Z.; Song, Z.; Bai, Y.; He, X.; Yu, S.; Zhang, S.; Gong, F. Remote Sensing of Global Sea Surface pH Based on Massive Underway Data and Machine Learning. Remote Sensing, 14, 2366, https://doi.org/10.3390/rs14102366, 2022.](https://doi.org/10.3390/rs14102366)
- [Keppler, L., Landschützer, P., Gruber, N., Lauvset, S. K., & Stemmler, I. Seasonal carbon dynamics in the near-global ocean. Global Biogeochemical Cycles, 34, e2020GB006571, https://doi.org/10.1029/2020GB006571, 2020.](https://doi.org/10.1029/2020GB006571)
- 600 Lan, X., Tans, P. & K.W. Thoning. Trends in globally-averaged CO<sub>2</sub> determined from NOAA Global Monitoring Laboratory measurements. <https://gml.noaa.gov/ccgg/trends/>, 2023.
- Landschützer, P., Gruber, N., Bakker, D. C., & Schuster, U. Recent variability of the global ocean carbon sink. *Global Biogeochemical Cycles*, 28(9), 927-949, <https://doi.org/10.1002/2014gb004853>, 2014.
- 605 Landschützer, P., Laruelle, G. G., Roobaert, A., & Regnier, P. A uniform pCO<sub>2</sub> climatology combining open and coastal oceans. *Earth System Science Data*, 12, 2537-2553, <https://doi.org/10.5194/essd-2020-90>, 2020.
- Lauvset, S. K., Gruber, N., Landschützer, P., Olsen, A., & Tjiputra, J. Trends and drivers in global surface ocean pH over the past 3 decades. *Biogeosciences*, 12, 1285-1298, <https://doi.org/10.5194/bg-12-1285-2015>, 2015.
- Lauvset, S. K., Key, R. M., Olsen, A., Van Heuven, S., Velo, A., Lin, X., ... & Watelet, S. A new global interior ocean mapped climatology: The 1 × 1 GLODAP version 2. *Earth System Science Data*, 8, 325-340, <https://doi.org/10.5194/essd-8-325-2016>, 2016.
- 610 Lauvset, S. K., Carter, B. R., Pérez, F. F., Jiang, L. Q., Feely, R. A., Velo, A., & Olsen, A. Processes driving global interior ocean pH distribution. *Global Biogeochemical Cycles*, 34(1), e2019GB006229, <https://doi.org/10.1029/2019gb006229>, 2020.
- Lauvset, S. K., Lange, N., Tanhua, T., Bittig, H. C., Olsen, A., Kozyr, A., ... & Key, R. M. GLODAPv2. 2022: the latest version of the global interior ocean biogeochemical data product. *Earth System Science Data*, 14, 5543-5572, <https://doi.org/10.5194/essd-14-5543-2022>, 2022.
- 615 Le Quéré, C., Takahashi, T., Buitenhuis, E. T., Rödenbeck, C., & Sutherland, S. C. Impact of climate change and variability on the global oceanic sink of CO<sub>2</sub>. *Global Biogeochemical Cycles*, 24, 4, <https://doi.org/10.1029/2009GB003599>, 2010.
- Lewis, E., Wallace, D., & Allison, L. J. (1998). Program developed for CO<sub>2</sub> system calculations (No. ORNL/CDIAC-105). Brookhaven National Lab., Dept. of Applied Science, Upton, NY (United States); Oak Ridge National Lab., Carbon Dioxide Information Analysis Center, TN (United States).
- 620 Li, L., Chen, B., Luo, Y., Xia, J., & Qi, D. Factors controlling acidification in intermediate and deep/bottom layers of the Japan/East Sea. *Journal of Geophysical Research: Oceans*, 127, e2021JC017712, <https://doi.org/10.1029/2021jc017712>, 2022.



- Luo, Y., Boudreau, B. P., & Mucci, A. Disparate acidification and calcium carbonate desaturation of deep and shallow waters of the Arctic Ocean. *Nature Communications*, 7, 12821, <https://doi.org/10.1038/ncomms12821>, 2016.
- Menemenlis, D., Campin, J. M., Heimbach, P., Hill, C., Lee, T., Nguyen, A., ... & Zhang, H. ECCO2: High resolution global ocean and sea ice data synthesis. *Mercator Ocean Quarterly Newsletter*, 31, 13-21, <https://doi.org/>, 2008.
- NASA Ocean Biology Processing Group. Aqua MODIS Level 3 Mapped Chlorophyll Data, Version R2022.0 [dataset]. NASA Ocean Biology Distributed Active Archive Center. <https://doi.org/10.5067/AQUA/MODIS/L3M/CHL/2022>, 2022a.
- 630 [NASA Ocean Biology Processing Group. Aqua MODIS Level 3 Mapped Photosynthetically Available Radiation Data, Version R2022.0 \[dataset\]. NASA Ocean Biology Distributed Active Archive Center. https://doi.org/10.5067/AQUA/MODIS/L3M/PAR/2022](https://doi.org/10.5067/AQUA/MODIS/L3M/PAR/2022), 2022b.
- NASA Ocean Biology Processing Group. Aqua MODIS Level 3 Mapped Downwelling Diffuse Attenuation Coefficient Data, Version R2022.0 [dataset]. NASA Ocean Biology Distributed Active Archive Center. <https://doi.org/10.5067/AQUA/MODIS/L3M/KD/2022>, 2022c.
- 635 [NASA Ocean Biology Processing Group. Aqua MODIS Level 3 Mapped Remote-Sensing Reflectance Data, Version R2022.0 \[dataset\]. NASA Ocean Biology Distributed Active Archive Center. https://doi.org/10.5067/AQUA/MODIS/L3M/RRS/2022](https://doi.org/10.5067/AQUA/MODIS/L3M/RRS/2022), 2022d.
- NASA Ocean Biology Processing Group. Aqua MODIS Level 3 Mapped Inherent Optical Properties Data, Version R2022.0 [dataset]. NASA Ocean Biology Distributed Active Archive Center. <https://doi.org/10.5067/AQUA/MODIS/L3M/IOP/2022>, 2022e.
- ~~NASA Ocean Biology Processing Group. Aqua MODIS Level 3 Mapped Photosynthetically Available Radiation Data, Version R2022.0 [dataset]. NASA Ocean Biology Distributed Active Archive Center. <https://doi.org/10.5067/AQUA/MODIS/L3M/PAR/2022>, 2022.~~
- 645 ~~NASA Ocean Biology Processing Group. Aqua MODIS Level 3 Mapped Particulate Organic Carbon Data, Version R2022.0 [dataset]. NASA Ocean Biology Distributed Active Archive Center. <https://doi.org/10.5067/AQUA/MODIS/L3M/POC/2022>, 2022.~~
- [Ólafsdóttir, S. R., Benoit-Cattin, A., and Danielsen, M.: Dissolved inorganic carbon \(DIC\), total alkalinity, temperature, salinity, nutrients and dissolved oxygen collected from discrete samples and profile observations during the R/Vs Arni Fridriksson and Bjarni Saemundsson Irminger Sea \(FX9\) time series cruises in the North Atlantic Ocean in from 2014-02-11 to 2022-08-09 \(NCEI Accession 0209072\), NOAA National Centers for Environmental Information \[data set\]. https://doi.org/10.25921/vjmy-8h90](https://doi.org/10.25921/vjmy-8h90), 2020a.
- 650 [Ólafsdóttir, S. R., Benoit-Cattin, A., and Danielsen, M.: Dissolved inorganic carbon \(DIC\), total alkalinity, temperature, salinity, nutrients and dissolved oxygen collected from discrete samples and profile observations during the R/Vs Arni Fridriksson and Bjarni Saemundsson time series IcelandSea \(LN6\) cruises in the North Atlantic Ocean from 2014-02-18 to 2022-08-16 \(NCEI Accession 0209074\), NOAA National Centers for Environmental Information \[data set\]. https://doi.org/10.25921/qhed-3h84](https://doi.org/10.25921/qhed-3h84), 2020b.
- 655 [Ólafsson, J.: Partial pressure \(or fugacity\) of carbon dioxide, dissolved inorganic carbon, temperature, salinity and other variables collected from discrete samples, profile and time series profile observations during the R/Vs Arni Fridriksson and Bjarni Saemundsson time series Iceland-Sea \(LN6\) cruises in the North Atlantic Ocean from 1985-02-22 to 2013-11-26 \(NCEI Accession 0100063\). NOAA National Centers for Environmental Information \[data set\]. https://doi.org/10.3334/cdiac/otg.carina\\_icelandsea](https://doi.org/10.3334/cdiac/otg.carina_icelandsea), 2012.
- 660 [Ólafsson, J.: Partial pressure \(or fugacity\) of carbon dioxide, dissolved inorganic carbon, temperature, salinity and other variables collected from discrete sample and profile observations using CTD, bottle and other instruments from ARNI FRIDRIKSSON and BJARNI SAEMUNDSSON in the North Atlantic Ocean from 1983-03-05 to 2013-11-13 \(NCEI](https://doi.org/10.3334/cdiac/otg.carina_icelandsea)
- 665

- Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., ... & Yool, A. Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature*, 437, 681-686, 670 <https://doi.org/10.1038/nature04095>, 2005.
- Qi, D., Ouyang, Z., Chen, L., Wu, Y., Lei, R., Chen, B., ... & Cai, W. J. Climate change drives rapid decadal acidification in the Arctic Ocean from 1994 to 2020. *Science*, 377, 1544-1550, <https://doi.org/10.1126/science.abo0383>, 2022.
- Sabine, C. L., & Tanhua, T. Estimation of anthropogenic CO<sub>2</sub> inventories in the ocean. *Annual Review of Marine Science*, 2, 175-198, <https://doi.org/10.1146/annurev-marine-120308-080947>, 2010.
- 675 Takahashi, T., Sutherland, S. C., Chipman, D. W., Goddard, J. G., Ho, C., Newberger, T., ... & Munro, D. R. Climatological distributions of pH, pCO<sub>2</sub>, total CO<sub>2</sub>, alkalinity, and CaCO<sub>3</sub> saturation in the global surface ocean, and temporal changes at selected locations. *Marine Chemistry*, 164, 95-125, <https://doi.org/10.1016/j.marchem.2014.06.004>, 2014.
- Terhaar, J., Kwiatkowski, L., & Bopp, L. Emergent constraint on Arctic Ocean acidification in the twenty-first century. *Nature*, 582, 379-383, <https://doi.org/10.1038/s41586-020-2360-3>, 2020.
- 680 Wolter, K., & Timlin, M. S. El Niño/Southern Oscillation behaviour since 1871 as diagnosed in an extended multivariate ENSO index (MEI. ext). *International Journal of Climatology*, 31, 1074-1087, <https://doi.org/10.1002/joc.2336>, 2011.
- Zhong, G., Li, X., Song, J., Qu, B., Wang, F., Wang, Y., ... & Duan, L. Reconstruction of global surface ocean pCO<sub>2</sub> using region-specific predictors based on a stepwise FFNN regression algorithm. *Biogeosciences*, 19, 845-859, <https://doi.org/10.5194/bg-19-845-2022>, 2022.
- 685 Zhong, G; Li, X., Song, J. Global ocean gridded seawater pH during 1992-2020 at 0-2000 m depth based on Stepwise FFNN algorithm 2023 version. Marine Science Data Center of the Chinese Academy of Sciences (<http://dx.doi.org/10.12157/IOCAS.20230720.001>), 2023