

# Coral Skeletal Proxy Records Database for the Great Barrier Reef, Australia

Ariella K. Arzey<sup>1,2</sup>, Helen V. McGregor<sup>1,2</sup>, Tara R. Clark<sup>1,3</sup>, Jody M. Webster<sup>4</sup>, Stephen E. Lewis<sup>5</sup>, Jennie Mallela<sup>6</sup>, Nicholas P. McKay<sup>7</sup>, Hugo W. Fahey<sup>1</sup>, Supriyo Chakraborty<sup>8</sup>, Tries B. Razak<sup>9, 10</sup> and Matt J. Fischer<sup>11</sup>

<sup>1</sup>Environmental Futures, School of Earth, Atmospheric and Life Sciences, University of Wollongong, Wollongong, 2522, Australia

<sup>2</sup>Securing Antarctica's Environmental Future, University of Wollongong, Wollongong, 2522, Australia

<sup>3</sup>Radiogenic Isotope Facility, School of the Environment, The University of Queensland, Brisbane, 4072, Australia

10 <sup>4</sup>Geocoastal Research Group, School of Geosciences, The University of Sydney, Camperdown, 2006, Australia

<sup>5</sup>Catchment to Reef Research Group, Centre for Tropical Water and Aquatic Ecosystem Research, James Cook University, Townsville, 4811, Australia

<sup>6</sup>Research School of Biology, The Australian National University, Canberra, 2601, Australia

<sup>7</sup>School of Earth and Sustainability, Northern Arizona University, Flagstaff, 86011, USA

15 <sup>8</sup>Indian Institute of Tropical Meteorology, Ministry of Earth Sciences (MoES), Pune, 411008, India

<sup>9</sup>Department of Marine Science and Technology, Faculty of Fisheries and Marine Sciences, IPB University, Bogor, 16680, Indonesia

<sup>10</sup>School of Coral Reef Restoration (SCORES), Faculty of Fisheries and Marine Science, IPB University, Bogor, 16680, Indonesia

20 <sup>11</sup>Environment Research and Technology Group, Australian Nuclear Science and Technology Organisation, Lucas Heights, 2234, Australia

*Correspondence to:* Ariella K. Arzey (aka548@uowmail.edu.au)

**Abstract.** The Great Barrier Reef (GBR), Australia has a long history of palaeoenvironmental coral research. However, it can be logistically difficult to find the relevant research and records, which are often unpublished or exist as 'grey literature'. This hinders researchers' ability to efficiently assess the current state of coral core studies on the GBR and thus identify any key knowledge gaps. This study presents the Great Barrier Reef Coral Skeletal Records Database (GBRCD), which compiles 208 records from coral skeletal research conducted since the early 1990s. The database includes records from the Holocene, from ~8,000 years ago, to the present day; from the northern, central, and southern GBR from inshore and offshore locations. Massive *Porites* spp. coral records comprise the majority (92.5 %) of the database, and the remaining records are from *Acropora*, *Isopora* or *Cyphastrea* spp. The database includes 78 variables, with Sr/Ca, U/Ca and Ba/Ca the most frequently measured. Most records measure data over 10 or more years and are at monthly or lower resolution. The GBRCD is machine readable and easily searchable so users can find records relevant to their research, for example, by filtering for site names, time period, or coral type. It is publicly available as comma-separated values (CSV) data and metadata files with entries linked by the unique record ID and as Linked Paleo Data (LiPD) files. The GBRCD is publicly available from the NOAA National Center for Environmental Information World Data Service for Paleoclimatology at <https://doi.org/10.25921/hqzk-8h74> (Arzey et al., 2024). The intention is to update the GBRCD annually, depending on the availability of relevant new GBR records or submission of legacy records to the GBRCD for archiving. The collection and curation of existing GBR coral research provides researchers with the ability to analyse common proxies such as Sr/Ca across multiple locations and/or examine regional to reef scale trends. The database is also suitable for multi-proxy comparisons and combination or composite analyses to determine overarching changes recorded by the proxies. This database represents the first comprehensive compilation of coral records from the GBR. It enables the investigation of multiple environmental factors via various proxy systems for the GBR, northeastern Australia and potentially the broader Indo-Pacific.

## 1 Introduction

Scleractinian coral skeletons have long been used to reconstruct past changes of reef environments and climates, and to understand reef responses to those changes (e.g. Druffel and Griffin, 1993; Webster et al., 2018; Thompson, 2022; Clark et al., 2017; Felis et al., 2014). Foundational to the field are studies on corals from the World Heritage listed Great Barrier Reef (GBR). Studies on GBR corals have established, for example, the relationships between coral skeletal Sr/Ca and  $\delta^{18}\text{O}$  with sea surface temperature (SST) (Weber and Woodhead, 1969; Weber and Woodhead, 1970; Weber and Woodhead, 1972; Weber, 1973; Mcculloch et al., 1994), coral luminescence with riverine inputs (Isdale, 1984; Boto and Isdale, 1985; Isdale et al., 1998), coral radiocarbon with oceanographic processes (Druffel and Griffin, 1993), the environmental signature recorded in coral density bands (Lough and Barnes, 1990; Lough and Barnes, 1997), evidence of sediment and nutrient exposure (Mcculloch et al., 2003; Wyndham et al., 2004; Sammarco et al., 1999) and a host of new analytical methods and approaches (Gagan et al., 1998; Sinclair et al., 1998; Barnes et al., 2003). In addition, palaeoclimate and palaeoenvironmental research based on GBR corals (typically massive *Porites* spp.) have advanced our knowledge on the most pressing threats to the GBR and coral reef ecosystems more broadly (Mcculloch et al., 2003; De'ath et al., 2009; Lough et al., 2015; Wei et al., 2009; Koop et al., 2001), tropical climate processes (Hendy et al., 2002; Lough, 2007; Druffel and Griffin, 1993), reef responses to climate and sea-level change (Webster et al., 2018; Sanborn et al., 2020; Yokoyama et al., 2011; Leonard et al., 2020), and coral bleaching (Suzuki et al., 2003; D'olivo and Mcculloch, 2017; De'ath et al., 2012).

Despite the long history of coral palaeoclimate and environmental research on the GBR, like other research fields, GBR coral research data are being lost to time (Vines et al., 2014). While there is a recent push to ensure that coral geochemical records are publicly available upon publication (Kaufman and Team, 2018; Khider et al., 2019; Dassié et al., 2017), many GBR coral core records are not yet publicly archived, and thus we risk losing these valuable datasets. The need to publicly archive GBR coral geochemical data is even more pressing because corals are under threat from human-induced stresses (Hughes et al.,

65 2017; Maynard et al., 2015; Guan et al., 2020; Ortiz et al., 2018). Long term monitoring of the GBR indicates reef-wide hard coral cover was down to 18 % in 2017 (Australian Institute of Marine Science, 2017). Since then hard coral cover has increased, but recovery has primarily been driven by fast growing *Acropora* species (Australian Institute of Marine Science, 2017, 2022), which are less suitable for producing continuous multi-century records. The future of GBR corals, and therefore the valuable geochemical records they contain, faces an uncertain future, which makes existing GBR records a precious and potentially  
70 finite resource.

This study presents a comprehensive database of GBR coral geochemical and luminescence records, and associated variables such as growth characteristics (where available), compiling both published records (201), and records published here for the first time (7 records). The GBR Coral Skeletal Records Database (GBRCD) transfers data into a standardised machine-readable format (structured using a consistent logic readable by and between machines) and makes the records available to researchers  
75 in a searchable database. The database consists of two main parts: 1) individual comma-separated values (CSV) files consisting of a single coral's geochemical and luminescence measurements and the associated age estimate, and 2) a metadata file for the entire suite of corals, which includes details of location, genus or species, analytical techniques, publications, cross-references, and other details. Alternatively, the GBRCD is available as Linked Paleo Data (LiPD) files and serialisations in Python, MATLAB and R, which includes the same fields standardised to the LiPD format and terminology. Basic analysis of the scope  
80 of the database is presented, as is a discussion of database caveats and future use. The GBRCD addresses both the potential loss of existing GBR coral data and provides the ability to easily explore what records exist. The GBRCD is a valuable resource for exploratory analysis to understand the GBR (and the wider Pacific) environment (e.g. Henley et al., 2024) and preserves the records for future use.


## 85 1.1 GBR Setting

The GBR is made up of more than 3,000 reefs that span 14 degrees of latitude, and the World Heritage Area comprises an area of 348,000 km<sup>2</sup>. The width of the GBR across the continental shelf changes with latitude, ranging from 50 to 250 km wide (Steinberg, 2007; Hopley et al., 2007). Thus, coral reefs in the GBR can be classified based on their proximity to the Queensland coast and sea floor depth, as either inner-shelf, mid-shelf, or outer-shelf reefs and, depending on their  
90 classification, these reefs will experience various gradients of terrigenous or open ocean influences. There is a seasonal atmospheric circulation that is characterised by an annual summer monsoon (December to March), which brings rainfall and increased river discharge to neighbouring GBR marine waters.

Average sea surface temperature (SST) varies latitudinally, with a 1.65°C difference between the northern region (11-17° S) and the southern region (20-24° S)(Fig. 1) for the 1981-2010 period (HadISST; Rayner et al., 2003). The timing of the annual  
95 minimum and maximum also varies slightly; as mean maximum SST in the 1981-2010 period (HadISST1.1) most frequently occurs in February in the central and southern GBR, and in January in the northern GBR, while minimum SSTs most frequently occur in August in the northern and southern GBR and in July in the central GBR.

The modern GBR coral reef ecosystem initiated ~8-9 thousand years ago (ka) following the Holocene marine transgression (Davies and Hopley, 1983; Davies et al., 1985; Dechnik et al., 2015; Hopley et al., 2007), although the timing of individual  
100 coral reef initiation and vertical accretion rates varied considerably (range of 1-16 m/1000 years) (Dechnik et al., 2017; Sanborn et al., 2020; Davies and Hopley, 1983; Hopley et al., 2007; Leonard et al., 2020; Ryan et al., 2018). Decreases in reef growth rate occurs where the reef surface is within ~2-3 m of sea level due to reduced growth and/or erosional loss (Davies and Hopley, 1983). Evidence from reef drilling suggests that there was little latitudinal variation in reef growth rate throughout the GBR, although the depth to the Pleistocene reef foundations in northern and southern GBR regions is shallower than the  
105 central region (e.g. Davies et al., 1985; Hopley et al., 2007). Additionally, there are reef-flat growth hiatuses in both the northern and southern GBR, but not in the central GBR, due to a relative fall in sea level of ~0.5 m in the other regions that

was not evident in the central GBR due to subsidence in the central region (Halifax Basin) from hydro-isostatic adjustment (Dechnik et al., 2017).



110 **Figure 1.** Map of locations with coral records in the Great Barrier Reef Coral Skeletal Records Database. Record locations are  
indicated by circles, select cities are indicated by orange squares, select rivers are indicated by dark blue lines, reefs are in light blue  
and the nominal regions of the north, central and south GBR are indicated by the dashed pink lines. Numbers identify rivers; 1 –  
115 Endeavour River, 2 – Daintree River, 3 – Herbert River, 4 – Burdekin River, 5 – Pioneer River and 6 – Fitzroy River. The drainage  
basin boundary associated with the included rivers is indicated by the dashed black lines. GIS layers © State of Queensland  
(Department of Resources) 2023. This material is licensed under a [Creative Commons - Attribution 4.0 International licence](https://creativecommons.org/licenses/by/4.0/). See  
Appendix G for full list of included GIS layers.

## 1.2 Palaeoclimate and palaeoenvironment reconstructed from GBR coral skeletons

Research on the GBR has a long history of discovery and scientific development, with ventures as far back as the 1928–1929  
GBR expedition lauded as having promoted the development of empirical and analytical approaches important to the  
120 foundation of modern coral reef science (Spencer et al., 2021). Coral skeletons provide in situ records of the reef environment  
as they incorporate trace elements, a variety of element isotopes, and organic materials in proportion to climate and  
environmental variations in the marine environment. These palaeoclimate and palaeoenvironment ‘proxies’ are a long-  
established method to assess the reef environment in the past and present, and provide a means to quantify relative changes

over space and time. Proxy records from coral skeletons offer the ability to extract high-resolution (weekly–monthly) information to assess a range of climate and environmental variables, most commonly SST and rainfall, and at variety of scales, for example near-weekly records that extend from the present day back several centuries prior to the instrumental record (e.g. Lewis et al., 2012; D'olivo and Mcculloch, 2022).

Beginning in the late 1960s, studies using GBR corals provided empirical evidence of a relationship between skeletal oxygen isotopes, luminescence, trace elements, radiocarbon and density and the coral's marine environment (Weber and Woodhead, 1969; Weber and Woodhead, 1970; Weber and Woodhead, 1972; Isdale, 1984; Druffel and Griffin, 1993; Lough and Barnes, 1990; Mcculloch et al., 2003) and were at the forefront of the development of new technology and methods to measure these coral properties (Isdale et al., 1998; Gagan et al., 1998; Sinclair et al., 1998; Barnes et al., 2003; D'olivo et al., 2018).

After the description of the utility of Sr/Ca as an SST proxy by Beck et al. (1992), coral skeleton proxy research in the GBR focused on development and calibration of proxy-SST relationships in the 1990s. GBR coral records from this period are generally short (<10 years) and were analysed using a variety of spectrometry methods (Alibert and Mcculloch, 1997; Druffel and Griffin, 1993, 1999; Mcculloch et al., 1994; Gagan et al., 1998; Sinclair et al., 1998). Research in the 1990s also included the first application of paired Sr/Ca and  $\delta^{18}\text{O}$  proxies to reconstruct the GBR palaeoenvironment (Mcculloch et al., 1994; Gagan et al., 1998) and the further exploration of coral luminescence as a proxy for river discharge (Isdale et al., 1998). These approaches were applied to extend our knowledge of GBR SST, riverine input and oceanographic variability prior to the 1800s (Lough, 2007, 2011b; Hendy et al., 2002; Hendy et al., 2003a; Calvo et al., 2007), and during the Holocene (Gagan et al., 1998; Leonard et al., 2016; Sadler et al., 2016b; Lough et al., 2014; Roche et al., 2014).

Since the 2000s, partially facilitated by the development of the Australian Institute of Marine Science luminometer and the Australian National University laser ablation methods, research on coral proxies in the GBR has focused on proxies for terrigenous inputs. These include indicators of terrestrial runoff and sediment exposure (e.g. Ba/Ca and rare earth elements (REEs) (Jupiter et al., 2008; Jupiter, 2008; Saha et al., 2021; Mcculloch et al., 2003; Lewis et al., 2012; Leonard et al., 2019)), land-use changes (e.g.  $\delta^{15}\text{N}$  and Mn (Lewis et al., 2007; Lewis et al., 2012; Erler et al., 2016; Erler et al., 2020; Marion et al., 2021)) and river flow patterns (e.g. Ba/Ca and luminescence (Lough et al., 2002; Lough, 2007; Lough et al., 2014; D'olivo and Mcculloch, 2022)). Additionally, there is continuing research that aims to explore new proxies (e.g. V/Ca (Saha et al., 2019a),  $\delta^{98}\text{Mo}$  (Wang et al., 2019)) as well as to develop proxies relevant to global issues such as ocean acidification (e.g.  $\delta^{11}\text{B}$  (Pelejero et al., 2005; Wei et al., 2009; D'olivo et al., 2015; Mcculloch et al., 2017)).

## 2 Overview of the GBR Coral Skeletal Records Database

### 2.1 Selection criteria for inclusion in GBRCDD

The GBRCDD was not assembled around a specific research goal unlike other coral palaeoclimate databases, such as the CoralHydro2k database (Walter et al., 2023), which was designed to support “the project's goal of reconstructing tropical hydroclimatic variability at seasonal and longer timescales”. The GBRCDD compiles and makes available coral geochemical and luminescence data from the GBR in a user-friendly format to facilitate their use in research. The GBRCDD is similar to the Coral Trait Database (Madin et al., 2016), but focuses on the GBR and coral proxy measurements (i.e. geochemical and luminescence measurements). Therefore, only three broad selection criteria are considered for including a record in the GBRCDD:

- Location. The GBRCDD compiles coral records from the GBR and, due to its regional context and latitudinal relevance, Flinders Reef.
- Length. The records are continuous (multi-year) measurements from coral skeletons.
- Age model. Each record includes a relevant chronology (i.e. an estimate of age as the independent variable) or has a chronology that could be recreated relatively easily based on available data.

165 All available geochemical or luminescence data were included in the GBRC D if they met these three criteria. No screening was applied on the quality of the data and no records were excluded based on analytical uncertainties, diagenesis screening, resolution (e.g. weekly, monthly, annual), significance of the correlation between proxy and target parameter or record length. It is at the discretion of researchers using data from the GBRC D to determine suitability for their aims.

Where available, additional variables, such as coral growth data (density, extension, etc.), are included alongside the  
170 geochemical and luminescence data for records in the GBRC D where the additional variable data are at the same resolution as the geochemical or luminescence measurements.

## 2.2 Data sources

The majority of GBR coral records are from peer-reviewed published studies (see Appendix Table A1). In many cases, the published records were publicly archived in repositories such as the NOAA National Centers for Environmental Information  
175 World Data Service for Paleoclimatology (NOAA/WDS Paleoclimatology) and World Data Center PANGAEA. The GBRC D includes digital object identifier (DOI) information linking back to these original sources and publications. In some cases, published studies included additional records that were not archived in public repositories. Instead, these data were obtained from the publication's supplemental material, or by reaching out to the corresponding authors. As part of the GBRC D compilation process, 114 previously unarchived records, as well as the seven previously unpublished records, were submitted  
180 to NOAA/WDS Paleoclimatology for archiving. These files are in the standard NOAA/WDS Paleoclimatology (tab-delimited) file format.

Seven records have not been published previously and are published in the GBRC D for the first time (Table 1 & Appendix A & C). These records are accompanied by their associated metadata record and appropriate background information (see Appendix A, C, D). If further information about the record is required, users are encouraged to reach out to the first  
185 author/contact listed in the GBRC D publication metadata.

**Table 1. Records first published in the GBRC D.**

<b>Record ID</b>	<b>Variables</b>	<b>Reef Location</b>
AR24DIP01	Ba/Ca; Mg/Ca; Sr/Ca; U/Ca; Y/Ca	Dip Reef
AR24OTI01	Sr/Ca	One Tree Island
AR24OTI02	Ba/Ca; B/Ca; Ca/Ca; Mg/Ca; Sr/Ca; U/Ca	One Tree Island
AR24OTI03	Ba/Ca; B/Ca; Ca/Ca; Mg/Ca; Sr/Ca; U/Ca	One Tree Island
AR24OTI04	Ba/Ca; B/Ca; Ca/Ca; Mg/Ca; Sr/Ca; U/Ca	One Tree Island
AR24OTI05	Ba/Ca; B/Ca; Ca/Ca; Mg/Ca; Sr/Ca; U/Ca	One Tree Island
AR24SLY01	$\delta^{13}\text{C}$ ; $\Delta^{14}\text{C}$ ; $\delta^{18}\text{O}$	Stanley Reef

Records included in the database are multi-year measurements of coral geochemistry and luminescence signals. In total, there  
190 are 78 variables available to investigate the GBR environment, or to determine new proxy-environment relationships (Table 2). The records include commonly measured isotopes and trace elements (TE) such as  $\delta^{18}\text{O}$ , Ba/Ca, Sr/Ca, and U/Ca, as well as less explored isotopes (such as  $\delta^{11}\text{B}$  and  $\delta^{15}\text{N}$ ) and emerging TE proxies such as Li/Mg and rare earth elements (REEs). Luminescence records are included in the database as coral luminescence is a proven hydrological proxy in the GBR (Lough 2011, Lough et al. 2014).

Care is advised for the use of all records in the GBRC D as the relevance of the measured variables may vary by location, or  
195 the sampling resolution of the records may not be suitable. Therefore, further processing needs be considered before integrating or comparing the records.

## 2.3 Database structure

The database is provided in a split form and includes a table of metadata describing each record, as well as individual CSV files that include the coral variables such as age, distance down core, measured trace elements and so on. It is also provided as LiPD serialisations with the metadata fields structured as per the LiPD format (see Table B1). The database is available in both formats to facilitate ease of use. The CSV files are easily accessible and do not need programming languages (e.g. R or python) to explore the datasets, whereas the LiPD files require programming languages to fully access and is designed as structured data to enable interoperability and quick analysis of palaeoclimate data (Mckay and Emile-Geay, 2016).

The metadata supplied alongside the coral record data provides a thorough background on each record, with assumptions and limitations made clear for researchers. While every best attempt has been made to provide information for all metadata fields that are relevant, there are some fields where the information was not available (i.e. flagged with NA). Metadata fields allow for filtering by, for example: site, coral genus or species, analysis method, proxy type, etc.

Individual CSV and LiPD files are provided for each coral record. In general, an individual record is defined as being from a single coral, whether a coral head was drilled directly, or the coral was retrieved from a core drilled into coral reef matrix (coral reef matrices can intersect individual coral heads as well as the reef framework in which they are preserved).

In some instances, multiple geochemical and/or luminescence records have been measured from the same coral head. In this situation, the multiple records have their own separate entries as they are usually either part of an included composite record, measured using different techniques, or have sufficient differences in the metadata or dataset to provide a good reason not to attempt to join the datasets. However, they are identified and cross-referenced in the database as being from the same coral or core.

## 2.4 Data processing

Published data, as was available or supplied, have been archived in the database format described here. Potentially abnormal data (e.g. values  $<0$  for Mg/Ca [after correcting against a method and/or lab standard]) were not removed from the datasets as they may be indicative of climate or environmental events, and it is up to the user to determine appropriate screening procedures when using the database.

A basic quality check was conducted on records published for the first time in the GBRCd (see Appendix C & F). This consisted of creating or adjusting the coral age model based on available information and checking for diagenetic alteration where there was a physical sample available.

Coral records that were archived or supplied with the ages in standard date format (e.g. DD/MM/YYYY) had all ages converted to decimal date in Common Era (CE)/Before Common Era (BCE) format. This means that there is no year zero, and negative (BCE) numbers start at “-1”. Since there are rounding issues with decimal dates assigned to the first day of the month (at midnight), 12 hours were added to each date to ensure the age remained assigned to the correct month of the year.

For archiving purposes, 16 records were assigned a new age model. This was primarily because the coral data were not available with an age model, or did not have one appropriate for archiving (e.g. date not assigned to each data point). A note is included in the relevant field about the chronology for records where the chronology was (re)created for archiving in the GBRCd and further details are provided in Appendix C. Nine fossil records were supplied or previously archived with chronologies relative to the record itself (i.e. 0 = first or youngest year of record). The available dating information was used to transform these chronologies to Common Era ages.

Before Common Era fossil coral data are included in the GBRCd. Absolute age estimates for these corals are based on uranium-thorium (U-Th, or U-series) dating and radiocarbon ( $^{14}\text{C}$ ) measurements, with basic information about the dating supplied in the GBRCd metadata. Five coral records dated by radiocarbon measurements were re-calibrated using CALIB rev. 8 (Stuiver and Reimer, 1993) and the Marine20 radiocarbon age calibration curve (Heaton et al., 2020) to standardise the age

estimates and details (including the local delta-R reservoir correction) are available in the dating\_notes field. Users of the GBRCDB should determine if the radiocarbon ages of the fossil records should be revised based on future updates to the calibration curves and/or reservoir ages. For records where U-Th dating information is reported for the first time in this study, information is provided in the GBRCDB metadata and complete relevant U-Th dating information (Dutton et al., 2017) is included in the Appendix (Table D1).

Relevant metadata was obtained from source publications, and where necessary subsequent or cited publications, and data archives. Coral IDs were also matched across publications using literature searches, the AIMS reef core database, and within the GBRCDB.

## 2.5 Database Records and Identification

Records are identified here as datasets from a single coral that may include one or more measured variables unified with a single age model. Similar to the Iso2k database protocols (Konecky et al., 2020), each record in the database was given a nine-digit alphanumeric ID. The dataset ID was created from the first two letters of the first author's last name, the last two numbers of the year of publication, the first three letters of the location (or in cases where appropriate, a commonly accepted shortening or three-letter acronym for the location was used), and two numbers that represent the core number. If data were not previously published, publication year was substituted with the year of this compilation. For reef locations in the GBR that are currently represented by a number rather than a name, and for records that are the composite of multiple locations, 'GBR' has been used as the three-letter reef location for core naming purposes.

However, the Iso2k nine-digit alphanumeric identifier was insufficient to separate records where researchers have published multiple papers from the same GBR location within the same year. To separate these records a 10<sup>th</sup> digit was added to the ID, adding an alphabetical end code of 'a', 'b' and so on following the order of publication (Table 2), similar to many journal standards for uniquely identifying references. Additionally, where the core record has been split due to notable differences in the record (for example due to data resolution or age model), a split identifier of '\_1' or '\_2' has been appended to the end of the identifier (Table 2).



**Table 2. Example of process of giving core records a unique identifier.**

GBRCD ID	Core ID from publication	Publication Reference	Location	Variables	Resolution	Split	Comments
SA16HER01a	AR HL 3	Sadler et al. (2016a)	Heron Reef	Sr/Ca	Biannual	Core 1	Suffix identifier for record as different coral(s) with publication(s) in the same year at the same reef location.
SA16HER01b	P HS 1	Sadler et al. (2016b)	Heron Reef	Sr/Ca	Bimonthly	Core 1	Suffix identifier for record as different coral(s) with publication(s) in the same year at the same reef location.
WE09ARL01_1	AREO 4	Wei et al. (2009)	Arlington Reef	Ba/Ca; $\delta^{11}\text{B}$ ; $\delta^{13}\text{C}$ ; $\delta^{18}\text{O}$ ; Mg/Ca; Sr/Ca; $\delta^{11}\text{B}$ pH	Annual	Core 1, split 1	Record split due to difference in measurement resolution.
WE09ARL01_2	AREO 4	Wei et al. (2009)	Arlington Reef	Ba/Ca; $\delta^{11}\text{B}$ ; $\delta^{13}\text{C}$ ; $\delta^{18}\text{O}$ ; Mg/Ca; Sr/Ca; $\delta^{11}\text{B}$ pH	>Annual	Core 1, split 2	Record split due to difference in measurement resolution.

## 2.6 Database Variables, Units and Standards

265 Records are all standardly formatted as CSV and LiPD files, with each record containing an age field and variable field(s) as per Table 3. Each GBRCD record includes between 1 and 33 variables.

The database collates 78 variables obtained from coral records: 72 measured and 6 calculated variables (Table 4).

Units reported in the database were standardised to enable interoperability among records. All trace element/calcium (TE/Ca) ratios were included as mmol/mol and, where relevant, the calibrations (proxy-SST) for the ratios included in the record metadata are in the same units. Isotopes were archived as permille (‰), and, where relevant, the calibrations for the ratios are in the same units. The majority of non-ratio elements were archived as parts per million. The exceptions to this were the rare earth elements (REEs) (lanthanum (La), cerium (Ce), praseodymium (Pr), neodymium (Nd), samarium (Sm), europium (Eu), gadolinium (Gd), terbium (Tb), dysprosium (Dy), holmium (Ho), erbium (Er), thulium (Tm), ytterbium (Yb), and lutetium (Lu)), yttrium (Y) and zirconium (Zr), which were archived as parts per billion (i.e.  $10^9$ ).

275 The list of included dataset field names for each record is provided in the metadata file (i.e. `meths_primaryVariableList` or `meths_additionalVariableList`; Table 4).

All records include an ‘Age’ field that is the age of each sample as determined by the coral age model. For records that are at greater than annual resolution the age supplied is representative of the grouping of years included in each data point. For example, the Druffel and Griffin (1999) biennial data lists the middle year of the sampled coral (i.e. October 1889 to October 1891 has the age 1890.8, while Hendy et al. (2002) quinquennial resolution data lists the middle year of the sampled coral (i.e. 1981 to 1985 has the age 1983). For sub-annual resolution records the tie-point within each month varies depending upon the

researchers' choice in the original publications. It is up to researchers to inspect the data to understand the 'Age' field for each record they intend to use.

285 **Table 3. Database record format. Describes the format of the CSV files of each record.**

Field name	Variable	Type	Description
Age	Age	numeric	Time data for the record. All time data are expressed as a decimal date. Annual or greater resolution default to [year] (equivalent of [year].0) unless original publication specified otherwise. Units: CE
[variable]	Data	numeric	A Nx1 vector of proxy data. Data type is specified in the list of proxy and additional variables listed in Table 4. Units are specified in Table 4.

**Table 4. Data variables. Describes variables included in the database, units, type, and brief description. The list of variables is alphabetical.**

Field name	Variable	Type	Units	Unit abbreviation	Description
Arag	$\Omega_{arag}$	calculated	dimensionless		Aragonite saturation state determined from calculated $\delta^{11}\text{B}$ pH.
Ba	Ba	measured	parts per million	ppm	Barium
Ba138Ca	Ba/Ca	measured	millimole/mole	mmol/mol	Barium/Calcium A secondary Ba/Ca measurement for laser ablation. This is a measure of the total Ba/Ca using the count intensities of the $\text{Ba}^{138}$ isotope.
BaCa	Ba/Ca	measured	millimole/mole	mmol/mol	Barium/Calcium For laser ablation this is a measure of the total Ba/Ca using the count intensities of the $\text{Ba}^{137}$ isotope.
BCa	B/Ca	measured	millimole/mole	mmol/mol	Boron/Calcium
BMg	B/Mg	measured	millimole/mole	mmol/mol	Boron/Magnesium
Ca	Ca	measured	parts per million	ppm	Calcium
CaCa	Ca/Ca	measured	millimole/mole	mmol/mol	A measure of the total Ca/Ca ratio using the count intensities of the Calcium <sup>48</sup> /Calcium <sup>43</sup> isotopes.
Calcn	Calcification rate	calculated	grams/centimetre/year	g/cm/yr	Annual coral calcification rate.
Cd	Cd	measured	parts per million	ppm	Cadmium
CdCa	Cd/Ca	measured	millimole/mole	mmol/mol	Cadmium/Calcium
Ce	Ce	measured	parts per million	ppm	Cerium (REE)
CeCa	Ce/Ca	measured	millimole/mole	mmol/mol	Cerium/Calcium

Field name	Variable	Type	Units	Unit abbreviation	Description
CeCe_anom	Ce/Ce*	calculated	dimensionless <sup>a</sup>		Cerium anomaly calculated <sup>b</sup> as: $\left(\frac{Ce}{Ce}\right) = \left(\frac{Ce_{sn}}{Pr_{sn} \times \left(\frac{Pr}{Nd}\right)_{sn}}\right) (1)$
d11B	$\delta^{11}\text{B}$	measured	permille	permil (‰)	$^{11}\text{B}/^{10}\text{B}$
d11B_pH	$\delta^{11}\text{B}$ pH	calculated	dimensionless		pH calculated from $\delta^{11}\text{B}$ .
d11B_pHcf_p res	$\delta^{11}\text{B}$ pH <sub>cf</sub>	calculated	dimensionless		pH of calcifying fluid calculated from $\delta^{11}\text{B}$ . Values are pressure corrected.
d13C	$\delta^{13}\text{C}$	measured	permille	permil (‰)	$^{13}\text{C}/^{12}\text{C}$
D14C	$\Delta^{14}\text{C}$	measured	permille	permil (‰)	$\delta^{14}\text{C}$ ( $^{14}\text{C}/^{12}\text{C}$ ) deviation from a standard and corrected for $\delta^{13}\text{C}$ .
d15N	CS- $\delta^{15}\text{N}$	measured	permille	permil (‰)	$^{15}\text{N}/^{14}\text{N}$  $\delta^{15}\text{N}$ of skeleton bound organic matter.
d18O	$\delta^{18}\text{O}$	measured	permille	permil (‰)	$^{18}\text{O}/^{16}\text{O}$
d18Osw	$\delta^{18}\text{O}_{\text{seawater}}$	calculated	permille	permil (‰)	$\delta^{18}\text{O}$ of seawater calculated from paired $\delta^{18}\text{O}$ and Sr/Ca.
d66Zn	$\delta^{66}\text{Zn}$	measured	permille	permil (‰)	$^{66}\text{Zn}/^{64}\text{Zn}$
d98Mo	$\delta^{98}\text{Mo}$	measured	permille	permil (‰)	$^{98}\text{Mo}/^{95}\text{Mo}$
Density	Density	measured	grams/centimetre	g/cm	Annual average coral density.
Distance	Distance	measured	millimetres	mm	Distance measured from outer edge of coral or core.
Dy	Dy	measured	parts per billion	ppb	Dysprosium (REE)
Er	Er	measured	parts per billion	ppb	Erbium (REE)
Eu	Eu	measured	parts per billion	ppb	Europium (REE)
EuCa	Eu/Ca	measured	millimole/mole	mmol/mol	Europium/Calcium
Extn	Extension rate	measured	millimetres/year	mm/yr	Annual linear extension rate of coral.
Fe	Fe	measured	parts per million	ppm	Iron
FeCa	Fe/Ca	measured	millimole/mole	mmol/mol	Iron/Calcium
Gd	Gd	measured	parts per billion	ppb	Gadolinium (REE)
GdCa	Gd/Ca	measured	millimole/mole	mmol/mol	Gadolinium/Calcium
Ho	Ho	measured	parts per billion	ppb	Holmium (REE)
La	La	measured	parts per billion	ppb	Lanthanum (REE)
LaCa	La/Ca	measured	millimole/mole	mmol/mol	Lanthanum/Calcium
LiCa	Li/Ca	measured	millimole/mole	mmol/mol	Lithium/Calcium
LiMg	Li/Mg	measured	millimole/mole	mmol/mol	Lithium/Magnesium
Lu	Lu	measured	parts per billion	ppb	Lutetium (REE)
LuCa	Lu/Ca	measured	millimole/mole	mmol/mol	Lutetium/Calcium

Field name	Variable	Type	Units	Unit abbreviation	Description
Lumin	Luminescence	measured	dimensionless		Coral luminescence as measured by luminometer.
LuminGB	Luminescence G/B	measured	dimensionless		Luminescence calculated from spectral green/blue ratio measured by spectral luminescence scanning.
LuminInd	Luminescence visual indices	measured	dimensionless		Annual coral luminescence indices as determined by visual assessment of annual luminescence lines.
LuminRange	Luminescence range	measured	dimensionless		Annual luminescence range as measured by luminometer; determined by difference between annual (summer) maximum luminescence and preceding annual (winter) minimum luminescence.
LuminSF	Luminescence sum	measured	dimensionless		Standardised annual fluorescence/luminescence calculated by interpolating and summing 12 values between annual minimum (winter) luminescence values.
MgCa	Mg/Ca	measured	millimole/mole	mmol/mol	Magnesium/Calcium
Mn	Mn	measured	parts per million	ppm	Manganese
MnCa	Mn/Ca	measured	millimole/mole	mmol/mol	Manganese/Calcium
Mo	Mo	measured	parts per million	ppm	Molybdenum
Nd	Nd	measured	parts per billion	ppb	Neodymium (REE)
NdYb	Nd/Yb	measured	dimensionless <sup>a</sup>		Neodymium/Ytterbium (REE)
P	P	measured	parts per million	ppm	Phosphorus
PbCa	Pb/Ca	measured	millimole/mole	mmol/mol	Lead/Calcium
PCa	P/Ca	measured	millimole/mole	mmol/mol	Phosphorus/Calcium
Pr	Pr	measured	parts per billion	ppb	Praseodymium (REE)
PrCa	Pr/Ca	measured	millimole/mole	mmol/mol	Praseodymium/Calcium
REEsCa	$\Sigma$ REE/Ca	measured	dimensionless <sup>a</sup>		Sum of REEs/Calcium
Sm	Sm	measured	parts per billion	ppb	Samarium (REE)
SmCa	Sm/Ca	measured	millimole/mole	mmol/mol	Samarium/Calcium
SrCa	Sr/Ca	measured	millimole/mole	mmol/mol	Strontium/Calcium
Tb	Tb	measured	parts per billion	ppb	Terbium (REE)
Th	Th	measured	parts per million	ppb	<sup>232</sup> Thorium
Ti	Ti	measured	parts per million	ppm	Titanium
Tm	Tm	measured	parts per billion	ppb	Thulium (REE)

Field name	Variable	Type	Units	Unit abbreviation	Description
UB	U/B	measured	millimole/mole	mmol/mol	Uranium/Boron Calculated from U/Ca and B/Ca.
UCa	U/Ca	measured	millimole/mole	mmol/mol	Uranium/Calcium
USr	U/Sr	measured	millimole/mole	mmol/mol	Uranium/Strontium Calculated from U/Ca and Sr/Ca.
VCa	V/Ca	measured	millimole/mole	mmol/mol	Vanadium/Calcium
Y	Y	measured	parts per billion	ppb	Yttrium (REY)
Yb	Yb	measured	parts per billion	ppb	Ytterbium (REE)
YbCa	Yb/Ca	measured	millimole/mole	mmol/mol	Ytterbium/Calcium
YCa	Y/Ca	measured	millimole/mole	mmol/mol	Yttrium/Calcium
YHo	Y/Ho	measured	parts per billion	ppb	Yttrium/Holmium as mass ratio.
Zn	Zn	measured	parts per million	ppm	Zinc
ZnCa	Zn/Ca	measured	millimole/mole	mmol/mol	Zinc/Calcium
Zr	Zr	measured	parts per billion	ppb	Zirconium

a – data considered dimensionless (pers. comm. N. Saha)

290 b – Ce/Ce\* formula (Eq. (1)) from Saha et al. (2019b) per Lawrence et al. (2006)

## 2.7 Coral Database Metadata

The GBRC D contains 104 metadata fields in the CSV version (reduced or combined into 76 fields in the LiPD version; Appendix Table B1) that enable identification, and investigation of the coral records. Information for the metadata fields have been sourced from publications, supplementary materials, data archive information and personal communications with researchers.

295

Nine fields are notes or free-form text fields that provide useful information for future investigations and understanding the record that cannot be succinctly reduced to controlled vocabulary and/or a numerical value.

The metadata included in the GBR coral record database was divided into six categories (Identifier, Geographic, Publication, Analysis, Calibration and Dating) based on best practice standards suggested by Marine Annually Resolved Proxy Archives (MARPA) (Dassié et al., 2017), Paleoclimate Community reporting Standard (PaCTS) 1.0 (Khider et al., 2019), Paleoenvironmental Standard Terms (PaST) Thesaurus (Morrill et al., 2021) and implemented in other palaeo-archive databases such as Iso2k and CoralHydro2k (Walter et al., 2023; Konecky et al., 2020).

300

### 2.7.1 Identifier Metadata

Identifier fields were used to distinguish and characterise each coral record and identify potential crossover (or duplication) between records in the database and other research (Table 5). Identifier fields include core names (as specified in the original publication(s)), International Generic Sample Number (IGSN), core collection time, coral archive species and the record period (maximum (max) and minimum (min) year). The metadata for the record period aims to maximise users' ability to find relevant records that may only include measurements for part of a year, for example, the max year for records ending in March 2000 or November 2000 would be 2000 for both.

310

The database includes 67 duplicate core names, which generally indicate different types of analyses on the same coral head, reflecting the development and use of new analysis techniques. For example, TE/Ca may be measured from the same core using different methods (e.g. inductively coupled plasma mass spectrometer (ICP-AES) and laser ablation inductively coupled

mass spectrometer (LA-ICP-MS)) or may be measurements of different variables (e.g. TE/Ca and luminescence). However, it also demonstrates the concentration of analyses on particular coral heads. For example, five of the 21 records from Magnetic Island in the central GBR are analyses from MAG01D and include an: ~weekly resolution multi-trace element LA-ICP-MS record, ~annual resolution stable isotope record, biennial resolution multi-trace element inductively coupled plasma mass spectrometer (ICP-MS) record, quinquennial resolution stable isotope and trace element record, and annual resolution luminescence record.

A notes field (cdata\_coralNotes; Table 5) is also included for additional information on the coral, including issues and observations for consideration if using the record.

**Table 5. Identifier metadata for the CSV version of the GBRCD for each coral record including database ID, publication core name, species, and time span.**

Field name	Variable	Type	Description
cdata_datasetID	Dataset ID	text	A specific identifier assigned to all records, with specifics for the publication and site.
cdata_coreName	Core name published	text	Core name as specified in text in publications. Allows for tracing of coral records through past and future publications. <sup>a</sup>
cdata_altCoreName	Core name alternate	text	Alternate core name to cdata_coreName if more than one was used in the publication or where a different name was specified in datasets or secondary sources.
cdata_IGSN	International Generic Sample Number	text	IGSN identifier for sample identification of the physical coral samples. At the time of publication this field is empty.
cdata_collectTime	Time collected	text	The year and month (if available) the coral core was collected from living coral. Format is [year CE-month]. Alternatively, if core was from non-living coral, it was described as “Fossil” in this field.
cdata_minYear	Min year	numeric	First/oldest year of record. Recorded as integer years CE. Coral age decimal values were rounded down to the calendar year integer. Negative values are years BCE.
cdata_maxYear	Max year	numeric	Last/youngest year of record. Recorded as integer years CE. Coral age decimal values were rounded down to the calendar year integer. Negative values are years BCE.
cdata_archiveSpecies	Coral species	text	Coral genus and species (if known). Records where species name was unknown or not given were written as ‘[Genus] sp.’ <sup>a</sup> Note that species level identification of coral species (particularly <i>Porites</i> spp.) is open to disagreement and thus carries a level of uncertainty.

Field name	Variable	Type	Description
cdata_isDatabaseDuplicate	Duplicate coral core name flag	logic	Indicates whether coral core name appears more than once in GBRCd. Flags if coral records were split, and/or where the core name (cdata_coreName & cdata_altCoreName) was identical between publications.
cdata_dataCoverageGroup	Record coverage	numeric	The group the record was sorted into based on the length of the record. Groups range from 1 to 3 as described in Table 6.
cdata_coralNotes	Coral notes	text	Any notes on coral and additional information that does not fit into other fields e.g. list of core names (if known) for composite records, record observations and considerations.

a – Descriptions based on Walter et al. (2023) CoralHydro2k database fields.

325

**Table 6. Coral record coverage group and description relating to the field cdata\_dataCoverageGroup.**

Category	Description
1	Record length >100 years
2	Record length 10–100 years
3	Record length <10 years

### 2.7.2 Geographic Metadata

Geographic fields (Table 7) provide further core identification information and give the physical location of the coral archive: geographic coordinates, site name and depth of the colony (termed ‘elevation’ below).

330 Coral records that did not have listed GPS coordinates for the data were assigned GPS coordinates using “Reefs and shoals – Queensland” dataset (EPSG:4283 - GDA94 coordinate reference system) from State of Queensland (Department of Natural Resources, Mines and Energy) 2021, available under a [Creative Commons - Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) licence. GPS coordinates were approximately matched to the supplied location if a sufficiently detailed map was included in the publication, or the coordinates approximating the reef centre were used. Where coordinates were supplied for the GBRCd, or the coordinates have been changed or updated from what was published in the original publication, this is noted in the geo\_notes field.

340 Elevation information is supplied for coral records in the geo\_elevation field. This value includes uncertainties inherent in the published data as elevation may describe depth below sea level to top or bottom of the coral and this is not always explicitly described. Likewise, the choice of height datum used to describe sea level is also not always explicitly stated. Fossil coral records from drill core matrices have the depth down core range (if known), as well as an estimated depth within that range, noted in the geo\_notes field. As there is uncertainty around the estimated depth down core, as well as uncertainty due to past changes in GBR sea level compared to present-day sea level (Lewis et al., 2013; Leonard et al., 2020; Hopley et al., 2007) it is left to the user to convert depth down core to an elevation relative to present-day sea level for fossil corals.

345 **Table 7. Geographic metadata for the CSV version of the GBRC D for record(s) describing the physical location of each coral archive sampling location.**

<b>Field name</b>	<b>Variable</b>	<b>Type</b>	<b>Description</b>
geo_latitude	Latitude	numeric	Latitude for the coral. All values are negative in the GBRC D due to the GBR’s location south of the equator. See geo_notes for considerations.
geo_longitude	Longitude	numeric	Longitude for coral. All values are positive in the GBRC D due to the GBR’s location within 180° east of the Prime Meridian. See geo_notes for considerations.
geo_siteName	Site	text	Structured location names. Format follows site name: [name] Reef/Island e.g. Abraham Reef or Havannah Island.
geo_siteName2	Site	text	Specific location name (if available) as supplied in publications, e.g. Geoffrey Bay, Heron Island Reef.
geo_elevation	Elevation	numeric	Elevation, if known, of corals in meters (m). Values are negative to indicate coral material was collected below present-day sea level or positive to indicate coral material was collected above present-day sea level. Where a range of values is reported the elevation is the average of the range.
geo_notes	Elevation notes	text	Any notes on coral geographic information. This includes reported specifics of coral elevation, such as the depth range listed or relationship to water height, information on fossil coral depths from drill core matrices (if available) and considerations.

### 2.7.3 Dating Metadata

350 The GBRC D includes 18 BCE records that have chronologies determined in part by U-Th or radiocarbon dating. The dating metadata (Table 8) provides a brief overview of information available in the related record publications and includes either the corrected U-Th age or uncalibrated <sup>14</sup>C age (± uncertainty) referenced to 1950 (yBP; years before present) and a summary of related information and/or assumptions.

The complete record (or all available) of essential information for reporting U-Th and radiocarbon dating can be found in the original publications. Dating information published for the first time in this study is supplied in Appendix Table D1.



355 **Table 8. Dating metadata for the CSV version of the GBRCd. Basic information for dated samples.**

Field name	Variable	Type	Description
dating_UThDate	U-Th Date	numeric	Uranium-thorium (U-Th) age corrected for initial thorium. If more than one age was measured, this field lists the youngest age, or the age used in the original record chronology. Units: yBP.
dating_UThDateUncertainty	U-Th Date Uncertainty	numeric	Uranium-thorium (U-Th) age (corrected) uncertainty. If more than one age was measured, this field lists the uncertainty for the youngest age, or the age used in the original record chronology. Units: yBP.
dating_14CDate	<sup>14</sup> C Date	numeric	Uncalibrated radiocarbon ( <sup>14</sup> C) age. If more than one age was measured, this field lists the youngest age, or the age used in the original record chronology. Units: yBP.
dating_14CDateUncertainty	<sup>14</sup> C Date Uncertainty	numeric	Uncalibrated radiocarbon ( <sup>14</sup> C) age uncertainty. If more than one age was measured, this field lists the uncertainty for the youngest age, or the age used in the original record chronology. Units: yBP.
dating_notes	Additional dating information	text	Any additional information relating to dating of corals such as the local delta-R reservoir correction used as well as assumptions used for database.

#### 2.7.4 Analysis Metadata

Analysis fields provide information about, or obtained, during measurement of coral variables (Table 9). It includes the list of proxies measured, information about sampling, analytical methods (acronyms given in Table 10), data resolution (as contained in the database record; Table 11) and analytical precision of measurements (annotations given in Table 12). It also includes information such as whether the international coral reference JcP-1 (Okai et al., 2002; Hathorne et al., 2013) was measured and information published about screening for diagenetic alteration of the corals.

Coral data resolution is described as the minimum, maximum, mean and median number of data points per year for each record. The measurements per year were calculated for complete calendar years based on all unique age dates for each record, as records were filtered assuming duplicate dates represent replicate samples. A nominal data resolution label was assigned to each record based on the calculated median resolution described in Table 11 and is a key field for filtering records in the database. The term nominal resolution suffix ‘\_uneven’ has been adopted from the CoralHydro2k database (Walter et al. 2023) to describe corals with a variable resolution (i.e. where the record’s min, max, mean, and median resolution are not the same). A notes field (meths\_methodNotes; Table 9) is also included for information on the measurement method(s) including the sampling method, and any additional methodology notes, as well as an identifier (“REEs measured”) for every record where any of the REEs (as listed in Sect. 2.6 above) were included in the record.

**Table 9. Analysis metadata for the CSV version of the GBRC. Description of fields relating to the measurement of coral data, including variables measured, analysis method and the age model used. Note ‘X’ is used to represent a list of variables as stated in the description.**

<b>Field name</b>	<b>Variable</b>	<b>Type</b>	<b>Description</b>
meths_primaryVariablesList	Proxy variables	text	List of proxy variables measured from coral archive. See Table 4 for complete list of variables.
meths_additionalVariablesList	Additional variables	text	List of variables that are not standard climate or environment proxy variables or are variables calculated from the measurement of proxy variables. See Table 4 for complete list of variables.
meths_XMethod	Method of X measurement	text	Method used for data measurement. (X = TE, Isotope, Lumin) e.g. ICP-MS. Where multiple TE or Isotope methods are used across variables, this field lists the Sr/Ca or $\delta^{18}\text{O}$ method and other methods in meths_altMethodInfo. See Table 10 for the complete list.
meths_XMethodMachine	Machine for X data measurement	text	Description of machine used for data measurement e.g. Finnigan MAT 251. (X = TE, Isotope, Lumin)
meths_altMethodInfo	Alternate proxy measurement information	text	List of proxy, method and machine used (if available) for data measurements not included in other method and machine information fields. Items are separated by “;”.
meths_methodNotes	Sampling notes	text	Any notes on methods including a brief description of sampling method, and other record measurement notes that do not fit in other fields.
meths_hasResolutionNominal	Nominal resolution	text	Nominal temporal resolution of the coral record. Nominal choice is based on median frequency of coral data. See Table 11 for term definitions.
meths_resolutionMin	Minimum resolution	numeric	Minimum temporal resolution of the record, expressed as data points per year.
meths_resolutionMax	Maximum resolution	numeric	Maximum temporal resolution of the record, expressed as data points per year.
meths_resolutionMean	Mean resolution	numeric	Mean temporal resolution of the proxy record, expressed as data points per year.
meths_resolutionMedian	Median resolution	numeric	Median temporal resolution of the record, expressed as data points per year.
meths_isAnomaly	Anomaly data flag	logic	Indicates whether [variable] values are considered anomaly data. Anomaly definition and therefore calculation method may vary by publication.

Field name	Variable	Type	Description
meths_chronologyNotes	Notes on age model	text	Brief notes on method used to determine coral data age model. Field includes the target SST dataset and interpolation information if supplied. SST datasets include AIMS (Australian Institute of Marine Science), Extended Reconstructed SST (ERSST), Comprehensive Ocean-Atmosphere Data Set (COADS), Integrated global ocean station system (IGOSS), Hadley Centre Sea Ice and SST (HadISST), NOAA/Reyn-Smith Optimum Interpolation SST (OISST). Also includes methods for determining the age model for fossil corals where the age model incorporates the absolute age (Table 8).
meths_coralExtensionRate	Extension rate	numeric	Average coral extension rate in mm/year. If a range was given in the publication, 'Extension rate' is the average of the range. <sup>a</sup>
meths_coralExtensionRateNotes	Extension rate notes	text	Coral extension rate given in the publication or determined from data. This field includes the units, uncertainty, or ranges in values which are not included in the previous field.
meths_tissueThickness	Tissue thickness	numeric	Average coral tissue thickness in millimetres (mm).
meths_jcpUsed	Flag for measurement of JcP-1	logic	Indicates whether the JcP-1 standard was measured.
meths_jcpSrCaValue	Measured value of JcP-1	numeric	Sr/Ca value measured for JcP-1 standard. Units: mmol/mol.
meths_XAnalyticalPrecision	Measured X analytical precision	numeric	Analytical error for measured proxy values based on a standard (X = Sr/Ca, U/Ca or $\delta^{18}\text{O}$ ).
meths_XAnalyticalPrecisionUnits	X analytical precision units	text	Units of analytical error. (X = Sr/Ca, U/Ca or $\delta^{18}\text{O}$ ). Units are relative standard deviation (RSD; %) for Sr/Ca and U/Ca, or permille (permil) for $\delta^{18}\text{O}$ .
meths_altPrecisionList	Alternate proxies with analytical precision data	text	List of proxies with supplied analytical precision in addition to Sr/Ca, U/Ca or $\delta^{18}\text{O}$ . See Table 12. Provided to enable comparing precision between studies and assess use of records for research. Items are separated by “;”.

Field name	Variable	Type	Description
meths_altAnalyticalPrecision	Measured analytical precision	text	List of analytical error value(s) based on a standard for alternate proxies. Values are numerical and are separated by “;” with the order matching meths_altPrecisionList. See Table 12 for the list.
meths_altAnalyticalPrecisionUnits	X analytical precision units	text	List of units for alternate proxy analytical error(s). Units are separated by “;” with the order matching meths_altPrecisionList. See Table 12 for the list of units.
meths_archiveDiagenesisCheck	Diagenesis screening information	text	Information about diagenesis screening as reported in publications, including the method(s) used to check diagenesis. Where screening information was supplied it is reported as ‘Yes; [method(s)]’. Methods used to check include X-Ray Diffraction (XRD), petrographic thin section (TS), scanning electron microscopy (SEM), ultraviolet light assessment (UV).

a – Descriptions based on Walter et al. (2023) CoralHydro2k database fields.

375

**Table 10. Description of abbreviations of methods used for the meths\_XMethod field given in Table 9.**

Method Acronym	Method
AMS	Accelerator Mass Spectrometry
GC	Gas counting
ICP-AES	Inductively Coupled Plasma Atomic Emission Spectrometry
ICP-OES	Inductively Coupled Plasma Optical Emission Spectrometry
ICP-MS	Inductively Coupled Plasma Mass Spectrometry
ID-TIMS	Isotope Dilution Thermal Ionisation Mass Spectrometry
IRMS	Isotope Ratio Mass Spectrometry
LA-ICP-MS	Laser Ablation Inductively Coupled Plasma Mass Spectrometry
LSA	Liquid Scintillation Analysis
MC-ICP-MS	Multicollector Inductively Coupled Plasma Mass Spectrometry
MS	Mass Spectrometry
PTIMS	Positive Thermal Ionisation Mass Spectrometry
Q-ICP-MS	Quadrupole Inductively Coupled Plasma Mass Spectrometry
SSAMS	Single Stage Accelerator Mass Spectrometry
TIMS	Thermal Ionisation Mass Spectrometry

**Table 11. Description of terms for nominal resolution**

Nominal resolution	Data points per year
weekly; weekly_uneven	>= 52 data points per year; “_uneven” was added to records with variable resolutions that typically have over 52 data points per year.
fortnightly; fortnightly_uneven	26 data points per year; “_uneven” was added to records with variable resolutions that typically have 26–51 data points per year.
monthly; monthly_uneven	12 data points per year; “_uneven” was added to records with variable resolutions that typically have 12–25 data points per year.
bimonthly; bimonthly_uneven	6 data points per year; “_uneven” was added to records with variable resolutions that typically have 6–11 data points per year. <sup>a</sup>
quarterly; quarterly_uneven	4 data points per year; “_uneven” was added to records with variable resolutions that typically have 4–5 data points per year. <sup>a</sup>
biannual; biannual_uneven	2 data points per year; “_uneven” was added to records with variable resolutions that typically have 2–3 data points per year. <sup>a</sup>
annual; annual_uneven	1 data point per year; “_uneven” was added to records with variable resolutions that typically have 1 data point per year, and are missing years of data.
>annual	Less than 1 data point per year. <sup>a</sup>

a – Descriptions based on Walter et al. (2023) CoralHydro2k database fields.

380

**Table 12. List of isotopes, trace elements and trace element ratios with analytical precision values supplied in meths\_altPrecisionList. The list of variables is formatted as per meths\_altPrecisionList. Units are as provided by original publication(s) or as supplied.**

Type	List	Units
Isotope	d11B, d13C, d14C, d15N	permille (permil)
Trace Element	Ba, Mn, Th, Y	RSD (%)
Trace Element Ratio	BaCa, Ba138Ca, BCa, BMg, LiMg, MgCa, MnCa, REEsCa, Sr88Ca <sup>a</sup> , YCa	RSD (%), mmol/mol

a – Sr88Ca = Sr<sup>88</sup>/Ca as per D'olivo and Mcculloch (2022).

### 2.7.5 Calibration Metadata

385 Calibration metadata (Table 13) provides information about the creation of proxy-environment equations. Due to the ongoing focus and broader regional significance of proxy-SST calibrations, detailed information has been provided for proxy-SST calibrations (intercept, slope, slope uncertainty and  $r^2$ ) for ‘raw’ (non-centred) data in associated fields for Sr/Ca, U/Ca and  $\delta^{18}\text{O}$ . Raw data are the data values after measurement processing, while centred (or anomalised) data are the raw data minus the data mean (which can be the mean of the whole record or part thereof). The calibration information for these SST proxies

390 was included due to their historical use and the body of published research on their feasibility and/or application for reconstructing SSTs in the GBR. Where a centred calibration was preferred or the only one published, the calibration information is separate from the more frequently used non-centred equations. As there were additional proxy-environmental variable calibrations published for the database records, other calibration equations are provided in calibration notes where applicable. Availability of additional proxy-environment calibrations that are not provided in the database are also flagged.

395

**Table 13. Calibration metadata for the CSV version of the GBRCd. Available information on the calibration of coral records to environmental variables, specifically SST. Note ‘X’ is used to represent a list of variables as stated in the description.**

<b>Field name</b>	<b>Variable</b>	<b>Type</b>	<b>Description</b>
calib_isSSTCalibrationCoral	SST calibration coral flag	logic	Indicates whether the coral was used in creating an SST calibration in the database. Includes records that have been used to create a composite record calibration.
calib_useSSTCalibration	SST calibration flag	logic	Indicates whether there is SST-Sr/Ca, SST-U/Ca, or SST- $\delta^{18}\text{O}$ calibration information in the database. Includes records for corals used to create calibration equations and where an SST calibration was applied (e.g. fossil corals).
calib_hasAlternateCalibration	Alternate calibration information flag	logic	Indicates whether there are additional or alternate calibration equation(s) available for the record that is not included in the database. Alternate calibration equations can include non-SST calibrations (e.g. luminescence-rainfall calibration equation), and where applicable, may be included in calibration notes.
calib_notes	Calibration notes	text	Unstructured description of calibration. Includes information about proxy-SST calibrations and proxy-other calibrations if relevant.
calib_isComposite	Composite calibration flag	logic	Indicates whether calibration is a composite from more than one record.
calib_compositeList	Composite calibration record list	text	List of coral IDs (if known) used for composite calibration.
calib_SSTdata	SST product	text	Description of SST dataset(s) used for proxy-SST calibration, and treatment where applicable. SST datasets include AIMS (Australian Institute of Marine Science), Extended Reconstructed SST (ERSST), Comprehensive Ocean-Atmosphere Data Set (COADS), Integrated global ocean station system (IGOSS), Great Barrier Reef Marine Park (GBRMPA), Hadley Centre Sea Ice and SST (HadISST), NOAA/Reyn-Smith Optimum Interpolation SST (OISST). SST dataset version information is included if the information was available.
calib_fitPeriod	Calibration fit period	text	Time period used to fit coral proxy-SST calibration if known. Takes the form of [min year]–[max year].
calib_hasCentredEquation	Centred equation flag	logic	Indicates whether there is an equation based on centred (data minus data mean for whole or specified period of data).

Field name	Variable	Type	Description
calib_centredEquationNotes	Centred equation information	text	Provides information on centred equation including method, equation (as provided in publication) and $r^2$ for relevant proxy-SST calibrations.
calib_method	Regression method	text	Regression method used for the proxy-SST regression(s) in the publication. If >1 method used, the field lists the methods used for the regression values in the following fields (i.e. calibX_equationIntercept, etc.). Regression methods include Ordinary Least Squares (OLS), Reduced Major Axis (RMA), and Weighted Least Squares (WLS).
calibX_equationIntercept	Proxy-SST intercept	numeric	The proxy-SST calibration intercept published for the coral record. Calibration equations take the form $\text{proxy} = \text{slope} * \text{SST} + \text{intercept}$ . (Units: [data units]/°C) <sup>a</sup> (X = 1 (Sr/Ca), 2 (U/Ca), 3 ( $\delta^{18}\text{O}$ ))
calibX_equationSlope	Proxy-SST slope	numeric	The proxy-SST calibration slope published for the coral record. Calibration equations take the form $\text{proxy} = \text{slope} * \text{SST} + \text{intercept}$ . (Units: [data units]/°C) <sup>a</sup> (X = 1 (Sr/Ca), 2 (U/Ca), 3 ( $\delta^{18}\text{O}$ ))
calibX_equationSlopeUncertainty	Proxy-SST slope uncertainty	numeric	The proxy-SST calibration slope uncertainty published for the coral record. Calibration equations take the form $\text{proxy} = \text{slope} * \text{SST} + \text{intercept}$ . (Units: [data units]/°C) <sup>a</sup> (X = 1 (Sr/Ca), 2 (U/Ca), 3 ( $\delta^{18}\text{O}$ ))
calibX_equationR2	Proxy-SST calibration $r^2$	numeric	The proxy-SST calibration $r^2$ value published for the coral record. <sup>a</sup> (X = 1 (Sr/Ca), 2 (U/Ca), 3 ( $\delta^{18}\text{O}$ ))

a – Descriptions based on Walter et al. (2023) CoralHydro2k database fields.

### 2.7.6 Publication Metadata

400 The GBRCDB includes metadata for relevant publications relating to the coral records. The bibliographic information is supplied for up to three related publications with relevant bibliographic information separated into separate fields for ease of filtering and includes (where applicable) the DOI for the publication and the archived data (Table 14). Another field for an alternate data URL related to the coral record is also provided to enable referencing the original data location when it has been provided in article supplements or similar. If a record (or part thereof) was featured in multiple publications, the order of the

405 bibliographic information was determined by the primary publication source of the data archived, and then the publications are generally in chronological order (oldest to newest) with preference for the most complete dataset or archive and methods description for the data.

**Table 14. Publication metadata. Publication details for one to three publications linked with the coral record. Note ‘X’ is used to represent number from the list of variables as stated in description.**

Field name	Variable	Type	Description
pubX_firstauthor	First author	text	First author of publication (X = 1, 2, 3). <sup>a</sup>
pubX_year	Publication year	numeric	Year of publication (X = 1, 2, 3). <sup>a</sup>
pubX_doi	DOI	text	DOI of publication (X = 1, 2, 3). <sup>a</sup>
pubX_citation	Full citation	text	Complete citation of publication (X = 1, 2, 3). <sup>a</sup> Form is based on the Harvard reference style.
pubX_authors	Full author list	text	Full list of authors (in the order published) from each publication (X = 1, 2, 3), including the first listed author.
pubX_title	Title	text	Title of publication (X = 1, 2, 3). <sup>a</sup>
pubX_journal	Journal	text	Journal of publication (X = 1, 2, 3). <sup>a</sup>
pubX_doiData	Data DOI	text	DOI for dataset(s) (X = 1,2,3).
pubX_altDataURL	Alternate Data URL	text	Alternate/additional DOI or links to published data related to the coral record.
pub_isCoreIDOtherStudy	Core used in other studies	logic	Indicates if coral core ID was known to be used in other studies. Other studies may or may not be included in the database as the study data might not have been available or might be data that does not meet the GBRC D selection criteria (e.g. coral growth studies). If the core ID supplied was ambiguous (i.e. was the site location name), then it was not linked with other studies. Flag is to best of knowledge and may not be definitive.
pub_coreIDOtherStudyList	Publications of alternate studies of core	text	List of references where core has been used. Reference is in short form and full list is in the Appendix Table A1. List is to best of knowledge and may not be definitive.

410 a – Descriptions based on Walter et al. (2023) CoralHydro2k database fields.

### 3 Database use and citation

The GBRC D is a comprehensive compilation of coral records from the GBR to date that span -5885 to 2017 CE. The database enables the investigation of multiple environmental factors via various coral proxy systems for the GBR, northeastern Australia and potentially the broader Indo-Pacific. Researchers can select from a list of 78 measured or calculated variables to investigate the GBR environment or determine new proxy-environment relationships.

415


The GBRC D is freely available on the NOAA/WDS Paleoclimatology website at <https://doi.org/10.25921/hqzk-8h74> (Arzey et al. 2024).

It is up to the user to determine applicable proxies for their investigation and filter records as needed. Best practice (Fig. 2) should begin with filtering by proxy or proxies of interest. Next the CSV dataset(s) should be joined with the metadata fields to enable further filtering of the database. It is then suggested to subsequently filter the data by other metadata of interest such as record coverage, resolution(s), and archive species. Users should read the field cdata\_coralNotes for all records of interest as this includes relevant information on caveats and assumptions (if any) that were noted for each record.

420

Sample code is available for R and Python.





425 **Figure 2. Flow chart of best practice handling of the two database file versions.**

As previously noted, care is advised for the use of all records in the GBRCDB as relevance of the measured variables may vary by location or resolution and so forth. Records have been collated for the GBRCDB and basic quality checks have been completed on the data to enable archiving for this study.

430 If researchers use the GBRCDB they should cite this study and the most recent version description of the GBRCDB. It is also best practice to cite the original publications of the data (Table A1) alongside the GBRCDB database. Citation information and reference links to the original publications and data archive are included in the metadata (i.e. Table 14).

## 4 Characterising the Database


### 4.1 Record coverage

435 The GBRCDB compiles 208 individual records published since the early 1990s from 58 named reefs and islands (Fig. 1). There are 193 (92.8 %) records from *Porites* spp., eight (3.8 %) from *Isopora* spp., six (2.9 %) from *Acropora* spp., and one (0.5 %) from a *Cyphastrea* sp. coral. The coral records include 78 variables, of which 72 are measured and six are calculated. The majority (116) of records are from the central GBR region (17 to 20° S), 74 from the southern GBR (20 to 24° S), and 18 records from the northern GBR (11 to 17° S).

440 Most records (58.2 %) have a maximum ('core top') age between 2000 CE and 2017 CE, and 73.1 % between 1990–2017 CE. Peak density of records occurs in the 1980s with  $\geq 105$  records in each year between 1980–1984 and nearly as many ( $\geq 100$ ) in the early 1990s (1990–1993) (Fig. 2). Only 59 (28.4 %) of the records' minimum age occurs before 1900 CE, and 18 of those are fossil coral records that range from ~2296 to 5885 years BCE. Of these fossil records, 16 (88.9 %) are from the central GBR and two are from the southern GBR.

445 Of the 140 records measuring trace elements in the GBRCDB, 49.3 % of records are a result of the development of laser ablation methods. There have been 69 LA-ICP-MS trace element records published and 71 trace element records analysed by other mass spectrometry methods. Many (54 records) of the non-LA-ICP-MS records are from studies that assess or use SST proxies, likely due to the expectation that trace element records from laser ablation methods have a poorer fit with SST than TE records derived from solution methods. For example, Wu et al. (2021b) LA-ICP-MS study found a notable poor periodicity and Sr/Ca-SST relationship in two out of six records, which led them to suggest the usefulness of LA-ICP-MS for Sr/Ca-SST analysis is potentially limited by the relatively larger LA-ICP-MS analytical error than other mass spectrometry methods.

450




**Figure 3. Temporal coverage of records in the GBRCD colour coded by (a) record coverage Group 1–3 as per Table 6 (1 – records >100 years; 2 – records >10 – <100 years; 3 – records <10 years), includes an inset that enlarges the period -6000 to -2000 CE, (b) subset of (a) showing only records since 1550 CE, (c) nominal resolution as per Table 11 (see figure legend) (records with “\_uneven” nominal resolution were combined with even resolution records), includes an inset that enlarges the period -6000 to -2000 CE, and (d) subset of (c) showing only records since 1550 CE. The ‘spikiness’ visible in (b) Group 1 & (d) >annual resolution records is due to the combination of biennial and quinquennial resolution records.**

The majority of records in the database include Sr/Ca measurements (118 of 208; Fig. 4), and 63 of these records are used in SST calibrations. The other variables measured most frequently include B/Ca (59 records), Ba/Ca (78 records. 4), Mg/Ca (73 records; Fig. 4), U/Ca (77 records; Fig. 4), and luminescence (57 records; Fig. 5). There are 51 isotope records, of which 27 include  $\delta^{18}\text{O}$  records (Fig. 5), 17  $\delta^{13}\text{C}$ , 15  $\delta^{11}\text{B}$ , 6  $\delta^{14}\text{C}$ , 5  $\delta^{15}\text{N}$  (Fig. 5), 1  $\delta^{66}\text{Zn}$  and 1  $\delta^{98}\text{Mo}$ . Additionally, there are 18 records measuring REEs (Fig. 5), noting this number excludes records that only measure yttrium and no REEs.


The number and spatial distribution of Sr/Ca records (Fig. 4C) reflects the dual utility of the ratio as an SST proxy and to determine the coral age model. Similarly, U/Ca (Fig. 4D) is also used as an SST proxy and to determine coral age models. When combined with other proxies (e.g. Sr/Ca), U/Ca may improve SST predictions (Wu et al., 2021b; Decarlo et al., 2016). The B/Ca proxy is commonly measured in addition to target proxies rather than being the focus of the study. However, B/Ca has more recently been paired with  $\delta^{11}\text{B}$  in studies examining pH, to determine the dissolved inorganic carbon in the calcifying fluid. Mg/Ca has been measured relatively frequently (Fig. 4B), and while known as an SST proxy in other genera, the Mg/Ca-SST relationship in *Porites* spp. is weak or results are highly variable (e.g. Sadler et al., 2014; Wu et al., 2021b), possibly due to the coral actively discriminating against  $\text{Mg}^{2+}$  during biomineralisation (Marchitto et al., 2018). Ba/Ca (Fig. 4A) is frequently measured in the GBR, primarily to assess terrigenous inputs and river flow, and is mainly measured by LA-ICP-MS (58 of 78 Ba/Ca records).

Despite the widespread analysis of coral  $\delta^{18}\text{O}$  across the tropical Pacific (~100  $\delta^{18}\text{O}$ -only and paired Sr/Ca-  $\delta^{18}\text{O}$  records (Walter et al., 2023)), there are few (only 27)  $\delta^{18}\text{O}$  records available for the GBR (Fig. 5A), especially considering the length

475 of time  $\delta^{18}\text{O}$  has been used as an SST or hydrological proxy. This is potentially due to the paired forcing (SST and hydrology) that determines the coral  $\delta^{18}\text{O}$  for nearshore corals, as well as the development and use of coral luminescence for assessing freshwater inputs into the GBR. Records including luminescence measurements make up 27.4 % of the GBRCD (Fig. 5B).



480 **Figure 4. Location map of GBRCD records for common major trace element ratios; (a) Ba/Ca, (b), Mg/Ca, (c) Sr/Ca and (d) U/Ca. The size of the point indicates the number of records, and is specific to each map.**



485 **Figure 5. Location map of GBRCD records for (a)  $\delta^{18}\text{O}$ , (b) luminescence (Lumin), (c)  $\delta^{15}\text{N}$  and (d) Rare Earth Elements (REEs). The size of the point indicates the number of records, and is specific to each map. Note that Lumin\* includes all the methods included in the GBRCD for measuring luminescence.**

## 5 Discussion

There has been a push for coral records to include a range of details to meet minimum reporting standards for palaeo-archive records (such as MARPA (Dassié et al., 2017) and PaCTS (Khider et al., 2019)). These community-led publications report the ideal metadata that should be supplied with the publication of each record and provide a useful guide for best practice. The call for minimum reporting standards for data and metadata is echoed here. A particular challenge in producing the GBRC D was that metadata was often spread across multiple publications. To improve traceability between publications, relevant metadata could be included in the supplement and in a publicly available data repository. The addition of IGSNs to capture the metadata associated with the physical samples would also improve the utility of the coral data (Dassié et al., 2017).

Another area for improvement is the reporting of screening corals for diagenesis. Diagenesis is a known source of error in geochemical analysis as it can remove the primary coral proxy signal (Hendy et al., 2007; Sayani et al., 2011; Mcgregor and Gagan, 2003; Nothdurft et al., 2007; Quinn and Taylor, 2006; Weerabaddana et al., 2024). Only 40 of 208 records in the GBRC D reported on diagenetic alteration. The majority fail to mention the screening method(s) or outcomes (see `meths_archiveDiagenesisCheck`) in the publication text, supplementary materials or could not be determined from cross-referencing sample IDs in other publications. There is no established standard of best practice for screening corals for diagenesis at present, though it is suggested that it is ideal to use petrographic analysis with a combination of methods such as XRD, and densitometry (Mcgregor and Abram, 2008). Other emerging approaches (e.g. Murphy et al., 2017; Takada et al., 2017) may also be useful for diagenesis screening. However, due to the strengths and weaknesses of the various methods as noted by Nothdurft and Webb (2009), examination by petrographic analysis (preferably using scanning electron microscopy) is required to confirm the type and level of diagenetic alteration (if any) present in a sample. Given the consequences of diagenesis for palaeoclimate and palaeoenvironment reconstructions from coral records we strongly advocate for authors to report their diagenetic screening methods in their publications or supplements.

X-radiographs showing sampling paths should also be routinely published with the data, either as a figure in the paper or as supplementary material. Pioneering work on corals from the GBR showed that it is essential for corals to be sampled along the maximum growth axis. Providing this information was previously common (Alibert and Mcculloch, 1997; Calvo et al., 2007; Pelejero et al., 2005; Chakraborty et al., 2000; DeLong et al., 2007), but x-radiographs have not been published consistently in recent years. Computed tomography (CT) scans of coral cores are sometimes used instead of, or in addition to, X-radiographs. Unlike X-radiographs, CT scans can be run on uncut coral cores and drill core matrices and similar to X-radiographs, can provide information about extension rates, density and calcification (Bosscher, 1993; Mollica et al., 2018; Crook et al., 2013), as well as growth morphologies and bioerosion (Prouty et al., 2017; Decarlo et al., 2015; Li et al., 2021). As with X-radiographs, the CT scan images providing relevant information should be included with the paper.

With the continued passage of time, it is likely there will be further loss of existing coral records, physical samples and associated data. Due to physical misplacement or destruction of data files, inability to read old computer files and/or retirement of researchers with existing records, the coral data underpinning early coral research may be lost forever. We urge researchers in possession of existing records to reach out to archival teams such as those sustaining existing databases (e.g. the GBRC D or the CoralHydro2k database (Walter et al., 2023)) to ensure records are archived for future use. An email address for submissions to the GBRC D is provided at the GBRC D GitHub and NOAA/WDS Paleoclimatology study page to facilitate capture of legacy or new coral data.

The GBRC D lends itself to meeting some of the suggestions put forward by Lough (2004) to improve the use of corals for palaeoclimatology. While not absolute replication, the GBRC D facilitates using multiple records to determine common environmental signals and identify non-climatic artifacts in those records. For example, the database enables combining multiple records to assess trends, similar to methods used by Hendy et al. (2002).

The long and comprehensive history of coral research on the GBR can make it difficult to find relevant research and records to understand the current state of coral palaeoclimate and palaeoenvironment studies on the GBR and thus identify gaps in existing research. The GBRCDB can be used to discern where future coral proxy records are needed. Additionally, the database includes 67 duplicate core names, which generally indicates different types of analyses on the same coral head and may provide an opportunity for comparisons between methods.

The GBRCDB can be used for analysis and examination of trends at local to regional scales. Records from the GBR database can be selectively used alongside other datasets to assess the past environment of Australia and/or the Pacific and identify trends and periods of note. It enables comparisons of different periods, such as the modern corals and older Holocene coral records, making it possible to identify notable changes between the time periods. Furthermore, records from the GBRCDB can be combined with new and existing models of the palaeoenvironment.

## 5.1 Database Availability and its Future

This is stage 1 of the GBRCDB.

The database is archived on the NOAA/WDS Paleoclimatology website. To access the database, visit <https://doi.org/10.25921/hqzk-8h74> (Arzey et al., 2024) and click on the “NOAA Study Page” link. Example scripts to help users access the database, in either format, are available on the NOAA/WDS Paleoclimatology study page. The example scripts and database information are also available on the GBRCDB GitHub (<https://github.com/arzeyak/GBR-Coral-Skeletal-Records-Database>; GBRCDB, 2024).

Future updates to the database are expected to occur on an annual basis. We welcome submission of historic GBR coral record data for inclusion in the GBRCDB, including existing unpublished datasets, supporting FAIR data principles (Wilkinson et al., 2016). Submitted records should meet general criteria outlined above in ‘Selection criteria for inclusion in GBRCDB’. For researchers that want to submit records (published or unpublished) to the GBRCDB, please check the GBRCDB GitHub for details on how to submit your records for the database via the email address available on the GBRCDB GitHub and NOAA/WDS Paleoclimatology study page.

## 6 Conclusions

The GBRCDB promotes efforts to ensure important research data is archived and that existing coral records are not lost to time. In conclusion:

1. The GBRCDB is a comprehensive database of coral records from the GBR produced and published since the early 1990s, as well as seven new records that are first published here. The database archives 208 coral records that are publicly available at NOAA/WDS Paleoclimatology.
2. Records have been standardised and converted to a machine-readable format that is easy to use to investigate the database, and to select records of interest for analysis. With detailed metadata in the GBRCDB there are many avenues for investigating the past and present GBR environment.
3. There is scope in the future for improving the archiving of coral research datasets by building on the GBRCDB and including the minimum recommended metadata information for traceability and use.

## Appendix A – GBRC D Records

565 **Table A1. Reference data for publications cited in the GBRC D. Primary and secondary variables are the variable name as per Table 4. Citations in the Record Publication(s) columns are listed in the order presented in the database (pub1; pub2; pub3). Citations in the Additional Core Publication(s) column are listed in alphanumerical order.**

Dataset ID	Core Name	Latitude	Longitude	Location	Primary Variables (Secondary Variables)	Record Publication(s)	Additional Core Publication(s)
AL03DAV01_1	Davies 2 (side)	-18.8	147.7	Davies Reef	Sr/Ca (Distance)	Alibert et al. (2003); Alibert and Mcculloch (1997)	Fallon (2000); Fallon et al. (2003)
AL03DAV01_2	Davies 2 (side)	-18.8	147.7	Davies Reef	Ba; Ba/Ca; Cd; Mn; Mn/Ca; P; Y (Distance)	Alibert et al. (2003)	Alibert and Mcculloch (1997); Fallon (2000); Fallon et al. (2003)
AL03PAN01_1	PAN 98-2 (B3)	-18.8	146.4	Pandora Reef	Sr/Ca (Distance)	Alibert et al. (2003)	
AL03PAN01_2	PAN 98-2 (B3)	-18.8	146.4	Pandora Reef	Ba/Ca; Mn; P (Distance)	Alibert et al. (2003)	
AL03PAN01_3	PAN 98-2 (B1)	-18.8	146.4	Pandora Reef	Ba/Ca; Cd; Mn; P; Y (Distance)	Alibert et al. (2003)	
AL03PAN01_4	PAN 98-2 (B1)	-18.8	146.4	Pandora Reef	Ba/Ca; Cd; Mn; Mn/Ca; P; Y (Distance)	Alibert et al. (2003)	
AR24DIP01	Dip-04	-18.414	147.448	Dip Reef	Ba/Ca; Mg/Ca; Sr/Ca; U/Ca; Y/Ca (Distance)	This study	
AR24OTI01	GUT12	-23.5053	152.0923	One Tree Island	Ba/Ca; Sr/Ca (Distance)	This study	
AR24OTI02	OTI 1-3	-23.499	152.072	One Tree Island	Ba/Ca; B/Ca; Ca/Ca; Mg/Ca; Sr/Ca; U/Ca (Distance)	This study	
AR24OTI03	OTI 3-1	-23.499	152.072	One Tree Island	Ba/Ca; B/Ca; Ca/Ca; Mg/Ca; Sr/Ca; U/Ca (Distance)	This study	
AR24OTI04	OTI4_6C (OTI-4-19- 6C)	-23.499	152.072	One Tree Island	Ba/Ca; B/Ca; Ca/Ca; Mg/Ca; Sr/Ca; U/Ca (Distance)	This study	

<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
AR24OTI05	OTI4_6D (OTI-4-22-6D)	-23.499	152.072	One Tree Island	Ba/Ca; B/Ca; Ca/Ca; Mg/Ca; Sr/Ca; U/Ca (Distance)	This study	
AR24SLY01	SR	-19.25	149.1167	Stanley Reef	$\delta^{13}\text{C}$ ; $\Delta^{14}\text{C}$ ; $\delta^{18}\text{O}$	This study	Chakraborty (1993); Chakraborty et al. (2000)
BR17HER01	GBR-637	-23.442	151.914	Heron Island	$\delta^{18}\text{O}$ ; Sr/Ca	Brenner et al. (2017)	
BR17HER02	GBR-641A	-23.442	151.914	Heron Island	$\delta^{18}\text{O}$ ; Sr/Ca	Brenner et al. (2017)	
BR17HER03	GBR-538	-23.442	151.914	Heron Island	$\delta^{18}\text{O}$ ; Sr/Ca	Brenner et al. (2017)	
BR17HER04	HER-1F (HER-OR)	-23.442	151.914	Heron Island	$\delta^{18}\text{O}$ ; Sr/Ca	Brenner et al. (2017)	
BR17HER05	HER-13 (HER12-13)	-23.442	151.914	Heron Island	$\delta^{18}\text{O}$ ; Sr/Ca	Brenner et al. (2017)	Linsley et al. (2019)
CA07FLI01	Flinders (FLI02A)	-17.73	148.43	Flinders Reef	$\delta^{18}\text{O}$ ; Sr/Ca	Calvo et al. (2007)	Pelejero et al. (2005)
CH93SLY01	SR	-19.25	148.1167	Stanley Reef	$\delta^{13}\text{C}$ ; $\delta^{18}\text{O}$ (Density; Distance)	Chakraborty (1993); Chakraborty et al. (2000)	
CH93SLY02	SR	-19.25	148.1167	Stanley Reef	$\delta^{13}\text{C}$ ; $\delta^{18}\text{O}$ (Density; Distance)	Chakraborty (1993); Chakraborty et al. (2000)	
DE14ARL01	10AR1	-16.6381	146.1036	Arlington Reef	$\delta^{13}\text{C}$ ; $\delta^{18}\text{O}$ ; Mg/Ca; Sr/Ca ( $\delta^{18}\text{O}_{\text{seawater}}$ ; Extension)	Deng et al. (2014)	
DE14ARL02	10AR2	-16.6381	146.1036	Arlington Reef	$\delta^{13}\text{C}$ ; $\delta^{18}\text{O}$ ; Mg/Ca; Sr/Ca ( $\delta^{18}\text{O}_{\text{seawater}}$ ; Extension)	Deng et al. (2014); Wang et al. (2019)	Chen et al. (2021); Xiao et al. (2020)
DO15GBR01	1709_6	-17.876	146.729	Reef 17-065	$\delta^{11}\text{B}$	D'olivo et al. (2015)	D'olivo et al. (2018)



<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
DO15HAV01	HAV09_3	-18.836	146.5259	Havannah Island	$\delta^{11}\text{B}$	D'olivo et al. (2015)	D'olivo and Mcculloch (2017); D'olivo et al. (2013); D'olivo et al. (2018)
DO15HAV02	HAV09_3	-18.836	146.5259	Havannah Island	$\delta^{11}\text{B}$	D'olivo et al. (2015); Chen et al. (2021)	D'olivo and Mcculloch (2017); D'olivo et al. (2013); D'olivo et al. (2018)
DO15HAV03	HAV06A	-18.84	146.54	Havannah Island	$\delta^{11}\text{B}$	D'olivo et al. (2015); Chen et al. (2021)	D'olivo et al. (2013)
DO15PAN01	PAN02 (PAN02A)	-18.81	146.43	Pandora Reef	$\delta^{11}\text{B}$	D'olivo et al. (2015); Chen et al. (2021)	D'olivo et al. (2013)
DO15RIB09	RIB09_3	-18.48	146.87	Rib Reef	$\delta^{11}\text{B}$	D'olivo et al. (2015)	D'olivo et al. (2013); D'olivo et al. (2018)
DO17HAV01	HAV09_3 (Path A)	-18.836	146.5259	Havannah Island	$\delta^{11}\text{B}$ (Extension)	D'olivo and Mcculloch (2017)	D'olivo et al. (2015); D'olivo et al. (2018)
DO17HAV02	HAV09_3 (Path B)	-18.836	146.5259	Havannah Island	B/Ca; $\delta^{11}\text{B}$ ; Li/Ca; Li/Mg; Mg/Ca; Sr/Ca; U/Ca (Extension)	D'olivo and Mcculloch (2017)	D'olivo et al. (2015); D'olivo et al. (2018)
DO17HAV03	HAV09_3 (Path C)	-18.836	146.5259	Havannah Island	B/Ca; $\delta^{11}\text{B}$ ; Li/Ca; Li/Mg; Mg/Ca; Sr/Ca; U/Ca (Extension)	D'olivo and Mcculloch (2017)	D'olivo et al. (2015); D'olivo et al. (2018)
DO17HAV04	HAV09_3 (Path D)	-18.836	146.5259	Havannah Island	B/Ca; $\delta^{11}\text{B}$ ; Li/Ca; Li/Mg; Mg/Ca; Sr/Ca; U/Ca (Extension)	D'olivo and Mcculloch (2017)	D'olivo et al. (2015); D'olivo et al. (2018)
DO18DAV01	DAV-1 (DAV09_2)	-18.83	147.63	Davies Reef	B/Ca; Li/Ca; Li/Mg; Mg/Ca; Sr/Ca; U/Ca	D'olivo et al. (2018)	

<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
DO18DAV02	DAV-2 (DAV13_2)	-18.8	147.63	Davies Reef	B/Ca; Li/Ca; Li/Mg; Mg/Ca; Sr/Ca; U/Ca	D'olivo et al. (2018)	Mcculloch et al. (2017); Thompson et al. (2022)
DO18DAV03	DAV-3 (DAV13_3)	-18.8	147.63	Davies Reef	B/Ca; Li/Ca; Li/Mg; Mg/Ca; Sr/Ca; U/Ca	D'olivo et al. (2018)	Mcculloch et al. (2017); Thompson et al. (2022)
DO18FIT01	FITZ (FZ04B)	-16.923	145.993	Fitzroy Island	B/Ca; Li/Ca; Li/Mg; Mg/Ca; Sr/Ca; U/Ca	D'olivo et al. (2018)	D'olivo and Mcculloch (2022)
DO18GBR01	CORE-17 (1709_6)	-17.876	146.729	Reef 17-065	B/Ca; Li/Ca; Li/Mg; Mg/Ca; Sr/Ca; U/Ca	D'olivo et al. (2018)	
DO18HAV01	HAV (HAV09_3)	-18.836	146.5259	Havannah Island	B/Ca; Li/Ca; Li/Mg; Mg/Ca; Sr/Ca; U/Ca	D'olivo et al. (2018)	D'olivo and Mcculloch (2017)
DO18LIZ01	LIZ (LIZ13_1)	-14.69	145.44	Lizard Island	B/Ca; Li/Ca; Li/Mg; Mg/Ca; Sr/Ca; U/Ca	D'olivo et al. (2018)	D'olivo et al. (2015)
DO18MYR01	MYR-1 (MYR5_271)	-18.27	147.38	Myrmidon Reef	B/Ca; Li/Ca; Li/Mg; Mg/Ca; Sr/Ca; U/Ca	D'olivo et al. (2018)	
DO18MYR02	MYR-2 (MYR13_2)	-18.27	147.38	Myrmidon Reef	B/Ca; Li/Ca; Li/Mg; Mg/Ca; Sr/Ca; U/Ca	D'olivo et al. (2018)	
DO18MYR03	MYR-3 (MYR13_5)	-18.27	147.38	Myrmidon Reef	B/Ca; Li/Ca; Li/Mg; Mg/Ca; Sr/Ca; U/Ca	D'olivo et al. (2018)	
DO18MYR04	MYR-4 (MYR13_7)	-18.27	147.38	Myrmidon Reef	B/Ca; Li/Ca; Li/Mg; Mg/Ca; Sr/Ca; U/Ca	D'olivo et al. (2018)	
DO18RIB01	RIB (RIB09_3)	-18.48	146.87	Rib Reef	Ba/Ca; B/Ca; B/Mg; Li/Ca; Li/Mg; Mg/Ca; Sr/Ca; U/Ca	D'olivo et al. (2018)	D'olivo et al. (2015)
DO22ARL01	AER04-1	-16.68	146.03	Arlington Reef	Ba <sup>138</sup> /Ca; Ba/Ca; Sr/Ca; U/Ca	D'olivo and Mcculloch (2022)	
DO22BRT01	BRT01	-18.14	146.73	Britomart Reef	Ba/Ca; Sr/Ca; U/Ca	D'olivo and Mcculloch (2022)	
DO22FIT01	FZ04B-1	-16.93	145.99	Fitzroy Island	Ba <sup>138</sup> /Ca; Ba/Ca; Sr/Ca; U/Ca	D'olivo and Mcculloch (2022)	D'olivo et al. (2018)

<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
DO22FIT02	FZ04-2	-16.93	145.99	Fitzroy Island	Ba <sup>138</sup> /Ca; Ba/Ca; Sr/Ca; U/Ca	D'olivo and Mcculloch (2022)	
DO22HIG01	HI03-1	-17.17	146	High Island	Ba <sup>138</sup> /Ca; Ba/Ca; Sr/Ca; U/Ca	D'olivo and Mcculloch (2022)	
DO22HMP01	HMP01B	-23.2	151	Humpy Island	Ba/Ca; Sr/Ca; U/Ca	D'olivo and Mcculloch (2022)	Lough (2007); Lough (2011a); Lough (2011b);
DO22MAG01	MAG01D	-19.15	146.87	Geoffrey Bay, Magnetic Island	Ba <sup>138</sup> /Ca; Ba/Ca; Sr/Ca; U/Ca	D'olivo and Mcculloch (2022)	Barnes and Lough (1992); De'ath et al. (2009); Erler et al. (2016); Lewis (2005); Lewis et al. (2007); Lough (2007); Lough (2011a); Lough (2011b); Lough and Barnes (1997)
DO22OYS01	OY04-3	-16.62	145.91	Oyster Reef	Ba <sup>138</sup> /Ca; Ba/Ca; Sr/Ca; U/Ca	D'olivo and Mcculloch (2022)	
DO22PAN01	PAN08-1	-18.82	146.43	Pandora Reef	Ba <sup>138</sup> /Ca; Ba/Ca; Sr/Ca; U/Ca	D'olivo and Mcculloch (2022)	D'olivo et al. (2013)
DO22PAN02	PAN08-2	-18.82	146.43	Pandora Reef	Ba <sup>138</sup> /Ca; Ba/Ca; Sr/Ca; U/Ca	D'olivo and Mcculloch (2022)	D'olivo et al. (2013)
DR95LMI01	Lady Musgrave	-23.9	152.38	Lady Musgrave Island	$\Delta 14C$	Druffel and Griffin (1995)	
DR99ABR01_1	Abraham-1 (99aust02a)	-22.1	152.48	Abraham Reef	$\delta^{13}C$ ; $\delta^{18}O$	Druffel and Griffin (1999); Druffel and Griffin (1995); Druffel and Griffin (1993)	Linsley et al. (2019)
DR99ABR01_2	Abraham-1 (99aust02a)	-22.1	152.48	Abraham Reef	$\Delta 14C$	Druffel and Griffin (1999); Druffel and Griffin (1995); Druffel and Griffin (1993)	Linsley et al. (2019)

<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
DR99ABR02	Abraham-2 (99aust02b)	-22.1	152.48	Abraham Reef	$\Delta^{14}\text{C}$	Druffel and Griffin (1999); Druffel and Griffin (1995); Druffel and Griffin (1993)	
DR99HER01	Heron (99aust03a)	-23.44	151.98	Heron Island	$\Delta^{14}\text{C}$	Druffel and Griffin (1999); Druffel and Griffin (1995)	
EL19OTI01	PET12	-23.4873	152.0814	Pete's Bay, One Tree Island	$\text{Ba}^{138}/\text{Ca}$ ; $\text{Ba}/\text{Ca}$ ; $\text{B}/\text{Ca}$ ; $\text{Cd}/\text{Ca}$ ; $\text{Fe}/\text{Ca}$ ; $\text{Li}/\text{Ca}$ ; $\text{Mg}/\text{Ca}$ ; $\text{Mn}/\text{Ca}$ ; $\text{Pb}/\text{Ca}$ ; $\text{P}/\text{Ca}$ ; $\text{Sr}/\text{Ca}$ ; $\text{U}/\text{Ca}$ ; $\text{Y}/\text{Ca}$	Ellis et al. (2019)	
ER16MAG01	MAG01D	-19.15	146.87	Geoffrey Bay, Magnetic Island	$\delta^{15}\text{N}$ ; $\delta^{18}\text{O}$ (Calcification; Density; Extension)	Erler et al. (2016); Erler et al. (2020)	Barnes and Lough (1992); D'olivo and Mcculloch (2022); De'ath et al. (2009); Lewis (2005); Lewis et al. (2007); Lough (2011a); Lough (2011b); Lough and Barnes (1997)

Dataset ID	Core Name	Latitude	Longitude	Location	Primary Variables (Secondary Variables)	Record Publication(s)	Additional Core Publication(s)
ER20HAV01	HAV01A	-18.837	146.548	Havannah Island	$\delta^{15}\text{N}$	Erler et al. (2020)	D'olivo and Mcculloch (2022); De'ath et al. (2009); Hendy et al. (2002); Hendy et al. (2003a); Hendy et al. (2003b); Hendy et al. (2007); Hendy et al. (2012); Isdale et al. (1998); Lough (2007); Lough (2011a); Lough (2011b); Lough and Barnes (1997); Lough et al. (2015); Mcculloch et al. (2003); Palmer et al. (2015)
ER20HAV02	Hav33a (Hav33b)	-18.837	146.548	Havannah Island	$\delta^{15}\text{N}$	Erler et al. (2020)	
ER20PAN01	Pan04b	-18.815	146.436	Pandora Reef	$\delta^{15}\text{N}$	Erler et al. (2020)	De'ath et al. (2009); Lough (2007); Lough and Barnes (1997)
ER20PAN02	Pan22b	-18.815	146.436	Pandora Reef	$\delta^{15}\text{N}$	Erler et al. (2020)	Cantin and Lough (2014); De'ath et al. (2009)
FA00PAN01	Pandora 1-98 (b)	-18.8	146.4	Pandora Reef	Ba/Ca; B/Ca; Mg/Ca; Sr/Ca; U/Ca (Distance)	Fallon (2000)	
FA03DAV01	Davies 2	-18.8	147.7	Davies Reef	Ba/Ca; B/Ca; Mg/Ca; Mn; Sr/Ca; U/Ca (Distance)	Fallon et al. (2003); Fallon (2000)	Alibert and Mcculloch (1997); Alibert et al. (2003)

<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
FA03DAV02	Davies 8	-18.8	147.7	Davies Reef	Ba/Ca; B/Ca; Mg/Ca; Mn; Sr/Ca; U/Ca (Distance)	Fallon et al. (2003); Fallon (2000)	Alibert and Mcculloch (1997)
FA03HAV01	Havannah Island	-18.843	146.537	Havannah Island	Ba/Ca; B/Ca; Mg/Ca; Mn; Sr/Ca; U/Ca (Distance)	Fallon et al. (2003); Fallon (2000)	
FA03MYR01	Myrmidon 2 (Myr-2)	-18.266	147.383	Myrmidon Reef	Ba/Ca; B/Ca; Mg/Ca; Mn; Sr/Ca; U/Ca (Distance)	Fallon et al. (2003); Fallon (2000)	Marshall and Mcculloch (2002)
FA03ORP01	Orpheus Island	-18.634	146.495	Orpheus Island	Ba/Ca; B/Ca; Mg/Ca; Mn; Sr/Ca; U/Ca (Distance)	Fallon et al. (2003); Fallon (2000)	
FA03PAN01	Pandora 1-98 (a)	-18.8	146.4	Pandora Reef	Ba/Ca; B/Ca; Mg/Ca; Mn; Sr/Ca; U/Ca (Distance)	Fallon et al. (2003); Fallon (2000)	
FA03WHE01	Wheeler Reef	-18.799	147.528	Wheeler Reef	Ba/Ca; B/Ca; Mg/Ca; Sr/Ca; U/Ca (Distance)	Fallon et al. (2003); Fallon (2000)	Alibert and Mcculloch (1997)
GA98ORP01	OR-1	-18.567	146.483	Iris Point, Orpheus Island	$\delta^{18}\text{O}$ ; Sr/Ca ( $\delta^{18}\text{O}_{\text{seawater}}$ )	Gagan et al. (1998); Gagan et al. (2002); Gagan et al. (1994)	Gagan et al. (2012)
HE02GBR01	NA	-17.78	146.13	Central GBR	$\delta^{18}\text{O}$ ; Sr/Ca; U/Ca ( $\delta^{18}\text{O}_{\text{seawater}}$ )	Hendy et al. (2002); Hendy et al. (2007)	
HE03GBR01	NA	-17.78	146.13	Central GBR	Luminescence visual indices	Hendy et al. (2003a)	

<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
HE03GPI01	GPI-02A-01 (GPI02A)	-18.733	146.567	Great Palm Island	Luminescence visual indices	Hendy et al. (2003a); Hendy et al. (2007); Leonard et al. (2016)	De'ath et al. (2009); Hendy et al. (2002); Hendy et al. (2003b); Leonard et al. (2016); Lough (2007); Lough (2011a); Lough (2011b); Lough and Barnes (1997)
IS98HAV02	Havannah (HAV01A)	-18.85	146.55	Havannah Island	Luminescence Sum	Isdale et al. (1998)	D'olivo and Mcculloch (2022); De'ath et al. (2009); Erler et al. (2020); Hendy et al. (2002); Hendy et al. (2003a); Hendy et al. (2003b); Hendy et al. (2007); Hendy et al. (2012); Lough (2007); Lough (2011a); Lough (2011b); Lough and Barnes (1997); Lough et al. (2015); Mcculloch et al. (2003); Palmer et al. (2015)
IS98PAN01	Pandora (PAN08B)	-18.82	146.43	Pandora Reef	Luminescence Sum	Isdale et al. (1998)	Hendy et al. (2002); Hendy et al. (2003a); Hendy et al. (2003b); Hendy et al. (2007); Hendy et al. (2012)
JU08KES01a	KIB	-20.9177	149.4178	Keswick Island	Luminescence	Jupiter et al. (2008)	

<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
JU08KES01b	KIA	-20.9177	149.4186	Keswick Island	Ce; Dy; Er; Eu; Gd; Ho; La; Lu; Nd; Pr; Sm; Tb; Tm; Y; Yb; Y/Ho	Jupiter (2008); Jupiter (2006)	
JU08KES02	KIC	-20.9177	149.4178	Keswick Island	Luminescence	Jupiter et al. (2008)	Marion et al. (2021)
JU08KES03	KIE	-20.9319	149.4267	Keswick Island	Ba/Ca; Luminescence	Jupiter et al. (2008)	
JU08RTI01a	RTC	-21.1755	149.2639	Round Top Island	Luminescence	Jupiter et al. (2008)	Marion et al. (2021)
JU08RTI01b	RTF	-21.1708	149.2644	Round Top Island	Ce; Dy; Er; Eu; Gd; Ho; La; Lu; Nd; Pr; Sm; Tb; Tm; Y; Yb; Y/Ho	Jupiter (2008); Jupiter (2006)	Marion et al. (2021)
JU08RTI02	RTH	-21.172	149.2639	Round Top Island	Ba/Ca; Luminescence	Jupiter et al. (2008)	Marion et al. (2021)
JU08RTI03	RTI	-21.1717	149.2634	Round Top Island	Luminescence	Jupiter et al. (2008)	
JU08SCA01	SCA	-20.8522	149.6021	Scawfell Island	Luminescence	Jupiter et al. (2008)	
JU08SCA02	SCB	-20.8522	149.6021	Scawfell Island	Luminescence	Jupiter et al. (2008)	
JU08SCA03	SCC	-20.8522	149.6021	Scawfell Island	Ba/Ca; Luminescence	Jupiter et al. (2008)	Marion et al. (2021)
LE05MAG01	MAG01D	-19.15	146.858	Geoffrey Bay, Magnetic Island	$\delta^{13}\text{C}$ ; $\delta^{18}\text{O}$ ( $\delta^{18}\text{O}_{\text{seawater}}$ )	Lewis (2005)	Barnes and Lough (1992); D'olivo and Mcculloch (2022); De'ath et al. (2009); Erler et al. (2016); Lewis et al. (2007); Lough (2007); Lough (2011a); Lough (2011b); Lough and Barnes (1997)
LE05NEL01	NEL01D	-19	147	Nelly Bay, Magnetic Island	$\delta^{13}\text{C}$ ; $\delta^{18}\text{O}$ ( $\delta^{18}\text{O}_{\text{seawater}}$ )	Lewis (2005)	Lewis et al. (2007); Lough et al. (2014)



<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
LE05NEL02	NEL03D	-19	147	Nelly Bay, Magnetic Island	$\delta^{13}\text{C}$ ; $\delta^{18}\text{O}$ ( $\delta^{18}\text{O}_{\text{seawater}}$ )	Lewis (2005)	Lough et al. (2014)
LE07MAG01	MAG01D	-19.15	146.87	Geoffrey Bay, Magnetic Island	Ba; Ba/Ca; /Ca; Mg/Ca; Mn; Pr; Sm; Sr/Ca; Th; U/Ca; Y; Y/Ho (Distance)	Lewis et al. (2007); Lewis (2005)	Barnes and Lough (1992); D'olivo and Mcculloch (2022); De'ath et al. (2009); Erler et al. (2016); Lewis (2005); Lough (2007); Lough (2011a); Lough (2011b); Lough and Barnes (1997)
LE07NEL01	NEL01D	-19	147	Nelly Bay, Magnetic Island	Ba; Ba/Ca; /Ca; Mg/Ca; Mn; Pr; Sm; Sr/Ca; Th; U/Ca; Y; Y/Ho (Distance)	Lewis et al. (2007); Lewis (2005)	Lough et al. (2014)
LE12CID01	CID-01A	-20.3	148.9	Cid Harbour Island	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Ca/Ca; Cd; Cd/Ca; Li/Ca; Mg/Ca; Mn; Mn/Ca; P; Pb/Ca; P/Ca; Sr/Ca; U/Ca; Y; Y/Ca (Distance)	Lewis et al. (2012)	De'ath et al. (2009); Lough (2007); Lough (2011a); Lough (2011b); Lough and Barnes (1997)
LE12CID02	CID-71B	-20.254	148.923	Cid Harbour Island	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Ca/Ca; Cd; Cd/Ca; Li/Ca; Mg/Ca; Mn; Mn/Ca; P; Pb/Ca; P/Ca; Sr/Ca; U/Ca; Y; Y/Ca (Distance)	Lewis et al. (2012)	

<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
LE12CID03	CID-73B	-20.253	148.92	Cid Harbour Island	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Cd; Cd/Ca; Ce; Ce/Ca; Eu; Eu/Ca; Gd; Gd/Ca; La; La/Ca; Li/Ca; Lu; Lu/Ca; Mg/Ca; Mn; Mn/Ca; P; Pb/Ca; P/Ca; Pr; Pr/Ca; Sm; Sm/Ca; Sr/Ca; U/Ca; Y; Yb; Yb/Ca; Y/Ca; Zn/Ca	Lewis et al. (2012)	
LE12COB01	COB-71A	-19.396	148.806	Cobham Reef	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Cd/Ca; Li/Ca; Mg/Ca; Mn/Ca; Pb/Ca; P/Ca; Sr/Ca; U/Ca; Y/Ca (Distance)	Lewis et al. (2012)	
LE12HAS01	HWD-73B	-20.26	149.096	Haslewood Island	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Cd; Cd/Ca; Ce; Ce/Ca; Eu; Eu/Ca; Gd; Gd/Ca; La; La/Ca; Li/Ca; Lu; Lu/Ca; Mg/Ca; Mn; Mn/Ca; P; Pb/Ca; P/Ca; Pr; Pr/Ca; Sm; Sm/Ca; Sr/Ca; U/Ca; Y; Yb; Yb/Ca; Y/Ca; Zn/Ca (Distance)	Lewis et al. (2012)	

<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
LE12HKO01	SNH-73A	-20.089	148.907	Stonehaven Bay, Hook Island	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Cd; Cd/Ca; Ce; Ce/Ca; Eu; Eu/Ca; Gd; Gd/Ca; La; La/Ca; Li/Ca; Lu; Lu/Ca; Mg/Ca; Mn; Mn/Ca; P; Pb/Ca; P/Ca; Pr; Pr/Ca; Sm; Sm/Ca; Sr/Ca; U/Ca; Y; Yb; Yb/Ca; Y/Ca; Zn/Ca (Distance)	Lewis et al. (2012)	
LE12NMI01	NMI-73A	-20.22	148.811	North Molle Island	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Cd; Cd/Ca; Ce; Ce/Ca; Eu; Eu/Ca; Gd; Gd/Ca; La; La/Ca; Li/Ca; Lu; Lu/Ca; Mg/Ca; Mn; Mn/Ca; P; Pb/Ca; P/Ca; Pr; Pr/Ca; Sm; Sm/Ca; Sr/Ca; U/Ca; Y; Yb; Yb/Ca; Y/Ca; Zn/Ca (Distance)	Lewis et al. (2012)	
LE12REP01	REP-71A	-20.618	148.865	Repulse Island	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Cd; Cd/Ca; Ce; Ce/Ca; La; La/Ca; Li/Ca; Lu; Lu/Ca; Mg/Ca; Mn; Mn/Ca; P; Pb/Ca; P/Ca; Pr; Pr/Ca; Sr/Ca; U/Ca; Y; Yb; Yb/Ca; Y/Ca; Zn/Ca (Distance)	Lewis et al. (2012)	

<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
LE12REP02	REP-72A	-20.617	148.864	Repulse Island	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Cd; Cd/Ca; Ce; Ce/Ca; La; La/Ca; Li/Ca; Lu; Lu/Ca; Mg/Ca; Mn; Mn/Ca; P; Pb/Ca; P/Ca; Pr; Pr/Ca; Sr/Ca; U/Ca; Y; Yb; Yb/Ca; Y/Ca; Zn/Ca (Distance)	Lewis et al. (2012)	
LE12REP03	REP-72B	-20.617	148.864	Repulse Island	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Ca/Ca; Cd; Cd/Ca; Li/Ca; Mg/Ca; Mn; Mn/Ca; P; Pb/Ca; P/Ca; Sr/Ca; U/Ca; Y; Y/Ca (Distance)	Lewis et al. (2012)	
LE12SHW01	SHW-82A	-20.48	149.069	Shaw Island	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Cd; Cd/Ca; Li/Ca; Mg/Ca; Mn; Mn/Ca; P; Pb/Ca; P/Ca; Sr/Ca; U/Ca; Y; Y/Ca (Distance)	Lewis et al. (2012)	
LE12SHW02	SHW-83B	-20.48	149.069	Shaw Island	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Cd; Cd/Ca; Li/Ca; Mg/Ca; Mn; Mn/Ca; P; Pb/Ca; P/Ca; Sr/Ca; U/Ca; Y; Y/Ca (Distance)	Lewis et al. (2012)	
LE12WHI01	WHI-34A	-20.169	148.957	Whitsunday Island	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Cd; Cd/Ca; Ce; Ce/Ca; La; La/Ca; Li/Ca; Mg/Ca; Mn; Mn/Ca; P; Pb/Ca; P/Ca; Sr/Ca; U/Ca; Y; Y/Ca (Distance)	Lewis et al. (2012)	De'ath et al. (2009)

<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
LE16GPI01	PAM2.0	-18.733	146.567	Great Palm Island	Luminescence visual indices	Leonard et al. (2016)	
LE16GPI02	PAM3.1	-18.733	146.567	Great Palm Island	Luminescence visual indices	Leonard et al. (2016)	
LE16GPI03	PAM5.0	-18.733	146.567	Great Palm Island	Luminescence visual indices	Leonard et al. (2016)	
LE18GFB01	GFB33B	-19.154	146.866	Geoffrey Bay, Magnetic Island	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Cd/Ca; Li/Ca; Mg/Ca; Mn/Ca; Pb/Ca; P/Ca; Sr/Ca; U/Ca; Y/Ca (Distance)	Lewis et al. (2018); D'olivo and Mcculloch (2022)	Saha et al. (2019b); Saha et al. (2021)
LE18GFB02	GFB34A	-19.164	146.853	Geoffrey Bay, Magnetic Island	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Cd/Ca; Li/Ca; Mg/Ca; Mn/Ca; Pb/Ca; P/Ca; Sr/Ca; U/Ca; Y/Ca (Distance)	Lewis et al. (2018); D'olivo and Mcculloch (2022)	
LE18HAV01	HAV32A	-18.835	146.544	Havannah Island	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Cd/Ca; Li/Ca; Mg/Ca; Mn/Ca; Pb/Ca; P/Ca; Sr/Ca; U/Ca; Y/Ca (Distance)	Lewis et al. (2018); D'olivo and Mcculloch (2022)	Lough et al. (2015)
LE18HAV02	HAV34A	-18.837	146.548	Havannah Island	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Cd/Ca; Li/Ca; Mg/Ca; Mn/Ca; Pb/Ca; P/Ca; Sr/Ca; U/Ca; Y/Ca (Distance)	Lewis et al. (2018); D'olivo and Mcculloch (2022)	Lough et al. (2015)
LE18NEL01	NEL29A	-19.165	146.854	Nelly Bay, Magnetic Island	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Cd/Ca; Li/Ca; Mg/Ca; Mn/Ca; Pb/Ca; P/Ca; Sr/Ca; U/Ca; Y/Ca (Distance)	Lewis et al. (2018); D'olivo and Mcculloch (2022)	Cantin and Lough (2014); De'ath et al. (2009); Lough et al. (2014)

<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
LE18NEL02	NEL39A	-19.169	146.85	Nelly Bay, Magnetic Island	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Cd/Ca; Li/Ca; Mg/Ca; Mn/Ca; Pb/Ca; P/Ca; Sr/Ca; U/Ca; Y/Ca (Distance)	Lewis et al. (2018); D'olivo and Mcculloch (2022)	Cantin and Lough (2014); De'ath et al. (2009)
LE18PAN01	PAN31B	-18.813	146.427	Pandora Reef	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Cd/Ca; Li/Ca; Mg/Ca; Mn/Ca; Pb/Ca; P/Ca; Sr/Ca; U/Ca; Y/Ca (Distance)	Lewis et al. (2018); D'olivo and Mcculloch (2022)	Cantin and Lough (2014); De'ath et al. (2009)
LE18PAN02	PAN36B	-18.813	146.427	Pandora Reef	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Cd/Ca; Li/Ca; Mg/Ca; Mn/Ca; Pb/Ca; P/Ca; Sr/Ca; U/Ca; Y/Ca (Distance)	Lewis et al. (2018); D'olivo and Mcculloch (2022)	Cantin and Lough (2014); De'ath et al. (2009)
LE18PEL01	PEL30A	-18.538	146.49	Pelorus Island	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Cd/Ca; Li/Ca; Mg/Ca; Mn/Ca; Pb/Ca; P/Ca; Sr/Ca; U/Ca; Y/Ca (Distance)	Lewis et al. (2018)	
LE18PEL02	PEL33B	-18.538	146.491	Pelorus Island	Ba <sup>138</sup> /Ca; Ba/Ca; B/Ca; Cd/Ca; Li/Ca; Mg/Ca; Mn/Ca; Pb/Ca; P/Ca; Sr/Ca; U/Ca; Y/Ca (Distance)	Lewis et al. (2018)	
LE19HIG01	HI 12.1	-17.161	146.007	High Island	Ba/Ca; Ce; Dy; Er; Eu; Gd; Ho; La; Lu; Nd; Pr; Sm; Tb; Tm; Y; Yb; Zr	Leonard et al. (2019)	

<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
LE19RUS01	FRI 12.1	-17.224	146.09	Russell Island	Ba/Ca; Ce; Dy; Er; Eu; Gd; Ho; La; Lu; Nd; Pr; Sm; Tb; Tm; Y; Yb; Zr	Leonard et al. (2019)	
LE19RUS02	FRI 12.3	-17.228	146.091	Russell Island	Ba/Ca; Ce; Dy; Er; Eu; Gd; Ho; La; Lu; Nd; Pr; Sm; Tb; Tm; Y; Yb; Zr	Leonard et al. (2019)	
LE19SUD01	SUD 12.1	-17.026	146.211	Sudbury Cay	Ba/Ca; Ce; Dy; Er; Eu; Gd; Ho; La; Lu; Nd; Pr; Sm; Tb; Tm; Y; Yb; Zr	Leonard et al. (2019)	
LO11ACH01	ACH01A	-18.95	146.65	Acheron Island	Luminescence Range	Lough (2011a); Lough (2011b)	Barnes and Lough (1992)
LO11BRO01	BRO01A	-18.15	146.28	Brook Island	Luminescence Range	Lough (2011a); Lough (2011b); D'olivo and Mcculloch (2022)	Barnes and Lough (1992); De'ath et al. (2009); Hendy et al. (2002); Hendy et al. (2003a); Hendy et al. (2003b); Hendy et al. (2007); Lough (2012); Lough (2007); Lough and Barnes (1997)
LO11CID01	CID01A	-20.27	148.93	Cid Harbour Island	Luminescence Range	Lough (2011a); Lough (2011b)	De'ath et al. (2009); Lewis et al. (2012); Lough (2007); Lough and Barnes (1997)
LO11CNR01	CNR01B	-15.13	145.32	Conical Rock Reef	Luminescence Range	Lough (2011a)	De'ath et al. (2009); Lough and Barnes (1997)

<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
LO11COO01	COO01E	-18.03	146.18	Coombe Island	Luminescence Range	Lough (2011a); Lough (2011b)	Barnes and Lough (1992); De'ath et al. (2009); Lough (2007); Lough and Barnes (1997)
LO11DUN01	DUN02A	-17.95	146.17	Dunk Island	Luminescence Range	Lough (2011a); Lough (2011b)	
LO11GPI01	GPI02A	-18.68	146.58	Great Palm Island	Luminescence Range	Lough (2011a); Lough (2011b)	De'ath et al. (2009); Hendy et al. (2002); Hendy et al. (2003a); Hendy et al. (2003b); Hendy et al. (2007); Lough (2007); Lough and Barnes (1997)
LO11HAV01	HAV01A	-18.85	146.55	Havannah Island	Luminescence Range	Lough (2011a); Lough (2011b); Palmer et al. (2015)	D'olivo and Mcculloch (2022); De'ath et al. (2009); Erler et al. (2020); Hendy et al. (2002); Hendy et al. (2003a); Hendy et al. (2003b); Hendy et al. (2007); Hendy et al. (2012); Isdale et al. (1998); Lough (2007); Lough and Barnes (1997); Lough et al. (2015); Mcculloch et al. (2003)



<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
LO11HKO01	HKO01B	-20.07	148.95	Hook Island	Luminescence Range	Lough (2011a); Lough (2011b)	Barnes and Lough (1992); De'ath et al. (2009); Lough (2007); Lough and Barnes (1997)
LO11HKO02	SNH01A	-20.1	148.9	Stonehaven Bay, Hook Island	Luminescence Range	Lough (2011a); Lough (2011b)	De'ath et al. (2009); Lough and Barnes (1997)
LO11HMP01	HMP01B	-23.2	151	Humpy Island	Luminescence Range	Lough (2011a); Lough (2011b); D'olivo and Mcculloch (2022)	Lough (2007);
LO11LUP01	LUP01C	-20.28	149.12	Lupton Island	Luminescence Range	Lough (2011a); Lough (2011b)	Barnes and Lough (1992); De'ath et al. (2009); Lough (2007); Lough and Barnes (1997)
LO11MAG01	MAG01D	-19.15	146.87	Geoffrey Bay, Magnetic Island	Luminescence Range	Lough (2011a); Lough (2011b); D'olivo and Mcculloch (2022)	Barnes and Lough (1992); De'ath et al. (2009); Erler et al. (2016); Lewis (2005); Lewis et al. (2007); Lough (2007); Lough and Barnes (1997)
LO11NMI01	NMI01B	-20.23	148.8	North Molle Island	Luminescence Range	Lough (2011a); Lough (2011b)	Barnes and Lough (1992); De'ath et al. (2009); Lough (2007); Lough and Barnes (1997)

<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
LO11NOR01	NOR01B	-17.2	146.1	Normanby Island	Luminescence Range	Lough (2011a); Lough (2011b); D'olivo and Mcculloch (2022)	Barnes and Lough (1992); De'ath et al. (2009); Lough (2007)
LO11PAN01	PAN07B	-18.82	146.43	Pandora Reef	Luminescence Range	Lough (2011a)	Barnes and Lough (1992); De'ath et al. (2009); Lough (2007)
LO11SMI01	SMI01C	-20.27	148.83	South Molle Island	Luminescence Range	Lough (2011a); Lough (2011b)	Barnes and Lough (1992); De'ath et al. (2009); Lough (2007); Lough and Barnes (1997)
LO14NEL01	NEL03A	-19	147	Nelly Bay, Magnetic Island	Luminescence Range	Lough et al. (2014)	
LO14NEL02	NEL03D	-19	147	Nelly Bay, Magnetic Island	Luminescence Range	Lough et al. (2014)	Lewis (2005)
LO14NEL03	NEL07B	-19	147	Nelly Bay, Magnetic Island	Luminescence Range	Lough et al. (2014)	
LO14NEL04	NEL07C	-19	147	Nelly Bay, Magnetic Island	Luminescence Range	Lough et al. (2014)	
LO14NEL05	NEL01D	-19	147	Nelly Bay, Magnetic Island	Luminescence Range	Lough et al. (2014)	Lewis (2005); Lewis et al. (2007)
LO14NEL06	NEL29A	-19.1646	146.8658	Nelly Bay, Magnetic Island	Luminescence Range	Lough et al. (2014)	Cantin and Lough (2014); De'ath et al. (2009); Lewis et al. (2018)
LO14NEL07	NEL35A	-19.164	146.854	Nelly Bay, Magnetic Island	Luminescence Range	Lough et al. (2014)	Cantin and Lough (2014); De'ath et al. (2009)
LO14NEL08	NEL39B	-19.169	146.8502	Nelly Bay, Magnetic Island	Luminescence Range	Lough et al. (2014)	Cantin and Lough (2014); De'ath et al. (2009)

<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
LO15HAV01	HAV01A	-18.85	146.55	Havannah Island	Luminescence Range	Lough et al. (2015)	D'olivo and Mcculloch (2022); De'ath et al. (2009); Erler et al. (2020); Hendy et al. (2002); Hendy et al. (2003a); Hendy et al. (2003b); Hendy et al. (2007); Hendy et al. (2012); Isdale et al. (1998); Lough (2007); Lough (2011a); Lough (2011b); Lough and Barnes (1997); Mcculloch et al. (2003); Palmer et al. (2015)
LO15HAV02	HAV31B	-18.837	146.5483	Havannah Island	Luminescence Range	Lough et al. (2015)	
LO15HAV03	HAV32A	-18.8351	146.544	Havannah Island	Luminescence Range	Lough et al. (2015)	Lewis et al. (2018)
LO15HAV04	HAV33A	-18.837	146.5483	Havannah Island	Luminescence Range	Lough et al. (2015)	
LO15HAV05	HAV34A	-18.837	146.5483	Havannah Island	Luminescence Range	Lough et al. (2015)	Lewis et al. (2018)
MA00MYR01	Myr-2	-18.266	147.383	Myrmidon Reef	Sr/Ca (Distance)	Marshall (2000); Marshall and Mcculloch (2002)	Fallon (2000); Fallon et al. (2003)
MA00MYR02	Myr2-16D	-18.266	147.383	Myrmidon Reef	$\delta^{13}\text{C}$ ; $\delta^{18}\text{O}$ (Distance)	Marshall (2000)	
MA00MYR03	Myr2-17D	-18.266	147.383	Myrmidon Reef	Ba/Ca; Mg/Ca; Sr/Ca; U/Ca (Distance)	Marshall (2000)	
MA00SLY01	St-1	-19.2892	148.0519	Stanley Reef	Sr/Ca (Distance)	Marshall (2000); Marshall and Mcculloch (2002)	

<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
MC03HAV01	Havannah	-18.843	146.537	Havannah Island	Ba/Ca	Mcculloch et al. (2003); Lewis et al. (2007); Walther et al. (2013)	
MC03PAN01	Pandora	-18.8	146.4	Pandora Reef	Ba/Ca	Mcculloch et al. (2003); D'olivo and Mcculloch (2022)	
MC17DAV01	Davies 13-2 (D-2)	-18.8	147.63	Davies Reef	Ba/Ca; B/Ca; B/Mg; $\delta^{11}\text{B}$ ; Li/Ca; Li/Mg; Mg/Ca; Sr/Ca; U/B; U/Ca; U/Sr ( $\delta^{11}\text{B pH}_{\text{cf}}$ )	Mcculloch et al. (2017); Thompson et al. (2022)	D'olivo et al. (2018)
MC17DAV02	Davies 13-3 (D-3)	-18.8	147.63	Davies Reef	Ba/Ca; B/Ca; B/Mg; $\delta^{11}\text{B}$ ; Li/Ca; Li/Mg; Mg/Ca; Sr/Ca; U/B; U/Ca; U/Sr ( $\delta^{11}\text{B pH}_{\text{cf}}$ )	Mcculloch et al. (2017); Thompson et al. (2022)	D'olivo et al. (2018)
PE05FLI01	FL02A (FLI02A)	-17.73	148.43	Flinders Reef	$\delta^{11}\text{B}$ ; $\delta^{13}\text{C}$ ( <i>Qarag</i> ; Calcification; Extension; $\delta^{11}\text{B pH}$ )	Pelejero et al. (2005)	Calvo et al. (2007)
RA17MYR01	myra-30	-18.278	147.379	Myrmidon Reef	Ba/Ca; Mg/Ca; Sr/Ca; U/Ca; Y/Ca	Razak et al. (2017)	
RA17MYR02	myra-31	-18.278	147.379	Myrmidon Reef	Ba/Ca; Mg/Ca; Sr/Ca; U/Ca; Y/Ca	Razak et al. (2017)	
RE19CLK01	CLK (CLK06A)	-11.97	143.28	Clerke Reef	Luminescence; Sr/Ca (Density; Distance)	Reed et al. (2019)	
RE19EEL01	EEL (EELB10)	-12.5	143.52	Eel Reef	$\delta^{13}\text{C}$ ; $\delta^{18}\text{O}$ ; Luminescence; Sr/Ca (Density; Distance)	Reed et al. (2019)	Barnes and Lough (1992); Lough and Barnes (2000); Lough et al. (2002)

<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
RE19GBR01	13-050 (050B06)	-13.33	143.95	Reef 13-050	$\delta^{13}\text{C}$ ; $\delta^{18}\text{O}$ ; Luminescence; Sr/Ca (Density; Distance)	Reed et al. (2019)	Barnes and Lough (1992); Lough and Barnes (1992); De'ath et al. (2009); Lough et al. (2002); Madin et al. (2016)
RE19NOM01	NOM (NOM05A)	-12.09	143.29	Nomad Reef	Luminescence; Sr/Ca (Density; Distance)	Reed et al. (2019)	
RE19POR01	POR (PORB03)	-12.591	143.408	Portland Roads	$\delta^{13}\text{C}$ ; $\delta^{18}\text{O}$ ; Luminescence; Sr/Ca (Density; Distance)	Reed et al. (2019)	Barnes and Lough (1992); Lough and Barnes (1992); De'ath et al. (2009); Lough et al. (2002); Madin et al. (2016)
RO14GKI01	GK2	-23.1505	150.9739	Great Keppel Island	Luminescence G/B	Rodriguez-Ramirez et al. (2014); Grove et al. (2015)	Saha et al. (2018b); Saha et al. (2019a); Saha et al. (2021)
RO14GKI02	GK3	-23.192	150.9627	Great Keppel Island	Luminescence G/B	Rodriguez-Ramirez et al. (2014)	
RO14KIN01	KR-MMA-1	-17.767	146.133	King Reef	$\delta^{18}\text{O}$ ; Sr/Ca ( $\delta^{18}\text{O}_{\text{seawater}}$ )	Roche et al. (2014)	
RO14KIN02	KR-AMA-2	-17.767	146.133	King Reef	$\delta^{18}\text{O}$ ; Sr/Ca ( $\delta^{18}\text{O}_{\text{seawater}}$ )	Roche et al. (2014)	
RO14MIA01	MI1	-23.155	150.9035	Miall Island	Luminescence G/B	Rodriguez-Ramirez et al. (2014); Grove et al. (2015)	
RO14MIA02	MI2	-23.1554	150.9034	Miall Island	Luminescence G/B	Rodriguez-Ramirez et al. (2014)	
RO14SQR01	SQ1	-23.0997	150.8862	Square Rocks	Luminescence G/B	Rodriguez-Ramirez et al. (2014); Grove et al. (2015)	

<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
RO14SQR02	SQ2	-23.0997	150.8862	Square Rocks	Luminescence G/B	Rodriguez-Ramirez et al. (2014); Grove et al. (2015)	
SA16HER01a	AR HL 3	-23.451	151.93	Heron Island	Sr/Ca	Sadler et al. (2016a)	
SA16HER01b	P HS 1	-23.451	151.93	Heron Island	Sr/Ca	Sadler et al. (2016b)	
SA16HER02a	AR HL 4	-23.451	151.93	Heron Island	Sr/Ca	Sadler et al. (2016a)	
SA16HER02b	P HS 3	-23.451	151.93	Heron Island	Sr/Ca	Sadler et al. (2016b)	
SA16HER03a	AR HL 6	-23.451	151.93	Heron Island	Sr/Ca	Sadler et al. (2016a)	
SA16HER03b	P HS 4	-23.451	151.93	Heron Island	Sr/Ca	Sadler et al. (2016b)	
SA16HER04a	AR HS 3	-23.451	151.93	Heron Island	Sr/Ca	Sadler et al. (2016a)	
SA16HER04b	P HS 7	-23.451	151.93	Heron Island	Sr/Ca	Sadler et al. (2016b)	
SA16HER05a	AR HS 5	-23.451	151.93	Heron Island	Sr/Ca	Sadler et al. (2016a)	
SA16HER05b	R9C (track 1)	-23.451	151.93	Heron Island	Sr/Ca	Sadler et al. (2016b)	
SA16HER06a	AR HS 6	-23.451	151.93	Heron Island	Sr/Ca	Sadler et al. (2016a)	
SA16HER06b	R15B3	-23.451	151.93	Heron Island	Sr/Ca	Sadler et al. (2016b)	
SA21GFB01	GFB33B	-19.1543	146.8656	Geoffrey Bay, Magnetic Island	Ba/Ca; Ce; Dy; Eu; Gd; Ho; La; Lu; Mn/Ca; Nd; Nd/Yb; Pr; Sm; Sr/Ca; Tb; Tm; Y; Yb; Y/Ca (Ce/Ce*; ΣREE/Ca)	Saha et al. (2021); Saha et al. (2019b)	Lewis et al. (2018)
SA21GKI01	GK2	-23.17	150.98	Great Keppel Island	Ba/Ca; Ce; Dy; Eu; Gd; Ho; La; Lu; Mg/Ca; Mn/Ca; Nd; Pr; Sm; Sr/Ca; Tb; Tm; U/Ca; V/Ca; Y; Yb; Y/Ca	Saha et al. (2021); Saha et al. (2018b); Saha et al. (2019a)	Rodriguez-Ramirez et al. (2014)

<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
SA21RAT01	RI2	-23.7663	151.3178	Rat Island	Ba/Ca; Ce; Dy; Eu; Gd; Ho; La; Lu; Mn/Ca; Nd; Pr; Sm; Sr/Ca; Tb; Tm; Y; Yb; Y/Ca	Saha et al. (2021); Saha et al. (2018a)	
WA13HAV01	Havannah (Hav08_2)	-18.8396	146.5502	Havannah Island	Ba/Ca	Walther et al. (2013); D'olivo and Mcculloch (2022)	
WA13MYR01	Myrmidon (MYR_S5)	-18.2614	147.3765	Myrmidon Reef	Ba/Ca	Walther et al. (2013)	
WA19ARL01	10AR2	-16.6381	146.1036	Arlington Reef	$\delta^{98}\text{Mo}$ ; Fe; Mn; Ti	Wang et al. (2019); Deng et al. (2014)	Chen et al. (2021); Xiao et al. (2020)
WE09ARL01_1	AREO 4	-16.667	146.109	Arlington Reef	Ba/Ca; $\delta^{11}\text{B}$ ; $\delta^{13}\text{C}$ ; $\delta^{18}\text{O}$ ; Mg/Ca; Sr/Ca ( $\delta^{11}\text{B}$ pH)	Wei et al. (2009); D'olivo and Mcculloch (2017); Chen et al. (2021)	Wei et al. (2015)
WE09ARL01_2	AREO 4	-16.674	146.109	Arlington Reef	Ba/Ca; $\delta^{11}\text{B}$ ; $\delta^{13}\text{C}$ ; $\delta^{18}\text{O}$ ; Mg/Ca; Sr/Ca ( $\delta^{11}\text{B}$ pH)	Wei et al. (2009); D'olivo and Mcculloch (2017); Chen et al. (2021)	Wei et al. (2015)
WU21CUR01	SEN01C	-23.7547	151.3169	South End, Curtis Island	B/Ca; Li/Ca; Mg/Ca; Sr/Ca; U/Ca	Wu et al. (2021b)	Cantin et al. (2018)
WU21HUM01	HUM03A	-23.4035	151.1472	Hummocky Island	B/Ca; Li/Ca; Mg/Ca; Sr/Ca; U/Ca	Wu et al. (2021b)	Cantin et al. (2018)
WU21HUM02	HUM04B	-23.4024	151.146	Hummocky Island	B/Ca; Li/Ca; Mg/Ca; Sr/Ca; U/Ca	Wu et al. (2021b)	
WU21MAS01a	MAS02A	-23.532	151.7403	Mast Head Island	B/Ca; Li/Ca; Mg/Ca; Sr/Ca; U/Ca	Wu et al. (2021b)	Cantin et al. (2018)
WU21MAS01b	MAS01E	-23.5325	151.746	Mast Head Island	$\Delta^{14}\text{C}$	Wu et al. (2021a); Cantin et al. (2018)	
WU21SHW01	SHW82C	-20.4796	149.0691	Shaw Island	B/Ca; Li/Ca; Mg/Ca; Sr/Ca; U/Ca	Wu et al. (2021b)	

<b>Dataset ID</b>	<b>Core Name</b>	<b>Latitude</b>	<b>Longitude</b>	<b>Location</b>	<b>Primary Variables (Secondary Variables)</b>	<b>Record Publication(s)</b>	<b>Additional Core Publication(s)</b>
WU21SMI01	SMI81A	-20.2588	148.8283	South Molle Island	B/Ca; Li/Ca; Mg/Ca; Sr/Ca; U/Ca	Wu et al. (2021b)	
XI20ARL01	10AR2	-16.6381	146.1036	Arlington Reef	$\delta^{13}\text{C}$ ; $\delta^{18}\text{O}$ ; $\delta^{66}\text{Zn}$ ; Sr/Ca; Zn	Xiao et al. (2020);	Chen et al. (2021); Deng et al. (2014); Wang et al. (2019)

Note: Additional publications referenced in the GBRC D include:

- Chakraborty and Ramesh (1993, 1997) for CH93SLY01 & CH93SLY02.
- Davies and Hopley (1983) for MA00MYR02 & MA00MYR03.
- Kamber et al. (2005) for JU08KES01b & JU08RTI01b
- Min et al. (1995) for HE02GBR01, LE07MAG01 & LE07NEL01.

570



## Appendix B – GBRC D to LiPD Field Translation

Table B1. GBRC D to LiPD time series (TS) field translation dictionary

GBRC D Field	LiPD Field	Notes
Age	year	LiPD standard age unit is yr AD so this description is used for the LiPD version rather than yr CE which is used for the CSV version. yr CE and yr AD are identical values.
cdata_datasetID	dataSetName	
cdata_coreName	paleoData_core	
cdata_altCoreName	NA	Joined with cdata_coreName, separated by “;”.
cdata_IGSN	hasIGSN	
cdata_collectTime	collectionYear	
cdata_minYear	minYear	Values for LiPD extracted from GBRC D record ‘Age’ field.
cdata_maxYear	maxYear	
cdata_archiveSpecies	paleoData_sensorSpecies	
cdata_isDatabaseDuplicate	paleoData_gbrIsDatabaseDuplicate	
cdata_dataCoverageGroup	paleoData_gbrDataCoverageGroup	
cdata_coralNotes	notes	
geo_latitude	geo_latitude	
geo_longitude	geo_longitude	
geo_siteName	geo_siteName	
geo_siteName2	geo_location	
geo_elevation	geo_elevation	
geo_elevationNotes	geo_notes	
dating_UThDate	paleoData_gbrUThDate	
Dating_UThDateError	paleoData_gbrUThDateUncertainty	
dating_14CDate	paleoData_gbr14CDate	
Dating_14CDateError	paleoData_gbr14CDateUncertainty	
dating_notes	paleoData_gbrDatingNotes	
meths_primaryVariablesList	NA	List of variables intrinsic to LiPD and therefore unnecessary to translate as a separate field in LiPD format.
meths_additionalVariablesList	NA	
meths_TEMethod	paleoData_measurementMethod	GBRC D CSV fields are split by measurement type whereas for the LiPD TS a single field name is used, and the information is
meths_TEMethodMachine	paleoData_measurementInstrument	
meths_isotopeMethod	paleoData_measurementMethod	
meths_isotopeMethodMachine	paleoData_measurementInstrument	

<b>GBRCD Field</b>	<b>LiPD Field</b>	<b>Notes</b>
meths_luminMethod	paleoData_measurementMethod	grouped with the variable. Where multiple methods and/or instruments have been used for one proxy both are available in the relevant field separated by ‘;’.
meths_luminMethodMachine	paleoData_measurementInstrument	
meths_altMethodInfo	paleoData_measurementMethod	
meths_methodNotes	paleoData_notes	
meths_hasResolutionNominal	paleoData_samplingResolution	
meths_resolutionMin	NA	Values intrinsic to LiPD format.
meths_resolutionMax	NA	
meths_resolutionMean	NA	
meths_resolutionMedian	NA	
meths_isAnomaly	paleoData_gbrIsAnomaly	
meths_chronologyNotes	paleoData_gbrChronologyNotes	
meths_coralExtensionRate	paleoData_coralExtensionRate	
meths_coralExtensionRateNotes	paleoData_coralExtensionRateNotes	
meths_tissueThickness	paleoData_coralTissueThickness	
meths_jcpMeasured	paleoData_jcpUsed	
meths_jcpSrCaValue	paleoData_jcpMeasured	
meths_SrCaAnalyticalPrecision	paleoData_uncertaintyAnalytical	
meths_SrCaAnalyticalPrecisionUnits	paleoData_uncertaintyAnalyticalUnits	GBRCD CSV fields are split by variables whereas for the LiPD TS a single field name is used (for the uncertainty and for the uncertainty units) and the information is grouped with the variable.
meths_UCaAnalyticalPrecision	paleoData_uncertaintyAnalytical	
meths_UCaAnalyticalPrecisionUnits	paleoData_uncertaintyAnalyticalUnits	
meths_d18OAnalyticalPrecision	paleoData_uncertaintyAnalytical	
meths_d18OAnalyticalPrecisionUnits	paleoData_uncertaintyAnalyticalUnits	
meths_altPrecisionList	N/A	
meths_altAnalyticalPrecision	paleoData_analyticalUncertainty	
meths_altAnalyticalPrecisionUnits	paleoData_uncertaintyAnalyticalUnits	
meths_archiveDiagenesisCheck	paleoData_measurementMaterialScreening	
calib_isSSTCalibration	gbrIsSstCalibration	
calib_useSSTCalibration	gbrUseSstCalibration	
calib_hasAlternateCalibration	gbrHasAlternateCalibration	
calib_notes	calibration_notes	
calib_isComposite	gbrIsCompositeCalibration	
calib_compositeList	gbrCompositeCalibrationList	
calib_SSTdata	calibration_targetDataset	
calib_fitPeriod	calibration_datasetRange	
calib_hasCentredEquation	N/A	The GBRCD field is a flag for information occurring in calib_centredEquationNotes and is not used in the LiPD format.
calib_centredEquationNotes	calibration_gbrCentredEquationNotes	

<b>GBRCD Field</b>	<b>LiPD Field</b>	<b>Notes</b>
calib_method	calibration_method	
calib1_equationIntercept	calibration_equationIntercept	
calib1_equationSlope	calibration_equationSlope	
calib1_equationSlopeUncertainty	calibration_equationSlopeUncertainty	GBRCD CSV fields are split by measurement type whereas for the LiPD TS a single field name is used (for the intercept, slope, slope uncertainty and $r^2$ ) and the information is grouped with the variable.
calib1_equationR2	calibration_equationR2	
calib2_equationIntercept	calibration_equationIntercept	
calib2_equationSlope	calibration_equationSlope	
calib2_equationSlopeUncertainty	calibration_equationSlopeUncertainty	
calib2_equationR2	calibration_equationR2	
calib3_equationIntercept	calibration_equationIntercept	
calib3_equationSlope	calibration_equationSlope	
calib3_equationSlopeUncertainty	calibration_equationSlopeUncertainty	
calib3_equationR2	calibration_equationR2	
pub1_firstauthor	pub1_firstAuthor	
pub1_year	pub1_year	
pub1_doi	pub1_doi	
pub1_citation	pub1_citation	
pub1_authors	pub1_author	
pub1_title	pub1_title	
pub1_journal	pub1_journal	
pub1_doiData	pub1_doiData	
pub1_altDataURL	pub1_altDataURL	
pub2_firstauthor	pub2_firstAuthor	
pub2_year	pub2_year	
pub2_doi	pub2_doi	
pub2_citation	pub2_citation	
pub2_authors	pub2_author	
pub2_title	pub2_title	
pub2_journal	pub2_journal	
pub2_altDataURL	pub2_altDataURL	
pub3_firstauthor	pub3_firstAuthor	
pub3_year	pub3_year	
pub3_doi	pub3_doi	
pub3_citation	pub3_citation	
pub3_authors	pub3_author	
pub3_title	pub3_title	
pub3_journal	pub3_journal	
pub3_altDataUrl	pub3_altDataUrl	
pub_isCoreIDOtherStudy	gbrIsCoreIDOtherStudy	
pub_coreIDOtherStudyList	gbrCoreIDOtherStudyList	

## 575 **Appendix C – GBRC D Additional Coral Information**

Several of the coral records were measured in specific contexts, had additional processing to recreate the original published age model in the GBR database or are published in the GBR database for the first time. This contextual and methodological information is summarised here and is included in the relevant notes field(s) in the GBR database metadata table.

### **C1 Split Datasets**

580 In the GBRC D four corals and their related records from three publications were given split IDs due to differences in the data, such as resolution and/or age model. The Alibert et al. (2003) (Davies-2; GBRC D ID: AL03DAV01\_1 & AL03DAV01\_2, PAN 98-2; AL03PAN01\_1 from AL03PAN01\_2, AL03PAN01\_3, & AL03PAN01\_4), Druffel and Griffin (1999) (Abraham-1; DR99ABR01\_1 & DR99ABR01\_2) and Wei et al. (2009) Arlington Reef (AREO 4; WE09ARL01\_1 & WE09ARL01\_2) coral records were split due to differences in the resolution of the data. AREO4 (WE09ARL01\_1 & WE09ARL01\_2) was split  
585 where the resolution changed from quinquennial (1800–1939) to annual (1940–2004) for all measured variables. The Abraham-1 (DR99ABR01\_1 & DR99ABR01\_2) coral record was previously archived separately ( $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  is ~annual, and the  $\Delta^{14}\text{C}$  is ~biennial), and this was maintained in the database. Davies-2 (AL03DAV01\_1 & AL03DAV01\_2) and PAN 98-2 (AL03PAN01\_1 from AL03PAN01\_2, AL03PAN01\_3 & AL03PAN01\_4) records were split due to differences in resolutions arising from two methods of measuring trace elements (thermal ionisation mass spectrometer (TIMS) vs laser  
590 ablation inductively coupled mass spectrometer (LA-ICP-MS)). Additionally, the LA-ICP-MS measurements from Alibert et al. (2003) were not combined into a single record as there were three distinct age models for the data and each of these was presented in the original publication.

### **C2 Additions, modifications and PhD theses**


The GBRC D includes 15 records from five publications where the age model was created for archiving in the GBRC D. As  
595 described in the main manuscript, a base level of information was required to archive each record in the GBRC D, notably each dataset should have an age model with an age assigned to each data point. A brief description of the method(s) used to create the age model for these records, as well any change(s) in the record compared to the previously published version, is described below. Additionally, as PhD theses can be more difficult to access than published journal articles, a brief description of relevant information is supplied for six records from three PhD theses.

#### **600 C2.1 Chakraborty (1993) – PhD thesis**


SR coral records (track 1 and track 2) (CH93SLY01 & CH93SLY02) were collected from Stanley Reef in December 1986 (Chakraborty et al., 2000). The coral was prepared and analysed for stable isotopes along two tracks, track 1 close to the central growth axis and track 2 ~20° off the axis, as described in Chakraborty et al. (2000). Methods were as described in Chakraborty and Ramesh (1993, 1997). The original coral age model was based on Comprehensive Ocean-Atmosphere Data Set (COADS)  
605 SST data, however, since COADS is missing months of data for the Stanley Reef region, the HadISST1.1 data (grid centred on 19.5° S, 148.5° E; Raynor et al. 2003) was used for archiving in the GBRC D. A new age model was created to obtain ages based on linear interpolation using QAnalySeries to align  $\delta^{18}\text{O}$  minima (maxima) with SST maxima (minima) for quarterly averaged HadISST data. The age model was based on available information; however, the records published in Chakraborty et al. (2000) have additional  $\delta^{18}\text{O}$  measurements not available to be archived in the GBRC D. There was no change in the SR  
610 track 1 age model, but for SR track 2, ~two years of data (summer 1978 to winter 1980) do not have sufficient data points to pair with the SST data and so these years were not included in the database.

## C2.2 Ellis et al. (2019)

PET12 (EL190TI01; Ellis et al., 2019) was harvested by J. Mallela in May 2012 from Pete's Bay in the One Tree Island lagoon. The original PET12 chronology published in Ellis et al. (2019) was based on Sr/Ca maxima and minima, and was applied to 3 pieces of PET-12 (PET12-1, 2 and 3) after data smoothing with a 10-point running mean and then interpolated to a monthly time series. LA-ICP-MS data from an additional piece of PET-12 (PET-12-4) was combined with the data from the other pieces (PET12-1, 2 and 3) (Fig. C1). To align the entire PET12 LA-ICP-MS dataset with the standard procedure used by other LA-ICP-MS datasets in the GBRC, the original data smoothed with a 10-point running mean was further reduced with a 10-point mean. A new chronology was then assigned to the dataset using QAnalySeries to tie Sr/Ca maxima (minima) with SST minima (maxima) using a weekly averaged NOAA Reyn\_SmithOiv2 weekly SST dataset (Reynolds et al., 2002) centred on 23.5° S, 152.5° E. Scanning electron microscopy (SEM) analysis of an alternate slice of the PET12 coral by A. Arzey indicated variable and patchy diagenetic alteration with evidence of micro-borers and secondary aragonite crystals generally 0-10 µm in length, but up to 20 µm length observed in some sections (Fig. C2).



625 **Figure C1. PET12 (EL190TI01) (a) X-radiograph (negative) of PET12 (EL190TI01) coral. Black boxes indicate the 4 pieces analysed and are labelled with the piece number (1–4 in the top of the box). (b) Photos of PET12 pieces 1–4 and dashed black lines indicate the track analysed by LA-ICP-MS for each section of the record included in EL190TI01. X-radiograph and photos supplied by J. Mallela.**



630

635

**Figure C2. PET12 (EL180TI01) SEM images from alternate slice of PET12 coral. (a) Approximate locations of SEM images labelled b–e on coral X-radiograph. X-radiograph by PRP Diagnostic Imaging Wollongong. (b) to (e) Select SEM images from PET12 (EL190TI01); (b) No secondary aragonite. Evidence of micro-borers (example noted by white arrow), (c) Low-level diagenetic alteration with nearly 100% cover of secondary aragonite needles  $\leq 5 \mu\text{m}$  and evidence of micro-borers, (d) Bioclastic cement plug (marked by white box) above relatively pristine coral surface with cement chips and evidence of micro-borers, (e) 90% cover of secondary aragonite needles of 5–20  $\mu\text{m}$ . Scale bar indicates 100  $\mu\text{m}$  (b–e).**

### C2.3 Fallon (2000) – PhD thesis

640

Pan 1-98b (FA00PAN01) is a *Porites* sp. coral collected from Pandora Reef in October 1998, and is an additional core from the same coral as Pan 1-98a (related record published in Fallon et al. (2003); FA03PAN01). Pan 1-98b was dead on top when collected and likely stopped recording environmental information in March/April 1998, as described in Fallon (2000). A 46 mm section from Pan 1-98b covering the period 1996–1998 was analysed by LA-ICP-MS as described in Fallon et al. (2003),

the method used an ArF excimer laser (193 nm) that was masked to illuminate a 50x500 µm rectangle on the coral surface and a laser pulsed at 5 Hz using a 50 mJ power setting.

#### **C2.4 Jupiter (2008)**

645 KIA (JU08KES01b) and RTF (JU08RTI01b) are *Porites* spp. coral cores collected in 2004 from Keswick Island and Round Top Island, respectively and were previously described in Jupiter (2006) (PhD thesis). Both corals' annual growth bands were bulk sampled and 3–5 consecutive years per decade were analysed for rare earth elements and Yttrium (REYs) for the period 1950–2002. Samples were analysed using a Thermo X-Series inductively coupled mass spectrometer (ICP-MS) at the University of Queensland.

#### **650 C2.5 Lewis (2005) – PhD thesis**

MAG01D (LE05MAG01) is a modern *Porites* sp. coral collected from Geoffrey Bay, Magnetic Island by the Australian Institute of Marine Science (AIMS) in 1987, while NEL01D (LE05NEL01) and NEL03D (LE05NEL02) are fossil *Porites* spp. corals collected from Nelly Bay Harbour, Magnetic Island in 2001, and were previously described in Lewis (2005). Top and bottom growth bands for NEL01D and NEL03D were U-Th dated at the ACQUIRE laboratory at the University of  
655 Queensland. Fragments (2 x 2 cm samples) of the bottom and top of MAG01D, NEL01D and NEL03D were platinum coated and examined for diagenesis using Scanning Electron Microscopy and additional thin sections were also examined with a Leica IM50 microscope. Analysis for diagenetic alteration of the corals noted no distinct differences between the modern and fossil corals. The corals were sampled by homogenising 5-yearly increments and were split into two batches. One batch was pre-treated with H<sub>2</sub>O<sub>2</sub> and was analysed for Ba/Ca, Mg/Ca, Sr/Ca, and Mn at the Queensland Health Scientific Services (QHSS)  
660 laboratory. The other batch was not pre-treated and was analysed for δ<sup>13</sup>C and δ<sup>18</sup>O with a Micromass PRISM III stable isotope mass spectrometer at the University of Wollongong.

#### **C2.6 Marshall (2000) – PhD thesis**

Myr-2 (MA00MYR01) is a *Porites* sp. coral collected from Myrmidon Reef in July 1996 (Marshall and Mcculloch, 2002). As  
665 the data was archived in the PhD thesis appendix without an age model, a new coral age model was created for archiving in the GBRCd. The new coral age model was based on published information from Marshall and Mcculloch (2002) and Marshall (2000). The original data chronology and calibration was based on Australian Institute of Marine Science (AIMS) weekly instrumental Myrmidon Reef SST data. However, the AIMS Myrmidon loggers' SST data only covered the period from 1988 to 1996, whereas the Myr-2 Sr/Ca record extends back to 1973, and is at approximately monthly resolution (<20 measurements  
670 per year). A new chronology was assigned to the data using QAnalySeries to align Sr/Ca minima (maxima) with SST maxima (minima) for HadISST grid square centred on 18.5° S, 147.5° E.

Myr2-16D (MA00MYR02) & Myr2-17D (MA00MYR03) are *Porites* sp. coral collected from the Myrmidon Hole 2 drill core matrix (Davies and Hopley, 1983; Marshall, 2000). Myr2-16D was analysed by Marshall (2000) for δ<sup>13</sup>C and δ<sup>18</sup>O at The Australian National University using a Finnigan MAT 251 mass spectrometer coupled with a Kiel device to assess the  
675 fluorescent banding in the coral and its potential link with precipitation/river flow. The δ<sup>18</sup>O data presented in Marshall (2000) have a correction of 0.011 ‰ per metre applied for ice volume that assumed sea level was 15–20 m below present-day sea level. The correction is not applied to the δ<sup>18</sup>O data archived in the GBRCd. Myr2-17D was analysed for B, Mg, Ca, Sr, Ba and U by LA-ICP-MS using a 193 nm λ ArF excimer laser connected to a VG Elemental Plasmaquad PQ2+ instrument with a 50 x 500 µm slit as described in Marshall (2000). TIMS U-series dates were available for both coral pieces, however the data  
680 for both corals did not include an age model. An age model for Myr2-16D and Myr2-17D was created for archiving using QAnalySeries to match δ<sup>18</sup>O or Sr/Ca maxima with generalised monthly SST minima (minimum SST occurring in July).

The Myr2-17D U-Th calibrated age published in Marshall (2000) was  $7880 \pm 60$  years [ago]. However, no date for the U-series analysis is supplied. Thus, to determine the age relative to the Common Era for archiving it is assumed that it was dated in the same year as publication (i.e. 2000), thus transforming Myr2-17D age to  $-5881 \pm 60$  CE (i.e.  $5881 \pm 60$  BCE).

685 Myr2-16D was in 3 parts and each section was previously U-Th dated separately giving a calibrated age range from  $7646 \pm 60$  years [ago] for the basal section to  $7968 \pm 100$  years [ago] for the top section (with a middle section dated as  $7846 \pm 30$  years [ago]) (Marshall, 2000). Density band counting of an alternate slice suggests a maximum of 20 years between the top and bottom of the coral piece. The U-series dates are not within error and/or suggest an age reversal in the Myrmidon Hole 2 drill core when considering the date of the top (youngest) section of Myr2-16D is older than Myr2-17D. In 2022 a piece of the  
690 middle section (~5–6 cm from the top of Myr2-16D) of the alternate slice was U-Th dated and used as the age of Myr2-16D for the GBRC. The 2022 U-series date was measured at the Radiogenic Isotope Facility, University of Queensland and gave a calibrated age of  $-5755 \pm 13$  CE (i.e.  $5755 \pm 13$  BCE; Table D1).

St-1 is a *Porites* sp. coral collected from Stanley Reef in January 1999 (Marshall 2000; Marshall & McCulloch 2002). The age model was created based on published information from Marshall & McCulloch (2002) and Marshall (2000). A chronology  
695 was assigned to the data for archiving in the GBRC using QAnalySeries to align Sr/Ca minima (maxima) with SST maxima (minima) for fortnightly averaged AIMS in situ logger data from Davies Reef and Hardy Reef. The original publications only used the Davies Reef SST data, which due to missing data can only create an age model for the period of August 1996 to January 1999. The AIMS Hardy Reef SST data closely resembles the Davies Reef SST data, and both are approximately equidistant from Stanley Reef making them a suitable dataset to estimate the regional SST seasonality. Using both SST datasets  
700 enables the St-1 age model to be extended to cover the period from February 1995 to January 1999.

### **C2.7 Razak et al. (2017)**

Myra-30 (RA17MYR01) and myra-31 (RA17MYR02) are both *Isopora palifera* corals harvested from Myrmidon Reef in May 2013 (Razak et al., 2017). The age models were created for archiving in the GBRC based on published information from Razak et al. (2017). A chronology was assigned using QAnalySeries to align Sr/Ca minima (maxima) with Myrmidon  
705 SST maxima (minima) for fortnightly averaged SSTs from a composite of 10 AIMS in situ loggers. Outliers were removed from the Sr/Ca records to match values reported in Razak et al. (2017). No outliers were removed from other measured trace element ratios.

### **C2.8 Sadler et al. (2016b)**

All modern and Holocene coral records from Heron Reef published in Sadler et al. (2016b) (SA16HER01b–SA16HER06b; modern: P HS 1, P HS 3, P HS 4 & P HS 7. Holocene: R9C, R15B3) were publicly archived with time series for all data points  
710 expressed as calendar year integers (i.e. all data from 2014 had an age listed as 2014). Therefore, to recreate the seasonal cycle (as presented in Sadler et al. 2016b) the chronology was assigned by tying Sr/Ca minima (maxima) to bimonthly averaged SST maxima (minima) using QAnalySeries. The original coral chronologies and Sr/Ca-SST calibration for the modern records were created using the Commonwealth Scientific and Industrial Research Organisation/Pacific Marine Environmental  
715 Laboratory Moored Autonomous pCO<sub>2</sub> (MAPCO 2) project in the Wistari Channel (Sadler et al., 2016b), but this was not available for the GBRC. For the GBRC, modern SST data was averaged from AIMS in situ temperature loggers. The AIMS data loggers from the region have patchy temporal coverage, so the bimonthly SST dataset is an average of data logger data from Erskine Island (x1), Halftide Rocks (x1), Halfway Island (x2), Heron Reef (x16), Square Rocks (x2), and Tryon Island (x2). For the Holocene corals the SST data used to reconstruct the age model was a generalised seasonal cycle based on the  
720 AIMS bimonthly averaged SST data for the Southern GBR region (SST minimum occurs in July–August). Holocene corals were dated at the Radiogenic Isotope Facility, University of Queensland as described in Sadler et al. (2016b). However, no



measurement date for U-Th analysis was published. To determine the coral ages relative to the Common Era the measurement date was determined using the Radiogenic Isotope Facility records (June 2015; Pers. comm. T. Clark).


### C2.9 Xiao et al. 2020

725 10AR2 (XI20ARL01) is a *Porites* sp. coral harvested from Arlington Reef in April 2010 (Deng et al., 2014). The age model was created for archiving in the GBRCDB based on information published in Xiao et al. (2020). A chronology was assigned using QAnalySeries to align Sr/Ca minima with the start of each year (i.e. January) to obtain a record with 10–16 data points per year.

## C3 Unpublished Records

### 730 C3.1 AR24DIP01

Dip-04 (AR24DIP01) is an *Isopora palifera* coral that was harvested from Dip Reef in January 1988, which was in an existing collection held at the Australian Institute of Marine Science (AIMS). The data were measured by T. Razak but have not been published previously. As per the methods described in Razak et al. (2017), ~500 µg of coral powder was milled at 1 mm increments with a 1–2 mm depth using a hand-held drill with a 1 mm drill bit (Fig. C3). Trace elements (Ba/Ca, Mg/Ca, Sr/Ca, 735 U/Ca, and Y/Ca) were measured using a Thermo X-Series II quadruple ICP-MS at the University of Queensland Radiogenic Isotope Facility following a modified method described in Nguyen et al. (2013). The age model for this study was created by aligning Sr/Ca minima (maxima) with SST maxima (minima) using QAnalySeries. The seasonal cycle pattern was cross-checked with U/Ca data. As no nearby in situ instrumental measurements of SST exist for the period included in the coral record, chronology was established using the HadISST dataset (grid square centred on 18.5° S, 147.5° E). No outliers were 740 removed from the dataset.




**Figure C3. X-radiograph (positive) of Dip-04 (AR24DIP01) coral. White dashed line indicates the track analysed with white solid line the end of record section. X-radiograph supplied by T. Razak.**


### 745 C3.2 AR24OTI01

GUT12 (AR24OTI01) is a *Porites* sp. coral that was harvested in May 2012 by J. Mallela from the Gutter in the One Tree Island lagoon. The data have not been published previously. GUT12 was previously sliced into ~7 mm thick slabs, and a section of GUT12 ~15 cm in length was selected for trace element analysis by A. Arzey (Fig. C4). This section was measured using the AIMS densitometer and examined for diagenetic alteration using a Phenom XL benchtop SEM at the University of 750 Wollongong by A. Arzey. Micro-borer holes and secondary aragonite crystals were present along most of the analysis track

with patches of trace amounts of secondary aragonite crystal ( $<2\ \mu\text{m}$ ) increasing along the measurement track until 100% of skeleton was covered in secondary aragonite crystals generally between 0–10  $\mu\text{m}$  length (Fig. C5). Overall diagenetic alteration is low level, and the coral skeleton preservation is fair to good. Coral powder was drilled continuously from a ledge at 0.3–0.6 mm increments using a 2 mm TiN-coated end mill to give a resolution of ~17-34 (median 24) samples per year. Mg/Ca and Sr/Ca ratios were measured by inductively coupled atomic emission spectrometry (ICP-AES) at the Australian Nuclear Science and Technology Organisation (ANSTO) with a Thermo Scientific iCAP 7600 series ICP-AES. Data was standardised using measured JCP-1. The coral age model was created using QAnalyzeSeries to align Sr/Ca minima (maxima) with SST maxima (minima) using the fortnightly averaged NOAA Reyn\_SmithOIv2 weekly SST dataset, grid centred on 23.5° S, 152.5° E.



760 **Figure C4. X-radiograph (positive) of GUT12 (AR24OTI01) coral. Dashed white line indicates the track analysed by ICP-AES for the record included in AR24OTI01. X-radiographs by PRP Diagnostic Imaging Wollongong.**




765 **Figure C5. GUT12 (AR24OTI01) SEM images (a) Approximate locations of SEM images labelled b–e on coral X-radiograph. (b) to (e) Select SEM images from GUT12; (b) Secondary aragonite needles  $\leq 5 \mu\text{m}$  patchy cover of coral surface (~50%) and evidence of micro-borers, (c) No secondary aragonite needles. Evidence of micro-borers, (d) low-level diagenetic alteration with 100% cover of secondary aragonite needles  $< 5 \mu\text{m}$ , (e) low-level diagenetic alteration with 100% cover of secondary aragonite needles of 5-15  $\mu\text{m}$ . Scale bar indicates 100  $\mu\text{m}$  (b-e).**


### C3.3 AR24OTI02 & AR24OTI03

770 OTI 1-3 (AR24OTI02) and OTI 3-1 (AR24OTI03) are *Porites* spp. corals that were harvested in 2004 from the One Tree Island lagoon by G. Marion. The corals were part of the Enrichment of Nutrients Coral Reef Experiment (ENCORE) (Koop et al., 2001; Steven and Atkinson, 2003), and were transplanted from the reef slope onto the reef flat on 17/01/1995 and stained with Alizarin Red S on 23/11/1995. OTI 1-3 (Fig. C6) was collected from the reference treatment patch reef at ambient levels


of nutrients, while OTI 3-1 (Fig. C7) was from the enriched nitrogen treatment patch reef. Both corals were measured by LA-  
775 ICP-MS as per methods described in Lewis et al. (2012); corals were analysed using a Varian 820 ICP-MS, with output data  
smoothed with a 10-point running mean followed by a 10-point mean. The original age model supplied by S. Lewis (where  
chronology assigned with Sr/Ca minima (maxima) matched to February 8<sup>th</sup> (August 8<sup>th</sup>)) was adjusted to improve correlation  
with SST by fitting data to match the annual maxima and minima of a fortnightly averaged NOAA Reyn\_SmithOiv2 weekly  
780 SST dataset centred on 23.5° S, 152.5° E grid. Each piece was examined for diagenetic alteration by A. Arzey, although due  
to the absence of information for both pieces, the original LA-ICP-MS track cannot be directly matched to the area examined  
by the SEM. OTI 1-3 and OTI 3-1 show highly variable diagenetic alteration (Fig. C8 & Fig. C9). Both corals show above  
average levels of diagenetic alteration compared to other corals examined, which may be due to their handling for, or the  
altered nutrient exposure from, the ENCORE experiment. There is scope for future research to determine the cause of the  
abnormal level of diagenetic alteration. Caution should be exercised if using OTI 1-3 and OTI 3-1 for climate assessment, due  
785 to the high levels of diagenesis and the corals' use in the ENCORE experiments.



**Figure C6. X-radiograph (positive) of alternate slice of OTI 1-3 (AR24OTI02) coral. Dashed white line indicates the estimated track analysed by LA-ICP-MS for the record included in AR24OTI02. X-radiographs by PRP Diagnostic Imaging Wollongong.**




790 **Figure C7. X-radiograph (positive) of alternate slice of OTI 3-1 (AR24OTI03) coral. Dashed white line indicates the estimated track analysed by LA-ICP-MS for the record included in AR24OTI03. X-radiographs by PRP Diagnostic Imaging Wollongong.**



795 **Figure C8. OTI 1-3 (AR24OTI02) SEM images. (a) Approximate locations of SEM images labelled b–e on coral X-radiograph. (b) to (e) Select SEM images from OTI 1-3; (b) low-level diagenetic alteration with 100% cover of secondary aragonite needles of 5–15 μm and evidence of micro-borers, (c) No secondary aragonite needles, with evidence of dissolution and micro-borers, (d) Evidence of micro-borers and possibly dissolution. Secondary aragonite needles and calcite crystals (0–10 μm), (e) 90% surface covered by secondary aragonite needles, left side of coral needles are  $\leq 5$  μm and right side of coral needles are  $\sim 5$ –15 μm. Evidence of micro-borers visible. Scale bar indicates 100 μm (b–e).**

800



805


**Figure C9.** OTI 3-1 (AR24OTI03) SEM images. (a) Approximate locations of SEM images labelled b–e on coral X-ray. (b) to (e) Select SEM images from OTI 3-1; (b) Secondary aragonite needles (0–10  $\mu\text{m}$ ) in lower left of image and possible low Mg-calcite cement on coral surface in upper right of image. Evidence of micro-borers, (c) No secondary aragonite. Evidence of micro-borers and possible minor dissolution of coral surface, (d) 100% cover with secondary aragonite needles (0–15  $\mu\text{m}$ ) with evidence of micro-borers. High Mg-calcite splays occur in pore space (marked by white box), (e) 100% cover with secondary aragonite needles (~5–15  $\mu\text{m}$ ) with evidence of micro-borers. Scale bar indicates 100  $\mu\text{m}$  (b–e).

#### C3.4 AR24OTI04 & AR24OTI05

810


OTI4\_6C (OTI-4-19; AR24OTI04) & OTI4\_6D (OTI-4-22; AR24OTI05) were collected from the One Tree Island hole 4 reef core matrix (Davies and Hopley, 1983) at a depth listed as between 5.72–7.47 m (OTI Hole 4 section 6) and estimated to be 6.2 m and 6.5 m depth down core, respectively. Both corals were measured by LA-ICP-MS as per methods described in Lewis et al. (2012) and as per AR24OTI02 & AR24OTI03 above. An age model was created by matching the Sr/Ca maxima with SST minima of a generalised SST time series (minimum SST occurs in August). The age model was further refined by adjusting for best fit of U/Ca maxima to SST minima. An off-cut of AR24OTI04 (Fig. C10) and alternate slice of AR24OTI05 were

815 dated in July 2022 at the Radiogenic Isotope Facility, University of Queensland to give an age of  $-3803 \pm 13$  CE &  $-3772 \pm 11$  CE respectively (Table D1). An alternate slice of AR24OTI04 was also dated in 2022, but the age used for AR24OTI04 is based on the off-cut of the slice that was measured with LA-ICP-MS as the relationship with the section dated for the alternate slice is not certain. SEM screening of AR24OTI04 coral off-cut indicated 100% cover of coral surface with secondary aragonite needles generally between 0–10  $\mu\text{m}$ , although small areas with crystals up to 20  $\mu\text{m}$  were observed (Fig. C11).



820

**Figure C10. X-radiograph (positive) of OTI4\_6C (OTI-4-19; AR24OTI04) coral. Piece is off-cut of the coral piece measured by LA-ICP-MS for the AR24OTI04 record. X-radiographs by PRP Diagnostic Imaging Wollongong.**



825 Figure C11. OTI4\_6C (OTI-4-19; AR24OTI04) SEM images. (a) Approximate locations of SEM images labelled b–e on coral X-ray.  
 (b) to (e) Select SEM images from OTI4\_6C (AR24OTI04); (b) 100% cover of secondary aragonite needles generally <10  $\mu\text{m}$ , but  
 section with uneven surface (marked by white box) indicates ~10–20  $\mu\text{m}$  crystals, (c) 100% cover of secondary aragonite needles of  
 <10  $\mu\text{m}$ , (d) 100% cover of secondary aragonite needles of 5–10  $\mu\text{m}$ . Includes secondary aragonite covering likely fungal growth or  
 part of borer tube that occurred when coral was alive (marked by white box), (e) 80% cover of trace (<5  $\mu\text{m}$ ) crystals of secondary  
 830 aragonite. Scale bar indicates 100  $\mu\text{m}$  (b–e).

### C3.5 AR24SLY01

The SR coral (AR24SLY01) was measured for radiocarbon by S. Chakraborty and R. Bhushan in 1991 at the Physical Research  
 Laboratory in Ahmedabad, India. Annual bands from the SR coral were analysed as per methods described in Chakraborty  
 (1993) for other coral samples. Briefly, the annual samples were cut from the slice using a <1 mm thickness small diamond  
 835 wheel operated by an electric hand drill. The cut bands were dried by heating overnight at 80–90° C and 15g of powder used  
 for the radiocarbon analysis.  $\beta$ -counting was carried out using a Packard Tri-Carb Liquid Scintillator Analyzer in low level  
 count mode. The coral X-ray was published in Chakraborty et al. (2000).



## Appendix D – Uranium-Thorium Dating Information

840 **Table D1. Uranium-thorium age data obtained using a Nu Plasma II Multicollector Inductively Coupled Plasma Mass Spectrometer of fossil corals from the Great Barrier Reef, Australia. Note: AA47\_OTI-4-19\_4\_Th and AA\_OTI4\_6C\_1 samples are alternate slices of the same coral piece linked to the GBRCD record AR24OTI04. (See next page)**

Notes on Table D1:

845 Values in parentheses are activity ratios calculated from atomic ratios using decay constants of Cheng et al. (2000). All values have been corrected for laboratory procedural blanks. All errors reported as 2-sigma. Uncorrected  $^{230}\text{Th}$  age (ka) was calculated using Isoplot/EX 3.75 program (Ludwig, 2003), where ka denotes thousand years.

To account for non-radiogenic or initial  $^{230}\text{Th}_0$ , two correction factors were applied:

- 850 a) A bulk-Earth correction assuming a non-radiogenic  $^{230}\text{Th}/^{232}\text{Th}_0$  activity value = 0.825+/-50%, with  $^{238}\text{U}$ ,  $^{234}\text{U}$ ,  $^{232}\text{Th}$  and  $^{230}\text{Th}$  in secular equilibrium.
- b) A two-component mixing correction value determined using the equation of Clark et al. (2014) that accounts for both detrital and hydrogenous  $^{230}\text{Th}/^{232}\text{Th}_0$  shown in Eq. (D1):

$$\left(\frac{^{230}\text{Th}}{^{232}\text{Th}}\right)_{\text{mix}} = \left(\frac{^{232}\text{Th}_{\text{live}}}{^{232}\text{Th}_{\text{dead}}}\right) \times \left(\frac{^{230}\text{Th}}{^{232}\text{Th}}\right)_{\text{live}} + \left(\frac{^{232}\text{Th}_{\text{dead}} - ^{232}\text{Th}_{\text{live}}}{^{232}\text{Th}_{\text{dead}}}\right) \times \left(\frac{^{230}\text{Th}}{^{232}\text{Th}}\right)_{\text{sed}} \quad (\text{D1})$$

855 Where  $^{232}\text{Th}_{\text{dead}}$  is the measured  $^{232}\text{Th}$  value (ppb) in the individual dead coral sample.  $^{232}\text{Th}_{\text{live}}$  is 0.77 ppb, being the mean  $^{232}\text{Th}$  value of 43 live *Porites* coral samples from the inshore Great Barrier Reef with a corresponding  $^{230}\text{Th}/^{232}\text{Th}_{\text{live}}$  activity ratio of  $1.066 \pm 0.063$  (20%) (Clark et al., 2012).  $^{230}\text{Th}/^{232}\text{Th}_{\text{sed}}$  activity ratio representative of the insoluble Th component incorporated post-mortem or as particulates during growth is assumed to be  $0.61 \pm 0.12$  (20%). This value is determined from the average y-intercept values of five  $^{230}\text{Th}/^{232}\text{Th}$  versus  $^{232}\text{Th}/^{238}\text{U}$  isochrons obtained from dead *Porites* corals sampled from the Palm Islands region (Clark et al., 2014).

The values in Table D1 are those determined after calculating the two-component correction. The ‘Bulk-Earth corr. age (ka)’ is supplied to indicate the difference in age calculated between the two corrections (~4–6 years).

860

GBRCD ID	Lab Sample ID (Sample ID)	Cleaning Method	Date of chemistry	U (ppm) <sup>232</sup> Th (ppb)	( <sup>230</sup> Th/ <sup>232</sup> Th)	( <sup>230</sup> Th/ <sup>238</sup> U)	$\delta^{234}\text{U}$	Uncorr. Age (ka)	Two-component corr. age (ka)	Bulk-Earth corr. age (ka)	Corr. Initial <sup>234</sup> U/ <sup>238</sup> U	Two-component corr. age (CE)
AR24OTI04	AA47_OTI4-19_4_Th (OTI4-19_4)	H <sub>2</sub> O <sub>2</sub>	4/7/2022	2.6683 ± 0.0014	0.2241 ± 0.00023	2153.51 ± 3.93	0.05961 ± 0.000010	1.1453 ± 0.013	5.82445 ± 0.013	5.8203 ± 0.0013	1.1477 ± 0.0017	-3803 ± 13
AR24OTI04	AA45_OTI4_6C_1_C (OTI4_6C_1)	H <sub>2</sub> O <sub>2</sub>	4/7/2022	2.6979 ± 0.0013	0.04085 ± 0.00012	11965.67 ± 36.99	0.05971 ± 0.000083	1.1457 ± 0.010	5.8339 ± 0.010	5.8301 ± 0.0010	1.1481 ± 0.0012	-3812 ± 10
AR24OTI04	AA73_OTI4_6C_1_U (OTI4_6C_1)	2 x H <sub>2</sub> O <sub>2</sub>	5/7/2022	2.7023 ± 0.0014	0.03442 ± 0.000069	14267.90 ± 34.66	0.05989 ± 0.000088	1.1462 ± 0.011	5.8491 ± 0.011	5.8453 ± 0.0011	1.1487 ± 0.0013	-3828 ± 11
AR24OTI04	AA88_OTI4_6C_1_D (OTI4_6C_1)	Water	5/7/2022	2.7092 ± 0.00085	0.03468 ± 0.000086	14211.60 ± 41.59	0.05996 ± 0.000096	1.1456 ± 0.011	5.8596 ± 0.011	5.8556 ± 0.0011	1.1481 ± 0.0010	-3838 ± 11
AR24OTI5	AA46_OTI4_6D_1_Th (OTI4_6D_1)	H <sub>2</sub> O <sub>2</sub>	4/7/2022	2.6694 ± 0.0016	0.03175 ± 0.00010	15130.62 ± 52.18	0.05931 ± 0.000094	1.1456 ± 0.011	5.7939 ± 0.011	5.7901 ± 0.011	1.1480 ± 0.0010	-3772 ± 11
MA00MYR02	AA61_MYR2_16D_1 (MYR2_16D_1)	H <sub>2</sub> O <sub>2</sub>	5/7/2022	2.4849 ± 0.0012	0.07492 ± 0.000079	7932.69 ± 12.56	0.07882 ± 0.000010	1.1444 ± 0.013	7.7762 ± 0.013	7.7698 ± 0.013	1.1477 ± 0.0011	-5755 ± 13

$$a - \delta^{234}\text{U} \left( \frac{^{234}\text{U}}{^{238}\text{U}} \right) \times 1000 \text{ (D2)}$$

## Appendix E – GBRC Data Resources

AIMS loggers: derived data based on Australian Institute of Marine Science data

<https://data.aims.gov.au/aimsrtids/datatool.xhtml>

Hadley Centre Sea Ice and Sea Surface Temperature (HadISST) (Rayner et al., 2003)

865 <https://coastwatch.pfeg.noaa.gov/erddap/griddap/erdHadISST.html>

NOAA REYN\_SmithOIV2 (Reynolds et al., 2002)

[http://iridl.ldeo.columbia.edu/SOURCES/.NOAA/.NCEP/.EMC/.CMB/.GLOBAL/.Reyn\\_SmithOIV2/](http://iridl.ldeo.columbia.edu/SOURCES/.NOAA/.NCEP/.EMC/.CMB/.GLOBAL/.Reyn_SmithOIV2/)

Reefs and shoals – Queensland: Queensland Department of Resources

© State of Queensland (Department of Resources) 2023

870 <http://qldspatial.information.qld.gov.au/catalogue/custom/search.page?q=%22Reefs and shoals - Queensland%22>

Drainage basins – Queensland

© State of Queensland (Department of Resources) 2023

<http://qldspatial.information.qld.gov.au/catalogue/custom/search.page?q=%22Drainage basins - Queensland%22>

Major watercourse lines – Queensland

875 © State of Queensland (Department of Resources) 2023

<http://qldspatial.information.qld.gov.au/catalogue/custom/search.page?q=%22Major watercourse lines - Queensland%22>

Populated places – Queensland

© State of Queensland (Department of Resources) 2023

<http://qldspatial.information.qld.gov.au/catalogue/custom/search.page?q=%22Populated%20places%20->

880 [%20Queensland%22](http://qldspatial.information.qld.gov.au/catalogue/custom/search.page?q=%22Populated%20places%20-%20Queensland%22)

Mainland – Queensland

© State of Queensland (Department of Resources) 2023

<http://qldspatial.information.qld.gov.au/catalogue/custom/search.page?q=%22Mainland%20-%20Queensland%22>

## 885 **Author Contributions**

AKA & HVM conceived the study and designed the database. AKA built and curated the database, and database testing was conducted by AKA & HWF. TRC contributed data analysis and advice for U-Th dating. AKA, SEL, JM, JMW, SC & TBR provided coral data and associated technical information. NPM facilitated converting the database to LiPD format, and AKA & HWF provided example scripts for using the database in R and Python. AKA & HVM wrote the manuscript with input from  
890 TRC, JMW, SEL, JM, NPM, HWF, SC, TBR & MJF.

## **Competing interests**

The authors declare that they have no conflict of interest.

## **Acknowledgements**

We would like to thank all the researchers whose publicly archived data or whose data were archived in supplementary material  
895 were included in the GBRC D, as well as the researchers who supplied us with their data (Appendix A).

We are thankful for the software and data sources that made this project possible. Sea surface temperature data (HadISST1.1) were provided by Met Office and accessed from <https://coastwatch.pfeg.noaa.gov/erddap/info/erdHadISST/index.html>.

Figure 1 was generated using QGIS 3.22.11 (<https://www.qgis.org>) with GIS layers sourced from State of Queensland (Department of Resources) found at <http://qldspatial.information.qld.gov.au/catalogue/>. The full list of GIS layers is available  
900 in Appendix E.

Figures 3–5 were generated using R 4.3.1 (<https://www.R-project.org/>) and the ggplot2 3.4.3, sf 1.0-14 and ozmaps 0.4.5 R packages found at <https://cran.r-project.org/web/packages/ggplot2/index.html>, <https://cran.r-project.org/web/packages/sf/index.html> & <https://cran.r-project.org/web/packages/ozmaps/index.html>.

We would also like to thank the NOAA National Centers for Environmental Information, especially Edward Gille, and LiPD  
905 teams for facilitating archiving and distribution of the GBRC D.

JM's fieldwork was undertaken with the permission of the Great Barrier Reef Marine Park Authority (GBRMPA Permit Number G12.35021.1) and One Tree Island Research Station.

## **Financial Support**

This research was funded by an Australian Government Research Training Program scholarship. This research was supported  
910 by an AINSE Ltd. Post Graduate Research Award (PGRA) (AKA). This research was funded by an Australian Research Council (ARC) Discovery Project DP200100206 to HVM, JMW and TRC. This work was supported by ARC SRIEAS Grant SR200100005 Securing Antarctica's Environmental Future and ARC Future Fellowship (FT140100286) to HVM. TRC was funded by an ARC Discovery Early Career Researcher Award (DECRA) DE180100017. JM was funded by an ARC DECRA. DE120101998.

915

## References

- GBRCD: GBR Coral Skeletal Records Database, GitHub repository [code]: <https://github.com/arzeyak/GBR-Coral-Skeletal-Records-Database>, last access: August 2024.
- Alibert, C. and McCulloch, M. T.: Strontium/calcium ratios in modern Porites corals From the Great Barrier Reef as a proxy for sea surface temperature: Calibration of the thermometer and monitoring of ENSO, *Paleoceanography*, 12, 345-363, <https://doi.org/10.1029/97PA00318>, 1997.
- Alibert, C., Kinsley, L., Fallon, S. J., McCulloch, M. T., Berkelmans, R., and McAllister, F.: Source of trace element variability in Great Barrier Reef corals affected by the Burdekin flood plumes, *Geochim. Cosmochim. Acta*, 67, 231-246, [https://doi.org/10.1016/S0016-7037\(02\)01055-4](https://doi.org/10.1016/S0016-7037(02)01055-4), 2003.
- Arzey, A. K., McGregor, H. V., Clark, T. R., Webster, J. M., Lewis, S. E., Mallela, J., McKay, N. P., Fahey, H. W., Chakraborty, S., Razak, T. B., and Fischer, M. J.: The Great Barrier Reef Coral Skeletal Records Database (2024) [dataset], <https://doi.org/10.25921/hqzk-8h74>, 2024.
- Australian Institute of Marine Science: Long-term Reef Monitoring Program - Annual Summary Report on coral reef condition for 2016/17, 2017.
- Australian Institute of Marine Science: Long-Term Monitoring Program Annual Summary Report of Coral Reef Condition 2021/22, 2022.
- Barnes, D. J. and Lough, J. M.: Systematic variations in the depth of skeleton occupied by coral tissue in massive colonies of Porites from the Great barrier reef, *J. Exp. Mar. Biol. Ecol.*, 159, 113-128, [https://doi.org/10.1016/0022-0981\(92\)90261-8](https://doi.org/10.1016/0022-0981(92)90261-8), 1992.
- Barnes, D. J., Taylor, R. B., and Lough, J. M.: Measurement of luminescence in coral skeletons, *J. Exp. Mar. Biol. Ecol.*, 295, 91-106, [https://doi.org/10.1016/s0022-0981\(03\)00274-0](https://doi.org/10.1016/s0022-0981(03)00274-0), 2003.
- Beck, J. W., Edwards, R. L., Ito, E., Taylor, F. W., Recy, J., Rougerie, F., Joannot, P., and Henin, C.: Sea-Surface Temperature from Coral Skeletal Strontium/Calcium Ratios, *Science*, 257, 644-647, <https://doi.org/10.1126/science.257.5070.644>, 1992.
- Bosscher, H.: Computerized tomography and skeletal density of coral skeletons, *Coral Reefs*, 12, 97-103, <https://doi.org/10.1007/BF00302109>, 1993.
- Boto, K. and Isdale, P.: Fluorescent bands in massive corals result from terrestrial fulvic acid inputs to nearshore zone, *Nature*, 315, 396-397, <https://doi.org/10.1038/315396a0>, 1985.
- Brenner, L. D., Linsley, B. K., and Potts, D. C.: A modern Sr/Ca- $\delta^{18}\text{O}$ -sea surface temperature calibration for *Isopora* corals on the Great Barrier Reef, *Paleoceanography*, 32, 182-194, <https://doi.org/10.1002/2016pa002973>, 2017.
- Calvo, E., Marshall, J. F., Pelejero, C., McCulloch, M. T., Gagan, M. K., and Lough, J. M.: Interdecadal climate variability in the Coral Sea since 1708 A.D, *Palaeogeogr. Palaeoclimatol.*, 248, 190-201, <https://doi.org/10.1016/j.palaeo.2006.12.003>, 2007.
- Cantin, N. E. and Lough, J. M.: Surviving Coral Bleaching Events: Porites Growth Anomalies on the Great Barrier Reef, *PLOS ONE*, 9, e88720, <https://doi.org/10.1371/journal.pone.0088720>, 2014.
- Cantin, N. E., Fallon, S. J., Wu, Y., and Lough, J. M.: Project ISP019: Calcification and geochemical signatures of industrial development of the Gladstone Harbour from century old coral skeletons, Australian Institute of Marine Science, Townsville, Qld, 40, 2018.
- Chakraborty, S.: Environmental significance of isotopic and trace elemental variations in banded corals, PhD Thesis (unpublished), The Maharaja Sayajirao University of Baroda, Vadodara, India, 119 pp., 1993.
- Chakraborty, S. and Ramesh, R.: Monsoon-induced sea surface temperature changes recorded in Indian corals, *Terra Nova*, 5, 545-551, <https://doi.org/10.1111/j.1365-3121.1993.tb00303.x>, 1993.

- Chakraborty, S. and Ramesh, R.: Environmental significance of carbon and oxygen isotope ratios of banded corals from Lakshadweep, India, *Quatern. Int.*, 37, 55-65, [https://doi.org/10.1016/1040-6182\(96\)00028-6](https://doi.org/10.1016/1040-6182(96)00028-6), 1997.
- 960 Chakraborty, S., Ramesh, R., and Lough, J. M.: Effect of intraband variability on stable isotope and density time series obtained from banded corals, *J. Earth Syst. Sci.*, 109, 145-151, <https://doi.org/10.1007/BF02719158>, 2000.
- Chen, X., Deng, W., Kang, H., Zeng, T., Zhang, L., Zhao, J.-x., and Wei, G.: A Replication Study on Coral  $\delta^{11}\text{B}$  and B/Ca and Their Variation in Modern and Fossil Porites: Implications for Coral Calcifying Fluid Chemistry and Seawater pH Changes Over the Last Millennium, *Paleoceanography and Paleoclimatology*, 36, e2021PA004319, <https://doi.org/10.1029/2021PA004319>, 2021.
- 965 Cheng, H., Edwards, R. L., Hoff, J., Gallup, C. D., Richards, D. A., and Asmerom, Y.: The half-lives of uranium-234 and thorium-230, *Chem. Geol.*, 169, 17-33, [https://doi.org/10.1016/S0009-2541\(99\)00157-6](https://doi.org/10.1016/S0009-2541(99)00157-6), 2000.
- Clark, T. R., Roff, G., Zhao, J.-x., Feng, Y.-x., Done, T. J., and Pandolfi, J. M.: Testing the precision and accuracy of the U–Th chronometer for dating coral mortality events in the last 100 years, *Quat. Geochronol.*, 23, 35-45, <https://doi.org/10.1016/j.quageo.2014.05.002>, 2014.
- 970 Clark, T. R., Roff, G., Zhao, J. X., Feng, Y. X., Done, T. J., McCook, L. J., and Pandolfi, J. M.: U-Th dating reveals regional-scale decline of branching *Acropora* corals on the Great Barrier Reef over the past century, *P. Natl. A. Sci. USA*, 114, 10350-10355, <https://doi.org/10.1073/pnas.1705351114>, 2017.
- Clark, T. R., Zhao, J.-x., Feng, Y.-x., Done, T. J., Jupiter, S., Lough, J., and Pandolfi, J. M.: Spatial variability of initial  $^{230}\text{Th}/^{232}\text{Th}$  in modern *Porites* from the inshore region of the Great Barrier Reef, *Geochim. Cosmochim. Acta*, 78, 99-118, <https://doi.org/10.1016/j.gca.2011.11.032>, 2012.
- 975 Crook, E. D., Cohen, A. L., Rebolledo-Vieyra, M., Hernandez, L., and Paytan, A.: Reduced calcification and lack of acclimatization by coral colonies growing in areas of persistent natural acidification, *P. Natl. A. Sci. USA*, 110, 11044-11049, <https://doi.org/10.1073/pnas.1301589110>, 2013.
- 980 D'Olivo, J. P. and McCulloch, M.: Impact of European settlement and land use changes on Great Barrier Reef river catchments reconstructed from long-term coral Ba/Ca records, *Sci. Total Environ.*, 830, 154461, <https://doi.org/10.1016/j.scitotenv.2022.154461>, 2022.
- D'Olivo, J. P. and McCulloch, M. T.: Response of coral calcification and calcifying fluid composition to thermally induced bleaching stress, *Sci. Rep.-UK*, 7, 2207, <https://doi.org/10.1038/s41598-017-02306-x>, 2017.
- 985 D'Olivo, J. P., McCulloch, M. T., and Judd, K.: Long-term records of coral calcification across the central Great Barrier Reef: assessing the impacts of river runoff and climate change, *Coral Reefs*, 32, 999-1012, <https://doi.org/10.1007/s00338-013-1071-8>, 2013.
- D'Olivo, J. P., McCulloch, M. T., Eggins, S. M., and Trotter, J.: Coral records of reef-water pH across the central Great Barrier Reef, Australia: assessing the influence of river runoff on inshore reefs, *Biogeosciences*, 12, 1223-1236, <https://doi.org/10.5194/bg-12-1223-2015>, 2015.
- 990 D'Olivo, J. P., Sinclair, D. J., Rankenburg, K., and McCulloch, M. T.: A universal multi-trace element calibration for reconstructing sea surface temperatures from long-lived *Porites* corals: Removing 'vital-effects', *Geochim. Cosmochim. Acta*, 239, 109-135, <https://doi.org/10.1016/j.gca.2018.07.035>, 2018.
- Dassié, E., DeLong, K. L., Kilbourne, K. H., Williams, B., Abram, N. J., Brenner, L. D., Brahmi, C., Cobb, K. M., Corrège, T., Dissard, D., Emile-Geay, J., Evangelista, H., Evans, M. N., Farmer, J., Felis, T., Gagan, M., Gillikin, D. P., Goodkin, N. F., Khodri, M., Lavagnino, A. C., LaVigne, M., Lazareth, C. E., Linsley, B., Lough, J., McGregor, H., Nurhati, I. S., Ouellette, G., Perrin, L., Raymo, M., Rosenheim, B., Sandstrom, M., Schöne, B. R., Sifeddine, A., Stevenson, S., Thompson, D. M., Waite, A., Wanamaker, A., and Wu, H.: Saving our marine archives, *Eos*, 98, 32-36, <https://doi.org/10.1029/2017EO068159>, 2017.

- 1000 Davies, P. J. and Hopley, D.: Growth fabrics and growth-rates of Holocene reefs in the Great Barrier-Reef, *BMR J. Aust. Geol. Geop.*, 8, 237-251, 1983.
- Davies, P. J., Marshall, J. F., and Hopley, D.: Relationships between reef growth and sea level in the Great Barrier Reef., *Proceedings of the Fifth International Coral Reef Congress, Tahiti, 27 May-1 June*, 95-103, 1985.
- De'ath, G., Lough, J. M., and Fabricius, K. E.: Declining Coral Calcification on the Great Barrier Reef, *Science*, 323, 116-119, <https://doi.org/10.1126/science.1165283>, 2009.
- 1005 De'ath, G., Fabricius, K. E., Sweatman, H., and Puotinen, M.: The 27-year decline of coral cover on the Great Barrier Reef and its causes, *P. Natl. A. Sci. USA*, 109, 17995-17999, <https://doi.org/10.1073/pnas.1208909109>, 2012.
- DeCarlo, T. M., Gaetani, G. A., Cohen, A. L., Foster, G. L., Alpert, A. E., and Stewart, J. A.: Coral Sr-U thermometry, *Paleoceanography*, 31, 626-638, <https://doi.org/10.1002/2015PA002908>, 2016.
- 1010 DeCarlo, T. M., Cohen, A. L., Barkley, H. C., Cobban, Q., Young, C., Shamberger, K. E., Brainard, R. E., and Golbuu, Y.: Coral macrobioerosion is accelerated by ocean acidification and nutrients, *Geology*, 43, 7-10, <https://doi.org/10.1130/g36147.1>, 2015.
- Dechnik, B., Webster, J. M., Davies, P. J., Braga, J.-C., and Reimer, P. J.: Holocene “turn-on” and evolution of the Southern Great Barrier Reef: Revisiting reef cores from the Capricorn Bunker Group, *Mar. Geol.*, 363, 174-190, <https://doi.org/10.1016/j.margeo.2015.02.014>, 2015.
- 1015 Dechnik, B., Webster, J. M., Webb, G. E., Nothdurft, L., and Zhao, J.-x.: Successive phases of Holocene reef flat development: Evidence from the mid- to outer Great Barrier Reef, *Palaeogeogr. Palaeoclimatol.*, 466, 221-230, <https://doi.org/10.1016/j.palaeo.2016.11.030>, 2017.
- DeLong, K. L., Quinn, T. M., and Taylor, F. W.: Reconstructing twentieth-century sea surface temperature variability in the southwest Pacific: A replication study using multiple coral Sr/Ca records from New Caledonia, *Paleoceanography*, 22, PA4212, <https://doi.org/10.1029/2007PA001444>, 2007.
- 1020 Deng, W., Wei, G., McCulloch, M., Xie, L., Liu, Y., and Zeng, T.: Evaluation of annual resolution coral geochemical records as climate proxies in the Great Barrier Reef of Australia, *Coral Reefs*, 33, 965-977, <https://doi.org/10.1007/s00338-014-1203-9>, 2014.
- 1025 Druffel, E. R. M. and Griffin, S.: Large variations of surface ocean radiocarbon: Evidence of circulation changes in the southwestern Pacific, *J. Geophys. Res.-Oceans*, 98, 20249-20259, <https://doi.org/10.1029/93JC02113>, 1993.
- Druffel, E. R. M. and Griffin, S.: Regional Variability of Surface Ocean Radiocarbon from Southern Great Barrier Reef Corals, *Radiocarbon*, 37, 517-524, <https://doi.org/10.1017/S0033822200031003>, 1995.
- Druffel, E. R. M. and Griffin, S.: Variability of surface ocean radiocarbon and stable isotopes in the southwestern Pacific, *J. Geophys. Res.-Oceans*, 104, 23607-23613, <https://doi.org/10.1029/1999JC900212>, 1999.
- 1030 Dutton, A., Rubin, K., McLean, N., Bowring, J., Bard, E., Edwards, R. L., Henderson, G. M., Reid, M. R., Richards, D. A., Sims, K. W. W., Walker, J. D., and Yokoyama, Y.: Data reporting standards for publication of U-series data for geochronology and timescale assessment in the earth sciences, *Quat. Geochronol.*, 39, 142-149, <https://doi.org/10.1016/j.quageo.2017.03.001>, 2017.
- 1035 Ellis, B., Grant, K., Mallela, J., and Abram, N.: Is XRF core scanning a viable method for coral palaeoclimate temperature reconstructions?, *Quatern. Int.*, 514, 97-107, <https://doi.org/10.1016/j.quaint.2018.11.044>, 2019.
- Erler, D. V., Farid, H. T., Glaze, T. D., Carlson-Perret, N. L., and Lough, J. M.: Coral skeletons reveal the history of nitrogen cycling in the coastal Great Barrier Reef, *Nat. Comm.*, 11, 1500, <https://doi.org/10.1038/s41467-020-15278-w>, 2020.
- Erler, D. V., Wang, X. T., Sigman, D. M., Scheffers, S. R., Martínez-García, A., and Haug, G. H.: Nitrogen isotopic composition of organic matter from a 168 year-old coral skeleton: Implications for coastal nutrient cycling in the Great Barrier Reef Lagoon, *Earth Planet. Sc. Lett.*, 434, 161-170, <https://doi.org/10.1016/j.epsl.2015.11.023>, 2016.

- Fallon, S. J.: Environmental Records from Corals and Coralline Sponges, PhD Thesis, Research School of Earth Sciences, Australian National University, <https://doi.org/10.25911/5d778a502c764>, 2000.
- 1045 Fallon, S. J., McCulloch, M. T., and Alibert, C.: Examining water temperature proxies in Porites corals from the Great Barrier Reef: a cross-shelf comparison, *Coral Reefs*, 22, 389-404, <https://doi.org/10.1007/s00338-003-0322-5>, 2003.
- Felis, T., McGregor, H. V., Linsley, B. K., Tudhope, A. W., Gagan, M. K., Suzuki, A., Inoue, M., Thomas, A. L., Esat, T. M., Thompson, W. G., Tiwari, M., Potts, D. C., Mudelsee, M., Yokoyama, Y., and Webster, J. M.: Intensification of the meridional temperature gradient in the Great Barrier Reef following the Last Glacial Maximum, *Nat. Comm.*, 5, 4102, <https://doi.org/10.1038/ncomms5102>, 2014.
- 1050 Gagan, M. K., Chivas, A. R., and Isdale, P. J.: High-resolution isotopic records from corals using ocean temperature and mass-spawning chronometers, *Earth Planet. Sc. Lett.*, 121, 549-558, [https://doi.org/10.1016/0012-821X\(94\)90090-6](https://doi.org/10.1016/0012-821X(94)90090-6), 1994.
- Gagan, M. K., Dunbar, G. B., and Suzuki, A.: The effect of skeletal mass accumulation in Porites on coral Sr/Ca and  $\delta^{18}O$  paleothermometry, *Paleoceanography*, 27, 16, <https://doi.org/10.1029/2011pa002215>, 2012.
- 1055 Gagan, M. K., Ayliffe, L. K., Opdyke, B. N., Hopley, D., Scott-Gagan, H., and Cowley, J.: Coral oxygen isotope evidence for recent groundwater fluxes to the Australian Great Barrier Reef, *Geophys. Res. Lett.*, 29, 1982, <https://doi.org/10.1029/2002GL015336>, 2002.
- Gagan, M. K., Ayliffe, L. K., Hopley, D., Cali, J. A., Mortimer, G. E., Chappell, J., McCulloch, M. T., and Head, M. J.: Temperature and Surface-Ocean Water Balance of the Mid-Holocene Tropical Western Pacific, *Science*, 279, 1014-1018, <https://doi.org/10.1126/science.279.5353.1014>, 1998.
- 1060 Grove, C. A., Rodriguez-Ramirez, A., Merschel, G., Tjallingii, R., Zinke, J., Macia, A., and Brummer, G.-J. A.: UV-Spectral Luminescence Scanning: Technical Updates and Calibration Developments, in: *Micro-XRF Studies of Sediment Cores: Applications of a non-destructive tool for the environmental sciences*, edited by: Croudace, I. W., and Rothwell, R. G., Springer Netherlands, Dordrecht, 563-581, [https://doi.org/10.1007/978-94-017-9849-5\\_23](https://doi.org/10.1007/978-94-017-9849-5_23), 2015.
- Guan, Y., Hohn, S., Wild, C., and Merico, A.: Vulnerability of global coral reef habitat suitability to ocean warming, acidification and eutrophication, *Glob. Change Biol.*, 26, 5646-5660, <https://doi.org/10.1111/gcb.15293>, 2020.
- 1065 Hathorne, E. C., Gagnon, A., Felis, T., Adkins, J., Asami, R., Boer, W., Caillon, N., D. C., Cobb, K. M., Douville, E., deMenocal, P., Eisenhauer, A., Garbe-Schönberg, D., Geibert, W., Goldstein, S., Hughen, K., Inoue, M., Kawahata, H., Kölling, M., Cornec, F. L., Linsley, B. K., McGregor, H. V., Montagna, P., Nurhati, I. S., Quinn, T. M., Raddatz, J., Rebaubier, H., Robinson, L., Sadekov, A., Sherrell, R., Sinclair, D., Tudhope, A. W., Wei, G., Wong, H., Wu, H. C., and
- 1070 You, C.-F.: Interlaboratory study for coral Sr/Ca and other element/Ca ratio measurements, *Geochem. Geophys. Geosy.*, 14, 3730-3750, <https://doi.org/10.1002/ggge.20230>, 2013.
- Heaton, T. J., Köhler, P., Butzin, M., Bard, E., Reimer, R. W., Austin, W. E. N., Bronk Ramsey, C., Grootes, P. M., Hughen, K. A., Kromer, B., Reimer, P. J., Adkins, J., Burke, A., Cook, M. S., Olsen, J., and Skinner, L. C.: Marine20—The Marine Radiocarbon Age Calibration Curve (0–55,000 cal BP), *Radiocarbon*, 62, 779-820, <https://doi.org/10.1017/RDC.2020.68>,
- 1075 2020.
- Hendy, E. J., Gagan, M. K., and Lough, J. M.: Chronological control of coral records using luminescent lines and evidence for non-stationary ENSO teleconnections in northeast Australia, *The Holocene*, 13, 187-199, <https://doi.org/10.1191/0959683603hl606rp>, 2003a.
- Hendy, E. J., Lough, J. M., and Gagan, M. K.: Historical mortality in massive Porites from the central Great Barrier Reef, Australia: evidence for past environmental stress?, *Coral Reefs*, 22, 207-215, <https://doi.org/10.1007/s00338-003-0304-7>,
- 1080 2003b.
- Hendy, E. J., Gagan, M. K., Lough, J. M., McCulloch, M., and deMenocal, P. B.: Impact of skeletal dissolution and secondary aragonite on trace element and isotopic climate proxies in Porites corals, *Paleoceanography*, 22, <https://doi.org/10.1029/2007PA001462>, 2007.



- 1085 Hendy, E. J., Gagan, M. K., Alibert, C. A., McCulloch, M. T., Lough, J. M., and Isdale, P. J.: Abrupt Decrease in Tropical Pacific Sea Surface Salinity at End of Little Ice Age, *Science*, 295, 1511-1514, <https://doi.org/10.1126/science.1067693>, 2002.
- Hendy, E. J., Tomiak, P. J., Collins, M. J., Hellstrom, J., Tudhope, A. W., Lough, J. M., and Penkman, K. E. H.: Assessing amino acid racemization variability in coral intra-crystalline protein for geochronological applications, *Geochim. Cosmochim. Acta*, 86, 338-353, <https://doi.org/10.1016/j.gca.2012.02.020>, 2012.
- 1090 Henley, B. J., McGregor, H. V., King, A. D., Hoegh-Guldberg, O., Arzey, A. K., Karoly, D. J., Lough, J. M., DeCarlo, T. M., and Linsley, B. K.: Highest ocean heat in four centuries places Great Barrier Reef in danger, *Nature*, 632, 320-326, 10.1038/s41586-024-07672-x, 2024.
- Hopley, D., Smithers, S. G., and Parnell, K.: *The Geomorphology of the Great Barrier Reef: Development, Diversity and Change*, Cambridge University Press, Cambridge, <https://doi.org/10.1017/CBO9780511535543>, 2007.
- 1095 Hughes, T. P., Kerry, J. T., Álvarez-Noriega, M., Álvarez-Romero, J. G., Anderson, K. D., Baird, A. H., Babcock, R. C., Beger, M., Bellwood, D. R., Berkemans, R., Bridge, T. C., Butler, I. R., Byrne, M., Cantin, N. E., Comeau, S., Connolly, S. R., Cumming, G. S., Dalton, S. J., Diaz-Pulido, G., Eakin, C. M., Figueira, W. F., Gilmour, J. P., Harrison, H. B., Heron, S. F., Hoey, A. S., Hobbs, J.-P. A., Hoogenboom, M. O., Kennedy, E. V., Kuo, C.-y., Lough, J. M., Lowe, R. J., Liu, G.,
- 1100 McCulloch, M. T., Malcolm, H. A., McWilliam, M. J., Pandolfi, J. M., Pears, R. J., Pratchett, M. S., Schoepf, V., Simpson, T., Skirving, W. J., Sommer, B., Torda, G., Wachenfeld, D. R., Willis, B. L., and Wilson, S. K.: Global warming and recurrent mass bleaching of corals, *Nature*, 543, 373-377, <https://doi.org/10.1038/nature21707>, 2017.
- Isdale, P.: Fluorescent bands in massive corals record centuries of coastal rainfall, *Nature*, 310, 578-579, <https://doi.org/10.1038/310578a0>, 1984.
- 1105 Isdale, P. J., Stewart, B. J., Tickle, K. S., and Lough, J. M.: Palaeohydrological variation in a tropical river catchment: a reconstruction using fluorescent bands in corals of the Great Barrier Reef, Australia, *The Holocene*, 8, 1-8, <https://doi.org/10.1191/095968398670905088>, 1998.
- Jupiter, S.: Coral rare earth element tracers of terrestrial exposure in nearshore corals of the Great Barrier Reef, *Proceedings of the 11th International Coral Reef Symposium*, Fort Lauderdale, Florida, 7-11 July, 102-106, 2008.
- 1110 Jupiter, S., Roff, G., Marion, G., Henderson, M., Schrammeyer, V., McCulloch, M., and Hoegh-Guldberg, O.: Linkages between coral assemblages and coral proxies of terrestrial exposure along a cross-shelf gradient on the southern Great Barrier Reef, *Coral Reefs*, 27, 887-903, <https://doi.org/10.1007/s00338-008-0422-3>, 2008.
- Jupiter, S. D.: *From Cane to Coral Reefs: Ecosystem Connectivity and Downstream Responses to Land Use Intensification*, University of California, Santa Cruz, 600 pp., 2006.
- 1115 Kamber, B. S., Greig, A., and Collerson, K. D.: A new estimate for the composition of weathered young upper continental crust from alluvial sediments, Queensland, Australia, *Geochim. Cosmochim. Acta*, 69, 1041-1058, <https://doi.org/10.1016/j.gca.2004.08.020>, 2005.
- Kaufman, D. S. and team, P. k. s.-i. e.: Technical note: Open-paleo-data implementation pilot – the PAGES 2k special issue, *Clim. Past*, 14, 593-600, <https://doi.org/10.5194/cp-14-593-2018>, 2018.
- 1120 Khider, D., Emile-Geay, J., McKay, N. P., Gil, Y., Garijo, D., Ratnakar, V., Alonso-Garcia, M., Bertrand, S., Bothe, O., Brewer, P., Bunn, A., Chevalier, M., Comas-Bru, L., Csank, A., Dassié, E., DeLong, K., Felis, T., Francus, P., Frappier, A., Gray, W., Goring, S., Jonkers, L., Kahle, M., Kaufman, D., Kehrwald, N. M., Martrat, B., McGregor, H., Richey, J., Schmittner, A., Scroxton, N., Sutherland, E., Thirumalai, K., Allen, K., Arnaud, F., Axford, Y., Barrows, T., Bazin, L., Pilaar Birch, S. E., Bradley, E., Bregy, J., Capron, E., Cartapanis, O., Chiang, H.-W., Cobb, K. M., Debret, M., Dommain,
- 1125 R., Du, J., Dyez, K., Emerick, S., Erb, M. P., Falster, G., Finsinger, W., Fortier, D., Gauthier, N., George, S., Grimm, E., Hertzberg, J., Hibbert, F., Hillman, A., Hobbs, W., Huber, M., Hughes, A. L. C., Jaccard, S., Ruan, J., Kienast, M., Konecky, B., Le Roux, G., Lyubchich, V., Novello, V. F., Olaka, L., Partin, J. W., Pearce, C., Phipps, S. J., Pignol, C., Piotrowska, N.,

- Poli, M.-S., Prokopenko, A., Schwanck, F., Stepanek, C., Swann, G. E. A., Telford, R., Thomas, E., Thomas, Z., Truebe, S., von Gunten, L., Waite, A., Weitzel, N., Wilhelm, B., Williams, J., Williams, J. J., Winstrup, M., Zhao, N., and Zhou, Y.: PaCTS 1.0: A Crowdsourced Reporting Standard for Paleoclimate Data, *Paleoceanography and Paleoclimatology*, 34, 1570-1596, <https://doi.org/10.1029/2019PA003632>, 2019.
- 1130 Konecky, B. L., McKay, N. P., Churakova, O. V., Comas-Bru, L., Dassié, E. P., DeLong, K. L., Falster, G. M., Fischer, M. J., Jones, M. D., Jonkers, L., Kaufman, D. S., Leduc, G., Managave, S. R., Martrat, B., Opel, T., Orsi, A. J., Partin, J. W., Sayani, H. R., Thomas, E. K., Thompson, D. M., Tyler, J. J., Abram, N. J., Atwood, A. R., Cartapanis, O., Conroy, J. L., Curran, M. A., Dee, S. G., Deininger, M., Divine, D. V., Kern, Z., Porter, T. J., Stevenson, S. L., von Gunten, L., and Iso2k Project, M.: The Iso2k database: a global compilation of paleo- $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  records to aid understanding of Common Era climate, *Earth Syst. Sci. Data*, 12, 2261-2288, <https://doi.org/10.5194/essd-12-2261-2020>, 2020.
- 1135 Koop, K., Booth, D., Broadbent, A., Brodie, J., Bucher, D., Capone, D., Coll, J., Dennison, W., Erdmann, M., Harrison, P., Hoegh-Guldberg, O., Hutchings, P., Jones, G. B., Larkum, A. W. D., O'Neil, J., Steven, A., Tentori, E., Ward, S., Williamson, J., and Yellowlees, D.: ENCORE: The Effect of Nutrient Enrichment on Coral Reefs. Synthesis of Results and Conclusions, *Mar. Pollut. Bull.*, 42, 91-120, [https://doi.org/10.1016/S0025-326X\(00\)00181-8](https://doi.org/10.1016/S0025-326X(00)00181-8), 2001.
- 1140 Lawrence, M. G., Greig, A., Collerson, K. D., and Kamber, B. S.: Rare Earth Element and Yttrium Variability in South East Queensland Waterways, *Aquat. Geochem.*, 12, 39-72, <https://doi.org/10.1007/s10498-005-4471-8>, 2006.
- Leonard, N. D., Welsh, K. J., Lough, J. M., Feng, Y. x., Pandolfi, J. M., Clark, T. R., and Zhao, J. x.: Evidence of reduced mid-Holocene ENSO variance on the Great Barrier Reef, Australia, *Paleoceanography*, 31, 1248-1260, <https://doi.org/10.1002/2016PA002967>, 2016.
- 1145 Leonard, N. D., Welsh, K. J., Nguyen, A. D., Sadler, J., Pandolfi, J. M., Clark, T. R., Zhao, J. x., Feng, Y. x., and Webb, G. E.: High resolution geochemical analysis of massive *Porites* spp. corals from the Wet Tropics, Great Barrier Reef: rare earth elements, yttrium and barium as indicators of terrigenous input, *Mar. Pollut. Bull.*, 149, 110634, <https://doi.org/10.1016/j.marpolbul.2019.110634>, 2019.
- 1150 Leonard, N. D., Lepore, M. L., Zhao, J.-x., Rodriguez-Ramirez, A., Butler, I. R., Clark, T. R., Roff, G., McCook, L., Nguyen, A. D., Feng, Y., and Pandolfi, J. M.: Re-evaluating mid-Holocene reef “turn-off” on the inshore Southern Great Barrier Reef, *Quaternary Sci. Rev.*, 244, 106518, <https://doi.org/10.1016/j.quascirev.2020.106518>, 2020.
- Lewis, S. E.: Environmental Trends in GBR lagoon and Burdekin River catchment during the mid-Holocene and since European settlement using *Porites* coral records, Magnetic Island, QLD, PhD Thesis, School of Earth Sciences, James Cook University, Townsville, QLD, 2005.
- 1155 Lewis, S. E., Shields, G. A., Kamber, B. S., and Lough, J. M.: A multi-trace element coral record of land-use changes in the Burdekin River catchment, NE Australia, *Palaeogeogr. Palaeoclimatol.*, 246, 471-487, <https://doi.org/10.1016/j.palaeo.2006.10.021>, 2007.
- 1160 Lewis, S. E., Sloss, C. R., Murray-Wallace, C. V., Woodroffe, C. D., and Smithers, S. G.: Post-glacial sea-level changes around the Australian margin: a review, *Quaternary Sci. Rev.*, 74, 115-138, <https://doi.org/10.1016/j.quascirev.2012.09.006>, 2013.
- Lewis, S. E., Lough, J. M., Cantin, N. E., Matson, E. G., Kinsley, L., Bainbridge, Z. T., and Brodie, J. E.: A critical evaluation of coral Ba/Ca, Mn/Ca and Y/Ca ratios as indicators of terrestrial input: New data from the Great Barrier Reef, Australia, *Geochim. Cosmochim. Acta*, 237, 131-154, <https://doi.org/10.1016/j.gca.2018.06.017>, 2018.
- 1165 Lewis, S. E., Brodie, J. E., McCulloch, M. T., Mallela, J., Jupiter, S. D., Williams, H. S., Lough, J. M., and Matson, E. G.: An assessment of an environmental gradient using coral geochemical records, Whitsunday Islands, Great Barrier Reef, Australia, *Mar. Pollut. Bull.*, 65, 306-319, <https://doi.org/10.1016/j.marpolbul.2011.09.030>, 2012.
- Li, Y., Liao, X., Bi, K., Han, T., Chen, J., Lu, J., He, C., and Lu, Z.: Micro-CT reconstruction reveals the colony pattern regulations of four dominant reef-building corals, *Ecol. Evol.*, 11, 16266-16279, <https://doi.org/10.1002/ece3.8308>, 2021.
- 1170

- Linsley, B. K., Dunbar, R. B., Dassié, E. P., Tangri, N., Wu, H. C., Brenner, L. D., and Wellington, G. M.: Coral carbon isotope sensitivity to growth rate and water depth with paleo-sea level implications, *Nat. Comm.*, 10, 2056, <https://doi.org/10.1038/s41467-019-10054-x>, 2019.
- 1175 Lough, J., Barnes, D., and McAllister, F.: Luminescent lines in corals from the Great Barrier Reef provide spatial and temporal records of reefs affected by land runoff, *Coral Reefs*, 21, 333-343, <https://doi.org/10.1007/s00338-002-0253-6>, 2002.
- Lough, J. M.: A strategy to improve the contribution of coral data to high-resolution paleoclimatology, *Palaeogeogr. Palaeoclimatol.*, 204, 115-143, [https://doi.org/10.1016/S0031-0182\(03\)00727-2](https://doi.org/10.1016/S0031-0182(03)00727-2), 2004.
- 1180 Lough, J. M.: Tropical river flow and rainfall reconstructions from coral luminescence: Great Barrier Reef, Australia, *Paleoceanography*, 22, PA2218, <https://doi.org/10.1029/2006pa001377>, 2007.
- Lough, J. M.: Great Barrier Reef coral luminescence reveals rainfall variability over northeastern Australia since the 17th century, *Paleoceanography*, 26, 14, <https://doi.org/10.1029/2010pa002050>, 2011a.
- Lough, J. M.: Measured coral luminescence as a freshwater proxy: comparison with visual indices and a potential age artefact, *Coral Reefs*, 30, 169-182, <https://doi.org/10.1007/s00338-010-0688-0>, 2011b.
- 1185 Lough, J. M. and Barnes, D. J.: Possible relationships between environmental variables and skeletal density in a coral colony from the central Great Barrier Reef, *J. Exp. Mar. Biol. Ecol.*, 134, 221-241, [https://doi.org/10.1016/0022-0981\(89\)90071-3](https://doi.org/10.1016/0022-0981(89)90071-3), 1990.
- Lough, J. M. and Barnes, D. J.: Comparisons of skeletal density variations in *Porites* from the central Great Barrier Reef, *J. Exp. Mar. Biol. Ecol.*, 155, 1-25, [https://doi.org/10.1016/0022-0981\(92\)90024-5](https://doi.org/10.1016/0022-0981(92)90024-5), 1992.
- 1190 Lough, J. M. and Barnes, D. J.: Several centuries of variation in skeletal extension, density and calcification in massive *Porites* colonies from the Great Barrier Reef: a proxy for seawater temperature and a background of variability against which to identify unnatural change, *J. Exp. Mar. Biol. Ecol.*, 211, 29-67, [https://doi.org/10.1016/S0022-0981\(96\)02710-4](https://doi.org/10.1016/S0022-0981(96)02710-4), 1997.
- Lough, J. M. and Barnes, D. J.: Environmental controls on growth of the massive coral *Porites*, *J. Exp. Mar. Biol. Ecol.*, 245, 225-243, [https://doi.org/10.1016/S0022-0981\(99\)00168-9](https://doi.org/10.1016/S0022-0981(99)00168-9), 2000.
- 1195 Lough, J. M., Lewis, S. E., and Cantin, N. E.: Freshwater impacts in the central Great Barrier Reef: 1648–2011, *Coral Reefs*, 34, 739-751, <https://doi.org/10.1007/s00338-015-1297-8>, 2015.
- Lough, J. M., Llewellyn, L. E., Lewis, S. E., Turney, C. S. M., Palmer, J. G., Cook, C. G., and Hogg, A. G.: Evidence for suppressed mid-Holocene northeastern Australian monsoon variability from coral luminescence, *Paleoceanography*, 29, 581-594, <https://doi.org/10.1002/2014pa002630>, 2014.
- 1200 Ludwig, K. R.: Mathematical–Statistical Treatment of Data and Errors for <sup>230</sup>Th/U Geochronology, in: Uranium-Series Geochemistry, edited by: Bourdon, B., Henderson, G. M., Lundstrom, C. C., and Turner, S. P., The Mineralogical Society of America, Washington, DC, 631-656, <https://doi.org/10.2113/0520631>, 2003.
- Madin, J. S., Anderson, K. D., Andreasen, M. H., Bridge, T. C. L., Cairns, S. D., Connolly, S. R., Darling, E. S., Diaz, M., Falster, D. S., Franklin, E. C., Gates, R. D., Harmer, A. M. T., Hoogenboom, M. O., Huang, D., Keith, S. A., Kosnik, M. A., 1205 Kuo, C.-Y., Lough, J. M., Lovelock, C. E., Luiz, O., Martinelli, J., Mizerek, T., Pandolfi, J. M., Pochon, X., Pratchett, M. S., Putnam, H. M., Roberts, T. E., Stat, M., Wallace, C. C., Widman, E., and Baird, A. H.: The Coral Trait Database, a curated database of trait information for coral species from the global oceans, *Scientific Data*, 3, 160017, <https://doi.org/10.1038/sdata.2016.17>, 2016.
- Marchitto, T. M., Bryan, S. P., Doss, W., McCulloch, M. T., and Montagna, P.: A simple biomineralization model to explain 1210 Li, Mg, and Sr incorporation into aragonitic foraminifera and corals, *Earth Planet. Sc. Lett.*, 481, 20-29, <https://doi.org/10.1016/j.epsl.2017.10.022>, 2018.

- Marion, G. S., Jupiter, S. D., Radice, V. Z., Albert, S., and Hoegh-Guldberg, O.: Linking isotopic signatures of nitrogen in nearshore coral skeletons with sources in catchment runoff, *Mar. Pollut. Bull.*, 173, 113054, <https://doi.org/10.1016/j.marpolbul.2021.113054>, 2021.
- 1215 Marshall, J. F.: Decadal-scale, high resolution records of sea surface temperature in the eastern Indian and south western Pacific Oceans from proxy records of the strontium/calcium ratio of massive porites corals, PhD thesis, Australian National University, <https://doi.org/10.25911/5d63bfacacc9f>, 2000.
- Marshall, J. F. and McCulloch, M. T.: An assessment of the Sr/Ca ratio in shallow water hermatypic corals as a proxy for sea surface temperature, *Geochim. Cosmochim. Acta*, 66, 3263-3280, [https://doi.org/10.1016/S0016-7037\(02\)00926-2](https://doi.org/10.1016/S0016-7037(02)00926-2), 2002.
- 1220 Maynard, J., van Hooidonk, R., Eakin, C. M., Puotinen, M., Garren, M., Williams, G., Heron, S. F., Lamb, J., Weil, E., Willis, B., and Harvell, C. D.: Projections of climate conditions that increase coral disease susceptibility and pathogen abundance and virulence, *Nat. Clim. Change*, 5, 688-694, <https://doi.org/10.1038/nclimate2625>, 2015.
- McCulloch, M., Fallon, S., Wyndham, T., Hendy, E., Lough, J., and Barnes, D.: Coral record of increased sediment flux to the inner Great Barrier Reef since European settlement, *Nature*, 421, 727, <https://doi.org/10.1038/nature01361>, 2003.
- 1225 McCulloch, M. T., D'Olivo, J. P., Falter, J., Holcomb, M., and Trotter, J. A.: Coral calcification in a changing World and the interactive dynamics of pH and DIC upregulation, *Nat. Comm.*, 8, 15686, <https://doi.org/10.1038/ncomms15686>, 2017.
- McCulloch, M. T., Gagan, M. K., Mortimer, G. E., Chivas, A. R., and Isdale, P. J.: A high-resolution Sr/Ca and  $\delta^{18}\text{O}$  coral record from the Great Barrier Reef, Australia, and the 1982–1983 El Niño, *Geochim. Cosmochim. Acta*, 58, 2747-2754, [https://doi.org/10.1016/0016-7037\(94\)90142-2](https://doi.org/10.1016/0016-7037(94)90142-2), 1994.
- 1230 McGregor, H. V. and Abram, N. J.: Images of diagenetic textures in Porites corals from Papua New Guinea and Indonesia, *Geochem. Geophys. Geosy.*, 9, Q10013, <https://doi.org/10.1029/2008GC002093>, 2008.
- McGregor, H. V. and Gagan, M. K.: Diagenesis and geochemistry of porites corals from Papua New Guinea: Implications for paleoclimate reconstruction, *Geochim. Cosmochim. Acta*, 67, 2147-2156, [https://doi.org/10.1016/S0016-7037\(02\)01050-5](https://doi.org/10.1016/S0016-7037(02)01050-5), 2003.
- 1235 McKay, N. P. and Emile-Geay, J.: Technical note: The Linked Paleo Data framework – a common tongue for paleoclimatology, *Clim. Past*, 12, 1093-1100, <https://doi.org/10.5194/cp-12-1093-2016>, 2016.
- Min, G. R., Edwards, R. L., Taylor, F. W., Recy, J., Gallup, C. D., and Beck, J. W.: Annual cycles of UCa in coral skeletons and UCa thermometry, *Geochim. Cosmochim. Acta*, 59, 2025-2042, [https://doi.org/10.1016/0016-7037\(95\)00124-7](https://doi.org/10.1016/0016-7037(95)00124-7), 1995.
- Mollica, N. R., Guo, W., Cohen, A. L., Huang, K.-F., Foster, G. L., Donald, H. K., and Solow, A. R.: Ocean acidification affects coral growth by reducing skeletal density, *P. Natl. A. Sci. USA*, 115, 1754-1759, <https://doi.org/10.1073/pnas.1712806115>, 2018.
- Morrill, C., Thrasher, B., Lockshin, S. N., Gille, E. P., McNeill, S., Shepherd, E., Gross, W. S., and Bauer, B. A.: The Paleoenvironmental Standard Terms (PaST) Thesaurus: Standardizing Heterogeneous Variables in Paleoscience, Paleooceanography and Paleoclimatology, 36, e2020PA004193, <https://doi.org/10.1029/2020PA004193>, 2021.
- 1245 Murphy, R., Webster, J. M., Nothdurft, L., Dechnik, B., and McGregor, H. V.: High-resolution hyperspectral imaging of diagenesis and clays in fossil coral reef material: a nondestructive tool for improving environmental and climate reconstructions, *Geochem. Geophys. Geosy.*, 18, 3209-3230, [10.1002/2017GC006949](https://doi.org/10.1002/2017GC006949), 2017.
- Nguyen, A. D., Zhao, J. x., Feng, Y. x., Hu, W. p., Yu, K. f., Gasparon, M., Pham, T. B., and Clark, T. R.: Impact of recent coastal development and human activities on Nha Trang Bay, Vietnam: evidence from a Porites lutea geochemical record, *Coral Reefs*, 32, 181-193, <https://doi.org/10.1007/s00338-012-0962-4>, 2013.
- 1250 Nothdurft, L. D. and Webb, G. E.: Earliest diagenesis in scleractinian coral skeletons: implications for palaeoclimate-sensitive geochemical archives, *Facies*, 55, 161-201, <https://doi.org/10.1007/s10347-008-0167-z>, 2009.

- Nothdurft, L. D., Webb, G. E., Bostrom, T., and Rintoul, L.: Calcite-filled borings in the most recently deposited skeleton in live-collected *Porites* (Scleractinia): Implications for trace element archives, *Geochim. Cosmochim. Acta*, 71, 5423-5438, <https://doi.org/10.1016/j.gca.2007.09.025>, 2007.
- 1255 Okai, T., Suzuki, A., Kawahata, H., Terashima, S., and Imai, N.: Preparation of a New Geological Survey of Japan Geochemical Reference Material: Coral JCp-1, *Geostandard. Newslett.*, 26, 95-99, <https://doi.org/10.1111/j.1751-908X.2002.tb00627.x>, 2002.
- Ortiz, J.-C., Wolff, N. H., Anthony, K. R. N., Devlin, M., Lewis, S., and Mumby, P. J.: Impaired recovery of the Great Barrier Reef under cumulative stress, *Science Advances*, 4, <https://doi.org/10.1126/sciadv.aar6127>, 2018.
- 1260 Palmer, J. G., Cook, E. R., Turney, C. S. M., Allen, K., Fenwick, P., Cook, B. I., O'Donnell, A., Lough, J., Grierson, P., and Baker, P.: Drought variability in the eastern Australia and New Zealand summer drought atlas (ANZDA, CE 1500–2012) modulated by the Interdecadal Pacific Oscillation, *Environ. Res. Lett.*, 10, 124002, <https://doi.org/10.1088/1748-9326/10/12/124002>, 2015.
- 1265 Pelejero, C., Calvo, E., McCulloch, M. T., Marshall, J. F., Gagan, M. K., Lough, J. M., and Opdyke, B. N.: Preindustrial to Modern Interdecadal Variability in Coral Reef pH, *Science*, 309, 2204-2207, <https://doi.org/10.1126/science.1113692>, 2005.
- Prouty, N. G., Cohen, A., Yates, K. K., Storlazzi, C. D., Swarzenski, P. W., and White, D.: Vulnerability of Coral Reefs to Bioerosion From Land-Based Sources of Pollution, *J. Geophys. Res.-Oceans*, 122, 9319-9331, <https://doi.org/10.1002/2017JC013264>, 2017.
- 1270 Quinn, T. M. and Taylor, F. W.: SST artifacts in coral proxy records produced by early marine diagenesis in a modern coral from Rabaul, Papua New Guinea, *Geophys. Res. Lett.*, 33, <https://doi.org/10.1029/2005GL024972>, 2006.
- Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., Rowell, D. P., Kent, E. C., and Kaplan, A.: Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century, *J. Geophys. Res.-Atmos.*, 108, 4407, <https://doi.org/10.1029/2002JD002670>, 2003.
- 1275 Razak, T. B., Mumby, P. J., Nguyen, A. D., Zhao, J.-X., Lough, J. M., Cantin, N. E., and Roff, G.: Use of skeletal Sr/Ca ratios to determine growth patterns in a branching coral *Isopora palifera*, *Mar. Biol.*, 164, 96, <https://doi.org/10.1007/s00227-017-3099-8>, 2017.
- Reed, E. V., Cole, J. E., Lough, J. M., Thompson, D., and Cantin, N. E.: Linking climate variability and growth in coral skeletal records from the Great Barrier Reef, *Coral Reefs*, 38, 29-43, <https://doi.org/10.1007/s00338-018-01755-8>, 2019.
- 1280 Reynolds, R. W., Rayner, N. A., Smith, T. M., Stokes, D. C., and Wang, W.: An Improved In Situ and Satellite SST Analysis for Climate, *J. Clim.*, 15, 1609-1625, [https://doi.org/10.1175/1520-0442\(2002\)015<1609:AIISAS>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<1609:AIISAS>2.0.CO;2), 2002.
- Roche, R. C., Perry, C. T., Smithers, S. G., Leng, M. J., Grove, C. A., Sloane, H. J., and Unsworth, C. E.: Mid-Holocene sea surface conditions and riverine influence on the inshore Great Barrier Reef, *The Holocene*, 24, 885-897, <https://doi.org/10.1177/0959683614534739>, 2014.
- 1285 Rodriguez-Ramirez, A., Grove, C. A., Zinke, J., Pandolfi, J. M., and Zhao, J.-x.: Coral Luminescence Identifies the Pacific Decadal Oscillation as a Primary Driver of River Runoff Variability Impacting the Southern Great Barrier Reef, *PLOS ONE*, 9, e84305, <https://doi.org/10.1371/journal.pone.0084305>, 2014.
- Ryan, E. J., Smithers, S. G., Lewis, S. E., Clark, T. R., and Zhao, J.-x.: The Variable Influences of Sea Level, Sedimentation and Exposure on Holocene Reef Development over a Cross-Shelf Transect, Central Great Barrier Reef, *Diversity*, 10, 110, <https://doi.org/10.3390/d10040110>, 2018.
- 1290 Sadler, J., Webb, G. E., Nothdurft, L. D., and Dechnik, B.: Geochemistry-based coral palaeoclimate studies and the potential of 'non-traditional' (non-massive *Porites*) corals: Recent developments and future progression, *Earth-Sci. Rev.*, 139, 291-316, <https://doi.org/10.1016/j.earscirev.2014.10.002>, 2014.

- Sadler, J., Nguyen, A. D., Leonard, N. D., Webb, G. E., and Nothdurft, L. D.: Acropora interbranch skeleton Sr/Ca ratios: Evaluation of a potential new high-resolution paleothermometer, *Paleoceanography*, 31, 505-517, <https://doi.org/10.1002/2015PA002898>, 2016a.
- Sadler, J., Webb, G. E., Leonard, N. D., Nothdurft, L. D., and Clark, T. R.: Reef core insights into mid-Holocene water temperatures of the southern Great Barrier Reef, *Paleoceanography*, 31, 1395-1408, <https://doi.org/10.1002/2016pa002943>, 2016b.
- 1295 Saha, N., Webb, G. E., Christy, A. G., and Zhao, J.-x.: Vanadium in the massive coral *Porites*: A potential proxy for historical wood clearing and burning, *Earth Planet. Sc. Lett.*, 527, 115793, <https://doi.org/10.1016/j.epsl.2019.115793>, 2019a.
- Saha, N., Webb, G. E., Zhao, J.-X., Leonard, N. D., and Nguyen, A. D.: Influence of marine biochemical cycles on seasonal variation of Ba/Ca in the near-shore coral *Cyphastrea*, Rat Island, southern Great Barrier Reef, *Chem. Geol.*, 499, 71-83, <https://doi.org/10.1016/j.chemgeo.2018.09.005>, 2018a.
- 1305 Saha, N., Rodriguez-Ramirez, A., Nguyen, A. D., Clark, T. R., Zhao, J. X., and Webb, G. E.: Seasonal to decadal scale influence of environmental drivers on Ba/Ca and Y/Ca in coral aragonite from the southern Great Barrier Reef, *Sci. Total Environ.*, 639, 1099-1109, <https://doi.org/10.1016/j.scitotenv.2018.05.156>, 2018b.
- Saha, N., Webb, G. E., Zhao, J.-X., Lewis, S. E., Nguyen, A. D., and Feng, Y.: Spatiotemporal variation of rare earth elements from river to reef continuum aids monitoring of terrigenous sources in the Great Barrier Reef, *Geochim. Cosmochim. Acta*, 299, 85-112, <https://doi.org/10.1016/j.gca.2021.02.014>, 2021.
- 1310 Saha, N., Webb, G. E., Zhao, J.-X., Nguyen, A. D., Lewis, S. E., and Lough, J. M.: Coral-based high-resolution rare earth element proxy for terrestrial sediment discharge affecting coastal seawater quality, Great Barrier Reef, *Geochim. Cosmochim. Acta*, 254, 173-191, <https://doi.org/10.1016/j.gca.2019.04.004>, 2019b.
- 1315 Sammarco, P. W., Risk, M. J., Schwarcz, H. P., and Heikoop, J. M.: Cross-continental shelf trends in coral  $\delta^{15}\text{N}$  on the Great Barrier Reef: further consideration of the reef nutrient paradox, *Mar. Ecol. Prog. Ser.*, 180, 131-138, <http://www.jstor.org/stable/24852096>, 1999.
- Sanborn, K. L., Webster, J. M., Webb, G. E., Braga, J. C., Humblet, M., Nothdurft, L., Patterson, M. A., Dechnik, B., Warner, S., Graham, T., Murphy, R. J., Yokoyama, Y., Obrochta, S. P., Zhao, J.-x., and Salas-Saavedra, M.: A new model of Holocene reef initiation and growth in response to sea-level rise on the Southern Great Barrier Reef, *Sediment. Geol.*, 397, 105556, <https://doi.org/10.1016/j.sedgeo.2019.105556>, 2020.
- 1320 Sayani, H. R., Cobb, K. M., Cohen, A. L., Elliott, W. C., Nurhati, I. S., Dunbar, R. B., Rose, K. A., and Zaunbrecher, L. K.: Effects of diagenesis on paleoclimate reconstructions from modern and young fossil corals, *Geochim. Cosmochim. Acta*, 75, 6361-6373, <https://doi.org/10.1016/j.gca.2011.08.026>, 2011.
- 1325 Sinclair, D. J., Kinsley, L. P. J., and McCulloch, M. T.: High resolution analysis of trace elements in corals by laser ablation ICP-MS, *Geochim. Cosmochim. Acta*, 62, 1889-1901, [https://doi.org/10.1016/S0016-7037\(98\)00112-4](https://doi.org/10.1016/S0016-7037(98)00112-4), 1998.
- Spencer, T., Brown, B., Hamylton, S., and McLean, R.: 'A Close and Friendly Alliance': Biology, Geology and the Great Barrier Reef Expedition of 1928–1929, in: *Oceanography and Marine Biology* edited by: Hawkins, S. J., 89-138, <https://doi.org/10.1201/9781003138846-2>, 2021.
- 1330 Steinberg, C.: Impacts of climate change on the physical oceanography of the Great Barrier Reef, in: *Climate change and the Great Barrier Reef: a vulnerability assessment*, The Great Barrier Reef Marine Park Authority, <https://hdl.handle.net/11017/536>, 2007.
- Steven, A. D. L. and Atkinson, M. J.: Nutrient uptake by coral-reef microatolls, *Coral Reefs*, 22, 197-204, <https://doi.org/10.1007/s00338-003-0303-8>, 2003.
- 1335 Stuiver, M. and Reimer, P. J.: Extended 14C Data Base and Revised CALIB 3.0 14C Age Calibration Program, *Radiocarbon*, 35, 215-230, <https://doi.org/10.1017/S0033822200013904>, 1993.

- Suzuki, A., Gagan, M. K., Fabricius, K., Isdale, P. J., Yukino, I., and Kawahata, H.: Skeletal isotope microprofiles of growth perturbations in *Porites* corals during the 1997–1998 mass bleaching event, *Coral Reefs*, 22, 357-369, <https://doi.org/10.1007/s00338-003-0323-4>, 2003.
- 1340 Takada, N., Suzuki, A., Ishii, H., Hironaka, K., and Hironiwa, T.: Thermoluminescence of coral skeletons: a high-sensitivity proxy of diagenetic alteration of aragonite, *Sci. Rep.-UK*, 7, 17969, 10.1038/s41598-017-18269-y, 2017.
- Thompson, D., McCulloch, M., Cole, J. E., Reed, E. V., D’Olivo, J. P., Dyez, K., Lofverstrom, M., Lough, J., Cantin, N., Tudhope, A. W., Cheung, A. H., Vetter, L., and Edwards, R. L.: Marginal Reefs Under Stress: Physiological Limits Render Galápagos Corals Susceptible to Ocean Acidification and Thermal Stress, *AGU Advances*, 3, e2021AV000509, <https://doi.org/10.1029/2021AV000509>, 2022.
- 1345 Thompson, D. M.: Environmental records from coral skeletons: A decade of novel insights and innovation, *WIREs Clim. Change*, 13, e745, <https://doi.org/10.1002/wcc.745>, 2022.
- Vines, Timothy H., Albert, Arianne Y. K., Andrew, Rose L., Débarre, F., Bock, Dan G., Franklin, Michelle T., Gilbert, Kimberly J., Moore, J.-S., Renaut, S., and Rennison, Diana J.: The Availability of Research Data Declines Rapidly with Article Age, *Curr. Biol.*, 24, 94-97, <https://doi.org/10.1016/j.cub.2013.11.014>, 2014.
- 1350 Walter, R. M., Sayani, H. R., Felis, T., Cobb, K. M., Abram, N. J., Arzey, A. K., Atwood, A. R., Brenner, L. D., Dassié, É. P., DeLong, K. L., Ellis, B., Emile-Geay, J., Fischer, M. J., Goodkin, N. F., Hargreaves, J. A., Kilbourne, K. H., Krawczyk, H., McKay, N. P., Moore, A. L., Murty, S. A., Ong, M. R., Ramos, R. D., Reed, E. V., Samanta, D., Sanchez, S. C., Zinke, J., and the, P. C. k. P. M.: The CoralHydro2k database: a global, actively curated compilation of coral  $\delta^{18}\text{O}$  and Sr/Ca proxy records of tropical ocean hydrology and temperature for the Common Era, *Earth Syst. Sci. Data*, 15, 2081-2116, <https://doi.org/10.5194/essd-15-2081-2023>, 2023.
- Walther, B. D., Kingsford, M. J., and McCulloch, M. T.: Environmental Records from Great Barrier Reef Corals: inshore versus offshore drivers, *PLoS One*, 8, e77091, <https://doi.org/10.1371/journal.pone.0077091>, 2013.
- Wang, Z., Li, J., Wei, G., Deng, W., Chen, X., Zeng, T., Wang, X., Ma, J., Zhang, L., Tu, X., Wang, Q., and McCulloch, M.: Biologically controlled Mo isotope fractionation in coral reef systems, *Geochim. Cosmochim. Acta*, 262, 128-142, <https://doi.org/10.1016/j.gca.2019.07.037>, 2019.
- 1360 Weber, J. N.: Incorporation of strontium into reef coral skeletal carbonate, *Geochim. Cosmochim. Acta*, 37, 2173-2190, [https://doi.org/10.1016/0016-7037\(73\)90015-X](https://doi.org/10.1016/0016-7037(73)90015-X), 1973.
- Weber, J. N. and Woodhead, P. M. J.: Factors affecting the carbon and oxygen isotopic composition of marine carbonate sediments—II. Heron Island, Great Barrier Reef, Australia, *Geochim. Cosmochim. Acta*, 33, 19-38, [https://doi.org/10.1016/0016-7037\(69\)90090-8](https://doi.org/10.1016/0016-7037(69)90090-8), 1969.
- 1365 Weber, J. N. and Woodhead, P. M. J.: Carbon and Oxygen Isotope Fractionation in the Skeletal Carbonate of Reef-Building Corals, *Chem. Geol.*, 6, 93-117, [https://doi.org/10.1016/0009-2541\(70\)90009-4](https://doi.org/10.1016/0009-2541(70)90009-4), 1970.
- Weber, J. N. and Woodhead, P. M. J.: Temperature dependence of oxygen-18 concentration in reef coral carbonates, *J. Geophys. Res.*, 77, 463-473, <https://doi.org/10.1029/JC077i003p00463>, 1972.
- 1370 Webster, J. M., Braga, J. C., Humblet, M., Potts, D. C., Iryu, Y., Yokoyama, Y., Fujita, K., Bourillot, R., Esat, T. M., Fallon, S., Thompson, W. G., Thomas, A. L., Kan, H., McGregor, H. V., Hinestrosa, G., Obrochta, S. P., and Loughheed, B. C.: Response of the Great Barrier Reef to sea-level and environmental changes over the past 30,000 years, *Nat. Geosci.*, 11, 426-432, <https://doi.org/10.1038/s41561-018-0127-3>, 2018.
- 1375 Weerabaddana, M. M., Thompson, D. M., Reed, E. V., Farfan, G. A., Kirk, J. D., Kojima, A. C., Dettman, D. L., de Brum, K., Kabua, E., and Edwards, F.: Impact of Intra-Skeletal Calcite on the Preservation of Coral Geochemistry and Implications for Paleoclimate Reconstruction, *Paleoceanography and Paleoclimatology*, 39, e2023PA004730, <https://doi.org/10.1029/2023PA004730>, 2024.

- Wei, G., McCulloch, M. T., Mortimer, G., Deng, W., and Xie, L.: Evidence for ocean acidification in the Great Barrier Reef of Australia, *Geochim. Cosmochim. Acta*, 73, 2332-2346, <https://doi.org/10.1016/j.gca.2009.02.009>, 2009.
- Wei, G., Wang, Z., Ke, T., Liu, Y., Deng, W., Chen, X., Xu, J., Zeng, T., and Xie, L.: Decadal variability in seawater pH in the West Pacific: Evidence from coral  $\delta^{11}\text{B}$  records, *J. Geophys. Res.-Oceans*, 120, 7166-7181, <https://doi.org/10.1002/2015JC011066>, 2015.
- Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-W., da Silva Santos, L. B., Bourne, P. E., Bouwman, J., Brookes, A. J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C. T., Finkers, R., Gonzalez-Beltran, A., Gray, A. J. G., Groth, P., Goble, C., Grethe, J. S., Heringa, J., 't Hoen, P. A. C., Hooft, R., Kuhn, T., Kok, R., Kok, J., Lusher, S. J., Martone, M. E., Mons, A., Packer, A. L., Persson, B., Rocca-Serra, P., Roos, M., van Schaik, R., Sansone, S.-A., Schultes, E., Sengstag, T., Slater, T., Strawn, G., Swertz, M. A., Thompson, M., van der Lei, J., van Mulligen, E., Velterop, J., Waagmeester, A., Wittenburg, P., Wolstencroft, K., Zhao, J., and Mons, B.: The FAIR Guiding Principles for scientific data management and stewardship, *Scientific Data*, 3, 160018, <https://doi.org/10.1038/sdata.2016.18>, 2016.
- Wu, Y., Fallon, S. J., Cantin, N. E., and Lough, J. M.: Surface ocean radiocarbon from a Porites coral record in the Great Barrier Reef: 1945-2017, *Radiocarbon*, 63, 1193-1203, <https://doi.org/10.1017/RDC.2020.141>, 2021a.
- Wu, Y., Fallon, S. J., Cantin, N. E., and Lough, J. M.: Assessing multiproxy approaches (Sr/Ca, U/Ca, Li/Mg, and B/Mg) to reconstruct sea surface temperature from coral skeletons throughout the Great Barrier Reef, *Sci. Total Environ.*, 786, 147393, <https://doi.org/10.1016/j.scitotenv.2021.147393>, 2021b.
- Wyndham, T., McCulloch, M., Fallon, S., and Alibert, C.: High-resolution coral records of rare earth elements in coastal seawater: biogeochemical cycling and a new environmental proxy, *Geochim. Cosmochim. Acta*, 68, 2067-2080, <https://doi.org/10.1016/j.gca.2003.11.004>, 2004.
- Xiao, H., Deng, W., Wei, G., Chen, J., Zheng, X., Shi, T., Chen, X., Wang, C., Liu, X., and Zeng, T.: A Pilot Study on Zinc Isotopic Compositions in Shallow-Water Coral Skeletons, *Geochem. Geophys. Geosy.*, 21, e2020GC009430, <https://doi.org/10.1029/2020GC009430>, 2020.
- Yokoyama, Y., Webster, J. M., Cotterill, C., Braga, J. C., Jovane, L., Mills, H., Morgan, S., Suzuki, A., and the, I. E. S.: IODP Expedition 325: Great Barrier Reefs Reveals Past Sea-Level, Climate and Environmental Changes Since the Last Ice Age, *Sci. Dril.*, 12, 32-45, <https://doi.org/10.2204/iodp.sd.12.04.2011>, 2011.