

glenglac: A database of global englacial temperatures

Mylène Jacquemart^{1,2}, Ethan Welty³, Marcus Gastaldello⁴, and Guillem Carcanade^{1,2}

¹Laboratory of Hydraulics, Hydrology and Glaciology (VAW), ETH Zurich, Switzerland

²Swiss Federal Institute of Forest, Snow and Landscape Research (WSL), Sion, Switzerland

³World Glacier Monitoring Service (WGMS), Department of Geography (GIUZ), University of Zürich (UZH), Switzerland

⁴Department of Geosciences, University of Fribourg (UniFr), Switzerland

Correspondence: Mylène Jacquemart (jacquemart@vaw.baug.ethz.ch)

Abstract. Measurements of englacial temperatures have been collected since the earliest years of glaciology, with the first measurements dating back to the mid-19th century. Although temperature is a defining characteristic of any glacier – and is notoriously laborious to collect – no effort had yet been made to gather all existing measurements.

In an attempt to make existing ice temperature data more accessible, we present glenglac, a [global database of englacial temperature measurements](#), [global englacial temperature database](#) compiled from 242 literature sources and nine data submissions and composed of 1142163 measurements of depth and temperature from 690 boreholes located on 186 glaciers outside of the ice sheets. Alongside recent compilations for the ice sheets (Løkkegaard et al., 2023; Vandecruix et al., 2023), most published englacial temperature measurements are now readily available to the research community.

10 Here, we review the variety of glacier thermal regimes that have been measured and summarize the spatial, temporal, and climatic coverage of measurements relative to global glacierized area. Measurements of cold and polythermal glacier ice greatly outnumber those of temperate ice. Overall, temperature has been measured in fewer than 1 % of all glaciers, and only 20 % of borehole locations have been measured more than once, highlighting the large potential to investigate changing temperature conditions by repeating past measurements. The database is developed on

15 GitHub (www.github.com/mjacqu/glenglac) and published to Zenodo (<https://doi.org/10.5281/zenodo.13334175>; ?) (<https://doi.org/10.5281/zenodo.13334175>; 11516611; Jacquemart et al., 2025). It consists of four relational tables and detailed [metadata](#). The GitHub repository also provides submission instructions

(including a spreadsheet template and validation tools), in the hopes that investigators can help us keep glenglac complete and current going forward. We hope that glenglac can help improve our understanding of glacier thermal regimes, help refine glacier thermodynamic models, or shed insight into hazardous glacier instabilities in a warming world.

1 Introduction

The englacial temperature is a defining characteristic of any glacier. It influences glacier flow dynamics and subglacial hydrology, it can be a decisive factor for glacier hazards, and it can serve as an archive of past climate.

To illustrate, ice viscosity and deformation rate depend directly on ice temperature (Deeley and Woodward, 1908; Glen, 1954; Cuffey and Paterson, 2010); only temperate (basal) ice permits glacier sliding over the bed (Cuffey and Paterson, 2010); impermeable cold ice can serve as a barrier for water, controlling sub- and englacial water flow and possibly contributing to the formation of hazardous water accumulations within the ice or in glacier sediment beds (Irvine-Fynn et al., 2011; Vincent et al., 2012; Gilbert et al., 2012; Kääb et al., 2018; Gilbert et al., 2018; Jacquemart et al., 2020; Kääb et al., 2021); ~~the warming of a formerly frozen bed can initiate basal sliding that can lead to large ice avalanches (Alean, 1985; Faillettaz et al., 2015; Troilo et al., 2021; Chiarle et al., 2023)~~. Under cold conditions, the variations of temperature with depth are a window into the past evolution of atmospheric temperatures, making englacial temperature changes an important climate variable (e.g., Gilbert et al., 2010). [Ice temperatures and glacier flow regimes also have implications for glacial archaeology, defining how well artifacts can survive in the ice \(Pilø et al., 2023\)](#). Finally, ice temperature records also serve to validate thermo-mechanical glacier models, which are key to an important tool for improving our understanding of glacier systems.

Glaciers are typically categorized either as temperate, cold, or polythermal. Temperate ice is at the pressure melting point, cold ice is below the pressure melting point, and polythermal glaciers contain both cold and temperate ice. The pressure melting point depends mostly on the ice overburden pressure (and to a lesser extent, the presence of air bubbles and other impurities), such that under temperate conditions ice temperature decreases with depth at around $6.5 \times 10^{-4} \text{ }^{\circ}\text{C}$ to $7.5 \times 10^{-4} \text{ }^{\circ}\text{C}$ per meter (Paterson, 1971; Harrison, 1972; Jania et al., 1996). More generally, the englacial temperature is determined by the complex interaction between the surface energy balance, the geothermal heat flux, and internal heating from ice deformation, basal friction, and refreezing of meltwater (Cuffey and Paterson, 2010). Available heat is transferred through the ice [and firn](#) via conduction and advection by ice and water flow. The superposition of these processes can lead glaciers to be fully cold or temperate, or simultaneously contain cold and temperate ice in a wide variety of spatial configurations (Blatter and Hutter, 1991; Irvine-Fynn et al., 2011).

[Ongoing human-driven climate change is leading to substantial changes of englacial temperatures, evidence of which is clear in \(few existing\) repeat measurements. At Dôme du Gouter in the French Alps, for example, a warming of \$1.5 \text{ }^{\circ}\text{C}\$ has been recorded at a depth of 50 m between 1994 and 2017 \(Vincent et al., 2020\). Long-term measurements at Golle Gnifetti \(Swiss Alps\) between 1991 and 2023 reveal the same amount of warming at a depth of 20 m \(Gastaldello and Hoelzle, 2024b\). This warming can have several consequences. For one, meltwater infiltrating cold firn can degrade or destroy the archive of past climatic conditions that can be stored in firn or ice cores \(Mattea et al., 2021; Gabrielli et al., 2016\). In steep terrain, warming at the glacier bed can lead to wide-spread sliding and destabilization of entire glaciers, increasing the probability of very large ice avalanches, similar to the one observed at Altels in Switzerland in 1895 \(Heim et al., 1895; Faillettaz et al., 2011\). Counterintuitively, warming](#)

can also lead some glaciers to cool, because their disappearing firn cover reduces the amount of englacial warming that stems from the latent heat release of refreezing meltwater (Gilbert et al., 2012; Huss and Fischer, 2016; Irvine-Fynn et al., 2011). A similar effect may also occur in areas still covered by firn, if the refreezing meltwater creates impermeable ice layers that prevent water percolation into the firn, thereby (locally) limiting the latent heat release (Vincent et al., 2020).

Measuring englacial temperatures is a laborious process; (deep) ice temperature measurements are therefore comparatively rare. Those that exist were typically collected for one of ~~two~~ several reasons: To gain an understanding of glacier dynamics and englacial temperatures directly (e.g., Agassiz, 1847; Blatter and Haeberli, 1984; Clarke et al., 1984; Copland et al., 2003; Ryser et al., 2013; Gilbert et al., 2010; Vincent et al., 2020; Troilo et al., 2021; Karušs et al., 2022) or in connection with the retrieval of ice cores used to reconstruct past climatic changes (e.g., Thompson et al., 1990, 2018; Kinnard et al., 2006; Schwikowski et al., 2013; Kinnard et al., 2020), or as part of operational glacier monitoring efforts to document current climate change (e.g., Hoelzle et al., 2020). The data resulting from such efforts are largely hidden away in scientific literature spanning more than a century, and are therefore not readily available as a community resource. At the same time, there is an increasing need to understand how englacial temperatures – and their changes over time – relate to glacier dynamics, climate change, and glacier hazards (Käab et al., 2018; Colgan et al., 2015; Gilbert and Vincent, 2013; Gilbert et al., 2015) (Gilbert and Vincent, 2013; Colgan et al., 2015; Gilbert et al., 2015; Käab et al., 2018; Machguth et al., 2023b). In order to make englacial temperature data from glaciers around the world more widely available, we have compiled a database of englacial temperatures glenglac, a global englacial temperature database, sourced – largely, but not exclusively – from published literature. In the following, we describe how englacial temperatures are measured in the field (Sec. 2.1), how we found and compiled these measurements (Sec. 2.2), and how the resulting glenglac database is structured and managed (Sec. 2.3). In Section 3, we present and discuss the content of glenglac (version 1.0), and close with instructions for how others can contribute additional data (Sec. 3.5). We hope that glenglac can serve as a community resource to help improve our understanding of ongoing changes in the cryosphere and that it will grow over time with the addition of past and future englacial temperature measurements.

2 Methods and Data

2.1 Measurement methods

Englacial temperature measurements are typically made by placing one or more thermistors in a borehole (Fig. 1). Boreholes are drilled with either mechanical (reviewed in Talalay, 2016) or thermal (reviewed in Talalay, 2020) drills, or a combination thereof. The advantage of mechanical drilling is that temperatures can be reliably measured after a short time. Thermal drilling (e.g., steam or hot water) significantly raises the temperature of the borehole, which subsequently needs time to re-adjust to the temperature of the surrounding ice (typically a several days to a few weeks Laternser, 1992; Miles et al., 2018). Depending on the measurement techniques and objectives, borehole temperatures are measured only once, or the thermistor chain is left in the hole (which is allowed to freeze up or



Figure 1. Photographs illustrating methods for measuring glacier temperatures: (a) drilling a shallow borehole with a handheld mechanical auger on Griesgletscher, Switzerland (photo by Matthias Huss), (b) drilling a deep borehole with a hot water drill and a large array of fuel drums and pumps to heat and pressurize the water on Rhonegletscher, Switzerland (photo by Raphael Moser), and (c) a typical string of thermistors used to measure temperature at different depths in a borehole (photo by Mylène Jacquemart)

kept open with a casing or fluid) and either remeasured manually or equipped with an automatic logger. A recent
90 innovation replaces discrete thermistors with a fibre optic cable (distributed temperature sensing (DTS), Law et al., 2021), which can provide measurements with an unprecedented vertical resolution, especially in deep boreholes where the required number of thermistors would be prohibitively expensive.

2.2 Data compilation

Most data included in glenglac are sourced from published literature and datasets. Publications were initially found by
95 searching Google Scholar (<https://scholar.google.com>) and Google Dataset Search (<https://datasetsearch.research.google.com>) for combinations of the English words glacier, ice, temperature, thermal, regime, englacial, borehole, drill(ing), measurement, and record, as well as 冰川温度钻孔 and скважина температуры ледника ("glacier bore-hole temperature" in Chinese and Russian, respectively). We then recursively sought out publications referenced
100 in previously-identified publications, striving to find the most complete and original data source for each measurement. Tracking down references was made more difficult by the practice of journals (primarily European and North American ones using Latin script) of not publishing references in their original form (e.g., "термический"), but only translations ("temperature") or phonetic transcriptions ("termicheskiy"). Finding these often involved reconstructing the original reference, since the modified form did not appear in search results. When we were unable to find the full text of a publication online or in print in nearby libraries, we requested it from the Swiss Library Service

105 Platform (SLSP) document delivery service through the University of Bern (ub.unibe.ch). In addition to extracting data from publications, we solicited data submissions on CRYOLIST (cryolist.org), at the 2023 Alpine Glaciology Meeting in Birmensdorf, Switzerland, and through personal communications. References to all the data sources can be found in the bibliography of this publication (see Appendix A for a list of glaciers and corresponding references) and within glenglaf itself (Sec. 2.3).

110 For this first version of glenglaf, we selected only firn and/or ice temperature measurements with a well-defined depth. This means that we did not take into account measurements made in tunnels dug into glaciers at undefined depths. We ~~largely omitted shallow measurements known to be only disregarded publications that provided measurements that were made exclusively~~ in seasonal snow (~~though occasionally including them if found alongside deeper measurements~~), but did not remove shallow measurement points of deeper measurements in firn or ice. 115 Where shallow measurements were provided alongside deeper ones, we did also include these. We also focused on glaciers and omitted measurements from the Antarctic and Greenland ice sheets, ~~in large part to a large degree~~ because data from the ice sheets have already been compiled by Løkkegaard et al. (2023) and Vandecrux et al. (2023).

120 For each measurement, we extracted depth and temperature data and their associated metadata (see below). Submitted or published data in machine-readable formats were added directly to the database, data published numerically in text or tables were transcribed with the help of Optical Character Recognition (OCR), and data represented graphically (e.g., as a plot of temperature versus depth) were digitized using the open-source software Plot Digitizer (<https://plotdigitizer.sourceforge.net/>). For plots that used points to indicate each measurement (Fig. 3a), we digitized the values at each point. For plots using a continuous line (Fig. 3b), such that the locations of the 125 measurements along the line are unknown, we used a point density sufficient to reproduce the original line.

130 Metadata associated with the temperature measurements were compiled from one or more sources to the best of our abilities (see Tables 1 – 4). For example, borehole coordinates were either extracted directly from text, digitized from a map with defined axes (e.g., latitude and longitude) using Plot Digitizer, digitized from a map (without such axes) visually georeferenced to a global basemap ~~using the~~ QGIS Freehand Raster Georeferencer plugin (<https://plugins.qgis.org/plugins/FreehandRasterGeoreferencer/>), or approximated on global basemaps with the help of terrain features, glacier morphology, or elevation information. Borehole elevation was published as text in most cases, though we did occasionally have to approximate elevation from provided contour lines or by comparison to an independent topographic map. Other metadata included the measurement and drilling dates, the drill type (mechanical, thermal, or combined), the depth of the firn-ice interface (i.e., whether the measurements are in firn, 135 ice, or both), whether the borehole reached the glacier bed, and the ~~accuracy uncertainty~~ of the temperature measurements. ~~Some columns were added later in the data compilation process, therefore not all fields are equally well populated.~~




Figure 2. Glenglat testing and publishing workflow. The input files (in gray) are those version-controlled in the Git repository.

2.3 Data structure and management

Glenglat is packaged and described following the Frictionless Tabular Data Package standard (Walsh et al., 2017),
140 version-controlled and tested on GitHub (<https://github.com/mjacqu/glenglat>), and published to Zenodo (<https://doi.org/10.5281/zenodo.13334175>). Data and metadata (Tables 1–4) are stored using common text file formats to ensure that they are human-readable, machine-actionable, and compatible with line-based version-control systems like Git.

The data are structured as a four-table relational database stored as CSV (comma-separated values) files (Tables 1–4). The source table (Tab. 1) contains a full reference to each data source that we used. Names in non-Latin scripts (Cyrillic, Hangul, and Chinese characters) are accompanied by a latinized form, and non-English titles are accompanied by an English translation. The borehole table (Tab. 2) describes the drill site – including the location,

elevation, drill method, and reported ~~accuracy~~uncertainty of the temperature measurements. The profile table (Tab. 3) describes each temperature-depth profile – including the timing of the measurement and whether or not the measurement was made after the borehole was known to have reached thermal equilibrium. Finally, the measurement table (Tab. 4) contains the measurements of temperature with depth. To improve manageability of the CSV files, data from boreholes with time-series measurements (i.e., with hundreds of profiles), the profile and measurement tables are stored in separate, source-specific CSV files. Supporting these tabular data are files that, for each source, document how and from where the data were extracted. For submissions, these include data files and email correspondence. For publications, these include the key text passages, tables, maps, or figures that served as the sources for the data. Additional files ~~document~~can be used to reproduce how numeric values were digitized from maps and figures using Plot Digitizer (*.xml) or georeferenced and digitized using QGIS (*.pgw, *.{png|jpg}.aux.xml, and *.geojson).

The tabular data are described in a single YAML (<https://yaml.org>) metadata file (datapackage.yaml). This file lists general attributes of the database – like name, description, version, license, and contributors – as well as a detailed description of the structure and content of each tabular data file. The CSV Dialect (Pollock, 2021) specifies how exactly the CSV files are structured, while the Table Schema (Walsh and Pollock, 2021) specifies the name and data type of each column, the constraints on each column's values, and the foreign-key relations between tables.

This metadata architecture allows data maintainers and contributors to use Frictionless Framework (<https://framework.frictionlessdata.io>) to test that the metadata is correctly structured and that the data are consistent with the metadata. These tests are run automatically in a continuous-integration pipeline on GitHub using Frictionless Repository (<https://repository.frictionlessdata.io>), ensuring the integrity of the database whenever any file is modified. Additional custom tests, which cannot be expressed by the metadata, further verify the integrity of the dataset, for example that all people who contributed data (referenced as a personal communication in table source) are listed as contributors in datapackage.yaml. Using software built on the Frictionless Tabular Data Package standard (<https://github.com/ezwelty/tablecloth>), we can also render the metadata as an interactive spreadsheet template with dropdown menus and real-time validation, lowering the bar for future data contributors.

We publish the database to Zenodo using a custom build process. The uploaded file archive and detailed Zenodo metadata are generated automatically from the contents of the GitHub repository and submitted using the Zenodo REST API (<https://developers.zenodo.org>). Zenodo manages the DOIs for glenglat, registering a concept DOI encompassing all versions (<https://doi.org/10.5281/zenodo.11516611>) and a version DOI for each new version (<https://doi.org/10.5281/zenodo.13334175>). To keep the structure simple and the download small, Zenodo releases contain only the CSV files (data/*.*), a license (LICENSE.md), simplified documentation (README.md), and a version of the metadata (datapackage.yaml) converted to JSON.

Table 1. Main columns of the source table (data/source.csv); a full description is provided in datapackage.yaml. Column names and categorical values closely match the Citation Style Language (CSL) 1.0.2 specification (Zelle et al., 2015). The primary key is indicated with a *.

Column	Type/Units	Description
id*	string	Unique identifier, constructed from author name and year (e.g., zagorodnov1981), referenced from other tables either formally in a foreign key or informally within free-form text.
author	string	Author names (optionally followed by their ORCID in parentheses) as a pipe-delimited list.
year	YYYY	Year issued (published, communicated, last updated)
type	string	Type (e.g., journal article, book chapter, dataset, personal communication)
title	string	Title of the work
url	string	URL (DOI if available)
language	string	Language as ISO 639-1 two-letter language code.
container_title	string	Title of the container (e.g., journal, book, data repository)

Table 2. Columns of the borehole table (data/borehole.csv). Primary keys are indicated with a *, foreign keys with a °.

Column	Type / Units	Description
id*	integer	Unique identifier
source_id°	string	Identifier of the source of the earliest temperature measurements (and the source of all borehole metadata, unless otherwise stated in notes).
glacier_name	string	Glacier or ice cap name (as reported).
glims_id	string	Global Land Ice Measurements from Space (GLIMS) glacier identifier.
location_origin	string	Origin of location (latitude, longitude): <ul style="list-style-type: none"> - submitted: Provided in data submission - published: Reported as numbers in original publication - digitized: Digitized from published map with complete axes - estimated: Estimated from published plot by comparing to a map - guessed: Estimated with difficulty (e.g., by comparing elevation to a map)
latitude	degrees	Latitude in the EPSG 4326 spatial reference system.
longitude	degrees	Longitude in the EPSG 4326 spatial reference system.
elevation_origin	string	Origin of elevation; same categories as for location_origin.
elevation	meters	Elevation above sea level of the drilling site.
mass_balance_area	string	Mass balance area: Ablation area, near the equilibrium line, or accumulation area
label	string	Borehole name (e.g., as labeled on a plot).
date_min	YYYY-MM-DD	Begin date (or first possible date) of drilling (e.g., 2019: 2019-01-01).
date_max	YYYY-MM-DD	End date (or last possible date) of drilling (e.g., 2019: 2019-12-31).
drill_method	string	Drilling method: mechanical, thermal, or combined.
ice_depth	meters	Starting depth of ice. Infinity (INF) indicates that ice was not reached.
depth	meters	Total borehole depth (not including drilling in the underlying bed).
to_bed	boolean	Whether the borehole reached the glacier bed.
temperature_uncertainty	degrees Celcius	Estimated temperature uncertainty (as reported).
notes	string	Additional remarks about the study site, the borehole, or the measurements therein.
curator	string	Names of people who added the data to the database, as a pipe-delimited list.
investigators	string	Names of people and/or agencies who performed the work, as a pipe-delimited list.
funding	string	Funding sources as a pipe-delimited list.

Table 3. Columns of the profile table (data/profile.csv and data/**/profile.csv). The ** indicates subdirectories that hold separate profile.csv and measurement.csv files from boreholes with many profiles (e.g., from automated loggers). The subdirectories are labeled source.id-glacier, where glacier is a simplified version of the glacier name (e.g. flowers2022-little-kluane). (Composite) primary keys are indicated with a *, foreign keys with a °.

Column	Type / Units	Description
borehole_id*°	integer	Borehole identifier.
id*	integer	Borehole profile identifier (starting from 1 for each borehole).
source_id°	integer	Source identifier.
measurement_origin	string	Origin of measurements: - submitted: Provided as numbers in data submission - published: Numbers read from original publication - digitized-discrete: Digitized from discrete points of depth versus temperature - digitized-continuous: Digitized from a continuous line plot
date_min	YYYY-MM-DD	First possible date of measurement (e.g., 2019: 2019-01-01).
date_max	YYYY-MM-DD	Last possible date of measurement (e.g., 2019: 2019-12-31).
time	hh:mm:ss	Measurement time.
utc_offset	hours	Time offset relative to Coordinated Universal Time (UTC).
equilibrated	boolean	Whether temperatures have equilibrated following drilling.
notes	string	Additional remarks about the profile or the measurements therein.

Table 4. Columns of the measurement table (data/measurement.csv and data/**/measurement.csv). The ** indicates subdirectories that hold separate profile.csv and measurement.csv files from boreholes with many profiles (e.g., from automated loggers). The subdirectories are labeled source.id-glacier, where glacier is a simplified version of the glacier name (e.g. flowers2022-little-kluane). (Composite) primary keys are indicated with a *, foreign keys with a °.

Column	Type / Units	Description
borehole_id*°	integer	Borehole identifier.
profile_id*°	integer	Borehole profile identifier.
depth*	meters	Depth below the glacier surface.
temperature	degrees Celsius	Measured temperature.




Figure 3. Screenshots of the digitization process with Plot Digitizer, where temperature versus depth is either plotted as (a) discrete points for each measurement or (b) a continuous line with unknown measurement locations. Data is from (a) 古里雅冰帽 (Guliya Ice Cap, GLIMS ID G081455E35226N) and (b) 天山1号冰川 (Urumqi Glacier No. 1, GLIMS ID G086810E43111N).

2.4 Errors from digitization and data reproduction

- 180 In addition to the errors uncertainty of the original measurement measurements, errors are introduced when the measurements are reproduced in tables or figures (henceforth "reproduction error"), and again when these reproductions are digitized (henceforth "digitization error"). Such errors can multiply if the data is shared between researchers or digitized from older publications and reprinted in subsequent publications. In 80 cases, we acquired the same temperature profile from two different sources (e.g., data submission and published figure, published figure and published table, published figure replaced by a data submission, table, or better figure), allowing us to assess the magnitude of reproduction errors (inclusive of digitization errors). In order to quantify the digitization error in isolation, 177 temperature profiles were digitized by two different people. This exercise also allowed us to refine our method by identifying avoidable human errors and software quirks (e.g. wrong scaling of an axis, misplaced points). We calculated both these errors from profile pairs as the difference between their temperatures after interpolating (but never extrapolating) the temperatures of one to match the depths of the other.
- 190

3 Results and Discussion

As far as we know, glenglac is the largest collection of englacial temperature measurements. It contains 1142163 measurements of depth and temperature, organized into 147583 profiles from 690 boreholes (Fig. 4). We included




Figure 4. Spatial distribution of temperature measurements recorded in glenglat. Boreholes are plotted in red, glaciers in light blue (according to the Randolph Glacier Inventory 7.0; RGI Consortium, 2023) in light blue. The Global Terrestrial Network for Glaciers (GTN-G) Glacier Regions (GTN-G, 2017) are shown in grey and numbered; these correspond to the region numbers in Fig. 6.

17873 profiles (for 79 boreholes) from nine data submissions. The remaining data were extracted from 175 primary
195 sources (see Tab. A1), with an additional 66 secondary sources helping to further populate the metadata. Non-English
sources make up 28 % (49) of all primary sources but 40 % (40) of those published before the year 2000.

3.1 Thermal regimes and borehole depths

A variety of thermal structures can be identified in the temperature profiles (Fig. 5). The borehole from the Devon
Ice Cap (Nunavut, Canada; data from Paterson and Clarke, 1978) is an example of fully cold conditions. At depth,
200 the temperature increases at a rate largely determined by the geothermal heat flux (heat conduction from the
Earth's interior). In contrast to the fully cold conditions, profiles can be fully temperate, such as on Hansbreen
(e.g., Svalbard; data from Jania et al., 1996) where the ice temperature decreases with depth in accordance with the
lowering of the pressure melting point. Between these two endmembers, there is a lot of variety. At Grenzgletscher




Figure 5. Measured profiles demonstrating the range of englacial temperatures and some typical profile shapes – from fully cold ([Devon Ice Cap](#)) ([Devon Ice Cap Paterson and Clarke, 1978](#)) to fully temperate ([Hansbreen](#)) ([Hansbreen Jania et al., 1996](#)). Measurements from accumulation areas are plotted with solid lines, those from ablation areas with dotted lines.

(Switzerland; data from Ryser et al., 2013; Hoelzle et al., 2011) and White Glacier (Axel Heiberg Island, Canada; 205 data from Blatter, 1987), for example, the ice in the accumulation area is colder than the ice in the ablation area, indicating that the cold ice is warmed (e.g., by shear heating and latent heat release) as it advects down from the accumulation area. The opposite is true at Kongsvegen (Svalbard; data from Björnsson et al., 1996), where the 210 latent heat release of refreezing meltwater and precipitation is large enough to eliminate the winter cold wave and create temperate firn and ice in the accumulation area. In the ablation area lower on the glacier, meltwater and precipitation can run off, allowing the near-surface ice to cool into a layer of cold ice superimposed on the temperate ice.

The temperatures measured in the boreholes range from temperate (i.e., at the pressure melting point) to -33.5°C (the lowest temperature at 15 m depth is -31.7°C). The majority (75%) of boreholes deeper than 15 m are in cold




Figure 6. Overview of borehole counts (all and those deeper than 15 m) and surface elevation, 15 m temperature, and maximum measured depth for each borehole (each represented by a dot) by region (see Fig. 4).

or polythermal ice (defined as those where the maximum measured temperature is colder than -0.5°C), with only about 7 % of all boreholes showing fully temperate conditions (i.e., the lowest measured temperature is warmer than -0.5°C). This is not surprising. For one, temperate (or even partially temperate) ice is of little interest to ice-core investigations because it does [not](#) retain a memory of past climatic conditions ([though research for dating temperate cores is ongoing](#)).

Secondly, temperate ice measurements are deemed less interesting, therefore, englacial temperature measurements are rarely carried out on glaciers that are assumed to be fully temperate, and if temperate conditions are measured, the results are rarely published. However, such measurements would be very valuable to train and calibrate models that predict glacier thermal regimes at regional to global scales. At present, it may be possible to train a model to accurately identify cold or polythermal glaciers, but it would be harder to constrain the boundaries – in terms of elevation, latitude, mean air temperature, etc. – between cold and temperate ice.

The median of the maximum-measured-depths in all boreholes in glenglac is 22 m (see Fig. 6). A total of 485 boreholes (70 %) were measured at depths greater than 15 m, and 148 (21 %) deeper than 100 m. Only 134 (19 %) are known to have reached the glacier bed, including the deepest, an 724 m ice core borehole drilled on Ледник Академии Наук (Akademii Nauk Ice Cap, Severnaya Zemlya, Russia; Kotlyakov et al., 2004).

Of 165 boreholes with a reported ice surface depth (an attribute that was added later and thus likely incomplete with respect to the available literature), 92 actually reached ice, whether at 0 m depth (for 68 boreholes) or below a maximum of 34 m of snow and firn on 慕士塔格冰川 (Muztagh Glacier, Xinjiang, China 李真[Li Zhen] et al., 2004). Unfortunately, whether a borehole was drilled into snow, firn, ice, or a combination thereof is not always known.

For applications where this is relevant, the stratigraphy can often be estimated – if not explicitly reported – from the depth of the borehole and its location on the glacier (e.g., ablation or accumulation area). Further complicating borehole stratigraphy, repeat measurements in the same borehole typically define depth relative to the glacier surface at the time of the initial drilling or thermistor installation, despite accumulation and ablation that may occur in the interim. This can lead to spurious and above-zero temperature measurements if thermistors melt out over time. Only in very rare cases (e.g., in Harrison et al., 1975) are the changes of the surface elevation recorded in detail.

3.2 Climatic conditions

Compared to the average climatic conditions at the locations of all glaciers in the Randolph Glacier Inventory 7.0 (RGI Consortium, 2023), the locations selected for englacial temperature measurements are biased towards cold and dry conditions (Fig. 7a). Most boreholes are in locations where the total annual precipitation is less than 1 m yr^{-1} and the mean annual air temperature is below -5°C . This focus on regions with a continental climate is again not surprising, because the high accumulation rates and warmer air temperatures of maritime climates are more likely to lead to temperate ice – which is considered less interesting – or high ice fluxes – which are not desirable for ice core measurements (e.g., Vance et al., 2016; Bohleber, 2019).

To explore the controls on englacial temperatures, we take the 15 m temperature – the depth at which seasonal temperature variations have mostly disappeared (see Fig.8) – as an indicator of the local glacier thermal regime. Comparing ~~these borehole temperatures~~ borehole temperatures measured since 1960 to the mean annual air temperature ~~taken from (Muñoz Sabater, 2019)~~ (MAAT) (ERA5-Land from Muñoz Sabater, 2019) of the ten years prior to the borehole measurement reveals that temperate ice can occur over a wide range of surface air temperatures (Fig. 7b). Englacial temperatures generally ~~increase with increasing decrease with decreasing~~ decrease with decreasing air temperature, but they are also consistently warmer than the temperatures at the surface. This is expected, given the numerous processes that can deliver heat into the glacier (latent heat release and geothermal, frictional, and strain heating). This warm bias is smallest at the coldest surface temperatures (presumably because there is little melt) – but can still be up to $+10^\circ\text{C}$ at -20°C . It reaches a maximum of $+15^\circ\text{C}$ at -15°C , before gradually decreasing towards warmer temperatures only because ice cannot be any warmer than the melting-point.

Temperate ice does not seem to occur where the mean annual air temperature is below $\sim -15^\circ\text{C}$. It is unclear, however, whether the absence of temperate ice in colder climates is real or due to the undersampling of temperate glaciers.

The trends and boundaries in Fig. 7b need to be evaluated with respect to possible caveats of the chosen dataset, and are intended only to show broad patterns. ERA5-Land (and other reanalysis products) may not be able to represent the true variability of temperature fields or precipitation patterns in complex alpine terrain. The observed 'warm-bias' could therefore partly be due to biases within this dataset, and the 2-m air temperature is in reality not the true surface boundary condition. Beyond this, the chosen lapse-rate (in our case $-6.5^\circ\text{C km}^{-1}$) changes the relationship between air temperature and englacial temperature, though we found that it mostly affects the




Figure 7. a) Distribution of 2000–2019 mean annual 2 m air temperature and total annual precipitation for the locations of glenglac boreholes (red) and locations of all RGI 7.0 glaciers (blue). b) Englacial temperatures ([1950–1960](#) to present) at 15 m depth versus mean annual air temperature at the borehole location for the ten years prior to the borehole measurement. Air temperatures were adjusted to the elevation of the borehole using a lapse-rate of $-6.5^{\circ}\text{C km}^{-1}$. The climate data is from ERA5-Land (Muñoz Sabater, 2019).

positions of the data points relative to the 1:1 line, while the overall shape remains stable. In reality, of course, englacial temperatures are controlled by much more than mean annual air temperature and precipitation, and the consideration of 15 m temperatures [further](#) ignores that: i) temperate ice can exist at deeper locations within the ice even under very cold conditions (e.g., due to shear heating Blatter, 1987), ii) ice temperatures can be controlled by glacier dynamics more than by climate (e.g., emergence of cold ice in the ablation area of a glacier in a temperate climate Ryser et al., 2013), and iii) ice at depth has a memory of past surface temperatures, which can lead to complex patterns of englacial temperatures that are not reflected in the 15 m temperature. Despite these caveats, the large number of englacial temperature measurements available in glenglac makes it possible, for the first time ever, to investigate global patterns of englacial temperatures and hopefully find more robust ways of predicting the thermal regime of all glaciers in a region or worldwide.

3.3 Spatial, temporal, and elevation distributions

The 690 boreholes in glenglac are located on 186 individual glaciers (based on their GLIMS IDs) scattered across the world (see Figs. 4 and 6). This represents less than 1 % of all glaciers worldwide, illustrating both how laborious englacial temperature measurements are ([see Appendix A](#)) and how interest in this glacier variable remains relatively




Figure 8. Examples of temperature profiles measured in the same borehole at different times of year, showing the elimination of seasonal surface temperature variations at a depth of around 15 m. Data from Vavilov Ice Cap (H. И. Барков [N. I. Barkov] et al., 1988), Austre Lovénbreen (孙维君[Sun Weijun] et al., 2016), Blichenok Glacier (Shiraiwa et al., 2001).

280 limited. Two thirds of all boreholes (68 %) are in either Arctic Canada (84), Svalbard and Jan Mayen (80), Central Europe (191 – albeit with 79 on a single glacier: Grenzgletscher), and South and Central Asia (112). Conversely, there are only five boreholes in all of Scandinavia or the Southern Andes, and none in Iceland or New Zealand.

If we envision using global or regional models to constrain thermal regimes spatially, by elevation, by temperature or precipitation regime, or based on glacier dynamics, we ideally need measurements that span the transition and range of conditions where glaciers exist. In this context it is important to understand what data is present in a possible training dataset. In the following, we therefore briefly discuss the temporal and spatial patterns of the data in glenglac.

The surface elevations of boreholes in glenglac range from 25 m (Erikbreen, Svalbard; Ødegård et al., 1992, (elevation above sea level assumed henceforth) to 7200 m asl (Erikbreen, Svalbard; Ødegård et al., 1992) to 7200 m asl (Dasoupu Glacier, China; Yao et al., 2002) (elevation above sea level assumed henceforth). Compared to the elevation distribution of glaciers worldwide (as represented by RGI 7.0), elevations above 2000 m are oversampled while elevations between 750 m and 1500 m are undersampled (Fig. 9). This sampling bias may have several causes. Topographic saddles and summits of very high elevation glaciers (> 4500 m) are of particular interest to ice core science because ice flow, accumulation, and melt are minimal and a maximum number of annual ice layers can be preserved. The middle elevations (2000 m to 4500 m) are likely oversampled because they circumvent many of the challenges of work at very high elevations, allowing for simpler (helicopter) logistics and reasonable working conditions. Additionally, this is the elevation range of glaciers




Figure 9. Elevation distribution of glenglac boreholes (red) compared to the elevation distribution of all RGI 7.0 glaciers (blue). Higher elevations, especially between 2000 m and 4500 m, are oversampled while low elevations are undersampled.

in Central Europe (region 11, see Figs. 4 and 6), which are historically overstudied compared to other regions. Below 300 2000 m, most of the data comes from Svalbard and Jan Mayen (region 7) and Arctic Canada (regions 3-4). Though there are many measurements from these regions, lower elevations remain undersampled, likely because i) there is a very large glacierized area in this elevation band, ii) there is lack of interest in measurements from temperate ice (e.g., the large low-lying glaciers along the west coast of Alaska), iii) the tongues of tidewater glaciers – which make up a large portion of this band – are notoriously difficult to access, and iv) when accessing land-terminating glaciers in these remote regions, working on their low-elevation tongues is easier than accessing the higher accumulation areas (750 m to 1500 m).

305 The earliest measurement in glenglac stems from 1842, when Louis Aggasiz and colleagues drilled a 60 m borehole in temperate ice on Unteraargletscher (Agassiz, 1847) in Switzerland (see Fig. 10). With the exception of other early outliers (1911, Vallot (1913); 1938, Hughes and Seligman (1939); 1948, Sharp (1951)), wide-spread measurement does not begin until the late 1950s – to a large part motivated by the 1957/1958 International Geophysical Year – by 310 when drilling technology and motorized transport were developed enough to allow increasingly ambitious expeditions to remote areas. Mid-century measurements likely remain underrepresented in glenglac because of early and obscure publications that were never indexed or published online. After sustained activity since the 1970s, measurements decline beginning in 2015, which may be the result of a lag between data collection and publishing, reduced interest and funding for complex field campaigns, a shift of focus from glaciers to ice sheets, or increased emphasis on modeling and remote sensing.

315 In only 156 boreholes (20 %) were temperatures measured more than once. The most frequently measured borehole is on Hintergrat Glacier (Italy; Carturan et al., 2023a), with five years of hourly measurements (2011-2016). The longest monitored borehole is CG05-1 on Grenzgletscher (Switzerland; Hoelzle, 2014; Darms, 2009; Hoelzle, 2017), drilled in 2005 and measured 4 times over 8 years (2007-2015). The lifetime of a single borehole is limited by creep closure, internal deformation, and other forces that inevitably lead to equipment failure, so to achieve longer records,




Figure 10. Number of measured boreholes (top) and number of measured glaciers (bottom) for each year. We assume that early measurements are still underrepresented, because some publications were not archived. The drop-off towards more recent years may be due to the lag between drilling and publication.

320 a new borehole is drilled and instrumented nearby. A cursory review of boreholes within \sim 100 m reveals only a few locations with multi-borehole records spanning more than 20 years, all of which are the result of deliberate repeat studies (Vincent et al., 2020; Hoelzle et al., 2011; Thompson et al., 2018; B. H. Михаленко [V. N. Mikhaleenko] et al., 2005b; Rabus and Echelmeyer, 2002), although clusters of more distant boreholes suggest opportunities for retroactive comparisons. The wide range of measurement dates in glenglat could present a challenge for training
325 or calibrating numerical models, as it requires longer model runs and inputs (e.g., climate reanalysis) from earlier periods with less and lower-quality data.

Conversely, the low percentage of boreholes that have been measured more than once indicate that there is a large potential for repeat measurements that would yield insight into how englacial temperatures have changed.

330 An identification procedure for high-value sites for repeat measurements could focus on sites with deep boreholes (e.g., more than 30 m), predominantly cold englacial temperatures, high measurement accuracies (low reported uncertainties), and a last measurement dating back around twenty years or more. Ignoring surge-type glaciers, a search of glenglat reveals high-value repeat-measurement sites in many different regions and climates. Some examples that fall into this category include Illimani Volcano (Bolivia, 138 m, measured in 1999), Vavilov Glacier (Russian Arctic, up to 460 m, last measured in 1985), or Åsgårdfonna (Svalbard, 185 m, measured in 1993).

335 3.4 Error analysis

In our analysis, we distinguish between three sources of uncertainty in the reported englacial temperatures: the reported uncertainty of the original measurements, errors introduced by their subsequent textual or graphical reproduction, and errors associated with the digitization of graphical reproductions.

340 The reported temperature accuracy addresses measurement error (mean 0.14°C), but not the additional errors
that may have crept in when the data was reproduced or when it was finally digitized for glenglac (

The measurement uncertainty encompasses a wide range of instrumental, methodological, and environmental factors. The design and calibration of a thermistor determines the precision and accuracy with which it measures temperature. Whether the measured temperature differs from the temperature of the glacier is primarily determined by the extent to which the borehole reached thermal equilibrium following drilling and sensor installation, or the method by which an equilibrium temperature is extrapolated from a sensor cooling curve. Secondary sources of thermal disturbance include "wind pumping" (air circulation within firn or snow caused by surface pressure changes; Clarke et al., 2013), sensor electronics (Ryser et al., 2013), and artificial thermal coupling with the surface (e.g. the black tubing used in Davis, 1966). Finally, the uncertainty in the measurement depth translates to an uncertainty in the temperature at that depth, especially in areas with steep temperature gradients (e.g., near the surface).

350 Temperature measurement uncertainty is reported for 65 % of the boreholes included in glenglac (0.01°C to 1.0°C ; see Fig. 11). Comparing It is rarely specified which sources of error the value represents, but sensor design and calibration are often the only factors discussed. Other glenglac attributes like drilling method, drilling and measurement dates, whether and how equilibrium was reported to have been reached, and repeat measurements from the same profile retrieved from different sources borehole are thus helpful for independently assessing measurement uncertainty. Contrary to the suggestion of Løkkegaard et al. (2023) that sensor accuracy has improved substantially since 1950, we found no significant change in reported temperature uncertainties over time. This could reflect increased caution in error reporting, an acknowledgment that sensor accuracy is only one component of the total uncertainty, or indeed that the accuracy of the typical englacial temperature measurement has not improved significantly.

360 Depth measurement uncertainty is so rarely reported that no provision was made to track it in glenglac. In many cases, it is also not specified from which reference surface depth was measured or whether this reference changed over time (e.g., published table versus published figure), we find that reproduction errors (standard deviation 0.48°C) are on par with the reported measurement errors at depths larger upper glacier surface at the time of installation, snow-ice boundary at the time of measurement). As the glacier surface changes, so does the depth of the sensors relative to this surface, and this can complicate the comparability of measurements through time.

370 Fig. 11 presents the reported measurement uncertainties alongside errors introduced by data reproduction and digitization. For the sake of comparison, we describe each distribution by the 0.5 (median), 0.68, and 0.95 quantiles (of the absolute values). For measurement uncertainties, these are 0.1°C , 0.2°C , and 0.3°C . Reproduction errors were calculated as the difference between temperatures for the same profile retrieved from a figure versus either a table or another figure (and thus include digitization error). They are typically smaller than measurement uncertainties (0.08°C , 0.14°C , 0.57°C), especially at depths greater than 15 m (standard deviation 0.18°C , median absolute deviation 0.06°C). Because of the steep temperature gradients 0.05°C , 0.11°C , 0.33°C). The largest of these are due to different measurements being included (the impact of which is greatest near the surface, the comparison of

different sources can be heavily influenced by whether and which near-surface measurements were included in each source. Comparing the result of where temperature gradients are the steepest), actual differences in the reported temperatures, or poor figure quality (e.g. non-square axes from hand drawing or bad scanning, high °C/pixel scale). Random digitization errors (isolated from reproduction errors) were calculated as the difference between temperatures for the same profile and figure digitized by two different people digitizing the same measurements; we find that digitization errors are even smaller (standard deviation 0.132 °C, median absolute deviation 0.025 °C) so as to be negligible. Overall, this indicates that the data available through glenglata is suitable for quantitative analysis as data collected directly, even though they have been drawn from one (. They are smaller (0.03 °C, 0.05 °C, 0.23 °C) and sensitive to the interaction between figure scale and temperature gradient. The °C/pixel size of the figures' digitized temperature axes – an imperfect proxy for their effective resolution – range from 0.002 °C to 0.109 °C (median 0.016 °C) and are weakly correlated with the digitization errors ($R^2 = 0.058$, slope = 1.29). However, the m/pixel size of the depth axis – once converted to °C based on the local slope of the profile – range from 0.000 °C to 1.007 °C (median 0.007 °C) and explain the observations much better ($R^2 = 0.483$, slope = 1.26).

A full assessment of uncertainty would require a careful evaluation of each profile (e.g. Løkkegaard et al., 2023), including a review of the original sources, since these often contain more details than could be integrated into glenglata. However, we can provide some general guidance to users: Favor data with low reported measurement uncertainties sourced from textual reproductions (measurement_origin "submitted" or "published"), inspect the source of digitized data (measurement_origin "digitized-discrete" or several iterations of) publications. "digitized-continuous"), favor profiles known to have reached thermal equilibrium (equilibrated), and avoid interpolation at depths with steep temperature gradients.

Reducing and quantifying uncertainties will be crucial for detecting and correctly interpreting the long-term thermal evolution of glaciers and ice sheets, so we urge future campaigns to carefully assess uncertainties and thoroughly and precisely describe their measurements. To eliminate reproduction and digitization errors in the future, new and existing measurements should be published somewhere as text, for example by submitting them directly to glenglata (see Sec. 3.5).

3.5 Future additions

We hope that glenglata can serve not only as a valuable resource for glaciological research today but also as a long-lived data repository for additional (past) past and future englacial temperature measurements. The dataset is currently hosted at <https://github.com/mjacqu/glenglata>. To encourage and facilitate submissions, we have included detailed instructions, a Microsoft Excel spreadsheet template, and a tutorial showing how to self-validate the data prior to submission using the Frictionless Python package. We have done our best to populate all fields of the database, but certain columns were added at a later stage of the database creation process, and we were not able to revisit every source every time. Therefore, not all columns are equally well populated. Additions to or refinements of existing entries are welcome any time, either by emailing the authors or by creating an issue at



Figure 11. Distribution of [the reported measurement uncertainties alongside distributions of the absolute values of the](#) reproduction errors (comparing the same measurement from different sources) and digitization errors (comparing the digitization of the same measurement by two different people). [These errors are on par with the reported accuracy of the original measurements.](#) The y-axis uses square-root scaling to accentuate differences near 0 °C. [For each distribution, horizontal lines are drawn at the 0.05, 0.5 \(median\), and 0.95 quantiles.](#)

410 <https://github.com/mjacqu/glenglat/issues>. Community members are also welcome to take on existing issues and contribute to the improvement of the dataset in this way. Anyone who submits data to glenglat will be invited to become a co-author on future releases of the dataset (see detailed authorship policy at <https://github.com/mjacqu/glenglat/tree/main?tab=readme-ov-file#authorship-policy>).

4 Conclusions & Outlook

Based on an extensive literature search and data submissions, we have created glenglat, the first (to our knowledge) englacial temperature database for all glaciers outside of the ice sheets. Together with the recent compilations of 415 deep boreholes in Greenland by Løkkegaard et al. (2023) and of shallow measurements for Greenland and Antarctica compiled in the SUMup collaborative database (Vandecrux et al., 2023), most published englacial temperature measurements are now readily available to researchers. Depending on community needs, it may be worth combining these datasets into one, for lower maintenance overhead, ease of use, and because the distinction between ice sheet and glacier will become increasingly arbitrary as glaciers detach from the retreating ice sheet margins. Subsurface 420 variables like density (as in SUMup) or stratigraphy (as in glenglat, but only for the depth of the snow/firn-ice

transition) may be worth adding, especially since these are often measured alongside temperature. Another enhancement would be to include qualitative temperature information, namely whether a borehole was measured as temperate (but no actual measurements were reported) and the presence and depth of a cold-temperate transition surface (often extracted from ice-penetrating radar profiles as an indicator of the glacier thermal regime; 425 e.g., Björnsson et al., 1996; Ødegård et al., 1992, 1997; Pettersson et al., 2003; Wilson et al., 2013)).

We believe that glenglac can contribute to better modelling and understanding of englacial temperatures, their spatial distribution, and their changes in a warming world – most directly as an unparalleled source of observational data for model training and validation. For measurements to better reflect global glacier conditions, however, we see a general need for more measurements (or reporting thereof) from temperate glaciers, warmer climates, elevations between 750 m and 1500 m, and underrepresented regions (e.g., Iceland, New Zealand, Scandinavia). Glenglac also presents new opportunities to investigate changes in englacial temperatures over time, both by making more evident the existence of repeat measurements and documenting a century of early measurements that could now be deliberately repeated.

5 Code and data availability

435 Glenglac is maintained as a Git repository hosted at <https://github.com/mjacqu/glenglac> and published to Zenodo (e.g., version 1.0.0-rc4, to which this manuscript refers: <https://doi.org/10.5281/zenodo.14696283>, Jacquemart et al., 2025). Glenglac is licensed under Creative Commons Attribution 4.0 International, though the repository's license does not extend to figures, tables, maps, or text extracted from publications. These are included in the sources folder for transparency and reproducibility. Full PDFs of the original sources will gladly be shared upon request at any 440 time.

Dataset citation: Jacquemart, M., Welty, E. et al., (2025). glenglac: Global englacial temperature database. Zenodo. <https://doi.org/10.5281/zenodo.11516611>. To cite a subset of the data, we recommend referencing this publication, the dataset, as well as the original source directly or as prefix to the glenglac citation (e.g. "Flowers et al. (2011), in: ...").

445 The GitHub repository contains all of the Python code for the tests, build process, and Zenodo publishing described above. It also contains a Jupyter notebook with tutorials on how to download the data from Zenodo, read the data into Python, and produce statistics and plots similar to those in this paper. It can be run in Google Colab (<https://colab.research.google.com>).

ERA-5 Land climate reanalysis data (Muñoz Sabater, 2019) was downloaded from the Copernicus Climate Change 450 Service.

6 References by glacier and region

Author contributions. MJ conceived the project, and together with EW, designed, implemented, and populated the database. EW managed the testing and publishing pipelines, and MJ and EW wrote the manuscript. MG curated the Colle Gnifetti dataset, including a (re-)evaluation of all associated uncertainties. GC re-digitized a large subset of figures from which data were sourced, providing the basis for the digitization error-assessment. All authors have read, edited, and approved the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. We thank the countless people who contributed to the planning, execution, and processing of the englacial measurements included in glenglat. As an (imperfect) proxy of who they are, we list the names of all authors who were involved in the publications from which we drew data in Appendix B. We are grateful to Gwenn Flowers, Shin Sugiyama, Tika Ram Gurung, Rainer Prinz, Martina Barandun, Olivier Gagliardini, Lonnie G. Thompson, and 张通 (Tong Zhang) for submitting data to us. Lander Van Tricht shared the results of his own literature search with us. Thank you to Daniel Farinotti and Matthias Huss for offering valuable suggestions and discussions along the way. We are particularly indebted to the librarians around the world who tracked down obscure references, including Lkhagvadulam Esterhammer and Monika Kriemler at the University of Zürich, staff at the Royal Danish Library in Copenhagen, and Tatiana A. Ustinova at the Russian State Library in Moscow. Finally, we thank William Colgan, Martin Hoelzle, and Baptiste Vandecrux (editor) for their constructive feedback.

Financial support: MJ was funded by the Swiss National Science Foundation project "PROGRES: Process-based modelling of global glacier changes" (grant 184634). EW was funded by the Federal Office of Meteorology and Climatology (MeteoSwiss) within the framework of the Global Climate Observing System (GCOS) in Switzerland. MC received funding through the "Measuring Firn Hydrology and Ice Dynamics at the Greenland Runoff Limit (MAGNOLIA)" project (funded by Swiss Polar Institute).

Appendix A: References by glacier and region

Table A1: Summary of englacial temperature measurements contained in glenglac, sorted by glacier region, then by glacier (as defined by groups of boreholes with matching glacier name or GLIMS ID), with borehole count, profile count, maximum depth, minimum temperature (at all depths **and at 15 m depth, if available**), range of years, and sources.

Region	Glacier names	GLIMS IDs	BH count	Profile count	Depth [m]	Temp. [°C] max	Year min	Year 15 m	Sources	
									min-max	
1	Black Rapids Glacier	G213683E63392N	3	6	12	-14.0	1973–1973	Harrison et al. (1975)		
1	Fox Glacier	G219698E61200N	6	6	48	-8.1	-7.6	1969–1969	Classen (1970)	
1	Jarvis Glacier	G214333E63481N	2	2995	72	-8.2	2017–2018	Lee (2019); Lee et al. (2020)		
1	Little Kluane Glacier	G220578E60873N	1	17 873	191	-0.3	2019–2021	Flowers (2022)		
1	McCall Glacier	G216152E69302N	27	56	180	-12.4	-4.8	1957–2008	Orvig and Mason (1963); Trabant et al. (1975); Rabus and Echelmeyer (2002); Weller et al. (2007); Delcourt et al. (2013)	
1	North Glacier	G220859E60905N	1	1	70	-3.0	-3.0	2011–2011	Wilson (2012); Wilson et al. (2013); Flowers (2022)	
1	Russell Glacier	G218192E61498N	1	1	460	-23.1	-22.9	2002–2002	Thompson et al. (2004); Urmann (2009)	
1	Seward Glacier	G219787E60289N	3	10	62	-13.0	1948–1950	Sharp (1951)		
1	South Glacier	G220869E60822N	1	14 914	82	-2.4	2011–2014	Flowers et al. (2011); Wilson et al. (2013); Flowers (2022)		
1	Steele Glacier	G219819E61242N	3	3	114	-6.7	-4.9	1972–1974	Jarvis and Clarke (1974); Clarke and Jarvis (1976)	
1	Trapridge Glacier	G219646E61222N	11	11	88	-8.4	-7.3	1972–1980	Jarvis and Clarke (1975); Clarke et al. (1984)	
2	Athabasca Glacier	G242719E52168N	13	16	198	-5.8	1967–1986	Paterson (1971, 1972)		
2	Blue Glacier	G236316E47813N	1	2	104	-0.2	-0.0	1969–1969	Harrison (1972)	
2	Upper Fremont Glacier	G250390E43132N	1	1	10	0.0	1990–1990	Naftz and Smith (1993)		
3	Agassiz Ice Cap	G288743E80950N	3	3	335	-24.6	-24.3	1977–1984	Clarke et al. (1987); Vinther et al. (2008)	
3	Devon Ice Cap	G278488E75058N	3	3	299	-23.2	-23.1	1972–2000	Paterson and Clarke (1978); Kinnard et al. (2006); Mankoff et al. (2022); Mankoff (2022e); Løkkegaard et al. (2023)	
		G277553E75571N								
3	John Evans Glacier	G285646E79663N	4	4	15	-12.2	-12.2	1997–1999	Copland et al. (2003)	
3	Laika Ice Cap	G280856E75887N	5	5	87	-11.2	-10.4	1975–1975	Blatter (1985); Blatter and Kappenberger (1988)	
3	McGill Ice Cap	G266878E79842N	2	2	38	-22.6	-22.0	1962–1962	Harrison (1963); Müller (1963a, b, 1976)	
		G269900E79733N								
3	Meighen Ice Cap	G260810E79982N	1	3	121	-23.1	-17.1	1965–1967	Koerner (1968); Paterson (1968)	
3	Prince of Wales Ice Cap	G279351E78361N	1	1	176	-21.3	-21.0	2005–2005	Mankoff et al. (2022); Mankoff (2022b); Løkkegaard et al. (2023)	
		G269329E79672N								
3	White Glacier	G269329E79672N	48	90	375	-20.4	-14.9	1959–1981	Müller (1961); Harrison (1963); Müller (1963a, 1976); Blatter (1985, 1987)	
4	Barnes Ice Cap	G287731E69650N	16	16	281	-11.1	-10.9	1973–1977	Hooke (1976); Classen (1977); Hooke et al. (1980); Gilbert et al. (2016)	
		G287718E69797N								
		G288059E69709N								
4	Penny Ice Cap	G294456E67304N	1	1	176	-12.8	-12.2	1996–1996	Mankoff et al. (2022); Mankoff (2022c); Løkkegaard et al. (2023)	
5	Flade Isblink Ice Cap	G344790E81287N	1	1	420	-17.4	2006–2006	Lemark (2010); Mankoff et al. (2022); Mankoff (2022d); Løkkegaard et al. (2023)		
5	Hans Tausen Ice Cap	G323085E82488N	1	1	341	-20.8	-20.0	1995–1995	Hammer (1995); Reeh (1995); Steffensen et al. (2001)	
5	Hare Glacier	G323403E82808N	6	6	286	-21.0	-16.9	1994–1995	Thomsen et al. (1996); Reeh et al. (2001)	
		G322065E82674N								
5	Hurlbut Gletscher	G292196E77316N	2	8	15	-18.2	1957–1957	ark (1959, 1960); Fristrup (1960a, b, 1961)		
5	Mitdluagkat Gletscher	G322197E65696N	1	1	15	-1.0	1958–1958	Fristrup (1960a, 1961); Hasholt (1987)		
5	Napassorsuaq Gletscher	G314725E60307N	1	1	7	-0.7	1957–1957	ark (1959, 1960); Fristrup (1960a, 1961); Weidick (1988)		
		G322065E82674N								
5	Nunatarssuaq Ice Cap	G292408E76864N	2	343	2	-8.6	2017–2017	Abermann et al. (2020); Prinz et al. (2022)		
5	Qaanaaq Glacier	G290838E77522N	3	3	13	-11.9	2014–2014	Tsutaki et al. (2017)		
5	Renland Ice Cap	G333444E71216N	1	1	300	-18.7	-18.4	1988–1988	Mankoff et al. (2022); Mankoff (2022a); Løkkegaard et al. (2023)	
		G308051E66298N								
5	Sermikavsk	G307173E71231N	1	1	7	-3.2	1957–1957	Møller (1959); Fristrup (1960a, 1961)		
5	Sukkertoppen Ice Cap	G307609E66296N	4	4	12	-4.6	1964–1964	Rundle (1965)		
		G308051E66298N								
5	Tuto Ramp	G291955E76463N	1	12	47	-22.1	-11.5	1961–1962	Davis (1967)	

Continued on next page

Table A1: Summary of englacial temperatures contained in glenglac, and their sources (continued).

Region	Glacier names	GLIMS IDs	BH	Profile	Depth	Temp.	Year	Sources	
			count	count	[m]	max	min	15 m	min–max
7	Amundsenisen Ледниково плато Амундсена	G015444E77229N	1	3	13	-9.5	1980–1980	В. С. Загороднов [V. S. Zagorodnov] (1981)	
7	Austfonna	G024340E79634N G025297E79771N G023619E79932N G024143E79973N	14	15	565	-16.4	-4.5	1987–1999	В. С. Загороднов [V. S. Zagorodnov] et al. (1990); Watts et al. (1997); Watanabe et al. (2001)
7	Austre Brøggerbreen	G011895E78886N	2	2	108	-4.2	-4.2	1990–1992	Hagen (1992); Björnsson et al. (1996)
7	Austre Grønfjordbreen	G014342E77910N	12	30	83	-8.5	-2.2	1966–2014	Е. М. Зингер [E. M. Singer] and В. И. Михаилев [V. I. Mikhailov] (1967); В. С. Загороднов [V. S. Zagorodnov] and И. А. Зотиков [I. A. Zotikov] (1981); Ю. Я. Мачерет [Y. Y. Macheret] et al. (1985); Kotlyakov et al. (2004); P. А. Чернов [R. A. Chernov] et al. (2015)
7	Austre Lovénbreen	G012161E78870N	3	17	20	-6.2	-3.2	2009–2011	孙维君 [Sun Weijun] et al. (2016)
7	Bertilbreen Ледник Бертиль	G016264E78699N	3	3	108	-11.5	-5.6	1980–1980	В. С. Загороднов [V. S. Zagorodnov] (1981)
7	Bogerbreen Ледник Багер	G015633E78130N	1	1	7	-7.3		1980–1980	В. С. Загороднов [V. S. Zagorodnov] (1981)
7	Erikbreen	G012478E79621N	5	10	20	-8.7	-3.5	1990–1992	Ødegård et al. (1992)
7	Finsterwalderbreen	G015235E77463N	2	2	189	-4.5	-3.1	1994–1995	Ødegård et al. (1997)
7	Fridtjofbreen	G014442E77835N	1	1	115	-5.2	-3.9	1981–1981	Ю. Я. Мачерет [Y. Y. Macheret] et al. (1985)
7	Hansbreen	G015592E77097N	6	6	330	-9.1	-3.5	1979–1995	Jania et al. (1996)
7	Høghetta	G016639E79309N	1	1	86	-13.3	-10.4	1987–1987	Kawamura et al. (1991)
7	Irenebreen	G012138E78665N	1	12	10	-5.8		2008–2009	Sobota (2011)
7	Kongsvegen	G013044E78792N	2	2	324	-5.6	-4.3	1990–1992	Hagen (1992); Björnsson et al. (1996)
7	Lomonosovfonna	G018042E78675N Ледниково Плато Ломоносова	4	15	122	-11.4	-2.4	1965–2013	Е. М. Зингер [E. M. Singer] et al. (1966); В. С. Загороднов [V. S. Zagorodnov] and И. А. Зотиков [I. A. Zotikov] (1981); van de Wal et al. (2002); Kotlyakov et al. (2004); Marchenko et al. (2017)
7	Midre Lovénbreen	G012039E78878N	2	2	133	-3.6	-3.3	1990–1992	Hagen (1992); Björnsson et al. (1996)
7	Nordenskiöldbreen Ледник Норденшельдъ	G017371E78745N	2	2	26	-7.1	-4.4	1965–1965	Е. М. Зингер [E. M. Singer] et al. (1966)
7	Scott Turnerbreen	G015894E78097N	2	2	54	-11.1	-7.2	1993–1993	Hodgkins et al. (1999)
7	Snfjellafonna	G013542E78988N	1	1	80	-3.4	-1.8	1992–1992	Kameda et al. (1993)
7	Vestfonna	G019951E79875N G020879E79901N G019797E80009N	6	11	200	-23.6	-2.7	1956–1995	Palosuo and Schytt (1960); Schytt (1964); В. М. Котляков [V. M. Kotlyakov] (1985); Palosuo (1987); Watanabe et al. (2001); Kotlyakov et al. (2004)
7	Waldemarbreene	G012079E78681N	3	23	10	-7.6		2007–2019	Sobota (2009); Karušs et al. (2022)
7	Werenskioldbreen	G015442E77070N	4	4	15	-4.2	0.0	1970–1970	Baranowski (1975)
7	Åsgårdfonna	G017048E79443N	2	2	182	-7.9	-6.1	1993–1993	Uchida et al. (1996)
8	Nigardsbreen	G007099E61715N	1	1	44	-0.8	-0.1	1987–1987	Kawamura et al. (1989)
8	Storglaciären	G018569E67903N	4	15	40	-9.0	-2.0	1965–2002	Schytt (1966, 1968); Pettersson et al. (2003)
9	Academy of Sciences Glacier Академии наук Ледник Академии Наук	G096063E80433N	3	3	743	-14.7		1986–2001	А. М. Саватюгин [A. M. Savatyugin] and В. С. Загороднов [V. S. Zagorodnov] (1988); В. С. Загороднов [V. S. Zagorodnov] (1989); Zagorodnov and Arkhipov (1990); Л. М. Саватюгин [L. M. Savatyugin] et al. (2001); Fritzschke et al. (2002); Kotlyakov et al. (2004)
9	Churlyanis Cupola Sedov Glacier Купол Чурляниса ледник Седова	G053403E80271N G053047E80333N G053032E80282N G052977E80310N	31	134	82	-27.5		1958–1959	Н. Г. Разумейко [N. G. Razumeiko] (1960, 1963)
9	Jackson Cupola Купол Джексона	G053200E80194N	1	13	20	-22.3		1959–1959	Н. Г. Разумейко [N. G. Razumeiko] (1963)
9	Salm Island Glacier ледник Остров Сальм	G059273E79977N	1	1	14	-18.0		2005–2005	Kubyshkin et al. (2006)
9	Shokalsky Glacier Ледник Шокальского	G062464E75974N G062675E76121N	16	38	30	-29.0		1958–1959	И. Ф. Хмелевской [I. F. Khmelevskoy] (1963, 1964)

Continued on next page

Table A1: Summary of englacial temperatures contained in glenglac, and their sources (continued).

Region	Glacier names	GLIMS IDs	BH	Profile	Depth	Temp.	Year	Sources	
			count	count	[m]	max	min	15 m	min–max
9	Vavilov Glacier	G095294E79482N	8	21	470	-25.0	-12.6	1974–1985	B. P. Барбаш [V. R. Barbash] et al. (1981); B. А. Морев [V. A. Morev] and B. А. Пухов [V. A. Pukhov] (1981); Н. И. Барков [N. I. Barkov] et al. (1988); B. А. Морев [V. A. Morev] et al. (1988); Kotlyakov et al. (2004)
	Vavilov Ice Cap	G096481E79287N							
	Купол Вавилова	G095612E79448N							
	Ледник Вавилова								
9	Vetreniy Ice Dome	G063846E80729N	1	1	308	-11.4	-6.8	1997–1997	Kotlyakov et al. (2004)
10	Belukha Glacier	G086577E49799N	1	1	75	-17.2		2001–2001	Olivier et al. (2003)
10	Blichenok Glacier	G160474E56097N	2	14	211	-23.2	-15.5	1996–1999	Shiraiwa et al. (1997, 2001)
10	Halasi Glacier	G087783E49100N	3	6	8	-4.0		1980–1980	王立伦[Wang Lilun] et al. (1983); 刘时银[Liu Shiyin] et al. (2012); Guo et al. (2015)
	哈拉斯冰川								
10	Khukh Nuru Uul glacier	G090853E48651N	1	1	70	-13.8	-12.6	2009–2009	Herren et al. (2013)
10	Malii Aktru Glacier	G087761E50048N	10	15	30	-11.4	-8.4	1980–1982	C. A. Никитин [S. A. Nikitin] (1986)
	ледник Малый Актру	G087720E50060N							
10	Sofiskiy Glacier	G087759E49791N	2	2	25	-0.3	-0.0	2000–2001	Fujii et al. (2002)
10	Tsambagarav Glacier	G090847E48595N	1	1	40	-13.4	-12.0	2008–2008	Liu et al. (2009); Davaa (2016); Khalzan et al. (2022)
10	Vodopadniy Glacier	G087789E50050N	1	2	12	-15.7		1981–1982	C. A. Никитин [S. A. Nikitin] (1986)
	ледник Водопадный								
10	Western Belukha Plateau	G086544E49802N	1	1	170	-15.7	-14.8	2003–2003	Takeuchi et al. (2004)
11	Alteisgletscher	G007671E46431N	3	22	21	-6.4	-0.4	1991–1991	Laternser (1992)
11	Breithornplateau	G007908E45948N	82	115	359	-17.0	-14.6	1975–2021	Haeberli (1976); Oeschger et al. (1977); Gäggeler et al. (1983); Blatter and Haeberli (1984); Haeberli and Funk (1991); Laternser (1992); Lüthi (1999); Keck (2001); Lüthi and Funk (2001); Suter et al. (2001, 2002); Schwerzmann et al. (2006b); Darms (2009); Eisen et al. (2009); Hoelzle (2009); Hoelzle et al. (2011); Diez et al. (2013); Ryser et al. (2013); Hoelzle (2014); Mayewski (2014); Hoelzle (2017); Bohleber et al. (2018); Hoelzle et al. (2020); Mattea (2020); swi (2021); Hoelzle (2022); Gastaldello and Hoelzle (2024a, b)
	Gornergletscher	G007800E45965N							
	Grenzgletscher	G007875E45922N							
11	Fieschergletscher	G008144E46504N	3	3	153	-6.8	-2.0	2003–2003	Schwerzmann et al. (2006b, a)
11	Glacier de Taconnaz	G006844E45863N	41	41	126	-15.1	-14.2	1911–2017	Valot (1913); Lliboutry et al. (1976); Jouzel et al. (1984); Suter et al. (2002); Vincent et al. (2007); Gilbert and Vincent (2013); Gilbert et al. (2015); Vincent et al. (2020)
	Glacier des Bossons	G006865E45868N							
	Taconnaz Glacier								
11	Glacier de Tête Rousse	G006819E45856N	19	41	70	-2.8	-2.7	2010–2023	Gilbert et al. (2012); Vincent et al. (2012); Gagliardini and Vincent (2023)
11	Glacier du Pelvoux	G006408E44900N	1	1	13	-0.2		1983–1983	Jouzel et al. (1984)
11	Glacier du Sex Rouge	G007212E46327N	2	2	35	-1.1	-0.5	2013–2014	Signer (2014)
11	Grubengletscher	G007996E46168N	6	6	46	-2.3	-2.3	1974–1975	Haeberli (1976)
11	Hintereisferner	G010752E46802N	6	70	15	-4.7	-0.3	1972–1976	Markl and Wagner (1977)
11	Hintergrat Glacier	G010554E46507N	1	41 581	10	-8.9		2011–2016	Gabrielli et al. (2016); Carturan et al. (2023a, b)
11	Jungfraufirn	G008032E46504N	10	33	20	-13.6	-3.6	1938–1991	Hughes and Seligman (1939); Laternser (1992); Suter et al. (2001)
11	Lysgletscher	G007846E45906N	6	6	22	-10.0	-5.5	1999–1999	Suter et al. (2002); Gastaldello and Hoelzle (2024a)
11	Mont Blanc	G006867E45829N	2	2	18	-17.1		1983–1998	Jouzel et al. (1984); Suter et al. (2002)
11	Sphinxgrat	G007985E46549N	1	1	10	-6.0		1981–1981	Haeberli and Alean (1985)
11	St. Annafirn	G008601E46597N	2	2	8	-1.3		2013–2014	Signer (2014)
11	Titlis-Gletscher	G008427E46774N	1	1	15	-0.7	-0.7	1979–1980	Haeberli and Alean (1985)
11	Unteraargletscher	G008187E46569N	1	1	60	0.0		1842–1842	Wild et al. (1842); Agassiz (1847)
11	Vadret da Morteratsch	G009927E46382N	1	1	42	-2.8	-2.2	2002–2002	Schwerzmann et al. (2006b)
11	Vadret dal Corvatsch	G009822E46416N	10	41	18	-9.1	-2.6	1999–2002	Hager (2002); Haeberli et al. (2004)
11	Vedretta Alta dell'Ortles	G010536E46513N	4	62 324	75	-9.4		2009–2016	Gabrielli et al. (2010, 2012, 2016); Carturan et al. (2023a, b)
12	Bezengi Glacier	G043100E43030N	1	1	80	-0.6	-0.5	1966–1966	T. В. Псарёва [T. V. Psareva] (1968); Khromova et al. (2022)
	Ледник Безенги								
12	Djankuat Glacier	G042766E43192N	5	5	55	-19.5	-0.5	1971–1971	gid (1978)
	Ледник Джанкуат								

Continued on next page

Table A1: Summary of englacial temperatures contained in glenglac, and their sources (continued).

Region	Glacier names	GLIMS IDs	BH	Profile	Depth	Temp.	Year	Sources	
			count	count	[m]	max	min	15 m	min–max
12	Garabashi Glacier Ледник Гарабаши	G042470E43307N	4	19	20	-13.0	1958–1988	М. Я. Плам [M. Ya. Plam] (1962); Б. С. Загороднов [V. S. Zagorodnov] et al. (1992)	
12	Mount Elbrus	G042429E43293N G042488E43308N	4	4	182	-17.3	-16.9	2004–2020	Б. Н. Михаленко [V. N. Mikhaleenko] et al. (2005a); Mikhaleenko et al. (2015); Б. Н. Михаленко [V. N. Mikhaleenko] et al. (2021)
13	Abramov Glacier	G071570E39610N	2	2	19	-0.3	-0.1	2013–2013	Barandun (2018, 2024)
13	Ashu-Tor Glacier ледник Ашу-Тор	G078182E42041N	1	1	20	-7.6	?–1962	А. Н. Диких [A. N. Dikikh] (1965); С. С. Кутузов [S. S. Kutuzov] (2012); Van Tricht et al. (2021); Алексей Викторович Цветков [Aleksey Viktorovich Tsvetkov] (2023)	
13	Batysh Sook Glacier	G077749E41787N	1	5	15	-6.3	-0.6	2013–2017	Barandun (2023)
13	Bogda Fan-Shaped Difffluence Glacier	G088313E43812N	1	4	20	-3.0	-1.5	1981–1981	仇家琪[Qiu Jiaqi] 和 邓养鑫[Deng Yangxin] (1983); 任贾文[Ren Jiawen] (1983); 刘时银[Liu Shiyin] et al. (2012); Guo et al. (2015)
13	Central Tuyuksu Glacier Ледник Туюксу Центральный	G077080E43049N	5	119	52	-16.1	1957–1959	Vilesov (1961); Е. Н. Вилесов [E. N. Vilesov] (1962a, b, c); Г. А. Цыкина [G. A. Tsykina] and Е. Н. Вилесов [E. N. Vilesov] (1963)	
13	Chongce Ice Cap 崇测冰帽	G081119E35239N	7	13	130	-16.4	-16.0	1987–2012	Huang (1990); Shao and Liu (1990); 周韬[Zhou Tao] (1990); Hou et al. (2018)
13	Crescent River Glacier No. 15 月牙河15号冰川	G087444E36402N	2	2	18	-8.2	-7.4	1988–1988	苏珍[Su Zhen] (1998); 刘时银[Liu Shiyin] et al. (2012); Guo et al. (2015)
13	Davydov Glacier Ледник Давыдова	G078204E41844N	1	4	30	-5.8	-2.5	1985–1985	Е. В. Василенко [E. V. Vasilenko] (1988)
13	Dunde Ice Cap	G096414E38091N	1	1	136	-7.3	-6.7	1984–1984	Thompson et al. (1990)
13	Geladandong Ice Cap	G091151E33199N	1	1	87	-12.1	-5.0	2004–2004	Wang and Pu (2005)
13	Grigoriev Glacier Grigoriev Ice Cap Ледник Григорьева	G077894E41995N G077923E41963N	14	1900	87	-22.9	-6.1	1962–2023	А. Н. Диких [A. N. Dikikh] (1965); Б. Н. Михаленко [V. N. Mikhaleenko] (1989); Thompson et al. (1993, 1997); С. М. Архипов [S. M. Arkhipov] et al. (2004); Б. Н. Михаленко [V. N. Mikhaleenko] et al. (2005b); Takeuchi et al. (2014); Kronenberg et al. (2022); Kronenberg (2022); Machguth et al. (2023a, b)
13	Guliya Ice Cap 古里雅冰帽	G081455E35226N G081480E35252N	7	7	310	-21.3	-17.8	1990–2015	姚檀栋[Yao Tandong] et al. (1992); Thompson et al. (1995a, 2018)
13	Guozha Glacier 郭扎冰川	G081064E35246N	1	2	12	-6.4		1987–1987	Shao and Liu (1990)
13	Halong Glacier 哈龙冰川	G099492E34764N	2	2	10	-7.3		1981–1981	王文颖[Wang Wenyng] (1987); 苏珍[Su Zhen] (1998); 刘时银[Liu Shiyin] et al. (2012); Guo et al. (2015)
13	July 1st Glacier	G097755E39237N	1	4	8	-8.9		2002–2002	Matsuda et al. (2004)
13	Laozugou Glacier	G096524E39457N	4	4	109	-10.3	-8.0	2010–2011	Wang et al. (2018)
13	Malan Glacier	G090770E35803N	1	1	100	-9.3	-4.7	1999–1999	Sun et al. (2021)
13	Meikuang Glacier 煤矿冰川	G094184E35669N	1	1	16	-7.0	-6.2	1989–1989	苏珍[Su Zhen] (1998); 刘时银[Liu Shiyin] et al. (2012); Guo et al. (2015)
13	Miaoergou Glacier	G094316E43053N	1	1	60	-8.3	-6.4	2005–2005	刘亚平[Liu Yaping] et al. (2006); Takeuchi et al. (2008); Liu et al. (2009); Jiao et al. (2023)
13	Muztagh Glacier 慕士塔格冰川	G075086E38293N	5	5	78	-26.2	-25.8	2002–2003	邬光剑[Wu Guangjian] et al. (2003); 李真[Li Zhen] et al. (2004)
13	Puruogangri Ice Cap	G089122E33894N	3	4	213	-9.9	-9.7	2000–2000	蒲健辰[Pu Jianchen] et al. (2002); Thompson et al. (2006); Liu et al. (2016)
13	Qiangtang No. 1 Glacier	G088700E33291N	1	1	109	-11.4	-7.7	2014–2015	Li et al. (2017)
13	Qingbingtan Glacier No. 72	G079894E41774N	3	3	10	-1.7		2008–2008	Wang et al. (2017)
13	Shule Nanshan Glacier	G097261E38695N	1	1	92	-9.9	-7.8	2007–2007	Liu et al. (2009)
13	South Inylchek Glacier	G079787E42137N	1	1	160	-12.0	-12.0	2000–2000	Aizen et al. (2001); Thompson (2022)
13	Urumqi Glacier No. 1 乌鲁木齐1号冰川	G086810E43111N G086801E43117N	24	26	107	-8.8	-7.4	1981–2006	任贾文[Ren Jiawen] et al. (1985); Huang (1990); Zhang et al. (1993); 李忠勤[Li Zhongqin] et al. (2011)
13	Xiao Dongkemadi Glacier 小冬克玛底冰川	G092063E33082N	1	2	15	-11.0	-7.3	1992–1993	蒲健辰[Pu Jianchen] et al. (1995)
13	Yanglong River Glacier No. 5 羊龙河5号冰川	G098570E39226N	3	27	16	-10.7	-8.2	1977–1977	任贾文[Ren Jiawen] and 黄茂桓[Huang Maohuan] (1981)
13	Zangser Kangri Glacier	G085843E34297N	1	1	127	-12.4	-12.3	2009–2009	An et al. (2016)

Continued on next page

Table A1: Summary of englacial temperatures contained in glenglac, and their sources (continued).

Region	Glacier names	GLIMS IDs	BH	Profile	Depth	Temp.	Year	Sources	
			count	count	[m]	max	min	15 m	min–max
14	Singhi Glacier 特拉木坎力冰川	G077054E35619N	1	1	10	-1.8	1987–1987	苏珍[Su Zhen] (1998); 刘时银[Liu Shiyin] et al. (2012); Guo et al. (2015)	
15	Baishui Glacier No. 1	G100187E27104N	4	13	21	-2.8	-0.5	1982–2010	Huang (1990); Du et al. (2013)
15	Dagongba Glacier	G101855E29563N	1	1	15	-0.9	1982–1982	Huang (1990)	
15	Dasuopu Glacier	G085752E28395N	2	2	168	-14.4	-13.8	1997–1997	Thompson et al. (2000); Yao et al. (2002); Thompson (2022)
15	East Rongbul Glacier	G086939E28060N	3	3	109	-10.9	-9.2	2002–2008	Hou et al. (2004, 2007); Zhang et al. (2013); 张通[Zhang Tong] (2022)
15	Gyabrag Glacier 加布拉冰川	G086633E28122N	1	1	69	-9.1	-7.7	2005–2005	Liu et al. (2009)
15	Khumbu Glacier	G086820E27978N	3	3	131	-3.3	-3.2	2017–2017	Miles et al. (2018, 2019); Hubbard et al. (2021)
15	Naimona'nyi Glacier	G081317E30454N	1	1	159	-9.6	-9.4	2006–2006	Thompson et al. (2018); Thompson (2022)
15	Rikha Samba Glacier	G083488E28819N	3	6 134	10	-10.6	2014–2015	Gilbert et al. (2020); Gurung (2022)	
15	Trambau Glacier	G086537E27874N	1	1	78	-1.3	2019–2019	Tsushima et al. (2021)	
15	Yala Glacier	G085612E28242N	2	2	60	-1.0	-1.0	1981–1982	Iida et al. (1984); Watanabe et al. (1984)
16	Illimani Volcano	G292220E16653S	1	1	138	-9.0	-8.1	1999–1999	Gilbert et al. (2010)
16	Mount Kilimanjaro	G037352E03058S	1	1	51	-1.6	-1.1	2000–2000	Thompson et al. (2002); Thompson (2022)
16	Nevado Huascarán	G282414E09082S	3	3	167	-9.0	-5.9	1993–2019	Thompson et al. (1995b); Thompson (2022); Thompson et al. (2023)
16	Nevado Sajama	G291113E18113S	1	1	132	-11.3	-10.6	1997–1997	Thompson et al. (1998); Vimeux et al. (2009); Thompson (2022)
16	Quelccaya Ice Cap	G289183E13941S	2	2	168	-7.2	0.0	1976–2003	Thompson (1980); Zagorodnov et al. (2005); Vimeux et al. (2009); Thompson (2015, 2022)
16	Volcán Coropuna	G287357E15537S	2	2	147	-11.1	-9.0	2003–2003	Zagorodnov et al. (2005, 2006); Vimeux et al. (2009); Thompson (2022)
17	Glaciar La Ollada	G289889E31964S	1	1	104	-18.5	-18.3	2005–2005	Schwerzmann et al. (2006b)
17	Glaciar Nef	G286668E46885S	1	1	13	-0.1	1996–1996	Matsuoka and Naruse (1999)	
17	Glaciar San Rafael	G286620E46550S	1	1	16	-11.9	-11.6	2005–2005	Vimeux et al. (2008)
17	Guanaco Glacier	G289989E29347S	1	4	112	-8.0	-6.2	2008–2011	Kinnard et al. (2020); Masiokas et al. (2020)
17	Pío XI glacier	G286372E49263S	1	1	40	-0.9	-0.7	2006–2006	Schwikowski et al. (2013)
17	Tapado Glacier	G290072E30145S	1	1	36	-12.4	-10.7	1992–1992	Ginot et al. (2006)
19	Bläskimen Island Ice Rise	G356949E70424S	1	1	19	-16.4	-16.3	2012–2014	Goel et al. (2017a, b)
19	Bruce Plateau	G295982E66134S	2	6	431	-15.8	-15.1	2010–2010	Zagorodnov et al. (2012)
19	Collins Ice Cap	G301284E62099S	7	16	30	-6.1	-2.0	1992–1992	韩建康[Han Jiankang] et al. (1995)
		G301112E62165S							
19	Dolleman Island	G299288E70606S	1	1	128	-17.2	-17.0	1986–1986	Nicholls and Paren (1993)
19	Dyer Plateau		1	1	104	-21.8	-21.4	1989–1989	Nicholls and Paren (1993)
19	James Ross Island Ice Cap	G302228E64270S	2	2	10	-14.2	1976–1977	Aristarain and Delmas (1981)	
19	Johnsons Glacier	G299645E62671S	2	2	153	-0.2	-0.2	2016–2016	Sugiyama et al. (2019); Sugiyama (2022)
19	King George Island Ice Cap	G301226E62159S	1	3	15	-1.8	1986–1986	任贾文[Ren Jiawen] (1990)	
19	Nelson Island Ice Cap	G301002E62274S	3	12	13	-1.6	1986–1986	任贾文[Ren Jiawen] (1990)	
		G300887E62269S							
19	Styx Glacier		1	1	210	-33.5	-31.7	2016–2016	Han et al. (2015); Yang et al. (2018)

Appendix B: Authors of all sources

Names of all the authors of all the sources used to compile the data in glenglat. Names in non-Latin script are followed by a
475 latinized form in square brackets. The list is sorted alphabetically by (Latin) family name, which is capitalized. The latinized
form preserves the order of the original, which is [family given] for Chinese, Japanese, and Korean names.

Jakob ABERMANN · Андрей Андреевич Абрамов [Andrey Andreevich ABRAMOV] · Louis AGASSIZ · AHN Jinho · 艾松
涛[AI Songtao] · Vladimir B. AIZEN · Jürg ALEAN · E. Calvin ALEXANDER JR. · Wenling AN · Katrine Krogh ANDERSEN
· Teruo AOKI · Alberto J. ARISTARAIN · C. M. Архипов [Serguei M. ARKHIPOV] · Andy ASCHWANDEN · Giovanni
480 BACCOLO · 保翰璋[BAO Hanzhang] · Weijia BAO · Martina BARANDUN · Stanisław BARANOWSKI · Carlo BARBANTE
· B. R. Барбаш [V. R. BARBASH] · Gonzalo BARCAZA · H. И. Барков [N. I. BARKOV] · Andreas BAUDER · A. Б. Бажев
[A. B. BAZHEV] · Emilie BEAUDON · Jürg BEER · Sultan BELEKOV · Carl BENSON · Steven BERNSEN · Giuliano
BERTAGNA · Etienne BERTHIER · Michele BERTÓ · Daniel BINDER · Robert G. BINGHAM · Helgi BJÖRNSSON ·
Heinz BLATTER · Norbert BLINDOW · K. В. Блинов [K. V. BLINOV] · Per Helge BØ · Carl Egede BØGGILD · Pascal
485 BOHLEBER · John F. BOLZAN · Francisca BOWN · Henry H. BRECHER · Donald BREN · M. BRIAT · Joel BROWN ·
Sabine BRÜTSCH · M. С. Бугорков [M. S. BUGORKOV] · I. V. BUZIN · Nicolas CAILLON · Seth W. CAMPBELL · R.
Kramer CAMPEN · Alberto CARTON · Luca CARTURAN · Gino CASASSA · Federico CAZORZI · Jorge Luis CEBALLOS
· Jérôme CHAPPELLAZ · Jizu CHEN · Xuejiao CHEN · 陈亚宁[CHEN Yaning] · P. A. Чернов [R. A. CHERNOV] ·
Taejin CHOI · Poul CHRISTOFFERSEN · Peter C. CHU · Garry K. C. CLARKE · David Farley CLASSEN · Henrik B.
490 CLAUSEN · David CLEMENS-SEWALL · Gary D. CLOW · Jihong COLE-DAI · William T. COLGAN · Sam G. COLLINS
· Luke COPLAND · Giulio COZZI · Michel CRESEVEUR · Dorthe DAHL-JENSEN · Giancarlo DALLA FONTANA · Gian
Andrea DARMS · Gombo DAVAA · Mary E. DAVIS · Robert M. DAVIS · Martine DE ANGELIS · Fabrizio DE BLASI
· Charlotte DELCOURT · Robert Jean DELMAS · 邓养鑫[DENG Yangxin] · Marc DESCLOITRES · Anja DIEZ · A. Н.
Диких [A. N. DIKIKH] · Roberto DINALE · Minghu DING · Д. Н. Дмитриев [D. N. DMITRIEV] · Zhiwen DONG · Julian
495 A. DOWDESWELL · Samuel H. DOYLE · Gianfranco DRAGÀ · Giuliano DREOSSI · DU Jiankuo · Wentao DU · 段克
勤[DUAN Keqin] · Inés Maria DUSSAILLANT · Mark DYURGEROV · Péteris DŽERIŅŠ · Keith A. ECHELMAYER · Anja
EICHLER · Olaf EISEN · Алексея Анатольевича Екайкина [Alexey Anatolyevich EKAYKIN] · Tobias ERHARDT · Björn
ERLINGSSON · Bakyt ERMENBAIEV · Nao ESASHI · Bernd ETZELMULLER · Stella EYRIKH · Xavier FAÏN · Ф. Ф.
Файзрахманов [F. F. FAIZRAKHMANOV] · Sonia FALOURD · Robert S. FAUSTO · Daniela FESTI · David A. FISCHER
500 · Hubertus FISCHER · David A. FISHER · Gwenn E. FLOWERS · Bernard FRANCOU · Regula FRAUNFELDER ·
Massimo FREZZOTTI · Børge FRISTRUP · Diedrich FRITZSCHE · Yoshiyuki FUJII · 藤田耕史[FUJITA Koji] · Takehiro
FUKUDA · Martin FUNK · Johannes J. FÜRST · Jacopo GABRIELI · Paolo GABRIELLI · Bogdan GADEK · Heinz
W. GÄGGELE · Olivier GAGLIARDINI · Stephan P. GALOS · Stéphane GARAMBOIS · Alex S. GARDNER · Marcus
GASTALDELLO · 耿志新[GENG Zhixin] · Christopher C. GERBI · Adrien GILBERT · Patrick GINOT · Yngvar GJESSING
505 · A. F. GLAZOVSKY · Vikram GOEL · Kumiko GOTO-AZUMA · Л. С. Говоруха [L. S. GOVORUKHA] · А. И. Громыко
[A. I. GROMYKO] · Niels S. GUNDESTRUP · 郭万钦[GUO Wanqin] · Tika Ram GURUNG · Robert J. GUSTAFSON ·
Hélène GUYARD · Wilfried HAEBERLI · Jon Ove HAGEN · Patrick HAGER · Claus Uffe HAMMER · Svein-Erik HAMRAN
· S.-E. HAMRANW · Hyangsun HAN · 韩建康[HAN Jiankang] · HAN Yeongcheol · Douglas R. HARDY · Joel HARPER ·

J. HARRISON · William Douglas HARRISON · Bent HASHOLT · H. HAUSMANN · Robert L. HAWLEY · HE Yuanqing

- 510 · Petra HEIDENWOLF · Keith A. HENDERSON · Pierre-Alain HERREN · Martin HEULE · Benjamin HILLS · Richard HODGKINS · Andrew HODSON · Martin HOELZLE · Georg HOFFMANN · Helene HOFFMANN · Coen HOFSTEDE · Per HOLMLUND · HONG Sang Bum · Roger LeB. HOOKE · 侯姗姗[HOU Shanshan] · 侯书贵[HOU Shugui] · Kate HRUBY · Houtse HSU · 黄茂桓[HUANG Maohuan] · Bryn HUBBARD · Hans-Wolfgang HUBBERTEN · T. P. HUGHES · Neil HUMPHREY · HUR Soon Do · Matthias HUSS · Philippe HUYBRECHTS · Makoto IGARASHI · Hajime IIDA · 515 Elisabeth ISAKSSON · Kaoru IZUMI · Mylène JACQUEMART · Youngjoon JANG · Jacek JANIA · Peter JANSSON · Gary T. JARVIS · Theo M. JENK · Jurijs JEŠKINS · Liming JIANG · Zongli JIANG · 焦克勤[JIAO Keqin] · Xiaoyu JIAO · 金会军[JIN Huijun] · Sigríður Jóhann JOHNSEN · Peter JONSSON · Daniel JOSWIAK · Bruno JOURDAIN · Jean JOUZEL · JUN Seong Joon · Andreas KÄÄB · Takao KAMEDA · Kokichi KAMIYAMA · 康世昌[KANG Schichang] · 康兴成[KANG Xingcheng] · Giovanni KAPPENBERGER · Nanna B. KARLSSON · Jānis KARUŠS · Sadao KAWAGUCHI · Toshiyuki 520 KAWAMURA · Rijan B. KAYASTHA · Lothar KECK · Natalie KEHRWALD · Donald V. KENNY · Purevdagva KHALZAN · Shfaqat A. KHAN · И. Ф. Хмелевской [I. F. KHMELEVSKOY] · Татьяна Емельяновна Хромова [Tatiana Emelyanova KHROMOVA] · Seong-Joong KIM · Christophe KINNARD · L. KLEIN · О. Л. Клементьев [O. L. KLEMENTIEV] · Roy M. KOERNER · Mika KOHNO · Shiro KOHSHIMA · П. А. Королев [P. A. KOROLEV] · B. C. Корякин [V. S. KORYAKIN] · Владимир М. Котляков [Vladimir M. KOTLYAKOV] · Anna KOZACHEK · Karl KRAINER · Karl KREUTZ · Marlene 525 KRONENBERG · Jean KRUG · Jumpei KUBOTA · N. V. KUBYSHKIN · A. B. Кудиков [A. V. KUDIKOV] · Christoph KULL · M. G. Кунакович [M. G. KUNAKHOVICH] · Станислав С. Кутузов [Stanislav S. KUTUZOV] · Olivier LAARMAN · Ang J. B. LAMA SHERPA · Kristaps LAMSTERS · Amaelle LANDAIS · Martin LATERNSER · Иван И. Лаврентьев [Ivan I. LAVRENTIEV] · Ю. А. Лаврушин [Yu. A. LAVRUSHIN] · Robert LAW · Emmanuel LE MEUR · LEE Hun-Gyu · Ian R. LEE · Eric LEFEBVRE · Anatoly LEGCHENKO · Michel LEGRAND · Andreas LEMARK · 李慧林[LI Huilin] · LI Shuang 530 · Yemeng LI · 李真[LI Zhen] · 李忠勤[LI Zhongqin] · Olav LIESTØL · S. LIM · Ping-Nan LIN · Song LINLIN · Владимир Я. Липенков [Vladimir Y. LIPENKOV] · Peter LISAGER · 刘潮海[LIU Chaohai] · K.-b. LIU · 刘雷保[LIU Leibao] · Lin LIU · Qiao LIU · 刘时银[LIU Shiyin] · 刘亚平[LIU Yaping] · 刘宇硕[LIU Yushuo] · LIU Zongxiang · Louis LLIBOUTRY · Albert LLUBERAS · Anja LØKKEGAARD · Martin Peter LÜTHI · Shelley MACDONELL · Joseph A. MACGREGOR · Юрий Яковлевич Мачерет [Yuri Yakovlevich MACHERET] · Horst MACHGUTH · Olivier MAGAND · Sudan B. MAHARJAN · 535 Volkmar MAIR · Philipp MALZ · Л. Н. Маневский [L. N. MANEVSKY] · Kenneth D. MANKOFF · Sergey MARCHENKO · B. A. Маркин [V. A. MARKIN] · Von G. MARKL · Tracy A. MASHIOTTA · Mariano H. MASIOKAS · Robert W. MASON · Valérie MASSON-DELMOTTE · Sumito MATOVA · Владимир Владимирович Мацковский [Vladimir Vladimirovich MATSKOVSKY] · 松田好弘[MATSUDA Yoshihiro] · Kenichi MATSUOKA · Enrico MATTEA · Paul A. MAYEWSKI · Lawrence R. MAYO · Ian McDOWELL · Toby MEIERBACHTOL · Hanno MEYER · Nikolay N. MIKHAILOV · Владимир 540 Н. Михаленко [Vladimir N. MIKHALENKO] · B. И. Михалёв [V. I. MIKHALEV] · Evan S. MILES · Katie E. MILES · Gifford H. MILLER · Heinz MILLER · Laurent MINGO · Hans-Erwin MINOR · Masahiro MINOWA · Morihiro MIYAHARA · Dariusz MOCHNACKI · J. Tyge MØLLER · Jangil MOON · John C. MOORE · B. A. Морев [V. A. MOREV] · M. MOSKALEVSKY · Ellen S. MOSLEY-THOMPSON · Hideaki MOTOYAMA · Fritz MÜLLER · Dhruba D. MULMI · Robert MULVANEY · A. Ya. MURAVIEV · Yaroslav D. MURAVYEV · Tavi MURRAY · A. MUTO · David L. NAFTZ · Олег 545 B. Нагорнов [Oleg V. NAGORNOV] · 中尾正[NAKAWO Masayoshi] · Fumio NAKAZAWA · Kazuhiko NAOKI · Chiyuki NARAMA · Hideki NARITA · Renji NARUSE · Francisco J. NAVARRO · Robbe NEYNS · K. W. NICHOLLS · Lindsey

NICHOLSON · H. NICOLET · Peter NIENOW · Станислав А. Никитин [Stanislav A. NIKITIN] · Fumihiko NISHIO · NIU
Hewen · Masashi NIWANO · Matt NOLAN · G. A. NOSENKO · Rune Strand ØDEGÅRD · Klaus OEGGL · H. OESCHGER ·
Atsumu OHMURA · Sachiko OKAMOTO · Ole B. OLESEN · Susanne OLIVIER · Yukihiko ONUMA · Anaïs J. ORSI · Svenn
550 ORVIG · Jaime OTERO · Alexander A. OVSYANNIKOV · Chloë Marie PAICE · Mark PALLETT · Anne PALMER · Erkki
PALOSUO · Finnur PÁLSSON · 庞洪喜[PANG Hongxi] · Tatyana PAPINA · J. G. PAREN · W. S. B. PATERSON · Nicolas
PATRIS · Frank PATTYN · Rickard PETTERSSON · E. C. PETTIT · Luc PIARD · Gianni PIFFER · Sam PIMENTEL
· Jean Francis PINGLOT · Pierre PITTE · M. Я. Плам [M. Ya. PLAM] · Veijo A. POHJOLA · Ulrich POLOM · Ivan A.
PONOMAREV · Victor POPOVNIN · Stacy E. PORTER · Philippe POSSENTI · M. POURCHET · Bernard POUYAUD ·
555 Susanne PREUNKERT · Rainer PRINZ · Т. В. Псарёва [T. V. PSAREVA] · 蒲健辰[PU Jianchen] · PU Tao · B. A. Пухов
[V. A. PUKHOV] · 秦天河[QIN Dahe] · 秦翔[QIN Xiang] · 仇家琪[QIU Jiaqi] · Duncan J. QUINCEY · Antoine RABATEL
· Bernhard T. RABUS · S. Olander RASMUSSEN · IO. B. Райковский [Yu. V. RAYKOVSKY] · Dominique RAYNAUD ·
Н. Г. Разумейко [N. G. RAZUMEIKO] · Niels REEH · 任贾文[REN Jiawen] · Patrick RIESEN · Mirko RINALDI · Andres
RIVERA · E. RÖSSLER · Hans RÖTHLISBERGER · O. B. Рототаева [O. V. ROTOTAEVA] · Nicolas ROUX · Ann V.
560 ROWAN · Lucas RUIZ · Arthur S. RUNDLE · Oleg RYBAK · Claudia RYSER · Yeongjun RYU · 坂井子[SAKAI Akiko] ·
Daiki SAKAKIBARA · Andrey N. SALAMATIN · Masaki SANO · Pamela SANTIBAÑEZ · Yota SATO · Kazuhide SATOW
· Rysbek SATYLKANOV · Matthias SAURER · A. M. Саватюгин [A. M. SAVATYUGIN] · Лев М. Саватюгин [Lev M.
SAVATYUGIN] · Takanobu SAWAGAKI · Ted A. SCAMBOS · Manuel SCHLÄPPI · Forrest SCHOESSOW · Christian G.
SCHOOF · Ulrich SCHOTTERER · Philipp SCHUPPLI · Aurel A. SCHWERZMANN · Margit SCHWIKOWSKI · Valter
565 SCHYTT · Takahiro SEGAWA · Julien SEGUINOT · 古基[SEKO Katsumoto] · Gerald SELIGMAN · Roberto SEPPI · 尚新春
[SHANG Xinchun] · Donghui SHANGGUAN · SHAO Wenzhang · Yaping SHAO · Martin J. SHARP · Robert P. SHARP ·
Tenzing C. SHERPA · Takayuki SHIRAIWA · M. Roxana SIERRA-HERNÁNDEZ · Marie-Louise SIGGAARD-ANDERSEN
· Nadia SIGNER · Е. М. Зингер [E. M. SINGER] · Anna SINISALO · С. А. Синкевич [S. A. SINKEVICH] · Delphine SIX ·
А. А. SKUTIN · Mark E. SMITH · Ireneusz SOBOTA · Сергея Альфредович Сократов [Sergey Alfredovich SOKRATOV] ·
570 Anne SOLGAARD · Johan Ludvig SOLLID · Alvaro SORUCO · Todd A. SOWERS · Nicole SPAULDING · Andrea SPOLAOR
· Francisco A. SQUEO · Mia STAMPE · Wolfgang STARZER · Bernhard STAUFFER · Jørgen Peder STEFFENSEN · Jakob
Friedrich STEINER · Barbara STENNI · 苏珍[SU Zhen] · Sonja SUCKRO · 杉山慎[SUGIYAMA Shin] · Huan SUN · Sainan
SUN · 孙维君[SUN Weijun] · Yafei SUN · Arzhan SURAZAKOV · Stephan SUTER · Keisuke SUZUKI · Toshitaka SUZUKI
· Anders M. SVENSSON · Akiyoshi TAKAHASHI · Shuhei TAKAHASHI · Shuhei TAKENAKA · Nozomu TAKEUCHI ·
575 Masumi TAKIKAWA · Khadga B. THAPA · Emmanuel THIBERT · Martijn P. A. THOMASSEN · David E. THOMPSON
· Lonnie G. THOMPSON · Henrik Højmark THOMSEN · 田立德[TIAN Lide] · Leonhard TOBLER · А. И. Толстой [A. I.
TOLSTOY] · David TONIDANDEL · Павел А. Торопов [Pavel A. TOROPOV] · Yoko TOYAMA · Dennis C. TRABANT ·
Д. С. Троицкий [L. S. TROITSKY] · Akane TSUSHIMA · 津俊[TSUTAKI Shun] · Алексей Викторович Цветков [Aleksey
Viktorovich TSVETKOV] · Г. А. Цыкина [G. A. TSYKINA] · S. TYUFLIN · Tsutomu UCHIDA · Jun UETAKE · Chiara
580 UGLIETTI · Norbert UNTERSTEINER · David URMANN · Рыскул Усубалиев [Ryskul USUBALIEV] · Joseph VALLOT
· Jonas VAN BREEDAM · Roderik S. W. VAN DE WAL · Brice VAN LIEFFERINGE · Ward J. J. VAN PEEL · Lander
VAN TRICHT · W. VAN WYCHEN · Е. В. Василенко [Evgeny V. VASILENKO] · Н. И. Васильев [N. I. VASILIEV] ·
Т. В. Васильева [T. V. VASILYEVA] · Geir VATNE · Carmen P. VEGA · Е. Н. Вилесов [E. N. VILESOV] · Françoise
VIMEUX · Christian VINCENT · Bo M. VINther · Диана Олеговна Владимирова [Diana Olegovna VLADIMIROVA]

585 · Edith VOGEL · Ulrich VON GRAFENSTEIN · H. R. VON GUNTEN · T. A. ВОСТОКОВА [T. A. VOSTOKOVA] · P. H.
Бострецов [R. N. VOSTRETSOV] · Mathias VUILLE · Edwin D. WADDINGTON · H. P. WAGNER · Stephan WAGNER
· Patrick WAGNON · Cameron P. WAKE · Chaomin WANG · Hansheng WANG · 王立伦[WANG Lilun] · Lin WANG · 王
宁练[WANG Ninglian] · Puyu WANG · WANG Shijin · Wenbin WANG · 王文颖[WANG Wenying] · WANG Xiaojun · 王叶
堂[WANG Yetang] · 王玉哲[WANG Yuzhe] · 王泽民[WANG Zemin] · Okitsugu WATANABE · Len G. WATTS · Matthias
590 WECHT · Junfeng WEI · Anker WEIDICK · Gunter WELLER · 温家洪[WEN Jiahong] · Gerd WENDLER · Jonas WICKY
· Johannes WILD · Frank WILHELM · Nat J. WILSON · Bjoern WOLD · WOONG Chung Ji · 邬光剑[WU Guangjian] ·
Lizong WU · Shuang-Ye WU · X. WU · Cunde XIAO · XIE Zichu · XIN Huijuan · 徐柏青[XU Baiqing] · 许君利[XU Junli]
· 徐跃通[XU Yuetong] · B. M. Яковлев [V. M. YAKOVLEV] · Kotaro YAMAGATA · Satoru YAMAGUCHI · Tetsuhide
595 YAMASAKI · Tetsuya YAMAZAKI · 闫明[YAN Ming] · Ji-Woong YANG · 杨梅学[YANG Meixue] · 杨志红[YANG Zhihong]
· 姚檀栋[YAO Tandong] · Xiaojun YAO · Chaolu YI · Yang YI · Yusuke YOKOYAMA · Yoshitaka YOSHIMURA · 尤根
祥[YOU Genxiang] · Pengchun YU · Виктор Сергеевич Загороднов [Victor Serafimovich ZAGORODNOV] · Thomas
ZANONER · Alexander ZAPF · Christian M. ZDANOWICZ · Harry ZEKOLLARI · Tiziana Lazzarina ZENDRINI · Piero
ZENNARO · 张金华[ZHANG Jinhua] · 张通[ZHANG Tong] · 张万昌[ZHANG Wanchang] · 张王滨[ZHANG Wangbin] · 章新
平[ZHANG Xinping] · 张寅生[ZHANG Yinsheng] · Liyun ZHAO · Ping ZHOU · 周韬[ZHOU Tao] · 朱国才[ZHU Guocai] · A.
600 Б. Журавлев [A. B. ZHURAVLEV] · И. А. Зотиков [I. A. ZOTIKOV]

References

- Danish scientific work in Greenland, 1957, *Polar Record*, 9, 452–454, <https://doi.org/10.1017/S0032247400066432>, 1959.
- Danish scientific work in Greenland, 1958, *Polar Record*, 10, 38–40, <https://doi.org/10.1017/S0032247400050609>, 1960.
- Ледник Джанкуат (Центральный Кавказ) [Djankuat Glacier (Central Caucasus)], Водно-ледовый и тепловой баланс горноледниковых бассейнов [Water-ice and heat balance of mountain glacier basins], Гидрометеоиздат [Gidrometeoizdat], http://geolibrary.ru/sites/default/files/2020-10/Djankuat_Glacier_1978.pdf, 1978.
- Journey through time - Topographic maps, <https://www.geocat.ch/geonetwork/srv/eng/catalog.search#/metadata/22287cd6-b75b-4caf-9413-aa3f196548b2>, 2021.
- Abermann, J., Steiner, J. F., Prinz, R., Wecht, M., and Lisager, P.: The Red Rock ice cliff revisited – six decades of frontal, mass and area changes in the Nunatarssuaq area, northwest Greenland, *Journal of Glaciology*, 66, 567–576, <https://doi.org/10.1017/jog.2020.28>, 2020.
- Agassiz, L.: De la température de l'intérieur du glacier [From the temperature of the interior of the glacier], in: Nouvelles études et expériences sur les glaciers actuels: leur structure, leur progression et leurs actions physiques sur le sol [New studies and experiments on current glaciers: their structure, their progression and their physical actions on the ground], pp. 419–434, Victor Masson, <https://gallica.bnf.fr/ark:/12148/bpt6k97697182/f454.item>, 1847.
- Aizen, V., Bren, D., Kreutz, K., and Wake, C.: Paleo-climate and glaciological reconstruction in Central Asia through the collection and analysis of ice cores and instrumental data from the Tien Shan, Tech. rep., Regents of The University of California Santa Barbara (US), 2001.
- Alean, J.: Ice avalanches: some empirical information about their formation and reach, *Journal of Glaciology*, 31, 324 – 333, <https://doi.org/https://doi.org/10.3189/S002214300006663>, 1985.
- An, W., Hou, S., Zhang, W., Wang, Y., Liu, Y., Wu, S., and Pang, H.: Significant recent warming over the northern Tibetan Plateau from ice core $\delta^{18}\text{O}$ records, *Climate of the Past*, 12, 201–211, <https://doi.org/10.5194/cp-12-201-2016>, 2016.
- Aristarain, A. J. and Delmas, R.: First glaciological studies on the James Ross Island Ice Cap, Antarctic Peninsula, *Journal of Glaciology*, 27, 371–379, <https://doi.org/10.3189/S0022143000011412>, 1981.
- Barandun, M.: A novel approach to estimate glacier mass balance in the Tien Shan and Pamir based on transient snowline observations, Ph.D. thesis, University of Fribourg > Department of Geosciences, <https://folia.unifr.ch/global/documents/307523>, 2018.
- Barandun, M.: Personal communication, 2023.
- Barandun, M.: Personal communication, 2024.
- Baranowski, S.: Glaciological investigations and glaciomorphological observations made in 1970 on Werenskiold Glacier and in its forefield, in: Results of investigations of the Polish Scientific Spitsbergen Expeditions 1970-1974, edited by Baranowski, S. and Jahn, A., vol. 1 of *Acta Universitatis Wratislaviensis*, pp. 69–94, 1975.
- Björnsson, H., Gjessing, Y., Hamran, S.-E., Hagen, J. O., Liestøl, O., Pálsson, F., and Erlingsson, B.: The thermal regime of sub-polar glaciers mapped by multi-frequency radio-echo sounding, *Journal of Glaciology*, 42, 23–32, <https://doi.org/10.3189/S0022143000030495>, 1996.

- Blatter, H.: On the thermal regime of Arctic glaciers: A study of the White Glacier, Axel Heiberg Island, and the Laika Glacier, Coburg Island, Canadian Arctic Archipelago, Tech. rep., Eidgenössische Technische Hochschule (ETH) Zürich > Geographisches Institut, 1985.
- Blatter, H.: On the thermal regime of an Arctic valley glacier: A study of White Glacier, Axel Heiberg Island, N.W.T., 640 Canada, *Journal of Glaciology*, 33, 200–211, <https://doi.org/10.3189/S002214300008704>, 1987.
- Blatter, H. and Haeberli, W.: Modelling temperature distribution in Alpine glaciers, *Annals of Glaciology*, 5, 18–22, <https://doi.org/10.3189/1984AoG5-1-18-22>, 1984.
- Blatter, H. and Hutter, K.: Polythermal conditions in arctic glaciers, *Journal of Glaciology*, 37, 261–269, <https://doi.org/10.3189/S0022143000007279>, 1991.
- 645 Blatter, H. and Kappenberger, G.: Mass balance and thermal regime of Laika Ice Cap, Coburg Island, N.W.T., Canada, *Journal of Glaciology*, 34, 102–110, <https://doi.org/10.3189/S002214300009126>, 1988.
- Bohleber, P.: Alpine Ice Cores as Climate and Environmental Archives, in: Oxford Research Encyclopedia of Climate Science, <https://doi.org/10.1093/acrefore/9780190228620.013.743>, 2019.
- 650 Bohleber, P., Erhardt, T., Spaulding, N., Hoffmann, H., Fischer, H., and Mayewski, P.: Temperature and mineral dust variability recorded in two low-accumulation Alpine ice cores over the last millennium, *Climate of the Past*, 14, 21–37, <https://doi.org/10.5194/cp-14-21-2018>, 2018.
- Carturan, L., De Blasi, F., Dinale, R., Dragà, G., Gabrielli, P., Mair, V., Seppi, R., Tonidandel, D., Zanoner, T., Zendrini, T. L., and Dalla Fontana, G.: Data from air, englacial and permafrost temperature measurements on Mt. Ortles (eastern European Alps), <https://doi.org/10.5281/ZENODO.8330289>, 2023a.
- 655 Carturan, L., De Blasi, F., Dinale, R., Dragà, G., Gabrielli, P., Mair, V., Seppi, R., Tonidandel, D., Zanoner, T., Zendrini, T. L., and Dalla Fontana, G.: Modern air, englacial and permafrost temperatures at high altitude on Mt Ortles (3905 m a.s.l.), in the eastern European Alps, *Earth System Science Data*, 15, 4661–4688, <https://doi.org/10.5194/essd-15-4661-2023>, 2023b.
- Chiarle, M., Viani, C., and Deline, P.: Large glacier failures in the Italian Alps over the last 90 years., *Geografia fisica e dinamica quaternaria*, 45, 19–40, 2023.
- 660 Clarke, G. K. C. and Jarvis, G. T.: Post-surge temperatures in Steele Glacier, Yukon Territory, Canada, *Journal of Glaciology*, 16, 261–268, <https://doi.org/10.3189/S0022143000031580>, 1976.
- Clarke, G. K. C., Collins, S. G., and Thompson, D. E.: Flow, thermal structure, and subglacial conditions of a surge-type glacier, *Canadian Journal of Earth Sciences*, 21, 232–240, <https://doi.org/10.1139/e84-024>, 1984.
- 665 Clarke, G. K. C., Fischer, D. A., and Waddington, E. D.: Wind pumping: A potentially significant heat source in ice sheets, in: The Physical Basis of Ice Sheet Modelling (Proceedings of the Vancouver Symposium, August 1987), Publication, pp. 169–180, International Association of Hydrological Sciences, <https://iahs.info/uploads/dms/7236.169-180-170-Clarke.pdf>, 1987.
- Classen, D. F.: Thermal drilling and deep ice-temperature measurements on the Fox Glacier, Yukon, Ph.D. thesis, University of British Columbia, 1970.
- 670 Classen, D. F.: Temperature profiles for the Barnes Ice Cap surge zone, *Journal of Glaciology*, 18, 391–405, <https://doi.org/10.3189/S0022143000021079>, 1977.
- Colgan, W., Sommers, A., Rajaram, H., Abdalati, W., and Frahm, J.: Considering thermal-viscous collapse of the Greenland ice sheet, *Earth's Future*, 3, 252–267, <https://doi.org/10.1002/2015EF000301>, 2015.

- Copland, L., Sharp, M. J., Nienow, P., and Bingham, R. G.: The distribution of basal motion beneath a High Arctic poly-
675 thermal glacier, *Journal of Glaciology*, 49, 407–414, <https://doi.org/10.3189/172756503781830511>, 2003.
- Cuffey, K. M. and Paterson, W. S. B.: *The Physics of Glaciers*, Academic Press, google-Books-ID: Jca2v1u1EKEC, 2010.
- Darms, G. A.: Firntemperaturen auf dem Colle Gnifetti: Zusammenstellung und Analyse bestehender und neuer Temperatur-
profile [Snow temperatures on Colle Gnifetti: Compilation and analysis of existing and new temperature profiles], Ph.D.
thesis, Universität Zürich > Geographisches Institut, https://uzb.swisscovery.slsp.ch/discovery/delivery/41SLSP_UZB:UZB/12464857880005508, 2009.
- 680 Davaa, G.: Glacier monitoring in Mongolia, <https://globalcryospherewatch.org/meetings/salekhard2016/presentations/3.1.3-e-G.Davaa-Mongolia%20cryospheric%20activities.pdf>, 2016.
- Davis, R. M.: Approach roads, Greenland, 1960-1964, Tech. rep., U.S. Cold Regions Research and Engineering Laboratory
(CRREL), <https://erdc-library.erdc.dren.mil/jspui/handle/11681/5708>, 1967.
- 685 Deeley, R. M. and Woodward, H.: The viscosity of ice, *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character*, 81, 250–259, <https://doi.org/10.1098/rspa.1908.0077>, 1908.
- Delcourt, C., Van Liefferinge, B., Nolan, M., and Pattyn, F.: The climate memory of an Arctic polythermal glacier, *Journal of Glaciology*, 59, 1084–1092, <https://doi.org/10.3189/2013JoG12J109>, 2013.
- 690 Di Stefano, E., Baccolo, G., Clemenza, M., Delmonte, B., Fiorini, D., Garzonio, R., Schwikowski, M., and Maggi, V.: Temporal markers in a temperate ice core: insights from ^3H and ^{137}Cs profiles from the Adamello Glacier, *The Cryosphere*, 18, 2865–2874, <https://doi.org/10.5194/tc-18-2865-2024>, 2024.
- Diez, A., Eisen, O., Hofstede, C., Bohleber, P., and Polom, U.: Joint interpretation of explosive and vibroseismic surveys on cold firn for the investigation of ice properties, *Annals of Glaciology*, 54, 201–210, <https://doi.org/10.3189/2013AoG64A200>, 2013.
- 695 Du, J., He, Y., Li, S., Wang, S., Niu, H., Xin, H., and Pu, T.: Mass balance and near-surface ice temperature structure of Baishui Glacier No.1 in Mt. Yulong, *Journal of Geographical Sciences*, 23, 668–678, <https://doi.org/10.1007/s11442-013-1036-4>, 2013.
- Eisen, O., Bauder, A., Lüthi, M., Riesen, P., and Funk, M.: Deducing the thermal structure in the tongue of Gornergletscher, Switzerland, from radar surveys and borehole measurements, *Annals of Glaciology*, 50, 63–70,
700 <https://doi.org/10.3189/172756409789097612>, 2009.
- Faillettaz, J., Sornette, D., and Funk, M.: Numerical modeling of a gravity-driven instability of a cold hanging glacier: reanalysis of the 1895 break-off of Altelgletscher, Switzerland, *Journal of Glaciology*, 57, 817–831, <https://doi.org/10.3189/002214311798043852>, 2011.
- Faillettaz, J., Funk, M., and Vincent, C.: Avalanching glacier instabilities: Review on processes and early warning perspectives, *Reviews of Geophysics*, 53, 203–224, <https://doi.org/10.1002/2014RG000466>, 2015.
- 705 Flowers, G.: Personal communication, 2022.
- Flowers, G. E., Roux, N., Pimentel, S., and Schoof, C. G.: Present dynamics and future prognosis of a slowly surging glacier, *The Cryosphere*, 5, 299–313, <https://doi.org/10.5194/tc-5-299-2011>, 2011.
- Fistrup, B.: Dänische glaziologische Untersuchungen im Internationalen Geophysikalischen Jahr [Danish glaciological investi-
710 gations in the International Geophysical Year], *Polarforschung*, 30, 3–11, <https://doi.org/10.2312/polarforschung.30.1-2.3>, 1960a.

- Fristrup, B.: Studies of four glaciers in Greenland, *Geografisk Tidsskrift* [Danish Journal of Geography], 59, 89–102, 1960b.
- Fristrup, B.: Danish glaciological investigations in Greenland, in: *Geology of the Arctic: Proceedings of the First International Symposium on Arctic Geology*, edited by Raasch, G. O., vol. 2, pp. 735–746, University of Toronto Press, 1961.
- 715 Fritzsch, D., Wilhelms, F., Savatyugin, L. M., Pinglot, J. F., Meyer, H., Hubberten, H.-W., and Miller, H.: A new deep ice core from Akademii Nauk Ice Cap, Severnaya Zemlya, Eurasian Arctic: first results, *Annals of Glaciology*, 35, 25–28, <https://doi.org/10.3189/172756402781816645>, 2002.
- Fujii, Y., Kameda, T., Nishio, F., Suzuki, K., Kohno, M., Nakazawa, F., Uetake, J., Savatyugin, L. M., Arkhipov, S. M., Ponomarev, I. A., and Mikhailov, N. N.: Outline of Japan-Russia joint Glaciological Research on Sofiyevkiy Glacier, Russian 720 Altai Mountains in 2000 and 2001, *Bulletin of Glaciological Research*, 19, 53–58, 2002.
- Gabrielli, P., Carturan, L., Gabrieli, J., Dinale, R., Krainer, K., Hausmann, H., Davis, M., Zagorodnov, V., Seppi, R., Barbante, C., Dalla Fontana, G., and Thompson, L. G.: Atmospheric warming threatens the untapped glacial archive of Ortles Mountain, South Tyrol, *Journal of Glaciology*, 56, 843–853, <https://doi.org/10.3189/002214310794457263>, 2010.
- Gabrielli, P., Barbante, C., Carturan, L., Cozzi, G., Dalla Fontana, G., Dinale, R., Dragà, G., Gabrieli, J., Kehrwald, N., Mair, 725 V., Mikhalenko, V., Piffer, G., Rinaldi, M., Seppi, R., Spolaor, A., Thompson, L. G., and Tonidandel, D.: Discovery of cold ice in a new drilling site in the eastern European Alps, *Geografia Fisica e Dinamica Quaternaria* [Physical Geography and Quaternary Dynamics], 35, 101–105, 2012.
- Gabrielli, P., Barbante, C., Bertagna, G., Bertó, M., Binder, D., Carton, A., Carturan, L., Cazorzi, F., Cozzi, G., Dalla Fontana, G., Davis, M., De Blasi, F., Dinale, R., Dragà, G., Dreossi, G., Festi, D., Frezzotti, M., Gabrieli, J., Galos, S. P., Ginot, P., Heidenwolf, P., Jenk, T. M., Kehrwald, N., Kenny, D., Magand, O., Mair, V., Mikhalenko, V., Lin, P. N., Oeggl, K., Piffer, G., Rinaldi, M., Schotterer, U., Schwikowski, M., Seppi, R., Spolaor, A., Stenni, B., Tonidandel, D., Uglietti, C., Zagorodnov, V., Zanoner, T., and Zennaro, P.: Age of the Mt. Ortles ice cores, the Tyrolean Iceman and glaciation of the highest summit of South Tyrol since the Northern Hemisphere Climatic Optimum, *The Cryosphere*, 10, 2779–2797, <https://doi.org/10.5194/tc-10-2779-2016>, 2016.
- 730 735 Gagliardini, O. and Vincent, C.: Personal communication, 2023.
- Gastaldello, M. and Hoelzle, M.: Personal communication, 2024a.
- Gastaldello, M. and Hoelzle, M.: Personal communication, 2024b.
- Gilbert, A. and Vincent, C.: Atmospheric temperature changes over the 20th century at very high elevations in the European Alps from englacial temperatures, *Geophysical Research Letters*, 40, 2102–2108, <https://doi.org/10.1002/grl.50401>, 2013.
- 740 Gilbert, A., Wagnon, P., Vincent, C., Ginot, P., and Funk, M.: Atmospheric warming at a high-elevation tropical site revealed by englacial temperatures at Illimani, Bolivia (6340 m above sea level, 16°S, 67°W), *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2009JD012961>, 2010.
- Gilbert, A., Vincent, C., Wagnon, P., Thibert, E., and Rabatel, A.: The influence of snow cover thickness on the thermal regime of Tête Rousse Glacier (Mont Blanc range, 3200 m a.s.l.): Consequences for outburst flood hazards and glacier 745 response to climate change, *Journal of Geophysical Research: Earth Surface*, 117, <https://doi.org/10.1029/2011JF002258>, 2012.
- Gilbert, A., Vincent, C., Gagliardini, O., Krug, J., and Berthier, E.: Assessment of thermal change in cold avalanching glaciers in relation to climate warming, *Geophysical Research Letters*, 42, 6382–6390, <https://doi.org/10.1002/2015GL064838>, 2015.

- Gilbert, A., Flowers, G. E., Miller, G. H., Rabus, B. T., Van Wychen, W., Gardner, A. S., and Copland, L.: Sensitivity of Barnes Ice Cap, Baffin Island, Canada, to climate state and internal dynamics, *Journal of Geophysical Research: Earth Surface*, 121, 1516–1539, <https://doi.org/10.1002/2016JF003839>, 2016.
- Gilbert, A., Leinss, S., Kargel, J., Kääb, A., Gascoin, S., Leonard, G., Berthier, E., Karki, A., and Yao, T.: Mechanisms leading to the 2016 giant twin glacier collapses, Aru Range, Tibet, *The Cryosphere*, 12, 2883–2900, <https://doi.org/10.5194/tc-12-2883-2018>, 2018.
- Gilbert, A., Sinisalo, A., Gurung, T. R., Fujita, K., Maharjan, S. B., Sherpa, T. C., and Fukuda, T.: The influence of water percolation through crevasses on the thermal regime of a Himalayan mountain glacier, *The Cryosphere*, 14, 1273–1288, <https://doi.org/10.5194/tc-14-1273-2020>, 2020.
- Ginot, P., Kull, C., Schotterer, U., Schwikowski, M., and Gäggeler, H. W.: Glacier mass balance reconstruction by sublimation induced enrichment of chemical species on Cerro Tapado (Chilean Andes), *Climate of the Past*, 2, 21–30, <https://doi.org/10.5194/cp-2-21-2006>, 2006.
- Glen, J. W.: The Stability of Ice-Dammed Lakes and other Water-Filled Holes in Glaciers, *Journal of Glaciology*, 2, 316–318, <https://doi.org/10.3189/S0022143000025132>, 1954.
- Goel, V., Brown, J., and Matsuoka, K.: In-situ temperatures in the top 20 m of firn near the summit of Blåskimen Island, Western Dronning Maud Land, <https://doi.org/10.21334/NPOLAR.2017.3845E964>, 2017a.
- Goel, V., Brown, J., and Matsuoka, K.: Glaciological settings and recent mass balance of Blåskimen Island in Dronning Maud Land, Antarctica, *The Cryosphere*, 11, 2883–2896, <https://doi.org/10.5194/tc-11-2883-2017>, 2017b.
- GTN-G: GTN-G Glacier Regions (GlacReg), <https://doi.org/10.5904/gtn-glacreg-2017-07>, 2017.
- Guo, W., Liu, S., Xu, J., Wu, L., Shangguan, D., Yao, X., Wei, J., Bao, W., Yu, P., Liu, Q., and Jiang, Z.: The second Chinese glacier inventory: data, methods and results, *Journal of Glaciology*, 61, 357–372, <https://doi.org/10.3189/2015JoG14J209>, 2015.
- Gurung, T. R.: Personal communication, 2022.
- Gäggeler, H., von Gunten, H. R., Rössler, E., Oeschger, H., and Schotterer, U.: 210Pb-dating of cold alpine firn/ice cores from Colle Gnifetti, Switzerland, *Journal of Glaciology*, 29, 165–177, <https://doi.org/10.3189/S002214300005220>, 1983.
- Haeberli, W.: Eistemperaturen in den Alpen [Ice temperatures in the Alps], *Zeitschrift für Gletscherkunde und Glazialgeologie*, 11, 203–220, 1976.
- Haeberli, W. and Alean, J.: Temperature and accumulation of high altitude firn in the Alps, *Annals of Glaciology*, 6, 161–163, <https://doi.org/10.3189/1985AoG6-1-161-163>, 1985.
- Haeberli, W. and Funk, M.: Borehole temperatures at the Colle Gnifetti core-drilling site (Monte Rosa, Swiss Alps), *Journal of Glaciology*, 37, 37–46, <https://doi.org/10.3189/S0022143000042775>, 1991.
- Haeberli, W., Frauenfelder, R., Kääb, A., and Wagner, S.: Characteristics and potential climatic significance of “miniature ice caps” (crest- and cornice-type low-altitude ice archives), *Journal of Glaciology*, 50, 129–136, <https://doi.org/10.3189/172756504781830330>, 2004.
- Hagen, J. O.: Temperature distribution in Brøggerbreen, Lovenbreen and Kongsvegen, Northwest Spitsbergen, in: Field Workshop on Glaciological Research in Svalbard: Current Problems at the Polish Polar Station, Hornsund, Spitsbergen 26-30 April 1992, 1992.

- Hager, P.: Glaziologische Untersuchungen am Gipfelgrat des Vadret dal Corvatsch: Thermik und Oberflächenprozesse, Ph.D. thesis, Universität Zürich > Geographisches Institut, 2002.
- Hammer, C. U.: Ice core drilling, in: Report on activities and results 1993-1995 for Hans Tavsen Ice Cap Project - Glacier and Climate Change Research, North Greenland, edited by Reeh, N., NMRs (Nordisk Minister Råd) miljøforskningsprogram - klimaforskning, pp. 16–28, Dansk Polar Center, 1995.
- 790 Han, Y., Jun, S. J., Miyahara, M., Lee, H.-G., Ahn, J., Woong, C. J., Hur, S. D., and Hong, S. B.: Shallow ice-core drilling on Styx Glacier, northern Victoria Land, Antarctica in the 2014-2015 summer, Journal of the Geological Society of Korea, 51, 343, <https://doi.org/10.14770/jgsk.2015.51.3.343>, 2015.
- Harrison, J.: Expedition area, Axel Heiberg Island, Canadian Arctic Archipelago, 1963.
- 795 Harrison, W. D.: Temperature of a temperate glacier, Journal of Glaciology, 11, 15–29, <https://doi.org/10.3189/S0022143000022450>, 1972.
- Harrison, W. D., Mayo, L. R., and Trabant, D. C.: Temperature measurements on Black Rapids Glacier, Alaska, 1973, in: Climate of the Arctic: Twenty-Fourth Alaska Science Conference, August 15-17, 1973: Fairbanks, edited by Weller, G. and Bowling, S. A., pp. 350–352, University of Alaska > Geophysical Institute, <https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=9dbf174017ca2945dec1f6266164b9707cdd33f3>, 1975.
- 800 Hasholt, B.: A new map of the Midluagkat glacier - a preliminary report, Geografisk Tidsskrift [Danish Journal of Geography], 87, 19–21, 1987.
- Heim, A., Pasquier, L. D., and Forel, F. A.: Die Gletscherlawine an der Altels am 11. September 1895, Zürcher und Furrer, google-Books-ID: wak3ZUof9yUC, 1895.
- 805 Herren, P.-A., Eichler, A., Machguth, H., Papina, T., Tobler, L., Zapf, A., and Schwikowski, M.: The onset of Neoglaciation 6000 years ago in western Mongolia revealed by an ice core from the Tsambagarav mountain range, Quaternary Science Reviews, 69, 59–68, <https://doi.org/10.1016/j.quascirev.2013.02.025>, 2013.
- Hodgkins, R., Hagen, J. O., and Hamran, S.-E.: 20th century mass balance and thermal regime change at Scott Turnerbreen, Svalbard, Annals of Glaciology, 28, 216–220, <https://doi.org/10.3189/172756499781821986>, 1999.
- 810 Hoelzle, M.: Englacial temperature, in: The Swiss Glaciers 2003/04 and 2004/05, edited by Bauder, A. and Rüegg, R., no. 125-126 in Glaciological Report, pp. 71–74, Swiss Academy of Sciences (SCNAT) > Cryospheric Commission (EKK), https://doi.glamos.ch/pubs/glrep/glrep_125-126.html, 2009.
- Hoelzle, M.: Englacial temperature, in: The Swiss Glaciers 2007/08 and 2008/09, edited by Bauder, A., Steffen, S., and Usselmann, S., no. 129-130 in Glaciological Report, pp. 69–73, Swiss Academy of Sciences (SCNAT) > Cryospheric Commission (EKK), 2014.
- 815 Hoelzle, M.: Englacial temperatures, in: The Swiss Glaciers 2013/14 and 2014/15, edited by Bauder, A., no. 135-136 in Glaciological Report, pp. 93–98, Swiss Academy of Sciences (SCNAT) > Cryospheric Commission (EKK), 2017.
- Hoelzle, M.: Englacial temperature, in: The Swiss Glaciers 2019/20 and 2020/21, edited by Bauder, A., Huss, M., and Linsbauer, A., no. 141-142 in Glaciological Report, pp. 115–119, Swiss Academy of Sciences (SCNAT) > Cryospheric Commission (EKK), 2022.
- 820 Hoelzle, M., Darms, G., Lüthi, M. P., and Suter, S.: Evidence of accelerated englacial warming in the Monte Rosa area, Switzerland/Italy, The Cryosphere, 5, 231–243, <https://doi.org/10.5194/tc-5-231-2011>, 2011.

- Hoelzle, M., Huss, M., Kronenberg, M., Machguth, H., and Mattea, E.: Englacial temperature, in: The Swiss Glaciers 2017/18 and 2018/19, edited by Bauder, A., Huss, M., and Linsbauer, A., no. 139-140 in Glaciological Report, pp. 111–115, Swiss Academy of Sciences (SCNAT) > Cryospheric Commission (EKK), 2020.
- Hooke, R. L.: Pleistocene ice at the base of the Barnes Ice Cap, Baffin Island, N.W.T., Canada, *Journal of Glaciology*, 17, 49–59, <https://doi.org/10.3189/S0022143000030719>, 1976.
- Hooke, R. L., Alexander Jr., E. C., and Gustafson, R. J.: Temperature profiles in the Barnes Ice Cap, Baffin Island, Canada, and heat flux from the subglacial terrane, *Canadian Journal of Earth Sciences*, 17, 1174–1188, <https://doi.org/10.1139/e80-124>, 1980.
- Hou, S., Qin, D., Jouzel, J., Masson-Delmotte, V., von Grafenstein, U., Landais, A., Caillon, N., and Chappellaz, J.: Age of Himalayan bottom ice cores, *Journal of Glaciology*, 50, 467–468, <https://doi.org/10.3189/172756504781829981>, 2004.
- Hou, S., Chappellaz, J., Jouzel, J., Chu, P. C., Masson-Delmotte, V., Qin, D., Raynaud, D., Mayewski, P. A., Lipenkov, V. Y., and Kang, S.: Summer temperature trend over the past two millennia using air content in Himalayan ice, *Climate of the Past*, 3, 89–95, <https://doi.org/10.5194/cp-3-89-2007>, 2007.
- Hou, S., Jenk, T. M., Zhang, W., Wang, C., Wu, S., Wang, Y., Pang, H., and Schwikowski, M.: Age ranges of the Tibetan ice cores with emphasis on the Chongce ice cores, western Kunlun Mountains, *The Cryosphere*, 12, 2341–2348, <https://doi.org/10.5194/tc-12-2341-2018>, 2018.
- Huang, M.: On the temperature distribution of glaciers in China, *Journal of Glaciology*, 36, 210–216, <https://doi.org/10.3189/S002214300000945X>, 1990.
- Hubbard, B., Miles, K., Doyle, S., Quincey, D., and Miles, E.: Ice temperature time-series from sensors installed in boreholes drilled into Khumbu Glacier, Nepal, in 2017 and 2018 as part of EverDrill research project, <https://doi.org/10.5285/32ECD5F4-1F00-4EEB-BD1E-5EAFFB60F556>, 2021.
- Hughes, T. P. and Seligman, G.: The temperature, melt water movement and density increase in the névé of an Alpine glacier, *Geophysical Journal International*, 4, 616–647, <https://doi.org/10.1111/j.1365-246X.1939.tb02922.x>, 1939.
- Huss, M. and Fischer, M.: Sensitivity of Very Small Glaciers in the Swiss Alps to Future Climate Change, *Frontiers in Earth Science*, 4, 2016.
- Iida, H., Watanabe, O., and Takikawa, M.: First results from Himalayan Glacier Boring Project in 1981–1982: Part II. Studies on internal structure and transformation process from snow to ice of Yala Glacier, Langtang Himal, Nepal, *Bulletin of Glacier Research*, 2, 25–33, 1984.
- Irvine-Fynn, T. D. L., Hodson, A. J., Moorman, B. J., Vatne, G., and Hubbard, A. L.: Polythermal glacier hydrology: a review, *Reviews of Geophysics*, 49, RG4002, <https://doi.org/10.1029/2010RG000350>, 2011.
- Jacquemart, M., Loso, M., Leopold, M., Welty, E., Berthier, E., Hansen, J. S., Sykes, J., and Tiampo, K.: What drives large-scale glacier detachments? Insights from Flat Creek glacier, St. Elias Mountains, Alaska, *Geology*, 48, 703–707, <https://doi.org/10.1130/G47211.1>, 2020.
- Jacquemart, M., Welty, E., Carcanade, G., Gastaldello, M., Van Tricht, L., Flowers, G. E., Sugiyama, S., Gurung, T. R., Prinz, R., Abermann, J., Steiner, J. F., Barandun, M., Gagliardini, O., Vincent, C., Thompson, L. G., Zhang, T., and Hoelzle, M.: glenglac: Global englacial temperature database, <https://doi.org/10.5281/zenodo.14696283>, 2025.
- Jania, J., Mochnacki, D., and Gądek, B.: The thermal structure of Hansbreen, a tidewater glacier in southern Spitsbergen, Svalbard, *Polar Research*, 15, 53–66, <https://doi.org/10.3402/polar.v15i1.6636>, 1996.

- Jarvis, G. T. and Clarke, G. K. C.: Thermal effects of crevassing on Steele Glacier, Yukon Territory, Canada, *Journal of Glaciology*, 13, 243–254, <https://doi.org/10.3189/S0022143000023054>, 1974.
- Jarvis, G. T. and Clarke, G. K. C.: The thermal regime of Trapridge Glacier and its relevance to glacier surging, *Journal of Glaciology*, 14, 235–250, <https://doi.org/10.3189/S0022143000021729>, 1975.
- 865 Jiao, X., Dong, Z., Baccolo, G., Chen, X., Qin, X., and Shao, Y.: Provenance of aeolian dust revealed by (^{234}U / ^{238}U) activity ratios in cryoconites from high-altitude glaciers in western China and its transport and settlement mechanisms, *Journal of Geophysical Research: Earth Surface*, 128, <https://doi.org/10.1029/2023JF007227>, 2023.
- Jouzel, J., Legrand, M., Pinglot, J. F., and Pourchet, M.: Chronologie d'un carottage de 20 m au col du Dôme (Massif du Mont Blanc) [Chronology of a core sample in the Col du Dôme (Mont Blanc massif)], *La Houille Blanche*, 70, 491–498, 870 <https://doi.org/10.1051/lhb/1984035>, 1984.
- Kameda, T., Takahashi, S., Goto-Azuma, K., Kohshima, S., Watanabe, O., and Hagen, J. O.: First report of ice core analyses and borehole temperatures on the highest icefield on western Spitsbergen in 1992, *Bulletin of Glacier Research*, pp. 51–61, 1993.
- 875 Karušs, J., Lamsters, K., Sobota, I., Ješkins, J., Džeriņš, P., and Hodson, A.: Drainage system and thermal structure of a high Arctic polythermal glacier: Waldemarbreen, western Svalbard, *Journal of Glaciology*, 68, 591–604, <https://doi.org/10.1017/jog.2021.125>, 2022.
- Kawamura, T., Fujii, Y., Satow, K., Kamiyama, K., Izumi, K., Kameda, T., Watanabe, O., Kawaguchi, S., Wold, B., and Gjessing, Y.: Glaciological characteristics of cores drilled on Jostedalsbreen, Southern Norway, *Proceedings of the NIPR Symposium on Polar Meteorology and Glaciology*, 2, 152–160, <https://doi.org/10.15094/00003576>, 1989.
- 880 Kawamura, T., Kameda, T., and Izumi, K.: Preliminary results of structural analyses of an 85.6m deep ice core retrieved from Hoghetta Ice Dome in Northern Spitsbergen, Svalbard, *Bulletin of Glacier Research*, pp. 77–83, 1991.
- Keck, L.: Climate significance of stable isotope records from Alpine ice cores, Ph.D. thesis, Ruprecht Karl University of Heidelberg > Combined Faculties for the Natural Sciences and for Mathematics, <https://archiv.ub.uni-heidelberg.de/volltextserver/1837/1/summary.pdf>, 2001.
- 885 Khalzan, P., Sakai, A., and Fujita, K.: Mass balance of four Mongolian glaciers: in-situ measurements, long-term reconstruction and sensitivity analysis, *Frontiers in Earth Science*, 9, 785 306, <https://doi.org/10.3389/feart.2021.785306>, 2022.
- Khromova, T. E., Nosenko, G. A., Glazovsky, A. F., Muraviev, A. Y., Nikitin, S. A., and Lavrentiev, I. I.: New inventory of Russian glaciers based on satellite data (2016–2019), *Water Resources*, 49, S55–S68, <https://doi.org/10.1134/S0097807822070065>, 2022.
- 890 Kinnard, C., Zdanowicz, C. M., Fisher, D. A., and Wake, C. P.: Calibration of an ice-core glaciochemical (sea-salt) record with sea-ice variability in the Canadian Arctic, *Annals of Glaciology*, 44, 383–390, <https://doi.org/10.3189/172756406781811349>, 2006.
- Kinnard, C., Ginot, P., Surazakov, A., MacDonell, S., Nicholson, L., Patris, N., Rabatel, A., Rivera, A., and Squeo, F. A.: Mass balance and climate history of a high-altitude glacier, desert Andes of Chile, *Frontiers in Earth Science*, 8, <https://doi.org/10.3389/feart.2020.00040>, 2020.
- Koerner, R. M.: Fabric analysis of a core from the Meighen Ice Cap, Northwest Territories, Canada, *Journal of Glaciology*, 7, 421–430, <https://doi.org/10.3189/S0022143000020621>, 1968.

- Kotlyakov, V. M., Arkhipov, S. M., Henderson, K. A., and Nagornov, O. V.: Deep drilling of glaciers in Eurasian Arctic as a source of paleoclimatic records, *Quaternary Science Reviews*, 23, 1371–1390, 900 https://doi.org/10.1016/j.quascirev.2003.12.013, 2004.
- Kronenberg, M.: Changing glacier firn in Central Asia and its impact on glacier mass balance, Ph.D. thesis, University of Fribourg > Department of Geosciences, 2022.
- Kronenberg, M., Lavrentiev, I., Machguth, H., Schuppli, P., and Wicky, J.: Gregoriev firn data, https://doi.org/10.5281/zenodo.7113282, 2022.
- Kubyshkin, N. V., Buzin, I. V., Skutin, A. A., and Glazovsky, A. F.: Determination of the area of generation of big icebergs in the Barents Sea – temperature distribution analysis, in: Proceedings of the Sixteenth (2006) International Offshore and Polar Engineering Conference, The International Society of Offshore and Polar Engineers, https://onepetro.org/ISOPEIOPEC/proceedings-abstract/ISOPE06/All-ISOPE06/9744, 2006.
- Kääb, A., Leinss, S., Gilbert, A., Bühler, Y., Gascoin, S., Evans, S. G., Bartelt, P., Berthier, E., Brun, F., Chao, W.-A., 910 Farinotti, D., Gimbert, F., Guo, W., Huggel, C., Kargel, J. S., Leonard, G. J., Tian, L., Treichler, D., and Yao, T.: Massive collapse of two glaciers in western Tibet in 2016 after surge-like instability, *Nature Geoscience*, 11, 114–120, https://doi.org/10.1038/s41561-017-0039-7, 2018.
- Kääb, A., Jacquemart, M., Gilbert, A., Leinss, S., Girod, L., Huggel, C., Falaschi, D., Ugalde, F., Petrakov, D., Chernomorets, S., Dokukin, M., Paul, F., Gascoin, S., Berthier, E., and Kargel, J. S.: Sudden large-volume detachments of low-angle 915 mountain glaciers – more frequent than thought?, *The Cryosphere*, 15, 1751–1785, https://doi.org/10.5194/tc-15-1751-2021, 2021.
- Laternser, M.: Firntemperaturmessungen in den Schweizer Alpen, Ph.D. thesis, Eidgenössische Technische Hochschule (ETH) Zürich, 1992.
- Law, R., Christoffersen, P., Hubbard, B., Doyle, S. H., Chudley, T. R., Schoonman, C. M., Bougamont, M., des 920 Tombe, B., Schilperoort, B., Kechavarzi, C., Booth, A., and Young, T. J.: Thermodynamics of a fast-moving Greenlandic outlet glacier revealed by fiber-optic distributed temperature sensing, *Science Advances*, 7, eabe7136, https://doi.org/10.1126/sciadv.abe7136, 2021.
- Lee, I.: Borehole tilt sensor data for Jarvis Glacier, Alaska (2017-2018), https://doi.org/10.18739/A2348GG12, 2019.
- Lee, I. R., Hawley, R. L., Bernsen, S., Campbell, S. W., Clemens-Sewall, D., Gerbi, C. C., and Hruby, K.: A novel tilt sensor 925 for studying ice deformation: application to streaming ice on Jarvis Glacier, Alaska, *Journal of Glaciology*, 66, 74–82, https://doi.org/10.1017/jog.2019.84, 2020.
- Lemark, A.: A study of the Flade Isblink ice cap using a simple ice flow model, Ph.D. thesis, University of Copenhagen, https://nbi.ku.dk/english/theses/masters-theses/andreas-lemark, 2010.
- Li, Y., Tian, L., Yi, Y., Moore, J. C., Sun, S., and Zhao, L.: Simulating the evolution of Qiangtang No. 1 Glacier in the central 930 Tibetan Plateau to 2050, *Arctic, Antarctic, and Alpine Research*, 49, 1–12, https://doi.org/10.1657/AAAR0016-008, 2017.
- Liu, L., Jiang, L., Sun, Y., Wang, H., Yi, C., and Hsu, H.: Morphometric controls on glacier mass balance of the Puruogangri Ice Field, central Tibetan Plateau, *Water*, 8, 496, https://doi.org/10.3390/w8110496, 2016.
- Liu, Y., Hou, S., Wang, Y., and Linlin, S.: Distribution of borehole temperature at four high-altitude alpine glaciers in Central Asia, *Journal of Mountain Science*, 6, 221–227, https://doi.org/10.1007/s11629-009-0254-9, 2009.

- 935 Lliboutry, L., Briat, M., Creseveur, M., and Pourchet, M.: 15m deep temperatures in the glaciers of Mont Blanc (French Alps), *Journal of Glaciology*, 16, 197–203, <https://doi.org/10.3189/S0022143000031531>, 1976.
- Løkkegaard, A., Mankoff, K. D., Zdanowicz, C., Clow, G. D., Lüthi, M. P., Doyle, S. H., Thomsen, H. H., Fisher, D., Harper, J., Aschwanden, A., Vinther, B. M., Dahl-Jensen, D., Zekollari, H., Meierbacholt, T., McDowell, I., Humphrey, N., Solgaard, A., Karlsson, N. B., Khan, S. A., Hills, B., Law, R., Hubbard, B., Christoffersen, P., Jacquemart, M., Seguinot, J., Fausto, R. S., and Colgan, W. T.: Greenland and Canadian Arctic ice temperature profiles database, *The Cryosphere*, 17, 3829–2023, <https://doi.org/10.5194/tc-17-3829-2023>, 2023.
- 940 Lüthi, M. P.: Experimental and numerical investigation of a firn covered cold glacier and a polythermal ice stream: case studies at Colle Gnifetti and Jakobshavns Isbræ, Ph.D. thesis, Eidgenössischen Technischen Hochschule (ETH) Zürich, 1999.
- Lüthi, M. P. and Funk, M.: Modelling heat flow in a cold, high-altitude glacier: interpretation of measurements from Colle 945 Gnifetti, Swiss Alps, *Journal of Glaciology*, 47, 314–324, <https://doi.org/10.3189/172756501781832223>, 2001.
- Machguth, H., Eichler, A., Schwikowski, M., Brütsch, S., Mattea, E., Kutuzov, S., Heule, M., Usualiev, R., Belekov, S., Mikhaleko, V. N., Hoelzle, M., and Kronenberg, M.: Firn core data from Grigoriev Ice Cap, 1962 to present, <https://doi.org/10.5281/zenodo.10082961>, 2023a.
- Machguth, H., Eichler, A., Schwikowski, M., Brütsch, S., Mattea, E., Kutuzov, S., Heule, M., Usualiev, R., Belekov, S., 950 Mikhaleko, V. N., Hoelzle, M., and Kronenberg, M.: Fifty years of firn evolution on Grigoriev ice cap, Tien Shan, Kyrgyzstan, *The Cryosphere*, 18, 1633–1646, <https://doi.org/10.5194/tc-18-1633-2024>, 2023b.
- Mankoff, K.: Renland88, https://github.com/GEUS-Glaciology-and-Climate/greenland_ice_borehole_temperature_profiles/tree/8b28d0e7333426426ac377615663ba735e5623f9/boreholes/Renland88, 2022a.
- Mankoff, K.: PrinceWales05, https://github.com/GEUS-Glaciology-and-Climate/greenland_ice_borehole_temperature_profiles/tree/8b28d0e7333426426ac377615663ba735e5623f9/boreholes/PrinceWales05, 2022b.
- 955 Mankoff, K.: Penny96, https://github.com/GEUS-Glaciology-and-Climate/greenland_ice_borehole_temperature_profiles/tree/8b28d0e7333426426ac377615663ba735e5623f9/boreholes/Penny96, 2022c.
- Mankoff, K.: FladeIsblink06, https://github.com/GEUS-Glaciology-and-Climate/greenland_ice_borehole_temperature_profiles/tree/8b28d0e7333426426ac377615663ba735e5623f9/boreholes/FladeIsblink06, 2022d.
- 960 Mankoff, K.: Devon98, https://github.com/GEUS-Glaciology-and-Climate/greenland_ice_borehole_temperature_profiles/tree/8b28d0e7333426426ac377615663ba735e5623f9/boreholes/Devon98, 2022e.
- Mankoff, K., Løkkegaard, A., Colgan, W., Thomsen, H., Clow, G., Fisher, D., Zdanowicz, C., Lüthi, M. P., Vinther, B., MacGregor, J. A., McDowell, I., Zekollari, H., Meierbacholt, T., Doyle, S., Law, R., Hills, B., Harper, J., Humphrey, N., Hubbard, B., Christoffersen, P., and Jacquemart, M.: Greenland deep ice temperature database, <https://doi.org/10.22008/FK2/3BVF9V>, 2022.
- 965 Marchenko, S., Pohjola, V. A., Pettersson, R., van Pelt, W. J. J., Vega, C. P., Machguth, H., Bøggild, C. E., and Isaksson, E.: A plot-scale study of firn stratigraphy at Lomonosovfonna, Svalbard, using ice cores, borehole video and GPR surveys in 2012–14, *Journal of Glaciology*, 63, 67–78, <https://doi.org/10.1017/jog.2016.118>, 2017.
- Markl, V. G. and Wagner, H. P.: Messungen von Eis- und Firntemperaturen am Hintereisferner (Ötztaler Alpen) [Measurements 970 of ice and firn temperatures on the Hintereisferner (Ötztal Alps)], *Zeitschrift für Gletscherkunde und Glazialgeologie*, 13, 261–265, 1977.

- Masiokas, M. H., Rabatel, A., Rivera, A., Ruiz, L., Pitte, P., Ceballos, J. L., Barcaza, G., Soruco, A., Bown, F., Berthier, E., Dussaillant, I., and MacDonell, S.: A review of the current state and recent changes of the Andean cryosphere, *Frontiers in Earth Science*, 8, <https://doi.org/10.3389/feart.2020.00099>, 2020.
- 975 Matsuda, Y., Sakai, A., Fujita, K., Nakawo, M., Duan, K., Pu, J., and Yao, T.: Glaciological observations on July 1st glacier in Qilian Mountains of west China during summer 2002, *Bulletin of Glaciological Research*, 21, 31–36, 2004.
- Matsuoka, K. and Naruse, R.: Mass balance features derived from a firn core at Hielo Patagónico Norte, South America, Arctic, Antarctic, and Alpine Research, 31, 333–340, <https://doi.org/10.1080/15230430.1999.12003318>, 1999.
- 980 Mattea, E.: Measuring and modelling changes in the firn at Colle Gnifetti, 4400 m a.s.l., Swiss Alps, Ph.D. thesis, University of Fribourg > Department of Geosciences, [https://bigweb.unifr.ch/Science/Geosciences/GeographyTechnical/Secretary/Pub/Publications/Geography/SelectedBachelorMasterThesis/2020/Mattea_E._\(2020\)_M_Measuring_modelling_changes_Colle_Gnifetti.pdf](https://bigweb.unifr.ch/Science/Geosciences/GeographyTechnical/Secretary/Pub/Publications/Geography/SelectedBachelorMasterThesis/2020/Mattea_E._(2020)_M_Measuring_modelling_changes_Colle_Gnifetti.pdf), 2020.
- 985 Mattea, E., Machguth, H., Kronenberg, M., van Pelt, W., Bassi, M., and Hoelzle, M.: Firn changes at Colle Gnifetti revealed with a high-resolution process-based physical model approach, *The Cryosphere*, 15, 3181–3205, <https://doi.org/10.5194/tc-15-3181-2021>, 2021.
- Mayewski, P.: Colle Gnifetti ice core (KCC) progress report (year one)—Arcadia ice core proposal: Initiatives on the science of the human past, Tech. rep., University of Maine, https://digitalcommons.library.umaine.edu/orsp_reports/29, 2014.
- 990 Mikhalenko, V., Sokratov, S., Kutuzov, S., Ginot, P., Legrand, M., Preunkert, S., Lavrentiev, I., Kozachek, A., Ekyakin, A., Fain, X., Lim, S., Schotterer, U., Lipenkov, V., and Toropov, P.: Investigation of a deep ice core from the Elbrus western plateau, the Caucasus, Russia, *The Cryosphere*, 9, 2253–2270, <https://doi.org/10.5194/tc-9-2253-2015>, 2015.
- Miles, K. E., Hubbard, B., Quincey, D. J., Miles, E. S., Sherpa, T. C., Rowan, A. V., and Doyle, S. H.: Polythermal structure of a Himalayan debris-covered glacier revealed by borehole thermometry, *Scientific Reports*, 8, 16825, <https://doi.org/10.1038/s41598-018-34327-5>, 2018.
- 995 Miles, K. E., Miles, E. S., Hubbard, B., Quincey, D. J., Rowan, A. V., and Pallett, M.: Instruments and methods: hot-water borehole drilling at a high-elevation debris-covered glacier, *Journal of Glaciology*, 65, 822–832, <https://doi.org/10.1017/jog.2019.49>, 2019.
- Muñoz Sabater, J.: ERA5-Land monthly averaged data from 1950 to present, <https://doi.org/10.24381/cds.68d2bb30>, 2019.
- Møller, J. T.: Glaciers in Upernivik Ø: With special reference to the periglacial phenomena, *Geografisk Tidsskrift [Danish Journal of Geography]*, 58, 30–53, 1959.
- 1000 Müller, F.: Englacial temperature measurements, in: Jacobsen-McGill Arctic Research Expedition to Axel Heiberg Island, Queen Elizabeth Islands: Preliminary Report of 1959–1960, edited by Müller, B. S., pp. 87–99, McGill University, 1961.
- Müller, F.: Englacial temperature measurements, in: Jacobsen-McGill Arctic Research Expedition 1959–1962: Preliminary report 1961–1962, Axel Heiberg Island Research Reports, pp. 81–89, McGill University, 1963a.
- 1005 Müller, F.: Accumulation studies, in: Jacobsen-McGill Arctic Research Expedition 1959–1962: Preliminary report 1961–1962, Axel Heiberg Island Research Reports, pp. 7–25, McGill University, 1963b.
- Müller, F.: On the thermal regime of a high-Arctic valley glacier, *Journal of Glaciology*, 16, 119–133, <https://doi.org/10.3189/S0022143000031476>, 1976.
- Naftz, D. L. and Smith, M. E.: Ice thickness, ablation, and other glaciological measurements on Upper Fremont Glacier, Wyoming, *Physical Geography*, 14, 404–414, <https://doi.org/10.1080/02723646.1993.10642488>, 1993.

- 1010 Nicholls, K. W. and Paren, J. G.: Extending the Antarctic meteorological record using ice-sheet temperature profiles, *Journal of Climate*, 6, 141–150, [https://doi.org/10.1175/1520-0442\(1993\)006<0141:ETAMRU>2.0.CO;2](https://doi.org/10.1175/1520-0442(1993)006<0141:ETAMRU>2.0.CO;2), 1993.
- Oeschger, H., Schotterer, U., Stauffer, B., Haeberli, W., and Röhlisberger, H.: First results from alpine core drilling projects, *Zeitschrift für Gletscherkunde und Glazialgeologie*, 13, 193–208, 1977.
- Olivier, S., Schwikowski, M., Brütsch, S., Eyrikh, S., Gäggeler, H. W., Lüthi, M., Papina, T., Saurer, M., Schotterer, U.,
1015 Tobler, L., and Vogel, E.: Glaciochemical investigation of an ice core from Belukha Glacier, Siberian Altai, *Geophysical Research Letters*, 30, <https://doi.org/10.1029/2003GL018290>, 2003.
- Orvig, S. and Mason, R. W.: Ice temperatures and heat flux, McCall Glacier, Alaska, in: General Assembly of Berkeley: Commission of Snow and Ice, vol. 61 of Publication, pp. 181–188, International Association of Scientific Hydrology, <https://iahs.info/uploads/dms/061021.pdf>, 1963.
- 1020 Palosuo, E.: A study of snow and ice temperatures on Vestfonna, Svalbard, 1956, 1957 and 1958, *Geografiska Annaler: Series A, Physical Geography*, 69, 431–437, <https://doi.org/10.2307/521356>, 1987.
- Palosuo, E. and Schytt, V.: Till Nordaustlandet med den svenska glaciologiska expeditionen [To Nordaustlandet with the Swedish glaciological expedition], *Eripainos Terrasta*, 72, 1–19, 1960.
- Paterson, W. S. B.: A temperature profile through the Meighen Ice Cap, in: General Assembly of Bern: Commission of Snow and Ice, vol. 79 of Publication, pp. 440–449, International Association of Scientific Hydrology, <https://iahs.info/uploads/dms/079041.pdf>, 1968.
1025
- Paterson, W. S. B.: Temperature measurements in Athabasca Glacier, Alberta, Canada, *Journal of Glaciology*, 10, 339–349, <https://doi.org/10.3189/S0022143000022036>, 1971.
- Paterson, W. S. B.: Temperature distribution in the upper layers of the ablation area of Athabasca Glacier, Alberta, Canada,
1030 *Journal of Glaciology*, 11, 31–41, <https://doi.org/10.3189/S0022143000022462>, 1972.
- Paterson, W. S. B. and Clarke, G. K. C.: Comparison of theoretical and observed temperature profiles in Devon Island Ice Cap, Canada, *Geophysical Journal International*, 55, 615–632, <https://doi.org/10.1111/j.1365-246X.1978.tb05931.x>, 1978.
- Pettersson, R., Jansson, P., and Holmlund, P.: Cold surface layer thinning on Storglaciären, Sweden, observed by repeated ground penetrating radar surveys, *Journal of Geophysical Research: Earth Surface*, 108,
1035 <https://doi.org/10.1029/2003JF000024>, 2003.
- Pilø, L., Reitmaier, T., Fischer, A., Barrett, J. H., and Nesje, A.: Ötzi, 30 years on: A reappraisal of the depositional and post-depositional history of the find, *The Holocene*, 33, 112–125, <https://doi.org/10.1177/09596836221126133>, 2023.
- Pollock, R.: CSV Dialect, <https://specs.frictionlessdata.io/csv-dialect/>, 2021.
- Prinz, R., Abermann, J., and Steiner, J.: Personal communication, 2022.
- 1040 Rabus, B. T. and Echelmeyer, K. A.: Increase of 10 m ice temperature: climate warming or glacier thinning?, *Journal of Glaciology*, 48, 279–286, <https://doi.org/10.3189/172756502781831430>, 2002.
- Reeh, N.: Summary of activities and preliminary results, in: Report on activities and results 1993–1995 for Hans Tausen Ice Cap Project - Glacier and Climate Change Research, North Greenland, edited by Reeh, N., NMRs (Nordisk Minister Råd) miljøforskningsprogram - klimaforskning, pp. 7–13, Dansk Polar Center, 1995.
- 1045 Reeh, N., Olesen, O. B., Thomsen, H. H., Starzer, W., and Bøggild, C. E.: Mass balance parameterisation for the Hans Tausen Iskappe, Peary Land, North Greenland, *Meddelelser om Grønland. Geoscience*, 39, <https://doi.org/10.7146/moggeosci.v39i.140219>, 2001.

- 1050 Rundle, A. S.: Glaciological investigations on Sukkertoppen Ice Cap, Southwest Greenland, summer 1964, Tech. rep., The Ohio State University Research Foundation, <http://hdl.handle.net/1811/51380>, 1965.
- Ryser, C., Lüthi, M., Blindow, N., Suckro, S., Funk, M., and Bauder, A.: Cold ice in the ablation zone: Its relation to glacier hydrology and ice water content, *Journal of Geophysical Research: Earth Surface*, 118, 693–705, <https://doi.org/10.1029/2012JF002526>, 2013.
- 1055 Schwerzmann, A., Funk, M., Blatter, H., Lüthi, M., Schwikowski, M., and Palmer, A.: A method to reconstruct past accumulation rates in Alpine firn regions: A study on Fiescherhorn, Swiss Alps, *Journal of Geophysical Research: Earth Surface*, 111, <https://doi.org/10.1029/2005JF000283>, 2006a.
- Schwerzmann, A. A., Blatter, H., and Funk, M.: Borehole analysis and flow modeling of firn-covered cold glaciers, Ph.D. thesis, Eidgenössische Technische Hochschule (ETH) Zürich, 2006b.
- 1060 Schwikowski, M., Schläppi, M., Santibañez, P., Rivera, A., and Casassa, G.: Net accumulation rates derived from ice core stable isotope records of Pío XI Glacier, Southern Patagonia Icefield, *The Cryosphere*, 7, 1635–1644, <https://doi.org/10.5194/tc-7-1635-2013>, 2013.
- Schytt, V.: Scientific results of the Swedish glaciological expedition to Nordaustlandet, Spitsbergen, 1957 and 1958, *Geografiska Annaler*, 46, 243–281, <https://doi.org/10.1080/20014422.1964.11881042>, 1964.
- 1065 Schytt, V.: Notes on glaciological activities in Kebnekaise, Sweden during 1965, *Geografiska Annaler: Series A, Physical Geography*, 48, 43–50, <https://doi.org/10.1080/04353676.1966.11879728>, 1966.
- Schytt, V.: Notes on glaciological activities in Kebnekaise, Sweden during 1966 and 1967, *Geografiska Annaler: Series A, Physical Geography*, 50, 111–120, <https://doi.org/10.1080/04353676.1968.11879777>, 1968.
- Shao, W. and Liu, Z.: Preliminary studies on the temperature in the surface layer of Guozha Glacier and Chongce Ice Cap in 1070 the West Kunlun Mountains, China, *Bulletin of Glacier Research*, 8, 87–91, 1990.
- Sharp, R. P.: Thermal regimen of firn on Upper Seward Glacier, Yukon Territory, Canada, *Journal of Glaciology*, 1, 476–487, <https://doi.org/10.3189/S0022143000026460>, 1951.
- Shiraiwa, T., Muravyev, Y. D., and Yamaguchi, S.: Stratigraphic features of firn as proxy climate signals at 1075 the summit ice cap of Ushkovsky Volcano, Kamchatka, Russia, *Arctic and Alpine Research*, 29, 414–421, <https://doi.org/10.1080/00040851.1997.12003262>, 1997.
- Shiraiwa, T., Murav'yev, Y. D., Kameda, T., Nishio, F., Toyama, Y., Takahashi, A., Ovsyannikov, A. A., Salamatin, A. N., and Yamagata, K.: Characteristics of a crater glacier at Ushkovsky volcano, Kamchatka, Russia, as revealed by the physical properties of ice cores and borehole thermometry, *Journal of Glaciology*, 47, 423–432, <https://doi.org/10.3189/172756501781832061>, 2001.
- 1080 Signer, N.: Analysis of ice temperatures of four selected very small glaciers in the Swiss Alps by means of modelling and ground penetrating radar, Ph.D. thesis, Universität Zürich, <https://lean-gate.geo.uzh.ch/typo3conf/ext/qfq/Classes/Api/download.php/mastersThesis/262>, 2014.
- Sobota, I.: The near-surface ice thermal structure of the Waldemarbreen, Svalbard, *Polish Polar Research*, 30, 317–338, 2009.
- Sobota, I.: Snow accumulation, melt, mass loss, and the near-surface ice temperature structure of Irenebreen, Svalbard, *Polar Science*, 5, 327–336, <https://doi.org/10.1016/j.polar.2011.06.003>, 2011.
- 1085

- Steffensen, J. P., Siggaard-Andersen, M.-L., Stampe, M., and Clausen, H. B.: Microparticles, soil derived chemical components and sea salt in the Hans Tausen Ice Cap ice core from Peary Land, North Greenland, *Meddelelser om Grønland*. Geoscience, 39, <https://doi.org/10.7146/moggeosci.v39i.140228>, 2001.
- Sugiyama, S.: Personal communication, 2022.
- 1090 Sugiyama, S., Navarro, F. J., Sawagaki, T., Minowa, M., Segawa, T., Onuma, Y., Otero, J., and Vasilenko, E. V.: Subglacial water pressure and ice-speed variations at Johnsons Glacier, Livingston Island, Antarctic Peninsula, *Journal of Glaciology*, 65, 689–699, <https://doi.org/10.1017/jog.2019.45>, 2019.
- Sun, H., Wang, N., and Hou, S.: Twentieth century warming reflected by the Malan Glacier borehole temperatures, northern Tibetan Plateau, Arctic, Antarctic, and Alpine Research, 53, 227–236, <https://doi.org/10.1080/15230430.2021.1974667>, 1095 2021.
- Suter, S., Laternser, M., Haeberli, W., Frauenfelder, R., and Hoelzle, M.: Cold firn and ice of high-altitude glaciers in the Alps: measurements and distribution modelling, *Journal of Glaciology*, 47, 85–96, <https://doi.org/10.3189/172756501781832566>, 2001.
- Suter, S., Minor, H.-E., and Ohmura, A.: Cold firn and ice in the Monte Rosa and Mont Blanc areas: spatial occurrence, 1100 surface energy balance and climatic evidence, Ph.D. thesis, Eidgenössische Technische Hochschule (ETH) Zürich, 2002.
- Takeuchi, N., Takahashi, A., Uetake, J., Yamazaki, T., Aizen, V. B., Joswiak, D., Surazakov, A., and Nikitin, S.: A report on ice core drilling on the western plateau of Mt. Belukha in the Russian Altai Mountains in 2003, *Polar Meteorology and Glaciology*, 18, 121–133, <https://doi.org/10.15094/00002977>, 2004.
- Takeuchi, N., Nakawo, M., Narita, H., and Han, J.: Miaoergou Glaciers in the Kalik Mountains, western China: Report of a 1105 reconnaissance for future ice core drilling and biological study, *Bulletin of Glaciological Research*, 26, 33–40, 2008.
- Takeuchi, N., Fujita, K., Aizen, V. B., Narama, C., Yokoyama, Y., Okamoto, S., Naoki, K., and Kubota, J.: The disappearance of glaciers in the Tien Shan Mountains in Central Asia at the end of Pleistocene, *Quaternary Science Reviews*, 103, 26–33, <https://doi.org/10.1016/j.quascirev.2014.09.006>, 2014.
- Talalay, P. G.: Mechanical Ice Drilling Technology, Springer, Singapore, <https://doi.org/10.1007/978-981-10-0560-2>, 2016.
- 1110 Talalay, P. G.: Thermal Ice Drilling Technology, Springer Geophysics, Springer, Singapore, <https://doi.org/10.1007/978-981-13-8848-4>, 2020.
- Thompson, L. G.: Glaciological investigations of the tropical Quelccaya Ice Cap, Peru, *Journal of Glaciology*, 25, 69–84, <https://doi.org/10.3189/S0022143000010297>, 1980.
- Thompson, L. G.: Curriculum vitae, https://research.bpcrc.osu.edu/Icecore/vitae/lgt_cv_old_format_2015.pdf, 2015.
- 1115 Thompson, L. G.: Personal communication, 2022.
- Thompson, L. G., Mosley-Thompson, E., Davis, M. E., Bolzan, J. F., Dai, J., Klein, L., Gundestrup, N., Yao, T., Wu, X., and Xie, Z.: Glacial stage ice-core records from the subtropical Dunde Ice Cap, China, *Annals of Glaciology*, 14, 288–297, <https://doi.org/10.3189/S0260305500008776>, 1990.
- Thompson, L. G., Mosley-Thompson, E., Davis, M., Lin, P. N., Yao, T., Dyurgerov, M., and Dai, J.: "Recent warming": 1120 ice core evidence from tropical ice cores with emphasis on Central Asia, *Global and Planetary Change*, 7, 145–156, [https://doi.org/10.1016/0921-8181\(93\)90046-Q](https://doi.org/10.1016/0921-8181(93)90046-Q), 1993.

- Thompson, L. G., Mosley-Thompson, E., Davis, M. E., Lin, P. N., Dai, J., Bolzan, J. F., and Yao, T.: A 1000 year climate ice-core record from the Guliya ice cap, China: its relationship to global climate variability, *Annals of Glaciology*, 21, 175–181, <https://doi.org/10.3189/S0260305500015780>, 1995a.
- 1125 Thompson, L. G., Mosley-Thompson, E., Davis, M. E., Lin, P. N., Henderson, K. A., Cole-Dai, J., Bolzan, J. F., and Liu, K.-b.: Late glacial stage and Holocene tropical ice core records from Huascarán, Peru, *Science*, 269, 46–50, <https://doi.org/10.1126/science.269.5220.46>, 1995b.
- Thompson, L. G., Mikhalenko, V., Mosley-Thompson, E., Durgerov, M., Lin, P. N., Moskalevsky, M., Davis, M. E., Arkhipov, S., and Dai, J.: Ice core records of recent climatic variability: Grigoriev and It-Tish ice caps in Central Tien Shan, Central 1130 Asia, *Материалы Гляциологических Исследований [Data of Glaciological Studies]*, 81, 100–109, 1997.
- Thompson, L. G., Davis, M. E., Mosley-Thompson, E., Sowers, T. A., Henderson, K. A., Zagorodnov, V. S., Lin, P.-N., Mikhalenko, V. N., Campen, R. K., Bolzan, J. F., Cole-Dai, J., and Francou, B.: A 25,000-year tropical climate history from Bolivian ice cores, *Science*, 282, 1858–1864, <https://doi.org/10.1126/science.282.5395.1858>, 1998.
- Thompson, L. G., Yao, T., Mosley-Thompson, E., Davis, M. E., Henderson, K. A., and Lin, P.-N.: A high- 1135 resolution millennial record of the South Asian monsoon from Himalayan ice cores, *Science*, 289, 1916–1919, <https://doi.org/10.1126/science.289.5486.1916>, 2000.
- Thompson, L. G., Mosley-Thompson, E., Davis, M. E., Henderson, K. A., Brecher, H. H., Zagorodnov, V. S., Mashiotta, T. A., Lin, P.-N., Mikhalenko, V. N., Hardy, D. R., and Beer, J.: Kilimanjaro ice core records: Evidence of Holocene climate change in tropical Africa, *Science*, 298, 589–593, <https://doi.org/10.1126/science.1073198>, 2002.
- 1140 Thompson, L. G., Mosley-Thompson, E. S., Zagorodnov, V., Davis, M. E., Mashiotta, T. A., and Lin, P.: 1500 years of annual climate and environmental variability as recorded in Bona-Churchill (Alaska) ice cores, <https://ui.adsabs.harvard.edu/abs/2004AGUFMPP23C..05T>, 2004.
- Thompson, L. G., Yao, T., Davis, M. E., Mosley-Thompson, E., Mashiotta, T. A., Lin, P.-N., Mikhalenko, V. N., and Zagorodnov, V. S.: Holocene climate variability archived in the Puruogangri Ice Cap on the central Tibetan Plateau, 1145 *Annals of Glaciology*, 43, 61–69, <https://doi.org/10.3189/172756406781812357>, 2006.
- Thompson, L. G., Yao, T., Davis, M. E., Mosley-Thompson, E., Wu, G., Porter, S. E., Xu, B., Lin, P.-N., Wang, N., Beaudon, E., Duan, K., Sierra-Hernández, M. R., and Kenny, D. V.: Ice core records of climate variability on the Third Pole with emphasis on the Guliya ice cap, western Kunlun Mountains, *Quaternary Science Reviews*, 188, 1–14, <https://doi.org/10.1016/j.quascirev.2018.03.003>, 2018.
- 1150 Thompson, L. G., Mosley-Thompson, E., Schoessow, F., Davis, M. E., Sierra-Hernández, M. R., and Beaudon, E.: The challenges, successes, and preliminary status report on the 2019 recovery of ice cores from Nevado Huascarán, Earth's highest tropical mountain, *Revista de Glaciares y Ecosistemas de Montaña*, 8, 31–42, 2023.
- Thomsen, H. H., Reeh, N., Olesen, O. B., and Jonsson, P.: Glacier and climate research on Hans Tausen Iskappe, North Greenland – 1995 glacier basin activities and preliminary results, *Bulletin Grønlands Geologiske Undersøgelse*, 172, 78–84, 1155 <https://doi.org/10.34194/bullggu.v172.6749>, 1996.
- Trabant, D., Harrison, W. D., and Benson, C.: Thermal regime of McCall Glacier, Brooks Range, Northern Alaska, in: *Climate of the Arctic: Twenty-Fourth Alaska Science Conference, August 15–17, 1973: Fairbanks*, edited by Weller, G. and Bowling, S. A., pp. 347–349, University of Alaska > Geophysical Institute, 1975.

- Troilo, F., Gottardelli, S., Giordan, D., Dematteis, N., Godio, A., and Vincent, C.: Geophysical and geomorphic recent surveys at Whymper hanging Glacier (Aosta Valley - Italy), pp. EGU21-5178, <https://doi.org/10.5194/egusphere-egu21-5178>, 2021.
- 1160 Tsushima, A., Miyahara, M., Yamasaki, T., Esashi, N., Sato, Y., Kayastha, R. B., Lama Sherpa, A. J. B., Sano, M., and Fujita, K.: Ice core drilling on a high-elevation accumulation zone of Trambau Glacier in the Nepal Himalaya, *Annals of Glaciology*, 62, 353–359, <https://doi.org/10.1017/aog.2021.15>, 2021.
- 1165 Tsutaki, S., Sugiyama, S., Sakakibara, D., Aoki, T., and Niwano, M.: Surface mass balance, ice velocity and near-surface ice temperature on Qaanaaq Ice Cap, northwestern Greenland, from 2012 to 2016, *Annals of Glaciology*, 58, 181–192, <https://doi.org/10.1017/aog.2017.7>, 2017.
- 1170 Uchida, T., Kamiyama, K., Fujii, Y., Takahashi, A., Suzuki, T., Yoshimura, Y., Igarashi, M., and Watanabe, O.: Ice core analyses and borehole temperature measurements at the drilling site on Asgardfonna, Svalbard, in 1993, *Memoirs of National Institute of Polar Research: Special issue*, 51, 377–386, 1996.
- Vallot, J.: Valeur et variation de la température profonde du glacier, au mont Blanc [Value and variation of the deep temperature of the glacier, at Mont Blanc], in: *Comptes rendus hebdomadaires des séances de l'Académie des Sciences* [Weekly reports of the sessions of the Academy of Sciences], vol. 156, pp. 1575–1578, Imprimerie Gauthier-Villars, <https://gallica.bnf.fr/ark:/12148/bpt6k3109m/f1577.item>, 1913.
- 1175 van de Wal, R. S. W., Mulvaney, R., Isaksson, E., Moore, J. C., Pinglot, J. F., Pohjola, V. A., and Thomassen, M. P. A.: Reconstruction of the historical temperature trend from measurements in a medium-length borehole on the Lomonosovfonna plateau, Svalbard, *Annals of Glaciology*, 35, 371–378, <https://doi.org/10.3189/172756402781816979>, 2002.
- 1180 Van Tricht, L., Huybrechts, P., Van Breedam, J., Fürst, J. J., Rybak, O., Satylkanov, R., Ermenbaiev, B., Popovnin, V., Neyns, R., Paice, C. M., and Malz, P.: Measuring and inferring the ice thickness distribution of four glaciers in the Tien Shan, Kyrgyzstan, *Journal of Glaciology*, 67, 269–286, <https://doi.org/10.1017/jog.2020.104>, 2021.
- Vance, T. R., Roberts, J. L., Moy, A. D., Curran, M. A. J., Tozer, C. R., Gallant, A. J. E., Abram, N. J., van Ommen, T. D., Young, D. A., Grima, C., Blankenship, D. D., and Siegert, M. J.: Optimal site selection for a high-resolution ice core record in East Antarctica, *Climate of the Past*, 12, 595–610, <https://doi.org/10.5194/cp-12-595-2016>, 2016.
- 1185 Vandecrux, B., Charles Amory, Andreas P. Ahlstrøm, Pete D. Akers, Mary Albert, and others: The SUMUp collaborative database: Surface mass balance, subsurface temperature and density measurements from the Greenland and Antarctic ice sheets (1912 - 2023), <https://doi.org/10.18739/A2M61BR5M>, 2023.
- Vilessov, E. N.: Temperature of ice in the lower parts of the Tuyuksu glaciers, in: General Assembly of Helsinki. 25-7 – 6-8 1960, Publication, pp. 313–324, International Association of Scientific Hydrology, <https://iahs.info/uploads/dms/1793.313-324-54-Vilessov-opt.pdf>, 1961.
- 1190 Vimeux, F., de Angelis, M., Ginot, P., Magand, O., Casassa, G., Pouyaud, B., Falourd, S., and Johnsen, S.: A promising location in Patagonia for paleoclimate and paleoenvironmental reconstructions revealed by a shallow firn core from Monte San Valentín (Northern Patagonia Icefield, Chile), *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2007JD009502>, 2008.

- Vimeux, F., Ginot, P., Schwikowski, M., Vuille, M., Hoffmann, G., Thompson, L. G., and Schotterer, U.: Climate variability during the last 1000 years inferred from Andean ice cores: A review of methodology and recent results, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 281, 229–241, <https://doi.org/10.1016/j.palaeo.2008.03.054>, 2009.
- Vincent, C., Le Meur, E., Six, D., Possenti, P., Lefebvre, E., and Funk, M.: Climate warming revealed by englacial temperatures at Col du Dôme (4250 m, Mont Blanc area), *Geophysical Research Letters*, 34, <https://doi.org/10.1029/2007GL029933>, 2007.
- Vincent, C., Descloitres, M., Garambois, S., Legchenko, A., Guyard, H., and Gilbert, A.: Detection of a sub-glacial lake in Glacier de Tête Rousse (Mont Blanc area, France), *Journal of Glaciology*, 58, 866–878, <https://doi.org/10.3189/2012JoG11J179>, 2012.
- Vincent, C., Gilbert, A., Jourdain, B., Piard, L., Ginot, P., Mikhalenko, V., Possenti, P., Le Meur, E., Laarman, O., and Six, D.: Strong changes in englacial temperatures despite insignificant changes in ice thickness at Dôme du Goûter Glacier (Mont Blanc area), *The Cryosphere*, 14, 925–934, <https://doi.org/10.5194/tc-14-925-2020>, 2020.
- Vinther, B. M., Clausen, H. B., Fisher, D. A., Koerner, R. M., Johnsen, S. J., Andersen, K. K., Dahl-Jensen, D., Rasmussen, S. O., Steffensen, J. P., and Svensson, A. M.: Synchronizing ice cores from the Renland and Agassiz ice caps to the Greenland ice core chronology, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2007JD009143>, 2008.
- Walsh, P. and Pollock, R.: Frictionless Table Schema, <https://specs.frictionlessdata.io/table-schema/>, 2021.
- Walsh, P., Pollock, R., and Keegan, M.: Frictionless Tabular Data Package, <https://specs.frictionlessdata.io/tabular-data-package/>, 2017.
- Wang, N. and Pu, J.: Strong influence of geothermal heat on the physical properties of glacier ice in the Tibetan Plateau, *Journal of Glaciology*, 51, 177–178, <https://doi.org/10.3189/S0022143000215207>, 2005.
- Wang, P., Li, Z., Li, H., Wang, W., Zhou, P., and Wang, L.: Characteristics of a partially debris-covered glacier and its response to atmospheric warming in Mt. Tomor, Tien Shan, China, *Global and Planetary Change*, 159, 11–24, <https://doi.org/10.1016/j.gloplacha.2017.10.006>, 2017.
- Wang, Y., Zhang, T., Ren, J., Qin, X., Liu, Y., Sun, W., Chen, J., Ding, M., Du, W., and Qin, D.: An investigation of the thermomechanical features of Laohugou Glacier No. 12 on Qilian Shan, western China, using a two-dimensional first-order flow-band ice flow model, *The Cryosphere*, 12, 851–866, <https://doi.org/10.5194/tc-12-851-2018>, 2018.
- Watanabe, O., Takenaka, S., Iida, H., Kamiyama, K., Thapa, K. B., and Mulmi, D. D.: First results from Himalayan Glacier Boring Project in 1981-1982: Part I. Stratigraphic analyses of full-depth cores from Yala Glacier, Langtang Himal, Nepal, *Bulletin of Glacier Research*, 2, 7–23, 1984.
- Watanabe, O., Motoyama, H., Igarashi, M., Kamiyama, K., Matoba, S., Goto-Azuma, K., Narita, H., and Kameda, T.: Studies on climatic and environmental changes during the last few hundred years using ice cores from various sites in Nordaustlandet, Svalbard (scientific paper), *Memoirs of National Institute of Polar Research: Special Issue*, 54, 227–242, 2001.
- Watts, L. G., Dowdeswell, J. A., and Murray, T.: The dynamics of Austfonna, Svalbard: two dimensional modelling of ice motion over a deformable substrate, *Материалы Гляциологических Исследований [Data of Glaciological Studies]*, 83, 10–21, 1997.

Weidick, A.: Gletschere i Sydgrønland: Historie, natur, omgivelser [Glaciers in South Greenland: history, nature, surroundings], no. 2 in Geologi i Grønland [Geology in Greenland], Grønlands Geologiske Undersøgelse [Geological Survey of Greenland], 1988.

- 1235 Weller, G., Nolan, M., Wendler, G., Benson, C., Echelmeyer, K., and Untersteiner, N.: Fifty years of McCall Glacier research: From the International Geophysical Year 1957–58 to the International Polar Year 2007–08, ARCTIC, 60, 101–110, <https://doi.org/10.14430/arctic280>, 2007.
- Wild, J., Nicolet, H., and Agassiz, L.: Carte du glacier inférieur de l'Aar, 1842.
- Wilson, N.: Characterization and interpretation of polythermal structure in two subarctic glaciers, Ph.D. thesis, Simon Fraser University, <https://summit.sfu.ca/item/12553>, 2012.
- 1240 Wilson, N. J., Flowers, G. E., and Mingo, L.: Comparison of thermal structure and evolution between neighboring subarctic glaciers, Journal of Geophysical Research: Earth Surface, 118, 1443–1459, <https://doi.org/10.1002/jgrf.20096>, 2013.
- Yang, J., Han, Y., Orsi, A. J., Kim, S., Han, H., Ryu, Y., Jang, Y., Moon, J., Choi, T., Hur, S. D., and Ahn, J.: Surface temperature in Twentieth Century at the Styx Glacier, northern Victoria Land, Antarctica, from borehole thermometry, 1245 Geophysical Research Letters, 45, 9834–9842, <https://doi.org/10.1029/2018GL078770>, 2018.
- Yao, T., Duan, K., Xu, B., Wang, N., Pu, J., Kang, S., Qin, X., and Thompson, L. G.: Temperature and methane changes over the past 1000 years recorded in Dasuopu Glacier (central Himalaya) ice core, Annals of Glaciology, 35, 379–383, <https://doi.org/10.3189/172756402781816997>, 2002.
- Zagorodnov, V. and Arkhipov, S.: Studies of structure, composition and temperature regime of sheet glaciers of Svalbard and 1250 Severnaya Zemlya: methods and outcomes, Bulletin of Glacier Research, 8, 19–28, 1990.
- Zagorodnov, V., Thompson, L. G., Ginot, P., and Mikhaleko, V.: Intermediate-depth ice coring of high-altitude and polar glaciers with a lightweight drilling system, Journal of Glaciology, 51, 491–501, <https://doi.org/10.3189/172756505781829269>, 2005.
- Zagorodnov, V., Nagornov, O., and Thompson, L. G.: Influence of air temperature on a glacier's active-layer temperature, 1255 Annals of Glaciology, 43, 285–291, <https://doi.org/10.3189/172756406781812203>, 2006.
- Zagorodnov, V., Nagornov, O., Scambos, T. A., Muto, A., Mosley-Thompson, E., Pettit, E. C., and Tyuflin, S.: Borehole temperatures reveal details of 20th century warming at Bruce Plateau, Antarctic Peninsula, The Cryosphere, 6, 675–686, <https://doi.org/10.5194/tc-6-675-2012>, 2012.
- Zelle, R. M., Wiernik, B. M., Bennet, F. G., D'Arcus, B., and Maier, D.: Citation Language Style: CSL 1.0.2. Specification, 1260 <https://docs.citationstyles.org/en/stable/specification.html>, 2015.
- Zhang, T., Xiao, C., Colgan, W., Qin, X., Du, W., Sun, W., Liu, Y., and Ding, M.: Observed and modelled ice temperature and velocity along the main flowline of East Rongbuk Glacier, Qomolangma (Mount Everest), Himalaya, Journal of Glaciology, 59, 438–448, <https://doi.org/10.3189/2013JoG12J202>, 2013.
- Zhang, W., Han, J., Xie, Z., Wang, X., Lluberas, A., and Goto-Azuma, K.: A preliminary study of ice texture and fabric on 1265 an ice core to the bedrock extracted from Glacier No. 1 at the headwater of Ulumqi River, Tianshan, China, Bulletin of Glacier Research, 11, 9–15, 1993.
- Ødegård, R. S., Hamran, S.-E., Bø, P. H., Etzelmüller, B., Vatne, G., and Sollid, J. L.: Thermal regime of a valley glacier, Erikbreen, northern Spitsbergen, Polar Research, 11, 69–79, <https://doi.org/10.3402/polar.v11i2.6718>, 1992.

- Ødegård, R. S., Hagen, J. O., and Hamranw, S.-E.: Comparison of radio-echo sounding (30–1000 MHz) and high-resolution borehole-temperature measurements at Finsterwalderbreen, southern Spitsbergen, Svalbard, *Annals of Glaciology*, 24, 262–267, <https://doi.org/10.3189/S0260305500012271>, 1997.
- 1270 А. М. Саватюгин [A. M. Savatyugin] and B. С. Загороднов [V. S. Zagorodnov]: Гляциологические исследования на ледниковом куполе Академии Наук [Glaciological studies on the Akademiya Nauk Ice Dome], Материалы Гляциологических Исследований [Data of Glaciological Studies], 61, 228–228, 1988.
- 1275 А. Н. Диких [A. N. Dikikh]: температурном режиме ледников плоских вершин (на примере ледника Григорьева) [The temperature regime of flat-top glaciers (using Grigoriev Glacier as an example)], in: Работы Тянь-Шаньской физико-географической станции [Glaciological studies in the Tien Shan], no. 11 in Гляциологические исследования на Тянь-Шане [Works of the Tien Shan physical-geographical station], pp. 32–35, 1965.
- 1280 Алексей Викторович Цветков [Aleksey Viktorovich Tsvetkov]: Неофициальный перечень высокогорных перевалов [Unofficial list of high mountain passes], <http://map.g-utka.ru>, 2023.
- 1285 В. А. Морев [V. A. Morev], О. Л. Клементьев [O. L. Klementev], Л. Н. Маневский [L. N. Manevsky], Ю. В. Райковский [Yu. V. Raykovsky], А. И. Толстой [A. I. Tolstoy], and В. М. Яковлев [V. M. Yakovlev]: Гляцио-буровые работы на Леднике Вавилова в 1979—1985 гг. [Glacio-drilling work on the Vavilov Glacier in 1979–1985], in: Географические и гляциологические в полярных странах [Geographical and glaciological studies in polar countries], edited by Korotkevich, E. S. and Petrov, V. N., pp. 25–32, Гидрометеоиздат [Gidrometeoizdat], 1988.
- 1290 В. А. Морев [V. A. Morev] and В. А. Пухов [V. A. Pukhov]: Экспериментальные работы по бурению холодных покровных ледников термобуровыми снарядами ААНИИ [Experimental work on drilling cold cover glaciers using AARI thermal drilling equipment], in: Исследования ледникового покрова и перигляциала Северной Земли [Studies of the ice cover and periglacial of Severnaya Zemlya], no. 367 in Труды ААНИИ [Proceedings of AARI], pp. 64–68, Арктического и Антарктического Научно-Исследовательского Института [Arctic and Antarctic Research Institute], 1981.
- 1295 В. М. Котляков [V. M. Kotlyakov]: Глубинное строение ледников [Deep structure of glaciers], in: Гляциология Шпицбергена [Glaciology of Spitsbergen], pp. 132–144, Наука, http://geolibRARY.ru/sites/default/files/2020-10/Glyatsiologiya_Shspitsbergena1985.pdf#page=71, 1985.
- 1300 В. Н. Михаленко [V. N. Mikhalenko]: Особенности массообмена ледников плоских вершин Внутреннего Тянь-Шаня [Properties of the mass-exchange of plateau glaciers in the inner Tian Shan], Материалы Гляциологических Исследований [Data of Glaciological Studies], 65, 86–92, 1989.
- 1305 В. Н. Михаленко [V. N. Mikhalenko], С. С. Кутузов [S. S. Kutuzov], И. И. Лаврентьев [I. I. Lavrentiev], М. Г. Кунакович [M. G. Kunakhovich], and Л. Г. Томпсон [L. G. Thompson]: Исследования западного ледникового плато Эльбруса: результаты и перспективы [Studies of the western Elbrus glacial plateau: results and prospects], Материалы Гляциологических Исследований [Data of Glaciological Studies], 99, 185–190, 2005a.
- В. Н. Михаленко [V. N. Mikhalenko], С. С. Кутузов [S. S. Kutuzov], Ф. Ф. Файзрахманов [F. F. Faizrakhmanov], О. В. Нагорнов [O. V. Nagornov], Л. Г. Томпсон [L. G. Thompson], М. Г. Кунакович [M. G. Kunakhovich], С. М. Архипов [S. M. Arkhipov], А. Н. Диких [A. N. Dikikh], and Р. Усубалиев [R. Usubaliev]: Сокращение оледенения Тянь-Шаня в XIX – начале XXI вв.: результаты кернового бурения и измерения температуры в скважинах [Reduction of glaciation in the Tien Shan in the 19th – early 21st centuries: results of core drilling and temperature measurements in wells], Материалы Гляциологических Исследований [Data of Glaciological Studies], 98, 175–182, 2005b.

- В. Н. Михаленко [V. N. Mikhalenko], С. С. Кутузов [S. S. Kutuzov], И. И. Лаврентьев [I. I. Lavrentiev], П. А. Торопов [P. A. Toropov], Д. О. Владимирова [D. O. Vladimirova], А. А. Абрамов [A. A. Abramov], and В. В. Мацковский [V. V. Matskovsky]: Гляциоклиматические исследования Института географии РАН в кратере Восточной вершины Эльбруса в 2020 г. [Glacioclimatological investigations of the Institute of Geography, RAS, in the crater of Eastern Summit of Mt. Elbrus in 2020], *Лёд и Снег* [Ice and Snow], 61, 149–160, <https://doi.org/10.31857/S2076673421010078>, 2021.
- 1310
Б. Р. Барбаш [V. R. Barbash], Л. С. Говоруха [L. S. Govorukha], and И. А. Зотиков [I. A. Zotikov]: О температурном состоянии толщи купола Вавилова [On the temperature state of the thickness of the Vavilov Ice Cap], in: Исследования ледникового покрова и перигляциала Северной Земли [Studies of the ice cover and periglacial of Severnaya Zemlya], no. 1315 367 in Труды ААНИИ [Proceedings of AARI], pp. 54–57, Арктического и Антарктического Научно-Исследовательского Института [Arctic and Antarctic Research Institute], 1981.
- Б. С. Загороднов [V. S. Zagorodnov]: Исследование строения и температурного режима Шпицбергенских ледников с помощью термобурения [Study of the structure and temperature regime of Svalbard glaciers using thermal drilling], Материалы Гляциологических Исследований [Data of Glaciological Studies], 41, 196–202, 1981.
- 1320
Б. С. Загороднов [V. S. Zagorodnov]: Температурный режим Ледника Академии Наук на Северной Земле [Temperature regime of the Academy of Sciences Glacier on Severnaya Zemlya], Материалы Гляциологических Исследований [Data of Glaciological Studies], 65, 134–138, 1989.
- Б. С. Загороднов [V. S. Zagorodnov], С. А. Синькевич [S. A. Sinkevich], and С. М. Архипов [S. M. Arkhipov]: Гидротермический режим ледораздельной области Восточного Ледяного Поля, О.Северо-Восточная Земля [Hydrothermal regime of the ice-divide area of Austfonna, Nordaustlandet], Материалы Гляциологических Исследований [Data of Glaciological Studies], 68, 133–141, 1990.
- 1325
Б. С. Загороднов [V. S. Zagorodnov], С. М. Архипов [S. M. Arkhipov], А. Б. Бажев [A. B. Bazhev], Т. А. Востокова [T. A. Vostokova], П. А. Королев [P. A. Korolev], О. В. Рототаева [O. V. Rototaeva], С. А. Синькевич [S. A. Sinkevich], and И. Ф. Хмелевской [I. F. Khmelevskoy]: Строение, состав и гидротермический режим ледника Гарабаши на Эльбрусе [Structure, composition and hydrothermal regime of the Garabashi Glacier at Elbrus], Материалы Гляциологических Исследований [Data of Glaciological Studies], 73, 109–117, 1992.
- Б. С. Загороднов [V. S. Zagorodnov] and И. А. Зотиков [I. A. Zotikov]: Керновое бурение на Шпицбергене [Core drilling in Svalbard], Материалы Гляциологических Исследований [Data of Glaciological Studies], 40, 157–163, 1981.
- Г. А. Цыкинаи [G. A. Tsykina] and Е. Н. Вилесов [E. N. Vilesov]: О температурном режиме Ледника Туюксу Центральный [About the temperature regime of the Central Tuyuksu Glacier], in: Исследования ледников и ледниковых районов [Research on glaciers and glacial areas], edited by Г. А. Авсяук [G. A. Avsyuk], vol. 3, pp. 56–66, Академия наук СССР [Academy of Sciences of the USSR], http://geolibrary.ru/sites/default/files/2020-11/IsLLeLegnR_3.pdf#page=56, 1963.
- 1335
Е. В. Василенко [E. V. Vasilenko]: Строение Ледника Давыдова по данным радиозондирования и термобурения [Structure of the Davydov Glacier according to radio sounding and thermal drilling data], Материалы Гляциологических Исследований [Data of Glaciological Studies], 62, 208–214, 1988.
- Е. М. Зингер [E. M. Singer], В. С. Корякин [V. S. Koryakin], Ю. А. Лаврушин [Yu. A. Lavrushin], В. А. Маркин [V. A. Markin], В. И. Михалёв [V. I. Mikhalev], and Л. С. Троицкий [L. S. Troitsky]: Исследование ледников Шпицбергена Советской экспедицией летом 1965 года [Exploration of Spitsbergen glaciers by the Soviet expedition in the summer of 1965], Материалы Гляциологических Исследований [Data of Glaciological Studies], 12, 59–72, 1966.

- 1345 Е. М. Зингер [E. M. Singer] and В. И. Михалёв [V. I. Mikhalev]: Аккумуляция снега на ледниках Шпицбергена [Snow accumulation on Spitsbergen glaciers], Материалы Гляциологических Исследований [Data of Glaciological Studies], 13, 86–100, 1967.
- Е. Н. Вилесов [E. N. Vilesov]: Температура льда: Стационарные наблюдения [Ice temperature: Stationary observations], vol. 1 of Материалы Гляциологических Исследований: Тянь-Шань Заилийский Алатау [Data of Glaciological Studies: Tien Shan Trans-Ili Alatau], Академия наук СССР > Институт Географии [Academy of Sciences of the USSR > Institute of Geography], <http://geolibrary.ru/sites/default/files/2024-02/%D0%97%D0%B0%D0%B8%D0%BB%D0%B8%D0%B9%D1%81%D0%BA%D0%B8%D0%B9%20%D0%90%D0%BB%D0%B0%D1%82%D0%80%D1%83.%20%D0%A2%D0%B5%D0%BC%D0%BF%D0%B5%D1%80%D0%80%D1%82%D1%83%D1%80%D0%80%D0%20%D0%BB%D1%8C%D0%B4%D0%80.%20%D0%92%D1%8B%D0%BF.1.pdf>, 1962a.
- 1350 Е. Н. Вилесов [E. N. Vilesov]: Температура льда: Стационарные наблюдения [Ice temperature: Stationary observations], vol. 2 of Материалы Гляциологических Исследований: Тянь-Шань Заилийский Алатау [Data of Glaciological Studies: Tien Shan Trans-Ili Alatau], Академия наук СССР > Институт Географии [Academy of Sciences of the USSR > Institute of Geography], 1962b.
- Е. Н. Вилесов [E. N. Vilesov]: Температура льда: Стационарные наблюдения [Ice temperature: Stationary observations], 1355 vol. 3 of Материалы Гляциологических Исследований: Тянь-Шань Заилийский Алатау [Data of Glaciological Studies: Tien Shan Trans-Ili Alatau], Академия наук СССР > Институт Географии [Academy of Sciences of the USSR > Institute of Geography], 1962c.
- И. Ф. Хмелевской [I. F. Khmelevskoy]: Температура снега, фирна и льда: Стационарные наблюдения на станции Барьер Сомнений и маршрутные исследования [Temperature of snow, firn and ice: Stationary observations at the station Barier Somneiy and en route researches], vol. 2 of Материалы Гляциологических Исследований: Новая Земля [Data of Glaciological Studies: Novaya Zemlya], Академия наук СССР > Институт Географии [Academy of Sciences of the USSR > Institute of Geography], 1963.
- 1360 И. Ф. Хмелевской [I. F. Khmelevskoy]: Температура снега, фирна и льда: Стационарные наблюдения на станции Ледораздельная [Temperature of snow, firn and ice: Stationary observations at the station Ledorazdelnaya], vol. 1 of Материалы Гляциологических Исследований: Новая Земля [Data of Glaciological Studies: Novaya Zemlya], Академия наук СССР > Институт Географии [Academy of Sciences of the USSR > Institute of Geography], 1964.
- Л. М. Саватюгин [L. M. Savatyugin], С. М. Архипов [S. M. Arkhipov], Н. И. Васильев [N. I. Vasiliev], Р. Н. Вострецов [R. N. Vostretsov], Д. Фритцше [D. Fritzsche], and Х. Миллер [H. Miller]: Российско-германские гляциологические исследования на Северной Земле и прилегающих островах в 2000 г. [Russian-German glaciological studies on Severnaya Zemlya and adjacent islands in 2000], Материалы Гляциологических Исследований [Data of Glaciological Studies], 91, 150–162, 2001.
- М. Я. Плам [M. Ya. Plam]: Температура фирна и льда [Temperature of firn and ice], Материалы Гляциологических Исследований: Эльбрус [Data of Glaciological Studies: Elbrus], pp. 5–59, 1962.
- 1365 Н. Г. Разумейко [N. G. Razumeiko]: Температура снега и льда: Стационарные исследования на куполе Чурляниса [Snow and ice temperature: Stationary studies of the Churlyanis Cupola], vol. 1 of Материалы Гляциологических Исследований: Земля Франца-Иосифа [Data of Glaciological Studies: Franz Josef Land], Академия наук СССР > Институт Географии [Academy of Sciences of the USSR > Institute of Geography], 1960.

- Н. Г. Разумейко [N. G. Razumeiko]: Температура снега и льда: Стационарные наблюдения на леднике Седова. Маршрутное термозондирование льда на куполах Чурляниса и Джэксона и леднике Седова [Snow and ice temperature: Stationary observations on the Sedov Glacier. En route thermosoundings of ice of the Churlyanis and Jackson cupolas and on the Sedov Glacier], vol. 2 of Материалы Гляциологических Исследований: Земля Франца-Иосифа [Data of Glaciological Studies: Franz Josef Land], Академия наук СССР > Институт Географии [Academy of Sciences of the USSR > Institute of Geography], 1963.
- 1385 Н. И. Барков [N. I. Barkov], К. В. Блинов [K. V. Blinov], М. С. Бугорков [M. S. Bugorkov], and Д. Н. Дмитриев [D. N. Dmitriev]: Геофизические исследования в скважине глубиной 460 м на Леднике Вавилова (Северная Земля) [Geophysical research in a well 460 m deep on the Vavilov Glacier (Severnaya Zemlya)], in: Географические и гляциологические в полярных странах [Geographical and glaciological studies in polar countries], pp. 14–24, Гидрометеоиздат [Gidrometeoizdat], 1988.
- 1390 Р. А. Чернов [R. A. Chernov], Т. В. Васильева [T. V. Vasilyeva], and А. В. Кудиков [A. V. Kudikov]: Температурный режим поверхности слоя ледника Восточный Грёнфьорд (Западный Шпицберген) [Temperature regime of upper layer of the glacier East Grönfjordbreen (West Svalbard)], Лёд и Снег [Ice and Snow], 131, 38, <https://doi.org/10.15356/2076-6734-2015-3-38-46>, 2015.
- 1395 С. А. Никитин [S. A. Nikitin]: Температурный режим ледника Малый Актру в период аблации [Temperature regime of Maly Aktru Glacier during the ablation period], Гляциология Сибири: Сборник статей [Glaciology of Siberia: Collection of articles], 3, 81–84, 1986.
- 1400 С. М. Архипов [S. M. Arkhipov], В. Н. Михаленко [V. N. Mikhalenko], М. Г. Кунакович [M. G. Kunakhovich], А. Н. Диких [A. N. Dikikh], and О. В. Нагорнов [O. V. Nagornov]: Термический режим, условия льдообразования и аккумуляция на леднике Григорьева (Тянь Шань) в 1962–2001 гг [Thermal regime, conditions of ice formation and accumulation on Grigoriev Glacier (Tien Shan) in 1962–2001], Материалы Гляциологических Исследований [Data of Glaciological Studies], 96, 77–83, 2004.
- 1405 С. С. Кутузов [S. S. Kutuzov]: Изменение площади и объёма ледников хр. Терской Ала-Тоо во второй половине XX в. [Changes in glacier area and volume in Terskey Ala-Too Range in the second half of the 20th century], Лёд и Снег [Ice and Snow], 52, 5–14, <https://doi.org/10.15356/2076-6734-2012-1-5-14>, 2012.
- 1410 Т. В. Псарёва [T. V. Psareva]: Экспериментальная 150-метровая буровая скважина на Леднике Безенги [Experimental 150-meter borehole on the Bezengi Glacier], Материалы Гляциологических Исследований [Data of Glaciological Studies], 14, 93–97, 1968.
- 1415 Ю. Я. Мачерет [Y. Y. Macheret], В. С. Загороднов [V. S. Zagorodnov], Е. В. Василенко [E. V. Vasilenko], А. И. Громыко [A. I. Gromyko], and А. Б. Журавлев [A. B. Zhuravlev]: Исследование природы внутренних радиолокационных отражения на субполярном леднике [On the nature of internal radio echo returns from a subpolar glacier], Материалы Гляциологических Исследований [Data of Glaciological Studies], 54, 120–130, 1985.
- 仇家琪[Qiu Jiaqi] and 邓养鑫[Deng Yangxin]: 天山博格达峰地区的雪崩[Snow Avalanche in Bogda Region, Tian Shan], 冰川冻土[Journal of Glaciology and Geocryology], 5, 227–234, <https://doi.org/10.7522/j.issn.1000-0240.1983.0055>, 1983.
- 任贾文[Ren Jiawen]: 天山博格达峰扇状分流冰川的冰层温度[The ice temperature of Bogda Fan-Shaped Diffusion Glacier in Bogda Area, Tian Shan], 冰川冻土[Journal of Glaciology and Geocryology], 5, 83–89, <https://doi.org/10.7522/j.issn.1000-0240.1983.0040>, 1983.

- 任贾文[Ren Jiawen]: 南极长城站附近地区冰川的温度状况[Temperature regime of the glaciers in the neighbourhood of Great Wall Station, Antarctica], 极地研究[Chinese Journal of Polar Research], 2, 22–27, 1990.
- 任贾文[Ren Jiawen], 张金华[Zhang Jinhua], and 黄茂桓[Huang Maohuan]: 天山乌鲁木齐河源1号冰川温度研究[A study of ice temperature in No. 1 Glacier in the Urumqi River headwaters, Tianshan], 冰川冻土[Journal of Glaciology and Geocryology], 7, 141–152, <https://doi.org/10.7522/j.issn.1000-0240.1985.0019>, 1985.
- 任贾文[Ren Jiawen] and 黄茂桓[Huang Maohuan]: 冰川活动层温度状况的传热学分析——以祁连山羊龙河5号冰川为例[Heat transfer within glacial active layer —— Taking No. 5 Glacier, Yanglong River, Qilian Shan as an example], 冰川冻土[Journal of Glaciology and Geocryology], 3, 23–28, <https://doi.org/10.7522/j.issn.1000-0240.1981.0042>, 1981.
- 刘亚平[Liu Yaping], 侯书贵[Hou Shugui], 任贾文[Ren Jiawen], 王叶堂[Wang Yetang], and 耿志新[Geng Zhixin]: 东天山庙儿沟平顶冰川钻孔温度分布特征[Temperature distribution characteristics of boreholes in Miaoergou flat-top glacier in East Tianshan Mountains], 冰川冻土[Journal of Glaciology and Geocryology], 28, 668–671, <https://doi.org/10.7522/j.issn.1000-0240.2006.0097>, 2006.
- 刘时银[Liu Shiyin], 郭万钦[Guo Wanqin], and 许君利[Xu Junli]: 中国第二次冰川编目数据集[The second glacier inventory dataset of China], <https://doi.org/10.3972/glacier.001.2013.db>, 2012.
- 周韬[Zhou Tao]: 西昆仑山崇测冰帽温度分布研究[Study on temperature distribution of Chongce ice cap in West Kunlun Mountains], 科学通报[Chinese Science Bulletin], 35, 212–215, <https://doi.org/10.1360/csb1990-35-3-212>, 1990.
- 姚檀栋[Yao Tandong], 焦克勤[Jiao Keqin], 章新平[Zhang Xinping], 杨志红[Yang Zhihong], and Thompson, L. G.: 古里雅冰帽冰川学研究[Glaciologic studies on Guliya Ice Cap], 冰川冻土[Journal of Glaciology and Geocryology], 14, 233–241, <https://doi.org/10.7522/j.issn.1000-0240.1992.0037>, 1992.
- 孙维君[Sun Weijun], 闫明[Yan Ming], 艾松涛[Ai Songtao], 朱国才[Zhu Guocai], 王泽民[Wang Zemin], 刘雷保[Liu Leibao], 徐跃通[Xu Yuetong], and 任贾文[Ren Jiawen]: 北极新奥尔松地区Austre Lovénbreen冰川温度变化特征 [Ice temperature characteristics of Austre Lovénbreen in Ny-Ålesund, Arctic Region], 武汉大学学报(信息科学版) [Geomatics and Information Science of Wuhan University], 41, 79–85, <https://doi.org/10.13203/j.whugis20150302>, 2016.
- 张通[Zhang Tong]: Personal communication, 2022.
- 李忠勤[Li Zhongqin], 李慧林[Li Huilin], and 陈亚宁[Chen Yaning]: Mechanisms and simulation of accelerated shrinkage of continental glaciers: A case study of Urumqi Glacier No. 1 in eastern Tianshan, Central Asia, Journal of Earth Science, 22, 423–430, <https://doi.org/10.1007/s12583-011-0194-5>, 2011.
- 李真[Li Zhen], 姚檀栋[Yao Tandong], 田立德[Tian Lide], 徐柏青[Xu Baiqing], 邬光剑[Wu Guangjian], and 朱国才[Zhu Guocai]: 慕士塔格冰川海拔7000m处冰芯钻孔温度[Ice-core borehole temperature at 7000 m on the Muztagh Glacier], 冰川冻土[Journal of Glaciology and Geocryology], 26, 284–288, <https://doi.org/10.7522/j.issn.1000-0240.2004.0047>, 2004.
- 王文颖[Wang Wenying]: Existing glaciers and their variations in the northeastern part of the Qinghai-Xizang Plateau, in: Reports on the northeastern part of the Qinghai-Xizang (Tibet) Plateau by Sino-W. German Scientific Expedition, edited by Hövermann, J. and 王文颖[Wang Wenying], pp. 22–37, Science Press, 1987.
- 王立伦[Wang Lilun], 刘潮海[Liu Chaohai], 康兴成[Kang Xingcheng], and 尤根祥[You Genxiang]: 我国阿尔泰山现代冰川的基本特征——以哈拉斯冰川为例[Fundamental features of modern glaciers in the Altay Shan of China —— Taking Halasi Glacier as an example], 冰川冻土[Journal of Glaciology and Geocryology], 5, 27–38, <https://doi.org/10.7522/j.issn.1000-0240.1983.0061>, 1983.

苏珍[Su Zhen]: 喀喇昆仑山—昆仑山地区冰川的物理与化学性质[Physical and chemical properties of glaciers in the Karakoram-Kunlun Mountains region], in: 喀喇昆仑山-昆仑山地区冰川与环境[Glaciers and environment in the Karakoram-Kunlun Mountains region], 青藏高原喀喇昆仑山-昆仑山地区科学考察丛书[Scientific Expedition Series in the Karakoram-Kunlun Mountains Region of the Tibetan Plateau], pp. 58–82, 科学出版社[Science Press], <https://book.sciencereading.cn/shop/book/Booksimple/show.do?id=B828976A04419443DB3FE50FBA719BB38000>, 1998.

1460 蒲健辰[Pu Jianchen], 姚檀栋[Yao Tandong], 张寅生[Zhang Yinsheng], 濑古胜基[Seko Katsuki], and 藤田耕史[Fujita Koji]: 冬
克玛底冰川和煤矿冰川的物质平衡(1992/1993年) [Mass balance on the Dongkemadi and Meikuang glaciers in 1992/1993],
1465 冰川冻土[Journal of Glaciology and Geocryology], 17, 138–143, <https://doi.org/10.7522/j.issn.1000-0240.1995.0018>, 1995.

蒲健辰[Pu Jianchen], 姚檀栋[Yao Tandong], 王宁练[Wang Ninglian], 段克勤[Duan Keqin], 朱国才[Zhu Guocai], and
杨梅学[Yang Meixue]: 青藏高原普若岗日冰原80m深冰层温度变化分析[The distribution of 80-m ice temperature
in Puruogangri Ice Field on the Tibetan Plateau], 冰川冻土[Journal of Glaciology and Geocryology], 24, 282–286,
<https://doi.org/10.7522/j.issn.1000-0240.2002.0053>, 2002.

1470 邬光剑[Wu Guangjian], 姚檀栋[Yao Tandong], 徐柏青[Xu Baiqing], 李真[Li Zhen], and 保翰璋[Bao Hanzhang]: 慕士塔格冰芯
钻孔温度测量结果[Ice-core borehole temperature in the Muztagh Ata, East Pamirs], 冰川冻土[Journal of Glaciology and
Geocryology], 25, 676–679, <https://doi.org/10.7522/j.issn.1000-0240.2003.0119>, 2003.

韩建康[Han Jiankang], 温家洪[Wen Jiahong], 尚新春[Shang Xinchun], and 金会军[Jin Huijun]: 南极洲乔治王岛柯林斯冰帽的
温度分布[Temperature distribution in Collins Ice Cap, King George Island, Antarctica], 极地研究[Chinese Journal of Polar
1475 Research], 7, 62–69, 1995.