

One year of high frequency monitoring of groundwater physico-chemical parameters in the Weierbach Experimental Catchment, Luxembourg.

Karl Nicolaus van Zweel^{1,2}, Laurent Gourdol², Jean François Iffly², Loïc Léonard², François Barnich², Laurent Pfister², Erwin Zehe³, and Christophe Hissler²

¹VTT Technical Research Centre Ltd, P.O. Box 1000, FI-02044 VTT, Finland

²Luxembourg Institute of Science and Technology (LIST), Avenue des Hauts-Fourneaux 5, L-4362 Esch-sur-Alzette, Luxembourg

³Karlsruher Institut für Technologie (KIT), Institut für Wasser und Gewässerentwicklung, Lehrstuhl für Hydrologie, Otto-Amman-Platz 1, 76131 Karlsruhe

Correspondence: Christophe Hissler (christophe.hissler@list.lu)

Abstract. The critical zone (CZ) is the ~~Earth's skin~~ skin of the Earth, where rock, water, air, and life interact~~-,~~ playing a pivotal role in sustaining ecological processes and life-supporting resources. Understanding these interactions, especially in forested headwater catchments, is ~~key-for-managing~~ essential to manage water resources, ~~predicting~~ predict environmental responses, and ~~assessing~~ assess human impacts. ~~We~~ Here, we present a novel dataset from the Weierbach Experimental Catchment in Luxembourg, derived from a year-long high-frequency monitoring campaign ~~that~~-focused on groundwater ~~physico-chemical parameters.~~ Through meticulous physicochemical parameters.

Through rigorous data collection and ~~rigorous~~-quality control, parameters such as electrical conductivity, dissolved oxygen, oxidation-reduction potential, and pH were measured, ~~offering new~~ providing insights into the CZ's hydrological and biogeochemical dynamics. ~~The dataset highlights the intricate interplay~~ Although the one-year data set offers valuable observations, it represents an initial step toward understanding long-term patterns. The data highlight the interaction between redox reactions, pH, and ~~ion-exchange processes, as well as the influence of seasonal variability and flowline interactions on solute transport.~~ By providing a detailed view of the catchment's response to hydrological changes seasonal hydrological variability, although these interpretations are limited by the temporal scope of the study.

By offering a detailed snapshot of the response of the catchment to hydrological variations, this dataset ~~fills a significant gap~~ contributes to addressing key gaps in CZ research ~~, offering a valuable resource and serves as a foundation~~ for advancing our understanding of ~~hydro-biogeochemical catchment processes.~~ The datasets introduced in this contribution may be leveraged by researchers and practitioners aiming hydrobiogeochemical processes at the catchment scale. Despite the limited observation period, the data set provides insights that can be integrated with long-term monitoring efforts. Researchers and practitioners can use these data to refine models, inform land management ~~practices, and foster a more holistic understanding of catchment~~ biogeochemistry decisions, and improve understanding of the biogeochemistry of the catchment.

1 Introduction

The critical zone (CZ) ~~stands-as-is~~ is the interface where rock, water, air, and living organisms interact, playing a vital role in supporting ecological processes and providing essential resources (Amundson et al., 2007; Anderson et al., 2012; Brantley et al., 2007). The dynamics within this “skin of the Earth” ~~are-is~~ are fundamental for managing water resources, predicting environmental responses to disturbances, and assessing the impacts of human activities, especially against the backdrop of climate change. Within the CZ, the fields of hydrology and biogeochemistry interconnect deeply, offering insights into ecological sustainability and resource management (Wymore et al., 2022).

The subsurface properties, especially ~~the~~ variations in permeability with depth, exert a significant control on the vertical connectivity within the CZ, affecting ~~the~~ water flow and chemical interactions. For example, in systems where vertical permeability decreases sharply, most water travels through only the ~~top-upper~~ layers of the subsurface, limiting its interaction with deeper, reactive rock formations ~~—~~ (Xiao et al., 2021). This results in a pronounced chemical contrast between shallow and deep waters, where shallow water, often younger and flowing rapidly through high-permeability soil layers, carries fewer geogenic solutes compared to ~~the~~ older, deeper water that has interacted more extensively with bedrock (Welch and Allen, 2014; Jin et al., 2010; Brantley et al., 2013). These differences are highly relevant, ~~as~~ because they shape the solute export patterns observed at ~~the~~ catchment outlets, influencing the concentration vs. discharge (c-Q) relationships. Understanding these subsurface interactions is important ~~—as-because~~ they not only determine stream chemistry but also provide insights into ~~hydro-biogeochemical~~ ~~hydrobiogeochemical~~ processes, which are essential ~~for-assessing-to-assess~~ ~~managing-water-resources~~ ~~effectively~~ ~~effectively~~ manage water resources.

One of the most accessible methods ~~to-understand-of understanding~~ the interactions occurring within the CZ is ~~by-observing~~ ~~to observe the~~ catchment outlets. Streams, acting as integrators of all processes within the CZ, reveal the cumulative effects of these interactions on ~~stream-behaviour-and-the behaviour of the stream and the~~ hydrochemical composition, thus serving as a window into the overall dynamics of the CZ (Brantley et al., 2017; Li et al., 2017; Riebe et al., 2017). This approach underpins the importance of analysing concentration vs. discharge (c-Q) patterns, which have been a focal point of research, providing ~~insights-into-information on~~ the hydrological and biogeochemical functioning of ~~the~~ catchments (Brooks et al., 2015; Zhi et al., 2019; Stewart et al., 2022). This line of inquiry underscores the complex interplay of ~~hydro-biogeochemical~~ ~~hydrobiogeochemical~~ processes within the CZ, often masked by spatial heterogeneities. Despite advancements, field measurements and their interpretations remain challenging, with water physico-chemistry data reflecting convoluted signatures that obscure direct cause-and-effect relationships (Weiler and McDonnell, 2006). Recent consensus, emerging from diverse measurements and observations, points towards the determination of stream chemistry by the properties of source waters, predominantly from shallow subsurface and/or deeper regolith and fractured bedrock flow paths. The contrasting chemical properties of these source waters are pivotal in shaping ~~the~~ solute export patterns (Stewart et al., 2022).

Given the complexity of interactions within the CZ, direct data supporting the nuanced understanding of subsurface and stream physico-chemical interactions are rare. This research gap emphasises the need for detailed datasets that can capture the intricacies of CZ dynamics ~~Addressing this need, our study presents a novel dataset centred on the~~ (Stewart et al., 2022).

Our study complements existing Critical Zone research efforts across Europe and beyond. Various observatories, such as the TERENO Wüstebach Catchment in Germany, have advanced our understanding of soil moisture dynamics, land-atmosphere exchange, and hydrological processes through dense sensor networks (?). Similarly, the SoilTrEC network of observatories across Switzerland, Austria, Greece, and the Czech Republic focusses on soil transformation, biogeochemical cycling, and the impacts of land use (?). The Calhoun CZO in the USA provides insights into deep soil biogeochemical processes, particularly carbon cycling and hydrological connectivity (?).

Our dataset from the Weierbach Experimental Catchment (WEC) ~~in Luxembourg. Chosen for its~~ provides high-resolution temporal data on groundwater physicochemical parameters, with a particular focus on redox dynamics and pH fluctuations. Although many observatories investigate both surface and subsurface processes, high-frequency monitoring of subsurface redox dynamics zonation remains limited. Our study offers initial insight into vertical biogeochemical dynamics within the subsurface, contributing to a better understanding of redox zonation, subsurface hydrological pathways, and the interplay between hydrological variability and biogeochemical processes in temperate forested catchments.

To address this gap, we present a novel dataset from the WEC in Luxembourg. With extensive prior research and its status as a CZ observatory, the WEC ~~offers an unparalleled opportunity to delve into the~~ provides a valuable opportunity to examine subsurface hydrological and biogeochemical processes shaping the CZ (Hissler et al., 2021).

This paper introduces our methodology for data collection, ~~focusing on the significance of subsurface architecture knowledge focussing on the importance of understanding subsurface architecture~~ and the exploration of hydrological dynamics through the ~~dataset~~data set. By presenting this dataset, we aim to contribute to ~~the a~~ broader understanding of complex ~~hydro-biogeochemical hydrobiogeochemical~~ processes within the CZ at specific subsurface locations within a long-term experimental catchment. This effort not only ~~enhances our comprehension of water resource dynamics but also underseores~~ improves our understanding of the dynamics of water resources, but also highlights the importance of integrating hydrological and biogeochemical data for environmental monitoring and management.

2 The Weierbach Experimental Catchment (WEC)

The catchment is located in the northwestern region of Luxembourg (Lat: 49.8273; Long: 5.7956) and has an area of 45 ha. The WEC is a forested headwater catchment within the Alzette River basin (Pfister et al., 2017). The catchment is located in the Ardennes massif, which is distinguished by an elevated subhorizontal plateau and deep V-shaped valleys. It is subdivided ~~in~~into three landscape units (Figure 1). The plateau area is about 31 ha and has slopes between 0 ° and 5 °. The hillslope area is about 14 ha and has slopes between 5 ° and 44 ° (Martínez-Carreras et al., 2016). The riparian area represents about 0.4 ha. The catchment's stream network consists of three tributaries that converge into a single stream and is bordered by a highly dynamic riparian zone composed primarily of litter degradation material and colluvium from the hillslopes (Antonelli et al.,

2020a, b). The riparian area in the catchment is narrow, flat, and ~~well-defined~~ well defined, with some areas where the riparian zone widens.

The ~~WEC's forest structure~~ forest structure of the Weierbach Experimental Catchment (WEC) is shaped by past and current management practices. Oak trees are evenly distributed, while beech ~~trees'~~ tree density increases from ~~plateau to the plateau to~~ the footslope. The primary plateau ~~has~~ hosts a dense mixed forest ~~with~~ composed of 78% ~~European-beech-trees~~ *Fagus sylvatica* and 22% ~~oak hybrids~~ (Fabiani et al., 2022), where the ~~*Quercus petraea* × *Q. robur*~~ (Fabiani et al., 2022). The understory is dominated by ~~blueberries, and~~ *Vaccinium myrtillus*, while the riparian zone ~~has~~ features various short plants (Martínez-Carreras et al., 2015). The secondary plateau is primarily ~~Norway spruce and Douglas fir~~ composed of *Picea abies* and *Pseudotsuga menziesii*. A detailed analysis of forest structure, including tree density and ecohydrological interactions, can be found in Fabiani et al. (2022); ?.

The WEC experiences a semi-marine climate with ~~even-precipitation-distribution~~ a uniform distribution of precipitation throughout the year, averaging 953 mm annually (2006–2014; Pfister et al. (2017)). Streamflow varies with seasons, with lower base flow during July to September due to higher evapotranspiration losses (annual average PET of 593 mm for the period ~~of~~ 2006-2014, from Pfister et al. (2017, 2023)). In dry conditions, intermittent streamflow occurs from upstream to downstream, ~~characterized~~ characterised by sharp, short-lasting discharge peaks (Wrede et al., 2015). In wet conditions, a double peak behaviour is observed, with a rapid discharge peak shortly after rainfall, followed by a broad ~~,~~ and long-lasting second peak that outweighs the first peak in volume (Martínez-Carreras et al., 2016; Scaini et al., 2018). Figure 2 displays the ~~daily-total~~ total daily precipitation in the WEC and the ~~daily-mean-discharge-at~~ mean daily discharge in SW1 from late 2008 to 2022.

The WEC regolith is ~~composed~~ made up of two parts called ~~the~~ solum, where pedogenic processes ~~are dominant and where~~ dominate and the biota play an important role ~~and,~~ and the subsolum, where the original rock structure or ~~the~~ fabric of the bedrock is preserved (Juilleret et al., 2016). The structure of these two parts ~~evolve~~ evolved according to the topography. The WEC's solum between 0 and 70 cm consists of a stony silt loam soil derived from periglacial solifluction deposits that is overlain on a weathered Devonian slate substratum. From the surface, a thin, dark organic surface horizon (O to A horizon, 0-8 cm on average) is followed by a cambic horizon (B horizon, 8-40 cm on average) with a well-developed angular blocky soil structure. The texture of the silt loam is inherited from the loess deposit and was mixed with slate clasts via solifluction, with the quantity of clasts increasing with depth. The subsolum is composed by slates, phyllites and quartz of Devonian age (Moragues-Quiroga et al., 2017). During the last 10 years, extensive drilling and geophysics (Gourdol et al., 2021) campaigns ~~has~~ have been conducted. The ~~WEC's~~ deep subsurface exploration of the WEC yielded more detail than is currently known. The upper part of the catchment can be subdivided between a primary plateau where the solum developed over a deep yellow-white slate saprolite and a secondary plateau where the solum directly developed over the black fractured slate bedrock. These distinct landscape units are separated by the black line in Figure 1, which represents the weathering front. The spatial occurrence of these two distinct regolith structures was observed using combined core drilling descriptions and electrical resistivity survey (Gourdol et al., 2021). In the hillslope and riparian areas, the subsolum is composed ~~on~~ of fractured and / or fresh bedrock ~~underneath~~ below the solum. A mineralogical and geochemical ~~characterization~~ characterisation of the WEC's regolith was conducted to gain a deeper understanding of the primary plateau regolith characteristics and evolution. This study allowed

observing three distinct compartments that have different geochemical —and mineralogical characteristics as a function of contrasting evolutions. These respective processes are related to different atmospheric deposition events, weathering ages and seasonal water saturation dynamics (Moragues-Quiroga et al., 2017).

2.1 Data Description

125 2.1.1 Fifteen years of groundwater monitoring

The WEC has been thoroughly equipped to continuously monitor water fluxes and ~~physico-chemical~~physicochemical parameters within various CZ compartments. ~~Additionally~~In addition, these compartments are sampled every two weeks at different locations to ~~analyze~~analyse the $\delta^{18}\text{O}$ and $\delta^2\text{H}$ isotopic composition of water, as well as its chemical composition, including rainfall, throughfall, groundwater, and streamwater. Some of these datasets have already been published (Hissler et al., 130 2021). The WEC monitoring infrastructure is continuously expanded and now encompasses an extensive network of 26 monitoring boreholes with varying depths, ranging from 1.95 meters to as deep as 40 meters below the surface. All the boreholes are spatially distributed ~~all-over~~throughout the catchment (Figure 1) and have screens that cover the range ~~from of 1 meter~~metre below the surface to the bottom. For this study, 9 boreholes were selected along one hillslope catena, representing one unique vegetation type, but several hydrogeological ~~positioning~~positions. The chosen boreholes include locations on the primary plateau (GW1 and GW5), in the middle of the hillslope (GW7 and GW9), and at low hillslope positions (GW2, GW10-12 135 and GW3). Groundwater levels in these boreholes are automatically recorded every 15 minutes using OTT Orphimedes and CTD (OTT, Aix-en-Provence, France). The relevant data are shown in Table 1. The depth reference for the water sampling is the top of the pipe, ~~whereas~~while it is the surface of the soil for the screen of the pipe. The standpipe represents the difference between the top of the pipe and the surface of the soil.

140 2.1.2 One year of physico-chemical parameters monitoring

A weekly frequency monitoring campaign was designed with the objective of achieving higher resolution ~~concerning key~~potential-on the potential key processes expected to occur in the subsurface of the catchment. The duration of the campaign spanned from 02/03/2021 to 14/03/2022, during which sampling was conducted at weekly intervals. The parameters measured during each sampling event included electrical conductivity (EC), dissolved oxygen (DO), oxidation-reduction potential (ORP), 145 corrected for probe bias to Eh, measures the ~~water's redox potential~~redox potential of water using a platinum electrode, commonly expressed relative to the standard hydrogen electrode (SHE) for comparison, and pH, all of which were analysed using a ~~flow-through~~flow cell system (Figure 3). The four parameters were recorded using two WTW® Multi 3630 IDS data loggers connected to the specific probes: the WTW® TetraCon® 925 probe for EC, the WTW® FDO® 925 probe for DO, the WTW® SenTix® ORP-T 900 probe for ORP, and the WTW® SenTix® 940 probe for pH. The pH probe ~~underwent calibration~~ 150 was calibrated in the field at each sampling event using HACH Singlet™ Solution Packs for a two-point calibration process (pH1=4.01 and pH2=7.00). In addition to the four parameters, the study also recorded pumping time, water level before and after pumping, as well as the quantity of water extracted from the well.

155 ~~Prior to measure~~ Before measuring the parameters and sampling a ~~borehole, a five-minute period was allowed for the probes~~
~~well, the system was flushed with one litre of demineralised Milli-Q water to remove potential contaminants. The probes~~
~~were then allowed a 5-minute period~~ to record the EC ~~datum~~-reading, which ranged from 0.1 to 1.7 $\mu\text{S}/\text{cm}$, ~~after flushing the~~
~~system with one litre of demineralized milliQ-water~~. Subsequently, the borehole was pumped ~~;~~ and the four parameters were
continuously recorded at ~~one-minute increments~~ increments of one minute. A groundwater sample was collected when the EC
reading remained stable for several minutes, under the assumption that the borehole demonstrated homogeneity. Measurements
were conducted in-line to minimise atmospheric contact and temperature changes. After collecting the water sample, a 20-
160 minute equilibration period was implemented, ~~wherein~~ in which the flow-through system was sealed ~~;~~ and the probes were
allowed to equilibrate with the solution in the cells. The final readings for EC, DO, ORP, and pH were then obtained. The
physico-chemical database consists of three subsets:

- The first subset includes a ~~high-frequency~~ high-frequency continuous pumping test ~~at borehole in~~ GW5 (Figure 1 and
Table 1). These tests were ~~conducted~~ carried out twice: once during the winter and again during the summer, covering
165 both wet and dry seasons. During these tests, the four physical parameters (EC, DO, ORP, and pH) were measured at
10-second intervals, while water samples were taken every minute ~~for measuring to measure~~ hydrochemistry.
- The second subset comprises time series data ~~representing that represent~~ the four parameters obtained from weekly
sampling events. In this dataset, each sample site is evaluated weekly for four physicochemical parameters: pH, EC,
ORP, and DO. The alignment of the ~~time-series~~ time-series data for all physico-chemical parameters was based on the
170 EC values. Consequently, the peaks in the EC for each time trend, corresponding to their respective sample sites, may
not always align due to variations in the time required for the probes to reach equilibrium after flushing the flow-through
cell setup with the ~~demineralized~~ demineralised milli-Q water. To standardise the data across various sampling sites,
a centring process was ~~employed~~ used. This process ~~entailed the identification of the involved~~ identifying the highest
EC value within each time series. Upon ~~locating~~ finding this maximal EC value, the time scale of the dataset was
175 subsequently adjusted. Specifically, the time series was ~~recalibrated~~ reconfigured such that the instance corresponding
to this peak EC value now serves as the reference starting point for each series. This adjustment ensures consistency in
the time frame ~~across for~~ all measurements, ~~thereby thus~~ facilitating more precise comparisons and analyses of ~~the~~ data
trends.
- The third subset comprises weekly physical parameters, which were measured after a 20-minute EC equilibration time.
180 It also contains related chemical analysis data from the water samples collected during these weekly events, along with
additional metadata recorded during the sampling process. This metadata includes information such as the water level
of the well before pumping, the water level of the well after pumping, the drawdown, and the volume of water removed
during each pumping event. Furthermore, this subset includes the daily mean water level obtained from the level logger
data.

185 2.1.3 Water sample preparation and analysis

All water samples were stored at 4 °C ~~prior to be~~ before being prepared for the different chemical analyses. They were filtered in the laboratory during the day ~~following the~~ after collection. A vacuum system and 0.45 µm cellulose acetate membrane filters, manufactured by Sartorius (type 111, 47 mm in diameter), were used. During ~~pre-processing~~ preprocessing of the samples, the EC was also measured ~~at~~ in the laboratory using a TetraCon® 925 probe. On the filtered water samples, the pH and alkalinity were measured using a Mettler Toledo equipment with 0.01N HCl up to pH 4.5; the dissolved silica (SiO₂), the ammonium (NH₄⁺) and the orthophosphate (HPO₄³⁻) concentrations and the absorbance at 254 nm (Abs254) were measured using spectrophotometry with a Skalar® continuous flow analyzer SAN++; the major cations (Na⁺, K⁺, Ca²⁺, Mg²⁺) and anions (Cl⁻, NO₃⁻, SO₄²⁻) concentrations were analyzed using ionic chromatography (Dionex ICS-5000 dual channel). On the filtered and acidified (1% HNO₃) water samples, the total content of major metals (Al, Fe, Mn, Ba, Sr) were analysed with a
190 Quadrupole ICP-MS (Agilent 7900) associated with an ISIS 3 (Agilent) injection system. The analyses were conducted in He mode and ¹⁰³Rh and ¹⁸⁵Re were used as internal standards. The calibration standards were prepared with Multi elements ICP standard solutions (CHEMLAB Analytical) diluted in 1% HNO₃. The analytical blank values were less than 1% of the lowest sample concentrations for all elements. For all analyses, the quantification limits are given in Table 2.

2.1.4 Data Quality

200 This study implements a comprehensive data quality control approach throughout its year-long ~~physico-chemical~~ physicochemical monitoring campaign, ensuring the reliability and accuracy of the ~~data collected~~ collected data. Parameters such as EC, DO, ORP, and pH were precisely measured using ~~standardized~~ standardised instrumentation, adhering to established protocols for environmental monitoring. Regular calibration of our equipment was ~~conducted~~ performed before each data collection to ensure ~~measurement accuracy~~ the accuracy of the measurement. This involved both factory standards and field adjustments to account
205 for environmental variables specific to the WEC. Meticulous sampling procedures were followed, complemented by established analytical techniques, to maintain methodological integrity. The ~~dataset's structured organization~~ structured organisation of the dataset, complete with detailed metadata, enhances its transparency and usability for further research. ~~Acknowledging~~ Recognising potential data variability challenges, such as probe equilibration times, which could be influenced by seasonal changes, we implemented procedures to ensure consistency across all measurements. We invite data users to engage with us
210 for validation of specific data subsets and to discuss any discrepancies, a step critical for maintaining the dataset's quality and integrity. These protocols collectively underscore the ~~dataset's robustness~~ robustness of the dataset, advocating for its proper citation and use in line with the highest standards of scientific inquiry.

2.1.5 Data Usage and Applications

~~When analysing~~ Understanding c-Q patterns in forested headwater catchments ~~, it is vital to understand the dynamics of~~
215 ~~biogeochemical processes taking place in the shallow horizons of the system~~ requires insight into the biogeochemical processes occurring in shallow subsurface horizons, such as redox reactions ~~or cation and anion exchanges in soil. Indeed, the varying~~

patterns of c-Q relationships observed in different catchments can be explained by the switching and ion exchange. The variability in c-Q relationships across catchments can often be linked to shifts in the dominance of different end-member source waters, which are driven by from shaped by subsurface biogeochemical heterogeneity (Zhi et al., 2019; Stewart et al., 220 2022). Our dataset potentially affords the capacity to quantify the relative contributions stemming from both

Our dataset provides an opportunity to explore the contributions of shallow and deep flowlines-flow paths during borehole pumping. This leverage is predicated upon the intrinsic course of electron acceptor succession (McMahon and Chapelle, 2008) ~~Consequently, this phenomenon instigates the formation and stratification~~ We propose that purging a borehole until the extracted water stabilizes reflects the natural mixing processes that occur when groundwater discharges into streams. By 225 conducting pumping tests under varying hydrological conditions and at multiple locations, we aim to investigate whether this approach can provide insights into the depth and spatial distribution of redox zones. As such, when borehole pumping is conducted contingent upon the hydrological state of the system, the composition of the extracted mixture can be computed based on the relative redox potential, which we hypothesize to be influenced by vertical permeability.

Since the succession of electron acceptors drives the stratification of the redox zone (McMahon and Chapelle, 2008), the 230 redox potential of extracted water (Eh) of said mixture. The provided database allows to address Gaps in the Current State of the Art on ~~may offer clues about the relative contributions of different flow paths. Although our dataset does not provide a direct map of redox zones, it offers high-resolution observations that may help assess their role in shaping groundwater chemistry and, by extension, stream composition.~~

The following section outlines how this dataset can potentially address key gaps in understanding redox processes, their 235 influence on c-Q relationships, and the broader implications for hydrobiogeochemical research.

– **High-Resolution Monitoring:** While Although previous studies have explored c-Q relationships and the influence of various water sources on stream chemistry, the novelty of this dataset lies in its this dataset provides high-resolution monitoring. By collecting data at frequent intervals, including the weekly measurement of four physical parameters, with a focus on capturing redox zone dynamics over space and time. By measuring four key parameters at weekly intervals, it of- 240 fers a level of detail that allows for a more nuanced understanding of dynamic catchment processes detailed perspective on these processes, complementing existing datasets that often emphasise other biogeochemical aspects. While not entirely novel in its approach, the dataset adds valuable insight into the spatio-temporal variability of redox conditions in forested catchments. This high-resolution approach addresses a gap where data may be limited to lower-frequency measurements. The temporal behaviour, at on different timescales, observed in GW5 is shown to illustrate the dynamic interplay between shallow and deep flowlines and how this influences the solute behaviour observed in the behavior observed in 245 groundwater (Figure 4 and Figure 5). This borehole was investigated in more detail and at higher frequency during two sequential sampling experiments conducted in high flow and low flow. It can be observed from these two graphs that the mixing ratio between the two flowlines (shallow and deep) changes as a function of the season in the catchment. The relationship between parameters that drive biogeochemical processes (EC, DO, ORPEh, ORP, Eh, pH – top of Figure 5) and the solute concentration (Ca^{2+} , Alkalinity, SO_4^{2-} and NO_3^- – bottom of Figure 5) can also be inferred and useful for 250 new hydrological and biogeochemical modelling approaches.

- **Quantifying Redox Reactions:** The dataset ~~significantly contributes to addressing a key gap in current knowledge related to the quantification of redox reactions within the shallow horizons of forested catchments~~gaps in understanding redox reactions in shallow forested catchments, as noted by ?, where "hot spots" and "hot moments" introduce significant variability. While the dataset provides empirical observations to inform future modelling efforts, it represents a step toward capturing this complexity rather than a definitive resolution of existing knowledge gaps. While previous research has acknowledged the importance of redox processes, this dataset provides the necessary empirical data to model and quantify these reactions accurately. As an example, understanding how ~~the~~ dissolved organic carbon (DOC), precipitation, and subsequent redox reactions affect the catchment's chemical composition represents a substantial advancement ~~(?)~~. In natural shallow systems, water can only be ~~oxidized~~oxidised/reduced under extremely oxidizing/reducing conditions, and the actual lower and upper Eh boundaries are governed by CO₂ reduction to CH₄ and the presence of O₂, respectively. Using Eh values, one can determine the dominant redox couple in a system. Because multiple electron donors can drive multiple redox reactions at the same time, systems rarely have strict redox boundaries in practice. The redox state of the soil and the unsaturated zone can vary greatly (Mansfeldt, 2003). It can be ~~oxidizing~~oxidising, weakly reducing, moderately reducing, or strongly reducing (≥ 400 , $200 - 400$, $100 - 200$, and -100 mV, respectively). Figure 6 shows the in-situ pH vs. Eh measurements taken throughout the one-year weekly campaign in the different boreholes. The defined zones, based on the redox zoning in soils (Zhang and Furman, 2021), indicate how the redox state of the soil and the unsaturated zone can vary greatly. Only boreholes GW1 and GW11 are seen to exhibit weakly reducing behaviour, with GW1 exhibiting a preference for moderately reducing conditions during the summer period. This behaviour can be understood, considering that GW11 represents the deepest currently observable flowline in the system (Figure 1-b), while GW1 corresponds to shallow groundwater on the plateau. GW1 represents vertically infiltrating water, influenced by redox processes associated with shallow and biologically active zones in the system.
- **Influence on pH and Ion Exchange:** The dataset ~~'s ability to capture the interplay between redox reactions and pH changes extends our understanding of CZ processes. Current research often focuses on individual aspects of the system, but this dataset enables a more holistic view by elucidating how redox conditions can influence pH and, consequently, the exchangeable pool of ions sorbed to the soil. This integrated approach addresses a gap in the current literature, which tends to treat these processes in isolation. Under steady-state chemical conditions, the composition of a cation exchanger will be the same as that of the groundwater around it~~provides high-resolution observations that can help identify the patterns of redox zone development and their potential influence on pH dynamics in shallow groundwater, as highlighted by ?. While it does not fully resolve these interactions, its spatial and temporal coverage could provide context for understanding how changes in water composition, such as acidification or vegetation shifts, impact groundwater chemistry through ion exchange processes. When the composition of the water changes due to acidification ~~, or or a~~ change in vegetation influence, the cation exchanger readjusts its composition to the new groundwater concentrations. As a result, the exchanger serves as a temporary buffer that can completely change the concentrations in water via a process known as ion chromatography (Appelo and Postma, 2004). The main sources of acidity in the most

near-surface environments are CO₂ from the atmosphere and organic acids formed by decaying DOC, resulting in an acid pH range of 5 ~~to~~ 6. Acid ~~sulfate~~ ~~sulphate~~ soils have lower pH values, with values as low as 4 occasionally recorded (DeLaune and Reddy, 2005). The lower bound for pH in natural shallow environments can then be assumed to be this low pH value.

- 290 – **Seasonal Variability and Flowline Dynamics:** ~~One of the dataset's unique strengths is its capacity to elucidate the~~ The dataset contributes to ongoing efforts to understand seasonal variability and flowline interactions within ~~the catchment.~~ ~~Previous studies may lack the temporal granularity needed to capture these nuances adequately. By demonstrating how the dominance of flowlines changes with~~ catchments, adding to the work of previous studies that have monitored these dynamics. While high-spatiotemporal-resolution datasets exist, this dataset provides complementary observations
- 295 that may help identify shifts in flowline dominance under varying hydrological conditions, ~~the dataset offers insights into hydrological dynamics that were previously underexplored in the literature.~~ Its focus on capturing redox zone development across different spatial and temporal scales offers an additional perspective on groundwater flow and chemistry in forested catchments. In Figure 7, boreholes GW3, GW7, GW9, GW10, and GW12 are grouped together as representing shallow flowlines. The shaded area in the graphs reflects the median trend of field parameters (indicated by
- 300 the dashed grey line), along with the 95% confidence interval derived from these five boreholes (depicted as a shaded grey area). On the other hand, the black line represents the deep flowline (GW11). The behaviour of both GW5 and GW2 is also depicted individually. It ~~'s~~ is evident that the dominance of one flowline over the other changes depending on the hydrological state of the system. Lastly, the behaviour of GW1 is also presented. It ~~'s~~ is apparent that this borehole is primarily influenced by chemical processes related to the consumption of dissolved oxygen and subsequent changes in
- 305 redox conditions.
- **Practical Implications:** Our findings on the relationship between stream chemistry and subsurface water chemistry have practical implications. ~~They suggest~~ It suggests that stream chemistry can serve as a valuable proxy for assessing and monitoring subsurface water chemistry. This bridges a gap in the current ~~state-of-the-art~~ state-of-the-art by highlighting the potential utility of readily available stream chemistry data for inferring subsurface conditions. To illustrate the respec-
- 310 tive influences of the shallow and deep flowlines on the temporal trend in the stream water at the outlet (labelled SW1 in Figure 1), a graph has been provided in Figure 6, where these influences are depicted in blue. This visual representation helps clarify how different flowlines contribute to changes in ~~stream-water chemistry~~ the chemistry of stream water over time.
- **Limitations and Future Work:** Although this high-frequency campaign spanned only one year, we captured two contrasting hydrological extremes: an exceptionally wet period and a severe drought, therefore covering a broad segment of the typical flow range of the catchment. However, these data cannot fully represent all possible interannual variability. To mitigate this limitation, we placed the one-year record in the context of our longer 10-year bi-weekly monitoring programme. This complementary dataset broadens the temporal scope and will enable future analyses to assess how often similar hydro-biogeochemical conditions recur and to detect potential shifts under changing climatic drivers.
- 315

In addition, while in-situ measurements of redox potential and dissolved oxygen offer valuable near-real-time insights, sensor drift remains a concern, particularly during prolonged deployment in varying temperature and flow regimes. We performed regular calibrations and spot checks to minimise errors, but subtle changes may still occur over time. Users of this dataset are encouraged to consider possible sensor bias when interpreting short-term fluctuations. Finally, weekly sampling, although substantially more frequent than traditional approaches, may still miss very rapid or localised events. Expanding temporal resolution during peak flow periods or employing automated samplers at multiple depths could help capture episodic phenomena and refine our understanding of the full biogeochemical complexity of the catchment.

3 Conclusions

The one-year monitoring campaign conducted within the WEC ~~has generated a rich dataset that extends beyond conventional hydrological and chemical monitoring efforts. By capturing the complex relationships among physical parameters, water chemistry~~ provides a dataset that contributes to ongoing efforts to understand the interactions between hydrological and biogeochemical processes in shallow groundwater. While similar datasets exist, this study adds to the existing body of work by offering high-resolution observations that may help to explore seasonal variability, flowline interactions, and redox reactions, ~~this dataset empowers researchers to more deeply explore the intricate processes governing redox zonation within the catchment. Moreover, its insights~~ zone development. The dataset does not provide a comprehensive understanding of these processes, ~~but offers empirical data that could support future research~~ into the influence of redox reactions on pH dynamics and ion exchange ~~widen our comprehension of how these interactions influence the catchment's biogeochemical landscape. As a dynamic dataset that incorporates both seasonal variability and flowline interactions, it emerges as a crucial tool for advancing our knowledge of CZ processes in forested headwater catchments. Weekly campaigns were executed to collect data on four physical parameters, including EC, DO, ORP, and pH from a comprehensive network of 10 sample points encompassing nine boreholes and one stream water. The temporal trends observed in DO and ORP through this monitoring campaign provide valuable insights into the intricate processes affecting redox conditions within the groundwater system. However, the true strength of this dataset resides in the continuous measurement of these physical parameters coupled with the concurrent collection of water samples at each monitoring point. This synergistic approach significantly enriches our understanding of the dynamic water quality characteristics within the WEC. The novelty and significance of this dataset are multifaceted: in forested catchments. These insights, though limited to a one-year period, provide a basis for considering key processes that can influence the hydrobiogeochemical behaviour of the catchment, as illustrated by the following aspects.~~

- **Modelling Redox Reactions in Shallow Horizons:** The dataset ~~offers essential parameters for accurately modeling~~ provides parameters that may be useful for investigating redox reactions within shallow horizons, ~~particularly highlighting the critical role of DOC in catalyzing chemical changes including observations related to DOC dynamics~~ during precipitation events. The ~~detailed documentation of the series of reactions initiated by the influx of DOC — its data capture processes such as DOC~~ oxidation, the subsequent release of CO₂, formation of carbonic acid, and the dissolution of bicarbonate minerals ~~— provides a comprehensive view of, which could help explore~~ the catchment's chemical responses

to rainfall, offering a nuanced understanding of the biogeochemical processes. While the dataset alone does not provide a comprehensive view of these processes, it may support further studies into the biogeochemical dynamics at play.

- 355 – **Influence of Redox Reactions on pH and Ion Exchange:** This dataset sheds light on the significant impact of redox reactions on pH levels and their potential cascading effects on the soil's ion exchange capacity. It allows for an in-depth examination of the complex relationship between redox changes and pH variations, which is vital for deciphering the broader implications of these shifts on the catchment's nutrient and solute dynamics, including the availability and mobility of essential nutrients and solutes. provides observations that may help to explore potential relationships between
- 360 redox reactions and pH variability in shallow groundwater. While the data are limited in scope and duration, they offer a foundation for investigating how changes in redox conditions might influence ion exchange processes and, consequently, solute concentrations. These observations contribute to ongoing efforts to better understand groundwater chemistry in forested catchments.
- 365 – **Hydrological Variability and Flowline Interactions:** The dataset uniquely captures the catchment's seasonal variability, emphasizing the intricate provides observations that can help investigate seasonal variability in the catchment, including potential differences in the dynamics between shallow and deep flowlines and their role in solute distribution. The detailed temporal analysis reveals how the interaction temporal patterns observed offer a basis for exploring how interactions between these flowlines evolves might evolve in response to seasonal variations, providing key insights into the contributing to a broader understanding of hydrological patterns and solute transport mechanisms that govern
- 370 the catchment's ecological and chemical processes in the catchment.

In conclusion, this dataset not only enriches the existing body of knowledge but also contributes to overcoming prevailing knowledge gaps. Through its high-resolution monitoring, emphasis on redox reactions, detailed exploration of while this high-frequency campaign lasted just one year, it included both a severe drought and an exceptionally wet period, effectively capturing two major flow extremes in the catchment. When combined with our ongoing 10-year bi-weekly dataset, these data

375 offer a more complete picture of the hydrological range and provide valuable insights into redox processes, pH dynamics, and comprehensive insights into seasonal and flowline dynamics, it substantially enhances our comprehension of CZ processes in forested headwater catchments. This is a substantial contribution for the scientific community, offering researchers and practitioners a robust foundation to seasonal flowline interactions. We believe these findings will help researchers refine predictive models, guide land management strategies, and foster a more integrated approach to catchment hydro-biogeochemistry.

380 By leveraging this dataset, the scientific community is poised to make significant strides in ecological conservation and resource management, contributing to a more sustainable and informed stewardship of natural landscapes decisions, and improve understanding of groundwater-driven ecosystem health.

Data availability. The database related to this data paper is publicly available at zenodo.org (<https://zenodo.org/records/10869166>) (Hissler et al., 2024). All datasets are cleaned, with outliers removed. A detailed description of all monitoring and sampling sites, including phys-

385 iographic characteristics, field equipment details, and sampling procedures, is presented in all shared .xlsx files. The hydrological time-series for each borehole and the discharge at the outlet are given in a separate file. Weekly timeseries of EC, DO, ORP, and pH and sample hydrochemistry are given in separate files. For more information on the available dataset, please contact the corresponding author (christophe.hissler@list.lu).

Author contributions. Karl Nicolaus van Zweel collected the majority of the field data as part of his Ph.D. research and contributed to the manuscript writing. Laurent Gourdol and Christophe Hissler provided significant assistance in data collection and manuscript preparation. All three authors participated in data processing. Jean-François Iffly was responsible for the field installations of the equipment and provided additional data. Loc Léonard conducted laboratory analyses and data capture, with the guidance of François Barnich. Laurent Pfister and Erwin Zehe offered technical input, reviewed the manuscript, and contributed to its writing. The collaborative efforts of all authors were invaluable in the completion of this manuscript.

395 *Competing interests.* The authors declare that they have no conflict of interest.

Disclaimer. TEXT

Acknowledgements. This work is part of the HYDRO-CSI project and was supported by the Luxembourg National Research Fund (FNR) in the framework of the FNR/PRIDE research programme (contract no. PRIDE15/10623093/HYDRO-CSI/Pfister). We ~~would like to~~ thank Jérôme Juilleret, ~~Viola Huck,~~ and Viola Huck for their contributions to the water sample preparation.

400 References

- Amundson, R., Richter, D. D., Humphreys, G. S., Jobbágy, E. G., and Gaillardet, J.: Coupling between biota and earth materials in the critical zone, *Elements*, 3, 327–332, 2007.
- Anderson, S. P., Anderson, R. S., and Tucker, G. E.: Landscape scale linkages in critical zone evolution, *Comptes Rendus. Géoscience*, 344, 586–596, 2012.
- 405 Antonelli, M., Glaser, B., Teuling, A. J., Klaus, J., and Pfister, L.: Saturated areas through the lens: 1. Spatio-temporal variability of surface saturation documented through thermal infrared imagery, *Hydrological Processes*, 34, 1310–1332, 2020a.
- Antonelli, M., Glaser, B., Teuling, A. J., Klaus, J., and Pfister, L.: Saturated areas through the lens: 2. Spatio-temporal variability of streamflow generation and its relationship with surface saturation, *Hydrological processes*, 34, 1333–1349, 2020b.
- Appelo, C. A. J. and Postma, D.: *Geochemistry, groundwater and pollution*, CRC press, 2004.
- 410 Brantley, S. L., Goldhaber, M. B., and Ragnarsdottir, K. V.: Crossing disciplines and scales to understand the critical zone, *Elements*, 3, 307–314, 2007.
- Brantley, S. L., Holleran, M. E., Jin, L., and Bazilevskaya, E.: Probing deep weathering in the Shale Hills Critical Zone Observatory, Pennsylvania (USA): the hypothesis of nested chemical reaction fronts in the subsurface, *Earth Surface Processes and Landforms*, 38, 1280–1298, 2013.
- 415 Brantley, S. L., McDowell, W. H., Dietrich, W. E., White, T. S., Kumar, P., Anderson, S. P., Chorover, J., Lohse, K. A., Bales, R. C., Richter, D. D., et al.: Designing a network of critical zone observatories to explore the living skin of the terrestrial Earth, *Earth Surface Dynamics*, 5, 841–860, 2017.
- Brooks, P. D., Chorover, J., Fan, Y., Godsey, S. E., Maxwell, R. M., McNamara, J. P., and Tague, C.: Hydrological partitioning in the critical zone: Recent advances and opportunities for developing transferable understanding of water cycle dynamics, *Water Resources Research*, 51, 6973–6987, 2015.
- 420 DeLaune, R. and Reddy, K.: Redox potential, *Encyclopedia of Soils in the Environment*, pp. 366–371, 2005.
- Fabiani, G., Schoppach, R., Penna, D., and Klaus, J.: Transpiration patterns and water use strategies of beech and oak trees along a hillslope, *Ecohydrology*, 15, e2382, 2022.
- Gourdol, L., Clément, R., Juilleret, J., Pfister, L., and Hissler, C.: Exploring the regolith with electrical resistivity tomography in large-scale surveys: Electrode spacing-related issues and possibility, *Hydrology and Earth System Sciences*, 25, 1785–1812, 2021.
- 425 Hissler, C., Martínez-Carreras, N., Barnich, F., Gourdol, L., Iffly, J. F., Juilleret, J., Klaus, J., and Pfister, L.: The Weierbach experimental catchment in Luxembourg: A decade of critical zone monitoring in a temperate forest-from hydrological investigations to ecohydrological perspectives, *Hydrological Processes*, 35, e14 140, 2021.
- Hissler, C., van Zweel, N., Gourdol, L., and Leonard, L.: One year of high frequency monitoring of groundwater physico-chemical parameters in the Weierbach Experimental Catchment, Luxembourg., <https://doi.org/10.5281/zenodo.10869166>, 2024.
- 430 Jin, L., Siegel, D. I., Lautz, L. K., Mitchell, M. J., Dahms, D. E., and Mayer, B.: Calcite precipitation driven by the common ion effect during groundwater–surface-water mixing: a potentially common process in streams with geologic settings containing gypsum, *Bulletin*, 122, 1027–1038, 2010.
- Juilleret, J., Dondeyne, S., Vancampenhout, K., Deckers, J., and Hissler, C.: Mind the gap: a classification system for integrating the subsolum into soil surveys, *Geoderma*, 264, 332–339, 2016.
- 435

- Li, L., Bao, C., Sullivan, P. L., Brantley, S., Shi, Y., and Duffy, C.: Understanding watershed hydrogeochemistry: 2. Synchronized hydrological and geochemical processes drive stream chemostatic behavior, *Water Resources Research*, 53, 2346–2367, 2017.
- Mansfeldt, T.: In situ long-term redox potential measurements in a dyked marsh soil, *Journal of Plant Nutrition and Soil Science*, 166, 210–219, 2003.
- 440 Martínez-Carreras, N., Wetzel, C., Frentress, J., Ector, L., McDonnell, J., Hoffmann, L., and Pfister, L.: Hydrological connectivity inferred from diatom transport through the riparian-stream system, *Hydrology and Earth System Sciences*, 19, 3133–3151, 2015.
- Martínez-Carreras, N., Hissler, C., Gourdol, L., Klaus, J., Juilleret, J., Iffly, J. F., and Pfister, L.: Storage controls on the generation of double peak hydrographs in a forested headwater catchment, *Journal of Hydrology*, 543, 255–269, 2016.
- McMahon, P. and Chapelle, F.: Redox processes and water quality of selected principal aquifer systems, *Groundwater*, 46, 259–271, 2008.
- 445 Moragues-Quiroga, C., Juilleret, J., Gourdol, L., Pelt, E., Perrone, T., Aubert, A., Morvan, G., Chabaux, F., Legout, A., Stille, P., et al.: Genesis and evolution of regoliths: Evidence from trace and major elements and Sr-Nd-Pb-U isotopes, *Catena*, 149, 185–198, 2017.
- Pfister, L., Martínez-Carreras, N., Hissler, C., Klaus, J., Carrer, G. E., Stewart, M. K., and McDonnell, J. J.: Bedrock geology controls on catchment storage, mixing, and release: A comparative analysis of 16 nested catchments, *Hydrological Processes*, 31, 1828–1845, 2017.
- Pfister, L., Bonanno, E., Fabiani, G., Gourdol, L., Hissler, C., Huck, V., Iffly, J. F., Keim, R., Martínez-Carreras, N., Mestdagh, X., et al.: 450 Fast motion view of a headwater creek—A hydrological year seen through time-lapse photography, *Hydrological Processes*, 37, e15 026, 2023.
- Riebe, C. S., Hahm, W. J., and Brantley, S. L.: Controls on deep critical zone architecture: A historical review and four testable hypotheses, *Earth Surface Processes and Landforms*, 42, 128–156, 2017.
- Scaini, A., Hissler, C., Fenicia, F., Juilleret, J., Iffly, J. F., Pfister, L., and Beven, K.: Hillslope response to sprinkling and natural rainfall using 455 velocity and celerity estimates in a slate-bedrock catchment, *Journal of Hydrology*, 558, 366–379, 2018.
- Stewart, B., Shanley, J. B., Kirchner, J. W., Norris, D., Adler, T., Bristol, C., Harpold, A. A., Perdrial, J. N., Rizzo, D. M., Sterle, G., et al.: Streams as mirrors: Reading subsurface water chemistry from stream chemistry, *Water Resources Research*, 58, e2021WR029 931, 2022.
- Weiler, M. and McDonnell, J. J.: Testing nutrient flushing hypotheses at the hillslope scale: A virtual experiment approach, *Journal of Hydrology*, 319, 339–356, 2006.
- 460 Welch, L. and Allen, D.: Hydraulic conductivity characteristics in mountains and implications for conceptualizing bedrock groundwater flow, *Hydrogeology Journal*, 22, 1003, 2014.
- Wrede, S., Fenicia, F., Martínez-Carreras, N., Juilleret, J., Hissler, C., Krein, A., Savenije, H. H., Uhlenbrook, S., Kavetski, D., and Pfister, L.: Towards more systematic perceptual model development: a case study using 3 Luxembourgish catchments, *Hydrological processes*, 29, 2731–2750, 2015.
- 465 Wymore, A. S., Yang, W. H., Silver, W. L., McDowell, W. H., and Chorover, J.: *Biogeochemistry of the Critical Zone*, Springer, 2022.
- Xiao, D., Brantley, S. L., and Li, L.: Vertical connectivity regulates water transit time and chemical weathering at the hillslope scale, *Water Resources Research*, 57, e2020WR029 207, 2021.
- Zhang, Z. and Furman, A.: Soil redox dynamics under dynamic hydrologic regimes-A review, *Science of the Total Environment*, 763, 143 026, 2021.
- 470 Zhi, W., Li, L., Dong, W., Brown, W., Kaye, J., Steefel, C., and Williams, K. H.: Distinct source water chemistry shapes contrasting concentration-discharge patterns, *Water Resources Research*, 55, 4233–4251, 2019.




Figure 1. Presentation of the Weierbach Experimental Catchment: (a) topography and Topography with water sampling location-locations, including contour lines representing 5-meter elevation intervals (altitudinal values not labeled; catchment elevation ranges from 450 to 500 m), and (b) regolith structure. The black contour line in the map (a) marks the chemical weathering front basement, derived from ERT surveys and core drillings (Gourdol et al (Gourdol et al., 2021), 2021).




Figure 2. Daily total precipitation in the Weierbach Experimental Catchment and the daily mean discharge at the catchment outlet from 2006 to 2022.




Figure 3. The schematic at the top illustrates the flow-through cell used to measure in-situ electrical conductivity (EC), dissolved oxygen (DO), oxidation-reduction potential (ORP), and pH. The ORP and pH probes were placed in parallel to avoid interference. Two WTW® Multi 3630 IDS data loggers were used to automatically record the probe data. The two photos at the bottom left and right show the setup used in the Weierbach Experimental Catchment and its field application.




Figure 4. High-frequency time series of the physico-chemical parameters recorded at GW5 on March 24th, 2021 (left) and August 2nd, 2022 (right).




Figure 5. Time series of the physico-chemical parameters (top) and the related solute concentrations (bottom) at GW5, collected during the weekly sampling campaigns from March 2021 to June 2022.




Figure 6. Eh-pH diagram depicting weekly in-situ field observations collected over the course of a year. The zones depicted on the graph are adapted from Zhang and Furman (2021). The upper and lower bounds represent the Eh - and pH stability ranges for water in natural environments. Adapted from Zhang and Furman (2021).




Figure 7. Time series of dissolved oxygen (DO), Eh, pH, and electrical conductivity (EC) from March 2021 to August 2022 for boreholes in the Weierbach Experimental Catchment. The grey dashed line and shaded area denotes the median and 5-95% bounds for the 'Shallow flowline' group, including GW3, GW7, GW9, GW10, and GW12. LOESS fits for SW1, GW1, GW2, GW5, and GW11 illustrate the variation in parameter trends across different locations within the catchment, highlighting differences between shallow and deep flowlines. GW11, located in the streambed, represents the deep flowline and the catchment's deepest point.

Table 1. Characteristics of the boreholes selected for the monitoring of the groundwater physico-chemical parameters during this study.

ID	X Wgs84- X WGS84	Y wgs84- Y WGS84	Sampling Depth	Standpipe	Diameter	Screen
GW1	5.79676	49.83466	220 cm	32 cm	50 mm	1 m depth to bottom
GW2	5.79623	49.83110	195 cm	35 cm	50 mm	1 m depth to bottom
GW3	5.79721	49.82832	222 cm	32 cm	50 mm	1 m depth to bottom
GW5	5.79715	49.83468	753 cm	33 cm	32 mm	1 m depth to bottom
GW7	5.79465	49.83225	555 cm	18 cm	32 mm	1 m depth to bottom
GW9	5.79713	49.83332	628 cm	28 cm	40 mm	1 m depth to bottom
GW10	5.79706	49.83146	299 cm	31 cm	40 mm	1 m depth to bottom
GW11	5.79709	49.83137	100 cm	30 cm	40 mm	no screen
GW12	5.79750	49.83220	305 cm	40 cm	40 mm	1 m depth to bottom

Table 2. Quantification limit (QL) of the chemical parameters considered in this study.

	SiO ₂	NH ₄ ⁺	HPO ₄ ³⁻	Na ⁺	K ⁺	Ca ²⁺	Mg ²⁺	Cl ⁻	NO ₃ ⁻	SO ₄ ²⁻	Al	Fe	Mn	Ba	Sr
	mg L ⁻¹										μg L ⁻¹				
QL	0.1	0.01	0.15	0.1	0.02	0.2	0.1	0.1	0.04	0.1	4.0	1.0	0.1	50	30