Response to essd-2024-264 RC2

- The referee's comments are in blue
- The authors' responses are shown in black

This study focuses on the development on a new high-resolution dataset from Landsat 8 for retrieval of sea ice concentration. There is novelty in this approach compared with more traditional passive microwave methods, and whilst the notion of sea ice concentration retrieval from Optical/IR methods is not intrinscially new, the research presented here demonstrates work of significant value – it is very nice to see research that transitions away from the MODIS sensor (which is approaching end of life and Terra no longer has a regular orbit), and the amount of data processed at high resolution is very novel, and quite exciting! I wholeheartedly agree with the authors thoughtful and well-presented argument of the importance of an independent from PMW SIC dataset given in the introduction.

The code used in this study is also on zenodo, this is really nice to see! I particularly was appreciative of the 'requirements.txt' outlining the specific versions of packages used in this study, this is excellent to see from the perspective of reproducibility. The code is organised with a logical structure and is sufficiently commented and documented for publication.

The dataset is also published as linked on zenodo, which is a suitable archive. The data is in a typical netcdf format with sensible file names, the time parameter is logical with a clearly defined epoch in the metadata, fill values indicated, and it is intuitive and easy to open and plot this data, which looked reasonable when I plotted some data in the Central Arctic.

This is a very good and innovative concept which has been robustly executed. The methodology is sound. On the whole, the manuscript is thorough, but at this stage there are some gaps in justification of particular choices and insufficient detail in the some of the explanations, particularly in regards to the visual inspection component of cloud masking, as I have outlined in my Specific Comments. Therefore, it is my view that this manuscript should be published subject to minor revisions. The manuscript would benefit from some additional proofing, I outline minor typographical corrections where I see them but do not intend for this to be exhaustive.

The authors sincerely appreciate for your positive evaluation of the manuscript and thank you for your detailed and constructive comments, which definitely helped the manuscript to be more complete in terms of justification of the particular choices made during the process of SIC production/validation and details of the processes undertaken to produce the Landsat-8 SIC dataset. Please check the response to the comments below.

Specific Comments:

General Introduction: There is, of course, already a 1km SIC dataset produced by the University of Bremen based on merging Optical/IR MODIS data with PWM methods. The value of your work is that is truly independent of PWM, and the UoB dataset isn't. Using MODIS in 2024 also has it's own problems. However, I think you need to write a paragraph on this dataset in the intro, and explain the niche of what your dataset does that this one doesn't. Certainly, without reference to this dataset I do not feel your argument can be considered complete.

Ludwig, V., G. Spreen, & L. T. Pedersen (2020). Evaluation of a New Merged Sea-Ice Concentration Dataset at 1 km Resolution from Thermal Infrared and Passive Microwave Satellite Data in the Arctic. Remote Sens. 12(19), 3183. doi:10.3390/rs12193183 [Article (PDF file)]

The authors sincerely thank you for your suggestion and sharing this great reference with us, which helped to make a strengthened introduction. The authors fully agree that for a complete argument of the necessity of a "fully independent" reference SIC, a discussion of the MODIS/AMSR2 merged SIC product should be included. In the revised manuscript, a paragraph discussing the dataset by Ludwig et al. (2020) and the need for a fully independent dataset despite the presence of an existent high-resolution and pan-Arctic SIC dataset was included.

[Added] 'Recently, efforts to leverage the advantages of both VIS/NIR sensors and PMW sensors for retrieving SIC have been explored through data merging techniques. Ludwig et al. (2020) used a combination of MODIS and AMSR2 measurements to construct high-resolution (1 km) and spatially continuous SIC data over pan-Arctic areas. This approach exploited the benefits of the 1 km resolution MODIS imagery while mitigating its inherent disadvantage of spatial discontinuity due to clouds by introducing the AMSR2 measurements. While the SIC dataset produced by Ludwig et al. (2020) is both high-resolution and covers pan-Arctic areas, due to the retrievals being dependent on the AMSR2 measurements, the product cannot be considered a fully independent reference data for PMW SIC validation. Therefore, it is still necessary to construct a dataset of Arctic SIC that is fully independent of PMW measurements.'

Reference:

Ludwig, V., Spreen, G., and Pedersen, L. T.: Evaluation of a New Merged Sea-Ice Concentration Dataset at 1km Resolution from Thermal Infrared and Passive Microwave Satellite Data in the Arctic, Remote Sens., 12(19), 3183, doi: <u>https://doi.org/10.3390/rs12193183</u>, 2020.

Line 107: I think there is a better way that 'the rest of this paper' but I like seeing an outline at this point in the paper. I'd probably also do a line break before to enhance clarity.

Thank you for this comment. Following your suggestion, we have rephrased 'the rest of this paper' into 'the remaining sections of this paper are' and have added a line break to enhance clarity in the revised manuscript.

[Old] (Line 107) 'The rest of this paper is organized as follows:...'

[New] 'The remaining sections of this paper are organized as follows:'

Line 117: This is probably a good time to mention the 'pole-hole' or unimaged portion of the central Arctic owing to Landsat-8 inclination. You could also do a diagram to illustrate which bit of the Arctic cant be imaged, but I leave this to your judgment. Noted that you explain this in line 130 but it really is quite important for it to go as soon as the sensor is introduced in my opinion.

The authors appreciate your valuable suggestions which led to a much-improved visibility and

clarity of Fig. 1. In the revised manuscript, illustration of the 'pole-hole' is added as the hatched area in Fig. 1 along with the caption explaining the pole hole. In order to incorporate your comment regarding the timing of bringing up the pole hole, in the revised manuscript, the authors restructured the paragraph to mention the pole hole right after the introduction of the OLI sensor as the following:

[Removed] (Line 130) 'It should also be noted that, due to the sun-synchronous orbit of Landsat-8 along with the narrow swath-width of OLI, Landsat-8 does not measure areas where latitude is higher than 82°N.'

[Added] '...with a spatial resolution of 30 m. It should be noted that areas with a latitude higher than 82°N in the northern hemisphere are not measured by Landsat-8 (i.e. the hatched area in Fig. 1) due to the orbit inclination of Landsat-8 and the relatively narrow swath width of the OLI.'



Figure 1: Footprints of the collected Landsat-8 images over each region of the pan-Arctic areas during the period of Jan. 2020 – Dec. 2022. The hatched region denotes the areas unmeasured by Landsat-8 due to its orbital inclination (i.e., pole hole). The regions of the pan-Arctic areas were distinguished using the region mask provided by Meier and Stewart (2023). The map projection is NSIDC Sea Ice Polar Stereographic North (EPSG: 3413) and the map was plotted using Python.

Line 122: I appreciate why Landsat-8 was chosen over Sentinel 2 given data coverage and robustness of cloud masking. However, I think you need to explicitly explain why you chose Landsat-8, as someone less familiar with Arctic sea ice research (and hasn't had the pleasure of trying to get Sen2Cor working over sea ice!) is going to be scratching their head wondering why you didn't choose Sentinel 2 (given the bands you use are overlapping, Landsat-8 band 5 and 6 approx. equal to S2 8a and 11, and S2 has a better resolution and repeat time). So, my advice here is to add in a couple of sentences explaining why you chose Landsat-8 and make an explicit reference to why you didn't choose Sentinel-2.

Thank you for mentioning this. The authors agree that an explanation for making the specific

choice of Landsat-8 over other available high-resolution measurements, such as Sentinel-2, would help justify this choice. To do this, an explanation of the more robust cloud mask of the Landsat-8 product relative to the Sentinel-2 product was given along with two citations supporting this argument (Zhu et al., 2015; Tarrio et al., 2020).

[Added] It is worth noting that the methods developed in this study (described in Section 3) utilize the NIR and SWIR bands for SIC retrieval and are therefore applicable to a wider range of high-resolution sensors that observe at similar bands, including the Multi-Spectral Instrument (MSI) onboard Sentinel-2. However, due to the more robust cloud mask performance of the Landsat-8 product, in this study, the Landsat-8 Collection 2 Level 1 product was selected to be used for the production of reference SIC data (Zhu et al., 2015; Tarrio et al., 2020).

References:

Zhu, Z., Wang, S., and Woodcock, C. E.: Improvement and expansion of the Fmask algorithm: cloud, cloud shadow, and snow detection for Landsats 4-7, 8, and Sentinel 2 images, Remote Sens. Environ., 159, 269-277, doi: <u>http://dx.doi.org/10.1016/j.rse.2014.12.014</u>, 2015.

Tarrio, K., Tang, X., Masek, J. G., Claverie, M., Ju, J., Qiu, S., Zhu, Z., Woodcock, C. E.: Comparison of cloud detection algorithms for Sentinel-2 imagery, Sci. Remote Sens. 2,, 100010, doi: <u>https://doi.org/10.1016/j.srs.2020.100010</u>, 2020.

Line 127: What method of cloud mask is implemented here? Is it appropriate? Why did you choose 10%?

The method of cloud mask implemented here is CFMask, which is the official cloud masking algorithm that generates the quality assessment band (i.e. an array containing masking information for clouds, cloud shadows, cirrus, and fill values) of Landsat-8.

The 10% threshold was given as a search criterion upfront to minimize the acquisition of cloud contaminated scenes. Admittedly, it would be ideal to give a 0% threshold as a criterion upfront for cloud cover, however, this leads to the removal of too many Landsat-8 scenes with clear sky portions available for SIC retrieval (Fig. S1, Table S1). In addition, since cloud pixels are removed from the application of CFMask prior to SIC production, the threshold for the Landsat-8 cloud cover during the acquisition of the data was relaxed to 10%.

Justification of the 10% threshold as well as the method of cloud mask implemented in this study was added in the revised manuscript. Fig. S1 and Table S1, which show the relationship between the number of available Landsat-8 scenes and the cloud cover threshold, were added in the supplements.

[Old] 'only Landsat-8 images with less than 10% daytime cloud cover (solar elevation higher than 15°) were collected.'

[New] 'only Landsat-8 images with less than 10% cloud cover (based on fractional cloud masked area from the quality assessment band of Landsat-8) during daytime (solar elevation higher than 15°) were collected. While the threshold of 0% cloud cover would ensure the acquisition of the least cloudy scenes, this also results in the loss of a considerable number of Landsat-8 scenes that contain clear-sky portions (see Fig. S1 and Table S1 in the supplements

for the number of available Landsat-8 scenes subject to different threshold values of cloud cover). Therefore, the threshold value for cloud cover was relaxed to 10% during the acquisition of Landsat-8 images.'

[Added] (Supplements) Figure S1 and Table S1 Number of available Landsat-8 images subject to varying cloud cover thresholds



Figure S1: Number of available Landsat-8 images (y-axis) subject to values of different cloud cover thresholds (x-axis) during the period of Jan. 2020 – Dec. 2022 over pan-Arctic areas.

Cloud cover threshold	0	5	10	15	20
Number of available	5,231	11,787	15,286	18,177	21,060

Table S1: Number of available Landsat-8 images subject to values of different cloud cover thresholds during the period of Jan. 2020 – Dec. 2022 over pan-Arctic areas.

Section 2.4: I agree with the study choice to use Ice Charts for validation, and I think you have done a good job in conveying why this is the most appropriate choice.

The authors thank you for your support in our choice of validation data.

Line 224: Great explanation!

Thank you for the positive comment.

Line 230: The phrase 'a visual inspection' is not exhaustive enough an explanation for this step. I want to know exactly what this process entailed, what images you looked at, what questions you asked, how you looked at them, how many images you looked at, whether anyone else

looked at them. I think a diagram of one typical judgement you made would be assistive. I need the thought process here to be outlined to the extent that I could replicate this step and generate the labels myself purely based on what is on the page, and it feels too muddled at the moment for me to plainly understand your meaning.

Thank you for pointing this out. The authors agree with your comment and the necessity of a more detailed description of the visual inspection process. In the revised manuscript, details of the visual inspection process, including the exact steps taken in the visual inspection process, an example case to show how the visual inspection was executed, who did the inspection, and the time it took to perform the inspection, are added in the appendix. A sentence to point the readers to the appendix was also added in Section 3.1.

[Added] 'Further details of this visual screening step are provided in Appendix B.'

[Added] Appendix B: Visual Inspection for Cloud Mask Quality Control

'In this section, a step-by-step description of the process taken to perform the visual inspection of Landsat-8 scenes is presented. As defined in Section 3.1, each pixel in a Landsat-8 scene can be sorted into the following four categories depending on the state of the cloud mask for the pixel: False negative (FN; cloud pixel mistaken as clear pixel), false positive (FP; clear pixel mistaken as cloud pixel), true negative (TN; clear pixel identified as clear pixel), and true positive (TP; cloud pixel identified as cloud pixel). It is noted that the pixels with FN are used to calculate SICs while the pixels with FP are not, indicating that the presence of FN pixels can directly introduce errors in the calculated SIC value. Therefore, visual inspection was performed very strictly to detect FN pixels.

Figure B1 outlines the steps taken to perform the visual inspection. The descriptions of each step are provided along with an example case of a Landsat-8 scene that is categorized into C1 during the section-wise inspection stage.

Step 1. Generating jpeg file of cloud mask (i.e., cloud mask image).

For each Landsat-8 scene, a false-colour image with each pixel classified as ice (white pixels in Fig. B2b, d), open water (blue pixels in Fig. B2b, d), cloud (grey pixels in Fig. B2b, d), and fill value (black pixels in Fig. B2b, d) is constructed using the OpenCV module in Python. Ice and open water pixels are then differentiated using the method described in Section 3.2. Cloud pixels are classified by masking the medium confidence cloud, high confidence cirrus, cloud shadow, and dilated cloud pixels identified by the quality assessment band (i.e., the cloud mask array produced by CFMask).

Step 2. Comprehensive inspection of cloud mask quality.

The cloud mask images generated in Step 1 are visually inspected against the truecolour image to identify sections populated with FN, FP, TN, or TP pixels. This is done in the following order: First, if no cloud pixels are observed from both the cloud mask image and the true-colour image (i.e., all pixels in the image are TN pixels), the scene is labelled as C4. Second, if any cluster of FN pixels is observed, the scene is labelled C1. Third, if any cluster of FP pixels is identified, the scene is labelled C2 and passed on to Step 3. If the clusters of cloud pixels in the cloud mask image are well corresponding to the position of clouds observed in the true-colour image (i.e., TP pixels), the scene is labelled C3 and passed on to Step 4 (Fig B2a, b). Step 3. Comprehensive inspection of cloud mask quality for C2.

For the scenes passed on to this step (i.e., scenes labelled C2 from Step 2), the cloud mask image is recreated using a higher confidence threshold (i.e., high confidence cloud, high confidence cirrus, cloud shadow, and dilated cloud pixels) for the quality assessment band. The new cloud mask image is visually inspected against the true-colour image, and if any cluster of FN pixels are observed, the confidence threshold for the quality assessment band is returned to its initial value (i.e., medium confidence cloud, high confidence cirrus, cloud shadow, and dilated cloud pixels). If the observed cluster of cloud pixels in the new cloud mask image is well corresponding to the position of clouds observed in the true-colour image, the higher confidence threshold is kept, and the scene is passed on to Step 4.

Step 4. Section-wise inspection of cloud mask quality.

In this step, the identified clusters of TP pixels are inspected in more detail. For each cluster of TP pixels observed, we zoom in (i.e. about 1000×1000 pixels; the full-size image is approximately 8000×8000 pixels) to the section of the cluster to check for the existence of FN pixels. If any FN pixels are found within the cluster, the scene is labelled C1 (Fig B2c, d).

An example of how a Landsat-8 scene may be categorized according to the process described in Fig B1 is presented using the case of a Landsat-8 scene acquired on Mar. 25, 2022, over the Barents Sea (Fig B2). First, from Step 2. (i.e., the comprehensive inspection step), visual inspection of the cloud mask image (Fig. B2b) against the true-colour image (Fig. B2a) shows that the position of clouds in the cloud mask array is generally well corresponding to those observed in the true-colour image. Therefore, at this step, this scene is labelled C3 and passed on to Step 4 as described above. Next, the section-wise inspection of the cloud mask quality is performed by zooming in to the cloud areas. This is illustrated in Fig. B2c and Fig. B2d, which is a zoomed in image of the area enclosed by the red rectangle in Fig. B2a and Fig. B2b. Inspection of this sub-section shows the presence of unmasked cloud shadow pixels, which results in the erroneous classification of ice as open water. Therefore, at this step, the label of this scene is changed to C1.

The visual inspection was done by HJ and it took approximately 5 - 10 minutes to inspect one Landsat-8 scene for cloud cover.'



Figure B1: Processing pipeline of the visual inspection step. Each Landsat-8 image is labelled as C1 (i.e., underestimated cloud cover), C2 (i.e., overestimated cloud cover), C3 (i.e., correctly estimated cloud cover for cloudy sky), or C4 (i.e., correctly estimated cloud cover for clear sky) depending on the observed dominance of true negative (TN; clear pixels identified as clear pixels), false negative (FN; cloud pixels mistaken as clear pixels), false positive (FP; clear pixels mistaken as cloud pixels), and true positive (TP; cloud pixels identified as cloud pixels) pixels.



Figure B2: The case of a Landsat-8 scene classified as C1 (i.e., underestimated cloud cover) from the visual inspection step. Shown in the panels are (a) full size true-colour image, (b) full size cloud mask array, (c) true-colour image of the area enclosed by the red rectangle in (a) and (b), and (d) cloud mask array of the area enclosed by the red rectangle in (a) and (b). The blue, white, gray, and black pixels in (b) and (d) are open water, ice, cloud, and fill value pixels, respectively. The scene is from Mar. 25, 2022, over the Barents Sea. The true-colour image is obtained from Earth Resources Observation and Science (EROS) Center (2020).

Line 270: Are these Top of Atmosphere (TOA) or not? In my view it is OK to use TOA for Arctic studies as the aerosol profiles are very poorly constrained and the magnitude of the atmospheric component is small compared to the bright surface, but this should be made very clear. If these are not TOA what processing was applied, and what is the justification for appropriateness?

Thank you for pointing this out. The values of Q_{DN} are values at TOA. This was made explicitly in the revised manuscript.

[Old] '...and Q_{DN} is the reflectivity...'

[New] '... and Q_{DN} is the TOA reflectivity...'

Line 355: This is robust error propagation and well justified.

Thank you for the positive comment.

Line 380: Thanks for making this clear, it's the right choice and good to see.

Thank you for the support of our choice.

Line 461: This is actually quite similar to how MOD29 is derived, which is considered to be one of the better cloud masks over sea ice. I leave it to your judgment if you feel inclusion of that would strengthen or weaken your case, but I think it is robust in any case.

Thank you for mentioning this. However, for this case, the authors feel it would be better to not include a discussion about the MOD29 for the sake bringing more focus to the validation results presented in Section 5.1 rather than the methods used to perform the validation.

Section 5.2: There does seem to be a notable bias between the ice chart and your dataset at lower SIC%. Can you comment on why this might be?

Thank you for this comment. There is indeed a bias in that the ice chart SICs are notably higher than the Landsat-8 SICs. This is one of the known characteristics of ice chart SICs as pointed out in the references by Tonboe et al. (2016) and Cheng et al. (2020), which were used in the original manuscript to explain the bias observed in Fig. 9.

- Tonboe et al. (2016) reports the overestimation of SIC by ice charts when compared with SIC retrievals from PMW measurements (most pronounced in the lower SIC range) and attributes this bias to "better-safe-than-sorry" approach often taken by the ice charting community.
- Cheng et al. (2020) also reports the overestimation of ice concentrations by ice analysts when compared to ice concentrations derived from automatic image segmentation.

In order to better explain this bias within the manuscript, comments regarding the higher bias at lower SIC range were inserted in the revised manuscript. In addition, possible causes of this bias, which are discussed in the studies of Tonboe et al. (2016) and Cheng et al. (2020), were also incorporated into the revised manuscript.

[Old] 'In addition, SIC from the ice chart was found to be positively biased to Landsat-8 SIC which is also supported by previous works of Tonboe et al. (2016) and Cheng et al. (2020).'

[New] 'In addition, SIC from the ice charts tends to be higher than that found from Landsat-8 SIC and the bias is more pronounced in the lower SIC range. This type of state-dependent overestimation of SIC from ice charts has been reported in previous works of Tonboe et al.

(2016) and Cheng et al. (2020), which shows that overestimation of SIC from ice charts is largest in the lower SIC range due to the "better-safe-than-sorry" practices of the ice charting community.'

References

Tonboe, R. T., Eastwood, S., Lavergne, T., Sørensen, A. M., Rathmann, N., Dybkjær, G., Pedersen, L. T., Høyer, J. L., and Kern, S.: The EUMETSAT sea ice concentration climate data record, The Cryosphere, 10, 2275-2290, doi: <u>https://doi.org/10.5194/tc-10-2275-2016</u>, 2016.

Cheng, A., Casati, B., Tivy, A., Zagon, T., Lemieux, J. F., Tremblay, L. B.: Accuracy and interanalyst agreement of visually estimated sea ice concentrations in Canadian Ice Service ice charts using single-polarization RADARSAT-2, The Cryosphere, 14, 1289-1310, doi: <u>https://doi.org/10.5194/tc-14-1289-2020</u>, 2020.

Line 578-59: always lovely to see DOIs but these should be referenced properly as datasets and cited in text.

Thank you for this catching. The in-text citations of each dataset are included in the revised manuscript.

Section 5.3: At what point in the season is this Landsat-8 data or is it the whole dataset? Would a more specific analysis around peak melt pond presence perhaps be better? (It is perfectly fine if you do not agree with this but adjust the manuscript to justify what choice was made and why in any case).

Thank you for this comment. The analysis presented in Section 5.3 was performed over six Landsat-8 scenes that were acquired over Jul. 2020, Aug. 2020, and Jul. 2021, which correspond to the melting season in the Arctic. A table showing the Landsat-8 scenes used in the evaluation is provided in Section S3 of the supplements. In the revised manuscript, a sentence pointing to Section 2.5, where the method used to select the six Landsat-8 scenes is explained, is added as well as the time periods of the scenes.

Filename (Melt Pond Fraction)	Filename (Landsat-8)	Time Difference [sec[
20210719_T45XVK_s2_mpf.nc	LC08_L1TP_173002_20210719_20210729_02_T1	-1539
20200806_T31XDL_s2_mpf.nc	LC08_L1GT_023243_20200806_20200916_02_T2	7268
20200806 T31XDL s2 mpf.nc	LC08 L1TP 023244 20200806 20200916 02 T1	7292
20200711_T13XEL_s2_mpf.nc	LC08_L1TP_065001_20200711_20200912_02_T1	-5904
20210719_T14XMQ_s2_mpf.nc	LC08_L1TP_068001_20210719_20210729_02_T1	-2378
20200711 T13XEL s2 mpf.nc	LC08 L1TP 081244 20200711 20200912 02 T1	5842

Table S3: List of Landsat-8 scenes and melt pond fraction datasets used for evaluation. Time differences between the two data for each scene are also tabulated.

[Added] 'As mentioned in Section 2.5, a total of six Landsat-8 scenes obtained from the periods of Jul. 2020, Aug. 2020, and Jul. 2021 were used in the evaluation.'

The method used to select the dataset for evaluation is described in detail in Section 2.5 of the original manuscript. The reason for using six datasets was because of the availability of the melt pond fraction dataset when subject to spatiotemporal collocation with the Landsat-8 SIC dataset. In the revised manuscript, this was rephrased to indicate that the data selection was not arbitrary, but rather limited by availability.

[Old] 'In this study, the total of six MPF datasets that are spatially overlapped with the coverage of Landsat-8 SIC dataset and have time difference of less than 3 hours with the Landsat-8 scene were selected for evaluation of variation in Landsat-8 SIC due to melt pond presence. The list of selected MPF datasets and the corresponding Landsat-8 scenes can be seen in Table S3 of the supplements.'

[New] 'In this study, each MPF dataset was tested for spatiotemporal overlap (time difference of less than 3 hours) with the coverage of Landsat-8 SIC. The total of six MPF datasets were found to be overlapping with the coverage of Landsat-8 SIC and thus available for use in the evaluation. The list of available MPF datasets and the corresponding Landsat-8 scenes can be seen in Table S3 of the supplements.'

Figures, Tables, and Equations:

I feel that your captions lack detail throughout. Figures and Tables should have terms used redefined so they can be interpretable independently of the text surrounding it. This will really help improve the clarity of the overall paper and should not be too arduous. You have done a very good job of defining terms within equations - I want to see the same clarity from the captions associated with your Tables and Figures.

The authors appreciate your comments regarding lacking information in the captions and wholeheartedly agree that the caption of the figures need to be edited for better clarity. In the revised manuscript, all the captions have been revised. Please look through the details of the changes made below.

Figure 1: Add reference to the mask you used to make this figure. Please also add in the projection and the software you used to make this map. I do not think a scale is necessary on a pan-Arctic map. I would also prefer to see your lat/lon labels appear in-front of the boxes, rather than behind, ie. The 70 deg is partially obscured in the Laptev sea. Please also reiterate the 'study period' with dates, the figure should be as self-contained as possible without reference to the text.

Thank you for the detailed comments. The revised caption now incorporates the reference of the region masks, the map projection, the software used to make the map, and specification of the 'study period'.

The 70 deg, which was obscured by the boxes, has been moved to the front of the boxes, and a hatched area corresponding to the pole hole was added to the figure.

[Old] Caption of Figure 1: 'Footprints of the collected Landsat-8 images over each sub-region over the pan-Arctic areas during the study period.'

[New] 'Footprints of the collected Landsat-8 images over each region of the pan-Arctic areas

during the period of Jan. 2020 – Dec. 2022. The hatched region denotes the areas unmeasured by Landsat-8 due to its orbital inclination (i.e. pole hole). The regions of the pan-Arctic areas were distinguished using the region mask provided by Meier and Stewart (2023). The map projection is NSIDC Sea Ice Polar Stereographic North (EPSG: 3413) and the map was plotted using Python.'



The old and new version of Fig. 1 can be seen as follows:

Figure 2: As with figure 1, add a citation for the regional mask and please also add in the projection and the software you used to make this map.

As was done with Fig. 1, the reference, the map projection, and the software used to make the map were added to the caption.

[Old] Caption of Fig. 2: 'Geographic distribution of the designated sub-regions of the Arctic Ocean, based on NSIDC Sea Ice Region Mask data.'

[New] 'Geographic distribution of the designated regions of the Arctic Ocean, based on NSIDC Sea Ice Region Mask data (Meier and Stewart, 2023). The map projection is NSIDC Sea Ice Polar Stereographic North (EPSG: 3413) and the map was plotted using Python.'

Figure 3: Really like to see diagrams like this, they're very helpful graphical illustrations of your work and improve readership comprehension. Good job!

Thank you for your encouragement. It's great to know that the diagram helped with readership comprehension.

Table 2: As with figures, tables should be as self-contained as possible. Please add a very brief explanation of C1, C2, C3, and C4 in your caption, as you have with Figure 5.

Thank you for this catching. Explanations of C1, C2, C3, and C4 have been added in the revised caption. In addition, as was done in Fig. 1 and Fig. 2, the 'study period' was specified in the revised caption.

[Old] Caption of Table 2: 'The number of Landsat-8 images for the four cloud mask categories over the twelve sub-regions over the Arctic Ocean during the study period.'

[New] 'The number of Landsat-8 images for the four cloud mask categories (i.e. C1: underestimated cloud cover, C2: overestimated cloud cover, C3: correctly estimated cloud cover for cloudy sky, and C4: correctly estimated cloud cover for clear sky) over the twelve regions of the Arctic Ocean during the periods of Jan. 2020 – Dec. 2022.'

Figure 4: Redefine $\rho 5$ in your caption please.

Thank you for this comment. Definitions of ρ_5 and NDSI (which was also not defined separately in the caption) are added in the revised caption.

[Old] Caption of Fig. 4: '...using p5 and NDSI criterion.'

[New] '...using ρ_5 and NDSI criterion, where ρ_5 and NDSI are the TOA reflectivity at band 5 of the OLI sensor and the Normalized Difference Snow Index, respectively.'

Figure 5: Redefine ρ 5 in your caption please.

As was done with Fig. 5, the definitions of ρ_5 and NDSI were added in the revised caption.

[Old] Caption of Fig. 5: '...derived from the selected scenes under perturbed thresholds for NDSI (red) and ρ_5 (blue).'

[New] '...derived from the selected scenes under perturbed thresholds for NDSI (red) and ρ_5 (blue), where ρ_5 and NDSI are the TOA reflectivity at band 5 of the OLI sensor and the Normalized Difference Snow Index, respectively.'

Figure 6: Please add a very brief explanation of C1, C2, C3, and C4 in your caption, as you have with Figure 5.

Thank you for this comment. Explanations of C1, C2, C3, and C4 were added in the revised caption.

[Old] Caption of Fig. 6: '... The black, red, blue, and green bars indicate values for categories C1, C2, C3, and C4, respectively.'

[New] '...The black, red, blue, and green bars indicate values for categories C1 (i.e., underestimated cloud cover), C2 (i.e., overestimated cloud cover), C3 (i.e., correctly estimated

cloud cover for cloudy sky), and C4 (i.e., correctly estimated cloud cover for clear sky), respectively.'

Figure 7: I think these plots warrant a scale bar, what do you think? I'd also prefer to see the dataset for the true colour images as a reference instead of a hyperlink to a USGS tool.

Thank you for this thoughtful comment. The authors do agree that addition of a scale bar would help with the interpretation of the figure and have added them in the revised figure. In addition, the hyperlink to a USGS tool was replaced with the reference to the dataset in the revised caption.

[Old] Caption of Figure 7: '... The true-color images were downloaded from United States Geological Survey Earth Explorer (https://earthexplorer.usgs.gov/, last access: May 22, 2024).'

[New] '... The true-color images were obtained from Earth Resources Observation and Science (EROS) Center (2020).'

The revised figures now include the scale length information as:



Figure 7: Example of (a, d, g, i) original Landsat-8 true-color image, (b, e, h, k) classification map of ice (white), open water (blue), and cloud (cyan, purple, and red), and (c, f, i, l) Landsat-8 SICs with 6.25 km resolution on (first row) Mar. 22, 2022 over the Kara Sea, (second row) Mar. 17, 2021 over the Barents Sea, (third row) Jun. 26, 2022 over the Kara Sea, and (fourth row) Jun. 15, 2022 over the Baeufort Sea. From top to bottom row, the select cases correspond to the cloud contamination categories of 1, 2, 3, and 4 respectively. SICs are not estimated over areas of cloud mask (cyan, purple and red pixels in the middle column), and SICs near the coastal area (6.25 km) are masked in this figure. The true-color images were obtained from Earth Resources Observation and Science (EROS) Center (2020).

Figure 8: Define sigmaSIC and p5 in your caption please.

Thank you for this comment. In the revised caption, definition of σ_{SIC} , ρ_5 , and NDSI were added.

[Old] Caption of Fig. 8: '(a) Uncertainties in Landsat-8 SICs and (b) contributions of the ρ_5 (blue) and the NDSI thresholds (red) to the estimated uncertainties for different SIC sub-range....'

[New] '(a) Uncertainties in Landsat-8 SICs (σ_{SIC}) and (b) contributions of the ρ_5 (blue) and the NDSI thresholds (red) to the estimated uncertainties for different SIC sub-range, where ρ_5 and NDSI are the TOA reflectivity at band 5 of the OLI sensor and the Normalized Difference Snow Index, respectively....'

Table 4: Very quick definition of CFMask please.

Thank you for this comment. Definition of CFMask is added in the revised caption.

[Old] Caption of Table 4: 'The number of cloud pixels that were undetected from CFMask...'

[New] 'The number of cloud pixels that were undetected from the C Function of Mask (CFMask)...'

Figure 9: Very nice to see error bars and the percentiles these confer to explicit. What R metric is this? Not clear if Pearson/Spearman/Coefficient of Determination or other, so make this explicit in your caption.

Thank you for the positive feedback on the error bars/percentiles. It is nice to hear that they were helpful for the interpretation of the figure. The R metric used in this figure is the Pearson correlation coefficient and in the revised caption, an explanation about the numerical values (including the Pearson correlation coefficient) that appear on the top of Fig. 9a is added.

[Old] Caption of Fig. 9: '...shown as the red vertical lines. (b) For the same SIC intervals...'

[New] '...shown as the red vertical lines. The values for number of data points (N), root-meansquare error (RMSE), bias, and Pearson correlation coefficient (R) are presented. (b) For the same SIC intervals...'

Figure 10: I'm not sure at all what this trendline represents, or if it has much physical meaning. Either add context to it in your caption or remove it.

Thank you for this suggestion. The authors agree that the trendline shown doesn't exhibit much physical meaning in the context of testing robustness of the Landsat-8 retrievals against melt ponds. In the revised figure, the trendline is removed following your suggestion.

The caption about the trend line is also removed and an explanation of the numerical values that are shown on the top-left of the figure is added in the revised caption.

The old and revised figures are found as follows:



And the corresponding captions has been changed as:

[Old] Caption of Fig. 10: '... are highlighted in red. The red line indicates the least-squares regression line.'

[New] '... are highlighted in red. The values for number of data points (N), Pearson correlation coefficient (R), and p-value for the correlation coefficient are presented.'

Figure 11: Put the USGS source as a reference. Define specifically what R we are talking about. A scale bar would be nice if it is easy to do, but not essential. Define BT and NT.

Thank you for this comment. In the same context as Fig. 7, the authors agree that a scale bar would indeed be helpful for the interpretation of the figure. In the revised figure, scale bars have been added.

As for the captions, similar to what was done for Fig. 7, the USGS source was added as a reference along with the reference for the passive microwave SIC datasets. Definitions of BT, NT, and the numerical values are also added in the revised caption.

[Old] Caption of Fig. 11: 'Geographical distributions of (a, i) original Landsat-8 true-color image, (e, m) Landsat-8 SIC, (b, j) SIC from BT algorithm, (f, n) SIC from NT algorithm, (c, k) difference in SICs between BT and Landsat-8, (g, o) difference in SICs between NT and Landsat-8 and scatterplot (d, l) between Landsat-8 SIC and SIC from BT and (h, p) between Landsat-8 SIC and SIC from NT. Upper two panels for July 21, 2022 (melting season) over the Laptev Sea and for March 4, 2020 over the Chukchi Sea, respectively. The true-color images

were downloaded from United States Geological Survey Earth Explorer (https://earthexplorer.usgs.gov/, last access: May 22, 2024).'

[New] 'Geographical distributions of (a, i) original Landsat-8 true-color image, (e, m) Landsat-8 SIC, (b, j) SIC from the Bootstrap (BT) algorithm, (f, n) SIC from the NASA Team (NT) algorithm, (c, k) difference in SICs between BT and Landsat-8, (g, o) difference in SICs between NT and Landsat-8 and scatterplot (d, l) between Landsat-8 SIC and SIC from BT and (h, p) between Landsat-8 SIC and SIC from NT. The values of root-mean-square error (RMSE), bias, and Pearson correlation coefficient (R) are presented with the scatter plots. Upper two panels for July 21, 2022 (melting season) over the Laptev Sea and for March 4, 2020 over the Chukchi Sea, respectively. The true-color images were obtained from Earth Resources Observation and Science (EROS) Center (2020) and the SIC retrievals from the BT and NT algorithms were obtained from Meier et al. (2021).'

The revised figure can be seen as:



Figure 11: Geographical distributions of (a, i) original Landsat-8 true-color image, (e, m) Landsat-8 SIC, (b, j) SIC from the Bootstrap (BT) algorithm, (f, n) SIC from the NASA Team (NT) algorithm, (c, k) difference in SICs between BT and Landsat-8, (g, o) difference in SICs between NT and Landsat-8 and scatterplot (d, l) between Landsat-8 SIC and SIC from BT and (h, p) between Landsat-8 SIC and SIC from NT. The values of root-mean-square error (RMSE), bias, and Pearson correlation coefficient (R) are presented with the scatter plots. Upper two panels for July 21, 2022 (melting season) over the Laptev Sea and for March 4, 2020 over the Chukchi Sea, respectively. The true-color images were obtained from Earth Resources Observation and Science (EROS) Center (2020) and the SIC retrievals from the BT and NT algorithms were obtained from Meier et al. (2021).

Figure 12: What are the units for (b)? Add a definition for 'in' and 'out' in (d).

Thank you for this catching. Since (b) is a map showing the number of Landsat-8 pixels located within each grid cell, the units for (b) should be "Number of Landsat-8 Pixels". This unit was added to the colorbar in the revised figure.

In addition, the "In" and "Out" designating grid cells located inside and outside the sub-region, respectively, were defined in the revised caption.

[Old] Caption of Fig. 12: 'Variables in the Landsat-8 SIC netCDF. The scene is from Jun. 12, 2021 over the Canadian Archipelago.'

[New] 'Variables in the Landsat-8 SIC netCDF. The scene is from Jun. 12, 2021 over the Canadian Archipelago. For the (d) sub-region mask, 'In' and 'Out' denote grid cells located inside and outside the designated region, respectively.'



The revised figure can be seen as:

Archipelago. For the (d) sub-region mask, 'In' and 'Out' denote grid cells located inside and outside the designated region, respectively.