

CLIMK–WINDS: A New Database of Extreme European Winter Windstorms

Clare Marie Flynn^{1, 2}, Julia Moemken³, Joaquim G. Pinto³, Michael Schutte¹, and Gabriele Messori^{1, 2, 4}

¹Department of Earth Sciences, Uppsala University, Uppsala, Sweden

²Swedish Centre for Impacts of Climate Extremes (climes), Uppsala University, Uppsala, Sweden

³Institute of Meteorology and Climate Research Troposphere Research, Karlsruhe Institute of Technology (KIT), Karlsruhe, Germany

⁴Department of Meteorology and Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden

Correspondence: Gabriele Messori (gabriele.messori@geo.uu.se)

Abstract. ~~European windstorms pose a significant threat to people, infrastructure and the natural environment. Several windstorms in the recent past have caused substantial damages, and losses associated with extreme windstorms may increase with climate change. Characterizing the footprints of destructive windstorms is thus key to providing quantitative estimates of storm-related economic losses. To that end, we have developed~~ The CLIMK–WINDS (CLimes IMK – WINDStorm) database is a new, publicly available database of extreme European windstorm footprints for the extended winter season during 1995–2015. In contrast to previously compiled European windstorm databases, ~~we include~~ it includes storm footprints derived from four different data sets, rather than a single source: the ERA5 reanalysis, the COSMO-REA6 reanalysis for Europe, the COSMO-Climate Limited-area Mode regional climate model driven by ERA5 on the EURO-CORDEX domain, and simulation output from the same model but on an enlarged Germany domain with higher horizontal resolution. The database includes ~~both~~ the footprints themselves, expressed as the relative daily maximum wind gusts associated with a storm event, ~~and~~ the daily maximum wind gusts in absolute magnitude associated with the footprints. ~~We derived and included the storm footprints associated with the 50 most extreme storms, or Top50 storms, identified within each of the four input data sets, and a measure of storm severity.~~ We applied a consistent methodology, the storm loss index, across input data sets for identifying storm footprints and assessing their severity. We identified and included the storm footprints associated with the 50 most severe storms, or Top50 storms, within each of the four input data sets. This enables a direct comparison between the footprints derived from the different input ~~sources~~ data sets, eases future efforts to extend the time record of the database or to include additional input data sets, and enables assessment of uncertainty in the footprints. Moreover, since we derived the Top50 storms from each input data set on its native horizontal resolution, the database also allows to characterize the impact that horizontal resolution can have on footprint identification and severity assessment. ~~Our database~~ We find that the choice of input data set – including the data’s horizontal resolution – can have major effects on extreme storm identification and characterization. Different storms were identified as belonging to the Top50 storms in the different data sets, and storm footprints for common storms displayed substantial variability across the data sets. A comparison of our database to two existing windstorm databases also highlights the important role of the footprint detection methodology. The CLIMK–WINDS database thus supports both the research com-

25 munity and the insurance industry in exploring the data set, [methodology](#) and resolution dependence of assessments of extreme storm hazards.

1 Introduction

Winter windstorms constitute the most costly natural hazards for Europe, posing a significant threat to people, infrastructure and the natural environment (Mitchell-Wallace et al., 2017; Priestley et al., 2018; Pinto et al., 2019; Walz and Leckebusch, 2019; Munich Re, 2022; Moemken et al., 2024b). Average annual windstorm insured losses in Europe are on the order of several billion USD, with total losses estimated to be well in excess of this. Windstorms also lead to important non-monetary losses such as casualties (Schwierz et al., 2010; Priestley et al., 2018; Gliksman et al., 2023). Windstorm damage scales non-linearly with storm ~~intensity~~[severity](#), and single storms of unusual severity can cause losses which exceed the long-term annual average. For example windstorm KYRILL, the most severe storm of the 2006/2007 winter season and one of the most severe within the past three decades, struck Europe in January 2007 (Fink et al., 2009; Priestley et al., 2018) and alone caused 5.8 billion USD in insured losses. It further led to 54 fatalities, significant disruptions in transportation, and electricity outages, among other impacts (Deutsche Rück, 2008; Munich Re, 2015). KYRILL was also the strongest storm within a series of intense windstorms that struck Europe during the 2006/2007 [winter](#) season, which ~~caused~~[saw](#) an estimated 10 billion USD in cumulative insured losses ~~for this season~~ as well as ~~additional~~[high numbers of](#) injuries and deaths (Pinto et al., 2014). Indeed, serial clustering of European winter windstorms, namely when multiple storms follow a similar track in quick succession (Dacre and Pinto, 2020), can magnify losses relative to individual storm events. Moreover, more severe storms are more likely to cluster than less severe storms (Pinto et al., 2014; Priestley et al., 2018). Though uncertainties remain in predictions of changes in windstorms with climate change, climate model evidence suggests an increase in both the frequency and strength of storms over northern and central Europe, particularly for more extreme windstorms, with a corresponding increase in storm losses (Pinto et al., 2007; Leckebusch et al., 2007; Schwierz et al., 2010; Pinto et al., 2012; Little et al., 2023; Severino et al., 2024). This occurs in spite of ~~the well known~~[a](#) general decrease in the total number of extratropical cyclones over the region (Ulbrich et al., 2009; Priestley and Catto, 2022). Characterizing and understanding extreme European winter windstorm losses is therefore a highly socioeconomically relevant goal.

To address this need, several publicly available and subscription-based storm loss databases have been created and maintained. The publicly available loss databases are often based on a storm severity index that incorporates meteorological indicators and empirical approximations for insured losses. A widely used index is the storm loss model based on Klawa and Ulbrich (2003), in which the predicted windstorm damage is proportional to the cube of the exceedance of the wind speed over a relative threshold value and to the population density (Klawa and Ulbrich, 2003; Pinto et al., 2012; Gliksman et al., 2023; Moemken et al., 2024b). Other approaches are also employed, such as statistical downscaling of wind data (van den Brink, 2019). Insurance and reinsurance companies also simulate storm losses ~~with catastrophe models~~. ~~Index-based loss estimates and industry-computed insured losses based on catastrophe modelling, however, both rely on using catastrophe models~~ (Moemken et al., 2024a). However, the loss estimates based on storm severity indices or on catastrophe models, both

require accurate wind speed or wind gust data with ~~sufficient~~-high spatial and temporal coverage. Windstorm damage is typically assumed to arise from the strongest wind speeds or gusts ~~encountered~~ during the storm, such that estimates of storm-related losses rely on identifying the locations impacted by a given storm as well as the peak winds as accurately as possible (Klawa and Ulbrich, 2003; Leekebusch et al., 2007; Pinto et al., 2012, 2014; Priestley et al., 2018; Cusack, 2023; Gliksman et al., 2023) (Klawa and Ulbrich, 2003; Leekebusch et al., 2007; Pinto et al., 2012, 2014; Priestley et al., 2018; Cusack, 2023; Gliksman et al., 2023; M

. The occurrence exceedance probability (OEP) and the annual exceedance probability (AEP), two metrics based on loss estimates that are commonly used in the insurance industry, ~~further~~-also rely on accurate wind data. The OEP and AEP represent the maximum loss event during a season and the total losses summed over all events during a season, respectively, and are used to assess return periods of extreme storms and the impact of storm clustering on losses (Priestley et al., 2018). The only loss data that are not sensitive to the quality of the meteorological data are industry reports of recorded insured losses, such as those collected in the PERILS database (<http://www.perils.org>). PERILS is based on data gathered by insurance and reinsurance companies, but only provides information on an annual subscription basis and for a subset of European countries (Gliksman et al., 2023; Moemken et al., 2024b).

Wind data of a sufficient quality are therefore crucial to producing reliable, openly accessible and comprehensive storm loss estimates and for the assessment of future risks. This wind information is often provided as a storm footprint, essentially a map of the peak winds and their magnitudes encountered during a storm event at locations affected by the storm. These are typically determined through use of a wind exceedance threshold such as that used in storm severity indices (Klawa and Ulbrich, 2003; Pinto et al., 2012, 2014; Priestley et al., 2018). Two publicly-available databases which provide storm footprints and the corresponding storm loss estimates at the time of writing are the eXtreme Wind Storms Catalogue (XWS; Roberts et al. (2014)) and the Copernicus Climate Change Service (C3S; C3S Climate Data Store (2022); van den Brink (2019)) ~~databases~~database. These contain the most ~~extreme~~-severe European winter windstorms over the past ~~approximately 40 years~~ decades and have proven useful in the study of extreme windstorm impacts; ~~more detail~~. Further details on these databases ~~is given~~-are provided in Section 2. However, the XWS and C3S databases are each derived from a single input data set; ~~;~~

the ERA-Interim (~~Dee et al., 2011~~)-reanalysis ~~reanalysis~~ (Dee et al., 2011) in the case of XWS and ERA5 (Hersbach et al., 2020) for C3S. While ERA5 is the current state-of-the-art reanalysis data set, there are indications that storm footprints based on ERA5 peak near-surface wind gusts may contain inaccuracies (Cusack, 2023), while ERA-Interim is known to have deficiencies in representing many storms during the later 20th and early 21st centuries (Moemken et al., 2024b). Furthermore, a downscaling approach involving the use of an atmospheric or statistical model was employed to derive the footprints and estimated losses for both databases, requiring intensive computational resources for database extension. Finally, different definitions were used in each database for the spatial extent of the footprint (Roberts et al., 2014; van den Brink, 2019), making the two not comparable, and thus hindering assessment of the uncertainty associated with footprint computation. Thus, while XWS and C3S constitute invaluable tools in the study of extreme windstorms, a need remains for a storm footprint database based on multiple meteorological data sets which are processed with a standardised methodology.

~~One of the identified challenges in terms of windstorm research is to have~~ Having a consistent, reliable and extendable database for historical windstorm risk is a key need for windstorm research (Pinto et al., 2019). In this manuscript, we present a

~~new database of the new CLIMK-WINDS (CLimes IMK – WINDStorm) database, which collects~~ the 50 most ~~extreme-severe~~ European winter windstorms derived from four different input data sets with four different native horizontal resolutions using a standardised methodology. The meteorological data comes from two reanalysis data sets, ERA5 (Hersbach et al., 2020) and COSMO-REA6 (Bollmeyer et al., 2015), and output from two regional climate model simulations, CCLM_ERA5_EUR-11 and CCLM_ERA5_CEU-3, as further described in Section 2. This database therefore complements the existing storm databases and expands the footprint data available for use by the scientific community and industry. Its design facilitates an assessment of the impact of different data sources and different horizontal resolution on identification of extreme storms and their ~~estimated losses~~~~severity – which we estimate here using an empirical storm loss model based on wind speed data.~~ Equally importantly, ~~an estimate of it also enables estimating~~ the uncertainty associated with the footprint itself~~could also be assessed given the diversity of input data sets used in our database, as similarly enabled by a new database for storm loss estimates from multiple perspectives (Moemken et al., 2024b).~~ This database ~~CLIMK-WINDS~~ may therefore be used by the scientific community and insurance industry to address the crucial question of ~~what "quality" of how differences in~~ wind gust data ~~is required for accurate and reliable~~ ~~affect the~~ characterization of extreme windstorms and their impacts, supporting ~~the need for better an~~ ~~uncertainty-aware~~ analysis of storm damage to ~~reduce their impact on society~~~~mitigate societal impacts.~~

Throughout the rest of this study, we use the terms "windstorm" and "storm" interchangeably to refer to the extreme storms ~~identified in each data source~~ that affected Europe during the extended winter season. Storm names, when they are given names rather than dates, are capitalised.

2 Methods

110 2.1 Overview of the database

~~Our European windstorm database was derived from~~ ~~CLIMK-WINDS builds on~~ four different input data sets with different horizontal resolutions ~~– (Sect. 2.2).~~ It consists of the windstorm footprints as represented by the relative daily maximum wind gusts during the storm event (unitless), their associated daily maximum wind gusts in absolute units (ms^{-1}), the loss index ($\text{LI}_{3\text{D}}$; unitless) integrated over a Core Europe region (42°N – 60°N , 10°W – 15°E), and the unintegrated loss index at each ~~grid point (unitless). The unintegrated loss index enables users to compute an integrated loss over a subset region of a given storm's footprint, such as at the country level. These quantities are described in Sect. 2.3.~~ We further provide the name and dates of occurrence of each windstorm, their ordinal rank, and their relative rank (also termed the normalized loss, ~~see Sect. 2.4).~~ Because the input data sets are on different native horizontal grid types and resolutions, we maintained the native grid information and created a separate database netCDF file for each input source rather than merging all into one file.

120 The database ~~is based on daily wind gust maxima derived from hourly wind gust data, and~~ covers extended winter seasons (October to March, ONDJFM) from January 1995 to December 2015, ~~a 20-year period covered by all 2015. This timespan was chosen to reflect the period common to both the four~~ input data sets ~~, and is based on daily wind gust maxima derived from hourly wind gust data used in CLIMK-WINDS and the two existing databases that we use for comparison, and to additionally include the year 2015 during which several storms common to all four input datasets were identified.~~ The fifty most ~~extreme~~

125 [severe](#) storms, or Top50 storms, were identified for each input data set based on the Core Europe integrated loss index and included in the database. The individual storms that make up the Top50 storms for each input [source data set](#) therefore differ, and in some cases the same storm may have somewhat different dates of occurrence across the input data sets.

2.2 Input Data Sets

Table 1 summarizes the input data sets used to create ~~the new database presented here~~[CLIMK-WINDS](#). These consist of
130 two reanalysis data sets, ERA5 (Hersbach et al., 2020) and COSMO-REA6 (Bollmeyer et al., 2015), and output from two regional climate model simulations on different domains, the CCLM_ERA5_EUR-11 and CCLM_ERA5_CEU-3 simulations, performed with the COSMO model ~~in climate mode~~(Rockel et al., 2008). These data sets differ in horizontal resolution, spatial domain sizes and types of horizontal grids, such as a regular latitude-longitude or curvilinear grid ([Figure 1](#)).

The publicly available ERA5 reanalysis ~~data set~~ from the European Centre for Medium-Range Weather Forecasts (ECMWF)
135 provides global data with a 0.25° horizontal resolution and with many parameters available ~~on an~~ [at](#) hourly temporal resolution (Hersbach et al., 2020). We analyse ERA5 storm footprints over $27-72^\circ$ N and 22° W- 45° E, [to approximately match the CORDEX EUR-11 domain \(Jacob et al., 2014, 2020\)](#). The reanalysis is based on the Integrated Forecasting System (IFS) Cy41r2 model (Hersbach et al., 2020), which was operational at ECMWF in 2016 (Bonavita et al., 2016). Data span 1940 to near-present. ERA5 is often used for evaluation and bench-marking of other data sets, such as model simulation output, due to
140 its global coverage, high temporal resolution, and observationally constrained, physically consistent atmospheric data.

The Hans-Ertel-Centre for Weather Research (HErZ) and the Deutscher Wetterdienst (DWD; German Weather Service) developed the COSMO-REA6 (Consortium for Small-scale Modelling-Reanalysis, 6 km) high-resolution regional reanalysis data set for Europe, with a 0.055° (approximately 6 km) horizontal resolution and 15-minute or hourly temporal resolutions (Bollmeyer et al., 2015). COSMO-REA6 is available over [January](#) 1995-August 2019, since it uses ERA-Interim reanalysis data
145 (Dee et al., 2011) for lateral boundary conditions and the latter is no longer produced. The spatial domain matches that of the CORDEX EUR-11 domain (Jacob et al., 2014, 2020), which covers approximately $27-72^\circ$ N and 22° W- 45° E. Unlike ERA5, COSMO-REA6 is based on the COSMO (Consortium for Small-scale Modelling) model in numerical weather prediction mode, developed by the DWD (Baldauf et al., 2011). It uses nudging of surface synoptic conditions, aircraft measurements, radiosondes, buoys, ship reports, and wind profilers ~~as in~~ [its](#) data assimilation scheme (Bollmeyer et al., 2015).

150 The CCLM_ERA5_CEU-3 simulations were run with the non-hydrostatic COSMO model [version 5.0](#) in climate mode (COSMO-Climate Limited-area Mode or COSMO-CLM) ~~regional climate model, version 5.0 (Rockel et al., 2008); (Rockel et al., 2008)~~ [. These were performed](#) by the DWD in collaboration with the Climate Limited-area Community (CLM-Community) collaborative network. COSMO-CLM is the climate version of the limited-area numerical weather prediction COSMO model used to produce COSMO-REA6, convection-permitting, and driven by ERA5 through direct downscaling. CCLM_ERA5_CEU-3
155 spans an enlarged Germany or COSMO-DE domain (approximately $45-58^\circ$ N and $1-20^\circ$ E) at 0.0275° (approximately 2.8 km) horizontal resolution (Baldauf et al., 2011; Brien et al., 2022). Output is available for the years 1979-2019 up to hourly resolution (<https://esgf.dwd.de/search/esgf-dwd/>). Output does not extend beyond 2019 as the DWD discontinued use of the COSMO-CLM model. The CCLM_ERA5_CEU-3 simulation output used to create the extreme storms database presented here

therefore ~~represents~~presents the highest horizontal resolution, but smallest spatial domain, ~~input data source of~~amongst the
160 four input data sets that we used.

Lastly, the CCLM_ERA5_EUR-11 simulations were performed by the Helmholtz Center Hereon in collaboration with the CLM-Community and EURO-CORDEX. The COSMO-CLM model in the same configuration as for the CCLM_ERA5_CEU-3 simulations was used, but on the CORDEX EUR-11 domain at a horizontal resolution of 0.11° (approximately 12 km). The European branch (EURO-CORDEX; Jacob et al. (2014, 2020)) of the Coordinated Regional Downscaling Experiment
165 (CORDEX; Giorgi and Gutowski (2015)), is a collaborative initiative that seeks to advance regional climate and Earth system science in Europe. It defined the CORDEX EUR-11 domain in 2013, intended to be used for simulations with a horizontal resolution of 0.11° . Output is available for the years 1979-2020 at up to hourly resolution.

2.3 Windstorm Footprint Identification

To identify the windstorm footprints within each input data set and to quantify their severity, we employed the storm loss model
170 known as ~~the~~-loss index (LI). This was developed by Pinto et al. (2012) and extended by Karremann et al. (2014), and is in turn based on the storm loss models developed by Klawns and Ulbrich (2003) and Leckebusch et al. (2007). The LI was originally developed based on near-surface daily maximum wind speed (Pinto et al., 2007), but we exchanged this for near-surface daily maximum wind gusts derived from hourly data. This is an unproblematic exchange, as wind speeds are often used as a proxy for wind gusts when gust data ~~were~~are unavailable, and maintains consistency across calculations and comparisons.

175 The LI is based on the assumption that storm damage occurs only for the highest 2% of local wind speeds, or wind gusts in our case (Klawns and Ulbrich, 2003; Pinto et al., 2012; Karremann et al., 2014). The footprint of an individual storm is therefore only those locations whose local daily maximum wind gust on the date of the storm exceeds the local 98th percentile of wind gusts~~computed over the full~~. We define this here using the 20-year ~~record~~. The period common to all data sets used in our analysis (including XWS and C3S, see Sect. 2.5), namely 1995 – 2014. The 98th percentile of wind gusts for each of the four
180 input data sets is shown in Figure 1. The storm footprint is expressed in our database in terms of the unitless relative wind gust; the absolute wind gusts associated with the footprint are also ~~given in our database~~provided. The relative wind gust is the ratio of the local daily maximum wind gust that occurred during a given storm to the local 98th percentile, which indicates the magnitude of the exceedance over the 98th percentile and is everywhere greater than 1.0 within the storm footprint (Pinto et al., 2007, 2012).

185 Individual storms must then be separated from each other in time and their severity assessed. This is accomplished through calculation of the LI itself, which relies on two additional assumptions. First, potential storm losses are assumed to increase with the cube of the maximum wind speed or gust, as this is proportional to wind ~~kinetic energy~~power. Second, storm losses are linked to the exposure/insured value, which can be approximated by population density (Pinto et al., 2012; Karremann et al., 2014). We used the Gridded Population of the World, version 4 (GPWv4) population density data set for the year 2020,
190 the latest year available, provided by the Center for International Earth Science Information Network (CIESIN) at Columbia University (CIESIN, 2018). These data have a horizontal resolution of 0.04° , and were regridded to the resolutions of the four input data sets before use (Figure 1 in the supplementary material). The LI was computed for each day, ~~just as the daily~~

~~maximum wind gust~~, and is defined as follows:

$$LI = \sum_{ij} \left[\left(\frac{v_{ij}}{v_{98_{ij}}} \right)^3 * POP_{ij} * I(v_{ij}, v_{98_{ij}}) \right] \quad (1)$$

195 where

$$I(v_{ij}, v_{98_{ij}}) = \begin{cases} 0, & \text{for } v_{ij} \leq v_{98_{ij}} \\ 1, & \text{for } v_{ij} > v_{98_{ij}} \end{cases}$$

and indicates whether or not the daily maximum wind gust at grid point ij falls within the storm footprint. POP_{ij} is the population density at grid point ij , and v_{ij} and $v_{98_{ij}}$ are the daily maximum wind gust and the 98th percentile wind gust at grid point ij , respectively (Pinto et al., 2012; Karremann et al., 2014). Differences in LI among the four data sets arise mostly in densely populated regions due to the respective horizontal resolution of the regridded population density data (Figure 2 in the supplementary material). An overlapping three-day sliding time window is applied to the LI in order to separate individual storms in time, and the temporal local maximum of each three-day window is assumed to be the individual storm event (Karremann et al., 2014):

$$LI_{3D} = \sum_{ij} \left[\left(\frac{max_{3D} v_{ij}}{v_{98_{ij}}} \right)^3 * POP_{ij} * I(v_{ij}, v_{98_{ij}}) \right] \quad (2)$$

205 The LI_{3D} provides the final, temporally separated storm footprints for individual storms. It should be noted, however, that if storms occur too closely together in time, such as storms LOTHAR and MARTIN in December 1999, it can be very difficult or impossible to separate the storms with the LI_{3D} ; ~~this~~. This seldom happened in the creation of our database, but when it did occur, the storms were typically counted as one "combined" storm in the database. The LI_{3D} can be integrated over any spatial domain of interest; to create our database, we integrated over the Core Europe region defined above, following Pinto et al. (2012). ~~It should be further noted that the~~ The unintegrated loss index included in the database refers to the summand inside the Σ operator in Equation 2 that is computed at each grid point before any integration is performed, and not to the LI_{3D} . This quantity is thus akin to a pre- LI_{3D} . The unintegrated loss index enables users to compute an integrated loss over a subset region of a given storm's footprint, such as at the country level. This should be regarded as a loss proxy, since the LI_{3D} builds upon an empirical storm loss model.

215 An example of a storm footprint as represented by the relative wind gusts is shown in Figure 2 for storm KYRILL for each of the four input data sets, as KYRILL was identified as the most ~~extreme~~ severe storm in all four ~~input sources~~. Figure 3 displays the absolute daily maximum wind gusts associated with the footprints for each of the four input sources as well as for XWS and C3S. Figures 2 and 3 clearly highlight the differences among the footprints for the same storm due to differences in input data sets ~~;~~ and, in the case of Figure ~~23~~, storm footprint definition.

220 2.4 Selection of Extreme Windstorms

We selected the 50 most ~~extreme~~ severe (Top50) storms over the 1995-2015 period derived from each input data set based on the ~~magnitude~~ 50 largest magnitudes of the LI_{3D} integrated over Core Europe. The ~~magnitudes were ranked from largest to smallest,~~

and the fifty largest magnitudes were chosen. The selected storms were manually checked to ensure that they represented either unique individual storms, or, in the case of storms that could not be sufficiently separated, a unique "combined" storm. The
225 dates of the Top50 storm occurrence for each input data set in the database are the mid-points of the three-day time windows
containing the fifty largest LI_{3D} magnitudes ~~were associated with unique dates per dataset, and the dates of the mid-point of
each window became the storm dates of occurrence for that dataset in the database.~~

Based on these dates, storm names were assigned to the individual storm events. Storm names were taken from several
sources, including the lists of named storms produced by the DWD and the Freie Universität Berlin for the years 1999-2015
230 (<https://www.wetterpate.de/namenslisten/tiefdruckgebiete/index.html>; in German), the past European winter windstorm docu-
mentation provided by Deutsche Rück for the years 1997-2015
(<https://www.deutscherueck.de/downloads>, in German), and from Wikipedia articles about European winter windstorms (pri-
marily English articles). Some storms, ~~primarily~~ chiefly those occurring earlier than 1997, appear to lack names given by a
meteorological service or research institution; ~~in~~ In these cases, we have taken the storm date of occurrence as the storm name
235 preceded by the lower case letter "u," and we have preferred the date as identified from ERA5 for the name, if the storm was
identified within ERA5 and at least one other data set. As mentioned above, some storms could not be sufficiently separated
with LI_{3D} in some input data sets, and were thus taken as a single, "combined" storm and the names of the individual storms
that could not be well separated were hyphenated to create the storm name. Some individual storms had two different names
given by different entities, ~~so these~~ These storms retained both names, one "main" name and a "secondary" name given in
240 parentheses (or connected by an underscore rather than parentheses within the database files for ease of coding). Lastly, two
different storms were given the same name, "Franz," and so we have identified one storm as FRANZ (2007-01-11) and the other
as FRANZ-II (1999-12-12); the hyphenated name in this one case does not indicate two storms that could not be ~~sufficiently~~
well separated.

2.5 Existing Windstorm Databases for Comparison

245 The XWS (Roberts et al., 2014) and C3S (C3S Climate Data Store, 2022; van den Brink, 2019) storm databases were used as
~~comparison data sets for the new database we describe here;~~ terms of comparison for CLIMK-WINDS. These characteristics
of these databases are also summarized in Table 1.

The publicly-available XWS database (<http://www.europeanwindstorms.org>) includes 50 of the most ~~extreme~~ severe Eu-
ropean storms within the extended winter season for the years 1979-2014, and includes storm footprints and storm severity
250 indices or loss estimates; ~~this database will not be extended.~~ There are currently no plans to extend this database beyond
2014. The footprints were computed through dynamically downscaling the ~~previous generation~~ ERA-Interim reanalysis data
set (Dee et al., 2011) to a horizontal resolution of 0.22° (approximately 25 km) with the UK Met Office Unified Model (Me-
tUM; Davies et al. (2005)) over a domain including western Europe and the eastern North Atlantic. The footprints are defined
as the maximum 3-second wind gusts at each grid point in the downscaled domain over a 72-hour period ~~;~~ but, rather than
255 taking the maximum gusts over the entire domain, all wind gusts outside of a 1000 km radius centered on the storm track are
neglected before taking the maximum. This is done to separate storms that occurred closely together in time from each other.

A meteorological severity index S_{ft} (Roberts et al., 2014) was computed for each storm as:

$$S_{ft} = (U_{max})^3 * N \quad (3)$$

where $(U_{max})^3$ is the cube of the storm's maximum near-surface wind speed as an indication of storm intensity and N is the footprint size index, defined as the number of 25 km footprint grid points over European and Scandinavian land for which the maximum wind gust exceeds 25 m s^{-1} . Though differently formulated, the S_{ft} index is based upon the winds associated with a storm, just as the LI and LI_{3D} . Of the 50 storms in the XWS database, 23 storms were included after consultation with the Willis Research Network and based on the extreme values of insured losses ~~and~~. These constitute the "insurance storms," while the remaining 27 storms were included based on the S_{ft} index. The insurance storms are provided with an insured loss amount in USD (indexed to the year 2012) in addition to the S_{ft} index. For comparison to our database, we ranked the 50 XWS storms based on their S_{ft} magnitudes. The XWS data were used directly and required no further processing, as this database has already restricted their footprints to the area affected by each storm.

The publicly-available C3S database (~~<https://eds.climate.copernicus.eu/>~~) (C3S Climate Data Store, 2022) was derived from ERA5 and includes significant winter storms from the years 1979-2021 for 21 countries in western, central, and northern Europe on a horizontal resolution of 1.0 km; southeastern European countries were not included. Though this database currently extends to 2021, it may be extended further as the ERA5 reanalysis will continue to be updated. Similarly to XWS, the C3S footprint was defined as the maximum 3-second near-surface wind gust over the 72-hour period capturing the storm; ~~but~~. However, the footprints were computed through statistically ~~downscaling ERA5 data instead~~ rather than dynamically downscaling reanalysis data – which in the case of C3S is ERA5. A multiple linear regression model to derive estimates of the strongest wind gusts during a storm period following van den Brink (2019) was developed and validated; ~~and is based on~~; the ERA5 wind gust data, wind gusts estimated from the wind speed ~~(also taken from ERA5)~~ shear between the 10 m and 100 m altitude levels ~~and~~ (also taken from ERA5), and weather observation station elevation height ~~to derive estimates of the strongest wind gusts during a storm period~~ are used as predictors. This method is valid only for land areas ~~and allowed the C3S database to estimate the strongest wind gusts for locations far from or in between weather observation stations~~. A total of 148 storm footprints are included in ~~this~~ the C3S database. However, the strongest wind gusts associated with a storm were estimated over all land areas within the full C3S domain, covering $35-70^\circ \text{ N}$ and $20^\circ \text{ W}-35^\circ \text{ E}$. Thus, the C3S data does not a priori distinguish between those grid points that are and are not impacted by a given storm. No cut-off criterion, such as the 1000 km radius centered on the storm track used in the XWS database, was applied. Though the footprint is often apparent as the area ~~crossing across~~ Europe with the ~~largest-strongest~~ wind gusts, this leaves ambiguity over precisely where the footprint begins and ends. Because the C3S database is derived from ERA5, we used the footprints we identified from the ERA5 data set in the course of creating CLIMK-WINDS as a spatial mask to "cut out" the C3S wind gusts belonging to a given storm's footprint, after first regridding the ERA5 footprints to the C3S horizontal grid. ~~We~~ Following this procedure, we then computed the LI_{3D} for each C3S storm ~~footprint using the regridded ERA5 footprints~~ to assess storm severity; ~~in order to maintain consistency between the footprint area and severity index.~~

290 ~~We selected only those storms from~~ Since the XWS and C3S databases are used as terms of comparison, we selected only
~~those storms~~ from the 1995-2015 period (1995-2014 for XWS) that were also found in at least one of the four input data sets
listed above, ~~as these databases were used for comparison only, leading to~~. This resulted in 23 extreme windstorms for XWS
and 30 for C3S. ~~It should be noted that these databases did~~ The XWS and C3S databases do not provide relative wind gust
295 only for the absolute wind gusts associated with the footprints.

2.6 Ordinal and Relative Ranking of Windstorms and Risk Metrics

In order to compare storm severity among the storms identified within our database, the Top50 storms were each assigned
an ordinal rank and a relative rank, also called the normalized loss index or simply normalized loss. The ordinal ranking is
straightforwardly based on the magnitude of the LI_{3D} , where the storm with the largest LI_{3D} was assigned rank 1, and the storm
300 with the smallest LI_{3D} was assigned rank 50. Ordinal ranks for the same storm identified in two or more input data sets can
differ.

The relative ~~ranks, or normalized losses as it will be referred~~ rank, which we also refer to in the following sections as relative
severity or normalized loss, is based on min-max scaling of the LI_{3D} :

$$normalizedloss_i = \frac{LI_{3D_i} - \min(LI_{3D})}{\max(LI_{3D}) - \min(LI_{3D})} \quad (4)$$

305 where i refers to the value for an individual storm. The normalized loss ranges from 0.0 for the storm with the smallest impact
(the storm with ordinal rank 50) to 1.0 for the storm with the most impact (the storm with ordinal rank 1); ~~the normalized~~
~~losses and the ordinal ranks vary inversely and monotonically with each other~~. The normalized loss thus expresses the severity
of each individual storm as relative to the severity of the most ~~extreme~~ severe storm (that with both ordinal and relative rank of
1) and provides an indication of how different the storms are from each other. The normalized losses and the ordinal ranks vary
310 inversely and monotonically with each other. These ordinal and relative ranks are included in the netCDF files that constitute
our extreme storms database. The ordinal ranks and normalized losses were also computed for the XWS and C3S storms ~~we~~
~~use for comparison below~~, though relative to a total storm number of 23 for XWS and 30 for C3S rather than ~~50~~. ~~We also~~ 50
(Sect. 2.5). We based the ranks and normalized losses on the S_R index for XWS, and on the LI_{3D} magnitudes corresponding to
the ERA5 footprints used to "cut out" the footprints for C3S.

315

In addition to ordinal ranks and normalised losses, we also compute the OEP and AEP risk metrics, namely the maximum
loss event during a season and the total losses summed over all events during a season. Their ratio informs on whether losses
during a season were dominated by a single, exceptionally damaging storm or by several less damaging storms. Here, we
compute the OEP and AEP using LI_{3D} , rather than from actual loss data.

3.1 Overview of the Identified Extreme Storms

Our database identified 76 unique storms within the Top50 storms across the four input data sets. Of these storms, 29 storms (approximately 38% of the unique storms) were identified as Top50 storms within all four input sources, constituting the common storms listed in Table 2. Table 2 also indicates whether these storms are found in the XWS and C3S databases.

325 The ~~supplemental-supplementary~~ file Top50Storms_All_Summary.csv summarizes all the Top50 storms ~~rather than only the common storms~~ identified within each of the input data sets and whether they are found in XWS and C3S, along with their ordinal ranks in each data set.

Despite occurring closely together in time, only storm LOTHAR belongs to the common storms while MARTIN does not, as it was not identified within the COSMO-REA6 and CCLM_ERA5_CEU-3 data sets; ~~this~~. This is likely because
 330 MARTIN could not be sufficiently separated from the first-occurring LOTHAR within the COSMO-REA6 data set, while MARTIN's more southerly storm track fell mostly outside the domain of CCLM_ERA5_CEU-3. The date of occurrence for MIKE-NIKLAS was identified as 2015-04-01 in CCLM_ERA5_EUR-11. Though technically outside the temporal domain we considered for the database, we decided to keep this storm for this input ~~source~~ data set as MIKE-NIKLAS was identified within the remaining input sources as occurring within the extended winter season we defined.

335 ~~This leaves~~ The remaining 47 storms (approximately 62% of the unique storms) ~~that were identified~~ were identified as Top50 in at least one of the input data sets but not all four (listed in the ~~supplemental-supplementary~~ file Top50Storms_All_Summary.csv). Of these 47 storms, ~~15 were identified~~ 18 were identified as Top50 in only one input source, ~~whereas the remaining 28 storms were identified in two or three input sources~~. Four storms were identified in ERA5 alone (BECKY, the BOXING DAY STORM, FRIEDHELM, and JETTE); six were identified in CCLM_ERA5_EUR-11 alone (DAGMAR, EBERHARD, JULIA,
 340 u19961106, u19961120, and u20000209) and in CCLM_ERA5_CEU-3 alone (DORIAN, ELIZABETH, EX-HURRICANE GONZALO, INGO, and QUINTEN); and two were identified in COSMO-REA6 alone (SUSANN and ORKUN). It is unexpected that CCLM_ERA5_CEU-3 rather than COSMO-REA6 displayed a similar number of storms identified only in that input data set as ERA5 and CCLM_ERA5_EUR-11, given its much smaller spatial domain. The remaining 29 storms within our database were identified in two or three of the input data sets, and no systematic pattern appears to exist in which combinations
 345 of input sources are preferred.

All common storms, with the exception of four storms, were identified as occurring on the same date across the four input data sets. The remaining four common storms displayed a discrepancy in date of occurrence of one day (Table 2). When considering all Top50 storms identified in at least two input data sets, approximately 74% of storms displayed no discrepancy in date of occurrence and approximately 26% of storms displayed a discrepancy of one day. One of the reasons for these
 350 discrepancies is the smaller spatial domain of CCLM_ERA5_CEU-3 compared to the others.

Only 12 of the common storms were also found in the suite of XWS storms examined here (approximately 52% of all the XWS storms used here), and 16 of these storms (or approximately 53% of all C3S storms used here) for C3S. When a common storm was also found within both the XWS and C3S databases, the dates of occurrence between CLIMK-WINDS, XWS

and C3S agreed with each other for all storms except ~~EMMA and XYNTHIA~~ for ANDREA (ULLI), EMMA, for which they
355 displayed a difference of one day GISELA-HEIDI and XYNTHIA. In the case of EMMA, XWS and C3S also disagreed with
each other. These discrepancies were of 1 or 2 days (Table 2). Relative to our database, XWS and C3S ~~displayed thus~~ most
often displayed no discrepancy in the date of occurrence for ~~both the common storms and for all.~~ The same holds for the other
Top50 ~~storms. The differences in dates of occurrence display no preference for before or after the dates in our database storms.~~

The ~~storms that are not common to all four data inputs~~ four input data sets in our database thus demonstrate some disagree-
360 ment on which storms are identified as belonging to the Top50, and ~~mild disagreements in the dates the storms occurred a~~
generally strong (albeit not perfect) agreement on the dates of the storms for those occurring in two or ~~three more~~ data sets.
~~Even among the common storms, mild variations in dates of occurrence were not eliminated.~~ These discrepancies exist despite
the use of a consistent methodology. This points to the important differences resulting from the use of different input data sets
when defining storm footprints and storm severity, while comparisons with XWS and C3S further highlight potential impacts
365 resulting from differences in storm footprint identification methodology.

3.2 Storms per Extended Winter Season

The distribution of storms per extended winter season over all Top50 storms identified from each of the four input data sets,
XWS and C3S is displayed in Figure 4. ~~The winters during the second ten years of our database exhibit less storm activity~~
~~relative to the first ten years; approximately 60% of all Top50 storms occurred during the first ten year period within each of~~
370 ~~the four input data sets, compared to approximately 40% during the second ten year period. However, important differences~~
~~among input data sets in how the storms are distributed across winters also appear. For instance, CCLM_ERA5_EUR-11 and~~
~~CCLM_ERA5_CEU-3 display a greater percentage of storms occurring during the first half of the timerecord (60% and 64%,~~
~~respectively) than do ERA5 and COSMO-REA6 (58% and 54%, respectively).~~

There is a large interannual variability, but no data set shows any significant trend in extreme storm occurrence over time.
375 Further, the input data sets demonstrate disagreement on the periods of low and high storm activity (Table 3). The input sources
agree best on the winters with no extreme storm activity: no storms were identified for the 1995/1996 and 2012/2013 extended
winter seasons from any of the four inputs, while one storm was identified within CCLM_ERA5_CEU-3 for the 2005/2006
(DORIAN) and 2008/2009 (QUINTEN) winters but not within the remaining three inputs. This level of agreement cannot
be found for other periods of low or high storm activity. There are no additional winters for which all four input sources
380 agree as belonging to periods of low storm activity, high storm activity, and very high storm activity, as indicated by winters
during which 2%, 8%, and 10+% of all storms occurred, respectively (Table 3). While ~~the three~~ data sets agree for one winter
within each of these periods, the remaining winters show agreement only among two input data sets or are unique to ~~that one~~
data set. However, even when different datasets agree on the categorisation of a specific winter, differences in the number of
detected storms can exist. For example, the ERA5, CCLM_ERA5_EUR-11, and CCLM_ERA5_CEU-3 data sets agree that the
385 1999/2000 winter exhibits very high extreme storm activity. However, 10% of all storms occurred during this winter in ERA5
and CCLM_ERA5_CEU-3, while 12% of all storms was obtained for CCLM_ERA5_EUR-11 ~~, demonstrating disagreement~~
~~on the number storms during this winter~~ (Figure 4; Table 3). It is notable that the two winters with very high activity displayed

by COSMO-REA6 occur during the second ten years of our database, whereas the very high activity winters for the remaining three input data sets occur during the first ten years.

390 The XWS and C3S databases display quite different distributions of storm frequencies compared to our database (Figure 4; Table 3). Though these two databases contained no extreme storms during the 1995/1996, ~~2005/2006,~~ and 2012/2013 winters, in agreement with our database, they ~~each contain four~~ also contain additional winters during which no extreme storms took place. ~~It is notable that XWS and C3S agree on three of these additional winters~~ which are not found in any of our four input data sets. They agree on two of these winters: 2003/2004 and 2010/2011 (Table 3). XWS and C3S further display no winters
395 that belong to the low storm activity category, and C3S displays no winters that belong to the high activity category. ~~XWS displays four high activity winters, agreeing with our database for two of these winters. Neither XWS nor~~ There is no clear difference in the level of agreement between our database and XWS or C3S ~~exhibit better agreement with our database,~~ given the similar mean absolute differences in storm percentage per winter ~~between XWS or C3S and our input data sets, and~~ and the large standard deviations (Table 4). The differences with our database partially reflect that we neglected storms identified
400 within XWS and C3S that were not identified within our database, and that XWS and C3S did not contain all the extreme storms identified within our database (nor all winters within the analysis period, in the case of XWS). The differences in storm frequency distribution and periods of high and low storm activity among our database, XWS, and C3S further highlight the influence of horizontal resolution, domain size, and methodology on extreme storm identification and characterization.

3.3 ~~Variations in~~ Relative Extreme Storm Severity

405 3.3.1 Relative Severity ~~per within and across~~ Winter ~~Season~~ Seasons

The interannual variability in winter severity within each data set is shown in Figure 5, where the proportion of total losses occurring during each winter is indicated. ~~In agreement with the storm frequency distributions discussed above, a larger proportion of total storm losses within each input data set took place during the first ten years of our database, with similar proportions of total losses during the first and second ten years as for storm frequencies. CCLM_ERA5_EUR-11 and CCLM_ERA5_CEU-3~~
410 ~~again display a greater proportion of total losses occurring during the first ten years than do ERA5 and COSMO-REA6. We~~ again see a large interannual variability, yet no significant trends in the percentage of normalised losses over time in any of the data sets. The proportion of total losses per winter within an input data set generally increases with the proportion of storms per winter, as ~~exhibited by the increases in the proportions of total losses with storm activity per winter for each input source~~ in also shown in Table 3. All four input data sets display a statistically significant Pearson's correlation coefficient ($p < .001$)
415 between 0.95 and 0.98 for the correlation between the percentage of all storms per winter and the percentage of total losses per winter. These statistics imply that ~~each input data set contains few winters during which the losses were,~~ for winters with several extreme storm occurrences, it is uncommon for the bulk of the losses to be caused by a single, exceptionally ~~strong~~ damaging storm. ~~storm rather than by several, weaker storms, as this would weaken the correlation.~~

damaging storm. This is further supported by the OEP/AEP ratio for each winter season and input data set displayed in
420 Figure 6. The OEP/AEP ratio indicates the degree to which a single storm event dominated that winter season's aggregated

loss (Priestley et al., 2018). A ratio much smaller than 1.0 indicates that multiple storm events contributed to the total seasonal losses, while a large ratio closer to 1.0 indicates that a single storm contributed most of the losses incurred during that season; a ratio exactly equal to 1.0 is achieved during winters containing only one storm. If we disregard the winters with a single storm, very few winters display a ratio greater than approximately 0.7 for each input source. This indicates that, for most winters that contained more than one extreme storm, the losses comprise large contributions from multiple storms. ~~Though the losses from a single storm during such winters sometimes contribute approximately 50-60% to the aggregated seasonal losses, it is unusual that an exceptionally severe storm contributes an overwhelming share of losses within our database (except for years with a single storm)~~ Single, exceptionally severe storms can contribute to a large part of the losses but rarely account for their near-totality.

~~However, some variability does exist~~ We also note some variability in average storm severity ~~during a given~~, both across different winters within each dataset and across datasets during the same winter. For example, during the 2006/2007 season, fewer storms caused a proportion of damages similar to the higher storm activity 1996/1997 and 2004/2005 winters in CCLM_ERA5_CEU-3, ~~and similar damage proportions to the higher storm activity 2007/2008 winter in~~. Similarly, ERA5 and displays twice the number of storms of CCLM_ERA5_EUR-11 (Table 3 in the 2002/2003 winter season, yet the proportions of

~~total losses in the two datasets are similar (cf. Figures 4 and 5). This implies that the storms identified within highlights both the variability in storm severity across winters for CCLM_ERA5_CEU-3 for the 2006/2007 season are more severe relative to the other input data sets for this winter, as well as to many other winters within and how the average storm severity per winter differs between CCLM_ERA5_CEU-3 EUR-11 and ERA5. As with the storm frequencies per winter, the~~ Thus, disagreements among input data sets for the severity of any given winter are also apparent in Figure 5 apparent, as was the case for storm frequencies per winter. As the same methodology was employed ~~, these to compute losses, these difference~~ could be due, at least in part, to differences in horizontal resolution and spatial domain impacting storm identification and severity assessment.

Again, ~~neither XWS nor C3S presented a better average comparison the level of agreement~~ with our database for the proportions of total losses per winter ~~, given the large standard deviations is comparable for XWS and C3S. While C3S shows a generally higher agreement, the improvement over XWS is small compared to the standard deviation of the mean differences~~

~~(Table 4). The storms within XWS for the 1997/1998 and 1998/1999 winters are likely weaker than the storms within the high activity category winters within our database given the smaller proportions of total XWS losses (Table 3). However, it is notable that the loss proportions for the XWS and C3S winters within the very high activity winter category are generally similar to those from our database, as are the remaining two high activity winters within XWS. Both XWS and C3S further exhibit two XWS also exhibits~~ winters of unremarkable storm activity, belonging to neither the ~~low nor high nor very~~ high activity categories, but large damages, ~~implying stronger~~. These winters thus displayed more severe storms compared to our database ~~for those winters (e.g., 2002/2003).~~ Variations in storm severity between XWS and C3S and relative to our database likely reflect differences in storm identification methodology that impact the storm severity assessments.

3.3.2 ~~Variations in~~ Relative Common Storm Severity

Differences in ~~We next consider~~ relative storm severity ~~among across~~ input data sets ~~are further highlighted in Figure 7~~ for the common storms ~~only~~ (Figure 7), and including XWS and C3S when these databases also identified one of the common storms. Storm XYNTHIA (2010-02-27) illustrates ~~this well, as XYNTHIA is least severe well the discrepancies between input data sets: it is amongst the least severe common storms~~ as derived from ERA5, ~~much stronger as derived from CCLM_ERA5_EUR-11 and XWS, and strongest as derived from~~ ~~but is much more severe in all other input datasets, most notably in COSMO-REA6 and CCLM_ERA5_CEU-3, which exhibit similar normalized losses~~ (Figure 7b). Storm KYRILL (2007-01-19) is instead an example of relatively good agreement among data sets, as it was the ~~strongest storm within our database~~ ~~most severe storm in all four input datasets to CLIMK-WINDS~~ and C3S in terms of ~~ordinal rank (Table 2)~~ and normalized losses (Figure 7), ~~a) and hence also first in ordinal rank (Table 2), and~~ and it was the ~~second-strongest~~ ~~second-most severe~~ storm within XWS.

~~No clear overarching trends~~ ~~Some differences~~ in storm severity with data set emerge in Figure 7, ~~but some tendencies are apparent. The~~. ~~Indeed~~, common storms tend to be stronger as derived from ERA5 and COSMO-REA6, in comparison to CCLM_ERA5_EUR-11 and CCLM_ERA5_CEU-3, as ~~implied by Figure 7 and~~ indicated by the the median and 75th percentile values of the common storm normalized losses for each input data set (Table 5). CCLM_ERA5_CEU-3, in turn, tends to exhibit the weakest storms, with smaller median, 25th percentile, and 75th percentile values than most of the other three input sources. However, the interquartile ranges for ~~all~~ the input data sets in Table 5 overlap with each other. Table 5 also demonstrates that the median and 25th percentile relative rank values for XWS fall within the ranges exhibited by ~~our database~~ ~~CLIMK-WINDS~~, while the 75th percentile is larger than ~~in for~~ any of the four ~~data sets input data sets that~~ we use. C3S ~~often exhibits the strongest or near-strongest common storms relative to our database and to XWS (Figure 5)~~ ~~exhibits the highest values for all percentiles~~. Its 25th percentile value is larger than all medians in Table 5 with the exception of COSMO-REA6, ~~suggesting statistically significantly larger storm severities and is also higher than the 75th percentile of CCLM_ERA5_CEU-3. This points to systematically higher normalised losses~~ compared to our database and XWS.

The substantial variation in storm severity with data set exhibited by the common storms, both within ~~our database and among our database, XWS, CLIMK-WINDS and when comparing CLIMK-WINDS to XWS~~ and C3S, ~~point points~~ to the impact of horizontal resolution and ~~storm footprint~~ identification methodology on ~~storm identification and~~ estimated severity. Additional causes, however, are likely also at play, such as the influence of the ~~configuration design~~ of the input data sets themselves on their representations of extreme ~~storms~~ ~~windspeeds~~. Because our database includes four different input data sets, this affords a unique perspective to investigate the cause(s) of these variations and their influence on extreme storm characterization and ~~impacts prediction~~ ~~related empirically-derived losses~~.

3.4 ~~Spatial Variability among Data Sets~~ ~~Extreme storm footprints~~

3.4.1 Spatial Variability of ~~Extreme~~ Storm Footprints

~~For the~~ ~~We conduct~~ a spatial comparison of the 29 common ~~storms, the storm footprints, using CCLM_ERA5_CEU-3 as reference. This is because it is the input data set with the highest native resolution and smallest native domain; CCLM_ERA5_CEU-3 is not assumed to be the most accurate or realistic data set. The ERA5, CCLM_ERA5_EUR-11, and COSMO-REA6 footprints~~

~~and corresponding absolute wind gusts, and the XWS and C3S absolute wind gusts,~~ were regridded using bilinear interpolation to the CCLM_ERA5_CEU-3 horizontal resolution and restricted to the enlarged Germany domain. The mean of the footprint differences over all ~~29~~ common storms was then taken for each input data set and ~~are presented as the mean~~
490 ~~footprint differences is presented~~ in Figure 8; ~~the same was done with the absolute wind gusts and the mean absolute wind gust~~
~~differences are presented in Figure 9. As these differences were not computed as absolute differences, the~~. The differences
between the CCLM_ERA5_CEU-3 input data set and the other data sets were first computed for each common storm, and then
averaged within each dataset across all common storms. The means presented in ~~Figures 8 and 9~~ Figure 8 carry the risk of
the cancellation of errors, which is particularly important for interpretation of mean differences of or near zero. ~~Comparisons~~
495 ~~were made to CCLM_ERA5_CEU-3 because this input data set has the highest native resolution and smallest native domain;~~
~~CCLM_ERA5_CEU-3 is not assumed to be the most accurate or realistic data set~~ We therefore also show the mean absolute
error in Figure 3 in the supplementary material. This analysis is not intended to evaluate which input source is the most accu-
rate, but rather seeks to characterize the main spatial differences in storm footprint arising among the input data sets used in
our database.

500 The footprint comparison between ERA5 and CCLM_ERA5_CEU-3 ~~stands out, as it~~ displays primarily positive mean dif-
ferences ~~over the enlarged Germany domain~~ with few exceptions, indicating that, on average, the footprint as derived from
CCLM_ERA5_CEU-3 is larger in magnitude than as derived from ERA5 (Figure 8). The footprint comparisons between
CCLM_ERA5_EUR-11 or COSMO-REA6 and CCLM_ERA5
_CEU-3 are more variable in terms of positive or negative mean ~~difference over this~~ differences across the domain. How-
505 ever, the mean footprint differences between CCLM_ERA5_CEU-3 and COSMO-REA6 tend to exhibit more locations with
negative sign or smaller-magnitude positive differences than do the differences with respect to CCLM_ERA5_EUR-11. The
footprint comparison between ERA5 and CCLM_ERA5_CEU-3 also stands out in terms of the larger magnitude of the differ-
ences, regardless of sign. Many locations within the domain display a footprint difference magnitude ~~between~~ above 0.2 ~~and~~
~~0.3~~, while the ~~smaller cluster of locations exhibiting larger magnitude~~ locations exhibiting negative differences within Czechia
510 and Slovakia ~~can reach a magnitude between~~ reach magnitudes below -0.2 ~~to~~ -0.4. This may be partially due to ERA5 hav-
ing the coarsest original native horizontal resolution ~~compared to CCLM_ERA5_CEU-3~~ of our input data sets. In compari-
son, the differences with CCLM_ERA5_EUR-11 and COSMO-REA6 tend to display magnitudes between -0.2 and 0.2 over
most of the enlarged Germany domain. The larger differences between CCLM_ERA5_CEU-3 and ERA5 become less evident
when considering mean absolute error (Figure 3 in the supplementary material), suggesting that CCLM_ERA5_EUR-11 and
515 COSMO-REA6 may display more cancellation of errors. It is also notable that CCLM_ERA5_CEU-3 and COSMO-REA6,
despite being derived from a common numerical model, still show considerable differences in their footprints.

These comparisons thus reveal substantial disagreements among the input data sets in the spatial structures of the common
storm footprints, though only the comparison between CCLM_ERA5_CEU-3 and ERA5 presents a domain-scale systematic
difference. ~~However, the mean differences can conceal other important disagreements among the footprints; these will be~~
520 ~~discussed in Section 3.4.3.~~

3.4.2 Spatial Variability of Associated Absolute Wind Gusts

~~Expectedly, the~~ We conduct the same comparison performed for the storm footprints for the absolute wind gusts. In this case, ~~we also include XWS and C3S. The~~ mean differences in the absolute wind gusts associated with the common storm footprints also reveal substantial disagreements among the input data sets, and ~~with between CLIMK_WINDS and XWS and C3S~~ (Figure 9). The mean ~~footprint differences and mean~~ absolute wind gust differences relative to CCLM_ERA5_CEU-3 for each input data set ~~are similar to each other in an overall sense generally resemble the footprint differences~~ (e.g. ~~;~~ a positive, larger magnitude difference in absolute wind gust in a region exhibiting a positive, larger magnitude difference in the footprint). ~~This is most notable over the northern portion of the domain (North and Baltic Seas, Denmark, and southern Sweden).~~ However, locations with differences of ~~opposing opposite~~ sign between the ~~footprint and~~ absolute wind gusts ~~and the footprint~~ are also apparent, as are locations where the ~~difference magnitudes are larger for the footprint but smaller for the absolute wind gusts, and vice versa. This seeming contradiction may be most readily seen in the comparison to ERA5. relative magnitudes of differences differ.~~ For example, the mean footprint differences between CCLM_ERA5_CEU-3 and ERA5 over northwestern Germany and the Netherlands are generally small and positive (between 0.0 and 0.1); ~~and~~; Figure 8), yet the mean absolute wind gust differences are ~~close to zero or more substantially negative (approximately -2.2 substantially negative over much~~ of the region (approximately -3 ms^{-1} or more ~~negative).~~ The mean absolute wind gust differences for CCLM; Figure 9). In terms of absolute wind gusts, ERA5_EUR-11 exhibit broader regions of positive difference (0.0 to approximately 4.4 ms^{-1} or greater) over much of continental Europe, in contrast to the mean footprint differences of small magnitude and variable sign. The mean footprint and absolute wind gust comparisons most resemble each other over the northern portion of the domain (North and Baltic Seas, Denmark, and southern Sweden) for each input data set. ERA5 no longer stands out over much of continental Europe as displaying ~~as displaying systematically~~ larger magnitude differences compared to the other two data sets when comparing the absolute wind gusts (see also Figure 4 in the supplementary material).

The mean footprint and absolute wind gust comparisons between CCLM_ERA5_CEU-3 and COSMO-REA6 resemble each other more than is the case for the other two input data sets, as would be expected since they are derived from a common numerical model. The differences identified between all data sets in terms of wind gusts could arise, for example, from different boundary conditions, different wind gust computations or parameterizations, or configuration options unique to a data set. Increasing horizontal resolution can also allow for greater spatial heterogeneity within a data set's wind gust field, and thus greater heterogeneity within the the threshold exceedances used to compute the footprints.

~~The comparisons for XWS or~~ The fact that some of the largest differences are found in the Alpine region also point to a role of spatial resolution, notably for the comparison between CCLM_ERA5_CEU-3 and COSMO-REA6 which issue from a common numerical model.

Notable differences also emerge in the comparison with XWS and C3S ~~further reflect impacts on the wind gust fields within these databases arising from different horizontal resolutions, the use of ERA-Interim or ERA5 to create the database, and the choice of storm identification methodology, and partially reflect that not, in this case with the added effect that the latter two~~

databases also present methodological differences to CLIMK-WINDS Figure 9). The results are also not fully comparable to the ones discussed above since not all common storms were captured by ~~these two databases. The XWS and C3S. The differences between CCLM ERA5 CEU-3 and XWS and C3S comparisons~~ do not resemble each other, nor do they resemble the comparisons within our own database. For example, ~~the comparison to C3S displays quite substantially primarily~~ positive differences over ~~almost the entire eastern half the eastern and southern parts~~ of the domain ~~not seen in the other data sets; these differences are much larger in magnitude than the positive differences exhibited by the comparison to CCLM ERA5 EUR-11, and lack the regions of very small to no difference that are also seen in the CCLM ERA5 EUR-11 comparison. The comparison to XWS is more variable in sign and magnitude over the central continental portion of the domain, but displays substantially and negative differences over northwestern parts. The latter instead display primarily positive differences for XWS. Similarly, XWS displays substantial~~ negative differences over Denmark, southern Sweden, Poland, and the Baltic Sea that are in stark contrast to ~~the remaining ERA5 and, to some extent, also to the other~~ data sets. The comparison to XWS indeed exhibits the broadest area of negative difference of any of the data sets, indicating that the absolute wind gusts from XWS tend to be ~~larger stronger~~ than those from CCLM ERA5 CEU-3. ~~This major difference also emerges clearly in the mean absolute errors (Figure 4 in the supplementary materials).~~

3.4.3 Windstorms KYRILL and ANDREA as Comparative Examples

~~As mentioned above, The above comparisons provide a mean picture, but~~ individual storm footprints and absolute wind gusts ~~contain manifestations of spatial variability across data sets additional to those display additional features not~~ apparent in the mean comparisons. ~~The comparisons between CCLM ERA5 CEU-3 and the remaining data sets for the individual We consider~~ storms KYRILL (2007-01-18) and ANDREA (2012-01-05) ~~provide such examples as cases in point.~~ KYRILL is the ~~strongest most severe~~ storm within our database and C3S, and the ~~second strongest second most severe~~ within XWS. ANDREA also falls within the top 10 most ~~extreme severe~~ storms for all data sets except XWS (Table 2), but ~~unlike KYRILL,~~ its relative storm ~~strengths are severity is~~ much more variable ~~than that of KYRILL~~ among input sources (Figure 7). ~~Most The~~ footprint and absolute wind gust comparisons for KYRILL (Figures 10 and 11) and ANDREA (Figures 12 and 13) depart substantially from the mean comparisons. ~~The ordinal and relative strengths for KYRILL and ANDREA serve to highlight that agreement in spatial variability among input data sets does not depend on storm severity, nor does it depend on agreement on storm severity among input data sets.~~

The KYRILL and ANDREA CCLM ERA5 CEU-3 footprints display ~~a the~~ signature of convective activity over continental Europe, albeit stronger in KYRILL, that is weaker or missing in the other data sets. This appears as the "straight line" maximum relative wind gusts apparent in Figures ~~2, 10, 10~~ and 12. ~~High resolution and a convection-permitting configuration, such as in The "straight line" maxima are also visible, although not as clearly, in the CCLM ERA5 CEU-3 , contribute to wind gusts~~ (Figure. 11 and 13) The higher resolution and convection-permitting configuration of CCLM ERA5 CEU-3, likely contribute to a better representation of the downdrafts and wind gusts arising from convective activity associated with winter windstorms ~~, causing the "straight line" relative wind gusts in the footprint~~ (Ludwig et al., 2015). Several storms in our database exhibit this signature in CCLM ERA5 CEU-3 and occasionally other input data sets. ERA5 typically misses it entirely, ~~with an increasing~~

ability to represent such mesoscale processes for CCLM with CCLM_ERA5_EUR-11 and COSMO-REA6 performing better, though not as well as CCLM_ERA5_CEU-3. Spatial variability arising from convective activity is entirely missing in the mean footprint comparison.

The mean comparisons also conceal the differences in footprint spatial extents or boundaries among data sets for the same storm, as appears in the KYRILL and ANDREA comparisons. Disagreements in extent typically occur in the eastern portion of and the southern boundary and southern portions of the domain, as highlighted by the bright red regions for KYRILL and ANDREA indicating that CCLM_ERA5_CEU-3 locates the footprint there but the other data set did not (Figures 10 and 12, 11, 12 and 13). In addition to footprint boundary disagreements, gaps within footprints containing locations that were not impacted by the surrounding storm exist for some data sets but not others. Such a gap is apparent in the comparison between CCLM_ERA5_CEU-3 and ERA5 for ANDREA over the Danish islands, Baltic Sea, southern Baltic Sea and northeastern Germany (Figure 12 Figures 12 and 13). The causes for disagreements in footprint extents and locations included in a footprint interior are unclear, but such disagreements are again likely related to horizontal resolution, footprint identification methodology, and data set configuration differences. Care should thus be taken when choosing an appropriate footprint source(s) for local or regional loss estimations.

Individual storms therefore present additional differences in their footprints and wind gusts across data sets that are not visible in a bulk comparison across multiple storms. The ordinal and relative ranks for KYRILL and ANDREA serve to highlight that these differences emerge independently of storm severity and of agreement on relative storm severity among input data sets.

4 Conclusions

We have presented and characterized CLIMK-WINDS: a new database of the Top50 most extreme European winter windstorms over the 1995-2015 extended winter seasons based on a consistent footprint identification and severity assessment methodology. The latter was based on an empirical storm loss model. We used four different meteorological input data sets, namely the ERA5, and COSMO-REA6 reanalyses, and the CCLM_ERA5_EUR-11, COSMO-REA6, and CCLM_ERA5_CEU-3 reanalyses or regional climate model simulation outputs. We applied the same footprint selection methodology to each input data set on its native horizontal resolution and spatial domain, thereby creating a consistent, systematic database of the most extreme storms over a 20-year time record severe storms identified across diverse input data source sets. This allows for a direct comparison of the effects of using different wind gust input data sources sets on the storm footprints, their severity, and their spatial characteristics. Equally importantly, the temporal range of the data sets can be easily extended when new data become available, and it is equally easy possible for users to include additional input data sets to which they may have access. Our database therefore provides complementary perspectives on extreme storm identification and severity assessment given the diverse horizontal resolutions and input sources used, and complements existing extreme European winter windstorm databases.

Our results highlight the major effects that horizontal resolution, choice of footprint identification methodology, and differing types of input data sets can have on extreme storm identification and characterization. The effect of horizontal resolution is most evident in the comparison of the CCLM ERA5 CEU-3 and CCLM ERA5 EUR-11 data, which originate from a common numerical model. The result of different methodological choices emerges from the comparison of the footprints within CLIMK-WINDS with those from the XWS and C3S data. Finally the data sets that we used to build our database include a global reanalysis, a regional reanalysis and regional climate models, elucidating the effect of differing types of input data sets. We found variability across input data sets in all aspects of our database, ~~from including~~ which storms were identified as belonging to the Top50 storms ~~to the quite substantial variability in the and the~~ spatial structure of the footprints ~~for of~~ storms identified in more than one input data set, ~~as well as~~. We found similar differences between our database and the XWS and C3S databases.

~~Our~~ The CLIMK-WINDS database identified a total of 76 ~~extreme storms, with unique storms ranking amongst the Top50 in at least one of our input data sets. There were~~ 29 common storms identified in all four input ~~sources. Another 47 storms were identified in at least one of the input sources but not all four. The strongest data sets. The most severe~~ storms, such as KYRILL, LOTHAR, and JEANETT, tended to be identified in all four input ~~sources~~ data sets, though the common storms also include many comparatively weaker storms. ~~However, the~~ Another 47 storms were identified as extreme storms in at least one of the input sources but not all four. Of these storms, 18 were identified in only one of the four datasets. Unexpectedly, CCLM ERA5 CEU-3 displayed more such storms than COSMO-REA6, despite its much smaller domain. This may be partly due to its higher resolution, leading it to identify as extreme some storms which do not display particularly strong wind gusts in the other datasets. For both the common storms and the storms identified in only two or three datasets, the severity of a given storm, as denoted by a storm's ordinal rank and normalized loss, often varied across the input data sets ~~in which it was identified for both the common storms and the storms identified in only two or three inputs. The variation in relative storm strengths could be severity was~~ substantial across input data sets for many storms, ~~though~~. Moreover, the storms as identified by C3S tended to be ~~stronger~~ relatively more severe than as identified in the other data sets. The input data sets, XWS, and C3S also disagreed on the winters with low and high storm activity ~~and~~, while a large, statistically significant positive correlation exists between the proportion of all storms per winter and the proportion of total losses per winter for each input data set, they disagree on how relatively damaging each winter was. Equally importantly, the CLIMK-WINDS footprints and associated absolute wind gusts themselves exhibited substantial variability across input data sets and with XWS and C3S, both in the mean comparisons and for individual storms. Comparisons of the footprint and absolute wind gust spatial variability for individual storms revealed ~~smaller-scale~~ smaller-scale features not present in the mean comparisons, such as the signatures of convective activity. ~~Mild disagreement of one day is~~ Disagreements are also seen for some storms in ~~their the~~ dates of occurrence, ~~but although these are typically of one day. Moreover,~~ the majority of storms within our database ~~agreed on their display the same~~ dates of occurrence in all input data sets and in XWS and C3S.

We underscore that our goal in creating ~~this the~~ the CLIMK-WINDS database was not to provide a record of accurate and inaccurate footprints nor to assess which input data set or horizontal resolution performed best at identification and characterization of extreme European winter storms. Rather, we seek to provide a diverse yet consistent record of storm footprints that allows

for different perspectives on to understand the uncertainties related to choice of input data set for these storms. A diversity of perspectives has been lacking for footprint data, and thus, by providing multiple sources but considering multiple input data sets processed using a consistent methodology, our database supports uncertainty quantification for a deeper understanding of extreme storms and their impacts. Indeed, uncertainty quantification for footprint data is a little-investigated topic, and our database could allow for assignment of an uncertainty range in the footprints, or the design and assessment of another uncertainty metric after the needs of specific end-users. Combining This could in the future be combined with an analysis of the uncertainties associated with the footprints and storm losses may lead storm loss models (Moemken et al., 2024a), leading to improved models of storm damages and improved understanding of how such storms these may change with climate change a changing climate. Storm footprints derived from multiple data sources on different resolutions could also present an advantage for many areas of research that currently give little weight to historical data uncertainty, such as storm clustering studies (Dacre and Pinto, 2020).

It is likely that the The footprint(s) that are most useful to a particular user of our the CLIMK-WINDS database could depend heavily on that user's specific goals; for. For example, interest in a storm affecting a mountainous region might call for footprints at higher horizontal resolution and neglect of those at coarser resolution. However, based on our comparisons here, it is most useful We nonetheless exhort users to consider the storms as identified from two or more input sources within our database, where possible, rather than a footprint from a single source. We exhort users to do so, in order to provide. This enables providing information on the uncertainties associated with the storms and their impacts and more reliable analysis. Indeed, our CLIMK-WINDS dataset and analysis highlight the value of considering multiple data sources when characterising extreme storms and their impacts.

5 Data availability

CLIMK-WINDS is publicly available online through Zenodo at <https://doi.org/10.5281/zenodo.10594399> (Flynn et al., 2024). The ERA5 reanalysis data (Hersbach et al., 2020) and the C3S winter windstorm indicators for Europe (the C3S windstorm database, (C3S Climate Data Store, 2022)), used as an input source for our database and as a comparison to our database, respectively, are publicly available online through the Copernicus Climate Change Service (<https://10.24381/cds.adbb2d47> and <https://doi.org/10.24381/cds.9b4ea013>). The CCLM_ERA5_CEU-3 simulation output (Brienen et al., 2022), used as an input source for our database, is publicly available online from the DWD Earth System Federation Grid nodes (<https://esgf.dwd.de/search/esgf-dwd/>). The COSMO-REA6 data set (Bollmeyer et al., 2015), used as an input source for our database, is publicly available from DWD (https://opendata.dwd.de/climate_environment/REA/). The XWS database (Roberts et al., 2014), used as a comparison to our database, is publicly available online (<https://www.europeanwindstorms.org/repository/>).

Author contributions. GM and JGP conceived of the database. CMF performed the data collection and analysis and wrote the initial paper draft. JM provided guidance to CMF in how to compute the storm footprints and loss indices, and how to rank the storms. MS and CMF

created and revised the figures and tables. GM revised the paper text. All authors discussed the results and contributed to drafting the manuscript.

Competing interests. The authors declare that they have no conflicts of interest.

690 *Acknowledgements.* This research has received funding from the European Union's Horizon research and innovation programme under European Research Council grant no. ~~101112727~~ [101112727](#) and from the Swedish Research Council Vetenskapsrådet under grant no. [2022-06599](#). JM was funded by the Bundesministerium für Bildung und Forschung (BMBF; German Ministry for Education and Research) under project "RegIKlim-NUKLEUS" (01LR2002B1). JGP thanks the AXA Research Fund for support. [All authors thank HR and one anonymous Reviewer for their constructive input on the study.](#)

- Baldauf, M., Seifert, A., Förstner, J., Majewski, D., Raschendorfer, M., and Reinhardt, T.: Operational Convective-Scale Numerical Weather Prediction with the COSMO Model: Description and Sensitivities, *Monthly Weather Review*, 139, 3887 – 3905, <https://doi.org/https://doi.org/10.1175/MWR-D-10-05013.1>, 2011.
- Bollmeyer, C., Keller, J. D., Ohlwein, C., Wahl, S., Crewell, S., Friederichs, P., Hense, A., Keune, J., Kneifel, S., Pscheidt, I., Redl, S., and Steinke, S.: Towards a high-resolution regional reanalysis for the European CORDEX domain, *Quarterly Journal of the Royal Meteorological Society*, 141, 1–15, <https://doi.org/https://doi.org/10.1002/qj.2486>, 2015.
- Bonavita, M., Hólm, E., Isaksen, L., and Fisher, M.: The evolution of the ECMWF hybrid data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 142, 287–303, <https://doi.org/https://doi.org/10.1002/qj.2652>, 2016.
- Brienen, S., Haller, M., Brauch, J., and Frueh, B.: HoKliSim-De COSMO-CLM climate model simulation data version V2022.01, https://doi.org/10.5676/DWD/HOKLISIM_V2022.01, [data set], 2022.
- C3S Climate Data Store: Winter windstorm indicators for Europe from 1979 to 2021 derived from reanalysis, Copernicus Climate Change Service (C3S) Climate Data Store (CDS), <https://doi.org/10.24381/cds.9b4ea013>, 2022.
- CIESIN: Gridded Population of the World, Version 4 (GPWv4): Population Density, Revision 11, Palisades, New York: NASA Socioeconomic Data and Applications Center (SEDAC), <https://doi.org/10.7927/H49C6VHW>, 2018.
- Cusack, S.: A long record of European windstorm losses and its comparison to standard climate indices, *Natural Hazards and Earth System Sciences*, 23, 2841–2856, <https://doi.org/10.5194/nhess-23-2841-2023>, 2023.
- Dacre, H. F. and Pinto, J. G.: Serial clustering of extratropical cyclones: A review of where, when and why it occurs., *NPJ Climate and Atmospheric Science*, 3 (1), 2397–3722, <https://doi.org/10.1038/s41612-020-00152-9>, 2020.
- Davies, T., Cullen, M. J. P., Malcolm, A. J., Mawson, M. H., Staniforth, A., White, A. A., and Wood, N.: A new dynamical core for the Met Office’s global and regional modelling of the atmosphere, *Quart. J. Roy. Meteor. Soc.*, 131, 1759–1782, 2005.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137, 553–597, <https://doi.org/https://doi.org/10.1002/qj.828>, 2011.
- Deutsche Rück: Sturmdokumentation 2007, Publication of the Deutsche Rück Reinsurance Company, Duesseldorf, Germany, https://www.deutscherueck.de/fileadmin/Downloads/Sturmdoku_2007_web.pdf (in German), last accessed: 2024-01-16, 2008.
- Fink, A. H., Brücher, T., Ermert, V., Krüger, A., and Pinto, J. G.: The European storm Kyrill in January 2007: synoptic evolution, meteorological impacts and some considerations with respect to climate change, *Natural Hazards and Earth System Sciences*, 9, 405–423, <https://doi.org/10.5194/nhess-9-405-2009>, 2009.
- Flynn, C. M., Moemken, J., Pinto, J. G., and Messori, G.: Storm Database Files for A New Database of Extreme European Winter Windstorms, <https://zenodo.org/uploads/10594399>, <https://doi.org/10.5281/zenodo.10594399>, version 1, Zenodo [data set], 2024.
- Giorgi, F. and Gutowski, W. J.: Regional Dynamical Downscaling and the CORDEX Initiative, *Annual Review of Environment and Resources*, 40, 467–490, <https://doi.org/10.1146/annurev-environ-102014-021217>, 2015.

- Gliksman, D., Averbek, P., Becker, N., Gardiner, B., Goldberg, V., Grieger, J., Handorf, D., Haustein, K., Karwat, A., Knutzen, F., Lentink, H. S., Lorenz, R., Niermann, D., Pinto, J. G., Queck, R., Ziemann, A., and Franzke, C. L. E.: Review article: A European perspective on wind and storm damage – from the meteorological background to index-based approaches to assess impacts, *Natural Hazards and Earth System Sciences*, 23, 2171–2201, <https://doi.org/10.5194/nhess-23-2171-2023>, 2023.
- 735 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/https://doi.org/10.1002/qj.3803>, 2020.
- 740 Jacob, D., Petersen, J., Eggert, B., Alias, A., Christensen, O. B., Bouwer, L. M., Braun, A., Colette, A., Déqué, M., Georgievski, G., Georgopoulou, E., Gobiet, A., Menut, L., Nikulin, G., Haensler, A., Hempelmann, N., Jones, C., Keuler, K., Kovats, S., Kröner, N., Kotlarski, S., Kriegsmann, A., Martin, E., van Meijgaard, E., Moseley, C., Pfeifer, S., Preuschmann, S., Radermacher, C., Radtke, K., Rechid, D., Rounsevell, M., Samuelsson, P., Somot, S., Soussana, J.-F., Teichmann, C., Valentini, R., Vautard, R., Weber, B., and Yiou, P.: EURO-CORDEX: new high-resolution climate change projections for European impact research, *Reg. Environ. Change*, 14, <https://doi.org/https://doi.org/10.1007/s10113-013-0499-2>, 2014.
- 745 Jacob, D., Teichmann, C., Sobolowski, S., Katragkou, E., Anders, I., Belda, M., Benestad, R., Boberg, F., Buonomo, E., Cardoso, R. M., Casanueva, A., Christensen, O. B., Christensen, J. H., Coppola, E., De Cruz, L., Davin, E. L., Dobler, A., Domínguez, M., Fealy, R., Fernandez, J., Gaertner, M. A., García-Díez, M., Giorgi, F., Gobiet, A., Goergen, K., Gómez-Navarro, J. J., Alemán, J. J. G., Gutiérrez, C., Gutiérrez, J., Güttler, I., Haensler, A., Halenka, T., Jerez, S., Jiménez-Guerrero, P., Jones, R. G., Keuler, K., Kjellström, E., Knist, S., Kotlarski, S., Maraun, D., van Meijgaard, E., Mercogliano, P., Montávez, J. P., Navarra, A., Nikulin, G., de Noblet-Ducoudré, N., Panitz, H.-J., Pfeifer, S., Piazza, M., Pichelli, E., Pietikäinen, J.-P., Prein, A. F., Preuschmann, S., Rechid, D., Rockel, B., Romera, R., Sánchez, E., Sieck, K., Soares, P. M. M., Somot, S., Srnc, L., Sørland, S. L., Termonia, P., Truhetz, H., Vautard, R., Warrach-Sagi, K., and Wulfmeyer, V.: Regional climate downscaling over Europe: perspectives from the EURO-CORDEX community, *Reg. Environ. Change*, 20, <https://doi.org/https://doi.org/10.1007/s10113-020-01606-9>, 2020.
- 755 Karremann, M., Pinto, J., Von Bomhard, P., and Klawa, M.: On the clustering of winter storm loss events over Germany, *Natural Hazards and Earth System Sciences*, 14, 2041–2052, 2014.
- Klawa, M. and Ulbrich, U.: A model for the estimation of storm losses and the identification of severe winter storms in Germany, *Natural Hazards and Earth System Sciences*, 3, 725–732, <https://doi.org/10.5194/nhess-3-725-2003>, 2003.
- 760 Leckebusch, G. C., Ulbrich, U., Fröhlich, L., and Pinto, J. G.: Property loss potentials for European midlatitude storms in a changing climate, *Geophysical Research Letters*, 34, <https://doi.org/https://doi.org/10.1029/2006GL027663>, 2007.
- Little, A. S., Priestley, M. D. K., and Catto, J. L.: Future increased risk from extratropical windstorms in northern Europe., *Nat. Commun.*, 14, <https://doi.org/https://doi.org/10.1038/s41467-023-40102-6>, 2023.
- Ludwig, P., Pinto, J. G., Hoepp, S. A., Fink, A. H., and Gray, S. L.: Secondary Cyclogenesis along an Occluded Front Leading to Damaging Wind Gusts: Windstorm Kyrill, January 2007, *Monthly Weather Review*, 143, 1417 – 1437, <https://doi.org/10.1175/MWR-D-14-00304.1>, 2015.
- 765 Mitchell-Wallace, K., Jones, M., Hillier, J., and Foote, M.: *Natural Catastrophe Risk Management and Modelling: A Practitioner’s Guide*, Wiley-Blackwell, 2017.

Moemken, J., Alifdini, I., Ramos, A. M., Georgiadis, A., Brocklehurst, A., Braun, L., and Pinto, J. G.: Insurance loss model vs. meteorological
770 loss index – how comparable are their loss estimates for European windstorms?, *Natural Hazards and Earth System Sciences*, 24, 3445–
3460, <https://doi.org/10.5194/nhess-24-3445-2024>, 2024a.

Moemken, J., Messori, G., and Pinto, J. G.: Windstorm losses in Europe – What to gain from damage datasets, *Weather and Climate Extremes*,
44, 100661, <https://doi.org/https://doi.org/10.1016/j.wace.2024.100661>, 2024b.

Munich Re: Geo Risks Research: Loss events in Europe 1980–2014: 10 costliest winter storms ordered by insured losses, Tech. rep., https://www.preventionweb.net/files/44281_19802014paketworldusde4zu3.pdf, last accessed: 2024-01-16, 2015.
775

Munich Re: Risks posed by natural disasters: Losses are trending upwards, [https://www.munichre.com/en/risks/
natural-disasters-losses-are-trending-upwards.html](https://www.munichre.com/en/risks/natural-disasters-losses-are-trending-upwards.html), last accessed: 2024-01-16, 2022.

Pinto, J. G., Fröhlich, E. L., Leckebusch, G. C., and Ulbrich, U.: Changing European storm loss potentials under modified climate condi-
tions according to ensemble simulations of the ECHAM5/MPI-OM1 GCM, *Natural Hazards and Earth System Sciences*, 7, 165–175,
780 <https://doi.org/10.5194/nhess-7-165-2007>, 2007.

Pinto, J. G., Karremann, M. K., Born, K., Della-Marta, P. M., and Klawns, M.: Loss potentials associated with European windstorms under
future climate conditions, *Climate Research*, 54, 1–20, 2012.

Pinto, J. G., Gómara, I., Masato, G., Dacre, H. F., Woollings, T., and Caballero, R.: Large-scale dynamics associated with clus-
tering of extratropical cyclones affecting Western Europe, *Journal of Geophysical Research: Atmospheres*, 119, 13,704–13,719,
785 <https://doi.org/https://doi.org/10.1002/2014JD022305>, 2014.

Pinto, J. G., Pantillon, F., Ludwig, P., Déroche, M.-S., Leoncini, G., Raible, C. C., Shaffrey, L. C., , and Stephenson, D. B.: From Atmo-
spheric Dynamics to Insurance Losses: An Interdisciplinary Workshop on European Storms, *B. Am. Meteorol. Soc.*, 100, ES175–ES178,
<https://doi.org/https://doi.org/10.1175/BAMS-D-19-0026.1>, 2019.

Priestley, M. D. K. and Catto, J. L.: Future changes in the extratropical storm tracks and cyclone intensity, wind speed, and structure, *Weather
790 and Climate Dynamics*, 3, 337–360, <https://doi.org/10.5194/wcd-3-337-2022>, 2022.

Priestley, M. D. K., Dacre, H. F., Shaffrey, L. C., Hodges, K. I., and Pinto, J. G.: The role of serial European windstorm clustering for extreme
seasonal losses as determined from multi-centennial simulations of high-resolution global climate model data, *Natural Hazards and Earth
System Sciences*, 18, 2991–3006, <https://doi.org/10.5194/nhess-18-2991-2018>, 2018.

Roberts, J. F., Champion, A. J., Dawkins, L. C., Hodges, K. I., Shaffrey, L. C., Stephenson, D. B., Stringer, M. A., Thornton, H. E., and
795 Youngman, B. D.: The XWS open access catalogue of extreme European windstorms from 1979 to 2012, *Natural Hazards and Earth
System Sciences*, 14, 2487–2501, <https://doi.org/10.5194/nhess-14-2487-2014>, 2014.

Rockel, B., Will, A., and Hense, A.: The Regional Climate Model COSMO-CLM (CCLM), *Meteorologische Zeitschrift*, 17, 347–348,
<https://doi.org/10.1127/0941-2948/2008/0309>, 2008.

Schwierz, C., Köllner-Heck, P., Zenklusen Mutter, E., Bresch, D. N., Vidale, P.-L., Wild, M., and Schär, C.: Modelling European winter wind
800 storm losses in current and future climate., *Climatic Change*, 101, 485–514, <https://doi.org/https://doi.org/10.1007/s10584-009-9712-1>,
2010.

Severino, L. G., Kropf, C. M., Afargan-Gerstman, H., Fairless, C., de Vries, A. J., Domeisen, D. I. V., and Bresch, D. N.: Projections and
uncertainties of winter windstorm damage in Europe in a changing climate, *Natural Hazards and Earth System Sciences*, 24, 1555–1578,
<https://doi.org/10.5194/nhess-24-1555-2024>, 2024.

805 Ulbrich, U., Leckebusch, G., and Pinto, J.: Extra-tropical cyclones in the present and future climate: a review., *Theor. Appl. Climatol.*, 96,
117–131, <https://doi.org/10.1007/s00704-008-0083-8>, 2009.

van den Brink, H. W.: An effective parametrization of gust profiles during severe wind conditions, *Environmental Research Communications*, 2, 011 001, <https://doi.org/10.1088/2515-7620/ab5777>, 2019.

Walz, M. A. and Leckebusch, G. C.: Loss potentials based on an ensemble forecast: How likely are winter windstorm losses similar to 1990?, *Atmospheric Science Letters*, 20, e891, <https://doi.org/https://doi.org/10.1002/asl.891>, 2019.

810

Table 1. Summary of input data sets used to create ~~the extreme windstorms database~~CLIMK-WINDS, or used for comparison to the database.

Data Set Name	Data Set Type	Domain	Domain Boundaries	Horizontal Resolution	Period Covered	Number of Storms Identified
ERA5	Global Reanalysis	Global (European sub-domain selected)	27 °N–72 °N, 22 °W–45 °E	°0.25	1940 – near present	50
CCLM_ERA5_EUR-11	Regional Climate Model	CORDEX EUR-11	~ 27 °N–72 °N, ~ 22 °W–45 °E	0.11° (~ 12 km)	1979 – 2020	50
COSMO-REA6	Regional Reanalysis	CORDEX EUR-11	~ 27 °N–72 °N, ~ 22 °W–45 °E	0.055° (~ 6 km)	1995 – 2019	50
CCLM_ERA5_CEU-3	Regional Climate Model	Enlarged Germany domain	~ 46 °N–58 °N, ~ 2 °E–20 °E	0.0275° (~ 2.8 km)	1979 – 2019	50
XWS	Derived from ERA-Interim Reanalysis and insurance data	Europe	~ 36 °N–68 °N, ~ 20 °W–40 °E	~ 25 km	1979 – 2014	23
C3S	Derived from statistical downscaling of ERA5	Europe	~ 30 °N–70 °N, ~ 25 °W–40 °E	1 km	1979 – 2021	30

Table 2. Summary of the Top50 extreme windstorms common to all of the four input data sets, with their dates of occurrence and ordinal ranks (rank 1 = most ~~extreme~~~~severe~~ storm, rank 50 = least ~~extreme~~~~severe~~ storm of the Top50 storms). A storm name in parentheses indicates that the same storm had two different names, while a hyphenated name indicates that two individual storms could not be effectively separated from each other within the database. For storms lacking given names and thus named after their date of occurrence, the ERA5 date was preferred. An "X" marks storms missing from a dataset; "–" indicates storms outside XWS's coverage period.

Storm Name	ERA5		CCLM_ERA5_EUR-11		COSMO-REA6		CCLM_ERA5_CEU-3		XWS		C3S	
	Date	Rank	Date	Rank	Date	Rank	Date	Rank	Date	Rank	Date	Rank
ANATOL	1999-12-03	18	1999-12-03	20	1999-12-03	14	1999-12-03	19	1999-12-03	3	1999-12-03	16
ANDREA (ULLI)	2012-01-05	4	2012-01-05	5	2012-01-05	5	2012-01-05	8	2012-01-03	12	2012-01-03	4
ANNA	2002-02-26	24	2002-02-26	17	2002-02-26	19	2002-02-26	25	X	X	2002-02-26	19
ARIANE	1997-02-13	32	1997-02-13	12	1997-02-13	10	1997-02-14	18	X	X	X	X
CARMEN	2010-11-12	33	2010-11-12	43	2010-11-12	48	2010-11-12	43	X	X	X	X
ELIVRA-FARAH	1998-03-04	20	1998-03-04	25	1998-03-04	22	1998-03-04	24	X	X	X	X
EMMA	2008-03-01	6	2008-03-01	19	2008-03-01	8	2008-03-01	4	2008-02-29	21	2008-03-02	6
FANNY	1998-01-04	12	1998-01-04	11	1998-01-04	11	1998-01-04	27	1998-01-04	20	1998-01-04	11
FRANZ	2007-01-11	7	2007-01-12	15	2007-01-11	13	2007-01-11	12	X	X	X	X
FRIDTJOF	2007-12-02	23	2007-12-02	23	2007-12-02	28	2007-12-02	37	X	X	X	X
GISELA-HEIDI	1997-02-25	9	1997-02-25	16	1997-02-25	9	1997-02-25	14	X	X	1997-02-23	8
GUNTER	2015-01-10	22	2015-01-10	21	2015-01-10	24	2015-01-10	16	–	–	2015-01-10	18
ILONA	2002-01-27	26	2002-01-27	28	2002-01-27	25	2002-01-27	17	X	X	2002-01-27	15
JEANETT	2002-10-27	5	2002-10-27	4	2002-10-27	3	2002-10-27	2	2002-10-27	1	2002-10-27	5
JOACHIM	2011-12-16	30	2011-12-16	31	2011-12-16	30	2011-12-16	31	2011-12-16	9	2011-12-16	23
KERSTIN-LIANE	2000-01-30	43	2000-01-30	44	2000-01-30	39	2000-01-30	33	X	X	X	X
KIRSTEN	2008-03-11	11	2008-03-11	7	2008-03-11	4	2008-03-11	13	X	X	2008-03-11	10
KYRILL	2007-01-18	1	2007-01-18	1	2007-01-18	1	2007-01-18	1	2007-01-18	2	2007-01-18	1
LARA	1999-02-05	40	1999-02-05	32	1999-02-05	33	1999-02-05	23	X	X	X	X
LOTHAR	1999-12-26	2	1999-12-26	3	1999-12-26	2	1999-12-26	3	1999-12-26	13	1999-12-26	2
MIKE-NIKLAS	2015-03-30	14	2015-04-01	10	2015-03-30	6	2015-30-30	6	–	–	X	X
NILS	2015-11-29	37	2015-11-29	50	2015-11-29	47	2015-11-30	38	–	–	X	X
NINA-ORALIE	2004-03-20	29	2004-03-20	36	2004-03-20	18	2004-03-20	26	X	X	X	X
ORATIA	2000-10-30	10	2000-10-30	22	2000-10-30	7	2000-10-30	42	2000-10-30	4	2000-10-30	9
u19950216	1995-02-16	47	1995-02-17	47	1995-02-17	37	1995-02-16	46	X	X	X	X
ULF	2005-02-13	48	2005-02-13	24	2005-02-13	35	2005-02-13	28	X	X	X	X
XAVER	2013-12-05	28	2013-12-05	26	2013-12-05	20	2013-12-05	20	2013-12-05	6	2013-12-05	22
XYLIA	1998-10-28	16	1998-10-28	9	1998-10-28	12	1998-10-28	10	1998-10-28	23	1998-10-28	14
XYNTHIA	2010-02-27	45	2010-02-27	29	2010-02-28	15	2010-02-28	11	2010-02-27	7	X	X

Table 3. Summary of the winters within each input data set, XWS, and C3S that belong to the periods of no storm activity (0% of all storms during one winter season), low storm activity (2% of all storms during one winter season), high storm activity (8% of all storms during one winter season), and very high storm activity (10+% of all storms during one winter season). The percentages of ~~total~~normalized storm losses that occurred during each winter for each data set are given in parentheses. Winters are listed in chronological order for each data set and section.

	ERA5 (%)	CCLM_ERA5_EUR-11 (%)	COSMO-REA6 (%)	CCLM_ERA5_CEU-3 (%)	XWS (%)	C3S (%)
No Storm Activity (0% of all storms)	1995/1996 (0%)	1995/1996 (0%)	1995/1996 (0%)	1995/1996 (0%)	1995/1996 (0%)	1995/1996 (0%)
	2005/2006 (0%)	2005/2006 (0%)	2005/2006 (0%)	2012/2013 (0%)	2003/2004 (0%)	2003/2004 (0%)
	2008/2009 (0%)	2008/2009 (0%)	2008/2009 (0%)		2005/2006 (0%)	2005/2006 (0%)
	2012/2013 (0%)	2012/2013 (0%)	2012/2013 (0%)		2010/2011 (0%)	2009/2010 (0%)
					2012/2013 (0%)	2010/2011 (0%)
Low Storm Activity (2% of all storms)						2012/2013 (0%)
						2015/2016 (0%)
	2003/2004 (1.8%)	2000/2001 (2.0%)	2002/2003 (3.5%)	2000/2001 (1.5%)		
	2009/2010 (1.3%)	2002/2003 (3.2%)	2004/2005 (1.6%)	2005/2006 (1.7%)		
	2015/2016 (1.4%)	2010/2011 (1.5%)	2010/2011 (1.4%)	2008/2009 (1.6%)		
High Storm Activity (8% of all storms)		2015/2016 (1.3%)		2009/2010 (2.3%)		
				2010/2011 (1.4%)		
	1994/1995 (6.9%)	2006/2007 (9.2%)	1994/1995 (7.3%)	1996/1997 (8.1%)	1997/1998 (4.5%)	
	1998/1999 (6.8%)	2007/2008 (8.0%)	1996/1997 (8.4%)		1998/1999 (4.4%)	
	2006/2007 (10.2%)	2013/2014 (7.9%)	2003/2004 (6.4%)		2011/2012 (7.2%)	
Very High Storm Activity (10+% of all storms)	2007/2008 (8.6%)		2007/2008 (8.7%)		2013/2014 (7.2%)	
	2013/2014 (8.0%)					
	1999/2000 (11.6%)	1996/1997 (11.4%)	2006/2007 (10.6%)	1999/2000 (10.1%)	1996/1997 (9.0%)	1996/1997 (10.3%)
		1999/2000 (13.9%)	2013/2014 (9.0%)	2003/2004 (7.9%)	1999/2000 (15.3%)	1999/2000 (15.5%)
				2004/2005 (8.7%)		2001/2002 (9.0%)
						2007/2008 (9.3%)
						2011/2012 (9.2%)
						2013/2014 (11.8%)

Table 4. Summary of the mean difference (± 1 standard deviation) between XWS or C3S and the input data sets within our database for the proportion of all storms per winter and the proportion of total normalized storm losses per winter. Differences are computed as the absolute value of the difference between XWS or C3S and the CLIMK-WINDS input data sets for each winter, and averaged across all winters.

		ERA5	CCLM_ERA5_EUR-11	COSMO-REA6	CCLM_ERA5_CEU-3
Proportion of All Storms	XWS	2.16% ($\pm 1.89\%$)	2.16% ($\pm 1.81\%$)	2.99% ($\pm 2.38\%$)	3.03% ($\pm 2.37\%$)
Per Winter	C3S	2.36% ($\pm 2.05\%$)	2.48% ($\pm 1.86\%$)	2.85% ($\pm 2.34\%$)	3.09% ($\pm 2.86\%$)
Proportion of Total Normalized Losses	XWS	2.50% ($\pm 2.19\%$)	2.22% ($\pm 2.44\%$)	2.79% ($\pm 2.92\%$)	2.78% ($\pm 2.36\%$)
Per Winter	C3S	1.83% ($\pm 1.45\%$)	1.89% ($\pm 1.36\%$)	2.31% ($\pm 2.19\%$)	2.69% ($\pm 2.38\%$)

Table 5. Summary of the 25th percentile value, median value, and 75th percentile value of the normalized losses (relative ranks) over all the common storms for each input data set, and over the common storms available from XWS and C3S. All quantities are unitless.

	ERA5	CCLM_ERA5_EUR-11	COSMO-REA6	CCLM_ERA5_CEU-3	XWS	C3S
25 th Percentile	0.15	0.18	0.19	0.16	0.17	0.36
Median	0.31	0.27	0.37	0.24	0.24	0.44
75 th Percentile	0.46	0.39	0.47	0.31	0.48	0.55




Figure 1. 98th percentile of the daily maximum wind gusts and spatial extent of the input data sets, shown as shaded areas. In clockwise order of the panels: ERA5, CCLM_ERA5_EUR-11, CCLM_ERA5_CEU-3, and COSMO-REA6. Since the CCLM_ERA5_CEU-3 data set spans only an enlarged Germany domain, its map covers a smaller geographical extent compared to the other three panels.

KYRILL Storm Footprint




Figure 2. Storm footprints, as represented by the locations for which the relative wind gust was greater than 1.0, for windstorm KYRILL (Jan 2007) for the four input data sets used here. For XWS and C3S it was not possible to compute or obtain the relative wind gusts. All footprints are plotted at their native horizontal resolution and for their native domain.




Figure 3. The daily maximum wind gusts (m s^{-1}) for windstorm KYRILL (Jan 2007) at each location associated with the storm footprints displayed in Fig. 2, or which were provided by the XWS and C3S databases. C3S wind gusts were plotted after masking has been performed (see text). All absolute wind gusts are plotted at their native horizontal resolution and for their native domain.




Figure 4. The percentage of extreme storms (y-axis) that occurred during each extended winter season (ONDJFM; x-axis) for each input data set (solid-colored lines), and XWS and C3S (dashed-colored lines) over the 20-year period covering the 1994/1995–2015/2016 extended winter seasons. The end-cap extended winter seasons 1994/1995 and 2015/2016 exclude storms that occurred in the years 1994 and 2016. Percentages are computed relative to the total number of extreme storms for each data set: 50 storms for ERA5, CCLM_ERA5_EUR-11, COSMO-REA6, and CCLM_ERA5_CEU-3; 23 for XWS; and 30 for C3S. No bars indicate that no extreme storms were identified in a given year and data set. Asterisks mark winters not covered by XWS.




Figure 5. The percentage of total normalized loss (y-axis) that was incurred per extended winter season (ONDJFM; x-axis) for each input data set (solid-colored lines), and XWS and C3S (dashed-colored lines) over the 20-year period covering the 1994/1995 – 2015/2016 extended winter seasons. The end-cap extended winter seasons 1994/1995 and 2015/2016 exclude storms that occurred in the years 1994 and 2016. Asterisks mark winters not covered by XWS.




Figure 6. The OEP/AEP ratio (y-axis; unitless) per extended winter season (ONDJFM; x-axis) for each input data set (solid-colored lines), and XWS and C3S (dashed-colored lines) over the 20-year period covering the 1994/1995 – 2015/2016 extended winter seasons. The ratio varies between undefined (no bar plotted for that data) for winters with no storm activity to a maximum of 1.0 for winters containing only a single storm. The end-cap extended winter seasons 1994/1995 and 2015/2016 exclude storms that occurred in the years 1994 and 2016. Asterisks mark winters not covered by XWS.




Figure 7. Bar plots indicating the normalized loss (x-axis; unitless) per common storm (y-axis) for each of the input data sets, and for XWS and C3S, when available. Storms are sorted into chronological descending order by normalized loss according to the ERA5 data set, with and separated into the most-recent Top 10 common storms at the (top panel), and least-recent at the remaining common storms (bottom panel). Common-Asterisks mark storms only that fall outside the XWS time period.

Common Storm Footprint Mean Difference




Figure 8. Mean storm footprint difference computed over the common storms only ~~between-, comparing~~ CCLM_ERA5_CEU-3 ~~and, in~~ ~~clockwise order, with~~ the ERA5, CCLM_ERA5_EUR-11, and COSMO-REA6 data sets ~~(shown in clockwise order across the panels)~~. The colors represent the mean difference in the relative wind (unitless). All panels are plotted at the CCLM_ERA5_CEU-3 horizontal resolution and on the enlarged Germany domain.




Figure 9. Mean difference in daily maximum wind gusts associated with the footprints computed over the common storms only between comparing CCLM_ERA5_CEU-3 and, in clockwise order, with the ERA5, CCLM_ERA5_EUR-11, XWS, C3S, and COSMO-REA6 data sets (shown in clockwise order across the panels). The colors represent the mean difference in the absolute wind gusts (ms^{-1}). All panels are plotted at the CCLM_ERA5_CEU-3 horizontal resolution and on the enlarged Germany domain.

KYRILL Storm Footprint Differences




Figure 10. Storm footprint for windstorm KYRILL (Jan 2007) derived from the CCLM_ERA5_CEU-3 data set and shown in Fig. 2, displayed in the top left panel. The three remaining panels display the difference between the CCLM_ERA5_CEU-3 footprint and the footprints derived from ERA5, COSMO-REA6, and CCLM_ERA5_EUR-11 after regridding to the CCLM_ERA5_CEU-3 resolution, in clockwise order. Colors for the CCLM_ERA5_CEU-3 footprint are the same as in Figure 2, while the colors representing the footprint differences displayed in the remaining three panels are defined by the color bar below the plot. All panels are plotted at the CCLM_ERA5_CEU-3 horizontal resolution. KYRILL is the strongest storm in our database.




Figure 11. The daily maximum wind gusts (ms^{-1}) for windstorm KYRILL (Jan 2007) associated with its footprint as derived from CCLM_ERA5_CEU-3 and shown in Fig. 3, displayed here in the top left panel. The five remaining panels display the difference in ms^{-1} between the CCLM_ERA5_CEU-3 absolute wind gusts and the wind gusts derived from ERA5, XWS, C3S, COSMO-REA6, and CCLM_ERA5_EUR-11 after regridding to the CCLM_ERA5_CEU-3 resolution, in clockwise order. Colors for the CCLM_ERA5_CEU-3 wind-gusts are the same as in Figure 3, while the colors representing the wind-gust differences displayed in the remaining five panels are defined by the color bar below the plot. All panels are plotted at the CCLM_ERA5_CEU-3 horizontal resolution. KYRILL is the strongest storm in our database.

ANDREA Storm Footprint Differences




Figure 12. Storm footprint for windstorm ANDREA (Jan 2012) derived from the CCLM_ERA5_CEU-3 data set in the top left panel. The three remaining panels display the difference between the CCLM_ERA5_CEU-3 footprint and the footprints derived from ERA5, COSMO-REA6, and CCLM_ERA5_EUR-11 after regridding to the CCLM_ERA5_CEU-3 resolution, in clockwise order. Colors for the CCLM_ERA5_CEU-3 footprint are the same as in Figure 2, while the colors representing the footprint differences displayed in the remaining three panels are defined by the color bar below the plot. All panels are plotted at the CCLM_ERA5_CEU-3 horizontal resolution. ANDREA is one of the stronger storms in our database.




Figure 13. The daily maximum wind gusts (ms^{-1}) for windstorm ANDREA (Jan 2012) associated with its footprint as derived from CCLM_ERA5_CEU-3, displayed here in the top left panel. The five remaining panels display the difference in ms^{-1} between the CCLM_ERA5_CEU-3 absolute wind gusts and the wind gusts derived from ERA5, XWS, C3S, COSMO-REA6, and CCLM_ERA5_EUR-11 after regridding to the CCLM_ERA5_CEU-3 resolution, in clockwise order. Colors for the CCLM_ERA5_CEU-3 wind gusts are the same as in Figure 3, while the colors representing the wind gust differences displayed in the remaining five panels are defined by the color bar below the plot. All panels are plotted at the CCLM_ERA5_CEU-3 horizontal resolution. ANDREA is one of the stronger storms in our database.