



Smoothed monthly Greenland ice sheet elevation changes during 2003-2023

Shfaqat A. Khan^{*1}, Helene Seroussi², Mathieu Morlighem³, William Colgan⁴, Veit Helm⁵, Gong Cheng³, Danjal Berg¹, Valentina R. Barletta¹, Nicolaj K. Larsen⁶, William Kochtitzky⁷, Michiel van den Broeke⁸, Kurt H. Kjær⁶, Andy Aschwanden⁹, Brice Noël¹⁰, Jason E. Box⁴, Joseph A. MacGregor¹¹, Robert S. Fausto⁴, Kenneth D. Mankoff^{12,13}, Ian M. Howat¹⁴, Kuba Oniszk¹, Dominik Fahrner⁴, Anja Løkkegaard⁴, Eigil Y. H. Lippert¹, Javed Hassan¹

¹DTU Space, Technical University of Denmark, Denmark

²Thayer School of Engineering, Dartmouth College, Hanover, NH, USA

³Department of Earth Sciences, Dartmouth College, Hanover, NH, USA

⁴Department of Glaciology and Climate, Geological Survey of Denmark and Greenland, Denmark

⁵Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung, Bremerhaven, Germany

⁶Globe Institute, University of Copenhagen, 1350 Copenhagen, Denmark

⁷School of Marine and Environmental Programs, University of New England, Biddeford, ME, USA

⁸Institute for Marine and Atmospheric Research, Utrecht University, The Netherlands

⁹University of Alaska Fairbanks, Fairbanks, AK, USA

¹⁰Laboratoire de Climatologie et Topoclimatologie, University of Liège, Liège, Belgium.

¹¹Cryospheric Sciences Lab, NASA Goddard Space Flight Center, Greenbelt, MD, USA

¹²NASA Goddard Institute for Space Studies, New York, NY, 10025 USA

¹³Autonomic Integra LLC, New York, NY, 10025 USA

¹⁴Byrd Polar Research Center and School of Earth Sciences, Ohio State University, Columbus, OH, USA

Correspondence to: Shfaqat Abbas Khan (abbas@space.dtu.dk)

Abstract. The surface elevation of the Greenland Ice Sheet is constantly changing due to the interplay between surface mass balance processes and ice dynamics, each exhibiting distinct spatiotemporal patterns. Here, we employ satellite and airborne altimetry data with fine spatial (1 km) and temporal (monthly) resolutions to document this spatiotemporal evolution from January 2003 to August 2023. To estimate elevation changes of the Greenland Ice Sheet (GIS), we utilize radar altimetry data from CryoSat-2 and EnviSat, laser altimetry data from the ICESat and ICESat-2, and laser altimetry data from NASA's Operation IceBridge Airborne Topographic Mapper. We produce continuous monthly ice surface elevation changes from January 2003 to August 2023 on a 1 km grid covering the entire GIS. We estimate cumulative ice loss of 4,352 Gt ± 315 Gt (12.1 ± 0.9 mm sea level equivalent) during this period, excluding peripheral glaciers. Between 2003 and 2023, the ice sheet land-terminating margin underwent a significant cumulative thinning of several meters. Ocean-terminating glaciers exhibited thinning between 20–40 m, with Jakobshavn Isbræ experiencing an exceptional thinning of nearly 70 m. This dataset of fine-



40 resolution altimetry data in both space and time will support studies of ice mass loss and useful for GIS ice sheet modelling.
To validate our monthly mass changes of the Greenland ice sheet, we use mass change from satellite gravimetry and mass
change from the Input-Output method. On multiannual timescales, there is a strong correlation between the time series, with
R values ranging from 0.88 to 0.92.

1 Introduction

Over the last three decades, satellite-based observations have revealed unprecedented details regarding the Greenland Ice
45 Sheet's (GIS) mass balance and its response to a warming climate. This wealth of satellite data has not only allowed for the
quantification of mass loss, but also offered insights into the complex interactions between atmospheric, oceanic, and
glaciological processes influencing the ice sheet's response (*Box et al., 2022; Khan et al., 2022a; Sasgen et al., 2020; van
den Broeke et al., 2016; Wood et al., 2021*). The Gravity Recovery and Climate Experiment (GRACE) and GRACE Follow-
On (FO) missions have played a crucial role in this endeavor, revealing a significant ice loss of $4,550 \pm 784$ gigatons (Gt)
50 (equivalent to 12.6 ± 2.2 mm of sea level equivalent, SLE) during the period from 2002 to 2019 (*Velicogna et al., 2020*). The
input-output method (IOM), which generates the longest continuous time series of mass change among the most commonly
used methods, indicates an accelerated mass loss for the GIS over the past four decades (*Mouginot et al., 2019; Mankoff et
al., 2021*). Notably, IOM permits separation of the total mass loss into its component processes, which indicates that ice
discharge remained relatively constant from 1972 to 2002 and thereafter escalated due to the acceleration of multiple outlet
55 glaciers.

Satellite and airborne altimetry, on the other hand, present a direct measurement approach for tracking changes in the ice
sheet thickness, expressed as changes in surface elevation at a finer spatial resolution of a few kilometers. Previous altimetry
studies have relied on observations from a single satellite mission or the fusion of multi-sensor data to estimate trends over
varying time intervals, typically ranging between 1 to 10 years (*Bamber and Dawson, 2020; Csatho et al., 2014; Gardner et
60 al., 2013; Helm et al., 2014; Hurkmans et al., 2014; Schenk et al., 2014; Shepherd et al., 2020; Benjamin Smith et al., 2021;
Benjamin Smith et al., 2019; Sorensen et al., 2018; Winstrup et al., 2024*). Few studies have produced sub-annual elevation
change estimates. Smith et al. (2023) used ICESat-2 to measure the net surface-height change of the GIS at 3-month
resolution to validate surface-height differences predicted by three combinations of climate- and firn-densification models.
Slater et al. (2021) used CryoSat-2 satellite altimetry during 2011-2020 to produce direct measurements of Greenland's
65 runoff variability, based on seasonal changes in the ice sheet's surface elevation. However, they relied on average values
over the entire ablation zone.

Typically, the spatial and temporal resolution of elevation change products is constrained by the resolution of satellite
ground tracks. Satellite altimetry generally provides 2-5 repeat measurements per year over the same location. To achieve
higher temporal resolution, such as monthly measurements, elevation changes must be averaged over a large area, resulting
70 in coarse spatial resolution. Alternatively, combining observations with a model representing ice surface changes can




75 term trends.

2. Input Data


2.1 Radar and laser altimetry data

To estimate the monthly mass changes of the Greenland Ice Sheet (GIS) from January 2003 to August 2023, we utilize the following datasets: (1) radar altimetry data from ESA's CryoSat-2 mission (Wingham et al., 2006), (2) radar altimetry data
80 from ESA's Environmental Satellite (EnviSat) (Rémy et al., 2015), (3) laser altimetry data from the Ice, Cloud, and land Elevation Satellite (ICESat) (Zwally et al., 2014), (4) laser altimetry data from the Ice, Cloud, and land Elevation Satellite-2 (ICESat-2) (Smith et al., 2021), and (5) laser altimetry data from NASA's Operation IceBridge Airborne Topographic Mapper (ATM) flights (Studinger, 2020) (see Figure 1).




85 **Figure 1. Temporal coverage of the airborne and satellite altimetry missions used to estimate monthly elevation changes.**

CryoSat-2: We incorporate all available CryoSat-2 data covering the Greenland ice sheet from November 2010 to August 2023. Data processing follows Khan et al. (2022), utilizing overlapping ground tracks to generate surface elevation time series. Points with series shorter than three years were excluded. The spatial coverage of CryoSat-2 elevation time series is illustrated in Figure 2, showing sparse data in regions with steep slopes, such as the central west Greenland terminus of
90 Jakobshavn Isbræ (Figure 2c). Overall, we use approximately 1.024 billion single point measurements to create about 10.5 million point time series.



95 **Figure 2.** (a) Spatial coverage of points with elevation time series from CryoSat-2. (b) Elevation time series in northeast Greenland. (c) Elevation time series at Jakobshavn Isbræ. The colorbar denotes ice surface elevation.

EnviSat: We use all available EnviSat data from August 2002 to March 2012, processed similarly to CryoSat-2 data. Overlapping ground tracks are employed to create surface elevation time series, excluding points with series shorter than
100 three years. In total, we utilize about 328.6 million single point measurements to generate approximately 2.0 million point time series (figure 3a).



105 **Figure 3. Spatial coverage of points with elevation time series from (a) EnviSat, (b) ICESat, and (c) ICESat-2.**

ICESat: We include all available ICESat data from February 2003 to September 2009 (Schenk and Csatho, 2012; Smith et al., 2020; Zwally et al., 2014), specifically using GLAS/ICESat L2 Global Antarctic and Greenland Ice Sheet Altimetry Data (HDF5), Version 34 (Zwally, 2014). Points with overlapping ground tracks are used to create surface elevation time series, excluding those shorter than three years. This results in the use of about 303.8 million single point measurements to produce approximately 3.4 million point time series. Figure 3b shows the coverage of ICESat point time series.

ICESat-2: All available ICESat-2 data from November 2010 to August 2023 is utilized, specifically ATLAS/ICESat-2 L3A Land Ice Height, Version 6 (Smith et al., 2023). Following the same method as ICESat and CryoSat-2, we use overlapping ground tracks to create surface elevation time series, excluding points with series shorter than three years. This results in about 3.209 billion single point measurements to create approximately 6.1 million point time series. Figure 3 illustrates the spatial coverage of ICESat, ICESat-2, and EnviSat data, showing that laser altimetry (ICESat, ICESat-2) better covers the ice sheet margins compared to radar altimetry (EnviSat, CryoSat-2).



120




Figure 4. (a) Spatial coverage of points with elevation time series from NASA's ATM flight. Elevation time series at northeast Greenland (b), Kangerlussuaq Glacier (c), Jakobshavn Isbræ (d), and Helheim Glacier (e).

NASA's ATM flights: To enhance data coverage near the ice margin, we supplement satellite altimetry with airborne
125 altimetry. We use all available laser altimetry data from NASA's Operation IceBridge ATM flights from April 2002 to April
2019 (Studinger, 2020), utilizing overlapping ground tracks to create surface elevation time series and excluding those
shorter than three years. This results in about 175.1 million single point measurements to create approximately 0.7 million
point time series. Figure 4a shows the coverage of ATM point time series. Notably, several main outlet glaciers, including
the main flowlines of Helheim Glacier (HG), Jakobshavn Isbræ (JI), Kangerlussuaq Glacier (KG), and the northeast
130 Greenland Ice Stream (NEGIS), were repeatedly overflown during 2003-2019.



3. Methods

135 3.1 Improvement compared to previous study

Here, we improve the method used by *Khan et al.* (2022a) and provide continuous monthly surface elevation changes during 2003–2023 on a 1x1 km grid covering the entire ice sheet. While *Khan et al.* (2022a) estimated elevation changes over a ten-year period (2011–2022) using annual mean elevation changes, our approach extends the temporal coverage to twenty years (2003–2023) and introduces monthly mean elevation changes. This expanded temporal range and improved monthly
140 resolution allow for a more detailed capture of fine-scale elevation dynamics. Additionally, this study incorporates ICESat and Envisat data, enhancing the comprehensiveness of the observational dataset. Here, we apply a revised seasonal function to more accurately capture the unequal distribution of surface thickening and thinning over the year, with approximately eight months of surface elevation increases and four months of decreases, closely aligning with observed seasonal trends. Together with recent sub-annual elevation change estimates, this dataset provides an essential foundation for detecting and
145 understanding short-term mass loss fluctuations on local and regional scales, as well as their influence on long-term trends.

3.2 Monthly elevation changes from ICESat

We use ICESat to estimate monthly elevation changes from February 2003 to September 2009 (*Zwally et al.*, 2014). To estimate elevation changes over the ice sheet we follow the procedure described by *Khan et al.* (2022a). We employ a regular grid with a 1x1 km resolution that spans the entire GIS. The center of each grid point is denoted as $C(x_0, y_0)$. For
150 every grid point, we select all ICESat data at coordinates $P(x_i, y_i, h_i, t_i)$, within 1000 m of C , where P includes elevation values h_i measured at time t_i . The index i denotes each specific data point.

Utilizing all available ICESat data collected between February 2003 to September 2009 (*Zwally et al.*, 2014), we generate surface elevation time series at each grid point C . To depict surface changes, we employ a 7th-order polynomial to characterize temporal elevation changes and a 3rd-order polynomial equation to describe the surface shape. Additionally, a
155 seasonal term is incorporated to address seasonal surface variations. For each grid point with the center at (x_0, y_0) , we identify the nearest data point within a 1000 m radius (x_i, y_i, h_i, t_i) and apply the 7th-order polynomial $H(t_i)_{poly}$, the 3rd-order surface topography H_{topo} , and the seasonal term $H(t_i)_{seasonal}$:

$$H(t_i) = H(t_i)_{poly} + H_{topo} + H(t_i)_{seasonal} \quad (1)$$

The 7th-order polynomial is:

$$160 \quad H(t_i)_{poly} = a_1 + a_2 t_i + a_3 t_i^2 + a_4 t_i^3 + a_5 t_i^4 + a_6 t_i^5 + a_7 t_i^6 + a_8 t_i^7, \quad (2)$$

where t_i is the time when the i -th measurement was observed. For simplicity, we used 1 January 2010 as reference time $t=0$. a_1 to a_8 are polynomial variables.

To describe the surface, we fit a 3rd-order polynomial to the observed elevations over the area of 1x1 km (*Csatho et al.*, 2014):



165
$$H_{topo} = a_9x + a_{10}y + a_{11}x^2 + a_{12}y^2 + a_{13}xy + a_{14}x^3 + a_{15}y^3 + a_{16}xy^2 + a_{17}x^2y, \quad (3)$$

where a_9 to a_{17} are parameters that describe the slope and concavity/convexity of the surface, x and y are coordinates of the ICESat data point, but in a system with x_0 and y_0 as the center, i.e., $x = x_i - x_0$ and $y = y_i - y_0$. The seasonal term is given by:

$$H(t_i)_{s1} = a_{18}\cos(\omega_1 t_i + a_{19}) + 0.5a_{18}\cos(\omega_2 t_i + 2a_{19} + 0.5\pi), \quad (4)$$

170 where a_{18} denotes the amplitude, a_{19} is the phase, and ω_1 and ω_2 are the frequencies of the annual (1 yr) and the semi-annual (0.5 yr) term, respectively.

Most previous studies employ a single cosine function to describe seasonal surface elevation changes at any point on the ice sheet:

$$H(t_i)_{s2} = A_i\cos(\omega_1 t_i + \varphi_i), \quad (5)$$

175 Here, we use the following function in equation 4 to better capture the unequal distribution of surface thickening and thinning over the year. The two functions, $H(t_i)_{s1}$ and $H(t_i)_{s2}$, are displayed in Figure 5.





Figure 5. Seasonal evolution with the black curve showing the seasonal signal used in this study and the red curve showing the cosine function used in previous studies.

180 The SMB and GRACE time series suggest ice mass gain over 8 months, from approximately mid-August to mid-April, and rapid mass loss during the remaining 4 months. To describe this behaviour, we combine annual and semi-annual functions. However, we predefine the parameters of the semi-annual term, so that the function reproduces surface elevation increases for approximately 8 months and surface lowering for 4 months. In principle, fitting an annual and semi-annual signal to surface elevations would require estimating four unknown parameters. However, the temporal resolution of satellite

185 altimetry data is often poor and does not strongly constrain all parameters. For instance, ICESat had a repeat track of 3 months, but many points have only 2-3 observations per year, preventing the separation of the annual and semi-annual signals over the entire ice sheet. Using the above equation, we only need to estimate two unknowns, A and φ .

For each grid point, we create a time series and use least squares adjustment to simultaneously estimate parameters a_1 to a_{19} .



190 **Figure 6. Surface elevation change time series derived from ICESat data for a single point. The location of the point is shown as a red dot on the map of Greenland. The solid red curves show the best-fitting 7th-order polynomial and the seasonal signal corrected for surface topography. The error bars denote observed elevations.**

Figure 6 displays a time series of surface elevation corrected for 3rd-order surface topography. The red curve shows the best-
195 fitting 7th-order polynomial + the seasonal term, $H(t_i)_{s1}$. We fit a polynomial only if the observed time series has a length of >3 yrs. Furthermore, we detect and remove outliers from each time series. We define an outlier as a point outside 2σ (standard deviation) of the residual signal (difference between observed elevation and predicted elevation from the polynomial fit). We use the parameters a_1 to a_{19} for each grid point to estimate elevation changes over the entire GIS for consecutive 1-month periods. Spatial coverage of ICESat data is shown in Figure 3b.

200




Figure 7. (a) Surface elevation change time series derived from ICESat data for same point as in figure 6. The solid curves show the best-fitting polynomial or order 1 to 7, and the seasonal signal. (b) differences between the observed elevations and the polynomial fits.

205

In Figure 7a, we present the best-fitting polynomials of orders 1 through 7, along with the seasonal component. The residuals (figure 7b), defined as the difference between the observed elevation and the polynomial fit, indicate that for polynomials of orders 5 to 7, the residuals are approximately ± 10 cm.

210 **ICESat-2:** Our method of deriving monthly surface elevation changes from ICESat-2 is identical to ICESat.

3.3 Monthly elevation changes from CryoSat-2

Our procedure for processing CryoSat-2 is identical to *Khan et al. (2022a)*. To estimate elevation changes using CryoSat-2 data, we follow the procedure of ICESat and ICESat-2 data, however, with a minor modification regarding the seasonal signal, $H(t_i)_{s1}$. We employ a regular grid with a 1x1 km resolution that spans the entire GIS. Utilizing all available CryoSat-2 data collected between November 2010 to October 2023, we generate surface elevation time series at each grid point.


215

For each grid point, we use point time series to estimate a 7th-order polynomial $H(t_i)_{poly}$, a 3rd-order surface topography H_{topo} , and a seasonal term $H(t_i)_{seasonal}$

Earlier studies (and this study for ICESat and ICESat-2) assume the shape of the surface remains constant (Schenk et al., 2014) throughout the studied period. However, near the ice margin, the shape of the surface may change significantly over



220 the course of 20 years. As a compromise, rather than fitting a polynomial to the entire 2010-2023 period, we consider two individual time sub-intervals separately, the first one between 2010-2017, and the second one between 2017-2023. During each sub-interval we assume the shape of the surface remains constant. This allows the shape of the surface to change over longer intervals.



225




230 **Figure 8. Surface elevation change time series derived from Cryosat-2 data for a single point. The location of the point is shown as a red dot on the map of Greenland. (a) The solid curves show the best-fitting 7th-order polynomial and the seasonal signal corrected for surface topography. The error bars denote observed elevations. (b) The solid curves show the best-fitting 7th-order polynomial corrected for the seasonal signal and surface topography. (c) The solid curves show the best-fitting 7th-order polynomial corrected for surface topography, however, a seasonal signal from ICESat/ICESat-2 is added.**

235 Figure 8a displays a time series of surface elevation corrected for 3rd-order surface topography. The red and blue curve shows the best-fitting 7th-order polynomial + the seasonal term. We fit a polynomial only if the observed time series has a length of >3 yrs. Furthermore, we detect and remove outliers from each time series. We use the parameters a_1 to a_{19} for each grid point and each (sub-)interval, e.g., 2010-2017 and 2017-2023, to estimate elevation changes over the entire GIS for consecutive 1-month periods. Spatial coverage of CryoSat-2 data is shown in Figure 2.

240 CryoSat-2 seasonal signal

245 The radar signal from CryoSat-2 might not be reflected by the snow surface but could instead penetrate through dry snow. There may be minor local discrepancies that may be attributed to the penetration of the radar signal into the snow. To address this issue, we remove a seasonal signal estimated from CryoSat-2, $H(t_i)_{seasonal,cryo}$, from point time series, and substitute it with a seasonal signal, $H(t_i)_{seasonal,ice1/ice2}$, estimated from ICESat and ICESat-2. The stepwise procedure is illustrated in Figure 9 and depicted for a point time series in Figure 8.



250 **Figure 9. The stepwise procedure on how a seasonal signal from CryoSat-2 data is replaced with a seasonal signal from ICESat and ICESat-2.**

In this study, we employ ICESat and ICESat-2 to obtain the seasonal signal between 2003-2009 and 2018-2023, respectively. For the 2009-2018 period, (when data from both missions is available), we derive the seasonal signal averaged from ICESat and ICESat-2.


255 **3.4 Monthly elevation changes from EnviSat**

To estimate elevation changes using EnviSat data, we follow the procedure of CryoSat-2 data. We employ a regular grid with a 1x1 km resolution that spans the entire GIS. Utilizing all available EnviSat data collected between August 2002 to March 2012, we generate surface elevation time series at each grid point. For each grid point, we use point time series to estimate a 7th-order polynomial $H(t_i)_{poly}$, a 3rd-order surface topography H_{topo} , and a seasonal term $H(t_i)_{seasonal,Envi}$

260 Next, we remove a seasonal signal estimated from EnviSat, $H(t_i)_{seasonal,envi}$, from point time series, and substitute it with a seasonal signal, $H(t_i)_{seasonal,ice1/ice2}$, estimated from ICESat and ICESat-2. Figure 3a shows spatial coverage of EnviSat time series.



3.5 Amplitude of the Seasonal signal



265

Figure 10. Amplitude of seasonal surface elevation changes from (a) ICESat, (b) ICESat-2 (c) CryoSat-2, and (d) Envisat.

The seasonal signal amplitudes derived from ICESat, ICESat-2, CryoSat-2, and Envisat satellite missions, are shown in Figure 10. Predominantly, the highest amplitudes are observed in southeast Greenland across all satellite missions. It is important to note that Envisat data is not inclusive of the ice sheet margin. The slight variations in amplitudes can be attributed, in part, to the differing time spans utilized for amplitude estimation.

270

In this study, we use seasonal signal from ICESat and ICESat-2. For the 2009-2018 period, we derive mean seasonal signal from ICESat and ICESat-2. Figure 10(e-f) show standard deviation of the amplitude at each grid point, i , from ICESat and ICESat-2, $\sigma_{i,icesat,amp}$ and $\sigma_{i,icesat2,amp}$, respectively. For the 2009-2018 period, we use standard error based on mean ICESat and ICESat-2 amplitudes.

275



The seasonal layer is the top layer on the ice sheet. For this layer we use a constant density of 315 ± 44 kg/m³ to convert snow volume to mass (Fausto et al., 2018).

3.6 Monthly elevation changes from NASA's Operation IceBridge ATM flights

280 During 2002-2019, NASA conducted annual airborne surveys with ATM over the GIS during the spring. These flights were mostly concentrated along the margins of the GIS. To estimate monthly changes, we use the same approach as used for satellite altimetry data described above. However, spring data alone do not allow us to extract a seasonal signal. Therefore, $H(t_i)_{seasonal}$ is not estimated for ATM data. Instead, we adopt a seasonal signal estimated from ICESat and ICESat-2 data. In addition, to allow the shape of the surface to change, we divide the study period into two separate (sub-)intervals, i.e.,
285 2002-2010 and 2010-2019. During each (sub-) interval we assume the shape of the surface remains constant. Figure 9 shows Spatial coverage of the ATM time series.

3.7 Monthly elevation changes and their uncertainty

For each grid point, we use point time series to estimate 7th-order polynomial, a 3rd-order surface topography, and a seasonal term from a synthesis of several sensors' datasets is somewhat different from other studies (Nilsson et al., 2022;
290 Simonsen et al., 2021). While other studies typically estimate the bias between sensors/missions, we do not merge data from different satellite missions and therefore do not estimate any biases. Instead, we estimate monthly elevation changes for each sensor's dataset independently as described in previous sections, and only merge the estimated monthly change rates from each dataset afterwards, when creating the multi-sensor monthly grid. Figure 11 illustrates this process for June 2008, for which we merged monthly elevation changes from NASA's ATM data, ICESat data and Envisat data. To detect local
295 outliers of monthly rates, we used planar regression in 20 km bins. Values that fell outside the 5- σ range are excluded. The estimated monthly elevation changes were used to interpolate elevation change rates onto a regular grid of 1x1 km. For each grid point, we use ordinary kriging to interpolate elevation change rates $dh_{i,krig}$ and the associated error $\sigma_{i,krig}$ (Khan et al., 2022a).

300




Figure 11. Surface elevation change during June 2008 from (a) ATM flights, (b) ICESat laser satellite altimetry, (c) EnviSat radar altimetry, (d) combined elevation changes from airborne and satellite data, (e) interpolated elevation changes onto a 1x1 km grid, and (f) uncertainties of the interpolated data.

305 3.8 Correction for elastic VLM, glacial isostatic adjustment, and firn compaction and associated uncertainties

3.8.1 Glacial Isostatic Adjustment

The observed ice surface elevation changes were corrected for bedrock movement caused by elastic vertical land motion (VLM) in response to present-day mass changes and long-term past ice changes (Glacial Isostatic Adjustment, GIA). To



310 correct for GIA, we use the GNET-GIA empirical model of Khan et al. (2016). For each grid point on a 0.5x0.5 km grid, we estimate the GIA uplift rate dh_{GIA} and the associated uncertainty σ_{GIA} retrieved from Khan et al. (2016). The GIA correction is between -3 and +16 mm/yr and the associated uncertainty, σ_{GIA} , is between 0 and 3 mm/yr (see figure 3 of Khan et al 2016). This correction of vertical land motion corresponds to total mass loss correction of 2.8 ± 1.2 Gt/yr or 56 Gt over 20 years.

3.8.2 Elastic vertical land motion

315 We correct for elastic VLM of the bedrock by convolving monthly mass loss estimates with the Green's functions derived by Wang et al. (2012) for elastic Earth model with refined crustal structure from Crust 2.0. (Laske et al. ,2012). For each grid point, we estimate the elastic uplift rate dh_{elas} and the associated uncertainty σ_{elas} . The uncertainties, σ_{elas} , are estimated by convolving uncertainties of monthly mass loss estimates with the Green's functions for elastic Earth model. The mean annual elastic correction varies between -2 to +37 mm/yr and the associated uncertainties are between 0 and 2 mm/yr. The average elastic correction over from 2003 to 2023 is 6.5 ± 0.4 Gt/yr.

320

3.8.3 Converting volume to mass

325 We convert volume to mass as described in Khan et al. (2022b). Conversion of the volume loss rate into the mass loss rate requires assumptions about density; therefore, using a constant ice sheet density would be inaccurate. Firn compaction must be taken into account to convert volume to mass correctly. Hence, elevation changes due to firn compaction are modeled with a simple firn model that includes melt and refreezing. It is forced by annual temperature, accumulation, melt, and refreezing from the regional climate model RACMO2.3p2 (Noël et al., 2018) at 5.5 km horizontal resolution. For each grid point, i , we estimate the firn compaction rate $dh_{i,firm}$ and the associated uncertainty $\sigma_{i,firm}$ as described by Khan et al 2022b. Uncertainties are estimated as described in Kuipers Munneke et al. (2015) (see their equations 8 and 9) using input fields from RACMO2.3p2. The average elastic correction over from 2003 to 2023 is 15.8 ± 2.4 Gt/yr.

330

Elevation change:

335 The interpolation was performed using the ordinary kriging method (Hurkmans et al., 2014; Nielsen et al., 2013). We first used the observed annual elevation change rates to estimate an empirical semi-variogram. Next, we fit an exponential model variogram for each 1-year interval (with a range of 65 km) to the empirical semi-variogram to take the spatial correlation of elevation change rates into account in the error budget. For each grid point, we interpolate (using kriging) elevation change rate $dh_{i,krig}$ and the associated error $\sigma_{i,krig}$.

The total elevation change rate for each grid point i is:

$$dh_i = dh_{i,krig} - dh_{i,elas} - dh_{i,GIA} - dh_{i,firm} \quad (6)$$



340 Assuming the uncertainty in each of these terms is independent, we estimate the total associated uncertainty by summing each uncertainty term in quadrature:

$$\sigma_i = \sqrt{\sigma_{i,krig}^2 + \sigma_{i,elas}^2 + \sigma_{i,GIA}^2 + \sigma_{i,firm}^2} \quad (7)$$

The total GIS mass change error for each month, t_n , is

$$\sigma_{month}(t_n) = \sum_{i=1}^{i_{max}} \sigma_i \quad (8)$$

345 Where i_{max} is the total number of grid cells that cover the GIS.

We generate a time series of cumulative GIS mass change by integrating our monthly time series of mass change over time.


We estimate the cumulative errors as the root sum square of errors,

$$\sigma_{cumul}(t) = \sqrt{\sum_{n=1}^{n_{max}} \sigma_{month}^2(t_n)} \quad (9)$$

350 Where n_{max} is the total number of months from January 2003 to July 2023 and is set to $n_{max}=247$ months for this study.

4. Result

Satellite and airborne altimetry (denoted as altimetry in figure 12) indicate an ice sheet mass loss of $4,352 \pm 315$ Gt (12.1 ± 0.9 mm SLE) from January 2003 through August 2023, excluding peripheral glaciers (PG). Our results are similar to previous studies (Bevis et al., 2019; Mankoff et al., 2021; Sasgen et al., 2020; Velicogna et al., 2020) suggesting enhanced ice loss during the summer months of 2010, 2011, 2012, and 2019 (Figure 12 and figure 13). Ice loss was slower in 2013-2018 and was followed by an increased ice loss rate during 2020–2023.



360 **Figure 12.** Time series of cumulative monthly ice mass change of the GIS from January 2003 to August 2023 in gigatons (left axis) and sea level rise equivalent (right axis). The purple curve displays GIS+PG mass change from satellite gravimetry adopted from Barletta et al. (2020). Brown curve shows GIS+PG mass change from IMBIE. The blue curve shows GIS mass change from the Input-Output method from Mankoff et al., (2021) extended to august 2024. The grey curve shows GIS mass change from satellite and airborne altimetry from this study. The shadings represent the associated uncertainties. Corrections for GIA (black dashed line), elastic deformation (red dashed line), and firn compaction (blue dashed line).

365 Time series of cumulative monthly ice mass change of the GIS from January 2003 to August 2023 along with the correction of GIA and elastic induced vertical land motion and firn compaction are displayed in figure 12.





Figure 13. Maps of annual elevation change rates from 2003 to 2023 from satellite and airborne altimetry from this study.

5. Validation

370 5.1 Mass change from satellite gravimetry and the Input-Output method

To validate our monthly Greenland ice sheet elevation changes, we use mass change from satellite gravimetry from Barlette et al., (2020) (denoted as “gravimetry”), mass change from the Input-Output method from Mankoff et al., (2021) (denoted as “IOM”), and mass change from IMBIE (IMBIE, 2023).



375

380

Figure 14. (a) Cumulative surface elevation change from altimetry from January 2003 through August 2023. Sub-panels show time-series of surface elevation changes (dh) for selected locations on Helheim Glacier (HG), Kangerlussuaq Glacier (KG), east interior GIS (East GIS), Rink Isbræ (RI), Zachariæ Isstrøm (ZI), Nioghalvfjærdsfjorden (NG), Humboldt Glacier (HBG), Upernavik Isstrøm (UI), Jakobshavn Isbræ (JI), and southern interior GIS (South GIS). (b) Cumulative surface elevation change inferred from gravimetry from January 2003 through August 2023. (c) Cumulative thinning inferred from altimetry along the GIS margin during 2003–2023.

385


Satellite gravimetry indicates an ice loss of $5,198 \pm 173$ Gt from January 2003 through August 2023, encompassing both the GIS and peripheral glaciers. The spatial resolution of satellite gravimetry, approximately ~ 300 km, prevents differentiation between the ice sheet and peripheral glaciers. The difference of 846 ± 359 Gt between satellite altimetry and gravimetry approximates the ice loss reported from Greenland's peripheral glaciers (Khan et al., 2022b). The cumulative surface elevation changes inferred by satellite gravimetry (Barletta et al., 2020, Barletta et al., 2024) suggest an increase in ice thickness throughout the high-level interior of Central and North Greenland (Figure 14b). This observation is inconsistent with our findings obtained from satellite altimetry measurements, which suggest that high-level thickening is limited to NE



390 Greenland and the saddle region between the main and south domes of the ice sheet. Satellite gravimetry does not capture the full magnitude of ice thinning around the periphery of the ice sheet due to the low spatial resolution.

The cumulative mass change from the IOM, as reported by Mankoff et al. (2021) and extended to August 2024, falls within the uncertainty range of mass change measured by altimetry (see Figure 12). Figure 12 also shows the cumulative mass change from IMBIE, which shows a larger mass change compared to altimetry due to the inclusion of peripheral glaciers. Overall, the cumulative mass changes from IMBIE, IOM, gravimetry, and altimetry over the past two decades agrees within 395 uncertainties.

To compare multiannual ice mass change variability, we first detrend the cumulative mass changes from IOM, gravimetry, and altimetry. Figure 15a shows the three detrended time series along with the correlation coefficients, R , between altimetry and gravimetry, altimetry and IOM, and IOM and gravimetry. On multiannual timescales, there is a strong correlation between the time series, with R values ranging from 0.88 to 0.92. Figures 15b-d display scatter plots of annual mass change 400 rates comparing altimetry vs. gravimetry, altimetry vs. IOM, and gravimetry vs. IOM. The correlation coefficients for these comparisons range from 0.58 to 0.80, with the best correlation observed between gravimetry and IOM. This is expected, as altimetry provides smoothed elevation changes, whereas IOM and gravimetry can detect rapid changes. In general, while the three methods agree on the total mass loss of the ice sheet, there is less agreement on the precise temporal and short-term distribution of this mass loss.



405

Figure 15: (a) Detrended cumulative mass change from altimetry, gravimetry and IOM. C correlation coefficients, R , between altimetry-gravimetry, altimetry-IOM and IOM-gravimetry. (b) comparison of annual mass change rates between (b) altimetry-gravimetry, (c) altimetry-IOM and (d) gravimetry-IOM. (b-d) axis units are Gt/yr.

410 5.2 Local elevation changes

Our altimetry observations, interpolated to a 1 km grid, enable a detailed examination of GIS mass change at both regional and individual glacier scales. Figure 14 illustrates the total (cumulative) surface elevation change from January 2003 through August 2023 along with point time series at selected outlet glaciers and regions, and cumulative thinning from altimetry around the GIS margin (Figure 14c).

415

A large spatial and temporal variability is observed, with the entirety of West Greenland exhibiting surface lowering extending deep inland. In contrast, the surface elevation of interior Northeast Greenland increased over the past two decades. Examining glacier-specific details, our altimetry time series reveals a net thinning of ~ 70 m near the terminus of Jakobshavn Isbræ. This thinning stopped and this sector underwent thickening during 2016–2018, followed by a return to thinning from 2019 to 2023, a phenomenon examined by *Khazendar et al. (2019)*.

420



Upernavik Isstrøm (UI) has experienced a ~30 m thinning over the past two decades, with intensified thinning 2010 to 2012. Northwest Greenland's Humboldt Glacier (HBG) has thinned by 23 m over the same period, at a constant rate. One of the two major glaciers in northeast Greenland, Nioghalvfjærdsfjorden Glacier (NG), also known as 79 North, exhibits a total thinning of 15 m, while the other, Zachariae Isstrøm (ZI), shows more substantial thinning of about 35 m. In the southeast, Kangerlussuaq Glacier (KG) has thinned by approximately 23 m, while Helheim Glacier (HG) has experienced a 28 m thinning from January 2003 through August 2023. This aligns with the findings of *Williams et al.* (2021), who observed a cumulative surface elevation change of approximately 15 m during the period 2011-2020. The two time series labelled as "East interior GIS" and "South interior GIS" are located in the interior of the ice sheet (about 150 km and 250 km from the ice margin, respectively) and depict a small thickening of a few meters over the past two decades.

6. Discussion

The observed spatial and temporal resolution of ice surface elevation changes from satellite altimetry is of critical importance for better quantifying the processes impacting ice sheet change, and for advancing and refining ice sheet modelling. This level of precision is crucial for accurately representing complex processes within ice sheet models, such as seasonal ice flow dynamics, and changes in surface mass balance (*Goelzer et al.*, 2017). Recent advances in machine learning and automatic differentiation tools enable transient calibration in ice sheet modelling, facilitating better estimation of unmeasurable parameters such as the basal sliding coefficient and ice rheology. The integration of fine-scale data into models enhances their ability to simulate realistic responses to climate variables, contributing to more accurate predictions of ice sheet behaviour and potential impacts on sea-level rise. Additionally, high-resolution satellite altimetry data helps identify local trends and patterns, allowing for a more nuanced understanding of regional variations in ice sheet dynamics (*Mankoff et al.*, 2019; *Mankoff et al.*, 2021). Ultimately, the incorporation of observed high-resolution data into ice sheet models is essential for improving the fidelity of simulations and enhancing our ability to assess the implications of climate change on ice sheet stability and sea-level rise (*Choi et al.*, 2023).

7. Limitations of the method and data

Temporal Resolution: We estimate monthly elevation changes by fitting a 7th-order polynomial and a seasonal signal to the observed elevation time series. This approach, which we refer to as "smoothed" elevation changes, results in smoothed data that cannot detect rapid elevation changes. The method's effectiveness is limited by the temporal resolution of available airborne and satellite altimetry data. For instance, ICESat provides 2-4 repeat measurements per year at the same location,



while ATM data offers only one measurement annually. This means that measurements should not be expected to resolve at high spatial resolution for ice-sheet changes at sub-seasonal time scales, only smoothed sub annual changes using some pre-defined seasonal model are possible.

Spatial Resolution: The spatial resolution of elevation changes estimated from satellite radar altimetry over ice sheet margins is constrained by the coarse spatial footprint of instruments like the EnviSat radar altimeters, leading to a lack of fine-scale detail. The steep and rugged topography of ice sheet margins causes signal scattering and reflection issues, resulting in inaccurate measurements. Figures 2 and 3a illustrate the poor spatial resolution along the ice margin, especially for EnviSat. Another limitation is that NASA's ATM flights cover main outlet glaciers but do not survey all glaciers annually. Generally, NASA's ATM flight coverage is denser in West Greenland than in East Greenland. Additionally, the Operation IceBridge campaign, with measurements conducted once per year, does not allow for the detection of seasonal signals. However, the ICESat and ICESat-2 missions significantly enhance our understanding of ice sheet elevation changes, including in the ice margin regions and seasonal signals.

In general, our method lacks detecting rapid ice elevation changes. For example, drainage of surface lakes on the Greenland Ice Sheet are not detected. A small area with elevation changes of up to tens of meters will be considered as an outlier. We detect local outliers of monthly rates, using planar regression in 20 km bins. Drainage of surface lakes, which often occurs over a small area will be detected as an outlier using our method. Also, Instances of rapid elevation changes occurring over extremely short time spans will go unnoticed through our approach. For instance, rapid accelerations and decelerations, as illustrated by *Vijay et al. (2021)*, taking place within a 2-month period are not fully captured by our surface elevation change product. Nevertheless, our method effectively identifies melt-driven rapid thinning episodes during the warm summers of 2010-2012 and 2019. Furthermore, in Greenland, 87% of the glacierized region terminates in the ocean leading to frontal ablation, involving both ice discharge and terminus retreat. We note that our mass loss estimate from satellite altimetry does not incorporate mass loss below sea level, constituting less than 10-20% of the overall frontal ablation, as indicated by *Kochitzky et al. (2023)* and *Greene et al. (2024)*.

8. Data products and availability

Our main products are gridded time series of monthly Digital Elevation Models and gridded monthly time series of surface elevation change (in water equivalent). However, we also deliver surface elevation change (in ice equivalent). This allows users to convert ice to mass using their own model for e.g. firn compaction. To make the data user friendly for the ice sheet modelling community, we deliver code that can create time series of elevations with respect to geoid model (often needed as input in numerical ice flow model).

To make our elevation change products useful for the ice sheet modeling community, we deliver a Digital Elevation Model (DEM). We use the DEM from the Greenland Ice Mapping Project (GIMP) (Howat et al., 2015), downscaled to a 1x1 km



grid resolution to align with the elevation change products developed in this study. Additionally, we convert it to a reference time of January 1, 2003, using the dh/dt product developed in this study.

485 **Table 1: Data products on a 1x1 km grid covering the Greenland Ice sheet.**

Data product	Spatial Resolution	Temporal Resolution	Temporal Coverage
Digital Elevation Models	1x1 km	One epoch*	2003.00
Monthly Elevation changes (in water equivalent)	1x1 km	Monthly	2003 to 2023
Monthly Elevation changes (in ice equivalent)	1x1 km	Monthly	2003 to 2023
Elastic VLM	1x1 km	Monthly	2003 to 2023
Firm compaction rate	1x1 km	Monthly	2003 to 2023
GIA	1x1 km	One epoch*	2003.00
Geoid model	1x1 km	One epoch**	2003.00

*) we assume vertical land motions due to GIA does not change over a 20 yrs period.

***) we assume the geoid does not change over a 20 yrs period.

All grid files use WGS 84 / NSIDC Sea Ice Polar Stereographic North (EPSG:3413)

Our main products are Digital Elevation Model and gridded monthly time series of surface elevation change (in water equivalent). However, we also deliver surface elevation change (in ice equivalent). This allows users to convert ice to mass
490 using their own model for e.g. firm compaction or GIA. To make the data user friendly for the ice sheet modelling community, we deliver geoid model (often needed as input in numerical ice flow model).

We provide a code snippet to estimate the parameters for equations 2 to 4 using the test input data.

495 **File naming and data format.**

We provide 2 zip files.

File 1: Greenland_geotiff_1kmgrid.zip

File 2: Greenland_netcdf_1kmgrid.zip

500

file 1 format: NetCDF (monthly dh/dt data)

file 2 format: geotiff (mean dh/dt from 2003-2023)

File 1 content: This is a zip file that contains the following netcdf files.

505

- **Greenland_DEM_1kmgrid.nc** (Digital Elevation Models)
- **Greenland_dhdt_elas_1kmgrid.nc** (monthly elastic Vertical Land Motion)



- **Greenland_dhdt_elas_err_1kmgrid.nc** (error associated with monthly elastic Vertical Land Motion)
- **Greenland_dhdt_firn_1kmgrid.nc** (monthly firn compaction rate)
- 510 • **Greenland_dhdt_firn_err_1kmgrid.nc** (error associated with monthly firn compaction rate)
- **Greenland_dhdt_GIA_1kmgrid.nc** (Glacial Isostatic Uplift rates)
- **Greenland_dhdt_GIA_err_1kmgrid.nc** (error associated with Glacial Isostatic Uplift rate)
- **Greenland_dhdt_icevol_1kmgrid.nc** (Monthly Elevation changes (in ice equivalent))
- **Greenland_dhdt_icevol_err_1kmgrid.nc** (error associated Monthly Elevation changes (in ice equivalent))
- 515 • **Greenland_dhdt_mass_1kmgrid.nc** (Monthly Elevation changes (in water equivalent))
- **Greenland_dhdt_mass_err_1kmgrid.nc** (error associated with Monthly Elevation changes (in water equivalent))
- **Greenland_geoid_1kmgrid.nc** (geoid height)

Each file contain a header with information about data types.

520 File 2 content: same as file 1 but with geotiff format.

9. Conclusion

The spatial and temporal resolution of elevation change products is limited by the resolution of satellite ground tracks. Satellite altimetry usually offers 2-5 repeat measurements per year over the same location. To achieve higher temporal resolution (e.g., 1 month), elevation changes must be averaged over a large area, which leads to coarse spatial resolution.

525 Alternatively, we model a seasonal signal (see Methods) and generate "smoothed" monthly elevation changes with higher spatial and temporal resolution. This allows us to produce an elevation change product with detailed insights into the dynamics of the Greenland Ice Sheet from January 2003 to August 2023. Our product reveals a cumulative ice loss of $4,352 \pm 315$ Gt, corresponding to a global mean sea level rise of 12.1 ± 0.9 mm over two decades and local thinning of up to 70 m near the terminus of JI. Validation of our monthly mass changes of the Greenland ice sheet, against mass change from

530 satellite gravimetry and the Input-Output method, suggests strong correlation with R values ranging from 0.88 to 0.92. Incorporating our new high-resolution data into ice sheet models can enhance our understanding of ice dynamics, thus improving predictions about the Greenland Ice Sheet's response to climate change and its impact on sea-level rise. Despite certain limitations, especially in capturing short, rapid elevation changes, our methodology offers a valuable tool for monitoring and analyzing the evolving dynamics of the Greenland Ice Sheet.

535



540 **Author contributions.**

SAK conceived the study, analysed most of the data. VH processed Cryosat-2 data. VB analysed GRACE and GRACE-FO data. KDM provided and prepared IOM data. Writing by SAK, HS, MM, WC, GC, DB, VRB, NKL, WK, MvdB, KHK, AA, BN, JEB, JAM, RSF, KDM, IMH, KO, DF, AL, EYHL, and JH. Project administration by SAK, WC, HS and NKL. IMH developed DEM model.

545

Competing interests.

Author Kenneth D. Mankoff is a member of the editorial board of the journal.

Acknowledgements

550 S A Khan, H Seroussi, M Morlighem, W Colgan, G Cheng, K Kjær, N K Larsen, and A Løkkegaard acknowledge support from the NOVO Nordisk foundation grant no NNF23OC00807040. This is a publication of the Center for Ice Sheet and Sea Level Predictions (CISP). B Noël was funded by the Fonds de la Recherche Scientifique de Belgique (F.R.S.-FNRS). R.S Fausto and J.E. Box acknowledges support from the Programme for monitoring of the Greenland ice sheet (PROMICE). K. Mankoff was supported by the NASA Modeling Analysis and Prediction program.

555

Data and code Availability

Monthly elevation change rates of the GIS from 2003 to 2023 is available at the following data repository <https://datadryad.org/stash/share/RfBPIGTZRn0Vs0u8hj1PSgide11IPAL-k8TvEOOEbDA> (Khan et al., 2024a). The DTU GRACE and GRACE-FO Greenland Mass Balance product is available at <https://doi.org/10.11583/DTU.12866579.v4> (Barletta et al., 2020). Code is available at <https://doi.org/10.5281/zenodo.13276108> (Khan et al., 2024b)

560

References

Bamber, J. L., and G. J. Dawson, Complex evolving patterns of mass loss from Antarctica's largest glacier, *Nature Geoscience*, 13(2), 127-131, doi:<https://doi.org/10.1038/s41561-019-0527-z>, 2020.

565 Barletta, V. R., L. S. Sørensen, and R. Forsberg, Scatter of mass changes estimates at basin scale for Greenland and Antarctica, *The Cryosphere*, 7(5), 1411-1432, doi:<https://doi.org/10.5194/tc-7-1411-2013>, 2013.

Barletta, V. R., L. S. Sørensen, and R. Forsberg, Gravitational mass balance of Greenland 2003 to present v2.1, <https://doi.org/10.11583/DTU.12866579.v4>, 2020.

570



- Barletta, V. R., Bordoni, A., & Khan, S. A., GNET derived mass balance and glacial isostatic adjustment constraints for Greenland. *Geophysical Research Letters*, 51, e2023GL106891. <https://doi.org/10.1029/2023GL106891>, 2024
- 575 Bevis, M., et al., Accelerating changes in ice mass within Greenland, and the ice sheet's sensitivity to atmospheric forcing, *Proceedings of the National Academy of Sciences*, 116(6), 1934-1939, doi:<https://doi.org/10.1073/pnas.1806562116>, 2019.
- Box, J. E., et al., Greenland ice sheet climate disequilibrium and committed sea-level rise, *Nature Climate Change*, 12(9), 808-813, doi:<https://doi.org/10.1038/s41558-022-01441-2>, 2022.
- 580 Brils, M., P. Kuipers Munneke, and M. R. van den Broeke, Spatial Response of Greenland's Firn Layer to NAO Variability, *Journal of Geophysical Research: Earth Surface*, 128(8), e2023JF007082, doi:<https://doi.org/10.1029/2023JF007082>, 2023.
- Caron, L., E. R. Ivins, E. Larour, S. Adhikari, J. Nilsson, and G. Blewitt, GIA Model Statistics for GRACE Hydrology, Cryosphere, and Ocean Science, *Geophysical Research Letters*, 45(5), 2203-2212, 585 doi:<https://doi.org/10.1002/2017GL076644>, 2018.
- Choi, Y., H. Seroussi, M. Morlighem, N. J. Schlegel, and A. Gardner, Impact of time-dependent data assimilation on ice flow model initialization and projections: a case study of Kjer Glacier, Greenland, *The Cryosphere*, 17(12), 5499-5517, doi:<https://doi.org/10.5194/tc-17-5499-2023>, 2023. 590
- Csatho, B. M., A. F. Schenk, C. J. van der Veen, G. Babonis, K. Duncan, S. Rezvanbehbahani, M. R. van den Broeke, S. B. Simonsen, S. Nagarajan, and J. H. van Angelen, Laser altimetry reveals complex pattern of Greenland Ice Sheet dynamics, *Proceedings of the National Academy of Sciences*, 111(52), 18478-18483, doi:<https://doi.org/10.1073/pnas.1411680112>, 2014. 595
- Fausto, R. S., et al., A Snow Density Dataset for Improving Surface Boundary Conditions in Greenland Ice Sheet Firn Modeling, *Frontiers in Earth Science*, 6, doi:<https://doi.org/10.3389/feart.2018.00051>, 2018.
- 600 Fettweis, X., J. E. Box, C. Agosta, C. Amory, C. Kittel, C. Lang, D. van As, H. Machguth, and H. Gallée, Reconstructions of the 1900–2015 Greenland ice sheet surface mass balance using the regional climate MAR model, *The Cryosphere*, 11(2), 1015-1033, doi:<https://doi.org/10.5194/tc-11-1015-2017>, 2017.
- Gardner, A. S., et al., A Reconciled Estimate of Glacier Contributions to Sea Level Rise: 2003 to 2009, *Science*, 340(6134), 852-857, doi:<https://doi.org/doi:10.1126/science.1234532>, 2013.



- 605 Goelzer, H., A. Robinson, H. Seroussi, and R. S. W. van de Wal, Recent Progress in Greenland Ice Sheet Modelling, *Current Climate Change Reports*, 3(4), 291-302, doi:<https://doi.org/10.1007/s40641-017-0073-y>, 2017.
- Greene, C. A., A. S. Gardner, M. Wood, and J. K. Cuzzone, Ubiquitous acceleration in Greenland Ice Sheet calving from 1985 to 2022, *Nature*, 625(7995), 523-528, doi:<https://doi.org/10.1038/s41586-023-06863-2>, 2024.
- 610 Hanna, E., J. M. Jones, J. Cappelen, S. H. Mernild, L. Wood, K. Steffen, and P. Huybrechts, The influence of North Atlantic atmospheric and oceanic forcing effects on 1900–2010 Greenland summer climate and ice melt/runoff, *International Journal of Climatology*, 33(4), 862-880, doi:<https://doi.org/10.1002/joc.3475>, 2013.
- 615 Helm, V., A. Humbert, and H. Miller, Elevation and elevation change of Greenland and Antarctica derived from CryoSat-2, *The Cryosphere*, 8(4), 1539-1559, doi:<https://doi.org/10.5194/tc-8-1539-2014>, 2014.
- Hurkmans, R. T. W. L., J. L. Bamber, C. H. Davis, I. R. Joughin, K. S. Khvorostovsky, B. S. Smith, and N. Schoen, Time-evolving mass loss of the Greenland Ice Sheet from satellite altimetry, *The Cryosphere*, 8(5), 1725-1740, doi:<https://doi.org/10.5194/tc-8-1725-2014>, 2014.
- 620 Howat, I., A. Negrete, and B. Smith, MEaSURES Greenland Ice Mapping Project (GIMP) Digital Elevation Model, Version 1.. Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Active Archive Center. <https://doi.org/10.5067/NV34YUIXLP9W>. [Date Accessed: 1 June 2024], 2015.
- 625 Khan, S. A., et al., Greenland Mass Trends From Airborne and Satellite Altimetry During 2011–2020, *Journal of Geophysical Research: Earth Surface*, 127(4), e2021JF006505, doi:<https://doi.org/10.1029/2021JF006505>, 2022a.
- Khan, S. A., W. Colgan, T. A. Neumann, M. R. van den Broeke, K. M. Brunt, B. Noël, J. L. Bamber, J. Hassan, and A. A. Bjørk, Accelerating Ice Loss From Peripheral Glaciers in North Greenland, *Geophysical Research Letters*, 49(12), e2022GL098915, doi:<https://doi.org/10.1029/2022GL098915>, 2022b.
- 630 Khan, S. A., et al., Geodetic measurements reveal similarities between post–Last Glacial Maximum and present-day mass loss from the Greenland ice sheet, *Science Advances*, 2(9), e1600931, doi:<https://doi.org/10.1126/sciadv.1600931>, 2016.
- 635 Khan, Shfaqat Abbas et al., Smoothed monthly Greenland ice sheet elevation changes during 2003–2023 [Dataset]. Dryad. <https://datadryad.org/stash/share/RfBPIGTZRn0Vs0u8hj1PSgje11IPAL-k8TvE00EBDA>, <https://doi.org/10.5061/dryad.s4mw6m9dh> (doi link not activated yet by DRYAD), 2024a.



- 640 Khan, S. A.. Matlab code related to paper: Smoothed monthly Greenland ice sheet elevation changes during 2003-2023 by
Khan et al, 2024. (in ESSD). In Smoothed monthly Greenland ice sheet elevation changes during 2003-2023.
Zenodo. <https://doi.org/10.5281/zenodo.13276108>, 2024b.
- 645 Khazendar, A., et al., Interruption of two decades of Jakobshavn Isbrae acceleration and thinning as regional ocean cools,
Nature Geoscience, 12(4), 277-283, doi:<https://doi.org/10.1038/s41561-019-0329-3>, 2019.
- Kochtitzky, W., L. Copland, M. King, R. Hugonnet, H. Jiskoot, M. Morlighem, R. Millan, S. A. Khan, and B. Noël, Closing
Greenland's Mass Balance: Frontal Ablation of Every Greenlandic Glacier From 2000 to 2020, *Geophysical Research
Letters*, 50(17), e2023GL104095, doi:<https://doi.org/10.1029/2023GL104095>, 2023.
- 650
- Laske, G., Masters, G., Reif, C., CRUST2.0: A New Global Crustal Model at 2_2 Degrees.
http://igppweb.ucsd.edu/_gabi/crust2.html, 2012.
- Mankoff, K. D., et al., Greenland Ice Sheet solid ice discharge from 1986 through 2017, *Earth Syst. Sci. Data*, 11(2), 769-
655 786, doi:<https://doi.org/10.5194/essd-11-769-2019>, 2019.
- Mankoff, K. D., et al., Greenland ice sheet mass balance from 1840 through next week, *Earth Syst. Sci. Data*, 13(10), 5001-
5025, doi:<https://doi.org/10.5194/essd-13-5001-2021>, 2021.
- 660 Mouginit, J., E. Rignot, A. A. Bjørk, M. van den Broeke, R. Millan, M. Morlighem, B. Noël, B. Scheuchl, and M. Wood,
Forty-six years of Greenland Ice Sheet mass balance from 1972 to 2018, *Proceedings of the National Academy of Sciences*,
116(19), 9239-9244, doi:<https://doi.org/10.1073/pnas.1904242116>, 2019.
- Nilsson, J., A. Gardner, L. Sandberg Sørensen, and R. Forsberg, Improved retrieval of land ice topography from CryoSat-2
665 data and its impact for volume-change estimation of the Greenland Ice Sheet, *The Cryosphere*, 10(6), 2953-2969,
doi:<https://doi.org/10.5194/tc-10-2953-2016>, 2016.
- Nilsson, J., A. S. Gardner, and F. S. Paolo, Elevation change of the Antarctic Ice Sheet: 1985 to 2020, *Earth Syst. Sci. Data*,
14(8), 3573-3598, doi:<https://doi.org/10.5194/essd-14-3573-2022>, 2022.
- 670
- Noël, B., W. J. van de Berg, S. Lhermitte, and M. R. van den Broeke, Rapid ablation zone expansion amplifies north
Greenland mass loss, *Science Advances*, 5(9), eaaw0123, doi:<https://doi.org/10.1126/sciadv.aaw0123>, 2019.



675 Noël, B., et al., Modelling the climate and surface mass balance of polar ice sheets using RACMO2 – Part 1: Greenland (1958–2016), *The Cryosphere*, 12(3), 811-831, doi:<https://doi.org/10.5194/tc-12-811-2018>, 2018.

680 Rémy, F., T. Flament, A. Michel, and D. Blumstein, Envisat and SARAL/AltiKa Observations of the Antarctic Ice Sheet: A Comparison Between the Ku-band and Ka-band, *Marine Geodesy*, 38(sup1), 510-521, doi:<https://doi.org/10.1080/01490419.2014.985347>, 2015.

Sasgen, I., B. Wouters, A. S. Gardner, M. D. King, M. Tedesco, F. W. Landerer, C. Dahle, H. Save, and X. Fettweis, Return to rapid ice loss in Greenland and record loss in 2019 detected by the GRACE-FO satellites, *Communications Earth & Environment*, 1(1), 8, doi:<https://doi.org/10.1038/s43247-020-0010-1>, 2020.

685 Schenk, T., B. Csatho, C. van der Veen, and D. McCormick, Fusion of multi-sensor surface elevation data for improved characterization of rapidly changing outlet glaciers in Greenland, *Remote Sensing of Environment*, 149, 239-251, doi:<https://doi.org/10.1016/j.rse.2014.04.005>, 2014.

690 Shepherd, A., et al., Mass balance of the Greenland Ice Sheet from 1992 to 2018, *Nature*, 579(7798), 233-239, doi:<https://doi.org/10.1038/s41586-019-1855-2>, 2020.

695 Simonsen, S. B., V. R. Barletta, W. T. Colgan, and L. S. Sørensen, Greenland Ice Sheet Mass Balance (1992–2020) From Calibrated Radar Altimetry, *Geophysical Research Letters*, 48(3), e2020GL091216, doi:<https://doi.org/10.1029/2020GL091216>, 2021.

Slater, T., et al., Increased variability in Greenland Ice Sheet runoff from satellite observations, *Nature Communications*, 12(1), 6069, doi:<https://doi.org/10.1038/s41467-021-26229-4>, 2021.

700 Smith, B. Medley, X. Fettweis, T. Sutterley, P. Alexander, D. Porter, and M. Tedesco, Evaluating Greenland surface-mass-balance and firn-densification data using ICESat-2 altimetry, *The Cryosphere*, 17(2), 789-808, doi:<https://doi.org/10.5194/tc-17-789-2023>, 2023.

705 Smith, B., B. M. Adusumilli, D. Csathó, H. A. Felikson, A. Fricker, N. Gardner, J. L. Holschuh, F. S. Paolo, M. R. Siegfried, and T. S. a. t. I.-S. Team, ATLAS/ICESat-2 L3A Land Ice Height, *Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Active Archive Center*, 5, doi:<https://doi.org/10.5067/ATLAS/ATL06.005>, 2021.



- Smith, B., et al., Land ice height-retrieval algorithm for NASA's ICESat-2 photon-counting laser altimeter, *Remote Sensing of Environment*, 233, 111352, doi:<https://doi.org/10.1016/j.rse.2019.111352>, 2019.
- 710 Sørensen, L., S. B. Simonsen, R. Forsberg, K. Khvorostovsky, R. Meister, and M. E. Engdahl, 25 years of elevation changes of the Greenland Ice Sheet from ERS, Envisat, and CryoSat-2 radar altimetry, *Earth and Planetary Science Letters*, 495, 234-241, doi:<https://doi.org/10.1016/j.epsl.2018.05.015>, 2018.
- Studinger, M., Updated 2020. IceBridge ATM L2 icesn elevation, slope, and roughness, version 2, *Boulder, Colorado, USA, NASA National Snow and Ice Data Center Distributed Active Archive Center*,
715 doi:<https://doi.org/10.5067/CPRXXK3F39RV>, 2020.
- van den Broeke, M. R., E. M. Enderlin, I. M. Howat, P. Kuipers Munneke, B. P. Y. Noël, W. J. van de Berg, E. van
720 Meijgaard, and B. Wouters, On the recent contribution of the Greenland ice sheet to sea level change, *The Cryosphere*, 10(5), 1933-1946, doi:<https://doi.org/10.5194/tc-10-1933-2016>, 2016.
- Velicogna, I., et al., Continuity of Ice Sheet Mass Loss in Greenland and Antarctica From the GRACE and GRACE Follow-On Missions, *Geophysical Research Letters*, 47(8), e2020GL087291, doi:<https://doi.org/10.1029/2020GL087291>, 2020.
- 725 Vijay, S., M. D. King, I. M. Howat, A. M. Solgaard, S. A. Khan, and B. Noël, Greenland ice-sheet wide glacier classification based on two distinct seasonal ice velocity behaviors, *Journal of Glaciology*, 67(266), 1241-1248, doi:<https://doi.org/10.1017/jog.2021.89>, 2021.
- Wang, H., L. Xiang, L. Jia, L. Jiang, Z. Wang, B. Hu, and P. Gao, Load Love numbers and Green's functions for elastic
730 Earth models PREM, iasp91, ak135, and modified models with refined crustal structure from Crust 2.0, *Computers & Geosciences*, 49, 190-199, doi:<https://doi.org/10.1016/j.cageo.2012.06.022>, 2012.
- Williams, J. J., N. Gourmelen, P. Nienow, C. Bunce, and D. Slater, Helheim Glacier Poised for Dramatic Retreat, *Geophysical Research Letters*, 48(23), e2021GL094546, doi:<https://doi.org/10.1029/2021GL094546>, 2021.
- 735 Wingham, D. J., et al., CryoSat: A mission to determine the fluctuations in Earth's land and marine ice fields, *Advances in Space Research*, 37(4), 841-871, doi:<https://doi.org/10.1016/j.asr.2005.07.027>, 2006.



740 Winstrup, M., Rannal, H., Larsen, S. H., Simonsen, S. B., Mankoff, K. D., Fausto, R. S., and Sørensen, L. S.: PRODEM:
Annual summer DEMs (2019–present) of the marginal areas of the Greenland Ice Sheet, *Earth Syst. Sci. Data Discuss.*
[preprint], <https://doi.org/10.5194/essd-2023-224>, in review, 2023.

745 Wood, M., et al., Ocean forcing drives glacier retreat in Greenland, *Science Advances*, 7(1), eaba7282,
doi:<https://doi.org/10.1126/sciadv.aba7282>, 2021.

Zwally, H. J., R. Schutz, J. C. Bentley, T. Bufton, J. Herring, J. S. Minster, and R. Thomas, GLAS/ICESat L2 Antarctic and
Greenland Ice Sheet Altimetry Data, *Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Active
Archive Center, Version 34*, doi:<https://doi.org/10.5067/ICESAT/GLAS/DATA225>, 2014.

750