

Bivalve monitoring over French coasts: multi-decadal records of carbon and nitrogen elemental and isotopic ratios ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$ and C:N) as ecological indicators of global change

5 Camilla Liénart¹, Alan Fournioux¹, Andrius Garbaras², Hugues Blanchet¹, Nicolas Briant³, Stanislas F. Dubois⁴, Aline Gangnery³, Anne Grouhel Pellouin³, Pauline Le Monier³, Arnaud Lheureux⁵, Xavier de Montaudouin¹, Nicolas Savoye¹

¹Université de Bordeaux, CNRS UMR 5805, Bordeaux INP, EPOC, Pessac, 33600, France

²Center for Physical Sciences and Technology, FTMC, Vilnius, 10257, Lithuania

³Ifremer Centre Atlantique, CCEM, Nantes, 44000, France

10 ⁴Ifremer, DYNECO, Plouzané, 29280, France

⁵MNHN, CNRS UMR 8067, SU, IRD 207, UCN, UA, BOREA, CP53, F-75005 Paris, France

Correspondence to: Camilla Liénart (camilla.lienart@u-bordeaux.fr)

Abstract

15 Recent changes in climate and environment, influenced by both global and local factors, have had profound impacts on coastal ecosystem functioning and trajectories. By examining archived samples from ongoing ecological monitoring efforts, particularly focusing on bivalves like mussels and oysters, we gain a valuable long-term perspective on how ecosystems are responding at various scales. We conducted analyses on carbon and nitrogen content (C, N) and elemental and isotopic ratios (C:N, $\delta^{13}\text{C}$, $\delta^{15}\text{N}$) of mussel and oyster soft tissues collected annually at 33 sites along the French coast from 1981 to 2021.

20 This extensive dataset (<https://doi.org/10.17882/100583>, Liénart et al., 2024a) offers a comprehensive view spanning multiple decades and ecosystems, allowing to track how coastal ecosystems and marine species record changing climate, physical-chemical environments and organic matter cycles. Additionally, these data are crucial for establishing isotope baselines for studying food webs. Ultimately, this data set provide valuable information for more effective ecosystem conservation and management strategies in our rapidly changing world.

25 1 Introduction

Over the past two decades, rapid and sometimes abrupt changes in climate and environmental conditions have significantly impacted ecosystems functioning with human activities widely acknowledge as a primary factor driving and modulating these changes (Cloern et al., 2016). At the land-ocean interface, coastal and estuarine environments are particularly vulnerable to the climate change, alongside with local human-induced pressures such as pollution and habitat destruction (Harley et al, 2006; 30 Lotze et al., 2006). In such dynamic environments, pressures vary simultaneously, with different directions and magnitudes, over different time (short vs long-term) and spatial (global vs local) scales (Cabral et al., 2019). Hence, a fundamental challenge persists in elucidating how the various facets of global change affect coastal ecosystems responses at local and regional scales, and consequently the services they provide to human societies. Recurrent ecological monitoring programs conducted at large spatial scales are valuable to assess ecosystem status and the pressures affecting important properties (e.g., biodiversity, carbon 35 and nutrient cycling), and to detect changes over multiple decades (Hofmann et al., 2013; Sukhotin and Berger, 2013). Yet, only a few ecological indicators are both integrative (i.e. measured in long-living species tissues) and sensitive enough (e.g. measured at molecular level such as metallothionein used as a biomarker of metal exposure, Amiard et al., 2006) to various disturbances while also exhibiting predictable responses with a low variability in its response (Dale and Beyeler, 2001; Niemi and McDonald, 2004).

40 Filter-feeding bivalves include important reef habitat forming species, which promote benthic-pelagic coupling and nutrient recycling (Ray and Fulweiler, 2020). Some of them such as mussels (*Mytilus* spp.) and oysters (*Crassostrea gigas*) represent an economical value through aquaculture and are widely used as bioindicators for climate and environmental change as well as contaminants monitoring (Kanduč et al., 2018; Karlson and Faxneld, 2021; Mazaleyrat et al., 2022; Briand et al., 2023; Chahouri et al., 2023; Liénart et al., 2024b). The use of ecological indicators is the basis of ecosystem monitoring to detect 45 early-warning signals in ecosystem changes or disturbances (Dale and Beyeler, 2001). Indeed, bivalve tissues records the environmental conditions of their sampling area over time. Since the end of the 1970s, the French monitoring network “ROCCH” (“Réseau d’Observation de la Contamination CHimique”, coordinated by Ifremer) monitor chemical contamination in the environment along French coastlines by using oysters and mussels as bioindicators (Briant et al., 2018; Chahouri et al., 2023). The ROCCH network focuses on chemical contaminants, but its archived samples represent a large sample bank that 50 can be useful for other ecological purposes, such as understanding long-term changes in ecosystem functioning and responses to global or local pressures.

Measuring elemental and isotopic composition of the main building blocks of life (i.e., carbon and nitrogen; C:N, $\delta^{13}\text{C}$, $\delta^{15}\text{N}$) is relevant to understand nutrients origin, to determine organisms diet and food webs structure, and can serve as indicators of water quality over space and time (Glibert et al., 2018). The C:N ratio is mostly an indicator of bivalve condition and 55 physiology, reflecting the balance between organisms’ requirements and elemental availability in the environment (i.e., ecological stoichiometry sensu Elser et al., 2003; N content increases (thus C:N ratio decreases) as protein content increases whereas C content increases (thus C:N ratio increases) as lipid or carbohydrate content increases). When measured over long-

term in bivalve tissues, such dataset allow scientists to understand ecosystems responses to changing physical-chemical environment and organic matter cycling and to track trends in climate changes and its effect on coastal ecosystems (Liénart et al., 2020, 2024b). We took advantage of the large-scale multi-decadal sample bank of the ROCCH to analyze C and N content and C:N, $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ ratios in mussels (*M. edulis*, *M. galloprovincialis*) and oysters (*C. gigas*) tissues from a set of 33 stations distributed along the French coasts. In this paper, we present a unique dataset of multi-decadal and multi-ecosystem carbon and nitrogen content and elemental and isotopic ratios from three widespread bivalve species. As long-term isotopic datasets reflect the impact of human activities, such as coastal pollution and habitat alteration, the data derived here can provide valuable information for conservation and management strategies for coastal areas, in helping make informed decisions to mitigate environmental threats and protect vulnerable ecosystems. Ultimately it could provide valuable input for developing predictive models of bivalve physiology (Emmery et al., 2011) or trophic ecology (Marín Leal et al., 2008) explaining ecosystem response to future environmental changes and possibly forecast potential impacts of climate change and human activities on coastal ecosystems. Overall, this long-term dataset of isotopic ratio in suspension-feeders tissues provides insights into ecosystem dynamics and is essential for advancing scientific understanding in the face of ongoing environmental challenges.

2 Methods and data

2.1 The ROCCH: network and sampling procedure

Over the last four decades, the French national monitoring network for chemical contaminant, the “ROCCH” (“Réseau d’Observation de la Contamination CHimique”), coordinated by Ifremer (“Institut français de recherche pour l’exploitation de la mer”, <https://littoral.ifremer.fr/Reseaux-de-surveillance/Environnement/ROCCH-Reseau-d-Observation-de-la-Contamination-CHimique-du-littoral>) has been annually sampling bivalves as bioindicators of chemical contamination. The number of stations has varied since the initiation of the network, so has the sampling frequency. Approximately 150 stations are nowadays monitored along the French coastlines, which begins in 1979 for historical stations. Sampling took place once a year during winter, currently mid-February, with a tolerance of one tidal cycle before and after the target date, meaning a 6 weeks amplitude spreading from the end of January to the beginning of March on the overall sites of the network. Three different species are targeted, the Pacific oyster *Crassostrea gigas* and the blue mussels *Mytilus edulis* in the English Channel and Atlantic facades, and *M. galloprovincialis* in the Mediterranean facade.

The sampling protocol has been identical since the start of the monitoring and was designed to acquire bivalve samples with consistent and homogeneous characteristics for measuring chemical concentrations of contaminants. Bivalves are collected alive at fixed points (maximum tolerance of 180 m around the selected point) chosen away from known anthropogenic discharges. Bivalves are sourced either from wild beds or from dedicated rearing facilities, ensuring that they remain on site for at least 6 months before being sampled. The selected individuals are adults of the same species and of uniform size (30 to 60 mm long for mussels, 90 to 140 mm long for oysters, i.e., 2 to 3 years old). A minimum of 50 mussels or 10 oysters was

90 required to constitute a representative pooled sample accounting for inter-individual variability and to get enough material for
chemical analysis and long-term storage. Bivalves are first depurated for 18 to 26 hours in decanted seawater collected near
the collection site. Next, once extracted from the shell, the whole bodies (i.e., total soft-tissues) of each individual are pooled
and drained all together for 30 minutes. The resulting pooled bivalve samples are placed in clean glass containers (washed and
baked for 8 hours at 450 °C) covered with calcined aluminum foil, and sealed with a plastic lid and frozen (-20 °C). Frozen
95 samples are sent to the central laboratory in Nantes (France) where the samples are thawed, grounded and homogenized in a
stainless steel-bladed blender and freeze-dried. After chemical analysis of a sample aliquot, the containers are indefinitely
stored at Nantes IFREMER center, at room temperature, protected from light, under moisture-regulation control to prevent
moisture pick-up. The original protocol (Grouhel 2023, in French) is available at <https://doi.org/10.13155/97878>.


2.2 Study sites and bivalve dataset

100 In the present dataset, bivalve samples were selected from 33 stations of the ROCCH network distributed along the three
French sea facades: the English Channel (E.C., 10 stations), the Atlantic Ocean (A.O., 17 stations) and the Mediterranean Sea
(M.S., 6 stations; Figure 1, Table 1). The stations were selected to span over a wide diversity of geomorphological and
environmental conditions encountered by bivalves along the French coastlines. Ecosystems vary from open-sea coasts (rocky
or sandy littoral shores), open or semi-enclosed bays and rias, lagoon (shallow water bodies almost closed by narrow
105 landforms), and estuary mouths, ranging from eutrophic to oligotrophic systems, under temperate oceanic or Mediterranean
climates, and along gradients of river influence from the main French rivers (Seine, Loire, Garonne, Dordogne, Rhône; Table
2). The complete dataset consist in 1141 bivalve winter samples for time series spanning 11 to 40 years (depending on stations)
over the period 1981-2021 (Table 1; <https://doi.org/10.17882/100583>, Liénart et al., 2024a).

2.3 Sample analysis and data quality

110 Aliquots of 400-700 µg of each dry and grounded archived sample of pooled bivalve tissues (not acidified) were analyzed for
carbon and nitrogen elemental and isotopic composition (C, N, $\delta^{13}\text{C}$, $\delta^{15}\text{N}$) at the Center for Physical Science and Technology
(Vilnius, Lithuania) with a Flash EA 1112 Series Elemental Analyzer (Thermo Finnigan) connected to a DeltaV Advantage
Isotope Ratio Mass Spectrometer (Thermo Fisher). Carbon and nitrogen elemental content are expressed in percent (%) of
mass ratio (g g^{-1}) dry weight, while C:N ratio is express in mol mol^{-1} . Isotope ratios are expressed using the conventional delta
115 notation: $\delta^{13}\text{C}_{\text{sample}}$ or $\delta^{15}\text{N}_{\text{sample}} = [(R_{\text{sample}}/R_{\text{standard}}) - 1]$, where $R = {}^{13}\text{C}/{}^{12}\text{C}$ or ${}^{15}\text{N}/{}^{14}\text{N}$, in per mil deviation (‰) from
international reference, Vienna Pee Dee Belemnite for $\delta^{13}\text{C}$ and atmospheric N_2 for $\delta^{15}\text{N}$. Mass spectrometer was calibrated
against external certified standards (Caffeine IAEA600: $\delta^{13}\text{C}$ -27.77 ‰; $\delta^{15}\text{N}$ 1.00 ‰, Graphite USGS24: $\delta^{13}\text{C}$ -16.05 ‰), and
internal standards (casein: $\delta^{13}\text{C}$ -23.30 ‰; $\delta^{15}\text{N}$ 6.30 ‰, glycine: $\delta^{13}\text{C}$ -45.20 ‰; $\delta^{15}\text{N}$ 3.00 ‰) controlled using sucrose IAEA-
CH-6, ammonium sulfate IAEA-N-1 and IAEA-N-2, Graphite USGS24 were added before and every 10 samples within each
120 batch of samples to control analytical performance.

Analytical precision was always better than 0.2 ‰ for $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ and 0.1 mol mol⁻¹ for C:N ratio (median of the standard deviations of each standard used for all batches: 0.12 ‰ for $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ and 0.04 mol mol⁻¹ for C:N ratio). Each sample was usually analyzed once and for some peculiar values that seemed too high/low or outside of the average trend, samples were reanalyzed to verify the values. If values were similar (< 0.5 ‰), we chose the average value of both analyses; otherwise, we
125 chose the value associated with the best analytical precision. The dataset presented here consists in a single value for a given year at a given station (the standard deviation associated with repeated measures is not presented). One data point was considered as an outlier in regard of its carbon isotope value (reanalysis confirmed the value): the station 14-PALL in 2014 (Figure 2) exhibited unexpected low values of $\delta^{13}\text{C}$ (-26.96 ‰) and high $\delta^{15}\text{N}$ (10.26 ‰) compare to the average values of this station ($\delta^{13}\text{C} = -18.71 \pm 0.75$ ‰; $\delta^{15}\text{N} = 8.41 \pm 0.93$ ‰), suggesting potential issues in the sampling and/or storing. This value
130 was therefore removed from average calculations and statistical tests. Within the C and N % datasets, there was a few exceptionally high/low values (Figure 2) for the stations 1-AMBL in 1990 (low C % only), 5-CLHV in 1994 (high C and N %), 18-BOUR in 1990 (low C and N %), 20-RIVE in 1989 (high C and N %) and 30-STMM in 2016 (high C and N %). These values must be considered with caution when interpreting or using C and N % data and were excluded from statistical calculations (see section 2.4). Nevertheless, these data pairs all led to consistent C:N ratio at these sites and years and were
135 kept for C:N statistical analyses. All the other values presented in this data set are considered analytically valid and were scientifically validated in regard of expert knowledge of each ecosystem.



140 **Figure 1: Map of the 33 stations from the three sea facades of France analyzed for bivalve's elemental and isotopic ratios. Bivalve species are the mussels *Mytilus edulis* and *Mytilus galloprovincialis* and the oyster *Crassostrea gigas*. The main rivers are presented in blue. Basemap source: Esri, HERE, © OpenStreetMap contributors, and the GIS User Community.**

150 **Table 1: Metadata description for the 33 stations analyzed for bivalve's carbon and nitrogen elemental and isotopic ratios. Species studied are the mussels *Mytilus edulis* and *Mytilus galloprovincialis* and the oyster *Crassostrea gigas*. Station 31-CART represent two stations merged over time (distant from 720m). The geographic coordinate system is WGS84.**

Sea Facade	Station ID	Station name	ROCCH codes	latitude	longitude	Species	Time series	n years	Missing years
English Channel	1-AMBL	Ambleteuse	002-P-032	50.807366667	1.595466667	<i>M. edulis</i>	1982-2021	36	1984, 1985, 1986, 1990
	2-PSQT	Pointe de St Quentin	006-P-009	50.2808	1.52853333	<i>M. edulis</i>	1988-2021	34	-
	3-VARE	Varengeville	008-P-013	49.9207688241	0.9820028963	<i>M. edulis</i>	1982-2021	37	1984, 1985, 1986
	4-ANTI	Antifer - digue	010-P-014	49.649085114	0.1503265738	<i>M. edulis</i>	1983-2021	36	1984, 1985, 1986
	5-CLHV	Cap de la Hève	010-P-055	49.5074135756	0.0619953224	<i>M. edulis</i>	1983-2017	29	1984, 1986, 1987, 1998, 1999, 2000
	6-VILL	Villerville	011-P-005	49.4040781596	0.1236660472	<i>M. edulis</i>	1982-2021	36	1984, 1985, 1986
	7-OUIS	Ouistreham	010-P-120	49.2940709028	-0.24800513	<i>M. edulis</i>	1994-2021	27	2004
	8-PBES	Port en Bessin	013-P-001	49.3515665687	-0.7531834463	<i>M. edulis</i>	1981-2021	37	1983, 1984, 1988, 2007
	9-BDVG	Bdv Grandcamp ouest	014-P-007	49.38633333	-1.101266667	<i>M. edulis</i>	1981-2021	37	1984, 1985, 1992, 1996
	10-PLAN	Pen al Lann	034-P-001	48.665109	-3.8944	<i>C. gigas</i>	1982-2021	37	1984, 1985, 1986
Atlantic Ocean	11-PASS	Le Passage (b)	039-P-007	48.391067	-4.384965	<i>C. gigas</i>	2001-2021	21	-
	12-PERS	Persuel	039-P-093	48.2938273007	-4.5500595819	<i>C. gigas</i>	2001-2021	21	-
	13-AULN	Aulne rive droite	039-P-124	48.281083	-4.260048	<i>C. gigas</i>	1982-2021	38	1984, 1985
	14-GUIL	Le Guilvin	060-P-001	47.56765	-2.9338	<i>C. gigas</i>	1982-2021	39	1986
	15-PEBE	Pen Bé	066-P-003	47.4306639649	-2.4679991263	<i>M. edulis</i>	1982-2021	40	-
	16-CHEM	Pointe de Chemoulin	070-P-102	47.234632	-2.297076	<i>M. edulis</i>	1981-2021	39	1982, 1983
	17-JOAL	Joalland (b)	070-P-006	47.1568267957	-2.2224877474	<i>C. gigas</i>	2010-2021	11	-
	18-BOUR	Bourgneuf - Coupelasse	071-P-065	47.0123255923	-2.0229805005	<i>C. gigas</i>	1984-2021	37	1985
	19-NOIR	Noirmoutier - Gresse-loup	071-P-068	46.95066	-2.146303	<i>C. gigas</i>	1983-2021	38	1985
	20-RIVE	Rivedoux	076-P-032	46.163319	-1.27077	<i>C. gigas</i>	1981-2021	40	1986
	21-PALL	Les Palles	080-P-004	45.9674774781	-1.1414394441	<i>C. gigas</i>	1982-2021	39	1983
	22-BOYV	Boyardville	080-P-033	45.9638097847	-1.2259409199	<i>C. gigas</i>	1984-2021	38	-
	23-PONT	Pontailac	084-P-015	45.6251372031	-1.0560967723	<i>C. gigas</i>	1984-2021	38	-
	24-FOSS	La Fosse	085-P-007	45.475634584	-0.9845921143	<i>C. gigas</i>	1981-2021	40	2018
	25-FERR	Cap Ferret	087-P-013	44.6439465896	-1.2412452987	<i>C. gigas</i>	1982-2021	39	2005
	26-JACQ	Les Jacquets	088-P-067	44.7222821428	-1.1945794691	<i>C. gigas</i>	1982-2021	38	1983, 1985
	27-COMP	Comprian	088-P-069	44.6839491697	-1.08457666	<i>C. gigas</i>	1982-2021	40	-
Mediterranean Sea	28-BANY	Banyuls - Labo Arago	094-P-008	42.4806130241	3.1388780428	<i>M.galloprovincialis</i>	1981-2021	32	1982, 1986, 1987, 1988, 1989, 1990, 2003
	29-HRLT	Embouchure de l'Hérault	095-P-026	43.2756325016	3.4405360127	<i>M.galloprovincialis</i>	1985-2021	31	1986, 1987, 1997, 1998, 2016
	30-STMM	Les Stes Maries de la mer	106-P-018	43.4439790562	4.4205519936	<i>M.galloprovincialis</i>	1982-2021	38	1984, 1986
	31-CART	Anse de Carteau	109-P-025 / -027	43.3756486632	4.8755622202	<i>M.galloprovincialis</i>	1982-2021	37	1986, 1996, 2007
	32-COUR	Cap Courronne	111-P-002	43.3239826009	5.0539002233	<i>M.galloprovincialis</i>	1983-2021	37	1986, 1996
	33-POME	Pomègues Est	111-P-025	43.2673170789	5.3005726112	<i>M.galloprovincialis</i>	1998-2021	24	-

Table 2: Environmental characteristics of the 33 stations studied for bivalves elemental and isotopic ratios. Average values for salinity, water temperature and chlorophyll a were calculated, when available, over the studied period from monitoring stations located nearby bivalves sampling stations, with data were retrieved from Surval (<https://surval.ifremer.fr> - Ifremer) and Somlit (<https://www.somlit.fr/> - INSU) databases. Average annual flow rates of the main rivers were calculated over the study period from the HydroPortail database (<https://hydro.eaufrance.fr/>). Trophic status was defined for each station based on literature (Liéart et al., 2017; Lheureux et al., 2023) and the knowledge of local experts and reflect the average status over the studied period.

Sea Facade	Station ID	Ecosystem	Ecosystem type	Tidal range	Salinity (mean±sd)	Water temperature (°C) (mean±sd)	Chlorophyll <i>a</i> (µg L ⁻¹) (mean±sd)	Trophic status	Main river influence	Annual flow rates (m ³ s ⁻¹) (mean±sd)	Additional local river influence
English Channel	1-AMBL	Eastern English Channel	Littoral	Megatidal	34 ± 1,0	12,7 ± 4,4	4,8 ± 4,6	Eutroph	Seine	499 ± 332	Liane
	2-PSQT	Eastern English Channel	Littoral	Megatidal	33 ± 1,5	13,0 ± 4,7	8,2 ± 7,8	Eutroph	Seine	449 ± 332	Somme
	3-VARE	Eastern English Channel	Littoral	Megatidal	33 ± 1,1	13,4 ± 4,5	2,2 ± 2,3	Eutroph	Seine	449 ± 332	-
	4-ANTI	Bay of Seine	Open bay	Macrotidal	32 ± 1,8	14,3 ± 4,3	4,8 ± 7,1	Eutroph	Seine	449 ± 332	-
	5-CLHV	Bay of Seine	Estuary mouth	Macrotidal	29 ± 3,5	12,9 ± 4,4	4,0 ± 4,7	Eutroph	Seine	449 ± 332	-
	6-VILL	Bay of Seine	Estuary mouth	Macrotidal	29 ± 3,5	12,9 ± 4,4	4,0 ± 4,7	Eutroph	Seine	449 ± 332	-
	7-OUIS	Bay of Seine	Open bay	Macrotidal	33 ± 0,8	15,2 ± 4,2	5,4 ± 5,2	Eutroph	Seine	449 ± 332	Orne
	8-PBES	Bay of Seine	Open bay	Macrotidal	34 ± 0,5	13,8 ± 4,4	2,8 ± 2,9	Eutroph	Seine	449 ± 332	-
	9-BDVG	Bay of Veys	Open bay	Macrotidal	33 ± 0,7	13,8 ± 4,2	2,6 ± 2,5	Eutroph	Seine	449 ± 332	Vire
	10-PLAN	Western English Channel	Ria	Macrotidal	35 ± 0,2	12,9 ± 2,2	0,9 ± 0,8	Mesotroph	-	-	-
Atlantic Ocean	11-PASS	Bay of Brest	Semi-enclosed ria	Macrotidal	27 ± 7,5	12,9 ± 4,1	NA ± NA	Mesotroph	Elorn	6 ± 5	-
	12-PERS	Bay of Brest	Semi-enclosed ria	Macrotidal	35 ± 0,6	13,4 ± 2,8	1,1 ± 0,9	Mesotroph	Aulne/Elorn	-	-
	13-AULN	Bay of Brest	Semi-enclosed ria	Macrotidal	23 ± 8,7	12,8 ± 4,4	2,3 ± 2,7	Mesotroph	Aulne	27 ± 29	-
	14-GUIL	Morbihan Gulf	Semi-enclosed bay	Macrotidal	33 ± 2	14,4 ± 3,7	1,8 ± 2,0	Mesotroph	Auray	2,8 ± 3,1	-
	15-PEBE	Bay of Vilaine	Semi-enclosed bay	Macrotidal	32 ± 3	14,8 ± 3,9	4,6 ± 4,5	Eutroph	Vilaine	27 ± 34	-
	16-CHEM	Loire Estuary	Estuary mouth	Macrotidal	31 ± 3,6	13,9 ± 3,6	3,4 ± 4,9	Mesotroph	Loire	833 ± 688	-
	17-JOAL	Loire Estuary	Estuary mouth	Macrotidal	31 ± 3,6	13,9 ± 3,6	3,4 ± 4,9	Mesotroph	Loire	833 ± 688	-
	18-BOUR	Bay of Bourgneuf	Semi-enclosed bay	Macrotidal	33 ± 2,3	13,8 ± 3,9	2,7 ± 3,9	Mesotroph	Loire	833 ± 688	-
	19-NOIR	Bay of Bourgneuf	Semi-enclosed bay	Macrotidal	33 ± 2,3	13,8 ± 3,9	2,7 ± 3,9	Mesotroph	Loire	833 ± 688	-
	20-RIVE	Charentais Sounds	Semi-enclosed sound	Macrotidal	33 ± 1,9	15,1 ± 4,3	2,2 ± 2,2	Mesotroph	Sèvre Niortaise	12 ± 13	-
	21-PALL	Charentais Sounds	Semi-enclosed sound	Macrotidal	27 ± 7,1	14,5 ± 4,8	4,6 ± 7,0	Mesotroph	Charente	65 ± 66	Sèvre Niortaise
	22-BOYV	Charentais Sounds	Semi-enclosed sound	Macrotidal	33 ± 2,5	14,9 ± 4,3	3,2 ± 4,1	Mesotroph	Charente	65 ± 66	Sèvre Niortaise
	23-PONT	Gironde Estuary	Estuary mouth	Macrotidal	26 ± 5	15,9 ± 4,2	1,9 ± 1,3	Eutroph	Garonne / Dordogne	539 ± 404	-
24-FOSS	Gironde Estuary	Estuary mouth	Macrotidal	26 ± 5	15,9 ± 4,2	1,9 ± 1,3	Eutroph	Garonne / Dordogne	259 ± 199	-	
25-FERR	Arcachon Lagoon	Semi-enclosed lagoon	Mesotidal	34 ± 0,8	15,2 ± 3,7	1,7 ± 1,1	Mesotroph	-	17 ± 14	-	
26-JACQ	Arcachon Lagoon	Semi-enclosed lagoon	Mesotidal	31 ± 2,8	15,8 ± 5,3	2,1 ± 1,1	Mesotroph	Leyre	17 ± 14	-	
27-COMP	Arcachon Lagoon	Semi-enclosed lagoon	Mesotidal	31 ± 2,8	16,0 ± 5,3	1,8 ± 1,0	Mesotroph	Leyre	17 ± 14	-	
Mediterranean Sea	28-BANY	Gulf of Lion	Open bay	Microtidal	38 ± 0,7	16,7 ± 3,9	0,6 ± 0,6	Oligotroph	-	-	Tech / TÉT
	29-HRLT	Gulf of Lion	Open bay	Microtidal	37 ± 2,4	17,7 ± 4,6	NA ± NA	Oligotroph	Hérault	35 ± 42	-
	30-STMM	Gulf of Lion	Open bay	Microtidal	NA ± NA	NA ± NA	NA ± NA	Oligotroph	Rhône	1669 ± 782	-
	31-CART	Gulf of Lion	Semi-enclosed bay	Microtidal	33 ± 3,7	16,5 ± 4,5	1,0 ± 1,1	Oligotroph	Rhône	1669 ± 782	-
	32-COUR	Gulf of Lion	Open bay	Microtidal	38 ± 0,3	17,0 ± 3,5	0,4 ± 0,4	Oligotroph	Rhône	1669 ± 782	-
	33-POME	Gulf of Lion	Open bay	Microtidal	38 ± 0,3	17,0 ± 3,5	0,4 ± 0,4	Oligotroph	Rhône	1669 ± 782	-

165 2.4 Statistical analyses

Statistical analyses were performed with the R software (R Core Team, 2022, 4.3.1 version).

Due to the nature of the dataset (i.e., only one of the three species was sampled for each sea façade, except for stations 15-PEBE and 16-CHEM from the Atlantic facade), it was not possible to test whether the observed differences in elemental and isotopic ratios were due to difference between species or sea facade, nor was it possible to ascertain if there was any interaction

170 between these two factors. Consequently, we have tested the effects of both factors (species and sea facade) independently. Therefore, particular attention should be given in interpreting the results of these tests. The effects of species (3 levels) and sea facade (3 levels) were tested independently for each elemental and isotopic variables over the entire dataset (all stations and time series) with non-parametric Kruskal-Wallis tests followed by Dunn *post-hoc* tests (R-package ‘PMCMRplus’ version 1.9.10, function ‘kruskalTest()’ and ‘kwAllPairsDunnTest()’, Pohlert, 2023) since normality and homogeneity of variance
175 (inspected using Shapiro-Wilk and Levene tests respectively) were not met, precluding the use of ANOVAs.

Monotonous temporal trends in bivalve elemental and isotopic times series were examined for each station using Mann-Kendall tests corrected for autocorrelation (R-package ‘modifiedmk’ version 1.6, function ‘mmkh()’, Patakamuri and O’Brien, 2021). Linear models (R-package ‘stats’ version 4.2.1, function ‘lm()’, R Core Team, 2022) were applied to calculate the value of the slope (in ‰ decade⁻¹) for each time series (note that all the time series do not have the same number of years). Complete
180 time series were tested for shifts allowing for the detection of one unique shift per time series (R-package ‘cpm’ version 2.3, function ‘detectChangePoint()’, Gordon, 2015).

3 Data characteristics and general overview

This dataset illustrates the temporal and spatial variability of carbon and nitrogen content (C and N %) and elemental and isotopic ratios (C:N, $\delta^{13}\text{C}$, $\delta^{15}\text{N}$) of three species of bivalve along the French coasts from 1981 to 2021. Overall, the average
185 $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of bivalves were significantly lower for the stations of the Mediterranean Sea compared with the stations of the English Channel and the Atlantic Ocean. We observed that stations closest to river mouths displayed the lowest average $\delta^{13}\text{C}$ values compared to those more distant from river influence. Over the last 10 to 40 years, nearly all stations exhibited a significant decrease in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$. There was no spatial nor temporal pattern in C:N but values differed between bivalve genus (mussel vs oyster). It is important to note that, due to the nature of the dataset (one species present for each sea facade),
190 it was not possible across all our study sites, to determine whether the differences observed in elemental and isotopic ratios between the three bivalve species and across the three sea facades were due to species effects or spatial differences. However, this could be assessed at the scale of the Atlantic facade (see sections 2.4 and 3.1).

3.1 Taxonomic patterns

Different species may exhibit different C and N elemental and isotopic ratios under the same environmental and growth
195 conditions (Mele et al., 2023), and within a single species, varying environmental conditions can lead to difference in these values (e.g. Magni et al., 2012; Briant et al., 2018). Overall, bivalve $\delta^{13}\text{C}$ ranged from -23.29 to -16.98 ‰, $\delta^{15}\text{N}$ from 2.47 to 13.15 ‰, C:N from 4.13 to 8.12 mol mol⁻¹, and C from 29.81 to 48.64 % and N from 5.81 to 10.57 % (out of extreme data, see 2.3). There was a significant difference in $\delta^{13}\text{C}$ and C content ($p < 0.001$) between *M. galloprovincialis* ($\delta^{13}\text{C}$: -20.05 ± 0.98 ‰; C: 36.10 ± 1.53 %) and the other two species *M. edulis* ($\delta^{13}\text{C}$: -19.38 ± 1.04 ‰; C: 39.80 ± 1.93 %) and *C. gigas*

200 ($\delta^{13}\text{C}$: $-19.25 \pm 1.06 \text{ ‰}$; C: $39.30 \pm 1.90 \%$), which were not significantly different (Figure 2). There was a significant difference in N content ($p < 0.001$) between *C. gigas* ($8.02 \pm 0.81 \%$) and the two other species *M. edulis* ($9.16 \pm 0.46 \%$) and *M. galloprovincialis* ($9.06 \pm 0.44 \%$; Figure 2). Finally, there was a significant difference in $\delta^{15}\text{N}$ ($p < 0.001$) between the three species of bivalves (*C. gigas*: $8.42 \pm 1.09 \text{ ‰}$; *M. edulis* $9.41 \pm 1.19 \text{ ‰}$; *M. galloprovincialis* $5.70 \pm 1.49 \text{ ‰}$) as well as in C:N ($p < 0.001$; *C. gigas*: $5.77 \pm 0.62 \text{ mol mol}^{-1}$; *M. edulis* $5.07 \pm 0.31 \text{ mol mol}^{-1}$; *M. galloprovincialis* $4.65 \pm 0.23 \text{ mol mol}^{-1}$; Figure 2). Since the mussel *M. galloprovincialis* is exclusively present in the Mediterranean Sea and the two other species, *M. edulis* and *C. gigas* are mostly present in the English Channel and Atlantic Ocean, respectively, it is difficult to separate the species factor from the spatial factor (see 3.2) in elemental and isotopic ratios (especially in $\delta^{13}\text{C}$ where *M. galloprovincialis* differs significantly). However, within the Atlantic facade where mostly oysters are sampled (15/17 stations), two stations are sampled for mussels (stations 15-PEBE and 16-CHEM), hence species effect can be tested for this facade. There was no significant differences in $\delta^{13}\text{C}$ between the two species (*C. gigas*: $-19.24 \pm 1.07 \text{ ‰}$; *M. edulis*: $-19.36 \pm 1.02 \text{ ‰}$), but $\delta^{15}\text{N}$, C and N contents, and C:N ratio differed significantly ($p < 0.001$) between *C. gigas* ($\delta^{15}\text{N}$: $8.43 \pm 1.12 \text{ ‰}$; C: $39.40 \pm 1.93 \%$; N: $8.06 \pm 0.82 \%$; C:N: $5.76 \pm 0.62 \text{ mol mol}^{-1}$) and *M. edulis* ($\delta^{15}\text{N}$: $10.54 \pm 0.88 \text{ ‰}$; C: $38.20 \pm 1.33 \%$; N: $9.34 \pm 0.39 \%$; C:N: $4.78 \pm 0.19 \text{ mol mol}^{-1}$). Note that the unequal sample sizes (15 stations for oyster, 2 stations for mussels) can reduce the power of the test and the ability to detect real differences.




Figure 2: Boxplots of $\delta^{13}\text{C}$, $\delta^{15}\text{N}$, C, N and C:N values of bivalves for the 33 stations of the three French sea facades over the period 1981-2021 (*Crassostrea gigas*, *Mytilus edulis*, *Mytilus galloprovincialis*). The lower and upper edges of the box indicate the 25th and 75th percentiles and the middle line the median. The whiskers indicate the maximum/minimum values and the open circle the extreme values. The black arrows and associated values represent extreme values out of graphical bounds.

3.2.1 Among sea facade

The three sea facades showed significant difference in bivalves $\delta^{13}\text{C}$ ($p < 0.001$). The Mediterranean Sea had lower values ($-20.05 \pm 0.98 \text{ ‰}$), than the other two sea facades, the English Channel ($-19.39 \pm 1.04 \text{ ‰}$) and the Atlantic Ocean ($-19.26 \pm 1.06 \text{ ‰}$) which were not significantly different (Figure 2). There was also a significant difference in bivalve $\delta^{15}\text{N}$ among the three sea facades ($p < 0.01$) showing a latitudinal gradient with high $\delta^{15}\text{N}$ values in the English Channel ($9.02 \pm 1.07 \text{ ‰}$), intermediate values in the Atlantic Ocean ($8.71 \pm 1.30 \text{ ‰}$), and low values in the Mediterranean Sea ($5.70 \pm 1.49 \text{ ‰}$; Figure 2). Bivalve C content and C:N ratio were also significantly different among the three sea facades ($p < 0.001$), with high values in the English Channel (C: $40.0 \pm 1.89 \text{ ‰}$; CN: $5.24 \pm 0.42 \text{ mol mol}^{-1}$) and Atlantic Ocean (C: $39.20 \pm 1.90 \text{ ‰}$; C:N: $5.63 \pm 0.67 \text{ mol mol}^{-1}$) and low values in the Mediterranean Sea (C: $36.10 \pm 1.53 \text{ ‰}$; C:N: $4.65 \pm 0.23 \text{ mol mol}^{-1}$; Figure 2). Finally, N content differed significantly ($p < 0.001$) between the Atlantic Ocean ($8.23 \pm 0.89 \text{ ‰}$) and the two other sea facades (English Channel: $8.94 \pm 0.69 \text{ ‰}$; Mediterranean Sea: $9.06 \pm 0.44 \text{ ‰}$), which is likely due to species effect (oysters vs mussels, see 3.1).

3.2.2 Among stations

The lowest $\delta^{13}\text{C}$ values were mostly observed in bivalves from the stations inside or close to the main river mouths: 5-CLHV ($-20.13 \pm 0.70 \text{ ‰}$) and 6-VILL ($-20.60 \pm 1.03 \text{ ‰}$) for the Seine river; 16-CHEM ($-20.15 \pm 0.74 \text{ ‰}$) and 17-JOAL ($-19.90 \pm 0.60 \text{ ‰}$) for the Loire river; 24-FOSS ($-20.71 \pm 1.04 \text{ ‰}$) for the Gironde estuary; 30-STMM ($-20.34 \pm 0.81 \text{ ‰}$) and 31-CART ($-20.58 \pm 1.14 \text{ ‰}$) for the Rhône river (Figure 2). The lowest $\delta^{15}\text{N}$ values were mostly observed for bivalve from the Mediterranean Sea stations (all below $< 6 \text{ ‰}$) with minimum average values for station 28-BANY ($4.27 \pm 0.94 \text{ ‰}$) and maximum for station 30-STMM ($5.91 \pm 0.75 \text{ ‰}$). The highest $\delta^{15}\text{N}$ values were observed for stations under the Loire and Seine River plumes ($> 9 \text{ ‰}$) e.g., 16-CHEM ($10.90 \pm 0.90 \text{ ‰}$), 17-JOAL ($9.79 \pm 0.43 \text{ ‰}$), 5-CLHV ($9.41 \pm 1.08 \text{ ‰}$), 6-VILL ($9.78 \pm 1.26 \text{ ‰}$), and locally under the influence of smaller rivers (Figure 2), e.g., 9-BDVG ($10.16 \pm 0.52 \text{ ‰}$), 13-AULN ($9.81 \pm 0.42 \text{ ‰}$), 15-PEBE ($10.19 \pm 0.71 \text{ ‰}$). This pattern is less clear for the Garonne and Dordogne rivers (Gironde estuary) where $\delta^{15}\text{N}$ shows intermediate values (23-PONT ($7.74 \pm 0.60 \text{ ‰}$), 24-FOSS ($8.04 \pm 0.73 \text{ ‰}$)). Bivalve C and N content and C:N ratio did not show any clear spatial pattern between stations and seemed mostly linked with genus/species effects (differing between facades, Figure 2; see also 3.1).

Interestingly, for stations 16-CHEM and 17-JOAL, both located at the mouth of the Loire river and sampled for mussels and oysters, respectively, the $\delta^{13}\text{C}$ signal did not differ significantly (mussels: $-19.95 \pm 0.74 \text{ ‰}$; oysters: $-19.90 \pm 0.60 \text{ ‰}$) over the same time period (2010-2021), despite being different species and genus. As for the comparison among the Atlantic façade (see 3.1), $\delta^{15}\text{N}$ values significantly differed between the two species (mussels: $10.30 \pm 0.44 \text{ ‰}$; oysters: $9.79 \pm 0.43 \text{ ‰}$), however, the C:N ratio was not significantly different between the two species for this specific area (mussels: $4.74 \pm 0.19 \text{ ‰}$;

250 oysters: 5.16 ± 0.64 ‰). This suggests that spatial location could be the main driver of bivalve carbon isotope signal but that nitrogen is likely influenced by both global and local processes, including bivalve physiology, that compensate for species difference.

3.2.3 Key messages


Bivalves' elemental and isotopic ratios were generally lower for the stations of the Mediterranean Sea compared to those
255 located in the Atlantic Ocean and English Channel. We found significant differences in carbon and nitrogen content and elemental and isotopic ratios between species (see 3.1), each of which is present in only one sea facade (with a few exceptions), nevertheless, this difference is also linked with spatial variability. Indeed, C and N baselines (for isotope signal) and trophic status (i.e., nutrient and chlorophyll amounts) vary between each sea facade (oligotrophic Mediterranean Sea vs. meso- to eutrophic Atlantic Ocean and English Channel; Table 2) but also locally (i.e., proximity to river mouth). To overcome the lack
260 of nutrients available in the Mediterranean Sea, some specific groups of phytoplankton (i.e., diazotrophs) uses ^{15}N -depleted atmospheric nitrogen as N source which decreases both the $\delta^{15}\text{N}$ of the overall particulate organic matter (Kerhervé et al., 2001; Wannicke et al., 2010; Landrum et al., 2011; Liénart et al., 2017) and, by trophic propagation, the low $\delta^{15}\text{N}$ signal is reflected in bivalves (Liénart et al., 2022, 2023). There is a clear difference in $\delta^{15}\text{N}$ baseline between the Mediterranean sea and the other French sea facades (Liénart et al., 2017). Generally, there was a clear difference in $\delta^{13}\text{C}$ signal close to river
265 mouths, decreasing along river plumes, and in $\delta^{15}\text{N}$ for some of the main (mostly eutrophicated) rivers. Enriched ^{15}N signal of nutrients coming from rivers with extensive watershed agricultural activities and urban outlets is reflected in the high $\delta^{15}\text{N}$ values of bivalves sampled close to river mouths (Fukumori et al., 2008; Thibault et al., 2020). Similarly, low $\delta^{13}\text{C}$ values for bivalves sampled close to river mouths are mostly linked to the inputs of continental particulate material bearing this specific signal (Liénart et al., 2017). $\delta^{13}\text{C}$ values of dissolved inorganic carbon (DIC) are lower in riverine/freshwater ecosystems
270 compared to marine environments (Mook and Rozanski, 2000), resulting in more negative $\delta^{13}\text{C}$ values in phytoplankton, which may be ingested locally by bivalves. Additionally, $\delta^{13}\text{C}$ of particulate organic carbon (POC) from terrestrial sources tends to be more negative (Liénart et al., 2017) and can be consumed by bivalves. However, whether terrestrial POC is a significant food source for bivalves remains debated (Malet et al., 2008; Marín Leal et al., 2008; Dubois et al., 2014). Similarly, the $\delta^{15}\text{N}$ signal of freshwater and marine phytoplankton differs due to the distinct nitrogen sources in these two environments. Therefore,
275 it is not surprising that the $\delta^{15}\text{N}$ signal in bivalves varies along a gradient related to proximity to river mouths.

3.3 Temporal patterns

3.3.1 Pluri-decadal $\delta^{13}\text{C}$ dataset

280 Bivalves $\delta^{13}\text{C}$ showed a significant decrease over the period 1981-2021 for 82 % of the stations (27/33 stations; Figure 3). The Mediterranean Sea exhibited a significant decrease for all stations (6/6 stations), the Atlantic Ocean for 82 % of the stations (14/17 stations) and 70 % of the stations in the English Channel (7/10 stations). The average decrease over the total period was consistent across all facades, averaging -0.58 ± 0.28 ‰ per decade, ranging from -0.22 to -1.21 ‰ per decade (i.e., -4.85 to -0.89 ‰ over the last 40 years) when the slopes were significant. This decrease in bivalves $\delta^{13}\text{C}$ was more pronounced for the shorter time series starting from late 1990's (17-JOAL, 2010-2021; 11-PASS and 12-PERS, 2001-2021; 33-POME, 1998-2021; 7-OUIS, 1994-2021; Figure 3). Overall, the slopes were more pronounced for the Mediterranean Sea and for the stations
285 in the western Bay of Seine, i.e., under the influence of the Seine plume (stations 7 to 10). Some of the stations showed strong interannual variability in bivalves $\delta^{13}\text{C}$ (e.g., 1-AMBL, 6-VILL, 24-FOSS, Figure 3).

Shifts in bivalve $\delta^{13}\text{C}$ time series were detected for 79 % of the stations (26/33 stations), equally represented in each sea facade. The shifts occurred mostly around the year 1999-2000 (6/26 stations) or after, around the years 2006 ± 1 year and 2012 ± 1 year, with no specific pattern per façade. In the Mediterranean Sea most of the shifts occurred before 2000. The earliest shift was in 1989 (31-CART) and two in 1999, the latest in 2014 (15-PEBE). Usually, no shift was detected at the stations close to
290 river mouths, i.e., Seine (5-CLHV, 6-VILL), Loire (16-CHEM, 17-JOAL), Gironde (24-FOSS) estuaries, and at some of the mouths of smaller rivers (e.g., 15-PEBE, Vilaine river; 29-HRLT, Hérault river).




295 **Figure 3: Bivalves $\delta^{13}\text{C}$ time series over the period 1981-2021 for the 33 stations of the three French sea facades (*Crassostrea gigas*, *Mytilus edulis*, *Mytilus galloprovincialis*). Black lines correspond to significant trends (Mann-Kendall tests, p -value < 0.05), the slope value (‰ per decade) appears in the upper right corner. Shifts are represented by the vertical dash line with the year mentioned on its down left. One outlier is shown in grey (21-PALL, -27‰) but was not considered for statistical analyses.**

3.3.2 Pluri-decadal $\delta^{15}\text{N}$ dataset

Bivalve $\delta^{15}\text{N}$ also showed a significant decrease over the period 1981-2021 for 64 % of the stations (22/33 stations; Figure 4).
300 The Mediterranean Sea exhibited a significant decrease for 83 % stations (5/6 stations), the English Channel for 80 % of the
stations (8/10 stations) and only 53 % of the stations in the Atlantic Ocean (9/17 stations). Only one station showed a significant
increase (25-FERR, 0.34 ‰ over 39 years). The average decrease was -0.44 ± 0.20 ‰ per decade, ranging from -0.18 to -0.84
‰ per decade (i.e., -0.72 to -3.35 ‰ over the last 40 years; Figure 4). However, this decrease was more pronounced in the
English Channel (-0.50 ± 0.21 ‰) in the eastern part and close the Seine river mouth (e.g., 3-VARE, 4-ANTI, 6-VILL), and
305 Mediterranean Sea (-0.54 ± 0.20 ‰) compared with the Atlantic Ocean (-0.32 ± 0.08 ‰) where the lowest slopes were near
the Loire river mouth and the inner two stations of the Arcachon lagoon. The interannual variability was relatively low for
 $\delta^{15}\text{N}$ with the exception of some stations (e.g., 1-AMBL, 6-VILL, 16-CHEM, 33-POME).

Shifts in bivalve $\delta^{15}\text{N}$ time series were detected for 70 % of the stations (23/33 stations), mostly in the English Channel and
Mediterranean Sea. The shifts occurred mostly around the year 1999-2000 (5/23 stations), then around the years 1995-1996
310 (4/23 stations) and 2004 ± 1 year (4/23 stations), with shifts mostly occurring before 2000 in the Mediterranean Sea. For each
sea facade, there were often shifts occurring the same year or in two close years for nearby stations (e.g., 15-PEBE, 2003 and
16-CHEM, 2004). The earliest shift was in 1991 (29-HRLT), the latest in 2013 (14-GUIL).



315 **Figure 4:** Bivalves $\delta^{15}\text{N}$ time series over the period 1981-2021 for the 33 stations of the three French sea facades (*Crassostrea gigas*, *Mytilus edulis*, *Mytilus galloprovincialis*). Black lines correspond to significant trends (Mann-Kendall tests, p-value < 0.05), the slope value (‰ per decade) appears in the upper right corner. Shifts are represented by the vertical dash line with the year mentioned on its down left.

320 3.3.3 Pluri-decadal C and N datasets

Bivalve C and N contents and C:N ratio showed significant trends for less than half of the stations (between 30 and 36 %) over the period 1981-2021 (C: 12/33 stations; N: 11/33 stations; C:N: 10/33 stations; Figure 5, 6 and 7). For C content, there was the same number of increasing and decreasing trends (both 18 %, 6/33 stations), with no spatial pattern, while N content showed only decreasing trends (33%, 11/33 stations). However, most of the significant trends were increasing for C:N ratio
325 (24 %, 8/33 stations), exclusively in the Atlantic Ocean (6/33 stations) and the English Channel (2/33 stations) and only 6 % (2/33 stations) of the stations showed decreasing trends, all in the Mediterranean Sea. The average increase in C:N ratio was $0.27 \pm 0.37 \text{ mol mol}^{-1}$ per decade, ranging from 0.05 to 0.38 mol mol^{-1} per decade (i.e., 0.20 to 1.53 mol mol^{-1} over the last 40 years; excluding 17-JOAL with a 1.45 mol mol^{-1} increase for 11 years; Figure 5). The average C:N decrease in the Mediterranean sea was $-0.05 \pm 0.02 \text{ mol mol}^{-1}$ per decade. Interannual variability in bivalve C:N was either very low for the
330 two mussel species (e.g., 7-OUIS, 16-CHEM, 33-POME), or very large for the oysters (e.g., 12-PERS, 19-NOIR, 22-BOYV). The decreasing trends in C:N ratio were mostly linked to decrease in bivalves N % (6/8 stations), whereas only 3/8 stations showed a significant increase in C %. Shifts in bivalve C, N and C:N time series were detected only for 18 to 33 % of the stations (C: 6/33 stations; N: 9/33 stations; C:N: 11/33 stations), without any peculiar spatial pattern. Most of the shifts occurred after the year 2000 (C: 4/33 stations; N: 5/33 stations; C:N: 7/11 stations). The earliest C:N shift was in 1989 (27-COMP), the
335 latest in 2014 (6-VILL, 22-BOYV). There was no apparent spatial pattern in the C, N and C:N shifts.






Figure 5: Bivalves C:N time series over the period 1981-2021 for the 33 stations of the three French sea facades (*Crassostrea gigas*, *Mytilus edulis*, *Mytilus galloprovincialis*). Black lines correspond to significant trends (Mann-Kendall tests, p-value < 0.05), the slope value (% per decade) appears in the upper right corner. Shifts are represented by the vertical dash line with the year mentioned on its down left.

340



345 **Figure 6: Bivalves C content (%) time series over the period 1981-2021 for the 33 stations of the three French sea facades (*Crassostrea gigas*, *Mytilus edulis*, *Mytilus galloprovincialis*). Black lines correspond to significant trends (Mann-Kendall tests, p-value < 0.05, calculated excluding extreme values mentioned in section 2.3), the slope value (‰ per decade) appears in the upper right corner.**

Shifts are represented by the vertical dash line with the year mentioned on its down left. Extreme values mentioned in section 2.3 are not represented and were excluded from the calculations of the slopes and shifts.



350 **Figure 7: Bivalves N content (%) time series over the period 1981-2021 for the 33 stations of the three French sea facades (*Crassostrea gigas*, *Mytilus edulis*, *Mytilus galloprovincialis*). Black lines correspond to significant trends (Mann-Kendall tests, p-value < 0.05,**

calculated excluding extreme values mentioned in section 2.3), the slope value (% per decade) appears in the upper right corner. Shifts are represented by the vertical dash line with the year mentioned on its down left. Extreme values mentioned in section 2.3 are not represented and were excluded from the calculations of the slopes and shifts.

3.3.4 Key messages

355 In summary, over the last 30 to 40 years, bivalve have shown a general decrease in both $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ for most stations of the
three French sea facades. These trends are most likely linked with global processes occurring at large spatial scale such as
climate change. Liénart et al. (2024) explained the decrease in $\delta^{13}\text{C}$ is primarily associated with the increase in atmospheric
CO₂ concentrations due to fossil-fuel burning, which generates a decrease in atmospheric CO₂ $\delta^{13}\text{C}$ values occurring since the
360 industrial revolution in 1850 (the Suess effect; Keeling 1979; Gruber et al., 1999). This decrease in $\delta^{13}\text{C}$ values propagates
along marine food webs and is visible in organisms' tissues (Schloesser et al. 2009; Liénart et al., 2022, 2024b). However,
other global factors such as rising temperatures or climate indices may also contribute to this overall decrease in $\delta^{13}\text{C}$ through
passive (e.g., Suess effect) or active processes related to bivalve physiology (e.g., change in fractionation; Liénart et al., 2024b).
Decreasing $\delta^{15}\text{N}$ trends are potentially linked to global factors, indirectly through a general decrease in nutrients and particles
365 inputs from rivers (Milliman et al., 2008; Bauer et al., 2013). Since organic particles generated in areas with high human
activities bears a high $\delta^{15}\text{N}$ signal, the decrease of such ^{15}N -enriched inputs is reflected in coastal water and marine fauna
(Connolly et al., 2013). However, such changes are strongly influenced by regional and local factors such as regional climate,
watershed activities and damming (Milliman et al., 2008) resulting in different slopes values in isotope time series close to
river mouths (more pronounced for both C and N isotopes). The observed decrease in both isotopic ratios likely results from
the cumulative, synergistic, or antagonistic effects of global and regional/local influences (Cabral et al., 2019). Such interacting
370 pressures are more likely to occur in regions strongly affected by climate change, such as the Mediterranean Sea (Tuel and
Eltahir, 2020). In this dataset, isotopes reveal both global (climate) effects, mostly through $\delta^{13}\text{C}$, and some local variability in
the organic matter used by bivalves, as shown by the different $\delta^{13}\text{C}$ (low) and $\delta^{15}\text{N}$ (high) signals along the river-mouth-to-sea
gradient. The decrease in bivalve $\delta^{13}\text{C}$ and in $\delta^{15}\text{N}$ over the past decades may indicate a decrease in nutrient and particles inputs
from the rivers over the same period. Finally, as an indicator of bivalve condition and physiology (Elser et al., 2003), the
375 increase in C:N ratio observed for most stations in the Atlantic and English Channel can be attributed to either an increase in
carbon content (i.e., increase in lipid or carbohydrate content), which is unlikely given rising temperatures and appear
significant for only few of our stations, or to a decrease in nitrogen content (i.e., proteins, amino acids), as observed in our data
and likely due to a decrease in N-nutrient availability in the environment which is consistent with the observed decrease in
 $\delta^{15}\text{N}$. The absence of trends at most of the stations and the large interannual variability of C:N ratio may result from the
380 complex interplay between global and local environmental effects and bivalves physiology on this parameter.

4 Data availability

The dataset presented in this article and the metadata associated can be freely accessed on the SEANOE open access repository (<https://www.seanoe.org/>) under the DOI <https://doi.org/10.17882/100583> (Liénart et al., 2024a). The data gives the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ (‰), C and N (%) and C:N ratio (mol mol^{-1}), for each station and year, with sampling date and species name. For each station, it contains the sea façade, station ID, station name, ROCCH codes, latitude and longitude (geographic coordinate system WGS84) as presented in Table 1. Carbon isotopic time-series are already published (Liénart et al., 2024b, except from station 17-JOAL) in a study focused on global effect on coastal ecosystems from large-scale anthropogenic and natural climate proxies, including the Suess effect, over the period 1981-2021. Bivalve $\delta^{13}\text{C}$ data and $\delta^{13}\text{C}$ corrected from the Suess effect are available under the permanent identifier <https://doi.org/10.6084/m9.figshare.24884871.v1> (Liénart et al., 2024c). The companion data from the ROCCH (contaminants) are available on the ‘Survival’ database (<https://surval.ifremer.fr>).

5 Conclusion and recommendation for use

By providing this dataset to the scientific community including caveats for interpreting such data spatially and over time, we expect it will be useful for numerous ecological studies. Such data is relevant to trace nutrients origin, to set accurate baselines to study organisms diets and food webs structure, and can be an indicator of water quality over space and time through bivalve physiology. It could also provide valuable input for developing predictive models of bivalve physiology (Emmery et al., 2011) or trophic ecology (Marín Leal et al., 2008). Multi-decadal time-series allow scientists to understand coastal ecosystems responses to global change through biological and biogeochemical processes. The carbon isotope dataset is already part of a study where trends are thoughtfully interpreted in regard of global proxies for climate and anthropogenic changes and corrected for the Suess effect (Liénart et al., 2024b). Part of the observed $\delta^{13}\text{C}$ trends were linked with the Suess effect, leading to shifts in isotope baseline over recent decades (i.e., Suess effect for $\delta^{13}\text{C}$) rather than specific changes in ecosystem functioning. Hence, we would like to draw users attention on the need to correct for the Suess effect before comparing biological samples collected one or more decades apart as recommended in literature (Dombrosky 2020; Clark et al. 2021; Liénart et al., 2024b). Additionally, in order to take into account the low $\delta^{13}\text{C}$ and high $\delta^{15}\text{N}$ signal observed near river mouths, we advise to compare slope values rather than absolute values when assessing temporal changes between stations. Overall, the complex interplay between global and regional/local effects needs to be considered when interpreting time series. This long-term dataset of elemental and isotopic values in suspension-feeders tissues provides insights into ecosystem dynamics and hold broader significance for advancing scientific understanding in the face of ongoing environmental challenges.

Author contribution

CL and NS are leaders of the project. NS, CL, AL, AGP, PLM, XdM, HB, SD and AIG participated in defining the scientific strategy and selection of archived samples for analysis. AF and NB prepared the samples for analysis. AnG carried out sample

analysis. PLM and AGP are the curator of the ROCCH sample archive. CL and AF performed the statistical analysis. CL prepared the manuscript and figures and integrated the final contribution from all co-authors.

415 **Competing interests**

The authors declare that they have no competing interests.

Disclaimer

420 Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Acknowledgements

425 This article is based on a collaborative work with the team of the ROCCH monitoring network coordinated by Ifremer. We thank the members of the ROCCH and of the different institutions, from the field workers and sample analysts to the coordinators, who made it possible to use these samples.

Financial support

This research was funded by the Office Français de la Biodiversité (OFB) within the frame of the research project EVOLECO-BEST.

430 **References**

- Amiard, J.C., Amiard-Triquet, C., Barka, S., Pellerin, J., and Rainbow, P.S.: Metallothioneins in aquatic invertebrates: Their role in metal detoxification and biomonitoring., *Aquatic Toxicology*, 76(2), 160-202. <https://doi.org/10.1016/j.aquatox.2005.08.015>, 2006.
- 435 Bauer, J. E., Cai, W.-J., Raymond, P., Bianchi, T. S., Hopkinson, C. S., and Regnier, P. G.: The changing carbon cycle of the coastal ocean., *Nature*, 504, 61–70, <https://doi.org/10.1038/nature12857>, 2013.
- Briand, M. J., Herlory, O., Briant, N., Brach-Papa, C., Boissery, P., and Bouchouca, M.: The French Mussel Watch: More than two decades of chemical contamination survey in Mediterranean coastal waters, *Mar. Pollut. Bull.*, 191, 114901, <https://doi.org/10.1016/j.marpolbul.2023.114901>, 2023.
- 440 Briant, N., Savoye, N., Chouvelon, T., David, V., Rodriguez, S., Charlier, K., Sonke, J. E., Chiffolleau, J. F., Brach-Papa, C., and Knoery J.: Carbon and nitrogen elemental and isotopic ratios of filter-feeding bivalves along the French coasts: An assessment of specific, geographic, seasonal and multi-decadal variations. *Sci. Total Environ.* 613–614: 196–207. doi:10.1016/j.scitotenv.2017.08.281, 2018.

- 445 Bristow, L. A., Jickells, T. D., Weston, K., Marca-Bell, A., Parker, R., and Andrews J. E.: Tracing estuarine organic matter sources into the southern North Sea using C and N isotopic signatures. *Biogeochemistry*, 113: 9–22, <https://doi.org/10.1007/s10533-012-9758-4>, 2013.
- Cabral, H., Fonseca, V., Sousa, T., and Leal, M. C.: Synergistic effects of climate change and marine pollution: An overlooked interaction in coastal and estuarine areas, *Int. J. Environ. Res. Public Health*, 16, 1–17, <https://doi.org/10.3390/ijerph16152737>, 2019.
- 450 Chahouri, A., Yacoubi, B., Moukrim, A., and Banaoui, A.: Bivalve molluscs as bioindicators of multiple stressors in the marine environment: Recent advances, *Cont. Shelf Res.*, 264, 105056, <https://doi.org/10.1016/j.csr.2023.105056>, 2023.
- Clark, C. T., Cape, M. R., Shapley, M. D., Mueter, F. J., Finney, B. P., and Misarti, N.: SuessR: Regional corrections for the effects of anthropogenic CO₂ on δ¹³C data from marine organisms, *Methods Ecol. Evol.*, 12, 1508–1520, <https://doi.org/10.1111/2041-210X.13622>, 2021.
- 455 Cloern, J. E., Abreu, P. C., Carstensen, J., Chauvaud, L., Elmgren, R., Grall, J., Greening, H., Johansson, J. O. R., Kahru, M., Sherwood, E. T., Xu, J., and Yin, K.: Human activities and climate variability drive fast-paced change across the world’s estuarine-coastal ecosystems, *Glob. Chang. Biol.*, 22, 513–529, <https://doi.org/10.1111/gcb.13059>, 2016.
- Connolly, R. M., Gorman, D., Hindell, J. S., Kildea, T. N., and Schlacher, T. A.: High congruence of isotope sewage signals in multiple marine taxa, *Mar. Pollut. Bull.*, 71, 152–158, <https://doi.org/10.1016/j.marpolbul.2013.03.021>, 2013.
- 460 Dale, V. H., and Beyeler, S. C.: Challenges in the development and use of ecological indicators. *Ecol. indic.*, 1(1), 3-10, [https://doi.org/10.1016/S1470-160X\(01\)00003-6](https://doi.org/10.1016/S1470-160X(01)00003-6), 2001.
- Dombrosky, J.: A ~1000-year ¹³C Suess correction model for the study of past ecosystems, *Holocene*, 30, 474–478, <https://doi.org/10.1177/0959683619887416>, 2020.
- 465 Dubois, S., Blanchet, H., Garcia, A., Massé, M., Galois, R., Grémare, A., Charlier, K., Guillou, G., Richard, P. and Savoye, N.: Trophic resource use by macrozoobenthic primary consumers within a semi-enclosed coastal ecosystem: stable isotope and fatty acid assessment. *Journal of Sea Research*, 88, 87-99. <https://doi.org/10.1016/j.seares.2014.01.004>, 2014.
- Elser, J. J., Acharya, K., Kyle, M., Cotner, J., Makino, W., Markow, T., Watts, T., Hobbie, S., Fagan, W., Schade, J., Hood, J., and Sterner, R. W.: Growth rate-stoichiometry couplings in diverse biota, *Ecol. Lett.*, 6, 936–943, <https://doi.org/10.1046/j.1461-0248.2003.00518.x>, 2003.
- 470 Emmery, A., Lefebvre, S., Alunno-Bruscia, M., and Kooijman, S. A. L. M.: Understanding the dynamics of δ¹³C and δ¹⁵N in soft tissues of the bivalve *Crassostrea gigas* facing environmental fluctuations in the context of Dynamic Energy Budgets (DEB)., *Journal of sea research*, 66(4), 361-371. <https://doi.org/10.1016/j.seares.2011.08.002>, 2011.
- Fry, B., and Sherr, E.: ¹³C measurements as indicators of carbon flow in marine and freshwater ecosystems. *Contrib. Mar.*

- Sci., 27:13–47, 1984.
- Fukumori, K., Oi, M., Doi, H., Takahashi, D., Okuda, N., Miller, T. W., Kuwae, M., Miyasaka, H., Genkai-Kato, M., Koizumi, Y., Omori, K., and Takeoka, H.: Bivalve tissue as a carbon and nitrogen isotope baseline indicator in coastal ecosystems, *Estuar. Coast. Shelf Sci.*, 79, 45–50, <https://doi.org/10.1016/j.ecss.2008.03.004>, 2008.
- Glibert, P. M., Middelburg, J. J., McClelland, J. W., and Vander Zanden, J. M.: Stable isotope tracers: Enriching our perspectives and questions on sources, fates, rates, and pathways of major elements in aquatic systems, *Limnol. Oceanogr.*, 1–32, <https://doi.org/10.1002/lno.11087>, 2018.
- 480 Gordon J. R.: Parametric and Nonparametric Sequential Change Detection in R: The cpm Package. *J. Stat. Software.* 66(3), 1-20. <http://www.jstatsoft.org/v66/i03/>, 2015.
- Grouhel A.: Prescriptions techniques pour l'échantillonnage de mollusques du réseau national d'observation des contaminants chimiques (ROCCH), RST- RBE/CCEM/ROCCH 23-02, <https://doi.org/10.13155/97878>, 2023.
- Gruber, N., Keeling, C. D., Bacastow, R. B., Guenther, P. R., Lueker, T. J., Wahlen, M., Meijer, H. A. J., Mook, W. G., and 485 Stocker, T. F: Spatiotemporal patterns of carbon-13 in the global surface ocean and the oceanic Suess effect, *Global Biogeochem. Cycles*, 13: 307–335, <https://doi.org/10.1029/1999GB900019>, 1999.
- Harley, C. D. G., Hughes, A. R., Hultgren, K. M., Miner, B. G., Sorte, C. J. B., Thornber, C. S., Rodriguez, L. F., Tomanek, L., and Williams, S. L.: The impacts of climate change in coastal marine systems, *Ecol. Lett.*, 9, 228–241, <https://doi.org/10.1111/j.1461-0248.2005.00871.x>, 2006.
- 490 Hofmann, G. E., Blanchette, C. A., Rivest, E. B., and Kapsenberg, L.: Taking the pulse of marine ecosystems: The importance of coupling long-term physical and biological observations in the context of global change biology. *Oceanography*, 26(3), 140–148, <https://www.jstor.org/stable/24862075>, 2013.
- Kanduč, T., Šlejkovec, Z., Falnoga, I., Mori, N., Budič, B., Kovačić, I., Pavičić – Hamer, D., and Hamer, B.: Environmental status of the NE Adriatic Sea, Istria, Croatia: Insights from mussel *Mytilus galloprovincialis* condition indices, stable isotopes 495 and metal(loid)s, *Mar. Pollut. Bull.*, 126, 525–534, <https://doi.org/10.1016/j.marpolbul.2017.09.052>, 2018.
- Karlson, A. M. L. and Faxneld, S.: Polycyclic aromatic hydrocarbons and stable isotopes of carbon and nitrogen in Baltic Sea blue mussels: Time series data 1981 – 2016, *Data Br.*, 35, 4–8, <https://doi.org/10.1016/j.dib.2021.106777>, 2021.
- Keeling, C. D.: The Suess effect: ¹³Carbon-¹⁴Carbon interrelations, *Environ. Int.*, 2, 229–300, [https://doi.org/10.1016/0160-4120\(79\)90005-9](https://doi.org/10.1016/0160-4120(79)90005-9), 1979.
- 500 Kerhervé, P., Minagawa, M., Heussner, S., and Monaco, A.: Stable isotopes (¹³C/¹²C and ¹⁵N/¹⁴N) in settling organic matter of the northwestern Mediterranean Sea: biogeochemical implications, *Oceanol. Acta*, 24, S77–S85, 2001.
- Landrum, J. P., Altabet, M. A., and Montoya, J. P.: Basin-scale distributions of stable nitrogen isotopes in the subtropical

North Atlantic Ocean: Contribution of diazotroph nitrogen to particulate organic matter and mesozooplankton, *Deep. Res. Part I Oceanogr. Res. Pap.*, 58, 615–625, <https://doi.org/10.1016/j.dsr.2011.01.012>, 2011.

505 Lheureux, A., David, V., Del Amo, Y., Soudant, D., Auby, I., Bozec, Y., Conan, P., Ganthy, F., Grégori, G., Lefebvre, A., Leynard, A., Rimmelin-Maury, P., Souchu, P., Vantrepote, V., Blondel, C., Cariou, T., Crispi, O., Cordier, M-A., Crouvoisier, M., Duquesne, V., Ferreira, S., Garcia, N., Gouriou, L., Grosteffan, E., Le Merrer, Y., Meteigner, C., Retho, M., Tournaire, M-P., and Savoye, N.: Trajectories of nutrients concentrations and ratios in the French coastal ecosystems: 20 years of changes in relation with large-scale and local drivers. *Sci. Tot. Envir.*, 857, 159619. <https://doi.org/10.1016/j.scitotenv.2022.159619>,
510 2023.

Liénart, C., Savoye, N., Bozec, Y., Breton, E., Conan, P., David, V., Feunteun, E., Grangeré, K., Kerhervé, P., Lebreton, B., Lefebvre, S., L'Helguen, S., Mousseau, L., Raimbault, P., Richard, P., Riera, P., Sauriau, P.-G., Schaal, G., Aubert, F., Aubin, S., Bichon, S., Boinet, C., Bourasseau, L., Bréret, M., Caparros, J., Cariou, T., Charlier, K., Claquin, P., Cornille, V., Corre, A.-M., Costes, L., Crispi, O., Crouvoisier, M., Czamanski, M., Del Amo, Y., Derriennic, H., Dindinaud, F., Durozier, M.,
515 Hanquiez, V., Nowaczyk, A., Devesa, J., Ferreira, S., Fournier, M., Garcia, F., Garcia, N., Geslin, S., Grossteffan, E., Gueux, A., Guillaudeau, J., Guillou, G., Joly, O., Lachaussée, N., Lafont, M., Lamoureux, J., Lecuyer, E., Lehodey, J.-P., Lemeille, D., Leroux, C., Macé, E., Maria, E., Pineau, P., Petit, F., Pujon-Pay, M., Rimelin-Maury, P., and Sultan, E.: Dynamics of particulate organic matter composition in coastal systems: A spatio-temporal study at multi-systems scale, *Prog. Oceanogr.*, 156, <https://doi.org/10.1016/j.pocean.2017.03.001>, 2017.

520 Liénart, C., Garbaras, A., Qvarfordt, S., Sysoev, A. Ö., Högländer, H., Walve, J., Schagerström, E., Eklöf, J., and Karlson, A. M.: Long-term changes in trophic ecology of blue mussels in a rapidly changing ecosystem, *Limnol. Oceanogr.*, 66, 694–710, <https://doi.org/10.1002/lno.11633>, 2020.

Liénart, C., Garbaras, A., Qvarfordt, S., Walve, J., and Karlson, A. M. L.: Spatio-temporal variation in stable isotope and elemental composition of key-species reflect environmental changes in the Baltic Sea, *Biogeochemistry*, 157, 149–170,
525 <https://doi.org/10.1007/s10533-021-00865-w>, 2022.

Liénart, C., Tedengren, M., Garbaras, A., Lim, H., Chynel, M., Robinson, C., Meziane, T., and Karlson, A. M. L.: Diet quality determines blue mussel physiological status: A long-term experimental multi-biomarker approach, *J. Exp. Mar. Bio. Ecol.*, 563, <https://doi.org/10.1016/j.jembe.2023.151894>, 2023.

Liénart, C., Fournioux, A., Garbaras, A., Blanche, H., Briant, N., Dubois, S., Gangnery, A., Grouhel, A., Le Monier, P.,
530 Lheureux, A., de Montaudouin, X., and Savoye, N. : Bivalve multi-decadal (1981-2021) records of carbon and nitrogen elemental and isotopic ratios ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$ and C:N) of 33 coastal sites of France. SEANO Sea scientific open data publication, <https://doi.org/10.17882/100583>, 2024a.

Liénart, C., Fournioux, A., Garbaras, A., Lheureux, A., Blanchet, H., Briant, N., Dubois, S. F., Gangnery, A., Grouhel Pellouin,

- A., Le Monier, P., de Montaudouin, X., and Savoye, N.: Bivalve tissues as a recorder of multidecadal global anthropogenic and climate-mediated change in coastal areas. *Limnol. Oceanogr. Letters*, <https://doi.org/10.1002/lol2.10399>, 2024b.
- Liénart, C., Fournioux, A., Garbaras, A., Lheureux, A., Blanchet, H., Briant, N., Dubois, S. F., Gangnery, A., Grouhel Pellouin, A., Le Monier, P., de Montaudouin, X., and Savoye, N. : Bivalves $\delta^{13}\text{C}$ time series (1981-2021) 32 stations France. Figshare, <https://doi.org/10.6084/m9.figshare.24884871.v1>, 2024c.
- Lotze, H. K., Lenihan, H. S., Bourque, B. J., Bradbury, R. H., Cooke, R. G., Kay, M. C., Kidwell, S. M., Kirby, M. X., Peterson, C. H., and Jackson, J. B.: Depletion, degradation, and recovery potential of estuaries and coastal seas. *Science*, 312(5781), 1806-1809, <https://doi.org/10.1126/science.1128035>, 2006.
- Magni, P., Rajagopal, S., Como, S., Jansen, J. M., van der Velde, G., and Hummel, H.: $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ variations in organic matter pools, *Mytilus* spp. and *Macoma balthica* along the European Atlantic coast. *Marine Biology*, 160, 541-552. <https://doi.org/10.1007/s00227-012-2110-7>, 2013.
- Malet, N., Sauriau, P. G., Ryckaert, M., Malestroit, P., and Guillou, G.: Dynamics and sources of suspended particulate organic matter in the Marennes-Oléron oyster farming bay: Insights from stable isotopes and microalgae ecology. *Estuarine, Coastal and Shelf Science*, 78(3), 576-586. <https://doi.org/10.1016/j.ecss.2007.11.001>, 2008.
- Marín Leal, J. C., Dubois, S., Orvain, F., Galois, R., Blin, J.-L., Ropert, M., Bataillé, M.-P., Ourry, A., and Lefebvre, S.: Stable isotopes ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$) and modelling as tools to estimate the trophic ecology of cultivated oysters in two contrasting environments. *Marine Biology*, 153, 673-688. <http://dx.doi.org/10.1007/s00227-007-0841-7>, 2008.
- Mazaleyrat, A., Normand, J., Dubroca, L., and Fleury, E.: A 26-year time series of mortality and growth of the Pacific oyster *C. gigas* recorded along French coasts, *Sci. Data*, 9, 392, <https://doi.org/10.1038/s41597-022-01511-2>, 2022.
- Mele, I., McGill, R. A., Thompson, J., Fennell, J., and Fitzer, S.: Ocean acidification, warming and feeding impacts on biomineralization pathways and shell material properties of *Magallana gigas* and *Mytilus* spp. *Marine Environmental Research*, 186, 105925. <https://doi.org/10.1016/j.marenvres.2023.105925>, 2023.
- Milliman, J. D., Farnsworth, K. L., Jones, P. D., Xu, K. H., and Smith, L. C.: Climatic and anthropogenic factors affecting river discharge to the global ocean, 1951-2000, *Glob. Planet. Change*, 62, 187-194, <https://doi.org/10.1016/j.gloplacha.2008.03.001>, 2008.
- Mook, W., and Rozanski, K.: Environmental isotopes in the hydrological cycle – Principles and applications. IAEA Publish, 39(1), 2. , 2000.
- Niemi, G. J., and McDonald, M. E.: Application of ecological indicators. *Annu. Rev. Ecol. Evol. Syst.*, 35, 89-111. <https://doi.org/10.1146/annurev.ecolsys.35.112202.130132>, 2004.

- Patakamuri, S., and O'Brien, N.: modifiedmk: Modified versions of Mann Kendall and Spearman's Rho Trend Tests, R package version 1.6, <https://CRAN.R-project.org/package=modifiedmk>, 2021.
- 565 Pohlert T.: *_PMCMRplus*: Calculate Pairwise Multiple Comparisons of Mean Rank Sums Extended, R package version 1.9.10, <https://CRAN.R-project.org/package=PMCMRplus>, 2023.
- R Core Team: R: A language and environment for statistical computing. R Foundation for Statistical Computing, <https://www.R-project.org/>, 2022.
- Ray, N. E., and Fulweiler, R. W.: Seasonal patterns of benthic-pelagic coupling in oyster habitats. *Mar. Ecol. Prog. Ser.*, 652, 570 95-109, <https://doi.org/10.3354/meps13490>, 2020.
- Schloesser, R. W., Rooker, J. R., Louchuarn, P., Neilson, J. D., and Secor, D. H.: Interdecadal variation in seawater $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ recorded in fish otoliths, *Limnol. Oceanogr.*, 54, 1665–1668, <https://doi.org/10.4319/lo.2009.54.5.1665>, 2009.
- Sukhotin, A., and Berger, V.: Long-term monitoring studies as a powerful tool in marine ecosystem research, *Hydrobio.*, 706, 1–9, <https://doi.org/10.1007/s10750-013-1456-2>, 2013.
- 575 Thibault, M., Duprey, N., Gillikin, D. P., Thébault, J., Douillet, P., Chauvaud, L., Amice, E., Munaron, J. M., and Lorrain, A.: Bivalve $\delta^{15}\text{N}$ isoscapes provide a baseline for urban nitrogen footprint at the edge of a World Heritage coral reef, *Mar. Pollut. Bull.*, 152, 110870, <https://doi.org/10.1016/j.marpolbul.2019.110870>, 2020.
- Tuel, A., and Eltahir, E. A.: Why is the Mediterranean a climate change hot spot?, *J. Clim.*, 33(14), 5829-5843. <https://doi.org/10.1175/JCLI-D-19-0910.1>, 2020.
- 580 Wannicke, N., Liskow, I., and Voss, M.: Impact of diazotrophy on N stable isotope signatures of nitrate and particulate organic nitrogen: case studies in the north-eastern tropical Atlantic Ocean, *Isotopes Environ. Health Stud.*, 46, 337–54, <https://doi.org/10.1080/10256016.2010.505687>, 2010.