



# Bivalve monitoring over French coasts: multi-decadal records of carbon and nitrogen elemental and isotopic ratios ( $\delta^{13}\text{C}$ , $\delta^{15}\text{N}$ and C:N) as ecological indicators of global change

5 Camilla Liénart<sup>1</sup>, Alan Fournioux<sup>1</sup>, Andrius Garbaras<sup>2</sup>, Hugues Blanchet<sup>1</sup>, Nicolas Briant<sup>3</sup>, Stanislas F. Dubois<sup>4</sup>, Aline Gangnery<sup>3</sup>, Anne Grouhel Pellouin<sup>3</sup>, Pauline Le Monier<sup>3</sup>, Arnaud Lheureux<sup>5</sup>, Xavier de Montaudouin<sup>1</sup>, Nicolas Savoye<sup>1</sup>

<sup>1</sup>Université de Bordeaux, CNRS UMR 5805, Bordeaux INP, EPOC, Pessac, 33600, France

<sup>2</sup>Center for Physical Sciences and Technology, FTMC, Vilnius, 10257, Lithuania

<sup>3</sup>Ifremer Centre Atlantique, CCEM, Nantes, 44000, France

10 <sup>4</sup>Ifremer, DYNECO, Plouzané, 29280, France

<sup>5</sup>MNHN, CNRS UMR 8067, SU, IRD 207, UCN, UA, BOREA, CP53, F-75005 Paris, France

*Correspondence to:* Camilla Liénart (camilla.lienart@u-bordeaux.fr)

## Abstract

15 Recent changes in climate and environment, influenced by both global and local factors, have had profound impacts on coastal ecosystem functioning and trajectories. By examining archived samples from ongoing ecological monitoring efforts, particularly focusing on bivalves like mussels and oysters, we gain a valuable long-term perspective on how ecosystems are responding at various scales. We conducted analyses on carbon and nitrogen elemental and isotopic ratios (C:N,  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ ) of mussel and oyster soft tissues collected annually at 33 sites along the French coast from 1981 to 2021. This extensive dataset  
20 (<https://doi.org/10.17882/100583>, Liénart et al., 2024a) offers a comprehensive view spanning multiple decades and ecosystems, allowing to track how coastal ecosystems and marine species record changing climate, physical-chemical environments and organic matter cycles. Additionally, these data are crucial for establishing isotope baselines for studying food webs. Ultimately, this data set provide valuable information for more effective ecosystem conservation and management strategies in our rapidly changing world.



## 25 1 Introduction

Over the past two decades, rapid and sometimes abrupt changes in climate and environmental conditions have significantly impacted ecosystems functioning with human activities widely acknowledged as a primary factor driving and modulating these changes (Cloern et al., 2016). At the land-ocean interface, coastal and estuarine environments are particularly vulnerable to the climate change, alongside with local human-induced pressures such as pollution and habitat destruction (Harley et al, 2006; 30 Lotze et al., 2006). In such dynamic environments, pressures vary simultaneously, with different directions and magnitudes, over different time (short vs long-term) and spatial (global vs local) scales (Cabral et al., 2019). Hence, a fundamental challenge persists in elucidating how the various facets of global change affect coastal ecosystems responses at local and regional scales, and consequently the services they provide to human societies. Recurrent ecological monitoring programs conducted at large 35 spatial scales are valuable to assess ecosystem status and the pressures affecting important properties (e.g., biodiversity, carbon and nutrient cycling), and to detect changes over multiple decades (Hofmann et al., 2013; Sukhotin and Berger, 2013). Yet, only a few ecological indicators are both integrative and sensitive enough to various disturbances while also exhibiting predictable responses with a low variability in its response (Dale and Beyeler, 2001; Niemi and McDonald, 2004).

Filter-feeding bivalves include important reef habitat forming species, which promote benthic-pelagic coupling and nutrient recycling (Ray and Fulweiler, 2020). Some of them such as mussels (*Mytilus* spp.) and oysters (*Crassostrea gigas*) represent 40 an economical value through aquaculture and are widely used as bioindicators for climate and environmental change as well as contaminants monitoring (Kanduč et al., 2018; Karlson and Faxneld, 2021; Mazaleyrat et al., 2022; Briand et al., 2023; Chahouri et al., 2023; Liénart et al., 2024b). The use of ecological indicators is the basis of ecosystem monitoring to detect early-warning signals in ecosystem changes or disturbances (Dale and Beyeler, 2001). Indeed, bivalve tissues records the environmental conditions of their sampling area over time. Since the end of the 1970s, the French monitoring network 45 “ROCCH” (“Réseau d’Observation de la Contamination CHimique”, coordinated by Ifremer) monitor chemical contamination in the environment along French coastlines by using oysters and mussels as bioindicators (Briand et al., 2018; Chahouri et al., 2023). The ROCCH network focuses on chemical contaminants, but its archived samples represent a large sample bank that can be useful for other ecological purposes, such as understanding long-term changes in ecosystem functioning and responses to global or local pressures.

50 Measuring elemental and isotopic composition of the main building blocks of life (i.e., carbon and nitrogen; C:N,  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ ) is relevant to understand nutrients origin, to determine organisms diet and food webs structure, and can serve as indicators of water quality over space and time (Glibert et al., 2018). The C:N ratio is mostly an indicator of bivalve condition and physiology, reflecting the balance between organisms’ requirements and elemental availability in the environment (i.e., ecological stoichiometry sensu Elser et al., 2003, N content is often used as a proxy for proteins, C for lipids). When measured 55 over long-term in bivalve tissues, such dataset allow scientists to understand ecosystems responses to changing physical-chemical environment and organic matter cycling and to track trends in climate changes and its effect on coastal ecosystems (Liénart et al., 2020, 2024b). We took advantage of the large-scale multi-decadal sample bank of the ROCCH to analyze C:N,



$\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  in mussels (*M. edulis*, *M. galloprovincialis*) and oysters (*C. gigas*) tissues from a set of 33 stations distributed along the French coasts. In this paper, we present a unique dataset of multi-decadal and multi-ecosystem carbon and nitrogen elemental and isotopic ratios from three widespread bivalve species. As long-term isotopic datasets reflect the impact of human activities, such as coastal pollution and habitat alteration, the data derived here can provide valuable information for conservation and management strategies for coastal areas, in helping make informed decisions to mitigate environmental threats and protect vulnerable ecosystems. Ultimately it could provide valuable input for developing predictive models of ecosystem response to future environmental changes and possibly forecast potential impacts of climate change and human activities on coastal ecosystems. Overall, this long-term dataset of isotopic values in suspension-feeders tissues provides insights into ecosystem dynamics and is essential for advancing scientific understanding in the face of ongoing environmental challenges.

## 2 Methods and data

### 2.1 The ROCCH: network and sampling procedure

Over the last four decades, the French national monitoring network for chemical contaminant, the “ROCCH” (“Réseau d’Observation de la Contamination CHimique”), coordinated by Ifremer (“Institut français de recherche pour l’exploitation de la mer”, <https://littoral.ifremer.fr/Reseaux-de-surveillance/Environnement/ROCCH-Reseau-d-Observation-de-la-Contamination-CHimique-du-littoral>) has been annually sampling bivalves as bioindicators of chemical contamination. The number of stations has varied since the initiation of the network, so has the sampling frequency. Approximately 150 stations are nowadays monitored along the French coastlines, which begins in 1979 for historical stations. Sampling took place once a year during winter, currently mid-February, with a tolerance of one tidal cycle before and after the target date, meaning a 6 weeks amplitude spreading from the end of January to the beginning of March on the overall sites of the network. Three different species are targeted, the Pacific oyster *Crassostrea gigas* and the blue mussels *Mytilus edulis* in the English Channel and Atlantic facades, and *M. galloprovincialis* in the Mediterranean facade.

The sampling protocol has been identical since the start of the monitoring and was designed to acquire bivalve samples with consistent and homogeneous characteristics for measuring chemical concentrations of contaminants. Bivalves are collected alive at fixed points (maximum tolerance of 180 m around the selected point) chosen away from known anthropogenic discharges. Bivalves are sourced either from wild beds or from dedicated rearing facilities, ensuring that they remain on site for at least 6 months before being sampled. The selected individuals are adults of the same species and of uniform size (30 to 60 mm long for mussels, 90 to 140 mm long for oysters, i.e., 2 to 3 years old). A minimum of 50 mussels or 10 oysters was required to constitute a representative pooled sample accounting for inter-individual variability and to get enough material for chemical analysis and long-term storage. Bivalves are then depurated for 18 to 26 hours in decanted seawater collected near the collection site. Next, once extracted from the shell, the whole bodies (i.e., total soft-tissues) of each individual are drained together for 30 minutes. The resulting samples are placed in clean glass containers (washed and baked for 8 hours at 450 °C)



90 covered with calcined aluminum foil, and sealed with a plastic lid and frozen ( $-20^{\circ}\text{C}$ ). Frozen samples are sent to the central laboratory in Nantes (France) where the samples are thawed, crushed in a stainless steel-bladed blender and freeze-dried. After chemical analysis of a sample aliquot, the containers are indefinitely stored at Nantes IFREMER center, at room temperature, protected from light, under moisture-regulation control to prevent moisture pick-up. The original protocol (Grouhel 2023, in French) is available at <https://doi.org/10.13155/97878>.

## 95 2.2 Study sites and bivalve dataset

In the present dataset, bivalve samples were selected from 33 stations of the ROCCH network distributed along the three French sea facades: the English Channel (E.C., 10 stations), the Atlantic Ocean (A.O., 17 stations) and the Mediterranean Sea (M.S., 6 stations; Figure 1, Table 1). The stations were selected to span over a wide diversity of geomorphological and environmental conditions encountered by bivalves along the French coastlines. Ecosystems vary from open-sea coasts (rocky or sandy littoral shores), open or semi-enclosed bays and rias, lagoon (shallow water bodies almost closed by narrow landforms), and estuary mouths, ranging from eutrophic to oligotrophic systems, under temperate oceanic or Mediterranean climates, and along gradients of river influence from the main French rivers (Seine, Loire, Garonne, Dordogne, Rhône; Table 2). The complete dataset consist in 1141 bivalve winter samples for time series spanning 11 to 40 years (depending on stations) over the period 1981-2021 (Table 1; <https://doi.org/10.17882/100583>, Liénart et al., 2024a).


## 105 2.3 Sample analysis and data quality

Aliquots of 400-700  $\mu\text{g}$  of each dry and grounded archived sample of bivalve tissues (not acidified) were analyzed for carbon and nitrogen elemental and isotopic composition (C, N,  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ ) at the Center for Physical Science and Technology (Vilnius, Lithuania) with a Flash EA 1112 Series Elemental Analyzer (Thermo Finnigan) connected to a DeltaV Advantage Isotope Ratio Mass Spectrometer (Thermo Fisher). Carbon and nitrogen elemental values are expressed in percent (%) dry weight. Isotope values are expressed using the conventional delta notation:  $\delta^{13}\text{C}_{\text{sample}}$  or  $\delta^{15}\text{N}_{\text{sample}} = [(R_{\text{sample}}/R_{\text{standard}}) - 1]$ , where  $R = {}^{13}\text{C}/{}^{12}\text{C}$  or  ${}^{15}\text{N}/{}^{14}\text{N}$ , in per mil deviation (‰) from international standards, Vienna Pee Dee belemnite for  $\delta^{13}\text{C}$  and atmospheric  $\text{N}_2$  for  $\delta^{15}\text{N}$ . Mass spectrometer was calibrated against external certified standards (Caffeine IAEA600, USGS24 graphite), and internal standards (casein, glycine) were added before and every 10 samples within each batch of samples to control analytical performance.

115 Analytical precision was always better than 0.2 ‰ for  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  and 0.1  $\text{mol mol}^{-1}$  for C:N ratio (median of the standard deviations of each standard used for all batches: 0.12 ‰ for  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  and 0.04  $\text{mol mol}^{-1}$  for C:N ratio). Each sample was usually analyzed once and for some peculiar values that seemed too high/low or outside of the average trend, samples were reanalyzed to verify the values. If values were similar ( $< 0.5$  ‰), we chose the average value of both analyses; otherwise, we chose the value associated with the best analytical precision. The dataset presented here consists in a single value for a given year at a given station (the standard deviation associated with repeated measures is not presented). One data point was



125 considered as an outlier (reanalysis confirmed the value): the station 14-PALL in 2014 exhibited unexpected low values of  $\delta^{13}\text{C}$  ( $-26.96\text{‰}$ ) and high  $\delta^{15}\text{N}$  ( $10.26\text{‰}$ ) compare to the average values of this station ( $\delta^{13}\text{C} = -18.71 \pm 0.75\text{‰}$ ;  $\delta^{15}\text{N} = 8.41 \pm 0.93\text{‰}$ ), suggesting potential issues in the sampling and/or storing. This value was therefore removed from average calculations and statistical tests. All the other values presented in this data set are considered analytically valid and were scientifically validated in regard of expert knowledge of each ecosystem.



130 **Figure 1:** Map of the 33 stations from the three sea facades of France analyzed for bivalve's elemental and isotopic ratios. Bivalve species are the mussels *Mytilus edulis* and *Mytilus galloprovincialis* and the oyster *Crassostrea gigas*. The main rivers are presented in blue. Basemap source: Esri, HERE, © OpenStreetMap contributors, and the GIS User Community.



135

**Table 1: Metadata description for the 33 stations analyzed for bivalve's carbon and nitrogen elemental and isotopic ratios. Species studied are the mussels *Mytilus edulis* and *Mytilus galloprovincialis* and the oyster *Crassostrea gigas*. Station 31-CART represent two stations merged over time (distant from 720m). The geographic coordinate system is WGS84.**

Sea Facade	Station ID	Station name	ROCCH codes	latitude	longitude	Species	Time series	n years	Missing years
English Channel	1-AMBL	Ambleteuse	002-P-032	50.807366667	1.595466667	<i>M. edulis</i>	1982-2021	36	1984, 1985, 1986, 1990
	2-PSQT	Pointe de St Quentin	006-P-009	50.2808	1.52853333	<i>M. edulis</i>	1988-2021	34	NA
	3-VARE	Varengeville	008-P-013	49.9207688241	0.9820028963	<i>M. edulis</i>	1982-2021	37	1984, 1985, 1986
	4-ANTI	Antifer - digue	010-P-014	49.649085114	0.1503265738	<i>M. edulis</i>	1983-2021	36	1984, 1985, 1986
	5-CLHV	Cap de la Hève	010-P-055	49.5074135756	0.0619953224	<i>M. edulis</i>	1983-2017	29	1984, 1986, 1987, 1998, 1999, 2000
	6-VILL	Villerville	011-P-005	49.4040781596	0.1236660472	<i>M. edulis</i>	1982-2021	36	1984, 1985, 1986
	7-OUIS	Ouistreham	010-P-120	49.2940709028	-0.24800513	<i>M. edulis</i>	1994-2021	27	2004
	8-PBES	Port en Bessin	013-P-001	49.3515665687	-0.7531834463	<i>M. edulis</i>	1981-2021	37	1983, 1984, 1988, 2007
	9-BDVG	Bdv Grandcamp ouest	014-P-007	49.38633333	-1.101266667	<i>M. edulis</i>	1981-2021	37	1984, 1985, 1992, 1996
	10-PLAN	Pen al Lann	034-P-001	48.665109	-3.8944	<i>C. gigas</i>	1982-2021	37	1984, 1985, 1986
Atlantic Ocean	11-PASS	Le Passage (b)	039-P-007	48.391067	-4.384965	<i>C. gigas</i>	2001-2021	21	NA
	12-PERS	Persuel	039-P-093	48.2938273007	-4.5500595819	<i>C. gigas</i>	2001-2021	21	NA
	13-AULN	Aulne rive droite	039-P-124	48.281083	-4.260048	<i>C. gigas</i>	1982-2021	38	1984, 1985
	14-GUIL	Le Guilvin	060-P-001	47.56765	-2.9338	<i>C. gigas</i>	1982-2021	39	1986
	15-PEBE	Pen Bé	066-P-003	47.4306639649	-2.4679991263	<i>M. edulis</i>	1982-2021	40	NA
	16-CHEM	Pointe de Chemoulin	070-P-102	47.234632	-2.297076	<i>M. edulis</i>	1981-2021	39	1982, 1983
	17-JOAL	Joalland (b)	070-P-006	47.1568267957	-2.2224877474	<i>C. gigas</i>	2010-2021	11	NA
	18-BOUR	Bourgneuf - Coupelasse	071-P-065	47.0123255923	-2.0229805005	<i>C. gigas</i>	1984-2021	37	1985
	19-NOIR	Noirmoutier - Gresse-loup	071-P-068	46.95066	-2.146303	<i>C. gigas</i>	1983-2021	38	1985
	20-RIVE	Rivedoux	076-P-032	46.163319	-1.27077	<i>C. gigas</i>	1981-2021	40	1986
	21-PALL	Les Palles	080-P-004	45.9674774781	-1.1414394441	<i>C. gigas</i>	1982-2021	39	1983
	22-BOYV	Boyardville	080-P-033	45.9638097847	-1.2259409199	<i>C. gigas</i>	1984-2021	38	NA
	23-PONT	Pontailiac	084-P-015	45.6251372031	-1.0560967723	<i>C. gigas</i>	1984-2021	38	NA
	24-FOSS	La Fosse	085-P-007	45.475634584	-0.9845921143	<i>C. gigas</i>	1981-2021	40	2018
	25-FERR	Cap Ferret	087-P-013	44.6439465896	-1.2412452987	<i>C. gigas</i>	1982-2021	39	2005
	26-JACQ	Les Jacquets	088-P-067	44.7222821428	-1.1945794691	<i>C. gigas</i>	1982-2021	38	1983, 1985
	27-COMP	Comprian	088-P-069	44.6839491697	-1.08457666	<i>C. gigas</i>	1982-2021	40	NA
Mediterranean Sea	28-BANY	Banyuls - Labo Arago	094-P-008	42.4806130241	3.1388780428	<i>M. galloprovincialis</i>	1981-2021	32	1982, 1986, 1987, 1988, 1989, 1990, 2003
	29-HRLT	Embouchure de l'Hérault	095-P-026	43.2756325016	3.4405360127	<i>M. galloprovincialis</i>	1985-2021	31	1986, 1987, 1997, 1998, 2016
	30-STMM	Les Stes Maries de la mer	106-P-018	43.4439790562	4.4205519936	<i>M. galloprovincialis</i>	1982-2021	38	1984, 1986
	31-CART	Anse de Carteau	109-P-025 / -027	43.3756486632	4.8755622202	<i>M. galloprovincialis</i>	1982-2021	37	1986, 1996, 2007
	32-COUR	Cap Couronne	111-P-002	43.3239826009	5.0539000233	<i>M. galloprovincialis</i>	1983-2021	37	1986, 1996
	33-POME	Pomègues Est	111-P-025	43.2673170789	5.3005726112	<i>M. galloprovincialis</i>	1998-2021	24	NA

140





145

**Table 2: Environmental characteristics of the 33 stations studied for bivalves elemental and isotopic ratios. Average values for salinity, water temperature and chlorophyll a were calculated, when available, over the studied period from monitoring stations located nearby bivalves sampling stations, with data were retrieved from Surval (<https://surval.ifremer.fr> - Ifremer) and Somlit (<https://www.somlit.fr/> - INSU) databases. Average annual flow rates of the main rivers were calculated over the study period from the HydroPortail database (<https://hydro.eaufrance.fr/>). Trophic status was defined for each station based on literature (Liénaert et al., 2017; Lheureux et al., 2023) and the knowledge of local experts and reflect the average status over the studied period.**

Sea Facade	Station ID	Ecosystem	Ecosystem type	Tidal range	Salinity (mean±sd)	Water temperature (°C) (mean±sd)	Chlorophyll a (µg L <sup>-1</sup> ) (mean±sd)	Trophic status	Main river influence	Annual flow rates (m <sup>3</sup> s <sup>-1</sup> ) (mean±sd)	Additional local river influence
English Channel	1-AMBL	Eastern English Channel	Littoral	Megatidal	34 ± 1,0	12,7 ± 4,4	4,8 ± 4,6	Eutroph	Seine	499 ± 332	Liane
	2-PSQT	Eastern English Channel	Littoral	Megatidal	33 ± 1,5	13,0 ± 4,7	8,2 ± 7,8	Eutroph	Seine	449 ± 332	Somme
	3-VARE	Eastern English Channel	Littoral	Megatidal	33 ± 1,1	13,4 ± 4,5	2,2 ± 2,3	Eutroph	Seine	449 ± 332	
	4-ANTI	Bay of Seine	Open bay	Macrotidal	32 ± 1,8	14,3 ± 4,3	4,8 ± 7,1	Eutroph	Seine	449 ± 332	
	5-CLHV	Bay of Seine	Estuary mouth	Macrotidal	29 ± 3,5	12,9 ± 4,4	4,0 ± 4,7	Eutroph	Seine	449 ± 332	
	6-VILL	Bay of Seine	Estuary mouth	Macrotidal	29 ± 3,5	12,9 ± 4,4	4,0 ± 4,7	Eutroph	Seine	449 ± 332	
	7-OUIS	Bay of Seine	Open bay	Macrotidal	33 ± 0,8	15,2 ± 4,2	5,4 ± 5,2	Eutroph	Seine	449 ± 332	Orne
	8-PBES	Bay of Seine	Open bay	Macrotidal	34 ± 0,5	13,8 ± 4,4	2,8 ± 2,9	Eutroph	Seine	449 ± 332	
	9-BDVG	Bay of Veys	Open bay	Macrotidal	33 ± 0,7	13,8 ± 4,2	2,6 ± 2,5	Eutroph	Seine	449 ± 332	Vire
	10-PLAN	Western English Channel	Ria	Macrotidal	35 ± 0,2	12,9 ± 2,2	0,9 ± 0,8	Mesotroph	NA	-	
Atlantic Ocean	11-PASS	Bay of Brest	Semi-enclosed ria	Macrotidal	27 ± 7,5	12,9 ± 4,1	NA ± NA	Mesotroph	Elorn	6 ± 5	
	12-PERS	Bay of Brest	Semi-enclosed ria	Macrotidal	35 ± 0,6	13,4 ± 2,8	1,1 ± 0,9	Mesotroph	Aulne/Elorn	-	
	13-AULN	Bay of Brest	Semi-enclosed ria	Macrotidal	23 ± 8,7	12,8 ± 4,4	2,3 ± 2,7	Mesotroph	Aulne	27 ± 29	
	14-GUIL	Morbihan Gulf	Semi-enclosed bay	Macrotidal	33 ± 2	14,4 ± 3,7	1,8 ± 2,0	Mesotroph	Auray	2,8 ± 3,1	
	15-PEBE	Bay of Vilaine	Semi-enclosed bay	Macrotidal	32 ± 3	14,8 ± 3,9	4,6 ± 4,5	Eutroph	Vilaine	27 ± 34	
	16-CHEM	Loire Estuary	Estuary mouth	Macrotidal	31 ± 3,6	13,9 ± 3,6	3,4 ± 4,9	Mesotroph	Loire	833 ± 688	
	17-JOAL	Loire Estuary	Estuary mouth	Macrotidal	31 ± 3,6	13,9 ± 3,6	3,4 ± 4,9	Mesotroph	Loire	833 ± 688	
	18-BOUR	Bay of Bourgneuf	Semi-enclosed bay	Macrotidal	33 ± 2,3	13,8 ± 3,9	2,7 ± 3,9	Mesotroph	Loire	833 ± 688	
	19-NOIR	Bay of Bourgneuf	Semi-enclosed bay	Macrotidal	33 ± 2,3	13,8 ± 3,9	2,7 ± 3,9	Mesotroph	Loire	833 ± 688	
	20-RIVE	Charentais Sounds	Semi-enclosed sound	Macrotidal	33 ± 1,9	15,1 ± 4,3	2,2 ± 2,2	Mesotroph	Sèvre Niortaise	12 ± 13	
	21-PALL	Charentais Sounds	Semi-enclosed sound	Macrotidal	27 ± 7,1	14,5 ± 4,8	4,6 ± 7,0	Mesotroph	Charente	65 ± 66	Sèvre Niortaise
	22-BOYV	Charentais Sounds	Semi-enclosed sound	Macrotidal	33 ± 2,5	14,9 ± 4,3	3,2 ± 4,1	Mesotroph	Charente	65 ± 66	Sèvre Niortaise
	23-PONT	Gironde Estuary	Estuary mouth	Macrotidal	26 ± 5	15,9 ± 4,2	1,9 ± 1,3	Eutroph	Garonne / Dordogne	539 ± 404	
	24-FOSS	Gironde Estuary	Estuary mouth	Macrotidal	26 ± 5	15,9 ± 4,2	1,9 ± 1,3	Eutroph	Garonne / Dordogne	259 ± 199	
	25-FERR	Arcachon Lagoon	Semi-enclosed lagoon	Mesotidal	34 ± 0,8	15,2 ± 3,7	1,7 ± 1,1	Mesotroph	Leyre	17 ± 14	
	26-JACQ	Arcachon Lagoon	Semi-enclosed lagoon	Mesotidal	31 ± 2,8	15,8 ± 5,3	2,1 ± 1,1	Mesotroph	Leyre	17 ± 14	
	27-COMP	Arcachon Lagoon	Semi-enclosed lagoon	Mesotidal	31 ± 2,8	16,0 ± 5,3	1,8 ± 1,0	Mesotroph	Leyre	17 ± 14	
Mediterranean Sea	28-BANY	Gulf of Lion	Open bay	Microtidal	38 ± 0,7	16,7 ± 3,9	0,6 ± 0,6	Oligotroph	-	-	Tech / Têt
	29-HRLT	Gulf of Lion	Open bay	Microtidal	37 ± 2,4	17,7 ± 4,6	NA ± NA	Oligotroph	Hérault	35 ± 42	
	30-STMM	Gulf of Lion	Open bay	Microtidal	NA ± NA	NA ± NA	NA ± NA	Oligotroph	Rhône	1669 ± 782	
	31-CART	Gulf of Lion	Semi-enclosed bay	Microtidal	33 ± 3,7	16,5 ± 4,5	1,0 ± 1,1	Oligotroph	Rhône	1669 ± 782	
	32-COUR	Gulf of Lion	Open bay	Microtidal	38 ± 0,3	17,0 ± 3,5	0,4 ± 0,4	Oligotroph	Rhône	1669 ± 782	
	33-POME	Gulf of Lion	Open bay	Microtidal	38 ± 0,3	17,0 ± 3,5	0,4 ± 0,4	Oligotroph	Rhône	1669 ± 782	

## 150 2.4 Statistical analyses

Statistical analyses were performed with the R software (R Core Team 2022, 4.3.1 version).

Due to the nature of the dataset (i.e., only one of the three species was sampled for each sea façade, except for stations 15-PEBE and 16-CHEM from the Atlantic facade), it was not possible to test whether the observed differences in elemental and isotopic values were due to difference between species or sea facade, nor was it possible to ascertain if there was any interaction



155 between these two factors. Consequently, we have tested the effects of both factors (species and sea facade) independently.  
Therefore, particular attention should be given in interpreting the results of these tests. The effects of species (3 levels) and  
sea façade (3 levels) were tested independently for each elemental and isotopic variables over the entire dataset (all stations  
and time series) with non-parametric Kruskal-Wallis tests followed by Dunn *post-hoc* tests (R-package ‘PMCMRplus’ version  
1.9.10) since normality and homogeneity of variance (inspected using Shapiro-Wilk and Levene tests respectively) were not  
160 met, precluding the use of ANOVAs.

Monotonous temporal trends in bivalve elemental and isotopic times series were examined for each station using Mann-  
Kendall tests corrected for autocorrelation (R-package ‘modifiedmk’ version 1.6, Patakamuri and O’Brien, 2021). Linear  
models were applied to calculate the value of the slope (in ‰ decade<sup>-1</sup>) for each time series (note that all the time series do not  
have the same number of years). Complete time series were tested for shifts allowing for the detection of one unique shift per  
165 time series (R-package ‘cpm’ version 2.3, Gordon, 2015).

### 3 Data characteristics and general overview

This dataset illustrates the temporal and spatial variability of carbon and nitrogen elemental and isotopic ratios (C:N,  $\delta^{13}\text{C}$ ,  
 $\delta^{15}\text{N}$ ) of three species of bivalve along the French coasts from 1981 to 2021. Overall, the average  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values of  
bivalves were significantly lower for the stations of the Mediterranean Sea compared with the stations of the English Channel  
170 and the Atlantic Ocean. We observed that stations closest to river mouths displayed the lowest average  $\delta^{13}\text{C}$  values compared  
to those more distant from river influence. Over the last 10 to 40 years, nearly all stations exhibited a significant decrease in  
 $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ . There was no spatial nor temporal pattern in C:N but values differed between bivalve genus (mussel vs oyster).  
It is important to note that, due to the nature of the dataset (one species present for each sea façade), it is impossible to determine  
if the differences observed in elemental and isotopic values between the three species of bivalves and between the three sea  
175 facades are due to species effect or spatial differences (see 2.4 and details hereafter).


#### 3.1 Taxonomic patterns

Overall, bivalve  $\delta^{13}\text{C}$  ranged from -23.29 to -16.98 ‰,  $\delta^{15}\text{N}$  from 2.47 to 13.15 ‰ and C:N from 4.13 to 8.12 mol mol<sup>-1</sup>. There  
was a significant difference in  $\delta^{13}\text{C}$  ( $p < 0.001$ ) between *M. galloprovincialis* (-20.05 ± 0.98 ‰) and the other two species *M.*  
*edulis* (-19.38 ± 1.04 ‰) and *C. gigas* (-19.25 ± 1.06 ‰), which were not significantly different (Figure 2). There was a  
180 significant difference in  $\delta^{15}\text{N}$  ( $p < 0.001$ ) between the three species of bivalves (*C. gigas*: 8.42 ± 1.09 ‰; *M. edulis* 9.41 ± 1.19  
‰; *M. galloprovincialis* 5.70 ± 1.49 ‰) and C:N ( $p < 0.001$ ; *C. gigas*: 5.77 ± 0.62 mol mol<sup>-1</sup>; *M. edulis* 5.07 ± 0.31 mol mol<sup>-1</sup>;  
*M. galloprovincialis* 4.65 ± 0.23 mol mol<sup>-1</sup>; Figure 2). Since the mussel *M. galloprovincialis* is exclusively present in the  
Mediterranean Sea and the two other species, *M. edulis* and *C. gigas* are mostly present in the English Channel and Atlantic  
Ocean, respectively, it is difficult to separate the species factor from the spatial factor (see 3.2) in elemental and isotopic values





185 (especially in  $\delta^{13}\text{C}$  where *M. galloprovincialis* differs significantly). However, within the Atlantic facade where mostly oysters are sampled (15/17 stations), two stations are sampled for mussels (stations 15-PEBE and 16-CHEM), hence species effect can be tested for this facade. There was no significant differences in  $\delta^{13}\text{C}$  between the two species (*-C. gigas*:  $-19.24 \pm 1.07$  ‰; *M. edulis*:  $-19.36 \pm 1.02$  ‰), but  $\delta^{15}\text{N}$  and C:N values differed significantly ( $p < 0.001$ ) between oysters ( $8.43 \pm 1.12$  ‰;  $5.76 \pm 0.62$  mol mol<sup>-1</sup>) and mussels ( $10.54 \pm 0.88$  ‰;  $4.78 \pm 0.19$  mol mol<sup>-1</sup>). Note that the unequal sample sizes (15 stations  
 190 for oyster, 2 stations for mussels) can reduce the power of the test and the ability to detect real differences.



195 **Figure 2: Boxplots of  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$  and C:N values of bivalves for the 33 stations of the three French sea facades over the period 1981-2021 (*Crassostrea gigas*, *Mytilus edulis*, *Mytilus galloprovincialis*). The lower and upper edges of the box indicate the 25th and 75th percentiles and the middle line the median. The whiskers indicate the maximum/minimum values and the open circle the extreme values.**

### 3.2 Spatial patterns

#### 3.2.1 Among sea facade

The three sea facades showed significant difference in bivalves  $\delta^{13}\text{C}$  ( $p < 0.001$ ). The Mediterranean Sea had lower values ( $-20.05 \pm 0.98$  ‰), than the other two sea facades, the English Channel ( $-19.39 \pm 1.04$  ‰) and the Atlantic Ocean ( $-19.26 \pm 1.06$  ‰) which were not significantly different (Figure 2). There was also a significant difference in bivalve  $\delta^{15}\text{N}$  among the three  
 200



205 sea facades ( $p < 0.01$ ) showing a latitudinal gradient with high  $\delta^{15}\text{N}$  values in the English Channel ( $9.02 \pm 1.07 \text{‰}$ ), intermediate values in the Atlantic Ocean ( $8.71 \pm 1.30 \text{‰}$ ), and low values in the Mediterranean Sea ( $5.70 \pm 1.49 \text{‰}$ ; Figure 2). Bivalve C:N ratios were also significantly different among the three sea facades ( $p < 0.001$ ), with high values in the English Channel ( $5.24 \pm 0.42 \text{ mol mol}^{-1}$ ) and Atlantic Ocean ( $5.63 \pm 0.67 \text{ mol mol}^{-1}$ ) and low values in the Mediterranean Sea ( $4.65 \pm 0.23 \text{ mol mol}^{-1}$ ; Figure 2).

### 3.2.2 Among stations

The lowest  $\delta^{13}\text{C}$  values were mostly observed in bivalves from the stations inside or close to the main river mouths: 5-CLHV ( $-20.13 \pm 0.70 \text{‰}$ ) and 6-VILL ( $-20.60 \pm 1.03 \text{‰}$ ) for the Seine river; 16-CHEM ( $-20.15 \pm 0.74 \text{‰}$ ) and 17-JOAL ( $-19.90 \pm 0.60 \text{‰}$ ) for the Loire river; 24-FOSS ( $-20.71 \pm 1.04 \text{‰}$ ) for the Gironde estuary; 30-STMM ( $-20.34 \pm 0.81 \text{‰}$ ) and 31-CART  
210 ( $-20.58 \pm 1.14 \text{‰}$ ) for the Rhône river (Figure 2). The lowest  $\delta^{15}\text{N}$  values were mostly observed for bivalve from the Mediterranean Sea stations (all below  $< 6 \text{‰}$ ) with minimum average values for station 28-BANY ( $4.27 \pm 0.94 \text{‰}$ ) and maximum for station 30-STMM ( $5.91 \pm 0.75 \text{‰}$ ). The highest  $\delta^{15}\text{N}$  values were observed for stations under the Loire and Seine River plumes ( $> 9 \text{‰}$ ) e.g., 16-CHEM ( $10.90 \pm 0.90 \text{‰}$ ), 17-JOAL ( $9.79 \pm 0.43 \text{‰}$ ), 5-CLHV ( $9.41 \pm 1.08 \text{‰}$ ), 6-VILL ( $9.78 \pm 1.26 \text{‰}$ ), and locally under the influence of smaller rivers (Figure 2), e.g., 9-BDVG ( $10.16 \pm 0.52 \text{‰}$ ), 13-AULN ( $9.81$   
215  $\pm 0.42 \text{‰}$ ), 15-PEBE ( $10.19 \pm 0.71 \text{‰}$ ). This pattern is less clear for the Garonne and Dordogne rivers (Gironde estuary) where  $\delta^{15}\text{N}$  shows intermediate values (23-PONT ( $7.74 \pm 0.60 \text{‰}$ ), 24-FOSS ( $8.04 \pm 0.73 \text{‰}$ )). Bivalve C:N ratios did not show any clear spatial pattern between stations and seemed mostly linked with genus/species effects (differing between facades, Figure 2; see also 3.1).

Interestingly, for stations 16-CHEM and 17-JOAL, both located at the mouth of the Loire river and sampled for mussels and  
220 oysters, respectively, the  $\delta^{13}\text{C}$  signal did not differ significantly (mussels:  $-19.95 \pm 0.74 \text{‰}$ ; oysters:  $-19.90 \pm 0.60 \text{‰}$ ) over the same time period (2010-2021), despite being different species and genus. As for the comparison among the Atlantic façade (see 3.1),  $\delta^{15}\text{N}$  values significantly differed between the two species (mussels:  $10.30 \pm 0.44 \text{‰}$ ; oysters:  $9.79 \pm 0.43 \text{‰}$ ), however, the C:N ratio was not significantly different between the two species for this specific area (mussels:  $4.74 \pm 0.19 \text{‰}$ ; oysters:  $5.16 \pm 0.64 \text{‰}$ ). This suggests that spatial location could be the main driver of bivalve carbon isotope signal but that  
225 nitrogen is likely influenced by both global and local processes, including bivalve physiology, that compensate for species difference.

### 3.2.3 Key messages

Bivalves' elemental and isotopic values were generally lower for the stations of the Mediterranean Sea compared to those located in the Atlantic Ocean and English Channel. We found significant differences in carbon and nitrogen elemental and  
230 isotopic values between species (see 3.1), each of which is present in only one sea facade (with a few exceptions), nevertheless,




235 this difference is also linked with spatial variability. Indeed, C and N baselines (for isotope signal) and trophic status (i.e.,  
nutrient and chlorophyll amounts) vary between each sea facade (oligotrophic Mediterranean Sea vs. meso- to eutrophic  
Atlantic Ocean and English Channel; Table 2) but also locally (i.e., proximity to river mouth). To overcome the lack of  
nutrients available in the Mediterranean Sea, some specific groups of phytoplankton (i.e., diazotrophs) uses  $^{15}\text{N}$ -depleted  
240 atmospheric nitrogen as N source which decreases both the  $\delta^{15}\text{N}$  of the overall particulate organic matter (Kerhervé et al.,  
2001; Wannicke et al., 2010; Landrum et al., 2011; Liénart et al., 2017) and, by trophic propagation, the low  $\delta^{15}\text{N}$  signal is  
reflected in bivalves (Liénart et al., 2022, 2023). There is a clear difference in  $\delta^{15}\text{N}$  baseline between the Mediterranean sea  
and the other French sea facades (Liénart et al., 2017). Generally, there was a clear difference in  $\delta^{13}\text{C}$  signal close to river  
mouths, decreasing along river plumes, and in  $\delta^{15}\text{N}$  for some of the main (mostly eutrophicated) rivers. Enriched  $^{15}\text{N}$  signal of  
245 nutrients coming from rivers with extensive watershed agricultural activities and urban outlets is reflected in the high  $\delta^{15}\text{N}$   
values of bivalves sampled close to river mouths (Fukumori et al., 2008; Thibault et al., 2020). Similarly, low carbon isotopic  
values for bivalves sampled close to river mouths are mostly linked to the inputs of continental particulate material bearing  
this specific signal (Liénart et al., 2017).

### 3.3 Temporal patterns

#### 245 3.3.1 pluri-decadal $\delta^{13}\text{C}$ dataset

Bivalves  $\delta^{13}\text{C}$  showed a significant decrease over the period 1981-2021 for 82 % of the stations (27/33 stations; Figure 3). The  
Mediterranean Sea exhibited a significant decrease for all stations (6/6 stations), the Atlantic Ocean for 82 % of the stations  
(14/17 stations) and 70 % of the stations in the English Channel (7/10 stations). The average decrease over the total period was  
consistent across all facades, averaging  $-0.58 \pm 0.28$  ‰ per decade, ranging from  $-0.22$  to  $-1.21$  ‰ per decade (i.e.,  $-4.85$  to  $-$   
250  $0.89$  ‰ over the last 40 years) when the slopes were significant. This decrease in bivalves  $\delta^{13}\text{C}$  was more pronounced for the  
shorter time series starting from late 1990's (17-JOAL, 2010-2021; 11-PASS and 12-PERS, 2001-2021; 33-POME, 1998-  
2021; 7-OUIS, 1994-2021; Figure 3). Overall, the slopes were more pronounced for the Mediterranean Sea and for the stations  
in the western Bay of Seine, i.e., under the influence of the Seine plume (stations 7 to 10). Some of the stations showed strong  
interannual variability in bivalves  $\delta^{13}\text{C}$  (e.g., 1-AMBL, 6-VILL, 24-FOSS, Figure 3).

255 Shifts in bivalve  $\delta^{13}\text{C}$  time series were detected for 79 % of the stations (26/33 stations), equally represented in each sea facade.  
The shifts occurred mostly around the year 1999-2000 (6/26 stations) or after, around the years  $2006 \pm 1$  year and  $2012 \pm 1$   
year, with no specific pattern per façade. In the Mediterranean Sea most of the shifts occurred before 2000. The earliest shift  
was in 1989 (31-CART) and two in 1999, the latest in 2014 (15-PEBE). Usually, no shift was detected at the stations close to  
river mouths, i.e., Seine (5-CLHV, 6-VILL), Loire (16-CHEM, 17-JOAL), Gironde (24-FOSS) estuaries, and at some of the  
260 mouths of smaller rivers (e.g., 15-PEBE, Vilaine river; 29-HRLT, Hérault river).



**Figure 3: Bivalves  $\delta^{13}\text{C}$  time series over the period 1981-2021 for the 33 stations of the three French sea facades (*Crassostrea gigas*, *Mytilus edulis*, *Mytilus galloprovincialis*). Black lines correspond to significant trends (Mann-Kendall tests,  $p$ -value  $< 0.05$ ), the slope value ( $\text{‰}$  per decade) appears in the upper right corner. Shifts are represented by the vertical dash line with the year mentioned on its down left. One outlier is shown in grey (21-PALL,  $-27\text{‰}$ ) but was not considered for statistical analyses.**



### 3.3.2 pluri-decadal $\delta^{15}\text{N}$ dataset


Bivalve  $\delta^{15}\text{N}$  also showed a significant decrease over the period 1981-2021 for 64 % of the stations (22/33 stations; Figure 4). The Mediterranean Sea exhibited a significant decrease for 83 % stations (5/6 stations), the English Channel for 80 % of the stations (8/10 stations) and only 53 % of the stations in the Atlantic Ocean (9/17 stations). Only one station showed a significant increase (25-FERR, 0.34 ‰ over 39 years). The average decrease was  $-0.44 \pm 0.20$  ‰ per decade, ranging from  $-0.18$  to  $-0.84$  ‰ per decade (i.e.,  $-0.72$  to  $-3.35$  ‰ over the last 40 years; Figure 4). However, this decrease was more pronounced in the English Channel ( $-0.50 \pm 0.21$  ‰) in the eastern part and close the Seine river mouth (e.g., 3-VARE, 4-ANTI, 6-VILL), and Mediterranean Sea ( $-0.54 \pm 0.20$  ‰) compared with the Atlantic Ocean ( $-0.32 \pm 0.08$  ‰) where the lowest slopes were near the Loire river mouth and the inner two stations of the Arcachon lagoon. The interannual variability was relatively low for  $\delta^{15}\text{N}$  with the exception of some stations (e.g., 1-AMBL, 6-VILL, 16-CHEM, 33-POME).

Shifts in bivalve  $\delta^{15}\text{N}$  time series were detected for 70 % of the stations (23/33 stations), mostly in the English Channel and Mediterranean Sea. The shifts occurred mostly around the year 1999-2000 (5/23 stations), then around the years 1995-1996 (4/23 stations) and  $2004 \pm 1$  year (4/23 stations), with shifts mostly occurring before 2000 in the Mediterranean Sea. For each sea facade, there were often shifts occurring the same year or in two close years for nearby stations (e.g., 15-PEBE, 2003 and 16-CHEM, 2004). The earliest shift was in 1991 (29-HRLT), the latest in 2013 (14-GUIL).

### 3.3.3 pluri-decadal C:N dataset

Bivalve C:N ratio showed significant trends for less than half of the stations (37 %) over the period 1981-2021 (10/33 stations; Figure 5). Most of the significant trends were increasing (24 %, 8/33 stations), exclusively in the Atlantic Ocean (6/33 stations) and the English Channel (2/33 stations). Only 6 % (2/33 stations) of the stations showed decreasing trends, all in the Mediterranean Sea. The average increase was  $0.27 \pm 0.37$  mol mol<sup>-1</sup> per decade, ranging from 0.05 to 0.38 mol mol<sup>-1</sup> per decade (i.e., 0.20 to 1.53 mol mol<sup>-1</sup> over the last 40 years; excluding 17-JOAL with a 1.45 mol mol<sup>-1</sup> increase for 11 years; Figure 5). The average decrease in the Mediterranean sea was  $-0.05 \pm 0.02$  mol mol<sup>-1</sup> per decade. Interannual variability in bivalve C:N was either very low (e.g., 7-OUIS, 16-CHEM, 33-POME) for the two mussel species, or very large (e.g., 12-PERS, 19-NOIR, 22-BOYV) for the oysters. The decreasing trends in C:N ratio were mostly linked to decrease in bivalves N % (6/8 stations), whereas only 3/8 stations showed a significant increase in C %.


Shifts in bivalve C:N time series were detected only for 33 % of the stations (11/33 stations), mostly in the Atlantic Ocean. Most of the shifts occurred after the year 2000 (7/11 stations). The earliest shift was in 1989 (27-COMP), the latest in 2014 (6-VILL, 22-BOYV). There was no apparent spatial pattern in the C:N shifts.



295

**Figure 4:** Bivalves  $\delta^{15}\text{N}$  time series over the period 1981–2021 for the 33 stations of the three French sea facades (*Crassostrea gigas*, *Mytilus edulis*, *Mytilus galloprovincialis*). Black lines correspond to significant trends (Mann-Kendall tests, p-value < 0.05), the slope value (‰ per decade) appears in the upper right corner. Shifts are represented by the vertical dash line with the year mentioned on its down left.





300

**Figure 5: Bivalves C:N time series over the period 1981-2021 for the 33 stations of the three French sea facades (*Crassostrea gigas*, *Mytilus edulis*, *Mytilus galloprovincialis*). Black lines correspond to significant trends (Mann-Kendall tests, p-value < 0.05), the slope value (% per decade) appears in the upper right corner. Shifts are represented by the vertical dash line with the year mentioned on its down left.**



### 305 3.3.4 Key messages

In summary, over the last 30 to 40 years, bivalve have shown a general decrease in both  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  for most stations of the three French sea facades. These trends are most likely linked with global processes occurring at large spatial scale such as climate change. Liénart et al. (2024) explained the decrease in  $\delta^{13}\text{C}$  is primarily associated with the increase in atmospheric  $\text{CO}_2$  concentrations due to fossil-fuel burning, which generates a decrease in atmospheric  $\text{CO}_2$   $\delta^{13}\text{C}$  values occurring since the industrial revolution in 1850 (the Suess effect; Keeling 1979; Gruber et al., 1999). This decrease in  $\delta^{13}\text{C}$  values propagates along marine food webs and is visible in organisms' tissues (Schloesser et al. 2009; Liénart et al., 2022, 2024b). However, other global factors such as rising temperatures or climate indices may also contribute to this overall decrease in  $\delta^{13}\text{C}$  through passive (e.g., Suess effect) or active processes related to bivalve physiology (e.g., change in fractionation; Liénart et al., 2024b). Decreasing  $\delta^{15}\text{N}$  trends are potentially linked to global factors, indirectly through a general decrease in nutrients and particles inputs from rivers (Milliman et al., 2008; Bauer et al., 2013). Since organic particles generated in areas with high human activities bears a high  $\delta^{15}\text{N}$  signal, the decrease of such  $^{15}\text{N}$ -enriched inputs is reflected in coastal water and marine fauna (Connolly et al., 2013). However, such changes are strongly influenced by regional and local factors such as regional climate, watershed activities and damming (Milliman et al., 2008) resulting in different slopes values in isotope time series close to river mouths (more pronounced for both C and N isotopes). The observed decrease in both isotopic ratios likely results from the cumulative, synergistic, or antagonistic effects of global and regional/local influences (Cabral et al., 2019). Such interacting pressures are more likely to occur in regions strongly affected by climate change, such as the Mediterranean Sea (Tuel and Eltahir, 2020). In this dataset, isotopes reveal both global (climate) effects, mostly through  $\delta^{13}\text{C}$ , and some local variability in the organic matter used by bivalves, as shown by the different  $\delta^{13}\text{C}$  (low) and  $\delta^{15}\text{N}$  (high) signals along the river-mouth-to-sea gradient. The decrease in bivalve  $\delta^{13}\text{C}$  and in  $\delta^{15}\text{N}$  over the past decades may indicate a decrease in nutrient and particles inputs from the rivers over the same period. Finally, as an indicator of bivalve condition and physiology (Elser et al., 2003), the increase in C:N ratio observed for most stations in the Atlantic and English Channel can be attributed to either an increase in carbon content (i.e., lipids), which is unlikely given rising temperatures and appear significant for only few of our stations, or to a decrease in nitrogen content (i.e., proteins, amino acids), as observed in our data and likely due to a decrease in N-nutrient availability in the environment which is consistent with the observed decrease in  $\delta^{15}\text{N}$ . The absence of trends at most of the stations and the large interannual variability of C:N ratio may result from the complex interplay between global and local environmental effects and bivalves physiology on this parameter.

## 4 Data availability

The dataset presented in this article and the metadata associated can be freely accessed on the SEANOE open access repository (<https://www.seanoe.org/>) under the DOI <https://doi.org/10.17882/100583> (Liénart et al., 2024a). The data gives the  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  (‰), C and N (%) and C:N ratio ( $\text{mol mol}^{-1}$ ), for each station and year, with sampling date and species name. For each station, it contains the sea façade, station ID, station name, ROCCH codes, latitude and longitude (geographic coordinate



system WGS84) as presented in Table 1. Carbon isotopic time-series are already published (Liénart et al., 2024b, except from station 17-JOAL) in a study focused on global effect on coastal ecosystems from large-scale anthropogenic and natural climate proxies, including the Suess effect, over the period 1981-2021. Bivalve  $\delta^{13}\text{C}$  data and  $\delta^{13}\text{C}$  corrected from the Suess effect are available under the permanent identifier <https://doi.org/10.6084/m9.figshare.24884871.v1> (Liénart et al., 2024c). The companion data from the ROCCH (contaminants) are available on the ‘Survival’ database (<https://surval.ifremer.fr>).

## 5 Conclusion and recommendation for use

By providing this dataset to the scientific community including caveats for interpreting such data spatially and over time, we expect it will be useful for numerous ecological studies. Such data is relevant to trace nutrients origin, to set accurate baselines to study organisms diets and food webs structure, and can be an indicator of water quality over space and time through bivalve physiology. Multi-decadal time-series allow scientists to understand coastal ecosystems responses to global change through biological and biogeochemical processes. The carbon isotope dataset is already part of a study where trends are thoughtfully interpreted in regard of global proxies for climate and anthropogenic changes and corrected for the Suess effect (Liénart et al., 2024b). Part of the observed  $\delta^{13}\text{C}$  trends were linked with the Suess effect, leading to shifts in isotope baseline over recent decades (i.e., Suess effect for  $\delta^{13}\text{C}$ ) rather than specific changes in ecosystem functioning. Hence, we would like to draw users attention on the need to correct for the Suess effect before comparing biological samples collected one or more decades apart as recommended in literature (Dombrosky 2020; Clark et al. 2021; Liénart et al., 2024b). Additionally, in order to take into account the low  $\delta^{13}\text{C}$  and high  $\delta^{15}\text{N}$  signal observed near river mouths, we advise to compare slope values rather than absolute values when assessing temporal changes between stations. Overall, the complex interplay between global and regional/local effects needs to be considered when interpreting time series. This long-term dataset of elemental and isotopic values in suspension-feeders tissues provides insights into ecosystem dynamics and hold broader significance for advancing scientific understanding in the face of ongoing environmental challenges.

### Author contribution

CL and NS are leaders of the project. NS, CL, AL, AGP, PLM, XdM, HB, SD and AIG participated in defining the scientific strategy and selection of archived samples for analysis. AF and NB prepared the samples for analysis. AnG carried out sample analysis. PLM and AGP are the curator of the ROCCH sample archive. CL and AF performed the statistical analysis. CL prepared the manuscript and figures and integrated the final contribution from all co-authors.

### 365 Competing interests

The authors declare that they have no competing interests.



## Disclaimer

370 Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Acknowledgements

This article is based on a collaborative work with the team of the ROCCH monitoring network coordinated by Ifremer. We  
375 thank the members of the ROCCH and of the different institutions, from the field workers and sample analysts to the coordinators, who made it possible to use these samples.

## Financial support

This research was funded by the Office Français de la Biodiversité (OFB) within the frame of the research project EVOLECO-  
380 BEST.

## References

- Bauer, J. E., Cai, W.-J., Raymond, P., Bianchi, T. S., Hopkinson, C. S., and Regnier, P. G.: The changing carbon cycle of the coastal ocean., *Nature*, 504, 61–70, <https://doi.org/10.1038/nature12857>, 2013.
- Briand, M. J., Herlory, O., Briant, N., Brach-Papa, C., Boissery, P., and Bouchouca, M.: The French Mussel Watch: More  
385 than two decades of chemical contamination survey in Mediterranean coastal waters, *Mar. Pollut. Bull.*, 191, 114901, <https://doi.org/10.1016/j.marpolbul.2023.114901>, 2023.
- Briant, N., Savoye, N., Chouvelon, T., David, V., Rodriguez, S., Charlier, K., Sonke, J. E., Chiffolleau, J. F., Brach-Papa, C., and Knoery J.: Carbon and nitrogen elemental and isotopic ratios of filter-feeding bivalves along the French coasts: An assessment of specific, geographic, seasonal and multi-decadal variations. *Sci. Total Environ.* 613–614: 196–207.  
390 [doi:10.1016/j.scitotenv.2017.08.281](https://doi.org/10.1016/j.scitotenv.2017.08.281), 2018.
- Bristow, L. A., Jickells, T. D., Weston, K., Marca-Bell, A., Parker, R., and Andrews J. E.: Tracing estuarine organic matter sources into the southern North Sea using C and N isotopic signatures. *Biogeochemistry*, 113: 9–22, <https://doi.org/10.1007/s10533-012-9758-4>, 2013.
- Cabral, H., Fonseca, V., Sousa, T., and Leal, M. C.: Synergistic effects of climate change and marine pollution: An overlooked  
395 interaction in coastal and estuarine areas, *Int. J. Environ. Res. Public Health*, 16, 1–17, <https://doi.org/10.3390/ijerph16152737>, 2019.
- Chahouri, A., Yacoubi, B., Moukrim, A., and Banaoui, A.: Bivalve molluscs as bioindicators of multiple stressors in the marine environment: Recent advances, *Cont. Shelf Res.*, 264, 105056, <https://doi.org/10.1016/j.csr.2023.105056>, 2023.



- Clark, C. T., Cape, M. R., Shapley, M. D., Mueter, F. J., Finney, B. P., and Misarti, N.: SuessR: Regional corrections for the effects of anthropogenic CO<sub>2</sub> on δ<sup>13</sup>C data from marine organisms, *Methods Ecol. Evol.*, 12, 1508–1520, <https://doi.org/10.1111/2041-210X.13622>, 2021.
- Cloern, J. E., Abreu, P. C., Carstensen, J., Chauvaud, L., Elmgren, R., Grall, J., Greening, H., Johansson, J. O. R., Kahru, M., Sherwood, E. T., Xu, J., and Yin, K.: Human activities and climate variability drive fast-paced change across the world's estuarine-coastal ecosystems, *Glob. Chang. Biol.*, 22, 513–529, <https://doi.org/10.1111/gcb.13059>, 2016.
- Connolly, R. M., Gorman, D., Hindell, J. S., Kildea, T. N., and Schlacher, T. A.: High congruence of isotope sewage signals in multiple marine taxa, *Mar. Pollut. Bull.*, 71, 152–158, <https://doi.org/10.1016/j.marpolbul.2013.03.021>, 2013.
- Dale, V. H., and Beyeler, S. C.: Challenges in the development and use of ecological indicators. *Ecol. indic.*, 1(1), 3–10, [https://doi.org/10.1016/S1470-160X\(01\)00003-6](https://doi.org/10.1016/S1470-160X(01)00003-6), 2001.
- Dombrosky, J.: A ~1000-year <sup>13</sup>C Suess correction model for the study of past ecosystems, *Holocene*, 30, 474–478, <https://doi.org/10.1177/0959683619887416>, 2020.
- Elser, J. J., Acharya, K., Kyle, M., Cotner, J., Makino, W., Markow, T., Watts, T., Hobbie, S., Fagan, W., Schade, J., Hood, J., and Sterner, R. W.: Growth rate-stoichiometry couplings in diverse biota, *Ecol. Lett.*, 6, 936–943, <https://doi.org/10.1046/j.1461-0248.2003.00518.x>, 2003.
- Fry, B., and Sherr, E.: <sup>13</sup>C measurements as indicators of carbon flow in marine and freshwater ecosystems. *Contrib. Mar. Sci.*, 27:13–47, 1984.
- Fukumori, K., Oi, M., Doi, H., Takahashi, D., Okuda, N., Miller, T. W., Kuwae, M., Miyasaka, H., Genkai-Kato, M., Koizumi, Y., Omori, K., and Takeoka, H.: Bivalve tissue as a carbon and nitrogen isotope baseline indicator in coastal ecosystems, *Estuar. Coast. Shelf Sci.*, 79, 45–50, <https://doi.org/10.1016/j.ecss.2008.03.004>, 2008.
- Glibert, P. M., Middelburg, J. J., McClelland, J. W., and Vander Zanden, J. M.: Stable isotope tracers: Enriching our perspectives and questions on sources, fates, rates, and pathways of major elements in aquatic systems, *Limnol. Oceanogr.*, 1–32, <https://doi.org/10.1002/lno.11087>, 2018.
- Gordon J. R.: Parametric and Nonparametric Sequential Change Detection in R: The cpm Package. *J. Stat. Software.* 66(3), 1–20. <http://www.jstatsoft.org/v66/i03/>, 2015.
- Grouhel A.: Prescriptions techniques pour l'échantillonnage de mollusques du réseau national d'observation des contaminants chimiques (ROCCH), RST- RBE/CCEM/ROCCH 23-02, <https://doi.org/10.13155/97878>, 2023.
- Gruber, N., Keeling, C. D., Bacastow, R. B., Guenther, P. R., Lueker, T. J., Wahlen, M., Meijer, H. A. J., Mook, W. G., and Stocker, T. F.: Spatiotemporal patterns of carbon-13 in the global surface ocean and the oceanic Suess effect, *Global Biogeochem. Cycles*, 13: 307–335, <https://doi.org/10.1029/1999GB900019>, 1999.



- Harley, C. D. G., Hughes, A. R., Hultgren, K. M., Miner, B. G., Sorte, C. J. B., Thornber, C. S., Rodriguez, L. F., Tomanek,  
430 L., and Williams, S. L.: The impacts of climate change in coastal marine systems, *Ecol. Lett.*, 9, 228–241,  
<https://doi.org/10.1111/j.1461-0248.2005.00871.x>, 2006.
- Hofmann, G. E., Blanchette, C. A., Rivest, E. B., and Kapsenberg, L.: Taking the pulse of marine ecosystems: The importance  
of coupling long-term physical and biological observations in the context of global change biology. *Oceanography*, 26(3), 140-  
148, <https://www.jstor.org/stable/24862075>, 2013.
- 435 Kanduč, T., Šlejkovec, Z., Falnoga, I., Mori, N., Budič, B., Kovačić, I., Pavičić – Hamer, D., and Hamer, B.: Environmental  
status of the NE Adriatic Sea, Istria, Croatia: Insights from mussel *Mytilus galloprovincialis* condition indices, stable isotopes  
and metal(loid)s, *Mar. Pollut. Bull.*, 126, 525–534, <https://doi.org/10.1016/j.marpolbul.2017.09.052>, 2018.
- Karlson, A. M. L. and Faxneld, S.: Polycyclic aromatic hydrocarbons and stable isotopes of carbon and nitrogen in Baltic Sea  
blue mussels: Time series data 1981 – 2016, *Data Br.*, 35, 4–8, <https://doi.org/10.1016/j.dib.2021.106777>, 2021.
- 440 Keeling, C. D.: The Suess effect: <sup>13</sup>Carbon-<sup>14</sup>Carbon interrelations, *Environ. Int.*, 2, 229–300, [https://doi.org/10.1016/0160-4120\(79\)90005-9](https://doi.org/10.1016/0160-4120(79)90005-9), 1979.
- Kerhervé, P., Minagawa, M., Heussner, S., and Monaco, A.: Stable isotopes (C-13/C-12 and N-15/N-14) in settling organic  
matter of the northwestern Mediterranean Sea: biogeochemical implications, *Oceanol. Acta*, 24, S77–S85, 2001.
- Landrum, J. P., Altabet, M. A., and Montoya, J. P.: Basin-scale distributions of stable nitrogen isotopes in the subtropical  
445 North Atlantic Ocean: Contribution of diazotroph nitrogen to particulate organic matter and mesozooplankton, *Deep. Res. Part  
I Oceanogr. Res. Pap.*, 58, 615–625, <https://doi.org/10.1016/j.dsr.2011.01.012>, 2011.
- Lheureux, A., David, V., Del Amo, Y., Soudant, D., Auby, I., Bozec, Y., Conan, P., Ganthy, F., Grégori, G., Lefebvre, A.,  
Leynard, A., Rimmelin-Maury, P., Souchu, P., Vantrepote, V., Blondel, C., Cariou, T., Crispi, O., Cordier, M-A., Crouvoisier,  
M., Duquesne, V., Ferreira, S., Garcia, N., Gouriou, L., Grosteffan, E., Le Merrer, Y., Meteigner, C., Retho, M., Tournaire,  
450 M-P., and Savoye, N.: Trajectories of nutrients concentrations and ratios in the French coastal ecosystems: 20 years of changes  
in relation with large-scale and local drivers. *Sci. Tot. Envir.*, 857, 159619. <https://doi.org/10.1016/j.scitotenv.2022.159619>,  
2023.
- Liénard, C., Savoye, N., Bozec, Y., Breton, E., Conan, P., David, V., Feunteun, E., Grangeré, K., Kerhervé, P., Lebreton, B.,  
Lefebvre, S., L’Helguen, S., Mousseau, L., Raimbault, P., Richard, P., Riera, P., Sauriau, P.-G., Schaal, G., Aubert, F., Aubin,  
455 S., Bichon, S., Boinet, C., Bourasseau, L., Bréret, M., Caparros, J., Cariou, T., Charlier, K., Claquin, P., Cornille, V., Corre,  
A.-M., Costes, L., Crispi, O., Crouvoisier, M., Czamanski, M., Del Amo, Y., Derriennic, H., Dindinaud, F., Durozier, M.,  
Hanquiez, V., Nowaczyk, A., Devesa, J., Ferreira, S., Fomier, M., Garcia, F., Garcia, N., Geslin, S., Grossteffan, E., Gueux,  
A., Guillaudeau, J., Guillou, G., Joly, O., Lachaussée, N., Lafont, M., Lamoureux, J., Lecuyer, E., Lehodey, J.-P., Lemeille,  
D., Leroux, C., Macé, E., Maria, E., Pineau, P., Petit, F., Pujon-Pay, M., Rimmelin-Maury, P., and Sultan, E.: Dynamics of





- 460 particulate organic matter composition in coastal systems: A spatio-temporal study at multi-systems scale, *Prog. Oceanogr.*, 156, <https://doi.org/10.1016/j.pocean.2017.03.001>, 2017.
- Liénart, C., Garbaras, A., Qvarfordt, S., Sysoev, A. Ö., Högländer, H., Walve, J., Schagerström, E., Eklöf, J., and Karlson, A. M.: Long-term changes in trophic ecology of blue mussels in a rapidly changing ecosystem, *Limnol. Oceanogr.*, 66, 694–710, <https://doi.org/10.1002/lno.11633>, 2020.
- 465 Liénart, C., Garbaras, A., Qvarfordt, S., Walve, J., and Karlson, A. M. L.: Spatio-temporal variation in stable isotope and elemental composition of key-species reflect environmental changes in the Baltic Sea, *Biogeochemistry*, 157, 149–170, <https://doi.org/10.1007/s10533-021-00865-w>, 2022.
- Liénart, C., Tedengren, M., Garbaras, A., Lim, H., Chynel, M., Robinson, C., Meziane, T., and Karlson, A. M. L.: Diet quality determines blue mussel physiological status: A long-term experimental multi-biomarker approach, *J. Exp. Mar. Bio. Ecol.*,  
470 563, <https://doi.org/10.1016/j.jembe.2023.151894>, 2023.
- Liénart, C., Fournioux, A., Garbaras, A., Blanche, H., Briant, N., Dubois, S., Gangnery, A., Grouhel, A., Le Monier, P., Lheureux, A., de Montaudouin, X., and Savoye, N. : Bivalve multi-decadal (1981-2021) records of carbon and nitrogen elemental and isotopic ratios ( $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$  and C:N) of 33 coastal sites of France. SEANOE Sea scientific open data publication, <https://doi.org/10.17882/100583>, 2024a.
- 475 Liénart, C., Fournioux, A., Garbaras, A., Lheureux, A., Blanchet, H., Briant, N., Dubois, S. F., Gangnery, A., Grouhel Pellouin, A., Le Monier, P., de Montaudouin, X., and Savoye, N.: Bivalve tissues as a recorder of multidecadal global anthropogenic and climate-mediated change in coastal areas. *Limnol. Oceanogr. Letters*, <https://doi.org/10.1002/lol2.10399>, 2024b.
- Liénart, C., Fournioux, A., Garbaras, A., Lheureux, A., Blanchet, H., Briant, N., Dubois, S. F., Gangnery, A., Grouhel Pellouin, A., Le Monier, P., de Montaudouin, X., and Savoye, N. : Bivalves  $\delta^{13}\text{C}$  time series (1981-2021) 32 stations France. Figshare,  
480 <https://doi.org/10.6084/m9.figshare.24884871.v1>, 2024c.
- Lotze, H. K., Lenihan, H. S., Bourque, B. J., Bradbury, R. H., Cooke, R. G., Kay, M. C., Kidwell, S. M., Kirby, M. X., Peterson, C. H., and Jackson, J. B.: Depletion, degradation, and recovery potential of estuaries and coastal seas. *Science*, 312(5781), 1806-1809, <https://doi.org/10.1126/science.1128035>, 2006.
- Mazaleyrat, A., Normand, J., Dubroca, L., and Fleury, E.: A 26-year time series of mortality and growth of the Pacific oyster  
485 *C. gigas* recorded along French coasts, *Sci. Data*, 9, 392, <https://doi.org/10.1038/s41597-022-01511-2>, 2022.
- Milliman, J. D., Farnsworth, K. L., Jones, P. D., Xu, K. H., and Smith, L. C.: Climatic and anthropogenic factors affecting river discharge to the global ocean, 1951-2000, *Glob. Planet. Change*, 62, 187–194, <https://doi.org/10.1016/j.gloplacha.2008.03.001>, 2008.



- 490 Niemi, G. J., and McDonald, M. E.: Application of ecological indicators. *Annu. Rev. Ecol. Evol. Syst.*, 35, 89-111.  
<https://doi.org/10.1146/annurev.ecolsys.35.112202.130132>, 2004.
- Patakamuri, S., and O'Brien, N.: modifiedmk: Modified versions of Mann Kendall and Spearman's Rho Trend Tests, R package version 1.6, <https://CRAN.R-project.org/package=modifiedmk>, 2021.
- R Core Team: R: A language and environment for statistical computing. R Foundation for Statistical Computing, <https://www.R-project.org/>, 2022.
- 495 Ray, N. E., and Fulweiler, R. W.: Seasonal patterns of benthic-pelagic coupling in oyster habitats. *Mar. Ecol. Prog. Ser.*, 652, 95-109, <https://doi.org/10.3354/meps13490>, 2020.
- Schloesser, R. W., Rooker, J. R., Louchuarn, P., Neilson, J. D., and Secor, D. H.: Interdecadal variation in seawater  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  recorded in fish otoliths, *Limnol. Oceanogr.*, 54, 1665–1668, <https://doi.org/10.4319/lo.2009.54.5.1665>, 2009.
- Sukhotin, A., and Berger, V.: Long-term monitoring studies as a powerful tool in marine ecosystem research, *Hydrobio.*, 706,  
500 1–9, <https://doi.org/10.1007/s10750-013-1456-2>, 2013.
- Thibault, M., Duprey, N., Gillikin, D. P., Thébault, J., Douillet, P., Chauvaud, L., Amice, E., Munaron, J. M., and Lorrain, A.: Bivalve  $\delta^{15}\text{N}$  isoscapes provide a baseline for urban nitrogen footprint at the edge of a World Heritage coral reef, *Mar. Pollut. Bull.*, 152, 110870, <https://doi.org/10.1016/j.marpolbul.2019.110870>, 2020.
- Tuel, A., and Eltahir, E. A.: Why is the Mediterranean a climate change hot spot?, *J. Clim.*, 33(14), 5829-5843.  
505 <https://doi.org/10.1175/JCLI-D-19-0910.1>, 2020.
- Wannicke, N., Liskow, I., and Voss, M.: Impact of diazotrophy on N stable isotope signatures of nitrate and particulate organic nitrogen: case studies in the north-eastern tropical Atlantic Ocean, *Isotopes Environ. Health Stud.*, 46, 337–54, <https://doi.org/10.1080/10256016.2010.505687>, 2010.