



A post-processed carbon flux dataset for 34 eddy covariance flux sites across the Heihe River Basin, China

Xufeng Wang¹, Tao Che^{1*}, Jingfeng Xiao², Tonghong Wang^{1,3}, Junlei Tan¹, Yang Zhang¹, Zhiguo Ren¹, Liying Geng¹, Haibo Wang¹, Ziwei Xu⁴, Shaomin Liu⁴, Xin Li⁵

- 5 ¹ Key Laboratory of Remote Sensing of Gansu Province/Heihe Remote Sensing Experimental Research Station/Key Laboratory of Cryospheric Science and Frozen Soil Engineering, Northwest Institute of Eco-Environment and Resources, Chinese Academy of Sciences, Lanzhou 730000, China.
- ² Earth Systems Research Center, Institute for the Study of Earth, Oceans, and Space, University of New Hampshire, Durham, NH 03824, USA.
- 10 ³ School of Geography and Environmental Sciences, Northwest Normal University, Lanzhou 730000, China.
- ⁴ State Key Laboratory of Remote Sensing Science, School of Geography, Beijing Normal University, Beijing 100101, China.
- ⁵ Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing 100101, China.
- 15 *Correspondence to:* Tao Che (chetao@lzb.ac.cn)

Abstract. The eddy covariance (EC) technique is currently the most widely used method for measuring carbon exchange between terrestrial ecosystems and the atmosphere at the ecosystem scale. Using this technique, a regional carbon flux network comprising a total of 34 sites has been established in the Heihe River Basin (HRB) in Northwest China. This network has been measuring the net ecosystem exchange (NEE) of CO₂ for a variety of vegetation types. In this study, we compiled and post-processed half-hourly flux data from these 34 EC flux sites in the HRB to create a continuous, homogenized time series dataset. We employed standardized processing procedures to fill data gaps in meteorological and NEE measurements at half-hourly intervals. NEE measurements were also partitioned into gross primary production (GPP) and ecosystem respiration (Reco). Furthermore, half-hourly meteorological and NEE data were aggregated to daily, weekly, monthly, and yearly timescales. As a result, we produced a continuous carbon flux and auxiliary meteorological dataset, which includes 18 sites with continuous multi-year observations and 16 sites observed only during the 2012 growing season, amounting to a total of 1,513 site-months. Using the post-processed dataset, we explored the temporal and spatial characteristics of carbon exchange in the HRB. In the diurnal variation curve, GPP, NEE, and Reco peak later for ecosystems in the artificial oasis (cropland and wetland) compared to those outside the artificial oasis (grassland, forest, woodland, and Gobi/desert). Seasonal NEE, GPP, and Reco peak in early July for grassland, forest, woodland, and cropland but remain close to zero throughout the year for gobi/desert. In the last decade, NEE of wetlands significantly increased, while NEE for other ecosystems did not exhibit significant trends. Annual NEE, GPP, and Reco are significantly higher for sites inside the artificial/natural oasis compared to those outside the oasis. This post-processed carbon flux dataset has numerous applications, including exploring the carbon exchange characteristics of alpine and arid ecosystems, analyzing ecosystem responses to climate extremes, conducting cross-site synthesis from regional to global scales, supporting regional and global upscaling studies, interpreting and calibrating remote sensing products, and evaluating and calibrating carbon cycle models.



40 **1 Introduction**

Terrestrial ecosystems absorb around 30% of anthropogenic carbon emissions (Friedlingstein et al., 2023) and thereby play a crucial role in the global carbon cycle. However, due to the complexity of the terrestrial ecosystems, efforts to quantify their carbon uptake capacity still face significant challenges. The eddy covariance (EC) technique is currently the most widely used method to measure the carbon exchange between terrestrial ecosystems and the atmosphere at the ecosystem scale (Baldocchi et al., 2001), providing insights into terrestrial carbon uptake capacity. Numerous regional and global carbon flux networks, such as FLUXNET, AmeriFlux, ICOS, AsiaFlux, TERN-OzFlux and ChinaFLUX, have been established to coordinate EC flux measurements across diverse terrestrial ecosystems. Despite the presence of over a thousand EC sites worldwide, these sites are predominantly located in North America, Europe, and East Asia (Pastorello et al., 2020). Many regions, like Northwest China and Central Asia, remain underrepresented, which hinders accurate quantification of carbon sinks in these areas and global-scale synthesis and upscaling studies.

The Heihe River Basin (HRB) is the second largest inland river basin in China and serves as an ideal experimental region for studying the carbon cycle in Northwest China (Cheng et al., 2014). Due to the significant gradients in elevation (ranging from 1,500 m a.s.l. downstream to 5,000 m a.s.l. upstream) and precipitation (from 50 mm downstream to 600 mm upstream), the HRB encompasses diverse landscapes, including snow/glacier, permafrost, alpine grassland, subalpine forest, irrigated cropland, riparian ecosystems, wetlands, and gobi/desert from the upstream to the downstream (Li et al., 2013). A carbon flux observation network in the HRB was established through two comprehensive field experiments: the Watershed Allied Telemetry Experimental Research (WATER) conducted from 2007 to 2010 (Li et al., 2009) and the Heihe Watershed Allied Telemetry Experimental Research (HiWATER) conducted from 2012 to 2017 (Li et al., 2013).

There are a total of 34 sites in the Heihe carbon flux network (Liu et al., 2018), among which 10 are long-term observation sites, while the rest are temporary sites that have been dismantled. The network started observing carbon flux data since 2008, and the quality controlled 30 min data is released annually on the National Tibetan Plateau Data Center. However, the released data contains numerous gaps due to instrument malfunctions and routine maintenance. Additionally, the net ecosystem exchange (NEE) has not been partitioned into gross primary productivity (GPP) and ecosystem respiration (Reco), two widely used carbon flux components in carbon cycle studies. These issues hinder the effective use of the dataset. To provide uniform and continuous carbon flux and auxiliary data, it is necessary to compile and post-process all the flux data in the HRB. Therefore, the objectives of this work are: 1) to effectively fill the gaps in carbon flux data and auxiliary meteorological data of the Heihe carbon flux network, and produce a high-quality, uniform, and continuous carbon flux dataset in the HRB; 2) to partition the half-hourly NEE measurements into GPP and Reco; and 3) to explore the diurnal, seasonal and inter-annual variability of carbon flux across diverse ecosystems in the HRB based on the gap-filled, partitioned dataset.

70 **2 Carbon flux network in the HRB**

The Heihe carbon flux network encompasses the main ecosystem types in the HRB, including alpine grassland, subalpine forest, wetland, irrigated cropland, riparian woody land, and gobi/desert (Fig. 1). Detailed information on these sites is provided in Table 1 and Table 2. The development of the Heihe carbon flux network has experienced three stages. The first stage spans from 2007 to 2011, during which the WATER experiment was conducted. During this period, the network comprised three sites: Arou, Guantan, and Yingke (Table 1) (Li et al., 2009). These three sites were dismantled in 2012.

The second stage spans from 2012 to 2015 when the HiWATER experiment was conducted (Li et al., 2013; Liu et al., 2018). During this period, the Heihe carbon flux network underwent comprehensive updates. In 2012, five new flux sites were established in and around the artificial oasis in the middle reaches of the HRB: Daman super site, Zhangye wetland site, Huazhaizi site, Bajitan site, and Shenshawo site (Table 1). Additionally, from May to September 2012, a flux matrix consisting of 17 sites was set up in the middle-stream artificial oasis (Table 2). In the summer of 2013, three flux sites were established in the upstream areas of the HRB: Arou super site, Dadongshu site, and Dshalong site (Table 1). Simultaneously, five sites



were set up in the downstream areas of the HRB: Sidaoqiao super site, Hunhelin site, Huyang site, Nongtian site, and Luodi site (Table 1).

85 The third stage spans from 2016 to the present, and the network was optimized to enhance its representativeness at the basin scale and to make maintenance more manageable (Liu et al., 2018). In 2016, four sites (Bajitan site, Shenshawo site, Luodi site, and Nongtian site) were dismantled, and two new flux sites were established: Huangmo site and Jingyangling site. Currently, ten sites are operational as long-term observing sites, with four in the upper reaches (Arou super site, Dशलалong site, Dadongshu site, and Jingyangling site), three in the middle reaches (Daman super site, Zhangye wetland site, and Huazhaizi site), and three in the lower reaches (Sidaoqiao super site, Hunhelin site, and Huangmo site).

90 At each site of the carbon flux network in the HRB, flux and auxiliary meteorological factors, including CO₂ flux (F_c), latent heat flux (LE), sensible heat flux (H), downward solar radiation (R_g), air temperature (T_a), soil temperature (T_s), relative humidity (RH), soil water content (SWC), precipitation (P), and atmospheric pressure (PA), are recorded half-hourly or processed to half-hourly data. The 30 min data has been performed quality control (Xu et al., 2020; Liu et al., 2018; Che et al., 2019; Liu et al., 2023) and released at the TPDC (<https://data.tpdc.ac.cn/zh-hans/topic/heihe>). This released half-hourly carbon
95 flux and auxiliary data include a lot of gaps, and the NEE data are not partitioned into GPP and Reco.




Fig. 1 Distribution of eddy covariance (EC) observation sites in the Heihe River Basin (HRB). (a) is EC sites distribution map in the Heihe River Basin with landcover as the background. (b) and (c) show the distribution of the EC sites in the matrix experiment area in middle reaches and core experiment area in the lower reaches. The photos from 1 to 34 illustrate the underlying landscapes of the EC sites.

100



Table 1 Information of long-term and short-term observing sites in HRB.

ID	Name	Site_ID	Longitude (°E)	Latitude (°N)	Landcover	Dominant plant	Elevation (m)	Period	Stream
1	Jingyangling site	JYL	101.116	37.8384	Alpine grassland	Kobresia pygmaea	3750m	2018- Now	Upper
2	Arou super site	ARS	100.4643	38.0473	Alpine grassland	Kobresia pygmaea	3033m	2013- Now	Upper
3	Dadongshu site	DDS	100.2421	38.0142	Alpine grassland	Kobresia pygmaea	4148m	2015- Now	Upper
4	Dashalong site	DSL	98.9406	38.8399	Alpine marshland	Kobresia pygmaea	3739m	2013- Now	Upper
5	Daman super site	DMS	100.3722	38.8555	Cropland	Seed corn	1556m	2012- Now	Middle
6	Zhangye wetland site	ZYW	100.4464	38.9751	Wetland	Reed	1460m	2012- Now	Middle
7	Huazhaizi site	HZZ	100.3186	38.7652	Desert	Salsola passerina	1731m	2012- Now	Middle
8	Sidaoqiao super site	SDQ	101.1374	42.0012	Woodland	Tamarisk	873m	2013- Now	Lower
9	Hunhelin site	HHL	101.1335	41.9903	Woodland	Populus euphratica and Tamarix	874m	2013- Now	Lower
10	Huangmo site	HMo	100.9872	42.1135	Desert	Reaumuria	1054m	2015- Now	Lower
11	Arou site	ARo	100.4646	38.0443	Alpine grassland	Kobresia pygmaea	3033m	2008-2011	Upper
12	Guantan site	GTa	100.2500	38.5333	Subalpine forest	Picea crassifolia	2835m	2010-2011	Upper
13	Yingke site	YKe	100.4103	38.8571	Cropland	Seed corn	1519m	2007-2011	Middle
14	Bajitan site	BJT	100.3042	38.9150	Gobi	/	1562m	2012-2014	Middle
15	Shenshawo site	SSW	100.4933	38.7892	Desert	/	1594m	2012-2015	Middle
16	Luodi site	LDi	101.1326	41.9993	Bare land	/	878m	2013-2015	Lower
17	Nongtian site	NTi	101.1338	42.0048	Cropland	Cucumis melo	875m	2013-2015	Lower
18	Huyanglin site	HYL	101.1239	41.9932	Woodland	Populus euphratica forest	876m	2013-2015	Lower

Table 2 Information of sites in the eddy covariance matrix experiment in the middle reaches of the HRB in 2012.

ID	Name	Site_ID	Longitude (°E)	Latitude (°N)	Landcover	Dominant plant	Elevation(m)	Period
19	EC Matrix 1	M01	100.35813	38.89322	Cropland	Vegetable	1552.75m	2012/6/4-9/17
20	EC Matrix 2	M02	100.35406	38.88695	Cropland	Seed corn	1559.09m	2012/6/3-9/21
21	EC Matrix 3	M03	100.37634	38.89053	Cropland	Seed corn	1543.05m	2012/6/3-9/18
22	EC Matrix 4	M04	100.35753	38.87752	Built-up	/	1561.87m	2012/5/31-9/17
23	EC Matrix 5	M05	100.35068	38.87574	Cropland	Seed corn	1567.65m	2012/6/3-9/18
24	EC Matrix 6	M06	100.35970	38.87116	Cropland	Seed corn	1562.97m	2012/5/28-9/21
25	EC Matrix 7	M07	100.36521	38.87676	Cropland	Seed corn	1556.39m	2012/5/29-9/18
26	EC Matrix 8	M08	100.37649	38.87254	Cropland	Seed corn	1550.06m	2012/5/28-9/21
27	EC Matrix 9	M09	100.38546	38.87239	Cropland	Seed corn	1543.34m	2012/6/4-9/17
28	EC Matrix 10	M10	100.39572	38.87567	Cropland	Seed corn	1534.73m	2012/6/4-9/17
29	EC Matrix 11	M11	100.34197	38.86991	Cropland	Seed corn	1575.65m	2012/5/29-9/18
30	EC Matrix 12	M12	100.36631	38.86515	Cropland	Seed corn	1559.25m	2012/5/28-9/21
31	EC Matrix 13	M13	100.37852	38.8607	Cropland	Seed corn	1550.73m	2012/5/27-9/20
32	EC Matrix 14	M14	100.35310	38.85867	Cropland	Seed corn	1570.23m	2012/5/30-9/21



33	EC Matrix 16	M16	100.36411	38.84931	Cropland	Seed corn	1564.31m	2012/6/6-9/17
34	EC Matrix 17	M17	100.36972	38.84510	Cropland	Orchard	1559.63m	2012/5/31-9/17

3 Data post-processing

In this work, the data post-process included three steps: 1) to perform quality control on the auxiliary meteorological data and fill the gaps in the meteorological data by combining meteorological reanalysis data; 2) to conduct quality control of the NEE measurements and to fill the flux data gaps; 3) to partition the half-hourly NEE measurements to GPP and Reco. The flow diagram for the data post-processing is shown in Fig. 2.




Fig. 2 Diagram for the meteorological and carbon flux data post-processing in the HRB.

3.1 Meteorological data post-processing

The half-hourly meteorological data underwent quality control to remove outliers, and only high-quality records were retained (Xu et al., 2020; Liu et al., 2018). In this work, we selected eight meteorological factors (Rg, Ta, Ts, RH, VPD, SWC, P, PA) measured at the flux sites in the HRB for post-processing, as these factors are highly related to carbon fluxes and are also available in the ERA5-Land (ECMWF Reanalysis v5) dataset. The ERA5-Land dataset is a global reanalysis dataset with spatial resolution of 0.1 degree and temporal resolution of 1 hour (Muñoz-Sabater et al., 2021). The corresponding factors were extracted from the ERA5-Land dataset according to the geographical coordinates of each site. To match the temporal resolution of the in-situ observed data, the extracted hourly ERA5-Land data was linearly interpolated to half-hourly data for Rg, Ta, Ts, RH, VPD, SWC, and PA. For precipitation (P), linear interpolation could result in overestimation of the yearly precipitation amount, and therefore the hourly precipitation was equally divided to two half-hours. After temporal matching between in-situ observations and extracted ERA5-Land data, a random forest (RF) model was trained for each factor. To test the accuracy of the RF model in meteorological gap-filling, 5-days of continuous artificial gaps were created in the meteorological factors, and these were then used to assess the performance of the gap-filling method. The RF model was able to accurately predict the missing meteorological observations for all variables except P using ERA5-land variables as input (Fig. 3).




Fig. 3 The performance of RF models in the gap-filling of the meteorological data. Rg: downward shortwave radiation; RH: relative humidity; SWC: soil water content; P: precipitation; PA: atmospheric pressure; VPD: vapor pressure deficit; Ta: air temperature; Ts: soil temperature. The suffix "_Obs" indicates the observed values. The suffix "_RF" indicates the random forest predicted values.

130



3.2 Carbon flux data post-processing

The original 10 Hz EC data were processed into 30-minute flux data by Xu and Liu (Xu et al., 2020; Liu et al., 2018).
135 Here, we further processed the 30-minute data to obtain continuous GPP, NEP, and Reco. First, outliers in the 30-minute NEE
data were excluded based on a three-times standard deviation criterion (Rousseeuw and Croux, 1993), which is a widely used
method in meteorological aberrant values detection.

Second, to further exclude poor-quality NEE data, u^* filtering was applied using the REddyProc package (Wutzler et al.,
2018), a post-processing tool for half-hourly EC measurements. During the night, stable stratification often occurs, leading to
140 underestimation of nighttime NEE. This issue was identified by examining the relationship between NEE and u^* . Nighttime
NEE values with u^* lower than a threshold u^* were filtered as invalid. A revised breakpoint detection method (Barr et al.,
2013) in the REddyProc package was used to determine the threshold u^* . After u^* filtering, the gaps in the half-hourly data
increased.

Third, it was necessary to fill these gaps to obtain continuous NEE data. In this study, both marginal distribution sampling
145 (MDS) (Reichstein et al., 2005) and RF were implemented to fill the gaps by combining gap-filled R_g , T_a , and VPD data with
valid NEE data. The MDS method fills half-hourly NEE gaps using different schemes depending on the availability of
meteorological data and is included in REddyProc package. For the RF method, a RF model is built using high-quality observed
NEE and auxiliary meteorological factors (R_g , T_a , and VPD). This model is then used to fill the NEE gaps by inputting the
gap-filled R_g , T_a , and VPD data. Both MDS and RF are effective in filling the gaps in NEE, with R^2 values of 0.77 for MDS
150 and 0.84 for RF between the filled and observed values (Fig. 4). While RF can fill all the gaps in NEE, MDS still leaves some
long gaps unfilled (Fig. 5).

Fourth, the gap-filled NEE data were partitioned into GPP and Reco, two critical variables in carbon cycle studies. The
NEE partitioning was also performed using the REddyProc package. During nighttime ($R_g < 10 \text{ W/m}^2$), NEE equals Reco
because there is no photosynthesis. The Lloyd-Taylor respiration function (Lloyd and Taylor, 1994) was fitted using nighttime
155 NEE and T_a . This fitted Lloyd-Taylor respiration function was then applied to estimate daytime Reco, with GPP calculated as
the difference between Reco and NEE during the daytime (Wang et al., 2012).

To improve user convenience, the dataset variables were aggregated at multiple time intervals, including daily ($_DD$),
weekly ($_WW$), monthly ($_MM$), and yearly ($_YY$). Variables such as GPP, Reco, NEE, and Precipitation were aggregated
over longer intervals using the sum. In contrast, variables like R_g , T_a , T_s , SWC, RH, VPD, PA, LE, and H were aggregated
160 using the average.

Considering the close performance of the MDS and RF methods, and the fact that the RF method can fill all gaps in the
data, the subsequent analysis of carbon flux in the HRB is based on the RF results.




Fig. 4 The performance of RF models in carbon flux gap-filling. The suffix "_Obs" indicates the observed values.

165 The suffix "_RF" indicates the gap-filled values by random forest method. The suffix "_MDS" indicates the gap-filled values by MDS method.




Fig. 5 Data gaps in NEE before and after gap-filling. The suffix "_Obs" indicates the observed values. The suffix "_RF"



indicates the gap-filled with random forest method. The suffix "_MDS" indicates the gap-filled with MDS method.

170 4 Dataset description

The post-processed dataset includes 34 sites with time spans ranging from a few months to ten years. For each site, the data comprises both original and gap-filled auxiliary meteorological factors (downward solar radiation: Rg, air temperature: Ta, soil temperature: Ts, relative humidity: RH, soil water content: SWC, precipitation: P, and atmospheric pressure: PA) and flux data (net ecosystem exchange: NEE, gross primary productivity: GPP, ecosystem respiration: Reco, latent heat flux: LE, and sensible heat flux: H). The data is provided at multiple temporal scales: half-hourly (_HH), daily (_DD), weekly (_WW), monthly (_MM), and yearly (_YY). The folder for each specific site is named according to the following convention: Site_ID + start year + end year + temporal scale suffix. The data is saved in the CSV format. Fields with the suffix "_ERA" indicate data extracted from ERA5-Land. Fields with the suffix "_F" represent gap-filled data. The explanation of the fields in the post-processed data is shown in Table 3.

180 **Table 3 Data fields in half-hourly dataset.**

Variables	Description	Unit	
TIMESTAMP_START	The initial time of observation	-	
Year	Year	-	
DoY	The day of the year	-	
Hour	Hour of the day	-	
NEE	Net ecosystem exchange	g C/m ² /30min	
LE	Latent heat flux	W m ⁻²	In-situ observed data
H	Sensible heat flux	W m ⁻²	
Rg	Downward shortwave radiation	W m ⁻²	
Ta	Air temperature	degC	
Ts_1	Surface soil temperature	degC	
RH	Relative humidity	%	
VPD	Saturated vapor pressure difference	hPa	
SWC_1	Surface soil water content	%	
P	Precipitation	mm	
PA	Atmospheric pressure	hPa	
uStar	Friction wind speed	ms ⁻¹	
LE_ERA	Latent heat flux	W m ⁻²	Extracted ERA data
H_ERA	Sensible heat flux	W m ⁻²	
Rg_ERA	Downward shortwave radiation	W m ⁻²	
Ta_ERA	Air temperature	degC	
Ts_1_ERA	Surface soil temperature	degC	
RH_ERA	Relative humidity	%	
VPD_ERA	Saturated vapor pressure difference	hPa	
SWC_1_ERA	Surface soil water content	%	
P_ERA	Precipitation	mm	
PA_ERA	Atmospheric pressure	hPa	
NEE_F_MDS	Gap-filled NEE with MDS method	g C/m ² /30min	Gap-filled data
NEE_F_RF	Gap-filled NEE with RF method	g C/m ² /30min	
NEE_F_fqc	Quality flag for Gap-filled NEE	-	
H_F_MDS	Gap-filled sensible heat flux with MDS method	W m ⁻²	
H_F_RF	Gap-filled sensible heat flux with RF method	W m ⁻²	
H_F_fqc	Quality flag for H	-	
LE_F_MDS	Gap-filled latent heat flux with MDS method	W m ⁻²	
LE_F_RF	Gap-filled latent heat flux with RF method	W m ⁻²	
LE_F_fqc	Quality flag for LE	-	
Rg_F_RF	Downward shortwave radiation	W m ⁻²	
Rg_F_fqc	Air temperature	-	



Ta_F_RF	Air temperature	degC	
Ta_F_fqc	Quality flag for Ta	-	
Ts_1_F_RF	Surface soil temperature	degC	
Ts_1_F_fqc	Quality flag for Ts	-	
RH_F_RF	Relative humidity	%	
RH_F_fqc	Quality flag for RH	-	
VPD_F_RF	Saturated vapor pressure difference	hPa	
VPD_F_fqc	Quality flag for VPD	-	
SWC_1_F_RF	Surface soil water content	%	
SWC_1_F_fqc	Quality flag for SWC	-	
P_F_RF	Precipitation	mm	
P_F_fqc	Quality flag for P	-	
PA_F_RF	Atmospheric pressure	hPa	
PA_F_fqc	Quality flag for PA	-	
GPP_F_RF	Gross primary production partitioned from NEE_F_RF	g C/m ² /30min	
Reco_F_RF	Ecosystem respiration partitioned from NEE_F_RF	g C/m ² /30min	Partitioned
GPP_F_MDS	Gross primary production partitioned from NEE_F_MDS	g C/m ² /30min	Carbon flux
Reco_F_MDS	Ecosystem respiration partitioned from NEE_F_MDS	g C/m ² /30min	

*fqc in HH data: 1 = measured; 0 = gap-filled; fqc in DD, WW, MM and YY data: indicating percentage of missed data (0-1).

5 Results

5.1 Diurnal variations of carbon fluxes for various ecosystem types in the HRB

185 To explore the temporal dynamics of the carbon fluxes for diverse ecosystem types in the HRB, the carbon fluxes were averaged for different ecosystems, including subalpine forest, alpine grassland, cropland, wetland, riparian woodland and gobi/desert. The averaged diurnal cycle curves and statistic metrics of these ecosystems in growing season (May to September) are shown in Fig. 6 and Table 4. Note that negative NEE values indicate net carbon uptake, while positive NEE indicate net carbon release. Over the course of the diurnal cycle, NEE, GPP and Reco varied greatly for subalpine forest, alpine grassland, cropland and wetland but slightly for gobi/desert. The half-hourly NEE reached minimum (i.e., the largest net carbon uptake) 190 at 11:30pm for subalpine forest, alpine grassland and riparian woodland, 12:30pm for cropland, 13:00pm for wetland and 11:30pm for gobi/desert. The GPP varied greatly in subalpine forest, alpine grassland, cropland, wetland and riparian woodland, and kept constant close to zero in gobi/desert. The half-hourly GPP reached maximum at 11:30pm for subalpine forest and alpine grassland, at 12:30 for riparian woodland, cropland and gobi/desert, at 13:00pm for wetland. The half-hourly Reco 195 reached maximum at 16:30pm for subalpine forest and gobi/desert, at 17:00 for riparian woodland, at 15:30pm for alpine grassland, at 16:00pm for cropland and wetland (Table 4 and Fig. 6).




Fig. 6 Diurnal variation in the growing season of carbon fluxes of different ecosystems in the HRB.

Table 4 Statistical metrics of carbon flux diurnal curves for ecosystems in the HRB. (NEE_{Min} is minimum NEE in diurnal course and NEE_{Min} Time is corresponding time. GPP_{Max} is maximum GPP in diurnal course and GPP_{Max} Time is the corresponding time. $Reco_{Max}$ is maximum Reco in diurnal course and $Reco_{Max}$ Time is the corresponding time. Unit for NEE_{Min} , GPP_{Max} and $Reco_{Max}$ is $gC\ m^{-2}\ 30min^{-1}$)

	Subalpine forest	Riparian woodland	Alpine grassland	Cropland	Wetland	Gobi/Desert
NEE_{Min}	-0.29	-0.081	-0.17	-0.40	-0.30	-0.017
NEE_{Min} Time	11:30	11:30	11:30	12:30	13:00	11:30
GPP_{Max}	0.34	0.113	0.22	0.49	0.35	0.0187
GPP_{Max} Time	11:30	12:30	11:30	12:30	13:00	12:30
$Reco_{Max}$	0.06	0.034	0.045	0.089	0.055	0.002
$Reco_{Max}$ Time	16:30	17:00	15:30	16:00	16:00	16:30

5.2 Seasonal variation of carbon fluxes for ecosystems in the HRB

The eighteen sites (Table 1) with more than one year of data were selected to explore the seasonal dynamics of carbon fluxes for different ecosystems in the HRB. These sites were grouped into six ecosystem types. The seasonal dynamics of carbon fluxes for these ecosystems are shown in Fig. 7.

Seasonal NEE varied significantly throughout the year for subalpine forest, alpine grassland, wetland, cropland, and riparian woodland but remained close to $0\ gC\ m^{-2}\ day^{-1}$ year-round for gobi/desert. During the non-growing season, NEE was close to zero for all ecosystems except forest. In the transition period from non-growing to growing season, NEE slightly increased and became positive. During the growing season, NEE was notably less than zero for all ecosystems except gobi/desert, indicating that these ecosystems except gobi/desert exhibited net carbon uptake. The minimum NEE for all ecosystems occurred in July, with values of $-5.62\ gC\ m^{-2}\ day^{-1}$ for subalpine forest, $-1.48\ gC\ m^{-2}\ day^{-1}$ for riparian woodland, $-4.08\ gC\ m^{-2}\ day^{-1}$ for alpine grassland, $-12.40\ gC\ m^{-2}\ day^{-1}$ for cropland, $-8.24\ gC\ m^{-2}\ day^{-1}$ for wetland, and $-0.49\ gC\ m^{-2}\ day^{-1}$



¹ for gobi/desert (Table 5).

215 Seasonal GPP also varied significantly throughout the year. During the non-growing season, GPP was very close to zero for all ecosystems except for subalpine forest. During the growing season, GPP was obviously higher than zero for all ecosystems except gobi/desert. Seasonal GPP reached its maximum value in July, with values of 8.11 gC m⁻² day⁻¹ for subalpine forest, 3.08 gC m⁻² day⁻¹ for riparian woodland, 6.62 gC m⁻² day⁻¹ for alpine grassland, 16.84 gC m⁻² day⁻¹ for cropland, 10.98 gC m⁻² day⁻¹ for wetland, and 0.75 gC m⁻² day⁻¹ for gobi/desert (Table 5).

220 Seasonal Reco followed a temporal pattern similar with seasonal GPP. Reco also reached its maximum in July, with values of 3.93 gC m⁻² day⁻¹ for subalpine forest, 1.98 gC m⁻² day⁻¹ for riparian woodland, 2.96 gC m⁻² day⁻¹ for alpine grassland, 5.64 gC m⁻² day⁻¹ for cropland, 3.41 gC m⁻² day⁻¹ for wetland, and 0.17 gC m⁻² day⁻¹ for gobi/desert (Table 5).

Table 5 Statistical metrics of carbon flux seasonal curves for ecosystems in the HRB. (NEE_{Mean} and NEE_{Min} are daily average and minimum NEE in over the year. GPP_{Mean} and GPP_{Max} are daily average and minimum GPP in over the year.

225 Reco_{Mean} and Reco_{Max} are daily average and minimum GPP in over the year. Unit for NEE_{Mean}, NEE_{Min}, GPP_{Mean}, GPP_{Max}, Reco_{Mean} and Reco_{Max} is gC m⁻² day⁻¹.)

	Subalpine forest	Riparian woodland	Alpine grassland	Cropland	Wetland	Gobi/Desert
NEE _{Mean}	-2.10	-0.34	-0.84	-1.75	-1.83	-0.25
NEE _{Min}	-5.62	-1.48	-4.08	-12.40	-8.24	-0.49
GPP _{Mean}	3.52	1.18	1.67	3.50	3.08	0.30
GPP _{Max}	8.11	3.08	6.62	16.84	10.98	0.63
Reco _{Mean}	1.42	0.84	0.83	1.75	1.25	0.05
Reco _{Max}	3.93	1.98	2.96	5.64	3.41	0.17




Fig. 7 Seasonal dynamics of carbon fluxes for different ecosystem types in the HRB.



230 5.3 Inter-annual variations of carbon fluxes for various ecosystem types in the HRB

To explore the inter-annual variations of carbon fluxes in different ecosystem types of the HRB, nine sites with more than seven years of data were selected. These sites were grouped into five ecosystem types. The yearly dynamics of GPP, Reco, and NEE is shown in Fig. 8, and the statistical metrics of GPP, Reco, and NEE are provided in Table 6.

235 The multi-year average NEE was $-123.43 \text{ gC m}^{-2} \text{ year}^{-1}$ in riparian woodland, $-307.84 \text{ gC m}^{-2} \text{ year}^{-1}$ in alpine grassland, $-638.77 \text{ gC m}^{-2} \text{ year}^{-1}$ in cropland, $-679.62 \text{ gC m}^{-2} \text{ year}^{-1}$ in wetland and $-92.04 \text{ gC m}^{-2} \text{ year}^{-1}$ in gobi/desert. Yearly NEE was the highest in cropland and the lowest in gobi/desert. Annual NEE of wetlands significantly increased during the last decade, while other ecosystems exhibited relatively stable NEE with slight inter-annual variations (Fig. 8a). The multi-year average GPP was $431.47 \text{ gC m}^{-2} \text{ year}^{-1}$ for riparian woodland, $609.22 \text{ gC m}^{-2} \text{ year}^{-1}$ for alpine grassland, $1269.19 \text{ gC m}^{-2} \text{ year}^{-1}$ for cropland, $1127.88 \text{ gC m}^{-2} \text{ year}^{-1}$ for wetlands, and $108.22 \text{ gC m}^{-2} \text{ year}^{-1}$ for gobi/desert. Over the last decade, the annual GPP of wetland and riparian woodland slightly increased, while the GPP of other ecosystems remained relatively stable (Fig. 8b). The multi-year average Reco was $308.03 \text{ gC m}^{-2} \text{ year}^{-1}$ for riparian woodland, $301.38 \text{ gC m}^{-2} \text{ year}^{-1}$ for alpine grassland, $630.42 \text{ gC m}^{-2} \text{ year}^{-1}$ for cropland, $448.26 \text{ gC m}^{-2} \text{ year}^{-1}$ for wetlands, and $16.18 \text{ gC m}^{-2} \text{ year}^{-1}$ for gobi/desert. The Reco in the HRB slightly increased for cropland and wetlands but remained relatively stable for other ecosystem types over the last decade (Fig. 8c).

245




Fig. 8 Yearly NEE, Reco and GPP for the main ecosystems in the HRB.

5.4 Spatial variations of carbon fluxes in the HRB



To examine the spatial patterns of carbon fluxes, the annual NEE, GPP and Reco of the eighteen sites with at least 1 year
 250 of data were compared. Annual NEE, GPP and Reco of the eighteen sites in the HRB are shown in Fig. 9. In the upper reaches
 of the river basin, annual NEE and GPP was the highest at the GTa site and the lowest at the DDS site. In the middle reaches,
 NEE and GPP was the highest at the ZYW, DMS and YKe sites, which are in the artificial oasis. In the lower reaches, NEE
 and GPP were higher at HHL, HYL, SDQ and NTi than at HMo and LDi. In the upper reaches, the NEE, GPP and Reco
 generally increased with elevation. In the middle and lower reaches, NEE, GPP and Reco of the sites inside the artificial/natural
 255 oasis were obviously higher than those of the sites outside the artificial/natural oasis.




Fig. 9 Yearly carbon fluxes of the 18 sites with more than 1 year of data for the upper, middle, and lower reaches of the HRB. The number under each site ID indicates the elevation of the site.

To explore the carbon flux change along the environmental gradients, we sorted the carbon flux sites based on air
 260 temperature, precipitation, soil water content, and downward shortwave radiation. We then explored the carbon flux variation
 in relation to these four factors. The yearly average air temperature increases from the upper reaches to the middle and lower
 reaches of the HRB. However, the GPP, Reco, and NEE did not show a similar gradient pattern with air temperature. The GPP,
 Reco, and NEE are significantly higher at DMS, YKe, GTa, and ZYW than at other sites, and the temperature at these sites is
 at an intermediate level among all the sites (Fig. 10a). In the upper reaches of the HRB, the NEE, GPP and Reco of the sites
 265 decreased as annual average temperature decreased, while in the middle and lower reaches of the HRB, NEE, GPP and Reco
 did not change with the temperature gradient. NEE, GPP and Reco generally follow the same spatial pattern, with higher
 carbon fluxes at sites with higher soil water content. The cropland and wetland with irrigation in the middle reaches have the
 highest GPP, Reco and NEE (Fig. 10b). Precipitation decreases from about 500 mm in the upper reaches to about 50 mm in
 the lower reaches of the HRB. The GPP, Reco, and NEE of these sites did not strictly increase or decrease with the spatial
 270 precipitation gradient (Fig. 10c). GPP, Reco, and NEE did not change with the gradient of Rg among the sites in the HRB (Fig.
 10d).




Fig. 10 Carbon flux spatial variation with the gradients of meteorological factors. Letters in the bracket stand for subregions in the HRB, U for upper reaches, M for middle reaches and L for lower reaches of the HRB.

275 6 Discussion

6.1 Carbon flux pattern and its drivers in the HRB

EC-based carbon flux data is perhaps the most effective data source to quantify the carbon sequestration capacity of ecosystems at the ecosystem scale. As the HRB is a typical inland river basin, the temporal and spatial patterns of carbon fluxes in this region provide insight into the carbon dynamics of inland river basins in the Northwest China and Central Asia more broadly.

The diurnal pattern of carbon flux shows that GPP and NEE peak around midday when downward shortwave radiation is at its highest, while Reco reaches its peak later in the afternoon when temperatures are at their maximum. This indicates that during the growing season, the diurnal NEE curve is primarily driven by GPP variations. The GPP diurnal pattern is largely influenced by downward shortwave radiation, whereas the Reco pattern is controlled by temperature (Kato et al., 2004). The half-hourly GPP and NEE of subalpine forest, riparian woodland, alpine grassland, and gobi/desert peak slightly earlier than those of cropland and wetlands. This difference is likely due to the fact that cropland and wetlands are not limited by water or



heat, whereas the other ecosystems experience stress from either heat or water (Lin et al., 2019).

290 The seasonal patterns of NEE, GPP, and Reco vary significantly among different ecosystems. In gobi/desert, NEE, GPP, and Reco remain close to zero throughout the year due to sparse vegetation coverage and low soil organic matter content. For alpine grassland, NEE, GPP, and Reco begin to increase later in spring and decrease earlier in autumn compared to subalpine forest, wetland, and riparian woodland. This can be attributed to the lower temperatures and higher elevations of alpine grassland (Wang et al., 2022). In cropland, NEE, GPP, and Reco also show a late start in spring and an early decline in autumn, mainly driven by agricultural management practices (Guo et al., 2021).

295 The upper reaches of the HRB are humid and cold, and the alpine grassland are weak carbon sinks with annual NEE range from $-108 \text{ gC m}^{-2} \text{ year}^{-1}$ to $-341.56 \text{ gC m}^{-2} \text{ year}^{-1}$. This is consistent with the previous studies on alpine grassland at the Haibei site (Zhao et al., 2005) and the Dangxiong site (Shi et al., 2006). The subalpine forest (*Picea crassifolia*) in the upper reaches is strong carbon sink, with NEE of $-767.01 \text{ gC m}^{-2} \text{ year}^{-1}$, which has been rarely reported. The carbon flux in the upstream region of the HRB is mainly stressed by low temperature (Sun et al., 2019).

300 The middle reaches of the HRB are dry and hot, with significant differences in carbon fluxes between sites inside and outside the artificial oasis. Sites within the artificial oasis have strong carbon uptake capacity, with NEE exceeding $-600 \text{ gC m}^{-2} \text{ year}^{-1}$ due to irrigation. In contrast, sites outside the artificial oasis have very weak carbon uptake capacity, with NEE less than $-100 \text{ gC m}^{-2} \text{ year}^{-1}$. Inside the artificial oasis, high temperatures and high soil water content promote vegetation growth, while outside the artificial oasis, high temperatures and low soil water content inhibit vegetation growth. Due to intensive irrigation, the carbon fluxes of sites inside the artificial oasis are decoupled from precipitation in this region (Wang et al., 305 2019).

The lower reaches are even drier and hotter than the middle reaches in the HRB. The NEE of the sites in this region ranges from $-49.72 \text{ gC m}^{-2} \text{ year}^{-1}$ to $-123.85 \text{ gC m}^{-2} \text{ year}^{-1}$. The natural oasis in the lower reaches consist of riparian ecosystems distributed along the main river channels. Vegetation in the natural oasis survives by relying on lateral water supply from the river channel and shallow groundwater. Vegetation in the natural oasis faces slightly water stress compared to vegetation in the artificial oasis in the middle reaches. The vegetation outside the natural oasis in the lower reaches faces more severe water stress than vegetation outside the artificial oasis in the middle reaches due to lower precipitation and soil water availability in the downstream region.

6.2 Possible sources of uncertainty in the carbon data in the HRB

315 Data post-processing can introduce uncertainties into carbon flux measurements. Due to instrument malfunctions and maintenance, data gaps are inevitable, yet continuous carbon flux data is essential for assessing an ecosystem's carbon uptake capacity. In post-processing, three key steps— u^* correction, gap-filling, and carbon flux partitioning—can result in uncertainties. The u^* correction can filter a large proportion of nighttime carbon flux data, with different methods yielding different u^* thresholds and varying proportions of filtered data. This correction can impact data availability for building lookup tables or training models in the gap-filling process.

320 The gap-filling process uses mathematical methods aided by meteorological data to fill in missing data, which can introduce significant uncertainties. The MDS and random forest methods are the two primary techniques currently used for data gap-filling (Zhu et al., 2022), and both are evaluated in this work for the HRB. The performance of MDS and random forest methods is very close, and both can effectively fill gaps in half-hourly NEE data. While previous studies have reported that MDS may systematically overestimate carbon emissions and underestimate CO_2 sequestration (Vekuri et al., 2023), we 325 did not observe this phenomenon in the HRB. However, MDS cannot effectively fill gaps longer than two weeks, whereas the RF method can fill all gaps if the corresponding auxiliary meteorological data are available.

The NEE partitioning method can also introduce uncertainties into GPP and Reco data (Tramontana et al., 2020). Although the night-time-based method is recommended as a standard by FLUXNET, it has some limitations. It only considers



330 temperature in the respiration estimation, neglecting other environmental factors that could influence respiration. Additionally,
it does not account for variations in respiration between nighttime and daytime under different light conditions (Oikawa et al.,
2017; Raj et al., 2016; Chen et al., 2024). Since direct measurement of GPP and Reco are not available, assessing uncertainties
in the NEE partitioning step remains challenging.

335 In addition to uncertainties introduced by data processing, harsh weather, complex terrain, and instrument maintenance
can contribute to uncertainties in carbon flux observations in the HRB. In the upstream regions, extremely cold temperatures
during the non-growing season can occasionally lead to frost formation on the gas analyzer, affecting the CO₂ concentration
signals. In the midstream and downstream areas, sandstorms can disrupt the optical path of the gas analyzer. Furthermore,
finding large, flat, ideal locations for EC instrumentation can be challenging for certain ecosystems. For example, subalpine
forest in the upstream area are primarily located on shaded hill slopes, suggesting that carbon fluxes at the GTa site may require
additional processing to account for terrain effects. Additionally, instrument degradation and updates can introduce
340 uncertainties or inconsistencies into the original observation data.

7 Data use guidelines

Data are fully public but should be appropriately referenced by citing this paper and the database (see Sect. 8). We
suggest that researchers planning to use this dataset as a core dataset for their analysis contact and collaborate with the first or
corresponding authors of this paper.

345 8 Data availability

The post-processed carbon flux and auxiliary data in the HRB is available at: <https://doi.org/10.11888/Terre.tpd.301321>
(Wang et al., 2024).

9 Conclusions

350 Over the past decade, a comprehensive carbon flux network has been established in the Heihe River Basin (HRB) in
Northwest China. In this study, carbon flux and auxiliary meteorological data from the network were post-processed to create
an analysis-ready dataset. This dataset encompasses 34 sites across six dominant ecosystems in the HRB: alpine grassland,
subalpine forest, cropland, wetland, riparian woodland, and gobi/desert. Eighteen of these sites have continuous multi-year
observations, while 16 sites were observed only during the 2012 growing season, totaling 1,513 site-months. Based on this
dataset, the following temporal and spatial characteristics of carbon exchange in the HRB were identified: 1) In the diurnal
355 variation curve, GPP, NEE, and Reco peak later for ecosystems in the artificial oasis (cropland and wetland) compared to those
outside the artificial oasis (grassland, forest, woodland, and gobi/desert). 2) Seasonal NEE, GPP, and Reco peak in early July
for grassland, forest, woodland, and cropland, while remain close to zero throughout the year for gobi/desert. 3) In the last
decade, NEE of wetlands significantly increased, while NEE for other ecosystems slightly fluctuated inter-annually. 4) NEE,
GPP, and Reco are significantly higher for sites inside the artificial/natural oasis compared to those outside. This post-
360 processed carbon flux dataset has many applications, e.g., exploring carbon exchange characteristic of alpine and arid
ecosystem, ecosystem responses to climate extremes, cross-site synthesis at regional to global scales, regional and global
upscaling studies, interpreting and calibrating remote sensing products, evaluating and calibrating carbon cycle models.

Author contributions

365 X.F. Wang, T. Che, J.F. Xiao, X. Li and S.M. Liu designed the research. X.F. Wang and T.H. Wang processed and
analyzed the data. J.L. Tan, Y. Zhang, Z.G. Ren, Z.W. Xu, L.Y. Geng and H.B. Wang collected the data, maintained carbon
flux network in HRB and processed the original data. X.F. Wang, T. Che, T.H. Wang and J.F. Xiao wrote the paper.



Competing interests

The contact author has declared that none of the authors has any competing interests.

Acknowledgements

370 We thank all the scientists, engineers, and students who contributed to the establishment and maintenance of the carbon flux network in the Heihe River Basin.

Financial support

This work has been supported by the Gansu Provincial Science and Technology Program (22ZD6FA005), the Foundation for Distinguished Young Scholars of Gansu Province (Grant No. 22JR5RA046), the Funds of National Natural Science Foundation of China (Grant No. 42371386 and U22A202690), and the Youth Innovation Promotion Association CAS 375 to X.W. (No. 2020422). J.F. Xiao was supported by University of New Hampshire via bridge support.

References

- Baldocchi, D., Falge, E., Gu, L., Olson, R., Hollinger, D., Running, S., Anthoni, P., Bernhofer, C., Davis, K., Evans, R., Fuentes, J., Goldstein, A., Katul, G., Law, B., Lee, X., Malhi, Y., Meyers, T., Munger, W., Oechel, W., Paw U, K. T., Pilegaard, K., Schmid, H. P., Valentini, R., Verma, S., Vesala, T., Wilson, K., and Wofsy, S.: FLUXNET: A New Tool to Study the Temporal and Spatial Variability of Ecosystem-Scale Carbon Dioxide, Water Vapor, and Energy Flux Densities, *Bulletin of the American Meteorological Society*, 82, 2415-2434, [https://doi.org/10.1175/1520-0477\(2001\)082<2415:FANTTS>2.3.CO;2](https://doi.org/10.1175/1520-0477(2001)082<2415:FANTTS>2.3.CO;2), 2001.
- 380 Barr, A., Richardson, A., Hollinger, D., Papale, D., Arain, M., Black, T., Bohrer, G., Dragoni, D., Fischer, M., and Gu, L.: Use of change-point detection for friction–velocity threshold evaluation in eddy-covariance studies, *Agricultural and forest meteorology*, 171, 31-45, 2013.
- Che, T., Li, X., Liu, S., Li, H., Xu, Z., Tan, J., Zhang, Y., Ren, Z., Xiao, L., and Deng, J.: Integrated hydrometeorological, snow and frozen-ground observations in the alpine region of the Heihe River Basin, China, *Earth System Science Data*, 11, 1483-1499, 2019.
- 390 Chen, H., Li, H., Wei, Y., McBean, E., Liang, H., Wang, W., and Huang, J. J.: Partitioning eddy covariance CO₂ fluxes into ecosystem respiration and gross primary productivity through a new hybrid four sub-deep neural network, *Agriculture, Ecosystems & Environment*, 361, 108810, <https://doi.org/10.1016/j.agee.2023.108810>, 2024.
- Cheng, G., Li, X., Zhao, W., Xu, Z., Feng, Q., Xiao, S., and Xiao, H.: Integrated study of the water–ecosystem–economy in the Heihe River Basin, *National science review*, 1, 413-428, 2014.
- 395 Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Bakker, D. C. E., Hauck, J., Landschützer, P., Le Quééré, C., Lujikx, I. T., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Anthoni, P., Barbero, L., Bates, N. R., Becker, M., Bellouin, N., Decharme, B., Bopp, L., Brasika, I. B. M., Cadule, P., Chamberlain, M. A., Chandra, N., Chau, T. T. T., Chevallier, F., Chini, L. P., Cronin, M., Dou, X., Enyo, K., Evans, W., Falk, S., Feely, R. A., Feng, L., Ford, D. J., Gasser, T., Ghattas, J., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, Ö., Harris, I., Hefner, M., Heinke, J., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Jacobson, A. R., Jain, A., Jarníková, T., Jersild, A., Jiang, F., Jin, Z., Joos, F., Kato, E., Keeling, R. F., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Körtzinger, A., Lan, X., Lefèvre, N., Li, H., Liu, J., Liu, Z., Ma, L., Marland, G., Mayot, N., McGuire, P. C., McKinley, G. A., Meyer, G., Morgan, E. J., Munro, D. R., Nakaoka, S. I., Niwa, Y., O'Brien, K. M., Olsen, A., Omar, A. M., Ono, T., Paulsen, M., Pierrot, D., Pockock, K., Poulter, B., Powis, C. M., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Rosan, T. M., Schwinger, J., Séférian, R., Smallman, T. L., Smith, S. M., Sospedra-Alfonso, R., Sun, Q., Sutton, A. J., Sweeney, C., Takao, S., Tans, P. P., Tian, H., Tilbrook, B., Tsujino, H., Tubiello, F., van der Werf, G. R., van Ooijen, E., Wanninkhof, R., Watanabe, M., Wimart-Rousseau, C., Yang, D., Yang, X., Yuan, W., Yue, X., Zaehle, S., Zeng, J., and Zheng, B.: Global Carbon Budget 2023, *Earth Syst. Sci. Data*, 15, 5301-5369, 10.5194/essd-15-5301-2023, 2023.
- 410 Guo, H., Li, S., Wong, F.-L., Qin, S., Wang, Y., Yang, D., and Lam, H.-M.: Drivers of carbon flux in drip irrigation maize fields in northwest China, *Carbon Balance and Management*, 16, 12, 10.1186/s13021-021-00176-5, 2021.
- Kato, T., Tang, Y., Gu, S., Hirota, M., Cui, X., Du, M., Li, Y., Zhao, X., and Oikawa, T.: Seasonal patterns of gross primary production and ecosystem respiration in an alpine meadow ecosystem on the Qinghai-Tibetan Plateau, *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/10.1029/2003JD003951>, 2004.
- 415 Li, X., Li, X., Li, Z., Ma, M., Wang, J., Xiao, Q., Liu, Q., Che, T., Chen, E., and Yan, G.: Watershed allied telemetry experimental research, *Journal of Geophysical Research: Atmospheres*, 114, 2009.
- Li, X., Cheng, G. D., Liu, S. M., Xiao, Q., Ma, M. G., Jin, R., Che, T., Liu, Q. H., Wang, W. Z., Qi, Y., Wen, J. G., Li, H.



- 420 Y., Zhu, G. F., Guo, J. W., Ran, Y. H., Wang, S. G., Zhu, Z. L., Zhou, J., Hu, X. L., and Xu, Z. W.: Heihe Watershed Allied
Telemetry Experimental Research (HiWATER): Scientific Objectives and Experimental Design, *Bulletin of the American
Meteorological Society*, 94, 1145-1160, 10.1175/Bams-D-12-00154.1, 2013.
- Lin, C., Gentine, P., Frankenberg, C., Zhou, S., Kennedy, D., and Li, X.: Evaluation and mechanism exploration of the
diurnal hysteresis of ecosystem fluxes, *Agricultural and Forest Meteorology*, 278, 107642,
<https://doi.org/10.1016/j.agrformet.2019.107642>, 2019.
- 425 Liu, S., Li, X., Xu, Z., Che, T., Xiao, Q., Ma, M., Liu, Q., Jin, R., Guo, J., and Wang, L.: The Heihe Integrated Observatory
Network: A basin-scale land surface processes observatory in China, *Vadose Zone Journal*, 17, 1-21, 2018.
- Liu, S., Xu, Z., Che, T., Li, X., Xu, T., Ren, Z., Zhang, Y., Tan, J., Song, L., Zhou, J., Zhu, Z., Yang, X., Liu, R., and Ma,
Y.: A dataset of energy, water vapor, and carbon exchange observations in oasis-desert areas from 2012 to 2021 in a typical
endorheic basin, *Earth Syst. Sci. Data*, 15, 4959-4981, 10.5194/essd-15-4959-2023, 2023.
- 430 Lloyd, J. and Taylor, J.: On the temperature dependence of soil respiration, *Functional ecology*, 315-323, 1994.
- Muñoz-Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo, G., Boussetta, S., Choulga, M.,
Harrigan, S., Hersbach, H., Martens, B., Miralles, D. G., Piles, M., Rodríguez-Fernández, N. J., Zsoter, E., Buontempo, C.,
and Thépaut, J. N.: ERA5-Land: a state-of-the-art global reanalysis dataset for land applications, *Earth Syst. Sci. Data*, 13,
4349-4383, 10.5194/essd-13-4349-2021, 2021.
- 435 Oikawa, P. Y., Sturtevant, C., Knox, S. H., Verfaillie, J., Huang, Y. W., and Baldocchi, D. D.: Revisiting the partitioning
of net ecosystem exchange of CO₂ into photosynthesis and respiration with simultaneous flux measurements of ¹³CO₂ and
CO₂, soil respiration and a biophysical model, *CANVEG, Agricultural and Forest Meteorology*, 234-235, 149-163,
<https://doi.org/10.1016/j.agrformet.2016.12.016>, 2017.
- Pastorello, G., Trotta, C., Canfora, E., Chu, H., Christianson, D., Cheah, Y.-W., Poindexter, C., Chen, J., Elbashandy, A.,
Humphrey, M., Isaac, P., Polidori, D., Reichstein, M., Ribeca, A., van Ingen, C., Vuichard, N., Zhang, L., Amiro, B., Ammann,
440 C., Arain, M. A., Ardó, J., Arkebauer, T., Arndt, S. K., Arriga, N., Aubinet, M., Aurela, M., Baldocchi, D., Barr, A.,
Beamesderfer, E., Marchesini, L. B., Bergeron, O., Beringer, J., Bernhofer, C., Berveiller, D., Billesbach, D., Black, T. A.,
Blanken, P. D., Bohrer, G., Boike, J., Bolstad, P. V., Bonal, D., Bonnefond, J.-M., Bowling, D. R., Bracho, R., Brodeur, J.,
Brümmer, C., Buchmann, N., Burbank, B., Burns, S. P., Buysse, P., Cale, P., Cavagna, M., Cellier, P., Chen, S., Chini, I.,
Christensen, T. R., Cleverly, J., Collalti, A., Consalvo, C., Cook, B. D., Cook, D., Coursolle, C., Cremonese, E., Curtis, P. S.,
445 D'Andrea, E., da Rocha, H., Dai, X., Davis, K. J., Cinti, B. D., Grandcourt, A. d., Ligne, A. D., De Oliveira, R. C., Delpiere,
N., Desai, A. R., Di Bella, C. M., Tommasi, P. d., Dolman, H., Domingo, F., Dong, G., Dore, S., Duce, P., Dufrêne, E., Dunn,
A., Dušek, J., Eamus, D., Eichelmann, U., ElKhidir, H. A. M., Eugster, W., Ewenz, C. M., Ewers, B., Famulari, D., Fares, S.,
Feigenwinter, I., Feitz, A., Fensholt, R., Filippa, G., Fischer, M., Frank, J., Galvagno, M., Gharun, M., Gianelle, D., Gielen,
B., Gioli, B., Gitelson, A., Godeed, I., Goeckede, M., Goldstein, A. H., Gough, C. M., Goulden, M. L., Graf, A., Griebel, A.,
450 Gruening, C., Grünwald, T., Hammerle, A., Han, S., Han, X., Hansen, B. U., Hanson, C., Hatakka, J., He, Y., Hehn, M.,
Heinesch, B., Hinko-Najera, N., Hörtnagl, L., Hutley, L., Ibrom, A., Ikawa, H., Jackowicz-Korczynski, M., Janouš, D., Jans,
W., Jassal, R., Jiang, S., Kato, T., Khomik, M., Klatt, J., Knohl, A., Knox, S., Kobayashi, H., Koerber, G., Kolle, O., Kosugi,
Y., Kotani, A., Kowalski, A., Kruijt, B., Kurbatova, J., Kutsch, W. L., Kwon, H., Launiainen, S., Laurila, T., Law, B., Leuning,
R., Li, Y., Liddell, M., Limousin, J.-M., Lion, M., Liska, A. J., Lohila, A., López-Ballesteros, A., López-Blanco, E., Loubet,
455 B., Loustau, D., Lucas-Moffat, A., Lüers, J., Ma, S., Macfarlane, C., Magliulo, V., Maier, R., Mammarella, I., Manca, G.,
Marcolla, B., Margolis, H. A., Marras, S., Massman, W., Mastepanov, M., Matamala, R., Matthes, G. H., Mazzenga, F.,
McCaughey, H., McHugh, I., McMillan, A. M. S., Merbold, L., Meyer, W., Meyers, T., Miller, S. D., Minerbi, S., Moderow,
U., Monson, R. K., Montagnani, L., Moore, C. E., Moors, E., Moreaux, V., Moureaux, C., Munger, J. W., Nakai, T., Neiryneck,
J., Nesci, Z., Nicolini, G., Noormets, A., Northwood, M., Nosetto, M., Nouvellon, Y., Novick, K., Oechel, W., Olesen, J. E.,
460 Ourcival, J.-M., Papuga, S. A., Parmentier, F.-J., Paul-Limoges, E., Pavelka, M., Peichl, M., Pendall, E., Phillips, R. P.,
Pilegaard, K., Pirk, N., Posse, G., Powell, T., Prasse, H., Prober, S. M., Rambal, S., Rannik, Ü., Raz-Yaseef, N., Rebmann, C.,
Reed, D., Dios, V. R. d., Restrepo-Coupe, N., Reverter, B. R., Roland, M., Sabbatini, S., Sachs, T., Saleska, S. R., Sánchez-
Cañete, E. P., Sanchez-Mejia, Z. M., Schmid, H. P., Schmidt, M., Schneider, K., Schrader, F., Schroder, I., Scott, R. L., Sedláč,
P., Serrano-Ortíz, P., Shao, C., Shi, P., Shironya, I., Siebicke, L., Šigut, L., Silberstein, R., Sirca, C., Spano, D., Steinbrecher,
465 R., Stevens, R. M., Sturtevant, C., Suyker, A., Tagesson, T., Takanashi, S., Tang, Y., Tapper, N., Thom, J., Tomassucci, M.,
Tuovinen, J.-P., Urbanski, S., Valentini, R., van der Molen, M., van Gorsel, E., van Huissteden, K., Varlagin, A., Verfaillie,
J., Vesala, T., Vincke, C., Vitale, D., Vygodskaya, N., Walker, J. P., Walter-Shea, E., Wang, H., Weber, R., Westermann, S.,
Wille, C., Wofsy, S., Wohlfahrt, G., Wolf, S., Woodgate, W., Li, Y., Zampedri, R., Zhang, J., Zhou, G., Zona, D., Agarwal,
D., Biraud, S., Torn, M., and Papale, D.: The FLUXNET2015 dataset and the ONEFlux processing pipeline for eddy
470 covariance data, *Scientific Data*, 7, 225, 10.1038/s41597-020-0534-3, 2020.
- Raj, R., Hamm, N. A. S., Tol, C. V. D., and Stein, A.: Uncertainty analysis of gross primary production partitioned from
net ecosystem exchange measurements, *Biogeosciences*, 13, 1409-1422, 10.5194/bg-13-1409-2016, 2016.
- Reichstein, M., Kätterer, T., Andrén, O., Ciais, P., Schulze, E.-D., Cramer, W., Papale, D., and Valentini, R.: Temperature
sensitivity of decomposition in relation to soil organic matter pools: critique and outlook, *Biogeosciences*, 2, 317-321, 2005.
- 475 Rousseeuw, P. J. and Croux, C.: Alternatives to the median absolute deviation, *Journal of the American Statistical
association*, 88, 1273-1283, 1993.
- Shi, P., Sun, X., Xu, L., Zhang, X., He, Y., Zhang, D., and Yu, G.: Net ecosystem CO₂ exchange and controlling factors
in a steppe—Kobresia meadow on the Tibetan Plateau, *Science in China Series D: Earth Sciences*, 49, 207-218, 2006.
- 480 Sun, S., Che, T., Li, H., Wang, T., Ma, C., Liu, B., Wu, Y., and Song, Z.: Water and carbon dioxide exchange of an alpine
meadow ecosystem in the northeastern Tibetan Plateau is energy-limited, *Agricultural and Forest Meteorology*, 275, 283-295,
2019.



- Tramontana, G., Migliavacca, M., Jung, M., Reichstein, M., Keenan, T. F., Camps-Valls, G., Ogee, J., Verrelst, J., and Papale, D.: Partitioning net carbon dioxide fluxes into photosynthesis and respiration using neural networks, *Global Change Biol.*, 26, 5235-5253, <https://doi.org/10.1111/gcb.15203>, 2020.
- 485 Vekuri, H., Tuovinen, J.-P., Kulmala, L., Papale, D., Kolari, P., Aurela, M., Laurila, T., Liski, J., and Lohila, A.: A widely-used eddy covariance gap-filling method creates systematic bias in carbon balance estimates, *Scientific Reports*, 13, 1720, 2023.
- Wang, H., Li, X., Xiao, J., Ma, M., Tan, J., Wang, X., and Geng, L.: Carbon fluxes across alpine, oasis, and desert ecosystems in northwestern China: The importance of water availability, *Sci Total Environ.*, 697, 133978, 2019.
- 490 Wang, S., Chen, W., Fu, Z., Li, Z., Wang, J., Liao, J., and Niu, S.: Seasonal and inter-annual variations of carbon dioxide fluxes and their determinants in an Alpine meadow, *Frontiers in Plant Science*, 13, 894398, 2022.
- Wang, X., Ma, M., Huang, G., Veroustraete, F., Zhang, Z., Song, Y., and Tan, J.: Vegetation primary production estimation at maize and alpine meadow over the Heihe River Basin, China, *International Journal of Applied Earth Observation and Geoinformation*, 17, 94-101, [10.1016/j.jag.2011.09.009](https://doi.org/10.1016/j.jag.2011.09.009), 2012.
- 495 Wang, X., Che, C., Xiao, J., Wang, T., Tan, J., Zhang, Y., Ren, Z., Geng, L., Wang, H., Xu, Z., Liu, S., and Li, X.: A post-processed carbon flux dataset for 34 eddy covariance flux sites across the Heihe River Basin, China, *National Tibetan Plateau Data Center [dataset]*, [10.11888/Terre.tpdc.301321](https://doi.org/10.11888/Terre.tpdc.301321), 2024.
- Wutzler, T., Lucas-Moffat, A., Migliavacca, M., Knauer, J., Sickel, K., Šigut, L., Menzer, O., and Reichstein, M.: Basic and extensible post-processing of eddy covariance flux data with REdDyProc, *Biogeosciences*, 15, 5015-5030, [10.5194/bg-15-5015-2018](https://doi.org/10.5194/bg-15-5015-2018), 2018.
- 500 Xu, Z., Liu, S., Zhu, Z., Zhou, J., Shi, W., Xu, T., Yang, X., Zhang, Y., and He, X.: Exploring evapotranspiration changes in a typical endorheic basin through the integrated observatory network, *Agricultural and Forest Meteorology*, 290, 108010, 2020.
- Zhao, L., LI, Y. N., Gu, S., ZHAO, X. Q., XU, S. X., and YU, G. R.: Carbon dioxide exchange between the atmosphere and an alpine shrubland meadow during the growing season on the Qinghai-Tibetan Plateau, *Journal of Integrative Plant Biology*, 47, 271-282, 2005.
- 505 Zhu, S., Clement, R., McCalmont, J., Davies, C. A., and Hill, T.: Stable gap-filling for longer eddy covariance data gaps: A globally validated machine-learning approach for carbon dioxide, water, and energy fluxes, *Agricultural and Forest Meteorology*, 314, 108777, 2022.

510