

The IAS2024 coastal sea level dataset and first evaluations

Fukai Peng^{1,*}, Xiaoli Deng², Yunzhong Shen³, Xiao Cheng¹

¹ School of Geospatial Engineering and Science, Sun Yat-Sen University, Guangzhou 510275, China

² School of Engineering, The University of Newcastle, University Drive, Callaghan, NSW 2308, Australia

5 ³ College of Surveying and Geo-informatics, Tongji University, Shanghai 200092, China

Correspondence to: Fukai Peng (pengfk@mail.sysu.edu.cn)

Abstract. A new dedicated 20-Hz coastal sea level dataset, called the International Altimetry Service 2024 (IAS2024, <https://doi.org/10.5281/zenodo.13208305>, Peng et al., 2024c), is presented for monitoring sea level changes along the world's coastlines. One of reasons for generating this dataset is that the quality of coastal altimeter data has been greatly improved with advanced coastal reprocessing strategies. In this study, the Seamless Combination of Multiple Retrackerers (SCMR) strategy is adopted to obtain the reprocessed Jason data from January 2002 to April 2022. The evaluation/validation results show that the IAS2024 20-Hz along-track coastal sea level dataset achieves good performance over global coastal oceans. The good consistency between IAS2024 and independent altimeter datasets, including the European Space Agency Climate Change Initiative version 2.4 (ESA CCI v2.4) 20-Hz along-track coastal sea level dataset and the Copernicus Marine Environment Monitoring Service Level-3 (CMEMS L3) 1-Hz along-track sea level dataset, is observed. The closure of sea level trend differences ($0.16 \pm 3.97 \text{ mm yr}^{-1}$) between IAS2024 and Permanent Service for Mean Sea Level (PSMSL) tide gauge data at the global scale is also achieved. Moreover, 1548 virtual stations have been constructed using the IAS2024 coastal sea level dataset, which will contribute to the analysis of coastal sea levels for the ocean community and to the risk management for the policy makers. Our study also found that no obvious variations exist in the linear sea level trends from offshore to the coast over the last 20 km coastal strip at the global scale. In addition, the vertical land motion (VLM) estimates from the combination of the IAS2024 dataset with the PSMSL tide gauge records agree well with the University of La Rochelle 7a (ULR7a) Global Navigation Satellite System (GNSS) solution, with mean difference of VLM estimates being $0.12 \pm 2.27 \text{ mm yr}^{-1}$, suggesting that altimeter-derived VLM estimates can be used as an independent data source to validate the GNSS solutions.

1 Introduction

25 Satellite altimetry has become a mature remote sensing technique over open oceans since the launch of high-precision satellite altimetry missions (e.g., Topex/Poseidon). It has been making great contributions to quantifying and monitoring sea level changes at both global and regional scales (Ablain et al., 2016; Prandi et al., 2021; Guérou et al., 2023). Among these missions, the Jason altimetry series are usually used as the reference missions for monitoring global sea levels considering their high precision and stable performance. During the past two decades, the precision of reprocessed 20-Hz along-track altimeter data in coastal zones from Jason missions has been remarkably increased thanks to the dedicated coastal retrackerers and improved

30

range/geophysical corrections (Cipollini et al., 2017; Vignudelli et al., 2019; Peng et al., 2024a), which makes it possible to construct altimeter-based virtual stations along the world’s coastlines (Benveniste et al., 2020; Cazenave et al., 2022).

The IAS (International Altimetry Service) Pilot Service was established in July 2023 as a service of the International Association of Geodesy (IAG) for providing information about altimetry data, geodetic and climate models, research and operational applications based on satellite altimetry technology innovations. One of its goals is to provide auxiliary data and algorithms to produce new products across the coastal zones including the coastal oceans, land-sea surfaces, estuaries and inland water bodies for applications in studies of the extremes, long-term climate change and environmental development.

In this paper, we present the IAS2024 20-Hz along-track coastal sea level dataset from Jason missions over the period of January 2002 and April 2022 reprocessed with the SCMR (Seamless Combination of Multiple Retracker) strategy. A total of 1548 virtual stations are built along the world’s coastlines, which can be used to monitor the coastal sea levels and **estimate the vertical land motions (VLM)**. In addition, the spatial variation of 20-Hz along-track sea level trends from offshore to nearshore in the last 20 km to the coast is analysed using the IAS2024 dataset. The quality of reprocessed altimeter data is validated against the monthly tide gauge records from the PSMSL (Permanent Service for Mean Sea Level, <https://psmsl.org/>, last accessed on September 29, 2024), the Copernicus Marine Environment Monitoring Service (CMEMS) L3 1-Hz along-track product (https://data.marine.copernicus.eu/product/SEALEVEL_GLO_PHY_L3_MY_008_062/description, last accessed on September 29, 2024) and the European Space Agency (ESA) Climate Change Initiative (CCI) version 2.4 20-Hz along-track coastal sea level dataset (<https://www.seanoe.org/data/00631/74354/>, last accessed on December 5, 2024).

Section 2 presents the details of the SCMR processing strategy and methods to construct the altimeter-based virtual stations and to assess the quality of reprocessed altimeter data. Sections 3 and 4 illustrate the cross-validation results against in-situ and independent altimeter datasets, as well as the application of coastal altimeter datasets. Sections 5 and 6 present the data availability and conclusions, respectively.

2 Computation of the IAS2024 coastal sea level dataset

2.1 The SCMR strategy

The IAS2024 coastal sea level dataset is generated using the SCMR (Peng et al., 2024b) strategy. The flow diagram is illustrated in Fig. 1, which starts from the waveform leading edge detection and ends with the seamless combination of sea surface height (SSH) estimates from multiple retracker. Four different waveform retracker are used in this study, which are the official Sensor Geophysical Data Record (SGDR) Maximum Likelihood Estimator 4-parameter (MLE4) retracker (Amarouche et al., 2004), the Adaptive Leading Edge Sub-waveform (ALES, Passaro et al., 2014), the Weighted Least Squares 3-parameter (WLS3, Peng and Deng, 2018) and the Modified Brown model 4-parameter (MB4) retracker (Poisson et al., 2018). The main steps are described in the following subsections.




Figure 1: Flow diagram of the SCMR processing strategy. It consists of four operating blocks from left to right: (1) Four retracker of the SGDR MLE4, ALES, WLS3 and MB4; (2) Estimating sigma0, SWH, and range estimates from different retracker; (3) Calculating SSH estimates for different retracker; and (4) Removing inter-retracker bias and combining SSH estimates from different retracker. The SSB corrections are individually computed for each retracker. The inter-retracker bias correction model is developed based on the negative correlation between SSH differences (Δh) and SWH differences (ΔH_s) with respect to the SGDR MLE4 retracker. The combination of SSH estimates from four retracker is conducted using the Dijkstra (1959) algorithm based on the situation that the altimeter along-track SSH estimates do not change significantly at the spatial scale of ~ 300 m.

2.1.1 Detection of leading edges

The leading edge is a sequence of waveform gates, where the power increases sharply and thus can be detected based on the differences between consecutive power of gates. The start gate of the leading edge is deemed as the first gate where the power difference of two consecutive gates is positive (Passaro et al., 2014). Similarly, the stop gate of the leading edge is deemed as the first gate where the power difference of two consecutive gates is negative (Passaro et al., 2014) or smaller than a tiny positive number (Wang et al., 2019). Moreover, the power of the start gate should be lower than a constant level, and the power difference between start and stop gates should be larger than a threshold value (Gommenginger et al., 2011). The main steps of the leading-edge detection method are as follows.

First, the intersection points between four horizontal lines with the power levels of 0.1, 0.2, 0.5 and 0.9 and the normalized waveform are derived. For each intersection point, the start and stop gates of the leading edge are searched forward and backward until satisfying the following equation,

$$P_{startgate} - P_{startgate-1} > 0.001 \quad (1)$$

$$P_{stopgate+1} - P_{stopgate} < 0.01$$

where the $P_{startgate}$ and $P_{stopgate}$ are the powers of the normalized waveform at the start and stop gates. If the difference (i.e., $P_{stopgate} - P_{startgate}$) is larger than 0.1 and the $P_{startgate}$ is lower than 0.2, the proportion of waveform from start to stop gates is selected as a potential leading edge. Here, the above threshold levels are selected based on previous studies and our empirical experience (Passaro et al., 2014; Peng and Deng, 2020a).

Second, once all potential leading edges are found, the 50% Threshold retracker (Gommenginger et al., 2011) is applied to each potential leading edge to calculate the range and corresponding SSH estimates. The Dijkstra (1959) algorithm is then applied to find the shortest path between offshore and nearshore along-track points, in which the edge weights are the height differences between connected nodes (Roscher et al., 2017). As a result, the optimal 20-Hz along-track SSH profile and the corresponding leading edge is determined (Peng et al., 2023). The advantage of this method is that it effectively avoids the perturbations before the true leading edge of the waveform (cf. Fig. 2 in Peng et al., 2024b).

2.1.2 Detection of land returns

When the altimeter approaches the coast, the contaminated waveforms appear with peaks that are caused by the high reflective areas inside the illuminated land surfaces or by the modification of the sea state close to the shoreline (Halimi et al., 2013). This type of waveforms is denoted as the Brown-peaky waveform, because the peaks caused by land returns are close to the waveform leading edge, while the waveform shape over oceans follows the Brown model (Fig. 4.5 in Gommenginger et al., 2011). Here, we detect the land returns following the idea of the adaptive peak detection (APD) method (Peng and Deng, 2018; Peng and Deng, 2020a) shown in Fig. 2, but with some modifications. The main steps are as follows.

Firstly, the forward moving average F_t and backward moving average B_t are derived using the normalized waveform,

$$\begin{aligned} F_t(i) &= \frac{1}{N} \sum_{j=0}^{N-1} P_t(i+j), \quad i = 1, 2, \dots, n-N+1 \\ B_t(i) &= \frac{1}{N} \sum_{j=0}^{N-1} P_t(i-j), \quad i = N, N+1, \dots, n \end{aligned} \quad (2)$$

where the N is the moving average step, which is selected as five through our trial-and-error test, n is the number of waveform gates, which is 104 for Jason missions.

Next, we define the Difference I as $(F_t - B_t)$. This is first used to determine the Leftfoot points, which correspond to the local maxima of Difference I whose height and prominence are larger than 0.12 using the MATLAB function “findpeaks”. This is then used to determine the Leftedge, Rightfoot and Rightedge points. The Leftedge and Rightedge points are the zero-crossing points of Difference I before and after the corresponding Leftfoot points. The Rightfoot points are the local minima between consecutive Leftfoot points (Fig. 2). If there exist multiple Leftfoot points, the waveform is denoted as the multi-peak waveform (Figs. 2a and 2b). Otherwise, the waveform is classified as the Brown-peaky waveform if the Difference I of the Rightfoot point is smaller than -0.2 (Figs. 2c and 2d).

Finally, the start gate of land returns is defined as the stop gate of the leading edge, while the stop gate of land returns is the last zero-crossing point between Rightfoot and Rightedge points (black line in Fig. 2d). Note that the modified APD method used here is dedicated for the Brown-peaky waveforms instead of the multi-peak waveforms.





Figure 2: Illustration of the adaptive peak detection (APD) method. The subplots (a) and (b) show the curve of Difference I and feature points detected by the APD method for multi-peak waveforms, respectively. The subplots (c) and (d) show the similar results for Brown-peaky waveforms. The Difference I is calculated as the difference between forward moving average and backward moving average of the normalized waveform. The black line corresponds to the gates affected by the land returns.

2.1.3 Waveform retracking

All retrackers included in Fig. 1 can process ocean waveforms. Of them, the ALES retracker can handle waveforms with land returns observed in the waveform trailing edge, the WLS3 retracker can cope with the Brown-peaky waveforms with the

125 downsized weights (i.e., 0.01) being assigned to land returns determined by the modified APD method, and the MB4 retracker can deal with the specular waveforms (Fig. 3). Note that the estimates from the SGDR MLE4 are directly provided by the official product, while the estimates from the ALES, WLS3 and MB4 retrackers are solved independently.



130 **Figure 3: An example of observed waveforms and fitted waveforms from different waveform retrackerers. The subplots (a)-(d) show the fitted results for different waveform types by methods of (a) SGDR MLE4, (b) ALES, (c) WLS3 and (d) MB4 retrackerers.**

2.1.4 Calculation of altimeter SSH

The altimeter SSH at each 20-Hz along-track point is calculated as follows,

135
$$SSH = h_{alt} - R_r - R_{cor} \quad (3)$$

where the h_{alt} is the height of the satellite above the Topex ellipsoid, the R_r is the distance between the altimeter and nadir sea surface derived from the waveform retracker, the R_{cor} include dry and wet tropospheric, ionospheric, and sea state bias corrections. **Note that the Topex ellipsoid is a reference ellipsoid which has an equatorial radius of 6378.1363 kilometre and a flattening coefficient of 1/298.257.** The appropriate corrections for each altimetry mission are investigated and summarized in Table 1.

Table 1. Range/geophysical corrections applied to Jason missions used in this study.

Corrections	Jason-1/2/3
Dry tropospheric correction	ECMWF (Hersbach et al., 2023)
Wet tropospheric correction	ECMWF*
Ionospheric correction	GIM (Komjathy and Born, 1999)
Sea state bias	Peng and Deng, (2020b)
Geocentric ocean and loading tide	FES2014 (Lyard et al., 2021)
Dynamic atmospheric correction	MOG2D (Carrère and Lyard, 2003)
Solid earth tide	Cartwright and Taylor (1971)
Pole tide	Desai et al., (2015)
Mean sea surface	CLS22 (Schaeffer et al., 2023)

The modelled wet tropospheric correction at zero altitude has been used instead of the GPD+ correction (Fernandes et al., 2016) considering that the GPD+ is unavailable beyond February 2021.

2.1.5 Seamless combination of SSH estimates from multiple retrackers

As illustrated by Fig.1, the coastal retrackers (i.e., ALES, WLS3 and MB4) are applied to all 20-Hz waveforms to **derive** range estimates, respectively. Combined with satellite altitude, range and geophysical corrections, the SSH estimates for these coastal retrackers and the SGDR MLE4 retracker are obtained. Considering the above retrackers can handle different types of waveforms as shown in Fig. 3 and achieve similar performance through our previous Monte Carlo simulation (Peng and Deng, 2018, Peng et al., 2021), it is possible to further improve the data availability by combining SSH estimates from these retrackers. However, these retrackers adopt different processing strategies, there inevitably exist systematic biases between SSH estimates from these retrackers, which would prohibit the seamless combination of SSH estimates. To solve this problem, we proposed an inter-retracker bias model to reduce the SSH bias with respect to the SGDR MLE4 from centimeter levels to millimeter levels (Peng et al., 2024b). Finally, the optimal SSH estimate at each along-track point is selected using the Dijkstra (1959) algorithm. The Dijkstra algorithm is used because it can automatically determine the along-track SSH profile with smooth

spatial variation (Roscher et al., 2017), which is consistent with the real situation that the altimeter along-track SSH estimates do not change significantly at the spatial scale of ~300 m (Cipollini et al., 2017). Moreover, the Dijkstra algorithm does not rely on the waveform classification results. For more detailed information about the use of Dijkstra algorithm, readers can refer to Peng et al. (2023).

160 The analytical form of the inter-retracker bias correction model is as follows (Peng et al., 2024b),

$$\Delta h = \rho \times \Delta H_s + c_b \quad (4)$$

$$h_{cor} = \hat{\rho} \times \Delta H_s + \hat{c}_b \quad (5)$$

where ΔH_s and Δh are the **significant wave height (SWH)** differences and SSH differences with respect to the SGDR MLE4 retracker (e.g., ALES minus SGDR MLE4), ρ and c_b are the linear regression slope and intercept parameters, which are unknown and need to be estimated. Once estimates of two parameters, $\hat{\rho}$ and \hat{c}_b , are determined, the inter-retracker bias correction, h_{cor} , can be calculated by substituting SWH differences ΔH_s into the Equation (5). After removing the inter-retracker bias, the most appropriate SSH profile between the offshore point and the point closest to the coast is derived by using the Dijkstra algorithm, in which the edge weight is defined as the absolute SSH difference between two connected nodes.

170 2.1.6 Calculation of altimeter SLA

The altimeter SLA estimate at each 20-Hz along-track point is calculated as follows,

$$SLA = SSH - G_{cor} - MSS \quad (6)$$

where the G_{cor} include geocentric ocean and loading tide corrections, dynamic atmospheric correction, solid earth tide and pole tide corrections, the MSS is the interpolated value from the CLS22 mean sea surface (MSS) model (Schaeffer et al., 2023) using the bilinear interpolation.

2.2 Assessment of the IAS2024 coastal sea level dataset

The assessment of the IAS2024 coastal sea level dataset is conducted as follows. Firstly, the data availability and precision of 20-Hz SLA estimates are calculated as a function of distance to the coast. The data availability is calculated as the percentage of available 20-Hz SLA estimates, while the data precision is represented as the median value of standard deviation (STD) of 20-Hz SLA estimates (Cipollini et al., 2017).

Secondly, the power spectrums of 20-Hz SLA estimates are computed with the periodogram method (Zanifé et al., 2003). The SLA power spectrum reflects the strength of SLA signals at different spatial scales and can be used to estimate the noise level from the high frequency part of the spectrum. In addition, the crossover analysis (Gaspar et al., 1994) is conducted to examine the spatial variation of data quality over global coastal oceans. For each single mission, the crossover point is the intersection **between the ascending and descending ground tracks**. The 20-Hz along-track points closest to the crossover point are used to

derive the collocated SLA estimates at the crossover point using the linear interpolation method. To reduce the effect of temporal variability, the collocated SLA estimates are only considered if their time lags are within one day.

Finally, the IAS2024 dataset is validated against monthly tide gauge records from the PSMSL (Holgate et al., 2013). To guarantee the robustness of the validation results, 549 tide gauges which have continuous data records over the period of 2002 and 2022 are used in this study (Fig. 4).




Figure 4: Distribution of (a) 549 tide gauges from PSMSL and (b) 1548 altimeter-based virtual stations from the IAS2024 dataset.

The main steps of the validation procedure are described as follows.

- 1) the Jason-3 ground tracks which achieve the maximum data points within 100 km to the coast are selected as the nominal ground track. The corresponding 20-Hz offshore distances are directly retrieved from the Jason-3 SGDR product, which are calculated with respect to the Global Self-consistent, Hierarchical, High-resolution Geography Database (GSHHG) dataset (<https://www.soest.hawaii.edu/pwessel/gshhg/>). After that, the single nominal ground track is divided into multiple ground track segments if the distance between consecutive 20-Hz along-track points is larger than 10 km. Only the ground track segments whose near-shore points with distances to the coast being smaller than 10 km and offshore points with distances to the coast being larger than 90 km are used in this study. It is found that there are 1458 ground track segments which fulfill the above criterion, and the distribution of closest distance to the coast for each ground track segment is shown in Fig. 5.
- 2) The SCMR-reprocessed 20-Hz along-track SLA estimates from the Jason missions are referenced to the corresponding ground track segments using the nearest neighborhood approach, and a 3-sigma filter is then applied to the along-track SLA estimates to remove the outliers. The inter-mission SLA biases between different Jason missions are estimated and removed using the overlapping time series method (Peng et al., 2022) before constructing SLA time series with a temporal sampling of 10 days. Then, the 3-sigma filter is again applied to the SLA time series. In this study, only time series with more than 80% of available data are used to derive the monthly SLA time series over the period of January 2002 and April 2022.
- 3) The linear sea level trends are derived from the de-seasoned monthly SLA time series with the Hector software (Bos et al., 2014), which can handle the time series with temporally correlated noise. The stochastic noise models used in

this study include the first- and fifth-order autoregressive (i.e., AR1 and AR5), and the autoregressive fractionally-integrated moving-average (ARFIMA) models. The most appropriate noise model is identified using the lowest mean value of both the Akaike Information Criterion (AIC; Akaike, 1992) and the Bayesian Information Criterion (BIC; Schwarz, 1978).

- 4) The monthly SLA time series of 20-Hz along-track points over the 0-20 km coastal strip are averaged to generate monthly SLA time series at altimeter-based virtual stations. The collocated virtual stations and tide gauges are derived if their distance is smaller than 200 km and the score between them reaches the maximum. The 200 km distance threshold is chosen due to the trade-off between including enough nearby virtual stations and retaining high correlation coefficients. The score is calculated using the correlation coefficient and root mean square (RMS) of the differences between the monthly SLA time series from the virtual station and tide gauge as follows,

$$score = score_{cc}^i \times 0.4 + score_{rms}^i \times 0.6 \quad (7)$$

$$score_{cc}^i = \frac{cc_i - \min(cc)}{\max(cc) - \min(cc)} \times 100$$

$$score_{rms}^i = \frac{\max(rms) - rms_i}{\max(rms) - \min(rms)} \times 100$$

The number of collocated stations, the correlation coefficients and RMS values of the differences between monthly SLA time series from collocated stations, as well as the differences of linear sea level trends derived from the monthly SLA time series are used for the validation.




Figure 5: Distribution of closest distance to the coast for 1548 ground track segments.

The ESA CCI v2.4 20-Hz and the CMEMS L3 1-Hz along-track sea level datasets are also interpolated into the ground track segments using the nearest neighbourhood approach. After that, the results from both the ESA CCI and CMEMS products are

compared with that from the IAS2024 dataset. For comparison, we only used ground track segments that have at least 10 common points with the ESA CCI and 2 common points with the CMEMS within the last 20 km to the coast. When it comes to the correlation coefficients, only those whose p-value are smaller than 0.05 are used.

240 2.3 Application of the IAS2024 coastal sea level dataset

Three applications of coastal altimeter datasets are presented in this study. Firstly, the coastal altimeter datasets can be used to build altimeter-based virtual stations (Benveniste et al., 2020; Cazenave et al., 2022), which is of great importance for monitoring the coastal sea level change and constraining the high-resolution ocean models. To examine the data quality of virtual stations obtained from different coastal strips, the monthly SLA time series of 20-Hz along-track points over 0-10 km, 5-15
245 km and 10-20 km are averaged for comparison. For clarity, these three types of virtual stations are denoted as onshore, near-shore and offshore virtual stations.

Secondly, the spatial variation of sea level trends can be analysed because the altimeter provides the along-track sea level trend profile, which would help us understand the relative contributions of local and remote processes on coastal sea levels. To address this issue, a linear regression is applied to the along-track sea level trends using the MATLAB function “robustfit”.
250 The trend differences between nearshore and offshore points in the last 20 km to the coast are then calculated to examine whether there exists significant change ($>\pm 2 \text{ mm yr}^{-1}$) of the along-track sea level trends.

Finally, the VLM estimates can be obtained by calculating the trend estimates from differenced monthly SLA time series between collocated virtual stations and tide gauges, which is used for the comparison with corresponding Global Navigation Satellite System (GNSS) solutions called the University of La Rochelle 7a (ULR7a) (Gravelle et al., 2023). The ULR7a solu-
255 tion is a preliminary version of the reanalysis of 21 years of GPS data from 2000 to 2020 that has been undertaken within the framework of the third data reprocessing campaign of the International GNSS Service (IGS), whose associated vertical velocity field is expressed in the International Terrestrial Reference Frame 2014 (ITRF 2014, Altamimi et al., 2016).

3 Evaluation

3.1 Data availability and precision

260 **Figure 6** shows the data availability and precision for the official SGDR MLE4 data and SCMR-reprocessed data over different Jason missions. These results are derived after removing the SLA estimates whose absolute values are greater than 1 m. As we can see from the graph, the results from different Jason missions are similar, demonstrating the stability of the Jason missions. Thanks to the Open Loop tracking mode (Biancamaria et al., 2018), the Jason-3 can recover more reliable SLA estimates than the other two satellites. The results also demonstrate that the SCMR strategy outperforms the official SGDR MLE4 over global
265 coastal oceans. The SCMR can remarkably increase the data availability over the entire 0-20 km distance bands, as well as the

data precision beyond 6 km to the coast. Note that the higher precision achieved by the official SGDR MLE4 within 0-6 km distance band can be attributed to the low data availability (Figs. 6a to 6c) and dedicated data editing strategy applied to the official SGDR product (Peng et al., 2024b). The improved percentage in terms of data availability and precision in different coastal strips (i.e., 0-5 km, 5-10 km and 10-20 km) are summarized in Table 2.

270 As can be seen from Table 2, the improved percentage for data availability increases with decreasing offshore distance. In contrast, the data precision shows a declining trend from offshore oceans towards the coast. This result demonstrates that the SCMR strategy can handle multiple waveform types and thus can recover more reliable SLA estimates than the official SGDR MLE4. Within distance 0-5 km to the coast, the data precision with respect to the official SGDR MLE4 is not improved, which is mainly attributed to the above reasons. In addition, the range estimates from contaminated waveforms and range/geophysical

275 corrections near the coast are inferior to those over open oceans, which results in a smaller or even negative improvement percentage of data precision. Therefore, a dedicated data editing strategy should be developed for the coastal altimeter data as what has been done in the SGDR product. Overall, the data availability from SCMR can retain more than 90% beyond 5 km offshore and more than 70%-80% in the last 5 km to the coast (Figs. 6a-6c). The data precision can retain at centimetre levels (~4.9 cm) over the 10-20 km distance band, increase to 7-9 cm over the 5-10 km distance band and rise up to decimetre levels (~20-23 cm) towards the coast (Figs. 5d-5f).

280 (~20-23 cm) towards the coast (Figs. 5d-5f).




Figure 6: Data availability (a-c) and precision (d-f) for official SGDR MLE4 data (in black) and SCMR-reprocessed data (in red). From left to right, the results for Jason-1, Jason-2 and Jason-3 are shown successively. The data availability is calculated as the percentage of available 20-Hz SLA estimates, while the data precision is represented as the median value of standard deviation (STD) of 20-Hz SLA estimates within one second in each 1-km distance band.

Table 2. Improved percentage of the SCMR against the official SGDR MLE4 in terms of data availability and precision over different coastal strips.

Satellite altimeters	0-5 km		5-10 km		10-20 km	
	Availability	Precision	Availability	Precision	Availability	Precision
Jason-1	53.0%	−72.4%	20.3%	7.0%	12.6%	9.2%
Jason-2	46.1%	−54.7%	20.7%	4.8%	14.7%	8.4%
Jason-3	53.2%	−51.9%	20.4%	10.1%	14.7%	10.5%

290

The power spectrum and crossover analysis (Fig. 7) further demonstrate the good performance of the SCMR strategy in improving the data availability and reducing noise levels at small spatial scales. Compared to the official SGDR MLE4, the SCMR achieves a reduction in the SLA spectra at the scales below 50 km, with the noise level at sub-kilometre scales being 23% less than that of the official SGDR MLE4 (Figs. 7a-7c). In addition, the crossover analysis shows that the SCMR significantly increases the number of collocated 20-Hz SLA estimates by 44%, while retaining smaller mean differences close to zero and slightly larger standard deviations around 14 cm (Figs. 7d-7f).

295




Figure 7: Power spectrum of 20-Hz SLA estimates (a-c) and the number, mean and standard deviation of differences from collocated 20-Hz SLA estimates at crossover points (d-f). J1, J2 and J3 represent Jason-1, Jason-2 and Jason-3 missions, respectively.

3.2 Comparison between IAS2024 and ESA CCI v2.4

In this section, the quality of the IAS2024 coastal dataset is investigated by comparing with the latest ESA CCI v2.4 coastal sea level dataset. The two datasets are compared on 807 ground track segments, where contain at least 10 common points for both IAS2024 and ESA CCI v2.4 datasets over the 0-20 km coastal strip. The selection of 10 common points is to guarantee the robustness of the results. To conduct the comparison between these two datasets, we first calculate the point-wise correlation coefficient and RMS of the differences between monthly SLA time series from these two datasets. Then, the averaged correlation coefficients and RMS values for different ground track segments are obtained.

Figure 8 shows the histograms of correlation coefficients and RMS values between detrended and de-seasoned monthly SLA time series from IAS2024 and ESA CCI v2.4 datasets for different ground track segments over global coastal oceans. As we can see from the graph, the consistency of the monthly SLA time series between these two datasets is good at the global scale, which is consistent with our previous study in the Australian coastal zone (Peng et al., 2022). The relatively high correlation coefficients (>0.4) are achieved in most cases. As a result, the mean correlation coefficient between IAS2024 and ESA CCI v2.4 datasets is 0.59, while the corresponding RMS values mostly varying between the range of 40 mm and 80 mm. The spatial distribution of both correlation coefficients and RMS values (Fig. 9) reveals that the discrepancy between IAS2024 and ESA CCI v2.4 datasets is only observed in the west coast of South America.




Figure 8: Distribution of averaged correlation coefficients (a) and RMS values (b) between detrended and de-seasoned monthly SLA time series from IAS2024 and ESA CCI v2.4 datasets for different ground track segments over the 0-20 km coastal strip.

320




Figure 9: Spatial distribution of averaged correlation coefficients (a) and RMS values (b) between detrended and de-seasoned monthly SLA time series from IAS2024 and ESA CCI v2.4 datasets for different ground track segments over the 0-20 km coastal strip.

325

In addition to the comparison of monthly SLA time series, we also analyse the sea level trends from these two datasets over the similar period. First, the point-wise sea level trends and trend uncertainties are derived from both IAS2024 and ESA CCI v2.4 datasets. Then, the trend differences and uncertainties at common points are averaged for each ground track segment. Finally, the histograms of averaged trend differences and trend uncertainties are shown in Fig. 10.

330

As we can see from the graph, the trend differences between them are predominantly within the range between -4 mm yr^{-1} and 6 mm yr^{-1} , with a mean of trend differences being $1.32 \pm 2.40 \text{ mm yr}^{-1}$ (Fig. 10a). The trend uncertainties for both datasets are similar, varying between 0.5 mm yr^{-1} and 2.0 mm yr^{-1} (Fig. 10b). This result implicates that the sea level trends derived from the IAS2024 dataset are generally higher than those from the ESA CCI v2.4 dataset, which is in line with our previous findings (Peng et al., 2022; Peng et al., 2024a). The reason behind this may be attributed to the different data processing techniques adopted, especially the methods used to estimate the inter-mission SLA biases (Peng et al., 2022).

335




Figure 10: Distribution of (a) averaged trend differences from de-seasoned monthly SLA time series between IAS2024 and ESA CCI v2.4 datasets, as well as (b) the trend uncertainties for different ground track segments over the 0-20 km coastal strip. The violet bars in subplot (b) are the overlaps between IAS2024 and ESA CCI.

At last, the tide gauge records from the PSMSL are deemed as the ground truth and used for the validation of these two datasets. Table 3 summarizes the number of collocated virtual stations and tide gauges for both IAS2024 and ESA CCI v2.4 datasets. In addition, the mean of correlation coefficients and RMS values, and the mean of trend differences between altimeter and tide gauge are listed in the Table. As illustrated by the table, the number of collocated stations for the IAS2024 dataset (290) is similar to that for the ESA CCI v2.4 dataset (293). This is because the same ground track segments are used for comparison. The comparable performance is also observed for these two datasets in terms of correlation coefficients and RMS values. However, these two datasets differ in the trend differences between altimeter and tide gauge at the global scale. As reported by previous studies (e.g., Waston et al., 2015; Wöppelmann and Marcos, 2016), the VLM would be cancelled out on average along the world’s coastlines. Therefore, it is reasonable to assume that the average trend difference between the altimeter and tide gauge is close to zero at the global scale. The result from the IAS2024 dataset is thus consistent with this assumption **even it is not a true global scale**, as the mean of trend differences is -0.26 ± 3.57 mm yr⁻¹, which demonstrates the good performance of the IAS2024 dataset in monitoring the coastal sea levels. However, a negative trend difference (-1.50 ± 3.31 mm yr⁻¹) is shown in Table 3 when validating the ESA CCI v2.4 dataset against the tide gauge records. This may be ascribed to the fact that there exists a systematic bias (1.32 ± 2.40 mm yr⁻¹) between sea level trends from IAS2024 and ESA CCI v2.4 datasets.

355

Table 3. Validation results of IAS2024 and ESA CCI v2.4 datasets against tide gauge records from PSMSL. Note that the correlation coefficients and RMS values are derived from the detrended and de-seasoned monthly SLA time series, while the sea level trends are derived from the de-seasoned monthly SLA time series.

Altimeter datasets	Number of collocated stations	Correlation coefficient with tide gauges	RMS (mm)	Trend difference (mm yr ⁻¹)
IAS2024	290	0.64	43.07	-0.26 ± 3.57
ESA CCI v2.4	293	0.61	38.45	-1.50 ± 3.31

360 Overall, the above comparison results reveal the good consistency between IAS2024 and ESA CCI v2.4 datasets over the same ground track segments, though there exists a systematic bias between their sea level trends. It is also noted that the IAS2024 dataset can achieve higher spatial coverage because the ESA CCI v2.4 dataset has no data in places such as Japan and the northern Europe (see Sections 4.1 and 4.2).

365 **3.3 Comparison between IAS2024 and CMEMS L3 along-track product**

The IAS2024 coastal dataset is also compared with the CMEMS L3 1-Hz along-track product. Note that 830 ground track segments **which** contain at least 2 common points for these two datasets over the 0-20 km coastal strip are used for comparison.

The selection of 2 common points is because the CMEMS L3 product provides the 1-Hz (~7 km along-track) instead of 20-Hz (~300 m along track) SLA data.

370 **Figure 11** shows the histograms of averaged correlation coefficients and RMS values between detrended and de-seasoned monthly SLA time series from the IAS2024 and CMEMS datasets for different ground track segments. As we can see from the graph, the correlation coefficients mostly vary between 0.4 and 0.8, and the RMS values are mainly smaller than 60 mm. This result is in line with the comparison result between IAS2024 and ESA CCI v2.4 datasets shown in Section 3.2, indicating the good consistency between IAS2024 and CMEMS at the global scale. The spatial distribution of both correlation coefficients and RMS values (Fig. 12) further reveals that the low correlation coefficients between IAS2024 and CMEMS datasets are again observed in the west coast of South America.

375




Figure 11: Same as Figure 7 but for the comparison between IAS2024 and CMEMS L3 along-track datasets.





Figure 12: Same as Figure 8 but for the comparison between IAS2024 and CMEMS L3 along-track datasets.

Figure 13 shows the histograms of averaged trend differences and uncertainties. As illustrated by the graph, most of the trend differences vary between ± 4 mm yr⁻¹, while the trend uncertainties are within the range of 0.5 mm yr⁻¹ and 2 mm yr⁻¹. The mean trend difference between IAS2024 and CMEMS is -0.18 ± 2.17 mm yr⁻¹, indicating the systematic bias between these

385

two datasets is relatively small. This result is in contrast with that between IAS2024 and ESA CCI v2.4 (see Section 3.2). The reason behind this requires further investigation, which would be our future work.



390 **Figure 13:** Same as Figure 9 but for the comparison between IAS2024 and CMEMS L3 along-track datasets. The violet bars in subplot (b) are the overlaps between IAS2024 and CMEMS.

In the end, the validation results against tide gauge records for IAS2024 and CMEMS are summarized in Table 4. As we can see from the table, the results for these two datasets are in good agreement. Moreover, the trend differences derived from these two datasets are similar and close to zero, which contrasts with the result from the ESA CCI v2.4 dataset (-1.50 ± 3.31 mm yr⁻¹). The reason behind this may be attributed to the fact that the systematic bias (-0.18 ± 2.17 mm yr⁻¹) of sea level trends between IAS2024 and CMEMS datasets is relatively small (Fig. 13a), while that for the ESA CCI v2.4 dataset is larger (1.32 ± 2.40 mm yr⁻¹, Fig. 10a).

400 **Table 4.** Same as Table 3 but for the comparison between IAS2024 and CMEMS L3 along-track datasets.

Altimeter datasets	Number of collocated stations	Correlation coefficient with tide gauges	RMS (mm)	Trend difference (mm yr ⁻¹)
IAS2024	438	0.66	45.50	0.31±3.95
CMEMS L3	413	0.64	45.20	0.36±3.72

4 Applications of coastal altimeter datasets

The coastal altimeter datasets are an important complement to the existing tide gauge networks considering their higher spatial coverage (Fig. 4). As we can see from the graph, the altimeter-based virtual stations are distributed along the world's coastlines, while the tide gauges, which have continuous data records over the period of 2002 and 2022, are mostly situated in Japan, Australia, North America and Western Europe. Moreover, the altimeter measures the absolute sea levels relative to the ITRF, which would not be affected by the VLM and thus would help us investigate the spatial variation of sea level changes from the offshore oceans to the coast.

In the following sections, we focus on three applications of coastal altimeter datasets. The first is to build altimeter-based virtual stations along the world's coastlines, which can be used to monitor the coastal sea level changes where tide gauges are unavailable. The second is to investigate whether there exist significant trend differences between offshore and nearshore oceans in the last 20 km to the coast. The former is of paramount importance for coastal management and risk adaptation (Fox-Kemper et al., 2021), while the latter would give us insights into the role of small-scale processes in changing the coastal sea level trends (Woodworth et al., 2019; Harvey et al., 2021; Cazenave et al., 2022). The last is to calculate the VLM estimates by combining virtual stations and tide gauges, which provides a way to validate the VLM estimates from GNSS stations (Wöppelmann and Marcos, 2016).

4.1 Altimeter-based virtual stations

To better monitor the coastal sea levels, it is expected that the virtual stations are located as close to the coast as possible. However, the precision of altimeter data degrades with decreasing offshore distance to the coast (Fig. 6). Therefore, it is necessary to investigate which coastal strip is preferable for building the virtual stations. Here, we generate three different sets of virtual stations following the methods described in Section 2.2 using the IAS2024 dataset, and refer them as the onshore, nearshore and offshore virtual stations according to their distances to the coast. Note that the onshore, nearshore and offshore stations are generated from the altimeter data within 0-10 km, 5-15 km and 10-20 km distance bands, respectively.

Figure 14 shows the comparison results between onshore virtual stations and tide gauges in terms of correlation coefficients and RMS values. As can be seen from the graph, the high correlation coefficients (>0.6) and low RMS values (<60 mm) are observed along the world's coastlines, which indicates the good quality of the onshore virtual stations. The better results can be achieved by the nearshore and offshore virtual stations in terms of higher number of virtual stations, while retaining the high correlation coefficients and low RMS values (Table 5). This suggest that the coastal sea level signals are highly correlated over the spatial scales of ~ 20 kilometres. Therefore, the correlation coefficient between nearshore and offshore time series can be an important indicator to evaluate the data quality.

It is also noted that the low correlation coefficients and high RMS values are observed at some onshore stations such as the east coast of North America and the west coast of South America (Fig. 14), which can be attributed to three reasons. Firstly, the VLM would affect the comparison between the tide gauge and altimeter at the local scale (Wöppelmann and Marcos, 2016).

Secondly, the collocated stations are usually tens to hundreds of kilometres away, from which they would measure different sea level signals (Vinogradov and Ponte, 2010). Finally, the coastal environments (e.g., bathymetry, sea states and morphology) can be quite different from the nearby offshore regions, which leads to the significant deterioration of range estimates and range/geophysical corrections (e.g., geocentric ocean and loading tide corrections, wet tropospheric correction, sea state bias correction and dynamic atmospheric correction) (Gommenginger et al., 2011; Abele et al., 2023; Peng et al., 2024b).

440




Figure 14: Comparison results in terms of (a) correlation coefficient and (b) root mean square (RMS) of the differences between monthly time series from onshore virtual stations and tide gauges. The onshore virtual stations are generated using the 20-Hz along-track altimeter data within the 0-10 km coastal strip.

445

Compared to the nearshore and offshore stations, the onshore stations are more likely affected by land contamination because they include the altimeter data within 5 km of the coast, leading to the smaller number of virtual stations (1359 vs 1548). However, the observation degradation seems not to be remarkable and the closure of the trend difference between satellite and tide gauge at the global scale is unchanged. This result implies that the altimeter data within 0-5 km coastal strip can be used when the dedicated editing strategy is taken. In addition, the sea level trends obtained from the altimeter data closer to the coastline are more consistent with those from tide gauge records at the global scale. Therefore, we suggest that the 5-15 km coastal strip is the most suitable for building the virtual stations considering the trade-off between the closest distances to the coast and the reliability of the data quality.

Table 5. Comparison results between altimeter-based virtual stations and tide gauges. The correlation coefficients and RMS values are derived from the monthly SLA time series between virtual stations and tide gauges, while the trend estimates are obtained from the de-seasoned SLA time series.

455

Statistics	Onshore	Nearshore	Offshore
Mean distance to the coast	6.6 km	8.6 km	12.9 km
Number of virtual stations	1359	1548	1548
Number of collocated stations	461	488	490
Correlation coefficient	0.80	0.81	0.82

RMS (mm)	51.75	49.79	47.72
Trend difference (mm yr ⁻¹)	0.16±3.97	0.18±3.90	0.20±3.88

4.2 Spatial variations of sea level trends towards the coast

As reported by previous studies (Deng et al., 2011; Benveniste et al., 2020; Harvey et al., 2021), the coastal sea level trends measured by tide gauges differ from altimeter-derived sea level trends offshore. However, it is uncertain whether the trend discrepancy is due to the different ocean dynamic processes measured by the tide gauge and altimeter or the VLM (Cazenave et al., 2022). Therefore, a good way to solve this issue is to explore the spatial variation of altimeter-derived sea level trends from offshore to nearshore in the last 20 km to the coast. For example, Cazenave et al. (2022) compared the coastal trends (within 4-5 km) with the offshore trends (within 15-17 km) and found no significant difference (within ± 2.0 mm yr⁻¹) at 78% of the 756 virtual stations. Note that the degradation of altimeter data within 5 km of the coast would affect the results from the above method. To avoid this problem, a robust linear regression is applied to the along-track trends using the MATLAB function “robustfit” in this study. This regression function is chosen because it can alleviate the impact of outliers on the estimation procedure from our experimental tests. The trend differences between nearshore and offshore are then calculated. If the absolute trend differences are smaller than 2.0 mm yr⁻¹, it is considered that there is no significant discrepancy between nearshore and offshore sea level trends. Otherwise, the increasing (decreasing) trend is detected if the trend difference is larger than 2.0 mm yr⁻¹ (smaller than -2.0 mm yr⁻¹).

Figure 15 presents four examples of 20-Hz along-track sea level trends against distance to the coast for both IAS2024 and ESA CCI v2.4 datasets. As we can see from the graph, the spatial variations of 20-Hz along-track trends show different patterns (i.e., constant, increasing or decreasing) depending on different geolocations. In addition, the smooth variations of 20-Hz along-track trends at the spatial scales of ~ 300 m is observed (Figs. 15a to 15c). Therefore, the abrupt fluctuation of 20-Hz along-track trends near the coast (Fig. 15d) can be attributed to the degradation of altimeter data.




Figure 15: Examples of 20-Hz along-track sea level trends against distance to the coast for both IAS2024 and ESA CCI v2.4 datasets. The subplots (a) to (d) show constant, increasing and decreasing trends, as well as abrupt fluctuation towards the coast. Horizontal dash lines indicate distances to the coast at 5 km (in black) and 10 km (in magenta).

The statistical results of trend differences over global coastal oceans for these two datasets are illustrated in Fig. 16. As demonstrated by the graph, there is no significant difference (within $\pm 2 \text{ mm yr}^{-1}$, i.e., of the maximum level of trend uncertainties) at 97% of the 1548 ground track segments. In the remaining 3%, we observe either an increasing trend (1.5%) or a decreasing trend (1.5%). This result is consistent with that from the ESA CCI v2.4 dataset, where 90% of the 807 ground track segments show no trend difference, and the percentage of decreasing trend (5%) is equal to that of increasing trend (5%). It is also observed that there is no any concentration of coastal trends departing from offshore trends in a particular region (Fig. 17), which means that the increasing or decreasing trends can occur along the world's coastlines.




Figure 16: Trend differences between nearshore and offshore for the IAS2024 and ESA CCI v2.4 datasets.





Figure 17: Trend differences between nearshore and offshore in the last 20 km to the coast for the IAS2024 (a) and ESA CCI v2.4 (b) datasets. Triangles and inverse triangles correspond to increasing and decreasing trends, respectively.

The trend differences in most places are insignificant, which may be ascribed to two aspects. Firstly, most of the trend uncertainties are relatively large ($\sim 0.5\text{--}2.0 \text{ mm yr}^{-1}$), which is insensitive to the small changes of sea level trends. Secondly, the small-scale processes (e.g., coastal currents, winds, waves, freshwater input in river estuaries) induced variations do not significantly affect the coastal sea level trends, because the sea level variation over longer time scales is more likely affected by the remote signals from open oceans (Han et al., 2019). It is also noted that the trend differences are remarkable in some places, which motivates us to investigate the reasons behind the trend variation. For instance, a recent study by Piccuch et al., (2018a) has demonstrated the role of river discharge in affecting the coastal sea levels over interannual or longer time scales. Moreover, the high-resolution ocean models (grid mesh smaller than 1 km) would contribute to the systematic quantification of coastal phenomena causing the reported trend to increase or decrease near the coast (Marti et al., 2019; Gouzenes et al., 2020; Dieng et al., 2021). However, the high-resolution ocean models are currently not available over global coastal oceans (Han et al., 2019; Melet et al., 2020; Laignel et al., 2023). Therefore, more efforts should be made in the future to improve our understanding of present-day coastal sea level changes and to improve the ability of climate models to simulate future sea levels in highly populated and vulnerable coastal regions of the world.

510 **4.3 Vertical land motion**

The combination of satellite radar altimetry with tide gauge data can be used to estimate the VLM (e.g., Wöppelmann and Marcos, 2016). Figure 18 shows the sea level trends at virtual stations and tide gauges, respectively. As we can see from the graph, the remarkable differences between the altimeter and tide gauge are concentrated in the coasts of the **east United States of America** and the northern Europe. This is because the northern Europe suffers from large uplift rates due to the Glacial Isostatic Adjustment (GIA), while **the U.S. east coast** is exposed to large coastal subsidence rates due to the subsurface fluid withdrawal **and the response to GIA** (Wöppelmann and Marcos, 2016; Piecuch et al., 2018b; Harvey et al., 2021).



The VLM (Fig. 19) is derived in this study from the difference of the de-seasoned monthly SLA time series between altimeter and tide gauge (i.e., ALT–TG) followed the method described in Section 2.3. As can be seen, the large uplift rates ($>2 \text{ mm yr}^{-1}$) are observed in the northern Europe and northwest coast of Canada, suggesting that the GIA-related radial crust displacement are significant in these regions. In addition, the upper lift rates are also observed along the southwest coast of South America. The large subsidence rates ($<-2 \text{ mm yr}^{-1}$), however, are presented along the coast of the northern Gulf of Mexico where the groundwater extraction is significant (Wöppelmann and Marcos, 2016; Harvey et al., 2021). Along the Australian coastline, both the uplift and subsidence rates are observed but dominated by subsidence rates along the northeast coast. When it comes to the western Europe and Japan, the VLM estimates show large variations dependent on different locations, with the overall median values being close to zero for different sets of virtual stations.




Figure 19: VLM estimates derived from the combination of virtual stations and tide gauges. The subplot (a) shows the results of onshore virtual stations, while the subplot (b) shows the boxplot of the VLM estimates from onshore, nearshore and offshore virtual stations. On each box, the central mark indicates the median, and the bottom and top edges of the box indicate the 25th and 75th percentiles, respectively. The whiskers present the most extreme data points outside the percentiles, while the outliers are plotted individually using the '+' symbol.

Figure 19b illustrates that the VLM estimates from onshore virtual stations are in accordance with those from nearshore and offshore stations with median values being close to zero. This result further demonstrates that nearshore and offshore trends in the last 20 km to the coast are identical in most cases, which is consistent with the results shown in Section 4.1. To assess the robustness of the results, the VLM estimates from GNSS observations are used for comparison. The PSMSL tide gauges and the nearest ULR7a GNSS stations are collocated if their distances are within 10 km. As a result, the number of collocated stations is 166 and the comparison results are shown in Fig. 20.




Figure 20: Comparison between ULR7a-derived VLM estimates from GNSS observations and IAS2024-based virtual stations. The subplots (a) to (c) show the scatterplot between IAS2024 onshore virtual stations and GNSS stations, the uncertainties of VLM from IAS2024 and GNSS, the boxplot of the difference of VLM for onshore, nearshore and offshore stations, respectively. In (a), the horizontal and vertical error bars represent the uncertainties of GNSS-derived and altimeter-derived VLM, respectively.

As can be seen, the VLM uncertainties from ULR7a are mostly within the range of 0 mm yr⁻¹ and 0.5 mm yr⁻¹, while the corresponding uncertainties from the combination of tide gauge and altimeter vary between 0.5 mm yr⁻¹ and 2 mm yr⁻¹ (Figs. 20a and 20b). The uncertainties from the combination of tide gauge and altimeter are slightly larger, which can be attributed

to the different spatial and temporal sampling manners between altimeter and tide gauge datasets, different geographical locations between altimeter-based virtual stations and tide gauges, as well as the short time span of the time series. Fig. 20c shows that the VLM estimates from the combination of altimeter and tide gauge agree well with those from GNSS observations in most cases with the differences being smaller than $\pm 1.5 \text{ mm yr}^{-1}$. The mean of the differences is 0.12 mm yr^{-1} with an STD of 2.27 mm yr^{-1} for onshore virtual stations, while the similar values are $0.14 (0.16) \text{ mm yr}^{-1}$ and $2.16 (2.16) \text{ mm yr}^{-1}$ for nearshore (offshore) virtual stations. Therefore, the results from altimeter-based virtual stations can be used as an independent source to validate the VLM estimates from GNSS observations.

5 Data availability

The IAS2024 coastal sea level dataset is available for open access at <https://doi.org/10.5281/zenodo.13208305> as network Common Data Form (netCDF) files (Peng et al., 2024c). This dataset includes monthly SLA time series at 20-Hz along-track points for different ground track segments and at onshore, nearshore and offshore virtual stations. Moreover, the corresponding SLA time series at 10-day cycle time scales. The FES2014 geocentric ocean and loading tide corrections, the MOG2D dynamic atmospheric correction, and the CLS22 MSS over the same time scales are also provided for studying oceanography and geodesy.

6 Conclusions

A new reprocessed 20-Hz coastal sea level dataset, namely IAS2024, for monitoring sea level changes along the world's coastlines has been presented, along with the first evaluations. The Seamless Combination of Multiple Retracker (SCMR) processing strategy is adopted to generate the coastal 20-Hz SLA estimates from Jason missions over the period of January 2002 and April 2022. The main conclusions of this study are summarized as follows.

The SCMR strategy has significantly increased the data availability when compared to the official SGDR MLE4 retracker, with the improvement percentage varying between 12.6% and 53.2% depending on different coastal strips. The moderate improvement (4.8%-10.1%) of data precision brought by the SCMR strategy is also observed, mainly because the SCMR can reduce the variability of the SLA spectrum below the wavelength of 50 km. As a result, the data availability of SCMR-reprocessed Jason data can retain more than 90% beyond 5 km offshore and more than 70%-80% onshore within 5 km to the coast.

In addition, the precision of 20-Hz SLA estimates can be retained at centimetre levels (5-9 cm) over the 5-20 km distance band and at decimetre levels (~20-23 cm) towards the coastline. These results suggest that the IAS2024 dataset generated with the SCMR strategy has the potential to provide reliable SLA estimates for monitoring the coastal sea level changes.

The performance of the IAS2024 dataset has been evaluated and validated through comparisons of the monthly SLA time series, as well as the sea level trends, with those from the PSMSL tide gauge records and independent altimeter datasets (i.e., ESA CCI v2.4 20-Hz and CMEMS L3 1-Hz along-track sea level datasets). The good consistency between different altimeter

590 datasets is observed over global coastal oceans in terms of high correlation coefficients (>0.4) and low RMS values (40 mm-60 mm). The sea level trends from the IAS2024 dataset are on average $1.32 \pm 2.40 \text{ mm yr}^{-1}$ higher than those from the ESA CCI v2.4 dataset, and are similar to those from CMEMS L3 product ($-0.18 \pm 2.17 \text{ mm yr}^{-1}$). These may be attributed to the different data processing techniques adopted, especially the methods used to estimate the inter-mission biases. The validation against tide gauges demonstrates that the IAS2024 and CMEMS datasets achieve the closure of trend differences ($0.16 \pm 3.97 \text{ mm yr}^{-1}$ and $0.36 \pm 3.72 \text{ mm yr}^{-1}$) at the global scale, which are very close to the theoretical value of zero, demonstrating the good performance of the IAS2024 dataset in monitoring the coastal sea levels. In contrast, a negative bias ($-1.50 \pm 3.31 \text{ mm yr}^{-1}$) is found for the ESA CCI v2.4 dataset.

595 Three applications of the IAS2024 dataset have been conducted in this study, which would give us insights into how dedicated coastal sea level datasets can be used for ocean communities and policymakers. Firstly, altimeter-based virtual stations are built along the world's coastline over 0-10 km (1359), 5-15 km (1548), 10-20 km (1548) coastal strips, which are denoted as onshore, nearshore and offshore virtual stations, respectively. The results show that virtual stations achieve much higher spatial coverage than the PSMSL tide gauges (1548 vs 549), which would contribute to the analysis of coastal sea level changes in places where tide gauge data are unavailable. It is also found that the onshore stations would be affected by the degradation of altimeter data within 5 km of the coast, leading a decrease of the number of virtual stations from 1548 to 1359.

600 Secondly, the spatial variations of the linear sea level trends are evaluated with both the IAS2024 and ESA CCI v2.4 datasets. The results from these two datasets show a similar pattern but with different magnitudes at the global scale, with the constant trend being dominant (97% for IAS2024 and 90% for ESA CCI v2.4). This result indicates that the local processes have little impact on coastal sea level changes over longer time scales, which is consistent with the findings revealed by Cazenave et al., (2022). Besides, the reasons leading to the increasing or decreasing trends found in some places should be further investigated with the use of high-resolution ocean models, which would be our future work. Finally, the VLM estimates derived from the combination of altimeter and tide gauge are consistent with those from the GNSS stations, with mean of differences of VLMs being $0.12 \pm 2.27 \text{ mm yr}^{-1}$ for onshore stations. Therefore, the altimeter-derived VLM estimates can be used as an independent data source for validating the GNSS solutions.

610 **Author contributions**

FKP conducted conceptualization, investigation, data curation, developed methodology to process the altimeter data using MATLAB software, validated and visualized the results, and funding acquisition. XLD helped with the formal analysis of the methods and results. YZS and XC provided the resources. All authors contributed to the manuscript writing.

615 **Competing interests**

The contact author has declared that none of the authors has any competing interests.

Disclaimer

Publisher’s note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

620 **Acknowledgments**

The authors are thankful to the space agencies for providing the high-quality altimeter data from Jason missions to users.

Financial support

This research is supported partially by the National Natural Science Foundation of China (Grant No. 42474022 and 42106175), the Natural Science Foundation of Guangdong Province (Grant No. 2022A1515011299), and the Fundamental Research Funds
625 for the Central Universities (Grant No. 02502150056). This research is also supported partially by the Australian Research Council's Discovery Projects funding scheme (Project No. DP220102969).

Review statement

References

630 Abele, A., Royston, S., and Bamber, J.: The impact of altimetry corrections of Sentinel-3A sea surface height in the coastal zone of the Northwest Atlantic, Remote Sensing, 15, 1132, <https://doi.org/10.3390/rs15041132>, 2023.

Ablain, M., Legeais, J. F., Prandi, P., Marcos, M., Fenoglio-Marc, L., Dieng, H. B., Benveniste, J., and Cazenave, A.: Satellite altimetry-based sea level at global and regional scales, Surveys in Geophysics, 38, 7–31, <https://doi.org/10.1007/s10712-016-9389-8>, 2017.

635 Akaike, H.: Information theory and an extension of the maximum likelihood principle, in: Breakthroughs in Statistics, edited by: Kotz, S., and Johnson, N. L., Springer, New York, 610–624, https://doi.org/10.1007/978-1-4612-0919-5_38, 1992.

Altamimi, Z., Rebischung, P., Metivier, L., Collilieux X.: ITRF2014: A new release of the International Terrestrial Reference Frame modeling nonlinear station motions, Journal of Geophysical Research-Solid Earth, 121, 6109–6131, <https://doi.org/10.1002/2016JB013098>, 2016.

- 640 Amarouche, L., Thibaut, P., Zanife, O.Z., Dumont, J.P., Vincent, P., Steunou, N.: Improving the Jason-1 ground retracking to better account for attitude effects, *Marine Geodesy*, 27, 171–197, <https://doi.org/10.1080/01490410490465210>, 2004.
- Benveniste, J., Birol, F., Calafat, F., Anny, C., Dieng, H., Gouzenes, Y., Legeais, J. F., Leger, F., Nino, F., Passaro, M., Schwatke, C., and Shaw, A.: Coastal sea level anomalies and associated trends from Jason satellite altimetry over 2002–2018, *Scientific Data*, 7, 357, <https://doi.org/10.1038/s41597-020-00694-w>, 2020.
- 645 Biancamaria, S., Schaedele, T., Blumstein, D., Frappart, F., Boy, F., Desjonqueres, J. D., Pottier, C., Blarel, F., and Nino, F.: Validation of Jason-3 tracking modes over French rivers, *Remote Sensing of Environment*, 209, 77–89, <https://doi.org/10.1016/j.rse.2018.02.037>, 2018.
- Carrère, L., and Lyard, F.: Modeling the barotropic response of the global ocean to atmospheric wind and pressure forcing-comparisons with observations, *Geophysical Research Letters*, 30, 1275, <https://doi.org/10.1029/2002GL016473>, 2003.
- 650 Cartwright, D. E., and Tayler, R. J.: New computations of the tide-generating potential, *Geophysical Journal International*, 23, 1, 45–73, <https://doi.org/10.1016/j.jog.2004.07.004>, 1971.
- Cazenave, A., Gouzenes, Y., Birol, F., Leger, F., Passaro, M., Calafat, F. M., Shaw, A., Nino, F., Legeais, J. F., Oelsmann, J., Restano, M., and Benveniste, J.: Sea level along the world’s coastlines can be measured by a network of virtual altimetry stations, *Communication Earth & Environment*, 3, 117, <https://doi.org/10.1038/s43247-022-00448-z>, 2022.
- 655 Cipollini, P., Calafat, F. M., Jevrejeva, S., Melet, A., and Prandi, P.: Monitoring sea level in the coastal zone with satellite altimetry and tide gauges, *Surveys in Geophysics*, 38, 33–57, <https://doi.org/10.1007/s10712-016-9392-0>, 2017.
- Davis, C. H.: Growth of the Greenland ice sheet: A performance assessment of altimeter retracking algorithms, *IEEE Transactions on Geoscience and Remote Sensing*, 33, 1108–1116, <https://doi.org/10.1109/36.469474>, 1995.
- Deng, X., Griffin, D. A., Ridgway, K., Church, J. A., Featherstone, W. E., White, N. J., and Cahill, M.: Satellite altimetry for geodetic, oceanographic, and climate studies in the Australian region, in: *Coastal altimetry*, edited by: Vignudelli, S., Cipollini, P., Kostianoy, A. G., and Benveniste, J., Springer, Berlin, Germany, 473–508, https://doi.org/10.1007/978-3-642-12796-0_18, 2011.
- 660 Desai, S., Wahr, J., and Beckley, B.: Revisiting the pole tide for and from satellite altimetry, *Journal of Geodesy*, 89, 1233–1243, <https://doi.org/10.1007/s00190-015-0848-7>, 2015.
- 665 Dieng, H. B., Cazenave, A., Gouzenes, Y., and Sow, B. A.: Trends and interannual variability of altimetry-based coastal sea level in the Mediterranean Sea: Comparison with tide gauges and models, *Advances in Space Research*, 68, 3279–3290, <https://doi.org/10.1016/j.asr.2021.06.022>, 2021.
- Dijkstra, E. W. E. W.: A note on two problems in connexion with graphs, *Numerische Mathematik*, 1, 269–271, <https://doi.org/10.1007/BF01386390>, 1959.
- 670 Fernandes, M. J., and Lázaro, C.: GPD plus wet tropospheric corrections for CryoSat-2 and GFO altimetry missions, *Remote Sensing*, 8, 851, <https://doi.org/10.3390/rs8100851>, 2016.
- Fox-Kemper, B., Hewitt, H. T., Xiao, C., Aðalgeirsdóttir, G., Drijfhout, S. S., Edwards, T. L., Golledge, N. R., Hemer, M., Kopp, R. E., Krinner, G., Mix, A., Notz, D., Nowicki, S., Nurhati, I. S., Ruiz, L., Sallée, J. -B., Slangen, A. B. A., and Yu

- Y.: Ocean, cryosphere and sea level change, In *Climate change 2021: The physical science basis*, edited by: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou B., Cambridge University Press, Cambridge, United Kingdom and New York, 1211–1362, <https://doi.org/10.1017/9781009157896.011>, 2021.
- Gaspar, P., Ogor, F., Letraon, P.Y., and Zanife, O.Z.: Estimating the sea state bias of the Topex and Poseidon altimeters from crossover differences, *Journal of Geophysical Research-Oceans*, 99, 24981–24994, <https://doi.org/10.1029/94JC01430>, 1994.
- Gommenginger, C., Thibaut, P., Fenoglio-Marc, L., Quartly, G., Deng, X., Gomez-Enri, J., Challenor, P., and Gao, Y.: Re-tracking altimeter waveforms near the coasts, In *Coastal altimetry*, edited by: Vignudelli, S., Cipollini, P., Kostianoy, A. G., and Benveniste, J., Springer, Berlin, Germany, 61–101, https://doi.org/10.1007/978-3-642-12796-0_4, 2011.
- Gouzenes, Y., Leger, F., Cazenave, A., Birol, F., Bonnefond, P., Passaro, M., Nino, F., Almar, R., Laurain, O., Schwatke, C., Legeais, J. F., and Benveniste, J.: Coastal sea level rise at Senetosa (Corsica) during the Jason altimetry missions, *Ocean Science*, 16, 1165–1182, <https://doi.org/10.5194/os-16-1165-2020>, 2020.
- Gravelle, M., Wöppelmann, G., Gobron, K., Altamimi, Z., Guichard, M., Herring, T., and Rebischung, P.: The ULR-repro3 GPS data reanalysis and its estimates of vertical land motion at tide gauges for sea level science, *Earth System Science Data*, 15, 497–509, <https://doi.org/10.5194/essd-15-497-2023>, 2023.
- Guérou, A., Meyssignac, B., Prandi, P., Ablain, M., Ribes, A., and Bignalet-Cazalet, F.: Current observed global mean sea level rise and acceleration estimated from satellite altimetry and the associated measurement uncertainty, *Ocean Science*, 19, 431–451, <https://doi.org/10.5194/os-19-431-2023>, 2023.
- Halimi, A., Mailhes, C., Tournet, J. Y., Thibaut, P., and Boy, F.: Parameter estimation for peaky altimetric waveforms, *IEEE Transactions on Geoscience Remote Sensing*, 51, 1568–1577, <https://doi.org/10.1109/TGRS.2012.2205697>, 2013.
- Han, W. Q., Stammer, D., Thompson, P., Ezer, T., Palanisamy, H., Zhang, X. B., Domingues, C. M., Zhang, L., and Yuan, D. L.: Impacts of basin-scale climate modes on coastal sea level: A review, *Surveys in Geophysics*, 40, 1493–1541, <https://doi.org/10.1007/s10712-019-09562-8>, 2019.
- Harvey, T. C., Hamlington, B. D., Frederikse, T., Nerem, R. S., Piecuch, C. G., Hammond, W. C., Blewitt, G., Thompson, P. R., Bekaert, D. P. S., Landerer, F. W., Reager, J. T., Kopp, R. E., Chandanpurkar, H., Fenty, I., Trossman, D., Walker, J. S., and Boening, C.: Ocean mass, steric dynamic effects, and vertical land motion largely explain US coast relative sea level rise, *Communications Earth & Environment*, 2, 233, <https://doi.org/10.1038/s43247-021-00300-w>, 2021.
- Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., Thépaut, J.-N.: ERA5 hourly data on single levels from 1940 to present. Copernicus Climate Change Service (C3S) Climate Data Store (CDS), <https://doi.org/10.24381/cds.adbb2d47>, 2023.

Holgate, S. J., Matthews, A., Woodworth, P. L., Rickards, L. J., Tamisiea, M. E., Bradshaw, E., Foden, P. R., Gordon, K. M., Jevrejeva, S., Pugh, J.: New data systems and products at the Permanent Service for Mean Sea Level, *Journal of Coastal Research*, 29, 493–504, <https://doi.org/10.2112/JCOASTRES-D-12-00175.1>, 2013.

- 710 Komjathy, A., and Born, C. H.: GPS-based ionospheric corrections for single frequency radar altimetry, *Journal of Atmospheric and Solar-Terrestrial Physics*, 61, 1197–1203, [https://doi.org/10.1016/S1364-6826\(99\)00051-6](https://doi.org/10.1016/S1364-6826(99)00051-6), 1999.
- Laignel, B., Vignudelli, S., Almar, R., Becker, M., Bentamy, A., Benveniste, J., Birol, F., Frappart, F., Idier, D., Salameh, E., Passaro, M., Menende, M., Simard, M., Turki, E. I., and Verpoorter, C.: Observation of the coastal areas, estuaries and deltas from space, *Surveys in Geophysics*, 44, 1309–1356, <https://doi.org/10.1007/s10712-022-09757-6>, 2023.
- 715 Lyard, F. H., Allain, D. J., Cancet, M., Carrère, L., and Picot, N.: FES2014 global ocean tide atlas: design and performance, *Ocean Science*, 17, 615–649, <https://doi.org/10.5194/os-17-615-2021>, 2021.
- Marti, F., Cazenave, A., Birol, F., Passaro, M., Léger, F., Niño, F., Almar, R., Benveniste, J., and Legeais, J. F.: Altimetry-based sea level trends along the coasts of Western Africa, *Advances in Space Research*, 68, 504–522, <https://doi.org/10.1016/j.asr.2019.05.033>, 2021.
- 720 Melet, A., Teatini, P., Le Cozannet, G., Jamet, C., Conversi, A., Benveniste, J., and Almar, R.: Earth observations for monitoring marine coastal hazards and their drivers, *Surveys in Geophysics*, 41, 1489–1539, <https://doi.org/10.1007/s10712-020-09594-5>, 2020.
- Passaro, M., Cipollini, P., Vignudelli, S., Quartly, G. D., and Snaith, H. M.: ALES: A multi-mission adaptive subwaveform retracker for coastal and open ocean altimetry, *Remote Sensing of Environment*, 145, 173–189, <https://doi.org/10.1016/j.rse.2014.02.008>, 2014.
- 725 Peng, F. K., and Deng, X. L.: A new retracking technique for Brown peaky altimetric waveforms, *Marine Geodesy*, 41, 99–125, <https://doi.org/10.1080/01490419.2017.1381656>, 2018.
- Peng, F. K., and Deng, X. L.: Validation of Sentinel-3A SAR mode sea level anomalies around the Australian coastal region, *Remote Sensing of Environment*, 237, 111548, <https://doi.org/10.1016/j.rse.2019.111548>, 2020a.
- 730 Peng, F. K., and Deng, X. L.: Improving precision of high-rate altimeter sea level anomalies by removing the sea state bias and intra-1-Hz covariant error, *Remote Sensing of Environment*, 251, 112081, <https://doi.org/10.1016/j.rse.2020.112081>, 2020b.
- Peng, F. K., Deng, X. L., and Cheng, X.: Quantifying the precision of retracked Jason-2 sea level data in the 0-5 km Australian coastal zone, *Remote Sensing of Environment*, 263, 112539, <https://doi.org/10.1016/j.rse.2021.112539>, 2021.
- 735 Peng, F. K., Deng, X. L., and Cheng, X.: Australian coastal sea level trends over 16 yr of reprocessed Jason altimeter 20-Hz data sets, *Journal of Geophysical Research-Oceans*, 127, e2021JC018145, <https://doi.org/10.1029/2021JC018145>, 2022.
- Peng, F. K., Deng, X. L., Jiang, M. F., Dinardo, S., and Shen, Y. Z.: A new method to combine coastal sea surface height estimates from multiple retrackers by using the Dijkstra algorithm, *Remote Sensing*, 15, 2329, <https://doi.org/10.3390/rs15092329>, 2023.

- Peng, F. K., Deng, X. L., and Shen, Y. Z.: Analyzing the coastal sea level trends from SCMR-reprocessed altimeter data: A case study in the northern South China Sea, *Advances in Space Research*, <https://doi.org/10.1016/j.asr.2024.06.036>, 2024a.
- Peng, F. K., Deng, X. L., and Shen, Y. Z.: Assessment of Sentinel-6 SAR mode and reprocessed Jason-3 sea level measurements over global coastal oceans, *Remote Sensing of Environment*, 311, 114287, <https://doi.org/10.1016/j.rse.2024.114287>, 2024b.
- Peng, F. K., Deng, X. L., Shen, Y. Z., and Cheng, X.: The IAS2024 coastal sea level dataset and first evaluations, Zenodo [data set], <https://doi.org/10.5281/zenodo.13208305>, 2024c.
- Piecuch, C. G., Bittermann, K., Kemp, A. C., Ponte, R. M., Little, C. M., Engelhart, S. E., and Lentz, S. J.: River-discharge effects on United States Atlantic and Gulf coast sea level changes, *Proceedings of the National Academy of Sciences of the United States of America*, 115, 7729–7734, <https://doi.org/10.1073/pnas.1805428115>, 2018a.
- Piecuch, C. G., Huybers, P., Hay, C. C., Kemp, A. C., Little, C. M., Mitrovica, J. X., Ponte, R. M. and Tingley, M. P.: Origin of spatial variation in US East Coast sea-level trends during 1900–2017, *Nature*, 564, 400–404, <https://doi.org/10.1038/s41586-018-0787-6>, 2018b.
- Poisson, J. C., Quartly, G. D., Kurekin, A. A., Thibaut, P., Hoang, D., and Nencioli, F.: Development of an Envisat altimetry processor providing sea level continuity between open ocean and Arctic leads, *IEEE Transactions on Geoscience and Remote Sensing*, 56, 5299–5319, <https://doi.org/10.1109/TGRS.2018.2813061>, 2018.
- Roscher, R., Uebbing, B., and Kusche, J.: STAR: Spatio-temporal altimeter waveform retracking using sparse representation and conditional random fields, *Remote Sensing of Environment*, 201, 148–164, <https://doi.org/10.1016/j.rse.2017.07.024>, 2017.
- Prandi, P., Meyssignac, B., Ablain, M., Spada, G., Ribes, A., and Benveniste, J.: Local sea level trends, accelerations and uncertainties over 1993–2019, *Scientific Data*, 8, 1, <https://doi.org/10.1038/s41597-020-00786-7>, 2021.
- Schaeffer, P., Pujol, M. I., Veillard, P., Faugere, Y., Dagneaux, Q., Dibarboure, G., Picot, N.: The CNES CLS 2022 mean sea surface: short wavelength improvements from CryoSat-2 and SARAL/Altika high-sampled altimeter data, 15, 1–20, <https://doi.org/10.3390/rs15112910>, 2023.
- Schwarz, G.: Estimating the dimension of a Model, *The Annals of Statistics*, 6, 461–464, <https://doi.org/10.1214/aos/1176344136>, 1978.
- Vignudelli, S., Birol, F., Benveniste, J., Fu, L. L., Picot, N., Raynal, M., and Roinard, H.: Satellite altimetry measurements of sea level in the coastal zone, *Surveys in Geophysics*, 40, 1319–1349, <https://doi.org/10.1007/s10712-019-09569-1>, 2019.
- Vinogradov, S. V., and Ponte, R. M.: Annual cycle in coastal sea level from tide gauges and altimetry, *Journal of Geophysical Research-Oceans*, 115, C04021, <https://doi.org/10.1029/2009JC005767>, 2010.
- Wang, X. F., Ichikawa, K., and Wei, D. N.: Coastal waveform retracking in the slick-rich Sulawesi Sea of Indonesia, based on variable footprint size with homogeneous sea surface roughness, *Remote Sensing*, 11, 1274, <https://doi.org/10.3390/rs11111274>, 2019.

- 775 Watson, C. S., White, N. J., Church, J. A., Burgette, R. J., Tregoning, P., and Coleman, R.: Absolute calibration in Bass Strait, Australia: Topex, Jason-1 and OSTM/Jason-2, Marine Geodesy, 34, 242–260, <https://doi.org/10.1080/01490419.2011.584834>, 2011.
- Watson, C. S., White, N. J., Church, J. A., King, M. A., Burgette, R. J., and Legresy, B.: Unabated global mean sea level rise over the satellite altimeter era, Nature Climate Change, 5, 565, <https://doi.org/10.1038/nclimate2635>, 2015.
- 780 Woodworth, P. L., Melet, A., Marcos, M., Ray, R. D., Wöppelmann, G., Sasaki, Y. N., Cirano, M., Hibbert, A., Huthnance, J. M., Monserrat, S., and Merrifield, M. A., Surveys in Geophysics, 40, 1351–1397, <https://doi.org/10.1007/s10712-019-09531-1>, 2019.
- Wöppelmann, G. and Marcos, M. Vertical land motion as a key to understanding sea level change and variability, Reviews of Geophysics, 54, 64–92, <https://doi.org/10.1002/2015RG000502>, 2016.