

IAPv4 ocean temperature and ocean heat content gridded dataset

Lijing Cheng^{1,11*}, Yuying Pan^{1,11}, Zhetao Tan^{1,11}, Huayi Zheng^{1,11}, Yujing Zhu^{1,11}, Wangxu Wei^{1,11}, Juan Du¹, Huifeng Yuan^{2,11}, Guancheng Li³, Hanlin Ye¹, Viktor Gouretski¹, Yuanlong Li^{4,11}, Kevin E. Trenberth^{5,6}, John Abraham⁷, Yuchun Jin^{4,11}, Franco Reseghetti⁸, Xiaopei Lin⁹, Bing Zhang^{4,11}, Gengxin Chen^{10,11}, Michael E. Mann¹², Jiang Zhu^{1,11}

¹ Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China, 100029.

² Computer Network Information Center, Chinese Academy of Sciences, Beijing, 100083.

³ Eco-Environmental Monitoring and Research Center, Pearl River Valley and South China Sea Ecology and Environment Administration, Ministry of Ecology and Environment, PRC, Guangzhou 510611, China.

⁴ Institute of Oceanography, Chinese Academy of Sciences, Qingdao, China.

⁵ National Center for Atmospheric Research, PO Box 3000, Boulder, CO 80307, USA.

⁶ University of Auckland, Auckland, New Zealand.

⁷ University of St. Thomas, School of Engineering, 2115 Summit Ave., St Paul, MN 55105, USA.

⁸ Istituto Nazionale di Geofisica e Vulcanologia, 40127, Bologna, Italy.

⁹ Frontier Science Center for Deep Ocean Multispheres and Earth System and Physical Oceanography Laboratory, Ocean University of China, Qingdao, China.

¹⁰ State Key Laboratory of Tropical Oceanography, South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou, China

¹¹ University of Chinese Academy of Sciences, Beijing, China.

¹² Dept. of Earth and Environmental Science, University of Pennsylvania, Philadelphia PA, USA

Correspondence to: Lijing Cheng (chenglij@mail.iap.ac.cn)

29 **Abstract.** Ocean observational gridded products are vital for climate monitoring, ocean
30 and climate research, model evaluation, and supporting climate mitigation and adaptation
31 measures. This paper describes the 4th version of the Institute of Atmospheric Physics
32 (IAPv4) ocean temperature and ocean heat content (OHC) objective analysis product. It
33 accounts for recent developments in quality control (QC) procedures, climatology, bias
34 correction, vertical and horizontal interpolation, and mapping and is available for the upper
35 6000 m (119 levels) since 1940 (more reliable after ~1957) for monthly and $1^\circ \times 1^\circ$
36 temporal and spatial resolutions. The IAPv4 is compared with the previous version, IAPv3,
37 and to the other data products, sea surface temperatures (SSTs), and satellite observations.
38 It has a slightly stronger long-term upper 2000 m OHC increase than IAPv3 for 1955-
39 2023, mainly because of newly developed bias corrections. IAPv4 OHC 0-2000 m trend is
40 also higher during 2005-2023 than IAPv3. The uppermost level of IAPv4 is consistent with
41 independent SST datasets. The month-to-month OHC variability for IAPv4 is desirably
42 less than IAPv3 and other OHC products investigated in this study, the trend of ocean
43 warming rate (i.e., warming acceleration) is more consistent with the net energy imbalance
44 at the top of the atmosphere than IAPv3, and the sea level budget can be closed within
45 uncertainty. The gridded product is freely accessible at:
46 <http://dx.doi.org/10.12157/IOCAS.20240117.002> for temperature data (Cheng et al.,
47 2024a) and <http://dx.doi.org/10.12157/IOCAS.20240117.001> for ocean heat content data
48 (Cheng et al., 2024b).

49

50 **1. Introduction**

51 Observational gridded products are essential for understanding the ocean, the
52 atmosphere, and climate change; they support policy decisions and social-economy
53 developments (Abraham et al., 2022; Abraham and Cheng, 2022; Cheng et al., 2022a). For
54 instance, many of the climate indicators used in the Working Group I report of the 6th
55 Intergovernmental Panel on Climate Change (IPCC-AR6-WG1) are based on gridded
56 products (Gulev et al., 2021; IPCC, 2021), mainly because the raw oceanic data suffer
57 from inhomogeneous data quality and irregular and incomplete data coverage (Abraham et
58 al., 2013; Boyer et al., 2016; Cheng et al., 2022a; Meyssignac et al., 2019).

59 As more than 90% of the Earth's energy imbalance (EEI) in the past half-century has
60 accumulated in the ocean, increasing ocean temperature (T) and ocean heat content (OHC)
61 are essential climate variables for monitoring, understanding, and projecting climate
62 change (e.g., Rhein et al., 2013; Hansen et al., 2011; Trenberth, 2022; Trenberth et al.,
63 2009; von Schuckmann et al., 2020; Cheng et al., 2022). OHC also impacts air-sea and ice-
64 sea interactions and thus exerts a considerable influence over the other components of the
65 climate system. It provides critical feedback through energy, water, and carbon cycles
66 (Cheng et al., 2022a; Trenberth, 2022; von Schuckmann et al., 2016). Substantial changes
67 in ocean temperatures also profoundly impact ocean biogeochemical processes and
68 ecosystems and are critical for ocean health and human society (Bindoff et al., 2019;
69 Cheng et al., 2022a).

70 Many gridded T/OHC datasets have been produced by independent groups, and most
71 of them are updated annually or more frequently (Cheng et al., 2022a; Good et al., 2013;
72 Hosoda et al., 2008; Ishii et al., 2017; Levitus et al., 2012; Li et al., 2017; Meyssignac et
73 al., 2019; Roemmich and Gilson, 2009). Most widely-used products are at $1^\circ \times 1^\circ$
74 horizontal resolution and monthly temporal resolution from near-surface to at least 2000 m
75 depth. Some products utilize all available *in situ* observations and span at least half a
76 century, prominent examples being the data products compiled by the Institute of
77 Atmospheric Physics (IAP) (Cheng and Zhu, 2016; Cheng et al., 2017) from 1940-present;
78 Japan Meteorological Agency (JMA) (Ishii et al., 2017) from 1955-present; National
79 Centers for Environmental Information (NCEI), National Oceanic and Atmospheric
80 Administration (NOAA) from 1950-present (Levitus et al., 2012); and University of
81 California since 1949 (Bagnell and DeVries, 2021). As Argo data has achieved near-global
82 upper 2000 m open ocean coverage since ~2005, many Argo-based or Argo-only gridded
83 products are available. Examples include gridded products from SCRIPPS after 2004
84 (Roemmich and Gilson, 2009); China Argo Real-time Data Center since 2005 (Li et al.,
85 2017); and Copernicus since 2005 (von Schuckmann and Le Traon, 2011). These products
86 usually span from ~2005 to the present for the upper ~2000 m. These data benefit from the
87 high quality of Argo data but are not fully resolving polar regions, shallow waters, and
88 regions with complex topography.

89 In 2016, the IAP group provided its first gridded product for the upper 700 m ocean
90 (Cheng and Zhu, 2016) by merging all available observations since 1960. With a revised
91 mapping method and a thorough evaluation process with synthetic observations, an update

92 (IAP version 3, IAPv3) became available in 2017 for the upper 2000 m ocean with data
93 since the 1950s (Cheng et al., 2017). The IAPv3 has supported scientific research, climate
94 assessment reports, and monitoring practices (Bindoff et al., 2019; Gulev et al., 2021;
95 WMO, 2022).

96 After the release of IAPv3, there has been progress with observation data quality
97 control and new/updated techniques for temperature data processing and reconstruction.
98 For example, Gouretski et al. (2022) found that old Nansen cast bottle data contained
99 systematic biases that impacted the T/OHC data before 1990. Revisions are also available
100 to the bias corrections for the Mechanical Bathythermographs (MBT) and eXpendable
101 Bathythermographs (XBT) data (Cheng et al., 2014; Gouretski and Cheng, 2020), mainly
102 impacting the data within 1940–2005. Tan et al. (2023) developed a new quality-control
103 system that advances the detection of outliers after accounting for the non-Gaussian
104 distribution of local temperatures in determining the local climatological range. The impact
105 of inhomogeneous vertical resolution of temperature profiles has been recognized
106 previously (Cheng and Zhu, 2014) and received more attention recently (Li et al., 2020)
107 with a new vertical interpolation approach (Barker and McDougall, 2020). Upgrading the
108 product with new developments is important to better support the ocean/climate research
109 and climate assessments.

110 This manuscript discusses the revisions to the IAP ocean objective analysis product
111 (IAPv4) since the publication of the IAPv3 (Cheng et al., 2017). The data and methods are
112 introduced in Section 2 and the results are presented in Section 3, with analyses of the
113 character of the IAPv4 on regional and global scales and at various time scales. The EEI
114 and sea level budgets based on the new data product are also investigated. A summary and
115 discussion are provided in Section 4, with some remaining issues and outlooks being
116 discussed.

117

118 **2. Data and Methods**

119 **2.1 Data source**

120 The majority of the *in situ* measurements used to create the data product come from
121 the World Ocean Database (WOD), downloaded in September 2023. Data from all
122 instrument types are used, including XBTs (Goni et al., 2019), Argo (Argo 2000),
123 Conductivity/Temperature/Depth profilers (CTDs), MBTs, bottles, moorings, gliders,
124 Animal Borne Ocean Sensors (McMahon et al., 2021) and others (Boyer et al., 2018) (Fig.


125 1). There is a total of 17,634,865 temperature profiles from January 1940 to September
126 2023 (Fig. 1a). MBT, XBT, Nansen Bottle and CTD data are the major instruments before
127 2000 (Fig. 1a, b). The spatial coverage of these data increased to >30% in 1960 and >70%
128 in the late 1960s for $1^\circ \times 1^\circ \times 1$ -year resolution. After 2005, there is a huge number of
129 GLD and APB data, and as they are mainly distributed in the polar regions (APB) and
130 coastal regions (GLD) (Fig. 1a), their spatial coverage is usually less than 5% for $1^\circ \times 1^\circ \times$
131 1 year resolution. By contrast, the Argo data cover most of the global open ocean since
132 ~2005 (Fig. 1b).

133 Argo data are processed following the recommendations of the Argo community.
134 Adjusted data are used where applicable. Both Delayed- and Real-Time Argo data have
135 been incorporated in IAPv4. As Real-Time Argo data have only passed automated, simple
136 QC tests in real-time, these data may still contain temperature, pressure, and salinity values
137 affected by unknown errors. However, through a sensitivity study, Cheng (2024) indicated
138 that including Real-Time Argo data does not bias the OHC calculation for the IAP
139 analysis. Nevertheless, IAP data are updated frequently (every 1-3 months): each time the
140 updated Argo data is used, the T/OHC fields are recalculated following the
141 recommendation by the Argo group (Wong et al., 2020). The data from the Argo floats in
142 the “grey list” have been removed from the calculation (<https://data-argo.ifremer.fr/>).

143 To complement the WOD with relatively less data in the Arctic and coastal regions of
144 the Northwest Pacific, this presented product also uses data from other sources. The
145 majority of these data are from the Chinese Academy of Sciences Ocean Science Data
146 Center (Zhang et al., 2024), and some data are rescued from the old documents of marine
147 surveys. All these data will be publicly available. There are a total of 85,990 additional
148 temperature profiles, about 0.50% of the data, which is expected to improve the
149 reconstruction in these data-sparse regions (compared with IAPv3 and other products).


150 The *in situ* data have been processed as described in a flow chart in Figure 2. In the
151 following sections, the key techniques of data processing are introduced.

152



154 **Figure 1: (a) Yearly number of temperature casts for different instruments; (b)**
 155 **percentage coverage (%) of ocean data for each instrument, calculated as the ratio**
 156 **between the number of $1^\circ \times 1^\circ \times 1$ year grid cells observed by each instrument and**
 157 **the total number of ocean grids; (c) number of subsurface temperature casts in 1-**
 158 **degree grid boxes from 1940 to 2023 collected by different instruments: CTD**
 159 **(Conductivity/Temperature/Depth), XBT (eXpendable BathyThermographs), MBT**
 160 **(Mechanical BathyThermograph), Bottle, APB (Animal mounted Pinniped Borne), PFL**
 161 **(Profiling Floats, i.e. Argo), GLD (Glider), MRB (Moored Buoy), and DRB (Drifting**
 162 **Buoy).**

163



164

165 **Figure 2: Flow chart of the IAP data reconstruction processes from the raw *in situ***
 166 **observations to gridded data (IAPv4) and OHC estimates.** The ellipses indicate the data
 167 (including data for error estimates), and the rectangle boxes show the techniques used to
 168 process the data.

169

170 2.2 Data quality control

171 The quality control (QC) procedure aims to identify spurious measurements (including
 172 outliers) and data with incorrect metadata through a set of quality checks and ensures the
 173 quality of the *in situ* dataset (Tan et al., 2022). There is growing evidence that QC is
 174 critical for accurate temperature/OHC reconstruction, as shown by Tan et al., (2023) where
 175 two different QC systems produced a difference of approximately 15 % (~7 %) in the OHC
 176 0-2000 m trend from 1955 to 1990 (2005-2021). Unfortunately, the impact of QC on OHC

177 estimates has not been evaluated in previous community-assessments on T/OHC
178 uncertainty (Boyer et al., 2016; Lyman et al., 2010). In this study, the QC procedure
179 follows the CAS-Ocean Data Center (CODC) Quality Control system, named CODC-QC
180 (Tan et al., 2023), where only the “good” data (flag=0) are used.

181 The CODC-QC system (Tan et al., 2023) has the following strengths, which make it
182 particularly suitable for T/OHC reconstruction:

183 1) A new local climatological range is defined in this CODC-QC system to identify
184 the outliers. Unlike many existing QC procedures, no assumption is made of a Gaussian
185 distribution law in the new approach, as the oceanic variables (e.g., temperature and
186 salinity) are typically skewed. Instead, the 0.5 % and 99.5 % quantiles are used as
187 thresholds in CODC-QC to define the local climatological parameter ranges.

188 2) Local climatological ranges change with time to account for the long-term trends of
189 ocean temperature accompanied by more frequent extreme events (e.g., Oliver et al., 2018;
190 Sun et al., 2023). Previously, the use of the static local ranges tended to remove too many
191 “extreme values” (at the tails of the temperature distributions) associated with climate
192 change in recent years that were actually real, leading to a QC-procedure related bias in the
193 gridded dataset and OHC estimate (Tan et al., 2023).


194 3) In addition, local climatological ranges for the vertical temperature gradient are
195 constructed to account for the variability of ‘vertical shape’, increasing the ability of the
196 scheme to identify spurious profiles.

197 4) The QC procedure is instrument-specific, accounting for characteristics inherent to
198 particular instrumentation types. For example, XBT digital recording systems are allowed
199 to continue to record beyond the rated terminal depth suggested by manufacturers (T7/DB
200 probes below 760 m; T4/T6 below 460 m; T5 below 1830 m). Below the rated maximum
201 depth, the XBT wire often breaks, leading to a characteristic change in recorded
202 temperature values. The new QC procedure effectively identifies such profiles.

203 5) The thorough evaluation of the QC procedure performance and the application of
204 the QC procedure to the manually QC-ed datasets (Thresher et al., 2008; Gouretski and
205 Koltermann, 2004) demonstrated the effectiveness of the proposed scheme in removing
206 spurious data and minimizing the percentage of mistakenly flagged good data.

207 Being applied to the entire temperature profile dataset the CODC-QC procedure
208 identifies 6.22 % of all temperature measurements as outliers. The rejection rates
209 (definition follows Tan et al., 2023) vary among instrumentation types (3.73 % for CTD,

210 1.97 % for Argo, 12.06 % for XBT, 4.93 % for MBT, 6.54 % for bottle, 5.92 % for APB,
211 4.54 % for DRB, 2.55 % for MRB). The overall percentage of outliers decreases over time
212 from ~5 % in the 1940s to ~2.5 % in the 2020s, reflecting the progressive improvement of
213 the instrumentation (Fig. 3). A rejection rate maximum (~12 %) during 2000~2010 is
214 linked to the XBT data, which are especially abundant in the 800–1100m layer and are
215 characterized by higher rejection rate below the maximum depth (Tan et al., 2023). The
216 generally higher rejection rate below 4000 meters is related to the gross errors (such as
217 measurements cooler than -2°C, big spikes, etc.) and the occurrence of the constant values
218 (recorded values don't change with depth). For example, the higher rejection rate within
219 2008-2009 below 4000 meters is because of the gross errors in the CTD data.



220
221 **Figure 3: The rejection rate (%) after CODC-QC as a function of calendar year and**
222 **depth.**

224 2.3 Bias correction

225 It is well known that data from several instrument types can exhibit biases both in
226 temperature and depth. Temperature profiles obtained using XBTs and MBTs provide an
227 example of biased data, especially because of uncertainties in the depth of measurement.
228 Gouretski and Koltermann (2007) demonstrated their significant impact on the magnitude
229 and variability of the global OHC estimates. That study triggered a series of publications
230 where different bias correction schemes have been suggested for XBT (Gouretski and

231 Reseghetti, 2010; Abraham et al., 2013; Cheng et al., 2016; Levitus et al., 2009; Wijffels et
232 al., 2008), MBT (Gouretski and Cheng 2020; Levitus et al., 2009) and other instruments
233 (Fig. 2). In the compilation of IAPv4, newly developed bias correction schemes are
234 applied.

235 The XBT temperature bias was found to be generally positive, as large as ~ 0.1 °C
236 before 1980 on the global 0–700 m average, diminishing to less than 0.05 °C after 1990
237 (Gouretski and Koltermann 2007; Wijffels et al., 2008). Here, we use an updated XBT bias
238 correction scheme (Cheng et al., 2014) to correct both depth and temperature biases in
239 XBT data, following the community recommendation (Cheng et al., 2016; Goni et al.,
240 2019). The depth and temperature biases depend on ocean temperature, probe type, and
241 time. An inter-comparison among several correction schemes rated the CH14 scheme the
242 most successful (Cheng et al., 2018). Using XBT and collocated CTD data, we updated the
243 CH14 scheme by re-calculating bias corrections between 1966–2016 and extending them
244 for the years 2017 to 2023.

245 Comparison with collocated reference CTD profiles recently revealed significant
246 biases in the old hydrographic profiles obtained by means of Nansen bottle casts
247 (Gouretski et al., 2022). Both depth and temperature measurements of bottle casts were
248 found to be biased, and the proposed correction scheme was also implemented in IAPv4.
249 The thermal bias is related to the time needed to bring the mercury thermometers in
250 equilibrium with the ambient temperature after the completion of the hydrographic cast.
251 The depth bias indicates an overestimation of the bottle depth due to the wire's deviation
252 from the vertical position and is mostly related to the hydrographic casts where the
253 thermometrical method of sample depth determination was not used. The correction
254 scheme includes a constant thermal bias of -0.02 °C and a depth- and time-variable depth
255 bias.

256 The MBT bias is as large as 0.28 °C before 1980 for the global average and reduces to
257 less than 0.18 °C after 1980 for the 0–200 m average. IAPv3 used Ishii and Kimoto, (2009)
258 (IK09) scheme to correct MBT bias, while a new scheme proposed by Gouretski and
259 Cheng, (2020) (GC20) is adopted in IAPv4. This shift is made because our assessment
260 indicates the under-correction of MBT bias by the IK09 scheme within the upper 120 m
261 and over-correction in the deeper layer, whereas GC20 corrects both depth and temperature
262 biases. GC20 also found the MBT bias to be country-dependent, explained in terms of
263 different instrumentation characteristics and working procedures. Therefore, the time-

264 varying bias corrections are applied separately for the MBT profiles obtained by ships
265 from the United States, Soviet Union/Russia, Japan, Canada, and Great Britain. Data from
266 all other countries are corrected using a globally averaged correction.

267 Finally, thermal biases were recently reported for the data obtained by different kinds
268 of data loggers attached to marine mammals (APB). Gouretski et al. (2024) analysed
269 temperature profiles obtained between 2004 and 2019 in the high and moderate latitudes of
270 both hemispheres. Comparison with the collocated reference CTD and Argo float data
271 revealed a systematic negative thermal offset (average value $-0.027\text{ }^{\circ}\text{C}$) for mammal
272 temperature profiles from SRDL (satellite-related data loggers). For the less accurate data
273 from TDR (Temperature-Depth-Recorders), the comparison revealed a small positive
274 temperature bias of $0.02\text{ }^{\circ}\text{C}$ and the depth (pressure) bias indicating depth overestimation.

275

276 **2.4 Climatology**

277 For IAP and other data product generators, horizontal interpolation (mapping) is
278 applied on a temperature anomaly field after removing a monthly climatology; thus, a pre-
279 defined climatology field with an annual cycle is mandatory (Fig. 2). The accuracy of the
280 climatology field is one of the key sources of uncertainty in reconstruction because the
281 error in climatology will propagate into the anomaly field, impact the spatial dynamical
282 consistency, and the accuracy of the reconstruction (Cheng and Zhu, 2015; Lyman and
283 Johnson, 2014; Boyer et al., 2016).

284 In IAPv4, the adjusted mapping procedure (see below) has been applied to reconstruct
285 the climatology field (Table 1). The merit of using IAP mapping for climatology is its
286 ability to better represent the spatial anisotropy of temperature variability (non-Gaussian
287 distribution). Unlike IAPv3, where the 1990–2005 reference period was used, IAPv4 uses
288 data between 2006 and 2020 to construct 12 monthly climatologies, taking advantage of
289 more reliable data combined with better and more homogeneous spatial and temporal
290 coverage in the last two decades (Table 1). Following the recommendation in Cheng and
291 Zhu, (2015), a relatively short period of 15-year is used because climatology constructed
292 with longer period of data will result in different baselines at different locations (i.e., the
293 baseline shifted to earlier years in the middle latitudes of the North Hemisphere and the
294 baseline shifted to more recent years in the Southern Hemisphere) and this inconsistency
295 will violate the spatial structure of the anomaly field (Cheng and Zhu, 2015). Recent

296 developments from other groups, such as Li et al., (2022), include the choice of a short-
297 period climatology.

298 IAPv4 used an 800 km influencing radii in climatology reconstruction, smaller than
299 the 20° for IAPv3, to more properly account for the rapid change of temperatures with
300 distance. There is a trade-off between data availability and the size of the influence radius.
301 Using radii smaller than 500 km does not ensure a global fractional coverage (defined as
302 the fraction of the total ocean area obtained by the mapping method) because of data
303 sparseness (Cheng, 2024). As our tests suggest, using 500~800 km results in very similar
304 reconstructions of climatology, therefore, 800 km is adopted.

305

306 **2.5 Vertical interpolation**

307 The vertical resolution of ocean temperature profiles changed dramatically over time
308 associated with instrument evolution and the increase of data storage capability. For
309 instance, the global mean vertical resolution at 500 m level changed from ~100 m in the
310 1960s to less than 10 m during the 2010s (Li et al., 2020). Vertical interpolation of the raw
311 profiles on standard levels is a critical process (Fig. 2): Cheng and Zhu (2014) indicated
312 that the use of linear or spline vertical interpolation methods can bias the temperature
313 reconstruction and OHC estimation (Barker and McDougall, 2020; Li et al., 2020; Li et al.,
314 2022). IAPv3 used the (Reiniger and Ross, 1968) (RR) method. Recently, Barker and
315 McDougall (2020) proposed a new approach using multiple Piecewise Cubic Hermite
316 Interpolating Polynomials (PCHIPs) to minimize the formation of unrealistic water masses
317 by the interpolation procedure.

318 Because the largest difference between interpolation methods is found mostly for the
319 low-resolution profiles (e.g., old Nansen casts), in practice, extremely low vertical
320 resolution profiles had to be removed to reduce the uncertainty in interpolation. In IAPv4,
321 this procedure is optimized compared to IAPv3, and only parts of profiles with a sufficient
322 vertical resolution are used. The thresholds for the vertical resolution are set by 50 m in the
323 upper 200m, 200m between 200 m and 1000 m, 500 m between 1000 m and 2000 m, and
324 600m between 2000 m and 6000 m. As no interpolation method can adequately interpolate
325 temperature for the vertical resolution beyond these thresholds, interpolation is not
326 performed in such cases to avoid errors (these extreme low-resolution data are not used in
327 further processing). Under this limitation for IAPv4, we still apply the RR method for
328 temperature profiles.

329 Finally, IAPv4 extends the set of standard vertical levels with a total of 119 levels
330 from 1 m to 6500 m (79 levels within the upper 2000 m) compared to 41 levels in IAPv3
331 between 1 m and 2000 m (Table 1). The increase in vertical resolution is critical for
332 accurately representing the mixed layer, as investigated below.

333

334 **2.6 Grid average and mapping**

335 The anomaly profiles are obtained by subtracting the monthly mean climatology from
336 the vertically interpolated profiles. These anomalies are then averaged (arithmetic mean)
337 into a $1^\circ \times 1^\circ$ grid at each standard level ($1^\circ \times 1^\circ$ gridded average field) (Fig. 2). Due to the
338 general data sparsity, variable time windows (larger than one month) are used for monthly
339 reconstructions to ensure a truly global analysis (Supplementary Table 1). This process
340 takes advantage of the larger persistence of anomalies (generally smaller monthly and
341 inter-annual variability) in the deep ocean than in the upper ocean and thus is physically
342 grounded. Specifically, after 2005, data within a three-month window are merged to
343 provide a monthly reconstruction for each layer of the upper 1950 m. Before 2005, a time-
344 varying and depth-varying time window is used, and it is generally smaller in the upper
345 ocean and wider in the deeper ocean (Supplementary Table 1). Below 2000 m, a 5-year
346 (60-month) window is adopted. The use of a time window will reduce the monthly
347 variance compared to other datasets, which is likely too high compared with independent
348 Earth's Energy Imbalance data at the top of the atmosphere (Trenberth et al., 2016).

349 Mapping interpolates the gridded (e.g., box-averaged) observations horizontally into a
350 spatially complete map (Fig. 2) because not all $1^\circ \times 1^\circ$ boxes are filled with data. (Fig. 2).
351 IAPv4 adopted a similar mapping approach (Ensemble Optimal Interpolation with dynamic
352 ensemble: EnOI-DE) as in IAPv3 introduced in Cheng and Zhu (2016) and Cheng et al.,
353 (2017) but with the following modifications:

354 1) the largest influence radius has changed from 20° in the upper 700 m (25° at 700–
355 2000 m) in IAPv3 to 2,000 km in the upper 700 m (2,500 km at 700–6000 m) in IAPv4, to
356 account for the reduced distance between two longitudes from tropics to the polar regions.
357 This change mainly helps to improve the reconstruction in the high-latitude regions;

358 2) The three iterative runs are taken to effectively bring in different scales of
359 variability with influencing radius changing from 2,000 km (2,500 km at 700–6000 m) to
360 800 km and 300 km, respectively, based on the tests presented in Cheng and Zhu (2016)
361 and Cheng et al., (2017);

362 3) For each month, IAPv3 used 40 model simulations (historical runs) from the
363 Coupled Model Intercomparison Project phase 5 (CMIP5) to provide a flow-dependent
364 ensemble, which is then constrained by observations to provide optimized spatial
365 covariance. IAP mapping uses model-based covariance because we argue that spatial
366 covariance can never be satisfactorily parametrized by some simple basic functions (such
367 as Gaussian) given its complexity. With model-based, flow-dependent, and dynamically-
368 consistent covariance, the IAP mapping provides a more realistic reconstruction than other
369 approaches based on Gaussian-based parameterized covariance, as evaluated by many
370 studies (Cheng et al., 2017; Cheng et al., 2020; Dangendorf et al., 2021; Nerem et al.,
371 2018).

372 4) The observation error variance (\mathbf{R}), which represents the error of the observations,
373 is updated in IAPv4 as follows. \mathbf{R} consists of both the instrumental error (\mathbf{R}_e) due to
374 inaccuracy and the representativeness error (\mathbf{R}_r) due to the need to represent the spatial (at
375 1° by 1° and 1 m standard grid depths) and temporal (1 month) averages from a limited
376 numbered of observations (Cheng and Zhu, 2016):

377
$$\mathbf{R} = \mathbf{R}_e + \mathbf{R}_r = (\sum_1^M \mathbf{E}i)/M + \sigma^2/M,$$


378 where M observations exist for a given grid cell. $\mathbf{E}i$ is the instrument's precision for
379 each individual observation, assuming random error (the basic assumption is that after bias
380 correction, the systematic errors can be eliminated). \mathbf{R}_e in each grid cell is set to the mean
381 of the typical precision of the different instruments contributing data in the cell, which is
382 set according to IQuOD (International Quality-Controlled Ocean Database) specification
383 (Cowley et al., 2021). σ^2 represents the variance of the various temperature measurements
384 against the monthly mean value. The data from 2005 to 2022 are used to calculate σ^2 in
385 each grid because of greater data abundance and quality compared to earlier times.

386 As the representativeness error (\mathbf{R}_r) is expected to be flow-dependent (i.e., the error is
387 expected to be higher in areas with a large gradient of the flow speed and regions of higher
388 variability), more observations are required to represent the mean value. Figure 4 shows a
389 larger variance (σ^2) in the boundary-current regions and near the Antarctic Circumpolar
390 Current (ACC) in the upper ocean (e.g., 10 m, 200 m, 500 m). At 200 m, it shows a larger
391 σ^2 in the Western Pacific Ocean, corresponding to the large thermocline variations at this
392 layer. Below 1000 m, larger σ^2 along the ACC frontal regions and in the North Atlantic
393 Ocean occur because of a stronger mixing and convection in these regions.

域代码已更改

域代码已更改

394 The uncertainty in the derived gridded reconstruction is also based on the EnOI
 395 framework formulated by Cheng and Zhu, (2016). The uncertainty accounts for
 396 instrumental, sampling and mapping errors. Other error sources, including the choice of
 397 climatology, vertical interpolation, bias corrections, and QC, are not considered in this
 398 uncertainty estimate. Therefore, a more thorough uncertainty quantification method is
 399 needed, and this is under development in a separate study.



400
 401 **Figure 4: Variance (σ^2) of ocean temperature at several representative layers. (a)**
 402 **10 m, (b) 200 m, (c) 500 m and (d) 1000 m. The unit is $^{\circ}\text{C}^2$.**
 403

404 **Table 1.** General information on IAPv4 and IAPv3 data products.

	IAPv3	IAPv4
Horizontal resolution	Global ($1^{\circ} \times 1^{\circ}$)	Global ($1^{\circ} \times 1^{\circ}$)
Vertical levels	41 levels from 1 m to 2000 m (1, 5, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 120, 140, 160, 180, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 1000, 1100, 1200, 1300, 1400, 1500, 1600, 1700, 1800, 2000)	119 levels from 1 m to 6000 m (1, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 220, 240, 260, 280, 300, 320, 340, 360, 380, 400, 425, 450, 475, 500, 525, 550, 575, 600, 625, 650, 675, 700, 750, 800, 850, 900, 950, 1000, 1050, 1100, 1150, 1200,

		1250, 1300, 1350, 1400, 1450, 1500, 1550, 1600, 1650, 1700, 1750, 1800, 1850, 1900, 1950, 2000, 2100, 2200, 2300, 2400, 2500, 2600, 2700, 2800, 2900, 3000, 3100, 3200, 3300, 3400, 3500, 3600, 3700, 3800, 3900, 4000, 4100, 4200, 4300, 4400, 4500, 4600, 4700, 4800, 4900, 5000, 5100, 5200, 5300, 5400, 5500, 5600, 5700, 5800, 5900, 6000)
Time period and resolution	1940–2022 (reliable data after 1955), monthly	1940–present (reliable data after 1955), monthly
Quality-control	WOD (Garcia et al., 2018)	CODC-QC (Tan et al., 2023)
Vertical interpolation	RR (Reiniger and Ross, 1968) interpolation	RR (Reiniger and Ross, 1968) interpolation
Climatology	IAP climatology: simple gridded average and then spatial interpolation with distance-weighted average	Improved IAP reconstruction with EnOI approach
XBT bias correction	CH14 (updated in 2018)	CH14 (revised and updated in 2023)
MBT bias correction	IK09 (Ishii and Kimoto, 2009)	GC20 (Gouretski and Cheng, 2020)
APB bias correction	None	GCR24 (Gouretski et al., 2024)
Bottle bias correction	None	GCT22 (Gouretski et al., 2022)
Mapping	EnOI-DE with influencing radius of 20, 8, 3 degrees, iteratively.	EnOI-DE with influencing radius of 2000, 800, 300 km, iteratively. Representative error updated with 2005–2022 observations. The radius of influence does not cross the land.
Uncertainty	Given by EnOI framework accounting for instrumental error and horizontal sampling/mapping error	Given by EnOI framework accounting for instrumental error and horizontal sampling/mapping error
DOI	/	http://dx.doi.org/10.12157/IOCAS.20240117.002 for temperature data (Cheng et al., 2024a) and http://dx.doi.org/10.12157/IOCAS.20240117.001 for ocean heat content data (Cheng et al., 2024b)

406 2.7 OHC calculation and volume correction

407 Based on the gridded temperature reconstruction (Table 1), OHC in each grid is
408 calculated as $OHC(x, y, z) = c_p \iiint_{V(x,y,z)} \rho T dV(x, y, z)$. following TEOS-10 standards,
409 where c_p is a constant of $\sim 3991.9 \text{ J (kg K)}^{-1}$ according to the new TEOS-10 standard
410 formulation as conservative temperature and absolute salinity are used, ρ is potential
411 density in kg m^{-3} , and T is conservative temperature measured in degrees Celsius (here it
412 is anomaly relative to the 2006–2020 baseline) (Cheng et al., 2022a).



413 As OHC is an integrated metric over a specific ocean volume, properly identifying
414 ocean volume is critical, especially in shallow waters. Previous studies found a 10–20 %
415 difference in the OHC trend in recent decades between different land-ocean masks (von
416 Schuckmann and Le Traon, 2011; Meyssignac et al., 2019; Savita et al., 2022).
417 Specifically, in marginal sea areas with complex topography, $1^\circ \times 1^\circ \times \Delta z$ grid boxes
418 (where Δz is the depth range of the grid box) near coasts and islands typically cover both
419 ocean and land areas but are assigned to represent land or ocean only. Thus, the gridded
420 ocean temperature datasets are subjected to errors from inaccurate land-sea attribution.
421 Here, we offer a volume correction (VC) for these grid boxes to improve the OHC
422 estimate, as follows.

423 For each $1^\circ \times 1^\circ \times \Delta z$ grid box, we introduce a VC factor (denoted as F_{VC}) to correct
424 the OHC values: $OHC_{VC}(x, y, z) = OHC(x, y, z) \times F_{VC}(x, y, z)$. First, we assume the
425 seawater volume distribution in 1 arc-minute topographic data of ETOPO1 as “truth”. No
426 correction is needed if a box is assigned to ocean according to ETOPO1 data, thus, $F_{VC}=1$.
427 If a fraction of a $1^\circ \times 1^\circ \times \Delta z$ grid box is land according to ETOPO1 and IAP data includes
428 T/OHC values, the F_{VC} is represented by the fraction of the ocean volume in this box
429 (illustrated in Fig. 5), and the volume for OHC calculations can be corrected with $F_{VC}(i)$. In
430 a grid box, if there is no IAP data (i.e., it is land according to the IAP mask), but this box
431 contains some ocean volume according to ETOPO1 data, we define $F_{VC}(a)$ again as the
432 fraction of the ocean volume in this box, and then this $F_{VC}(a)$ is added to the adjacent grid
433 boxes where there are values in IAP data. If all the adjacent grid boxes contain no data, the
434 volume is equally redistributed to the diagonal boxes (Fig. 5). The volume is discarded if
435 there is no data in all adjacent and diagonal boxes.

436 With this approach, the VC factor in each grid box is a sum of two components: a
437 local adjustment $F_{VC(i)}$ and a redistribution from the adjacent grids:

$$438 F_{VC(a)}: F_{VC} = F_{VC(i)} + F_{VC(a)},$$

439 To avoid misidentification of sea ice, we performed VC only on the global grid points
440 within 60 °S to 60 °N. Eventually, we obtained a three-dimensional FVC that fits the IAP
441 grids (119 × 360 × 180; depth coverage to 6000 m) and used it to compute OHC. The VC
442 applied to ~15% of all the 1° × 1° × Δz grid boxes of IAPv4 ocean grid boxes (with $F_{VC} \neq$
443 1) for the entire 0-6000 m ocean and ~10% grid boxes of the upper 2000 m. Since the open
444 ocean accounts for the vast majority of the global ocean volume, the influence of the VC
445 method on the global OHC trend is small. For example, the upper 2000 m OHC trend with
446 VC is ~0.15% (~0.45%) smaller than without VC from 1958-2023 (2005-2023) for IAPv4.
447 However, it can significantly affect regional OHC estimates, especially in regions with
448 complex topography. For example, the Maritime Continent region's 0-2000 m OHC trend
449 is reduced by 6.9% (4.2%) after applying VC from 1958-2023 (2005-2023) (Jin et al.,
450 2024).



451

452 **Figure 5:** An example explaining the Volume Correction algorithm. (a) Bathymetry
453 derived from ETOPO1. (b) Bathymetry in IAPv4 analysis.

454

455 2.8 Independent datasets for comparison and evaluation

456 Four Sea Surface Temperature (SST) datasets are used to evaluate the upper-most
457 layer (1 m) of IAPv4, including Extended Reconstructed SST version 5 (ERSST5) (Huang
458 et al., 2017); Japan Meteorological Agency Centennial Observation-Based Estimates of
459 SSTs version 1 (COBE1) (Ishii et al., 2005), and its version 2: COBE2 (Hirahara et al.,
460 2014); Hadley Centre Sea Ice and Sea Surface Temperature dataset (HadISST) (Rayner et
461 al., 2003). The anomalies relative to a 2006-2020 average were computed by removing the
462 monthly climatology. Measurements of SST are made *in situ* by means of thermometers or
463 retrieved remotely from infrared and passive microwave radiometers on satellites
464 (Kennedy, 2014; O'Carroll et al., 2019). Satellite SST observations began in the early

465 1980s. *In situ* SST observations go back to the 19th century and involve many different
466 measurement methods, including wooden and later insulated metal buckets to collect water
467 samples, engine room inlet measurements, and sensors on moored and drifting buoys
468 (Kennedy, 2014). The subsurface temperatures are collected as “profiles” which contain
469 multiple measurements at discrete vertical levels. Because of the differences in observation
470 systems, SSTs are fundamentally different in their temporal and spatial coverage and
471 temporal extent compared to subsurface observations on which OHC estimates rely. SST
472 measurements also have different uncertainty sources and error structures; thus, the two
473 systems are typically treated as independent data sources and have been used for cross-
474 validation (Gouretski et al., 2012).

475 An independent *in situ* observation dataset in the Labrador Sea is used to evaluate
476 IAPv4. This dataset, provided by the Bedford Institute of Oceanography (BIO)
477 (Yashayaev, 2007; Yashayaev and Loder, 2017), includes independently validated and
478 bias-corrected data from multi-section hydrological surveys (i.e., AR7W) in the Labrador
479 Sea, spanning from 1896 to 2020 (this study used 1960-2020 data). These data have not
480 been incorporated into the WOD.

481 The capability of the new product to close the sea level budget and the Earth’s energy
482 budget also provides tools for validation. A superior dataset should be capable of closing
483 the sea level and the Earth’s energy budgets. The total sea level change has been monitored
484 via satellite altimetry since 1993 (from the University of Colorado
485 <https://sealevel.colorado.edu/>). The ocean mass change is derived from JPL RL06.1Mv3
486 Mascon Solution GRACE and GRACE-FO data since 2002 (Watkins et al., 2015). For
487 long-term total sea level change since the 1950s, we use a tide-gauge-based reconstruction
488 (Frederikse et al., 2020). During the same period, the estimates of the Greenland ice sheet,
489 Antarctic ice sheet, land water storage, and glacier ice melt contributions from Frederikse
490 et al., (2020) are used to derive ocean mass change. To derive steric sea level, IAP salinity
491 data is used (Cheng et al. 2020). The temperature and salinity data are converted to steric
492 sea level based on the Thermodynamic Equation Of Seawater – 2010 (TEOS-10) standard
493 (McDougall and Barker, 2011).

494 For the energy budget, the ice, land, and atmosphere heat content changes are from
495 (von Schuckmann et al., 2023) from 1960 to the present. Because of the less reliable data
496 before the 1990s for land, sea ice and ice sheets, the other set of land–atmosphere–ice data
497 from 2005–19 is used as in Trenberth, (2022) to investigate the recent changes. The net

498 radiation change at the top of the atmosphere is based on Clouds and Earth's Radiant
499 Energy Systems (CERES) Energy Balanced and Filled (EBAF) data from Loeb et al.,
500 (2021) and Loeb et al., (2018) and Deep-C data from the University of Reading (Liu and
501 Allan, 2022; Liu et al., 2017).

502 Several gridded ocean T/OHC gridded products are used here for inter-comparison,
503 including the IAPv3 (Cheng et al., 2017), the EN4 ocean objective analysis product from
504 the UK Met Office Hadley Centre (Good et al., 2013); the ocean objective analysis product
505 (Ishii et al., 2017) (termed "ISH" hereafter) from JMA, an Argo-only gridded product from
506 SCRIPPS (Roemmich and Gilson, 2009) (termed "RG" hereafter), and an OHC product
507 based on random forest regressions (termed "RFROM" hereafter) using in situ training
508 data from Argo and other sources on a 7-day \times $1/4^\circ \times 1/4^\circ$ grid with latitude, longitude,
509 time, SSH, and SST as predictors (Lyman and Johnson, 2023). Several datasets available
510 in IPCC-AR6 (Gulev et al., 2023) are used for comparison, including: the PMEL product
511 from Lyman and Johnson, (2014); Machine learning based reconstruction of OHC by
512 Bagnell and DeVries, (2021); BOA product based on refined Barnes successive corrections
513 by the China Argo Real-time Data Center (Li et al., 2017); International Pacific Research
514 Center (IPRC) (2005-2020), von Schuckmann and Le Traon 2011 (KvS11); Green function
515 based OHC estimate derived from SST (Zanna et al., 2019).

516

517 **2.9 Trend calculation and uncertainty estimates**

518 The trends in this study have been estimated by a LOWESS approach (Cheng et al.,
519 2022b), i.e., we apply a locally weighted scatterplot smoothing (LOWESS) to the time
520 series (25-year window, equal to an effective 15-years smoothing), and then the OHC
521 difference between the first and the end year is used to calculate the trend. This approach
522 provides an effective method to quantify the local trend by minimizing the impact of year-
523 to-year variability and start/end points.

524 Throughout this paper, the 90 % confidence interval is shown. The uncertainty of
525 trend also follows the approach in Cheng et al., (2022a) based on a Monte Carlo
526 simulation. First, a surrogate OHC series is formed by simulating a new residual series
527 (after removing the LOWESS smoothed time series) based on the AR(1) process and
528 adding it to the LOWESS line. Then a LOWESS trendline is estimated for each surrogate.
529 This process is repeated 1000 times, and 1000 trendlines are available. The 90 %
530 confidence interval for the trendline is calculated based on ± 1.65 times the standard

531 deviation of all 1000 trendlines of the surrogates. Secondly, the uncertainty in the rate of
532 the OHC is estimated by the 1000 LOWESS trendlines: 1) calculating the rate based on the
533 difference between the first and last annual mean value of the LOWESS trendline in a
534 specific period; 2) calculating ± 1.65 times the standard deviation of the 1000 rate values.


535

536 **3. Results**

537 **3.1 Climatological annual cycle**

538 The annual cycle of the OHC above 2000 m of IAPv4 is compared with IAPv3, ISH,
539 EN4, RG and RFROM (Fig. 6 and Fig. 7) for 2006–2020. There is a consistent annual
540 cycle among different datasets for the global and hemispheric oceans. Globally, the ocean
541 releases heat from boreal spring to autumn and accumulates heat from boreal autumn to
542 spring, which is dominated by the southern hemisphere due to its larger ocean surface area
543 (Fig. 6). The two hemispheres show opposite annual variations in OHC, associated with
544 the annual change of solar radiation and different distribution of land and sea. For the
545 global OHC above 2000 m, IAPv4 shows a positive peak in April and a dip in August,
546 with the magnitude of OHC variation of 60.4 ZJ for IAPv4 (66.9 ZJ for IAPv3), consistent
547 with other datasets: 53.2 ZJ for ISH, 58.1 ZJ for EN4, 69.2 ZJ for RG and 56.6 ZJ for
548 RFROM (where 1 ZJ = 10^{21} J).

549 There are some unphysical variations in the OHC annual variations for IAPv3 (blue
550 lines). For example, the global OHC shows large spikes in January and December, and a
551 big shift from September to October, by contrast, the other three data products show much
552 smoother changes (Fig. 6a). The IAPv3 Arctic OHC (north of 69.5 °N) shows different
553 phase change compared with the other datasets together with a big shift from September to
554 December, and the magnitude of variability is much larger in IAPv3 than other datasets
555 (Fig. 6d). The improvement in IAPv4 is mainly because of the methodology
556 improvements: IAPv3 used 1990–2005 data to construct climatology which suffered from
557 errors related to sparse data coverage, use of “degree distance” instead of “km distance”,
558 and other error sources. Therefore, the IAPv4 analysis presents a physically tenable OHC
559 seasonal variation.



560

561


562 **Figure 6: Annual cycle of OHC of upper 2000 m for (a) the global oceans, (b) the**
 563 **Southern Hemisphere, (c) the Northern Hemisphere and (d) the oceans north of**
 564 **69.5°N. Five different data products are presented, including IAPv4 (red), IAPv3 (black),**
 565 **ISH (purple), EN4 (green), RG (orange), and RFROM (pink).**

566

567 IAPv4 OHC data shows significant improvements in the Arctic region, reflected in
 568 both the spatial distribution and seasonal variation of OHC. In IAPv3, the maximum upper
 569 2000 m OHC occurs in December, and the minimum OHC occurs in August. However, for
 570 IAPv4, the maximum amounts to 2.9 ZJ in October and decreases to a minimum of -3.4 ZJ
 571 in April. [This estimate of the Arctic annual cycle is consistent with a constrained Arctic](#)
 572 [OHC estimate with atmospheric data by enforcing energy budget closure \(Mayer et al.,](#)
 573 [2019\). Furthermore, the spread of the OHC annual cycle in the Arctic region across](#)
 574 [different datasets is reduced from 5.2 ZJ to 2.5 ZJ, indicating a smaller uncertainty. The](#)
 575 [spatial OHC anomaly distribution in the Arctic region of the IAPv4 is more spatially](#)
 576 [homogeneous than IAPv3, and IAPv3 appears as rays emerging from the pole which are](#)
 577 [not physical \(Fig. 7\). IAPv4 displays a consistent seasonal variation north of 69.5 °N](#)
 578 [mainly because of the changes of the influencing radius from “degrees” to “kilometers”.](#)

设置了格式: 英语(英国)

删除了: T



581 **Figure 7: Seasonal distribution of monthly mean upper 2000 m OHC anomalies**
 582 **and root mean square error (RMSE) of OHC 0-2000 m between gridded data and in**
 583 **situ observations. For OHC anomalies, four months are shown: March, June,**
 584 **September, and December. The OHC anomalies are relative to the 2006 – 2020**
 585 **annual mean. The upper and lower panels are for IAPv3 and IAPv4 products,**
 586 **respectively. The panels in the last column are for annual RMSE for IAPv3 (upper) and**
 587 **IAPv4 (lower), respectively.**


588

589 3.2 Mixed layer depth

590 Mixed layer depth (MLD) provides a crucial parameter of upper ocean dynamics
 591 relevant for upper-deeper ocean and air-sea interactions. Spatial distributions of the MLD
 592 in March and August are shown in Fig. 8 for IAPv4, based on criteria of $\Delta T = 0.2$ °C
 593 temperature for the 10 m depth temperature. As expected, the seasonal variations of the
 594 MLD are generally opposite in the northern and southern hemispheres. The MLD shows a
 595 much stronger seasonal variation in the subtropics and midlatitudes (for example, 20°~70°
 596 in both hemispheres) than in other regions (including the tropics, for example,
 597 20°S~20°N), which is manifested as shallower MLD (~20 m) in summer due to strong
 598 surface heating that increases stratification, and deeper MLD in winter (>70 m) because of
 599 surface cooling and increased surface wind creating stronger mixing.

600 In the north hemisphere, the maximum MLD occurs during the wintertime in the
 601 subpolar North Atlantic deep water formation regions (40 °N ~ 65 °N), with values over

602 500 m in the Iceland Basin. In comparison, in the midlatitudes, the maximum of MLD is
 603 generally less than 125 m in the wintertime. The MLD minimum in the north hemisphere is
 604 in the summertime, and the values are mostly within 20 m depth. In the Southern
 605 Hemisphere, the MLD maximum values (deeper than 300 m) occur between 45 °S and
 606 60 °S of the Southern Ocean (north of the Antarctic Circumpolar Current) in the boreal
 607 summer where the year-round intense westerly winds are located. The minimum MLD in
 608 this region in the boreal winter is less than 70 m. The seasonal variation of the MLD is
 609 well established by previous studies (Chu and Fan, 2023; de Boyer Montégut et al., 2004;
 610 Holte et al., 2017), and this evaluation confirms that IAPv4 temperature data is capable of
 611 reasonably representing the MLD. However, as pointed out by de Boyer Montégut (2004),
 612 the MLD estimated from the average temperature profiles might lead to an underestimation
 613 of MLD by ~25% compared to the MLD computed from individual profiles based on the
 614 same 0.2 °C criterion method. This potential issue needs further investigation.



615
 616 **Figure 8: Spatial pattern of the climatological mean MLD (left panels) and zonal**
 617 **mean MLD (right panels) in March (top) and August (bottom) estimated from the**


618 **IAPv4.** Here, the MLD is calculated using the temperature difference criterion of $\Delta T =$
619 $0.2\text{ }^{\circ}\text{C}$ between the surface and 10-meter depth.

620

621 **3.3 Sea surface temperature**

622 IAPv4 and IAPv3 temperature time series at 1 m depth (Fig. 9) are compared with
623 four independent SST data products (ERSST5, HadISST, COBE1, and COBE2). All data
624 products including IAPv4 show robust sea surface warming in the global ocean and four
625 main basins since 1955 (Fig. 9). Since the HadISST and COBE2 data did not include the
626 year 2023, we compare the long-term SST trend during 1955–2022 using these products
627 (Fig. 9f). The global-mean IAPv4 SST rate between 1955 and 2022 is $1.01 \pm 0.15\text{ }^{\circ}\text{C}$
628 century^{-1} (90 % CI), which is within the range of the SST products (ranging from 0.78 to
629 $1.05\text{ }^{\circ}\text{C century}^{-1}$). The 1955–2022 trend of IAPv4 SST is slightly weaker than IAPv3 for
630 the global ocean ($1.11 \pm 0.16\text{ }^{\circ}\text{C century}^{-1}$) and all the ocean basins. The largest difference
631 between IAPv4 and other SST products comes mainly from the Pacific and the Southern
632 Ocean before 1980, associated with sparser in situ observations for both SST and
633 subsurface temperature data.


634 The spatial distribution of long-term SST trends over the 1955–2022 period provides
635 insights into the data consistencies and differences. First, IAPv4 shows a pattern of SST
636 consistent with other datasets (Fig. 10). More rapid warming is found in the poleward
637 western boundary currents regions, such as the East Australian Current and the Gulf
638 Stream. The warmer ocean in the upwelling areas, such as the Tropical Eastern Pacific and
639 Gulf of Guinea, are identified by all data products. The surface warming in the South
640 Indian for IAPv4 data is weaker than for IAPv3, ERSST5, and COBE2 but is more
641 consistent with HadISST and COBE1. The surface cooling to the south of $60\text{ }^{\circ}\text{S}$ can also
642 be found in all the datasets but with some discrepancies in magnitude and locations related
643 to data sparsity. The tropical Pacific SST trends are mostly insignificant in the eastern and
644 south-eastern Pacific Ocean because of the strong inter-annual and decadal fluctuations
645 (Figure not shown).



646

647 **Figure 9: Global and basin time series of SST change for IAPv4, compared with**
 648 **ERSST/HadISST/COBE1/COBE2 and IAPv3 from 1955 to present. (a) Global, (b)**
 649 **Pacific, (c) Atlantic, (d) Indian and (e) Southern oceans (South of 30 °S) (units: °C). (f)**
 650 **shows the warming rate from 1955 to 2022 The pink thin line is the monthly time series of**
 651 **IAPv4 SST and other time series are annual time series of different datasets. The vertical**
 652 **scales are different for different panels. All anomaly time series are relative to a 2006–**
 653 **2020 baseline.**

654



655

656 **Figure 10: Spatial maps of the SST long-term trends during the 1955–2022 period.** (a)
 657 IAPv4, (b) IAPv3, (c) ERSST5, (d) HadISST, (e) COBE1 and (f) COBE2 (units: $10^{-2} \text{ }^{\circ}\text{C}$
 658 yr^{-1}). The contour line interval is $0.5 \times 10^{-2} \text{ }^{\circ}\text{C yr}^{-1}$. The stippling indicates the regions with
 659 signals that are not statistically significant (90 % CI).

660

661 3.4 Global OHC time series


662 Global OHC time series for 0–700 m, 700–2000 m, 0–2000 m, and 2000–6000 m
 663 layers of IAPv4 (Fig. 11) for 1955–2023 versus IAPv3 show a robust ocean warming, with
 664 a linear warming rate of $4.4 \pm 0.2 \text{ ZJ yr}^{-1}$ (0–700 m), $2.0 \pm 0.1 \text{ ZJ yr}^{-1}$ (700–2000 m), and
 665 $6.4 \pm 0.3 \text{ ZJ yr}^{-1}$ (0–2000 m). The long-term warming revealed by IAPv4 is greater than
 666 IAPv3 ($4.1 \pm 0.2 \text{ ZJ yr}^{-1}$ for 0–700 m, $1.9 \pm 0.1 \text{ ZJ yr}^{-1}$ for 700–2000 m and 6.0 ± 0.3
 667 ZJ yr^{-1} for 0–2000 m). Before ~1980, bottle bias correction reduces the time-varying
 668 systematic warm bias in Nansen bottle data and leads to a stronger warming rate from
 669 1955–1990. The updated MBT and XBT corrections are mainly responsible for the

670 difference between 1980 and 2000 (Cheng et al., 2014; Gouretski and Cheng, 2020). Data
671 QC impacts the intra-seasonal and inter-annual variation of the OHC time series (Tan et
672 al., 2023). Also, because of the application of Bottle/XBT/MBT corrections, IAPv4 shows
673 a stronger upper 2000 m ocean warming trend than most of the other available products
674 assessed in Fig. 12.

675 From 2005–2023, the new IAPv4 product shows stronger warming than IAPv3. The
676 mean upper 2000 m warming rate is $10.7 \pm 1.0 \text{ ZJ yr}^{-1}$ for IAPv4 and $9.6 \pm 1.1 \text{ ZJ yr}^{-1}$ for
677 IAPv3 (Fig. 11), mainly because of the replacement of the WOD-QC system by the new
678 CODC-QC system in IAPv4. Tan et al., (2023) indicated that the WOD-QC system had
679 removed more extreme higher temperature values in the regions of warm eddies and
680 marine heat waves than CODC-QC. The IAPv3 700–2000 m OHC shows a much bigger
681 drop in 2018 than IAPv4 (Fig. 11b), while the IAPv4 indicates an approximately linear
682 700–2000 m warming since 2005, resulting in stronger 700–2000 m warming in IAPv4
683 ($3.6 \pm 0.5 \text{ ZJ yr}^{-1}$) than in IAPv3 ($2.9 \pm 0.5 \text{ ZJ yr}^{-1}$). Compared with other available
684 products shown in Fig. 12, IAPv4 shows a similar OHC 0–2000 m trend to RFROM from
685 2005–2023, but with stronger warming trends than the two Argo-based products (BOA and
686 SCRIPPS). From 1993–2023, IAPv4 showed a stronger OHC 0–2000 m trend than NCEI,
687 Ishii, OPEN, and Zanna data and a slightly weaker trend than PMEL and RFROM (Fig.
688 12).

689 Since the 1990s, the World Ocean Circulation Experiment (WOCE) provided a global
690 network of abyssal ocean observations, sustained by repeated hydrological lines and a
691 deep-Argo program (Katsumata et al., 2022; Roemmich et al., 2019; Sloyan et al., 2019).
692 These high-quality data provide an opportunity to estimate deep OHC changes below 2000
693 m in this study. IAPv4 provides a new OHC estimate below 2000 m by collecting 5 years
694 of data centered on each month. The result (Fig. 11d) indicates a robust abyssal (2000–
695 6000 m) ocean warming trend since ~1993 of $2.0 \pm 0.3 \text{ ZJ yr}^{-1}$. This is higher (within the
696 uncertainty range) than the previous estimate of $1.17 \pm 0.5 \text{ ZJ yr}^{-1}$ in Purkey and Johnson
697 (2010) but consistent with the recent assessment showing the acceleration of deep ocean
698 warming in the Southwest Pacific Ocean (Johnson et al., 2019).

699



700


701 **Figure 11: Global OHC time series for 0–700 m (a), 700–2000 m (b), 0–2000 m (c) and**
 702 **2000–6000 m (d).** All-time series are relative to a 1981–2010 baseline. The shading

703 indicates the 90 % confidence interval. The vertical scales are different for different panels.

704

The unit is ZJ.

705



706

707 **Figure 12: A comparison of annual mean OHC 0-2000 m time series from different**
 708 **data products.** Solid and dashed lines represent direct and indirect estimates, respectively,
 709 and shading indicates the IAPv4 90% confidence interval (pink shading). OHC anomalies
 710 are relative to a 2005–2019 baseline. The plot is updated from Cheng et al. (2022a).

711

712 Another feature of IAPv4 is the suppression of month-to-month noise compared to
 713 many available data products. Trenberth et al. (2016) noted that the month-to-month
 714 variation (quantified by the standard deviation of the monthly $dOHC/dt$ time series) in all
 715 *in situ*-based OHC records is much larger than implied by the CERES records, suggesting
 716 that the OHC variation on this time scale is most likely spurious. Therefore, the magnitude
 717 of the month-to-month variation in the OHC record can be used as a benchmark of the data
 718 quality. The standard deviation of the CERES record is 0.67 Wm^{-2} from 2005 to 2023
 719 (Loeb et al., 2018). While IAPv4, IAPv3, ISH, EN4, BOA, NCEI, and SIO data show a
 720 standard deviation of $dOHC/dt$ time series of 3.52, 3.52, 7.49, 8.79, 10.05, 11.29, 10.00
 721 Wm^{-2} , respectively for the upper 2000 m (Table 2). Note that differentiation to get the rate
 722 of change amplifies noise, and applying a 12-month running smoother significantly knocks
 723 down the noise so that the IAPv4 standard deviation becomes 0.75 Wm^{-2} , the smallest
 724 among the datasets investigated in this study (Table 2) and is the most physically plausible
 725 time series from this noise-level perspective. In addition, Lyman and Johnson's (2013) data

726 suggest a yearly variance ratio of 1.3 between annual RFROM and CERES data from 2008
 727 to 2021. Using the yearly mean OHCT indicates a ratio of 1.4 at the same period between
 728 IAPv4 and CERES, which is similar to that of RFROM.

729

730 **Table 2. Characteristics of Month-to-month variation of OHCT compared with**
 731 **CERES.** Comparisons of different ocean gridded products: the monthly standard deviation
 732 (std dev) of the monthly rates of change of OHC (W m^{-2}); the corresponding standard
 733 deviation of the 12-month running mean (13-points are used, with start-point and endpoint
 734 weighted by 0.5), and the linear trend with 90% confidence limits (Wm^{-2}) (global surface
 735 area). The values are for 2005–2022. The OHC trend for CERES is calculated as the mean
 736 of net TOA radiation flux within 2005–2022 multiplied by 0.9, assuming 90% of the EEI
 737 stored in the ocean.


Source	Std dev	Std dev (12 month)	OHC Trend (2005–2022)
IAPv4	3.52	0.75	0.66 ± 0.04
IAPv3	3.52	0.79	0.56 ± 0.03
ISH	7.49	1.35	0.63 ± 0.05
EN4	8.79	1.03	0.67 ± 0.04
BOA	10.05	1.16	0.60 ± 0.07
NECI	11.29	1.11	0.61 ± 0.07
SIO	10.00	1.24	0.56 ± 0.08
CERES	0.67	0.33	0.77

738

739 3.5 Regional OHC trends

740 For 1960–2023 (Fig. 13), the IAPv4 trends are slightly weaker than IAPv3 in the
 741 Pacific Ocean but slightly higher in the Atlantic Ocean (Fig. 13), with more than 95 % of
 742 the ocean area showing a warming trend. The polar regions also show remarkable
 743 differences compared to IAPv3 (Section 3.1), mainly because of the change of covariance,
 744 which improves the spatial reconstruction in the polar regions. The IAPv4 shows stronger
 745 warming near the boundary currents regions, mainly because of the improved QC that does
 746 not flag high-temperature anomalies. Nevertheless, the pattern of trends is very similar in
 747 the two versions of data, indicating the robustness of the ocean warming pattern. The
 748 Atlantic Ocean (within 50°S – 50°N) and the Southern Ocean store more heat than the
 749 other basins, probably associated with the deep convection and subduction processes


750 effectively transporting heat into the deep layers (Cheng et al., 2022a). The cold spots
 751 mainly include the Northwest Pacific and subpolar North Atlantic Ocean. In particular, the
 752 so-called “warming hole” in the subpolar North Atlantic Ocean can extend to at least 800
 753 m and is responsible for decreased OHC in this region. Some studies have linked this
 754 fingerprint to the slowdown of AMOC (Rahmstorf et al, 2015; Caesar et al., 2018).



756 **Figure 13: Spatial pattern of the OHC trends for 0–300 m, 0–700 m and 0–2000 m,**
 757 **700–2000 m from 1960 to 2023.** The left panels show IAPv3, the middle panels are
 758 IAPv4; the right panels are the difference between IAPv4 and IAPv3.


759 For 1991–2023 (Fig. 14), the IAPv4 and IAPv3 pattern is also consistent. A trend
 760 pattern mimicing a negative Pacific Decadal Variability (PDV) phase appears in the Pacific
 761 for the 0–300 m, 0–700 m, and 0–2000 m OHCs. There is a contrast between the warming

762 trend of the tropical western Pacific and the cooling trend of the tropical eastern Pacific.
 763 Some studies have linked this pattern to the natural climate mode (PDV) (England et al.,
 764 2014), but some suggest it is a forced change driven by greenhouse gas increases (Fasullo
 765 and Nerem, 2018; Mann, 2021). Below 700 m, the 1960–2023 and 1991–2023 trend
 766 patterns are similar because deep ocean warming mainly occurs after 1990. Broad warming
 767 in most regions, but subtropical oceans in the West Pacific and South Indian oceans show a
 768 cooling, which is likely related to the subtropical gyre intensification in the North but a
 769 spin-down in the North Pacific Ocean (Zhang et al., 2014).



771 **Figure 14: Spatial pattern of the OHC trends for 0–300 m, 0–700 m, 0–2000 m and**
 772 **700–2000 m from 1991 to 2023.** The left panels show IAPv3, the middle panels are
 773 IAPv4; the right panels are the difference between IAPv4 and IAPv3.

774 Furthermore, the reconstruction of IAPv4 is compared with completely independent
 775 observations in the central Labrador Sea (see Data and Methods section for details;
 776 Yashayaev, 2007; Yashayaev and Loder, 2017) for the 200-2000 m mean temperature time
 777 series (Fig. 15). The direct observations show a substantial decadal variation in the central
 778 Labrador Sea, with negative anomalies 1970-2003 and 2015-2020, and positive anomalies
 779 1963-1972 and 2004-2014. Reconstructed based on data from WOD, IAPv4 can well
 780 represent this decadal variability. The largest difference occurs in 1989, where direct
 781 observations show nearly zero anomaly while IAPv4 shows a big negative anomaly; this
 782 difference is likely caused by using a time window in IAPv4, which has a smoothing effect
 783 on the time series.



784
 785 **Figure 15: Comparison of IAPv4 data with independent observations in the central**
 786 **Labrador Sea (304 -310 °E, 55-61 °N) from 1960 to 2020.** The 200-2000 m averaged
 787 temperature anomaly time series is shown, and the baseline is 1960-2020. The inner box
 788 shows the locations of the independent observations in black dots (showing a total of
 789 49,849 profiles).

790
 791 **3.6 Ocean meridional heat transport**

792 The ocean meridional heat transport (MHT) is fundamental to maintaining the earth's
 793 energy balance. Thus, its change and stability are key to the climate system and its
 794 variability. The direct observations of ocean MHT are only possible in several cross-basin
 795 sections such as RAPID. The ocean MHT can be derived from the OHC and air-sea heat

796 flux data (Trenberth and Fasullo, 2017; Trenberth et al., 2019) as follows: we integrate the
797 OHCT, air-sea heat flux and heat gain/loss by sea ice changes from the North Pole
798 southward in the Atlantic Ocean, and solve the energy budget equation, the residual at each
799 latitude is the MHT, i.e.,

$$800 \quad MHT(\varphi) = \int_{\varphi}^{90} \left[Fs + \frac{dOHC}{dt} + Q_{ice} \right] a d\varphi$$

801 Where a is the Earth's radius, φ is latitude, Fs is net surface heat flux, and Q_{ice} is
802 the heat inferred from the changes of sea ice mass. Consistent with Trenberth et al. (2019),
803 this study uses the sea ice volume data from the Pan-Arctic Ice Ocean Modeling and
804 Assimilation System (PIOMAS; Schweiger et al. 2011), and assumes a constant latent heat
805 of fusion of $3.34 \times 10^5 \text{ J kg}^{-1}$ and a density of ice of 900 kg m^{-3} . Both Fs and OHCT are
806 important for the MHT derivation: the integrated air-sea heat flux dominates the magnitude
807 of the MHT, while the OHCT dominates the variability of the MHT (Liu et al., 2020).

808 The comparison between OHC-derived MHT and RAPID data allows one to check
809 the consistency among various observations. Here, we calculate the Atlantic MHT from
810 April 2004 to December 2022 using IAPv4 OHC and air-sea net heat flux data (Fs) derived
811 by TOA net energy flux and atmospheric heat divergence (Fig. 16). Fs is an average of
812 three available products including MAYER2021 (Mayer et al., 2021) TF2018 (Trenberth et
813 al., 2019) and the DEEP-C Version 5.0 from Reading University (Liu and Allan, 2022; Liu
814 et al., 2020). The data are adjusted following Trenberth et al. (2019) approach to ensure
815 zero MHT on the Antarctica coast. The inferred time series of MHT at 26.5°N from other
816 OHC data sets (IAPv3, Ishii, and EN4) are also shown in Fig. 16, compared with the
817 RAPID observations (Johns et al., 2023).

818 The Inferred long-term mean (April 2004–December 2022) MHT from the updated
819 IAPv4 OHCT (solid red line with the mean transport of 1.18 PW) is identical to the
820 RAPID observation of 1.18 ± 0.19 PW. Different OHC datasets cause different inter-
821 annual variability in the MHT. It is shown that, from 2008 to 2020, the RAPID MHT
822 agrees best with the IAPv4 estimates with a correlation of 0.52. By comparison, the
823 correlation coefficients between RAPID and IAPv3, EN4, and Ishii are 0.33, 0.51, and
824 0.49, respectively. Over the entire period of 2005~2022, the IAPv4 lies mostly within the
825 RAPID uncertainty envelope.

删除了: and

删除了: contributed

删除了: mass


删除了: question

删除了: Here we

设置了格式: 上标

设置了格式: 上标

设置了格式: 上标



831

832 **Figure 16: Derived meridional heat transport at 26.5 °N.** The 12-month running mean
 833 northward MHT across 26.5 °N of different data sets compared with results from the
 834 RAPID array in PW. The error bars for RAPID in grey are 1.64 σ .

删除了: M


835

836 3.7 Inter-annual variability

837 The year-to-year variation of OHC is strongly influenced by ENSO from global to
 838 regional scales (Cheng et al., 2019; Roemmich and Gilson, 2011). To demonstrate the
 839 change of OHC associated ENSO, Figure 17 shows a Hovmöller diagram of the zonal
 840 upper 2000 m OHC and its change (time derivative of OHC: $d(\text{OHC})/dt$) in the tropical
 841 Pacific Ocean from 1985 to 2023, compared with the Oceanic Niño Index (ONI). It is
 842 evident that both OHC and OHCT are closely correlated with ENSO.

843 Before the onset of El Niño events, there is an accumulation of heat ($d(\text{OHC})/dt > 0$)
 844 in the southern and equatorial tropical Pacific ocean region (20 °S– 5 °N). The positive
 845 tropical Pacific $d\text{OHC}/dt$ leads ONI by ~15 months (with peak correlation >0.5), making it
 846 a precursor of El Niño (Cane and Zebiak 1985; McPhaden, 2012; Lian et al., 2023). In
 847 contrast, heat is redistributed ($d(\text{OHC})/dt < 0$) from the tropical Pacific (20 °S – 5 °N) to
 848 the North Pacific (5 °N – 25 °N) during and after El Niño (Cheng et al., 2019), with a
 849 maximum correlation >0.8 at 5 months after the El Niño peak. The magnitude of the
 850 prominent change can reach up to 50 Wm^{-2} during the 1997–1998 and 2015–2016 extreme
 851 El Niño events. For the other moderate El Niño events, the regional Pacific OHC change
 852 varies around 10–20 Wm^{-2} (Mayer et al., 2018). This typical heat recharge-discharge
 853 paradigm is crucial in ENSO evolution (Jin, 1997). Correspondingly, the zonal OHC
 854 anomalies in the Pacific Ocean show a warming state ($\text{OHC} > 0$) between ~20 °N and
 855 ~5 °S before the peak of El Niño events (with peak correlation >0.7 at 5 months before El

857 Niño peak), followed by a period of cooling ($OHC < 0$) after the peak of El Niño (with
 858 peak correlation >0.7 at 12 months after El Niño peak). These variations are all physically
 859 meaningful and indicate that IAPv4 represents regional inter-annual variability, especially
 860 associated with ENSO.



861
 862 **Figure 17:** Hovmöller diagrams illustrating the zonal mean (top) upper 2000 m $d(OHC)/dt$
 863 (Wm^{-2}) and (bottom) OHC (ZJ) in each 1° latitude band within $25^\circ S \sim 25^\circ N$ in the
 864 tropical Pacific basin using IAPv4 data. The ONI is shown in green. Vertical dashed lines
 865 denote the peak time of each Niño event.

866

867 3.8 Ocean and Earth Energy Budget

868 The EEI provides a critical quantifier of the Earth's energy flow and climate change.
 869 It is also policy-relevant because it clearly shows the need to stabilize the climate system.
 870 With new T/OHC data, we re-assess the Earth's energy inventory since 1960. The land,
 871 atmosphere, and ice contributions are from the estimates obtained by von Schuckmann et
 872 al. (2023) for 1960-2023 and by Trenberth (2022) for 2015-2019.


873 It is evident that the earth has been accumulating heat since 1960. The Earth's heat
874 inventory is 524.0 ± 95.6 ZJ from 1960 to 2023 and 260.3 ± 25.3 ZJ from 2005–2023
875 based on our data. The upper 700 m ocean, 700–2000 m, 2000 m-bottom, land, ice, and
876 atmosphere contribute to 59.3%, 24.1%, 7.4%, 5.2%, 2.9%, and 1.1% of the total EEI,
877 respectively, since 1960. The relative contribution has changed with time; for instance,
878 since 1993, the contributions are 53.7% (0–700 m ocean), 24.8% (700–2000 m ocean),
879 12.8% (2000 m–bottom ocean), 4.1% (land), 3.2% (ice), and 1.4% (atmosphere). The land
880 and ice contribution has increased in the recent two decades because of accelerated land
881 and sea ice melting (Comiso et al., 2017; Hugonnet et al., 2021; Minière et al., 2024).
882 From 2005–2019, more reliable land–atmosphere–ice datasets in Trenberth (2022) suggest
883 a non–ocean contribution of 13.4 ZJ. Combined with the results for OHC with IAPv4, the
884 accumulated EEI is 182.5 ZJ with the ocean heat uptake of 169.1 ± 19.7 for 2005–19,
885 consistent with the value of 186.4 ± 23.1 ZJ using the non–ocean contribution data by von
886 Schuckmann et al. (2023).

887 The derived energy inventory has been compared with satellite–based observations at
888 the top of the atmosphere (TOA). Two comparisons are made: (1) integrate the TOA EEI
889 to compare with the energy inventory (Fig. 18); (2) take the time derivative of the annual
890 OHC to compare it with the TOA net radiation flux (Fig. 19). Here we always assume 90%
891 of EEI is stored in the ocean and leads to an increase of OHC (Trenberth et al. 2009;
892 Hansen et al., 2011; von Schuckmann et al., 2020).

893 The first approach avoids calculating the time derivative of OHC, which exacerbates
894 noise in the time series. The net CERES change has been adjusted to 0.71 Wm^{-2} within
895 2005–2015, here we adjust the trend of the integrated CERES data to the IAPv4 OHC
896 trend to make it consistent and then compare the variability difference (Fig. 18). The
897 RMSE between DeepC and IAPv4 is 17.9 ZJ and 15.5 ZJ between CERES and IAPv4. The
898 comparison also indicates that the heat inventory shows a stronger heat increase from 2000
899 to 2005 but too slow heat accumulation during 2005–2010 compared with DeepC and
900 CERES (Fig. 18). This might be due to the data gaps before the Argo network was fully
901 established. DeepC and CERES show stronger heat accumulation since ~2015 than the
902 heat inventory, probably associated with the accelerated abyssal ocean warming found by
903 the Deep-Argo program (Johnson et al., 2019). Furthermore, IAPv4 OHC shows a slightly
904 higher (but consistent within the uncertainty range) Earth's heat uptake compared to von

905 Schuckmann et al. (2023) results by 76.2 ZJ from 1960 to 2020, mainly because the
906 correction of Nansen bottle biases and the updates of XBT and MBT biases in IAPv4 data.

907 The second approach to compare OHC with satellite-based EEI is to calculate the
908 time derivative of OHC. To suppress the month-to-month noises, we estimate annual
909 OHC based on one-year data centered on June (Fig. 19a) and December (Fig. 19b)
910 separately, and then $dOHC/dt$ is calculated with a forward derivative approach based on
911 the annual time series. The annual mean of EEI time series is also used here for
912 comparison (Fig. 19). The IAPv4 and CERES estimates show inter-annual variability with
913 a correlation of 0.44 (the correlation is statistically significant at 90% confidence level,
914 where autocorrelation reduction is taken into account). The higher correlation of IAPv4
915 versus CERES than IAPv3 increases confidence for the new data (correlation of only ~ 0.15
916 for IAPv3). The trend of $dOHC/dt$ is $0.36 \text{ Wm}^{-2} \text{ dec}^{-1}$ from 2005 to 2023, within the
917 uncertainty range of the CERES record ($0.50 \pm 0.47 \text{ Wm}^{-2} \text{ dec}^{-1}$ in Loeb et al., 2021).
918 However, it should be noted that the calculation of $dOHC/dt$ is sensitive to the choices of
919 methods, data products, and time periods because of the noises and variability in the OHC
920 time series. A careful analysis of the trend of $dOHC/dt$ (and EEI) is a research priority.




921

922 **Figure 18: The global energy budget from 1960 to 2023.** The atmosphere, land, and ice
923 heat inventory is from von Schuckmann et al., (2023). Integrated EEI from DEEP-C

删除了: A

925 (1985–2018) (Liu and Allan, 2022) and CERES (2001–2023) (Loeb et al., 2021) dataset
 926 are presented by dashed lines for comparison, with the trend adjusted to the IAP estimate
 927 to account for the arbitrary choice of integration constant. 95% Confidence Interval is
 928 presented assuming the independency of different budget components.

929



930

931 **Figure 19: Annual ocean heating rate compared with CERES data.** Both annual OHC
 932 and CERES EEI data are centred in June. The long-term mean is removed for all-time
 933 series.

934

935 3.9 Steric sea level and sea level budget

936 The updated IAPv4 data is used to assess the sea level budget for 1960-2023 in
 937 combination with other data, including IAP salinity data, glaciers, Greenland, Antarctic ice
 938 sheets mass loss from Frederikse et al. (2020) and altimetry sea level record (see Methods
 939 section for details). From 1960 to 2023, the observed GMSL rise is $2.07 \pm 0.55 \text{ mm yr}^{-1}$
 940 (Frederikse et al., 2020), which is derived by combining tide-gauge observations with
 941 estimates of local vertical land motion from permanent Global Positioning System stations
 942 and the difference between tide-gauge and satellite altimetry observations (Frederikse et
 943 al., 2018). During the same period, the sum of contributors (Glaciers, Greenland and
 944 Antarctic ice sheets, land water storage, and steric sea level) yields a mean sea level rise of
 945 $1.87 \pm 0.42 \text{ mm yr}^{-1}$. Thus, the sea level budget can be closed within a 90% confidence
 946 interval. This updated estimate indicates that the steric sea level, Antarctic ice sheet,

删除了: ,

删除了: and

删除了: contributions

删除了: .

951 Greenland ice sheet, glaciers, and land water storage contribute to the total sea level with
952 47.3%, 8.6%, 18.0%, 29.1%, and -3.1%, respectively for 1960-2023.

953 To isolate the contribution of the IAPv4 to the sea level budget, we replace the steric
954 sea level estimate in Frederikse et al., (2020) with IAPv4 and re-assess the sea level budget
955 for 1960-2018, 1993-2018 and 2005-2018, and the other components are identical to
956 Frederikse et al. (2020). Two metrics are used to quantify the performance of sea level
957 budget closure: the mean residual error and the **temporal** root mean square error (RMSD)
958 between the observed GMSL and the sum of contributions. We find that the residual sea
959 level budget based on IAPv4 is 0.20 ± 0.53 , 0.11 ± 0.34 , 0.47 ± 0.56 mm yr⁻¹ for 1960-
960 2018, 1993-2018 and 2005-2018, respectively. These mean residual errors are all smaller
961 than presented in Frederikse et al., (2020), which showed a residual error of 0.29 ± 0.57 ,
962 0.20 ± 0.34 and 0.54 ± 0.58 mm yr⁻¹ for 1960-2018, 1993-2018 and 2005-2018,
963 respectively. The RMSD using IAPv4 (or using steric sea level in Frederikse et al., 2020)
964 is 5.59 (5.35), 4.89 (5.33) and 4.21 (4.51) mm for the above-mentioned three periods,
965 respectively. Therefore, both metrics show that IAPv4 data improves the sea level budget
966 in three typical periods.


967 A similar test is done with the IPCC-AR6 sea level budget estimate (Gulev et al.,
968 2021): the thermosteric sea level estimate in IPCC-AR6 is replaced by IAPv4, and the sea
969 level budget is re-assessed for 1993-2018. IAPv4 suggests a larger thermosteric sea level
970 rise of 1.43 ± 0.16 for 1993-2018 than IPCC (1.31 ± 0.36 mm yr⁻¹) from 1993-2018.
971 Replacing the thermosteric sea level estimate by IAPv4 reduces the mean residual error
972 from 0.40 ± 0.57 to 0.28 ± 0.48 mm yr⁻¹. This suggests ~~that the stronger warming since the~~
973 1993 revealed by IAPv4 **than assessed in IPCC-AR6 (Gulve et al., 2021)**, seems more
974 realistic.

975 After 2002, the GRACE satellite supported the direct observation of barystatic sea
976 level, which is the sum of the sea level change due to the land water storage, Antarctica ice
977 sheet, Greenland ice sheet, and glaciers. The sea level budget can be obtained by
978 comparing altimetry-based GMSL with the barystatic sea level observed by GRACE and
979 the steric sea level. It is evident that the sea level budget can be closed between 2002 and
980 2015 with ± 5 mm residual errors (Fig. 20b). However, after ~2015, the sum of steric and
981 barystatic sea level is smaller than the total sea level rise for all ocean temperature
982 products. Previous studies have attributed this misclosure to salinity data biases (Barnoud
983 et al., 2021), altimetry data errors (Barnoud et al., 2023), and GRACE data errors (Wang et

删除了: again

985 al., 2021). The steric sea level inferred from IAPv4 showed a lower residual (~5 mm)
986 between 2005–2023 than ISH and EN4 data (10–20 mm), indicating that the temperature
987 data might be partly responsible for lack of closure of sea level budget since ~2015. This
988 suggests [again](#) that the stronger warming in recent years, as indicated by IAPv4, is more
989 realistic. As discussed in Section 3.4, the QC is mainly responsible for the increased
990 warming of IAPv4 compared with IAPv3 since ~2015 (Fig. 11).

991 Many traditional QC procedures use a static climatological range check to filter out
992 outliers, which does not account for the increase of extreme events with climate change
993 and removes too many extreme (positive) values during the recent period. Thus, we
994 strongly recommend that data product generation groups revisit the QC procedure.
995 Furthermore, as the stronger long-term OHC trends since ~1960 in IAPv4 than in IAPv3
996 are mainly attributed to the bias corrections for Nansen Bottle, MBT, and XBT data, it is
997 also recommended that international groups to revisit the biases in ocean observations.
998



999

1000 **Figure 20: (a) The sea level budget from 1960 to 2023.** Observed global mean sea level
 1001 for 1960–2023 and the individual contributions from land water storage, Antarctica,
 1002 Greenland and Glaciers (Frederikse et al., 2020). The budget is relative to a 1960 baseline.
 1003 Here, the land water storage and Glaciers data are through 2018, and a linear extrapolation
 1004 is made for 2019–2023. Antarctica ice sheet and Greenland ice sheet changes are estimated
 1005 by GRACE after 2018. Tide gauge after 2018 are updated by altimetry. Altimetry sea level
 1006 is shown in red dashed line for comparison. (b) Sea level budget residual time series since
 1007 2005. The residual of GMSL minus barystatic and steric sea level. The seasonal cycle is
 1008 reduced based on 2005–2015 climatology. 6-month running means are shown here to
 1009 reduce the noise.

删除了: .A

删除了: smooth

删除了: is applied

1013

1014 **4. Data availability**

1015 IAPv4 global ocean temperature product is available at
1016 <http://dx.doi.org/10.12157/IOCAS.20240117.002> (Cheng et al., 2024a) and
1017 <http://www.ocean.iap.ac.cn/>.

1018 IAPv4 global ocean heat content product is available at
1019 <http://dx.doi.org/10.12157/IOCAS.20240117.001> (Cheng et al., 2024b) and
1020 <http://www.ocean.iap.ac.cn/>.

1021 The code used in this paper includes data quality control, and the resultant dataset is
1022 available at <http://www.ocean.iap.ac.cn/>.

1023

1024 The data used in this study (but not generated by this work) are listed below. IAP data are
1025 available at <http://www.ocean.iap.ac.cn/>. The NCEI/NOAA data are available at
1026 (<https://www.ncei.noaa.gov/products/climate-data-records/global-ocean-heat-content>). ISH
1027 data from (<https://climate.mri-jma.go.jp/pub/ocean/ts/v7.2/>). The EN4 data
1028 (<https://www.metoffice.gov.uk/hadobs/en4/index.html>) For SST: ERSSTv5
1029 (<https://www1.ncdc.noaa.gov/pub/data/cmb/ersst/v5/netcdf/>); COBE2
1030 (<https://psl.noaa.gov/data/gridded/data.cobe2.html>); and HadSST3
1031 (<https://www.metoffice.gov.uk/hadobs/hadsst3/data/download.html>). For sea level data:
1032 AVISO+ GMSL ([https://www.aviso.altimetry.fr/en/data/products/ocean-indicators-](https://www.aviso.altimetry.fr/en/data/products/ocean-indicators-products/mean-sea-level.html#c15723)
1033 [products/mean-sea-level.html#c15723](https://www.aviso.altimetry.fr/en/data/products/ocean-indicators-products/mean-sea-level.html#c15723)), JPL GRACE ([https://grace.jpl.nasa.gov/data/get-](https://grace.jpl.nasa.gov/data/get-data/jpl_global_mascons/)
1034 [data/jpl_global_mascons/](https://grace.jpl.nasa.gov/data/get-data/jpl_global_mascons/)), the data in Frederikse et al., (2020) from
1035 (<https://zenodo.org/records/3862995>). The data in von Schuckmann et al., (2023)
1036 (https://www.wdc-climate.de/ui/entry?acronym=GCOS_EHI_1960-2020). Argo data were
1037 collected and made freely available by the International Argo Program and the national
1038 programs that contribute to it (<https://argo.ucsd.edu>, <https://www.ocean-ops.org>). DEEP-C
1039 data from <https://doi.org/10.17864/1947.000347>; CERES data ([tool.larc.nasa.gov/ord-tool/jsp/EBAFTOA41Selection.jsp](https://ceres-
1040 <a href=)); **GIOMAS ice volume data**
1041 **from (https://psc.apl.washington.edu/zhang/Global_seaice/data.html)**, SCRIPPS data from
1042 (http://sio-argo.ucsd.edu/RG_Climatology.html); BOA data from
1043 (<https://argo.ucsd.edu/data/argo-data-products/>).

1044

删除了：PIOMAS

删除了： <http://psc.apl.uw.edu/research/projects/arctic-sea-ice-volumeanomaly/>

1048 **5. Summary and Discussion**

1049 This paper introduces a new version of the ocean temperature and heat content
1050 gridded products and describes the data source and data processing techniques in detail.
1051 The key technical advances include the new QC, new or updated XBT/MBT/Bottle/APB
1052 bias corrections, new ocean temperature climatology, improved mapping approach, and
1053 grid-cell ocean volume corrections. These data and technical advances allow a better
1054 estimate of long-term ocean temperature and heat content changes since the mid-1950s
1055 from the sea surface down to 2000 m. We show that the new data product could better
1056 close the sea level and energy budgets than IAPv3. For rates of change, compared with
1057 CERES, the IAPv4 also shows a better correlation from 2005 to 2023 than IAPv3.

1058 Despite several marked improvements, issues needing further investigation remain.
1059 Although inter-annual and decadal-scale changes of satellite-based EEI and observational
1060 OHC are generally consistent, a mismatch remains between EEI and OHC for their month-
1061 to-month variation, as the monthly variation of OHC is still much larger than implied by
1062 EEI. There are several possibilities, in our opinion: first, there is substantial heat storage
1063 and release for land and ice monthly, which needs to be accurately quantified; second, the
1064 accuracy of OHC estimate on a monthly basis still needs to be improved for month-to-
1065 month variation because of the limited data coverage; third, the EEI observed by CERES
1066 also suffers from sampling biases on a monthly basis (Loeb et al., 2009). Thus, a better
1067 understanding of the monthly variation of OHC and EEI is still a research priority. Besides,
1068 the failure to close the 2015-2023 sea level budget indicates that the underlying data still
1069 has bias problems, which need to be explored and resolved.

1070 Second, the application of CODC-QC in IAPv4 leads to a stronger ocean warming
1071 rate in the past decade than WOD-QC used in IAPv3 because WOD-QC removes more
1072 positive temperature anomalies than CODC-QC. This could imply that the rate of increase
1073 in OHC is still slightly underestimated and deserves an in-depth investigation. Several
1074 fundamental questions must be answered: first, are there still real temperature extremes
1075 being removed by CODC-QC, such as in small warm/cold eddies? Are the extremes well
1076 sampled by the current observation system? If not, what is the impact? Moreover, it is clear
1077 that the high latitudes where sea ice occurs are not well sampled and need more attention.

1078 Third, during the development of the data product, we discovered that much metadata
1079 relating to the profiles in the World Ocean Database is missing and that much existing
1080 metadata is incorrect, also giving rise to duplicate profiles, putting a strain on the overall

1081 quality of a database of oceanic observations. More than ever, long-term concerted efforts
1082 are needed to eliminate duplicate profiles and identify and correct missing metadata using
1083 statistical methods, expert control, or machine learning techniques. For example, the
1084 International Quality-Controlled Database (IQuOD) group is coordinating some activities
1085 related to data processing techniques, uncertainty quantification, and improving the overall
1086 quality of ocean data (Cowley et al., 2021).

1087 Fourth, the deep ocean changes below 2000 m are estimated based on the currently
1088 available data, including data from hydrological sections and Deep-Argo. IAP mapping
1089 technique is applied. Because of the lack of independent observations with global ocean
1090 coverage, evaluating the deep ocean change estimate is still dicey. Thus, the below-2000 m
1091 estimate should be used with caution, as also indicated in previous estimates (Purkey and
1092 Johnson, 2010; Desbruyères et al., 2017; Good et al., 2013). A community-agreed
1093 evaluation approach for the deep ocean changes is critically needed. Besides, other
1094 mapping techniques deserving further investigation include interpolation on isopycnal
1095 surfaces (Palmer and Haines, 2009).

1096 Furthermore, the quantification of uncertainty for *in situ* measurements, gridded
1097 T/OHC values, and the global OHC estimates need to be improved. IAPv4 only accounts
1098 for the instrumental error and sampling/mapping error. In the future, comprehensive
1099 quantification of other uncertainty sources will be made, including the choice of
1100 climatology, vertical interpolation, XBT/MBT/APB/Bottle corrections, etc. It is also
1101 necessary to analyze the correlation between these error sources. This also helps to
1102 understand regions with larger uncertainty for OHC estimates, which supports the design
1103 of the global ocean observing system.

1104

1105 **Author contributions.** L.C. has worked on this study's conceptualization, coordination,
1106 methodologies, and writing the manuscript. Z.T. worked on *in situ* observation collections,
1107 metadata format, and the automated quality control procedure (CODC-QC) development.
1108 Y.P. has worked on calculating and comparing the OHC annual cycle, the mixed layer
1109 depth, and the MHT among different data sets. V.G. worked on bias correction schemes for
1110 MBT, APB, and bottle data and on developing the automated quality control procedure.
1111 H.Y. worked on the analysis of inter-annual variability. J.D. has worked on OHC trend
1112 calculation and analysis. G.L. worked on SST calculation and its analysis. H. Z. worked on
1113 global energy and sea level budget calculations and analyses. Y.L. and Y.J. worked on the

1114 volume correction. All authors have contributed to formal analysis, data validation, and
1115 editing of the original draft.

1116

1117 **Acknowledgement and Funding.** The IAP/CAS analysis is supported by the National
1118 Natural Science Foundation of China (Grant no. 42122046, 42076202, 42206208,
1119 42261134536), Strategic Priority Research Program of the Chinese Academy of Sciences
1120 (Grant no. XDB42040402), the new Cornerstone Science Foundation through the
1121 XPLOER PRIZE, DAMO Academy Young Fellow, Youth Innovation Promotion
1122 Association, Chinese Academy of Sciences, National Key Scientific and Technological
1123 Infrastructure project “Earth System Science Numerical Simulator Facility” (EarthLab),
1124 the Young Talent Support Project of Guangzhou Association for Science and Technology.
1125 The calculations in this study were carried out on the ORISE Supercomputer. Some data
1126 were collected onboard of R/V Shiyan 6 implementing the Open Research Cruise
1127 NORC2022-10+NORC2022-303 supported by NSFC Shiptime Sharing Projects
1128 42149910. NCAR is sponsored by the US National Science Foundation. We acknowledge
1129 the World Climate Research Programme’s Working Group on Coupled Modelling, which
1130 is responsible for CMIP, and we thank the climate modeling groups for producing and
1131 making their model output available through the Earth System Grid Federation. The Argo
1132 Program is part of the Global Ocean Observing System.

1133

1134 **Competing interests.** The contact author has declared that none of the authors has any
1135 competing interests.

1136

1137 **References**

- 1138 Abraham, J. P., Cheng, L., Mann, M. E., Trenberth, K., and von Schuckmann, K.: The
1139 ocean response to climate change guides both adaptation and mitigation efforts.
1140 Atmospheric and Oceanic Science Letters, 15, 100221,
1141 <https://doi.org/10.1016/j.aosl.2022.100221>, 2022.
- 1142 Abraham, J. P., and Cheng, L.: Intersection of Climate Change, Energy, and Adaptation.
1143 Energies. 15(16), 5886; <https://doi.org/10.3390/en15165886>, 2022.
- 1144 Abraham, J. P., Baringer, M., Bindoff, N. L., Boyer, T., Cheng, L. J., Church, J. A.,
1145 Conroy, J. L., Domingues, C. M., Fasullo, J. T., Gilson, J., Goni, G., Good, S. A.,

1146 Gorman, J. M., Gouretski, V., Ishii, M., Johnson, G. C., Kizu, S., Lyman, J. M.,
1147 Macdonald, A. M., Minkowycz, W. J., Moffitt, S. E., Palmer, M. D., Piola, A. R.,
1148 Reseghetti, F., Schuckmann, K., Trenberth, K. E., Velicogna, I., and Willis, J. K.: A
1149 review of global ocean temperature observations: Implications for ocean heat content
1150 estimates and climate change, *Rev. Geophys.*, 51, 450–483,
1151 <https://doi.org/10.1002/rog.20022>, 2013.

1152 Argo: Argo float data and metadata from Global Data Assembly Centre (Argo GDAC).
1153 SEANOE, 2000.

1154 Bagnell, A., and DeVries, T.: 20(th) century cooling of the deep ocean contributed to
1155 delayed acceleration of Earth's energy imbalance. *Nat. Comm.*, 12, 4604,
1156 <https://doi.org/10.1038/s41467-021-24472-3>, 2021.

1157 Barker, P. M., and McDougall, T. J.: Two Interpolation Methods Using Multiply-Rotated
1158 Piecewise Cubic Hermite Interpolating Polynomials. *J. Atmos. Ocean Technol.*, 37,
1159 605-619, <https://doi.org/10.1175/JTECH-D-19-0211.1>, 2020.

1160 Barnoud, A., Pfeffer, J., Cazenave, A., Fraudeau, R., Rousseau, V., and Ablain, M.:
1161 Revisiting the global mean ocean mass budget over 2005–2020. *Ocean Science*, 19,
1162 321-334, <https://doi.org/10.5194/os-19-321-2023>, 2023.

1163 Barnoud, A., Pfeffer, J., Guérou, A., Frery, M.-L., Siméon, M., Cazenave, A., Chen, J.,
1164 Llovel, W., Thierry, V., Legeais, J.-F., and Ablain, M.: Contributions of Altimetry
1165 and Argo to Non-Closure of the Global Mean Sea Level Budget Since 2016.
1166 *Geophys Res Lett*, 48, e2021GL092824, <https://doi.org/10.1029/2021GL092824>,
1167 2021.

1168 Bindoff, N. L., Cheung, W. W. L., Kairo, J. G., Aristegui, J., Guinder, V. A., Hallberg, R.,
1169 Hilmi, N., Jiao, N., and Karim, M. S.: Changing Ocean, Marine Ecosystems, and
1170 Dependent Communities. In *IPCC Special Report on the Ocean and Cryosphere in a
1171 Changing Climate* [H.-O. Pörtner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, M.
1172 Tignor, E. Poloczanska, K. Mintenbeck, A. Alegria, M. Nicolai, A. Okem, J. Petzold,
1173 B. Rama, N.M. Weyer (eds.)], Cambridge University Press, Cambridge, UK and
1174 New York, NY, USA, pp. 447–587. <https://doi.org/10.1017/9781009157964.007>,
1175 2019.

1176 Boyer, T., Domingues, C. M., Good, S. A., Johnson, G. C., Lyman, J. M., Ishii, M.,
1177 Gouretski, V., Willis, J. K., Antonov, J., Wijffels, S., Church, J. A., Cowley, R., and
1178 Bindoff, N. L.: Sensitivity of Global Upper Ocean Heat Content Estimates to

1179 Mapping Methods, XBT Bias Corrections, and Baseline Climatologies, *J. Climate*,
1180 29, 4817–4842, <https://doi.org/10.1175/JCLI-D-15-0801.1>, 2016.

1181 Caesar, L., Rahmstorf, S., Robinson, A., Feulner, G., and Saba, V.: Observed fingerprint of
1182 a weakening Atlantic Ocean overturning circulation. *Nature*, 556, 191-196,
1183 <https://doi.org/10.1038/s41586-018-0006-5>, 2018.

1184 Cane, M. A., and Zebiak, S. E.: A theory for El Niño and the southern oscillation. *Science*,
1185 228, 1085-1087, <https://doi.org/10.1126/science.228.4703.1085>, 1985.

1186 Cheng, L.: Sensitivity of Ocean Heat Content to Various Instrumental Platforms in Global
1187 Ocean Observing System. *Ocean-Land-Atmosphere Research*, 0,
1188 <https://doi.org/10.34133/olar.0037>, 2024a.

1189 Cheng, L., and Zhu, J.: Uncertainties of the Ocean Heat Content Estimation Induced by
1190 Insufficient Vertical Resolution of Historical Ocean Subsurface Observations. *J.*
1191 *Atmos. Ocean Technol.*, 31, 1383-1396, [https://doi.org/10.1175/JTECH-D-13-](https://doi.org/10.1175/JTECH-D-13-00220.1)
1192 [00220.1](https://doi.org/10.1175/JTECH-D-13-00220.1), 2014.

1193 Cheng, L., and Zhu, J.: Influences of the Choice of Climatology on Ocean Heat Content
1194 Estimation. *J. Atmos. Ocean Technol.*, 32, 388-394, [https://doi.org/10.1175/JTECH-](https://doi.org/10.1175/JTECH-D-14-00169.1)
1195 [D-14-00169.1](https://doi.org/10.1175/JTECH-D-14-00169.1), 2015.

1196 Cheng, L., and Zhu, J.: Benefits of CMIP5 Multimodel Ensemble in Reconstructing
1197 Historical Ocean Subsurface Temperature Variations. *J. Climate*, 29, 5393-5416,
1198 <https://doi.org/10.1175/JCLI-D-15-0730.1>, 2016.

1199 Cheng, L., Zhu, J., Cowley, R., Boyer, T., and Wijffels, S.: Time, Probe Type, and
1200 Temperature Variable Bias Corrections to Historical Expendable Bathythermograph
1201 Observations. *J. Atmos. Ocean. Technol.*, 31, 1793-1825,
1202 <https://doi.org/10.1175/jtech-d-13-00197.1>, 2014.

1203 Cheng, L., Abraham, J., Goni, G., Boyer, T., Wijffels, S., Cowley, R., Gouretski, V.,
1204 Reseghetti, F., Kizu, S., Dong, S., Bringas, F., Goes, M., Houpert, L., Sprintall, J.,
1205 and Zhu, J.: XBT Science: Assessment of Instrumental Biases and Errors, *B. Am.*
1206 *Meteorol. Soc.*, 97, 924–933, <https://doi.org/10.1175/BAMS-D-15-00031.1>, 2016.

1207 Cheng, L., Trenberth, K. E., Fasullo, J., Boyer, T., Abraham, J., and Zhu, J.: Improved
1208 estimates of ocean heat content from 1960 to 2015. *Sci. Adv.*, 3, e1601545,
1209 <https://doi.org/10.1126/sciadv.1601545>, 2017.

1210 Cheng, L., Trenberth, K. E., Fasullo, J. T., Mayer, M., Balmaseda, M., and Zhu, J.:
1211 Evolution of Ocean Heat Content Related to ENSO. *J. Climate*, 32, 3529-3556,
1212 <https://doi.org/10.1175/jcli-d-18-0607.1>, 2019.

1213 Cheng, L., Trenberth, K. E., Gruber, N., Abraham, J. P., Fasullo, J. T., Li, G., Mann, M. E.,
1214 Zhao, X., and Zhu, J.: Improved Estimates of Changes in Upper Ocean Salinity and
1215 the Hydrological Cycle. *J. Climate*, 33, 10357-10381, <https://doi.org/10.1175/JCLI->
1216 [D-20-0366.1](https://doi.org/10.1175/JCLI-D-20-0366.1), 2020.

1217 Cheng, L., von Schuckmann, K., Abraham, J. P., Trenberth, K. E., Mann, M. E., Zanna, L.,
1218 England, M. H., Zika, J. D., Fasullo, J. T., Yu, Y., Pan, Y., Zhu, J., Newsom, E. R.,
1219 Bronselaer, B., and Lin, X.: Past and future ocean warming. *Nat. Rev. Earth Env.*, 3,
1220 776-794, <https://doi.org/10.1038/s43017-022-00345-1>, 2022a.

1221 Cheng, L., Foster, G., Hausfather, Z., Trenberth, K. E., and Abraham, J.: Improved
1222 quantification of the rate of ocean warming. *J. Climate*, 35, 4827-4840,
1223 <https://doi.org/10.1175/jcli-d-20-0366.1>, 2022b.

1224 Cheng, L., Tan, Z., Pan, Y., Zheng, H., Zhu, Y., Wei, W., Du, J., Li, G., Ye, H., Gourteski,
1225 V.: IAP temperature 1° gridded analysis product (IAPv4),
1226 <http://dx.doi.org/10.12157/IOCAS.20240117.002>, 2024a.

1227 Cheng, L., Tan, Z., Pan, Y., Zheng, H., Zhu, Y., Wei, W., Du, J., Li, G., Ye, H., Gourteski,
1228 V.: IAP global ocean heat content 1° gridded analysis product (IAPv4),
1229 <http://dx.doi.org/10.12157/IOCAS.20240117.001>, 2024b.

1230 Chu, P. C., and Fan, C.: Global climatological data of ocean thermohaline parameters
1231 derived from WOA18. *Scientific Data*, 10, 408, <https://doi.org/10.1038/s41597-023->
1232 [02308-7](https://doi.org/10.1038/s41597-023-02308-7), 2023.

1233 Comiso, J. C., Meier, W. N., and Gersten, R.: Variability and trends in the Arctic Sea ice
1234 cover: Results from different techniques. *J. Geophys. Res.- Oceans*, 122, 6883-6900,
1235 <https://doi.org/10.1002/2017JC012768>, 2017.

1236 Cowley, R., Killick, R. E., Boyer, T., Gouretski, V., Reseghetti, F., Kizu, S., Palmer, M.
1237 D., Cheng, L., Storto, A., Le Menn, M., Simoncelli, S., Macdonald, A. M., &
1238 Domingues, C. M.: International Quality-Controlled Ocean Database (iQuOD) v0.1:
1239 The Temperature Uncertainty Specification. *Front. Mar. Sci.*, 8,
1240 <https://doi.org/10.3389/fmars.2021.689695>, 2021.

1241 Dangendorf, S., Frederikse, T., Chafik, L., Klinck, J. M., Ezer, T., and Hamlington, B. D.:
1242 Data-driven reconstruction reveals large-scale ocean circulation control on coastal
1243 sea level. *Nat. Clim. Change*, 11, 514-520, <https://doi.org/10.1038/s41558-021->
1244 [01046-1](https://doi.org/10.1038/s41558-021-01046-1), 2021.

1245 de Boyer Montégut, C., Madec, G., Fischer, A. S., Lazar, A., and Iudicone, D.: Mixed
1246 layer depth over the global ocean: An examination of profile data and a profile-based

1247 climatology. *J. Geophys. Res.- Oceans*, 109, <https://doi.org/10.1029/2004JC002378>,
1248 2004.

1249 Desbruyères, D., McDonagh, E. L., King, B. A. & Thierry, V. Global and full-depth ocean
1250 temperature trends during the early twenty-first century from Argo and repeat
1251 hydrography. *J. Clim.* 30, 1985–1997, 2017.

1252 England, M. H., McGregor, S., Spence, P., Meehl, G. A., Timmermann, A., Cai, W.,
1253 Gupta, A. S., McPhaden, M. J., Purich, A., & Santoso, A.: Recent intensification of
1254 wind-driven circulation in the Pacific and the ongoing warming hiatus. *Nat. Clim.*
1255 *Change*, 4, 222-227, <https://doi.org/10.1038/nclimate2106>, 2014.

1256 Fasullo, J. T., and Nerem, R. S.: Altimeter-era emergence of the patterns of forced sea-
1257 level rise in climate models and implications for the future. *P. Natl. Acad. Sci.*, 115,
1258 12944-12949, <https://doi.org/10.1073/pnas.1813233115>, 2018.

1259 [Frederikse, T., Jevrejeva, S., Riva, R. E. M. and Dangendorf, S.: A Consistent Sea-Level](#)
1260 [Reconstruction and Its Budget on Basin and Global Scales over 1958–2014. *J.*](#)
1261 [*Climate*, 31, 1267–1280, <https://doi.org/10.1175/JCLI-D-17-0502.1>, 2018.](#)

1262 Frederikse, T., Landerer, F., Caron, L., Adhikari, S., Parkes, D., Humphrey, V. W.,
1263 Dangendorf, S., Hogarth, P., Zanna, L., Cheng, L., and Wu, Y.-H.: The causes of sea
1264 level rise since 1900. *Nature*, 584, 393-397, <https://doi.org/10.1038/s41586-020->
1265 2591-3, 2020.

1266 Garcia, H. E., Boyer, T. P., Locarnini, R. A., Baranova, O. K., and Zweng, M. M.: *World*
1267 *Ocean Database 2018: User’s Manual.*, T. E. A.V. Mishonov, NOAA, Silver Spring,
1268 MD., Ed. , 2018

1269 Goni, G. J., Sprintall, J., Bringas, F., Cheng, L., Cirano, M., Dong, S., Domingues, R.,
1270 Goes, M., Lopez, H., Morrow, R., Rivero, U., Rossby, T., Todd, R. E., Trinanes, J.,
1271 Zilberman, N., Baringer, M., Boyer, T., Cowley, R., Domingues, Hutchinson, K.,
1272 Kramp, M., Mata, M. M., Reseghetti, F., Sun, C., Bhaskar Tvs U., Volkov, D.: More
1273 Than 50 Years of Successful Continuous Temperature Section Measurements by the
1274 Global Expendable Bathythermograph Network, Its Integrability, Societal Benefits,
1275 and Future. *Fron. Mar. Sci.*, 6, <http://dx.doi.org/10.3389/fmars.2019.00452>, 2019.

1276 Good, S. A., Martin, M. J., and Rayner, N. A.: EN4: Quality controlled ocean temperature
1277 and salinity profiles and monthly objective analyses with uncertainty estimates. *J.*
1278 *Geophys. Res. Oceans*, 118, 6704-6716, <https://doi.org/10.1002/2013jc009067>, 2013.

1279 Gouretski, V., and Koltermann, K. P.: How much is the ocean really warming? *Geophys.*
1280 *Res. Lett.*, 34, L01610, <https://doi.org/10.1029/2006GL027834>, 2007.

1281 Gouretski, V. and Reseghetti, F.: On depth and temperature biases in bathythermograph
1282 data: Development of a new correction scheme based on analysis of a global ocean
1283 database, *Deep Sea Res.*, 57, 6, 812-833, <https://doi.org/10.1016/j.dsr.2010.03.011>,
1284 2010.

1285 Gouretski, V., and Cheng, L.: Correction for Systematic Errors in the Global Dataset of
1286 Temperature Profiles from Mechanical Bathythermographs. *J. Atmos. Ocean.
1287 Technol.*, 37, 841-855, <https://doi.org/10.1175/jtech-d-19-0205.1>, 2020.

1288 Gouretski, V., Cheng, L., and Boyer, T.: On the Consistency of the Bottle and CTD Profile
1289 Data. *J. Atmos. Ocean Technol.*, 39, 1869-1887, [https://doi.org/10.1175/JTECH-D-
22-0004.1](https://doi.org/10.1175/JTECH-D-
1290 22-0004.1), 2022.

1291 Gouretski, V., Roquet, F., and Cheng, L.: Measurement biases in ocean temperature
1292 profiles from marine mammal data loggers. *J. Atmos. Ocean Technol.*, submitted,
1293 2024.

1294 Gouretski, V., Kennedy, J., Boyer, T., and Köhl, A.: Consistent near-surface ocean
1295 warming since 1900 in two largely independent observing networks. *Geophys. Res.
1296 Lett.*, 39, <https://doi.org/10.1029/2012GL052975>, 2012.

1297 Gouretski V, Koltermann K P. 2004. WOCE global hydrographic clima- tology. *Berichte
1298 des BSH*, 35: 1–52.

1299 Gulev, S. K., Thorne, P. W., Ahn, J., Dentener, F. J., Domingues, C. M., Gerland, S.,
1300 Gong, D., Kaufman, D. S., Nnamchi, H. C., Quaas, J., Rivera, J. A., Sathyendranath,
1301 S., Smith, S. L., Trewin, B., Schuckmann, K. von, and Vose, R. S.: Changing State
1302 of the Climate System Supplementary Material, in: *Climate Change 2021: The
1303 Physical Science Basis. Contribution of Working Group I to the Sixth Assessment
1304 Report of the Intergovernmental Panel on Climate Change*, edited by: Masson-
1305 Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N.,
1306 Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews,
1307 J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B.,
1308 Cambridge University Press, Cambridge, United Kingdom and New York, NY,
1309 USA, 287– 422, <https://doi.org/10.1017/9781009157896.004>, 2021.

1310 Hakuba, M. Z., Frederikse, T., and Landerer, F. W.: Earth’s Energy Imbalance From the
1311 Ocean Perspective (2005–2019), *Geophys. Res. Lett.*, 48, e2021GL093624,
1312 <https://doi.org/10.1029/2021GL093624>, 2021.

1313 Hansen, J., Sato, M., Kharecha, P., and von Schuckmann, K.: Earth's energy imbalance and
1314 implications. *Atmos. Chem. Phys.*, 11, 13421-13449, [https://doi.org/10.5194/acp-11-](https://doi.org/10.5194/acp-11-13421-2011)
1315 [13421-2011](https://doi.org/10.5194/acp-11-13421-2011), 2011.

1316 Hirahara, S., Ishii, M., and Fukuda, Y.: Centennial-Scale Sea Surface Temperature
1317 Analysis and Its Uncertainty. *J. Climate*, 27, 57-75, [https://doi.org/10.1175/JCLI-D-](https://doi.org/10.1175/JCLI-D-12-00837.1)
1318 [12-00837.1](https://doi.org/10.1175/JCLI-D-12-00837.1), 2014.

1319 Holte, J., Talley, L. D., Gilson, J., and Roemmich, D.: An Argo mixed layer climatology
1320 and database. *Geophys. Res. Lett.*, 44, 5618-5626,
1321 <https://doi.org/10.1002/2017GL073426>, 2017.

1322 Hosoda, S., Ohira, T., and Nakamura, T.: Monthly mean dataset of global oceanic
1323 temperature and salinity derived from Argo float observations. *JAMSTEC Report of*
1324 *Research and Development*, 8, <https://doi.org/10.5918/jamstecr.8.47>, 2008.

1325 Huang, B., Thorne, P. W., Banzon, V. F., Boyer, T., Chepurin, G., Lawrimore, J. H.,
1326 Menne, M. J., Smith, T. M., Vose, R. S., and Zhang, H.-M.: Extended Reconstructed
1327 Sea Surface Temperature, Version 5 (ERSSTv5): Upgrades, Validations, and
1328 Intercomparisons. *J. Climate*, 30, 8179-8205, [https://doi.org/10.1175/JCLI-D-16-](https://doi.org/10.1175/JCLI-D-16-0836.1)
1329 [0836.1](https://doi.org/10.1175/JCLI-D-16-0836.1), 2017.

1330 Hugonnet, R., McNabb, R., Berthier, E., Menounos, B., Nuth, C., Girod, L., Farinotti, D.,
1331 Huss, M., Dussaillant, I., Brun, F., and Kääb, A.: Accelerated global glacier mass
1332 loss in the early twenty-first century. *Nature*, 592, 726-731,
1333 <https://doi.org/10.1038/s41586-021-03436-z>, 2021.

1334 IPCC: Annex I: Observational Products [Trewin, B. (ed.)], pp. 2061–2086 pp. , 2021

1335 Ishii, M., and Kimoto, M.: Reevaluation of historical ocean heat content variations with
1336 time-varying XBT and MBT depth bias corrections. *J. Oceanogr.*, 65, 287-299,
1337 <https://doi.org/10.1007/s10872-009-0027-7>, 2009.

1338 Ishii, M., Shouji, A., Sugimoto, S., and Matsumoto, T.: Objective analyses of sea-surface
1339 temperature and marine meteorological variables for the 2⁰th century using ICOADS
1340 and the Kobe Collection. *Int. J. Climatol.*, 25, 865-879,
1341 <https://doi.org/10.1002/joc.1169>, 2005.

1342 Ishii, M., Y. Fukuda, S. Hirahara, S. Yasui, T. Suzuki, and K. Sato: Accuracy of Global
1343 Upper Ocean Heat Content Estimation Expected from Present Observational Data
1344 Sets. *Sola*, 13, 163-167, <https://doi.org/10.2151/sola.2017-030>, 2017.

1345 Jin, F.-F.: An Equatorial Ocean Recharge Paradigm for ENSO. Part I: Conceptual Model.
1346 J. Atmos. Sci., 54, 811-829, <https://doi.org/10.1175/1520->
1347 [0469\(1997\)054%3C0811:AEORPF%3E2.0.CO;2](https://doi.org/10.1175/1520-0469(1997)054%3C0811:AEORPF%3E2.0.CO;2), 1997.

1348 Jin, Y., Li, Y., Cheng, L., Duan, J., Li, R., & Wang, F.: Ocean heat content increase of the
1349 Maritime Continent since the 1990s. Geophysical Research Letters, 51,
1350 e2023GL107526. <https://doi.org/10.1029/2023GL107526>, 2024.

1351 Johns, W. E., Elipot, S., Smeed, D. A., Moat, B., King, B., Volkov, D. L., and Smith, R. H.:
1352 Towards two decades of Atlantic Ocean mass and heat transports at 26.5° N.
1353 Philosophical Transactions of the Royal Society A: Mathematical, Physical and
1354 Engineering Sciences, 381, 20220188, 2023.

1355 Johnson, G. C., Purkey, S. G., Zilberman, N. V., and Roemmich, D.: Deep Argo Quantifies
1356 Bottom Water Warming Rates in the Southwest Pacific Basin. Geophys. Res. Lett.,
1357 46, 2662-2669, <https://doi.org/10.1098/rsta.2022.0188>, 2019.

1358 Katsumata, K., Purkey, S. G., Cowley, R., Sloyan, B. M., Diggs, S. C., Moore, T. S.,
1359 Talley, L. D., and Swift, J. H.: GO-SHIP Easy Ocean: Gridded ship-based
1360 hydrographic section of temperature, salinity, and dissolved oxygen. Scientific Data,
1361 9, 103, <https://doi.org/10.1038/s41597-022-01212-w>, 2022.

1362 Kennedy, J.: A review of uncertainty in in situ measurements and data sets of sea surface
1363 temperature. Rev. Geophys., 52, 1-32, <https://doi.org/10.1002/2013RG000434>, 2014.

1364 Levitus, S., Antonov, J. I., Boyer, T. P., Locarnini, R. A., Garcia, H. E., and Mishonov, A.
1365 V.: Global ocean heat content 1955–2008 in light of recently revealed
1366 instrumentation problems. Geophys. Res. Lett., 36,
1367 <https://doi.org/10.1029/2008GL037155>, 2009

1368 Levitus, S., Antonov, J. I., Boyer, T. P., Baranova, O. K., Garcia, H. E., Locarnini, R. A.,
1369 Mishonov, A. V., Reagan, J. R., Seidov, D., Yarosh, E. S., and Zweng, M. M.: World
1370 ocean heat content and thermosteric sea level change (0–2000 m), 1955–2010,
1371 Geophys. Res. Lett., 39, L10603, <https://doi.org/10.1029/2012GL051106>, 2012.

1372 Li, G., Cheng, L., Zhu, J., Trenberth, K. E., Mann, M. E., and Abraham, J. P.: Increasing
1373 ocean stratification over the past half-century. Nat. Clim. Change, 10, 1116-1123,
1374 <https://doi.org/10.1038/s41558-020-00918-2>, 2020.

1375 Li, H., Xu, F., Zhou, W., Wang, D., Wright, J. S., Liu, Z., and Lin, Y.: Development of a
1376 global gridded Argo data set with Barnes successive corrections. J. Geophys. Res.
1377 Oceans, 122, 866-889, <https://doi.org/10.1002/2016JC012285>, 2017.

1378 Li, Y., Church, J. A., McDougall, T. J., and Barker, P. M.: Sensitivity of Observationally
1379 Based Estimates of Ocean Heat Content and Thermal Expansion to Vertical
1380 Interpolation Schemes. *Geophys. Res. Lett.*, 49, e2022GL101079,
1381 <https://doi.org/10.1029/2022GL101079>, 2022.

1382 Lian, T., Wang, J., Chen, D., Liu, T. and Wang, D.: A Strong 2023/24 El Niño is Staged by
1383 Tropical Pacific Ocean Heat Content Buildup. *Ocean-Land-Atmosphere Research*, 2,
1384 0011, <https://doi.org/10.34133/olar.0011>, 2023.

1385 Liu, C., and Allan, R.: Reconstructions of the radiation fluxes at the top of atmosphere and
1386 net surface energy flux: DEEP-C version 5.0. . a. A. University of Reading Dataset,
1387 Ed., <https://doi.org/10.17864/1947.000347>, 2022

1388 Liu, C., Allan, R. P., Mayer, M., Hyder, P., Loeb, N. G., Roberts, C. D., Valdivieso, M.,
1389 Edwards, J. M., and Vidale, P.-L.: Evaluation of satellite and reanalysis-based global
1390 net surface energy flux and uncertainty estimates. *J. Geophys. Res.- Atmospheres*,
1391 122, 6250-6272, <https://doi.org/10.1002/2017JD026616>, 2017.

1392 Liu, C., Allan, R. P., Mayer, M., Hyder, P., Desbruyères, D., Cheng, L., Xu, J., Xu, F., and
1393 Zhang, Y.: Variability in the global energy budget and transports 1985–2017, *Clim.*
1394 *Dynam.*, 55, 3381– 3396, <https://doi.org/10.1007/s00382-020-05451-8>, 2020.

1395 Loeb, N. G., B. A. Wielicki, D. R. Doelling, G. L. Smith, D. F. Keyes, S. Kato, N. Manalo-
1396 Smith, and T. Wong: Toward Optimal Closure of the Earth's Top-of-Atmosphere
1397 Radiation Budget. *J. Climate*, 22, 748–766, <https://doi.org/10.1175/2008JCLI2637.1>,
1398 2009.

1399 Loeb, N. G., Johnson, G. C., Thorsen, T. J., Lyman, J. M., Rose, F. G., and Kato, S.:
1400 Satellite and Ocean Data Reveal Marked Increase in Earth's Heating Rate. *Geophys.*
1401 *Res. Lett.*, 48, <https://doi.org/10.1029/2021gl093047>, 2021.

1402 Loeb, N. G., Thorsen, T. J., Norris, J. R., Wang, H., and Su, W.: Changes in Earth's energy
1403 budget during and after the “Pause” in global warming: An observational
1404 perspective, *Climate*, 6, 62, <https://doi.org/10.3390/cli6030062>, 2018.

1405 Lyman, J. M., and Johnson, G. C.: Estimating Global Ocean Heat Content Changes in the
1406 Upper 1800 m since 1950 and the Influence of Climatology Choice. *J. Climate*, 27,
1407 1945-1957, <https://doi.org/10.1175/JCLI-D-12-00752.1>, 2014.

1408 Lyman, J. M., Good, S. A., Gouretski, V. V., Ishii, M., Johnson, G. C., Palmer, M. D.,
1409 Smith, D. M., and Willis, J. K.: Robust warming of the global upper ocean. *Nature*,
1410 465, 334-337, <https://doi.org/10.1038/nature09043>, 2010.

1411 Lyman, J. M., and G. C. Johnson, 2023: Global High-Resolution Random Forest
1412 Regression Maps of Ocean Heat Content Anomalies Using In Situ and Satellite Data.
1413 J. Atmos. Oceanic Technol., 40, 575–586, [https://doi.org/10.1175/JTECH-D-22-](https://doi.org/10.1175/JTECH-D-22-0058.1)
1414 [0058.1](https://doi.org/10.1175/JTECH-D-22-0058.1).

1415 Mann, M.E., Beyond the Hockey Stick: Climate Lessons from The Common Era, Proc.
1416 Natl. Acad. Sci., 118 (39) e2112797118, <https://doi.org/10.1073/pnas.2112797118>,
1417 2021.

1418 McDougall T. J. and P. M. Barker, 2011: Getting started with TEOS-10 and the Gibbs
1419 Seawater (GSW) Oceanographic Toolbox, 28pp., SCOR/IAPSO WG127, ISBN 978-
1420 0-646-55621-5.

1421 Mayer, J., Mayer, M., and Haimberger, L.: Consistency and Homogeneity of Atmospheric
1422 Energy, Moisture, and Mass Budgets in ERA5. J. Climate, 34, 3955-3974,
1423 <https://doi.org/10.1175/JCLI-D-20-0676.1>, 2021.

1424 Mayer, M., Alonso Balmaseda, M., and Haimberger, L.: Unprecedented 2015/2016 Indo-
1425 Pacific Heat Transfer Speeds Up Tropical Pacific Heat Recharge. Geophys. Res.
1426 Lett., 45, 3274-3284, <https://doi.org/10.1002/2018GL077106>, 2018.

1427 [Mayer, M., S. Tietsche, L. Haimberger, T. Tsubouchi, J. Mayer, and H. Zuo, 2019: An](https://doi.org/10.1175/JCLI-D-19-0233.1)
1428 [Improved Estimate of the Coupled Arctic Energy Budget. J. Climate, 32, 7915–7934,](https://doi.org/10.1175/JCLI-D-19-0233.1)
1429 <https://doi.org/10.1175/JCLI-D-19-0233.1>.

1430 McPhaden, M. J.: A 2^{1st} century shift in the relationship between enso sst and warm water
1431 volume anomalies. Geophys. Res. Lett., 39, 9706,
1432 <https://doi.org/10.1029/2012GL051826>, 2012.

1433 McMahon, C. R., Roquet, F., Baudel, S., Belbeoch, M., Bestley, S., Blight, C., Boehme,
1434 L., Carse, F., Costa, D. P., Fedak, M. A., Guinet, C., Harcourt, R., Heslop, E.,
1435 Hindell, M. A., Hoenner, X., Holland, K., Holland, M., Jaine, F. R. A., Jeanniard du
1436 Dot, T., . . . Woodward: Animal Borne Ocean Sensors – AniBOS – An Essential
1437 Component of the Global Ocean Observing System. Front. Mar. Sci., 8,
1438 <https://doi.org/10.3389/fmars.2021.751840>, 2021.

1439 Meyssignac, B., Boyer, T., Zhao, Z., Hakuba, M. Z., Landerer, F. W., Stammer, D., Köhl,
1440 A., Kato, S., L’Ecuyer, T., Ablain, M., Abraham, J. P., Blazquez, A., Cazenave, A.,
1441 Church, J. A., Cowley, R., Cheng, L., Domingues, C. M., Giglio, D., Gouretski, V.,
1442 Ishii, M., Johnson, G. C., Killick, R. E., Legler, D., Llovel, W., Lyman, J., Palmer,
1443 M. D., Piotrowicz, S., Purkey, S. G., Roemmich, D., Roca, R., Savita, A.,
1444 Schuckmann, K. von, Speich, S., Stephens, G., Wang, G., Wijffels, S. E., and

1445 Zilberman, N.: Measuring Global Ocean Heat Content to Es-
1446 timate the Earth Energy Imbalance, *Front. Mar. Sci.*, 6, 432, <https://doi.org/10.3389/fmars.2019.00432>, 2019.

1447 Minière, A., von Schuckmann, K., Sallée, J.-B., and Vogt, L.: Robust acceleration of Earth
1448 system heating observed over the past six decades. *Sci. Rep.*, 13, 22975,
1449 <https://doi.org/10.1038/s41598-023-49353-1>, 2024

1450 Nerem, R. S., Beckley, B. D., Fasullo, J. T., Hamlington, B. D., Masters, D., and Mitchum,
1451 G. T.: Climate-change–driven accelerated sea-level rise detected in the altimeter era.
1452 *P. Natl. Acad. Sci.*, 115, 2022–2025, <https://doi.org/10.1073/pnas.1717312115>, 2018.

1453 O’Carroll, A. G., Armstrong, E. M., Beggs, H. M., Bouali, M., Casey, K. S., Corlett, G. K.,
1454 Dash, P., Donlon, C. J., Gentemann, C. L., Høyer, J. L., Ignatov, A., Kabobah, K.,
1455 Kachi, M., Kurihara, Y., Karagali, I., Maturi, E., Merchant, C. J., Marullo, S.,
1456 Minnett, P. J., Pennybacker, M., Ramakrishnan, B., Ramsankaran, R. Santoleri, R.,
1457 Sunder, S., Saux Picart, S. Vázquez-Cuervo, J., Wimmer, W.: Observational Needs
1458 of Sea Surface Temperature. *Front. Mar. Sci.*, 6,
1459 <https://doi.org/10.3389/fmars.2019.00420>, 2019.

1460 Oliver, E. C. J., Benthuisen, J.A., Darmaraki, S., Donat, M. G., Hobday, A. J., Holbrook,
1461 N. J., Schlegel, R.W., and Sen Gupta A., Marine Heatwaves. *Annual review of*
1462 *marine science* **13**, 313–342, [https://doi.org/10.1146/annurev-marine-032720-](https://doi.org/10.1146/annurev-marine-032720-095144)
1463 [095144](https://doi.org/10.1146/annurev-marine-032720-095144), 2021.

1464 Purkey, S. G., and Johnson, G. C.: Warming of Global Abyssal and Deep Southern Ocean
1465 Waters between the 1990s and 2000s: Contributions to Global Heat and Sea Level
1466 Rise Budgets. *J. Climate*, 23, 6336–6351, <https://doi.org/10.1175/2010jcli3682.1>,
1467 2010.

1468 Palmer, M. D., and K. Haines: Estimating Oceanic Heat Content Change Using Isotherms.
1469 *J. Climate*, 22, 4953–4969, <https://doi.org/10.1175/2009JCLI2823.1>, 2009.

1470 Rahmstorf, S., Box, J., Feulner, G., Mann, M.E., Robinson, A., Rutherford, S.,
1471 Schaffernicht, E. Exceptional 2⁰h-Century slowdown in Atlantic Ocean
1472 overturning, *Nature Climate Change*, 5, 475–480, 2015.

1473 Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., Rowell, D.
1474 P., Kent, E. C., and Kaplan, A.: Global analyses of sea surface temperature, sea ice,
1475 and night marine air temperature since the late nineteenth century. *J. Geophys. Res.-*
1476 *Atmospheres*, 108, [https://doi.org/https://doi.org/10.1029/2002JD002670](https://doi.org/10.1029/2002JD002670), 2003.

1477 Reiniger, R. F., and Ross, C. K.: A method of interpolation with application to
1478 oceanographic data. *Deep Sea Research*, 15, 185-193, <https://doi.org/10.1016/0011->
1479 [7471\(68\)90040-5](https://doi.org/10.1016/0011-7471(68)90040-5), 1968.

1480 Rhein, M., S.R. Rintoul, S. Aoki, E. Campos, D. Chambers, R.A. Feely, S. Gulev, G.C.
1481 Johnson, S.A. Josey, A. Kostianoy, C. Mauritzen, D. Roemmich, L.D. Talley and F.
1482 Wang, 2013: Observations: Ocean. In: *Climate Change 2013: The Physical Science*
1483 *Basis. Contribution of Working Group I to the Fifth Assessment Report of the*
1484 *Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner,
1485 M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley
1486 (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York,
1487 NY, USA.

1488 Roemmich, D., and Gilson, J.: The 2004–2008 mean and annual cycle of temperature,
1489 salinity, and steric height in the global ocean from the Argo Program. *Prog.*
1490 *Oceanogr.*, 82, 81-100, <https://doi.org/10.1016/j.pocean.2009.03.004>, 2009.

1491 Roemmich, D., and Gilson, J.: The global ocean imprint of ENSO. *Geophys. Res. Lett.*, 38,
1492 <https://doi.org/10.1029/2011GL047992>, 2011.

1493 Roemmich, D., Alford, M. H., Claustre, H., Johnson, K., King, B., Moum, J., Oke, P.,
1494 Owens, W. B., Pouliquen, S., Purkey, S., Scanderbeg, M., Suga, T., Wijffels, S.,
1495 Zilberman, N., Bakker, D., Baringer, M., Belbeoch, M., Bittig, H. C., Boss, E., . . .
1496 Yasuda, I.: On the Future of Argo: A Global, Full-Depth, Multi-Disciplinary Array.
1497 *Front. Mar. Sci.*, 6, 2019.

1498 Savita, A., and Coauthors: Quantifying Spread in Spatiotemporal Changes of Upper-Ocean
1499 Heat Content Estimates: An Internationally Coordinated Comparison. *J. Climate*, 35,
1500 851–875, <https://doi.org/10.1175/JCLI-D-20-0603.1>, 2022.

1501 [Schweiger, A., R. Lindsay, J. Zhang, M. Steele, H. Stern, and R. Kwok: Uncertainty in](https://doi.org/10.1029/2011JC007084)
1502 [modeled Arctic sea ice volume. *J. Geophys. Res.*, 116, C00D06,](https://doi.org/10.1029/2011JC007084)
1503 <https://doi.org/10.1029/2011JC007084>, 2011.

1504 Sloyan, B. M., Wanninkhof, R., Kramp, M., Johnson, G. C., Talley, L. D., Tanhua, T.,
1505 McDonagh, E., Cusack, C., O'Rourke, E., McGovern, E., Katsumata, K., Diggs, S.,
1506 Hummon, J., Ishii, M., Azetsu-Scott, K., Boss, E., Ansorge, I., Perez, F. F., Mercier,
1507 H., . . . Campos, E.: The Global Ocean Ship-Based Hydrographic Investigations
1508 Program (GO-SHIP): A Platform for Integrated Multidisciplinary Ocean Science.
1509 *Front. Mar. Sci.*, 6, 2019.

1510 Sun, D., Li, F., Jing, Z. et al. Frequent marine heatwaves hidden below the surface of the
1511 global ocean. *Nat. Geosci.* 16, 1099–1104, [https://doi.org/10.1038/s41561-023-](https://doi.org/10.1038/s41561-023-01325-w)
1512 [01325-w](https://doi.org/10.1038/s41561-023-01325-w), 2023.

1513 Su, H. et al. OPEN: a new estimation of global ocean heat content for upper 2000 meters
1514 from remote sensing data. *Remote Sens.*, <https://doi.org/10.3390/rs12142294>, 2020.

1515 Tan, Z., Zhang, B., Wu, X., Dong, M., and Cheng, L.: Quality control for ocean
1516 observations: From present to future. *Science China Earth Sciences*, 65, 215-233,
1517 <https://doi.org/10.1007/s11430-021-9846-7>, 2022.

1518 Tan, Z., Cheng, L., Gouretski, V., Zhang, B., Wang, Y., Li, F., Liu, Z., & Zhu, J.: A new
1519 automatic quality control system for ocean profile observations and impact on ocean
1520 warming estimate. *Deep Sea Research Part I: Oceanographic Research Papers*, 194,
1521 103961, <https://doi.org/10.1016/j.dsr.2022.103961>, 2023.

1522 Trenberth, K. E.: *The Changing Flow of Energy Through the Climate System*.
1523 Cambridge University Press., <https://doi.org/10.1017/9781108979030>, 2022.

1524 Trenberth, K. E., and Fasullo, J. T.: Atlantic meridional heat transports computed from
1525 balancing Earth's energy locally. *Geophys. Res. Lett.*, 44, 1919-1927,
1526 <https://doi.org/10.1002/2016gl072475>, 2017.

1527 Trenberth, K. E., Fasullo, J. T., and Kiehl, J.: Earth's Global Energy Budget *Bull. Am.*
1528 *Meteorol. Soc.*, 90, 311-324, <https://doi.org/10.1175/2008bams2634.1>, 2009.

1529 Trenberth, K. E., Fasullo, J. T., Von Schuckmann, K., and Cheng, L.: Insights into Earth's
1530 Energy Imbalance from Multiple Sources. *J. Climate*, 29, 7495-7505,
1531 <https://doi.org/10.1175/jcli-d-16-0339.1>, 2016.

1532 Trenberth, K. E., Zhang, Y., Fasullo, J. T., and Cheng, L.: Observation-Based Estimates of
1533 Global and Basin Ocean Meridional Heat Transport Time Series. *J. Climate*, 32,
1534 4567-4583, <https://doi.org/10.1175/jcli-d-18-0872.1>, 2019.

1535 von Schuckmann, K., and Le Traon, P. Y.: How well can we derive Global Ocean
1536 Indicators from Argo data? *Ocean Sci.*, 7, 783-791, [https://doi.org/10.5194/os-7-783-](https://doi.org/10.5194/os-7-783-2011)
1537 [2011](https://doi.org/10.5194/os-7-783-2011), 2011.

1538 von Schuckmann, K., Cheng, L., Palmer, M. D., Hansen, J., Tassone, C., Aich, V.,
1539 Adusumilli, S., Beltrami, H., Boyer, T., Cuesta-Valero, F. J., Desbruyères, D.,
1540 Domingues, C., García-García, A., Gentine, P., Gilson, J., Gorfer, M., Haim-berger,
1541 L., Ishii, M., Johnson, G. C., Killick, R., King, B. A., Kirchengast, G.,
1542 Kolodziejczyk, N., Lyman, J., Marzeion, B., Mayer, M., Monier, M., Monselesan, D.
1543 P., Purkey, S., Roemmich, D., Schweiger, A., Seneviratne, S. I., Shepherd, A., Slater,

1544 D. A., Steiner, A. K., Straneo, F., Timmermans, M.-L., and Wijffels, S. E.: Heat
1545 stored in the Earth system: where does the energy go?, *Earth Syst. Sci. Data*, 12,
1546 2013–2041, <https://doi.org/10.5194/essd-12-2013-2020>, 2020.

1547 von Schuckmann, K., Palmer, M. D., Trenberth, K. E., Cazenave, A., Chambers, D.,
1548 Champollion, N., Hansen, J., Josey, S. A., Loeb, N., Mathieu, P.-P., Meyssignac, B.,
1549 and Wild, M.: An imperative to monitor Earth's energy imbalance, *Nat. Clim.*
1550 *Change*, 6, 138–144, <https://doi.org/10.1038/nclimate2876>, 2016.

1551 von Schuckmann, K., Minière, A., Gues, F., Cuesta-Valero, F. J., Kirchengast, G.,
1552 Adusumilli, S., Straneo, F., Ablain, M., Allan, R. P., Barker, P. M., Beltrami, H.,
1553 Blazquez, A., Boyer, T., Cheng, L., Church, J., Desbruyeres, D., Dolman, H.,
1554 Domingues, C. M., García-García, A., Giglio, D., Gilson, J. E., Gorfer, M., Haimberger,
1555 L. Hakuba, M. Z., Hendricks, S., Hosoda, S., Johnson, G. C., Killick, R., King, B.,
1556 Kolodziejczyk, N., Korosov, A., Krinner, G., Kuusela, M., Landerer, F. W., Langer, M.,
1557 Lavergne, T., Lawrence, I., Li, Y., Lyman, J., Marti, F., Marzeion, B., Mayer, M.,
1558 MacDougall, A. H., McDougall, T., Monselesan, D. P., Nitzbon, J., Otosaka, I., Peng, J.,
1559 Purkey, S., Roemmich, D., Sato, K., Sato, K., Savita, A., Schweiger, A., Shepherd, A.,
1560 Seneviratne, S. I., Simons, L., Slater, D. A., Slater, T., Steiner, A. K., Suga, T., Szekely,
1561 T., Thiery, W., Timmermans, M. L., Vanderkelen, I., Wijffels, S. E., Wu, T., Zemp, M.:
1562 Heat stored in the Earth system 1960–2020: where does the energy go? *Earth Syst.*
1563 *Sci. Data*, 15, 1675–1709, <https://doi.org/10.5194/essd-15-1675-2023>, 2023.

1564 Wang, F., Shen, Y., Chen, Q., and Sun, Y.: Reduced misclosure of global sea-level budget
1565 with updated Tongji-Grace2018 solution. *Sci Rep-Uk*, 11, 17667,
1566 <https://doi.org/10.1038/s41598-021-96880-w>, 2021.

1567 Watkins, M. M., Wiese, D. N., Yuan, D. N., Boening, C., and Landerer, F. W.: Improved
1568 methods for observing Earth's time variable mass distribution with GRACE using
1569 spherical cap mascons. *J. Geophys. Res.- Solid Earth*, 120, 2648–2671,
1570 <https://doi.org/10.1002/2014JB011547>, 2015.

1571 Wijffels, S. E., Willis, J., Domingues, C. M., Barker, P., White, N. J., Gronell, A.,
1572 Ridgway, K., and Church, J. A.: Changing Expendable Bathythermograph Fall Rates
1573 and Their Impact on Estimates of Thermosteric Sea Level Rise. *J. Climate*, 21, 5657–
1574 5672, <https://doi.org/10.1175/2008jcli2290.1>, 2008.

1575 WMO: State of the Global Climate 2021, WMO-No. 1290, 2022.

- 1576 Wong, A. P. S., Wijffels, S. E., Riser, S. C., et al.: Argo Data 1999–2019: Two Million
1577 Temperature-Salinity Profiles and Subsurface Velocity Observations From a Global
1578 Array of Profiling Floats, <https://doi.org/10.3389/fmars.2020.00700>, 2020.
- 1579 Zanna, L., Khatiwala, S., Gregory, J. M., Ison, J. & Heimbach, P. Global reconstruction of
1580 historical ocean heat storage and transport. *Proc. Natl Acad. Sci. USA* 116, 1126–
1581 1131, <https://doi.org/10.1073/pnas.1808838115>, 2019.
- 1582 Zhang, B. et al. CAS-Ocean Data Center, Global Ocean Science Database (CODCv1):
1583 temperature. Marine Science Data Center of the Chinese Academy of Science,
1584 doi:10.12157/IOCAS.20230525.001 (2024).
- 1585 Zhang, X., Church, J. A., Platten, S. M., and Monselesan, D.: Projection of subtropical
1586 gyre circulation and associated sea level changes in the Pacific based on CMIP3
1587 climate models. *Clim. Dyn.*, 43, 131-144, [https://doi.org/10.1007/s00382-013-1902-](https://doi.org/10.1007/s00382-013-1902-x)
1588 x, 2014.
- 1589