

1 Countrywide Digital Surface Models and Vegetation Height Models 2 from Historical Aerial Images

3 Mauro Marty¹, Livia Piermattei^{1,2}, Lars T. Waser¹, Christian Ginzler¹

4 ¹ Land Change Science Research Unit, Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf,
5 8903, Switzerland

6 ²Department of Geography, University of Zurich, Zurich, 8057, Switzerland

7 *Correspondence to:* Mauro Marty (mauro.marty@wsl.ch)

8 **Abstract.** Historical aerial images, captured by film cameras in the previous century, are valuable resources for quantifying
9 Earth's surface and landscape changes over time. In the post-war period, these images were often acquired to create
10 topographic maps, resulting in the acquisition of large-scale aerial photographs with stereo coverage. Photogrammetric
11 techniques applied to these stereo images enable the extraction of 3D information to reconstruct digital surface models (DSMs)
12 and orthoimages.

13 Here, we present a highly automated photogrammetric approach for generating countrywide DSMs of Switzerland, at a 1 m
14 resolution, from approximately 3240,000 scanned aerial stereo images acquired between 1979 and 2006, with known exterior
15 and interior orientation. We derived four countrywide DSMs for the epochs 1979–1985, 1985–1991, 1991–1998, and 1998–
16 2006. From the DSMs, we generated corresponding countrywide vegetation height models (VHMs). We assessed the quality
17 of the historical DSMs at the country scale and within six representative study sites, evaluating the vertical accuracy and the
18 completeness of image-matching across different land cover types.

19 Mean completeness ranged from 64% for 'glacial and perpetual snow' to 98% for 'sealed surfaces', with a value of 93% for
20 the 'closed forest' class. Across Switzerland, the median elevation accuracy of the historical DSMs compared with a reference
21 digital terrain model (DTM) on sealed surface points ranged from 0.028 to 0.1653 m, with a normalised median absolute
22 deviation (NMAD) of around 0.84 m and a maximum root mean square error (RMSE) of 1.203-90 m. Similar accuracies are
23 obtained when comparing ~~The same analysis between historical DSMs with measured~~ geodetic points ~~and historical DSMs~~
24 ~~showed higher accuracies, with median values of ≤ 0.05 m and an NMAD < 1 m.~~

25 The VHMs generated in this study enabled the detection of major changes in forest areas due to windstorm damage, forest
26 dynamics, and growth. This work demonstrates the feasibility of generating accurate, very high-resolution DSM time series
27 (spanning three decades) and VHMs from historical aerial images of the entire surface of Switzerland in a highly automated
28 manner. The VHMs are already being used to estimate countrywide biomass changes. The countrywide DSMs and VHMs for
29 the four epochs, along with auxiliary data, are available online at <https://doi.org/10.16904/envidat.528> (Marty et al., 2024) and
30 can be used to quantify long-term elevation changes and related processes across different surfaces.

1 Introduction

Photographs taken by film cameras on aeroplanes throughout the last century represent a unique and invaluable information source for quantifying past changes in the Earth's surface and landscape. Various institutions and mapping agencies have recognised this value by preserving and scanning these photographs and making them publicly available.

After the Second World War, large-scale aerial photographs were frequently taken to create topographic maps. To ensure stereoscopic coverage and enable the extraction of three-dimensional (3D) information, image blocks were acquired in parallel strips, with each photograph overlapping the next along the same flight line (forming a stereo pair). Due to the stereoscopic acquisition and very high resolution of these images, digital surface models (DSMs) and orthoimages at a metric or sub-metric resolution can be generated using digital photogrammetry and computer vision techniques. This capability has led to the use of historical aerial images to assess long-term surface elevation changes across various geoscience fields.

Multi-temporal DSMs derived from historical aerial images have frequently been used in cryospheric research for assessments of changes in glacier elevation (e.g., Magnússon et al., 2016; Belart et al., 2020; Denzinger et al., 2021), in periglacial studies (e.g., Cusicanqui et al., 2021; Fleischer et al., 2021), in geomorphology (e.g., Micheletti et al., 2015; Piermattei et al., 2023; Schwat et al., 2023), in volcanology (e.g., Gomez et al., 2015), in analyses of land-use change (e.g., Nebiker et al., 2014; Bolles et al., 2018) and in archaeology (e.g., Risbøl et al., 2015; Peppas et al., 2018). Most studies have been focused on local scales and have involved processing a small set of scanned aerial images, with a few considering larger scales (Geyman et al., 2022; Muhammed et al., 2023) or even regional scales such as Greenland (Korsgaard et al., 2016).

In this study, we generated countrywide DSMs of Switzerland using historical aerial images acquired from the 1980s to the early 2000s. Additionally, as we carried out the study within the framework of the Swiss National Forest Inventory (NFI), we produced corresponding countrywide vegetation height models (VHMs). The Swiss NFI aims to collect objective information on forests at the national level for scientific, political, social and economic purposes (Abegg et al., 2023). Data collection within the context of the NFI is mainly based on a systematic terrestrial sample inventory, which is intended to provide results for Switzerland as a whole, as well as for larger regions and individual cantons. Since the early days of the NFI, aerial image interpretation and remote sensing data have been used. Over the decades, the image data has evolved from analogue black-and-white images to digital RGB true colour aerial images, and finally to the digital colour infrared (CIR) sensor data used today. In addition to the application of stereo-image-based analysis on various sample grids, comprehensive datasets have been developed and produced.

The use of historical aerial images for 3D forest reconstruction is limited, with most studies focused on 2D changes in tree and canopy cover at local scales using orthoimages (e.g., Kadmon and Harari-Kremer, 1999; Morgan and Gergel, 2010; Kulha et al., 2018) and texture analysis (Hudak and Wessman, 1998; Kupidura et al., 2019). In Central Switzerland, Waser et al. (2008a, b) used scanned colour infrared aerial images to generate DSMs and quantify tree/shrub cover changes between 1997 and 2002. Recent research has focused on methods based on artificial intelligence (AI) to extract information on woody vegetation along alpine treeline ecotones, tree cover (Wang et al., 2022), and forest cover (Hufkens et al., 2020) from historical aerial

64 orthoimages. The study by Vége and St-Onge (2008) was among the first where forest canopy growth was quantified using
65 DSM time series from historical aerial images. As changes in the height of forest canopies over time represent a key aspect of
66 forest dynamics, further studies have been conducted to explore the use of historical images for tracking long-term changes in
67 canopy height (Nurminen et al., 2015; Božek et al., 2019; Hufkens et al., 2020) and forest stand age (Vastaranta et al., 2015)
68 and for describing the evolution of forest successional stages in tropical forests (Berveglieri et al., 2016, 2018).
69 Here, we present a highly automated approach to process scanned aerial images (hereafter referred to as ‘historical aerial
70 images’) for generating countrywide DSMs and VHMs in Switzerland. Expanding on the work by Heisig and Simmen (2021),
71 who generated a single countrywide orthoimage of Switzerland from images taken between 1985 and 1991, we processed all
72 scanned images available from swisstopo, the Swiss Federal Office of Topography (swisstopo, 2024a), acquired over
73 Switzerland between 1979 and 2006. Countrywide coverage was obtained using image acquisition intervals of approximately
74 seven years, leading to four countrywide DSM products for the epochs 1979–1985, 1985–1991, 1991–1998, and 1998–2006.
75 To derive VHMs, we normalised the elevation values from the historical DSMs by subtracting the available countrywide
76 digital terrain model (DTM; swissALTI3D version 2017, swisstopo). We assessed the quality of the generated DSMs by
77 evaluating the completeness of the image-matching process across different land cover types and the vertical accuracy of the
78 historical DSMs by comparing elevation values with: (i) the reference DTM [on sealed surface points](#), (ii) geodetic elevation
79 points on a national scale, and (iii) manual stereoscopic elevation measurements from six representative study sites in
80 Switzerland. We close the article by highlighting the potential applications of these datasets in research on forest dynamics
81 and management.

82 2 Dataset and methods

83 2.1 Historical aerial stereo imagery of Switzerland [and study sites selection](#)

84 DSMs and VHMs were generated from [approximately 32’000](#) historical images [acquired with an RC10 panchromatic aerial](#)
85 [camera manufactured by Wild Heerbrugg \(currently owned by Leica Geosystems, Heerbrugg, Switzerland\). Multiple airborne](#)
86 [campaigns were conducted from 1979 to 2006](#) covering the entire 41,285 km² of Switzerland, with an elevation range between
87 197 and 4634 m a.s.l. [Figure 1 shows the image footprints of the stereo pairs used to generate the DSMs, along with the](#)
88 [corresponding year of acquisition and the total number of images per epoch. Each campaign spans approximately 7 years to](#)
89 [cover the entire country. Image acquisition was guided by the 1:25000 map sheet units \(12 x 17.5 km\) representing the](#)
90 [topographic maps of Switzerland \(swisstopo, 2024b\). The first countrywide coverage was obtained from the acquisition](#)
91 [campaign that took place between 1979 and 1985 \(epoch 1\), followed by subsequent campaigns from 1985 to 1991 \(epoch 2\),](#)
92 [1991 to 1998 \(epoch 3\), and 1998 to 2006 \(epoch 4\) \(Fig. 12\).](#)

93 [The primary purpose of these photograph acquisitions was to update the topographic maps produced by swisstopo. The](#)
94 [photographs were acquired with approximately 70–80% front overlap and 30–40% side overlap. The average flight height for](#)

all datasets was approximately 4,000 m above the ground, with an image scale of about 1:25,000. The photographs were scanned with a resolution of 14 μm using a Leica DSW700 scanner (swisstopo). This resulted in an average ground sample distance (GSD) of around 0.35 m in the scanned images, with larger variations in high-elevation areas. Historical aerial images are freely available on swisstopo's website (swisstopo, 2024a) and are provided with exterior orientation parameters, as well as camera calibration protocols containing interior orientation parameters such as focal length and fiducial mark values.




Figure 1: Stereo footprint coverage over Switzerland for the four epochs and the number of images used for the respective digital surface model (DSM) generation (barplot, bottom). There is an overlap of acquisition years between adjacent epochs, but images from the same year cover different forest production regions in each epoch.

In the Swiss NFI, Switzerland is characterised by different types of landscapes and is divided into five main forest production regions: the Jura in the northwest (1), the Swiss Plateau (2), the Prealps (3) between the Swiss Plateau and the Alps, the Alps (4/4a), and the Southern Alps (5) (Fig. 24). These regions exhibit different climatic and topographic characteristics, such as land cover classes, elevation ranges, tree species and forest structures (Tab. 1). ~~Six r~~Representative study sites of approximately 210 km² ~~each~~ (i.e., one map sheet; ~~see section 2.2.2~~) were selected within these regions (Fig. 24). In the Alps region, an additional study site encompassing the tongue of the Grosser Aletsch Glacier was included. ~~The data used to generate the DSMs consists of approximately 40,000 panchromatic images acquired with an RC10 aerial camera manufactured by Wild Heerbrugg (currently owned by Leica Geosystems, Heerbrugg, Switzerland). Multiple airborne campaigns were conducted from 1979 to 2006, with each campaign spanning approximately seven years to cover the entire country. The first countrywide coverage was obtained from the acquisition campaign that took place between 1979 and 1985 (epoch 1), followed by subsequent campaigns from 1985 to 1991 (epoch 2), 1991 to 1998 (epoch 3), and 1998 to 2006 (epoch 4) (Fig. 2).~~




Figure 24: The Swiss NFI forest production regions (1 = Jura, 2 = Swiss Plateau, 3 = Pre-Alps, 4 = Alps, 5 = Southern Alps) in Switzerland and the respective location of the study site for each region, along with the generated orthoimage for epoch 1 (1979–1985). An additional study site (4a) was selected for the Alps forest production region.

The primary purpose of these photograph acquisitions was to update the topographic maps produced by swisstopo. The photographs were acquired with approximately 70–80% front overlap and 30–40% side overlap. The average flight height for all datasets was approximately 4,000 m above the ground, with an image scale of about 1:25,000. The photographs were scanned with a resolution of 14 μm using a Leica DSW700 scanner (swisstopo). This resulted in an average ground sample distance (GSD) of around 0.35 m in the scanned images, with larger variations in high elevation areas. Historical aerial images are freely available on swisstopo's website (swisstopo, 2024a) and are provided with exterior orientation parameters, as well

126 as camera calibration protocols containing interior orientation parameters such as focal length and fiducial mark values. Figure
127 2 shows the image footprints of the stereo pairs used to generate the DSMs, along with the corresponding year of acquisition.

128 **Table 2: Forest production region characteristics. Topographic parameters are based on the swissALTI3D digital terrain model**
129 **(DTM) version 2017. Forest parameters are based on the Swiss National Forest Inventory (NFI).**

Forest production region		Area	Elevation range	Mean slope	Forest cover	Forest type			
No.	Name	(km ²)	(m a.s.l.)	(°)	(%)	pure coniferous (%)	mixed coniferous (%)	mixed deciduous (%)	pure deciduous (%)
1	Jura	4935.4	244–1677	12.3	39.5	28.6	25.5	18.8	22.3
2	Swiss Plateau	9412.8	311–1685	7.2	24.2	37.8	22.9	13.6	22.1
3	Prealps	6608.1	370–2899	19.8	32.5	53.7	22.4	8.2	9.6
4	Alps	16782.6	370–4634	28.9	21.0	71.0	9.6	5.8	8.6
5	Southern Alps	3545.7	192–3899	31.6	39.2	31.7	7.4	5.2	51.9




Figure 2: Stereo footprint coverage over Switzerland for the four epochs and the number of images used for the respective digital surface model (DSM) generation (barplot, bottom). There is an overlap of acquisition years between adjacent epochs, but images from the same year cover different forest production regions in each epoch.

2.2 DSM and VHM generation

The scanned input images, auxiliary data, and the steps used to generate the DSMs and the VHMs are illustrated in Fig. 3. The countrywide DSMs and VHMs are provided at 1 m spatial resolution in the Swiss projected coordinate system CH1903+ / LV95 (EPSG code 2056), and height information is supplied in LN02 (EPSG code 5728) where applicable.

138 The scanned historical images were processed using the commercial photogrammetric software SocetSet (v5.6.0) by BAE
139 Systems (Falls Church, USA). With known interior and exterior orientation parameters, the historical images were imported
140 to the software through the “frame import dialogue”. The import process was separately done for all images originating from
141 the same camera. During the frame import process, camera calibration values were entered, and fiducial marks were primarily
142 detected using automatic fiducial detection, implemented in the frame import workflow. Where this automatic approach failed,
143 manual measurement of fiducial marks was conducted. Subsequently, the DSMs can be generated using the Next-Generation
144 Automatic Terrain Extraction (NGATE) package implemented in SocetSet (DeVenecia et al., 2007). The exterior orientation
145 information for the images from epochs 1, 2 and 4 was derived by swisstopo within the framework of the Swiss Land
146 Use/Cover Statistics program around 2004. This image orientation was performed using ground control points (GCPs), visible
147 in the historical imagery, and in more recent already-oriented digital stereo images available from swisstopo. The elevation of
148 the GCPs was stereoscopically extracted from the digital images. This image orientation resulted in accurate horizontal
149 positioning but biases in the Z direction, ranging from 1 to 5 m, randomly distributed across Switzerland. Therefore, the DSMs
150 from these epochs were vertically corrected, as described in section 2.2.2. In contrast, the images for epoch 2 were recently
151 oriented using a highly automated workflow, as described by Heisig and Simmen (2021). ~~In their study reported the~~
152 ~~documented~~ planimetric residuals of the GCPs for the different triangulation blocks, which ~~are in the~~ ranged from ~~of 2~~ to 3
153 pixels (0.6-0.9 m RMSE). Additionally, ~~—and a comparison between the derived historical orthomosaic and to a reference~~
154 ~~orthomosaic at sample points evenly distributed over Switzerland revealed leads to a median difference of 1 m, with no~~
155 ~~systematic bias across Switzerland (Heisig and Simmen, 2021). Although there are no documented values for no values are~~
156 ~~documented on the~~ planimetric accuracies for the images ~~from of~~ epochs 1, 3 and 4, a comparisons of the ortho mosaics from
157 ~~those of those epochs to one an orthomosaic derived from epoch 23 show similar results.~~




Figure 3: Workflow of historical image processing and countrywide digital surface model (DSM) and vegetation height model (VHM) generation.

2.2.1 Image-matching and DSMs

The DSM generation process using the NGATE package followed a highly automated workflow. To handle the DSM reconstruction process for approximately 3240,000 images, image-matching was performed on distributed machines for single stereo footprints (i.e., stereo pairs). In the initial step, the footprint for each image was calculated, and the footprints of adjacent images along the same flight line (belonging to the same acquisition year) were intersected to create stereo footprints. In total, 20,346 stereo footprints were processed. All stereo footprints were stored in a table on a PostGIS database, which was used to control the image-matching process running on distributed virtual machines. Since stereo pairs were formed only from the neighbouring images along the same flight line, multi-image-matching was not involved in this process.

NGATE provides several predefined correlation/matching strategies for different types of image content, such as urban, flat and low-contrast areas. To ensure the extraction of accurate and complete elevation information across different land cover types, two strategies were adopted, namely urban and low contrast. The urban strategy utilises internal correlation parameters that account for strong elevation discontinuities, which represents the most prominent characteristics of forest canopies. In contrast, while the low-contrast strategy aims to identify congruent points on highly homogeneous surfaces, such as found

on meadows or bare land. While image-matching in NGATE is conducted on every pixel of the input imagery, the resulting output file represents a regular grid of three-dimensional (XYZ) coordinates. Optionally, this grid can include information about successful image-matching or interpolation for the given pixel, as indicated by the Figure of Merit value, a product derived by the software. The spatial resolution of the output DSM is user-defined and, in our case, was set to 1 m, approximately three times the GSD of the scanned images.

The image-matching using the two strategies resulted in two separate DSMs for each stereo footprint. These two DSMs were combined into one DSM by including all successfully matched points from the urban strategy and, as forests typically exhibit large elevation discontinuities. Any filling remaining gaps in this DSM were filled with correlated points from the low-contrast strategy, where available.

2.2.2 DSM merging and vertical correction

The 20,346 DSMs resulting from image-matching of the single stereo footprints were merged to the level of the 1:25,000 map sheets, which were already used to guide the image acquisition campaigns (see section 2.1). For each map sheet geometry, all DSMs from the same epoch that corresponded to individual stereo footprints of single stereo footprints DSMs belonging to the same epoch were selected based on via spatial intersection. They were then merged and cropped, merged and cut to fit the extent of the map sheet, which typically included extent (approximately around 50 stereo footprints in flat areas and up to 150 in high-alpine regions). Since adjacent stereo footprints spatially overlapped spatially, the pixels in these overlapping areas pixels with identical x and y coordinates on these locations often included multiple elevation values. To address this, the elevation values which were merged generalised using the median height. Grid cells without any successfully matched elevation information were interpolated using a triangular irregular network (TIN) surface.

The exterior orientation of the images from epochs 1, 3 and 4 resulted in a vertical offset of up to 5 m between the DSMs and the reference DTM on stable terrain (i.e., sealed surfaces and grass/herb points; see section 2.3.1). Therefore, the DSMs from these epochs were vertically corrected. For consistency, this correction was also applied to the DSM from epoch 2, resulting in neglectable changes due to the already precise horizontal and vertical orientation.

The vertical correction was performed on the merged map sheet DSMs as illustrated in Fig. 4. The correction of approximately 20,000 DSMs of single stereo footprints was performed at the level of 1:25,000 map sheet units, provided by swisstopo (swisstopo, 2024b). These map sheets were originally used to plan airborne image acquisition campaigns. For each map sheet geometry, all DSMs of single stereo footprints belonging to the same epoch were selected via spatial intersection, merged and cut to the map sheet extent (approximately 50 stereo footprints in flat areas and up to 150 in high-alpine regions). Since adjacent stereo footprints always overlapped spatially, pixels on these locations often included multiple elevation values, which were generalised using the median height. Grid cells without any successfully matched elevation information were interpolated using a triangular irregular network (TIN) surface.

The vertical correction of the generated DSM for each map sheet and epoch is illustrated in Fig. 4. The workflow consisted of calculating the difference between the reference DTM (swissALTI3D version 2017; see section 2.2.3) and the generated DSM

at points corresponding to the classes ‘sealed surface’ and ‘grass/herb’ in the first and third campaigns of the Swiss Land Use/Cover Statistics program (1979–1985 and 2004–2009, respectively; see section 2.3.1). With this approach, only points with an unchanged land cover classification were used. Outliers were filtered out based on mean ± 1 standard deviation (Std), calculated within a moving window of 50×50 pixels. The median of the remaining elevation differences was then calculated for each of the 1,200 regular grid cells per map sheet. The number of regular grid cells was determined through an empirical approach, ensuring that each grid cell contained enough difference values to yield a valid median and that there were sufficient grid cells per map sheet to enable the interpolation of a continuous correction surface using spline interpolation. Finally, a coarse correction surface (spline interpolation) was derived from these median values and this correction surface was subsequently then applied to the resulting DSMs at the map sheet unit level to correct elevation bias.




Figure 4: Illustration of the workflow to correct the elevation bias in the historical digital surface models (DSMs) for each map sheet. The correction is based on the elevation difference between the reference digital terrain model (DTM) and the historical DSM for sample points classified as sealed surfaces and grass/herb.

2.2.3 Countrywide DSMs and VHM

After correcting for randomly distributed elevation biases (section 2.2.2) were corrected for in each map sheet DSM and of each epoch (as described in section 2.2.2), the individual all the corrected map sheet DSMs from belonging to the same epoch were mosaiced combined into a single one DSM as a Geotiff file, covering the whole of Switzerland.

226 The four countrywide VHMs were derived by calculating the difference between the historical DSMs and the reference DTM
227 (swissALTI3D version 2017, swisstopo; see workflow in Fig. 3). This process resulted in a normalised DSM (nDSM),
228 representing the heights of all ~~existing~~ objects above ground. Note that the reference DTM was resampled from the default
229 resolution of 2 m to 1 m to match the resolution of the generated DSMs. The swissALTI3D DTM is based on airborne laser
230 scanning data for elevations up to 2000 m a.s.l., with an overall point density of 0.5 points/m² and a vertical accuracy of ±0.5
231 m (standard deviation). Above 2000 m a.s.l., elevation information is derived by image-matching and manual editing, yielding
232 a documented vertical accuracy of 1–3 m on average (swisstopo, 2024c).
233 nDSM raster cells with values > 60 m were considered outliers due to erroneous image-matching and thus were replaced by
234 ‘no data’. Next, the VHM (i.e., the nDSM of vegetated areas) was generated by assigning a height above ground of 0 m to all
235 non-vegetated objects. To achieve this, a binary non-vegetation mask (0 = non-vegetation and 1 = vegetation) was created by
236 combining information from the topographic landscape model (TLM; swisstopo, 2024d) and a normalised difference
237 vegetation index (NDVI) map obtained from aerial orthoimages of the years 2009–2015. All areas classified in the TLM as
238 building, street, and water surface, as well as pixels with an NDVI < 0.1, were set to 0. To retain vegetation close to buildings
239 and streets in the VHMs, single trees and hedges from the TLM were buffered by 15 m and set to 1. This non-vegetation mask
240 was multiplied by the nDSMs. The resulting VHMs have a spatial resolution of 1 m, consistent with the historical DSMs.

241 2.3 Quality assessment of the DSMs

242 The quality of the generated DSMs was assessed by analysing: (1) the completeness achieved through the image-matching
243 process over different land cover types and (2) the elevation accuracy based on elevation differences to reference datasets (see
244 section 2.3.1). The elevation differences were calculated consistently as the ~~historical DSMs reference data minus the historical~~
245 ~~DSMs the reference data~~. Therefore, positive differences indicate that historical DSMs had ~~positive negative~~ biases, i.e., that
246 the reference data was ~~lower higher~~ than the historical DSM. Quality assessment was performed and reported for the six
247 selected areas and the countrywide DSMs.

248 2.3.1 Reference data

249 The reference dataset used to assess the completeness of the image-matching process over different land cover classes consisted
250 of sample point interpretation on a regular 100 m grid across Switzerland. These sample points were collected by the Swiss
251 Federal Statistical Office in the framework of the Swiss Land Cover/Use Statistics program (BFS, 2024) (Fig. A1). These
252 countrywide land cover statistics campaigns took place in 1979–1985, 1992–1997, 2004–2009 and 2013–2018. The
253 completeness for epoch 1 was assessed, and the reference data from the 1979–1985 campaign was used. This dataset comprised
254 121,896 points for the 6 selected areas. The 27 specific land cover classes were grouped into 6 categories: (1) sealed surface,
255 (2) bare land, (3) grass/herb, (4) shrub, (5) closed forest, and (6) glacial and perpetual snow.

256 The vertical accuracy of the DSMs was assessed by comparing the elevation with three reference datasets (~~i.e., DSM minus~~
257 ~~reference data~~). First, the sealed surface points that remained unchanged between the 1979–1985 and 2004–2009 campaigns

were used. At these points, the elevation differences between ~~the reference DTM and~~ the DSMs ~~and the reference DTM~~ were calculated. The reference DTM was the same as the one used for generating the VHM (see section 2.2.3).

The second [reference](#) dataset comprised 565 selected geodetic control points (i.e., measured with GNSS or triangulation) distributed throughout Switzerland. This dataset is monitored and maintained by swisstopo; the documented vertical accuracy ranges from 0.003 to 0.1 m, with 42% of the points showing accuracies of 0.05 m or better. For the remaining ones, an accuracy of 0.1 m is documented (swisstopo, 2023). These geodetic points enabled a completely independent estimation of the vertical accuracy of the derived DSMs.

As a third reference dataset, 16,650 stereoscopic elevation measurements conducted by a stereo-image interpreter at the Swiss Federal Institute for Forest, Snow and Landscape Research (WSL) were used. The data collection scheme was the same as in the NFI (Ginzler, 2019). These measurements were derived only for epoch 1 and within the 666 forest plots of 50 × 50 m area intersecting the 6 study sites (Fig. 1). For each forest plot, 25 elevation measurements were extracted on ~~the one of the same overlapping stereo pairs as used to create for generating~~ the historical DSM (Fig. A1). In addition, the six land cover classes used to assess completeness were assigned to each stereo measurement. The ‘closed forest’ land cover class was further divided into coniferous or deciduous trees. [It is worth noting that this reference data was used to evaluate agreement rather than serving as a measure of accuracy, since the stereoscopic measurements have some inherent inaccuracies and the measurement was done on one stereo pair only.](#)

2.3.2 DSM completeness

Completeness is an essential quality measure for photogrammetric DSM generation. Here, it was computed as the percentage of successfully matched points out of the potential total number of matched points within a defined spatial unit. Completeness was calculated within circles with a radius of 5 m centred around points of the land cover statistics dataset. Then, the mean completeness was calculated for the six land cover classes (i.e., sealed surface, bare land, grass/herb, shrub, closed forest, and glacial and perpetual snow). Completeness values were calculated ~~for all four epochs from the DSMs of~~ the six study sites [before cropping to the actual map sheet extent \(see section 2.2.2\)](#)~~for all four epochs.~~

2.3.3 Vertical accuracy

Vertical accuracy was calculated as the difference between the [DSMs and the](#) reference dataset (see section 2.3.1)~~and the DSMs~~. First, the elevation differences between the ~~reference DSMs and the reference~~ DTM ~~and the DSMs~~ were calculated at points classified as sealed surfaces. [Sealed surface points closer than 10 m to historical and current houses and the forest boundary were removed.](#) In addition, points classified as grass/herb and bare land were used to quantify the elevation bias of the DSMs to terrain slope, aspect and elevation. Sealed surface points were not used in this investigation because they were mainly located in flat areas. In a second analysis, the elevation values of the DSMs were compared with approximately 500 independent geodetic points. Finally, > 16,000 stereoscopic elevation measurements were used to assess the performance of the image-matching workflow on deciduous and coniferous trees, the primary target objects. [It is worth noting that this latter](#)

comparison was used to evaluate agreement rather than serving as a measure of accuracy, since the stereoscopic measurements have some inherent inaccuracies.

To calculate the statistical measures, differences greater than ± 50 m between the DSMs and the reference measurements ~~and~~ ~~DSMs~~ were excluded because such values indicated a failure in image-matching due to cloud cover or saturated images, yielding unrealistic elevation values. From the remaining differences, robust statistics, such as the median and the normalised median absolute deviation (NMAD) were calculated, as biases in spatial datasets are often not normally distributed (Höhle and Höhle, 2009). The NMAD is defined as $1.4826 \cdot \text{MAD}$ (~~median mean~~ absolute deviation). Root mean square error (RMSE) values were additionally calculated, as a standard quality metric to facilitate the comparison of our results with those from other studies. RMSE values were determined after applying a robust outlier removal (~~median mean~~ ± 35 NMADStd).

3 Results

3.1 Generation of DSMs

Four countrywide DSMs were generated successfully with the NGATE package of SocetSet. Data voids are related to the lack of stereo images (Fig. 2) or to unsuccessful image-matching in saturated or cloudy areas. Figure 5 shows the hillshaded DSM for the epoch 1979–1985 for the six study sites.





Figure 5: Hillshaded digital surface models (DSMs) for the six selected study sites and epoch 1 (1979–1985). The year of the DSM is 1982 for sites 1 and 2, 1980 for site 3, 1985 for sites 4 and 4a, and 1983 for site 5.

307 **3.2 DSM completeness**

308 Figure 6 shows the DSM completeness percentages for epoch 1 and the six land cover classes: sealed surface, bare land,
309 grass/herb, shrub, closed forest, and glacial and perpetual snow. The number of sample points for each class is also provided
310 in the figure. Completeness results for epochs 2, 3 and 4 are provided in Appendix A (Fig. A2), and Table A1 shows the
311 corresponding mean values. The lowest completeness occurred in areas classified as glacial and perpetual snow, with a mean
312 completeness of 65% across the four epochs. This result was expected, due to the texture less black-and-white images of these
313 surfaces, which resulted in poor image-matching performance. The highest completeness was achieved for sealed surfaces,
314 with a mean of 98%. For the classes bare land, shrub and closed forest, the mean completeness was approximately 90%,
315 whereas the grass/herb class had a mean of 79%. However, the completeness for the grass/herb class was higher for epochs 2,
316 3 and 4 (Fig. A2), with a minimum value of 84%.

317






318 **Figure 6: Distribution of the mean completeness values (as percentages) for epoch 1 across the six study sites and the six land cover**
319 **classes – sealed surface, bare land, grass/herb, shrub, closed forest, and glacial and perpetual snow. The snow class is only present**
320 **in study sites 4, 4a and 5. Note that for some classes and study sites, the number of points is minimal. For example, there are only 18**
321 **points in study site 5, which is equivalent to 0.15% for the class ‘glacial and perpetual snow’, and therefore the horizontal bar is not**
322 **visible.**

324 **3.3 DSM vertical accuracy assessment**

325 The vertical accuracy of the DSMs for each epoch was assessed by comparing them with three reference elevation datasets
326 (see section 2.3.1). Figure 7a shows the spatial distributions of the sealed surface points and the reference geodetic points,
327 while Table 2 summarises the statistical measures. Sealed surface points exhibit a clustered distribution in low-elevation areas
328 (i.e., regions 1, 2 and 3) and a sparser distribution in the Alps and Southern Alps (regions 4 and 5, respectively). No clear
329 systematic difference is observed when visualising the positive and negative elevation differences at sealed surface points

between the ~~DSM from the first epoch~~DTM and the ~~DTM~~DSM from the first epoch. However, plotting the frequency distribution of elevation differences across the entire country reveals a ~~posi~~negatively skewed distribution for all epochs (Fig. 7b). ~~Points on the sealed surface that were suboptimally located, such as those near urban areas and the forest edge, were removed. However, several values still exceed 10 m.~~Positive-Negative differences, where the reference DTM is lower than the historical DSM, may result in the suboptimal location of sealed surface points close to the urban area and forest edge, which could explain the presence of several values < -10 m. The ~~positive~~ negative skew is also reflected in the median values of the elevation differences (Table 2), with values between -0.083 and -0.165 m and a ~~maximum~~ # NMAD of ~~0.861-41~~approximately ~~1.5~~ m.

In the analysis of the vertical accuracy of the DSMs in comparison to independent geodetic points, the median is around zero, with an NMAD of < 1 m (Table 2). Note that the number of geodetic points is two orders of magnitude smaller than the sealed surface points and more evenly distributed over the country. Additionally, the number of geodetic points varies slightly across the different epochs due to the spatial coverage of the image-matching results (e.g., cloud cover and image saturation). The differences between the DSMs and geodetic points ~~and DSMs~~ have a more symmetric distribution for all four epochs, with few positive differences > 10 m (Fig. 7c).






Figure 7: (a) Spatial distribution of the sealed surface points and the geodetic points used for the vertical accuracy assessment of the historical digital surface models (DSMs). Sealed surface points are depicted in blue and red, representing the positive and negative differences, respectively, between [the DSM for epoch 1 and the reference digital terrain model \(DTM, *swissALTI3D*\) and the DSM for epoch 1](#). The inset shows the distribution of the sealed surface points and the geodetic points for study site 2, overlaid on the orthophoto for the year 2021 (swisstopo, 2023). Frequency distributions of the elevation difference in metres between the [DSM and the DTM \(*swissALTI3D*\) and the DSM](#) (b) on sealed surface points and (c) [between the DSM and the geodetic points for the whole of Switzerland for the four epochs](#). The vertical dashed line indicates the median value. The bin size is 0.5 m. [The histogram shows the distribution before the filtering used for the root mean square error \(RMSE\) calculation.](#)

357 **Table 2: Elevation difference between the historical digital surface models (DSMs) and the digital terrain model (DTM) ~~and the~~**
 358 **historical digital surface models (DSMs) over the sealed surface points and the geodetic points over Switzerland for the four epochs.**
 359 **The number of points before and after outlier filtering is reported; the latter number was used for the root mean square error**
 360 **(RMSE) calculation.**
 361

Epochs	Accuracy on sealed surface points					Accuracy on geodetic points				
	No. points (±50 m)	Median (m)	NMAD (m)	No. points after filter (± 5 NMAD)	RMSE (m)	No. points (±50 m)	Median (m)	NMAD (m)	No. points after filtering (± 5 NMAD)	RMSE (m)
Epoch 1	5490190 509	0.10- 0.28	0.81± 08	506638047 36993	1.14± 572.9 8	551	-0.05	0.96	498532	1.35± 75
Epoch 2	7168511 1761	0.16- 0.35	0.86± 11	661409922 1107496	1.20± 623.1 2	542	-0.03	0.82	471526	1.17± 25
Epoch 3	4927510 8358	0.08- 0.41	0.68± 14	446719511 4104032	0.95± 753.4 0	536	-0.02	0.78	468517	1.05± 91
Epoch 4	4645110 2338	0.13- 0.53	0.85± 41	424999054 88157	1.19± 143.9 0	525	-0.00	0.98	470509	1.27± 33

362

363 The elevation difference between the historical DSMs ~~DTM~~ and the DTM ~~the historical DSMs~~ over grass/herb and bare land

364 points shows no significant bias with slope, aspect or elevation, as shown in Figure 8. However, for elevations above 3000 m

365 a.s.l. and slopes greater than 45°, the NMAD increases drastically, exceeding 2 m for all four countrywide DSMs. However,

366 points within these intervals represent only approximately 5% (slope) and 1% (elevation), as shown in the histograms in Fig.

367 8. The elevation bias dependence on aspect shows a potential shift, with median values below 0.5 m and a constant NMAD of

368 ± 0.8 m for all datasets and aspect intervals.

369 The agreement between the stereoscopic measurement (see section 2.3.3) taken within the NFI plots and the historical DSM

370 of epoch 1 is reported in Table 3 for the different land cover classes, including a division of the closed forest class into

371 deciduous and coniferous trees. Figure 9 shows one NFI plot with the stereo measurement location and elevation values

372 compared with the DSM profile. As shown in Table 3, the closest agreement is observed for the sealed surface class, with a

373 median difference of -0.19 m and an NMAD of 0.63 m, comparable to the results obtained when analysing the difference

374 between the ~~DSM~~~~DTM~~ and ~~D~~~~TMSM~~ for this land cover class (see Table 2). For the grass/herb class, the differences to the

375 stereo measurements exhibit a similar offset but with a larger spread, with a median of -0.15 m and a ~~an~~ NMAD of about 1 m.


376 Sample points from the shrub and bare land classes show a wider spread, with NMAD values of 1.62 m and 1 m, respectively,

377 and larger systematic biases, with median values greater than -0.4 m. The weakest agreement is found for the closed forest

378 class, where deciduous and coniferous trees yield comparable results, with median values of about -1.8 m and NMAD values

379 of up to 3.77 m.

380






Figure 8: Elevation differences in grass/herb and bare land points as a function of slope (a), aspect (b), and elevation (c). The histogram on the right shows the number of points for each interval. Elevation differences are calculated as the historical digital surface models (DSMs) reference digital terrain model (DTM) minus reference digital terrain model (DTM) historical digital surface models (DSMs). Slope, aspect and elevations were derived from the reference DTM.

Table 3: Elevation differences over different land cover classes between the historical digital surface model (DSM) and the stereo measurements and the historical digital surface model (DSM) from epoch 1.

Land cover class	No. points	Median (m)	NMAD (m)	No. points after filtering (± 53 NMADStd)	RMSE (m)
Closed forest, deciduous trees	2917	-1.83	2.75	284755	4.2535
Closed forest, coniferous trees	2478	-1.77	3.77	243541	5.3647
Grass/herb	6792	-0.15	1.02	65086634	1.316
Shrub	424	-0.41	1.62	4115	2.1635
Sealed surfaces	408	-0.19	0.63	379401	0.86124
Bare land	1231	-0.74	1.00	121719	1.413
Glacial and perpetual snow	567	-0.17	2.15	53940	2.699




Figure 9: Stereo measurements within a Swiss National Forest Inventory (NFI) forest plot on the historical orthophoto of epoch 1 (left). On the right, a comparison is depicted between the digital surface model (DSM) profile (green line, interval of 1 m) and the elevation values of the stereo measurements.

3.4 VHMs and forest change

Based on the four countrywide historical DSMs for Switzerland, four VHMs were generated by subtracting a countrywide DTM and applying a non-vegetation mask (see section 2.2.3). Figure 10 shows the VHMs and the forest structure for the six study sites for epoch 2. Since the VHMs are direct derivatives of the DSMs, the statement above about the quality remains valid.

Forest dynamics, such as growth, cutting and regrowth between observation epochs are evident over the six study sites, as shown by the height differences of the VHMs between consecutive epochs (Fig. A3). A detailed look at the VHM profiles of study sites 2 and 4 (Fig. 10b) shows minimal change in forest height in mature forest stands (VHM > 20 m), indicating consistency among the VHMs. However, forest growth, cuts or damage—followed by regrowth can be detected, for example, between epoch 3 and epoch 4 (cut or damage) and 2022 (regrowth) (Fig. 10b, profile 2, 1250 m - 1500 m Fig. 10). The Swiss Plateau forest production region, where study site 2 is located, was heavily affected by storm damage. Figure 11 shows a small

area of the VHM in region 2 for epochs 1, 3 and 4, overlaid by polygons indicating areas damaged by the two largest storm events recorded in Switzerland: Vivian (25–27 February 1990) and Lothar (26 December 1999). The VHMs effectively capture the storm-affected areas. The VHM of 1982 depicts forest cover in all storm-affected areas, while the 1994 VHM clearly shows the damage caused by storm Vivian. The areas impacted by storm Lothar show mature forest stands in the 1994 VHM, while the heavily damaged areas are visible in the 2002 VHM.




Figure 10: Illustration of (a) the vegetation height models (VHMs) of epoch 2 (1985–1991) for the six selected study sites. The VHMs correspond to the years: 1987 for site 1, 1988 for site 2, 1986 for sites 3 and 4a, 1991 for sites 4, and 1989 for site 5. (b) The profile of the VHMs from historical aerial images for the four epochs and from a modern airborne survey (2022) for study sites 2 (acquisition period 2018 and 2022) and 4 (acquisition period 2022). The 2 km profile line is shown in red in (a).

Besides these qualitative results, these historical VHMs have been used in various studies (Price et al., 2020; Ginzler et al., 2021). Ginzler et al. (2021) investigated forest dynamics in Switzerland between 1980 and 2010. Their analysis was based on aggregated average vegetation heights for 100 × 100 m cells derived from the VHMs. They quantified forest dynamics by

retrieving the absolute height change over the entire period (sum of absolute differences) and the classified height change between periods (height gain or loss). The results showed high forest dynamics in the Swiss Plateau and some pre-alpine areas in the northern Alps, while significant parts of the Jura and southern Switzerland showed little change during these 30 years. Price et al. (2020) used the historical VHM to map and monitor woody above-ground biomass (AGB) dynamics across Switzerland over 35 years (1983–2017). They found a consistent relationship between vegetation height derived from the VHM and NFI measures of woody AGB across four inventory periods.




Figure 11: Small extent of region 2 (Swiss Plateau), showing the vegetation height models (VHM) of epochs 1, 3, and 4, overlaid by polygons indicating areas heavily affected by two major storm events in Switzerland (Vivian: 25 to 27.02.1990 and Lothar: 26.12.1999).

4 Discussion

4.1 Potential of countrywide historical DSMs and VHM of Switzerland

This study underscores the great value of historical aerial images for reconstructing accurate and high-resolution DSMs over approximately three decades. For the epochs 1979–1985, 1985–1991, 1991–1998 and 1998–2006, we successfully processed four countrywide DSMs and VHM at a resolution of 1 m. The quality of the generated models is sufficient for various geoscience applications, such as the quantification of natural hazards, mass movements, and sediment erosion, and the monitoring of human interference (e.g., deforestation) and urban development. [For urban analysis, we recommend using the DSM and normalising it again if needed, as pixel heights larger than 60 m in the VHM were removed.](#) Our historical VHM products show a high level of consistency, suggesting that changes in forest height are real and significant and thus are valuable tools for understanding forest dynamics in Switzerland. This includes the tracking of forest growth and changes in forest cover over time, the differentiation between disturbed (due to management practices or storm events) and undisturbed forests, and the detection of regrowth after disturbance. The clear delineation of storm-damaged areas (e.g., from storms Vivian and Lothar) in our VHM data demonstrates the capability of the models to identify disturbances accurately. By analysing the temporal

changes in these areas, we can differentiate between disturbances caused by natural events and those resulting from forest management activities. Therefore, insight can be gained into post-storm regrowth and forest management activities across the forest production regions, and their effectiveness can be quantified, for example in terms of protection forest function. This dataset offers great added value to the Swiss NFI forest management data and supports future management strategies.

4.2 Quality of the DSMs

The completeness of the image-matching across epochs remains nearly constant (Fig. A12). However, an improvement in completeness is observed for grass and herbs, as well as glacial and perennial snow classes, in relation to Epoch 1, although this may vary depending on the study area.~~shows a tendency steady to increase over the four epochs, especially starting from epoch 2 (TabFig. A12) and, particularly in areas classified as grasses and herbs (TabFig. A12) The biggest improvement This improvement is strongly related to advancements in camera technology (models and lenses), image-matching results, and the quality of the exterior orientation provided in this study.~~

In areas with successful image-matching, countrywide accuracy analysis comparing derived DSM elevations to swissALTI3D DTM elevations at sealed surface points shows sub-meter values~~metric to sub-metric accuracy~~ with a maximum $\pm$ uncertainty (NMAD) of $0.86 \leq 1.412$ m.~~This value is roughly 2.5 times the GSD, although the DSMs were generated at 1 m resolution. A. However, a~~ comparison with an independent dataset such as the geodetic points distributed across Switzerland shows comparable even higher, decametric accuracy.

The positive negative bias, where reference data was lower than the historical DSMs, may have resulted from changes in the suboptimal location of sealed surface points in near urban areas and forest edges (Fig. 12a), which could explain the presence of several values more positive negative than -10 m (Fig. 7b). The comparison with geodetic points also shows more positive outliers with values above 10 m (Fig. 7c), likely related to the suboptimal location of reference geodetic points on objects such as rooftops and statues, which cannot be precisely reconstructed from historical DSM photographs (Fig. 12b).




Figure 12: Example of suboptimal locations of (a) sealed-surface points and (b) geodetic points used for the accuracy assessment. Negative Positive biases over these points are shown in red blue, whilst negative ones are shown in red.

An elevation bias in areas with slopes greater than 45° and elevations above 3000 m a.s.l. (Fig. 8a, c) is present in many photogrammetric products from airborne and satellite sources (e.g., Piermattei et al., 2019) and is more pronounced in historical images with 8-bit radiometric resolution and lower accuracy in exterior orientation. Steep slopes and mountainous areas often experience strong shadows, overexposed photographs on snow-covered surfaces, and clouds over mountain tops, all affecting image-matching results and vertical accuracy, and increasing noise in the DSMs.

Comparing the agreement between manual stereo measurements and the historical DSM of epoch 1, the closed forest land cover class shows the largest bias among all classes. However, similar patterns are visible when comparing manual stereo measurements with a DSM from images acquired by a modern near-infrared Leica ADS Sensor with a GSD of 0.5 m (Ginzler et al., 2015). Ginzler et al. (2015) achieved median values of -0.34 m (coniferous) and -0.79 m (deciduous), with NMAD

values in the same range as in this study. ~~Compared to this study, the greater bias in the historical data could be due to the more challenging visual capture of canopy objects in manual stereo measurements on 8-bit panchromatic images. Also, the measurement was only performed on one stereo model, while the DSM in this study was created from the fusion of several overlapping pairs. Compared to this study, The larger bias in historical data might be due to the more demanding visual detection of canopy top objects during manual stereo measurements on panchromatic 8-bit imagery and the coarser resolution of the images. This observation is supported by Ginzler et al. (2015), where stereo measurements on imagery with 0.25 m GSD reduced median values to 0.08 m (coniferous) and 0.16 m (deciduous).~~ All other land cover classes show median values between ~0.1 m and 0.7 m and NMAD values between ~1 m and 1.5 m in both studies.

4.3 Challenges of image processing and limitations of our dataset

Challenges in working with airborne historical aerial images can include difficulties accessing the archive, the high cost of images, ~~deteriorated the quality deterioration film negatives low quality of scanning film negatives~~, and the poor availability of metadata such as exterior and interior orientation. In Switzerland, swisstopo provides free access to a comprehensive archive of scanned historical aerial images covering the entire country, along with calibration protocols and exterior orientation parameters for each image, essential for photogrammetric processing. However, the orientation accuracy of historical images varies between epochs, requiring a vertical correction of the DSMs due to offsets of several metres. Often, historical images lack calibration protocols and exterior orientation information, requiring ground control points to calculate the camera location. This has led the scientific community to adopt ‘structure from motion’ photogrammetric approaches for processing historical aerial images (e.g., Muhammed et al., 2023), as these approaches do not require approximated values, such as camera calibration information. Additionally, pipelines have been developed to minimise the effort of collecting GCPs (e.g., Knuth et al., 2023).

To handle ~~a~~ countrywide datasets acquired over three decades, the processing of approximately ~~3240~~ 3240,000 images, as well as ~~the~~ post-processing, including the vertical correction of about 20,000 ~~stereo footprint~~ DSMs, was done at the map sheet level. Vertical correction was required to minimise offsets caused by an inaccurate exterior orientation of the images (see section 2.2.2). Each corrected map sheet DSM was merged without further consideration of the edges of adjacent map sheets, potentially creating small jumps at the edges. These jumps are negligible at the country scale, but further co-registration should be considered for local and catchment studies involving more than one map sheet.

Another challenge of historical black-and-white imagery with 8-bit radiometric resolution is oversaturation in snow-covered or heavily illuminated areas, leading to no or erroneous image-matching results, particularly in mountainous regions. This is evident in the completeness results for points classified as glacial and perpetual snow (see section 3.2). Therefore, for studies related to glaciers and snow, the noise and the voids in the generated DSMs must be addressed with proper void filling and filtering. We used a simple TIN interpolation to fill the voids, providing a binary mask indicating interpolated areas, which can be used to remove these areas for further analysis. For glaciological studies or applications requiring precise dates, we do not have the exact information on the dates of each DSM/pixel, but we provide the year of the survey campaign per map sheet.

5 Data availability


Datasets can be accessed from EnviDat (<https://doi.org/10.16904/envidat.528>, Marty et al., 2024). The following files are available for the four epochs: countrywide digital surface model (DSM), hillshaded DSM, and vegetation height models (VHMs). A metadata shapefile is provided with information about the acquisition year of the photographs used here; the geometry corresponds to the 1:25,000 map sheets used during this study. In addition, a binary raster is provided indicating successfully matched pixels (1) and interpolated pixels (0), and a non-vegetation mask raster ‘TLM_Vegetation_mask’ is provided, indicating vegetated areas (1) and non-vegetated areas (0). All the raster files are provided as GeoTIFF files.

6 Conclusions

Millions of aerial photographs dating from the First World War to the early 2000s represent an enormous, yet largely untapped, resource for geoscientists, with significant potential for documenting and quantifying surface changes over the last century. In this study, we used freely available historical stereo images covering the whole of Switzerland, processed using photogrammetric software. These images were provided with both exterior and interior orientation, allowing us to derive four countrywide DSMs at a 1 m spatial resolution across four epochs, with survey campaigns spanning approximately seven years. Our developed workflow achieved ~~sub-metric~~ [meter to sub-meter](#) accuracy and high completeness of the DSMs in nearly all regions of Switzerland, demonstrating the feasibility of capturing continuous surface change patterns at a high spatial resolution over different land cover classes. [The vertical accuracy \(NMAD\) in areas below 2000 m a.s.l. and with a slope of less than 20° ranges from 1.57 to 2.06 times the GSD \(on average 0.35 m\).](#) However, lower completeness and increased noise were present in snow-covered areas, due to image saturation and reduced performance of the image-matching algorithm in these areas. As our study was conducted as part of the Swiss NFI, we also generated four countrywide VHMs using an available countrywide DTM. These VHMs have been used to quantify forest changes (Ginzler et al., 2021) and monitor above-ground biomass (Price et al., 2020). The high consistency of the VHMs indicates their potential for future research into forest ecosystem dynamics, services and management.

Future work can build upon these datasets by integrating modern data to extend the observation period, thereby enhancing the detection of climate signals. Additionally, refining time series methods for very-high-resolution datasets will further improve our understanding of surface changes and related processes.

Finally, this study highlights the importance of recognising the value of historical aerial imagery datasets. We urge mapping agencies, governmental authorities and military services to unlock these archives and make them freely accessible online. The creation of countrywide historical aerial image DSMs is feasible only when such data is readily available, as demonstrated by the open access provided by swisstopo in Switzerland.



544
545 **Figure A1: Illustration of the distribution of the Swiss land cover statistic points and the Swiss National Forest Inventory (NFI)**
546 **forest plots with the stereo measurements. The background is the generated orthophoto from the first epoch (1979–1985).**




Figure A2: Distribution of the mean completeness values, as percentages, for (a) epoch 2, (b) epoch 3, and (c) epoch 4, across the six study sites and the six land cover classes – sealed surface, bare land, grass/herb, shrub, closed forest, and glacial and perpetual snow. The last class is only present in study sites 4, 4a and 5.

551 **Table A1: Mean completeness values (as percentages) for each epoch over the six study sites, and for the six land cover classes. For**
552 **each class, the number of points used to assess the accuracy is provided. n.a. indicates not available.**

Study site	Sealed surface		Bare land		Grass/herb		Shrub		Closed forest		Glacial and perpetual snow	
	Mean (%)	Count (No.)	Mean (%)	Count (No.)	Mean (%)	Count (No.)	Mean (%)	Count (No.)	Mean (%)	Count (No.)	Mean (%)	Count (No.)
Epoch 1												
1	95	1312	84	221	53	17428	87	286	84	17899	n.a.	n.a.
2	98	3416	86	565	58	19879	92	586	93	12254	n.a.	n.a.
3	97	726	91	1172	74	20202	94	409	88	14877	n.a.	n.a.
4	99	173	95	19892	99	8785	99	3244	96	8496	51	1543
4a	99	284	94	12759	91	8119	98	1838	98	5346	64	9774
5	100	190	94	9045	99	3550	95	7125	98	15193	82	18
Epoch 2												
1	97	1526	90	287	84	21520	93	331	92	22974	n.a.	n.a.
2	99	3669	97	606	93	22699	98	646	97	13934	n.a.	n.a.
3	98	776	93	956	94	21609	96	434	88	16075	n.a.	n.a.
4	99	176	94	17811	96	8638	98	3139	95	8477	67	951
4a	100	296	92	11497	99	7163	98	1593	95	5287	78	9155
5	100	198	93	9699	99	3768	96	7330	97	16393	76	22
Epoch 3												
1	94	1947	87	342	91	24952	96	408	91	25816	n.a.	n.a.
2	99	3718	95	618	92	23317	95	639	92	13841	n.a.	n.a.
3	98	819	91	1400	95	23259	88	514	94	17229	n.a.	n.a.
4	99	163	93	17761	96	8437	98	3158	96	8239	68	1312
4a	99	360	94	13907	98	8868	98	1964	97	6060	85	12399
5	100	233	87	10475	97	4106	88	8122	94	17776	87	22
Epoch 4												
1	99	1748	96	281	89	22408	99	348	98	22718	n.a.	n.a.
2	99	3784	97	560	86	21622	96	670	94	15078	n.a.	n.a.
3	98	840	94	1715	95	23288	97	553	94	17637	n.a.	n.a.
4	99	147	93	15428	98	7680	96	3008	92	7927	85	953
4a	99	365	90	12840	97	8811	96	1862	96	5623	85	11283
5	99	154	81	4235	93	2175	84	4497	88	11808	27	5

553

554




Figure A3: The height difference between the vegetation height models (VHM) of consecutive epochs for each study site (1–5). The black line in study sites 2 and 4 indicates the location of the profile shown in Figure 10.

558 **Supplement**

559 No supplemental material.

560 **Author Contributions**

561 MM and CG designed the study. MM processed and generated all the data. LP and MM analysed the dataset. LP prepared all
562 the figures and tables. LP and MM wrote the manuscript. All authors contributed to the discussion and editing of the text.

563 **Competing interests**

564 The authors declare that they have no conflicts of interest.

565 **Acknowledgements**

566 This study was supported by the Swiss National Forest Inventory (NFI). We thank the Federal Statistical Office FSO and the
567 Federal Office of Topography swisstopo for providing access to the historical photographs. We are grateful to Holger Heisig
568 (swisstopo) for the successful semi-automatic orientation of thousands of historical images and fruitful and interesting
569 discussions about this source of data and to Dr Melissa Dawes for professional language editing.

570 **Funding**

571 This study was carried out in the framework of the Swiss National Forest Inventory (NFI), a cooperative effort between the
572 Swiss Federal Institute for Forest, Snow and Landscape Research (WSL) and the Swiss Federal Office for the Environment
573 (FOEN).

574 **References**

575 Abegg, M., Bösch, R., Kükenbrink, D., and Morsdorf, F.: Tree volume estimation with terrestrial laser scanning—testing for
576 bias in a 3D virtual environment, *Agric. For. Meteorol.*, 331, 109348, <https://doi.org/10.1016/j.agrformet.2023.109348>, 2023.
577
578 Belart, J.M., Magnússon, E., Berthier, E., Gunnlaugsson, Á.Þ., Pálsson, F., Aðalgeirsdóttir, G., Jóhannesson, T., Thorsteinsson,
579 T., and Björnsson, H.: Mass balance of 14 Icelandic glaciers, 1945–2017: spatial variations and links with climate, *Front. Earth*
580 *Sci.*, 8, 163, <https://doi.org/10.3389/feart.2020.00163>, 2020.
581

582 Berveglieri, A., Imai, N.N., Tommaselli, A.M., Casagrande, B. and Honkavaara, E.: Successional stages and their evolution
583 in tropical forests using multi-temporal photogrammetric surface models and superpixels, *ISPRS-J. Photogramm. Remote*
584 *Sens.*, 146, 548-558, <https://doi.org/10.1016/j.isprsjprs.2018.11.002>, 2018.

585

586 Berveglieri, A., Tommaselli, A.M.G., Imai, N.N., Ribeiro, E.A.W., Guimaraes, R.B. and Honkavaara, E.: Identification of
587 successional stages and cover changes of tropical forest based on digital surface model analysis, *IEEE J. Sel. Top. Appl. Earth*
588 *Obs. Remote Sens.*, 9(12), 5385-5397, <https://doi.org/10.1109/JSTARS.2016.2606320>, 2016.

589

590 BFS. Bundesamt für Statistik, Arealstatistik Schweiz – Nomenklatur 2004.
591 <https://www.bfs.admin.ch/bfs/de/home/statistiken/raum-umwelt/nomenklaturen/arealstatistik.html>, last access: 29 August
592 2024.

593

594 Bolles, K.C. and Forman, S.L.: Evaluating landscape degradation along climatic gradients during the 1930s dust bowl drought
595 from panchromatic historical aerial photographs, United States Great Plains. *Front. Earth Sci.*, 6, 153,
596 <https://doi.org/10.3389/feart.2018.00153>, 2018.

597

598 Božek, P., Janus, J. and Mitka, B.: Analysis of changes in forest structure using point clouds from historical aerial photographs,
599 *Remote Sens.*, 11(19), p.2259, <https://doi.org/10.3390/rs11192259>, 2019.

600

601 Cusicanqui, D., Rabatel, A., Vincent, C., Bodin, X., Thibert, E., and Francou, B.: Interpretation of volume and flux changes
602 of the Laurichard rock glacier between 1952 and 2019, French Alps, *J. Geophys. Res. Earth Surf.*, 126(9), p.e2021JF006161,
603 <https://doi.org/10.1029/2021JF006161>, 2021.

604

605 DeVenecia K., Walker S, and Bingcai, Z.: New Approaches to Generating and Processing High Resolution Elevation Data
606 with Imagery, *Photogrammetric Week '07*, 297-308, Wichmann Verlag, Heidelberg, 2007.

607

608 Denzinger, F., Machguth, H., Barandun, M., Berthier, E., Girod, L., Kronenberg, M., Usabaliev, R., and Hoelzle, M.: Geodetic
609 mass balance of Abramov Glacier from 1975 to 2015, *J. Glaciol.*, 67(262), 331-342, <http://doi.org/10.1017/jog.2020.108>,
610 2021.

611

612 Fleischer, F., Haas, F., Piermattei, L., Pfeiffer, M., Heckmann, T., Altmann, M., Rom, J., Stark, M., Wimmer, M.H., Pfeifer,
613 N., and Becht, M.: Multi-decadal (1953–2017) rock glacier kinematics analysed by high-resolution topographic data in the
614 upper Kaunertal, Austria, *Cryosphere*, 15(12), 5345-5369, <https://doi.org/10.5194/tc-15-5345-2021>, 2021.

615

616 Geyman, E.C., J. J. van Pelt, W., Maloof, A.C., Faste Aas, H., and Kohler, J.: Historical glacier change on Svalbard predicts
617 doubling of mass loss by 2100, *Nature*, 601, 374–379, <https://doi.org/10.1038/s41586-021-04314-4>, 2022.

618

619 Ginzler, C. and Hobi, M.L.: Countrywide stereo-image matching for updating digital surface models in the framework of the
620 Swiss National Forest Inventory, *Remote Sens.*, 7, 4343–4370, <https://doi.org/10.3390/rs70404343>, 2015.

621

622 Ginzler, C., Price, B., Weber, D., Hobi, M., and Marty, M.: Dynamik der Vegetationshöhen im Schweizer Wald, *Swiss Forestry*
623 *J.*, 172(5), 310-317, <https://doi.org/10.3188/szf.2021.0310>, 2021.

624

625 Gomez, C., Hayakawa, Y., and Obanawa, H.: A study of Japanese landscapes using structure from motion derived DSMs and
626 DEMs based on historical aerial photographs: New opportunities for vegetation monitoring and diachronic geomorphology,
627 *Geomorphology*, 242, 11-20, <https://doi.org/10.1016/j.geomorph.2015.02.021>, 2015.

628

629 Heisig, H. and Simmen, J.L.: Re-engineering the past: countrywide geo-referencing of archival aerial imagery, *J. Photogramm.*
630 *Remote Sens. Geoinf. Sci.*, 89(6), 487-503, <https://doi.org/10.1007/s41064-021-00162-z>, 2021.

631

632 Höhle, J. and Höhle, M.: Accuracy assessment of digital elevation models by means of robust statistical methods, *ISPRS-J.*
633 *Photogramm. Remote Sens.*, 64(4), 398-406, <https://doi.org/10.1016/j.isprsjprs.2009.02.003>, 2009.

634

635 Hufkens, K., de Haulleville, T., Kearsley, E., Jacobsen, K., Beeckman, H., Stoffelen, P., Vandeloos, F., Meeus, S., Amara,
636 M., Van Hirtum, L., and Van den Bulcke, J.: Historical aerial surveys map long-term changes of forest cover and structure in
637 the Central Congo Basin, *Remote Sens.*, 12(4), 638, <https://doi.org/10.3390/rs12040638>, 2020.

638

639 Hudak, A.T. and Wessman, C.A.: Textural analysis of historical aerial photography to characterise woody plant encroachment
640 in South African Savanna, *Remote Sens. Environ.*, 66(3), 317-330, [https://doi.org/10.1016/S0034-4257\(98\)00078-9](https://doi.org/10.1016/S0034-4257(98)00078-9), 1998.

641

642 Kadmon, R. and Harari-Kremer, R.: Studying long-term vegetation dynamics using digital processing of historical aerial
643 photographs, *Remote Sens. Environ.*, 68(2), 164-176, [https://doi.org/10.1016/S0034-4257\(98\)00109-6](https://doi.org/10.1016/S0034-4257(98)00109-6), 1999.

644

645 Knuth, F., Shean, D., Bhushan, S., Schwat, E., Alexandrov, O., McNeil, C., Dehecq, A., Florentine, C., and O’Neel, S.:
646 Historical Structure from Motion (HSfM): Automated processing of historical aerial photographs for long-term topographic
647 change analysis, *Remote Sens. Environ.* 285, 113379, <https://doi.org/10.1016/j.rse.2022.113379>, 2023.

648

649 Korsgaard, N.J., Nuth, C., Khan, S.A., Kjeldsen, K.K., Bjørk, A.A., Schomacker, A., and Kjær, K.H.: Digital elevation model
650 and orthophotographs of Greenland based on aerial photographs from 1978–1987, *Sci. Data*, 3, 160032,
651 <https://doi.org/10.1038/sdata.2016.32>, 2016.

652

653 Kulha, N., Pasanen, L., and Aakala, T.: How to calibrate historical aerial photographs: a change analysis of naturally dynamic
654 boreal forest landscapes, *Forests*, 9(10), 631, <https://doi.org/10.3390/f9100631>, 2018.

655

656 Kupidura, P., Osńska-Skotak, K., Lesisz, K., and Podkowa, A.: The efficacy analysis of determining the wooded and shrubbed
657 area based on archival aerial imagery using texture analysis, *ISPRS Int. J. Geo-Inf.*, 8(10), 450,
658 <https://doi.org/10.3390/ijgi8100450>, 2019.

659

660 Marty, M., Piermattei, L., Ginzler, C., and Waser, L.T.: Countrywide DSM and VHM from historical aerial images, *Envidar*
661 [data set], <https://doi.org/10.16904/envidat.528>, 2024.

662

663 Magnússon, E., Muñoz-Cobo Belart, J., Pálsson, F., Ágústsson, H., and Crochet, P.: Geodetic mass balance record with
664 rigorous uncertainty estimates deduced from aerial photographs and lidar data – case study from Drangajökull ice cap, NW
665 Iceland, *Cryosphere*, 10(1), 159-177, <https://doi.org/10.5194/tc-10-159-2016>, 2016.

666

667 Micheletti, N., Lane, S.N., and Chandler, J.H.: Application of archival aerial photogrammetry to quantify climate forcing of
668 alpine landscapes, *Photogram. Rec.*, 30, 143-165, <https://doi.org/10.1111/phor.12099>, 2015.

669

670 Morgan, J.L. and Gergel, S.E.: Quantifying historic landscape heterogeneity from aerial photographs using object-based
671 analysis, *Landscape Ecol.*, 25, 985-998, <https://doi.org/10.1007/s10980-010-9474-1>, 2010.

672

673 Muhammed, M.A., Hailu, B.T., Miehe, G., Wraase, L., Nauss, T. and, Zeuss, D.: High-resolution digital elevation models and
674 orthomosaics generated from historical aerial photographs (since the 1960s) of the Bale Mountains in Ethiopia, *Earth Syst.*
675 *Sci. Data*, 15(12), 5535-5552, <https://doi.org/10.5194/essd-15-5535-2023>, 2023.

676

677 Nebiker, S., Lack, N., and Deuber, M.: Building change detection from historical aerial photographs using dense image
678 matching and object-based image analysis, *Remote Sens.*, 6(9), 8310-8336, <https://doi.org/10.3390/rs6098310>, 2014.

679

680 Nurminen, K., Litkey, P., Honkavaara, E., Vastaranta, M., Holopainen, M., Lyytikäinen-Saarenmaa, P., Kantola, T., and
681 Lyytikäinen, M.: Automation aspects for the georeferencing of photogrammetric aerial image archives in forested scenes,
682 *Remote Sens.*, 7(2), 1565-1593, <https://doi.org/10.3390/rs70201565>, 2015.

683

684 Peppa, M.V., Mills, J.P., Fieber, K.D., Haynes, I., Turner, S., Turner, A., Douglas, M., and Bryan, P.G.: Archaeological feature
685 detection from archive aerial photography with a SfM-MVS and image enhancement pipeline, *Int. arch.*
686 *Photogramm. Remote Sens. Spat. Inf. Sci.*, 42, 869-875, <https://doi.org/10.5194/isprs-archives-XLII-2-869-2018>, 2018.

687

688 Piermattei, L., Heckmann, T., Betz-Nutz, S., Altmann, M., Rom, J., Fleischer, F., Stark, M., Haas, F., Ressler, C., Wimmer,
689 M.H., and Pfeifer, N.: Evolution of an Alpine proglacial river during 7 decades of deglaciation, *Earth Surf. Dyn.*, 11(3), 383-
690 403, <https://doi.org/10.5194/esurf-11-383-2023>, 2023.

691

692 Piermattei, L., Marty, M., Ginzler, C., Pöchtrager, M., Karel, W., Ressler, C., Pfeifer, N., and Hollaus, M.: Pléiades satellite
693 images for deriving forest metrics in the Alpine region, *Int. J. Appl. Earth Obs. Geoinf.*, 80, 240-256,
694 <https://doi.org/10.1016/j.jag.2019.04.008>, 2019.

695

696 Price, B., Waser, L.T., Wang, Z., Marty, M., Ginzler, C. and Zellweger, F.: Predicting biomass dynamics at the national extent
697 from digital aerial photogrammetry, *Int. J. Appl. Earth Obs. Geoinf.*, 90, 102116, <https://doi.org/10.1016/j.jag.2020.102116>,
698 2020.

699

700 Risbøl, O., Briese, C., Doneus, M., and Nesbakken, A.: Monitoring cultural heritage by comparing DEMs derived from
701 historical aerial photographs and airborne laser scanning, *J. Cult. Herit.*, 16(2), 202-209,
702 <https://doi.org/10.1016/j.culher.2014.04.002>, 2015.

703

704 Schwat, E., Istanbuluoglu, E., Horner-Devine, A., Anderson, S., Knuth, F., and Shean, D.: Multi-decadal erosion rates from
705 glacierized watersheds on Mount Baker, Washington, USA, reveal topographic, climatic, and lithologic controls on sediment
706 yields, *Geomorphology*, 438, 108805, <https://doi.org/10.1016/j.geomorph.2023.108805>, 2023.

707

708 Swisstopo, swissALTI3D version 2017: [https://backend.swisstopo.admin.ch/fileservice/sdweb-docs-prod-swisstopoch-](https://backend.swisstopo.admin.ch/fileservice/sdweb-docs-prod-swisstopoch-files/files/2023/11/14/cf4990a1-6bfb-4331-b990-a894d686266e.pdf)
709 [files/files/2023/11/14/cf4990a1-6bfb-4331-b990-a894d686266e.pdf](https://backend.swisstopo.admin.ch/fileservice/sdweb-docs-prod-swisstopoch-files/files/2023/11/14/cf4990a1-6bfb-4331-b990-a894d686266e.pdf), last access: 29 August 2024, 2017.

710

711 Swisstopo geodetic points: <https://opendata.swiss/en/dataset/lagefixpunkte-lfp1-landesvermessung>, last access: 29 August
712 2024, 2023.

713

714 Swisstopo, aerial photographs: <https://www.swisstopo.admin.ch/en/analogue-aerial-photographs>, last access: 29 August 2024,
715 2024a.

716

Swisstopo, National Map 1:25'000: <https://www.swisstopo.admin.ch/en/national-map-1-25000>, last access: 29 August 2024, 2024b.

Swisstopo, swissALTI3D technical information: <https://www.swisstopo.admin.ch/en/height-model-swissalti3d>, last access: 29 August 2024, 2024c.

Swisstopo, The topographic Landscape Model TLM: <https://www.swisstopo.admin.ch/en/information-topographic-landscape-model>, last access: 29 August 2024, 2024d.

Vastaranta, M., Niemi, M., Wulder, M.A., White, J.C., Nurminen, K., Litkey, P., Honkavaara, E., Holopainen, M., and Hyypä, J.: Forest stand age classification using time series of photogrammetrically derived digital surface models, *Scand. J. For. Res.*, 31(2), 194-205, <https://doi.org/10.1080/02827581.2015.1060256>, 2016.

Véga, C. and St-Onge, B.: Height growth reconstruction of a boreal forest canopy over a period of 58 years using a combination of photogrammetric and lidar models, *Remote Sens. Environ.*, 112(4), 1784-1794, <https://doi.org/10.1016/j.rse.2007.09.002>, 2008.

Wang, Z., Ginzler, C., Eben, B., Rehush, N., and Waser, L.T.: Assessing changes in mountain treeline ecotones over 30 years using CNNs and historical aerial images, *Remote Sens.*, 14, 2135, <https://doi.org/10.3390/rs14092135>, 2022.

Waser, L.T., Baltsavias, E., Ecker, K., Eisenbeiss, H., Ginzler, C., Kuchler, M., Thee, P., and Zhang, L.: High-resolution digital surface models (DSMs) for modelling fractional shrub/tree cover in a mire environment, *Int. J. Remote Sens.*, 29(5), 1261-1276, <https://doi.org/10.1080/01431160701736422>, 2008a.

Waser, L.T., Baltsavias, E., Ecker, K., Eisenbeiss, H., Feldmeyer-Christe, E., Ginzler, C., and Zhang, L.: Assessing changes of forest area and shrub encroachment in a mire ecosystem using digital surface models and CIR aerial images, *Remote Sens. Environ.*, 112(5), 1956-1968, <https://doi.org/10.1016/j.rse.2007.09.015>, 2008b.

Waser, L.T., Fischer, C., Wang, Z., and Ginzler, C.: Wall-to-wall forest mapping based on digital surface models from image-based point clouds and a NFI forest definition, *Forests*, 6, 4510–4528, <https://doi.org/10.3390/f6124386>, 2015.