

Insights into the North Hemisphere daily snowpack at high resolution from the new Crocus-ERA5 product

Silvana Ramos Buarque ¹, Bertrand Decharme ¹, Alina L. Barbu ¹, and Laurent Franchisteguy ²

¹CNRM UMR 3589, Météo-France/CNRS, Université fédérale de Toulouse, Toulouse, France

²Direction des Systèmes d'Observation, Météo-France, Toulouse, France

Correspondence: S. Ramos Buarque (silvana.buarque@meteo.fr)

Abstract. This article provides a detailed analysis of the Crocus-ERA5 snow product covering the Northern Hemisphere from 1950 to 2022. It assesses the product's performance in terms of snow depth and extentcover compared to in situ observations and satellite data. Compared to its predecessor, Crocus-ERA-Interim, Crocus-ERA5 benefits from improved spatial resolution and better atmospheric data assimilation, resulting in more accurate snowpack estimates, especially induring spring in Eurasia. The findings show a good match with observations, though biases remain, particularly inboreal forest areas and some Arctic regions, where the model tends to overestimate spring melt. In low-vegetation areas as tundra, Crocus-ERA5 may introduce biases due to its limited consideration of inter-annual vegetation changes, leading in inaccuracies in the simulation of snowmelt. The production of this snow dataset is motivated by its use by theresponds to the request of the continental cryosphere community ,and i. In particular the collaboration between the French and Canadian government institutions CNRM (National Center for Meteorological Research(CNRM) and ECCC (Environment and Climate Change Canada(ECCC) ,which hashave been involved in monitoring Arctic snow cover monitoringas part of the "Terrestrial Snow" section of the Arctic Report Card since 2017. The Crocus-ERA5 product is freely available on a daily basis and at 0.25° resolution over the 1950-07-01 to 2023-06-30 period (Decharme et al., 2024, <https://doi.org/10.5281/zenodo.14513248>).

1 Introduction

The Arctic is particularly vulnerable to global climate change. Changes in atmospheric circulation can result in altered precipitation patterns affecting the amount and type of precipitation that contributes to snow accumulation (Ramos Buarque and Salas y Melia, 2018). Whereas changes in sea ice can also indirectly affect the land surface snow cover by altering the surface albedo and heat exchange processes (Pörtner et al., 2019). Over the past few decades, the Arctic has experienced warming at a rate approximately twice as fast as the global average, resulting in unquestionable alterations to the Arctic cryosphere, as documented in the Intergovernmental Panel on Climate Change (IPCC) Special Report on Oceans and Cryosphere in a Changing Climate (Meredith et al., 2022). In addition to the above-average surface temperatures, an unprecedented (since instrumental records began) geographic spread of heat waves and warm spells occurred (Dunn et al., 2023). As a result, the rise in global temperatures linked to oceanic warming has led to a reduction in sea ice extent.

~~This decline is particularly pronounced~~ The pronounced decline in sea ice extent during late summer and early autumn months, which
25 ~~leads in an~~ increases the ocean's heat storage capacity, ~~of the ocean, thereby delaying~~ which in turn delays the formation of new ice at
the onset of the cold season. ~~Simultaneously on land,~~ This additional ocean heat also influences adjacent land areas, contributing to
unprecedented high near-surface permafrost temperature. ~~have reached unprecedented levels, resulting in the formation of deeper active layers~~
~~than previously observed with significant~~ As a result, the active layer – the seasonally thawed layer of soil above the permafrost – is
becoming deeper, with serious implications for soil stability, surface hydrology, and carbon emissions, ~~which in turn can contribute~~
30 ~~to climate warming~~ all of which further intensify climate warming. On land, terrestrial snow cover - which now vanishes entirely
from Arctic land surfaces during summer - plays a critical role in this interconnected system. Snow acts as both an insulator
for the underlying permafrost and a reflective surface that regulates land surface temperatures. Its disappearance exposes the
ground to direct solar radiation, accelerating permafrost thaw and deepening active layers. In late spring, early snow loss re-
duces surface albedo, allowing more solar energy to be absorbed and amplifying surface warming. This accelerates snowmelt
35 and reinforces the snow-albedo feedback, a key driver of permafrost thaw, shifts in runoff patterns, changes in wildlife behav-
ior, and heightened wildfire risk. Snow loss during this period also enhances carbon uptake in subsequent months, affecting
Arctic ecosystem productivity. Moreover, snow responds rapidly to temperature fluctuations, particularly in early autumn and
early spring, making it a highly sensitive and visible indicator of Arctic climate change. Variations in snow cover not only
influence the timing and volume of snow melt into rivers and streams but also affect permafrost stability and the functioning of
40 Arctic ecosystems. Together, these interlinked changes in sea ice, snow, and permafrost create reinforcing feedback loops that
amplify the pace and impacts of climate change in the Arctic. ~~Finally, snow depth and overall snow cover duration is significantly impacted.~~
~~For example, since the 1970s, snow cover has declined significantly in the northern hemisphere.~~

The aforementioned evidences clearly illustrates the intricate nature of these changes and the complex interactions between
the various components of the Arctic climate system. In this context, the Arctic Report Card (ARC, <https://arctic.noaa.gov/>
45 [report-card/](https://arctic.noaa.gov/)), published annually by the National Oceanic and Atmospheric Administration (NOAA) of the United States,
enables the global scientific community to monitor and document these changes. This report presents a comprehensive and
up-to-date assessment of the current state of the Arctic, based on the most recent scientific data. The ARC addresses a range
of environmental aspects pertaining to the Arctic, including air and ocean temperatures, precipitation, Greenland ice sheet,
sea ice extent, snow, permafrost, vegetation, etc. The objective of the report is to provide policymakers, scientists, and the
50 general public with information about the rapid and important changes occurring in the Arctic as a result of climate warming.
It identifies significant events of the past year, year-to-year variability and long-term trends, providing robust monitoring of
the evolution of this critical region for the global climate. By providing accurate and timely information, the ARC plays an
important role in raising awareness and supporting informed decision-making on the environmental challenges facing the
Arctic.

55 ~~The motivation for the present study stems from the fact that,~~ Since 2017, the French National Center for Meteorological Research
(CNRM) ~~is~~ has been involved in the "Terrestrial Snow" contribution to the ARC through a collaboration led by the Environ-
nement et Changement Climatique Canada (ECCC) Institute to monitor the evolution of snow cover and mass each year (e.g.
Mudryk et al., 2023). ~~The CNRM contribution to the "Terrestrial Snow" ARC~~ CNRM's role is to provide a daily snowpack product de-

rived from the Crocus complex snow scheme (Brun et al., 1992, 1989) ~~in order to take part of~~ contributing to the monitoring and
60 ~~improving advancement~~ of knowledge on the snow cover and mass of the North Hemisphere (NH). ~~Despite it~~ In addition to its use
in the ARC, from 2017 to 2020, this product has also been used in several scientific studies (Mudryk et al., 2015; Mortimer
et al., 2020; Kouki et al., 2023).

~~The CNRM contribution to the "Terrestrial Snow" ARC is to provide a daily snowpack product derived from the Crocus complex snow scheme (Brun et al., 1992, 1989) in order to take part of monitoring and improving of knowledge on the snow cover and mass of the North Hemisphere (NH).~~ Recently,
65 our snow product has been updated using the same Crocus configuration but driven by the fifth generation of the ECMWF
global atmospheric reanalysis ERA5 (Hersbach et al., 2020) , referred to as Crocus-ERA5. Although this product has been
little evaluated, it has been used since 2021 in the ARC report (Mudryk et al., 2023) and in several scientific studies (Derksen
and Mudryk, 2023; Mudryk et al., 2024).

The aim of this study is to provide a brief evaluation and insight into this Crocus-ERA5 daily snow product, which ~~will be used~~
70 ~~in a number of~~ support numerous studies on the evolution of the Arctic snow cover in the coming years. This product is the result
of a novel rerun of the Crocus snowpack model whose innovation is threefold: (i) the use of the improved ECMWF reanalysis
ERA5 as atmospheric forcing, offering higher spatial and temporal resolution, which enhances the accuracy of snowpack simu-
lations; (ii) comprehensive documentation of this update, highlighting both the strengths and limitations of the two most widely
used variables in the community across the vast expanse of the Northern Hemisphere; and (iii) validation through comparison
75 with fully independent references, including in-situ observations and multi-source data analysis.

The data and methods used to evaluate the new Crocus-ERA5 product are presented in Section 2. The main findings of this
study are presented in Section 3. The Crocus-ERA5 product is first compared with its predecessor, Crocus forced by ERA-
Interim, hereafter referred to as ERAI, to assess the progress (or lack thereof) between successive generations of these snow
products. In the Arctic region, validation of the snowpack benefits of various observations and estimates, including in situ and
80 satellite data, so Crocus-ERA is then evaluated against a range of in situ and satellite data. Finally, a brief discussion and the
main conclusions are presented in section 4.

2 Data and Methods

The performance of snowpack modelling in the context of climate change, can be ~~summarized by the~~ effectively characterized
using two mainkey variables ~~used as indicators of climate change because of their interactions and feedbacks with surface energy:~~ snow depth and
85 snow cover. These variables are widely used as climate change indicators because of their strong interactions and feedbacks
with the surface energy balance.

Snow depth ~~is a~~ plays a critical ~~variable for~~ role in both seasonal and long-term evolution of frozen ground and permafrost.
Changes in snow depth have significant implications in atmospheric circulation and ~~potentially may impact the~~ influence weather in
other parts of the world. For instance, large snow depths in western Russia have been associated with local cyclonic circulation
90 ~~as the, where~~ northerly flow onto the west of the trough favour the advection of cold air into the regions, ~~thereby preserving the mantle~~
~~snowy~~ helping to preserve the snow cover. Locally, over regions at higher latitudes or altitudes the dynamics of snow depth may

favour more prolonged ~~periods of snow covers~~ snow cover periods and thus be, further modulated by local environmental conditions. For instance, ~~e~~Extreme weather events such as blizzards, can ~~also~~ lead to rapid changes in snow depth over short ~~period~~ time scales. For these reasons, a major ~~Due to its climatic importance, considerable~~ effort has been made over the years by various countries in the northern hemisphere to carry out extensive and consistent measurements of snow depth.

Snow cover extent (SCE) is an equally important ~~proxy for~~ climate change proxy in the Northern Hemisphere. Its seasonal cycle drives a number of important energy and water cycle processes. ~~The Arctic land surface is subject to many processes arising from seasonal~~ In Arctic regions, snow cover dynamics ~~that~~ drive the seasonal thermal regime of the ground, with implications for the carbon cycle, permafrost, and terrestrial and freshwater ecosystems. ~~Namely~~ In particular, ~~in~~ during spring ~~when the observed, the~~ retrat of snow cover ~~extent decreases, it~~ directly influence the extent magnitude of the climatic ~~cooling~~ warming.

Another key snow-related variable widely used by the scientific community is snow water equivalent (SWE), which represents the amount of water contained in the snowpack. SWE is particularly valuable because it provides a direct measure of the snowpack's total water content, which is crucial for hydrological applications such as water resource management and runoff prediction. In the Crocus-ERA5 model, snow depth accounts for the density of individual snow layers, while SWE is derived from both snow depth and density. Snow depth is much easier to observe through remote sensing and ground measurements, making it a valuable proxy for validating snowpack models, whereas SWE remains more difficult to measure directly. Satellite-based snow depth relies on SWE estimates and assumed or modeled snow density. This approach introduces significant uncertainties due to spatial variability in snow density, retrieval errors, and sensor limitations. These factors can lead to biases in satellite-derived snow depth estimates. In contrast, in-situ measurements offer reliable, consistent data that are essential for validating snow models like Crocus-ERA5. As a result, this study prioritizes snow depth, which provides broader and more reliable observational coverage. Accurate snow depth data are essential not only for refining SWE estimates but also for enhancing hydrological, flood, and climate models. Recognizing this importance, many Northern Hemisphere countries have maintained long-term, consistent snow depth monitoring programs.

For these reasons, in order to assess the reliability of the Crocus-ERA5 results, this study focuses on both snow depth and snow cover, comparing them with independent observations.

2.1 Atmospheric Forcings

The Crocus snowpack model is forced, in this study, to simulate snow evolution with the ERA5 atmospheric reanalysis (Hersbach et al., 2020), produced by the European Centre for Medium-Range Weather Forecasts (ECMWF). As the latest global meteorological reanalysis, ERA5 represents a significant advancement over ERAI (Dee et al., 2011). Table 1 summarizes the main differences between ERAI and ERA5 reanalyses. One of the most notable improvements is the spatial resolution, with ERA5 offering a resolution of approximately 31 km, compared to 80 km in ERAI. Beyond the finer spatial and temporal resolution, ERA5 also benefits from over a decade of advancements in model physics, core dynamics, and data assimilation.

The ERA5 reanalysis includes an advanced land data assimilation system that has been enhanced to improve the representation of hydrological variables such as soil moisture and snow depth. Both ERAI and ERA5 use a single-layer snow scheme that assumes uniform density and temperature throughout the entire snowpack. Compared to ERAI, ERA5 improves the repre-

130 sensation of key physical processes, including snow accumulation, melting, wind-driven redistribution as interactions with the ground by incorporating better data assimilation techniques. However, ERA5 does not account for vertical variations within the snowpack, such as temperature gradients or changes in snow properties at different depths - features typically represented in multi-layer snow models. Consequently, snow parameters derived from ERA5, including snow depth and SWE, are less detailed and refined than those produced by multi-layer models that explicitly resolve multiple snowpack layers to capture more detailed variations in snow properties.

Table 1. Key differences between ERA-Interim (ERA-I) and ERA5 reanalysis products.

Feature	ERA-I	ERA5
Time coverage	1979–2019	1950–present
Horizontal resolution	0.75° (80 km)	0.25° (31 km)
Temporal resolution	6-hourly data	Hourly data
Vertical levels	60 levels (top at 0.1 hPa)	137 levels (top at 0.01 hPa)
Data assimilation system	4D-Var (IFS Cycle 31r2)	4D-Var (IFS Cycle 41r2)
Land surface model	TESSEL	HTESSEL (updated version)
Snowpack structure	Single-layer scheme	Single-layer scheme
Advantages	Longer-established dataset, widely validated in past studies	Higher resolution, ore detailed atmospheric processes, updated physics, improved land-surface interactions

2.2 Crocus-ERA5 Framework

Crocus is coupled to the ISBA (Interactions between Soil–Biosphere–Atmosphere) land surface model (Vionnet et al., 2012; Brun et al., 2013) and embedded in the SURFEX numerical platform (Masson et al., 2013, <https://www.umr-cnrm.fr/surfex/>).

135 This coupling, as described by Vionnet et al. (2012), allows Crocus to interact with other surface components such as soil, low vegetation, and atmosphere within a consistent framework. Within this setup, Crocus is forced by key atmospheric variables, including 2m air temperature, humidity, wind speed, incoming shortwave and longwave radiation, and precipitation (both rain and snow). Integrated into SURFEX, Crocus benefits from a shared surface energy balance and atmospheric forcing structure, enabling more realistic simulations of snowpack evolution in complex surface environments. A full description of the

140 Crocus model and the technical configuration of the Crocus Northern Hemisphere Snowpack product can be found in Vionnet et al. (2012) and Brun et al. (2013), respectively.

The growing needs of the scientific community for improved snow modeling began gaining momentum in the early 2000s, driven by an increased focus on the impacts of climate change and water resources. Since the early 2010, the availability of high-resolution atmospheric reanalysis datasets has encouraged the use of one-dimensional multi-layer mass and energy balance models over large areas, improving the understanding of snowpack evolution and the simulation of snow cover and

145

albedo. The integration of these enhanced atmospheric reanalysis forcings has further increased model accuracy, particularly in representing snow season dynamics and refining albedo estimates through explicit modeling of snow grain types.

Until 2020, the Crocus model was driven by meteorological forcing (temperature, precipitation, humidity, wind, etc.) derived from the ERAI global atmospheric reanalysis. Building on the availability of more advanced reanalysis products, the Crocus-
150 ERA5 product was developed. All snow characteristics – A set of seven snow-related variables – snow depth, snow water equivalent, liquid water content in the snowpack, snow albedo, snow surface temperature, snowpack internal temperature, and snow cover fraction – is freely available on a daily basis at <https://doi.org/10.5281/zenodo.14513248> (Decharme et al., 2024).

The CROCUS-ERA5 dataset also benefits from adopting the standards of the Coupled Model Intercomparison Project Phase 6 (CMIP6), including metadata structured according to ISO 19115. Additionally, CMIP6 data follows the Climate and Fore-
155 cast (CF) metadata conventions (CF-1.7, CMIP-6.2) and complies with the Data Reference Syntax (DRS) for systematic file organization and naming, promoting interoperability.

2.3 Snowpack Modeling

The Crocus multi-layer snow model simulates snow albedo, heat transfer and phase change, snow mass, snow density, and snow grain metamorphism based on experimental laws (Brun et al., 1992, 1989). The number of layers is variable from 1 to 50
160 depending on snow depth and stratification.

~~The model representation of the snowpack does not account for forest, i.e. it only concerns the snowpack of open areas, where only low vegetation is modelled, as in~~
The snowpack model does not explicitly represent forested areas. Instead, these zones are treated as low vegetation, following the approach of Brun et al. (2013). This simplification was a deliberate choice, as the model does not simulate snow-forest interactions. While this improves internal consistency, it comes with some limitations. Boreal Forests significantly
165 impact snow processes through canopy interception, sublimation, and shading effects. Ignoring these mechanisms can lead to overestimation of snow accumulation at ground level, as it neglects snowfall interception by tree canopies and underestimates sublimation of intercepted snow driven by longwave (infrared) radiation. Forest shading also reduces surface albedo, slowing snowmelt. These interactions are complex and remain challenging to simulate accurately, which underscores the trade-off between model simplicity and physical realism. Additionally, in Crocus-ERA5, the snowpack is simulated on an idealised
170 herbaceous surface with a climatological physiography (e.g. no inter-annual variability in leaf area index).

Rising temperatures and increasingly frequent winter thaws are reshaping snowpack structure, leading to notable changes in snow distribution. The Crocus-ERA5 snowpack model quantifies these changes by explicitly simulating snowpack processes, including snow cover fraction. It distinguishes snow-covered areas from snow-free regions, such as vegetation or bare soil within the grid, preventing unrealistic vegetation effects on snow dynamics. This precise partitioning allows for a more ac-
175 curate representation of snow-melt and rain absorption processes, improving the model's reliability in capturing snow cover evolution.

To refine snowpack representation, the model dynamically divides the snowpack into layers, with a minimum of 3 and a maximum of 50, allowing it to capture thermal gradients and conserve mass and heat during melting or accumulation. Processes such solar absorption, water retention, heat transfer, and compaction are calculated based on snow density. In thin snowpacks,

180 solar energy can penetrate to the ground, influencing albedo, whereas deep snowpacks minimize this effect, acting as insulators that reduce heat conduction, suppress turbulent transfer due to low surface roughness, and modify the net radiation budget because of snow's high albedo.

The model estimates land surface albedo by incorporating snow cover's effect on the surface radiation budget, diagnosing the effective total surface albedo (Boone et al., 2017) . This is determined by evaluating the difference between downward
185 radiation components and total surface net radiative fluxes. In snow-covered areas, particularly those overlaying vegetation, the effective surface albedo is significantly altered due to upwelling shortwave and longwave radiative fluxes.

2.4 Analysis Methods

~~Furthermore,~~ Evaluate changes in the estimation of the snowpack from Crocus stand alone involves variations influenced by climate change that are directly taken into account by atmospheric forcing. The two variables retained in our study, snow depth
190 and snow cover, are strongly linked to each other and reflect the direct response of atmospheric forcing providing essential information on the performance of the snowpack modeling. For this reason, the new Crocus-ERA5 product with a horizontal resolution of $1/4^\circ$ is first compared with the previous Crocus-ERA5 version with a resolution of $1/2^\circ$ over the period 1979-2018. The comparison focuses on the monthly variability of snow depth anomalies. It evaluates changes in Arctic snow cover variability due to changes in atmospheric forcing, and to a lesser extent, spatial resolution, separately for North America
195 and Eurasia, as Eurasia, North America, and Greenland are affected differently by the components of the cryosphere. Within the Arctic Circle, the Eurasia pan-region includes Scandinavia (Norway, Sweden, Finland) and northern Russia, while North America includes northern Alaska (USA) and northern Canada.

In this study, the assessment of the Crocus-ERA5 daily snow depth is done against the harmonized dataset of in-situ observations across North America and Eurasia providing a comprehensive view of historical snow conditions. In Brun et al. (2013),
200 this *Crocus-ERA5* daily snow product was validated against local observations from over 1000 monitoring stations in northern Eurasia. Assessing the timing of snow onset and melt simulated by large-scale models is especially challenging due to the rapid and highly variable response of snow to temperature fluctuations. The ability of the Crocus-ERA5 to map snow depth (or mass, related to depth by the density) is thus evaluated against these in-situ observations in terms of time-average and local daily variability. To identify particular errors in the biases of Crocus-ERA5, statistics were calculated for diagnostics derived from
205 continuous snow depth information, such as duration and first/last day of continuous snow following the method described in many previous studies (Brun et al., 2013; Schellekens et al., 2017; Decharme et al., 2016, 2019). Additional diagnostics include the averaged and annual maximum snow depths, the date of maximum snow depth (expressed as days since January 1st) as the number of snow-covered days per year.

Over the years, significant advancements in SCE analysis, particularly through multi-sensor approaches, have greatly improved mapping by reducing the uncertainties associated with sparse, single-sensor data. Issues related to mismatches in temporal and spatial scales - which historically made large-scale validation increasingly reliant on modeling frameworks and gridded interpolations - have thus been reduced.

210

215 The use of SCE provides a distinct advantage, as it represents a cumulated variable that can be reliably derived from satellite observations over large regions. Additionally, expressing snow cover as a percentage (%) offers a valuable advantage for validation purposes, providing a simple, spatially explicit measure of snow presence or absence over a given area. SCE and snow cover percentage can thus be directly compared with satellite-derived estimates, making them practical and robust metrics for evaluating the spatial accuracy of snowpack models.

In this study, SCE is characterized by representing its monthly standardized anomalies for 2000-2022, which directly addresses the main objective of validating the seasonality and inter-annual variability of snow cover changes over large areas.

220 2.5 Observational Data

The in-situ dataset includes over 2,000 stations with daily observations spanning the period from 1950 to 2012. Most measurements are taken at synoptic stations following World Meteorological Organization (WMO) standards, typically representing A series of daily observations were collated for over 2,000 stations spanning the period from 1950 to 2012. Given that a significant number of measurements are conducted at synoptic stations in accordance with the standards set forth by the World Meteorological Organization, the vast majority of the observations pertain to bare ground or open areas with regular grass cutting (Schellekens et al., 2017, <https://doi.org/10.5194/essd-9-389-2017-supplement>). This extensive dataset that was carefully processed to ensure consistency and minimize elevation-related biases offers valuable spatial and temporal coverage, providing is particularly valuable due to its extensive spatial and temporal coverage that offer robust insights into historical snow depth variability and trends. Decharme et al. (2019) described the processing to this dataset namely the difference between outlined specific selection criteria, including: (i) a local and model local-model elevation inferior to of less than 100 m, (ii) a number of days a minimum of 100 days with a non-zero depth measurement greater than 100 days over the whole period over the full period and, (iii) at least eight snowy days per year.

The ability of the Crocus-ERA5 to map snow cover extent is evaluated against the Interactive Multisensor Snow and Ice Mapping System (IMS) satellite data (U.S. National Ice Center, 2008, <https://nsidc.org/data/g02156/versions/1>). The IMS resolution and accuracy have advanced due to the integration of various technologies and data sources, including snow cover analysis incorporates estimates from NOAA AVHRR, MODIS, VIIRS, and Sentinel satellites, alongside in-situ observations and NCEP model data (Helfrich et al., 2007). By integrating time-sequenced satellite images and has become a more efficient tool for validating snow cover over large areas like the Northern Hemisphere imagery, IMS enhances the differentiation between snow and cloud (Estilow et al., 2015), ensuring superior spatial and temporal coverage for more precise and reliable snow cover estimate. The adoption of advanced multi-sensor fusion techniques in 2020 has set IMS apart, allowing it to combine multiple satellite sources and models for an all-weather, day-and-night analysis. This makes IMS more robust and comprehensive than other snow cover analyses, which often lack the same level of real-time updates and multi-source data integration. Additionally, in this study, the period covered by IMS complements that of in-situ observations (1950–2012). To get as close as possible to model resolution, snow cover from IMS is used here at a spatial resolution of 24 km for the period 2000–2022 To align as closely as possible with model resolution, IMS snow cover data is used at a spatial resolution of 24 km for the period 2000–2022.

While Crocus-ERA5 provides valuable insights into snowpack dynamics, it faces limitations in accurately representing different land cover types. Forests are not explicitly represented in Crocus. Although this simplification affects average snow depth estimates, it has a limited impact on inter-annual variability. In forested areas, snow generally melts slightly earlier, but this timing shift does not significantly alter year-to-year trends. Beyond forests, Crocus-ERA5 faces limitations in open areas with low vegetation, such as grasslands and tundra, where it may not accurately capture wind-driven snow redistribution. This can introduce biases in snow depth and duration estimates, affecting the dataset's applicability in regions where blowing snow and drift processes play a dominant role in snowpack evolution.

Despite these challenges, Crocus-ERA5 enhances snow cover simulation by explicitly modeling snowpack processes and distinguishing snow-covered and snow-free areas, surpassing the capabilities of satellite observations alone. While satellites rely on indirect measurements and assumptions, Crocus-ERA5 uses physically based simulations to represent snowpack dynamics more accurately. Its detailed physical representation, combined with increasing spatial and temporal resolution, improves snowpack dynamics simulation, enhances accuracy, and extends dataset applicability across diverse landscapes

3.1 Comparison with the Crocus-ERA5 product

The variability of monthly snow depth is highlighted by the changes in its monthly anomalies over time, both displayed annually and seasonally, for both the North American and Eurasian regions (Figure 1). Anomalies are calculated in relation to the average for the period 1979-2018, which encompasses both products in order to ensure its temporal and spatial consistency. Negatives anomalies indicate lower snow depth, while positive values indicate higher snow depth. In the Northern High latitudes, the snow depth typically increases during the winter months as a result of the accumulation of snowfall. The snow depth from both the Crocus-ERA5 and Crocus-ERA5 products are generally well in phase for both the North American and Eurasian regions. In Eurasia, the decrease in snow depth that started at the beginning of the 21st century is very similar for both simulations, however Crocus-ERA5 shows more pronounced negative anomalies (brown), particularly for 2000-2010. In North America, however, the similarity between the snow depth anomalies of both simulations is more nuanced, and there is a significant inter-annual variability throughout the full data series.

The snow depth on the ground is closely related with the extent of snow cover, particularly in terms of its spatial distribution. Furthermore, the anomaly of snow cover varies considerably depending on the geographic location and local climate conditions. In the northernmost regions of North America and Eurasia where the largest tundra biome is found, the presence of snow cover during the winter months acts as an insulator for the underlying soil limiting the flow of carbon from the northern permafrost. In particular, variations in snow cover in Eurasia can influence the thermal regime of permafrost, which may have consequences for the release of greenhouse gases. In the late spring (April to June), a decrease in snow thickness is observed as temperatures rise, leading to snow melt as in autumn (September to November), the return of colder conditions favours the onset of new snow accumulation, marking the beginning of the seasonal increase in snow cover. Figure 2 shows the standardized anomalies in snow cover extent for land areas situated within the Arctic Circle (latitudes > 60° N) for the period of late spring, from 1979 to 2018, for both the North American and Eurasian Arctic regions. Solid

black and red lines depict 5-yr running means, providing a smoothed trend of variability, whereas filled circles highlight standardized anomalies relative to the last year 2018. Across ~~Over these both~~ regions, the Crocus-ERA5 and the Crocus-ERA1 snow cover products exhibit a closely similar inter-annual variability, with an overall declining trend throughout late spring and early autumn months. However, after the early 2000s, ~~although the~~ North American Arctic anomalies for Crocus-ERA5 in May and June ~~seem to be appear~~ less negative after the beginning of the 2000s. In Eurasia, around 2012, a particularly pronounced reduction in snow cover occurred during late spring and early summer (April-June), preceded by another noticeable decline in the autumn months (September-November) around 2009. These patterns highlight the ongoing and broad seasonal impacts of Arctic warming on snow cover dynamics. The inter-annual variability in Crocus-ERA5 and Crocus-ERA1 is very similar. However, over Eurasia in late spring, Crocus-ERA5 anomalies are slightly more pronounced, in contrast to the North American Arctic, where Crocus-ERA1 anomalies are less pronounced. In autumn, differences in the inter-annual variability are less evident. The yearly standardized anomalies relative to 2018 (filled circles) display a broadly similar pattern with consistent inter-annual variability. They illustrate how the monthly snow cover state in each year compares to 2018 and deviates from the smoothed long-term climate trend. These deviations from the long-term mean are generally larger during the onset of accumulation, particularly in Eurasia (Figure 2, panels g-l). This highlights that early-season snow cover is more sensitive to short-term atmospheric variability, whereas late-spring snow melt is more strongly influenced by long-term climate trends.

The ~~monthly~~ April-May standard deviation of snow depth from its mean on the land surface on the period 1979-2018, highlight the ~~main differences between~~ variability of both products (Figure 3). There is a ~~more significant divergence for~~ clear stability in snow depth distribution, i.e. deviation varies similarly around the mean, for both regions Eurasia ~~than for and~~ North America. In Eurasia, the shift of the CROCUS-ERA5 distribution toward higher values of both the mean and standard deviation suggests a proportional relationship, where greater snow depth corresponds to increased variability. In North America, while the CROCUS-ERA5 standard deviation also increases with the mean, the relationship is less pronounced. In April, the mean monthly snow depth is higher for Crocus-ERA5 compared to Crocus-ERA1 by about 15% (from 0.44 to 0.52 m) in Eurasia and by 8% (from 0.42 to 0.46 m) in North America. The standard deviation also increases by similar proportions, with a rise of approximately 15% (from 0.24 to 0.29 m) in Eurasia and 7% (from 0.24 to 0.26 m) in North America. This significant changes in the statistics, mean and standard deviation, in same proportions suggests that Crocus-ERA5 shifts towards larger values. In May, the Crocus-ERA5 snow depth is higher by about 26% (from 0.17 to 0.23 m) in Eurasia and 16% (from 0.21 to 0.25 m) in North America. The standard deviation likewise increases, with a rise of approximately 24% (from 0.21 to 0.28 m) in Eurasia and 10% (from 0.24 to 0.26 m) in North America. For North America, this increase in mean coupled with a smaller rise in standard deviation suggests that the data is becoming more consistent around higher values, with lower values showing less variability. This significant increase in the Crocus-ERA5 statistics (mean and standard deviation) in similar proportions, with the same spread of points and clusters, aligns with increased snow accumulation and reduced melting. These are linked, respectively, to higher precipitation - particularly in mountainous and mid-to-high latitude regions - and colder temperatures in ERA5 (Wang et al., 2019). Another indirect conclusion is that wind redistribution, which primarily affects snow depth, does not change significantly, as snow depth variability remains stable.




Figure 1. (Panels left) Time series of monthly snow depth anomalies from January 1979 to December 2018, calculated relative to the mean snow depth for 1979-2018. Positive anomalies are shown in blue-green, while negative anomalies are shown in brown. Adapted from Figure 1 in Mudryk et al. (2020). (Panels right) Monthly climatology of snow depth anomalies for Crocus-ERA5 (black line) and IMS (red line). The x-axis represents climatological values, while the y-axis lists months from January (top) to December (bottom).

3.2 Snow depth

The annual cycle of snow depth is characterized by a gradual accumulation of continental snow from October to March, followed by a rapid ablation of the snowpack during the spring (Figure 1). The snow that persists at the advent of spring incorporates precipitation that occurred during the preceding winter, thereby reflecting the atmospheric variability of winter conditions. In order to represent the onset of the spring period, the amount of snow is verified by calculating the time-averaged snow depth on 10 March for two climatological periods for both the Crocus-ERA5 and the in-situ observations products. Two tier of climatic normals have been highlighted: the period 1950-1980 is used to define the past climate and the period 1980-




Figure 2. Monthly time series of snow cover extent (SCE) shown as standardized anomalies relative to the period 1979-2018. Spring and Autumn are represented respectively for months of April-May-June, from (a) to (f), and September-October-November, from (g) to (l). Land surface averages for North America and Eurasia refer to latitudes above 60° N. Solid black and red lines depict 5-yr running means providing a smoothed trend of variability. Filled circles are used to highlight anomalies relative to the last year 2018. Adapted from Figure 1 in Mudryk et al. (2023).

2012 is used to define the present-day climate (Figure 4). The present-day period cover the start of the acceleration in global warming and particularly the rapid warming of the Arctic that has led to the thawing permafrost.

The spatial distribution of the snow depth is generally well reproduced by Crocus-ERA5. Overall, Crocus-ERA5 generally reproduces the snow depth pattern well, as indicated by the similarity between modeled (shaded pattern) and observed (circles) values. Regions where the colors closely match suggest strong agreement between the two datasets. The long-term mean of snow depth on 10 March exceeds 0.3 m for latitudes above 60° in Eurasia and above 45° in North America , with localized variations reflecting regional climatic influences. Crocus-ERA5 tends to slightly overestimate snow depth, particularly in localized areas of




Figure 3. Monthly standard deviation of snow depth from its mean on the land surface for Eurasia (left) and North America (right) with respect to latitudes above 60° N. Representation of the months **(a)** April and **(b)** May.

the central Russian Plain, east of the Ural Mountains, as well as in parts of the Central Siberian Plateau. In Eurasia, maximum of snow depths are found in the north-eastern regions of European Russia and the central Siberian plateau, and minimums in the eastern part of Siberia. Crocus-ERA5 slightly overestimates the snow depth in the central part of the Russian plain to the east of the Ural mountains. For the present period, two stations with strong values (> 0.75 m) differ strongly from Crocus-ERA5, one at the end of the Ural Mountains and the other at the mouth of the Ob Gulf. Otherwise there is good agreement on both sides of the Ural Mountains and on the Siberian Plain for both periods, where also some stations show strong values (> 0.75 m) compared to Crocus-ERA5 (< 0.50 m) in the modern period. Weak snow depths (< 0.30 m) are well simulated by Crocus-ERA5 on the Verkhoïansk and Tcherski mountains, where the large decline in snow depth during the summer months is well represented. This performance has been already highlighted by Brun et al. (2013) with the previous Crocus-ERA1 product in terms of low density, which is explained by the ability of Crocus to simulate the metamorphism of the snowpack layers into depth hoar under extreme temperature gradients. Further west, in the East Siberian lowlands, deviations from observations occur at very low values. Wherever the relief is not significant, in the plain but around the mountain terrain, the observed snow depth compares reasonably well with in situ observations. This was highlighted by Mudryk et al. (2024) in its evaluation of gridded snow water equivalent (SWE) products for their ability to represent climatology, variability and trends in regions spanning the Northern Hemisphere.




Figure 4. Time-averaged snow depth from Crocus-ERA5 (2D field) for the two climatological periods 1950-1980 and 1981-2012 on 10 March superimposed on the corresponding time-averaged observed snow depth (circles).

The upper part of Figure 4 shows the spatial distribution of the time-averaged snow depth in early spring, as simulated by Crocus-ERA5 for the period 1950-2022. The figure depicts the long-term daily snow depth on 10 March for both the Canadian and Eurasian domains. The bottom panels show the results for a selection of stations representing diverse geographical regions. The first Canadian station on Figure 4 (station S2) is situated at an elevated altitude (1323 m) in the vicinity of the Canadian Rocky Mountains. Others stations are located in the Canadian Shield region, which is characterised by numerous hills and glacier-carved lakes. The observed and simulated time series of snow depth on 10 March are in very good agreement over the Canadian Rocky Mountains for the present-day period (station S2), which is characterised by a high variability. In the northern continental part, between 65° and 70° latitude (stations S29 and S25), snow depths exhibit less variability around 0.4 m but also show excellent agreement with observations. In the northern Hudson Bay Lowlands (station S12), there is also a high degree of agreement between the two time series, although less agreement is revealed for the past climate (before 1980). Nevertheless, confidence in the Crocus-ERA5 snow depth in past climates is however supported by its close phase with in situ observations in the Arctic region (S37 and S38). However, in these very high-latitude regions, snow depths are overestimated by Crocus-ERA5, particularly in the north of Baffin Island which is divided into numerous peninsulas (S30).




Figure 5. Location of stations on the spatial distribution of time-averaged snow depth in March for the period 1950-2022 and the two domains: Canada and Eurasia Northern North America (stations on Canada only (top left) and top right, respectively) and Eurasia (top right) Under Below each domain (column) the time series of snow depth for each station is shown for both Crocus-ERA5 (grey lines) and observations (orange circle with dotted lines) on 10 March, early spring.

Each time series panel corresponds to a station point on the map (top). The x-axis starts at the beginning of the longest available time series.

In Eurasia (left panels of Figure 4), in the part of the Ural Mountains near the Kara Sea (station S19), time series of snow depth on 10 March show an overestimation of Crocus-ERA5 in the past climate (1960-1984) with a phase close to the observations. However, in the subsequent present period (1984-2012), Crocus-ERA5 and the in situ observations demonstrated a high degree of agreement. Near the city of Norilsk (station S28), Crocus-ERA5 exhibits three distinct behaviours: an overestimation for the period 1950-1967, followed by an underestimation for 1968-2000, ending with an excellent agreement in phase and magnitude for 2000-2012. On the Central Siberian Plateau, in a station located in a latitudinal valley near the city of Tura (station S16, 277 m), Crocus-ERA5 overestimates snow depth but is remarkably in phase for the entire period of 1950-2015. This significant overestimation of Crocus-ERA5 in relation to observations associated to a remarkable agreement in phase is also shown in the data from the S8 station (682 m), located in the Kolyma mountain. In regions along the Laptev and Siberian Seas (stations S34 and S39), where snow depth varies below 0.5 m, Crocus-ERA5 exhibits a slight tendency to overestimate observations. However, both the model and observations remain in relatively close alignment. Finally, there is a noticeable match in phase and magnitude to the west of the Kamchatka Peninsula (station S2), where deep snow depths generally exceed 1m. Figure 6 compares the Crocus-ERA5 and observations for the period 1950-2012 in terms of average duration, maximum, first and last day of continuous snow on the ground. A day with snow on the ground is defined as a day with more than 1 cm of snow (Brun et al., 2013; Decharme et al., 2019). Overall, Crocus-ERA5 demonstrates a satisfactory level of concordance with the observations. This underlines the efficacy of the model in replicating the seasonal snow cycle in the northern hemisphere, despite the persistence of certain biases. Crocus-ERA5 simulates predominately a shorter snow season in the Arctic compared to observations, while in the sub-Arctic plains it predicts a longer snow season. However, these biases are relatively small in both regions (~2 days, Figure 6.a). Two regions differ significantly from the observations. Around the Rocky Mountains, the Crocus-ERA5 show many more continuous snow days than observations. Additionally, in Norway, the biases alternate strongly between positive and negative values (~20 days). The biases in the first and last snowy days of the continuous snow period indicate that the snow season generally starts later and ends earlier in the Crocus-ERA5 estimates (Figures 6.c and 6.d). The largest discrepancies in snow peak between Crocus-ERA5 estimates and measurements are scattered in the Arctic region, where ERA5 has difficulty producing accurate estimates due to incomplete or sparse observational data. In the western part of Eurasia below 60° latitude, the bias in the maximum date of snowfall is small, approximately 2 days (Figure 6.b and Table 2), allowing accurate estimation and forecasting of changes in snow cover.

Table 2 shows the statistics of some Crocus-ERA5 variables related to the station measurements for the two periods 1950-1980 and 1981-2012. The lowest inter-annual correlations (R) are for the date of maximum snow depth with 0.55 and 0.60 for the periods 1950-1980 and 1981-2012 respectively, although this date remains quite close on average for both in situ observations and Crocus-ERA5, whatever the period. This point is relevant because of long-term trends in climate change may shift the maximum date of snowfall. Our results show that the climate has remained rather stable in the years leading up to 2012. All others variables exhibits significant correlations with $R > 0.8$. The strongest correlations ($R \geq 0.96$) are for the number of days with snow and the duration of continuous snow cover, reflecting the quality of both the Crocus snow model and the precipitation estimated by ERA5.




Figure 6. Bias between measurement stations and Crocus-ERA5 for 1950-2012 concerning their temporal phases: (a) Duration of continuous snow, (b) Maximum date of snowfall, (c) First day of continuous snow and (d) Last day of continuous snow.

Table 2. Statistics of biases between monitoring stations and Crocus-ERA5 for two levels of climatic normals: the period 1950-1980 related to the past climate and the period 1981-2012 related to the present climate.

Dataset Variables	1950-1980					1981-2012				
	Cocus Mean	Obs Mean	Bias	R	RMSE	Cocus Mean	Obs Mean	Bias	R	RMSE
averaged snow depth (m)	0.07	0.07	0.01	0.82	0.06	0.07	0.08	-0.00	0.84	0.06
annual maximum snow depth (m)	0.29	0.30	-0.01	0.80	0.20	0.31	0.34	-0.02	0.84	0.19
date of maximum snow depth*	197.84	199.72	-1.88	0.55	35.24	198.34	200.25	-1.91	0.60	32.30
number of days with snow per year	85.97	86.54	-0.57	0.97	18.49	88.66	90.71	-2.06	0.98	17.17
first day of continuous snow*	112.69	104.85	7.84	0.86	20.12	109.90	101.70	8.20	0.84	20.21
last day of continuous snow*	238.70	249.95	-11.25	0.88	21.92	240.44	252.43	-11.98	0.88	21.36
duration of continuous snow	78.44	75.19	3.25	0.96	23.48	80.99	78.74	2.25	0.96	22.08

*number of days since 1st August




Figure 7. (Panels left)Time series of monthly snow cover extent (SCE) anomalies from January 2000 to December 2022, calculated relative to the mean snow depth for 2000-2022. Positive anomalies are shown in blue-green, while negative anomalies are shown in brown. (Panels right) Monthly climatology of snow cover extentSCE for Crocus-ERA5 (black line) and IMS (red line). The x-axis represents climatological values, while the y-axis lists months from January (top) to December (bottom).

3.3 Snow Cover ~~Extent~~

The monthly ~~snow cover~~SCE anomalies relative to the ~~climatology for the~~period 2000-2022 as its climatology were assessed in terms of their seasonality and inter-annual variability (Figure 7). Although the differences are striking with a high degree of inter-annual variability, three points stand out : (i) the good agreement from 2007 ~~for the months of~~ January to May, ~~overlapping~~overlaps with the spring period (MAM), when ~~the strong~~SCE decreases ~~in snow extent occurs~~sharply – the agreement in March is especially remarkable ; (ii) the disagreement ~~for the month of~~ June when ~~the snow cover continues to decrease~~SCE anomalies is still significant in IMS ; and (iii) the good~~broad~~ agreement for the autumn (SON), ~~although there are~~with consistent inter-annual variability even there are some differences in magnitude.

~~In spring, the seasonal cycle of snow cover is well reproduced, with only~~ Minor differences in small orders of magnitude, that ~~can~~ lead to shifts between positive and negative values, ~~can be seen here or there (e.g. April 2000, 2005, 2008 and 2015).~~ ~~Note that the years 2001, 2002 and 2006 show significant~~ Some zones of positive (green) and negative (brown) SCE anomalies in Crocus-ERA5, but the ~~signal remains consistent with IMS~~can be detected as a ten-year alternation for both Crocus-ERA5 and IMS in autumn. In June, ~~snow~~ coverSCE anomalies are positive and much more pronounced in the IMS ~~dataset for most of the climatological period~~indicating that snow remains longer on the land surface (Figure 7, left panel); this is even more evident in the climatology, where the Crocus-ERA5 SCE (black curve) drops sharply to near zero while the IMS curve (red) still shows significant snow cover (Figure 7, right panel).

~~snow cover~~SCE varies considerably with geographical location and local climatic conditions, particularly in the northernmost regions when vegetation emerges in early summer. ~~it~~However because Crocus-ERA5 does not take into account interactions with vegetation anomalies, ~~which it~~ can significantly alter these dynamics in early summer - a limitation particularly evident in Figure 9, for spring (MAM). As a result, snowmelt in Crocus-ERA5, the snow melting occurs more rapidly than in IMS over northern western Canada, southern eastern Canada and eastern Siberia. ~~By limiting incident radiations at the snow surface, the presence of the boreal forest prolongs the duration of snow cover in comparison to areas devoid of such vegetation.~~ Furthermore, since Crocus-ERA5 only represents open areas with low vegetation, it may fail to accurately capture wind-driven snow redistribution, resulting in bi- ases in snow depth and duration estimates in regions where blowing snow and drifting processes strongly influence snowpack evolution.

Figure 8 illustrates the evolution of the monthly anomalies of continental ~~snow cover~~SCE inacross the Northern Hemisphere from Crocus-ERA5 and IMS for the period 2000-2022 ~~is shown in Figure 8.~~ The correlation between the two datasets is 0.72, ~~with a p-value of 0.87,~~ indicating ~~that the two curves come~~they originate from the same distribution. Both time series show very similar seasonal variability, but there are differences in the magnitude of the peaks, which may explain the difference in trends. The long-term trend of ~~snow cover~~SCE in the northern hemisphere from 2000 to 2022 is slightly downward in both time series although the Crocus-ERA5 exhibits a steeper decline compared to IMS with respectively a slope of -1.9×10^{-12} and -4.5×10^{-13} . ~~However, the Mean Absolute Error (MAE) between the two curves is of 0.066 showing that the average absolute difference between the two series is relatively small.~~ In physical terms, these slopes suggest a gradual trend in snow cover anomalies over 2000–2022~~that is,~~ evolving slowly over very large timescales. ~~It might make sense as a~~This highlights ~~of these minimal~~the progressive nature of changes.




Figure 8. Monthly Time series from January 2000 to December 2022 of monthly ~~snow cover fraction~~SCE standardized anomalies presented as a continuous line graph to illustrate trends and variations over the specified period.

The ~~overall~~ Northern hemisphere-wide ~~area covered by snow~~snow cover fraction from Crocus-ERA5 is compared to ~~those that~~ from the IMS long-term analysis to assess biases (Figure 9). During autumn (SON) and winter (DJF), above 60°N, there is strong agreement between Crocus-ERA5 and IMS, as most of continental North America and Eurasia is entirely covered with snow. However, below 60°N, in mountainous regions where snow cover is highly uneven, a positive bias is found between Crocus-ERA5 and IMS. This may be related to a poor representation of the topography at 0.25° resolution and thus to an overly simplistic relationship between simulated snow height and diagnosed snow cover used in Crocus in mountainous regions at this resolution. Snow extent seasonality is closely related to air surface temperature and radiative forcing, and in autumn and winter (low radiative forcing) is influenced by both temperature and precipitation. On the plains, and with accurate atmospheric forcing, Crocus-ERA5 shows excellent performance.

During spring (MAM), snow thickness decreases as temperatures (and radiative forcing) increase, leading to snow thinning and eventual melting. The spring thaw typically starts at lower latitudes and elevations and gradually moves northwards and to higher elevations as the season progresses. In Crocus-ERA5, ~~the snow melts~~ing seems to occur more rapidly than in IMS over northern western Canada, southern eastern Canada and eastern Siberia. This ~~weakness is well known for models that do not take into account~~ the influence of the bias is partly due to the model's lack of representation of boreal forests ~~-on the timing of at high latitudes (below 60°N), which delay~~ snowmelt by limiting ~~incident~~incoming radiation at the snow surface (Decharme et al., 2019). The presence of the boreal forest prolongs the duration of snow cover in comparison to areas devoid of such vegetation. In Crocus-ERA5 the snowpack is simulated as an open-field surface so it does not take into account such process. Moreover, Crocus-ERA5 assumes a fixed herbaceous, so it cannot capture impact of tundra emergence above 60°N, which alter snowpack dynamics in early summer.

In summer (JJA), the negative effect of high altitude on the agreement between the two snow cover products is reduced by the absence of snow. There are negative biases in the northernmost part of the Arctic, excluding Greenland, ~~due to the~~which




Figure 9. Northern hemisphere distributions of mean seasonal snow extent fraction biases of Crocus-ERA5 compared to IMS estimates in winter (DJF), spring (MAM), summer (JJA) and autumn (SON) over the period 2000-2022.

correspond to the lack of snow cover over the melt zone in Crocus-ERA5, largest region of the tundra biome, however the differences during the melt seasons remain inferior to below 10%.

445 In autumn (SON), the emergence of continental snow cover shows significant bias only over mountains range as Yablonoi and Rocky Mountains.

4 Conclusions and Perspectives

This paper proposes a synthetic assessment of current Crocus-ERA5 snowpack over land, is based on the two variables most commonly used representative of snow characteristics and commonly used to monitor snow, snow depth and snow cover. The challenge was to

450 validate model simulations against independent data in order to minimize the uncertainty attributed to the interdependence between explained and independent variables. The strengths of this study include direct comparisons of Crocus-ERA5 with homogeneous long-term time series from in situ snow depth observations and gridded satellite multi-sensor analyses of snow cover fraction. Comparisons reveal a strong overall performance of Crocus-ERA5 in capturing snow dynamics in the Northern Hemisphere.

455 Sensitivity to atmospheric forcing was assessed by comparing ~~the Crocus-ERA5 product with its predecessor,~~ Crocus-ERA-Interim (Crocus-ERA-I), for the common period ~~from~~ 1979 to 2018. The ERA5 forcing ~~includes~~ uses an advanced land data assimilation system, ~~and i.~~ Its higher spatial resolution of 0.25° ~~in latitude/longitude offers~~ provides more detailed representations of topography and land cover ~~compared to~~ than ERA-I, which has a resolution of 0.75°. ~~These improvements translate.~~ This led to snow depth increase and snow cover decrease in Crocus-ERA5 compared to Crocus-ERA-I. Regarding snow depth, Crocus-ERA5
460 is in agreement with Crocus-ERA-I in terms of inter-annual variability and seasonal cycles in Eurasia and North America. It is noteworthy that Crocus-ERA5 tends to produce stronger anomalies, which are consistent with improvements in the atmospheric forcing. A persistent negative trend in snow depth over Eurasia is evident since 2000, while North America shows greater inter-annual variability. The statistics (mean and standard deviation) show a shift in the CROCUS-ERA5 distribution toward higher values. The standard deviation varies similarly around the mean for both Crocus-ERA-I and Crocus-ERA5 distributions, in both regions, Eurasia and to a lesser extent, North America. The Crocus-ERA5 shows generally higher values snow depth than Crocus-ERA-I in spring, especially in Eurasia. The distributions of snow depth highlight differences between both simulations for April and May, with a more significant difference observed in Eurasia than in North America. This comparison between the two products is important for the community that has been using Crocus-ERA-I and is in the process of migrating to Crocus-ERA5.

The ~~analysis of the~~ climatological periods within 1950–1980 and 1981–2012 reveals ~~notable~~ strong agreement in snow depth be-
470 tween Crocus-ERA5 and ~~independent~~ in-situ data. The ability of Crocus-ERA5 to represent daily snow depth (or mass, related to depth by density) is assessed against relative to long-term snow depth on 10 March for various in-situ stations covering different geographic regions in ~~both~~ Canada and Eurasia. ~~South of 70°N in North America,~~ The long-term Crocus-ERA5 snow depth ~~estimates from Crocus-ERA5~~ aligns remarkably well with in-situ observations, demonstrating its ability to capture inter-annual variability. The most significant differences in periodicity and/or magnitude are found in the Arctic Circle, likely due to the challenges of achieving accuracy in reanalysis
475 when using incomplete or sparse observational data in the region. In Eurasia, the results are more mixed and not directly linked to latitude variations. Despite this, Crocus-ERA5 generally captures inter-annual variability well, providing valuable insights into snow depth patterns and trends. Anywhere, Crocus-ERA5 demonstrates a strong ability to reproduce inter-annual variability. In Canada, the simulation performs well across a range of environments, including complex regions as mountains, hills and glacier-carved lakes. In Eurasia, although there is a slight overestimation of snow depth in earlier decades, Crocus-ERA5 shows good phase and amplitude agreement in snow occurrence, particularly
480 from the 1990s onward.

An important limitation of the Crocus-ERA5 product was highlighted by comparing the simulated snow cover with estimates from the IMS multi-sensor product. ~~Crocus-ERA5 simulates the Northern Hemisphere snow cover on an "open field" surface neglecting the boreal forests.~~ Compared to the IMS multi-sensor product, Crocus-ERA5 performs less effectively in early summer due to its idealized herbaceous surface with fixed climatological properties and its omission of forest-snow interactions. This hypothesis leads to a significant simplification intro-

485 ~~duces a bias during spring melt, which occurs too quickly in Crocus-ERA5 where the boreal forest is present~~ causing snow disappear too early
in regions where vegetation is evolving. For example, Crocus-ERA5 fails to capture the impact of tundra emergence at higher
latitudes (above 60°N). This underscores that Crocus-ERA5 should not be used to quantify the absolute value of spring snow cover in areas where forests
dominate. However, using this product to quantify the evolution of inter-annual variability in snow cover or its long-term evolution seems quite acceptable. As
490 June especially in Arctic circle. From early autumn to late winter, the snow cover fraction in Crocus-ERA5 shows a positive
bias compared to the IMS dataset in mountainous regions below 60°N, likely due to coarse topographic representation.

To effectively assess water resources and forecast spring runoff, it is essential to measure the total water ~~volume~~ stored in
snowpack, ~~with the~~ using Snow Water Equivalent (SWE) ~~being as~~ a key indicator. SWE ~~is more accurate than snow depth and/or cover,~~
~~as it accounts for~~ reflects snow density and compaction. ~~Furthermore, SWE has a profound impact on and~~ plays a critical role in Arctic
495 amplification and climate change ~~that is altering the~~ by influencing snowmelt timing and extent. ~~of snowmelt. Under these conditions,~~
~~a spatially comprehensive and temporally continuous monitoring system is essential for tracking the total amount of water stored in solid form within the~~
~~snowpack, particularly in regions with limited in situ data.~~ There are major differences in time and spatial resolution between existing
SWE products (Mudryk et al., 2024), which significantly ~~restrain~~ limits their usefulness in studies of the cryosphere and climate
change ~~studies. The~~. Long-term time series ~~of SWE obtained through~~ from climate models or ~~derived directly from~~ remote sensing ~~has~~
500 ~~adata have~~ spatial accuracy often limited, especially in areas where snow processes, such as melting and compaction, are
not well-represented. Winkler et al. (2021) proposed a method to estimate SWE using only snow depths ~~and their changes~~, which
outperforms models relying on empirical regressions. Fontrodona-Bach et al. (2023) regionalized this method to create the NH-
SWE dataset for the Northern Hemisphere. Shao et al. (2022) developed a high-precision SWE (~~Snow Water Equivalent~~) product
by integrating various existing SWE data sources into a Ridge Regression Model (RRM), which uses machine learning. The
505 temporal resolution of the RRM SWE product is daily, and the spatial resolution is 10 km. The study demonstrated that this
method is effective for creating SWE products on a global scale, offering seamless spatial and temporal coverage. The RRM
SWE product minimizes dependence on a single SWE dataset, optimally utilizing multiple SWE sources and considering
altitude. While this paper focuses on direct spatio-temporal validation of the Crocus-ERA5 snow product using independent
data, an interesting future direction would be to evaluate SWE from Crocus-ERA5 against these newly ~~developed~~ datasets. It
510 should be noted, however, that recently Mudryk et al. (2024) showed that Crocus-ERA5 was one of the most effective product
for reproducing SWE in the Northern Hemisphere, at least in plain areas.

5 Data availability

The new **Crocus-Era5** dataset is free to access and available at <https://doi.org/10.5281/zenodo.14513248> (Decharme et al.,
2024). The dataset is provided over the period 1950-2023 in netcdf format and contains modelled daily snow depth, snow water
515 equivalent, liquid water content in the snowpack, snow albedo, snow surface temperature, snowpack internal temperature, and
snow cover fraction. The previous **Crocus-Era-Interim** dataset is free to access and available at [https://doi.org/10.5281/zenodo.](https://doi.org/10.5281/zenodo.14513040)
14513040 (Decharme, 2024). The dataset is provided over the period 1979-2019 in netcdf format and contains modelled daily

snow depth and snow water equivalent, as well as monthly snow surface temperature, snowpack internal temperature, and snow cover fraction.

520 *Author contributions.* BD defined the scientific framework, performed the numerical simulations and supervised the findings. SRB defined the methodology and selected the relevant datasets, pre-processed the observational data, developed the analytic calculations, performed the analysis and wrote the manuscript. AB and LF performed the ERA5 and the ERA-Interim atmospheric forcing, respectively.

Competing interests. The authors declare that they have no competing interests.

Acknowledgements. We acknowledge the National Snow and Ice Data Center for making the Interactive Multisensor Snow and Ice Mapping
525 System (IMS, U.S. National Ice Center 2008) data available and making this comparison possible.

References

- Boone, A., Samuelsson, P., Gollvik, S., Napoly, A., Jarlan, L., Brun, E., and Decharme, B.: The interactions between soil–biosphere–atmosphere land surface model with a multi-energy balance (ISBA-MEB) option in SURFEXv8–Part 1: Model description, *Geoscientific Model Development*, 10, 843–872, 2017.
- 530 Brun, E., Martin, E., Simon, V., Gendre, C., and Coleou, C.: An energy and mass model of snow cover suitable for operational avalanche forecasting, *Journal of glaciology*, 35, 333–342, <https://doi.org/10.3189/S0022143000009254>, 1989.
- Brun, E., David, P., Sudul, M., and Brunot, G.: A numerical model to simulate snow-cover stratigraphy for operational avalanche forecasting, *Journal of Glaciology*, 38, 13–22, <https://doi.org/10.3189/S0022143000009552>, 1992.
- Brun, E., Vionnet, V., Boone, A., Decharme, B., Peings, Y., Valette, R., Karbou, F., and Morin, S.: Simulation of northern Eurasian local snow
535 depth, mass, and density using a detailed snowpack model and meteorological reanalyses, *Journal of Hydrometeorology*, 14, 203–219, <https://doi.org/10.1175/JHM-D-12-012.1>, 2013.
- Decharme, B.: Crocus-ERA-Interim daily snow product over the Northern Hemisphere at 0.5° resolution [Data set], <https://doi.org/10.5281/zenodo.14513040>, 2024.
- Decharme, B., Brun, E., Boone, A., Delire, C., Le Moigne, P., and Morin, S.: Impacts of snow and organic soils parameterization on northern
540 Eurasian soil temperature profiles simulated by the ISBA land surface model, *The Cryosphere*, 10, 853–877, <https://doi.org/10.5194/tc-10-853-2016>, 2016.
- Decharme, B., Delire, C., Minvielle, M., Colin, J., Vergnes, J.-P., Alias, A., Saint-Martin, D., Séférian, R., Sénési, S., and Voldoire, A.: Recent changes in the ISBA-CTrip land surface system for use in the CNRM-CM6 climate model and in global off-line hydrological applications, *Journal of Advances in Modeling Earth Systems*, 11, 1207–1252, <https://doi.org/10.1029/2018MS001545>, 2019.
- 545 Decharme, B., Barbu, A., and Ramos Buarque, S.: Crocus-ERA5 daily snow product over the Northern Hemisphere at 0.25° resolution, <https://doi.org/10.5281/zenodo.14513248>, 2024.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J. J.,
550 Park, B. K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J. N., and Vitart, F.: The ERA-Interim reanalysis: Configuration and performance of the data assimilation system, *Quarterly Journal of the royal meteorological society*, 137, 553–597, <https://doi.org/10.1002/qj.828>, 2011.
- Derksen, C. and Mudryk, L.: Assessment of Arctic seasonal snow cover rates of change, *The Cryosphere*, 17, 1431–1443, <https://doi.org/10.5194/tc-17-1431-2023>, 2023.
- 555 Dunn, R. J. H., Miller, J. B., Willett, K. M., Gobron, N., Ades, M., Adler, R., Alexe, M., Allan, R. P., Anderson, J., Anneville, O., Aono, Y., Arguez, A., Arosio, C., Augustine, J. A., Azorin-Molina, C., Barichivich, J., Barnes, J. E., Beck, H. E., Bellouin, N., Benedetti, A., Blagrove, K., Blenkinsop, S., Bock, O., Bodin, X., Bosilovich, M., Boucher, O., Buechler, D., Buehler, S. A., Campos, D., Carrea, L., Chang, K.-L., Christiansen, H. H., Christy, J. R., Chung, E.-S., Ciasto, L. M., Clingan, S., Coldewey-Egbers, M., Cooper, O. R., Cornes, R. C., Covey, C., Créatux, J.-F., Crimmins, T., Cropper, T., Crotwell, M., Culpepper, J., Cusicanqui, D., Davis, S. M., de Jeu, R. A. M.,
560 Degenstein, D., Delaloye, R., Dokulil, M. T., Donat, M. G., Dorigo, W. A., Dugan, H. A., Durre, I., Dutton, G., Duveiller, G., Estilow, T. W., Estrella, N., Fereday, D., Fioletov, V. E., Flemming, J., Foster, M. J., Franz, B., Frith, S. M., Froidevaux, L., Füllekrug, M., Garforth, J., Garg, J., Gibbes, B., Goodman, S., Goto, A., Gruber, A., Gu, G., Hahn, S., Haimberger, L., Hall, B. D., Harris, I., Hemming, D. L.,

- Hirschi, M., peng Ho, S., Holzworth, R., Hrbáček, F., Hu, G., Hurst, D. F., Inness, A., Isaksen, K., John, V. O., Jones, P. D., Junod, R., Kääb, A., Kaiser, J. W., Kaufmann, V., Kellerer-Pirklbauer, A., Kent, E. C., Kidd, R., Kipling, Z., Koppa, A., Kraemer, B. M., Kramarova, N., Kruger, A., Fuente, S. L., Laas, A., Lan, X., Lang, T., Lantz, K. O., Lavers, D. A., Leblanc, T., Leibensperger, E. M., Lennard, C., Liu, Y., Loeb, N. G., Loyola, D., Maberly, S. C., Madelon, R., Magnin, F., Matsuzaki, S.-I., May, L., Mayer, M., McCabe, M. F., McVicar, T. R., Mears, C. A., Menzel, A., Merchant, C. J., Meyer, M. F., Miralles, D. G., Moesinger, L., Monet, G., Montzka, S. A., Morice, C., Mrekaj, I., Mühle, J., Nance, D., Nicolas, J. P., Noetzli, J., Noll, B., O’Keefe, J., Osborn, T. J., Park, T., Parrington, M., Pellet, C., Pelto, M. S., Petersen, K., Phillips, C., Pierson, D., Pinto, I., Po-Chedley, S., Pogliotti, P., Polvani, L., Preimesberger, W., Price, C., Pulkkanen, M., Randel, W. J., Rémy, S., Ricciardulli, L., Richardson, A. D., Robinson, D. A., Rocha, W., Rodell, M., Rodriguez-Fernandez, N., Rosenlof, K. H., Rozanov, A., Rozkošný, J., Rusanovskaya, O. O., Rutishauser, T., Sabeerali, C. T., Sánchez-Lugo, A., Sawaengphokhai, P., Schenzinger, V., Schlegel, R. W., Schmid, M., Schneider, U., Sezaki, F., Sharma, S., Shi, L., Shimaraeva, S. V., Silow, E. A., Simmons, A. J., Smith, S. L., Soden, B. J., Sofieva, V., Sparks, T. H., Sreejith, O., Stackhouse, P. W., Stauffer, R., Steinbrecht, W., Steiner, A. K., Stradiotti, P., Streletskiy, D. A., Surendran, D. E., Thackeray, S. J., Thibert, E., Timofeyev, M. A., Tourpali, K., Tye, M. R., van der A, R., van der Schalie, R., van der Schrier, G., van Vliet, A. J., Verburg, P., Vernier, J.-P., Vimont, I. J., Virts, K., Vivero, S., Vömel, H., Vose, R. S., Wang, R. H. J., Wang, X., Warnock, T., Weber, M., Wiese, D. N., Wild, J. D., Williams, E., Wong, T., Woolway, R. I., Yin, X., Zeng, Z., Zhao, L., Zhou, X., Ziemke, J. R., Ziese, M., Zotta, R. M., Zou, C.-Z., Allen, J., Camper, A. V., Haley, B. O., Hammer, G., Love-Brotak, S. E., Ohlmann, L., Noguchi, L., Riddle, D. B., and Veasey, S. W.: Global Climate, Bulletin of the American Meteorological Society, 104, S11 – S145, <https://doi.org/10.1175/BAMS-D-23-0090.1>, 2023.
- Estilow, T. W., Young, A. H., and Robinson, D. A.: A long-term Northern Hemisphere snow cover extent data record for climate studies and monitoring, *Earth System Science Data*, 7, 137–142, <https://doi.org/10.5194/essd-7-137-2015>, 2015.
- Fontrodona-Bach, A., Schaeffli, B., Woods, R., Teuling, A. J., and Larsen, J. R.: NH-SWE: Northern Hemisphere Snow Water Equivalent dataset based on in situ snow depth time series, *Earth System Science Data*, 15, 2577–2599, <https://doi.org/10.5194/essd-15-2577-2023>, 2023.
- Helfrich, S. R., McNamara, D., Ramsay, B. H., Baldwin, T., and Kasheta, T.: Enhancements to, and forthcoming developments in the Interactive Multisensor Snow and Ice Mapping System (IMS), *Hydrological Processes: An International Journal*, 21, 1576–1586, <https://doi.org/10.1002/hyp.6720>, 2007.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., et al.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.3189/10.1002/qj.3803>, 2020.
- Kouki, K., Luojus, K., and Riihelä, A.: Evaluation of snow cover properties in ERA5 and ERA5-Land with several satellite-based datasets in the Northern Hemisphere in spring 1982–2018, *The Cryosphere*, 17, 5007–5026, <https://doi.org/10.5194/tc-17-5007-2023>, 2023.
- Masson, V., Le Moigne, P., Martin, E., Faroux, S., Alias, A., Alkama, R., Belamari, S., Barbu, A., Boone, A., Bouysse, F., Brousseau, P., Brun, E., Calvet, J.-C., Carrer, D., Decharme, B., Delire, C., Donier, S., Essaouini, K., Gibelin, A.-L., Giordani, H., Habets, F., Jidane, M., Kerdraon, G., Kourzeneva, E., Lafaysse, M., Lafont, S., Lebeaupin Brossier, C., Lemonsu, A., Mahfouf, J.-F., Marguinaud, P., Mokhtari, M., Morin, S., Pigeon, G., Salgado, R., Seity, Y., Taillefer, F., Tanguy, G., Tulet, P., Vincendon, B., Vionnet, V., and Voldoire, A.: The SURFEXv7.2 land and ocean surface platform for coupled or offline simulation of earth surface variables and fluxes, *Geoscientific Model Development*, 6, <https://doi.org/10.5194/gmd-6-929-2013>, 2013.
- Meredith, M., Sommerkorn, M., Cassota, S., Derksen, C., Ekaykin, A., Hollowed, A., and Kofinas, G.: Polar Regions, in: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate, edited by: Pörtner, H.-O., Roberts, D. C., Masson-Delmotte, V., Zhai, P.,

- Tignor, M., Poloczanska, E., Mintenbeck, K., Alegría, A., Nicolai, M., Okem, A., Petzold, J., Rama, B., and Weyer, N. M., p. 203–320, Cambridge University Press, <https://doi.org/10.1017/9781009157964.005>, 2022.
- Mortimer, C., Mudryk, L., Derksen, C., Luoju, K., Brown, R., Kelly, R., and Tedesco, M.: Evaluation of long-term Northern Hemisphere snow water equivalent products, *The Cryosphere*, 14, 1579–1594, <https://doi.org/10.5194/tc-14-1579-2020>, 2020.
- 605 Mudryk, L., Derksen, C., Kushner, P. J., and Brown, R.: Characterization of Northern Hemisphere Snow Water Equivalent Datasets, 1981–2010, *Journal of Climate*, 28, 8037 – 8051, <https://doi.org/10.1175/JCLI-D-15-0229.1>, 2015.
- Mudryk, L., Santolaria-Otín, M., Krinner, G., Ménégos, M., Derksen, C., Brutel-Vuilmet, C., Brady, M., and Essery, R.: Historical Northern Hemisphere snow cover trends and projected changes in the CMIP6 multi-model ensemble, *The Cryosphere*, 14, 2495–2514, <https://doi.org/10.5194/tc-14-2495-2020>, 2020.
- 610 Mudryk, L., Elias Chereque, A., Derksen, C., Luoju, K., and Decharme, B.: NOAA Arctic Report Card 2023: Terrestrial Snow Cover, <https://doi.org/10.25923/xqwa-h543>, 2023.
- Mudryk, L., Mortimer, C., Derksen, C., Elias Chereque, A., and Kushner, P.: Benchmarking of SWE products based on outcomes of the SnowPEX+ Intercomparison Project, *EGUsphere*, 2024, 1–28, <https://doi.org/10.5194/egusphere-2023-3014>, 2024.
- Pörtner, H.-O., Roberts, D. C., Masson-Delmotte, V., Zhai, P., Tignor, M., Poloczanska, E., Weyer, N., et al.: The ocean and cryosphere in a changing climate, IPCC special report on the ocean and cryosphere in a changing climate, 1155, 10–1017, <https://doi.org/10.1017/9781009157964>, 2019.
- 615 Ramos Buarque, S. and Salas y Melia, D.: Link between the North Atlantic Oscillation and the surface mass balance components of the Greenland Ice Sheet under preindustrial and last interglacial climates: a study with a coupled global circulation model, *Climate of the Past*, 14, 1707–1725, <https://doi.org/10.5194/cp-14-1707-2018>, 2018.
- 620 Schellekens, J., Dutra, E., Martínez-de La Torre, A., Balsamo, G., Van Dijk, A., Sperna Weiland, F., Minvielle, M., Calvet, J.-C., Decharme, B., Eisner, S., et al.: A global water resources ensemble of hydrological models: the earthH2Observe Tier-1 dataset, *Earth System Science Data*, 9, 389–413, <https://doi.org/10.5194/essd-9-389-2017>, 2017.
- Shao, D., Li, H., Wang, J., Hao, X., Che, T., and Ji, W.: Reconstruction of a daily gridded snow water equivalent product for the land region above 45° N based on a ridge regression machine learning approach, *Earth System Science Data*, 14, 795–809, <https://doi.org/10.5194/essd-14-795-2022>, 2022.
- 625 U.S. National Ice Center: IMS Daily Northern Hemisphere Snow and Ice Analysis at 1 km, 4 km, and 24 km Resolutions, Version 1, <https://doi.org/10.7265/N52R3PMC>, 2008.
- Vionnet, V., Brun, E., Morin, S., Boone, A., Faroux, S., Le Moigne, P., Martin, E., and Willemet, J.-M.: The detailed snowpack scheme Crocus and its implementation in SURFEX v7.2, *Geoscientific model development*, 5, 773–791, <https://doi.org/10.5194/gmd-5-773-2012>, 2012.
- 630 Wang, C., Graham, R. M., Wang, K., Gerland, S., and Granskog, M. A.: Comparison of ERA5 and ERA-Interim near-surface air temperature, snowfall and precipitation over Arctic sea ice: effects on sea ice thermodynamics and evolution, *The Cryosphere*, 13, 1661–1679, 2019.
- Winkler, M., Schellander, H., and Gruber, S.: Snow water equivalents exclusively from snow depths and their temporal changes: the Δ_{SNOW} model, *Hydrology and Earth System Sciences*, 25, 1165–1187, <https://doi.org/10.5194/hess-25-1165-2021>, 2021.