



Residential heating emissions for the Western Balkans

Christian Asker¹, Eef van Dongen¹, and Olivier Tasse¹

¹Swedish Meteorological and Hydrological Institute, SMHI FoU, 601 76 Norrköping, Sweden

Correspondence: Christian Asker (christian.asker@smhi.se)

Abstract. Air pollution adversely affects health, ecosystems and infrastructure. In the *Western Balkans* (Albania, Bosnia and Herzegovina, Kosovo¹, Montenegro, Republic of North Macedonia and Serbia), the air pollution situation is more adverse than in the European Union in general. Understanding the air quality situation requires high quality emission data, with high resolution spatial distribution, especially for enabling remediation efforts.

5 In this work we have calculated air pollution emissions from heating of individual housing units in the Western Balkan region. The basis for the dataset is a geographical dataset of buildings detected from satellite imagery by artificial intelligence (AI) methods. The building data has been combined with geospatial landuse datasets as well as statistical data for heating needs for residential buildings in the countries included, and finally with emission factors to calculate the heating emissions.

The resulting datasets provides high-resolution heating emission data for common pollutants and are published as open data
10 (Asker, 2024) . When comparing national totals for emissions, the datasets in this work are comparable to other, spatially coarser datasets, though the agreement strongly depends on the fuel usage data for each country/region.

1 Introduction

The Western Balkans (in this work denoted *WB*) region has longstanding issues with poor air quality (UNEP), especially during the winter season. The region has a population of 17 million (World Bank). While traffic, industries, power production and
15 residential heating are all considered important sources of air pollution in the region, there is a lack of detailed emission data for all these sectors. Detailed emission data for the most important sectors is vital in order to understand the air pollution situation and in order to be able to improve the situation. High spatial resolution provides not only a more detailed picture of the sources of air pollution, but may improve the quality of atmospheric dispersion modeling and help find emission hotspots, which in turn can facilitate effective remediation efforts.

20 Emission data from district heating facilities is available for at least some of the cities in the WB. Data for heating emissions from individual housing units (1-3 family houses) is very limited however, even though they are expected to have an important contribution to the total emissions.

The emissions from individual housing units are typically released at a fairly low height (around roof height) and are therefore not dispersed effectively, especially during cold weather and temperature inversions, leading to a high contribution to
25 the concentration of pollutants locally. This makes a detailed description of these emissions even more important.

¹All references to Kosovo in this document shall be understood to be in the context of the United Nations Security Council resolution 1244 (1999)



In this work we present a methodology and resulting dataset for emissions from heating of individual housing units in the WB. The emission dataset has high spatial resolution and covers both urban and rural areas. The dataset is compared with datasets for regional (European) scale and suggestions for improving the methodology are discussed.

2 Input data

30 2.1 Buildings

The starting point is a building dataset for the WB, from the Global ML Building Footprints project (Microsoft), denoted *ML-Buildings* onwards in this report. For the WB region, the dataset contains about 30 million building polygons. For parts of the Republic of North Macedonia and south-eastern Serbia, there are gaps in the MLBuildings dataset. The gaps were replaced by building polygons from the Openstreetmap project (osm). It is worth nothing that for those areas, the Openstreetmap has fairly
35 low level of coverage of buildings and therefore do not compensate fully for the lack of data in the MLBuildings dataset. An example of building polygons for Novi Travnik (Bosnia and Herzegovina) can be seen in Figs. 1 - 2.

2.2 Landuse data

Corine land cover (CLC) (European Environment Agency, 2020a) is a geographic landuse dataset covering Europe, in vector
40 format, having a *minimal mapping unit* (MMU) of 25 hectares. Urban Atlas (UA) (European Environment Agency, 2020b) is a detailed landuse dataset covering european municipalities having at least 100000 residents, with an MMU of 0.25 hectares. These two datasets were merged into a single dataset covering the whole geographical region.

2.3 Energy data

The following fuels have been included:

- 45 – wood
- wood residue
- pellets
- coal
- lpg (liquefied petroleum gas)
- 50 – natural gas
- oil

For some regions only a subset of these fuels have been used since the energy usage data available do not contain all fuels for all regions, nor are all fuels used in all regions.




Figure 1. Example of the MLBuildings dataset for Novi Travnik (Bosnia and Herzegovina).

3 Methodology

55 The gis operations described below used the following softwares: QGIS (QGIS Development Team, 2021), Python + GeoPandas (Jordahl et al., 2020) and GDAL (GDAL/OGR contributors, 2024).

3.1 Filtering buildings

Starting from the MLBuildings + Openstreetmap dataset, the aim was to remove all buildings not 1-3 family housing units.

60 The buildings are all polygons, allowing for calculation of the building footprint area. Incorrect and duplicate geometries where detected and corrected. Then all buildings having a footprint area less than $50 m^2$ or more than $200 m^2$ where removed.

The next step was to merge the landuse data from CLC and UA datasets into a single dataset covering the whole region. The CLC dataset was cutout for the WB region, and the attributes for landuse-classes adjusted to match that of the UA dataset. Further, cutouts were made in the CLC datasets for all cities where UA data was available. Finally the two datasets where

65 merged into a single one.




Figure 2. Example of the MLBuildings dataset for Novi Travnik (Bosnia and Herzegovina). The map shows the same area as in Fig. 1, but with UrbanAtlas and Corine land cover merged as background.

The merged landuse dataset was then joined onto the buildings dataset, so that each building polygon also had attributes for landuse. The following landuse-classes² were assumed to possibly contain some degree of 1-3 family housing units:

- 11100 Continuous urban fabric
- 70 – 11200 Discontinuous urban fabric
- 11210 Discontinuous Dense Urban Fabric (S.L.: 50% - 80%)
- 11220 Discontinuous Medium Density Urban Fabric (S.L.: 30% - 50%)
- 11230 Discontinuous Low Density Urban Fabric (S.L.: 10% - 30%)
- 11240 Discontinuous very low density urban fabric (S.L. < 10%)
- 75 – 11300 Isolated structures
- 21000 Arable land (annual crops)
- 21100 Non-irrigated arable land
- 21200 Permanently irrigated land

²The classes listed here are a combination of the UA and CLC datasets.



- 21300 Rice fields
- 80 – 22000 Permanent crops
- 22100 Vineyards
- 22200 Fruit trees and berry plantations
- 22300 Olive groves
- 23000 Pastures
- 85 – 23100 Pastures
- 24000 Complex and mixed cultivation patterns
- 24100 Annual crops associated with permanent crops
- 24200 Complex cultivation patterns
- 24300 Land principally occupied by agriculture with significant areas of natural vegetation

90 All buildings belonging to landuse-classes other than the above were removed from the dataset. While some of the above landuse classes are mainly agricultural lands, from random sampling of the dataset it can be seen that those landuse classes may contain housing, in particular for areas close to smaller settlements, though the extent of this is different in different regions of the WB region. An example image of the buildings dataset after filtering is shown in figure 3.

95 Comparing the resulting dataset with statistics in the *Typology of Residential Buildings in Bosnia and Herzegovina* (Arnautović-Aksić et al., 2016) and *National Typology of Residential Buildings in Serbia* (Jovanović Popović and Ignjatović, 2013), showed that the number of buildings in the dataset were still quite large. Especially for landuse categories where there is a considerable mix of housing and smaller farm buildings this is likely to be a problem, leading to a possible overestimation of energy usage for such areas.

100 For such areas in Bosnia-Herzegovina and Serbia, by using nearest-neighbour analysis, all buildings that were within 20 meters from another building were selected, for the landuse categories 23100, 24200 and 24300. Of these selected buildings, half were randomly removed. Though these operations reduced the number of buildings for Bosnia-Herzegovina and Serbia, the dataset still contained more buildings than the *Typology of Residential Buildings in Bosnia and Herzegovina*, which may lead to an overestimation of energy usage and therefore emissions there. This is further discussed in the results section.

105 For the region surrounding Strumica in North Macedonia, an issue was discovered during the building analysis. In this area, the MLBuildings AI software had detected a large number of large buildings outside urban areas. Most of these were in fact greenhouses and agricultural lands that had been covered with semi-transparent cloth, leading to the erroneously detected buildings. An example of this is shown in Fig. 4.

110 In order to solve this issue, the bounding boxes, (width, length and angle) of each building polygon were calculated. Then, all buildings having very long-narrow shapes ($width/length < 0.3$) were removed, which removed these structures without removing buildings that should be kept in the dataset. This filtering step was verified by visual inspection of the dataset to make sure that only the right objects were removed.

115 Finally, since the area of each individual house is hard to estimate correctly from the MLBuildings dataset, the polygon geometries are not needed and therefore we convert the polygon layer into centroid points. The outline area size of each individual building is not used to calculate the heating energy needs.




Figure 3. Example of the MLBuildings dataset for Novi Travnik (Bosnia and Herzegovina). The image show the same area as in figures 1 and 2 (above), after the buildings have been filtered to only keep 1-3 family housing units. See text for more details.

3.2 Energy estimation

In each building point, energy consumption for the different fuels are assigned based on available statistical data. The data provide the energy usage per fuel per 1-3 family housing units. The level of detail, in terms of geographical regions and fuels, varies between the different datasources. Each building point in the dataset is processed to calculate the energy usage.

- 120 – Albania: Region-wise energy usage data (3 climate zones), from (Novikova et al., 2015b).
- Bosnia Herzegovina: Entity-based energy usage statistics used, from (Arnautović-Aksić et al., 2016).
- Kosovo: the same data as for Serbia has been used.
- Montenegro: national totals from Eurostat . We assume that all biomass fuels are used for individual housing. The resulting fuel-usage per housing unit is then adjusted for different regions based on degree-days from (Novikova et al., 2018).
- 125 – North Macedonia: statistics for each administrative region (Simovski, 2019).
- Serbia: The same data for the whole country have been used, based on (Jovanović Popović and Ignjatović, 2013).

The full table of energy data used is shown in appendix A1.



Figure 4. False positive building detection on agricultural lands surrounding Strumica, before filtering of such features.



3.3 Emissions to air

130 Once the energy use for each fuel has been assigned to all building points, the resulting emissions for the relevant pollutants may be calculated. The methodology follows the EEA guidebook (European Environment Agency, 2023), for emission factors as well as the composition of heating appliances. For each building point, emissions are calculated for each fuel, based on the composition of heating appliances and the emission factors for each appliance. The following pollutants have been included in the dataset: PM_{10} , $PM_{2.5}$, NO_x and SO_x .

135 The MLbuildings dataset contains data based on satellite imagery captured during multiple years, from 2014 and onwards. Therefore the buildings dataset in itself do not correspond to a well-defined year. Similarly, the energy usage data represents multiple years, 2015-2020, while the appliance composition data in the EEA guidebook is for the year 2010. The yearly change in heating emissions from 1-3 family housing units is not expected to vary considerably between the years in the period 2010-2020. We therefore set the year to be 2019, since it is the most recent year in the period that was not affected by the covid-pandemic.

140 Finally, the point-wise emissions were rasterized into grids for each substance, with 500x500 m horizontal resolution, in the ETRS89-extended / LAEA Europe projection (EPSG:3035). While the calculations of energy and subsequently emissions are done for centroid points for all buildings, the data is rasterized into cells since it might otherwise seem to be more detailed and accurate than it actually is. Further, when using the resulting dataset for atmospheric dispersion modeling or similar, it is computationally more efficient to treat the data as rasters.

145 4 Results and discussion

4.1 Buildings per country

Table 1 lists the number of buildings together with estimates from other sources. Also listed is the percentage of *urban buildings*, calculated from the number of buildings on urban landuse categories (landuse code ≤ 11240 in section 2.2 above). The other sources (references are listed in the table) are different statistical reports and not from national databases of building permits or similar. It is also worth noting that
150 in the different references, the categorization of buildings may differ. As far as possible we have compared the number of buildings in this work to 1-3 family housing units in the references.

For several countries the agreement is fairly good, especially for Albania, Republic of North Macedonia and Serbia. For Kosovo and especially for Bosnia and Herzegovina, the methodology used in this work leads to considerably more buildings, 9 and 30 % more respectively.
155 The resulting number of buildings for Bosnia and Herzegovina may be overestimated, though it is uncertain since there is a lack of independent housing data. It is also uncertain what could cause such overestimation.

Inspection of a random selection of locations suggests that the high number of buildings for Bosnia and Herzegovina is due to the dataset having too many buildings in rural areas where there are smaller farms. This is also supported by the percentage of urban buildings in the dataset, which is considerably lower for Bosnia and Herzegovina (and Montenegro) than for the other countries. This may have a larger effect
160 in Bosnia and Herzegovina than in the other countries. However, the percentage of urban buildings may very well be different in the different WB countries. There are also uncertainties in the references for the number of residential buildings. In order to determine the accuracy of the number of buildings, more independent datasets are required.



Region	buildings in this work	% urban**	buildings comparison	ref
Albania	593970	64.6	576096	(Novikova et al., 2015b)
Bosnia-Herzegovina	1090970	49.8	841543	(Arnautović-Aksić et al., 2016)
Kosovo*	420976	74.1	386046	(agency of statistics, 2022)
Montenegro	179297	55.6	158176	(Novikova et al., 2015a)
Republic of North Macedonia	422274	74.4	426344	(Gjorgjievska, 2021; mkl, 2021)
Serbia	2156724	74.6	2186246	(Jovanović Popović and Ignjatović, 2013)

Table 1. Total number of 1-3 family houses for the WB countries, obtained from the methodology in this work, compared to other statistical sources. * All references to Kosovo in this document shall be understood to be in the context of the United Nations Security Council resolution 1244 (1999). **: percentage of buildings in this work that are on urban landuse categories. See text for more details.

4.2 Energy

165 In figure 5 the total energy is shown. All building points have been rasterized to a 500x500 m grid. It is worth noting some of the the different patterns that can be seen in this image, in particular one in Serbia: The northern part lies on a plateau and have large agricultural areas with settlements in fairly concentrated villages, leading to a "dotted" pattern in the image. The south-eastern part of Serbia, on the other hand, is more mountainous and have smaller agricultural areas scattered between the mountains and the settlements are also more scattered into smaller pieces. This leads to a more continuous pattern in the image.

170 When comparing the countries in the Western Balkans in this image, it is worth noting that the calculated result in figure 5 depend on both the number of buildings and the statistical data for heating energy needs.

An uncertainty not included in this study is the fraction of residential buildings that are not in use. This has not been included due to a lack of available data.

4.3 Emission data

175 Maps of emission data for PM_{2.5}, NO_x and SO_x are shown in figures 6, 7 and 8. Note that the scales in the figures are different. PM₁₀ is not shown, since it is very similar to PM_{2.5}, which is expected for heating emissions due to the fact that most of the particles from combustion in PM₁₀ are fine particles (PM_{2.5}). Both PM_{2.5} and NO_x show similar patterns, while for SO_x the emissions in Montenegro, Albania and Republic of Northern Macedonia are much lower compared to the other countries. This is an effect of the heating energy data for different fuels, which is very low or zero for coal in these three countries, leading to very low SO_x emissions.

180 4.4 Comparison to other emission datasets

The resulting emission data for NO_x and PM_{2.5} were compared to CAMS-reg 4.2, *sector C* (Kuenen et al., 2021, 2022) as well as EDGAR, *sector buildings* (Muntean et al., 2018)³. The CAMS and EDGAR datasets are regional and therefore have a coarser resolution (0.05x0.1 and 0.1x0.1 degrees⁴, respectively) than the dataset in this work which is gridded to 500x500 m.

³ Accessed through https://edgar.jrc.ec.europa.eu/country_profile/, on 2024-09-10.

⁴ 0.1x0.1 degrees corresponds to about 8x11 km for the WB region.



185 The CAMS emissions were reprojected to the ETRS89-extended / LAEA Europe projection (EPSG:3035) and resampled to 1x1 km grid size. Then the total emissions for the different regions were calculated using *zonal statistics* in QGIS. The resulting national statistics is shown in table 2.

Region	Emission NOx (t)			Emission PM2.5 (t)			Emission SOx (t)		
	EDGAR	CAMS	this work	EDGAR	CAMS	this work	EDGAR	CAMS	this work
Albania	1041	382	796	2348	3482	4103	1283	499	67
Bosnia-Herzegovina	2182	1859	5275	6755	16800	46376	3852	5900	20488
Kosovo*	-	479	1798	-	5461	8882	-	1135	9375
Montenegro	-	228	401	-	2008	5316	-	644	88
Republic of North Macedonia	755	644	703	3269	6081	7619	1047	988	167
Serbia	-	2371	9212	-	23414	45505	-	9699	48028
Serbia and Montenegro	11197	3078	11312	40055	30883	59703	14771	11478	57492

Table 2. Comparison of emission totals for different regions in the Western Balkans, CAMS-regional 4.2 (Kuenen et al., 2021, 2022) EDGAR (Muntean et al., 2018) Note: for EDGAR the region *Serbia and Montenegro* is still used, which also contains Kosovo*. The equivalent have been calculated also for CAMS and this work, to facilitate comparison. * *All references to Kosovo in this document shall be understood to be in the context of the United Nations Security Council resolution 1244 (1999)*

190 For NOx, the overall agreement between the two regional dataset and between this work and the regional datasets are similar, though the results in this work are closer to the EDGAR dataset than to CAMS-reg. The largest difference is found for Bosnia and Herzegovina, where this work has more than twice the emissions compared to CAMS-reg and EDGAR.

195 While for NOx the EDGAR dataset has higher emissions than CAMS-reg, for PM2.5 the case is the opposite except for *Serbia and Montenegro* where EDGAR is about a third higher. For PM2.5 the emissions in this work are considerably higher than both regional datasets. Since this was not the case for NOx, the discrepancy is not only a result of the number of buildings, but must also be due to different fuel usage and/or emission factors.

200 Turning to SOx, finally, the data in this work is rather non-consistent when comparing to the regional datasets. While for some countries it is much lower, it is considerably higher for other countries. This is a direct effect of the fuel data, as mentioned above. For Albania, Montenegro and Republic of North Macedonia, there is no coal usage reported (see section 2.3), which lead to very low SOx emissions, while for Bosnia and Herzegovina as well as Serbia, the coal usage lead to high emissions of SOx, in this case much higher than the regional datasets.




Figure 5. Estimated total energy use for heating in individual houses, rasterized to 500x500m. All references to Kosovo in this document shall be understood to be in the context of the United Nations Security Council resolution 1244 (1999)




Figure 6. Calculated emissions of for the Western Balkans for PM_{2.5}, rasterized to 500x500m. All references to Kosovo in this document shall be understood to be in the context of the United Nations Security Council resolution 1244 (1999)




Figure 7. Calculated emissions of for the Western Balkans for NO_x, rasterized to 500x500m.




Figure 8. Calculated emissions of for the Western Balkans for SO_x, rasterized to 500x500m. All references to Kosovo in this document shall be understood to be in the context of the United Nations Security Council resolution 1244 (1999)



5 Conclusions

A complete picture of the air quality situation in a country, region or city requires not only monitoring data, but also an accurate description of the emission sources. Atmospheric dispersion modelling may provide maps of the concentrations for different pollutants and also validate the emission-data. In this work we have developed a dataset for emissions from heating of 1-3 family housing for the whole of the Western Balkans, at high spatial resolution.

While there are multiple sources of uncertainties in the dataset, the basic methodology is

1. obtaining building data, in this case from the MLBuildings dataset
2. filtering of the buildings data, to obtain only the relevant housing units.
3. calculating the heating energy based on available data for energy usage
4. calculating resulting emissions using the EEA emission database

This methodology may be applied wherever the above datasources are available.

Possible improvements in the methodology include better filtering of buildings dataset and, importantly, reliable energy usage data for each region. While the methodology is tailor-made for heating from 1-3 family housing units, other residential housing (apartment buildings) is not included. There are also errors in the MLBuildings dataset, which may be reduced in the future as higher-quality satellite imagery as well as even better building detection AI methods emerge.

Comparing the national totals from the dataset developed in this work with regional emission datasets suggests that the results have acceptable quality and may be used as a basis for emissions analysis, development of air quality action plans as well as input for atmospheric dispersion modeling, which in turn may validate the emission data or lead to improvements in the methodology. It is the hope of the authors that the dataset may help in improving the air pollution situation in the Western Balkans.

Data availability. The emission GIS-files (rasters) created in this work are publicly available at <https://doi.org/10.5281/zenodo.13906810> (Asker, 2024).



Appendix A: Energy use per country and region

Country	region/entity	wood	wood-residue	pellets	coal	lpg	natural gas	oil
Albania	climate zone A	1.2853	0.0	0.0	0.0	0.0	2.6509	0.0
Albania	climate zone B	1.8538	0.0	0.0	0.0	0.0	3.0575	0.0
Albania	climate zone C	8.1657	0.0	0.0	0.0	0.0	7.7732	0.0
Bosnia and Herzegovina	Federation BiH	8.428	2.744	0.0	5.292	0.196	2.94	0.0
Bosnia and Herzegovina	Republika Srpska	13.054	1.926	0.0	5.992	0.428	0.0	0.0
Bosnia and Herzegovina	Brčko district	12.524	1.212	0.0	6.262	0.202	0.0	0.0
Kosovo*	-	3.93	0.0	0.0	6.71	0.0	4.23	1.71
Montenegro	climate zone I	7.26	0.52	0.26	0.0	0.0	0.0	0.0
Montenegro	climate zone II	11.31	0.81	0.41	0.0	0.0	0.0	0.0
Montenegro	climate zone III	15.16	1.08	0.55	0.0	0.0	0.0	0.0
Republic of North Macedonia	Northeastern	7.10	0.00621	0.671668	0.0	0.006869	0.07089	0.0124768
Republic of North Macedonia	Skopje	3.02	0.01081	0.410650	0.05500	0.154395	0.0	0.111276
Republic of North Macedonia	Southwestern	10.32	0.01127	0.547852	0.0	0.078288	0.0	0.022153
Republic of North Macedonia	Eastern	11.04	0.00575	0.413996	0.0	0.118660	0.0	0.101164
Republic of North Macedonia	Southeastern	8.18	0.0	0.329859	0.0	0.045015	0.42060	0.048908
Republic of North Macedonia	Pelagonia	6.33	0.0	0.845203	0.0	0.092912	0.0	0.0
Republic of North Macedonia	Vardar	6.36	0.0	1.047899	0.0	0.060023	0.0	0.0
Republic of North Macedonia	Polog	8.59	0.00253	1.324693	0.050417	0.096611	0.0	0.282571
Serbia	-	3.93	0.0	0.0	6.71	0.0	4.23	1.71

Table A1. Energy usage (MWh/y) per residential building for each country and region. For Albania the regions follow (Novikova et al., 2015b) while for Montenegro (Novikova et al., 2015a). *All references to Kosovo in this document shall be understood to be in the context of the United Nations Security Council resolution 1244 (1999)

Author contributions. CA: conceptualization, methodology, data processing, analysis, visualization, writing. EvD: methodology, verification of emission calculations, writing. OT: project management, writing

225 *Competing interests.* The contact author has declared that none of the authors has any competing interests.

Acknowledgements. The authors would like to thank the Swedish Environmental Protection Agency (SEPA) and environmental agencies in the Western Balkan countries for collaboration.



References

- Open Street Map, <http://www.openstreetmap.org>, accessed: 2024-09-10.
- 230 SUMMARY OF THE RESEARCH ON HEATING PRACTICES IN THE RESIDENTIAL SECTOR OF NORTH MACEDONIA, Tech. rep., RES Foundation, 2021.
- agency of statistics, K.: Results of the Survey on Income and Living Conditions (SILC), 2022, Tech. rep., Kosovo agency of statistics, 2022.
- Arnautović-Aksić, D., Burazor, M., Delalić, N., Gajić, D., Gvero, P., Kadrić, D., Kotur, M., Salihović, E., Todorović, D., and Zagora, N.:
235 TYPOLOGY OF RESIDENTIAL BUILDINGS IN BOSNIA AND HERZEGOVINA, Tech. rep., Arhitektonski fakultet Univerziteta u Sarajevu, ISBN: 978-9958-691-51-5, 2016.
- Asker, C.: Emissions for individual housing in the Western Balkans, <https://doi.org/10.5281/zenodo.13906810>, 2024.
- European Environment Agency: Corine Land Cover (CLC) 2018, <https://doi.org/10.2909/71c95a07-e296-44fc-b22b-415f42acdf0>, 2020a.
- European Environment Agency: Urban Atlas 2018, <https://doi.org/10.2909/fb4dffa1-6ceb-4cc0-8372-1ed354c285e6>, 2020b.
- European Environment Agency: EMEP/EEA air pollutant emission inventory guidebook 2023 – Technical guidance to prepare national
240 emission inventories, Publications Office of the European Union, <https://doi.org/doi/10.2800/795737>, 2023.
- GDAL/OGR contributors: GDAL/OGR Geospatial Data Abstraction software Library, <https://doi.org/10.5281/zenodo.5884351>, 2024.
- Gjorgjievska, T.: TOTAL RESIDENT POPULATION, HOUSEHOLDS AND DWELLINGS IN THE REPUBLIC OF NORTH MACEDONIA, Census 2021, Tech. rep., State Statistical Office, Republic of North Macedonia, ISBN 978-608-227-344-0, 2021.
- Jordahl, K., den Bossche, J. V., Fleischmann, M., Wasserman, J., McBride, J., Gerard, J., Tratner, J., Perry, M., Badaracco, A. G., Farmer,
245 C., Hjelle, G. A., Snow, A. D., Cochran, M., Gillies, S., Culbertson, L., Bartos, M., Eubank, N., maxalbert, Bilogur, A., Rey, S., Ren, C., Arribas-Bel, D., Wasser, L., Wolf, L. J., Journois, M., Wilson, J., Greenhall, A., Holdgraf, C., Filipe, and Leblanc, F.: geopandas/geopandas: v0.8.1, <https://doi.org/10.5281/zenodo.3946761>, 2020.
- Jovanović Popović, M. and Ignjatović, D.: National Typology of Residential Buildings in Serbia, Tech. rep., Faculty of Architecture, University of Belgrade, ISBN: 978-86-7924-102-3, 2013.
- 250 Kuenen, J., Dellaert, S., Visschedijk, A., Jalkanen, J.-P., Super, I., and Denier van der Gon, H.: Copernicus Atmosphere Monitoring Service regional emissions version 4.2 (CAM5-REG-v4.2), Tech. rep., Copernicus Atmosphere Monitoring Service, doi:10.24380/0vzb-a387, 2021.
- Kuenen, J., Dellaert, S., Visschedijk, A., Jalkanen, J.-P., Super, I., and Denier van der Gon, H.: CAM5-REG-v4: a state-of-the-art high-resolution European emission inventory for air quality modelling, Earth System Science Data, 14, 491–515, <https://doi.org/10.5194/essd-14-491-2022>, 2022.
- 255 Microsoft: Global ML Building Footprints, <https://github.com/microsoft/GlobalMLBuildingFootprints>.
- Muntean, M., Janssens-Maenhout, G., Song, S., Giang, A., Selin, N. E., Zhong, H., Zhao, Y., Olivier, J. G., Guizzardi, D., Crippa, M., Schaaf, E., and Dentener, F.: Evaluating EDGARv4.tox2 speciated mercury emissions ex-post scenarios and their impacts on modelled global and regional wet deposition patterns, Atmospheric Environment, 184, 56–68, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2018.04.017>,
260 2018.
- Novikova, A., Csoknyai, T., Miljanic, Z., Gligorić, B., Vušanović, I., and Szalay, Z.: Tipologija fonda stambenih zgrada Crne Gore - modeliranje njihove transformacije u budućnosti sa niskim nivoom ugljenika, Tech. rep., Regional Environmental Center, 2015a.
- Novikova, A., Szalay, Z., Simaku, G., Thimjo, T., Plaku, T., and Csoknyai, T.: The typology of the residential building stock in Albania and the modelling of its low-carbon transformation, Tech. rep., Regional Environmental Center for Central and Eastern Europe (REC), 2015b.



- 265 Novikova, A., Csoknyai, T., and Szalay, Z.: Low carbon scenarios for higher thermal comfort in the residential building sector of South Eastern Europe, *Energy Efficiency*, 11, 845–872, 2018.
- QGIS Development Team: QGIS Geographic Information System, <https://www.qgis.org>, 2021.
- Simovski, A.: ENERGY CONSUMPTION IN HOUSEHOLDS, 2019, Tech. rep., State Statistical Office, ISBN 978-608-227-337-2, 2019.
- UNEP: Air pollution is responsible for up to one in five premature deaths in 19 Western Balkan cities, <https://www.unep.org/news-and-stories/press-release/air-pollution-responsible-one-five-premature-deaths-19-western>.
- 270 World Bank: World Bank population total, <https://data.worldbank.org/indicator/SP.POP.TOTL>.