

1 **First comprehensive stable isotope dataset of diverse**
2 **water units in a permafrost-dominated catchment on the**
3 **Qinghai–Tibet Plateau**

4

5 Yuzhong Yang^{1,2}, Qingbai Wu^{1,2}, Xiaoyan Guo³, Lu Zhou¹, Helin Yao¹, Dandan⁴
6 Zhang, Zhongqiong Zhang^{1,2}, Ji Chen^{1,2}, Guojun Liu^{1,2}

7 ¹ State Key Laboratory of Frozen Soil Engineering, Northwest Institute of Eco-Environment and
8 Resources, Chinese Academy of Sciences, Lanzhou, China

9 ² Qinghai-Beiluhe Plateau Frozen Soil Engineering Safety National Observation and Research Station,
10 China

11 ³ Key Laboratory of Ecohydrology of Inland River Basin, Northwest Institute of Eco-Environment and
12 Resources, Chinese Academy of Sciences, Lanzhou, China

13 ⁴ College of Energy and Power Engineering, Lanzhou University of technology, Lanzhou, China

14

15 *Correspondence to:* Yuzhong Yang (yangyuzhong08@lzb.ac.cn)

16 **Definition or description of permafrost associated terms**

17 ***Thermokarst lake:*** A lake occupying a closed depression formed by settlement of the ground following
18 thawing of ice-rich permafrost or the melting of massive ice.

19 ***Ground ice:*** A general term referring to all types of ice contained in freezing and frozen ground

20 ***Pore ice:*** It termed interstitial or ‘cement’ ice, is the bonding material that holds soil grains together.

21 ***Segregated ice:*** It is formed by the migration of pore water to the ‘frozen fringe’ where it forms
22 discrete lenses or layers.

23 ***Excess ice:*** defined as volume of ice in the ground that exceeds the total pore volume that the ground
24 would have under natural unfrozen conditions.

25 ***Active layer:*** It is usually identified as a ground or rock above the permafrost table which undergoes
26 seasonal freezing in winter.

27 **Abstract**

28 Considered as the Asian water tower, the Qinghai–Tibet Plateau (QTP) processes substantial
29 permafrost, where its hydrological environments are spatially differed and can be easily disturbed by
30 changing permafrost and melting ground ice. Permafrost degradation compels melting permafrost to
31 become an important source of surface runoff, changes the storage of groundwater, and greatly influences
32 the hydrological processes in permafrost regions. However, the evidences linking permafrost degradation
33 and hydrological processes on the QTP are lacking, which increase the uncertainties of the evaluation
34 results of changing permafrost on the water resources. Stable isotopes offer valuable information on the
35 connections between changing permafrost (ground ice) and water components. It is therefore particularly
36 important to observe the changes in the stable isotopes of different waterbodies, which can vary over
37 hourly to annual timescales and truly capture the thawing signals and reflect the influence of permafrost
38 (ground ice) on the regional hydrological processes. The Beiluhe Basin (BLH) in the hinterland of QTP
39 were selected, which well integrates all the water components related to hydrological cycles, and is an
40 ideal site to study hydrological effect of permafrost change. This paper presents the temporal data of
41 stable isotopes ($\delta^{18}\text{O}$, δD , and d-excess) in different water bodies (precipitation, stream water,
42 thermokarst lake, and groundwater) in the BLH produced between 2017 and 2022. In special, the first
43 detailed stable isotope data of ground ice at 17 boreholes and 2 thaw slumps are presented. A detailed
44 description of the sampling processes, sample pretreating processes, and isotopic data quality control is
45 given. The data firstly described the full seasonal isotope amplitude in the precipitation, stream, and
46 thermokarst lakes, and delineated the depth isotopic variability in ground ice. Totally, 554 precipitation
47 samples, 2402 lakes/ponds samples, 675 stream water samples, 102 supra-permafrost water samples, and
48 19 sub-permafrost water samples were collected during six years' continuous sampling work.
49 Importantly, 359 ground ice samples at different depths from 17 boreholes and 2 profiles were collected.
50 This first data set provides a new basis for understanding the hydrological effects of permafrost
51 degradation on the QTP. It also provides supports on the cryospheric study on the Northern Hemisphere.
52 The dataset provided in this paper can be obtained at <https://doi.org/10.5281/zenodo.10684110> (Yang,
53 2024).

54

55 **1 Introduction**

56 Recognized as the main components of cryosphere, permafrost plays critical roles in climate change,
57 evolution of ecosystem, water cycle, and human activities (Brown et al., 1997). Throughout the past
58 several decades, the thermal stability of permafrost has suffered serious threats (Cheng et al., 2019;
59 Douglas et al., 2021; Biskaborn et al., 2019) caused by continuous global warming (IPCC, 2019). Latest
60 IPCC report indicates that up to 24-69% of permafrost will disappear by 2100 (IPCC, 2019). Warming
61 and thawing of permafrost and an overall reduction in the ice content have been predicted under future
62 climate change scenarios (IPCC, 2019). Dramatic permafrost degradation and ground ice melting has
63 changed the regional hydrological processes (Yang et al., 2011; Quinton and Baltzer, 2013; Rogger et
64 al., 2017), enhanced the hydraulic connections (Connon et al., 2014; Cheng and Jin, 2013; Zhang et al.,
65 2013), and compel ground ice to become an important source of surface runoff and lakes (Yang et al.,
66 2019; Zhang et al., 2005; Lawrence and Slater, 2005). Accordingly, clarifying the influence of degrading
67 permafrost on the ecohydrology and water resources is of great significance to the protection of eco-
68 environment and effective utilization of fresh water in permafrost regions in the world.

69 The Qinghai–Tibet Plateau (QTP) is known as the “Asia Water Tower”, which is considered as the
70 headwater regions of many large rivers in Asia (Immerzeel et al., 2010). As the world’s largest high-
71 altitude permafrost regions (Cheng et al., 2019), the QTP contains as many as 1.06×10^6 km² of permafrost
72 and 12700 km³ of ground ice (Cheng et al., 2019). Extensive development of permafrost and substantial
73 reserves of ground ice has exerted critical roles in climate change, ecosystem transition, water resource,
74 carbon budget, and infrastructure of QTP (Zhao et al., 2020; Liu et al., 2022a; 2022b). Accordingly, the
75 QTP has been becoming a hot region for scientists from different research fields (Wang et al., 2006;
76 Yang et al., 2019; Zhao et al., 2021). During recent decades, the QTP has been experiencing severe
77 warming over the past 50 years (Yao et al., 2013; Ran et al., 2022; Kuang and Jiao, 2016), which leads
78 to accelerated permafrost degradation (Wu and Zhang, 2010; Zhao et al., 2021), and thereafter greatly
79 affected the plateau water-eco environment-carbon cycle systems (Wang et al., 2023a; Yi et al., 2014;
80 Liu et al., 2022a).

81 So far, due to the harsh climate conditions, inconvenient transportations, and high experimental
82 costs of site-specific field data, there has been a lack of comprehensive research on different water bodies

83 in permafrost regions over a long time on the QTP, making it challenging to study the water cycle and
84 hydrological processes associated with changing permafrost. In addition, traditional method (e.g.,
85 modelling, GRACE satellite technique) is thus difficult to delineate the processes of ice-water transition
86 truly and comprehensively, greatly increasing the uncertainties of evaluation results about the impacts of
87 permafrost degradation on the hydrological processes (Guo et al., 2017). Hydrogen and oxygen stable
88 isotopes ($\delta^{18}\text{O}$, δD) are widely existing natural tracers, which are considered to be ideal tools to identify
89 temporal-spatial patterns of precipitation-river-lake-groundwater systems (Knapp et al., 2019; Narancic
90 et al., 2017; Vystavna et al., 2021) and therefore to delineate hydrological connectivity under degrading
91 permafrost (Wang et al., 2022; Streletskiy et al., 2015; Yang et al., 2019). Furthermore, the stable
92 isotopes can well document the signals of ice-water phase transition and freezing history, making them
93 provide convenient means for investigating of ground ice evolution (Michel, 2011; Lacelle et al., 2013;
94 Porter et al., 2019) in permafrost.

95 Accordingly, continued observations of the stable isotope data, required to understand the changes
96 of hydrological processes and water vapor cycles linked with permafrost degradation and ground ice melt,
97 are therefore of great importance. However, the acquisition of long time series stable isotopic data in
98 permafrost-dominated catchment on the QTP is challenging, especially the stable isotope records of
99 thermokarst lakes/ponds and ground ice on the QTP are extremely scarce. It greatly limits the deep
100 understanding of the hydrological processes under thawing permafrost.


101 In this paper, we provide information on the study site and full documentation of the water
102 components in a typical permafrost watershed (Beiluhe Basin, BLH) on the QTP. The data sets presented
103 here, including the stable isotopes of daily precipitation, monthly isotope data of surface waters (stream
104 and thermokarst lakes/ponds) and groundwater, and ground ice within 20 m in depth, will be of great
105 value for tracking water vapor cycles, for capturing the signals of permafrost thawing and delineating the
106 hydrological routines of permafrost meltwater, and in continuing baseline studies for future permafrost
107 degradation trend analysis and water resources evaluations on the QTP. Special emphasis is given to the
108 critical role of BLH for research in the hinterland of QTP to diagnose the effect of thawing permafrost.

109 2 Study area

110 A typical permafrost catchment, namely the Beiluhe Basin (BLH; Fig. 1), was selected to
111 comprehensively observe the hydrological processes under changing permafrost. The BLH is situated in
112 the interior of the QTP, with elevations of 4,500 to 4,600 m.s.l. It is considered as a core region of the
113 Hoh Xil Nature Reserve region and provides the best habitats for wild animals on the QTP. The BLH is
114 also identified as one of the most fragile and sensitive ecosystems in the world due to the diversities in
115 the ecosystems, which including swamp meadow, alpine meadow, degrading alpine meadow, alpine
116 steppe, desert alpine grassland, sparse grassland (Yin et al., 2017). According to the meteorological
117 station of BLH, between 2017 and 2022, the annual mean air temperature ranged between -3.57 °C (2019)
118 and -2.43 °C (2022), the annual precipitation ranged between 394 mm (2020) and 556 mm (2018), the
119 duration of negative air temperature exceeds 200 d.

120 The BLH is closely connected with the Source Area of Yangtze River (i.e., the Tuotuohe River),
121 and is characterized by a complex hydrological system of streams (Yang et al., 2017), thermokarst lakes
122 (Yang et al., 2016; Niu et al., 2011), groundwater (springs), as well as abundant ground ice (Yang et al.,
123 2013; 2016). Thermokarst lakes are widely distributed in the basin, with a total lake-number of more
124 than 1200 (Luo et al., 2015) which are showing gradual increase trend. In addition, controlled by the
125 piedmont faults of Gushan Mountain (Fig. 1) in the BLH, the natural springs are extensively exposed on
126 the ground, which are the main sources of small streams. The connectivity of lakes, streams, groundwater,
127 as well as melting water from permafrost exerted important roles on how ecological and hydrological
128 systems are propagated in this basin.

129 The BLH is located in the zone of continuous ice-rich permafrost in the Changtang Basin. The
130 permafrost thickness is approximately 20–80 m thick. Mean annual ground temperature (MAGT) at 15
131 m depth ranges from -1.8 to -0.5°C and the active layer thickness is 1.6–3.4 m (Wu et al., 2015). Ground
132 ice is abundant in this region, and as high as 70% of this area has a volumetric ice content (VIC) higher
133 than 30% (Luo et al., 2015). Most of the ground ice in the BLH is identified as excess ice (Niu et al.,
134 2002), which could melt out to recharge supra-permafrost water (springs) or even surface water (Yang
135 et al., 2016). Accordingly, the BLH is a natural laboratory to conduct field hydrological observations,
136 the observation data can facilitate the developments of human infrastructure and ecological restoration
137 of QTP.



138
 139 **Figure 1: (a) Location of the Beiluhe Basin on the QTP, (b) Distribution of our study area in the Beiluhe Basin,**
 140 **and (c) the specific sampling sites of different water components in the BLH. Supra-PW denotes the Supra-**
 141 **permafrost water.**

142 **3 General design of the monitoring network**

143 From 2017 to 2022, we set up sampling sites of precipitation, stream, thermokarst lake/pond,
144 groundwater (including supra-permafrost water and sub-permafrost water), and ground ice in the BLH
145 basin (Fig. 1). The precipitation stable isotope sampling site was setup at the BLH frozen soil station
146 (Fig. 1). A rain gauge was installed to collect daily rain, and a steel plate was put on the roof to obtain
147 enough snow samples. In addition, we selected a typical small stream (defined as Gushan Mountain
148 Stream, GMS) in the BLH Basin, which originates from four natural springs in foothill of the Gushan
149 Mountain (Fig. 2; Fig. S1). This stream is 4.8 km in length. The vegetation along this stream is mainly
150 composed of deserted steppe. A total of 25 fixed points along the stream were selected to collect water
151 samples during the ice-free seasons between June and October. Furthermore, a typical thermokarst lake
152 belt located in the southwestern of the BLH station on the QTP were selected to observe lake water
153 balance (Fig. 1). For the groundwater observation, we selected two areas with substantial natural opening
154 springs occurring, i.e., springs along the both sides of the observation stream (named as GSHQ) and
155 spring in the source area of this stream (named as GSYTQ) (Fig. 1; 2). Given the intermittent occurrence
156 of these springs among different years and their unstable isotopic signals, we identified them as supra-
157 permafrost water. In addition, a perennial spring (CSQ; Fig. 1) for domestic water supply behind the
158 BLH station (Fig. 1), with its aquifer depth (reaching 92 m) being deeper than the permafrost thickness
159 (~50m) in the BLH, is selected to conduct continuous sampling work. In regards to the small fluctuations
160 in water level all the year and little interannual differences in stable isotopes of spring, we identified it
161 as the observation site of sub-permafrost water. In order to detect the permafrost changes and clarify the
162 characteristics of ground ice conditions, 17 boreholes (20 m in depth) were drilled in the BLH basin (Fig.
163 1). All visible ice samples were collected in the field.

164 Meanwhile, an auto meteorological station is set up in the center of the BLH since 2005. Air
165 temperature is measured in a solar radiation shield at 2.0 m above the ground surface. The precipitation
166 amount from nearby meteorological station was measured using a T200B rain/snow gauge (Geonor,
167 Norway), and data were recorded every 30 min. The meteorological data have high quality and continuity
168 with very limited missing data due to regular maintenance by Beiluhe Frozen soil station.

169

170

171

Table 1 Location information on the sampling sites in the Beiluhe Basin

Sampling sites	Precipitation	Stream	Thermokarst lakes/ponds	Springs	Ground ice
Latitude/°	N 34.83	N 34.82~34.84	N 34.82~34.83	N 34.83~34.84	N 34.82~34.83
Longitude/°	E 92.94	E 92.92~92.93	E 92.89~92.93	E 92.92~92.93	E 92.93~92.89
Altitude/m	4628	4668~4697	4704~4752	4752~4771	4629~4691

172

4. Sample collection and processing

173

4.1 Sampling and preservation

174

4.1.1 Precipitation sampling work

175

According to the International Atomic Energy Agency/Global Network of Isotopes in Precipitation (IAEA/GNIP) precipitation sampling guide, a precipitation collector was manually constructed in an open area near the BLH meteorological station. To avoid the contamination of water vapor from evaporation of shallow soil and surface water, and the mixing of windblown snow, this collector was installed 2 m above the ground. We define one complete precipitation day beginning at 20:00 on one day, and ending at 20:00 in the next day, then the one sample was collected. All the rainfall samples were immediately collected after the end of precipitation to minimize the effects of evaporation. Hail and snow samples were filled in pre-cleaned plastic bags, the air was pressed out from the bags and sealed to avoid the water vapor exchange, and all samples in sealed bags were melted at room temperature (25 °C). In order to clarify the changes comprehensively and accurately in the precipitation isotopes in the BLH Basin, we tried to collect all samples during every precipitation event, including light rain and short-time events (usually with precipitation amount of less than 5 mm). Accordingly, a wide mouth stainless steel plate (400 mm×600 mm) was used to collect enough samples of light rain and short-time rain/snow events for analysis.

189


Regarding preserving samples, 100 ml high-density polyethylene (HDPE) bottles were used. Before the sampling, the bottles were washed three times with rain water and then rapidly filled. Totally, 554 precipitation samples were collected, including 224 rain samples, 203 snow samples, 85 hail samples, and 42 sleet samples.

192

193 4.1.2 Stream, thermokarst lakes/ponds, and groundwater sampling

194 Samples of thermokarst lakes/ponds (Fig. 2) were collected by hand using a self-made water sample
195 collector at monthly intervals during ice-free seasons (between May and October) from 2017 to 2022 in
196 the BLH Basin (Fig. 1). During the observation periods, the occurrence numbers of thermokarst lakes
197 dynamically changed among different sampling years (Table 2) due to the interannual variations in the
198 precipitation, active layer thickness, supra-permafrost water, as well as near-surface ground ice. Some of
199 the sampled lakes disappeared in the next sampling year and additional new lakes emerged. Accordingly,
200 we obtained sufficient lake water samples to constrain the seasonal changes in the lake water hydrology
201 and try to clarify the influence of permafrost and climate on the water balance of thermokarst lakes in
202 this region. Influenced by the Covid-19 and lockdown policies between August, 2022 to December, 2022
203 in China, only two months' sampling work (June and July) was conducted in 2022. Lake water samples
204 were taken at the center of lakes from 20–40 cm below water surface. The running water samples of
205 stream water samples were collected at each fixed point 20-30 cm beneath the water surface. In addition,
206 the supra-permafrost water and sub-permafrost water were randomly collected using a man-made water
207 ladle at the location where the springs gushing out during each field work. The water ladle was washed
208 using the spring water before sampling.

209 Totally, as many as 2402 thermokarst lakes/ponds samples, 675 stream water samples (Table 2),
210 102 supra-permafrost water samples, and 19 sub-permafrost water samples were collected during six
211 years' continuous sampling work.



212

213

214

215

Figure 2: (a) General conditions of Gushan Mountain Stream (GMS) and distribution of springs; (b) Typical feature of one spring gushing out from sand sediment; (c) Overview picture of GMS; and (d) Sampling thermokarst lakes in the BLH.

Table 2 Sampling descriptions of surface water in the BLH

Sampling Information	Number of samples	
	Thermokarst lake/pond	Stream
Jun-17	23	25
Jul-17	76	25
Aug-17	74	25
Sep-17	99	25
Oct-17	72	25
May-18	74	N.A
Jun-18	14	25
Jul-18	45	25
Aug-18	110	25
Sep-18	93	25
Oct-18	106	25
May-19	80	N.A
Jun-19	115	25
Jul-19	134	25
Aug-19	87	25
Sep-19	85	25
Oct-19	110	25
Jun-20	86	25
Jul-20	124	25
Aug-20	116	25
Sep-20	93	25
May-21	73	25
Jun-21	70	25
Jul-21	100	25
Aug -21	100	25
Sep-21	94	25
Jun-22	75	25
Jul-22	74	25
Total sample size	2402	675

218 **4.1.3 Ground ice sampling**

219 To clarify the characteristics of ground ice and its role on the local hydrological cycles and regional
220 eco-environment, we have designed 17 boreholes (~20 m in depth) in the BLH basin (Fig. 1). A total of
221 12 boreholes were drilled near the Qinghai-Tibet Highway (QTH) in 2014, and 5 boreholes were
222 distributed in the center of BLH basin, which were drilled between 2011 and 2021. In addition, 2 thaw
223 slumps were dug (Fig. 1). Frozen soil cores were extracted from different depths using a mechanical
224 drilling rig with a drilling diameter of 157 mm (Fig. 3). All visible ground ice samples were collected
225 immediately after the core barrel was pulled out. During sampling work, the disposable PE gloves were
226 used, and the exterior of each sample was removed to avoid contamination from mud and the surplus
227 water in the borehole. Totally, 355 ground ice samples were collected from 17 boreholes and 4 samples
228 were obtained from 2 profiles (Fig. 3; Table 3).



229

230 **Figure 3: Field permafrost drilling work and various types of ground ice obtained during drilling.**

231

Table 3 Borehole drilling and ground ice sampling information in the BLH

Borehole name	Drilling time	Depth range of ice sampling/m	Ground ice types	Sample number
BLH-L-1	Aug-2014	4.8-14.9	Pore ice/segregated ice/excess ice	10
BLH-L-2	Aug-2014	2.7-14.3	Pore ice/segregated ice/excess ice	28
BLH-L-3	Aug-2014	2.9-14.8	Pore ice/segregated ice/excess ice	20
BLH-L-4	Aug-2014	2.55-14.2	Pore ice/segregated ice/excess ice	34
BLH-L-5	Aug-2014	2.3-14.0	Pore ice/segregated ice/excess ice	15
BLH-L-6	Aug-2014	2.6-14.3	Pore ice/segregated ice/excess ice	11
BLH-R-1	Aug-2014	3.0-12.9	Pore ice/segregated ice/excess ice	10
BLH-R-2	Aug-2014	1.9-14.9	Pore ice/segregated ice/excess ice	20
BLH-R-3	Aug-2014	1.25-8.1	Pore ice/segregated ice/excess ice	17
BLH-R-4	Aug-2014	1.8-11.9	Pore ice/segregated ice/excess ice	32
BLH-R-5	Aug-2014	1.7-13.8	Pore ice/segregated ice/excess ice	36
BLH-R-6	Aug-2014	2.1-14.6	Pore ice/segregated ice/excess ice	22
DZK	Aug-2012	0.0-20.55	Pore ice/segregated ice/excess ice	27
ZK-1	Aug-2011	12.4-17.4	Pore ice/segregated ice/ Pure ice layer	28
ZK-2	Aug-2011	3.0-7.2	Pore ice/segregated ice/excess ice	15
ZK-3	Aug-2011	2.6-12.8	Pore ice/segregated ice/excess ice	13
ZK-4	Aug-2011	2.2-5.5	Pore ice/segregated ice/excess ice	17
Z	Oct-2021	2.0-3.0	Thaw slump ice	2
FBX	Oct-2021	2.0-3.0	Thaw slump ice	2

234 4.1.4 Sample storage

235 **Liquid water storage:** All the samples were transferred to 100 ml high-density polyethylene
236 (HDPE) bottles. The sample bottles were filled up without bubbles and sealed with parafilm. The
237 collection date sample types (precipitation, lake water, stream water, groundwater) were labelled. For
238 the precipitation samples, the precipitation types (rain, snow, hail) were recorded. All the samples were
239 stored at 4°C and shipped to the State Key Laboratory of Frozen Soil Engineering (SKLFSE) in
240 Northwest Institute of Eco-Environment and Resources, Chinese Academy of Sciences (CAS), China.

241 **Ground ice storage:** All the treated raw frozen soil samples were immediately preserved in HDPE
242 bottles. The massive ice and pure ice layers were sealed in the pre-cleaned plastic bags. The depths and
243 drilling site information were recorded. All the frozen soil and ground ice samples were kept frozen at
244 -4°C in the field to avoid sublimation of the ice and evaporation of the water in the soil.

245 4.2 Sample pretreatment and stable isotope analysis

246 Before analyzing, each liquid sample was pretreated to remove the impurities through 0.22- μ m
247 disposable membrane filters. The frozen soil samples and pure ground ice samples were allowed to
248 completely melt at 4 °C in sealed plastic bags. The supernatant water from thawed soil and meltwater
249 from ground ice were also filtered through a 0.22- μ m membrane. The processed liquid water samples
250 were filled in 2 ml analytical vial and were stored in a cold room (4 °C) in the dark for the stable isotopes
251 ($\delta^{18}\text{O}$ and δD) analysis within 1 week.

252 The $\delta^{18}\text{O}$ and δD ratios were measured at SKLFSE, using an Isotopic Liquid Water and Water Vapor
253 Analyzer (Picarro L2130-i, U.S.) based on the wavelength-scanned cavity ring down spectroscopy
254 technique. The guaranteed instrument precision was 0.025 ‰ for the $\delta^{18}\text{O}$ value measurements and 0.1 ‰
255 for the δD value measurements. The isotopic values were reported using notation representing the per
256 mille (‰) relative difference with respect to the IAEA Vienna Standard Ocean Water (VSMOW)
257 standard following Eq. (1):

$$258 \quad \delta = (R_{\text{sa}}/R_{\text{st}} - 1) \times 1000 \text{ ‰}$$

259 R_{sa} is the respective isotope ratio measurement for the samples and R_{st} is the defined isotope ratio of a
260 standard sample.

261 **4.3 Quality control of data**

262 **4.3.1 Sampling errors**

263 The precipitation samples were transferred to HDPE bottles immediately. If multiple rain/snow
264 events occurred during one sampling day, the water sample from every single precipitation event was
265 collected separately. At the end of one complete sampling day, all the samples collected from single
266 event were mixed. If the precipitation types changed during one sampling day, different samples were
267 collected separately. The final complete samples were kept cool at 4 °C. All we have done is to avoid the
268 influence of evaporation on enrichment of D and ¹⁸O and ensure the originality of samples.

269 During the sampling work of thermokarst lakes/ponds and streams, we do our best to control the
270 sampling time at the same period during every month (controlling the sampling time within one week,
271 i.e., between 17th and 22th in every month) to make sure that all the samples can represent the average
272 level of the whole month. The sampling HDPE bottles were precleaned three times using the raw water.
273 Lake water was taken at the center of lakes from 20–40 cm beneath water surface. The running water
274 samples of stream were collected at each fixed point 20-30 cm beneath the water surface.

275 **4.3.2 Analytic errors**

276 Before we started to analyze the samples, we firstly prepared 14 distilled water samples with the
277 same stable isotopes to test the stability of our analyzer. The precisions of the $\delta^{18}\text{O}$ and δD values were
278 calculated by calculating the 1-sigma standard deviation of groups of 12 injections and then calculating
279 the average of these standard deviations. In order to ensure the data quality, the “high precision” mode
280 was employed during analysis. Under this mode, the analyzing time for each injection is about 8.75
281 minutes. The drift of the analyzer was determined by taking the mean of these same 12 groups of
282 measurements and calculating the difference between the maximum and minimum means. All these
283 measured precision and drift values were less than those of the guaranteed precision (0.025‰ and 0.1‰
284 for $\delta^{18}\text{O}$ and δD) and drift values (0.2‰ and 0.8‰ for $\delta^{18}\text{O}$ and δD), indicating that the analyzer achieve
285 both a good repeatability and a good reproducibility. If the measured precision and drift values were not
286 passed the guaranteed values, the comprehensive inspection of the analyzer was conducted, i.e., the
287 instrument analyzing system, the vaporizer module, as well as the quality of dry nitrogen. After
288 completing all checking processes, we repeated the analysis of 14 distilled/tap water samples and
289 calculated the drift values until they passed the guaranteed values. The results were normalized to the V-

290 SMOW-SLAP scale by analyzing internal standards before and after each set of ten samples. Five
291 laboratory standards (provided by LICA United Technology Limited, Beijing, China) with given isotopic
292 values were inserted before 10 samples, which were used for instrument calibration: with $\delta^{18}\text{O}$ values of
293 -21.28‰ , -16.71‰ , -11.04‰ , -7.81‰ , and -2.99‰ , and δD values of -165.7‰ , -123.8‰ , -79.6‰ ,
294 -49.2‰ , -9.9‰ . The best-fit linear relationship between the five known calibration values and the
295 analyzer's reported values was determined. The slope and intercept of the best-fit line through these
296 points are used to calibrate the results of our samples.

297 To avoid memory effects, the first three results of measurements were discarded and arithmetic
298 mean values were calculated from the last three injections. During the analyzing process, the real-time
299 data of water concentration of all injections were controlled within a range between 19000 ppm and
300 20000 ppm and with a standard deviation of less than 200 ppm. Once the water concentration values
301 appear to decrease, the work was stopped and the syringe was detached to wash using the deionized
302 water. All measurements were post-processed with the Picarro ChemCorrect™ software to monitor the
303 organic contamination and correct the data.

304 **5 General characteristics of stable isotopes in different water components**

305 **5.1 Variations in the stable isotopes of different water components**

306 **5.1.1 Precipitation**

307 The stable isotopes of precipitation exhibit a remarkable seasonal variability during six years'
308 observations (Fig. 4). The $\delta^{18}\text{O}$ and δD of the local precipitation in the BLH Basin ranged from -30.4‰
309 to 6.2‰ and from -238.0‰ to 65.4‰ , respectively. The d-excess ranged between -37.5‰ and 44.5‰ .
310 The amount-weighted average values of annual precipitation are -10.9‰ , -72.1‰ , and 15.4‰ for $\delta^{18}\text{O}$,
311 δD , and d-excess, respectively. As shown, the $\delta^{18}\text{O}$ and δD display distinct seasonal patterns with high
312 values in summer and low values in winter (Fig. 2; Fig. S2), it is due to the changes in moisture sources
313 and the influence of local climate conditions (Guo et al., 2022; Tian et al., 2005; Guan et al., 2013;
314 Bershaw et al., 2012).


315 5.1.2 Surface water bodies

316 For comparison, the $\delta^{18}\text{O}$ and δD of thermokarst lakes/ponds are more positive than those of
317 precipitation due to strong evaporation and resultant enrichments of heavier isotopes in lake water (Yang
318 et al., 2016; Narancic et al., 2017; Ala-aho et al., 2018). The $\delta^{18}\text{O}$ ranged from -14.4‰ to 5.7‰ (mean:
319 -6.0‰), the δD is between -104.1‰ and 22.6‰ (mean: -48.0‰), and the d-excess is ranged from -35.8‰
320 to 21.8‰ (mean: -0.1‰), respectively. Similarly, the isotopic patterns of thermokarst lakes/ponds
321 exhibited strong seasonal variations (Fig. 4; Fig. S3), which is due to the changes in source waters (i.e.,
322 precipitation, meltwater of thawing permafrost/ground ice, groundwater) and alternations of evaporation
323 degrees due to air temperature fluctuations (Narancic et al., 2017; Yang et al., 2021; Aichner et al., 2022;
324 Zhu et al., 2022). Generally, the heavy isotope contents of lakes/ponds are lower in August and
325 September (Fig. 4; Fig. S3), which is attributed to the recharges of monsoonal precipitation and water
326 with more negative isotopes fed by melting ground ice (Gibson et al., 2015; Yang et al., 2021). In
327 comparison, majority of isotope values of lakes/ponds are positive in May, June, July, and October (Fig.
328 4; Fig. S3) due to evaporation and recharge of isotopic-enriched precipitation water.

329 For the streams, the isotope values varied from -13.7‰ to -7.2‰ ($\delta^{18}\text{O}$, mean: -11.1‰) and from -
330 83.8‰ to -53.3‰ (δD , mean: -73.6‰), and the d-excess is ranged from -0.6‰ to 25.6‰ (mean: 15.0‰),
331 respectively. The mean values are equivalent to the average values of annual precipitation in the BLH.
332 Compared with thermokarst lakes/ponds, the $\delta^{18}\text{O}$ values of stream water exhibited relatively stable
333 patterns (Fig. 4) due to short residence time (Yang et al., 2021; Wang et al., 2023b; Song et al., 2017),
334 which indicates weak evaporation. However, the stream isotopes also represented seasonal variations
335 during six year's observation (Fig. 4; Fig. S4), lower values were prevailing in August and September.
336 The temporal changes of stream isotopes are mainly influenced by the seasonal variability of evaporation
337 (Yang et al., 2017) and differences in the source water, i.e., alternative replenishment of precipitation,
338 melting ground ice, and groundwater (Streletskiy et al., 2015; Yang et al., 2019; Ala-aho et al., 2018).

339 The two kinds of supra-permafrost water (i.e., GSHQ and GSYTQ) exhibited similar seasonal trend
340 (Fig. 4). For comparison, the GSHQ displayed relatively more positive isotopic peaks during whole
341 sampling periods (Fig. 4), with $\delta^{18}\text{O}$ ranging from -13.3‰ to -5.8‰ (mean: -11.2‰), the δD is ranging
342 between -86.7‰ and -39.0‰ (mean: -74.2‰), and the d-excess varying from 6.5 to 22.4‰ (mean:
343 15.1‰), respectively. The isotopes of GSYTQ varied from -13.5‰ to -8.4‰ (mean: -11.4‰), the δD is

344 ranging between -83.2‰ and -50.6‰ (mean: -73.8‰), and the d-excess is varying from 4.6 to 25.1‰
 345 (mean: 16.9‰). The isotopic peaks of the two types of springs lagged behind those of precipitation (Fig.
 346 4), indicating replenishments of precipitation via infiltration. By contrast, the stable isotopes of sub-
 347 permafrost water are more negative than those of supra-permafrost water, ranging between -12.7‰ and
 348 -11.1‰ (mean: -11.8‰) for $\delta^{18}\text{O}$, from -83.7‰ to -77.7‰ (mean: -80.7‰) for δD , and from 10.9‰ to
 349 17.7‰ for d-excess (mean: 13.5‰). In addition, they kept nearly stable over long time series (Fig. 4),
 350 suggesting unchanged sources of isotopically light water (e.g., monsoonal precipitation, meltwater from
 351 thawing permafrost, etc) and insignificant influence by precipitation.



352
 353 **Figure 4: Temporal variations in the $\delta^{18}\text{O}$ of different water components in the BLH. The numbers denote**
 354 **the observation months of thermokarst lakes/ponds. The dots with different colours represent event values,**
 355 **while the red dotted line denote the monthly average values. GSHQ and GSYTQ denotes the springs along**
 356 **the both sides of the observation stream and spring in the source area of this stream, respectively.**

357 5.1.3 Ground ice


358 The distributions of stable isotope dots of all cores are scattered along depths (Fig. 5). Generally,
 359 the $\delta^{18}\text{O}$ ranging from -15.0‰ to -8.3‰ (mean: -12.2‰), from -113.7‰ to -66.4‰ (mean: -94.4‰)
 360 for δD , and between -13.4‰ to 15.5‰ (mean: 3.1‰) for d-excess, respectively. Comparing with the
 361 precipitation, majorities of the $\delta^{18}\text{O}$ values of ground ice are lower than the precipitation, indicating

362 multi-sources of initial water during ice formation under variable climatic conditions and complex
363 geological contexts on the QTP (Michel, 2011; Yang et al., 2017; 2023; Murton, 2013).

364 Specifically, the stable isotopes of ground ice varied between different boreholes (Fig. 5; Table 4).
365 It is attributed to the influences of initial source water and complex ice formation mechanism. In addition,
366 the isotopic patterns along depths showed marked differences between boreholes (Fig. 5), suggesting
367 influence of lithology on the water migration and freezing fractionation of stable isotopes (Yang et al.,
368 2020; Lacelle, 2014; Fisher et al., 2021). Remarkably, the thaw slump ice was isotopically lighter than
369 those of drilling ground ice (Fig. 5; Table 4), it is due to the considerable differences in the initial source
370 water and freezing processes. The thaw slump ice is considered to replenished by winter snowmelt water
371 via cracks and freezing quickly (Fritz et al., 2011; Porter et al., 2020). However, the pore ice with lower
372 isotopic values in these boreholes is suffered isotope fractionation due to freeze-thaw under climate
373 transitions (Wetterich et al., 2014; Yang et al., 2023).

Table 4 General stable isotope composition of ground ice in the Beiluhe Basin

Borehole name	Stable isotopes of ground ice								
	$\delta^{18}\text{O}/\text{‰}$			$\delta\text{D}/\text{‰}$			d-excess/ ‰		
	Max	Min	Mean	Max	Min	Mean	Max	Min	Mean
BLH-L-1	-11.7	-14.0	-13.0	-91.0	-102.5	-96.8	9.9	2.8	7.0
BLH-L-2	-10.5	-14.5	-13.2	-86.2	-110.2	-101.9	10.6	-2.5	3.7
BLH-L-3	-11.39	-15.0	-12.8	-92.7	-113.7	-100.6	11.4	-7.1	1.7
BLH-L-4	-9.1	-14.6	-12.3	-80.4	-108.2	-95.5	13.9	-13.3	3.0
BLH-L-5	-10.2	-14.1	-12.7	-89.0	-108.6	-100.2	8.1	-7.3	1.7
BLH-L-6	-10.1	-13.3	-11.6	-86.9	-105.4	-96.1	4.4	-10.3	-2.9
BLH-R-1	-9.3	-13.1	-11.9	-80.3	-100.6	-90.8	9.3	-6.1	4.4
BLH-R-2	-10.0	-14.3	-12.5	-80.9	-102.8	-93.5	15.5	-0.9	6.7
BLH-R-3	-11.5	-14.0	-12.8	-90.8	-103.0	-97.5	11.5	-1.6	4.9
BLH-R-4	-11.0	-14.0	-12.7	-94.6	-102.3	-98.5	11.5	-8.5	3.0
BLH-R-5	-9.6	-13.9	-12.4	-84.8	-103.3	-96.3	11.0	-7.8	3.0
BLH-R-6	-8.4	-14.2	-11.8	-75.2	-108.3	-93.1	9.2	-9.2	1.6
DZK	-8.3	-12.3	-10.8	-66.4	-91.8	-85.1	8.0	-2.6	1.3
ZK-1	-10.3	-12.2	-11.8	-83.6	-89.8	-88.1	8.4	-0.8	6.3
ZK-2	-9.7	-13.1	-11.8	-78.8	-102.1	-93.6	7.4	-13.4	1.2
ZK-3	-9.2	-13.2	-11.4	-74.3	-103.6	-90.9	11.8	-13.2	0.7
ZK-4	-9.1	-12.9	-10.9	-69.9	-96.4	-84.4	9.7	-9.1	3.0
TSI	-14.3	-14.7	-14.5	-93.2	-96.3	-94.9	21.2	20.2	20.9



375

376 **Figure 5: Variations in the stable isotopes of ground ice along depths in the BLH. The blue line denotes the**
 377 **amount-weighted average $\delta^{18}\text{O}$ value of precipitation in the BLH.**

378 5.2 $\delta^{18}\text{O}$ - δD relations and hydrological connections


379 5.2.1 $\delta^{18}\text{O}$ - δD relationships of different water components

380 The local meteoric water line (LMWL), determined by three different methods, i.e., ordinary least
 381 square regression using the daily isotopic data, the arithmetic mean isotopic values, and the amount-
 382 weighted multi-monthly mean isotopic values during six years (2017-2022). They are expressed as:
 383 $\text{LMWL}_{\text{event}}: \delta\text{D}=7.97\delta^{18}\text{O}+15.26$ ($r^2=0.96$), $\text{LMWL}_{\text{montlyly}}: \delta\text{D}=8.06\delta^{18}\text{O}+12.58$ ($r^2=0.93$), $\text{LMWL}_{\text{PWA}}:$
 384 $\delta\text{D}=7.78\delta^{18}\text{O}+8.78$ ($r^2=0.92$). The slope is nearly identical to that of the global meteoric water line
 385 (GMWL; Craig, 1961). However, the intercepts are quietly different (Fig. 6) due to the influences of
 386 precipitation amounts and the exceptional meteorological conditions (Barešić et al., 2006; Hughes and
 387 Crawford, 2012; Kern et al., 2016).

388 The $\delta^{18}\text{O}$ - δD diagrams of lakes, streams, and groundwater were built using the monthly stable
 389 isotopic values, and defined as local evaporation line (LELs). The LELs observed during six years are
 390 calculated as: $\delta\text{D}=5.88\delta^{18}\text{O}-12.80$ ($r^2=0.95$), $\delta\text{D}=4.89\delta^{18}\text{O}-19.41$ ($r^2=0.83$), $\delta\text{D}=5.69\delta^{18}\text{O}-10.50$ ($r^2=0.85$)

391 (supra-permafrost water), and $\delta D=3.54\delta^{18}O-39.06$ ($r^2=0.92$) (sub-permafrost water), respectively. The
 392 slopes of the three LELs are all lower than those of LMWL (Fig. 6), and ranging between 4 and 6,
 393 indicating strong evaporation (Cui et al., 2017; Yang et al., 2019) due to lower relative humidity (Clark
 394 and Fritz, 1997). Interestingly, the correlation coefficients of streams and supra-permafrost water are
 395 much lower (less than 0.9) and the slopes are smaller than those of precipitation and lakes/ponds (Fig.
 396 6), which may be affected by the transitions of source water during warm seasons and the evaporative
 397 concentration of isotopes.

398 The $\delta^{18}O$ - δD relationship for ground ice was established using the stable isotopic values of the ice
 399 samples, and the correlation is defined as the freezing line (FL; Souchez et al., 2000). In this study, the
 400 freezing line of the ground ice at 16 borehole sites were calculated as: $\delta D=5.36\delta^{18}O-29.15$ ($r^2=0.73$),
 401 which is significantly different from the LMWL (Fig. 6). The difference reflects the freezing
 402 characteristics of liquid water under different conditions (Lacelle, 2011). Our freezing slope in between
 403 6.2 and 7.3 were usually obtained during equilibrium freezing Rayleigh-type fractionation (Lacelle,
 404 2011). The lower correlation coefficient (Fig. 6) suggests variable freezing rates (Souchez et al., 2000),
 405 kinetic isotopic fractionation during ice formation (Souchez et al., 2000), as well as the influence of the
 406 initial source water of the ground ice at different sites (Lacelle, 2011; Yang et al., 2017).




407
 408 **Figure 6: The relation between δD and $\delta^{18}O$ of different water components in the BLH. The Wt. avg. SP, Wt.**
 409 **avg. WP, Wt. avg. AP, LEL, and FL denotes the weighted average value of summer precipitation, weighted**
 410 **average value of winter precipitation, weighted average value of annual precipitation, local evaporation line**
 411 **of surface water components, and freezing line of ground ice, respectively.**

412 5.2.2 Hydrological connections between various water components

413 Majority of the stable isotopes of stream lie on the LMWL (Fig. 6) and embrace in the range of
414 supra-permafrost water (Fig. 7), in addition, the mean value is close to the amount-weighted average
415 value of annual/summer precipitation, indicating the direct recharge of precipitation and supra-
416 permafrost waters. However, some of the isotopic points do not lie on the LMWL, exhibit a clear
417 evaporative effect. The supra-permafrost water and sub-permafrost water display concentrated isotopic
418 patterns comparing with precipitation, reflecting relatively stable recharge sources. In addition, the
419 scattered isotopic dots of supra-permafrost water rather than sub-permafrost water indicated changeable
420 sources and climate conditions. For comparison, some of the isotope points of the supra-permafrost water
421 are overlapping with those of precipitation and stream water, suggesting important replenishment of
422 precipitation and stream. However, the isotopic cluster of sub-permafrost water is significantly deviated
423 from the LMWL and all the isotope values are lower than the annual average value of modern
424 precipitation, suggesting the recharge signal of past water with negative $\delta^{18}\text{O}$ (δD) values under cold
425 climate conditions. The LEL of thermokarst lakes/ponds significantly deviated from LMWL (Fig. 6; 7),
426 partial of the isotopic dots overlapped with precipitation, groundwater, and ground ice, indicating the
427 hydrological connections between them (Yang et al., 2016; 2017).

428 The cluster of ground ice is partly overlapped with precipitation, groundwater, lakes, and stream
429 (Fig. 7). It is indicative of mutual replenishment relations between them. Some of the isotope dots are
430 more positive than the summer precipitation, implying the recharge from evaporative active layer water.
431 A clear freezing slope is shown, indicating typical freezing of liquid water (Jouzel and Souchez, 1982;
432 Souchez and Jouzel, 1984; Lacelle et al., 2011; Perşoiu and Pazdur., 2011). However, the d-excess values
433 of ground ice are lower than those of river water and the amount-weighted average value of
434 annual/summer precipitation (Fig. 7), suggesting the important recharge of active layer water (subjected
435 to evaporation) to the near-surface ground ice (Yang et al., 2013; Throckmorton et al., 2016). In addition,
436 the thaw slump ice exhibited more negative isotopes, which is even lower than the amount-weighted
437 average value of winter precipitation (Fig. 7), indicating the main recharge of snowmelt water (Yang et
438 al., 2020; Opel et al., 2018).



439

440 **Figure 7: Hydrological connections between different water components.**

441 **6 Data availability**

442 The dataset provided in this paper can be obtained at <https://doi.org/10.5281/zenodo.10684110> (Yang,
443 2024). The link will become publicly available until full publication.

444 **7 Conclusions**

445 From 2017 to 2022, we constructed the first stable isotope monitoring network in a typical
446 permafrost-dominated watershed (namely the Beiluhe Basin, BLH) in central Qinghai-Tibet Plateau
447 (QTP). Totally, we obtained 554 precipitation samples, 2402 lakes/ponds samples, 675 stream water
448 samples, 102 supra-permafrost water samples, and 19 sub-permafrost water samples. Importantly, 359
449 ground ice samples at different depths from 17 boreholes and 2 profiles were collected, which is the first
450 detailed isotopic data of permafrost ice on the QTP. The following findings are drawn:

451 1) The stable isotopes of precipitation display distinct seasonal patterns with high $\delta^{18}\text{O}$ (δD) values
452 in summer and low $\delta^{18}\text{O}$ (δD) values in winter. The slope of LMWL is reflected the global mean.
453 However, the intercepts are quietly different due to the influences of precipitation amounts and the
454 exceptional meteorological conditions.

455 2) The thermokarst lakes/ponds and streams exhibit remarkable seasonal patterns in stable isotopes,
456 which is due to the transition of source water and evaporation differences. The lower isotopic values in

457 August and September are attributed to the recharges of monsoonal precipitation and melting ground ice.
458 Evaporation enrichment and recharges of precipitation with heavier isotopes greatly influenced the
459 isotopic patterns in May, June, July, and October. The slopes of the three LELs are all lower than those
460 of LMWL, indicating strong evaporation due to lower relative humidity. The supra-permafrost water was
461 recharged by precipitation via infiltration. By contrast, the sub-permafrost water was replenished by
462 unchanged sources of isotopically lighter water during cold periods.

463 3) The stable isotopes of ground ice varied between different boreholes. It is attributed to the
464 influences of initial source water and complex ice formation mechanism. The near-surface ground ice
465 was closely related to the recent precipitation and active layer hydrology, however, the deep-layer ground
466 ice exhibited complicated formation mechanism. In addition, variability in the isotopic patterns along
467 depths suggested influence of lithology on the water migration and freezing fractionation of stable
468 isotopes. The freezing line of the ground ice is significantly different from the LMWL, reflected the
469 freezing characteristics of liquid water under different conditions.

470 This first comprehensive data set provides a new basis for studying the isotopic hydrology and
471 exploring the hydrological effects of degrading permafrost on the QTP. It also enriches the cryospheric
472 database of the Northern Hemisphere.

473 **Author contributions**

474 YY and QW conceived the idea of the study. YY designed the isotope observation network and
475 completed the manuscript. XG and ZZ analyzed water samples and plotted figures. LZ, HY, and DZ
476 participated the field work. JC and GL provided and analyzed the meteorological data.

477 **Acknowledgements**

478 This work was supported by the Key Research Program of Frontier Sciences, CAS (Grant No. ZDBS–
479 LY–DQC026), the National Natural Science Foundation of China (Grant No. 41871062; 42177073), and
480 the foundation of the State Key Laboratory of Frozen Soil Engineering (Grant No. SKLFSE–ZT–202118).
481 We give special thanks to Hongtan, Yandong Hou, Jing Zhan, Siru Gao, Guanli Jiang, and Peng Zhang
482 for their kind help during field sampling work.

References

- 485 [1] Aichner, B., Dubbert, D., Kiel, C., Kohnert, K., Ogashawara, I., Jechow, A., ... & Berger, S. A. 2022. Spatial and seasonal patterns of water isotopes in northeastern German lakes. *Earth System Science Data*, 14(4), 1857-1867.
- [2] Ala-aho, P., Soulsby, C., Pokrovsky, O., Kirpotin, S.N., Karlsson, J.P., Serikova, S., Manasypov, R., Lim, A., Krickov, I., Kolesnichenko, L.G., 2018. Permafrost and lakes control river isotope composition across a boreal-arctic transect in the Western Siberia lowlands. *Environmental Research Letters*, 13(3): 034028.
- 490 [3] Barešić, J., Horvatinčić, N., Krajcar Bronić, I., Obelić, B., & Vreča, P. 2006. Stable isotope composition of daily and monthly precipitation in Zagreb. *Isotopes in Environmental and Health Studies*, 42(3), 239-249.
- [4] Bershaw, J.; Penny, S.M.; Garzzone, C.N. 2012. Stable isotopes of modern water across the Himalaya and eastern Tibetan Plateau: Implications for estimates of paleoelevation and paleoclimate. *J. Geophys. Res. Atmos*, 117.
- [5] Biskaborn, B.K., Smith, S.L., Noetzli, J., Matthes, H., Vieira, G., Streletskiy, D.A., ... & Lantuit, H., 2019. Permafrost is warming at a global scale. *Nature Communications*, 10(1), 264.
- 495 [6] Brown, J., Sidlauskas, F. J., & Delinski, G. F. 1997. (Eds.): *Circum-Arctic map of permafrost and ground ice conditions*. Washington, DC: U.S. Geological Survey in Cooperation with the Circum-Pacific Council for Energy and Mineral Resources, Circum-Pacific Map Series CP-45, scale 1:10,000,000, 1 sheet.
- [7] Cheng, G., Zhao, L., Li, R., Wu, X., Sheng, Y., Hu, G., Zou, D., Jin, H., Li, X., Wu, Q., 2019. Characteristic, changes and impacts of permafrost on Qinghai-Tibet Plateau. *Chinese Science Bulletin*, 64(27), 2783-2795.
- 500 [8] Cheng, G.D., Jin, H.J. 2013. Permafrost and Groundwater on the Qinghai-Tibet Plateau and in Northeast China. *Hydrogeology Journal*, 21 (1): 5-23.
- [9] Clark, I., Fritz, P. 1997. *Environmental Isotopes in Hydrogeology*. Lewis Publishers, Boca Raton – New York.
- [10] Cannon RF, Quinton WL, Craig JR, Hayashi M. 2014. Changing hydrologic connectivity due to permafrost thaw in the lower Liard River valley, NWT, Canada. *Hydrological Processes*, 28(14): 4163–4178.
- 505 [11] Craig, H. 1961. Isotopic variations in meteoric waters. *Science*, 133(3465), 1702-1703.
- [12] Cui, J., Tian, L., Biggs, T. W., & Wen, R. 2017. Deuterium-excess determination of evaporation to inflow ratios of an alpine lake: Implications for water balance and modeling. *Hydrological Processes*, 31(5), 1034-1046.
- [13] Douglas, T. A., Hiemstra, C. A., Anderson, J. E., Barbato, R. A., Bjella, K. L., Deeb, E. J., ... & Wagner, A. M., 2021. Recent degradation of Interior Alaska permafrost mapped with ground surveys, geophysics, deep drilling, and repeat
- 510 airborne lidar. *The Cryosphere*, 15(8), 3555-3575.
- [14] Fisher, D.A., Lacelle, D., Pollard, W., Faucher, B., 2021. A model for stable isotopes of residual liquid water and ground ice in permafrost soils using arbitrary water chemistries and soil-specific empirical residual water functions. *Permafrost and Periglacial Processes*, 32(2): 248-260.
- [15] Fritz, M., Wetterich, S., Meyer, H., Schirrmeister, L., Lantuit, H., & Pollard, W. H. 2011. Origin and characteristics of massive ground ice on Herschel Island (western Canadian Arctic) as revealed by stable water isotope and hydrochemical signatures. *Permafrost and Periglacial Processes*, 22(1), 26-38.
- 515 [16] Gibson, J. J., Birks, S. J., Yi, Y., & Vitt, D. 2015. Runoff to boreal lakes linked to land cover, watershed morphology and permafrost thaw: a 9-year isotope mass balance assessment. *Hydrological Processes*, 29(18), 3848-3861.
- [17] Guan, H., Zhang, X., Skrzypek, G., Sun, Z. and Xu, X., 2013. Deuterium excess variations of rainfall events in a coastal area of South Australia and its relationship with synoptic weather systems and atmospheric moisture sources. *Journal of Geophysical Research: Atmospheres*, 118(2), 1123–1138.
- 520 [18] Guo, D., Wang, H., Wang, A., 2017. Sensitivity of historical simulation of the permafrost to different atmospheric forcing data sets from 1979 to 2009. *Journal of Geophysical Research: Atmospheres*, 122(22), 12-269.
- [19] Guo, X., Feng, Q., Si, J., & Zhang, X. 2022. Considerable influences of recycled moistures and summer monsoons to local precipitation on the northeastern Tibetan Plateau. *Journal of Hydrology*, 605, 127343.
- 525 [20] Hughes, C. E., & Crawford, J. 2012. A new precipitation weighted method for determining the meteoric water line for hydrological applications demonstrated using Australian and global GNIP data. *Journal of Hydrology*, 464, 344-351.
- [21] Immerzeel, W. W., Van Beek, L. P., & Bierkens, M. F. 2010. Climate change will affect the Asian water towers. *Science*,

- 328(5984), 1382-1385.
- 530 [22] IPCC, 2019: Climate Change and Land: an IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems.
- [23] Jouzel, J., Souchez, R. A. 1982. Melting–refreezing at the glacier sole and the isotopic composition of the ice. *Journal of Glaciology*, 28(98), 35-42.
- 535 [24] Kern, Z., Harmon, R. S., & Fórizs, I. 2016. Stable isotope signatures of seasonal precipitation on the Pacific coast of central Panama. *Isotopes in Environmental and Health Studies*, 52(1-2), 128-140.
- [25] Knapp, J.L., Neal, C., Schlumpf, A., Neal, M., Kirchner, J.W., 2019. New water fractions and transit time distributions at Plynlimon, Wales, estimated from stable water isotopes in precipitation and streamflow. *Hydrology and Earth System Sciences*, 23(10): 4367-4388.
- 540 [26] Kuang, X., & Jiao, J. J. 2016. Review on climate change on the Tibetan Plateau during the last half century. *Journal of Geophysical Research: Atmospheres*, 121(8), 3979-4007.
- [27] Lacelle, D. 2011. On the $\delta^{18}\text{O}$, δD and D-excess relations in meteoric precipitation and during equilibrium freezing: theoretical approach and field examples. *Permafrost and Periglacial Processes*, 22(1), 13-25.
- [28] Lacelle, D., Fontaine, M., Forest, A. P., & Kokelj, S. 2014. High-resolution stable water isotopes as tracers of thaw unconformities in permafrost: A case study from western Arctic Canada. *Chemical Geology*, 368, 85-96.
- 545 [29] Lacelle, D., Vasil'chuk, Y. K., 2013. Recent progress (2007–2012) in permafrost isotope geochemistry. *Permafrost and Periglacial Processes*, 24(2), 138-145.
- [30] Lawrence, D. M., Slater, A. G., 2005. A projection of severe near-surface permafrost degradation during the 21st century. *Geophysical Research Letters*, 32(24).
- 550 [31] Liu, F., Qin, S., Fang, K., Chen, L., Peng, Y., Smith, P., & Yang, Y. 2022a. Divergent changes in particulate and mineral-associated organic carbon upon permafrost thaw. *Nature Communications*, 13(1), 5073.
- [32] Liu, L., Zhuang, Q., Zhao, D., Zheng, D., Kou, D., & Yang, Y. 2022b. Permafrost Degradation Diminishes Terrestrial Ecosystem Carbon Sequestration Capacity on the Qinghai-Tibetan Plateau. *Global Biogeochemical Cycles*, 36(2), e2021GB007068.
- 555 [33] Luo, J., Niu, F., Lin, Z., Liu, M., & Yin, G. 2015. Thermokarst lake changes between 1969 and 2010 in the Beilu River basin, Qinghai–Tibet plateau, China. *Science Bulletin*, 60(5), 556-564.
- [34] Michel, F. A., 2011. Isotope characterisation of ground ice in northern Canada. *Permafrost and Periglacial Processes*, 22(1), 3-12.
- [35] Murton, J.B., 2013. Ground Ice and Cryostratigraphy. In: Shroder, J.F. (Ed.), *Treatise on Geomorphology*. Academic Press, San Diego. 173-201.
- 560 [36] Narancic, B., Wolfe, B.B., Pienitz, R., Meyer, H., Lamhonwah, D., 2017. Landscape-gradient assessment of thermokarst lake hydrology using water isotope tracers. *Journal of Hydrology*, 545: 327-338.
- [37] Niu, F., Lin, Z., Liu, H., & Lu, J. 2011. Characteristics of thermokarst lakes and their influence on permafrost in Qinghai–Tibet Plateau. *Geomorphology*, 132(3-4), 222-233.
- 565 [38] Opel, T., Meyer, H., Wetterich, S., Laepple, T., Dereviagin, A., Murton, J., 2018. Ice wedges as archives of winter paleoclimate: A review. *Permafrost and Periglacial Processes*, 29(3): 199-209.
- [39] Perşoiu, A., & Pazdur, A. 2011. Ice genesis and its long-term mass balance and dynamics in Scărișoara Ice Cave, Romania. *The Cryosphere*, 5(1), 45-53.
- [40] Porter, T. J., & Opel, T. 2020. Recent advances in paleoclimatological studies of Arctic wedge-and pore-ice stable-water isotope records. *Permafrost and Periglacial Processes*, 31(3), 429-441.
- 570 [41] Porter, T.J., Schoenemann, S.W., Davies, L.J., Steig, E.J., Bandara, S., Froese, D.G., 2019. Recent summer warming in northwestern Canada exceeds the Holocene thermal maximum. *Nature Communications*, 10(1): 1631.
- [42] Quinton WL, Baltzer LJ. 2013. The active-layer hydrology of a peat plateau with thawing permafrost (scotty creek, canada). *Hydrogeology Journal*, 21 (1): 201-220.
- 575 [43] Ran, Y., Li, X., Cheng, G., Che, J., Aalto, J., Karjalainen, O., ... & Chang, X. 2022. New high-resolution estimates of the permafrost thermal state and hydrothermal conditions over the Northern Hemisphere. *Earth System Science Data Discussions*, 14 (2), 865–884.
- [44] Rogger M, Chirico G, Hausmann H, Krainer K, Brückl E, Stadler P, Blöschl G. 2017. Impact of mountain permafrost on

- flow path and runoff response in a high alpine catchment. *Water Resources Research*, 53(2):1288-1308.
- 580 [45] Schwamborn G, Meyer H, Chirrneister L, G, F., 2014. Past freeze and thaw cycling in the margin of the El'gygytgyn Crater deduced from a 141m long permafrost record. *Climate of the Past*, 10: 1109-1123.
- [46] Song, C., Wang, G., Liu, G., Mao, T., Sun, X., & Chen, X. (2017). Stable isotope variations of precipitation and streamflow reveal the young water fraction of a permafrost watershed. *Hydrological Processes*, 31(4), 935-947.
- [47] Souchez, R. A., & Jouzel, J. 1984. On the isotopic composition in δD and $\delta^{18}O$ of water and ice during freezing. *Journal of Glaciology*, 30(106), 369-372.
- 585 [48] Souchez, R., Jouzel, J., Lorrain, R., Sleewaegen, S., Stiévenard, M., & Verbeke, V. 2000. A kinetic isotope effect during ice formation by water freezing. *Geophysical Research Letters*, 27(13), 1923-1926.
- [49] Streletskiy, D.A., Tananaev, N.I., Opel, T., Shiklomanov, N.I., Nyland, K.E., Streletskaia, I.D., Shiklomanov, A.I., 2015. Permafrost hydrology in changing climatic conditions: seasonal variability of stable isotope composition in rivers in discontinuous permafrost. *Environmental Research Letters*, 10(9): 095003.
- 590 [50] Throckmorton, H. M., Newman, B. D., Heikoop, J. M., Perkins, G. B., Feng, X., Graham, D. E., ... & Wilson, C. J. 2016. Active layer hydrology in an arctic tundra ecosystem: quantifying water sources and cycling using water stable isotopes. *Hydrological Processes*, 30(26), 4972-4986.
- [51] Tian, L., Yao, T., White, J.W.C., Yu, W., Wang, N., 2005. Westerlies moisture transport to the middle of Himalayas revealed from the high deuterium excess. *Chinese Science Bulletin*, 50(10), 1026-1030.
- 595 [52] Vasil'chuk, Y.K., Lawson, D.E., Yoshikawa, K., Budantseva, N.A., Chizhova, J.N., Podborny, Y.Y., Vasil'chuk, A.C., 2016. Stable isotopes in the closed-system Weather Pingo, Alaska and Pestsovoye Pingo, northwestern Siberia. *Cold Regions Science and Technology*, 128: 13-21.
- [53] Vystavna, Y., Harjung, A., Monteiro, L. R., Matiatos, I., & Wassenaar, L. I. 2021. Stable isotopes in global lakes integrate catchment and climatic controls on evaporation. *Nature Communications*, 12(1), 7224.
- 600 [54] Wang, G., Li, Y., Wu, Q., & Wang, Y. 2006. Impacts of permafrost changes on alpine ecosystem in Qinghai-Tibet Plateau. *Science in China Series D: Earth Sciences*, 49, 1156-1169.
- [55] Wang, S., He, X., Kang, S., Fu, H., & Hong, X. 2022. Estimation of stream water components and residence time in a permafrost catchment in the central Tibetan Plateau using long-term water stable isotopic data. *The Cryosphere*, 16(12), 5023-5040.
- 605 [56] Wang, S., He, X., Kang, S., Hong, X., Fu, H., Xue, Y., ... & Guo, H. 2023b. Assessment of streamwater age using water stable isotopes in a headwater catchment of the central Tibetan Plateau. *Journal of Hydrology*, 618, 129175.
- [57] Wang, T., Yang, D., Yang, Y., Zheng, G., Jin, H., Li, X., ... & Cheng, G. 2023a. Unsustainable water supply from thawing permafrost on the Tibetan Plateau in a changing climate. *Science Bulletin*, S2095-9273.
- [58] Wetterich, S., Tumskey, V., Rudaya, N., Andreev, A. A., Opel, T., Meyer, H., ... & Hüls, M. 2014. Ice Complex formation in arctic East Siberia during the MIS3 Interstadial. *Quaternary Science Reviews*, 84, 39-55.
- 610 [59] Wu, Q., & Zhang, T. 2010. Changes in active layer thickness over the Qinghai-Tibetan Plateau from 1995 to 2007. *Journal of Geophysical Research: Atmospheres*, 115(D9).
- [60] Yang, K., Ye, B., Zhou, D., Wu, B., Foken, T., Qin, J., & Zhou, Z. 2011. Response of hydrological cycle to recent climate changes in the Tibetan Plateau. *Climatic Bhang*, 109, 517-534.
- 615 [61] Yang, Y., Guo, X., Wu, Q., Jin, H., & Liu, F. 2023. Formation processes of shallow ground ice in permafrost in the Northeastern Qinghai-Tibet Plateau: A stable isotope perspective. *Science of The Total Environment*, 863, 160967.
- [62] Yang, Y., Wu, Q., Jiang, G., Zhang, P., 2017. Stable Isotopic Stratification and Growth Patterns of Ground Ice in Permafrost on the Qinghai-Tibet Plateau, China. *Permafrost and Periglacial Processes*, 28(1), 119-129.
- [63] Yang, Y., Wu, Q., Jiang, G., Zhang, P., 2020. Ground ice at depths in the Tianshuihai Lake basin on the western Qinghai-Tibet Plateau: An indication of permafrost evolution. *Science of the Total Environment*, 729, 138966.
- 620 [64] Yang, Y., Wu, Q., Jin, H., Wang, Q., Huang, Y., Luo, D., Gao, S., Jin, X., 2019. Delineating the hydrological processes and hydraulic connectivities under permafrost degradation on Northeastern Qinghai-Tibet Plateau, China. *Journal of hydrology*, 569, 359-372.
- [65] Yang, Y., Wu, Q., Liu, F., & Jin, H. 2021. Spatial-temporal trends of hydrological transitions in thermokarst lakes on Northeast Qinghai-Tibet Plateau based on stable isotopes. *Journal of Hydrology*, 597, 126314.
- 625 [66] Yang, Y., Wu, Q., Yun, H., 2013. Stable isotope variations in the ground ice of Beiluhe Basin on the Qinghai-Tibet

Plateau. *Quaternary International*, 313, 85-91.

- [67] Yang, Y., Wu, Q., Yun, H., Jin, H., Zhang, Z., 2016. Evaluation of the hydrological contributions of permafrost to the thermokarst lakes on the Qinghai–Tibet Plateau using stable isotopes. *Global and Planetary Change*, 140, 1-8.
- 630 [68] Yao, T., Masson-Delmotte, V., Gao, J., Yu, W., Yang, X., Risi, C., Sturm, C., Zhao, H., He, Y. and Ren, W., 2013. A review of climatic controls on $\delta^{18}\text{O}$ in precipitation over the Tibetan Plateau: Observations and simulations. *Reviews of Geophysics*, 51(4), 525–548.
- [69] Yao, T., Qin, D., Shen, Y., Zhao, L., Wang, N., Lu, A., 2013. Cryospheric changes and their impacts on regional water cycle and ecological conditions in the Qinghai-Tibetan Plateau. *Chin. J. Nat.*, 35(3), 179-186.
- 635 [70] Yi, S., Wang, X., Qin, Y., Xiang, B., & Ding, Y. 2014. Responses of alpine grassland on Qinghai–Tibetan plateau to climate warming and permafrost degradation: a modeling perspective. *Environmental Research Letters*, 9(7), 074014.
- [71] Yin, G., Niu, F., Lin, Z., Luo, J., & Liu, M. 2017. Effects of local factors and climate on permafrost conditions and distribution in Beiluhe basin, Qinghai-Tibet Plateau, China. *Science of the Total Environment*, 581, 472-485.
- [72] Zhang X, He J, Zhang J, Polyakov I, Gerdes R, Inoue J, Wu P. 2013. Enhanced poleward moisture transport and amplified northern high-latitude wetting trend. *Nature Climate Change*, 3: 47-51.
- 640 [73] Zhang, T., Frauenfeld, O.W., Serreze, M.C., Etringer, A., Oelke, C., McCreight, J., Barry, R.G., Gilichinsky, D., Yang, D., Ye, H., Ling, F., Chudinova, S., 2005. Spatial and temporal variability in active layer thickness over the Russian Arctic drainage basin. *Journal of Geophysical Research: Atmospheres*, 110(D16), 1-14.
- [74] Zhao, L., Zou, D., Hu, G., Du, E., Pang, Q., Xiao, Y., Li, R., Sheng, Y., Wu, X., Sun, Z., Wang, L., Wang, C., Ma, L., Zhou, H., Liu, S., 2020. Changing climate and the permafrost environment on the Qinghai–Tibet (Xizang) Plateau. *Permafrost and Periglacial Processes*, 31(3), 396-405.
- 645 [75] Zhao, L., Zou, D., Hu, G., Wu, T., Du, E., Liu, G., ... & Cheng, G. 2021. A synthesis dataset of permafrost thermal state for the Qinghai–Tibet (Xizang) Plateau, China. *Earth System Science Data*, 13(8), 4207-4218.
- [76] Zhu, G., Liu, Y., Shi, P., Jia, W., Zhou, J., Liu, Y., ... & Zhao, K. 2022. Stable water isotope monitoring network of different water bodies in Shiyang River basin, a typical arid river in China. *Earth System Science Data*, 14(8), 3773-3789.
- 650