

The JapanFlux2024 dataset for eddy covariance observations covering Japan and East Asia from 1990 to 2023

- Masahito Ueyama¹, Yuta Takao¹, Hiromi Yazawa², Makiko Tanaka², Hironori Yabuki³, Tomo'omi Kumagai⁴, Hiroki Iwata⁵, Md. Abdul Awal⁶, Mingyuan Du⁷, Yoshinobu Harazono⁸, Yoshiaki Hata⁴,
5 Takashi Hirano⁹, Tsutomu Hiura⁴, Reiko Ide¹⁰, Sachinobu Ishida¹¹, Mamoru Ishikawa¹², Kenzo Kitamura¹³,
Yuji Kominami¹⁴, Shujiro Komiya¹⁵, Ayumi Kotani¹⁶, Yuta Inoue¹⁴, Takashi Machimura¹⁷, Kazuho Matsumoto¹⁸,
Yojiro Matsuura¹⁴, Yasuko Mizoguchi¹⁹, Shohei Murayama²⁰, Hirohiko Nagano²¹, Taro Nakai²², Tatsuro Nakaji²³,
10 Ko Nakaya²⁴, Shinjiro Ohkubo²⁵, Takeshi Ohta²⁶, Keisuke Ono²⁷, Taku M. Saitoh²⁸, Ayaka Sakabe²⁹, Takanori Shimizu¹⁴,
Seiji Shimoda³⁰, Michiaki Sugita³¹, Kentaro Takagi³², Yoshiyuki Takahashi¹⁰, Naoya Takamura⁴, Satoru Takanashi¹⁹, Takahiro Takimoto²⁷,
Yukio Yasuda¹⁴, Qinxue Wang¹⁰, Jun Asanuma³³, Hideo Hasegawa²¹, Tetsuya Hiyama³⁴, Yoshihiro Iijima³⁵, Shigeyuki Ishidoya²⁰,
Masayuki Itoh³⁶, Tomomichi Kato⁹, Hiroaki Kondo²⁰, Yoshiko Kosugi²⁹, Tomonori Kume³⁷, Takahisa Maeda²⁰,
Shoji Matsuura²⁷, Trofim Maximov³⁸, Takafumi Miyama¹⁴, Ryo Moriwaki³⁹, Hiroyuki Muraoka⁴, Roman Petrov³⁸,
Jun Suzuki⁴⁰, Shingo Taniguchi⁴¹, & Kazuhito Ichii²
- 15 ¹Graduate School of Agriculture, Osaka Metropolitan University, Sakai 599-8531, Japan
²Center for Environmental Remote Sensing (CEReS), Chiba University, Chiba 263-8522, Japan
³National Institute of Polar Research (NIPR), Tokyo 190-8518, Japan
⁴Graduate School of Agricultural and Life Sciences, The University of Tokyo, Tokyo 113-8657, Japan
⁵Department of Environmental Science, Faculty of Science, Shinshu University, Matsumoto 390-8621, Japan
20 ⁶Department of Crop Botany, Bangladesh Agricultural University, Mymensingh 2202, Bangladesh
⁷Xingjiang Institute of Ecology and Geography, Chinese Academy of Sciences, Xinjiang 830011, China
⁸International Arctic Research Center, University of Alaska Fairbanks, Fairbanks, AK 99775, USA
⁹Research Faculty of Agriculture, Hokkaido University, Sapporo 060-8589, Japan
¹⁰National Institute for Environmental Studies, Tsukuba 305-8506, Japan
25 ¹¹Graduate School of Science and Technology, Hirosaki University, Hirosaki 036-8561, Japan
¹²Faculty of Earth Environmental Science, Hokkaido University, Sapporo 060-0810 Japan
¹³Kyushu Research Center, Forestry and Forest Products Research Institute, Kumamoto 860-0862, Japan
¹⁴Forestry and Forest Products Research Institute, Tsukuba 305-8687, Japan
¹⁵Department of Biogeochemical Processes, Max Planck Institute for Biogeochemistry, Jena 07745, Germany
30 ¹⁶Graduate School of Bioagricultural Sciences, Nagoya University, Nagoya 464-8601, Japan
¹⁷Graduate School of Engineering, Osaka University, Suita 565-0871, Japan
¹⁸Faculty of Agriculture, Iwate University, Morioka 020-8550, Japan
¹⁹Kansai Research Center, Forestry and Forest Products Research Institute, Kyoto 612-0855, Japan
²⁰National Institute of Advanced Industrial Science and Technology (AIST), Tsukuba 305-8569, Japan
35 ²¹Institute of Science and Technology, Niigata University, Niigata 950-2181, Japan
²²School of Forestry and Resource Conservation, National Taiwan University, Taipei 106319, Taiwan
²³Sapporo Experimental Forest, Hokkaido University, Sapporo 060-0809, Japan
²⁴Sustainable System Research Laboratory, Central Research Institute of Electric Power Industry, Abiko 270-1194, Japan
²⁵Forestry Research Institute, Forest Research Department, Hokkaido Research Organization, Bibai 079-0198, Japan
40 ²⁶Professor emeritus, Nagoya University, Nagoya 464-8601 Japan
²⁷Institute for Agro-Environmental Sciences, National Agriculture and Food Research Organization (NARO), Tsukuba 305-8604 Japan
²⁸Center for Environmental and Societal Sustainability, Gifu University, Gifu 501-1193, Japan

- ²⁹Graduate School of Agriculture, Kyoto University, Kyoto 606-8501 Japan
- 45 ³⁰Memuro Research Station, Hokkaido Agricultural Research Center, NARO, (HARC/M /NARO), Memuro 082-0081, Japan
- ³¹Institute of Life and Environmental Sciences, University of Tsukuba, Tsukuba 305-8572, Japan
- ³²Field Science Center for Northern Biosphere, Hokkaido University, Toikanbetsu, 098-2943, Japan
- ³³Center for Research in Isotopes and Environmental Dynamics (CRiED), University of Tsukuba, Tsukuba 305-8572, Japan
- ³⁴Institute for Space-Earth Environmental Research, Nagoya University, Nagoya 464-8601, Japan
- 50 ³⁵Department of Geography, Tokyo Metropolitan University, Tokyo 192-0397 Japan
- ³⁶Research Institute for Sustainable Humanosphere, Kyoto University, Uji 611-0011, Japan
- ³⁷Kasuya Research Forest, Kyushu University, Fukuoka 811-2415, Japan
- ³⁸Institute for Biological Problems of Cryolithozone, Yakutsk 677980, Russia
- ³⁹Faculty of Engineering, Ehime University, Matsuyama 790-8577, Japan
- 55 ⁴⁰Faculty of Agriculture, Shinshu University, Nagano 399-4598, Japan
- ⁴¹Faculty of Agriculture, University of the Ryukyus, Okinawa 903-0213, Japan

Correspondence to: Masahito Ueyama (mueyama@omu.ac.jp)

Abstract. Eddy covariance observations play a pivotal role in understanding the land–atmosphere exchange of energy, water, carbon dioxide (CO₂), and other trace gases, as well as the global carbon cycle and earth system. To promote the networking of individual measurements and the sharing of data, FLUXNET links regional networks of researchers studying land–atmosphere processes. JapanFlux was established in 2006 as a country branch of AsiaFlux. Despite the growing number of shared data globally, the availability in Asia is currently limited. In this study, we developed an open dataset of the eddy covariance observations for Japan and East Asia, called JapanFlux2024, that was conducted by researchers affiliated with Japanese research institutions. The data was processed using selected standard methods from the FLUXNET community, with adaptations specific to the JapanFlux2024 dataset. Here, we present the data description and data processing and show the value of processed fluxes of sensible heat, latent heat, and CO₂. The dataset will facilitate important studies for Japan and East Asia, such as land-atmosphere interactions, improvement of process models, and upscaling fluxes using machine learning and remote sensing technology as well as bridge collaborations between Asia and FLUXNET.

70

1 Introduction

The global network of micrometeorological flux observations, FLUXNET (Delwiche et al., 2024; <https://fluxnet.org/>), plays a pivotal role in multi-disciplinary fields, such as land–atmosphere interactions, global biogeochemical cycles, and earth system science (Baldocchi et al., 2024; Bonan et al., 2012). FLUXNET started in 1997 as a global network of eddy covariance observations that provides data on land–atmosphere exchanges of energy, water, carbon dioxide (CO₂), methane (CH₄), and other trace gases by measuring direct turbulent transfer. The quasi-continuous eddy covariance observations revealed variations of land–atmosphere exchange at the diurnal, seasonal, interannual, and decadal scales, ranging from site (Takamura et al., 2023; Ueyama et al., 2024f) to global scales (Beer et al., 2010; Keenan et al., 2023; Ueyama et al., 2020a).

80 The eddy flux communities have developed publicly open databases to promote the multidisciplinary sciences. FLUXNET has
periodically released the open datasets for eddy covariance observations: La Thuile Database (252 sites in 2007; Verma et al.,
2014; <https://fluxnet.org/data/la-thuille-dataset/>), and FLUXNET2015 (212 sites in 2015; Pastorello et al., 2020). Together with
the global carbon project (Friedlingstein et al., 2023; www.globalcarbonproject.org), FLUXNET also provided a topical
dataset, FLUXNET-CH₄ (Delwiche et al., 2021), which promotes understanding of wetland CH₄ emissions across the globe
85 (Knox et al., 2019; Ueyama et al., 2023). Multiple open databases for the environmental sciences have also been developed
for understanding CO₂ fluxes in high-latitude ecosystems (Virkkala et al., 2021) and soil respiration (Bond-Lamberty et al.,
2020).

Asia has ca. ~60% of the total world population, and thus humans have been intensively modifying forest land cover in this
90 region for food and energy production. Such land use changes in combination with climate change are likely to impact the
regional and global carbon and water cycling. These issues are the greatest environmental concerns for the survival of the
human population. Flux studies using eddy covariance observations were conducted since the early 1990s for agricultural
fields, wetlands, lakes, plantations, primary and secondary forests, disturbed ecosystems, and urban areas. In Asia, although
private databases for eddy covariance measurements were developed (Hirata et al., 2008; Ichii et al., 2017; Saigusa et al.,
95 2013), no open databases have yet been developed except the AsiaFlux database (<https://asiaflux.net/>), which does not provide
consistent gap-filling and flux partitioning.

JapanFlux (<https://www.japanflux.org/>) was established in 2006 as a national branch of AsiaFlux (Kang and Cho, 2021;
Mizoguchi et al., 2009) for the promotion of a network of micrometeorological measurements by researchers affiliated with
100 Japanese research institutions. The mission of JapanFlux is to promote micrometeorological measurements and their
collaborations with each other, researchers from other countries, and other research fields (e.g., remote sensing and modeling).
Measurements by Japanese institutions have been conducted in Japan and other regions of East Asia (Mizoguchi et al., 2009;
Saigusa et al., 2013) since the early 1990s for understanding energy, water, carbon, and greenhouse gas exchanges at various
land surfaces.

105

In this study, we developed JapanFlux2024, the first publicly open dataset by JapanFlux that consists of micrometeorological
data measured since the early 1990s. The data is processed with the selected standard methods employed by the FLUXNET
community. The dataset is prepared with consistent post-processing, such as gap-filling and flux partitioning, and provides
data at various temporal resolutions of half-hourly/hourly, daily, weekly, monthly, and annual intervals. The dataset consists
110 of data collected at 83 sites with 683 site-years. The dataset promotes collaborations between researchers in Japan and other
countries and improves our understanding of land-atmosphere interactions.

2 Data and methods

The JapanFlux2024 dataset is processed using selected standard methods from the FLUXNET community, with adaptations specific to the JapanFlux2024 dataset. According to the processing strategy of Pastorello et al., (2020), the JapanFlux2024 dataset was developed in four steps: (1) data submission by site teams, (2) formatting data in a FLUXNET format, (3) gap-filling and flux partitioning, and (4) preparing subsets and complete datasets (Fig. 1). Meta data files, so-called Biological, Ancillary, Disturbance, and Metadata (BADM), were also prepared. The data are available from the data portal (<https://ads.nipr.ac.jp/japan-flux2024/>) under the data management system, Arctic and Antarctic Data archive System (ADS). Under the ADS, a digital object identifier (DOI) was provided for each site (Table 1). The processing pipeline mentioned in this data paper represent steps downstream of “Filling gaps in meteorology with ERA5” in Fig. 1.




Figure 1. Flow chart of data processing in the JapanFlux2024 dataset. Details in each step and meaning of abbreviations are shown in the text.

Table 1. Information about sites included in the JapanFlux2024 dataset.

Site Code (BAD M)	Asia Flux ID	Country	Country ID	Site Name	Latitude (degree)	Longitude (degree)	Elevation (m)	Köppen climate	IGBP (land use)	Status	Years	Reference	doi
RU-Tur	TUR	Russia	RU	Tura	64.20888	100.463555	250	Dfc	DNF	Ongoing	04	Nakai et al. (2008)	Matsuura and Morishita (2025)
RU-NeB		Russia	RU	Neleger Burnt Forest	62.325937	129.487342	221	Dfd	GRA	Completed	99-00	Iwahana et al. (2005)	Machimura (2025a)
RU-NeF		Russia	RU	Neleger larch forest	62.315615	129.499964	223	Dfd	DNF	Completed	99-06	Iwahana et al. (2005)	Machimura (2025b)
RU-NeC		Russia	RU	Neleger Cutover	62.314844	129.500075	221	Dfd	OSH	Completed	01-06	Iwahana et al. (2005)	Machimura (2025c)
RU-SkP	YLF	Russia	RU	Yakutsk Spasskaya Pad larch	62.25471	129.618543	217	Dfc	DNF	Ongoing	04-14	Ohta et al. (2008)	Maximov et al. (2025b)
RU-Sk2	YPF	Russia	RU	Yakutsk Spasskaya Pad Pine	62.241291	129.651336	216	Dfc	ENF	Completed	04-08	Hamada et al. (2004)	Kotani et al. (2025)
RU-USk		Russia	RU	Ulakhan Sykkhan Alas	62.150995	130.527517	143		GRA	Completed	00	Yabuki et al. (2004)	Yabuki et al. (2025)
RU-Ege		Russia	RU	Elgeei forest station	60.01551563	133.8240123	203	Dfd	DNF	Ongoing	10-18	Kotani et al. (2014)	Maximov et al. (2025a)
MN-Skt	SKT	Mongolia	MN	Southern Khentei Taiga	48.351861	108.654333	1630	Dwc	DNF	Completed	03-06	Li et al. (2005b)	Asanuma (2025b)
MN-Udg		Mongolia	MN	Udleg practice forest	48.25638888	106.8511111	1342	Dwc	DNF	Ongoing	10-12	Miyazaki et al. (2014)	Ishikawa (2025)
MN-Nkh		Mongolia	MN	Nalaikh grassland	47.693592	107.489342	1531	BSk	GRA	Completed	15-20	Wang et al. (2023)	Wang et al. (2025b)
MN-Hst		Mongolia	MN	Hustai grassland	47.594131	105.856439	1227	BSk	GRA	Completed	15-20	Wang et al. (2023)	Wang et al. (2025a)

MN-Kbu	KBU	Mongolia	MN	Kherlenbay an Ulaan	47.2 1397 2	108.73 7333	1235	Bsk	GRA	Completed	03-09	Li et al. (2005a)	Asanuma (2025a)
CN-Lsh	LSH	China	CN	Laoshan	45.2 7983 9	127.57 8206	340	Cfc	DNF	Ongoing	02-06	Wang et al. (2005)	Saigusa and Wang (2025)
JP-Sb1		Japan	JP	Sarobetsu Mire Moss	45.1 0472 2	141.68 8194	6	Dfb	WET	Completed	07-10	Hirano et al. (2016)	Hirano (2025a)
JP-Sb2		Japan	JP	Sarobetsu Mire Sasa	45.1 0361 1	141.68 0833	4	Dfb	WET	Completed	07-10	Hirano et al. (2016)	Hirano (2025b)
JP-Tef	TSE	Japan	JP	CC-LaG Teshio Experiment al Forest	45.0 5580 8	142.10 7122	79.4 7	Dfb	DNF	Ongoing	01-23	Takagi et al. (2009)	Takagi and Takahashi (2025)
JP-MBF	MBF	Japan	JP	Moshiri Birch Forest Site	44.3 8416 667	142.31 86111	596	Af	DBF	Completed	03-11	Nakai et al. (2006)	Nakai et al (2025a)
JP-MMF	MMF	Japan	JP	Moshiri Mixd Forest Site	44.3 2194 444	142.26 13889	343	Af	MF	Completed	03-11	Nakai et al. (2006)	Nakai et al (2025b)
JP-BBY	BBY	Japan	JP	Bibai bog	43.3 2296	141.81 079	17	Dfb	WET	Completed	12-21	Ueyama et al. (2020c)	Ueyama et al. (2025a)
JP-Km1		Japan	JP	Kushiro Mire: Onnenai Fen	43.1 0751 1	144.33 0906	4.9	Dfb	WET	Completed	94-96, 98	Miyata et al. (2001)	Harazono and Miyata (2025a)
JP-Km2		Japan	JP	Kushiro Mire: Akanuma Bog	43.1	144.35	7	Dfb	WET	Completed	98-99	Miyata et al. (2001)	Harazono and Miyata (2025b)
JP-Spp	SAP	Japan	JP	Sapporo forest meteorology research site	42.9 8684 31	141.38 53305	174	Dfb	DBF	Ongoing	00-18	Yamanoi et al. (2015)	Mizoguchi and Kitamura (2025)
CN-In4		China	CN	Inner Mongolia maize	42.9 4413 333	120.72 66222	354	Bsk	CRO	Completed	94	Li et al. (2000)	Harazono and Takagi (2025d)

CN-In5		China	CN	Inner Mongolia no grazing	42.9 3415 833	120.70 90778	355	Bsk	GRA	Completed	92-94	Li et al. (2000)	Harazono and Takagi (2025e)
CN-In6		China	CN	Inner Mongolia heavy grazing	42.9 3401 389	120.71 15472	355	Bsk	GRA	Completed	92-94	Li et al. (2000)	Harazono and Takagi (2025f)
CN-In8		China	CN	Inner Mongolia medium grazing	42.9 3396 667	120.71 05306	355	Bsk	GRA	Completed	92, 94	Li et al. (2000)	Harazono and Takagi (2025h)
CN-In2		China	CN	Inner Mongolia grassland	42.9 3396 389	120.71 09639	355	Bsk	GRA	Completed	91	Li et al. (2000)	Harazono and Takagi (2025b)
CN-In7		China	CN	Inner Mongolia light grazing	42.9 3391 944	120.70 96056	355	Bsk	GRA	Completed	92-94	Li et al. (2000)	Harazono and Takagi (2025g)
CN-In1		China	CN	Inner Mongolia dune	42.9 2970 833	120.70 735	356	Bsk	BSV	Completed	90-91	Li et al. (2000)	Harazono and Takagi (2025a)
CN-In4		China	CN	Inner Mongolia maize	42.9 4413 333	120.72 66222	354	Bsk	CRO	Completed	94	Li et al. (2000)	Harazono and Takagi (2025d)
JP-Tmk	TM K	Japan	JP	Tomakomai Flux Research Site	42.7 3697 2	141.51 6944	140	Dfb	DNF	Completed	01-03	Hirano et al. (2003)	Hirata and Hirano (2025)
JP-Tmd	TM K	Japan	JP	Tomakomai Flux Research Site Disturbed	42.7 3591 1	141.52 3147	117	Dfb	DBF	Ongoing	05-23	Hirano et al. (2017)	Hirano and Hirata (2025)
JP-Toc		Japan	JP	Tomakomai Crane site	42.7 0972 7	141.56 5898	96	Dfb	DBF	Ongoing	10-14	Nakamura et al. (2014)	Nakaji et al. (2025)
JP-Tom	TOE	Japan	JP	Tomakomai Experimental Forest	42.6 9890 6	141.57 1488	90	Dfb	DBF	Completed	99-13	Shibata et al. (2005)	Nakaji (2025)

JP-Srk	SRK	Japan	JP	Shirakami Beech Forest Site	40.5 6548 5	140.12 7794	340	Dfa	DBF	Ongoin g	10-16	Ishida et al. (2009)	Ishida (2025)
JP-Api	API	Japan	JP	Appi forest meteorology research site	40.0 0135 8581 5243	140.93 658591 8296	831	Dfa	DBF	Ongoin g	00-22	Yasuda et al. (2012)	Yasuda (2025a)
JP-Mra	MRA	Japan	JP	Muramatsu Agricultural Field	37.6 9027 5	139.19 4429	43	Cfa	CRO	Ongoin g	23	Boiarskii and Hasegawa (2019)	Nagano and Hasegawa (2025)
CN-HaM	QHB	China	CN	Qinghai Flux Research Site	37.6 0743 2	101.33 2	3250	BSk	GRA	Ongoin g	01-14	Du et al. (2021)	Du et al. (2025)
JP-NsM	NSS	Japan	JP	Nasu Research Station, Manure Application Plot	36.9 1583 333	139.93 58333	320	Cfa	GRA	Completed	04-15	Matsuura et al. (2023)	Matsuura (2025a)
JP-NsC	NSS	Japan	JP	Nasu Research Station, Chemical Fertilizer Plot	36.9 15	139.93 66667	320	Cfa	GRA	Completed	04-15	Matsuura et al. (2023)	Matsuura (2025b)
JP-Kzw	KZW	Japan	JP	Karuizawa	36.4 0666 7	138.57 25	1385	Dfb	DBF	Completed	01-08	Nakaya et al. (2006)	Nakaya et al. (2025)
JP-Tkb		Japan	JP	Tsukuba Experimental Watershed	36.1 7337 9	140.17 6634	341	Cfa	ENF	Ongoin g	14, 18-21	Iida et al. (2020)	Shimizu et al. (2025b)
JP-Tak	TKY	Japan	JP	Takayama deciduous broadleaf forest site	36.1 4616 667	137.42 31111	1425	Dfb	DBF	Ongoin g	98-21	Murayama et al. (2024a)	Murayama et al. (2025b)
JP-Ta2	TKC	Japan	JP	Takayama evergreen coniferous forest site	36.1 3972 2	137.37 0833	800	Dfb	ENF	Ongoin g	05-22	Saitoh et al. (2010)	Saitoh and Tamagawa (2025)

JP-Tgf	TGF	Japan	JP	Terrestrial Environme nt Research Center, University of Tsukuba	36.1 1353	140.09 488	27	Cfa	GRA	Compl eted	02-22	Shimoda et al. (2005)	Asanuma and Shimoda (2025)
JP-KaP		Japan		Kasumigau ra lotus paddy	36.0 8	140.24	3	Cfa	CRO	Compl eted	97-98	Takagi et al. (2003)	Harazono and Takagi (2025)
JP-Mse	MSE	Japan	JP	Mase paddy flux site	36.0 5393	140.02 693	11	Cfa	CRO	Ongoin g	01-09	Saito et al. (2005)	Ono (2025)
JP- SwL	SWL	Japan	JP	Suwa Lake Site	36.0 4657 222	138.10 83528	758	Dfc	WAT	Ongoin g	15-23	Iwata et al. (2018)	Iwata (2025b)
JP-KaL		Japan	JP	Koshin, Lake Kasumigau ra	36.0 3777 8	140.40 4167	0.26(at the wate r level of Y.P. 1.1 m)	Cfa	WAT	Ongoin g	07-22	Sugita et al. (2020)	Sugita (2025)
JP-Nsb		Japan	JP	NIAES Soybean	36.0 2430 3	140.11 4975	24	Cfa	CRO	Compl eted	90	Harazono et al. (1992)	Harazono (2025a)
JP-Yrp		Japan	JP	Yawara Rice paddy	36.0 0766 667	140.03 01752	23	Cfa	CRO	Compl eted	93-95	NA	Harazono (2025b)
JP- Kwg	KW G	Japan	JP	Kawagoe forest meteorolog y research site	35.8 725	139.48 69	41	Cfa	DBF	Compl eted	97-02	Yasuda et al. (1998)	Yasuda (2025b)
JP-Shn		Japan	JP	Shinshu University Experiment al Forest Site	35.8 6575 5	137.93 2563	775	Dfa	MF	Ongoin g	14-19	NA	Iwata and Suzuki (2025)
JP- Nkm	NK M	Japan	JP	Nishikoma Site	35.8 0806 4	137.83 3883	2641	Dfb	ENF	Ongoin g	18-23	NA	Iwata (2025a)

JP-Fmt		Japan	JP	Field Museum Tama Hills	35.6 3874 5	139.37 9748	168	Cfa	MF	Ongoing	13-23	Matsuda et al. (2015)	Takagi and Matsuda (2025)
JP-Kgu		Japan	JP	Kugahara urban residencial area	35.5 8285 9	139.69 3543	18.5	Cfa	URB	Completed	01-02	Moriwaki and Kanda (2004)	Kanda and Moriwaki (2025)
JP-Fjy	FJY	Japan	JP	Fujiyoshida forest meteorology research site	35.4 5454	138.76 225	1043	Cfa	ENF	Ongoing	00-21	Mizoguchi et al. (2012)	Takanashi et al. (2025a)
JP-Fhk	FHK	Japan	JP	Fuji Hokuroku Flux Observation Site	35.4 4355 577	138.76 46931	1100	Cfa	DNF	Ongoing	06-23	Takahashi et al. (2015)	Takahashi et al. (2025)
JP-Hrt		Japan	JP	Hiratsuka Rice Paddy	35.3 6277 8	139.33 8056	6.98	Cfa	CRO	Completed	13	Komiya (2015)	Komiya (2025a)
JP-SMF	SMF	Japan	JP	Seto Mixed Forest Site	35.2 6152 8	137.07 875	212	Cfa	MF	Completed	02-16	Matsumoto et al. (2008)	Kotani and Ohta (2025)
JP-Nuf		Japan	JP	Nagoya University Forest	35.1 5241 667	136.97 18889	66	Cfa	DBF	Completed	00-01	Hiyama et al. (2005)	Awal and Ohta (2025a)
JP-Tdf		Japan	JP	Toyota Deciduous Forest	35.0 3588 889	137.18 57778	104	Cfa	DBF	Completed	02-04	Awal et al. (2010)	Awal and Ohta (2025b)
JP-Yms	YMS	Japan	JP	Yamashiro forest meteorology research site	34.7 9027 8	135.84 0939	220	Cfa	DBF	Ongoing	00-23	Komina mi et al. (2008)	Takanashi et al. (2025b)
JP-Nap		Japan	JP	Nunoike Agricultural Pond	34.7 7485	134.89 2442	40	Cfa	WAT	Completed	21-23	NA	Sakabe and Itoh (2025)
JP-Ako	AKO	Japan	JP	Akou green belt	34.7 3519 2	134.37 4798	10.5	Cfa	EBF	Completed	00-03	Kosugi et al. (2005)	Kosugi and Takanashi (2025)

JP-Sac	SAC	Japan	JP	Sakai City Office	34.5 7391 389	135.48 28889	17	Cfa	URB	Ongoing	08-23	Ueyama and Takano (2022)	Ueyama (2025d)
JP-Ozm	IZM	Japan	JP	Oizumi Urban Park	34.5 6346 9	135.53 3483	22	Cfa	URB	Completed	15-16	Ueyama and Ando (2016)	Ueyama (2025a)
JP-Om1	OM1	Japan	JP	B11 building in Osaka Metropolitan University	34.5 4717 7	135.50 2861	27	Cfa	URB	Ongoing	14-23	Ueyama and Ando (2016)	Ueyama (2025b)
JP-Om2	OM2	Japan	JP	Farm field in Osaka Metropolitan University	34.5 4245 2	135.50 8227	50	Cfa	GRA	Ongoing	22-23	NA	Ueyama (2025c)
JP-Hc3		Japan	JP	Hachihama Experimental Farm: Double Crop	34.5 3967 2	133.91 1731	-0.25	Cfa	CRO	Completed	05-09	Takimoto et al. (2010)	Takimoto and Iwata (2025b)
JP-Hc1		Japan	JP	Hachihama Experimental Farm	34.5 3789 167	133.92 67972	0	Cfa	CRO	Completed	96	Harazono et al. (1998)	Harazono (2025i)
JP-Hc2	HCH	Japan	JP	Hachihama Experimental Farm	34.5 3751 8	133.92 7545	-1	Cfa	CRO	Completed	99-08	Ohtaki (1984)	Takimoto and Iwata (2025a)
JP-Khw	KHW	Japan	JP	Kahoku Experiment watershed	33.1 3658	130.70 834	196	Cfa	ENF	Ongoing	00-03, 07-21	Shimizu et al. (2015)	Kitamura et al (2025)
JP-Ynf	YNF	Japan	JP	Yona-Field Tower Site	26.7 51	128.21 2667	213	Cfa	EBF	Ongoing	13-22	Matsumoto et al. (2023)	Matsumoto et al. (2025)
TH-Kog		Thailand	TH	Kog-Ma Watershed	18.8	98.9	1265	Af	EBF	Completed	05-13	Kume et al. (2007)	Kumagai and Takamura (2025a)
TH-Mae		Thailand	TH	Mae Moh plantation	18.3 8333 333	99.716 66667	380	Aw	DBF	Completed	05-16	Igarashi et al. (2015)	Kumagai and Takamura (2025b)

TH-Kms		Thailand	TH	Kamphaeng Saen Rice Paddy	14.09167	99.984167	4.74	Aw	CRO	Completed	14	Komiya (2015)	Komiya (2025b)
KH-Kmp		Cambodia	KH	Kampong Thom Lowland Dry Evergreen Forest	12.74457978	105.4785661	95	Am	EBF	Ongoing	11-14	Kabeya et al. (2021)	Shimizu et al. (2025a)
MY-LHP	LHP	Malaysia	MY	Lambir Hills National Park	4.201007	114.039079	140	Af	EBF	Completed	09-19	Takamura et al. (2023)	Kumagai et al. (2025)
ID-Pag		Indonesia	ID	Palangkaraya Undrained Forest	-2.323916667	113.9043917	22	Am	EBF	Ongoing	04-19	Hirano et al. (2024)	Hirano and Ohkubo (2025c)
ID-PaB		Indonesia	ID	Palangkaraya Drained Burnt forest	-2.340796	114.0379	14	Am	OSH	Completed	04-17	Ohkubo et al. (2021)	Hirano and Ohkubo (2025a)
ID-PaD	PDF	Indonesia	ID	Palangkaraya Drained forest	-2.346070697	114.036408	26	Am	EBF	Completed	01-17	Hirano et al. (2024)	Hirano and Ohkubo (2025b)

2.1 Data collections

130 We collected the micrometeorological measurement data from the site teams, which were identified using the web pages for AsiaFlux (<https://www.asiaflux.net/>, last access: 11 July 2024) and JapanFlux (<https://www.japanflux.org/>, last access: 11 July 2024). We also collected information on previous studies that reported micrometeorological measurements from domestic researcher connections and literature surveys. The collected data were from eddy covariance observations that were operated by the site teams affiliated with Japanese research institutes and universities. By this criterion, the dataset covers not only Japan but also other countries, such as Russia, China, Mongolia, Cambodia, Thailand, Malaysia, and Indonesia. Most of the sites were established for long-term monitoring of CO₂ fluxes, but intensive observations for about a week in the 1990s were also included in the dataset. Since the data format differed in each team, we reformatted the file to the FLUXNET format (<https://ameriflux.lbl.gov/data/aboutdata/data-variables/>, last access: 11 July 2024) after consultation with each site team. Generally, non-gap-filled data were provided by the site teams, but some teams provided gap-filled meteorological and flux data in addition to the non-gap-filled data. The JapanFlux2024 dataset differs from datasets such as FLUXNET2015 in that it

provided site principal investigators (PIs) with increased flexibility in data screening. When clear anomalies were identified, quality control procedures were applied by the management team in collaboration with the respective site PI.

The dataset consists of data from 83 sites with 683 site-years, of which 52 sites are located in Japan (Fig. 2; Table 1). The dataset includes 43 forest sites, 15 grassland sites, 5 wetland sites, 10 cropland sites, 3 lake and pond sites, and 4 sites in urban landscapes. Sites that suffered from various types of disturbance are also included: wind damage by typhoon (JP-Tmd, JP-Spp), fire (RU-NeB, ID-PaB), harvesting (RU-NeC, JP-Tef), thinning (JP-Fhk), insect outbreak (JP-Api), drainage (ID-Pag), and mowing (JP-NsC, JP-NsM, JP-Tgf, JP-Om2). The data records started in 1990 at a soybean cropland in Japan (Harazono et al., 1992), their number increased in the early 2000s, and peaked at 34 sites in 2008, 2014 and 2015 (Fig. 3). More recently, the number of data records gradually declined owing to site closure or the fact that the data have not been processed yet. The longest record was 24 years (JP-Tak and JP-Yms; both deciduous broadleaf forests) (Fig. 4). There are 26 sites with observation records of CO₂ flux for more than 10 years and 6 sites with those for more than 20 years (JP-Tef, JP-Tmk/JP-Tmd, JP-Api, JP-Fjy, JP-Tak, JP-Yms). Note that JP-Tmk and JP-Tmd represent a continuous observation series, although they are assigned different site IDs. At 12 sites, data records are available for less than one year. Data for CH₄ flux are available at six sites (JP-BBY, JP-SwL, JP-Nap, JP-Hrt, JP-Sac, JP-Om1).




Figure 2. Distributions of the sites that constitute the JapanFlux2024 database on a land cover map provided by the MOD12 product (version 6.1; Sulla-Menashe et al., 2019): a map of Asia region (a) and an enlarged map showing Japan (b).




Figure 3. Number of site data records in each year. Land cover types: evergreen broadleaf forest (EBF), evergreen needleleaf forest (ENF), deciduous broadleaf forest (DBF), deciduous needleleaf forest (DNF), mixed forest (MF), grassland (GRA), open shrubland (OSH), cropland (CRO), wetland (WET), urban (URB), lake (WAT), and barren sparse vegetation (BSV).




Fig. 4. The number of site records with different durations of data records. Sites affected by disturbance that changed the vegetation type during the observational period were classified according to the dominant land cover type: JP-Tef as DNF, JP-Tmd as DBF, and ID-PaB as OSH. Land cover type abbreviations are as in Fig. 3.

2.2 Gap-filling meteorological variables

As with the FLUXNET2015 dataset (Pastorello et al., 2020; Vuichard and Papale, 2015), the meteorological variables were filled using the European Center for Medium-Range Weather Forecasts Reanalysis v5 (ERA5) data (Hersbach et al., 2020). Instead of using ERA5, we used the gap-filled meteorology if the site teams had filled the gaps. If meteorological variables for multiple sensors or positions were available, these variables were prioritized and aggregated; if data were missing in the highest priority dataset, they were filled with values from the second-highest priority dataset, or, if that were also unavailable, based on the priority order. The gaps in the aggregated meteorological variables were then filled with ERA5 data because measured variables were less biased than ERA5, even when measured at different locations within a site. Air temperature, relative humidity, wind speed, downward shortwave radiation, downward longwave radiation, precipitation, and barometric pressure were filled using ERA5 after correcting biases at each site each year. Linear regression for meteorological variables (except precipitation) between observations and ERA5 was determined and then applied to correct site-specific biases in ERA5 to fill the data gaps. Water vapor pressure was calculated from the relative humidity, and the gaps in relative humidity were filled using the gap-filled water vapor pressure and air temperature, rather than directly filling the relative humidity. If all meteorological variables were missing in some years when constructing the linear regression, the bias was corrected using a regression for the entire multi-year data record. For precipitation (denoting rainfall plus snowfall), we determined the ratio of the annual precipitation between observations and ERA5 during the period when observed precipitation were available, and then filled hourly or half-hourly precipitation after multiplying the ratio to ERA5-based precipitation. If only the rainfall was measured, the correction ratio was determined using liquid precipitation which was defined as precipitation when relative humidity was below the critical relative humidity (RH_{cri} ; %): $RH_{cri} = 92.5 - 7.5T$, where T is the air temperature (Matsuo and Sasyo, 1981).

2.3 Gap-filling and flux partitioning

Gap-filling and flux partitioning were conducted based on REddyProc (version 1.3.2; Wutzler et al., 2018). First, the friction velocity (u^*) threshold was determined for the identified low-turbulence conditions during the nighttime using the moving-point method (Papale et al., 2006). The u^* threshold was determined from the temperature sensitivity of nighttime net ecosystem exchange (NEE) by seasonal clustering, an approach that is widely used in the FLUXNET community. In the moving point method, the u^* threshold was first determined for each of the four seasons, and the maximum value among them was used for the entire year. Thus, the determined u^* threshold was conservative (Papale et al., 2006). In this dataset, we determined the u^* threshold each year to consider its potential shift over the years, which is termed as a Variable u^* Threshold (vUT). The vUT differs slightly from the definition of the Variable u^* Threshold (VUT) in FLUXNET2015 (Pastorello et al., 2020) (Table 2), where VUT in FLUXNET2015 was determined by pooling data from each year along with data from the immediately preceding and following years (if available). The u^* threshold was determined with 100 bootstrap replicates, where reference (original data obtained without using a bootstrapped sample), the 5th, 50th, and 95th percentiles of the estimated u^* threshold were used for subsequent data filtering, gap-filling, and flux partitioning. Here, the nighttime was

205 defined as downward shortwave radiation $< 10 \text{ W m}^{-2}$, and was further confirmed using exact solar time at the site location. On the basis of the estimated u^* threshold, nighttime CO_2 fluxes and/or NEE were eliminated. This dataset does not include the estimation of NEE using the constant u^* threshold (CUT), nor the advanced uncertainty estimation provided with the _REF suffix, as implemented in the ONEFLUX pipeline (Pastorello et al., 2020). For urban sites, the threshold was generally not used for two reasons (e.g., Liu et al., 2012; Ueyama and Ando, 2016): 1) nighttime CO_2 fluxes were not expected to correlate with air temperature, making it difficult to evaluate the correct u^* threshold, and 2) the surface layer was often unstable even at night. Consequently, the u^* filtering was not applied for highly urbanized sites (JP-Sac and JP-Kgu).

Table 2 List of variable basenames not used in FLUXNET2015, along with their descriptions and related sites.

Basename	Description	Sites
pre-processing variables		
SW_IN_SLOPE_PI	Slope-normal incoming shortwave radiation	JP-Nkm
NETRAD_SLOPE_PI	Slope- normal net radiation	JP-Nkm
USTAR_QC_FP	Friction velocity qualified with footprint	JP-Ozm
H_QC_FP	Sensible heat flux qualified with footprint	JP-Ozm, JP-Khw
LE_QC_FP	Latent heat flux qualified with footprint	JP-Ozm, JP-Khw
FC_QC_FP	CO_2 flux qualified with footprint	JP-Ozm, JP-Khw, JP-Sac
NEE_QC_FP	NEE flux qualified with footprint	JP-Khw
post-processing variables		
TA_multiple	Air temperature by multiple sensors or positions	CN-In1, CN-In2, CN-In3, CN-In4, CN-In5, CN-In6, CN-In7, CN-In8, CN-Lsh, JP-BBY, JP-Fhk, JP-Fjy, JP-Hc1, JP-Ozm, JP-Khw, JP-KaP, JP-Km1, JP-Km2, JP-Kzw, JP-MBF, JP-MMF, JP-Nsb, JP-Nuf, JP-Spp, JP-Tgf, JP-Tak, JP-Tmk, JP-Tef, JP-Ynf, JP-Yrp, MN-Udg, MY-LHP, RU-Ege, RU-SkP, RU-Sk2, TH-Kog, TH-Mae, RU-USk

RH_multiple	Relative humidity by multiple sensors or positions	CN-In1, CN-In2, CN-In3, CN-In4, CN-In5, CN-In6, CN-In7, CN-In8, CN-Lsh, JP-BBY, JP-Fhk, JP-Fjy, JP-Hc1, JP-KaP, JP-Km1, JP-Km2, JP-Kzw, JP-MBF, JP-MMF, JP-Nsb, JP-Nuf, JP-Spp, JP-Tgf, JP-Tak, JP-Tmk, JP-Tef, JP-Ynf, JP-Yrp, MN-Udg, MY-LHP, RU-Ege, RU-SkP, RU-Sk2, TH-Mae, RU-USk
SW_IN_multiple	Incoming shortwave radiation by multiple sensors or positions	JP-Fhk, JP-Spp, JP-Tgf, JP-Tmk, JP-Tef, JP-Ynf
P_multiple	Precipitation by multiple sensors or positions	JP-BBY, JP-Khw
WS_IN_multiple	Wind speed by multiple sensors or positions	CN-In1, CN-In2, CN-In3, CN-In4, CN-In5, CN-In6, CN-In7, CN-In8, CN-HaM, JP-Fhk, JP-Hc1, JP-KaP, JP-Km1, JP-Km2, JP-Kzw, JP-MBF, JP-MMF, JP-Nsb, JP-Spp, JP-SMF, JP-Tgf, JP-Tef, JP-Yms, JP-Ynf, JP-Yrp, MN-Udg, TH-Kog, TH-Mae, RU-USk
G_multiple	Ground heat flux by multiple sensors or positions	JP-Sac, JP-Spp, JP-Ynf, MN-Udg
NETRAD_F_MDS	Net radiation filled with MDS	JP-Tef
PPFD_IN_F_MDS	PPFD filled with MDS	CN-Lsh, CN-HaM, JP-Km2, JP-MBF, JP-MMF, JP-Nkm, JP-Tgf, RU-SkP, TH-Kog
NEE_vUT	Gap-filled NEE with the variable u^* threshold	ALL sites
RECO_NT_vUT	RECO with the variable u^* threshold based on the nighttime approach	ALL sites
GPP_NT_vUT	GPP with the variable u^* threshold based on the nighttime approach	ALL sites
RECO_DT_vUT	RECO with the variable u^* threshold based on the daytime approach	ALL sites
GPP_DT_vUT	GPP with the variable u^* threshold based on the daytime approach	ALL sites

215 Gaps in sensible heat flux (H), latent heat flux (LE), and NEE were filled using marginal distribution sampling (MDS) based on REdDyProc. In MDS, a look-up-table (LUT) with air temperature, downward shortwave radiation, and vapor pressure

deficit (VPD) was created for a 7-day window. When data gaps could not be filled with this window, they were filled in the following order: (1) LUT was applied with a 14-day window, (2) the mean diurnal variation method (Falge et al., 2001) was applied with a 1- or 2-day window, and (3) LUT was applied with a 21-day window, which was increased with a 7-day step until 70-day window if not enough data points were available. NEE were filled using MDS with the four different u^* thresholds (reference, 5th, 50th, and 90th percentiles values), whereas H and LE were filled without applying the u^* threshold. In addition to the fluxes, net radiation, soil temperature, ground heat flux, and photosynthetically photon flux density (PPFD) were also filled using MDS. In the data collected from the site teams, energy imbalance correction (Twine et al., 2000) was not applied for H and LE at any sites; thus, the gap-filled H and LE were not corrected for the energy balance closure.

Using REdDyProc, NEE was partitioned into gross primary productivity (GPP) and ecosystem respiration (RECO) using two methods: nighttime flux partitioning and daytime flux partitioning. In the nighttime partitioning method, nighttime NEE was parameterized on the basis of the temperature response function (Lloyd and Taylor, 1994) with a 7-day window, and then this function was used to calculate daytime and nighttime RECO. GPP was determined by subtracting RECO from NEE. In the daytime partitioning method (Lasslop et al., 2010), the common rectangular hyperbolic light-response curve was determined with a 4-day window, where the function accounted for the VPD effect on the initial slope of the light-response curve and the temperature effect of respiration. GPP and RECO using the daytime partitioning method were calculated based on a fitted model that combines a light-response curve and a temperature-dependent respiration model; thus, the daytime method did not directly add up observed NEE (Wutzler et al., 2018). Using the two methods, fluxes were partitioned for NEE with different u^* thresholds.

2.4 Site-specific considerations

For the sites with heterogeneous land surfaces—JP-Khw and JP-Ozm—the dominant land surface fluxes were extracted using wind sectors. JP-Khw is an evergreen needleleaf plantation forest consisting of *Cryptomeria japonica* (sugi) and *Chamaecyparis obtusa* (hinoki), but evergreen or deciduous broadleaf trees grow in gaps in some wind sectors. The H, LE, and CO₂ fluxes for sugi, which occupies the dominant wind sector area (the right-bank side), were extracted as quality control with footprint, "_QC_FP" (Table 2). To extract these flux data, daytime fluxes for a wind sector on the right bank were selected, but nighttime fluxes for all wind sectors were used to increase data availability because there were no clear differences in nighttime fluxes among wind sectors. Gap-filling and flux partitioning were done only for the extracted data. JP-Ozm is located at the edge of an urban park; thus, measured flux representing this park (Ueyama and Ando, 2016) were selected and designated "_QC_FP" in addition to the variables for measured fluxes representing both sectors of the urban park and other land covers. Gap-filling and flux partitioning were done only for the extracted data, which represented the urban park. These extracted flux data ("_QC_FP") were included in the ALLVARS files (described in section 2.5) in addition to measured fluxes for all sectors, and the gap-filled extracted fluxes were included in the COREVARS files (described in section 2.5).

Flux partitioning and gap-filling for JP-Nkm, located on a complex mountainous terrain, were conducted using slope-normal shortwave radiation instead of downward shortwave radiation. Horizontally observed incident shortwave radiation was converted to radiation normal to the slope on the basis of the tilt and azimuth angles of the slope and the solar altitude and azimuth angles (Hammerle et al., 2007; Nie et al., 1992) as follows. Horizontally observed incident shortwave radiation was partitioned into direct and diffuse components using observed diffuse fraction (BF5, Delta-T Devices, UK), and the direct component was converted to that normal to the slope surface. The diffuse component was assumed to be isotropic. The total incident shortwave radiation normal to the slope surface was calculated as the sum of the direct component converted as above and the original diffuse component. When diffuse fraction was not observed, it was estimated from the relationship between the diffuse fraction and cloudiness; the latter was defined as the ratio of observed incident shortwave radiation to extraterrestrial radiation (Wang et al., 2018). The slope-normal shortwave radiation was included as a variable, SW_IN_SLOPE_PI_1_1_1, in ALLVARS.

For tropical ecosystems (TH-Kms, TH-Kog, TH-Mae, ML-LHP, ID-PaB), nighttime-based flux partitioning failed because little seasonality in temperature hampered the determination of a significant relationship between nighttime CO₂ flux and temperature. For these sites, only daytime partitioning was provided in the dataset. In a subtropical forest (KH-Kmp), the determination of the u^* threshold failed; thus, the u^* threshold was estimated using gap-filled u^* by the site team instead of measured u^* with data gaps. Since the data quality of u^* for KH-Kmp seemed reasonable, we were unable to find out why REdDyProc failed to determine the u^* thresholds with measured u^* in KH-Kmp. The u^* threshold for ID-Pag and ID-PaD could not be determined for several years; so, constant u^* thresholds across these years were determined with REdDyProc and applied for the subsequent data processing.

Low availability of nighttime data due to the limited fetch in JP-Ako (Kosugi et al., 2005) hampered determination of the u^* threshold, gap-filling flux for CO₂ flux, and flux partitioning with REdDyProc. Consequently, no aggregated fluxes longer than half-hourly data for CO₂ flux, GPP, and RECO were provided in the dataset.

Fluxes were not partitioned for lakes and a pond (JP-SwL, JP-KaL, JP-Nap) and an urban center (JP-Sac). For the lakes and pond, gap-filling H, LE, and CO₂ flux was based on MDS. For JP-Sac, gap-filling for H and LE was also based on MDS, but MDS was not applied to CO₂ flux because it was controlled by traffic volume and air temperature (Ueyama and Takano, 2022). Gap-filling for CO₂ flux at JP-Sac was conducted by the site team on the basis of random forest regression (Ueyama and Takano, 2022), and was included as FCO2_F_PI in COREVARS and ALLVARS. The u^* threshold was not applied for JP-Nap, because the moving point method (Papale et al., 2006) developed for terrestrial ecosystems was not applicable to the pond.

In this dataset, CH₄ fluxes were not gap-filled, because (1) consistent gap-filling was not possible because of missing important variables, such as water table depth, and (2) inconsistent processes control CH₄ emissions on different land surfaces, such as a rice paddy (JP-Hrt), bog (JP-BBY; Ueyama et al., 2020c, 2022b), lake (JP-SwL; Iwata et al., 2018), pond (JP-Nap), and urban landscapes (JP-Sac, JP-Om1; Takano and Ueyama, 2021). If the gap-filled CH₄ fluxes were provided by the site team (i.e., JP-BBY), the data were included as FCH4_F_PI in COREVARS; otherwise, non-gap-filled data were included in ALLVARS.

290

2.5 Data format

The dataset was prepared in a format partially compatible with FLUXNET format, although the content and split of variables between ALLVARS and COREVARS were slightly different to FLUXNET2015 (Pastorello et al., 2020) (Table 2), which consists of files separated by sites, temporal aggregation (i.e., half-hourly/hourly, daily, weekly, monthly, and annual), and data product, i.e., ALLVARS and COREVARS, as described later. The separated files for ALLVARS and COREVARS were combined into two zip files for each site.

295

The following file naming rules (Pastorello et al., 2020) were followed.

300 [SITE_ID]_JapanFlux2024_[DATA_PRODUCT]_[RESOLUTION]_[FIRST_YEAR]-[LAST_YEAR]_[SITE_VERSION]-
[CODE_VERSION].csv

[SITE_ID] is the site ID, the CC-SSS format: CC is a two-letter country code, and SSS is the three-character site code. [Data_PRODUCT] represents the data types: ALLVARS, COREVARS, AUXMETEO, AUXNEE, or ERA5. COREVARS is the data type representing selected data variables, including basic micrometeorological data and fluxes, and quality information flags. ALLVARS is a data file representing all variables of data products, including variables listed in COREVARS, original data before the processing pipeline, and internal variables. AUXMETEO includes auxiliary variables related to the meteorological downscaling of ERA5. ERA5 includes the meteorological data from ERA5 for 1990–2024. [RESOLUTION] is the temporal resolution of the data products: HH (half-hourly time step), HR (hourly time step), DD (daily time step), WW (weekly time step), MM (monthly time step), and YY (annual time step). [FIRST_YEAR] is the first year in the file, and [LAST_YEAR] is the last year in the file. The first and last years are based on the years in which the micrometeorological measurements are conducted, except for ERA5, where the first year is 1990 and the last year is 2024 for all sites. [SITE_VERSION] is the version of the original dataset, and [CODE_VERSION] is the code of the data processing pipeline used to process the dataset.

315

The COREVARS file included variables for basic meteorology and turbulent fluxes. The gap-filled meteorological variables of air temperature, incoming shortwave radiation, incoming longwave radiation, relative humidity, VPD, atmospheric pressure, precipitation, wind speed, net radiation, ground heat flux, soil temperature, PPFD, CO₂ concentration, and soil water content were included. If the original data provided by a site team included wind direction, outgoing shortwave radiation, outgoing longwave radiation, outgoing PPFD, u^* , and shortwave radiation, incoming, potential (top of atmosphere), non-gap-filled data for these variables were included. Gap-filled soil temperature and soil water content were measured at the shallowest depth, while CO₂ concentration was gap-filled for the highest altitude. A quality information flag was assigned for gap-filled variables, where 0 is the original data, 1 is a gap-filled value of the most reliable quality (calculated using a 14-day window), 2 is a gap-filled value of the medium quality (calculated using a 14- to 56-day window), and 3 is the gap-filled value of the least reliable quality (calculated using a window longer than 56 days) (Wutzler et al., 2018). If gap-filled CH₄ flux data were provided by the site team (i.e., at JP-BBY), they were included in COREVARS. The COREVARS file was provided with five temporal resolutions (half-hourly/hourly, daily, weekly, monthly, and annual aggregations).

The ALLVARS file includes original unprocessed data, internal variables (aggregated meteorological variables measured at different locations or with different sensors), and meteorological data from ERA5, in addition to processed variables included in COREVARS. The ALLVARS file is provided with the five temporal resolutions listed above.

For NEE, GPP, and RECO, the unit was $\mu\text{mol m}^{-2} \text{s}^{-1}$ for half-hourly and hourly timescales, $\text{g C m}^{-2} \text{d}^{-1}$ for the daily, weekly, and monthly timescales, and $\text{g C m}^{-2} \text{yr}^{-1}$ for the annual timescale. For CH₄ flux, the unit for the half-hourly and hourly timescales was $\text{nmol m}^{-2} \text{s}^{-1}$, whereas the units for the other timescales were as the same as those for CO₂ fluxes. The units of precipitation were mm for the half-hourly and hourly timescales, mm d^{-1} for the daily, weekly, and monthly timescales, and mm yr^{-1} for the annual timescale. The units of other variables followed FLUXNET format (<https://ameriflux.lbl.gov/data/aboutdata/data-variables/>), which did not change with the timescales.

The ERA5 file contains the data for air temperature (TA_ERA5; °C), relative humidity (RH_ERA5; %), VPD (VPD_ERA5; hPa), vapor pressure (e_ERA5; hPa), saturation vapor pressure (e_sat_ERA5; hPa), wind speed (WS_ERA5; m s^{-1}), atmospheric pressure (PA_ERA5; kPa), incoming shortwave radiation (SW_ERA5; W m^{-2}), incoming longwave radiation (LW_ERA5; W m^{-2}), and precipitation (P_ERA5; mm). The ERA5 file is provided with the five temporal resolutions listed above. The variables in the ERA5 file were not corrected for the bias in comparison to the site data.

Two auxiliary files—for meteorology and u^* -threshold—are provided. The AUXMETEO file includes the following statistics for downscaling ERA5 to the site scale: the linear slope between the measured data and ERA5 (ERA_SLOPE), intercept (ERA_INTERCEPT), root mean square error (ERA_RMSE), and correlation coefficient (ERA_CORRELATION). These statistics are included for each year and for all years when measurements were conducted. The TIMESTAMP column in the

350 AUXMETEO file represents the year for the statistics, where -9999 represents the statistics for the entire year. The AUXNEE file includes the u^* threshold in each year, with the reference threshold and the 5th, 50th, and 95th percentiles of the estimated u^* threshold.

The dataset also includes the BADM files, which are used in the FLUXNET community. Six BADM files are provided: (1) 355 general information, (2) instrument, (3) instrument operations, (4) vegetation cover, (5) soil, and (6) disturbance and management.

3 Database summary

3.1 CO₂ flux

360 Based on the dataset constructed, mean seasonalities in NEE, GPP, and RECO were as expected from the biomes and mean climatology (Figs. 5, 6). In northern boreal forests in Siberia (RU-Tur, RU-NeF, RU-SkP, RU-Sk2), the magnitude of the flux was generally low, and growing seasons when GPP was not negligible were short. In the southern Eurasian boreal forests in Siberia and Mongolia (RU-Ege, MN-Udg, MN-Skt), the magnitudes of CO₂ fluxes were greater than those in the above northern boreal forests. Inland grasslands in Mongolia (MN-Nkh, MN-Hst, MN-Kbu) had smaller CO₂ flux magnitudes than 365 the nearby forests (MN-Udg, MN-Skt). For temperate forest and grassland sites, the dataset showed known seasonality with spring onset, summer peak, and autumn senescence with low fluxes in winter. Among forest sites, seasonal variations became smaller in the subtropics (JP-Ynf), and clear seasonality disappeared in the tropics (KH-Kmp, MY-LHP, ID, Pag, ID-PaD, ID-PaB) as the climate became warmer. Among rice paddies, single-cropping sites had a single peak (JP-Mse, JP-Hc2), but a double cropping site had two peaks (JP-Hc3) in GPP, RECO, and NEE (Fig. 6). For lakes (JP-SwL, JP-KaL), a pond (JP-Nap), 370 and an urban center (JP-Sac), CO₂ fluxes showed smaller seasonality than those at vegetation surfaces.

Some data for CO₂ fluxes raise suspicions. First, markedly negative NEE values in harsh winters were estimated for MN-Skt and MN-Kbu (Figs. 5, 6), which could be caused by an artifact known for the open path sensor (Burba et al., 2008). The artificially negative NEE caused a considerable positive GPP in winter. Data users should be cautious about the data for MN- 375 Skt and MN-Kbu. Second, the daytime partitioning method extrapolated the relationship obtained during the growing season to winters when NEE was not measured. The result was erroneous estimation of GPP and RECO (e.g., JP-Nkm in Fig. 5). Using the nighttime approach, GPP and RECO were not estimated for the period when NEE was not measured. Despite these suspicious data, the fluxes partitioned using the nighttime and daytime methods were generally consistent across the sites.

380 The spatial variabilities in annual NEE, GPP, and RECO were also consistent with earlier reports for Asian ecosystems (Fig. 7; Table 3). In Asia, the spatial variabilities in GPP and RECO are explained mostly by mean annual air temperature (Hirata et al., 2008; Kato and Tang, 2008; Saigusa et al., 2013; Yu et al., 2013). Except in disturbed forests and croplands, GPP and

RECO increased linearly with mean annual air temperature (Fig. 7). Correlations of GPP and RECO with the annual sum of precipitation were lower than with mean annual air temperature. No clear correlation was found between annual NEE and mean annual air temperature or annual sum of precipitation, but the maximum CO₂ sink (i.e., negative NEE) with each temperature range appeared to be increased by temperature up to the annual mean temperature range until approximately 10°C (Fig. 7a). Except for disturbed forests and urban sites, most ecosystems were estimated to be a CO₂ sink of up to 1.0 kg C m⁻² yr⁻¹.




Figure 5. Mean seasonality of GPP, RECO, and NEE across forest sites. GPP and RECO were partitioned using the daytime method (DT, solid lines) or the nighttime method (NT, dashed lines). The seasonality is shown when NEE was measured, and

those for GPP and RECO are shown when the partitioning was successful. The seasonality is the ensemble mean of the daily fluxes with for each day of year for all years. The sites are ordered according to latitude from high to low. The mean seasonality is shown for sites having the data at least one growing season.





Figure 6. Mean seasonality of GPP, RECO, and NEE across sites other than forests. Designations are as in Fig. 5.



400

Fig. 7. Relationships of annual NEE (a, b), GPP (c, d), and RECO (e, f) to mean climate of annual mean air temperature (a, c, e) and annual sum of precipitation (b, d, f). GPP and RECO were estimated using the daytime method. The stars represent fluxes obtained at disturbed forests, where a disturbed forest was defined as a forest that experienced disturbance within the last 10 years. The annual fluxes were calculated based on the sum of mean seasonality shown in Figs. 5 and 6; missing measurements during the winter in high-latitudes were gap-filled as zero. Since sites of JP-Spp, JP-Tmd, and JP-Tef experienced significant disturbance (windthrow or clearcut) during the measurement period, data obtained within 10 years after a disturbance were classified as disturbed forests (DIS). The lines represent linear regressions with shading showing the confidence intervals ($p < 0.05$), that was determined excluding the DIS data. The annual CO_2 flux for JP-Sac ($5.8 \text{ kg C m}^{-2} \text{ yr}^{-1}$) is not shown due to the totally different carbon budget in the urban center compared to those in ecosystems. The values are shown in Table 3. Land cover type abbreviations are in Fig. 3.

410

Table 3. Summary of mean annual air temperature (TA), annual sum of precipitation (PREC), mean annual downward shortwave radiation (Rsd), mean annual carbon fluxes (NEE, GPP, RECO), and mean annual latent heat flux (LE), mean annual sensible heat flux (H), evapotranspiration (ET), and land cover. The statistics were calculated for observation years; for disturbed sites, the data were considered separately for the periods before, during, and after disturbance. Disturbed ecosystems were defined as those that experienced disturbance within the last 10 years. GPP, RECO, NEE, LE, and ET for boreal forests in Russia that lacked winter measurements (RU-Tur, RU-NeB, RU-NeC, RU-NeF, RU-SkP, RU-Ege, RU-Sk2) were considered zero. GPP, RECO, and NEE at MN-Skt and MN-Kbu were also considered zero during winter, when daily mean air temperature was below -5 °C (indicated by asterisks in the table), to mitigate the influence of the negative values of CO₂ fluxes caused by an artifact associated with an open-path sensor. The N/A values were listed because missing observations, even after gap-filled fluxes, prevented the calculation of annual fluxes or because the standard flux partitioning was not available for pond, lakes, and urban landscapes.

Site ID	disturbance	Land cover	TA	PREC	Rsd	NEE	GPP	RECO	LE	H	ET
			°C	mm yr ⁻¹	W m ⁻²	g C m ⁻² yr ⁻¹	g C m ⁻² yr ⁻¹	g C m ⁻² yr ⁻¹	W m ⁻²	W m ⁻²	mm yr ⁻¹
RU-Tur		DNF	-7.0	264	93	-83	180	76	N/A	N/A	N/A
RU-NeB	fire	GRA	-12.3	244	117	37	57	85	N/A	N/A	N/A
RU-NeF		OSH	-8.6	175	117	-166	653	455	10	N/A	132
RU-NeC	clearcut	DIS	-10.9	224	112	12	147	170	13	N/A	162
RU-SkP		DNF	-5.7	238	118	-139	523	375	21	N/A	265
RU-Sk2		ENF	-8.7	328	117	-194	522	283	16	N/A	203
RU-USk		GRA	-11.6	289	110	N/A	N/A	N/A	15	N/A	193
RU-Ege		DNF	-6.3	248	122	-225	715	466	20	N/A	259
MN-Skt		DNF	-1.7	279	169	-722 *	1058 *	324 *	15	45	189
MN-Udg		DNF	-0.6	303	157	-431	617	413	19	29	246
MN-Nkh		GRA	-1.9	163	171	-83	603	516	20	25	253
MN-Hst		GRA	1.6	197	181	76	541	468	18	27	228
MN-Kbu		GRA	0.7	204	183	-433	647 *	248 *	9 *	30	116
CN-Lsh		DNF	4.4	443	144	-60	1606	1255	25	36	324
JP-Sb1		WET	6.0	794	145	-288	1098	638	N/A	N/A	N/A
JP-Sb2		WET	5.5	840	140	-313	1476	935	N/A	N/A	N/A

JP-Tef	before	DIS	5.9	926	125	192	882	898	18	10	232
	clearcut	GRA	5.9	926	125	-118	1363	1114	27	13	343
	after	MIX	5.9	926	125	-26	1238	1034	21	17	268
JP-MBF		DBF	3.9	1373	134	-442	1233	862	37	20	472
JP-MMF		MF	5.4	1092	134	-689	1537	861	43	21	537
JP-BBY		WET	7.2	953	143	-118	785	610	41	12	524
JP-Spp	before	DIS	7.4	1215	145	235	1366	1554	35	16	444
	windthrow	DBF	7.4	1215	145	-42	1557	1432	40	15	500
JP-Tmk		DNF	6.6	1092	133	-270	1727	1401	45	31	568
JP-Tmd	windthrow	GRA	7.0	1738	139	421	1176	1249	N/A	N/A	N/A
	after	DBF	7.0	1738	139	89	1225	1147	34	21	427
JP-Toc		DBF	7.6	1342	137	-556	1729	1189	33	30	418
JP-Tom		DBF	6.9	1173	128	-249	916	642	N/A	40	N/A
JP-Srk		DBF	8.1	2669	129	-847	1408	600	41	-1	509
JP-Api		DBF	6.3	1509	150	-375	1307	958	18	14	235
CN-HaM		GRA	-1.1	97	200	-77	862	618	31	23	389
JP-NsM		GRA	12.2	1658	150	-251	1989	1779	55	6	704
JP-NsC		GRA	12.2	1658	150	-376	2098	1880	53	7	674
JP-Kzw		DBF	7.0	1524	165	-155	919	709	15	32	187
JP-Tkb		ENF	13.3	1514	159	-599	2490	1781	45	-7	579
JP-Tak		DBF	6.8	2483	146	-342	1024	597	11	26	135
JP-Ta2		ENF	9.8	1760	148	-695	1990	1247	43	16	546
JP-Tgf		GRA	14.3	1141	153	-291	2307	1851	53	19	681
JP-KaP		CRO	14.9	561	155	-774	1127	584	70	7	894
JP-Mse		CRO	13.9	1407	154	-197	1004	763	67	6	858
JP-SwL		WAT	11.8	1499	178	-287	N/A	N/A	80	18	1021
JP-KaL		WAT	16.0	1575	163	-826	N/A	N/A	59	21	759
JP-Kwg		DBF	15.2	1492	151	-214	1631	1393	N/A	18	N/A
JP-Shn		MF	12.3	1713	167	-242	1612	1211	53	43	675
JP-Nkm		ENF	0.5	2544	162	-381	1442	725	36	-3	445

JP-Fmt		MF	15.0	1611	158	35	2720	1652	71	35	913
JP-Kgu	urbanization	URB	16.5	1400	149	N/A	N/A	N/A	27	41	344
JP-Fjy		ENF	9.9	1989	165	-404	1772	1270	39	20	501
JP-Fhk	before	DIS	9.6	1846	168	-74	1945	1873	43	43	554
	thinning	DNF	9.6	1846	168	-433	1914	1619	40	40	510
JP-SMF		MF	14.8	1543	165	-142	1587	1059	51	23	658
JP-Nuf		DBF	15.4	1465	156	-327	1590	1139	22	16	277
JP-Tdf		DBF	14.8	2039	155	-601	1559	840	45	19	584
JP-Yms		DBF	15.0	1384	159	-223	1644	1400	63	30	805
JP-Nap		WAT	16.5	1083	176	-48	N/A	N/A	60	9	773
JP-Ako		EBF	15.3	739	169	N/A	N/A	N/A	27	47	347
JP-Sac	urbanization	URB	16.4	1594	159	5807	N/A	N/A	28	43	354
JP-Ozm	urbanization	URB	16.5	1828	150	793	1485	2353	52	23	673
JP-Om1	urbanization	URB	17.5	1202	165	1032	515	1622	23	39	294
JP-Om2	mowing	GRA	16.9	1466	166	430	2634	2937	71	9	908
JP-Hc3		CRO	15.8	1136	175	-663	1265	625	51	10	659
JP-Hc2		CRO	15.7	1141	161	-132	890	697	60	10	772
JP-Khw		ENF	15.2	2294	158	-906	2359	1689	81	20	1044
JP-Ynf		EBF	20.9	2611	159	-374	2200	1808	65	5	834
TH-Kog		EBF	19.9	2004	183	-1301	2078	1178	73	23	935
TH-Mae		DBF	25.3	1333	205	-579	2215	1783	69	36	888
KH-Kmp		EBF	26.9	1786	206	-72	3842	3044	106	19	1368
MY-LHP		EBF	26.2	2752	184	-989	3431	2552	89	28	1157
ID-Pag		EBF	26.1	2639	200	-183	3840	3486	111	29	1430
ID-PaB	fire	OSH	26.4	2642	197	-110	1746	1450	90	27	1164
ID-PaD		EBF	26.2	2543	197	179	3100	2997	94	29	1215

In the developed dataset, annual CO₂ fluxes tended to differ by land cover type (Fig. 8). Forest ecosystems included in the datasets had, on average, similar CO₂ sinks. Among the forest ecosystems, the mean CO₂ sink tended to be highest in ENF. GPP and RECO in temperate managed grasslands were higher than those in natural grasslands in Mongolia and Russia. The annual CO₂ sink also tended to be greater in managed grasslands compared to natural grasslands, except for a frequently mowed site (JP-Om2), which exhibited net annual CO₂ emissions. Disturbed forests, on average, acted as a small CO₂ source. CO₂ emissions in urban centers (JP-Sac; 5.8 kg C m⁻² yr⁻¹; not included in Fig. 8a) were considerably higher than those from natural or agricultural ecosystems. The annual GPP was highest in EBF among forest ecosystems, followed by ENF, DBF, and DNF. RECO was highest in EBF, whereas those in ENF, DBF, and DNF were similar to each other. Annual GPP and RECO varied greatly among grasslands because they included inland dry grasslands and Japan's weedy grasslands (Fig. 8b, c).




Figure 8. Boxplots for annual NEE, GPP, and RECO by land cover type. GPP and RECO were estimated using the daytime method. Fluxes at urban sites are not shown because the range of urban CO₂ emissions was totally different from those for vegetation or lakes. Since flux partitioning was not conducted for lakes and ponds, partitioned fluxes for these water surfaces were not shown. Land cover type abbreviations are in Fig. 3, although the grassland category was separated into natural

grassland (GRAn) and managed grassland (GRAM). The definition of DIS was the same as in Fig. 7, where all data from RU-NeC, RU-NeB, and ID-PaB are also classified as DIS. The box represents the interquartile range (25th to 75th percentiles), the whiskers represent the maximum and minimum values, excluding outliers shown by circles, and the orange bar represents the median value.

3.2 Energy fluxes

Mean annual energy fluxes represented in the dataset were explained better by air temperature than precipitation (Fig. 9; Table 3). The mean annual LE increased with the mean annual air temperature; their strong linear correlation could be explained by a close coupling between transpiration and photosynthesis (Medlyn et al., 2011), where spatial variations in annual GPP were strongly correlated with annual air temperature (Fig. 7c). Evaporation could also be enhanced under high air temperature and resulting high VPD conditions (Zhang et al., 2015). The dataset included mostly ecosystems around the Pacific Ocean, which were especially densely distributed in Japan, whereas water-limited inland ecosystems were scarce. Consequently, the correlation between LE and precipitation was weaker than those reported in a literature survey for Asia (Kang and Cho, 2021). Under similar climate conditions, LE was lower and H was higher in urban landscapes in comparison with vegetation surfaces, in agreement with a previous report (Ueyama et al., 2021). Mean annual H did not change with air temperature or precipitation, possibly be caused by missing high-latitude observations owing to missing winter data (e.g., RU-Tur, RU-SkP, Ru-Ege) (Fig. 5). Negative H values in high-latitude ecosystems were observed owing to decreased available energy associated with snow albedo (Nakai et al., 2013; Ueyama et al., 2020b).




Figure 9. Relationships of annual energy fluxes of latent heat flux (LE) (a, b) and sensible heat flux (H) (c, d) to mean climate of annual mean air temperature (a, c) and annual sum of precipitation (b, d). The classification of the disturbed forest (DIS) is as in Fig. 8. Annual H were only calculated for the case where there were no missing data in the mean seasonality, whereas the missing LE data during the winter were considered to be zero for boreal forests in Russia. The lines represent linear regressions, with shading showing the confidence intervals ($p < 0.05$), that was determined excluding the data from DIS, urban areas (URB), and lakes and ponds (WAT). The values are shown in Table 3. Land cover type abbreviations are in Fig. 3.

4 Data availability and data use guidelines

The dataset associated with this publication can be found at the ADS website (<https://ads.nipr.ac.jp/japan-flux2024/>), where individual site data have their own DOIs. All data are available under the CC BY 4.0 copyright policy with appropriate citations of this paper. We suggest that researchers planning to use this dataset as a core dataset for their analysis contact and collaborate with database developers and relevant site teams. As in the data policy of FLUXNET2015, in case of a synthesis using both CC BY 4.0 and other private data, all data should be treated as Tier Two of the FLUXNET data policy (data producers must have opportunities to collaborate and consult with data users).

5 Conclusions

The JapanFlux2024 dataset is the first public dataset that includes as much data as possible, both old and new, as an activity of JapanFlux. The dataset is consistent with previous synthesis studies in Asia in terms of seasonalities in CO_2 and energy

fluxes across Japan and East Asia, but substantially increased the number of the data, 83 sites with 683 site-years from 1990 to 2023. The dataset will facilitate important studies in East Asia including Japan, such as those on land-atmosphere interactions, improvement of process models, and upscaling fluxes using machine learning. Since the dataset is processed in
480 line with reference to selected procedures from the FLUXNET standard dataset, the JapanFlux2024 dataset will bridge collaborations between researchers from Asia and FLUXNET.

Author contributions

The JapanFlux2024 dataset was conceptualized by MU. The standardized dataset was prepared by MU and YT, and the
485 metadata was compiled by HY and TH in collaboration with the data contributors. The data distribution website was developed by a team led by HY. KI contributed to the editing of the manuscript. The remaining co-authors contributed eddy covariance data to the dataset and/or participated in editing the manuscript.

Competing interests

490 The authors declare that they have no conflict of interest.

Acknowledgements

The development of the database was supported by the digital biosphere project under KAKENHI (21H05316, to T. Kumagai was supported), PAWCs project under KAKENHI (19H05668), JSPS A3 Foresight Program Number (JPJSA3F20220002),
495 and the Arctic Challenge for Sustainability II (ArCS II; JPMXD1420318865). The CH-Lsh data were provided by Dr. N. Saigusa of the National Institute of Environmental Studies and H. Wang of the Chinese Academy of Sciences. Observations at JP-Tmk were supported by Dr. R. Hirata of the National Institute of Environmental Studies. Observations of JP-Kgu were supported by Dr. M. Kanda of the Institute of Science Tokyo. Observations of JP-Tdf were supported by Professor emeritus S. Hattori of Nagoya University. Observations of JP-Ynf were supported by Dr. S. Tanigushi of the University of the Ryukyus.
500 M. Ueyama was supported by KAKENHI (18H03362, 24K03065). S. Ishida was supported by KAKENHI (25450201). K. Ichii, H. Yazawa, and M. Tanaka were supported by KAKENHI (22H05711, 22H05004, 24H01504) and Environment Research and Technology Development Fund (JPMEERF24S12207). H. Iwata was supported by KAKENHI (17H05039, 21H02315, 23K21248, 23KK0194). M. Sugita was supported by KAKENHI (15K01159, 20H01384, 23K20125). T. Shimizu was supported by KAKENHI (20H0309). Multidisciplinary observations at Takayama sites (JP-Tak and JP-Ta2) have been
505 supported jointly by H. Muraoka (KAKENHI 21H05316, 21H05312, 19H03301), T.M. Saitoh (KAKENHI 18780113, 21241009, 22248017, 23710005, 24241008, 26241005, 26292092, 15H04512, 20H03041, 20K06144, 21H02245, 21H05316, 23K11395, 24K01818, 24K00986, the Environment Research and Technology Development Fund JPMEERF20232M01 of the Environmental Restoration and Conservation Agency provided by the Ministry of the Environment of Japan, the Global Environment Research Coordination System from the Ministry of the Environment, Japan MAFF2254), H. Kondo, S.
510 Murayama, S. Ishidoya and T. Maeda (KAKENHI 24241008, 24310017, 15H02814, 18H03365, 19H01975, 22H00564,

22H05006, Global Environment Research Coordination System from the Ministry of the Environment, Japan MAFF0751, MAFF1251, MAFF2254, the Global Environment Research Fund of the Ministry of the Environment, Japan S-1: Integrated Study for Terrestrial Carbon Management of Asia in the 21st Century Based on Scientific Advancement). JP-Spp, JP-API, JP-Fjy, JP-Yms, and JP-Khw were supported by KAKEN (16K07789), Research revolution 2002: Global Warming Initiatives (FY2002-2006) by the Ministry of Education, Culture, Sports, Science and Technology of Japan, Commissioned project study from the Ministry of Agriculture, Forestry and Fisheries (JPJ005317), Environment Research and Technology Development Fund (S-1), and Research Coordination System (MAFF0751, 1251, 2254) from the Ministry of the Environment of Japan, and Research grants (#199903, #200303, #201802) from the Forestry and Forest Products Research Institute. JP-Tom was supported by KAKENHI (11213204, 14656059, 16208014, 2331001513, 2529207903) and by the Ministry of the Environment (0708BD437, D-0909) to T. Hiura. JP-Mse was supported by the Global Environmental Research Fund (S-1) of the Ministry of Environment of Japan, a research project entitled “Development of technologies for mitigation and adaptation to climate change in agriculture, forestry and fisheries” by the Ministry of Agriculture, Forestry and Fisheries of Japan, and KAKENHI (19H03077, 19H03085, 23H02341). K. Matsumoto was supported by KAKENHI (25304027, 16H02762, 21H02238, 22K05752, 24H01520). T. Maximov was partly supported by the project “Study of biogeochemical cycles and adaptive reactions of plants of boreal and arctic ecosystems of northeastern Russia” (AAAA-A21-121012190034-2) of the Ministry of Education and Science of Russia. We thank the two anonymous reviewers and Dr. D. Papale for their constructive comments and suggestions.

References

- Asanuma, J.: JapanFlux2024 MN-Kbu Kherlenbayan Ulaan, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121004>, 2025a.
- Asanuma, J.: JapanFlux2024 MN-Skt Southern Khentei Taiga, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121011>, 2025b.
- Asanuma, J., Shimoda, S.: JapanFlux2024 JP-Tgf Terrestrial Environment Research Center, University of Tsukuba, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121013>, 2025.
- Awal, M. A., and Ohta, T.: JapanFlux2024 JP-Nuf Nagoya University Forest, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121008>, 2025a.
- Awal, M. A., and Ohta, T.: JapanFlux2024 JP-Tdf Toyota Deciduous Forest, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121012>, 2025b.
- Awal, M. A., Ohta, T., Matsumoto, K., Toba, T., Daikoku, K., Hattori, S., Hiyama, T., and Park, H.: Comparing the carbon sequestration capacity of temperate deciduous forests between urban and rural landscapes in central Japan, *Urb. For. Urb. Greening*, 9, 261–170, <https://doi.org/10.1016/j.ufug.2010.01.007>, 2010.
- Baldocchi, D., Novick, K., Keenan, T., and Torn, M.: AmeriFlux: Its Impact on our understanding of the ‘breathing of the biosphere’, after 25 years, *Agric. For. Meteorol.*, 348, 109929, <https://doi.org/10.1016/j.agrformet.2024.109929>, 2024.

- 545 Beer, C., Reichstein, M., Tomelleri, E., Ciais, P., Jung, M., Carvalhais, N., Rödenbeck, C., Arain, M. A., Baldocchi, D., Bonan, G. B., Bondeau, A., Cescatti, A., Lasslop, G., Lindroth, A., Lomas, M., Luysaert, S., Margolis, H., Oleson, K. W., Rouspard, O., Veenendaal, E., Viovy, N., Williams, C., Ian Woodward, F., and Papale, D.: Terrestrial Gross Carbon Dioxide Uptake: Global Distribution and Covariation with Climate. *Science*, 329, 834-838, <https://doi.org/10.1126/science.1184984>, 2010.
- 550 Boiarskii, B., and Hasegawa, H.: Comparison of NDVI and NDRE Indices to Detect Differences in Vegetation and Chlorophyll Content, *Internatl. Conf. Appl. Sci., Tech. Engin. J. Mech. Cont.& Math. Sci.*, 4, 20-29, <https://doi.org/10.26782/jmcms.spl.4/2019.11.00003>, 2019.
- Bonan, G. B., Oleson, K. W., Fisher, R. A., Lasslop, G., Reichstein, M.: Reconciling leaf physiological traits and canopy flux data: Use of the TRY and FLUXNET databases in the Community Land Model version 4, *J. Geophys. Res. Biogeosci.*, 117, G02026, <https://doi.org/10.1029/2011JG001913>, 2012.
- 555 Bond-Lamberty, B., Christianson, D. S., Malhotra, A., Pennington, S. C., Sihi, D., Agha-Kouchak, A., Anjileli, H., Arain, M. A., Armesto, J. J., Ashraf, S., Ataka, M., Baldocchi, D., Black, T. A., Buchmann, N., Carbone, M. S., Chang, S.-C., Crill, P., Curtis, P. S., Davidson, E. A., Desai, A. R., Drake, J., El-Madany, T. S., Gavazzi, M., Görres, C.-M., Gough, C. M., Goulden, M., Gregg, J., del Arroyo, O. G., He, J.-S., Hirano, T., Hopple, A., Hughes, H., Järveoja, J., Jassal, R., Jian, J., Kan, H., Kaye, J., Kominami, Y., Liang, N., Lipson, D., Macdonald, C., Maseyk, K., Mathes, K., Mauritz, M., Mayes, M.
- 560 A., McNulty, S., Miao, G., Migliavacca, M., Miller, S., Miniati, C. F., Nietz, J. G., Nilsson, M. B., Noormets, A., Norouzi, H., O'Connell, C. S., Osborne, B., Oyónarte, C., Pang, Z., Peichl, M., Pendall, E., Perez-Quezada, J. F., Phillips, C. L., Phillips, R. P., Raich, J. W., Renchon, A. A., Ruehr, N. K., Sánchez-Cañete, E. P., Saunders, M., Savage, K. E., Schrumpf, M., Scott, R. L., Seibt, U., Silver, W. L., Sun, W., Szutu, D., Takagi, K., Takagi, M., Teramoto, M., Tjoelker, M. G., Trumbore, S., Ueyama, M., Vargas, R., Varner, R. K., Verfaillie, J., Vogel, C., Wang, J., Winston, G., Wood, T. E., Wu, J., Wutzler, T., Zeng, J., Zha, T., Zhang, Q., and Zou, J.: COSORE: A community database for continuous soil respiration and other soil-atmosphere greenhouse gas flux data. *Glob. Change Biol.*, 26, 7268-7283, <https://doi.org/10.1111/gcb.15353>, 2020.
- Burba, G., McDermitt, D. K., Grelle, A., Anderson, D. J., Xu, L.: Addressing the influence of instrument surface heat exchange on the measurements of CO₂ flux from open-path gas analyzers, *Glob. Change Biol.*, 14, 1854-1876, <https://doi.org/10.1111/j.1365-2486.2008.01606.x>, 2008.
- 570 Delwiche, K. B., Knox, S. H., Malhotra, A., Fluet-Chouinard, E., McNicol, G., Feron, S., Ouyang, Z., Papale, D., Trotta, C., Canfora, E., Cheah, Y.-W., Christianson, D., Alberto, M. C. R., Alekseychik, P., Aurela, M., Baldocchi, D., Bansal, S., Billesbach, D. P., Bohrer, G., Bracho, R., Buchmann, N., Campbell, D. I., Celis, G., Chen, J., Chen, W., Chu, H., Dalmagro, H. J., Dengel, S., Desai, A. R., Detto, M., Dolman, H., Eichelmann, E., Euskirchen, E., Famulari, D., Fuchs, K., Goeckede, M., Gogo, S., Gondwe, M. J., Goodrich, J. P., Gottschalk, P., Graham, S. L., Heimann, M., Helbig, M., Helfter, C., Hemes, K. S., Hirano, T., Hollinger, D., Hörtnagl, L., Iwata, H., Jacotot, A., Jurasinski, G., Kang, M., Kasak, K., King, J., Klatt, J., Koebisch, F., Krauss, K. W., Lai, D. Y. F., Lohila, A., Mammarella, I., Manca, G., Marchesini, L. B., Matthes, J. H.,

- Maximon, T., Merbold, L., Mitra, B., Morin, T. H., Nemitz, E., Nilsson, M. B., Niu, S., Oechel, W. C., Oikawa, P. Y.,
580 Ono, K., Peichl, M., Peltola, O., Reba, M. L., Richardson, A. D., Riley, W., Runkle, B. R. K., Ryu, Y., Sachs, T., Sakabe,
A., Sanchez, C. R., Schuur, E. A., Schäfer, Karina, V. R., Sonnentag, O., Sparks, J. P., Stuart-Haëntjens, E., Sturtevant, C.,
Sullivan, R. C., Szutu, D. J., Thom, J. E., Torn, M. S., Tuittila, E.-S., Turner, J., Ueyama, M., Valach, A. C., Vargas, R.,
Varlagin, A., Vazquez-Lule, A., Verfaillie, J. G., Vesala, T., Vourlitis, G. L., Ward, E. J., Wille, C., Wohlfahrt, G., Wong,
G. X., Zhang, Z., Zona, D., Windham-Myers, L., Poulter, B., and Jackson, R. B.: FLUXNET-CH₄: A global, multi-
585 ecosystem dataset and analysis of methane seasonality from freshwater wetlands. *Earth System Science Data*, 13, 3607-
3689, <https://doi.org/10.5194/essd-13-3607-2021>, 2021.
- Delwiche, K. B., Nelson, J., Kowalska, N., Moore, C. E., Shirkey, G., Tarin, T., Cleverly, J. R., and Keenan, T. F.: Charting
the Future of the FLUXNET Network, *Bull. Ameri. Meteorol. Soc.*, 105, E466–E473, <https://doi.org/10.1175/BAMS-D-23-0316.1>, 2024.
- 590 Du, M., Kato, T., Tang, Y., Li, Y., Gu, S., Zhao, L., and Zhang, F.: JapanFlux2024 CN-HaM Qinghai Flux Research Site, 1.00,
Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102214>, 2025.
- Du, M., Li, Y., Zhang, F., Zhao, L., Li, H., Gu, S., Yonemura, S., and Tang, Y.: Characteristics and scenarios projection of
NEE change in an alpine meadow on the Tibetan Plateau, *Internatl. J. Glob. Warming (IJGW)*, 24,307-
325, <https://doi.org/10.1504/IJGW.2021.116711>, 2021.
- 595 Falge, E., Baldocchi, D., Olson, R., Anthoni, P., Aubinet, M., Bernhofer, C., Burba, G., Ceulemans, R., Clement, R., Dolman,
H., Granier, A., Gross, P., Grünwald, T., Hollinger, D., Jensen, N.-O., Katul, G., Keronen, P., Kowalski, A., Lai, C. T.,
Law B. E., Meyers, T., Moncrieff, J., Moors, E., Munger, W., Pilegaard, K., Rannik, Ü., Rebmann, C., Suyker, A.,
Tenhunen, J., Tu, K., Verma, S., Vesala, T., Wilson, K., and Wofsy, S.: Gap filling strategies for defensible annual sums
of net ecosystem exchange, *Agric. For. Meteorol.*, 107, 43-69, [https://doi.org/10.1016/S0168-1923\(00\)00225-2](https://doi.org/10.1016/S0168-1923(00)00225-2), 2001.
- 600 Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Bakker, D. C. E., Hauck, J., Landschützer, P., Le Quéré, C.,
2399 Luijkx, I. T., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson,
R. B., 2400 Alin, S. R., Anthoni, P., Barbero, L., Bates, N. R., Becker, M., Bellouin, N., Decharme, B., Bopp, L., Brasika,
I. B. M., 2401 Cadule, P., Chamberlain, M. A., Chandra, N., Chau, T.-T.-T., Chevallier, F., Chini, L. P., Cronin, M., Dou,
X., Enyo, K., 2402 Evans, W., Falk, S., Feely, R. A., Feng, L., Ford, D. J., Gasser, T., Ghattas, J., Gkritzalis, T., Grassi,
605 G., Gregor, L., Gruber, 2403 N., Gürses, Ö., Harris, I., Hefner, M., Heinke, J., Houghton, R. A., Hurtt, G. C., Iida, Y.,
Ilyina, T., Jacobson, A. R., Jain, A. 2404 K., Jarníková, T., Jersild, A., Jiang, F., Jin, Z., Joos, F., Kato, E., Keeling, R. F.,
Kennedy, D., Klein Goldewijk, K., Knauer, 2405 J., Korsbakken, J. I., Körtzinger, A., Lan, X., Lefèvre, N., Li, H., Liu, J.,
Liu, Z., Ma, L., Marland, G., Mayot, N., McGuire, 2406 P. C., McKinley, G. A., Meyer, G., Morgan, E. J., Munro, D. R.,
Nakaoka, S., Niwa, Y., O'Brien, K. M., Olsen, A., Omar, A. 2407 M., Ono, T., Paulsen, M., Pierrot, D., Pocock, K., Poulter,
610 B., Powis, C. M., Rehder, G., Resplandy, L., Robertson, E., 2408 Rödenbeck, C., Rosan, T. M., Schwinger, J., Séférian,
R., Smallman, T. L., Smith, S. M., Sospedra-Alfonso, R., Sun, Q., 2409 Sutton, A. J., Sweeney, C., Takao, S., Tans, P. P.,
Tian, H., Tilbrook, B., Tsujino, H., Tubiello, F., van der Werf, G. R., van Ooijen, E., Wanninkhof, R., Watanabe, M.,

- Wimart-Rousseau, C., Yang, D., Yang, X., Yuan, W., Yue, X., Zaehle, S., Zeng, 2411 J., and Zheng, B.: Global Carbon Budget 2023, *Earth Syst. Sci. Data*, 15, 5301–5369, <https://doi.org/10.5194/essd-15-5301-2023>, 2023.
- 615 Hamada, S., Ohta, T., Hiyama, T., Kuwada, T., Takahashi, A., and Maximov, T. C.: Hydrometeorological behaviour of pine and larch forests in eastern Siberia, *Hydrol. Proc.*, 18, 23-29, <https://doi.org/10.1002/hyp.1308>, 2004.
- Hammerle, A., Haslwanter, A., Schmitt, M., Bahn, M., Tappeiner, U., Cernusca, A., and Wohlfahrt, G.: Eddy covariance measurements of carbon dioxide, latent and sensible energy fluxes above a meadow on a mountain slope, *Bound.-Layer Meteorol.*, 122, 397-416, <https://doi.org/10.1007/s10546-006-9109-x>, 2007.
- 620 Harazono, Y.: JapanFlux2024 JP-Nsb NIAES Soybean, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102240>, 2025a.
- Harazono, Y.: JapanFlux2024 JP-Yrp Yawara Rice paddy, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121019>, 2025b.
- Harazono, Y., and Miyata, A.: JapanFlux2024 JP-Km1 Kushiro Mire: Onnenai Fen, 1.00, Arctic Data archive System (ADS),
- 625 Japan, <http://doi.org/10.17592/001.2024102239>, 2025a.
- Harazono, Y., and Miyata, A.: JapanFlux2024 JP-Km2 Kushiro Mire: Akanuma Bog, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121026>, 2025b.
- Harazono, Y., Kim, J., Miyata, A., Choi, T., Yun, J.-I., and Kim, J.-W.: Measurement of energy budget components during the International Rice Experiment (IREX) in Japan, *Hydrol. Proc.*, 12, 2081-2092, [https://doi.org/10.1002/\(SICI\)1099-1085\(19981030\)12:13/14<2081::AID-HYP721>3.0.CO;2-M](https://doi.org/10.1002/(SICI)1099-1085(19981030)12:13/14<2081::AID-HYP721>3.0.CO;2-M), 1998.
- 630 Harazono, Y., and Takagi, K.: JapanFlux2024 CN-In1 Inner Mongolia dune, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102241>, 2025a.
- Harazono, Y., and Takagi, K.: JapanFlux2024 CN-In2 Inner Mongolia grassland, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102242>, 2025b.
- 635 Harazono, Y., and Takagi, K.: JapanFlux2024 CN-In3 Inner Mongolia soybean, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102243>, 2025c.
- Harazono, Y., and Takagi, K.: JapanFlux2024 CN-In4 Inner Mongolia maize, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102244>, 2025d.
- Harazono, Y., and Takagi, K.: JapanFlux2024 CN-In5 Inner Mongolia no grazing, 1.00, Arctic Data archive System (ADS),
- 640 Japan, <http://doi.org/10.17592/001.2024102245>, 2025e.
- Harazono, Y., and Takagi, K.: JapanFlux2024 CN-In6 Inner Mongolia heavy grazing, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102246>, 2025f.
- Harazono, Y., and Takagi, K.: JapanFlux2024 CN-In7 Inner Mongolia light grazing, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102247>, 2025g.
- 645 Harazono, Y., and Takagi, K.: JapanFlux2024 CN-In8 Inner Mongolia medium grazing, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102248>, 2025h.

- Harazono, Y.: JapanFlux2024 JP-Hc1 Hachihama Experimental Farm: the International Rice Experiment, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102237>, 2025i.
- Harazono, Y., Takagi, K., and Miyata, A.: JapanFlux2024 JP-KaP Kasumigaura lotus paddy, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102238>, 2025.
- Harazono, Y., Yamada, C., and Nishizawa, T.: Characteristics of Aerodynamic Parameters and Turbulent Transport of Momentum and CO₂ Over a Soybean Canopy, *Bull. Environ. Res. Cent., the University of Tsukuba*, 16, 13-25, 1992.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis. *Quart. J. Roy. Meteorol. Soc.* 146(730), 1999–2049. <https://doi.org/10.1002/qj.3803>, 2020.
- Hirano, T.: JapanFlux2024 JP-Sb1 Sarobetsu Mire Moss, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102216>, 2025a.
- Hirano, T.: JapanFlux2024 JP-Sb2 Sarobetsu Mire Sasa, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102217>, 2025b.
- Hirano, T., and Hirata, R.: JapanFlux2024 JP-Tmd Tomakomai Flux Research Site Disturbed, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102230>, 2025.
- Hirano, T., Hirata, R., Fujinuma, Y., Saigusa, N., Yamamoto, S., Harazono, Y., Takada, M., Inukai, K., Inoue, G.: CO₂ and water vapor exchange of a larch forest in northern Japan. *Tellus*, 55B, 244-257, <https://doi.org/10.3402/tellusb.v55i2.16753>, 2003.
- Hirano, T., Ohkubo, S.: JapanFlux2024 ID-PaB Palangkaraya Drained Burnt forest, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102213>, 2025a.
- Hirano, T., and Ohkubo, S.: JapanFlux2024 ID-PaD Palangkaraya drained forest, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102232>, 2025b.
- Hirano, T., Ohkubo, S.: JapanFlux2024 ID-Pag Palangkaraya Undrained Forest, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102233>, 2025c.
- Hirano, T., Ohkubo, S., Itoh, M., Tsuzuki, H., Sakabe, A., Takahashi, H., Kusin, K., and Osaki, M.: Large variation in carbon dioxide emissions from tropical peat swamp forests due to disturbances, *Comm. Earth & Environ.*, 5, 221, <https://doi.org/10.1038/s43247-024-01387-7>, 2024.
- Hirano, T., Suzuki, K., and Hirata, R.: Energy balance and evapotranspiration changes in a larch forest caused by severe disturbance during an early secondary succession, *Agric. For. Meteorol.*, 232, 457-468, <https://doi.org/10.1016/j.agrformet.2016.10.003>, 2017.

- 680 Hirano, T., Yamada, H., Takada, M., Fujimura, Y., Fujita, H., and Takahashi, H.: Effects of the expansion of vascular plants in Sphagnum-dominated bog on evapotranspiration, *Agric. For. Meteorol.*, 220, 90-100, <https://doi.org/10.1016/j.agrformet.2016.01.039>, 2016.
- Hirata, R., and Hirano, T.: JapanFlux2024 JP-Tmk Tomakomai Flux Research Site, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102222>, 2025.
- 685 Hirata, R., Saigusa, N., Yamamoto, S., Ohtani, Y., Ide, R., Asanuma, J., Gamo, M., Hirano, T., Kondo, H., Kosugi, Y., Li, S. -G., Nakai, Y., Takagi, K., Tani, M., and Wang, H.: Spatial distribution of carbon balance in forest ecosystems across East Asia, *Agric. Meteorol. For.*, 148, 761-775, <https://doi.org/10.1016/j.agrformet.2007.11.016>, 2008.
- Hiyama, T., Kochi, K., Kobayashi, N. and Sirisampan, S.: Seasonal variation in stomatal conductance and physiological factors observed in a secondary warm-temperate forest, *Ecol. Res.*, 20, 333-346, [https://link.springer.com/chapter/10.1007/4-431-](https://link.springer.com/chapter/10.1007/4-431-29361-2_10)
- 690 29361-2_10, 2005.
- Ichii, K., Ueyama, M., Kondo, M., Saigusa, N., Kim, J., Alberto, M. C., Ardö, J., Euskirchen, E., Kang, M., Hirano, T., Joiner, J., Kobayashi, H., Marchesini, L. B., Merbold, L., Miyata, A., Saitoh, T. M., Takagi, K., Varlagin, A., Bret-Harte, M. S., Kitamura, K., Kosugi, Y., Kotani, A., Kumar, K., Li, S. -G., Machimura, T., Matsuura, Y., Mizoguchi, Y., Ohta, T., Mukherjee, S., Yanagi, Y., Yasuda, Y., Zhang, Y., and Zhao, F.: New data-driven estimation of terrestrial CO₂ fluxes in
- 695 Asia using a standardized database of eddy covariance measurements, remote sensing data, and support vector regression, *J. Geophys. Res. Biogeosci.*, 122, 767-795, <https://doi.org/10.1002/2016JG003640>, 2017.
- Igarashi, Y., Katul, G. G., Kumagai, T., Yoshifuji, N., Sato, T., Tanaka, N., Tanaka, K., Fujinami, H., Suzuki, M., and Tantasirin, C.: Separating physical and biological controls on long-term evapotranspiration fluctuations in a tropical deciduous forest subjected to monsoonal rainfall, *J. Geophys. Res.: Biogeosci.*, 120, 1262-1278, <https://doi.org/10.1002/2014JG002767>, 2015.
- 700 Iida, S., Shimizu, T., Shinohara, Y., Takeuchi, S., and Kumagai, T.: The necessity of sensor calibration for the precise measurement of water fluxes in forest ecosystems, In: Levina, D.F., Carlyle-Moses, D.E., Iida, S., Michalzik, B., Nanko, K., Tischer, A. (eds) *Forest-Water Interactions. Ecological Studies*, vol 240. Springer, Cham. [https://doi.org/10.1007/978-](https://doi.org/10.1007/978-3-030-26086-6_2)
- 705 Ishida, S.: JapanFlux2024 JP-Srk Shirakami Beech Forest Site, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102218>, 2025.
- Ishida S., Ito, D., and Matsuura, Y.: Overview of Shirakami Flux Tower and General Meteorological Conditions between July and October, 2008, *Shirakami Kenkyu.* 6, 18 - 25, 2009. (in Japanese with English abstract)
- Ishikawa, M.: JapanFlux2024 MN-Udg Udleg practice forest, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121017>, 2025.
- 710 Iwahana, G., Machimura, T., Kobayashi, Y., Fedorov, A. N., Konstantinov, P. Y., and Fukuda, M.: Influence of forest clear-cutting on the thermal and hydrological regime of the active layer near Yakutsk, eastern Siberia, *J. Geophys. Res.*, 110, G02004, <https://doi.org/10.1029/2005JG000039>, 2005.

- Iwata, H.: JapanFlux2024 JP-Nkm Nishikoma Site, 1.00, Arctic Data archive System (ADS), Japan, 715 <http://doi.org/10.17592/001.2024102231>, 2025a.
- Iwata, H.: JapanFlux2024 JP-SwL Suwa Lake Site, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.20241022-019>, 2025b.
- Iwata, H., and Suzuki, J.: JapanFlux2024 JP-Shn Shinshu University Experimental Forest Site, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121009>, 2025.
- 720 Iwata, H., Hirata, R., Takahashi, Y., Miyabara, Y., Itoh, M., and Iizuka, K.: Partitioning eddy-covariance methane fluxes from a shallow lake into diffusive and ebullitive fluxes, *Bound.-Layer Meteorol.*, 169, 413-428, <https://doi.org/10.1007/s10546-018-0383-1>, 2018.
- Kabeya, N., Shimizu, A., Shimizu, T., Iida, S., Tamai, K., Miyamoto, A., Chann, S., Araki, M., and Ohnuki, Y.: Long-term Hydrological Observations in a Lowland Dry Evergreen Forest Catchment Area of the Lower Mekong River, Cambodia, 725 *Japan Agricultural Res. Quart.*, 55, 177-190, <https://doi.org/10.6090/jarq.55.177>, 2021.
- Kanda, M., and Moriwaki, R.: JapanFlux2024 JP-Kgu Kugahara urban residential area, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121018>, 2025.
- Kang, M., and Cho, S.: Progress in water and energy flux studies in Asia: A review focused on eddy covariance measurements, *J. Agric. Meteorol.*, 77, 2-23, <https://doi.org/10.2480/agrmet.D-20-00036>, 2021.
- 730 Kato, T., and Tang, Y.: Spatial variability and major controlling factors of CO₂ sink strength in Asian terrestrial ecosystems: evidence from eddy covariance data, *Glob. Change Biol.*, 14, 2333-2348, <https://doi.org/10.1111/j.1365-2486.2008.01646.x>, 2008.
- Keenan, T. F., Luo, X., Stocker, B. D., De Kauwe, M. G., Medlyn, B. E., Prentice, I. C., Smith, N. G., Terrer, C., Wang, H., Zhang, Y., and Zhou, S.: A constraint on historic growth in global photosynthesis due to rising CO₂, *Nat. Clim. Chang.*, 735 13, 1376-1381, <https://doi.org/10.1038/s41558-023-01867-2>, 2023.
- Kitamura, K., Shimizu, T., Kominami, Y., Hagino, K., Mizoguchi, Y., Tamai, K., Shimizu, A., Ohnuki, Y., Kobayashi, M.: JapanFlux2024 JP-Khw Kahoku Experiment watershed, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121005>, 2025.
- Knox, S. H., Jackson, R. B., Poulter, B., McNicol, G., Fluet-Chouinard, E., Zhang, Z., Hugelius, G., Bousquet, P., Canadell, 740 J. G., Saunois, M., Papale, D., Chu, H., Keenan, T. F., Baldocchi, D., Torn, M. S., Mammarella, I., Trotta, C., Aurela, M., Bohrer, G., Campbell, D. I., Cescatti, A., Chamberlain, S., Chen, J., Chen, W., Dengel, S., Desai, A. R., Euskirchen, E., Friborg, T., Gasbarra, D., Godeed, I., Goeckede, M., Heimann, M., Helbig, M., Hirano, T., Hollinger, D. Y., Iwata, H., Kang, M., Klatt, J., Krauss, K. W., Kutzbach, L., Lohila, A., Mitra, B., Morin, T. H., Nilsson, M. B., Niu, S., Noormets, A., Oechel, W. C., Peichl, M., Peltola, O., Reba, M. L., Richardson, A. D., Runkle, B. R., Ryu, Y., Sachs, T., Schäfer, K. 745 B. R., Schmid, H. P., Shurpali, N., Sonnentag, O., Tang, A. C. I., Ueyama, M., Vargas, R., Vesala, T., Ward, E. J., Windham-Myers, L., Wohlfahrt, G., and Zona, D., 2019. FLUXNET-CH₄ synthesis activity: objectives, observations, and future directions, *Bull. Ameri. Meteorol. Soc.*, 100, 2607-2632, <https://doi.org/10.1175/BAMS-D-18-0268.1>, 2019.

- Kominami, Y., Jomura, M., Dannoura, M., Goto, Y., Tamai, K., Miyama, T., Kanazawa, Y., Kaneko, S., Okumura, M., Misawa, N., Hamada, S., Sasaki, T., Kimura, H., and Ohtani, Y.: Biometric and eddy-covariance-based estimates of carbon balance for a warm-temperate mixed forest in Japan, *Agric. For. Meteorol.*, 148, 723-737, <https://doi.org/10.1016/j.agrformet.2008.01.017>, 2008.
- Komiya, S.: Methane and carbon dioxide dynamics in temperate and tropical rice paddy fields, Ph.D. Dissertation, Meiji University, pp. 118, 2015.
- Komiya, S.: JapanFlux2024 JP-Hrt Hiratsuka Rice Paddy, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121020>, 2025a.
- Komiya, S.: JapanFlux2024 TH-Kms Kamphaeng Saen Rice Paddy, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121021>, 2025b.
- Kosugi, Y., Takanashi, S.: JapanFlux2024 JP-Ako JP-Ako Akou green belt, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121001>, 2025.
- Kosugi, Y., Tanaka, H., Takanashi, S., Matsuo, N., Ohte, N., Shibata, S., and Tani, M.: Three years of carbon and energy fluxes from Japanese evergreen broad-leaved forest, *Agric. For. Meteorol.*, 132, 329-343, <https://doi.org/10.1016/j.agrformet.2005.08.010>, 2005.
- Kotani, A., Ohta, T.: JapanFlux2024 JP-SMF Seto Mixed Forest Site, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102234>, 2025.
- Kotani, A., Ohta, T., Yabuki, H., Maximov, T., Petrov, R.: JapanFlux2024 RU-Sk2 Yakutsk Pine, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102236>, 2025.
- Kotani, A., Kononov, A. V., Ohta, T., and Maximov, T. C.: Temporal variations in the linkage between the net ecosystem exchange of water vapour and CO₂ over boreal forests in eastern Siberia, *Ecohydrology*, 7, 209-225, <https://doi.org/10.1002/eco.1449>, 2014.
- Kumagai, T., Hata, Y., Matsumoto, K., Kume, T.: JapanFlux2024 MY-LHP Lambir Hills National Park, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102206>, 2025.
- Kumagai, T., Takamura, N.: JapanFlux2024 TH-Kog Kog-Ma Watershed, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121023>, 2025a.
- Kumagai, T., Takamura, N.: JapanFlux2024 TH-Mae Mae Moh plantation, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121024>, 2025b.
- Kume, T., Takizawa, H., Yoshifuji, N., Tanaka, K., Tantasirin, C., Tanaka, N., and Suzuki, M.: Impact of soil drought on sap flow and water status of evergreen trees in a tropical monsoon forest in northern Thailand, *For. Ecol. Management*, 238, 220-230, <https://doi.org/10.1016/j.foreco.2006.10.019>, 2007.
- Lasslop, G., Reichstein, M., Papale, D., Richardson, A. D., Arneeth, A., Barr, A., Stoy, P., and Wohlfahrt, G.: Separation of net ecosystem exchange into assimilation and respiration using a light response curve approach: critical issues and global evaluation, *Glob. Change Biol.*, 16, 187-208, <https://doi.org/10.1111/j.1365-2486.2009.02041.x>, 2010.

- Li, S.-G., Asanuma, J., Eugster, W., Kotani, A., Davaa, G., Oyunbaatar, D., and Sugita, M.: Net ecosystem carbon dioxide exchange over grazed steppe in central Mongolia, *Glob. Change Biol.*, 11, 1941–1955, <https://doi.org/10.1111/j.1365-2486.2005.01047.x>, 2005a.
- 785 Li, S.-G., Asanuma, J., Kotani, A., Eugster, W., Davaa, G., Oyunbaatar, D., and Sugita, M.: Year-round measurements of net ecosystem CO₂ flux over a montane larch forest in Mongolia, *J. Geophys. Res.*, 110, D0930, <https://doi.org/10.1029/2004JD005453>, 2005b.
- Li, S. G., Harazono, Y., Oikawa, T., Zhao, H. L., He, Z. Y., and Chang, X. L.: Grassland desertification by grazing and the resulting micrometeorological changes in Inner Mongolia, *Agric. For. Meteorol.*, 102, 125-137, [https://doi.org/10.1016/S0168-1923\(00\)00101-5](https://doi.org/10.1016/S0168-1923(00)00101-5), 2000.
- 790 Liu, H.Z., Feng, J.W., Järvi, L., and Vesala, T.: Four-year (2006-2009) eddy covariance measurements of CO₂ flux over an urban area in Beijing. *Atmos. Chem. Phys.* 12, 7881–7892, <https://doi.org/10.5194/acp-12-7881-2012>, 2012.
- Lloyd, J. and Taylor, J. A.: On the temperature dependence of soil respiration. *Funct. Ecol.* 8, 315-323, <https://doi.org/10.2307/2389824>, 1994.
- 795 Machimura, T.: JapanFlux2024 RU-NeB Neleger Burnt Forest, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102210>, 2025a.
- Machimura, T.: JapanFlux2024 RU-NeC Neleger Cutover, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102211>, 2025b.
- Machimura, T.: JapanFlux2024 RU-NeF Neleger larch forest, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102212>, 2025c.
- 800 Matsuda, K., Watanabe, I., Mizukami, K., Ban, S., and Takahashi, A.: Dry deposition of PM_{2.5} sulfate above a hilly forest using relaxed eddy accumulation, *Atmo. Environ.*, 107, 255-261, <https://doi.org/10.1016/j.atmosenv.2015.02.050>, 2015.
- Matsumoto, K., Taniguchi, S., Takashima, A.: JapanFlux2024 JP-Ynf Yona-Field Tower Site, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102225>, 2025.
- 805 Matsumoto, K., Ohta, T., Nakai, T., Kuwada, T., Daikoku, K., Iida, S., Yabuki, H., Kononov, A. V., van der Molen, M. K., Kodama, Y., Maximov, T. C., Dolman, A. J., and Hattori, S.: Energy consumption and evapotranspiration at several boreal and temperate forests in the Far East, *Agric. For. Meteorol.*, 148, 1978-1989, <https://doi.org/10.1016/j.agrformet.2008.09.008>, 2008.
- Matsumoto, K., Terasawa, K., Taniguchi, S., Ohashi, M., Katayama, A., Kume, T., and Takashima, A.: Spatial and seasonal variations in soil respiration in a subtropical forest in Okinawa, Japan, *Ecol. Res.*, 38, 367-490, <https://doi.org/10.1111/1440-1703.12386>, 2023.
- 810 Matsuo, T., Sasyo, Y.: Non-melting phenomena of snowflakes observed in subsaturated air below freezing level, *J. Meteorol. Soc. Jpn.*, 59, 26-32, https://doi.org/10.2151/jmsj1965.59.1_26, 1981.

- Matsuura, S., Mori, A., Miyata, A., Hatano, R.: Effects of farmyard manure application and grassland renovation on net ecosystem carbon balance in a temperate grassland: analysis of 11-year eddy covariance data, *J. Agric. Meteorol.*, 79, 2-17, <https://doi.org/10.2480/agrmet.D-22-00007>, 2023.
- Matsuura, S.: JapanFlux2024 JP-NsM Nasu Research Station, Manure Application Plot, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121028>, 2025a.
- Matsuura, S.: JapanFlux2024 JP-NsC Nasu Research Station, Manure Application Plot, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121029>, 2025b.
- Matsuura, Y., and Morishita, T.: JapanFlux2024 RU-Tur Tura, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102223>, 2025.
- Maximov, T., Kotani, A., Petrov, R., Hiyama, T., and Ohta, T.: JapanFlux2024 RU-Ege Elgeei forest station, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102229>, 2025a.
- Maximov, T., Kotani, A., Petrov, R., Iijima, Y., Yabuki, H., Hiyama, T., and Ohta, T.: JapanFlux2024 RU-SkP Yakutsk Spasskaya Pad larch, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102235>, 2025b.
- Miyata, A., Harazono, Y., Kim, J., Terai, H., Takahashi, H., and Nishio, F.: Carbon dioxide and methane fluxes at Kushiro Mire, *Proc. International Workshop for Advanced Flux Network and Flux Evaluation*, 29-32, 2001.
- Miyazaki, S., Ishikawa, M., Baatarbileg, N., Damdinsuren, S., Ariuntuya, N., and Jambaljav, Y.: Interannual and seasonal variations in energy and carbon exchanges over the larch forests on the permafrost in northeastern Mongolia, *Polar Sci.*, 8, 166-182, <https://doi.org/10.1016/j.polar.2013.12.004>, 2014.
- Mizoguchi, Y., and Kitamura, K.: JapanFlux2024 JP-Spp Sapporo forest meteorology research site, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121010>, 2025.
- Mizoguchi, Y., Miyata, A., Ohtani, Y., Hirata, R., and Yuta, S.: A review of tower flux observation sites in Asia, *J. For. Res.*, 14, 1-9, <https://doi.org/10.1007/s10310-008-0101-9>, 2009.
- Mizoguchi, Y., Ohtani, Y., Takanashi, S., Iwata, H., Yasuda, Y., and Nakai, Y.: Seasonal and interannual variation in net ecosystem production of an evergreen needleleaf forest in Japan, *J. For. Res.*, 17, 283–295, <https://doi.org/10.1007/s10310-011-0307-0>, 2012.
- Moriwaki, R., and Kanda, M.: Seasonal and diurnal fluxes of radiation, heat, water vapor, and carbon dioxide over a suburban area, *J. Appl. Meteorol.*, 43, 1700-1710, <https://doi.org/10.1175/JAM2153.1>, 2004.
- Murayama, S., Kondo, H., Ishidoya, S., Maeda, T., Saigusa, N., Yamamoto, S., Kamezaki, K., and Muraoka, H.: Interannual variation and trend of carbon budget observed for more than two decades at Takayama in a cool-temperate deciduous forest in central Japan, *J. Geophys. Res.: Biogeosci.*, 129, e2023JG007769, <https://doi.org/10.1029/2023JG007769>, 2024a.
- Murayama, S., Kondo, H., Muraoka, H., Ishidoya, S., and Maeda, T.: JapanFlux2024 JP-Tak Takayama deciduous broadleaf forest site, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102221>, 2025b.

- Medlyn, B. E., Duursma, R. A., Eamus, D., Ellsworth, D. S., Prentice, I. C., Barton, C. V., Crous, K. Y., De Angelis, P., Freeman, M., and Wingate, L.: Reconciling the optimal and empirical approaches to modelling stomatal conductance, *Glob. Change Biol.*, 17, 2134-2144, <https://doi.org/10.1111/j.1365-2486.2010.02375.x>, 2011.
- 850 Nagano, H., and Hasegawa, H.: JapanFlux2024 JP-Mra Muramatsu agricultural field, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102209>, 2025.
- Nakai, T., Kim, Y., Busey, R. C., Suzuki, R., Nagai, S., Kobayashi, H., Park, H., Sugiura, K., and Ito, A.: Characteristics of evapotranspiration from a permafrost black spruce forest in interior Alaska, *Polar Sci.*, 7, 136-148, <https://doi.org/10.1016/j.polar.2013.03.003>, 2013.
- 855 Nakai, T., Ohta, T., Kodama, Y., Sumida, A., Toda, M., and Hara, T.: JapanFlux2024 JP-MBF Moshiri Birch Forest Site, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102207>, 2025a.
- Nakai, T., Ohta, T., Kodama, Y., Sumida, A., Toda, M., and Hara, T.: JapanFlux2024 JP-MMF Moshiri Mixed Forest Site, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.20241022-008>, 2025b.
- Nakai, T., van der Molen, M. K., Gash, J. H. C., and Kodama, Y.: Correction of sonic anemometer angle of attack errors, *Agric. For. Meteorol.*, 136, 19-30, <https://doi.org/10.1016/j.agrformet.2006.01.006>, 2006.
- 860 Nakai, Y., Matsuura, Y., Kajimoto, T., Abaimov, A. P., Yamamoto, S., and Zyryanova, O. A.: Eddy covariance CO₂ flux above a Gmelin larch forest on continuous permafrost in Central Siberia during a growing season, *Theo. Appl. Climatol.*, 93, 133-147, <https://doi.org/10.1007/s00704-007-0337-x>, 2008.
- Nakaji, T.: JapanFlux2024 JP-Tom Tomakomai Experimental Forest, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121015>, 2025.
- 865 Nakaji, T., Nakamura, M., and Ide, R.: JapanFlux2024 JP-Toc Tomakomai Crane site, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121014>, 2025.
- Nakamura, M., Nakaji, T., Muller, O., and Hiura, T.: Different initial responses of the canopy herbivory rate in mature oak trees to experimental soil and branch warming in a soil-freezing area, *OIKOS*, 124, 8, 1071-1077, <https://doi.org/10.1111/oik.01940>, 2014
- 870 Nakaya, K., Suzuki, C., Kobayashi, T., Ikeda, H., and Yasuike, S.: Application of a displaced-beam small aperture scintillometer to a deciduous forest under unstable atmospheric conditions, *Agric. For. Meteorol.*, 136, 45–55, <https://doi.org/10.1016/j.agrformet.2005.12.009>, 2006.
- Nakaya, K., Suzuki, C., Kobayashi, T., Ikeda, H., and Yasuike, S.: JapanFlux2024 JP-Kzw Karuizawa, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102205>, 2025.
- 875 Nie, D., Demetriades-Shah, T., and Kanemasu, E. T.: Surface energy fluxes on four slope sites during FIFE 1988, *J. Geophys. Res. Atmos.*, 97, 18641-18649, <https://doi.org/10.1029/91JD03043>, 1992.
- Ohta, T., Maximov, T. C., Dolman, A. J., Nakai, T., van der Molen, M. K., Kononov, A. V., Maximov, A. P., Hiyama, T., Iijima, Y., Moors, E. J., Tanaka, H., Toba, T., and Yabuki, H.: Interannual variation of water balance and summer

- evapotranspiration in an eastern Siberian larch forest over a 7-year period (1998–2006), *Agric. For. Meteorol.*, 148, 1941–1953, <https://doi.org/10.1016/j.agrformet.2008.04.012>, 2008.
- Ohtaki, E.: Application of an infrared carbon dioxide and humidity instrument to studies of turbulent transport. *College of Liberal Art. Sci.*, 29, 85-107, 1984.
- Ohkubo, S., Hirano, T., and Kusin, K.: Assessing the carbon dioxide balance of a degraded tropical peat swamp forest following multiple fire events of different intensities, *Agric. For. Meteorol.*, 306, 108448, <https://doi.org/10.1016/j.agrformet.2021.108448>, 2021.
- Ono, K.: JapanFlux2024 JP-Mse Mase paddy flux site, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121007>, 2025.
- Papale, D., Reichstein, M., Aubinet, M., Canfora, E., Bernhofer, C., Kutsch, W., Longdoz, B., Rambal, S., Valentini, R., Vesala, T., and Yakir, D.: Towards a standardized processing of Net Ecosystem Exchange measured with eddy covariance technique: algorithms and uncertainty estimation, *Biogeosciences*, 3, 571-583, <https://doi.org/10.5194/bg-3-571-2006>, 2006.
- Pastorello, G., Trotta, C., Canfora, E., Chu, H., Christianson, D., Cheah, Y.-W., Poindexter, C., Chen, J., Elbashandy, A., Humphrey, M., Isaac, P., Polidori, D., Ribeca, A., van Ingen, C., Zhang, L., Amiro, B., Ammann, C., Arain, M. A., Ardö, J., Arkebauer, T., Arndt, S. K., Arriga, N., Aubinet, M., Aurela, M., Baldocchi, D., Barr, A., Beamesderfer, E., Marchesini, L. B., Bergeron, O., Beringer, J., Bernhofer, C., Berveiller, D., Billesbach, D., Black, T. A., Blanken, P. D., Bohrer, G., Boike, J., Bolstad, P. V., Bonal, D., Bonnefond, J.-M., Bowling, D. R., Bracho, R., Brodeur, J., Brümmer, C., Buchmann, N., Burban, B., Burns, S. P., Buysse, P., Cale, P., Cavagna, M., Cellier, P., Chen, S., Chini, I., Christensen, T. R., Cleverly, J., Collalti, A., Consalvo, C., Cook, B. D., Cook, D., Coursolle, C., Cremonese, E., Curtis, P. S., D’Andrea, E., da Rocha, H., Dai, X., Davis, K. J., De Cinti, B., de Grandcourt, A., De Ligne, A., De Oliveira, R. C., Delpierre, N., Desai, A. R., Di Bella, C. M., di Tommasi, P., Dolman, H., Domingo, F., Dong, G., Dore, S., Duce, P., Dufrêne, E., Dunn, A., Dušek, J., Eamus, D., Eichmann, U., ElKhidir, H. A. M., Eugster, W., Ewenz, C. M., Ewers, B., Famulari, D., Fares, S., Feigenwinter, I., Feitz, A., Fensholt, R., Filippa, G., Fischer, M., Frank, J., Galvagno, M., Gharun, M., Gianelle, D., et al.: The FLUXNET2015 dataset and the ONEFlux processing pipeline for eddy covariance data, *Sci. Data*, 7, 225, <https://doi.org/10.1038/s41597-020-0534-3>, 2020.
- Saigusa, N., Li, S.-G., Kwon, H., Takagi, K., Zhang, L.-M., Ide, R., Ueyama, M., Asanuma, J., Choi, Y.-J., Chun, J. H., Han, S.-J., Hirano, T., Hirata, R., Kang, M., Kato, T., Kim, J., Li, Y.-N., Maeda, T., Miyata, A., Mizoguchi, Y., Murayama, S., Nakai, Y., Ohta, T., Saitoh, T. M., Wang, H.-M., Yu, G.-R., Zhang, Y.-P., and Zhao, F.-H.: Dataset of CarboEastAsia and uncertainties in the CO₂ budget evaluation caused by different data processing, *J. For. Res.*, 18, 41-48, <https://doi.org/10.1007/s10310-012-0378-6>, 2013.
- Saigusa, N., and Wang, H.: JapanFlux2024 CN-Lsh Laoshan, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121006>, 2025.

- Saito, M., Miyata, A., Nagai, H., and Yamada, T.: Seasonal variation of carbon dioxide exchange in rice paddy field in Japan, *Agric. For. Meteorol.*, 135, 93–109, <https://doi.org/10.1016/j.agrformet.2005.10.007>, 2005.
- Saitoh, T. M., and Tamagawa, I.: JapanFlux2024 JP-Ta2 Takayama evergreen coniferous forest site, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102220>, 2025.
- Saitoh, T. M., Tamagawa, I., Muraoka, H., Lee, N.-Y. M., Yashiro, Y., and Koizumi, H.: Carbon dioxide exchange in a cool-temperate evergreen coniferous forest over complex topography in Japan during two years with contrasting climates, *J. Plant Res.*, 123, 473–483, <https://doi.org/10.1007/s10265-009-0308-7>, 2010.
- Sakabe, A., and Itoh, M.: JapanFlux2024 JP-Nap Nunoiike Agricultural Pond, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121025>, 2025.
- Shibata, H., Hiura, T., Tanaka, Y., Takagi, K., and Koike, T.: Carbon cycling and budget in a forested basin of southwestern Hokkaido, northern Japan, *Ecol. Res.*, 20, 325–331, <https://doi.org/10.1007/s11284-005-0048-7>, 2005.
- Shimizu, T., Iida, S., Kabeya, N., and Iwagami, S.: JapanFlux2024 JP-Tkb Tsukuba Experimental Watershed, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121022>, 2025b.
- Shimizu, T., Kabeya, N., Iida, S., Tamai, K., Shimizu, A., Chann, S., and Saing, S.: JapanFlux2024 KH-Kmp Kampong Thom Lowland Dry Evergreen Forest, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102203>, 2025a.
- Shimizu, T., Kumagai, T., Kobayashi, M., Tamai, K., Iida, S., Kabeya, N., Ikawa, R., Tateishi, M., Miyazawa, Y., and Shimizu, A.: Estimation of annual forest evapotranspiration from a coniferous plantation watershed in Japan (2): Comparison of eddy covariance, water budget and sap-flow plus interception loss, *J. Hydrol.*, 522, 250–264, <https://doi.org/10.1016/j.jhydrol.2014.12.021>, 2015.
- Shimoda, S., Mo, W., and Oikawa, T.: The effects of characteristics of Asian Monsoon climate on interannual CO₂ exchange in a humid temperate C₃/C₄ co-occurring grassland, *SOLA*, 1, 169–172, <https://doi.org/10.2151/sola.2005-044>, 2005.
- Sugita, M., Ogawa, S., and Kawade, M.: Wind as a main driver of spatial variability of surface energy balance over a shallow 10²-km² scale lake: Lake Kasumigaura, Japan, *Water Resources Research*, 56, e2020WR027173, <https://doi.org/10.1029/2020WR027173>, 2020.
- Sugita, M.: JapanFlux2024 JP-KaL Koshin, Lake Kasumigaura, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102230>, 2025.
- Sulla-Menashe, D., Gray, J. M., Abercrombie, S. P., and Friedl, M. A.: Hierarchical mapping of annual global land cover 2001 to present: The MODIS Collection 6 Land Cover product, *Remote Sens. Environ.* 222, 183–194, <http://dx.doi.org/10.1016/j.rse.2018.12.013>, 2019.
- Takagi, K., Fukuzawa, K., Liang, N., Kayama, M., Nomura, M., Hojo, H., Sugata, S., Shibata, H., Fukuzawa, T., Takahashi, Y., Nakaji, T., Oguma, H., Mano, M., Akibayashi, Y., Murayama, T., Koike, T., Sasa, K., and Fujinuma, Y.: Change in CO₂ balance under a series of forestry activities in a cool-temperate mixed forest with dense undergrowth, *Glob. Change Biol.*, 15, 1275–1288, <https://doi.org/10.1111/j.1365-2486.2008.01795.x>, 2009.

- Takagi, K., and Matsuda, K.: JapanFlux2024 JP-Fmt Field Museum Tama Hills, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121003>, 2025.
- 950 Takagi, K., Miyata, A., Harazono, Y., Ota, N., Komine, M., and Yoshimoto, M.: An alternative approach to determining zero-plane displacement, and its application to a lotus paddy field, *Agric. For. Meteorol.*, 115, 173–181, [https://doi.org/10.1016/S0168-1923\(02\)00209-5](https://doi.org/10.1016/S0168-1923(02)00209-5), 2003.
- Takagi, K., and Takahashi, Y.: JapanFlux2024 JP-Tef CC-LaG Teshio Experimental Forest, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121016>, 2025.
- Takahashi, Y., Liang, N., Ide, R., Hatsumi, K., Yamao, Y., and Hirose, Y.: JapanFlux2024 JP-Fhk Fuji Hokuroku Flux Observation Site, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121002>, 2025.
- 955 Takahashi, Y., Saigusa, N., Hirata, R., ide, R., Fujinuma, Y., Okano, T., and Arase, T.: Characteristics of temporal variations in ecosystem CO₂ exchange in a temperate deciduous needle-leaf forest in the foothills of a high mountain, *J. Agric. Meteorol.*, 71, 302–317, <https://doi.org/10.2480/agrmet.D-14-00009>, 2015.
- Takamura, N., Hata, Y., Matsumoto, K., Kume, T., Ueyama, M., and Kumagai, T.: El Niño–Southern Oscillation forcing on carbon and water cycling in a Bornean tropical rainforest, *Proc. Natl. Acad. Sci. U. S. A.*, 120, e2301596120, <https://doi.org/10.1073/pnas.2301596120>, 2023.
- 960 Takanashi, S., Kominami, Y., and Miyama, T.: JapanFlux2024 JP-Fjy Fujiyoshida forest meteorology research site, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102202>, 2025a.
- Takanashi, S., Kominami, Y., and Miyama, T.: JapanFlux2024 JP-Yms Yamashiro forest meteorology research site, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102224>, 2025b.
- 965 Takano, T., and Ueyama, M.: Spatial variations in daytime methane and carbon dioxide emissions in two urban landscapes, Sakai, Japan. *Urb. Clim.*, 36, 100798, <https://doi.org/10.1016/j.uclim.2021.100798>, 2021.
- Takimoto, T., and Iwata, T., 2024, JapanFlux2024 JP-Hc2 Hachihama Experimental Farm, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102249>, 2025a.
- 970 Takimoto, T., and Iwata, T.: JapanFlux2024 JP-Hc3 Hachihama Experimental Farm: Double Crop, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102228>, 2025b.
- Takimoto, T., Iwata, T., Yamamoto, S., and Miura, T.: Characteristics of CO₂ and CH₄ flux at barley-rice double cropping field in southern part of Okayama, *J. Agric. Meteorol.*, 66, 181–191, <https://doi.org/10.2480/agrmet.66.3.5>, 2010.
- Twine, T. E., Kustas, W. P., Norman, J. M., Cook, D. R., Houser, P. R., Meyers, T. P., Prueger, J. H., Starks, P. J., and Wesely, M. L.: Correcting eddy-covariance flux underestimates over a grassland, *Agric. For. Meteorol.*, 103, 279–300, [https://doi.org/10.1016/S0168-1923\(00\)00123-4](https://doi.org/10.1016/S0168-1923(00)00123-4), 2000.
- 975 Ueyama, M.: JapanFlux2024 JP-Ozm Oizumi Urban Park, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024072201>, 2025a.
- Ueyama, M.: JapanFlux2024 JP-Om1 B11 building in Osaka Metropolitan University, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024072203>, 2025b.

- 980 Ueyama, M.: JapanFlux2024 JP-Om2 Farm field in Osaka Metropolitan University, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024072204>, 2025c.
- Ueyama, M.: JapanFlux2024 JP-Sac Sakai City Office, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102215>, 2025d.
- Ueyama, M., and Ando T.: Diurnal, weekly, seasonal and spatial variabilities in carbon dioxide flux in different urban
985 landscapes in Sakai, Japan, *Atmos. Chem. Phys.*, 16, 14727–14740, <https://doi.org/10.5194/acp-16-14727-2016>, 2016.
- Ueyama, M., Ichii, K., Kobayashi, H., Kumagai, T., Beringer, J., Merbold, L., Euskirchen, E., Hirano, T., Belelli M. L., Baldocchi, D., Saitoh, T., Mizoguchi, Y., Ono, K., Kim, J., Varlagin, A., Kang, M., Shimizu, T., Kosugi, Y., Bret-Harte, M., Machimura, T., Matsuura, Y., Ohta, T., Takagi, K., Takanashi, S., and Yasuda, Y.: Inferring CO₂ fertilization effect based on global monitoring land-atmosphere exchange with a theoretical model, *Environ. Res. Lett.*, 15, 084009,
990 <https://doi.org/10.1088/1748-9326/ab79e5>, 2020a.
- Ueyama, M., Hirano, T., and Kominami, Y.: JapanFlux2024 JP-BBY Bibai bog, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024072202>, 2025e.
- Ueyama, M., Iwata, H., Nagano, H., Kuku, N., and Harazono, Y.: Anomalous wet summers and rising atmospheric CO₂ concentrations increase the CO₂ sink in a poorly drained forest on permafrost, *Proc. Natl. Acad. Sci. U. S. A.*, 121, e2414539121 <https://doi.org/10.1073/pnas.2414539121>, 2024f.
- 995 Ueyama, M., Knox, S. H., Delwiche, K. B., Bansal, S., Riley, W. J., Baldocchi, D., Hirano, T., McNicol, G., Schafer, K., Windham-Myers, L., Poulter, B., Jackson, R. B., Chang, K.-Y., Chen, J., Chu, H., Desai, A. R., Gogo, S., Iwata, H., Kang, M., Mammarella, I., Peichl, M., Sonnentag, O., Tuittila, E.-S., Ryu, Y., Euskirchen, E. S., Göckede, M., Jacotot, A., Nilsson, M. B., and Sachs, T.: Modeled production, oxidation, and transport processes of wetland methane emissions in temperate,
1000 boreal, and Arctic regions, *Glob. Change Biol.*, 29, 2313–2334, <https://doi.org/10.1111/gcb.16594>, 2023.
- Ueyama, M., and Takano, T.: A decade of CO₂ flux measured by the eddy covariance method including the COVID-19 pandemic period in an urban center in Sakai, Japan, *Environ. Pollution*, 119210, <https://doi.org/10.1016/j.envpol.2022.119210>, 2022a.
- Ueyama, M., Taguchi, A., and Takano, T.: Water vapor emissions from urban landscapes in Sakai, Japan, *Journal of Hydrology*,
1005 598, 126384, <https://doi.org/10.1016/j.jhydrol.2021.126384>, 2021.
- Ueyama, M., Yamamori, T., Iwata, H., and Harazono, Y.: Cooling and moistening of the planetary boundary layer in interior Alaska due to a postfire change in surface energy exchange, *J. Geophys. Res. Atmos.*, 125, e2020JD032968, <https://doi.org/10.1029/2020JD032968>, 2020b.
- Ueyama, M., Yazaki, T., Hirano, T., and Endo, R.: Partitioning methane flux by the eddy covariance method in a cool temperate
1010 bog based on a Bayesian framework, *Agric. For. Meteorol.*, 316, 108852, <https://doi.org/10.1016/j.agrformet.2022.108852>, 2022b.
- Ueyama, M., Yazaki, T., Hirano, T., Futakuchi, and Okamura, M.: Environmental controls on methane fluxes in a cool temperate bog, *Agric. For. Meteorol.*, 281, 107852, <https://doi.org/10.1016/j.agrformet.2019.107852>, 2020c.

- Verma, M., Friedl, M. A., Richardson, A. D., Kiely, G., Cescatti, A., Law, B. E., Wohlfahrt, G., Gielen, B., Rouspard, O.,
1015 Moors, E. J., Toscano, P., Vaccari, F. P., Gianelle, D., Bohrer, G., Varlagin, A., Buchmann, N., van Gorsel, E., Montagnani,
L., and Propastin, P.: Remote sensing of annual terrestrial gross primary productivity from MODIS: an assessment using
the FLUXNET La Thuile data set, *Biogeosciences*, 11, 2185-2200, <https://doi.org/10.5194/bg-11-2185-2014>, 2014.
- Virkkala, A.-M., Natali, S. M., Rogers, B. M., Watts, J. D., Savage, K., Connon, S. J., Mauritz, M., Schuur, E. A. G., Peter,
D., Minions, C., Nojeim, J., Commane, R., Emmerton, C. A., Goeckede, M., Helbig, M., Holl, D., Iwata, H., Kobayashi,
1020 H., Kolari, P., López-Blanco, E., Marushchak, M. E., Mastepanov, M., Merbold, L., Parmentier, F.-J. W., Peichl, M., Sachs,
T., Sonnentag, O., Ueyama, M., Voigt, C., Aurela, M., Boike, J., Celis, G., Chae, N., Christensen, T. R., Bret-Harte, M. S.,
Dengel, S., Dolman, H., Edgar, C. W., Elberling, Bo., Euskirchen, E., Grelle, A., Hatakka, J., Humphreys, E., Järveoja, J.,
Kotani, A., Kutzbach, L., Laurila, T., Lohila, A., Mammarella, I., Matsuura, Y., Meyer, G., Nilsson, M. B., Oberbauer, S.
F., Park, S.-J., Petrov, R., Prokushkin, A. S., Schulze, C., St. Louis, V. L., Tuittila, E.-S., Tuovinen, J.-P., Quinton, W.,
1025 Varlagin, A., Zona, D., and Zyryanov V. I.: The ABCflux database: Arctic-Boreal CO₂ flux observations and ancillary
information aggregated to monthly time steps across terrestrial ecosystems. *Earth System Science Data*, 14, 179-208,
<https://doi.org/10.5194/essd-14-179-2022>, 2021.
- Vuichard, N., and Papale, D.: Filling the gaps in meteorological continuous data measured at FLUXNET sites with ERA-
Interim, *Earth Syst. Sci. Data*, 7, 157–171, <https://doi.org/10.5194/essd-7-157-2015>, 2015.
- 1030 Wang, Q., Peng, X., Okadera, T., Watanabe, M., Saito, Y., and Batkhishig, O.: JapanFlux2024 MN-Nkh Nalaikh grassland,
1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102227>, 2025a.
- Wang, Q., Peng, X., Okadera, T., Watanabe, M., Saito, Y., and Batkhishig, O.: JapanFlux2024 MN-Hst Hustai grassland, 1.00,
Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102226>, 2025b.
- Wang, H., Sun, F., Wang, T., Liu, W.: Estimation of daily and monthly diffuse radiation from measurements of global solar
1035 radiation a case study across China, *Renewable Energy*, 126, 226-241, <https://doi.org/10.1016/j.renene.2018.03.029>, 2018.
- Wang, H., Zu, Y., Saigusa, N., Yamamoto, S., Kondo, H., Yang, F., and Wang, W.: CO₂, water vapor and energy fluxes in a
larch forest in northeast China, *J. Agr. Meteorol.*, 60, 549–552, <https://doi.org/10.2480/agrmet.549>, 2005.
- Wang, Q., Peng, X., Watanabe, M., Batkhishig, O., Okadera, T., and Saito, Y.: Carbon budget in permafrost and non-
permafrost regions and its controlling factors in the grassland ecosystems of Mongolia, *Glob. Ecol. Conserv.*, 41, e02373,
1040 <https://doi.org/10.1016/j.gecco.2023.e02373>, 2023.
- Wutzler, T., Lucas-Moffat, A., Migliavacca, M., Knauer, J., Sickel, K., Šigut, L., Menzer, O., and Reichstein, M.: Basic and
extensible post-processing of eddy covariance flux data with REdDyProc, *Biogeosciences*, 16, 5015-5030,
<https://doi.org/10.5194/bg-15-5015-2018>, 2018.
- Yabuki, H., Ishii, Y., Ohata, T.: Comparison of water and heat balance on grassland and forest in Central Yakutia, East Siberia,
1045 In: *Proceedings 6th international study Conference on GEWEX in Asia and GAME (GAME CD-ROM Publ. 11)*,
T1HY30Jul04115511, 2004.

- Yabuki, H., Ishii, Y., Ohata, T.: JapanFlux2024 RU-Usk Ulakhan Sykkhan Alas, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024121027>, 2025.
- Yamanoi, K., Mizoguchi, Y., and Utsugi, H.: Effects of a windthrow disturbance on the carbon balance of a broadleaf deciduous forest in Hokkaido, Japan, *Biogeosciences*, 12, 6837–6851, <https://doi.org/10.5194/bg-12-6837-2015>, 2015.
- 1050 Yasuda, Y.: JapanFlux2024 JP-Api Appi forest meteorology research site, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102201>, 2025a.
- Yasuda, Y.: JapanFlux2024 JP-Kwg Kawagoe forest meteorology research site, 1.00, Arctic Data archive System (ADS), Japan, <http://doi.org/10.17592/001.2024102204>, 2025b.
- 1055 Yasuda, Y., Saito, T., Hoshino, D., Ono, K., Ohtani, Y., Mizoguchi, Y., and Morisawa, T.: Carbon balance in a cool–temperate deciduous forest in northern Japan: seasonal and interannual variations, and environmental controls of its annual balance, *J. For. Res.*, 17, 253–267, <https://doi.org/10.1007/s10310-011-0298-x>, 2012.
- Yasuda, Y., Watanabe, T., Ohtani, Y., Okano, M., and Nakayama, K.: Seasonal variation of CO₂ flux over a broadleaf deciduous forest, *J. Japan Soc. Hydrol. & Water Resour*, 11, 575–585, <https://doi.org/10.3178/jjshwr.11.575>, 1998.
- 1060 Yu, G.-R., Zhu, X.-J., Fu, Y.-L., He, H.-L., Wang, Q.-F., Wen, X.-F., Li, X.-R., Zhang, L.-M., Zhang, L., Su, W., Li, S.-G., Sun, X.-M., Zhang, Y.-P., Zhang, J.-H., Yan, J.-H., Wang, H.-M., Zhou, G.-S., Jia, B.-R., Xiang, W.-H., Li, Y.-N., Zhao, L., Wang, Y.-F., Shi, P.-L., Chen, S.-P., Xin, X.-P., Zhao, F.-H., Wang, Y.-Y., Tong, C.-L.: Spatial patterns and climate drivers of carbon fluxes in terrestrial ecosystems of China, *Glob. Change Biology*, 19, 798–810, <https://doi.org/10.1111/gcb.12079>, 2013.
- 1065 Zhang, Y., Peña-Arancibia¹, J. L., McVicar, T. R., Chiew, F. H. S., Vaze¹, J., Liu, C., Lu, X., Zheng, H., Wang, Y., Liu, Y. Y., Miralles, D. G., Pan, M.: Multi-decadal trends in global terrestrial evapotranspiration and its components, *Sci. Rep.*, 6, 19124, <https://doi:10.1038/srep19124>, 2016.