

1 **Crop-specific Management History of Phosphorus Fertilizer Input (CMH-P) in the**
2 **Croplands of United States: Reconciliation of Top-down and Bottom-up data Sources**

3 Peiyu Cao^{1,2,3,†}, Bo Yi^{1,†}, Franco Bilotto^{2,3}, Carlos Gonzalez Fischer^{2,3}, Mario Herrero^{2,3}, Chaoqun Lu¹

4 ¹Department of Ecology, Evolution, and Organismal Biology, Iowa State University, Ames, Iowa 5011,
5 USA

6 ²Department of Global Development, College of Agriculture and Life Sciences, Cornell University,
7 Ithaca, New York, USA

8 ³Cornell Atkinson Center for Sustainability, Cornell University, Ithaca, New York, USA

9 † These authors contributed equally to this work

10

11 *Correspondence to:* Chaoqun Lu (clu@iastate.edu)

12

13 **Abstract**

14 Understanding and assessing the spatiotemporal patterns in crop-specific phosphorus (P) fertilizer
15 management is crucial for promoting crop yield and mitigating environmental problems. The existing P
16 fertilizer dataset, derived from sales data, depicts an average application rate on total cropland at the
17 county level but overlooks cross-crop variations. Conversely, the survey-based dataset offers crop-
18 specific application details at the state level yet lacks inter-state variability. By reconciling these two
19 datasets, we developed long-term gridded maps to characterize crop-specific P fertilizer application rates,
20 timing, and methods across the contiguous US at a resolution of 4 km × 4 km from 1850 to 2022. We
21 found that P fertilizer application rate on fertilized area in the US increased from 0.9 g P m⁻² yr⁻¹ in 1940
22 to 1.9 g P m⁻² yr⁻¹ in 2022, with substantial variations among crops. However, approximately 40% of
23 cropland nationwide has remained unfertilized in the recent decade. The hotspots for P fertilizer use have
24 shifted from the southeastern and eastern US to the Midwest and the Great Plains over the past century,
25 reflecting changes in cropland area, crop choices, and P fertilizer use across different crops. Pre-planting
26 (fall and spring) and broadcast application are prevalent among corn, soybean, and cotton in the Midwest
27 and the Southeast, indicating a high P loss risk in these regions. In contrast, wheat and barley in the Great
28 Plains receive the most intensive P fertilizer at planting and via non-broadcast application. The P fertilizer
29 management dataset developed in this study can advance our comprehension in agricultural P budget and

30 facilitate the refinement in P fertilizer best management practices to optimize crop yield and reduce P
31 loss. Datasets are available at <https://doi.org/10.5281/zenodo.10700822> (Cao et al., 2024).

32 1 Introduction

33 Phosphorus (P) is fundamental for life on Earth, serving as a crucial component of genetic material,
34 cellular membranes, and adenosine triphosphate for energy storage. The application of P has facilitated
35 unprecedented increases in food, feed, fiber, and fuel production, and is one of the cornerstones of
36 modern agriculture (Tilman et al., 2002). Before the 19th century, the major P sources for agricultural
37 land were animal and human excreta, along with slaughterhouse by-products (Cordell et al., 2009;
38 Bouwman et al., 2013). Starting around the mid-to-late 19th century, the production of mineral P
39 fertilizers from phosphate rock grew rapidly after the mid-20th century (Lu and Tian, 2017). The
40 application of mineral P fertilizer increased from 1.0 Tg P yr⁻¹ to 1.7 Tg P yr⁻¹ from 1960 to 2017 in the
41 US (Samreen, 2019), rectifying the P deficiency of soils. However, P application was found to exceed the
42 crops needs by up to 50% in many regions across the US (Glibert, 2020; Sabo et al., 2021). A substantial
43 part of surplus P, defined as the difference between input and removal by crops, can be lost through
44 soluble P in runoff and subsurface flow, and particulate P in soil erosion. These losses can accumulate
45 along transport pathways such as soils, riparian areas, streams, and wetlands, leading to long-term impacts
46 on P loading (Sharpley et al., 2013; Stackpoole et al., 2019). Increased P loading has contributed to the
47 harmful algal blooms and large hypoxia zones, which degrade aquatic ecosystems and harm coastal
48 economies by destroying habitats, disrupting the food web, and damaging tourism and fisheries. To
49 improve P use efficiency in agriculture and mitigate the environmental impacts of excessive P, it is
50 essential to understand the spatial distribution and temporal dynamics of P fertilizer use.

51 Developing a contemporary P fertilizer dataset is challenging due to incomplete data from multiple
52 sources and the lack of information on crop-specific applications. Previous studies have developed
53 historical county-level P fertilizer consumption in the US from 1945 to 2017, following a top-down
54 approach that relies on state-level fertilizer sales data and county-level fertilizer expenditure data
55 (Alexander and Smith, 1990; Falcone, 2021; Brakebill and Gronberg, 2017). In these studies, the average
56 P fertilizer application was estimated by dividing the consumption by the total cropland area within each
57 county. These top-down P fertilizer databases utilize a single value for average P fertilizer use,
58 overlooking cross-crop variations. Additionally, the percentage of fertilized area relative to the total
59 planting area varies significantly among different crops (USDA-ERS, 2019). As not all planting areas are
60 fertilized, distributing total P fertilizer application on the total planting area has underestimated the actual
61 application rate in the fertilized fields. Characterizing the spatial and temporal heterogeneity of crop-

62 specific P fertilizer application rate due to different P demands across crop types can offer deeper insights
63 into P use efficiency, budget trajectories, and P loading analysis (Sabo et al., 2021; Stackpoole et al.,
64 2019; Swaney and Howarth, 2019). P fertilizer management practices, such as application timing and
65 method, also differ among crop types and are crucial for optimal nutrient management. For example, over
66 30% of rice fields in the US received injected P fertilizer, whereas around 40% of corn fields received
67 broadcasting P fertilizer (USDA-ERS, 2024), implying high potential P loss by runoff and erosion from
68 corn fields. A bottom-up approach, based on crop-specific P fertilizer application rates and management
69 practices on the treated areas, can help to improve the performance of models and develop P fertilizer
70 conserving strategies. However, to the best of our knowledge, there is a lack of comprehensive bottom-up
71 databases that provide long-term, spatially explicit, crop-specific P fertilizer management data across the
72 US.

73 By combining the top-down (total P consumption and average P application rate) and bottom-up (crop-
74 specific P application rate) data sets, we developed a spatially explicit time-series database to characterize
75 agricultural P fertilizer application rate, timing, and method in the contiguous US (CONUS) at 4 km
76 resolution from 1850 to 2022. The main objectives of this study are 1) to characterize the spatiotemporal
77 patterns of P fertilizer application rates across the US over the last 170 years by considering P fertilizer
78 management differences among crops; 2) to investigate the spatial patterns of P fertilizer application
79 timing and method.

80 2 Methods

81 We reconstructed the annual state-level crop-specific P fertilizer (hereafter referred to as P) application
82 rate from 1850 to 2022 using the same methodology in Cao et al. (2018) by integrating and gap-filling
83 multiple sources. Subsequently, the crop-specific P fertilizer application rate was adjusted to match the
84 state-level total P consumption. Using the same approach in Zhang et al. (2021), we further downscaled
85 the application rate to county-level during 1930-2022 based on county-level P consumption and cropland
86 acreage of each crop type (Ye et al., 2024). We split the annual P application rate generated above into
87 four application timings and three application methods according to the statewide crop-specific survey
88 data during the study period. The datasets of crop-specific P fertilizer management (application rate,
89 timing, and method) generated above were then spatialized into gridded maps based on annual time-series
90 maps of crop area and type at the spatial resolution of 1 km × 1 km across the CONUS (Ye et al., 2024)
91 (Fig. 1).

Deleted: 1970

Deleted: contiguous US

94 2.1 Historical P fertilizer use rate reconstruction

95 2.1.1 P fertilizer consumption

96 We obtained the historical P consumption from 1850 to 2022 for the CONUS by harmonizing the national
97 P consumption data from Mehring et al. (1957) for 1850-1951, USDA (1971) for 1952-1959, USDA-ERS
98 (2019) for 1960-2015, and FAO (2021) for 2016-2022.

Deleted: contiguous US

99 We integrated the annual state-level P consumption from multiple sources that cover different periods
100 during 1930-2016 (Table S1). We gap-filled the unavailable state-level P consumption data for the
101 periods pre-1930 and 2017-2022 by one-way interpolation (Eq. 1) using the national P consumption
102 generated above as a reference. Whereas the periods 1970-1975 and 1978-1987 were gap-filled by
103 distance-weighted interpolation (Eq. 2). The state-level P consumption generated above includes all
104 crops, cropland pasture, permanent pasture, and non-farm land (Table S2). By harmonizing and linearly
105 interpolating the ratio of P consumption of these lands to total consumption from multi-sources, we
106 calculated the P consumption of croplands, cropland pasture, permanent pasture, and non-farm from 1850
107 to 2022 in each state respectively (See supplementary material for details). We calculated the state-level P
108 application rate of cropland by dividing the P fertilizer consumption of cropland by the total cropland area
109 of each state.

110 Based on state fertilizer sales data provided by AAPFCO (2022) and county-level fertilizer expenditure
111 data from the USDA Census, the county-level P consumption was estimated every 5 years from 1969 to
112 2017 with 1987-2016 annually interpolated (Falcone, 2021; NuGIS, 2022). The missing years were
113 interpolated by Equation (2) during the periods of 1970-1986 and 2013-2016, and by Equation (1) after
114 2017 using the state-level P consumption generated above as reference. The state shares of different lands
115 were applied to estimate the P consumption of these lands in each county.

$$116 \text{ Interpolated } data_{i+k} = \frac{\text{Referenced trend}_{i+k}}{\text{Referenced trend}_i} \times \text{Raw data}_i, \quad (1)$$

$$117 \text{ Interpolated } data_{i+k} = \frac{\text{Referenced trend}_{i+k} \times \text{Raw data}_i}{\text{Referenced trend}_i} \times \frac{k-i}{j-i} + \frac{\text{Referenced trend}_{i+k} \times \text{Raw data}_j}{\text{Referenced trend}_j} \times \frac{j-k}{j-i}, \quad (2)$$

118 Where *Raw data* is the raw data that contains missing values, *Referenced trend* is the complete data
119 from which the inter-annual variations that raw data can refer to, *i* and *j* are the beginning and ending
120 year of the gap, *i + k* is the *k*th missing year. Equation 1 was used when the beginning or ending year is
121 unavailable, whereas Equation 2 was used when both years are available.

123 2.1.2 Referenced state-level crop-specific P application rate

124 The national P application rates of 9 major crop types, including corn, soybean, winter wheat, spring
125 wheat, cotton, sorghum, rice, barley, and durum wheat, from 1927 to 2022 were obtained by integrating
126 multiple data sources (Table S4). In contrast to the state-level P application rate generated in section
127 2.1.1, reflecting the inter-annual variation of each state, the national crop-specific P application rate
128 characterizes the variation of each crop at the national scale. We gap-filled the national crop-specific P
129 application rate for the period of 1850-2022 by using state-level P application rates as a reference. For the
130 period before 1927, when national crop-specific P application rates were unavailable, Equation (1) was
131 used to retrieve the P application rate of each crop. For the period from 1927 to 2022, the cubic spline
132 interpolation method was used to gap-fill P application rates when raw data were missing in less than 3
133 consecutive years. While Equation (2) was applied in gap-filling when missing data were found in more
134 than 3 consecutive years.

135 Four regression models, quadratic, cubic, exponential, and logarithmic functions, were built between the
136 interpolated national crop-specific P application rates and raw state-level crop-specific P application rates
137 of 9 crops from 1954 to 2022. The best-fit model was used to adjust the national crop-specific P
138 application rates (Cao et al., 2018). Finally, the interpolated national crop-specific P application rates
139 from 1850 to 1953 with no adjustment and from 1954 to 2022 with adjustment jointly served as the
140 referenced state-level crop-specific P application rate trend.

141 2.1.3 State- and county-level crop-specific P application rates

142 We obtained the state-level crop-specific P application rates of 9 crops from 1954 to 2022 from the same
143 data sources as national crop-specific P application rates (Table S4). ~~This includes the information of P~~
144 application rates in the fertilized ~~croplands and percentage of fertilized croplands~~. Due to the lack of
145 information to identify the fertilized cropland spatially, the P application rates were adjusted by
146 multiplying use rates with fertilized cropland percentage. For winter wheat, spring wheat, and durum
147 wheat, only the total P consumption of these three wheat types was available at the state level for the
148 period of 1954-1989. The wheat types planted in each state were determined based on the Agricultural
149 Chemical Use Survey (USDA-NASS, 2021). We calculated the fractions of P consumption for each
150 wheat type to the total P consumption of all wheat types in each state in 1990. This fraction was used to
151 estimate the P consumption of each wheat type for the period of 1954-1989. The P application rate of
152 each wheat type was then calculated as P consumption divided by the planting area of the corresponding
153 wheat type.

154 For the period from 1850 to 1953, the state-level P application rates of 9 crops were gap-filled by Eq. (1)
155 using the referenced P application rate generated in section 2.1.2. Whereas Eq. (2) and the cubic spline

Deleted: assembled

Deleted:), which represents

Deleted: cropland

159 method were used to gap-fill the missing years between 1954 and 2022 for missing years over or less 3
 160 consecutive years, respectively. The P consumption of cropland pasture calculated in section 2.1.1 was
 161 divided by the area in each state to generate the cropland pasture P application rate. The P consumption of
 162 all other crops in each state was calculated by subtracting the P consumption of 9 crops, cropland pasture,
 163 permanent pasture, and non-farm from state total P consumption. The P use rate of “Other Crops” was
 164 generated by dividing the P consumption by the area of Other Crops. Due to the mismatch between state
 165 total P consumption from top-down sales data and crop-specific P consumption from the bottom-up
 166 survey, the summed P consumption of 9 major crops exceeds the state total P amount in some states (Fig.
 167 S1), resulting in a negative rate of Other Crops. We adjusted the crop-specific application rates of major
 168 crops to match the state total P consumption, by assuming that total P consumption data from top-down
 169 source is more reliable. First, we reconstructed the positive application rates of Other Crops in each state.
 170 If the 10-year moving average of the positive application rates of the Other Crops was available, we used
 171 it to replace the negative rates of the Other Crops. Otherwise, if the moving average was unavailable, we
 172 interpolated the gaps using the area-weighted mean of Other Crops across all states within the
 173 corresponding region as the reference trend. The selection of Eq. (1) and Eq. (2) for interpolation depends
 174 on the availability of the beginning and ending year of the gap. After excluding the P fertilizer
 175 consumption of cropland pasture, Other Crops, permanent pasture, and non-farm uses from the state total
 176 P consumption, we used the remaining total consumption to scale the crop-specific P fertilizer application
 177 rates for major crops. Specifically, for certain crops that exhibit abnormal change trends in some states
 178 due to inadequate survey data (e.g., corn in Illinois), we manually adjusted the rates for these crops to
 179 align with the differences (Fig. S2).

180 By assuming the relative ratio of P application rate among crop types in counties follow their state-level
 181 patterns in the same year, the crop-specific P application rate generated above was downscaled from state
 182 level to county level using Eq. (3) from 1970 to 2022. The P consumption of each crop within a given
 183 county was calculated by multiplying the state-level P application rate by the planting acreage. A scaler
 184 was then calculated by dividing the county total P consumption by the summation of P consumption of all
 185 crop types to adjust the state-level P use rates for each crop within this county.

$$186 \quad P rate_i^{ct} = \frac{P cons_{ct}}{\sum_{j=1}^{11} P rate_j^{st} \times Area_j^{ct}} \times P rate_i^{st} \quad (3)$$

187 where $P rate_i^{ct}$ is the P application rate of crop type i in a given county, $P cons_{ct}$ is annual county P
 188 consumption, $P rate_j^{st}$ is the P application rate of crop type j in state st , $Area_j^{ct}$ is county-level planting
 189 area of crop type j , crops include 9 crops aforementioned, cropland pasture, and Other Crops.

Deleted: . When

Deleted: is

Deleted: were replaced by the average. When

Deleted: is

Deleted: by

Deleted: (Fig. 3).

Deleted: application

Deleted: By

Deleted: scaled

Deleted: of

Deleted: to align with the differences

201 2.2 P fertilizer application timing

202 By using the same approach as Cao et al. (2018), we estimated the P use at four application timings: fall
203 (previous year), spring (before planting), at planting, and after planting of 9 major crops in each state
204 from 1996 to 2013 from a statewide survey by USDA-ERS (2021) (Table S5). The raw data includes
205 crop-specific P fertilizer application rates and percentages of the fertilized cropland for each of the 4
206 timings in each state. We calculated the P fertilizer consumption at each timing by multiplying the
207 application rate with the area percentage and total cropland area. The fraction of the P fertilizer
208 consumption at each timing was used to split the annual P fertilizer application rate generated in Sect. 2.1
209 into 4 application timings. The years before 1996 and after 2013 were assumed to adopt the same
210 application timing strategy of years 1996 and 2013, respectively. We linearly interpolated the fractions of
211 missing years between 1996 and 2013. The average application timing fraction based on the fraction of
212 the abovementioned 8 major crops (excluding winter wheat), peanuts, and oats was used for cropland
213 pasture and Other Crops.

Deleted: P

Deleted: at

Deleted: Due to the lack of spatial information to locate

Deleted: fertilized area, all cropland was assumed to be fertilized at a lower application rate

Deleted: rates

Deleted: for 4 timings.

Deleted: application rate in

214 2.3 P fertilizer application method

215 USDA-ERS (2021) reported the percentages of fertilized cropland by 5 P application methods for each
216 crop during 1996-2013 based on a statewide survey (Table S5). For the years before 1996 and after 2013,
217 we assume farmers adopt the same application method strategy of years 1996 and 2013, respectively. Due
218 to the low adoption rate of the two mixed methods (Mixed method with incorporation and Mixed method
219 without incorporation, < 5%), we regrouped all 5 methods into 3 types: No Broadcast (e.g., chisel, knifed
220 in, and banded in), Incorporation (Broadcast with incorporation and Mixed method with incorporation),
221 and No Incorporation (Broadcast without incorporation and Mixed method without incorporation). We
222 calculated the fraction of fertilized cropland by each method to total fertilized cropland to split the annual
223 P application rate into 3 application methods. The average application method fraction of 8 major crops
224 (excluding winter wheat), peanuts, and oats was used for cropland pasture and other crops.

225 2.4 Developing gridded maps for characterizing P fertilizer management history

226 To characterize the variation in spatial P fertilizer management information, we assigned the state-level
227 (1850-1929) and county-level (1930-2021) crop-specific P fertilizer management data generated above to
228 1 km × 1 km gridded maps based on historical crop type distribution maps of the CONUS from 1850 to
229 2022 developed by Ye et al. (2024). It is worth noting that the P fertilizer management information
230 remains consistent for the same crop within a given county but varies across crops, while 1-km annual
231 crop type and area maps help add spatial heterogeneity of P fertilizer input within a county. The crop type

Deleted: For

Deleted: analysis

Deleted: contiguous US

243 distribution maps were developed using satellite images and imputed county-level planting area of each
244 crop type from the USDA-National Agricultural Statistics Service (2022). We timed the gridded P
245 application rate with crop density maps to convert the unit of P use rate from g P per cropland area to g P
246 per land area. The crop density maps were reconstructed by integrating various sources of inventory and
247 satellite data, representing the percentage of cropland within each pixel. More details about the land cover
248 maps can be found in Ye et al. (2024). We then resampled the P fertilizer management maps a $4 \text{ km} \times 4$
249 km resolution for display purposes. To examine the regional discrepancy of P fertilizer management in
250 the study area, we partitioned the CONUS into 7 regions according to the US-FNCA (2022), including
251 the Northwest (NW), the Southwest (SW), the Northern Great Plains (NGP), the Southern Great Plains
252 (SGP), the Midwest (MW), the Northeast (NE), and the Southeast (SE).

Deleted: at

Deleted: contiguous US

253 3 Results

254 3.1 Magnitude and spatiotemporal patterns of P fertilizer uses

255 The amount of total P consumption in the US kept a moderate increase trend from $0.002 \text{ Tg P yr}^{-1}$ in 1850
256 to 0.3 Tg P yr^{-1} in 1930, followed by a rapid rise to 2.2 Tg P yr^{-1} by 1980. After a swift fall to 1.6 Tg P^{-1}
257 in 1987, P consumption experienced large inter-annual fluctuations, reaching 1.7 Tg P^{-1} in 2022 (Fig. 2a).
258 In 1980, corn was the primary consumer of P fertilizer use (43% of national consumption), followed by
259 Other Crops (17%), soybean (11%), and winter wheat (10%). Conversely, other crop types accounted for
260 less than 10% of total use. In 2022, corn remained the dominant P fertilizer consumer (37%). However,
261 the shares of Other Crops and soybean increased to 23% and 19% in 2022, respectively, while the shares
262 of other crops diminished or remained stagnant (Fig. 2b & Fig S3). The P application rate on fertilized
263 areas rapidly increased from $0.9 \text{ g P m}^{-2} \text{ yr}^{-1}$ in 1940 to $2.5 \text{ g P m}^{-2} \text{ yr}^{-1}$ in 1979, then declined to 1.9 g P
264 $\text{m}^{-2} \text{ yr}^{-1}$ in 2022. In contrast, the P application rate on all cropland gradually increased from a low level of
265 $0.3 \text{ g P m}^{-2} \text{ yr}^{-1}$ in 1940, reaching its peak at $1.2 \text{ g P m}^{-2} \text{ yr}^{-1}$ in 1979 and leveling off to $1.1 \text{ g P m}^{-2} \text{ yr}^{-1}$ in
266 2022. It exhibited a smaller range of fluctuations over time. Correspondingly, a dramatic elevation in P
267 application rate was found among various crops from 1940 to 1980, with increments ranging from 0.5 g P
268 $\text{m}^{-2} \text{ yr}^{-1}$ in durum wheat to $2.4 \text{ g P m}^{-2} \text{ yr}^{-1}$ in corn (Fig. 2c). From 1980 to 2020, large decreases in
269 application rates were found in corn, winter wheat, sorghum, and cropland pasture, while large increases
270 were found in spring wheat, rice, and durum wheat. As an increasing proportion of total cropland received
271 P fertilizer from 1940 to 2022, the gap between P fertilizer use rate that on all cropland and on fertilized
272 area has been narrowing for most crops except for soybean and cropland pasture.

Deleted: conversely

Deleted: 2c

Deleted: 2b

278 Geospatially, as the P fertilizer consumption declined in the southeastern and eastern US and increased in
279 the Midwest and the Northern Great Plains since 1900, the hotspot of P use has shifted correspondingly
280 (Fig. 3-4). Low application rates ($< 0.4 \text{ g P m}^{-2} \text{ yr}^{-1}$) were common in the eastern US before 1940. The
281 application rates in the Midwest and west coast showed remarkable increases to above $1.0 \text{ g P m}^{-2} \text{ yr}^{-1}$ by
282 1980. After 2000, the east of the Northern Great Plains and the Midwest became the US hotspots,
283 displaying the most intensive P fertilizer use.

284 The P use in the Midwest and the Northern Great Plains is dominated by the nine major crops, whereas in
285 other regions, like the Northwest, Southwest, and Northeast, Other Crops account for a considerable share
286 of P use (Fig. 4). Owing to their wide cultivation, corn and soybean are the primary recipients of P
287 nationwide in the most recent decade (the 2020s). The intense P fertilizer use is concentrated in the
288 Midwest and the Northern Great Plains for corn ($> 0.8 \text{ g P m}^{-2} \text{ yr}^{-1}$) and for soybean ($0.5\text{-}1.2 \text{ g P m}^{-2} \text{ yr}^{-1}$)
289 (Fig. 5). In comparison, the P uses of the rest seven major crops are mainly distributed in different
290 regions. Low-level of application rate ($< 0.5 \text{ g P m}^{-2} \text{ yr}^{-1}$) is applied to cotton in the Southeast and the
291 Southern Great Plains. Sorghum is planted mainly in the Southern Great Plains with application rate < 0.2
292 $\text{g P m}^{-2} \text{ yr}^{-1}$. Rice is highly concentrated along the rice-belt and part of California with a relatively high
293 application rate ($0.5\text{-}0.8 \text{ g P m}^{-2} \text{ yr}^{-1}$). P fertilizer applied to barley, spring wheat, and durum wheat is
294 distributed in the Northern Great Plains at a moderate rate ($0.3\text{-}0.8 \text{ g P m}^{-2} \text{ yr}^{-1}$). Winter wheat has a wider
295 spatial distribution with a low application rate, except for some regions in Kansas, Oklahoma, and
296 Montana ($0.3\text{-}0.5 \text{ g P m}^{-2} \text{ yr}^{-1}$).

297 3.2 Patterns of P fertilizer application timings

298 Nationwide, corn, soybean, and cotton producers favor fall and spring applications before planting.
299 Conversely, producers of all three wheats and barley apply a large portion of annual P fertilizer at
300 planting (Fig. 6). The timing of P application varies significantly across the CONUS (Fig. S4). Fall
301 application prevails in the Midwest and the Southern Great Plains ($> 40\%$), especially in Iowa ($> 60\%$)
302 and Illinois ($> 50\%$) (Fig. S4a). Relatively high portions of P fertilizer, up to 20%, are also applied in fall
303 in the Southeast, the eastern Northern Great Plains, and the Northwest. In comparison, P applied in spring
304 before planting dominates across the nation, especially in the east of the US (Fig. S4b). Intense P
305 application ($> 50\%$) at planting is prevalent in the Northeast, the Northwest, and both the north part of the
306 Northern Great Plains and the Southern Great Plains (Fig. S4c). Application after planting is the least
307 popular application timing ($< 20\%$) in the nation, which mainly occurs in the Southern Great Plains, the
308 Southeast, and some other states (e.g., Michigan, Nebraska, and Washington) (Fig. S4d). In contrast to the
309 wider distribution of different timing ratios, the hotspots of P application rate for 4 timings were found in
310 the Midwest, the Great Plains, and the rice-belt due to generally low application rate in other regions (Fig.

Deleted: contiguous US

Deleted: 7

Deleted: 7a

Deleted: 7b

Deleted: 7c

Deleted: 7d.

317 7). Intense P fertilizer was applied in the fall in the Midwest ($> 0.6 \text{ g P m}^{-2}$) (Fig. 7a), particularly in Iowa
318 and Illinois. Spring application was concentrated in the corn-belt and rice belt with rates greater than 0.5 g
319 P m^{-2} (Fig. 7b). Farmers in the Northern Great Plains, Kansas, Indiana, and Wisconsin favored application
320 at planting (Fig. 7c). After planting applications were minimal ($< 0.2 \text{ g P m}^{-2}$) in the rice-belt and
321 Nebraska (Fig. 7d).

Formatted: Font color: Text 1

3.3 Patterns of P fertilizer application methods

Deleted: 2

323 Nationally, broadcast application is popular among corn, soybean, cotton, and rice. In contrast, the non-
324 broadcast method (e.g., injection and side-dress) dominates among three wheat types, sorghum, and
325 barley (Fig. 6). The adoption of the P application method differs substantially among regions (Fig. S5).

Deleted: .

Deleted: 8

326 Non-broadcast is predominantly used in Wisconsin, Michigan, the Great Plains, and the Northwest (Fig.
327 S5a). Broadcast with incorporation is widespread in the CONUS. However, the adoption rate is relatively

Deleted: 8a

Deleted: contiguous US.

Deleted: 8b

328 low ($< 40 \%$) in most of the region (Fig. S5b). In comparison, high P application by broadcast without
329 incorporation ($> 50\%$) is mainly distributed in the Midwest and the Southeast (Fig. S5c). Due to the
330 intense use of P fertilizer in the corn-belt and rice-belt, the hotspots of P application rate ($> 0.6 \text{ g P m}^{-2}$)
331 for 3 methods were found in various regions within these two belts (Fig. 8). Non-broadcast application is
332 prevalent in the Northern Great Plains, Kansas, and Minnesota (Fig. 8a). Intense application of P fertilizer
333 via broadcast with incorporation was observed in Minnesota and Illinois (Fig. 8b). The corn-belt and rice-
334 belt received most of their P fertilizer through broadcast without incorporation (Fig. 8c).

335 4 Discussion

336 4.1 Adjustments and improvements in state-level crop-specific P application rate

337 The national total P consumption obtained from the gap-filled bottom-up data in this study, summed from
338 all major crops, cropland pasture, permanent pasture, and non-farm use, aligns well with diverse top-
339 down data sources both in magnitude and inter-annual variations (Fig. S6). However, the bottom-up
340 source displays a larger P consumption of certain crops in certain states (e.g., corn in Illinois),
341 contributing to the divergences between these two approaches, notably after 2010 (Fig. S1&S2). These
342 overestimations may be caused by distorted crop-specific P application rate and/or fertilized area
343 percentage, derived from an inadequate survey pool. By modifying the surveyed crop-specific P
344 application rate at the state level, we matched the state total P consumption between bottom-up and top-
345 down approaches (Fig. 4). Despite the bottom-up source offering insights into cross-crop variations of P
346 application rate, it overlooks the inter-state variability. Based on the total P consumption and crop-
347 specific planting area in each county, we scaled the P application rate of each crop from state level to

Deleted: S3

355 county level, which portrays greater variability across counties. Particularly, the ranges are wider for corn,
356 soybean, winter wheat, sorghum, and barley ($0-6 \text{ g P m}^{-2} \text{ yr}^{-1}$) than those for spring wheat, cotton, rice,
357 durum, cropland pasture, and Other Crops (Fig. 9). In addition, downscaling state-level P application rate
358 to the county level augments the clarity of the geospatial pattern (Fig. 10). Top-down sources calculated
359 average P use rate in each county by dividing the total P consumption by all cropland areas, yielding in a
360 uniform value within each county but contrasting patterns across counties (Fig. 10a, d, g). Conversely,
361 our map based on bottom-up sources at the state level detailed spatial heterogeneity in intensive
362 agricultural regions, highlighting the cross-crop differences in P fertilizer use (Fig. 10b, e, h). By
363 combining these two sources, our map characterizes spatial variability across counties and crop types
364 (Fig. 10c, f, j). It highlights the region with intense P use, indicated by the top-down source, but also
365 differentiates P application rates among crops within each county, indicated by the bottom-up source.
366 This is particularly evident in the southern part of Missouri and the boundary between Minnesota and
367 Dakotas (Fig. 10c&j). Accurate information on fertilizer management is essential for improving
368 agricultural sustainability (Dhillon et al., 2017). Different crops have distinct P needs, and tailoring P use
369 based on these needs can enhance the efficiency of P fertilizer utilization, maximizing crop yield while
370 mitigating environmental impacts (Sabo et al., 2021). Moreover, detailed information on crop-specific P
371 fertilizer management is important for assessing P losses attributed to runoff, erosion, and leaching,
372 contributing to the development of agricultural policies (Daloğlu et al., 2012). Given the significance of
373 crop-specific information, we advocate for the incorporation of cross-crop variations into the
374 development of P fertilizer datasets.

375 4.2 Temporal and spatial dynamics of P fertilizer management

376 Concurrent with the historical changes in US cropland since 1850, P use has experienced different stages
377 of change similar to nitrogen fertilizer use (Cao et al., 2018), influenced by various factors. From 1850 to
378 1940, the primary crops, corn, cotton, and winter wheat, were mainly concentrated in the eastern US. The
379 constrained production of phosphate rock and low demand by limited crop productivity contributed to the
380 low level of P consumption and application rate. As cropland expanded to the Midwest and the Great
381 Plains from 1940 to 1980, the consumption of P fertilizer peaked after a sharp increase, driven by the
382 rising application rate and percentage of fertilized area across various crops (Fig. 2-5). The major
383 contributors to P consumption during this period were corn in the Midwest and spring wheat and winter
384 wheat in the Great Plains. Following a brief decline in the 1980s, due to improved fertilizer use efficiency,
385 increased use of animal manure, and farm crisis (Scholz et al., 2013; Bouwman et al., 2017; Zhang et al.,
386 2018), P consumption has stabilized with annual fluctuations primarily caused by changes in grain
387 demand and fertilizer prices (US-EPA, 2024). Throughout this period, P consumption continued to

Deleted: ,

Deleted: ,

Deleted: impacted

391 decline in the eastern US while increasing or leveling off in other regions, driven by the continued
392 expansion of corn and soybean at the expense of other crops (Fig. 2-5). Another possible contributing
393 factor to the decline in P consumption is that the generous high-rate P application over a half-century has
394 raised soil P level so much that it made it possible to have lower application and still meet crop demands
395 (Sabo et al., 2021; Bian et al., 2022).

396 In the past decade, the average percentage of P fertilized area in the US was around 60% (including
397 cropland and pasture), notably lower than that for nitrogen fertilizer. (Fig. ~~S7~~). The percentage of
398 fertilized area varies among crops, ranging from 42% for soybean to 89% for spring wheat. Estimating P
399 use efficiency and P losses in agricultural systems highly ~~relies~~ on the precise application rate of P
400 fertilizer (Solangi et al., 2023). It is noteworthy that, ~~when we develop the environmental assessments that~~
401 ~~are sensitive to P fertilizer application rates, the results might be biased without considering the fertilized~~
402 ~~area percentage, especially for~~ the crops with lower fertilized area percentages, such as soybean, cotton,
403 and sorghum.

404 Despite the application of P fertilizer after planting is strongly recommended for improving P fertilizer
405 use efficiency and minimizing P losses to the environment, this application timing remains the least
406 popular choice for major crops in the US. Notably, rice in the US rice belt, sorghum in the Southern Great
407 Plains, and cotton along the southwest coast were major contributors to post-planting applications. In
408 contrast, both fall and spring applications before planting, leaving P susceptible to loss (King et al., 2018),
409 have been widely adopted across multiple crops in the ~~CONUS~~ due to lower fertilizer prices, the
410 availability of labor, and the ease of operating equipment (Carver et al., 2022). Winter wheat in the
411 Southern Great Plain and the Northwest received over 40% of its annual P fertilizer in the fall, potentially
412 contributing to boosting yield. However, corn and soybean farmers in the Midwest, cotton farmers in the
413 Southwest and north of Texas, and sorghum farmers in the Southern Great Plains favor fall application,
414 implying a high potential risk for P loss (Nelson et al., 2023; Yuan et al., 2013). Except for winter wheat,
415 spring wheat, and durum wheat, all other crops receive more than a quarter of their annual P fertilizer in
416 spring before application. Despite being closer to the planting date, the P fertilizer applied during early
417 spring may be prone to loss via runoff, erosion, and leaching during intense rainfall (Williams and King,
418 2020; Algoazany et al., 2007). Application at planting is more prevalent among winter wheat and spring
419 wheat in the Southern Great Plains and the Northern Great Plains, respectively.

420 Non-broadcast application is commonly found for winter wheat, durum wheat, and barley in the
421 Northwest and Northern Great Plains, and for spring wheat, cotton, and sorghum in the Southern Great
422 Plains. In addition, corn farmers in Wisconsin, Michigan, and the Northeast apply most of their annual P
423 fertilizer using the non-broadcast method. The non-broadcast has been considered as a more conservative

Deleted: S4

Deleted: rely

Deleted: , may impact the estimate of these application rate-sensitive assessments without considering the fertilized area percentage.

Deleted: contiguous US

430 management to prevent P loss (Carver et al., 2022; Smith et al., 2016). However, broadcasting, including
431 post-incorporation and non-incorporation, remains widespread across the US, particularly in the Midwest
432 (hotspot for P fertilizer use) and the Southeast.

433 4.3 Uncertainty

434 The uncertainties of this database are mainly from several aspects: (1) The reconstructed P fertilizer
435 management data extends back to 1850. However, compared to the national P use information, finer scale
436 sources at the state- and county-level are only available from the 1930s onwards. Due to the absence of
437 earlier data, we interpolated the state-level P fertilizer consumption use back to 1850 by assuming they
438 have the consistent interannual variations with the national data. This approach to addressing the
439 temporal gaps may introduce larger uncertainties in the state-level temporal trajectories before
440 the 1930s; (2) Limited information on P use in cropland pasture and permanent pasture at finer temporal
441 and spatial resolution, contributing to uncertain estimates for Other Crops; (3) Due to the lack of
442 information on where croplands are fertilized, we assumed all the croplands in each state were fertilized
443 but at a lower rate by multiplying the rates in the fertilized cropland with the percentage of fertilized
444 cropland. This could lead to underestimation of P fertilizer use rate in fertilized areas and overestimation
445 in non-fertilized area, especially when the state-level fertilized cropland percentage is low. (4)
446 Adjustments were made on crop-specific P fertilizer use rates at the state level to reconcile top-down and
447 bottom-up data sources. However, the paucity of detailed crop-specific information may introduce biases
448 in our adjustments made for certain crops; (5) The composition of the Other Crops differs across states.
449 All crop types under Other Crops within each state receive equal P application rate, which may bias the
450 application rate for some crop types; (6) Due to the lack of finer spatial resolution information, we
451 assumed the crop-specific P application timing and method are identical within each state. However, the
452 spatial heterogeneity of application timing and method may be overlooked. Therefore, a finer resolution
453 of spatial and temporal survey capturing crop-specific P application rate, timing, and method will be
454 invaluable for enhancing our understanding of the spatiotemporal patterns of P fertilizer management
455 information in the US.

Deleted: 2

Deleted: 3

Deleted: 4

Deleted: .

456 5 Data availability

457 The P fertilizer management dataset is publicly available via ZENODO at
458 <https://doi.org/10.5281/zenodo.10700822> (Cao et al., 2024).

463 6 Conclusion

464 By harmonizing various data sources, we reconstructed a long-term spatially explicit P fertilizer
465 management dataset at 4 km ×4 km resolution from 1850 to 2022 in the CONUS. We discussed the
466 divergence between top-down (total P consumption) and bottom-up (crop-specific P fertilizer use) data
467 sources, underscoring the necessity to improve crop-specific management information in future surveys.
468 The newly developed dataset, leveraging the strengths of both data sources, highlights cross-crop
469 variabilities in the long-term use of P fertilizer among counties. The results reveal a substantial increase in
470 P fertilizer consumption and application rate from 1850 to 2022, notably during 1940-1980. However, the
471 magnitude and long-term changing trend differed significantly across crop types. It is worth noting that
472 approximately 40% of cropland in the US does not receive P fertilizer inputs. Since 1850, the hotspots of
473 P fertilizer use have shifted from the southeastern and eastern US to the Midwest and the Great Plains,
474 driven by changes in cropland distribution and P fertilizer application rate across different crop types.
475 Additionally, P fertilizer application timing and method vary substantially across crop types and regions.
476 Corn, soybean, and cotton in the Midwest and the Southeast receive over 60% of their annual P fertilizer
477 at pre-planting and through broadcasting. Conversely, winter wheat, spring wheat, durum wheat, and
478 barley in the Great Plains and the Northwest predominantly receive their annual P fertilizer at- and post-
479 planting, and via non-broadcasting. Promoting efficient P fertilizer management, encompassing the proper
480 application rate, timing, and method, is essential for enhancing P use efficiency and thus contributes to
481 economic, social, and environmental sustainability and profitability.

Deleted: contiguous US.

482 Author contributions

483 CL, PC, and BY conceptualized the paper and developed the methodology. PC and BY reconstructed the
484 dataset. PC and BY prepared the manuscript with contributions from all the co-authors.

485 Competing interests

486 At least one of the (co-)authors is a member of the editorial board of Earth System Science Data.

487 Acknowledgments

488 This work is supported by the Iowa Nutrient Research Center, the ISU College of Liberal Arts and
489 Sciences Dean's Faculty Fellowship, and the NSF CAREER grant (1945036).

491 **References**

- 492 Alexander, R. B. and Smith, R. A.: County-level estimates of nitrogen and phosphorus fertilizer use in the
493 United States, 1945 to 1985, US Department of the Interior, US Geological Survey, 1990.
- 494 Algoazany, A. S., Kalita, P. K., Czapar, G. F., and Mitchell, J. K.: Phosphorus Transport through
495 Subsurface Drainage and Surface Runoff from a Flat Watershed in East Central Illinois, USA, *J Environ*
496 *Qual*, 36, 681–693, <https://doi.org/https://doi.org/10.2134/jeq2006.0161>, 2007.
- 497 Association of American Plant Food Control Officials (AAPFCO): Commercial Fertilizers, available at:
498 <http://www.aapfco.org/publications.html>, last access: 20 December 2021, 2022.
- 499 Bian, Z., Pan, S., Wang, Z., Yao, Y., Xu, R., Shi, H., Kalin, L., Anderson, C., Justic, D., Lohrenz, S., and
500 Tian, H.: A Century-Long Trajectory of Phosphorus Loading and Export From Mississippi River Basin to
501 the Gulf of Mexico: Contributions of Multiple Environmental Changes, *Global Biogeochem Cycles*, 36,
502 e2022GB007347, <https://doi.org/https://doi.org/10.1029/2022GB007347>, 2022.
- 503 Bouwman, [A. F.](#), [Beusen, A. H. W.](#), [Lassaletta, L.](#), [Van Apeldoorn, D. F.](#), [Van Grinsven, H. J. M.](#), [Zhang,](#)
504 [J.](#), & [Ittersum Van, M. K.](#): [Lessons from temporal and spatial patterns in global use of N and P fertilizer](#)
505 [on cropland](#), *Sci Rep* 7, 40366, <https://doi.org/10.1038/srep40366>, 2017.
- 506 [Bouwman, L.](#), Goldewijk, K. K., Van Der Hoek, K. W., Beusen, A. H. W., Van Vuuren, D. P., Willems,
507 J., Rufino, M. C., and Stehfest, E.: Exploring global changes in nitrogen and phosphorus cycles in
508 agriculture induced by livestock production over the 1900-2050 period, *Proc Natl Acad Sci U S A*, 110,
509 20882–7, <https://doi.org/10.1073/pnas.1012878108>, 2013.
- 510 Brakebill, J. W. and Gronberg, J. M.: County-level estimates of nitrogen and phosphorus from
511 commercial fertilizer for the conterminous United States, 1987-2012, US Geological Survey Data release,
512 2017.
- 513 Cao, P., Lu, C., and Yu, Z.: Historical nitrogen fertilizer use in agricultural ecosystems of the contiguous
514 United States during 1850–2015: application rate, timing, and fertilizer types, *Earth Syst Sci Data*, 10,
515 969–984, <https://doi.org/10.5194/essd-10-969-2018>, 2018.
- 516 Cao, P., Yi, B., Bilotto, F., Gonzalez Fischer, C., Herrero, M., and Lu, C.: Annual crop-specific
517 management history of phosphorus fertilizer input (CMH-P) in the croplands of United States from 1850
518 to 2022: Application rate, timing, and method, *Zenodo*, <https://doi.org/10.5281/zenodo.10700822>, 2024.

Deleted: ..

520 Carver, R. E., Nelson, N. O., Roozeboom, K. L., Kluitenberg, G. J., Tomlinson, P. J., Kang, Q., and Abel,
521 D. S.: Cover crop and phosphorus fertilizer management impacts on surface water quality from a no-till
522 corn-soybean rotation, *J Environ Manage*, 301, 113818,
523 <https://doi.org/https://doi.org/10.1016/j.jenvman.2021.113818>, 2022.

524 Cordell, D., Drangert, J. O., and White, S.: The story of phosphorus: Global food security and food for
525 thought, *Global Environmental Change*, 19, 292–305, <https://doi.org/10.1016/j.gloenvcha.2008.10.009>,
526 2009.

527 Daloglu, I., Cho, K. H., and Scavia, D.: Evaluating Causes of Trends in Long-Term Dissolved Reactive
528 Phosphorus Loads to Lake Erie, *Environ Sci Technol*, 46, 10660–10666,
529 <https://doi.org/10.1021/es302315d>, 2012.

530 Dhillon, J., Torres, G., Driver, E., Figueiredo, B., and Raun, W. R.: World Phosphorus Use Efficiency in
531 Cereal Crops, *Agron J*, 109, 1670–1677, <https://doi.org/https://doi.org/10.2134/agronj2016.08.0483>,
532 2017.

533 Falcone, J. A.: Estimates of county-level nitrogen and phosphorus from fertilizer and manure from 1950
534 through 2017 in the conterminous United States, Open-File Report, Reston, VA, 20 pp.,
535 <https://doi.org/10.3133/ofr20201153>, 2021.

536 FAO (Food and Agriculture Organization of the United Nations): FAO online database, available at:
537 <http://www.fao.org/faostat/en/#data/RF>, last access: 10 August 2021, 2021.

538 Glibert, P. M.: From hogs to HABs: impacts of industrial farming in the US on nitrogen and phosphorus
539 and greenhouse gas pollution, Springer International Publishing, 139–180 pp.,
540 <https://doi.org/10.1007/s10533-020-00691-6>, 2020.

541 King, K. W., Williams, M. R., LaBarge, G. A., Smith, D. R., Reutter, J. M., Duncan, E. W., and Pease, L.
542 A.: Addressing agricultural phosphorus loss in artificially drained landscapes with 4R nutrient
543 management practices, *J Soil Water Conserv*, 73, 35, <https://doi.org/10.2489/jswc.73.1.35>, 2018.

544 Lu, C. and Tian, H.: Global nitrogen and phosphorus fertilizer use for agriculture production in the past
545 half century: Shifted hot spots and nutrient imbalance, *Earth Syst Sci Data*, 9, 181–192,
546 <https://doi.org/10.5194/essd-9-181-2017>, 2017.

547 Mehring, A. L., Adams, J. R., and Jacob, K. D.: Statistics on Fertilizers and Liming Materials in the
548 United States, USDA-Agricultural Research Service, Statistical Bulletin No. 191, Washington, D.C.,
549 USA, 1957.

550 Nelson, N. O., Roozeboom, K. L., Yeager, E. A., Williams, J. R., Zerger, S. E., Kluitenberg, G. J.,
551 Tomlinson, P. J., Abel, D. S., and Carver, R. E.: Agronomic and economic implications of cover crop and
552 phosphorus fertilizer management practices for water quality improvement, *J Environ Qual*, 52, 113–125,
553 <https://doi.org/https://doi.org/10.1002/jeq2.20427>, 2023.

554 Nutrient Use Geographic Information System (NuGIS): No Title, available at: <https://nugis.tfi.org/>, last
555 access: 20 December 2022, 2022.

556 Sabo, R. D., Clark, C. M., Gibbs, D. A., Metson, G., Todd, M. J., LeDuc, S. D., Greiner, D., Fry, M. M.,
557 Polinsky, R., Yang, Q., Tian, H., and Compton, J. E.: Phosphorus Inventory for the Conterminous United
558 States (2002–2012), *J Geophys Res Biogeosci*, n/a, e2020JG005684,
559 <https://doi.org/https://doi.org/10.1029/2020JG005684>, 2021.

560 Samreen, S.: Phosphorus Fertilizer: The Original and Commercial Sources, edited by: Zhang, S. K. E.-T.,
561 IntechOpen, Rijeka, Ch. 6, <https://doi.org/10.5772/intechopen.82240>, 2019.

562 [Scholz, R. W., Ulrich, A. E., Eilittä, M., & Roy, A.: Sustainable use of phosphorus: a finite resource, *Sci.*](#)
563 [*Total Environ.*, 461, 799–803, 2013.](#)

564 Sharpley, A., Jarvie, H. P., Buda, A., May, L., Spears, B., and Kleinman, P.: Phosphorus Legacy:
565 Overcoming the Effects of Past Management Practices to Mitigate Future Water Quality Impairment, *J*
566 *Environ Qual*, 42, 1308–1326, <https://doi.org/https://doi.org/10.2134/jeq2013.03.0098>, 2013.

567 Smith, D. R., Harmel, R. D., Williams, M., Haney, R., and King, K. W.: Managing Acute Phosphorus
568 Loss with Fertilizer Source and Placement: Proof of Concept, *Agricultural & Environmental Letters*, 1,
569 150015, <https://doi.org/https://doi.org/10.2134/acl2015.12.0015>, 2016.

570 Solangi, F., Zhu, X., Khan, S., Rais, N., Majeed, A., Sabir, M. A., Iqbal, R., Ali, S., Hafeez, A., Ali, B.,
571 Ercisli, S., and Kayabasi, E. T.: The Global Dilemma of Soil Legacy Phosphorus and Its Improvement
572 Strategies under Recent Changes in Agro-Ecosystem Sustainability, *ACS Omega*, 8, 23271–23282,
573 <https://doi.org/10.1021/acsomega.3c00823>, 2023.

574 Stackpoole, S. M., Stets, E. G., and Sprague, L. A.: Variable impacts of contemporary versus legacy
575 agricultural phosphorus on US river water quality, *Proc Natl Acad Sci U S A*, 116, 20562–20567,
576 <https://doi.org/10.1073/pnas.1903226116>, 2019.

577 Swaney, D. P. and Howarth, R. W.: Phosphorus use efficiency and crop production: Patterns of regional
578 variation in the United States, 1987–2012, *Science of the Total Environment*, 685, 174–188,
579 <https://doi.org/10.1016/j.scitotenv.2019.05.228>, 2019.

580 Tilman, D., Cassman, K., Matson, P., Naylor, R., and Polasky, S.: Agricultural sustainability and
581 intensive production practices, *Nature* 418, 671–677, <https://doi.org/10.1038/nature01014>, 2002.

582 U.S. Fourth National Climate Assessment: No Title, available at: <http://www.globalchange.gov/nca4>, last
583 access: 20 December 2022, 2022.

584 USDA (U.S. Department of Agriculture): Consumption of Commercial Fertilizers, Primary Plant
585 Nutrients, and Micronutrients, 1850–1969, USDA-Statistical Reporting Service, Crop Reporting Board,
586 Statistical Bulletin No. 472, Washington, D.C., USA, 1971.

587 Tailored Reports: Crop Production Practices: <https://data.ers.usda.gov/reports.aspx?ID=17883>.

588 USDA-ERS (U.S. Department of Agriculture-Economic Research Service): Fertilizer Use and Price,
589 available at: <https://www.ers.usda.gov/data-products/arms-farm-financial-and-cropproduction-practices/>
590 (last access: 10 August 2021), 2019.

591 USDA-NASS (U.S. Department of Agriculture-National Agricultural Service), S.: Agricultural Chemical
592 Use Program, available at: [https://www.nass.usda.gov/Surveys/Guide_
593 to_NASS_Surveys/Chemical_Use/index.php](https://www.nass.usda.gov/Surveys/Guide_to_NASS_Surveys/Chemical_Use/index.php), last access: 17 August 2021, 2021.

594 Williams, M. R. and King, K. W.: Changing Rainfall Patterns Over the Western Lake Erie Basin (1975–
595 2017): Effects on Tributary Discharge and Phosphorus Load, *Water Resour Res*, 56, e2019WR025985,
596 <https://doi.org/https://doi.org/10.1029/2019WR025985>, 2020.


597 Yuan, Y., Locke, M. A., Bingner, R. L., and Rebich, R. A.: Phosphorus losses from agricultural
598 watersheds in the Mississippi Delta, *J Environ Manage*, 115, 14–20,
599 <https://doi.org/https://doi.org/10.1016/j.jenvman.2012.10.028>, 2013.

600 Zhang, J., Gilbert, D., Gooday, A. J., Levin, L., Naqvi, S. W. A., Middelburg, J. J., Scranton, M., Ekau,
601 W., Peña, A., Dewitte, B., Oguz, T., Monteiro, P. M. S., Urban, E., Rabalais, N. N., Ittekkot, V., Kemp,
602 W. M., Ulloa, O., Elmgren, R., Escobar-Briones, E., and Van der Plas, A. K.: Natural and human-induced
603 hypoxia and consequences for coastal areas: synthesis and future development, *Biogeosciences*, 7, 1443–
604 1467, <https://doi.org/10.5194/bg-7-1443-2010>, 2010.

605 Zhang, J., Cao, P., and Lu, C.: Half-Century History of Crop Nitrogen Budget in the Conterminous
606 United States: Variations Over Time, Space and Crop Types, *Global Biogeochem Cycles*, 35,
607 e2020GB006876, <https://doi.org/https://doi.org/10.1029/2020GB006876>, 2021.



608 [Zhang, W., & Tidgren, K.: The current farm downturn vs the 1920s and 1980s farm crises: An economic](#)
609 [and regulatory comparison, *Agric. Econ. Rev.*, 78\(4\), 396-411, 2018.](#)

Deleted: ¶



613


614 Figure 1. Diagram for P fertilizer management dataset development. The upper blue box represents the
 615 development of state-level crop-specific P fertilizer application rate based on the bottom-up dataset. The
 616 lower green box represents the development of county-level P fertilizer application rate development by
 617 reconciling the top-down and bottom-up dataset.



Deleted:

618
 619 Figure 2. Time-series of P fertilizer consumption and average application rates for all crops (a), and P
 620 fertilizer consumption (b) and application rates (c) for 11 specific crops in the contiguous US. All
 621 cropland is the total planting area, while the fertilized area is the proportion of the cropland that receives
 622 P fertilizer. In panel (c), light-colored bars denote the application rate on fertilized area and dark-colored
 623 bars show the modified application rate with the assumption that the county-level P fertilizer consumption
 624 was distributed on all the croplands. Both start from zero on the y-axis.

- Deleted: P fertilizer
- Deleted: for 11 specific crops (b), and
- Deleted: shares across 11 crops
- Deleted: b),
- Deleted: cropland.




631
 632 Figure 3. Spatial distribution of P fertilizer application rates in the 1990s, 1940s, 1960s, 1980s, 2000s,
 633 and 2020s in the contiguous US at a resolution of 4-km x 4-km, with regions framed as NW (Northwest),
 634 NGP (Northern Great Plains), SGP (Southern Great Plains), SW (Southwest), MW (Midwest), SE


635 (Southeast), and NE (Northeast). The maps generated for 1900, 1940, and 1960 relied on state-level crop-
 636 specific data. Subsequent maps, post-1960, utilized county-level crop-specific data. The values on the
 637 map represent the P fertilizer use rate on all land areas and can be converted to P fertilizer use rate on per
 638 unit cropland area by lining up with our crop type and area database (Ye et al., 2024).

Formatted: Font color: Blue


639



640 Figure 4. Time-series of P fertilizer consumption by each state and 9 major crops from 1950 to 2022 in
 641 the contiguous US. The top-left figure illustrates the scales of x-axis and y-axis. The solid black line in
 642 each subplot represents total P fertilizer consumption, and the stacked area represents P fertilizer
 643 consumption by different crops. NW is the Northwest, NGP is the Northern Great Plains, SGP is the
 644 Southern Great Plains, SW is the Southwest, MW is the Midwest, SE is the Southeast, NE is the
 645 Northeast.





646
 647 Figure 5. Spatial distribution of P fertilizer application rates for 9 major crops in 2020 at 4-km x 4-km
 648 resolution, with regions framed as NW (Northwest), NGP (Northern Great Plains), SGP (Southern Great
 649 Plains), SW (Southwest), MW (Midwest), SE (Southeast), and NE (Northeast). [The values on the map](#)
 650 [represent the P fertilizer use rate on all land areas and can be converted to P fertilizer use rate on per unit](#)
 651 [cropland area by lining up with our crop type and area database \(Ye et al., 2024\)](#)



652
 653 Figure 6. The share of each application timing and method for 9 major crops in the US. FAL is fall
 654 application in previous year. SPR is spring application before planting. ATP is application at planting.
 655 AFP is application after planting. NBC is non-broadcast. BWJ is broadcast with injection, which is mix or
 656 inject after broadcast. BNJ is broadcast with no injection.

Deleted:




658

659 Figure 7. Spatial distribution of P fertilizer application rates at four application timings across the
 660 contiguous US in 2020.


Deleted: the fractions of four

Deleted: in




665

666 Figure 8. Spatial distribution of P fertilizer application rates in three application methods across the
 667 contiguous US in 2020.



Deleted: the fractions of three


Deleted: in



676

677 Figure 9. Comparison between state-level (red line) and county-level average (black boxplot) crop-
 678 specific P fertilizer application rate in primary crop-planting states in 2015. The red line indicates the
 679 state-level P fertilizer application rate. The box plot shows the distribution of county-level P fertilizer
 680 application rate (dots are outliers).

Deleted: 1



684

685 Figure 10. Comparison of spatial distribution of P fertilizer application rate in the contiguous US in 2016.
 686 NuGIS (a, d, g) represents the average application rate derived from county-level sales data. State (b, d, h)
 687 and county (c, f, i) data used for plotting represent the crop-specific P fertilizer application rate at state-
 688 and county-level developed in this study, respectively. To make it comparable, the same cropland map
 689 was used to mask out the cropland extent for NuGIS. Two red boxes in Fig d were zoomed in to
 690 demonstrate more details in the top and bottom panels.