



1 **Crop-specific Management History of Phosphorus Fertilizer Input (CMH-P) in the**
2 **Croplands of United States: Reconciliation of Top-down and Bottom-up data Sources**

3 Peiyu Cao^{1, 2, 3, †}, Bo Yi^{1, †}, Franco Bilotto^{2, 3}, Carlos Gonzalez Fischer^{2, 3}, Mario Herrero^{2, 3}, Chaoqun Lu¹

4 ¹Department of Ecology, Evolution, and Organismal Biology, Iowa State University, Ames, Iowa 5011,
5 USA

6 ²Department of Global Development, College of Agriculture and Life Sciences, Cornell University,
7 Ithaca, New York, USA

8 ³Cornell Atkinson Center for Sustainability, Cornell University, Ithaca, New York, USA

9 † These authors contributed equally to this work

10

11 *Correspondence to:* Chaoqun Lu (clu@iastate.edu)

12

13 **Abstract**

14 Understanding and assessing the spatiotemporal patterns in crop-specific phosphorus (P) fertilizer
15 management is crucial for promoting crop yield and mitigating environmental problems. The existing P
16 fertilizer dataset, derived from sales data, depicts an average application rate on total cropland at the
17 county level but overlooks cross-crop variations. Conversely, the survey-based dataset offers crop-
18 specific application details at the state level yet lacks inter-state variability. By reconciling these two
19 datasets, we developed long-term gridded maps to characterize crop-specific P fertilizer application rates,
20 timing, and methods across the contiguous US at a resolution of 4 km × 4 km from 1850 to 2022. We
21 found that P fertilizer application rate on fertilized area in the US increased from 0.9 g P m⁻² yr⁻¹ in 1940
22 to 1.9 g P m⁻² yr⁻¹ in 2022, with substantial variations among crops. However, approximately 40% of
23 cropland nationwide has remained unfertilized in the recent decade. The hotspots for P fertilizer use have
24 shifted from the southeastern and eastern US to the Midwest and the Great Plains over the past century,
25 reflecting changes in cropland area, crop choices, and P fertilizer use across different crops. Pre-planting
26 (fall and spring) and broadcast application are prevalent among corn, soybean, and cotton in the Midwest
27 and the Southeast, indicating a high P loss risk in these regions. In contrast, wheat and barley in the Great
28 Plains receive the most intensive P fertilizer at planting and via non-broadcast application. The P fertilizer
29 management dataset developed in this study can advance our comprehension in agricultural P budget and



30 facilitate the refinement in P fertilizer best management practices to optimize crop yield and reduce P
31 loss. Datasets are available at <https://doi.org/10.5281/zenodo.10700822> (Cao et al., 2024).

32 1 Introduction

33 Phosphorus (P) is fundamental for life on Earth, serving as a crucial component of genetic material,
34 cellular membranes, and adenosine triphosphate for energy storage. The application of P has facilitated
35 unprecedented increases in food, feed, fiber, and fuel production, and is one of the cornerstones of
36 modern agriculture (Tilman et al., 2002). Before the 19th century, the major P sources for agricultural
37 land were animal and human excreta, along with slaughterhouse by-products (Cordell et al., 2009;
38 Bouwman et al., 2013). Starting around the mid-to-late 19th century, the production of mineral P
39 fertilizers from phosphate rock grew rapidly after the mid-20th century (Lu and Tian, 2017). The
40 application of mineral P fertilizer increased from 1.0 Tg P yr⁻¹ to 1.7 Tg P yr⁻¹ from 1960 to 2017 in the
41 US (Samreen, 2019), rectifying the P deficiency of soils. However, P application was found to exceed the
42 crops needs by up to 50% in many regions across the US (Glibert, 2020; Sabo et al., 2021). A substantial
43 part of surplus P, defined as the difference between input and removal by crops, can be lost through
44 soluble P in runoff and subsurface flow, and particulate P in soil erosion. These losses can accumulate
45 along transport pathways such as soils, riparian areas, streams, and wetlands, leading to long-term impacts
46 on P loading (Sharpley et al., 2013; Stackpoole et al., 2019). Increased P loading has contributed to the
47 harmful algal blooms and large hypoxia zones, which degrade aquatic ecosystems and harm coastal
48 economies by destroying habitats, disrupting the food web, and damaging tourism and fisheries. To
49 improve P use efficiency in agriculture and mitigate the environmental impacts of excessive P, it is
50 essential to understand the spatial distribution and temporal dynamics of P fertilizer use.

51 Developing a contemporary P fertilizer dataset is challenging due to incomplete data from multiple
52 sources and the lack of information on crop-specific applications. Previous studies have developed
53 historical county-level P fertilizer consumption in the US from 1945 to 2017, following a top-down
54 approach that relies on state-level fertilizer sales data and county-level fertilizer expenditure data
55 (Alexander and Smith, 1990; Falcone, 2021; Brakebill and Gronberg, 2017). In these studies, the average
56 P fertilizer application was estimated by dividing the consumption by the total cropland area within each
57 county. These top-down P fertilizer databases utilize a single value for average P fertilizer use,
58 overlooking cross-crop variations. Additionally, the percentage of fertilized area relative to the total
59 planting area varies significantly among different crops (USDA-ERS, 2019). As not all planting areas are
60 fertilized, distributing total P fertilizer application on the total planting area has underestimated the actual
61 application rate in the fertilized fields. Characterizing the spatial and temporal heterogeneity of crop-



62 specific P fertilizer application rate due to different P demands across crop types can offer deeper insights
63 into P use efficiency, budget trajectories, and P loading analysis (Sabo et al., 2021; Stackpoole et al.,
64 2019; Swaney and Howarth, 2019). P fertilizer management practices, such as application timing and
65 method, also differ among crop types and are crucial for optimal nutrient management. For example, over
66 30% of rice fields in the US received injected P fertilizer, whereas around 40% of corn fields received
67 broadcasting P fertilizer (USDA-ERS, 2024), implying high potential P loss by runoff and erosion from
68 corn fields. A bottom-up approach, based on crop-specific P fertilizer application rates and management
69 practices on the treated areas, can help to improve the performance of models and develop P fertilizer
70 conserving strategies. However, to the best of our knowledge, there is a lack of comprehensive bottom-up
71 databases that provide long-term, spatially explicit, crop-specific P fertilizer management data across the
72 US.

73 By combining the top-down (total P consumption and average P application rate) and bottom-up (crop-
74 specific P application rate) data sets, we developed a spatially explicit time-series database to characterize
75 agricultural P fertilizer application rate, timing, and method in the contiguous US at 4 km resolution from
76 1850 to 2022. The main objectives of this study are 1) to characterize the spatiotemporal patterns of P
77 fertilizer application rates across the US over the last 170 years by considering P fertilizer management
78 differences among crops; 2) to investigate the spatial patterns of P fertilizer application timing and
79 method.

80 2 Methods

81 We reconstructed the annual state-level crop-specific P fertilizer (hereafter referred to as P) application
82 rate from 1850 to 2022 using the same methodology in Cao et al. (2018) by integrating and gap-filling
83 multiple sources. Subsequently, the crop-specific P fertilizer application rate was adjusted to match the
84 state-level total P consumption. Using the same approach in Zhang et al. (2021), we further downscaled
85 the application rate to county-level during 1970-2022 based on county-level P consumption and cropland
86 acreage of each crop type (Ye et al., 2024). We split the annual P application rate generated above into
87 four application timings and three application methods according to the statewide crop-specific survey
88 data during the study period. The datasets of crop-specific P management (application rate, timing, and
89 method) generated above were then spatialized into gridded maps based on annual time-series maps of
90 crop area and type at the spatial resolution of 1 km \times 1 km across the contiguous US (Ye et al., 2024)
91 (Fig. 1).



92 2.1 Historical P fertilizer use rate reconstruction

93 2.1.1 P fertilizer consumption

94 We obtained the historical P consumption from 1850 to 2022 for the contiguous US by harmonizing the
95 national P consumption data from Mehring et al. (1957) for 1850-1951, USDA (1971) for 1952-1959,
96 USDA-ERS (2019) for 1960-2015, and FAO (2021) for 2016-2022.

97 We integrated the annual state-level P consumption from multiple sources that cover different periods
98 during 1930-2016 (Table S1). We gap-filled the unavailable state-level P consumption data for the
99 periods pre-1930 and 2017-2022 by one-way interpolation (Eq. 1) using the national P consumption
100 generated above as a reference. Whereas the periods 1970-1975 and 1978-1987 were gap-filled by
101 distance-weighted interpolation (Eq. 2). The state-level P consumption generated above includes all
102 crops, cropland pasture, permanent pasture, and non-farm land (Table S2). By harmonizing and linearly
103 interpolating the ratio of P consumption of these lands to total consumption from multi-sources, we
104 calculated the P consumption of croplands, cropland pasture, permanent pasture, and non-farm from 1850
105 to 2022 in each state respectively (See supplementary material for details). We calculated the state-level P
106 application rate of cropland by dividing the P fertilizer consumption of cropland by the total cropland area
107 of each state.

108 Based on state fertilizer sales data provided by AAPFCO (2022) and county-level fertilizer expenditure
109 data from the USDA Census, the county-level P consumption was estimated every 5 years from 1969 to
110 2017 with 1987-2016 annually interpolated (Falcone, 2021; NuGIS, 2022). The missing years were
111 interpolated by Equation (2) during the periods of 1970-1986 and 2013-2016, and by Equation (1) after
112 2017 using the state-level P consumption generated above as reference. The state shares of different lands
113 were applied to estimate the P consumption of these lands in each county.

$$114 \text{ Interpolated } data_{i+k} = \frac{\text{Referenced trend}_{i+k}}{\text{Referenced trend}_i} \times \text{Raw data}_i, \quad (1)$$

$$115 \text{ Interpolated } data_{i+k} = \frac{\text{Referenced trend}_{i+k} \times \text{Raw data}_i}{\text{Referenced trend}_i} \times \frac{k-i}{j-i} + \frac{\text{Referenced trend}_{i+k} \times \text{Raw data}_j}{\text{Referenced trend}_j} \times \frac{j-k}{j-i}, \quad (2)$$

116 Where *Raw data* is the raw data that contains missing values, *Referenced trend* is the complete data
117 from which the inter-annual variations that raw data can refer to, *i* and *j* are the beginning and ending
118 year of the gap, *i + k* is the *k*th missing year. Equation 1 was used when the beginning or ending year is
119 unavailable, whereas Equation 2 was used when both years are available.



120 2.1.2 Referenced state-level crop-specific P application rate

121 The national P application rates of 9 major crop types, including corn, soybean, winter wheat, spring
122 wheat, cotton, sorghum, rice, barley, and durum wheat, from 1927 to 2022 were obtained by integrating
123 multiple data sources (Table S4). In contrast to the state-level P application rate generated in section
124 2.1.1, reflecting the inter-annual variation of each state, the national crop-specific P application rate
125 characterizes the variation of each crop at the national scale. We gap-filled the national crop-specific P
126 application rate for the period of 1850-2022 by using state-level P application rates as a reference. For the
127 period before 1927, when national crop-specific P application rates were unavailable, Equation (1) was
128 used to retrieve the P application rate of each crop. For the period from 1927 to 2022, the cubic spline
129 interpolation method was used to gap-fill P application rates when raw data were missing in less than 3
130 consecutive years. While Equation (2) was applied in gap-filling when missing data were found in more
131 than 3 consecutive years.

132 Four regression models, quadratic, cubic, exponential, and logarithmic functions, were built between the
133 interpolated national crop-specific P application rates and raw state-level crop-specific P application rates
134 of 9 crops from 1954 to 2022. The best-fit model was used to adjust the national crop-specific P
135 application rates (Cao et al., 2018). Finally, the interpolated national crop-specific P application rates
136 from 1850 to 1953 with no adjustment and from 1954 to 2022 with adjustment jointly served as the
137 referenced state-level crop-specific P application rate trend.

138 2.1.3 State- and county-level crop-specific P application rates

139 We obtained the assembled state-level crop-specific P application rates of 9 crops from 1954 to 2022
140 from the same data sources as national crop-specific P application rates (Table S4), which represents the
141 P application rates in the fertilized cropland. Due to the lack of information to identify the fertilized
142 cropland spatially, the P application rates were adjusted by multiplying use rates with fertilized cropland
143 percentage. For winter wheat, spring wheat, and durum wheat, only the total P consumption of these three
144 wheat types was available at the state level for the period of 1954-1989. The wheat types planted in each
145 state were determined based on the Agricultural Chemical Use Survey (USDA-NASS, 2021). We
146 calculated the fractions of P consumption for each wheat type to the total P consumption of all wheat
147 types in each state in 1990. This fraction was used to estimate the P consumption of each wheat type for
148 the period of 1954-1989. The P application rate of each wheat type was then calculated as P consumption
149 divided by the planting area of the corresponding wheat type.

150 For the period from 1850 to 1953, the state-level P application rates of 9 crops were gap-filled by Eq. (1)
151 using the referenced P application rate generated in section 2.1.2. Whereas Eq. (2) and the cubic spline
152 method were used to gap-fill the missing years between 1954 and 2022 for missing years over or less 3



153 consecutive years, respectively. The P consumption of cropland pasture calculated in section 2.1.1 was
154 divided by the area in each state to generate the cropland pasture P application rate. The P consumption of
155 all other crops in each state was calculated by subtracting the P consumption of 9 crops, cropland pasture,
156 permanent pasture, and non-farm from state total P consumption. The P use rate of “Other Crops” was
157 generated by dividing the P consumption by the area of Other Crops. Due to the mismatch between state
158 total P consumption from top-down sales data and crop-specific P consumption from the bottom-up
159 survey, the summed P consumption of 9 major crops exceeds the state total P amount in some states (Fig.
160 S1), resulting in a negative rate of Other Crops. We adjusted the crop-specific application rates of major
161 crops to match the state total P consumption. When the 10-year moving average of the positive
162 application rates of the Other Crops is available, the negative rates of the Other Crops were replaced by
163 the average. When the moving average is unavailable, we interpolated the gaps by using the area-
164 weighted mean of Other Crops across all states within the corresponding region (Fig. 3). The application
165 of Eq. (1) and Eq. (2) for interpolation depends on the availability of the beginning and ending year of the
166 gap. By excluding the P consumption of cropland pasture, Other Crops, permanent pasture, and non-farm
167 from state total P consumption, we scaled the crop-specific P application rates of major crops to align
168 with the differences. Specifically, for certain crops that exhibit abnormal change trends in some states due
169 to inadequate survey data (e.g., corn in Illinois), we manually adjusted the rates for these crops to align
170 with the differences (Fig. S2).

171 By assuming the relative ratio of P application rate among crop types in counties follow their state-level
172 patterns in the same year, the crop-specific P application rate generated above was downscaled from state
173 level to county level using Eq. (3) from 1970 to 2022. The P consumption of each crop within a given
174 county was calculated by multiplying the state-level P application rate by the planting acreage. A scaler
175 was then calculated by dividing the county total P consumption by the summation of P consumption of all
176 crop types to adjust the state-level P use rates for each crop within this county.

$$177 \quad P rate_i^{ct} = \frac{P cons_{ct}}{\sum_{j=1}^{11} P rate_j^{st} \times Area_j^{ct}} \times P rate_i^{st} \quad (3)$$

178 where $P rate_i^{ct}$ is the P application rate of crop type i in a given county, $P cons_{ct}$ is annual county P
179 consumption, $P rate_j^{st}$ is the P application rate of crop type j in state st , $Area_j^{ct}$ is county-level planting
180 area of crop type j , crops include 9 crops aforementioned, cropland pasture, and Other Crops.

181 2.2 P fertilizer application timing

182 By using the same approach as Cao et al. (2018), we estimated four P application timings: fall (previous
183 year), spring (before planting), at planting, and after planting of 9 major crops in each state from 1996 to



184 2013 from a statewide survey by USDA-ERS (2021) (Table S5). The raw data includes crop-specific P
185 application rates and percentages of the fertilized cropland at 4 timings in each state. Due to the lack of
186 spatial information to locate the fertilized area, all cropland was assumed to be fertilized at a lower
187 application rate by multiplying the application rates with the area percentage for 4 timings. The fraction
188 of the application rate in each timing was used to split the annual P application rate generated in Sect. 2.1
189 into 4 application timings. The years before 1996 and after 2013 were assumed to adopt the same
190 application timing strategy of years 1996 and 2013, respectively. We linearly interpolated the fractions of
191 missing years between 1996 and 2013. The average application timing fraction based on the fraction of
192 the abovementioned 8 major crops (excluding winter wheat), peanuts, and oats was used for cropland
193 pasture and Other Crops.

194 2.3 P fertilizer application method

195 USDA-ERS (2021) reported the percentages of fertilized cropland by 5 P application methods for each
196 crop during 1996-2013 based on a statewide survey (Table S5). Due to the low adoption rate of the two
197 mixed methods (Mixed method with incorporation and Mixed method without incorporation, < 5%), we
198 regrouped all 5 methods into 3 types: No Broadcast (e.g., chisel, knifed in, and banded in), Incorporation
199 (Broadcast with incorporation and Mixed method with incorporation), and No Incorporation (Broadcast
200 without incorporation and Mixed method without incorporation). We calculated the fraction of fertilized
201 cropland by each method to total fertilized cropland to split the annual P application rate into 3
202 application methods. The average application method fraction of 8 major crops (excluding winter wheat),
203 peanuts, and oats was used for cropland pasture and other crops.

204 2.4 Developing gridded maps for characterizing P fertilizer management history

205 For spatial analysis, we assigned the state-level and county-level crop-specific P management data
206 generated above to 1 km × 1 km gridded maps based on historical crop type distribution maps of the
207 contiguous US from 1850 to 2022 developed by Ye et al. (2024). The crop type distribution maps were
208 developed using satellite images and imputed county-level planting area of each crop type from the
209 USDA-National Agricultural Statistics Service (2022). We timed the gridded P application rate with crop
210 density maps to convert the unit of P use rate from g P per cropland area to g P per land area. The crop
211 density maps were reconstructed by integrating various sources of inventory and satellite data,
212 representing the percentage of cropland within each pixel. More details about the land cover maps can be
213 found in Ye et al. (2024). We then resampled the P management maps at a 4 km × 4 km resolution for
214 display purposes. To examine the regional discrepancy of P management in the study area, we partitioned
215 the contiguous US into 7 regions according to the US-FNCA (2022), including the Northwest (NW), the



216 Southwest (SW), the Northern Great Plains (NGP), the Southern Great Plains (SGP), the Midwest (MW),
217 the Northeast (NE), and the Southeast (SE).

218 3 Results

219 3.1 Magnitude and spatiotemporal patterns of P fertilizer uses

220 The amount of total P consumption in the US kept a moderate increase trend from 0.002 Tg P yr⁻¹ in 1850
221 to 0.3 Tg P yr⁻¹ in 1930, followed by a rapid rise to 2.2 Tg P yr⁻¹ by 1980. After a swift fall to 1.6 Tg P yr⁻¹
222 in 1987, P consumption experienced large inter-annual fluctuations, reaching 1.7 Tg P yr⁻¹ in 2022 (Fig. 2a).
223 In 1980, corn was the primary consumer of P fertilizer use (43% of national consumption), followed by
224 Other Crops (17%), soybean (11%), and winter wheat (10%). conversely, other crop types accounted for
225 less than 10% of total use. In 2022, corn remained the dominant P fertilizer consumer (37%). However,
226 the shares of Other Crops and soybean increased to 23% and 19% in 2022, respectively, while the shares
227 of other crops diminished or remained stagnant (Fig. 2c). The P application rate on fertilized areas rapidly
228 increased from 0.9 g P m⁻² yr⁻¹ in 1940 to 2.5 g P m⁻² yr⁻¹ in 1979, then declined to 1.9 g P m⁻² yr⁻¹ in
229 2022. In contrast, the P application rate on all cropland gradually increased from a low level of 0.3 g P m⁻²
230 yr⁻¹ in 1940, reaching its peak at 1.2 g P m⁻² yr⁻¹ in 1979 and leveling off to 1.1 g P m⁻² yr⁻¹ in 2022. It
231 exhibited a smaller range of fluctuations over time. Correspondingly, a dramatic elevation in P application
232 rate was found among various crops from 1940 to 1980, with increments ranging from 0.5 g P m⁻² yr⁻¹ in
233 durum wheat to 2.4 g P m⁻² yr⁻¹ in corn (Fig. 2b). From 1980 to 2020, large decreases in application rates
234 were found in corn, winter wheat, sorghum, and cropland pasture, while large increases were found in
235 spring wheat, rice, and durum wheat. As an increasing proportion of total cropland received P fertilizer
236 from 1940 to 2022, the gap between P fertilizer use rate that on all cropland and on fertilized area has
237 been narrowing for most crops except for soybean and cropland pasture.

238 Geospatially, as the P fertilizer consumption declined in the southeastern and eastern US and increased in
239 the Midwest and the Northern Great Plains since 1900, the hotspot of P use has shifted correspondingly
240 (Fig. 3-4). Low application rates (< 0.4 g P m⁻² yr⁻¹) were common in the eastern US before 1940. The
241 application rates in the Midwest and west coast showed remarkable increases to above 1.0 g P m⁻² yr⁻¹ by
242 1980. After 2000, the east of the Northern Great Plains and the Midwest became the US hotspots,
243 displaying the most intensive P fertilizer use.

244 The P use in the Midwest and the Northern Great Plains is dominated by the nine major crops, whereas in
245 other regions, like the Northwest, Southwest, and Northeast, Other Crops account for a considerable share
246 of P use (Fig. 4). Owing to their wide cultivation, corn and soybean are the primary recipients of P



247 nationwide in the most recent decade (the 2020s). The intense P fertilizer use is concentrated in the
248 Midwest and the Northern Great Plains for corn ($> 0.8 \text{ g P m}^{-2} \text{ yr}^{-1}$) and for soybean ($0.5\text{-}1.2 \text{ g P m}^{-2} \text{ yr}^{-1}$)
249 (Fig. 5). In comparison, the P uses of the rest seven major crops are mainly distributed in different
250 regions. Low-level of application rate ($< 0.5 \text{ g P m}^{-2} \text{ yr}^{-1}$) is applied to cotton in the Southeast and the
251 Southern Great Plains. Sorghum is planted mainly in the Southern Great Plains with application rate < 0.2
252 $\text{g P m}^{-2} \text{ yr}^{-1}$. Rice is highly concentrated along the rice-belt and part of California with a relatively high
253 application rate ($0.5\text{-}0.8 \text{ g P m}^{-2} \text{ yr}^{-1}$). P fertilizer applied to barley, spring wheat, and durum wheat is
254 distributed in the Northern Great Plains at a moderate rate ($0.3\text{-}0.8 \text{ g P m}^{-2} \text{ yr}^{-1}$). Winter wheat has a wider
255 spatial distribution with a low application rate, except for some regions in Kansas, Oklahoma, and
256 Montana ($0.3\text{-}0.5 \text{ g P m}^{-2} \text{ yr}^{-1}$).

257 3.2 Patterns of P fertilizer application timings

258 Nationwide, corn, soybean, and cotton producers favor fall and spring applications before planting.
259 Conversely, producers of all three wheats and barley apply a large portion of annual P fertilizer at
260 planting (Fig. 6). The timing of P application varies significantly across the contiguous US (Fig. 7). Fall
261 application prevails in the Midwest and the Southern Great Plains ($> 40\%$), especially in Iowa ($> 60\%$)
262 and Illinois ($> 50\%$) (Fig. 7a). Relatively high portions of P fertilizer, up to 20%, are also applied in fall
263 in the Southeast, the eastern Northern Great Plains, and the Northwest. In comparison, P applied in spring
264 before planting dominates across the nation, especially in the east of the US (Fig. 7b). Intense P
265 application ($> 50\%$) at planting is prevalent in the Northeast, the Northwest, and both the north part of the
266 Northern Great Plains and the Southern Great Plains (Fig. 7c). Application after planting is the least
267 popular application timing ($< 20\%$) in the nation, which mainly occurs in the Southern Great Plains, the
268 Southeast, and some other states (e.g., Michigan, Nebraska, and Washington) (Fig. 7d).

269 3.2 Patterns of P fertilizer application methods

270 Nationally, broadcast application is popular among corn, soybean, cotton, and rice. In contrast, the non-
271 broadcast method (e.g. injection and side-dress) dominates among three wheat types, sorghum, and barley
272 (Fig. 6). The adoption of the P application method differs substantially among regions (Fig. 8). Non-
273 broadcast is predominantly used in Wisconsin, Michigan, the Great Plains, and the Northwest (Fig. 8a).
274 Broadcast with incorporation is widespread in the contiguous US. However, the adoption rate is relatively
275 low ($< 40\%$) in most of the region (Fig. 8b). In comparison, high P application by broadcast without
276 incorporation ($> 50\%$) is mainly distributed in the Midwest and the Southeast (Fig. 8c).



277 4 Discussion

278 4.1 Adjustments and improvements in state-level crop-specific P application rate

279 The national total P consumption obtained from the gap-filled bottom-up data in this study, summed from
280 all major crops, cropland pasture, permanent pasture, and non-farm use, aligns well with diverse top-
281 down data sources both in magnitude and inter-annual variations (Fig. S3). However, the bottom-up
282 source displays a larger P consumption of certain crops in certain states (e.g., corn in Illinois),
283 contributing to the divergences between these two approaches, notably after 2010 (Fig. S1&S2). These
284 overestimations may be caused by distorted crop-specific P application rate and/or fertilized area
285 percentage, derived from an inadequate survey pool. By modifying the surveyed crop-specific P
286 application rate at the state level, we matched the state total P consumption between bottom-up and top-
287 down approaches (Fig. 4). Despite the bottom-up source offering insights into cross-crop variations of P
288 application rate, it overlooks the inter-state variability. Based on the total P consumption and crop-
289 specific planting area in each county, we scaled the P application rate of each crop from state level to
290 county level, which portrays greater variability across counties. Particularly, the ranges are wider for corn,
291 soybean, winter wheat, sorghum, and barley ($0\text{--}6\text{ g P m}^{-2}\text{ yr}^{-1}$) than those for spring wheat, cotton, rice,
292 durum, cropland pasture, and Other Crops (Fig. 9). In addition, downscaling state-level P application rate
293 to the county level augments the clarity of the geospatial pattern (Fig. 10). Top-down sources calculated
294 average P use rate in each county by dividing the total P consumption by all cropland areas, yielding in a
295 uniform value within each county but contrasting patterns across counties (Fig. 10a, d, g). Conversely,
296 our map based on bottom-up sources at the state level detailed spatial heterogeneity in intensive
297 agricultural regions, highlighting the cross-crop differences in P fertilizer use (Fig. 10b, e, h). By
298 combining these two sources, our map characterizes spatial variability across counties and crop types
299 (Fig. 10c, f, j). It highlights the region with intense P use, indicated by the top-down source, but also
300 differentiates P application rates among crops within each county, indicated by the bottom-up source.
301 This is particularly evident in the southern part of Missouri and the boundary between Minnesota and
302 Dakotas (Fig. 10c&j). Accurate information on fertilizer management is essential for improving
303 agricultural sustainability (Dhillon et al., 2017). Different crops have distinct P needs, and tailoring P use
304 based on these needs can enhance the efficiency of P fertilizer utilization, maximizing crop yield while
305 mitigating environmental impacts (Sabo et al., 2021). Moreover, detailed information on crop-specific P
306 fertilizer management is important for assessing P losses attributed to runoff, erosion, and leaching,
307 contributing to the development of agricultural policies (Daloğlu et al., 2012). Given the significance of



308 crop-specific information, we advocate for the incorporation of cross-crop variations into the
309 development of P fertilizer datasets.

310 4.2 Temporal and spatial dynamics of P fertilizer management

311 Concurrent with the historical changes in US cropland since 1850, P use has experienced different stages
312 of change similar to nitrogen fertilizer use (Cao et al, 2018), influenced by various factors. From 1850 to
313 1940, the primary crops, corn, cotton, and winter wheat, were mainly concentrated in the eastern US. The
314 constrained production of phosphate rock and low demand by limited crop productivity contributed to the
315 low level of P consumption and application rate. As cropland expanded to the Midwest and the Great
316 Plains from 1940 to 1980, the consumption of P fertilizer peaked after a sharp increase, driven by the
317 rising application rate and percentage of fertilized area across various crops (Fig. 2-5). The major
318 contributors to P consumption during this period were corn in the Midwest and spring wheat and winter
319 wheat in the Great Plains. Following a brief decline in the 1980s, P consumption has stabilized with
320 fluctuations impacted by changes in grain demand and fertilizer prices (US-EPA, 2024). Throughout this
321 period, P consumption continued to decline in the eastern US while increasing or leveling off in other
322 regions, driven by the continued expansion of corn and soybean at the expense of other crops (Fig. 2-5).
323 Another possible contributing factor to the decline in P consumption is that the generous high-rate P
324 application over a half-century has raised soil P level so much that it made it possible to have lower
325 application and still meet crop demands (Sabo et al., 2021; Bian et al., 2022).

326 In the past decade, the average percentage of P fertilized area in the US was around 60% (including
327 cropland and pasture), notably lower than that for nitrogen fertilizer. (Fig. S4). The percentage of
328 fertilized area varies among crops, ranging from 42% for soybean to 89% for spring wheat. Estimating P
329 use efficiency and P losses in agricultural systems highly rely on the precise application rate of P fertilizer
330 (Solangi et al., 2023). It is noteworthy that the crops with lower fertilized area percentages, such as
331 soybean, cotton, and sorghum, may impact the estimate of these application rate-sensitive assessments
332 without considering the fertilized area percentage.

333 Despite the application of P fertilizer after planting is strongly recommended for improving P fertilizer
334 use efficiency and minimizing P losses to the environment, this application timing remains the least
335 popular choice for major crops in the US. Notably, rice in the US rice belt, sorghum in the Southern Great
336 Plains, and cotton along the southwest coast were major contributors to post-planting applications. In
337 contrast, both fall and spring applications before planting, leaving P susceptible to loss (King et al., 2018),
338 have been widely adopted across multiple crops in the contiguous US due to lower fertilizer prices, the
339 availability of labor, and the ease of operating equipment (Carver et al., 2022). Winter wheat in the



340 Southern Great Plain and the Northwest received over 40% of its annual P fertilizer in the fall, potentially
341 contributing to boosting yield. However, corn and soybean farmers in the Midwest, cotton farmers in the
342 Southwest and north of Texas, and sorghum farmers in the Southern Great Plains favor fall application,
343 implying a high potential risk for P loss (Nelson et al., 2023; Yuan et al., 2013). Except for winter wheat,
344 spring wheat, and durum wheat, all other crops receive more than a quarter of their annual P fertilizer in
345 spring before application. Despite being closer to the planting date, the P fertilizer applied during early
346 spring may be prone to loss via runoff, erosion, and leaching during intense rainfall (Williams and King,
347 2020; Algoazany et al., 2007). Application at planting is more prevalent among winter wheat and spring
348 wheat in the Southern Great Plains and the Northern Great Plains, respectively.

349 Non-broadcast application is commonly found for winter wheat, durum wheat, and barley in the
350 Northwest and Northern Great Plains, and for spring wheat, cotton, and sorghum in the Southern Great
351 Plains. In addition, corn farmers in Wisconsin, Michigan, and the Northeast apply most of their annual P
352 fertilizer using the non-broadcast method. The non-broadcast has been considered as a more conservative
353 management to prevent P loss (Carver et al., 2022; Smith et al., 2016). However, broadcasting, including
354 post-incorporation and non-incorporation, remains widespread across the US, particularly in the Midwest
355 (hotspot for P fertilizer use) and the Southeast.

356 4.3 Uncertainty

357 The uncertainties of this database are mainly from several aspects: (1) Limited information on P use in
358 cropland pasture and permanent pasture at finer temporal and spatial resolution, contributing to uncertain
359 estimates for Other Crops; (2) Adjustments were made on crop-specific P fertilizer use rates at the state
360 level to reconcile top-down and bottom-up data sources. However, the paucity of detailed crop-specific
361 information may introduce biases in our adjustments made for certain crops; (3) The composition of the
362 Other Crops differs across states. All crop types under Other Crops within each state receive equal P
363 application rate, which may bias the application rate for some crop types; (4) Due to the lack of finer
364 spatial resolution information, we assumed the crop-specific P application timing and method are
365 identical within each state. However, the spatial heterogeneity of application timing and method may be
366 overlooked. Therefore, a finer resolution of spatial and temporal survey capturing crop-specific P
367 application rate, timing, and method will be invaluable for enhancing our understanding of the
368 spatiotemporal patterns of P fertilizer management information in the US.



369 5 Data availability

370 The P fertilizer management dataset is publicly available via ZENODO at
371 <https://doi.org/10.5281/zenodo.10700822> (Cao et al., 2024).

372 6 Conclusion

373 By harmonizing various data sources, we reconstructed a long-term spatially explicit P fertilizer
374 management dataset at 4 km ×4 km resolution from 1850 to 2022 in the contiguous US. We discussed the
375 divergence between top-down (total P consumption) and bottom-up (crop-specific P fertilizer use) data
376 sources, underscoring the necessity to improve crop-specific management information in future surveys.
377 The newly developed dataset, leveraging the strengths of both data sources, highlights cross-crop
378 variabilities in the long-term use of P fertilizer among counties. The results reveal a substantial increase in
379 P fertilizer consumption and application rate from 1850 to 2022, notably during 1940-1980. However, the
380 magnitude and long-term changing trend differed significantly across crop types. It is worth noting that
381 approximately 40% of cropland in the US does not receive P fertilizer inputs. Since 1850, the hotspots of
382 P fertilizer use have shifted from the southeastern and eastern US to the Midwest and the Great Plains,
383 driven by changes in cropland distribution and P fertilizer application rate across different crop types.
384 Additionally, P fertilizer application timing and method vary substantially across crop types and regions.
385 Corn, soybean, and cotton in the Midwest and the Southeast receive over 60% of their annual P fertilizer
386 at pre-planting and through broadcasting. Conversely, winter wheat, spring wheat, durum wheat, and
387 barley in the Great Plains and the Northwest predominantly receive their annual P fertilizer at- and post-
388 planting, and via non-broadcasting. Promoting efficient P fertilizer management, encompassing the proper
389 application rate, timing, and method, is essential for enhancing P use efficiency and thus contributes to
390 economic, social, and environmental sustainability and profitability.

391 Author contributions

392 CL, PC, and BY conceptualized the paper and developed the methodology. PC and BY reconstructed the
393 dataset. PC and BY prepared the manuscript with contributions from all the co-authors.

394 Competing interests

395 At least one of the (co-)authors is a member of the editorial board of Earth System Science Data.

396 Acknowledgments



397 This work is supported by the Iowa Nutrient Research Center, the ISU College of Liberal Arts and
398 Sciences Dean's Faculty Fellowship, and the NSF CAREER grant (1945036).

399 References

400 Alexander, R. B. and Smith, R. A.: County-level estimates of nitrogen and phosphorus fertilizer use in the
401 United States, 1945 to 1985, US Department of the Interior, US Geological Survey, 1990.

402 Algoazany, A. S., Kalita, P. K., Czapar, G. F., and Mitchell, J. K.: Phosphorus Transport through
403 Subsurface Drainage and Surface Runoff from a Flat Watershed in East Central Illinois, USA, *J Environ*
404 *Qual*, 36, 681–693, <https://doi.org/https://doi.org/10.2134/jeq2006.0161>, 2007.

405 Association of American Plant Food Control Officials (AAPFCO): Commercial Fertilizers, available at:
406 <http://www.aapfco.org/publications.html>, last access: 20 December 2021, 2022.

407 Bian, Z., Pan, S., Wang, Z., Yao, Y., Xu, R., Shi, H., Kalin, L., Anderson, C., Justic, D., Lohrenz, S., and
408 Tian, H.: A Century-Long Trajectory of Phosphorus Loading and Export From Mississippi River Basin to
409 the Gulf of Mexico: Contributions of Multiple Environmental Changes, *Global Biogeochem Cycles*, 36,
410 e2022GB007347, <https://doi.org/https://doi.org/10.1029/2022GB007347>, 2022.

411 Bouwman, L., Goldewijk, K. K., Van Der Hoek, K. W., Beusen, A. H. W., Van Vuuren, D. P., Willems,
412 J., Rufino, M. C., and Stehfest, E.: Exploring global changes in nitrogen and phosphorus cycles in
413 agriculture induced by livestock production over the 1900-2050 period., *Proc Natl Acad Sci U S A*, 110,
414 20882–7, <https://doi.org/10.1073/pnas.1012878108>, 2013.

415 Brakebill, J. W. and Gronberg, J. M.: County-level estimates of nitrogen and phosphorus from
416 commercial fertilizer for the conterminous United States, 1987-2012, US Geological Survey Data release,
417 2017.

418 Cao, P., Lu, C., and Yu, Z.: Historical nitrogen fertilizer use in agricultural ecosystems of the contiguous
419 United States during 1850–2015: application rate, timing, and fertilizer types, *Earth Syst Sci Data*, 10,
420 969–984, <https://doi.org/10.5194/essd-10-969-2018>, 2018.

421 Cao, P., Yi, B., Bilotto, F., Gonzalez Fischer, C., Herrero, M., and Lu, C.: Annual crop-specific
422 management history of phosphorus fertilizer input (CMH-P) in the croplands of United States from 1850
423 to 2022: Application rate, timing, and method, *Zenodo*, <https://doi.org/10.5281/zenodo.10700822>, 2024.

424 Carver, R. E., Nelson, N. O., Roozeboom, K. L., Kluitenberg, G. J., Tomlinson, P. J., Kang, Q., and Abel,
425 D. S.: Cover crop and phosphorus fertilizer management impacts on surface water quality from a no-till



- 426 corn-soybean rotation, *J Environ Manage*, 301, 113818,
427 <https://doi.org/https://doi.org/10.1016/j.jenvman.2021.113818>, 2022.
- 428 Cordell, D., Drangert, J. O., and White, S.: The story of phosphorus: Global food security and food for
429 thought, *Global Environmental Change*, 19, 292–305, <https://doi.org/10.1016/j.gloenvcha.2008.10.009>,
430 2009.
- 431 Daloğlu, I., Cho, K. H., and Scavia, D.: Evaluating Causes of Trends in Long-Term Dissolved Reactive
432 Phosphorus Loads to Lake Erie, *Environ Sci Technol*, 46, 10660–10666,
433 <https://doi.org/10.1021/es302315d>, 2012.
- 434 Dhillon, J., Torres, G., Driver, E., Figueiredo, B., and Raun, W. R.: World Phosphorus Use Efficiency in
435 Cereal Crops, *Agron J*, 109, 1670–1677, <https://doi.org/https://doi.org/10.2134/agronj2016.08.0483>,
436 2017.
- 437 Falcone, J. A.: Estimates of county-level nitrogen and phosphorus from fertilizer and manure from 1950
438 through 2017 in the conterminous United States, Open-File Report, Reston, VA, 20 pp.,
439 <https://doi.org/10.3133/ofr20201153>, 2021.
- 440 FAO (Food and Agriculture Organization of the United Nations): FAO online database, available at:
441 <http://www.fao.org/faostat/en/#data/RF>, last access: 10 August 2021, 2021.
- 442 Glibert, P. M.: From hogs to HABs: impacts of industrial farming in the US on nitrogen and phosphorus
443 and greenhouse gas pollution, Springer International Publishing, 139–180 pp.,
444 <https://doi.org/10.1007/s10533-020-00691-6>, 2020.
- 445 King, K. W., Williams, M. R., LaBarge, G. A., Smith, D. R., Reutter, J. M., Duncan, E. W., and Pease, L.
446 A.: Addressing agricultural phosphorus loss in artificially drained landscapes with 4R nutrient
447 management practices, *J Soil Water Conserv*, 73, 35, <https://doi.org/10.2489/jswc.73.1.35>, 2018.
- 448 Lu, C. and Tian, H.: Global nitrogen and phosphorus fertilizer use for agriculture production in the past
449 half century: Shifted hot spots and nutrient imbalance, *Earth Syst Sci Data*, 9, 181–192,
450 <https://doi.org/10.5194/essd-9-181-2017>, 2017.
- 451 Mehring, A. L., Adams, J. R., and Jacob, K. D.: Statistics on Fertilizers and Liming Materials in the
452 United States, USDA-Agricultural Research Service, Statistical Bulletin No. 191, Washington, D.C.,
453 USA, 1957.
- 454 Nelson, N. O., Roozeboom, K. L., Yeager, E. A., Williams, J. R., Zerger, S. E., Kluitenberg, G. J.,
455 Tomlinson, P. J., Abel, D. S., and Carver, R. E.: Agronomic and economic implications of cover crop and



- 456 phosphorus fertilizer management practices for water quality improvement, *J Environ Qual*, 52, 113–125,
457 <https://doi.org/https://doi.org/10.1002/jeq2.20427>, 2023.
- 458 Nutrient Use Geographic Information System (NuGIS): No Title, available at: <https://nugis.tfi.org/>, last
459 access: 20 December 2022, 2022.
- 460 Sabo, R. D., Clark, C. M., Gibbs, D. A., Metson, G., Todd, M. J., LeDuc, S. D., Greiner, D., Fry, M. M.,
461 Polinsky, R., Yang, Q., Tian, H., and Compton, J. E.: Phosphorus Inventory for the Conterminous United
462 States (2002-2012), *J Geophys Res Biogeosci*, n/a, e2020JG005684,
463 <https://doi.org/https://doi.org/10.1029/2020JG005684>, 2021.
- 464 Samreen, S.: Phosphorus Fertilizer: The Original and Commercial Sources, edited by: Zhang, S. K. E.-T.,
465 IntechOpen, Rijeka, Ch. 6, <https://doi.org/10.5772/intechopen.82240>, 2019.
- 466 Sharpley, A., Jarvie, H. P., Buda, A., May, L., Spears, B., and Kleinman, P.: Phosphorus Legacy:
467 Overcoming the Effects of Past Management Practices to Mitigate Future Water Quality Impairment, *J*
468 *Environ Qual*, 42, 1308–1326, <https://doi.org/https://doi.org/10.2134/jeq2013.03.0098>, 2013.
- 469 Smith, D. R., Harmel, R. D., Williams, M., Haney, R., and King, K. W.: Managing Acute Phosphorus
470 Loss with Fertilizer Source and Placement: Proof of Concept, *Agricultural & Environmental Letters*, 1,
471 150015, <https://doi.org/https://doi.org/10.2134/aer2015.12.0015>, 2016.
- 472 Solangi, F., Zhu, X., Khan, S., Rais, N., Majeed, A., Sabir, M. A., Iqbal, R., Ali, S., Hafeez, A., Ali, B.,
473 Ercisli, S., and Kayabasi, E. T.: The Global Dilemma of Soil Legacy Phosphorus and Its Improvement
474 Strategies under Recent Changes in Agro-Ecosystem Sustainability, *ACS Omega*, 8, 23271–23282,
475 <https://doi.org/10.1021/acsomega.3c00823>, 2023.
- 476 Stackpole, S. M., Stets, E. G., and Sprague, L. A.: Variable impacts of contemporary versus legacy
477 agricultural phosphorus on US river water quality, *Proc Natl Acad Sci U S A*, 116, 20562–20567,
478 <https://doi.org/10.1073/pnas.1903226116>, 2019.
- 479 Swaney, D. P. and Howarth, R. W.: Phosphorus use efficiency and crop production: Patterns of regional
480 variation in the United States, 1987–2012, *Science of the Total Environment*, 685, 174–188,
481 <https://doi.org/10.1016/j.scitotenv.2019.05.228>, 2019.
- 482 Tilman, D., Cassman, K., Matson, P., Naylor, R., and Polasky, S.: Agricultural sustainability and
483 intensive production practices, *Nature* 418, 671–677, <https://doi.org/10.1038/nature01014>, 2002.
- 484 U.S. Fourth National Climate Assessment: No Title, available at: <http://www.globalchange.gov/nca4>, last
485 access: 20 December 2022, 2022.



486 USDA (U.S. Department of Agriculture): Consumption of Commercial Fertilizers, Primary Plant
487 Nutrients, and Micronutrients, 1850–1969, USDA-Statistical Reporting Service, Crop Reporting Board,
488 Statistical Bulletin No. 472, Washington, D.C., USA, 1971.

489 Tailored Reports: Crop Production Practices: <https://data.ers.usda.gov/reports.aspx?ID=17883>.

490 USDA-ERS (U.S. Department of Agriculture-Economic Research Service): Fertilizer Use and Price,
491 available at: <https://www.ers.usda.gov/data-products/arms-farm-financial-and-cropproduction-practices/>
492 (last access: 10 August 2021), 2019.

493 USDA-NASS (U.S. Department of Agriculture-National Agricultural Service), S.: Agricultural Chemical
494 Use Program, available at: [https://www.nass.usda.gov/Surveys/Guide_](https://www.nass.usda.gov/Surveys/Guide_to_NASS_Surveys/Chemical_Use/index.php)
495 [to_NASS_Surveys/Chemical_Use/index.php](https://www.nass.usda.gov/Surveys/Guide_to_NASS_Surveys/Chemical_Use/index.php), last access: 17 August 2021, 2021.


496 Williams, M. R. and King, K. W.: Changing Rainfall Patterns Over the Western Lake Erie Basin (1975–
497 2017): Effects on Tributary Discharge and Phosphorus Load, *Water Resour Res*, 56, e2019WR025985,
498 <https://doi.org/https://doi.org/10.1029/2019WR025985>, 2020.

499 Yuan, Y., Locke, M. A., Bingner, R. L., and Rebich, R. A.: Phosphorus losses from agricultural
500 watersheds in the Mississippi Delta, *J Environ Manage*, 115, 14–20,
501 <https://doi.org/https://doi.org/10.1016/j.jenvman.2012.10.028>, 2013.

502 Zhang, J., Gilbert, D., Gooday, A. J., Levin, L., Naqvi, S. W. A., Middelburg, J. J., Scranton, M., Ekau,
503 W., Peña, A., Dewitte, B., Oguz, T., Monteiro, P. M. S., Urban, E., Rabalais, N. N., Ittekkot, V., Kemp,
504 W. M., Ulloa, O., Elmgren, R., Escobar-Briones, E., and Van der Plas, A. K.: Natural and human-induced
505 hypoxia and consequences for coastal areas: synthesis and future development, *Biogeosciences*, 7, 1443–
506 1467, <https://doi.org/10.5194/bg-7-1443-2010>, 2010.


507 Zhang, J., Cao, P., and Lu, C.: Half-Century History of Crop Nitrogen Budget in the Conterminous
508 United States: Variations Over Time, Space and Crop Types, *Global Biogeochem Cycles*, 35,
509 e2020GB006876, <https://doi.org/https://doi.org/10.1029/2020GB006876>, 2021.

510






512 Figure 1. Diagram for P fertilizer management dataset development. The upper blue box represents the
 513 development of state-level crop-specific P fertilizer application rate based on the bottom-up dataset. The
 514 lower green box represents the development of county-level P fertilizer application rate development by
 515 reconciling the top-down and bottom-up dataset.



516
 517 Figure 2. Time-series of P fertilizer consumption and P fertilizer application rates for all crops (a), and for
 518 11 specific crops (b), and P fertilizer consumption shares across 11 crops (c) in the contiguous US. All
 519 cropland is the total planting area, while the fertilized area is the proportion of the cropland that receives
 520 P fertilizer. In panel (b), light-colored bars denote the application rate on fertilized area and dark-colored



521 bars show the application rate on all cropland.




522


523 Figure 3. Spatial distribution of P fertilizer application rates in the 1990s, 1940s, 1960s, 1980s, 2000s,
524 and 2020s in the contiguous US at a resolution of 4-km x 4-km, with regions framed as NW (Northwest),



525 NGP (Northern Great Plains), SGP (Southern Great Plains), SW (Southwest), MW (Midwest), SE
 526 (Southeast), and NE (Northeast). The maps generated for 1900, 1940, and 1960 relied on state-level crop-
 527 specific data. Subsequent maps, post-1960, utilized county-level crop-specific data.




528
 529 Figure 4. Time-series of P fertilizer consumption by each state and 9 major crops from 1950 to 2022 in
 530 the contiguous US. The top-left figure illustrates the scales of x-axis and y-axis. The solid black line in
 531 each subplot represents total P fertilizer consumption, and the stacked area represents P fertilizer
 532 consumption by different crops. NW is the Northwest, NGP is the Northern Great Plains, SGP is the
 533 Southern Great Plains, SW is the Southwest, MW is the Midwest, SE is the Southeast, NE is the
 534 Northeast.



535


536 Figure 5. Spatial distribution of P fertilizer application rates for 9 major crops in 2020 at 4-km x 4-km
537 resolution, with regions framed as NW (Northwest), NGP (Northern Great Plains), SGP (Southern Great
538 Plains), SW (Southwest), MW (Midwest), SE (Southeast), and NE (Northeast).



539

540 Figure 6. The share of each application timing and method for 9 major crops in the US. FAL is fall
541 application in previous year. SPR is spring application before planting. ATP is application at planting.
542 AFP is application after planting. NBC is non-broadcast. BWJ is broadcast with injection, which is mix or
543 inject after broadcast. BNJ is broadcast with no injection.

544



545

546 Figure 7. Spatial distribution of the fractions of four P fertilizer application timings in the contiguous US.


547

548

549

550


551



552

553 Figure 8. Spatial distribution of the fractions of three P fertilizer application methods in the contiguous
554 US.


555





557 Figure 9. Comparison between state-level (red line) and county-level average (black boxplot)
558 crop-specific P fertilizer application rate in primary crop-planting states in 2015. The red line indicates the
559 state-level P fertilizer application rate. The box plot shows the distribution of county-level P fertilizer
560 application rate (dots are outliers).

561



562

563 Figure 10. Comparison of spatial distribution of P fertilizer application rate in the contiguous US in 2016.
564 NuGIS (a, d, g) represents the average application rate derived from county-level sales data. State (b, d, h)
565 and county (c, f, i) data used for plotting represent the crop-specific P fertilizer application rate at state-
566 and county-level developed in this study, respectively. To make it comparable, the same cropland map
567 was used to mask out the cropland extent for NuGIS. Two red boxes in Fig d were zoomed in to
568 demonstrate more details in the top and bottom panels.