

Imputation of missing ~~IPCC AR6 data on~~ land carbon sequestration data in the AR6 Scenario Database

Ruben Prütz^{1, 2, 3}, Sabine Fuss^{1, 2}, Joeri Rogelj^{3, 4, 5}

5 ¹Geography Department, Humboldt-Universität zu Berlin, Berlin, Germany

²Mercator Research Institute on Global Commons and Climate Change (MCC), Berlin, Germany

³Grantham Institute for Climate Change and the Environment, Imperial College London, London, United Kingdom

⁴Centre for Environmental Policy, Imperial College London, London, United Kingdom

⁵Energy, Climate and Environment Program, International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria

10 *Correspondence to:* Ruben Prütz (Pruetz@mcc-berlin.net)

Abstract. The AR6 Scenario Database is a vital repository of climate change mitigation pathways used in the latest IPCC assessment cycle. In its current version, ~~several~~many scenarios in the database lack information about the level of ~~gross anthropogenic~~ carbon ~~dioxide~~ removal ~~on~~via land sinks, as net negative CO₂ emissions and gross removals on land are not always separated and consistently reported across models. This makes scenario analyses focusing on carbon ~~removals~~dioxide removal challenging. We test and compare the performance of different regression models to impute missing data on land carbon sequestration for the global level and for R10 regions from available data on net CO₂ emissions in agriculture, forestry, and other land use. We find that a ~~gradient boosting~~k-nearest neighbors regression performs best among the tested regression models and provide two publicly available imputation ~~dataset~~datasets [<https://doi.org/10.5281/zenodo.10696654>] (Prütz et al., 2024) on carbon ~~dioxide~~ removal ~~on~~via land sinks for 404-incomplete global scenarios (n=404) and incomplete regional R10 scenario variants (n=2358) in the AR6 Scenario Database. We discuss the limitations of our approach, ~~it~~the use ~~of~~our datasets for secondary assessments of AR6 scenario ensembles, and how this approach compares to other recent AR6 data ~~re-analyses~~reanalyses.

1 Introduction

25 ~~Climate change mitigation pathways, created with integrated assessment models (IAMs), have come to take up a critical role in the assessment work of Working Group III of the Intergovernmental Panel on Climate Change (IPCC) (Riahi et al., 2022; Guivareh et al., 2022). The AR6 Scenario Database hosted by the International Institute for Applied Systems Analysis (IIASA) contains climate change mitigation pathways compiled for and considered in the Working Group III Contribution to the IPCC Sixth Assessment Report (Byers et al., 2022; Kikstra et al., 2022).~~

30 ~~In these pathways, carbon dioxide removal (CDR) from the atmosphere is primarily represented by bioenergy with carbon capture and storage (BECCS) and by carbon sequestration on land—primarily via afforestation and reforestation (Riahi et al., 2022). Among the scenarios in the AR6 Scenario Database that passed the vetting process (n=1202) (see Guivareh et al. (2022)~~

for details about the AR6 scenario vetting process), 419 pathways miss the variable for carbon sequestration on land ('Carbon Sequestration|Land Use'), which complicates secondary analyses that investigate CDR implications across scenarios and models. This gap requires the use of proxy data and interim solutions. Two such interim solutions to account for this data gap are documented in the literature, including the use of net negative CO₂ emissions in agriculture, forestry, and other land use (AFOLU) as a proxy variable for land-based CDR (Warszawski et al., 2021; Schleussner et al., 2022; Prütz et al., 2023) or scenario filtering and exclusion (Prütz et al., 2023). Both approaches have limitations in depicting gross carbon removals on land adequately and comprehensively (Ganti et al., 2024). A more recent approach is based on a re-analysis of land CO₂ fluxes using the earth system model OSCAR v3.2 (Gidden et al., 2023). While the AR6 re-analysis dataset by Gidden et al. manages to resolve several of the data issues linked to carbon removal on land, it still combines gross and net CO₂ emissions on land in their land-based CDR variable, resulting in both positive and negative CDR values, which conflicts with the concept and clean definition of gross CDR. Also, while being very comprehensive, the re-analyzed dataset by Gidden et al. is limited to a subset (n=914) of all global and vetted scenarios (n=1202) of the AR6 Scenario Database. Figure 1 compares the available land carbon sequestration data of the AR6 Scenario Database to the re-analyzed variable by Gidden et al. and the net negative AFOLU CO₂ proxy, showing the discrepancy of the net negative AFOLU CO₂ proxy and the negative values for land-based CDR of the re-analysis.

Here, we test and compare the performance of several different regression models to impute missing data on gross land carbon sequestration. Climate change mitigation pathways, created with integrated assessment models (IAMs), have come to take up a critical role in the assessment work of Working Group III of the Intergovernmental Panel on Climate Change (IPCC) (Guivarch et al., 2022b; Riahi et al., 2022). The AR6 Scenario Database hosted by the International Institute for Applied Systems Analysis (IIASA) contains climate change mitigation pathways compiled for and considered in the Working Group III Contribution to the IPCC Sixth Assessment Report (Byers et al., 2022; Kikstra et al., 2022).

In these pathways, carbon dioxide removal (CDR) from the atmosphere is primarily represented by bioenergy with carbon capture and storage (BECCS) and by carbon sequestration in land sinks – primarily via afforestation and reforestation (Riahi et al., 2022). Among the global scenarios in the AR6 Scenario Database that passed the vetting process (n=1202) (see Guivarch et al. (2022b) for details about the AR6 scenario vetting process), 419 pathways miss the variable for carbon sequestration on land ('Carbon Sequestration|Land Use'), which complicates secondary analyses that investigate CDR implications across scenarios and models. A range of different secondary scenario ensemble evaluations based on data from the AR6 Scenario Database have been published in recent years, e.g., assessing the arising gap in CDR deployment (Lamb et al., 2024), determining the level and composition of residual emissions (Lamb, 2024), analysing the removal per land unit (Zhao et al., 2024), evaluating the attainability of mitigation scenarios (Warszawski et al., 2021), classifying emission pathways reflecting the climate objectives of the Paris Agreement (Schleussner et al., 2022), or exploring scenario characteristics driving CDR deployment (Prütz et al., 2023). All these analyses rely on proxy data or interim solutions to address the limited data availability of land carbon sequestration in the AR6 Scenario Database.

Two such interim solutions to account for this data gap are documented in the literature, including the use of net negative CO₂ emissions in agriculture, forestry, and other land use (AFOLU) as a lower bound proxy variable for CDR via land sinks (Prütz et al., 2023; Schleussner et al., 2022; Warszawski et al., 2021) or criteria-based scenario filtering and exclusion to ensure a consistent selection of scenarios with similar reporting of CDR via land sinks (Prütz et al., 2023). Both approaches have limitations in depicting CDR via land sinks adequately and comprehensively (Ganti et al., 2024). A more recent approach is based on a reanalysis of land CO₂ fluxes using the reduced-complexity compact earth system model OSCAR v3.2 (Gidden et al., 2023). While the AR6 reanalysis dataset by Gidden et al. manages to resolve several of the data issues linked to CDR via land sinks – specifically, aligning the removal baseline and improving the consistency across scenarios – it still combines gross and net CO₂ fluxes on land in their land sink CDR variable, resulting in both positive and negative CDR values, which conflicts with the concept and clean definition of anthropogenic CDR from the atmosphere (Matthews et al., 2021). In the AR6 Scenario Database, CDR is conventionally reported in positive numbers. Also, while being very comprehensive, the reanalyzed dataset by Gidden et al. is limited to a subset (n=914) of all global and vetted scenarios (n=1202) of the AR6 Scenario Database, while also providing reanalysed scenario data for five sub-global macro regions (R5 level). Figure 1 compares the available land sink CDR data of the AR6 Scenario Database to the reanalyzed variable by Gidden et al. and the net-negative AFOLU CO₂ proxy, showing the discrepancy of the net-negative AFOLU CO₂ proxy and the negative values for land sink CDR of the reanalysis.

Here, we test and compare the performance of several different regression models to impute missing data on land carbon sequestration (Land CDR) based on available data on net CO₂ emissions in AFOLU– for both global scenarios and the R10 regions in the AR6 Scenario Database. We use the best performing regression model to impute missing data for 404 global scenarios and 2358 sub-global scenario variants across the R10 regions and provide two imputation datasets, which are made publicly available. Lastly, we discuss our approach's use cases and limitations and detail how our approach compares to the two above-mentioned interim solutions and the recent re-analysis of the AR6 land carbon removal data. reanalysis of the AR6 Land CDR data. In the following, we refer to CDR via land sinks or carbon sequestration on land as Land CDR. Table 1 gives an overview and description of key variables in this analysis.

90





Table 1. Overview of the analysis variables

<u>Variable</u>	<u>Description</u>
<u>'Carbon Sequestration Land Use'</u>	<u>This variable from the AR6 Scenario Database is defined as the “total carbon dioxide sequestered through land sinks (e.g., afforestation, soil carbon enhancement, biochar)”. This is the target variable that we impute for incomplete scenarios. In this analysis, we refer to this variable as AR6 Land CDR.</u>
<u>'AR6 Reanalysis OSCARv3.2 Carbon Removal Land Direct'</u>	<u>This variable from the reanalysis by Gidden et al. is intended to depict CDR through land sinks, similar to the AR6 Land CDR. However, the baseline CO₂ flux substantially differs compared to the AR6 Land CDR, as the data was aligned to national greenhouse gas inventories. This variable contains both positive and negative values, which suggests that it is showing net instead of gross removal. In this analysis, we refer to this variable as Gidden et al. Land CDR (direct).</u>
<u>'Emissions CO₂ AFOLU'</u>	<u>This variable from the AR6 Scenario Database is defined as the net “CO₂ emissions from agriculture, forestry and other land use (IPCC category 3)”. This is the predictor variable that we use to predict the target variable. In this analysis, we refer to this variable as net AFOLU CO₂ emissions.</u>
<u>'Emissions CO₂ AFOLU' < 0 </u>	<u>This variable shows the net CO₂ removal from agriculture, forestry and other land use, based on the negative values in the variable net AFOLU CO₂ emissions. This variable has been used in several studies as lower bound proxy for Land CDR. We refer to this variable as net negative AFOLU CO₂.</u>
<u>'Imputed Carbon Sequestration Land Use'</u>	<u>This is one of two variables in the imputation datasets provided in this analysis. This variable contains the predicted values from our data imputation without further adjustment.</u>



95 **Figure 1:** Comparison of available AR6 land-based Land CDR data ('Carbon Sequestration|Land Use') with the land carbon removal reanalysis-Land CDR reanalysis by Gidden et al. ('AR6 Reanalysis|OSCARv3.2|Carbon Removal|Land|Direct') and the AR6 net negative AFOLU CO₂ emissions (based on negative values in 'Emissions|CO₂|AFOLU') as a conservative lower bound proxy for land-based Land CDR across AR6 scenario categories. Only scenarios available for all three variables were considered in the figure (scenarios n=725). The Land CDR scenarios in the reanalysis by Gidden et al. are aligned with national greenhouse gas inventories, shown by the difference in baseline in 2020 compared to the other two variables. The solid lines show the median across scenarios while the shaded area shows the min-max range. Note: We follow the convention of the AR6 Scenario Database, to report CDR in positive numbers.

2 Methods

105 **Overview.** In our analysis, we used different regression models to predict missing AR6 data on gross land carbon sequestration (dependent Land CDR (target variable: 'Carbon Sequestration|Land Use')) for 404 global scenarios and 2358 sub-global scenario variants across R10 regions based on available scenario data on AFOLU CO₂ emissions (independent predictor variable: 'Emissions|CO₂|AFOLU'). AFOLU CO₂ emissions were chosen as predictor variable due to good data availability in the AR6 Scenario Database and because this variable is conceptually most closely related to Land CDR among the variables

110 in the AR6 Scenario Database – the variable for AFOLU CO₂ emissions represents the net emission fluxes corresponding to the gross variable for Land CDR. The AR6 R10 region classification comprises 10 macro regions plus one additional region for “rest of the world”, as shown in Figure 3a. As an initial step, we selected all vetted scenarios from the AR6 Scenario Database for which both the ~~independent~~predictor and the ~~dependent~~target variable are available at the global level (n=783) and across the R10 regions (n=6162). Among the vetted global scenarios (n=1202) in the AR6 Scenario Database, 15 scenarios from the model REMIND 1.6 do not report AFOLU CO₂ emissions, which is why we could not include these scenarios in our imputation. Among the vetted regional scenario variants across the R10 regions (n=8531) in the AR6 Scenario Database, 11

115 regional variants of scenario EN_INDCi2100 from the model GEM-E3 V2021 do not report AFOLU CO₂ emissions, which is why we could not include these scenario variants in our imputation.

We ~~then~~ split ~~this dataset~~ both the global and the regional scenario datasets into training and testing sets (9:1) for our regression analysis: to have a large dataset for training the models while still having a sufficiently large testing dataset to evaluate the prediction performance and to validate the models. The training set was used to fit the ~~dependent~~predictor variable to the ~~independent~~target variable to train the regression ~~model~~models, and the testing set was then used to evaluate the prediction performance of the trained regression ~~model~~models. The regression models were separately trained on the global scenario data and the regional scenario variants as the scale of Land CDR deployment differs substantially between the global and the regional level. Regional scenario variants for model training were not further split into training bins depending on their respective R10 region, and we did not split scenarios into different training bins based on their AR6 scenario categories to

125 keep the number of training data as large as possible to optimize the models’ performances. The AR6 scenario category (C1-8) classification is based on the scenarios’ global warming level from low warming of 1.5 °C with no or limited temporary temperature overshoot (C1) to high warming of more than 4 °C within this century (C8) (Guivarch et al., 2022a).

Regression models. We considered and compared four commonly used regression models in our analysis: gradient boosting, decision tree, random forest, and a k-nearest ~~neighbor~~neighbors regression model. In the initial stage, a more extensive set of ~~different~~commonly used regression models, including linear regression and multilayer perceptron regression, were tested, among which the four models mentioned above were selected for further hyperparameter tuning due to their superior performance compared to other regression models in the initial set. ~~We used grid search for our regression model hyperparameter optimization, using the machine learning library scikit learn (Pedregosa et al., 2011), based on the performance evaluation metrics described below.~~

135 For all models, we use the machine learning scikit-learn library for Python (Pedregosa et al., 2011). In the following, the four considered regression models are briefly described, while more detail is provided in the referenced seminal works and the scikit-learn documentation of the respective models including the mathematical representations of the underlying algorithms. A decision tree model is a supervised learning method to predict a target variable based on decision rules derived from a predictor variable. The model produces piecewise approximations of the target variable through a series of binary data splits

140 – similar to a tree structure with multiple branches. For continuous predictor variables, as in our case net AFOLU CO₂ emissions, the decision tree iteratively selects thresholds to split the data of the predictor into two groups. At each node of the

tree, multiple potential splits are evaluated, and the imposed threshold that minimizes the prediction error is selected to increase the accuracy of the prediction of the target variable. These nodes represent decision points where the data is divided based on whether it falls above or below the selected threshold. This process continues recursively, splitting the data at each node until the tree reaches the leaf nodes. At the leaf nodes, the tree makes a final prediction of the target variable – in our case, the expected Land CDR (Breiman et al., 1984). A gradient boosting regression model is an ensemble method which sequentially combines multiple simple models, called weak learners (typically decision trees as described above), which correct the previous models' predictions to reduce the error and improve the final model (Friedman, 2001). A random forest model is also an ensemble method which combines multiple decision trees as described above, but unlike gradient boosting, the trees in a random forest model run in parallel instead of sequentially. Each decision tree works independently, and their individual predictions are averaged to produce the final prediction (Breiman, 2001). The k-nearest neighbors model is not based on decision trees. Instead, it uses the proximity (similarity) of a scenario to a number (k) of neighbors (similar scenarios) to make predictions. For a given scenario, the model identifies the k nearest data points of the predictor variable in the feature space and then averages the target variable values of these neighbors to come up with a prediction (Goldberger et al., 2004).

Performance optimization and evaluation. From the machine learning library scikit-learn, we also used grid search for our regression model hyper-parameter optimization, and bootstrapping to estimate the variability in prediction performance for different subsamples of our training and testing data (Pedregosa et al., 2011). Grid search is an algorithm commonly used for regression problems, which allows to efficiently run regression models in different setups using all possible hyper-parameter combinations to eventually select the best performing model setup. The selection of hyper-parameter options for the model optimization was driven by the observed model performance and computation time. We used bootstrapping to explore how the prediction performance of our optimized regression models varied based on different resamples (n=1000) of the training and testing data, allowing us to better evaluate the robustness of the perceived performance of the tested models.

The model performance was ~~evaluated~~ determined based on four widely applied evaluation metrics, namely R-squared, mean absolute error, median absolute error, and maximum absolute error. These four evaluation metrics are briefly described in the following. R-squared was used to explore how well the tested regression model captured the relationship between the ~~dependent and independent~~ predicted variable, ~~while mean, median and maximum absolute errors were used to evaluate and the actual variable in the validation dataset, indicating the goodness of their fit.~~ R-squared can range from zero to one with higher values indicating better fits. The other three evaluation metrics instead indicate absolute error, meaning the absolute difference between the ~~dependent~~ predicted and the ~~independent~~ actual variable in the validation dataset throughout 2020-2100. – lower error values indicate more accurate predictions of the regression models. As the absolute error differs across variable pairs of the global scenarios (n=79) and the regional scenario variants (n=617) in the training and testing dataset, we reported the mean, median and maximum error across the considered scenarios. Mean and median error are useful to estimate the prediction models' overall performance whereas the maximum error is used to indicate the extreme in lower-end performance, based on the most inaccurately predicted scenario. Ultimately, the best performing model (~~gradient boosting regression~~ k-nearest neighbors) was used to impute the missing ~~gross land carbon sequestration~~ Land CDR data for 404 incomplete ~~scenarios~~

global scenarios (n=404) and incomplete sub-global scenario variants (n=2358) across the R10 regions in the AR6 Scenario Database.

For two time steps of two imputed scenarios, negative values of up to $-3 \text{ Mt CO}_2 \text{ yr}^{-1}$ were predicted, which is conceptually false and likely explained by the remaining. The performance of the four considered regression models and the selection of the k-nearest neighbors model error. These values are only slightly below the conceptual minimum and, therefore, set to zero is discussed further under Results.

Data postprocessing. For all imputed scenarios, the predicted ~~dependent~~target variable was compared to their ~~independent~~predictor variable to identify cases where imputed CDR on land is smaller than the respective net negative AFOLU CO₂ emissions, as this conceptual error was partly also perceived in the AR6 Scenario Database. The two imputation ~~dataset~~ containsdatasets for the global scenarios and R10 regional variants contain two data sheets. The first data sheet contains unadjusted imputation outputs. In contrast, the second sheet accounts for the conceptual error described above by replacing conceptually ~~false~~inconsistent predictions with their respective net negative AFOLU CO₂ emissions as a conservativelower bound proxy for land-basedLand CDR – implications are explained in the discussion section. The code to implement the analysis and the global and regional imputation ~~dataset~~datasets are publicly available at [\[https://doi.org/10.5281/zenodo.1069665413373539\]](https://doi.org/10.5281/zenodo.1069665413373539).

3 Results

Figure ~~2 shows~~2a and 3a show the performance of the four tested regression models along the four above-described evaluation metrics based on the testing set used for the regression model validation- for the global scenarios and their regional scenario variants. Overall, the ~~gradient boosting~~k-nearest neighbors regression model performs best, as it ~~describes the relationship between~~resembles the ~~dependent and independent~~actual variable most accurately, while keeping mean, median and absolute difference between the ~~two variables~~predicted variable and the actual variable comparatively low throughout 2020-2100- ~~While the k nearest neighbor regression- for both the global scenarios and the regional scenario variants. It also shows relatively low variance in performance across the bootstrapping results (Figure 2a and 3a). While the gradient boosting regression performs comparatively well or- for the prediction of the global scenarios and slightly better concerning the mean and median~~maximum absolute error- (Figure 2a), the k-nearest neighbors regression outperforms the gradient boosting regression outperforms the k nearest neighbor regression- regarding R-squared and the maximum absolute error. Also, the performance of the mean and median absolute error for the prediction of the incomplete regional scenario variants (Figure 3a). While the overall performance of these two regression models is similar, the k-nearest neighbors model was chosen to produce the two imputation datasets of this study, as the gradient boosting regression is most consistent when varying the ratio between the ~~training and the testing set~~model partly predicted slightly negative values in the target variable, which is conceptually inconsistent with a clean definition of Land CDR, which should have a uniform removal sign. The other two regression models perform less well than the k-nearest neighbors and gradient boosting and k nearest neighbor regressions. Overall, all models

show a slight performance drop for R-squared around 2020-2060, with more stable or increased performance thereafter – we have found no convincing explanation for this slight temporal variation in performance.

210




Figure 2: Performance of tested regression models to predict missing AR6 land removal data based on the used regression validation dataset (scenarios n=79).

215

Figure 3 shows the carbon removal on land across the scenarios. In absolute terms, the mean, median, and maximum errors are larger for the evaluated global scenarios than for their regional scenario variants – this is expected due to the substantially higher levels of Land CDR deployment on the global level compared to the R10 regions. On the global level, mean error is consistently below 200 MtCO₂ yr⁻¹ and for the median error consistently below 40 MtCO₂ yr⁻¹ – on the R10 region level we see mean error consistently below 15 MtCO₂ yr⁻¹ and median error close to zero. For both the global and regional level, the mean and median absolute difference between the predicted variable and the actual variable is judged to be reasonably low, based on the k-nearest neighbors regression, and the absolute difference between the actual and predicted variable is substantially smaller than between the actual variable and the net negative AFOLU CO₂ emissions as a lower bound proxy for comparison (see also Figure 2b and 3b). As a point of reference, the median Land CDR deployment across available scenarios of the scenario categories C1-8 in the AR6 Scenario Database is 1253 MtCO₂ yr⁻¹ for 2020-2060 and 3570 MtCO₂ yr⁻¹ for 2060-2100 – across the R10 regions median deployment is 39 MtCO₂ yr⁻¹ and 179 MtCO₂ yr⁻¹ respectively. This means that the median

220

error is around 1% of the median Land CDR deployment throughout the timeseries for the global scenarios and even lower for the regional scenario variants in the regression validation dataset. However, while the regression model seems to perform well overall based on the regression evaluation dataset, the observed maximum error suggests substantially worse performance in extreme cases, when looking at the scenario with the highest absolute error (Figure 2a and 3a).





Figure 2. Prediction performance for the global scenario data. Panel (a) shows the performance of tested regression models to predict missing AR6 land removal data based on the used regression validation dataset (scenarios $n=79$). Performance across the four evaluation metrics is shown as the median (solid line) and 5-95 percentile range (shaded area) of the bootstrapping results ($n=1000$) for each of the four tested regression models. The performance results refer to the comparison between the predicted variable compared to the actual variable in the regression validation dataset. Panel (b) shows the actual ('Carbon Sequestration|Land Use') versus predicted Land CDR and the AR6 net negative AFOLU CO₂ emissions (based on negative values in 'Emissions|CO₂|AFOLU') as a lower bound proxy for Land CDR across AR6 scenario categories in the regression validation dataset (scenarios $n=79$). The predicted data in the figure is based on the k-nearest neighbors


regression. The solid lines show the median across scenarios while the shaded area shows the 5-95 percentile range. Note: We follow the convention of the AR6 Scenario Database, to report CDR in positive numbers.

240 Figure 2b and 3b show Land CDR across the global scenarios and their regional scenario variants in the regression validation dataset, considering the actual AR6 variable for ~~carbon sequestration on land~~ Land CDR, the predicted ~~carbon sequestration on land~~ Land CDR using the ~~gradient boosting~~ nearest neighbors regression, and the net negative AFOLU CO₂ emissions as a ~~conservative~~ lower bound proxy for comparison. Considering the scenarios in ~~this~~ the global and regional regression validation ~~dataset~~ datasets, the predicted variable appears to be a better proxy variable for missing AR6 ~~land carbon sequestration~~ Land CDR than the net negative AFOLU CO₂ emissions proxy, as the predicted variable better resembles the shape of the actual variable and shows less absolute error throughout 2020-2100. While the predicted variable resembles the actual variable well across all eight AR6 scenario categories, Figure 32b suggests some variance in performance across these categories – for C8 scenarios, the drop in resemblance of the actual variable is most visible. This is at least partly due to the small number of underlying scenarios of this category in the regression validation dataset at the global level (n=2). The prediction performance across the different R10 regions is comparatively consistent, as shown in Figure 3b.

245

250





255 **Figure 3: Actual**, Prediction performance for the R10 regions scenario data. Panel (a) shows the performance of tested regression models to predict missing AR6 land removal data based on the used regression validation dataset (regional scenario variants n=617). Performance across the four evaluation metrics is shown as the median (solid line) and 5-95 percentile range (shaded area) of the bootstrapping results (n=1000) for each of the four tested regression models. The performance results refer to the comparison between the predicted variable compared to the actual variable in the regression validation dataset. Panel (b) shows the actual ('Carbon Sequestration|Land Use') versus predicted land-based Land CDR and the AR6 net negative AFOLU CO₂ emissions (based on negative values in 'Emissions|CO₂|AFOLU') as a conservative lower bound proxy for land-based Land CDR across AR6 scenario categories in the regression validation dataset (scenarios regional scenario variants n=79617). The predicted data in the figure is based on the gradient-boosting-regression-k-nearest

260

neighbors regression. The solid lines show the median across scenarios while the shaded area shows the 5-95 percentile range. Note: We follow the convention of the AR6 Scenario Database, to report CDR in positive numbers.

4 Discussion and conclusion

265 In this study, we tested and compared four regression models to impute missing AR6 scenario data on land-carbon
sequestration|Land CDR based on available data on net AFOLU CO₂ emissions. The tested gradient-boosting|k-nearest
neighbors regression model performed best and was used to impute the missing land-carbon-sequestration|Land CDR data for
404-incomplete global scenarios: (n=404) and incomplete sub-global scenario variants (n=2358) across the R10 regions. The
global and regional imputation dataset ~~is~~ datasets are publicly available at:
270 [<https://doi.org/10.5281/zenodo.1069665413373539>].

While we effectively resemble and impute land carbon sequestration data for 404 incomplete scenarios, our imputed dataset
does not account for perceived land sequestration related data issues in the AR6 Scenario Database beyond data availability.
The use of the variable 'Carbon Sequestration|Land Use' is further complicated as different reporting methodologies were used
across IAMs, and land CO₂ fluxes are not always consistently and explicitly split into emissions and removals (Ganti et al.,
275 2024). Different baselines for today's land removal are also perceived across scenarios, as shown in Figure 1. For several
scenarios in the AR6 Scenario Database, net negative AFOLU CO₂ emissions are larger than the reported carbon sequestration
on land, which indicates conceptual errors as carbon sequestration on land is perceived to be a gross variable (Byers et al.,
2022; Prütz et al., 2023). The issue of inconsistent removal baselines and net removal being larger than gross removal (which
is the case for less than a quarter of all scenarios) is partly also perceived in our imputed dataset, as we use data from the AR6
280 Scenario Database to train our model.

To address the latter problem, we provide an unadjusted imputation dataset as well as an adjusted imputation dataset for which
we replaced conceptually false predictions (net removal being larger than gross removal) with their respective net negative
AFOLU CO₂ emissions as a conservative proxy for land-based CDR. We emphasize that our imputed dataset is imperfect and
that the remaining data issues highlighted above must be considered when using our data imputation. Nevertheless, Figure 3
285 shows that our imputed land-based CDR variable is a markedly better proxy than the use of net negative CO₂ emissions, which
was partly used in previous studies (Schleussner et al., 2022; Warszawski et al., 2021; Prütz et al., 2023) — both in terms of
resembling the removal curve and reducing absolute error. Our imputation is also a better alternative to omitting a large part
of the scenario space that does not report carbon sequestration on land.

We While we effectively resemble and impute Land CDR data for incomplete scenarios, our imputed datasets do not resolve
290 underlying inconsistencies in the reporting of Land CDR in the AR6 Scenario Database. The use of the variable 'Carbon
Sequestration|Land Use' is further complicated as different reporting methodologies were used across IAMs, and land CO₂
fluxes are not always consistently and explicitly split into net negative CO₂ emissions and gross removals (Ganti et al., 2024;
Prütz et al., 2023). Different baselines for today's land removal are also perceived across scenarios, as shown in Figure 1. For
several scenarios in the AR6 Scenario Database, net negative AFOLU CO₂ emissions are larger than the reported Land CDR,

295 which indicates conceptual errors as Land CDR is a gross variable, which can only be larger or equal to net negative AFOLU CO₂ emissions (Byers et al., 2022; Prütz et al., 2023). The issues of inconsistent removal baselines and net negative CO₂ emissions being larger than gross removal are partly also perceived in our imputed datasets, as we use data from the AR6 Scenario Database to train our model.

300 To address the latter problem, we provide an unadjusted imputation dataset as well as an adjusted imputation dataset for which we replaced conceptually inconsistent predictions (net negative CO₂ emissions being larger than gross removal) with their respective net negative AFOLU CO₂ emissions as a lower bound proxy for Land CDR in the global (n=106) and regional (n=1594) imputation dataset.

305 We emphasize that our global and regional imputed datasets are imperfect and that the persisting issue of net negative CO₂ emissions and gross removals on land not always being separated and consistently reported across models, must be considered when using our data imputation. Nevertheless, Figure 2b and 3b show that our imputed Land CDR variable is a markedly better proxy than the use of net-negative CO₂ emissions, which was partly used in previous studies (Prütz et al., 2023; Schleussner et al., 2022; Warszawski et al., 2021) – both in terms of resembling the removal curve and reducing absolute error. Our imputation is also a better alternative to omitting a large part of the scenario space that does not report Land CDR.

310 Concerning use cases, we believe our global and regional imputed datasets on land carbon removal is Land CDR are most useful for analyses that aim to use the largest possible set of both original and imputed global scenarios (n=783+404) or regional R10 scenario variants (n=6162+2358) and a uniform carbon removal sign. However, the re-Such scenario ensemble assessments are relevant to better understand a range of different aspects concerning Land CDR in climate change mitigation pathways. Several specific use cases have been highlighted above, including an assessment of the arising gap in CDR deployment (Lamb et al., 2024), an analysis mentioned of residual emissions including the land sector (Lamb, 2024),

315 estimations of land-per-removal (Zhao et al., 2024) or evaluations of the attainability of mitigation scenarios, which rely on Land CDR (Warszawski et al., 2021).

So far, such analyses rely on insufficient proxy data or interim solutions to address the limited data availability of land carbon sequestration in the AR6 Scenario Database and could benefit from the here provided more comprehensive dataset on Land CDR across scenarios. Based on the evaluation of mean, median, and maximum absolute error of the here used regression

320 model, it is advisable to use our dataset for analyses that rely on a large ensemble of scenarios, e.g., all scenarios of a certain scenario category or even several categories. This is because the prediction results from the regression model are more reliable for scenario ensembles than for individual scenarios, which may show larger error, as shown by the maximum error in Figure 2a and 3a – arguably, it is generally advisable to aim to use scenario ensembles instead of individual scenarios to better capture uncertainties and diverse underlying assumptions, which may lead to more robust and credible analysis outcomes (Guivarch et al., 2022b).

325 The reanalysis discussed above by Gidden et al. is perceived to be more useful/suitable in terms of consistency and accuracy of today's removals and for direct comparisons of scenario data and national greenhouse gas inventories (NGHGI). While our imputation dataset contains Land CDR data for the scenario starting year 2020, today's and also historical emissions and removals are better captured and more comprehensively discussed by the Global Carbon Project (Friedlingstein

et al., 2023) – the merit of our imputation dataset lies in the future timesteps of scenarios. Ultimately, we hope this study can
330 be a valuable and complementary addition to the existing approaches addressing the ~~land carbon sequestration~~ Land CDR data
gap in the AR6 Scenario Database.

Acknowledgments

We thank the Integrated Assessment Modeling Consortium (IAMC) and the International Institute for Applied Systems
Analysis (IIASA) for their valuable work on collecting and hosting quantitative integrated assessment scenarios from and for
335 the research community. We also thank the scikit-learn team for providing the free software machine learning library, that
made this analysis easily implementable. The authors acknowledge funding from the European Union’s Horizon 2020 research
and innovation programme under grant agreement No 101003687 (PROVIDE), and No 951542 (GENIE). ~~R.P.R.~~ R.P.R.
acknowledges funding from the European Union’s Horizon 2020 research and innovation programme under grant agreement
No 101081521 (UPTAKE). ~~J.R.J.R.~~ J.R.J.R. acknowledges funding from the European Union’s Horizon 2020 research and innovation
340 programme under grant agreement No 101003536 (ESM2025). S.F. acknowledges funding from the German Ministry for
Education and Research under grant agreement No 01LS2101F (CDRSynTra).

Code and data availability

~~The analysis code and the imputed dataset are publicly available at: [https://doi.org/10.5281/zenodo.10696654] (Prütz et al.,
2024).~~
345 The analysis code and the global and regional imputed datasets are publicly available at:
[https://doi.org/10.5281/zenodo.13373539] (Prütz et al., 2024).

Competing interests. The authors declare that they have no conflict of interest.

Author contributions. ~~R.P.R.~~ R.P.R. led the study and conceptualization, with supervision by ~~S.F.S.F.~~ S.F.S.F. and ~~J.R. R.P.J.R.~~ R.P.
implemented the analysis and wrote the original draft. All authors reviewed and edited the paper.

350 References

Byers, E., Krey, V., Kriegler, E., Riahi, K., Schaeffer, R., Kikstra, J., Lamboll, R., Nicholls, Z., Sandstad, M., Smith, C., van
der Wijst, K., Loeoeq, F., Portugal-Pereira, J., Saheb, Y., Stromann, A., Winkler, H., Auer, C., Brutschin, E., Lepault, C.,
Müller-Casseres, E., Gidden, M., Huppmann, D., Kolp, P., Marangoni, G., Werning, M., Calvin, K., Guivarch, C., Hasegawa,
T., Peters, G., Steinberger, J., Tavoni, M., van Vuuren, D., Al-Khourdajie, A., Forster, P., Lewis, J., Meinshausen, M., Rogelj,
355 J., Samset, B., and Skeie, R.: AR6 Scenarios Database, <https://doi.org/10.5281/zenodo.5886912>, April 2022.
~~Ganti, G., Gasser, T., Bui, M., Geden, O., Lamb, W., Minx, J., Schleussner, C. F., and Gidden, M.: Carbon dioxide removal
deployment consistent with global climate objectives, (Preprint), <https://doi.org/https://doi.org/10.21203/rs.3.rs-3719978/v1>,~~

2024.

360 Gidden, M. J., Gasser, T., Grassi, G., Forsell, N., Janssens, I., Lamb, W. F., Minx, J., Nicholls, Z., Steinhauser, J., and Riahi, K.: Aligning climate scenarios to emissions inventories shifts global benchmarks, *Nature*, 624, 102–108, <https://doi.org/10.1038/s41586-023-06724-y>, 2023.

365 Guivarch, C., Le Gallie, T., Bauer, N., Fragkos, P., Huppmann, D., Jaxa-Rozen, M., Keppo, I., Kriegler, E., Krisztin, T., Marangoni, G., Pye, S., Riahi, K., Schaeffer, R., Tavoni, M., Trutnevyte, E., van Vuuren, D., and Wagner, F.: Using large ensembles of climate change mitigation scenarios for robust insights, *Nat. Clim. Chang.*, 12, 428–435, <https://doi.org/10.1038/s41558-022-01349-x>, 2022.

370 Kikstra, J. S., Nicholls, Z. R. J., Smith, C. J., Lewis, J., Lamboll, R. D., Byers, E., Sandstad, M., Meinshausen, M., Gidden, M. J., Rogelj, J., Kriegler, E., Peters, G. P., Fuglestad, J. S., Skeie, R. B., Samset, B. H., Wienpahl, L., van Vuuren, D. P., van der Wijst, K. I., Al Khourdajie, A., Forster, P. M., Reisinger, A., Schaeffer, R., and Riahi, K.: The IPCC Sixth Assessment Report WGIII climate assessment of mitigation pathways: from emissions to global temperatures, *Geosci. Model Dev.*, 15, 9075–9109, <https://doi.org/10.5194/gmd-15-9075-2022>, 2022.

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D., Brucher, M., Perrot, M., and Duchesnay, E.: Scikit-learn: Machine Learning in Python, *J. Mach. Learn. Res.*, 2011.

375 Prütz, R., Strefler, J., Rogelj, J., and Fuss, S.: Understanding the carbon dioxide removal range in 1.5 °C compatible and high overshoot pathways, *Environ. Res. Commun.*, 5, 41005, <https://doi.org/10.1088/2515-7620/accd8a>, 2023.

Prütz, R., Fuss, S., and Rogelj, J.: Imputation of missing IPCC AR6 data on land carbon sequestration, <https://doi.org/10.5281/zenodo.10696654>, February 2024.

380 Riahi, K., Schaeffer, R., Arango, J., Calvin, K., Guivarch, C., Hasegawa, T., Jiang, K., Kriegler, E., Matthews, R., Peters, G. P., Rao, A., Robertson, S., Sebbit, A. M., Steinberger, J., Tavoni, M., and van Vuuren, D. P.: Mitigation pathways compatible with long-term goals, in: IPCC: Climate Change 2022: Mitigation of Climate Change. Contribution of Working Group III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Shukla, P. R., Skea, J., Slade, R., Khourdajie, A. Al, Diemen, R. van, McCollum, D., Pathak, M., Some, S., Vyas, P., Fradera, R., Belkacemi, M., Hasija, A., Lisboa, G., Luz, S., and Malley, J., Cambridge University Press, Cambridge / New York, 2022.

385 Schleussner, C. F., Ganti, G., Rogelj, J., and Gidden, M. J.: An emission pathway classification reflecting the Paris Agreement climate objectives, *Commun. Earth Environ.*, 3, 135, <https://doi.org/10.1038/s43247-022-00467-w>, 2022.

390 Warszawski, L., Kriegler, E., Lenton, T. M., Gaffney, O., Jacob, D., Klingensfeld, D., Koide, R., Costa, M. M., Messner, D., Nakicenovic, N., Schellnhuber, H. J., Schlosser, P., Takeuchi, K., Van Der Leeuw, S., Whiteman, G., and Rockström, J.: All options, not silver bullets, needed to limit global warming to 1.5 °C: a scenario appraisal, *Environ. Res. Lett.*, 16, 64037, <https://doi.org/10.1088/1748-9326/abf6ee>, 2021.

Breiman, L.: Random Forests, *Mach Learn.* 45, 5–32, <https://doi.org/10.1023/A:1010933404324>, 2001.

Breiman, L., Friedman, J. H., Olshen, R. A., and Stone, C. J.: Classification And Regression Trees, Routledge, <https://doi.org/10.1201/9781315139470>, 1984.

395

Byers, E., Krey, V., Krieglner, E., Riahi, K., Schaeffer, R., Kikstra, J., Lamboll, R., Nicholls, Z., Sandstad, M., Smith, C., van der Wijst, K., Lecocq, F., Portugal-Pereira, J., Saheb, Y., Stromann, A., Winkler, H., Auer, C., Brutschin, E., Lepault, C., Müller-Casseres, E., Gidden, M., Huppmann, D., Kolp, P., Marangoni, G., Werning, M., Calvin, K., Guivarch, C., Hasegawa, T., Peters, G., Steinberger, J., Tavoni, M., van Vuuren, D., Al-Khourdajie, A., Forster, P., Lewis, J., Meinshausen, M., Rogelj, J., Samset, B., and Skeie, R.: AR6 Scenarios Database, <https://doi.org/10.5281/zenodo.5886912>, April 2022.

400

Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Bakker, D. C. E., Hauck, J., Landschützer, P., Le Quéré, C., Luijkx, I. T., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Anthoni, P., Barbero, L., Bates, N. R., Becker, M., Bellouin, N., Decharme, B., Bopp, L., Brasika, I. B. M., Cadule, P., Chamberlain, M. A., Chandra, N., Chau, T.-T.-T., Chevallier, F., Chini, L. P., Cronin, M., Dou, X., Enyo, K., Evans, W., Falk, S., Feely, R. A., Feng, L., Ford, D. J., Gasser, T., Ghattas, J., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, Ö., Harris, I., Hefner, M., Heinke, J., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Jacobson, A. R., Jain, A., Jarník, T., Jersild, A., Jiang, F., Jin, Z., Joos, F., Kato, E., Keeling, R. F., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Körtzinger, A., Lan, X., Lefèvre, N., Li, H., Liu, J., Liu, Z., Ma, L., Marland, G., Mayot, N., McGuire, P. C., McKinley, G. A., Meyer, G., Morgan, E. J., Munro, D. R., Nakaoka, S.-I., Niwa, Y., O'Brien, K. M., Olsen, A., Omar, A. M., Ono, T., Paulsen, M., Pierrot, D., Pockock, K., Poulter, B., Powis, C. M., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Rosan, T. M., Schwinger, J., Séférian, R., et al.: Global Carbon Budget 2023, *Earth Syst Sci Data*, 15, 5301–5369, <https://doi.org/10.5194/essd-15-5301-2023>, 2023.

405

410

415

Friedman, J. H.: Greedy function approximation: A gradient boosting machine., *The Annals of Statistics*, 29, 1189–1232, <https://doi.org/10.1214/aos/1013203451>, 2001.

Ganti, G., Gasser, T., Bui, M., Geden, O., Lamb, W. F., Minx, J. C., Schuessner, C.-F., and Gidden, M. J.: Evaluating the near- and long-term role of carbon dioxide removal in meeting global climate objectives, *Commun Earth Environ*, 5, 377, <https://doi.org/10.1038/s43247-024-01527-z>, 2024.

420

Gidden, M. J., Gasser, T., Grassi, G., Forsell, N., Janssens, I., Lamb, W. F., Minx, J., Nicholls, Z., Steinhauser, J., and Riahi, K.: Aligning climate scenarios to emissions inventories shifts global benchmarks, *Nature*, 624, 102–108, <https://doi.org/10.1038/s41586-023-06724-y>, 2023.

425

Goldberger, J., Hinton, G. E., Roweis, S., and Salakhutdinov, R. R.: Neighbourhood Components Analysis, in: Advances in Neural Information Processing Systems, 2004.

430 Guivarch, C., Kriegler, E., Portugal-Pereira, J., Bosetti, V., Edmonds, J., Fishedick, M., Havlik, P., Jaramillo, P., Krey, V., Lecocq, F., Lucena, A., Meinshausen, M., Mirasgedis, S., O'Neill, B., Peters, G. P., Rogelj, J., Rose, S., Saheb, Y., Strbac, G., Strøm, A. H., and Zhou, N.: IPCC, 2022: Annex III: Scenarios and modelling methods, in: IPCC: Climate Change 2022: Mitigation of Climate Change. Contribution of Working Group III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Shukla, P. R., Skea, J., Slade, R., Khourdajie, A. Al, van Diemen, R., McCollum, D., Pathak, M., Some, S., Vyas, P., Fradera, R., Belkacemi, M., Hasija, A., Lisboa, G., Luz, S., and Malley, J., 2022a.

435 Guivarch, C., Le Gallic, T., Bauer, N., Fragkos, P., Huppmann, D., Jaxa-Rozen, M., Keppo, I., Kriegler, E., Krisztin, T., Marangoni, G., Pye, S., Riahi, K., Schaeffer, R., Tavoni, M., Trutnevyte, E., van Vuuren, D., and Wagner, F.: Using large ensembles of climate change mitigation scenarios for robust insights, Nat Clim Chang, 12, 428–435, <https://doi.org/10.1038/s41558-022-01349-x>, 2022b.

440 Kikstra, J. S., Nicholls, Z. R. J., Smith, C. J., Lewis, J., Lamboll, R. D., Byers, E., Sandstad, M., Meinshausen, M., Gidden, M. J., Rogelj, J., Kriegler, E., Peters, G. P., Fuglestvedt, J. S., Skeie, R. B., Samset, B. H., Wienpahl, L., van Vuuren, D. P., van der Wijst, K.-I., Al Khourdajie, A., Forster, P. M., Reisinger, A., Schaeffer, R., and Riahi, K.: The IPCC Sixth Assessment Report WGIII climate assessment of mitigation pathways: from emissions to global temperatures, Geosci Model Dev, 15, 9075–9109, <https://doi.org/10.5194/gmd-15-9075-2022>, 2022.

445 Lamb, W. F.: The size and composition of residual emissions in integrated assessment scenarios at net-zero CO₂, Environmental Research Letters, 19, 044029, <https://doi.org/10.1088/1748-9326/ad31db>, 2024.

450 Lamb, W. F., Gasser, T., Roman-Cuesta, R. M., Grassi, G., Gidden, M. J., Powis, C. M., Geden, O., Nemet, G., Pratama, Y., Riahi, K., Smith, S. M., Steinhauser, J., Vaughan, N. E., Smith, H. B., and Minx, J. C.: The carbon dioxide removal gap, Nat Clim Chang, 14, 644–651, <https://doi.org/10.1038/s41558-024-01984-6>, 2024.

455 Matthews, J. B. R., Möller, V., Van Diemen, R., Fuglestvedt, J. S., Masson-Delmotte, V., Méndez, C., Semenov, S., and Reisinger, A.: Annex VII: Glossary., in: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, 2215–2256, <https://doi.org/10.1017/9781009157896.022>, 2021.

460 Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R.,
Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D., Brucher, M., Perrot, M., and Duchesnay, E.: Scikit-learn: Machine
Learning in Python, Journal of Machine Learning Research, 2011.

465 Prütz, R., Strefler, J., Rogelj, J., and Fuss, S.: Understanding the carbon dioxide removal range in 1.5 °C compatible and high
overshoot pathways, Environ Res Commun, 5, 41005, <https://doi.org/10.1088/2515-7620/acdbba>, 2023.

Prütz, R., Fuss, S., and Rogelj, J.: Imputation of missing IPCC AR6 data on land carbon sequestration,
<https://doi.org/10.5281/zenodo.13373539>, February 2024.

470 Riahi, K., Schaeffer, R., Arango, J., Calvin, K., Guivarch, C., Hasegawa, T., Jiang, K., Kriegler, E., Matthews, R., Peters, G.
P., Rao, A., Robertson, S., Sebbit, A. M., Steinberger, J., Tavoni, M., and van Vuuren, D. P.: Mitigation pathways compatible
with long-term goals, in: IPCC: Climate Change 2022: Mitigation of Climate Change. Contribution of Working Group III to
the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Shukla, P. R., Skea, J., Slade, R.,
Khourdajie, A. Al, Diemen, R. van, McCollum, D., Pathak, M., Some, S., Vyas, P., Fradera, R., Belkacemi, M., Hasija, A.,
Lisboa, G., Luz, S., and Malley, J., Cambridge University Press, Cambridge / New York, 2022.

475 Schleussner, C.-F., Ganti, G., Rogelj, J., and Gidden, M. J.: An emission pathway classification reflecting the Paris Agreement
climate objectives, Commun Earth Environ, 3, 135, <https://doi.org/10.1038/s43247-022-00467-w>, 2022.

480 Strefler, J., Bauer, N., Humpenöder, F., Klein, D., Popp, A., and Kriegler, E.: Carbon dioxide removal technologies are not
born equal, Environmental Research Letters, 16, 74021, <https://doi.org/10.1088/1748-9326/ac0a11>, 2021.

485 Warszawski, L., Kriegler, E., Lenton, T. M., Gaffney, O., Jacob, D., Klingensfeld, D., Koide, R., Costa, M. M., Messner, D.,
Nakicenovic, N., Schellnhuber, H. J., Schlosser, P., Takeuchi, K., Van Der Leeuw, S., Whiteman, G., and Rockström, J.: All
options, not silver bullets, needed to limit global warming to 1.5 °C: a scenario appraisal, Environmental Research Letters, 16,
64037, <https://doi.org/10.1088/1748-9326/abfecc>, 2021.

Zhao, X., Mignone, B. K., Wise, M. A., and McJeon, H. C.: Trade-offs in land-based carbon removal measures under 1.5 °C
and 2 °C futures, Nat Commun, 15, 2297, <https://doi.org/10.1038/s41467-024-46575-3>, 2024.

-