

# Constructing a measurement-based spatially explicit inventory of US oil and gas methane emissions (2021)

Mark Omara<sup>1,2</sup>, Anthony Himmelberger<sup>2</sup>, Katlyn MacKay<sup>1</sup>, James P. Williams<sup>1</sup>, Joshua Benmergui<sup>1,2,3</sup>, Maryann Sargent<sup>3</sup>, Steven C. Wofsy<sup>3</sup>, Ritesh Gautam<sup>1,2</sup>

5

<sup>1</sup>Environmental Defense Fund, New York, NY, USA 10010

<sup>2</sup>MethaneSAT, LLC, Austin, TX, USA 78701

<sup>3</sup>School of Engineering and Applied Science/Department of Earth and Planetary Science, Harvard University, Cambridge, MA, USA 02138

10 *Correspondence to:* Mark Omara ([momara@edf.org](mailto:momara@edf.org)) and Ritesh Gautam ([rgautam@edf.org](mailto:rgautam@edf.org))

## Abstract

Accurate and comprehensive quantification of oil and gas methane emissions is pivotal in informing effective methane mitigation policies, while also supporting the assessment and tracking of progress towards emissions reduction targets set by governments and industry. While national bottom-up source-level inventories are useful for understanding the sources of methane emissions, they are often unrepresentative across spatial scales, and their reliance on generic emission factors produces underestimations when compared with measurement-based inventories. Here, we compile and analyze previously reported ground-based facility-level methane emissions measurements ( $n = 1,540$ ) in the major US oil and gas producing basins and develop representative methane emission profiles for key facility categories in the US oil and gas supply chain, including well sites, natural gas compressor stations, processing plants, crude oil refineries, and pipelines. We then integrate these emissions data with comprehensive spatial data on national oil and gas activity to estimate each facility's mean total methane emissions and uncertainties for the year 2021, from which we develop a mean estimate of annual national methane emissions, resolved at  $0.1^\circ \times 0.1^\circ$  spatial scales ( $\sim 10 \text{ km} \times 10 \text{ km}$ ). From this measurement-based methane emissions inventory (EI-ME), we estimate total US national oil/gas methane emissions of approximately 16 Tg (95% confidence interval of 14-18 Tg) in 2021, which is  $\sim 2$  times greater than the EPA Greenhouse Gas Inventory. Our estimate represents a mean gas production-normalized methane loss rate of 2.6%, consistent with recent satellite-based estimates. We find significant variability in both the magnitude and spatial distribution of basin-level methane emissions, ranging from  $<1\%$  production-normalized methane loss rates in the gas-dominant Appalachian and Haynesville regions to  $>3\text{-}6\%$  in oil-dominant basins, including the Permian, Bakken, and the Uinta. Additionally, we present and compare novel comprehensive wide-area airborne remote sensing data and results of total area methane emissions and the relative contributions of diffuse and concentrated methane point sources as quantified using MethaneAIR in 2021. The MethaneAIR assessment showed reasonable agreement with independent regional methane quantification results in sub-regions of the Permian and Uinta basins and indicated that diffuse area sources accounted for the majority of total oil and gas emissions in these two regions. Our assessment offers key insights into plausible underlying drivers of basin-to-basin variabilities in oil and gas methane emissions, emphasizing the importance of integrating measurement-based data in developing high-resolution, spatially explicit

35

methane inventories in support of accurate methane assessment, attribution, and mitigation. The high-resolution spatially explicit EI-ME inventory is publicly available at <https://doi.org/10.5281/zenodo.10734300> (Omara et al. 2024)

## 40 1. Introduction

Accurate characterization of oil and gas methane emissions across spatial scales – from the facility-level to the basin- and national-level – is an essential component of methane reduction programs, integral to mitigating the near-term catastrophic impacts of human-induced global warming (IPCC, 2021). As governments, industry, and various stakeholders publicly commit to cut their methane emissions footprint (OGCI, 2021; GMP, 2021), accurate methane inventories will play a crucial role in the development and implementation of effective methane reduction approaches as well as tracking progress toward emission reduction targets.

At the national level, methane inventories are typically developed using “bottom-up” methods, for example, these methods are used by most countries that report annual greenhouse gas inventories to the UNFCCC (UNFCCC, 2023). “Bottom-up” methane inventories are developed by applying generic, or in some cases, empirically determined, component- or source-level emission factors to national oil and gas activity data (EPA, 2022). While these inventories are useful as first-order estimates of the emission sources, they often lack the accuracy needed to characterize methane emissions, their sources, and their trends over time at the facility-scale to the basin-level.

In addition, scores of recent studies at specific oil and gas basins (Zhang et al., 2020), countries (Alvarez et al., 2018; Shen et al., 2021; Zavala-Araiza et al., 2021; Johnson et al., 2023), and globally (Shen et al., 2023) have consistently found an underestimation in bottom-up inventories when compared to measurement-based inventories, pointing to a need for improvements in the bottom-up methane inventory methodologies. Furthermore, satellite-based quantification of regional, national, and global methane emissions has emerged as crucial tools for assessing the accuracies of methane inventories (Jacob et al., 2022; Shen et al., 2023). However, when Bayesian inversion models are used for methane flux quantification, spatially explicit methane inventories are needed as a priori information (Shen et al., 2021, 2023). Past efforts have produced such a priori information by spatially disaggregating methane emissions inventories reported to the UNFCCC (Maasackers et al., 2023; Scarpelli et al., 2022; EDGAR, 2023) which, as noted above, can have large underestimation and uncertainties in both the magnitude and spatial distributions of oil and gas methane emissions.

In this work, we utilize previous peer-reviewed facility-level measurement data ( $n = 1,540$ ) for methane emissions at oil and gas facilities in the major US oil and gas production basins to develop an improved assessment of national, basin-level, and facility-level methane emissions based on oil and gas activity in 2021. Our measurement-based inventory differs from other “bottom-up” inventories that use generic emission factors (e.g., EPA GHGI) in that we leverage empirical observations to derive insights on facility-level methane emission distributions useful for estimating population mean total methane emissions. Our contributions are three-fold: First, we develop statistically robust facility-level methane emission models based on measurement data collected in the years post-2011 (when EPA’s New Source Performance Standards for the oil and gas industry were first proposed) through 2020. We use these models to estimate national methane emissions, on both an absolute basis (Tg/year) and production-normalized

basis (% emitted relative to methane production). Second, we extend this approach to assess variability and underlying drivers of oil and gas methane emissions and methane loss rates across the major US oil and gas basins. As part of this assessment, we present and compare the quantification of total area methane emissions and the relative contributions of diffuse area emissions versus large concentrated methane sources in the Permian and Uinta Basins, based on new remote sensing measurements by MethaneAIR (Staebell et al., 2021; Chulakadabba et al., 2023; Chan Miller et al., 2023), an airborne precursor to MethaneSAT ([www.methanesat.org](http://www.methanesat.org)). Finally, we construct a high-resolution spatially explicit oil and gas methane emissions inventory for 2021, aggregated at  $0.1^\circ \times 0.1^\circ$  (~10 km  $\times$  10 km) spatial scales, and use these results to characterize the spatial patterns in national emissions.

## 2. Methods

### 2.1. Oil and gas activity data

We follow the procedure developed by Omara et al. (2022) to assess the total number and site-level production characteristics of actively producing onshore oil and gas well sites in the US in 2021. Aggregation of wellhead data to well site data (a well site can have multiple wellheads) is needed because (i) methane measurement-based data are reported at the well site level and (ii) production data are reported on a monthly basis for each producing wellhead. Briefly, we use the monthly well-level oil and gas production data as reported by Enverus Prism (Enverus, 2024), which aggregates public and proprietary data on monthly well-level production. For each actively producing well, we derive average well-level oil (barrels per day, bpd; 1 barrel crude oil ~ 0.136 tonnes), gas (1 thousand cubic feet per day, Mcfd;  $1 \text{ ft}^3 = 0.0283 \text{ m}^3$ ), and combined oil and gas (barrels of oil equivalent per day; 1 boed = 6 Mcfd gas) production rates based on the reported number of production days in the year, and assuming 365 calendar days in the calendar year if production days were not reported, which occurred at <5% of producing wells (Supplementary Fig. 10). We filtered the data for only the actively producing wells ( $n = 824,003$ ) and used geospatial clustering approaches, described in detail in Omara et al. (2022), to derive well site attributes (e.g., number of wells per site, site-level oil, gas, and boed production). Based on this analysis, we estimate a total of 660,149 actively producing onshore oil and gas well sites in the US in 2021 (Table 1), indicating an average of 1.2 wellheads per well site. Finally, we differentiate between low-production ( $\leq 15$  boed) and non-low production ( $> 15$  boed) oil and gas well sites based on their average site-level boed production rates in 2021. Our assessment indicates that low production well sites accounted for 82% of the total number of US onshore actively producing well sites in 2021 (Table 1). We consider these spatial data as comprehensive for the US oil and gas production well sites as it is consistent with the official gross oil and gas production reported by the US Energy Information Administration for 2021 (e.g., the sum of gross gas production from spatially explicit well-level production data from Enverus Prism is consistent with the total of ~42 Tcf of US natural gas gross withdrawals reported by the US Energy Information Administration, [https://www.eia.gov/dnav/ng/ng\\_prod\\_sum\\_dc\\_NUS\\_mmcf\\_a.htm](https://www.eia.gov/dnav/ng/ng_prod_sum_dc_NUS_mmcf_a.htm))

We estimate the total number of operational gathering and transmission compressor stations, natural gas processing plants, and crude oil refineries, based on spatial data reported by Enverus Prism (Enverus, 2024), supplemented with additional spatial data from the Oil and Gas Infrastructure Mapping (OGIM) database (Omara et al., 2023), which consolidates public-domain data on oil and gas infrastructure locations and facility characteristics.

110 For gathering and transmission pipelines, we estimate total pipeline miles based on the Enverus Prism (Enverus, 2024).  
In addition, we assess methane emissions associated with gas flaring activity, leveraging the natural gas flaring  
detections dataset based on VIIRS (visible infrared imaging radiometer suite) instruments onboard the Suomi National  
Polar-orbiting Partnership (NPP) and NOAA-20 satellites to identify the locations of gas flaring facilities or clusters  
of facilities and associated gas flared volumes (Elvidge et al., 2015). Table 1 shows the summary statistics for the oil  
115 and gas activity data used in this study.

## 2.2. Facility-level measurement-based methane emissions data

We begin by performing a comprehensive data review and assessment of previously published peer-reviewed  
data on facility-level methane emissions measurements for US oil and gas basins, leveraging Google Scholar search  
120 results based on keywords that reflect geography of interest (oil and natural gas methane emissions in the US),  
measurement methods (ground-based facility-level methods, OTM-33A, tracer flux, mobile transects), and major oil  
and natural gas facility categories (production well sites, natural gas gathering and transmission compressor stations,  
processing facilities, pipelines, crude oil refineries). We focus on ground-based measurement studies that report total  
facility-level methane emissions quantification for well sites, natural gas gathering and boosting compressor stations,  
125 natural gas transmission compressor stations, and natural gas processing plants.

These ground-based measurement approaches include the dual tracer downwind mobile measurements  
(Mitchell et al., 2015; Omara et al., 2016, 2018), the EPA Other Test Method (OTM-33A) downwind stationary  
measurements (Brantley et al., 2014; Robertson et al., 2017, 2020), and downwind mobile measurements with  
Gaussian plume transport modelling (Caulton et al., 2019; Omara et al., 2018). Omara et al. (2018) provides a detailed  
130 overview of these ground-based measurement methods. Other recent published studies that used chamber flux  
quantification approaches and reported only wellhead methane emissions quantification (e.g., wellhead methane  
emissions in Deighton et al. (2020) and Riddick et al. (2019)) are not included, as unquantified methane sources (e.g.,  
from separators, tanks, pneumatic devices, etc) likely lead to a low bias in facility-level total methane emissions.  
However, we use the total facility-level methane emissions data reported by Zimmerle et al. (2021) for natural gas  
135 gathering and boosting stations, based on aggregation of each facility's onsite component-level measurements  
performed using high flow sampler following leak detection with an infrared camera. We acknowledge possible low  
bias in this dataset given the limitations of facility-level measurements using high-flow samplers, including inability  
to access all methane emitting sources and/or to quantify large emission sources beyond the high-flow sampler  
capacity (Zimmerle et al., 2021). Finally, given their large size and difficulty of quantifying facility-wide emissions  
140 with ground-based measurement approaches, we use available measurement-based methane emissions data for crude  
oil refineries based on aerial remote sensing methods (Lavoie et al., 2017; Duren et al., 2019).

145

**Table 1.** Oil and gas activity data and estimates of total methane emissions

Facility category	Facility sub-category	Units	Activity data	Measurement-based data sources (sample size) <sup>a</sup>	Estimated total methane emissions, 2021 (Tg, 95% CI)	EPA GHGI, 2021 (Tg) <sup>b</sup>
Well sites	Low production	# of well sites	541,987	<i>n</i> = 1,153, see footnote for study references	4.3 (2.9-6.0)	3.4
	Non-low production	# of well sites	118,162		5.1 (3.6-7.4)	
Natural gas compressor stations	Gathering and boosting stations	# of stations	4,651	<i>n</i> = 116 (Mitchell et al. (2015)), <i>n</i> = 180 (Zimmerle et al. (2020))	1.6 (0.9-3.0)	1.4
	Transmission stations	# of stations	2,107	<i>n</i> = 47 (Subramanian et al. (2015))	1.7 (0.7-4.5)	
Natural gas processing plants	--	# of plants	908	<i>n</i> = 16 (Mitchell et al. (2015))	1.6 (0.7-3.7)	0.51
Crude oil refineries <sup>b</sup>	--	# of refineries	143	<i>n</i> = 28 (see footnote)	0.14(0.1-0.18)	0.03
Pipelines	Gathering pipelines	Pipeline miles	367,717	EPA Greenhouse Gas Inventory (EPA, 2022)	0.13 (0.12-0.14)	0.13
	Transmission pipelines	Pipeline miles	552,150		0.47 (0.46-0.48)	0.17
Natural gas flaring detections	# of flaring detections	# of detections	3,153	<i>n</i> = 3,153; Elvidge et al. (2015)	0.56 (0.55-0.57)	-
	Estimated gas flared volumes	MMcf/year	344,217	Elvidge et al. (2015)		
<b>Total estimated methane emissions</b>					15.7 (14.1-18.0)	8.3 (7.0-9.6)

**a** Measurements at well sites include 1,153 facility-level measurements from nine studies in eight basins or production regions in the US. Studies include Brantley et al. (2014), Robertson et al. (2017), Robertson et al. (2020), Omara et al. (2016), Omara et al. (2018), Caulton et al. (2019), Rella et al. (2015), Lan et al. (2015), and Yacovitch et al. (2015). For crude oil refineries, available facility-level measurements are based on aerial remote sensing quantification (Duren et al., 2019; Lavoie et al., 2017). **b** EPA GHGI total includes 0.5 Tg methane from natural gas distribution, LNG storage, and other sources not shown in this table.


For non-low production well sites, we use previously published facility-level measurement data collected in eight US basins, including the Barnett ( $n = 254$ ; Brantley et al. 2014; Lan et al., 2015; Rella et al., 2015; Yacovitch et al., 2015), Denver-Julesburg ( $n = 46$ ; Robertson et al., 2017; Brantley et al., 2014; Omara et al., 2018), Eagle Ford ( $n = 3$ ; Brantley et al., 2014); Fayetteville ( $n = 47$ ; Robertson et al., 2017), Marcellus Shale ( $n = 572$ ; Omara et al., 2016; Omara et al., 2018; Caulton et al., 2019), Permian ( $n = 72$ ; Robertson et al., 2020), Uinta ( $n = 31$ ; Robertson et al., 2017; Omara et al., 2018), and Upper Green River ( $n = 129$ ; Brantley et al., 2014; Robertson et al., 2017). The consolidated site-level measurement data included data collected in the years post-2011 (when EPA’s New Source Performance Standards for the oil and gas industry were first proposed) through 2020. We only focus on data from studies that reported total facility-level emissions quantification in addition to the production characteristics (i.e., gas and/or oil production rates). We use each study’s reported facility-level methane loss rate, computed as methane emissions relative to methane production at each facility, in our modelling of methane emissions. Where methane loss rates were not reported, we compute the percent methane loss rates as follows, based on the reported average gas production rate at the time of measurement:

$$\text{methane loss rate [unitless]} = CH_4 \left[ \frac{kg}{h} \right] \times \frac{1}{Gas [Mcf/d]} \times \frac{1 Mcf}{19.2 [kg CH_4]} \times \frac{1}{\sigma_{CH_4}} \times \frac{24h}{1d}$$

where  $CH_4$  [kg/h] is the measured facility-level methane emission rate in kg/h,  $Gas[Mcf/d]$  is the reported gas production rate in thousand cubic feet [Mcf] per day,  $19.2 kg/Mcf$  is the methane density at 60 °F (15.5 °C) and 1 atm, and  $\sigma_{CH_4}$  is the assumed methane content of the produced natural gas (we assume an average of 80% methane content in the produced natural gas).

For low-production well sites ( $\leq 15$  boed), we use the same facility-level methane emissions data and emissions assessment methods as described in detail in Omara et al. (2022). Briefly, we use the reported empirical observations ( $n = 240$ ; Omara et al., 2022) in a hybrid Monte-Carlo and non-parametric probabilistic model that simultaneously estimates the frequency of below-detection-limit sites, the frequency of high-emitting sites representing the top 5% of emitting facilities based on absolute methane emissions, and the distribution of high-emitter methane emissions, while accounting for the weakly observed positive relationship between emission rates and production rates for the bottom 95% of emitting well sites. We integrate this model with spatially explicit activity data on low-production oil and gas well sites in 2021 (Enverus, 2024) to estimate their total methane emissions.

For non-low production well sites ( $>15$  boed), we use the reported site-level measurement data described above and shown in Fig. 1a, which indicates an inverse relationship between production-normalized methane loss rates and facility-level gas production rates (Omara et al., 2018). The measurement-based data includes measurements that were reported as zeros or below the method detection limits of 0.036 kg/h (Robertson et al. 2017; Brantley et al., 2014) for the OTM-33A methods and 0.01 kg/h (Omara et al., 2016) for the dual tracer flux quantification.



185

190

195

**Figure 1.** Previously reported facility-level measurement-based methane emissions data. **a.** Facility-level methane emissions data (percent methane loss rate) as a function of gas production rate ( $n = 961$  non-low production well sites). The bottom  $x$ -axis shows the log-normalized gas production rates, with dashed vertical lines delineating the seven production cohorts used to model total methane emissions. Sample sizes for each production cohort are shown at the bottom of the Fig. panel above the bottom  $x$ -axis tick labels. The top  $x$ -axis shows the same production data in Mcfd. Each point is color-coded by basin and sized by the quantified methane emission rate in kg/h. Not shown are measurements that were reported as below the method detection limits. **b.** Absolute methane emission rate data (kg/h) for gathering and boosting (G&B) compressor stations ( $n = 295$ ), transmission and storage (T&S) compressor stations ( $n = 47$ ), natural gas processing plants ( $n = 16$ ), and crude oil refineries ( $n = 28$ ). The swarm plots show individual facility-level measurements, while the notched box plots show the distribution, where the boxes represent the 25<sup>th</sup> and 75<sup>th</sup> percentiles and the whiskers extend to  $1.5\times$  the interquartile range.

200

Figure 1b shows the previously reported facility-level measurements at midstream/downstream facilities, including natural gas gathering and boosting compressor stations, transmission compressor stations, processing plants, and crude oil refineries. In all cases, we use the average facility-level methane emissions data as reported, acknowledging inherent limitations in these measurement approaches (e.g., pseudo-random facility-level measurements with small sample sizes in ground-based approaches, or difficulty quantifying large emitters using high flow samplers in component-level measurements, etc.) likely increases the uncertainties in our estimates of total, regional, and national methane emissions.

205

### 2.3. Facility-level methane emissions model development and estimation of total national methane emissions

210

Our approach for estimating regional and national oil and gas methane emissions builds upon previous works that used data from hundreds to thousands of ground-based facility-level measurements (Zavala-Araiza et al., 2015; Alvarez et al., 2018; Omara et al. 2018; Omara et al., 2022) in combination with robust probabilistic models integrated with oil and gas activity data. Zavala-Araiza et al. (2015) and Alvarez et al. (2018) demonstrated that measurement-

based inventories developed using these methods produce total methane emission results that are in good agreement, within statistical uncertainty, of independent airborne measurements of total area methane emissions.

For non-low production well sites (average facility-level production rates > 15 boed), we begin by evaluating facility representativeness on the basis of (i) geographical diversity of measurements, (ii) distribution of facility-level production rates of measurements compared with the national population of well site facilities, and (iii) the distribution of facility-level methane emission rates across basins (Supplemental Fig. 3). Our measurement data, while limited in sample size, covers eight major US oil and gas basins with diverse oil and gas production characteristics, including the Appalachian, the Permian, Uinta, Barnett, Fayetteville, Greater Green River, and Denver-Julesburg. The wide range in basin-level gas-to-oil ratios (~1 to 800 Mcf/barrel) is well represented in the data for the sampled basins (Supplementary Fig. 3b).

In addition, the distribution of facility-level natural gas production rates shows reasonable overlap with that for the national population of non-low production facilities, and the broad range in distribution of facility-level production rates across the national population of sites (~90 Mcfd to >50,000 Mcfd) is well represented in the sampled sites (Supplementary Fig. 3c). However, the distribution of production rates for the sampled sites suggests potential bias toward higher-producing sites relative to the national distribution (Supplementary Fig. 3c). We account for any such potential biases by developing emission models based on production-normalized methane loss rate distributions (methane emitted relative to methane produced) across seven cohorts of specific gas production rates (further details below).

We develop and use probabilistic emission rate distributions based on production-normalized methane loss rates, which shows a wide range <0.01% to >90% (Figure 1a) across all basins (Supplementary Fig. 3d), reflecting, in part, the diversity in production characteristics within and across basins. We use production-normalized methane loss rate distributions because (i) the empirical data across a wide diversity of oil and gas production facilities suggests an inverse relationship in which high-producing facilities exhibit comparatively lower methane loss rates, and vice versa (Figure 1a) and (ii) the consolidated dataset includes measurements collected in earlier years before 2021. By using the production-normalized methane loss rate distribution models for specific cohorts of facility-level production rates, we do not model any particular site that is active in 2021 as exhibiting the same emission rate size as observed when measurements were taken in the past, as the empirical data and the model constrains facility-level methane loss rates to production levels, which will be time-variant. As such, we provide a necessary constraint on our estimates, effectively adjusting modelled facility-level methane emission rates if production rates have substantially changed over time.

To estimate regional methane emissions for non-low production well sites, we group the data for the empirical facility-level methane loss rates into seven log-normalized gas production (Mcfd) cohorts, shown in Fig. 1a and delineated by dashed vertical lines ( $\log\text{-Mcfd} \leq 5$ , 5-6, 6-7, 7-8, 8-9, 9-10 and  $\log\text{-Mcfd} > 10$ ). We use one log-e space (between  $\log\text{-Mcfd} \leq 5$  to  $\log\text{-Mcfd} > 10$ ) to develop these production cohorts, given the inverse relationship between facility-level methane loss rate and production rates, and selected to provide sufficient sample sizes for emissions distribution modelling for each production cohort (Fig. 1a). For each cohort, we simulate the frequency of finding a site emitting below the method detection limits (reported as zeros or below the method detection limit) through a




random bootstrapping procedure, repeated  $10^4$  times, with replacement. From this simulation, we develop a frequency distribution for the sites below the detection limits ( $f_{BDL}$ ), which averaged roughly 20% to 30% for all cohorts, with the exception of the last production cohort ( $>10$  Mcfd), where the frequency drops to roughly 10 to 20% (Supplementary Fig. 1).

For the measured oil and gas production well sites with emissions above the method detection limits, we begin by applying a log-transformation to the reported facility-level methane loss rates in each cohort and assessing the goodness of fit for the empirical distributions to a lognormal distribution, using the Kolmogorov-Smirnov test with significance established at  $p < 0.05$ . For all seven cohorts, we find that the lognormal distribution assumption is valid, with  $p > 0.05$  (Supplementary Fig. 2). For each cohort's empirical distribution, we assume a univariate normal likelihood with mean  $\mu$  and standard deviation  $\sigma$  and use Bayesian models with weakly informative priors to estimate  $\mu$  and  $\sigma$ , for example, as  $\mu \sim \text{Normal}(-10, 5)$  and  $\sigma \sim \text{HalfNormal}(3)$  for the first cohort of non-low production sites. For Bayesian inference, we draw 5,000 posterior samples from the posterior distribution using the PyMC3 (Salvatier et al., 2016) implementation of the No-U-Turn Sampler (NUTS) algorithm (Hoffman and Gelman, 2014) from which we estimate  $\mu$  and  $\sigma$ , as well as the 94% highest posterior density intervals (HPD). Note that the mean facility-level methane loss rate for each cohort can be computed as  $\exp(\mu + 0.5\sigma^2)$ . From the posterior results, we generate 5,000 predictions of the facility-level methane loss rate for each measured well site within each production cohort. Fig. 2 shows the cumulative probability distribution function for the observed data and 500 random samples from the model predictions.

We follow similar Bayesian modelling procedure to develop predictions of emission distributions (kg/h/facility), conditional on empirical data, for the gathering and boosting compressor stations, transmission compressor stations, natural gas processing plants, and crude oil refineries. For these facility categories, specifically for natural gas compressor stations and processing plants, we use the measured mean absolute methane emissions data as is (kg/h/facility) in our models, as we lack natural gas capacity or throughput information for the national population of facilities.

We then proceed as follows to estimate methane emissions for the total population of facilities: for every facility in each facility category and/or production cohort, we randomly draw an emission rate from the modelled posterior predictions (Fig. 2). For non-low production oil and gas production facilities, we randomly draw a methane loss rate estimate which is then multiplied by the facility's average methane production rate to estimate methane emissions in kg/h. As some facilities can have emissions below the method detection limits, we decrement the total estimated emission rate based on a randomly sampled frequency of BDL sites ( $f_{BDL}$ ), randomly drawn from the modelled distributions. We repeat this procedure 500 times and develop a methane emission distribution for the total methane emissions for each facility category or production cohort.

280



**Figure 2.** Empirical cumulative distribution functions of observed data and model predictions. Empirical CDFs are shown in solid black lines while thin colored lines show 500 random samples drawn from the model predictions. Sample sizes and data sources for empirical data are shown in Table 1. **a.** Non-low production well sites (modelled as facility-level methane loss rates), the Fig. shows the CDF for the  $5 < \log\text{-Mcf/d} < 6$  production cohort. Supplementary Fig. 2 shows the CDFs for all seven non-low production cohorts in Fig. 1. **b.** Low-production well sites (kg/h/site), **c.** Gathering and boosting compressor stations, **d.** Transmission compressor stations, **e.** Natural gas processing plants, **f.** Crude oil refineries.


285  
 290  
 295

Given the scarcity of facility-level measurements for gathering and transmission pipelines, we use the emission factors estimated by the US EPA Greenhouse Gas Emission Inventory (EPA, 2022; 285 kg methane/mile/year and 582 kg methane/mile/year, respectively) and assume normal distributions of emission factors with 50% uncertainty. Our use of EPA’s GHGI emission factors for these emissions sources makes it possible to provide a more complete spatially explicit inventory of oil and gas methane emissions (inclusive of gathering and transmission pipelines for which we have geospatial activity data), but likely increases uncertainties in our total methane emission estimates given potential underestimation in the GHGI emission factors.

We also estimate the methane emissions associated with gas flaring activities using location-specific gas flaring data from the VIIRS instrument (Elvidge et al., 2015) and apply average effective methane destruction removal efficiency of 91% (Plant et al., 2022; 95% confidence interval of ~90—92%).

300

Finally, we combine the emission distributions for all facility categories and sources using Monte Carlo methods to estimate the mean total national methane emissions and the 95% confidence interval based on the 2.5th and the 97.5th percentiles of the modelled distributions. Fig. 3 shows a general schematic of the emissions model development and estimation of total methane emissions.



**Figure 3.** General schematic for model development and estimation of total methane emissions, given activity data for each facility category.

#### 2.4. Spatial allocation of estimated methane emissions and basin-level methane loss rates

305 For each facility with known location (latitude, longitude), our assessment includes 500 different estimates of likely facility-level methane emission rates (in kg/h), from which we derive 500 different estimates of total national methane emissions. We use 500 simulation results for each facility as a reasonable simulation size that is not too computationally intensive to implement but that also gives sufficient statistical power to develop robust model uncertainty assessment. We use a search algorithm to identify a random sample of the facility-level emission rate distribution that most closely matches the computed mean estimate for the population of facilities. We use a similar approach to select a random sample of the facility-level emissions distributions representing uncertainties in the total emission estimates (i.e., the distribution that most closely matches the lower bound and upper bound of the 95<sup>th</sup> percent confidence intervals on the total estimated methane emissions). We then aggregate total mean methane emissions (and associated upper and lower bound estimates) on regular grids of  $0.1^\circ \times 0.1^\circ$  decimal degrees ( $\sim 10 \text{ km} \times 10 \text{ km}$ ), to produce spatially explicit oil and gas methane emission inventories and related uncertainties on the total methane emissions within each grid.

315

320

Our spatial allocation of estimated total oil and gas methane emissions is dependent, in part, on the completeness and spatial accuracy of oil and gas infrastructure locations for specific regions and oil and gas basins, for which related uncertainties are difficult to quantify based on available information. Our spatial allocation provides

325 the mean methane emissions estimates for the year 2021 aggregated at each  $0.1^\circ \times 0.1^\circ$  grid ( $\sim 10 \text{ km} \times 10 \text{ km}$ ) and are  
not intended to characterize methane emissions at a specific point in time, where substantial short-term variability in  
emissions may occur in part due to the stochastic character of facility-level methane emissions.

We compute basin-level and national methane loss rates as the ratio of estimated basin-level methane  
emissions to gross methane production in 2021, based on gross natural gas production data from Enverus Prism  
330 (Enverus, 2024) and an assumed average methane content of 80% in natural gas. Our assumption of an average 80%  
methane content in natural gas is informed by regional estimates of methane composition in natural gas based on the  
EPA GHGI (EPA, 2022). We acknowledge that uncertainties in methane composition across basins likely increases  
uncertainties in our overall methane loss rate calculations. Further studies on basin-level methane composition are  
needed to constrain these uncertainties. This methane intensity metric allows for a direct comparison of estimated  
335 methane losses relative to gross methane production across different basins. While our use of gross methane  
production accounts for emissions from associated gas produced during oil operations, the results are not intended to  
represent lifecycle emission intensities, which are outside the scope of this work.

## 2.5. Model uncertainties and limitations

340 In our modelling, we use the average facility-level emissions data as is, while assuming facility emissions  
arise from an underlying methane emissions distribution that is statistically described by lognormal distributions. The  
implementation of these probabilistic models produces emission distribution models (Fig. 2) that account for  
uncertainties in each facility's measured average methane emission rate and facility-to-facility variability in methane  
emissions within and across multiple oil and gas production regions. The 95% confidence intervals obtained through  
345 the Monte Carlo methods above reflect these uncertainties, as well as the model uncertainties in predictions of  
emissions distributions, given the limited sample sizes used herein. Additional uncertainties that are difficult to  
quantify include uncertainties in the oil and gas activity data and uncertainties in the potential impacts of recently  
promulgated federal/state-specific regulations or operator-specific practices regarding regular facility-level methane  
emissions monitoring and repair. In addition, due to lack of comprehensive spatially explicit activity data, our  
350 measurement-based inventory does not include methane emissions from downstream natural gas distribution, LNG  
storage, post-meter emissions, and abandoned oil and gas wells. The EPA GHGI (EPA, 2022) estimates these sources  
account for  $\sim 0.5$  to 1 Tg a year of total methane emissions, the vast majority of these would be distributed in urban  
locations outside of major oil and gas production regions.

## 3. Results and discussion


### 3.1 Total national oil and gas supply chain methane emissions

We estimate a measurement-based methane emission inventory (EI-ME) of total national oil and gas methane  
emissions for the onshore US as 15.7 Tg (95% confidence interval of 14 – 18 Tg or -10%/+15% uncertainty; Table 1;  
Fig. 4) for the year 2021. Our central estimate and confidence bounds are in reasonable agreement with recent  
measurement-based facility-level emission estimates (Alvarez et al., 2018; Rutherford et al., 2021 (production sector

360 only)) and satellite-derived oil and gas methane emissions, including GOSAT (Lu et al., 2022, 2023) and TROPOMI  
(Shen et al., 2022) quantification (Fig. 3b). In addition, consistent with previous findings (Alvarez et al., 2018;  
Rutherford et al., 2021; Shen et al., 2022), our central estimate is significantly greater than inventories developed  
using the traditional “bottom-up” source-level emission factor approaches: we find a factor of 1.9× and 1.8× greater  
total methane emissions than is estimated by the EPA Greenhouse Gas Inventory (EPA, 2022) and EDGAR v8  
365 (EDGAR, 2023) inventories for the year 2021. (Fig. 4).

We attribute the largest discrepancy between our measurement-based estimates and the EPA GHGI to the  
estimated emissions for the oil and gas production sector, which we estimate accounts for approximately 60% of the  
total onshore methane emissions for a total of ~9 Tg in 2021, or roughly 2.6× greater than the EPA GHGI’s estimate  
for the production-related methane emissions (Fig. 4; Table 1). These results are in reasonable agreement with  
370 previous measurement-based inventories (Alvarez et al., 2018; Rutherford et al., 2021; Omara et al., 2022) and, as has  
been noted elsewhere (Alvarez et al., 2018; Rutherford et al., 2021; Omara et al., 2018), likely reflect the use of  
emission factors in the EPA GHGI that do not adequately characterize the contributions of high-emitting methane  
sources that have been consistently observed in measurement-based studies. Furthermore, within the oil and gas  
production sector, we find that the low-production well site cohort (<15 boed) accounts for roughly one-half of total  
375 production site methane emissions in 2021, consistent with recent findings based on 2019 oil and gas activity (Omara  
et al. 2022). As Table 1 shows, the estimated total methane emissions from the low-production well site cohort alone  
are ~26% more than the total methane emissions from all low-production and non-low production well sites based on  
the EPA GHGI.

In 2021, we estimate a national methane loss rate of 2.6% (95% CI: 2.3 – 2.9%) relative to gross natural gas  
380 production, assuming an average of 80% methane content in natural gas (Methods). Our average methane loss rate  
assessment is in reasonable agreement with recent satellite-derived estimates (Shen et al., 2022 using TROPOMI and  
Lu et al., 2023 using GOSAT). Lu et al. (2023) reports a steadily declining national methane loss rate between 2010  
(~3.7%) and 2019 (~2.5%) and attributes these trends to two likely factors: (i) a slower increase/decrease in absolute  
methane emissions compared to the increase in methane production during this period and (ii) the impact of national  
385 regulations, such as the EPA’s New Source Performance Standards, promulgated in 2012, which focused on reducing  
emissions from newly constructed well sites, among other requirements. As we discuss further below, we find  
significant variability in the total methane emissions as well as spatial distributions of the estimated emissions at  
regional/basin-level for the of oil and gas activity in 2021.



390

**Figure 4.** Comparison of this study’s national estimate of total methane emissions from the oil and gas supply chain with previous measurement-based estimates. The first three bars show the oil and gas methane emissions estimated from facility-level measurements (this study, Alvarez et al. 2018) and production-sector-only methane emissions estimate by Rutherford et al. (2021) using models developed from component-level measurement data. Blue bars show the estimated emissions for the production sector, gold bars show the estimated emissions for the midstream and downstream facilities (compressor stations, processing plants, refineries, gathering and transmission pipelines). Error bars show the estimated 95% confidence bounds on the mean total methane emissions estimates. This study’s estimate of total national methane emissions include ~0.1 Tg/year of estimated methane emissions for Alaska. The green bars and the red bars show the satellite-derived estimates for the contiguous US based on GOSAT and TROPOMI observations, respectively. The last two bars show the “bottom-up” inventories from EPA GHGI and EDGAR v8 for the contiguous US. In all cases, the years for which methane emissions are estimated are shown on the top x-axis.

395

400

### 3.2. Variability in estimated basin-level methane emissions

Among the major oil and gas production basins, we identify the Permian, Appalachian, Anadarko, Eagle Ford, Haynesville, and the Barnett basins as the top six methane emitting basins, with estimated mean total basin-level methane emissions ranging from approximately 70 t/h to 340 t/h (Table 2, Fig. 5). These six basins account for 72% of onshore total combined oil and gas production (boe), and 52% of estimated total oil and gas methane emissions. Among these basins, we estimate considerable variability in gas production-normalized methane loss rates, with the lowest mean methane loss rates of <1% in the Appalachian and the Haynesville basins and the highest mean methane


405

410 **Table 2.** Top six methane emitting basins’ production, loss rate, and comparison with the EPA GHGI (traditional bottom-up inventory) and Lu et al. 2023<sup>a</sup> (satellite-derived estimates).

Basin	Basin area (km <sup>2</sup> )	Well site count (% from low-prod.)	Total annual (2021) gas production (Tcf <sup>b</sup> )	EI-ME methane emissions, 2021 (t/hr, 95% CI)   % from well sites	EPA GHGI methane emissions, 2020 (t/hr)	EI-ME methane loss rate, 2021 (% , 95% CI) <sup>c</sup>	EPA GHGI methane loss rate, 2020 (%)	Lu et al. GOSAT methane loss rate, 2019 (%) <sup>d</sup>	Shen et al. TROPOMI methane loss rate, 2019 (%) <sup>e</sup>
Permian	165,325	129,364 (78%)	6.5	335 (274 - 428)   69%	106	2.6 (2.1-3.3)	1.0	2.7 (1.6 - 3.0)	3.5 - 4.6
Appalachian	415,446	167,132 (97%)	12.7	231 (165 - 324)   75%	145	0.92 (0.66-1.30)	0.68	0.45 (0.40 - 0.47)	0.46
Anadarko	42,479	24,180 (64%)	1.9	119 (93 -166)   55%	33	3.2 (2.5-4.4)	1.0	3.4 (2.1 - 3.6)	1.5
Eagle Ford	50,179	24,377 (54%)	2.3	90 (73 - 119)   75%	29	2.0 (1.7-2.7)	0.69	1.1 (0.78 - 1.3)	2.0
Haynesville	28,922	23,895 (78%)	4.8	75 (59 - 95)   69%	30	0.80 (0.63-1.0)	0.41	1.2 (0.89 - 1.2)	1.0
Barnett	68,146	25,760 (79%)	0.92	74 (57 - 96)   68%	35	4.1 (3.1-5.3)	2.0	4.0 (3.3 - 4.1)	2.6

Note the differences in the temporal resolution of the studies used for the comparison, specifically the EPA GHGI basin-level estimates are based on Maasakkers et al. (2023) for the year 2020 (the latest year for which spatially explicit estimates are available), Lu et al. (2023) GOSAT estimates are for the year 2019, and Shen et al. (2022) TROPOMI estimates are based on satellite observation data aggregated over the period between May 2018 and February 2020. **a** Loss rates calculated assuming 90% methane content in each basin, for ease of comparison with Lu et al. **b** Tcf = Trillion cubic feet **c**. Methane loss rate calculated using 2021 production data from Enverus Prism **d**. Methane loss rate calculated using 2019 production data from Enverus Prism. **e**. For the Permian, Shen et al. (2022) reports posterior emissions in the range of 2.9 to 3.7 Tg/year representing a production-normalized methane loss rate of 3.5% to 4.6%.

420  
 loss rates of 3-4% in the Permian, Anadarko, and the Barnett (Table 2). The basin-level differences in methane loss rates among basins are consistent with the GOSAT-derived estimates for 2019 (Lu et al. 2023, Table 2), except for the Appalachian and the Eagle Ford Basins, where this study’s estimates are roughly 2× greater (Table 2). As with our findings on the comparative assessments with the EPA GHGI at the national level, our basin-level methane emissions estimates are consistently greater than the EPA GHGI estimates (Maasakkers et al., 2023) by a factor of 1.7× (Appalachian) to ~4× (Anadarko).



430 **Figure 5.** Basin-level differences in modelled mean total methane emissions and comparison with the EPA GHGI (Maasackers et al., 2023), TROPOMI-derived estimates (Shen et al., 2022) and GOSAT-derived estimates (Lu et al., 2023). See Supplementary Fig. 11 for a similar chart showing the state-by-state breakdown for the top 10 US emitting states based on the EI-ME inventory estimates.

435 The confluence of various possible factors, including the spatial density and characteristics of methane emitting oil and gas infrastructure and basin-level operational characteristics (gas-dominant versus oil-dominant, intensive flaring versus basins with negligible flaring, etc) contribute to the differences in the modelled basin-level methane emissions. In each basin, we estimate a predominant contribution of total methane emissions from well site infrastructure, ranging from 55% to 75% of the total basin-level methane emissions (Table 2; Fig. 5). Well site infrastructure characteristics vary significantly among basins, for example, the Appalachian Basin is characterized by a large population of old, leak-prone low-producing gas well sites (Omara et al., 2016; Deighton et al., 2020; Riddick et al., 2019) even as more than 95% of the gas produced comes from ~3% of well sites that are unconventional non-low production well sites (Enverus, 2024). This contrasts with the San Joaquin Basin, where well site infrastructure is dominated by low-producing oil pump jacks with limited onsite processing equipment, which in turn contrasts with the oil-dominant Bakken, dominated by high-producing horizontally-drilled well site facilities, typically with multiple wellheads and auxiliary processing equipment including separators, storage tanks, and flare stacks. Such basin-level 445 oil and gas infrastructure characteristics likely contribute to the modelled emission differences, given that the empirical



data synthesized herein reveal the weakly correlated methane emission profiles with well site production characteristics (loss rates, Fig. 1a) and infrastructure category (absolute emissions, Fig. 1b, Fig. 2).

Furthermore, the magnitude of modelled methane emissions varies by basin-level operational characteristics. For example, the Permian Basin with significant new oil and gas development stands in contrast to the relatively mature basins such as the Barnett or Uinta with steadily declining gas production and aging well site infrastructure. As Lu et al. (2023) observed, high methane loss rates tend to be associated with oil-dominant basins where production activities are focused on oil production even as substantial associated gas is co-produced along with oil (e.g., Permian, Eagle Ford, Bakken). In these basins, potentially higher methane emissions may occur due to venting and/or inefficient flaring of the co-produced gas, especially when there is insufficient infrastructure to gather, process, and transport to market the associated gas production, as has been postulated for the Permian Basin (Lyon et al., 2021; Varon et al., 2023; Lu et al., 2023). As noted previously, basin-level differences in total methane emissions could also be impacted by federal/state-level regulations of oil and gas methane emissions and/or operator-specific practices, affecting both the magnitude and temporal variability in emissions. While our methods are based on insights derived from empirical observations and robust modelling to estimate methane emissions specific to oil and gas activity in 2021, we lack sufficient data to characterize the impacts of specific regulations or operator practices. Further studies are needed to assess oil and gas methane emission trends and corresponding underlying drivers.


### **3.3. Sub-basin methane assessment and comparison with emissions quantification using MethaneAIR**

This study's EI-ME inventory provides methane emission estimates at geolocated oil and gas facilities, making it possible to develop aggregate methane emissions estimates across sub-basin to basin and national levels. We compare our sub-basin estimates for the Delaware portion of the Permian Basin and Uinta Basin with new remote sensing-based quantification by MethaneAIR (Staebell et al., 2021; Chulakadabba et al., 2023; Chan Miller et al., 2023), an airborne precursor to MethaneSAT satellite, which is scheduled to launch in 2024. MethaneAIR and MethaneSAT missions are managed by MethaneSAT LLC ([www.methanesat.org](http://www.methanesat.org)), which is a wholly owned subsidiary of Environmental Defense Fund. Both MethaneAIR and MethaneSAT are designed to produce quantitative data on total regional methane emissions while spatially disaggregating diffuse area emissions and detecting high-emitting point sources. Detailed description of the MethaneAIR instrument technical specifications, instrument calibration, retrieval methods and point source detections and validation can be found in recent works by Staebell et al., 2021, Conway et al. (2023), Chulakadabba et al., 2023, El Abbadi et al. (2023), Chan Miller et al. (2023), and Omara et al. (2023).

In August 2021, MethaneAIR flew a ~10,000 km<sup>2</sup> area in the Delaware sub-basin of the Permian Basin (research flight RF-06) and Uinta Basin (research flight RF-08) and produced quantification of total area methane emissions using a geostatistical inverse modelling (GIM) framework (based on Miller et al., 2020). The GIM framework was applied to inversion of the column mean methane dry air mole fraction retrieved using MethaneAIR measurements flying at 40,000 ft above ground aboard the NCAR GV aircraft ([https://www.eol.ucar.edu/field\\_projects/methaneair](https://www.eol.ucar.edu/field_projects/methaneair)). For MethaneAIR and MethaneSAT, the GIM framework is specialized to exploit the instrument's high spatial resolution, wide spatial coverage, and high precision, and ingests

high-emitting point source detections which are quantified using the modified integrated mass enhancement method (Chulakadabba et al., 2023). As such, remote sensing measurements by MethaneAIR at 40,000 ft above ground produce a high resolution, spatially explicit quantification of the total area methane emissions as well as high-emitting methane point sources emitting above ~200 kg/h.

We compare these new MethaneAIR total area methane quantification with the EI-ME modelled results, as well as the results from previous peer-reviewed studies in overlapping domains with the Permian RF-06 and the Uinta RF-08 regions. For both regions, we find good agreement, within uncertainty bounds, of the MethaneAIR quantification with other studies (Fig. 6), with emission rates quantification that fall within a representative range of mean total sub-basin methane emissions of 80 – 100 t/h and 17 – 24 t/h for RF-06 Permian and RF-08 Uinta, respectively (horizontal dashed lines in Fig. 6).




**Fig. 6.** Comparison of the EI-ME inventory with MethaneAIR and other peer-reviewed studies for two sub-regions of the Permian and Uinta Basins. Bars are color-coded by emission quantification method (MethaneAIR – hatched green bar; EI-ME – hatched dark green; PermianMAP airborne studies – purple; PermianMAP or tower-based study – red color; TROPOMI studies – dark blue bars). Lin et al. (2021) reports total Uinta Basin methane emissions estimates; we adjusted their estimate by the ratio of gas production in RF-08 region to total gas produced in Uintah and Duchesne counties in 2021 (RF-08 accounts for 74% of the total production in the two counties). For all other studies, we use only the reported emission estimates that overlap the MethaneAIR target boundaries. The dashed horizontal lines show a representative range of sub-basin methane emissions, computed via a bootstrapping procedure of all previously reported methane emissions (including the EI-ME results) to derive a lower bound and upper bound on the mean total methane emissions, based on the 2.5<sup>th</sup> and 97.5<sup>th</sup> percentiles of the resulting bootstrap distribution. Map credit: ESRI, 2023.

Based on MethaneAIR quantification for these two regions, we estimate that diffuse area emissions (which are assessed using the GIM modelling framework) account for the majority of methane emissions in both sub-basins, representing 63% and 88% of the total area methane emissions in the RF-06 and RF-08 regions, respectively. The remainder (37% and 12% of the total in RF-06 and RF-08 respectively), is attributable to the quantified high-emitting methane point sources with facility-specific methane emission rates in excess of ~200 kg/h/facility. These results are in reasonable agreement with the EI-ME results –averaged over the year—for the same spatial domains, in which oil and gas methane sources with mean methane emission rates <200 kg/h account for 85% and 90% of the total estimated methane emissions for RF-06 and RF-08, respectively. Furthermore, Cusworth et al. (2022) reports similar results for the same regions overlapping these domains in the Permian and Uinta, finding that methane sources below 200 kg/h account for 70% and 88% of total area emissions, which were quantified based on area-inversion of TROPOMI satellite observations and point source detections by AVIRIS-NG in 2021 and 2020 for RF-06 and RF-08, respectively. Furthermore, for a different sub-region of the Permian Basin, Kunkel et al. (2023) observed that facility sized emission sources with rates below 280 kg/h contribute 67% of the total emission rate from all sources with rates above 10 kg/h. At the national level, Omara et al. (2022) previously showed that the large population of low producing well sites (also known as marginal wells), with population average methane emissions rates of ~1 kg/h/site, account for roughly one-half of all production site methane emissions. Taken together, these results underscore the importance of small methane emitting sources dispersed across areas that, while individually emitting at low rates, nevertheless can contribute, in aggregate, a disproportionate fraction of regional total methane emissions. Williams et al. (2024) expands on these assessments, providing a detailed look at facility-level methane emissions distributions at the basin- and national-level.


### 3.4. Variability in estimated spatial distribution of methane emissions

Our spatially explicit EI-ME inventory suggests that basin-level differences also manifest as differences in the spatial distribution of total methane emissions. On average, we find oil and gas methane emission hotspots in every major US oil and gas production basin, including the Permian (the largest oil producing basin in the US located in western Texas and southern New Mexico), the Appalachian (Pennsylvania, Ohio, West Virginia, New York), Anadarko (Oklahoma, Texas), Eagle Ford (Texas), Bakken (North Dakota), and the Haynesville basins (Texas, Louisiana; Fig. 6)., Supplementary Fig. 4). Our analysis suggests methane emission hotspots tend to concentrate in high oil and gas production areas, for example, as evidenced by the two large hotspots in the rapidly developing, high-producing Delaware and Midland sub-basins of the Permian Basin in (Fig. 7a; see Supplementary Fig. 9 for maps representing lower bounds and upper bounds on spatially gridded emissions), consistent with spatial distributions for the satellite-observed methane emissions quantification in this region (Zhang et al., 2020; Varon et al., 2023). In addition, as with the total basin-level emissions, methane emissions spatial distributions are functions of oil and gas activity and their related facility-level emission characteristics. For example, substantial low-production oil and gas well site activity yields modelled methane emission hotspots in the southwestern tip of the Appalachian Basin (Fig. 7a), even as this region is not an oil and gas production hotspot (Supplementary Fig. 5). Furthermore, our analysis suggests spatial correlation of methane emission hotspots with intensive gas flaring activity, particularly for the oil

545




550



555

560

565




570

575 **Figure 7.** Estimated spatial distribution of total methane emissions and comparison with the EPA GHGI estimates. **a.**  
This study’s assessment of spatial distribution of US total oil and gas methane emissions, showing the estimates for  
the contiguous US (excluding Alaska, the total estimated methane emissions is 15.6 Tg in 2021). For visualization  
and comparison with the EPA GHGI inventory, the total methane emissions are gridded to  $0.1^\circ \times 0.1^\circ$  spatial scales  
(~10 km  $\times$  10 km). Major basin boundaries are outlined using black polygons. **b.** Estimated spatial distribution of total  
580 oil and gas methane emissions based on the EPA GHGI (2020; Maasackers et al., 2023). Note that the EPA GHGI  
data shown here is for the year 2020, the latest year for which spatially explicit data are available. **c.** Difference in  
spatially explicit methane emissions between this study’s estimates and the EPA GHGI. Warmer colors indicate  
comparatively higher estimates from this study relative to the EPA GHGI. We acknowledge the comparison is limited  
by the different time periods in the two studies – 2021 in this study versus 2020 for the EPA GHGI. Nevertheless, as  
585 both studies report annual averages, it is unlikely that significant differences in aggregate spatial distribution would  
have occurred between 2020 and 2021 to alter the main conclusions from this analysis. For the EI-ME, uncertainty  
estimates for each grid (i.e., lower bound and upper bound on mean estimates are presented in map form in  
Supplementary Fig. 9). Map credit: ESRI, 2023. Basin boundaries based on US EIA basin boundaries data  
(<https://www.eia.gov/maps/maps.php>)

590 producing basins with substantial associated gas production, including the Permian, the Eagle Ford, and the Bakken  
regions (Fig. 4, Supplementary Fig. 5).

We further assess variability in the spatial distribution of modelled methane loss rates, which reveals areas  
(25  $\times$  25 km<sup>2</sup> grids) in each major basin where methane loss rates are <0.25-1% of methane production. These areas,  
595 in general, are characterized by significant unconventional oil and gas production, for example, in the Appalachian  
Basin (north-eastern Pennsylvania and the tri-state corner of southern Pennsylvania, eastern Ohio, and northern West  
Virginia) as well as in the Permian Delaware and Midland sub-basins, and parts of the Haynesville, Eagle Ford, and  
the Bakken (Fig. 8). We also estimate areas with excessive methane loss rates >10% of methane production (Fig. 8)  
in each major producing basin, particularly in the Appalachian Basin, in the Michigan Basin, and in the greater  
600 Anadarko area of Missouri (Fig. 8). High methane loss rates are likely linked to the predominance of old, leak-prone  
low-producing well sites (e.g., in parts of the Appalachian and San Joaquin basins—although, note the overall lack of  
absolute methane emissions [kg/h] with site age in Supplementary Fig. 8; also see Brantley et al., 2014) or may be  
associated with modelled midstream infrastructure emissions from sources not collocated with significant oil and gas  
production.

605 The updated 2020 gridded EPA GHGI inventory for oil and gas systems (Maasackers et al., 2023) uses the  
same source of oil and gas activity data as this study (Enverus, 2024), and allocates GHGI emissions to specific  
emission source categories using infrastructure locations and methane emission scaling factors (e.g., scaled using well  
count, oil, and/or gas production for well sites depending on source category). The estimated methane emission  
hotspots (Fig. 7b) are in reasonable agreement with this study’s estimated spatial distributions (Fig. 7a,  $r = 0.64$ ), with  
610 notable exceptions in parts of the Michigan Basin (Michigan), the Appalachian Basin (Pennsylvania, Ohio, West  
Virginia) Basin, the Powder River Basin (Wyoming), the Barnett (east Texas), the Permian (west Texas), and the San  
Joaquin (southern California) Basins (Fig. 7b). In parts of these basins, strong methane hotspots appear in regions that  
likely reflect a dependence of emissions spatial allocation on spatial density of infrastructure (Supplementary Fig. 5).  
This differs with this study’s spatial allocation which leverages not just infrastructure locations, but simultaneously  
615 integrates the empirically observed facility-level methane emissions characteristics (Fig. 1, Fig. 2), which can vary



**Figure 8.** Estimated mean spatial distribution of production-normalized methane loss rates. For ease of visualization, we aggregate our facility-level methane inventories to a coarser spatial grid ( $25 \times 25 \text{ km}^2$ ) and normalize each grid's total estimated methane emissions relative to total methane production to derive spatially explicit methane loss rates, assuming 80% methane content in natural gas. Major basin boundaries are outlined in black and mean basin-level methane loss rates are shown as %. Map credit: ESRI, 2023. Basin boundaries based on US EIA basin boundaries data (<https://www.eia.gov/maps/maps.php>).

620

625

among populations of the same facility category (e.g., the distinction between emission profiles for low/non-low production well sites or among different production cohorts of the non-low production well site category).

In general, we find large differences in the magnitude of methane emissions in all of major basins shown in Fig. 7 when comparing the spatially explicit methane emissions in the GHGI and this study's estimates (Fig. 7c, Supplementary Fig. 6). We also find large differences in the spatial distributions of the methane emissions when comparing this study's spatially explicit emissions inventory with the EDGAR v8 inventory (EDGAR, 2023; Supplementary Fig. 7). Note that the EDGAR v8 total methane emissions are similar in magnitude to the EPA GHGI inventory estimates (Fig. 1), although emissions spatial allocation methods are primarily dependent on scaling by oil production characteristics, such that large methane hotspots are estimated in the oil-dominant basins of the Permian, the Bakken, and the Eagle Ford Basins (Supplementary Fig. 7).

630

635

Our results suggest both an underestimation in the magnitude of spatially explicit emissions in key US oil and gas basins as well as potentially unrepresentative spatial distributions of these emissions in the EDGAR v8 and

the EPA GHGI gridded inventories. These results carry important implications for the use of traditional “bottom-up” inventories as a priori information in Bayesian inversions of satellite observations for methane quantification, since both the magnitude and spatial allocation of emissions could influence the posterior results from these modelling systems under certain observational data constraints, such as insufficient observational data density (Shen et al., 2022). 640

#### 4. Data availability

EI-ME\_v1.0 can be accessed at <https://doi.org/10.5281/zenodo.10734300> (Omara et al. 2024) in an open-access GeoPackage file format. The GeoPackage file includes estimates for Alaska, while a .netcdf file is also provided with gridded emission results for only the lower 48 US states to facilitate easier comparison with recent satellite-derived methane emissions estimates for the contiguous US (Shen et al., 2022; Lu et al., 2023). 645

#### 5. Code availability

Python 3.7 code used for emissions modelling, extrapolation to population of facilities, and data visualization is available from the corresponding author upon reasonable request. 650

#### 7. Conclusions

Accurate and comprehensive assessment of oil and gas methane emissions is pivotal in informing effective methane mitigation policies. In this study, we develop robust statistical models based on measured facility-level methane emissions and integrate these models with comprehensive oil and gas activity data for onshore US oil and gas facilities to estimate total national oil and gas methane emissions for the year 2021. We estimate a total of ~16 (14 – 18) Tg of oil and gas methane emissions in 2021, representing a mean methane loss rate of 2.6% of gross gas production. Our national methane emission estimate, while in reasonable agreement with previous measurement-based estimates using facility-level measurements and satellite observations, are nevertheless roughly a factor of 2× greater than official inventories from the EPA Greenhouse Gas Inventory (GHGI). This improved assessment of national methane emissions underscores the importance of integrating measurement-based data to develop robust methane emission inventories which, as we show in this work, exhibit substantial variability in both the magnitude and spatial distribution of total methane emissions across major oil and gas basins. 655 660

Further improvements to methane emission inventories are possible through greater integration of measurement-based data including remote sensing approaches that can provide comprehensive area-wide total methane emissions, quantification of high-emitting methane point sources, as well as high-resolution spatial disaggregation of total methane emissions. In this study, we present the first set of such remote sensing quantification, based on MethaneAIR measurements in sub-basins of the Permian and Uinta and demonstrate reasonable agreement with several previous peer-reviewed assessments of total area methane emissions over similar spatial domains and time periods. These comprehensive area wide assessments also enable a detailed characterization of the importance of small methane sources dispersed across regions viz-a-viz large concentrated methane point sources, revealing their relative importance and variability across unique US oil and gas producing basins. 670

The EI-ME inventory provides a detailed characterization of total methane emissions by key facility categories at the national level as well as at the regional/basin-level, thus helping provide policy-relevant information that is important in developing and tracking effective methane mitigation strategies. The quantified uncertainties in our methane emission estimates could be improved upon in future studies through additional peer-reviewed data collection efforts, which are needed to develop further insights in response to ongoing methane mitigation efforts. There is a research need to develop robust statistical methods for effective integration of lower-detection-limit ground-based facility-level methane emissions data (such as data synthesized herein) with the growing number of airborne facility-level measurement studies, which generally have higher method detection limits (e.g., airborne methane remote sensing data in Duren et al., 2019; Cusworth et al., 2022; Sherwin et al., 2024). As demonstrated herein, improved integrated assessments of facility-level, regional, and national methane emission inventories, based on measurement data, support ongoing efforts to accurately quantify methane emissions, identify key methane sources and regions for targeted methane reductions, and track progress toward methane reduction goals.

685

#### **Author contributions**

MO and RG conceptualized the study. MO developed the facility-level methane emission models. Formal data analysis, interpretation, and visualization were performed by MO, AH and RG, with assistance from KM and JPW. MethaneAIR GIM modelling and high emitting methane point source assessments were performed by JB, MS and SCW. MO wrote the manuscript with contributions from all co-authors.

690

#### **Competing interests**

The authors declare that they have no conflict of interest.

#### **Acknowledgments**

This work was made possible by support from the Bezos Earth Fund. We thank Madeleine A. O'Brien for providing the OGIM database and related geospatial data gridding script. We are grateful to Daniel Zavala-Araiza for providing comments.

#### **References**

- Alvarez, R. A., Zavala-Araiza, D., Lyon, D. R., Allen, D. T., Barkley, Z. R., Brandt, A. R., Davis, K. J., Herndon, S. C., Jacob, D. J., Karion, A., Kort, E. A., Lamb, B. K., Lauvaux, T., Maasakkers, J. D., Marchese, A. J., Omara, M., Pacala, S. W., Peischl, J., Robinson, A. L., Shepson, P. B., Sweeney, C., Townsend-Small, A., Wofsy, S. C., Hamburg, S. P. Assessment of Methane Emissions from the U.S. Oil and Gas Supply Chain. *Science*, 361, 186–188, <https://doi.org/10.1126/science.aar7204>, 2018.
- Brantley, H. L., Thoma, E. D., Squier, W. C., Guven, B. B., Lyon, D. Assessment of Methane Emissions from Oil and Gas Production Pads Using Mobile Measurements. *Environ. Sci. Technol.*, 48 (24), 14508–14515, <https://doi.org/10.1021/es503070q>, 2014.
- Caulton, D. R., Lu, J. M., Lane, H. M., Buchholz, B., Fitts, J. P., Golston, L. M., Guo, X., Li, Q., McSperritt, J., Pan, D., Wendt, L., Bou-Zeid, E., Zondlo, M. A. Importance of Superemitter Natural Gas Well Pads in the Marcellus Shale. *Environ. Sci. Technol.*, 53(9), 4747–4754, <https://doi.org/10.1021/acs.est.8b06965>, 2019.

710



Chan Miller, C., Roche, S., Wilzewski, J. S., Liu, X., Chance, K., Souri, A. H., Conway, E., Luo, B., Samra, J., Hawthorne, J., Sun, K., Staebell, C., Chulakadabba, A., Sargent, M., Benmergui, J. S., Franklin, J. E., Daube, B. C., Li, Y., Laughner, J. L., ... Wofsy, S. C. Methane retrieval from MethaneAIR using the CO<sub>2</sub> Proxy Approach: A demonstration for the upcoming MethaneSAT mission [Preprint]. *Gases/Remote Sensing/Data Processing and Information Retrieval*, <https://doi.org/10.5194/egusphere-2023-1962>, 2023.

715

Chulakadabba, A., Sargent, M., Lauvaux, T., Benmergui, J. S., Franklin, J. E., Chan Miller, C., Wilzewski, J. S., Roche, S., Conway, E., Souri, A. H., Sun, K., Luo, B., Hawthorne, J., Samra, J., Daube, B. C., Liu, X., Chance, K. V., Li, Y., Gautam, R., ... Wofsy, S. C. Methane Point Source Quantification Using MethaneAIR: A New Airborne Imaging Spectrometer [Preprint]. *Gases/Remote Sensing/Validation and Intercomparisons*, <https://doi.org/10.5194/egusphere-2023-822>, 2023.

720

Conway, E. K., Souri, A. H., Benmergui, J., Sun, K., Liu, X., Staebell, C., Chan Miller, C., Franklin, J., Samra, J., Wilzewski, J., Roche, S., Luo, B., Chulakadabba, A., Sargent, M., Hohl, J., Daube, B., Gordon, I., Chance, K., Wofsy, S. Level0-to-Level1B processor for MethaneAIR [Preprint]. *Gases/Remote Sensing/Data Processing and Information Retrieval*, <https://doi.org/10.5194/amt-2023-111>, 2023.

725

Cusworth, D. H.; Thorpe, A. K.; Ayasse, A. K.; Stepp, D.; Heckler, J.; Asner, G. P.; Miller, C. E.; Yadav, V.; Chapman, J. W.; Eastwood, M. L.; Green, R. O.; Hmiel, B.; Lyon, D. R.; Duren, R. M. Strong Methane Point Sources Contribute a Disproportionate Fraction of Total Emissions across Multiple Basins in the United States. *Proc. Natl. Acad. Sci.* 119 (38), e2202338119. <https://doi.org/10.1073/pnas.2202338119>, 2022.

730

Deighton, J. A., Townsend-Small, A., Sturmer, S. J., Hoschouer, J., Heldman, L. Measurements Show That Marginal Wells Are a Disproportionate Source of Methane Relative to Production. *J. Air Waste Manag. Assoc.*, 70 (10), 1030–1042, <https://doi.org/10.1080/10962247.2020.1808115>, 2020

Duren, R. M., Thorpe, A. K., Foster, K. T., Rafiq, T., Hopkins, F. M., Yadav, V., Bue, B. D., Thompson, D. R., Conley, S., Colombi, N. K., Frankenberg, C., McCubbin, I. B., Eastwood, M. L., Falk, M., Herner, J. D., Croes, B. E., Green, R. O., Miller, C. E. California's methane super-emitters. *Nature*, 575(7781), 180–184, <https://doi.org/10.1038/s41586-019-1720-3>, 2019.

735

EDGAR (Emissions Database for Global Atmospheric Research) Community GHG Database, a collaboration between the European Commission, Joint Research Centre (JRC), the International Energy Agency (IEA), and comprising IEA-EDGAR CO<sub>2</sub>, EDGAR CH<sub>4</sub>, EDGAR N<sub>2</sub>O, EDGAR F-GASES version 8.0, European Commission, JRC (Datasets), [https://edgar.jrc.ec.europa.eu/report\\_2023](https://edgar.jrc.ec.europa.eu/report_2023), 2023.

740

El Abbadi, S., Chen, Z., Burdeau, P., Rutherford, J., Chen, Y., Zhang, Z., Sherwin, E., Brandt, A. Comprehensive evaluation of aircraft-based methane sensing for greenhouse gas mitigation [Preprint]. *Engineering*, <https://doi.org/10.31223/X51D4C>, 2023.

Elvidge, C., Zhizhin, M., Baugh, K., Hsu, F.-C., Ghosh, T. Methods for Global Survey of Natural Gas Flaring from Visible Infrared Imaging Radiometer Suite Data. *Energies*, 9(1), 14, <https://doi.org/10.3390/en9010014>, 2015.

745

Enverus Prism, [www.enverus.com](http://www.enverus.com), last accessed February 06, 2024

EPA: United States Environmental Protection Agency, Inventory of US Greenhouse Gas Emissions and Sinks, <https://www.epa.gov/ghgemissions/inventory-us-greenhouse-gas-emissions-and-sinks> (last access: 20 December 2023), 2022.

750

GMP: Global Methane Pledge, <https://www.ccacoalition.org/en/resources/global-methane-pledge> (last access: 12 December 2023), 2021.

Hoffman, M. D., Gelman, A. The No-U-Turn Sampler: Adaptively Setting Path Lengths in Hamiltonian Monte Carlo. *J. Mach. Learn. Res.*, 15 (1), 1593-1623, <https://api.semanticscholar.org/CorpusID:12948548>, 2014.

755

IPCC: The Intergovernmental Panel on Climate Change, Sixth Assessment Report. Climate Change 2021: The Physical Science Basis, Summary for Policy Makers, [https://www.ipcc.ch/report/ar6/wg1/downloads/report/IPCC\\_AR6\\_WGI\\_SPM.pdf](https://www.ipcc.ch/report/ar6/wg1/downloads/report/IPCC_AR6_WGI_SPM.pdf) (last access: 12 December 2023), 2021.

Jacob, D. J., Varon, D. J., Cusworth, D. H., Dennison, P. E., Frankenberg, C., Gautam, R., Guanter, L., Kelley, J., McKeever, J., Ott, L. E., Poulter, B., Qu, Z., Thorpe, A. K., Worden, J. R., Duren, R. M. Quantifying

- 760 methane emissions from the global scale down to point sources using satellite observations of atmospheric methane. *Atmos. Chem. Phys.*, 22(14), 9617–9646, <https://doi.org/10.5194/acp-22-9617-2022>, 2022.
- Johnson, M. R., Conrad, B. M., Tyner, D. R. Creating Measurement-Based Oil and Gas Sector Methane Inventories Using Source-Resolved Aerial Surveys. *Commun. Earth Environ.*, 4 (1), 139, <https://doi.org/10.1038/s43247-023-00769-7>, 2023.
- 765 Kunkel, W. M., Carre-Burritt, A. E., Aivazian, G. S., Snow, N. C., Harris, J. T., Mueller, T. S., Roos, P. A., Thorpe, M. J. Extension of Methane Emission Rate Distribution for Permian Basin Oil and Gas Production Infrastructure by Aerial LiDAR. *Environ. Sci. Technol.*, 57 (33), 12234–12241, <https://doi.org/10.1021/acs.est.3c00229>, 2023.
- 770 Lan, X., Talbot, R., Laine, P., Torres, A. Characterizing Fugitive Methane Emissions in the Barnett Shale Area Using a Mobile Laboratory. *Environ. Sci. Technol.*, 49(13), 8139–8146, <https://doi.org/10.1021/es5063055>, 2015.
- Lavoie, T. N., Shepson, P. B., Gore, C. A., Stirm, B. H., Kaeser, R., Wulle, B., Lyon, D., Rudek, J. Assessing the Methane Emissions from Natural Gas-Fired Power Plants and Oil Refineries. *Environ. Sci. Technol.*, 51(6), 3373–3381, <https://doi.org/10.1021/acs.est.6b05531>, 2017.
- 775 Lin, J. C.; Bares, R.; Fasoli, B.; Garcia, M.; Crosman, E.; Lyman, S. Declining Methane Emissions and Steady, High Leakage Rates Observed over Multiple Years in a Western US Oil/Gas Production Basin. *Sci. Rep.* 2021, 11 (1), 22291. <https://doi.org/10.1038/s41598-021-01721-5>.
- 780 Lu, X., Jacob, D. J., Wang, H., Maasackers, J. D., Zhang, Y., Scarpelli, T. R., Shen, L., Qu, Z., Sulprizio, M. P., Nesser, H., Bloom, A. A., Ma, S., Worden, J. R., Fan, S., Parker, R. J., Boesch, H., Gautam, R., Gordon, D., Moran, M. D., Reuland, F., Villasana, C. A. O., Andrews, A. Methane Emissions in the United States, Canada, and Mexico: Evaluation of National Methane Emission Inventories and 2010–2017 Sectoral Trends by Inverse Analysis of in Situ (GLOBALVIEWplus CH<sub>4</sub>; ObsPack) and Satellite (GOSAT) Atmospheric Observations. *Atmospheric Chem. Phys.*, 22 (1), 395–418, <https://doi.org/10.5194/acp-22-395-2022>, 2022.
- 785 Lu, X., Jacob, D. J., Zhang, Y., Shen, L., Sulprizio, M. P., Maasackers, J. D., Varon, D. J., Qu, Z., Chen, Z., Hmiel, B., Parker, R. J., Boesch, H., Wang, H., He, C., Fan, S. Observation-Derived 2010–2019 Trends in Methane Emissions and Intensities from US Oil and Gas Fields Tied to Activity Metrics. *Proc. Natl. Acad. Sci.*, 120 (17), e2217900120, <https://doi.org/10.1073/pnas.2217900120>, 2023.
- 790 Lyon, D. R., Hmiel, B., Gautam, R., Omara, M., Roberts, K. A., Barkley, Z. R., Davis, K. J., Miles, N. L., Monteiro, V. C., Richardson, S. J., Conley, S., Smith, M. L., Jacob, D. J., Shen, L., Varon, D. J., Deng, A., Rudelis, X., Sharma, N., Story, K. T., Brandt, A. R., Kang, M., Kort, E. A., Marchese, A. J., Hamburg, S. P. Concurrent Variation in Oil and Gas Methane Emissions and Oil Price during the COVID-19 Pandemic. *Atmospheric Chem. Phys.*, 21 (9), 6605–6626, <https://doi.org/10.5194/acp-21-6605-2021>, 2021.
- 795 Maasackers, J. D., McDuffie, E. E., Sulprizio, M. P., Chen, C., Schultz, M., Brunelle, L., Thrush, R., Steller, J., Sherry, C., Jacob, D. J., Jeong, S., Irving, B., Weitz, M. A Gridded Inventory of Annual 2012–2018 U.S. Anthropogenic Methane Emissions. *Environ. Sci. Technol.*, 57 (43), 16276–16288, <https://doi.org/10.1021/acs.est.3c05138>, 2023.
- 800 Maasackers, J. D., Jacob, D. J., Sulprizio, M. P., Scarpelli, T. R., Nesser, H., Sheng, J., Zhang, Y., Lu, X., Bloom, A. A., Bowman, K. W., Worden, J. R., Parker, R. J. 2010–2015 North American methane emissions, sectoral contributions, and trends: a high-resolution inversion of GOSAT observations of atmospheric methane, *Atmos. Chem. Phys.*, 21, 4339–4356, <https://doi.org/10.5194/acp-21-4339-2021>, 2021.
- Miller, S. M., Saibaba, A. K., Trudeau, M. E., Mountain, M. E., and Andrews, A. E.: Geostatistical inverse modeling with very large datasets: an example from the Orbiting Carbon Observatory 2 (OCO-2) satellite, *Geosci. Model Dev.*, 13, 1771–1785, <https://doi.org/10.5194/gmd-13-1771-2020>, 2020.
- 805 Mitchell, A. L., Tkacik, D. S., Roscioli, J. R., Herndon, S. C., Yacovitch, T. I., Martinez, D. M., Vaughn, T. L., Williams, L. L., Sullivan, M. R., Floerchinger, C., Omara, M., Subramanian, R., Zimmerle, D., Marchese, A. J., Robinson, A. L. Measurements of Methane Emissions from Natural Gas Gathering Facilities and Processing Plants: Measurement Results. *Environ. Sci. Technol.*, 49 (5), 3219–3227, <https://doi.org/10.1021/es5052809>, 2015.

- Nesser, H., Jacob, D. J., Maasackers, J. D., Lorente, A., Chen, Z., Lu, X., Shen, L., Qu, Z., Sulprizio, M. P., Winter, M., Ma, S., Bloom, A. A., Worden, J. R., Stavins, R. N., Randles, C. A. High-resolution US methane emissions inferred from an inversion of 2019 TROPOMI satellite data: contributions from individual states, urban areas, and landfills, *Atmos. Chem. Phys.*, 24, 5069–5091, <https://doi.org/10.5194/acp-24-5069-2024>, 2024.
- OGCI: The Oil and Gas Climate Initiative Reporting Framework. Version 3.5, [https://www.ogci.com/wp-content/uploads/2021/11/OGCI\\_Reporting\\_Framework\\_ver\\_3.5\\_Nov\\_2021.pdf](https://www.ogci.com/wp-content/uploads/2021/11/OGCI_Reporting_Framework_ver_3.5_Nov_2021.pdf) (last access: 12 December 2023), 2021.
- Omara, M., Gautam, R., O'Brien, M. A., Himmelberger, A., Franco, A., Meisenhelder, K., Hauser, G., Lyon, D. R., Chulakadabba, A., Miller, C. C., Franklin, J., Wofsy, S. C., Hamburg, S. P. Developing a spatially explicit global oil and gas infrastructure database for characterizing methane emission sources at high resolution. *Earth Syst. Sci. Data*, 15(8), 3761–3790, <https://doi.org/10.5194/essd-15-3761-2023>, 2023.
- Omara, M., Gautam, R., Himmelberger, A. et al. Measurement-Based Spatially Explicit Methane Emission Inventory (EI-ME) <https://doi.org/10.5281/zenodo.10734300>, 2024.
- Omara, M., Zavala-Araiza, D., Lyon, D. R., Hmiel, B., Roberts, K. A., Hamburg, S. P. Methane emissions from US low production oil and natural gas well sites. *Nat. Commun.*, 13(1), 2085, <https://doi.org/10.1038/s41467-022-29709-3>, 2022.
- Omara, M., Sullivan, M. R., Li, X., Subramanian, R., Robinson, A. L., Presto, A. A. Methane Emissions from Conventional and Unconventional Natural Gas Production Sites in the Marcellus Shale Basin. *Environ. Sci. Technol.*, 50 (4), 2099–2107, <https://doi.org/10.1021/acs.est.5b05503>, 2016.
- Omara, M., Zimmerman, N., Sullivan, M. R., Li, X., Ellis, A., Cesa, R., Subramanian, R., Presto, A. A., Robinson, A. L. Methane Emissions from Natural Gas Production Sites in the United States: Data Synthesis and National Estimate. *Environ. Sci. Technol.*, 52 (21), 12915–12925, <https://doi.org/10.1021/acs.est.8b03535>, 2018.
- Plant, G., Kort, E. A., Brandt, A. R., Chen, Y., Fordice, G., Gorchoy Negron, A. M., Schwietzke, S., Smith, M., Zavala-Araiza, D. Inefficient and unlit natural gas flares both emit large quantities of methane. *Science*, 377(6614), 1566–1571, <https://doi.org/10.1126/science.abq0385>, 2022.
- Rella, C. W., Tsai, T. R., Botkin, C. G., Crosson, E. R., Steele, D. Measuring Emissions from Oil and Natural Gas Well Pads Using the Mobile Flux Plane Technique. *Environ. Sci. Technol.*, 49 (7), 4742–4748, <https://doi.org/10.1021/acs.est.5b00099>, 2015.
- Riddick, S. N., Mauzerall, D. L., Celia, M. A., Kang, M., Bressler, K., Chu, C., Gum, C. D. Measuring Methane Emissions from Abandoned and Active Oil and Gas Wells in West Virginia. *Sci. Total Environ.*, 651, 1849–1856, <https://doi.org/10.1016/j.scitotenv.2018.10.082>, 2019.
- Robertson, A. M., Edie, R., Field, R. A., Lyon, D., McVay, R., Omara, M., Zavala-Araiza, D., Murphy, S. M. New Mexico Permian Basin Measured Well Pad Methane Emissions Are a Factor of 5–9 Times Higher Than U.S. EPA Estimates. *Environ. Sci. Technol.*, 54 (21), 13926–13934, <https://doi.org/10.1021/acs.est.0c02927>, 2020.
- Robertson, A. M., Edie, R., Snare, D., Soltis, J., Field, R. A., Burkhart, M. D., Bell, C. S., Zimmerle, D., Murphy, S. M. Variation in Methane Emission Rates from Well Pads in Four Oil and Gas Basins with Contrasting Production Volumes and Compositions. *Environ. Sci. Technol.*, 51 (15), 8832–8840, <https://doi.org/10.1021/acs.est.7b00571>, 2017.
- Rutherford, J. S., Sherwin, E. D., Ravikumar, A. P., Heath, G. A., Englander, J., Cooley, D., Lyon, D., Omara, M., Langfitt, Q., Brandt, A. R. Closing the Methane Gap in US Oil and Natural Gas Production Emissions Inventories. *Nat. Commun.*, 12 (1), 4715, <https://doi.org/10.1038/s41467-021-25017-4>, 2021.
- Salvatiery, J., Wiecki, T. V., Fonnesbeck, C. Probabilistic Programming in Python Using PyMC3. *PeerJ Comput. Sci.*, 2, e55, <https://doi.org/10.7717/peerj-cs.55>, 2016.
- Scarpelli, T. R., Jacob, D. J., Grossman, S., Lu, X., Qu, Z., Sulprizio, M. P., Zhang, Y., Reuland, F., Gordon, D., Worden, J. R. Updated Global Fuel Exploitation Inventory (GFEI) for methane emissions from the oil, gas, and coal sectors: Evaluation with inversions of atmospheric methane observations. *Atmos. Chem. Phys.*, 22(5), 3235–3249, <https://doi.org/10.5194/acp-22-3235-2022>, 2022.

- 855 Shen, L., Jacob, D. J., Gautam, R., Omara, M., Scarpelli, T. R., Lorente, A., Zavala-Araiza, D., Lu, X., Chen, Z., Lin, J. National quantifications of methane emissions from fuel exploitation using high resolution inversions of satellite observations. *Nat. Commun.*, 14(1), 4948, <https://doi.org/10.1038/s41467-023-40671-6>, 2023.
- Shen, L., Zavala-Araiza, D., Gautam, R., Omara, M., Scarpelli, T., Sheng, J., Sulprizio, M. P., Zhuang, J., Zhang, Y., Qu, Z., Lu, X., Hamburg, S. P., and Jacob, D. J.: Unravelling a large methane emission discrepancy in Mexico using satellite observations, *Remote Sens. Environ.*, 260, 112461, <https://doi.org/10.1016/j.rse.2021.112461>, 2021.
- 860 Shen, L., Gautam, R., Omara, M., Zavala-Araiza, D., Maasackers, J. D., Scarpelli, T. R., Lorente, A., Lyon, D., Sheng, J., Varon, D. J., Nesser, H., Qu, Z., Lu, X., Sulprizio, M. P., Hamburg, S. P., Jacob, D. J. Satellite Quantification of Oil and Natural Gas Methane Emissions in the US and Canada Including Contributions from Individual Basins. *Atmospheric Chem. Phys.*, 22 (17), 11203–11215, <https://doi.org/10.5194/acp-22-11203-2022>, 2022.
- 865 Sherwin, E.D., Rutherford, J.S., Zhang, Z. et al. US oil and gas system emissions from nearly one million aerial site measurements. *Nature* 627, 328–334 (2024). <https://doi.org/10.1038/s41586-024-07117-5>, 2024.
- 870 Staebell, C., Sun, K., Samra, J., Franklin, J., Chan Miller, C., Liu, X., Conway, E., Chance, K., Milligan, S., Wofsy, S. Spectral calibration of the MethaneAIR instrument. *Atmos. Meas. Tech.*, 14(5), 3737–3753, <https://doi.org/10.5194/amt-14-3737-2021>, 2021.
- 875 Subramanian, R., Williams, L. L., Vaughn, T. L., Zimmerle, D., Roscioli, J. R., Herndon, S. C., Yacovitch, T. I., Floerchinger, C., Tkacik, D. S., Mitchell, A. L., Sullivan, M. R., Dallmann, T. R., Robinson, A. L. Methane Emissions from Natural Gas Compressor Stations in the Transmission and Storage Sector: Measurements and Comparisons with the EPA Greenhouse Gas Reporting Program Protocol. *Environ. Sci. Technol.*, 49 (5), 3252–3261, <https://doi.org/10.1021/es5060258>, 2015.
- UNFCCC. The United Nations Framework Convention on Climate Change: GHG data from UNFCC. <https://unfccc.int/topics/mitigation/resources/registry-and-data/ghg-data-from-unfccc> (last access: 12 December 2023)
- 880 Varon, D. J., Jacob, D. J., Hmiel, B., Gautam, R., Lyon, D. R., Omara, M., Sulprizio, M., Shen, L., Pendergrass, D., Nesser, H., Qu, Z., Barkley, Z. R., Miles, N. L., Richardson, S. J., Davis, K. J., Pandey, S., Lu, X., Lorente, A., Borsdorff, T., ... Aben, I. Continuous weekly monitoring of methane emissions from the Permian Basin by inversion of TROPOMI satellite observations. *Atmos. Chem. Phys.*, 23(13), 7503–7520, <https://doi.org/10.5194/acp-23-7503-2023>, 2023.
- 885 Williams, J. P., Regehr, A., Kang, M. Methane Emissions from Abandoned Oil and Gas Wells in Canada and the United States. *Environ. Sci. Technol.*, 55 (1), 563–570, <https://doi.org/10.1021/acs.est.0c04265>, 2024.
- Williams, J.P., Omara, M., Himmelberger, A., Zavala-Araiza, D., MacKay, K., Benmergui J., Sargent, M., Wofsy, S., Hamburg, S.P., Gautam, R. Small emission sources disproportionately account for a large majority of total methane emissions from the US oil and gas sector. [Preprint] <https://doi.org/10.5194/egusphere-2024-1402>, 2024.
- 890 Yacovitch, T. I., Herndon, S. C., Pétron, G., Kofler, J., Lyon, D., Zahniser, M. S., Kolb, C. E. Mobile Laboratory Observations of Methane Emissions in the Barnett Shale Region. *Environ. Sci. Technol.*, 49 (13), 7889–7895, <https://doi.org/10.1021/es506352j>, 2015.
- 895 Zavala-Araiza, D.; Lyon, D. R.; Alvarez, R. A.; Davis, K. J.; Harriss, R.; Herndon, S. C.; Karion, A.; Kort, E. A.; Lamb, B. K.; Lan, X.; Marchese, A. J.; Pacala, S. W.; Robinson, A. L.; Shepson, P. B.; Sweeney, C.; Talbot, R.; Townsend-Small, A.; Yacovitch, T. I.; Zimmerle, D. J.; Hamburg, S. P. Reconciling Divergent Estimates of Oil and Gas Methane Emissions. *Proc. Natl. Acad. Sci.* 112 (51), 15597–15602. <https://doi.org/10.1073/pnas.1522126112>, 2015.
- 900 Zavala-Araiza, D., Omara, M., Gautam, R., Smith, M. L., Pandey, S., Aben, I., Almanza-Veloz, V., Conley, S., Houweling, S., Kort, E. A., Maasackers, J. D., Molina, L. T., Pusuluri, A., Scarpelli, T., Schwietzke, S., Shen, L., Zavala, M., and Hamburg, S. P.: A tale of two regions: methane emissions from oil and gas production in offshore/onshore Mexico, *Environ. Res. Lett.*, 16, 024019, <https://doi.org/10.1088/1748-9326/abceeb>, 2021.
- Zhang, Y., Gautam, R., Pandey, S., Omara, M., Maasackers, J. D., Sadavarte, P., Lyon, D., Nesser, H., Sulprizio, M. P., Varon, D. J., Zhang, R., Houweling, S., Zavala-Araiza, D., Alvarez, R. A., Lorente, A., Hamburg, S.

905 P., Aben, I., Jacob, D. J. Quantifying Methane Emissions from the Largest Oil-Producing Basin in the United States from Space. *Sci. Adv.*, 6 (17), eaaz5120, <https://doi.org/10.1126/sciadv.aaz5120>, 2020.

Zimmerle, D., Vaughn, T., Luck, B., Lauderdale, T., Keen, K., Harrison, M., Marchese, A., Williams, L., Allen, D. Methane Emissions from Gathering Compressor Stations in the U.S. *Environ. Sci. Technol.*, 54 (12), 7552–7561, <https://doi.org/10.1021/acs.est.0c00516>, 2020.