



High-resolution inventory and classification of retrogressive thaw slumps in West Siberia

Nina Nesterova^{1,2}, Ilia Tarasevich^{3,4}, Marina Leibman³, Artem Khomutov³, Alexander Kizyakov⁴, Ingmar Nitze¹, Guido Grosse^{1,2}

¹Permafrost Research Section, Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Potsdam, 14473, Germany

²Institute of Geosciences, University of Potsdam, Potsdam, 14476, Germany

³Earth Cryosphere Institute, Tyumen Scientific Centre SB RAS, 625026, Tyumen, Russia

⁴Cryolithology and Glaciology Department, Faculty of Geography, Lomonosov Moscow State University, Moscow, 119991, Russia

Correspondence to: Nina Nesterova (nina.nesterova@awi.de)

Abstract. Permafrost thaw disrupts ecosystems, hydrology, and biogeochemical cycles, reinforcing climate change through a positive permafrost-carbon feedback loop. Thaw can be gradual, deepening the active layer, or abrupt, triggering thermokarst, thermo-erosion, or thermodenudation. Retrogressive thaw slumps (RTSs) are a key manifestation of abrupt permafrost thaw. Yet, their distribution, scale, and environmental controls in the West Siberian Arctic remain poorly understood, further complicated by their rapid evolution. This study presents an extensive update of the West Siberian RTS inventory through manual mapping using high-resolution, multi-source, multi-year recent (2016-2023) satellite basemaps (ESRI, Google Earth, and Yandex Maps). We developed an RTS classification capturing key environmental parameters, including morphology, spatial organization, terrain position, and associated relief-forming concurrent processes. The dataset comprises 6168 classified RTS landforms, integrating newly mapped sites with previously reported occurrences to provide a comprehensive view of a 445226 km² region covering the Yamal, Gydan, and Tazovsky peninsulas. The collected data underwent manual filtering and verification, leveraging local field experience and observations from key sites to reduce uncertainty and minimize false positives. Accuracy analysis, performed by comparing the dataset with various field datasets collected across the peninsulas, confirmed high accuracy (>90%) for RTS identification. The dataset likely underestimated the distribution of small RTSs due to the resolution limitations of remote sensing data, hence generally providing a conservative estimate. This dataset serves as a valuable resource for diverse research fields, including ecology, biogeochemistry, geomorphology, climatology, permafrost science, and natural hazard assessment. Additionally, it provides a crucial reference dataset for machine learning applications, enhancing upcoming remote sensing classification and predictive modeling approaches.



1 Introduction

Permafrost is any ground that stays below 0°C for two or more consecutive years (Harris et al., 1988). It constitutes about 15% of the Northern Hemisphere landmass (Obu et al., 2021) and is experiencing significant warming and reduction in extent due to global warming (AMAP, 2017; Biskaborn et al., 2019; Smith et al., 2022). Permafrost thaw not only affects the high-latitude northern ecosystems and hydrological cycle but also releases carbon into the atmosphere and hydrosphere, contributing to global climate change with a positive feedback loop (Schuur et al., 2015). However, permafrost carbon emissions are still poorly integrated into global climate models (Miner et al., 2022). Furthermore, permafrost degradation manifests itself both gradually and abruptly. Gradual thaw slowly deepens the active layer over time (Brown et al., 2000; Luo et al., 2016; Vasiliev et al., 2020), while abrupt thaw in ice-rich permafrost triggers rapid thermokarst or thermo-erosion processes, leading to the formation of various landforms. Prime examples of such abrupt thaw events are specific types of permafrost-region landslides termed retrogressive thaw slumps (RTSs) (Nesterova et al., 2024).

RTSs are slope failures formed due to the thaw of exposed ice-rich permafrost (Fig. 1) (Mackay, 1966). These dynamic features can develop in a polycyclic fashion (Lantuit et al., 2005). Usually, the initial stages involve active ice ablation and downslope mudflows, followed by a stage of stabilization and colonization with pioneer vegetation (Mackay, 1966; Kerfoot, 1969; Leibman and Kizyakov, 2007). Active RTS can be considered as one of the clear indicators of permafrost response to increased air temperatures and higher summer precipitation. RTS occurrence significantly impacts the environment by altering the vegetation, topography, hydrology, as well as carbon fluxes (Lantz et al., 2009; Thienpoint et al., 2013; Cassidy et al., 2017). The prediction of RTS occurrence and activity is challenging due to heterogeneous ground ice distribution (Pollard and French, 1980) across the Arctic, limited observational field data (Jones et al., 2019), and the lack of models capable of simulating RTS initiation and dynamics.

The West Siberian Arctic with its continuous permafrost distribution is characterized by a high abundance of RTS. The prevalence of massive ground ice (Baulin et al., 1967; Streletskaia et al., 2013; Leibman and Kizyakov, 2007; Badu, 2015) that often occurs close to the surface contributes to the widespread abundance of RTSs (Khomutov et al., 2017). So far, the majority of RTS studies in the north of West Siberia have only been based on fieldwork at local key sites (Leibman and Kizyakov, 2007; Leibman et al., 2015; Khomutov et al., 2017; Novikova et al., 2018; Streletskaia et al., 2018; Babkina et al., 2019). Long-term field observations at the research station “Vaskiny Dachi” in Central Yamal reported the activation of rapid thaw processes after the extreme summer warmth of 2012 (Khomutov et al., 2017; Babkina et al., 2019).



Figure 1 RTS in Central Yamal, West Siberia, Russia. Photo taken in August 2021 by Nina Nesterova.

The vast majority of novel large-scale RTS studies utilize automated mapping with remote sensing data. This automated approach has some limitations for West Siberia so far, including using only a moderate spatial resolution of 30m not sufficient for detecting smaller RTS, only a partial cover of the West Siberian Arctic, the lack of high-resolution ground truth data, a large amount of false positive detection, and further feature interpretation ambiguities (Nitze et al., 2018; Runge et al., 2022; Nitze et al., 2024). Furthermore, the polycyclic nature of RTS development results in highly complex spatial patterns characterized by multiple overlapping or nested RTSs (Nesterova et al., 2024), which introduces further difficulties in highly automated mapping efforts. New cutting-edge panarctic datasets building on automated detection methods are being released (DARTS, Nitze et al., 2024b) but still have some limitations in accuracy on the local to regional scale.

In contrast, manual mapping of RTSs with high-resolution imagery by experts with regional knowledge can provide higher accuracy and decrease the amount of false positive detections (Lewkowicz and Way, 2019; Ward Jones et al., 2019; Nitze et al., 2024). A first manually mapped inventory of RTSs in the West Siberian Arctic was performed using the Yandex Maps high to moderate resolution satellite basemap representing the 2016-2018 period (Nesterova et al., 2021). The dataset reports 439 RTSs over both the Yamal and Gydan peninsulas. Due to the different spatial resolutions of satellite images used in the basemap (ranging from 0.4 to 15 m), the results tend to underestimate modern RTS distribution, particularly in areas where



only lower resolution imagery was available. Therefore, there was still no full understanding of the scale of thaw slumping in the West Siberian Arctic, its distribution, and environmental parameters, which are further complicated by the rapid evolution of RTSs.

We here provide an extensive update of the West Siberian RTS inventory, by performing manual RTS mapping in the north of West Siberia using multi-source and multi-year satellite basemaps (high-resolution ESRI, Google Earth, and Yandex Maps satellite basemaps). We further added all the RTS locations reported for this region in the literature so far. The collected dataset was manually filtered and compared to field data. This multi-source approach, in combination with regional field experience and field observations, gathered earlier at various key sites, helped us to minimize the uncertainty and decrease the number of false positive detections. We additionally developed a classification to describe each RTS, capturing their main environmental parameters such as morphology, spatial organization, terrain position, and concurrent relief-forming processes.

2 Methodology

Our approach includes four main steps: (1) visual identification of RTS and manual RTS point collection, (2) classification and parameter attribution, (3) iterative correction loop, and (4) final accuracy assessment (Fig. 2). Manual RTS point collection, classification, and correction were performed in QGIS software version 3.14. Accuracy analysis, plotting, and statistical calculations were performed using Python version 3.12.7.




Figure 2 Workflow overview. Rectangles with rounded corners present the datasets, and rectangles with sharp corners present the curation steps. The four main stages are numbered: 1 – Visual RTS identification and manual point collection stage, 2 – RTS point classification and parameter attribution, 3 – Iterative correction, and 4 – Accuracy assessment.



2.1 RTS point mapping

The study area in the north of West Siberia is 445 226 km² and includes the Yamal, Gydan, and Tazovsky peninsulas (Fig. 3). To ensure the completeness of the RTS dataset we reviewed previously published RTS datasets for the region, all of which were mapped using automated methods (Fig. 3). We manually filtered RTS datasets from Nitze et al. (2018), Runge et al. (2022), Bernhard et al. (2022), Huang et al. (2023), and Nitze et al. (2024b) to verify the presence of RTS and ensure that only true positives were included. This verification was conducted using the same available datasets that we later used for manual point collection, as described further below.




Figure 3 Study area in West Siberia with RTS datasets previously published in the literature. ESRI basemap has the following credits: Esri, DigitalGlobe, GeoEye, i-cubed, USDA FSA, USGS, AEX, Getmapping, Aerogrid, IGN, IGP, swisstopo, and the GIS User Community. Please note, that none of the external datasets fully covers the entire study area.



For our visual identification and manual collection of RTS points, we created a regular grid of 3.9 * 3.9 km cells covering the entire study area (Fig. 4a). This cell size was chosen as the optimal for visual inspection of the area and progress tracking. The ESRI satellite basemap was used as the primary source of information for RTS point collection due to the best quality of its recent very high-resolution imagery. This included high-resolution imagery (up to 0.31 m) largely with low cloudiness and an almost complete absence of visual artifacts. In rare cases, when the ESRI basemap did not fulfill visual quality criteria, such as no clouds, summer time of the image acquisition, and no artifacts, we used the Yandex Maps satellite basemap instead. In exceptional cases when neither the ESRI nor the Yandex basemaps fulfilled the visual quality criteria, we additionally checked the Google satellite basemap.




Figure 4 Manual mapping of RTSs in West Siberia: (a) Example of a grid cell with manually mapped RTSs (orange dots); (b) Coverage of the study area by high-resolution satellite images from different years in the ESRI basemap in km²; (c) Example of a lake shore RTS (marked by yellow point) on ESRI basemap imagery and typical visual RTS indicators: 1 – headwall, 2 – mudflow, 3 – contrasting colors of the disturbed slump floor with bare ground and the surrounding intact tundra vegetation; (d) Example of coastal RTS (marked by yellow point) on ESRI basemap imagery affected by coastal thermo-erosion, with white bracket indicating the full elongated extent of the coastal landform considered to be a single RTS in our inventory dataset. ESRI basemap used in (a), (c), and (d) has the following credits: Esri, DigitalGlobe, GeoEye, i-cubed, USDA FSA, USGS, AEX, Getmapping, Aerogrid, IGN, IGP, swisstopo, and the GIS User Community.

The majority of the high-resolution satellite images used in the ESRI basemap mosaic are recent Maxar images obtained after 2015 (Fig. 4b). Over a third of the study area is covered by satellite images from 2023 (Fig. 4b). Since the ESRI basemap was utilized as the primary source, all metadata related to the satellite images in the mosaic for identifying the RTS is stored within the inventory dataset's metadata. Yandex Maps basemap presents a mosaic of various satellite imagery taken in 2016-2018



with spatial resolutions ranging from 0.4 up to 15 m. The majority of images are dated July 2017 (Nesterova et al., 2021). For the Google satellite image layer, no individual image metadata was provided. RTSs were identified at a 1:1000 mapping scale in the satellite imagery based on visual indicators such as a clear outline of the headwall, the presence of a mudflow, and the sharp contrast in colors between the disturbed slump floor with bare ground and the adjacent intact tundra vegetation (Fig.4c). Thus, stabilized RTSs were also identified when the indicators were still visible. For each identified feature, we created a point in the location of the RTS within the visible outlines of the RTS. Each digitized point represented one feature that would be classified (see Sect. 2.2). Due to the complex nature of coastal RTSs sometimes stretching along coastal segments (Fig. 4d), we decided to identify each elongated contour with visible semicircles embedded inland as one feature. Such contours were often separated from each other by little streams or watercourses. This approach allowed us to utilize a single technique for all coastal RTSs, regardless of their size and shape. The RTS points underwent two visual corrections by the first author. To differentiate the process of coastal erosion from thermodenudation (Günther et al., 2012; Nesterova et al., 2024) and thereby distinguish other coastal landforms from RTSs, a special correction was applied to all coastal RTSs and thermoterrace RTSs (see Sect. 2.2). This involved verifying the headwall retreat of the RTS outline using the ESRI Wayback Machine - a digital archive of the World Imagery basemap of different versions providing multi-temporal imagery (ESRI Wayback Imagery, 2024). The same verification procedure was applied for the identification of RTS in the southernmost part of our West Siberian study area, where no reliable data on massive ground ice distribution is available and thus permafrost landforms can have different origins. The literature specifies the limits of massive ground ice extent in the north of West Siberia only very approximately (Baulin and Danilova, 1998).

2.2 Classification

We classified each RTS point based on terrain position, morphology, spatial organization, and concurrent cryogenic processes (Fig. 5). The four main criteria had a total of 15 parameters.




Figure 5 RTS classification scheme with four main criteria (shown as grey blocks) and 15 variables.



The terrain position of an RTS is defined based on the location of the object to either some hydrological feature (*sea coast*, *river bank*, *lakeshore*, and *gully*) or just *slope* when there was no visible hydrological feature. The location *lake* was selected for RTSs even on the former shores of drained lakes.

We further defined three types of RTS morphologies: *thermocirque*, *thermoterrace*, or a *combination* of these two (Nesterova et al., 2024). *Thermocirque* generally presents a horseshoe-like RTS shape (Fig. 6a), while *thermoterrace* is applied to an elongated RTS with mostly straight headwall outlines parallel to a coastline or riverbank (Fig. 6b). The combination of these two morphologies sometimes occurs when the elongated RTS landform also contains circular isometric curves of headwall outlines (Fig. 6c). It is usually formed when a thermocirque merges with a thermoterrace or when multiple thermocirques merge in one elongated landform. The complicated shapes of these combined RTS features make it highly challenging to distinguish between individual elongated and horseshoe-like RTSs (Fig. 6c). Our decision tree to define the morphology of RTS is shown in Supplement.



Figure 6 Examples of the three main RTS morphologies mapped in West Siberia: (a) Two thermocirques (yellow dots); (b) A single large thermoterrace (yellow dot); (c) A combined RTS morphology of merged thermocirques or merged thermocirque and a thermoterrace (yellow dot). ESRI basemap used has the following credits: Esri, DigitalGlobe, GeoEye, i-cubed, USDA FSA, USGS, AEX, Getmapping, Aerogrid, IGN, IGP, swisstopo, and the GIS User Community.

Due to the polycyclic nature of RTS development, these landforms can exhibit a very complex spatial organization of nested and amalgamated RTSs (Nesterova et al., 2024). We identified two types of RTS spatial organization: single landforms and complex landforms. RTS can be classified as a single landform when its outline is distinct and clearly defined and there is no more than one actively thawing zone within this outline (Fig. 7a). RTS can be classified as a complex landform when its boundary is difficult to define and/or there are two or more actively thawing zones (Fig. 7b). All the RTSs with combined morphologies were marked as complex landforms.

The influence of concurrent (happening in parallel to RTS development) processes on RTS development is described in Nesterova et al. (2024). For each mapped RTS, we noted the possible presence of 5 concurrent processes: *lateral thermo-erosion*, *coastal thermo-erosion*, *ice wedge erosion*, *nivation*, and *thermokarst subsidence*. *Lateral thermo-erosion* was identified by the rugged outline of the RTS and visible traces of erosive channels (Fig. 8a). The *Coastal thermo-erosion* classifier includes not only the sea cost erosion but also river and lakeshore erosion. It was determined by a sharp dark outline of the RTS base along the coastline of a waterbody and the absence of sediment accumulation in the water (Fig. 8a). We have



noted *ice wedge erosion* when an RTS headwall had a jagged outline resembling the adjacent polygonal surface of undisturbed tundra (Fig. 8b). *Nivation* was detected as white patches of snowpacks that stayed over the summer within RTS (Fig. 8a). *Thermokarst subsidence* appears as small thermokarst ponds filled with water. It is noticeable as black patches within the RTS outline (Fig. 8b).




Figure 7 Examples of the different spatial organization of RTSs: (a) Single RTS landform with a distinct outline (yellow dot); (b) Complex RTS landform (yellow dot) with multiple nested active (1) and stabilized (2) RTSs within one contour. ESRI basemap used has the following credits: Esri, DigitalGlobe, GeoEye, i-cubed, USDA FSA, USGS, AEX, Getmapping, Aerogrid, IGN, IGP, swisstopo, and the GIS User Community.




Figure 8 Examples of RTS with concurrent processes: (a) Stabilized RTS (yellow dot) at a riverbank. The white arrow (1) points to the clear dark boundary between the RTS and the waterbody, which together with the absence of sediment accumulation, indicates ongoing coastal thermo-erosion at the slump base. The purple arrows (2) point to the rugged outline of RTS and traces of erosive channels, indicating lateral thermo-erosion. The green arrow (3) points to the white patch of the remaining snowpack (nivation). (b) Stabilized RTS (yellow dot) at a lakeshore. The light blue arrows (4) point to the polygonal surface around the RTS and (5) the jagged outline of the headwall suggesting ice-wedge degradation. The orange arrows (6) point to the small black patches of thermokarst ponds within the RTS. ESRI basemap used has the following credits: Esri, DigitalGlobe, GeoEye, i-cubed, USDA FSA, USGS, AEX, Getmapping, Aerogrid, IGN, IGP, swisstopo, and the GIS User Community.



2.3 Accuracy assessment

2.3.RTS location accuracy

We compared the RTS point locations of our dataset with two sets of ground truth field data to estimate the accuracy of our mapped RTS point locations.

The first set of RTS locations was collected for the Vaskiny Dachi Research Station in Central Yamal by Khomutov et al. (2024) and included 158 points. The authors used satellite images of QuickBird-2 for 2010, GeoEye-1 and WorldView-2 for 2013, and WorldView-2, 3 for 2018, as well as the results of long-term field observation to map RTSs. Since the RTS mapping protocols can significantly affect the results (Nitze et al., 2024), we have adjusted these ground truth points to align with our mapping protocol in which one point stands for one RTS landform. When comparing our points to the ground truth collected, we observed inconsistencies in mapping RTS points. For example, while the ground truth dataset might contain two or three points for an RTS landform, our approach would place only one. To account for these differences, we recalibrated the dataset and calculated accuracy statistics for both the original (unadjusted) and adjusted RTS points (Table 1).

Two RTS surveys were conducted during helicopter flights in 2020 and 2023. We manually identified the exact locations of aerial photos and created another RTS dataset. We then used it to perform an accuracy analysis in the central Gydan Peninsula (Fig 9b, c). These points were also adjusted to our RTS mapping protocol, and the accuracy statistics were calculated for both versions (Table 1). The performance of our dataset was evaluated using precision, recall, and F1-score, which integrates both measures. In this context, precision refers specifically to the metric used in the F1-score calculation and should not be confused with measurement precision, as no measurements were performed. Precision is calculated as the proportion of correctly identified (true positive) RTS points when compared to the ground truth RTS points, among all mapped RTS points in the dataset. Recall represents the proportion of correctly identified RTS points relative to the total number of RTS points in the ground truth dataset. The F1-score is the harmonic mean of precision and recall, providing a balanced evaluation of both false positives and false negatives.




Figure 9 Field validation: (a) Locations of the Vaskiny Dachi research station with field survey area on Yamal Peninsula and helicopter survey area on central Gydan Peninsula, basemap: ESRI; (b) Photo of RTSs from the helicopter taken by Artem Khomutov, July 2023; (c) the same RTSs marked with the yellow point on the ESRI basemap, WorldView-2 24 July 2019. ESRI basemap used in (a) and (c) has the following credits: Esri, DigitalGlobe, GeoEye, i-cubed, USDA FSA, USGS, AEX, Getmapping, Aerogrid, IGN, IGP, swisstopo, and the GIS User Community.

Table 1 Number of RTSs used for the location accuracy analysis. The unadjusted number of RTSs represents the initial amount of RTSs in the ground truth datasets. The adjusted number of RTSs represents the amount of RTSs in the ground truth datasets adapted to the RTS mapping protocol applied for manual collection.

Vaskiny Dachi Research Station Survey 2024		Gydan Helicopter Survey 2020		Gydan Helicopter Survey 2023	
unadjusted	adjusted	unadjusted	adjusted	unadjusted	adjusted
158	132	60	39	12	12

2.3.2 Classification accuracy

To assess the subjectivity of the classification, we conducted an experiment in which five co-authors of this study were tasked with classifying a subsample of 120 randomly stratified RTS points that equally covered all three types of morphology. The decision-tree schemes and the collection of screenshots of different RTSs were used as supportive materials (see Supplement).



We calculated the proportion of the same classifications by 5 co-authors compared to the original dataset and Jensen-Shannon distances explaining the deviation of classifications.

3. Results

3.1 RTS points

The dataset is presented in a GeoPackage vector file of point geometry with 6168 RTS point locations. Mapped RTSs were distributed unevenly, covering Tazovsky Peninsula where no RTS were found, the Yamal Peninsula except its northern part, and covering the Gydan Peninsula except its southern part (Fig.10a). RTSs were significantly clustered according to Ripley's K function on a wide range of distances (p-value=0.001). Ripley's K function determines whether spatial points have a random, dispersed, or cluster distribution over a certain distance or scale (Dixon, 2001). The majority of areas of both peninsulas had less than 20 RTSs per 30*30 km hexagon grid cell, indicating distinct hotspots of RTS occurrence with more than 100 RTSs per grid cell. The main areas with high RTS density were the western part of central Yamal and the area between the southern-western and north-eastern parts of central Gydan. On Gydan, they clustered along a distinct linear feature on its southern edge, south of which RTSs abruptly become almost absent (Fig.10b).




Figure 10 Distribution of all mapped RTSs: (a) Manually mapped RTSs (purple dots); (b) Density map of RTSs per 30 × 30 km hexagonal grid cell. Projection: WGS 84 UTM Zone 43. Basemap: OSM Standard.



3.2 Terrain position

More than 75% of all RTSs were found at lakeshores (Fig.11a). The high-density areas of lakeshore RTSs correspond to RTS occurrence hotspots in the western part of central Yamal and the area between the south-western and north-eastern parts of central Gydan (Fig.11c).

The density of RTSs at the sea coasts was mostly less than 10 RTSs per grid cell. The highest density of coastal RTSs was found along the northern shores of Yuribei Bay in south-western Yamal (Fig.11b). For RTSs along river banks, gullies, and slopes, the predominating values of density were less than 10 RTSs per grid cell, not showing any spatial clustering (Appendix A).




Figure 11 Distribution of all mapped RTSs: (a) Manually mapped RTSs classified by location; Density maps of RTSs per 30 × 30 km hexagonal grid cell located along the (b) seacoast and (c) lakeshores. Projection: WGS 84 UTM Zone 43. Basemap: OSM Standard.



3.3 Morphology

The majority (72%) of RTSs were classified as thermocirques, one-quarter of all RTSs are combined landforms, and less than 3% were classified as thermoterraces (Fig.12a). The majority of RTSs in all categories have a spatial density of less than 15 RTSs per grid cell.




Figure 12 Distribution of all mapped RTSs: (a) Manually mapped RTSs classified by morphology; Density maps of RTSs per 30 × 30 km hexagonal grid cell classified as (b) thermocirque, (c) thermoterrace, and (d) a combination of both. Projection: WGS 84 UTM Zone 43. Basemap: OSM Standard.



Thermocirques were highly concentrated in hotspot areas of general RTS abundance (Fig. 10b). Combination landforms followed the high RTS abundance pattern mostly in the Gydan Peninsula but less so on the Yamal Peninsula. In contrast, thermoterraces lacked distinct high-density hotspots.

3.4 Spatial organization

More than half of all RTSs (64%) were classified as complex landforms and slightly more than one-third (36%) as single landforms (Fig. 13). Both complex and single landforms followed the general spatial distribution patterns, with high-density areas being located in the western part of the central Yamal Peninsula and the southern-western and north-eastern parts of the central Gydan Peninsula. The most frequent density range for both classes was less than 10 RTSs per grid cell.




Figure 13 Distribution of all mapped RTSs: (a) Manually mapped RTSs classified by spatial organization; Density maps of RTS per 30 × 30 km hexagonal grid cell classified by spatial organization as (b) single or (c) complex landforms. Projection: WGS 84 UTM Zone 43. Basemap: OSM Standard.

3.5 Concurrent processes

More than half (53.8%) of all RTSs were found to have at least one concurrent process detected, more than a third (33.4%) of all RTSs showed only one process detected, while much fewer RTSs demonstrated two or more processes detected at the same time (Fig. 14a). Lateral thermo-erosion and thermokarst were two very abundant RTS-concurrent processes (Fig. 14b). For the



cases where only one process was detected per RTS, there was a predominance of thermokarst (38%) followed by lateral thermo-erosion processes (30%) (Fig. 14c). Using chord diagrams (Fig. 14d, e, f) allowed a depiction of the co-occurrence of concurrent processes estimated for the cases when two, three, or four processes were detected for RTS. In general, the co-occurrence of the concurrent processes shows different results depending on the cases of the amount of the processes detected. There was a clear trend of the co-occurrence of nivation and lateral thermo-erosion among all 3 cases (Fig. 14d, e, f). The co-occurrence of lateral thermo-erosion and ice-wedge erosion gradually increased with more processes detected. The co-occurrence of the nivation and the coastal thermo-erosion, when only 2 processes are detected, was relatively low but increased significantly with more processes detected. The presence of thermokarst processes, in general, decreased with more processes detected.




Figure 14 Results of concurrent processes detected for each RTS: (a) Pie-chart of the number of concurrent processes detected for each RTS; (b) Histogram representing the total count of all concurrent processes identified in mapped RTS; (c) Histogram representing the distribution of concurrent processes when only 1 process per RTS was detected. Chord diagrams representing the occurrence of concurrent processes in the case when (d) two concurrent processes were detected, (e) three concurrent processes were detected, and (f) four concurrent processes were detected. The size of the outer frame corresponds to the count of each concurrent process. The lines connecting color-coded concurrent processes stand for the co-occurrence: the thicker the line, the higher the co-occurrence.




Figure 15 Density maps of RTs per 30×30 km hexagonal grid cell classified by the presence of concurrent process: (a) Lateral thermo-erosion; (b) Coastal thermo-erosion; (c) Ice wedge erosion; (d) Nivation; (e) Thermokarst subsidence. Projection: WGS 84 UTM Zone 43. Basemap: OSM Standard.

RTs attributed with concurrent processes exhibit low densities, with fewer than 5 RTs per grid cell, regardless of the type of concurrent process (Fig. 15). RTs with lateral thermo-erosion detected had higher densities in the western part of the central Yamal Peninsula and the central and northern Gydan Peninsula, with a hotspot in central Gydan Peninsula (Fig. 15a). RTs with concurrent coastal thermo-erosion had higher densities in the western part of central Yamal and the north-western Gydan peninsulas, with three hotspots located at south-western part of central Yamal Peninsula, and northern and north-western Gydan Peninsula. (Fig. 15b). In general, the spatial distribution of RTs with coastal thermo-erosion did not follow the main spatial patterns detected in the Fig. 10b. RTs with ice wedge erosion had higher densities on the northern Gydan Peninsula and rather lower densities on the Yamal Peninsula, with one hotspot located on central Yamal Peninsula (Fig. 15c).



The spatial distribution of RTSs with concurrent ice wedge erosion also did not follow the main spatial patterns detected in Fig. 10b. RTSs with nivation had higher densities in central and northern Gydan Peninsula (more than 30 RTSs per grid cell) and rather lower (less than 15 RTSs per grid cell) densities on Yamal Peninsula (Fig. 15d). There were four hotspots: one on central Yamal Peninsula and three on central Gydan Peninsula. The spatial distribution of RTSs with nivation also did not follow the main spatial patterns detected in Fig. 10b. RTSs with concurrent thermokarst did follow the main spatial patterns detected in Fig. 10b and thus had higher densities and some hotspots in the western part of central Yamal Peninsula and the area between the southern-western and north-eastern parts of central Gydan Peninsula (Fig. 15e).

4. Discussion

4.1 Data limitations

The manual collection of RTS points using the ESRI satellite basemap was effective across a large region but also had several limitations. One challenge was the resolution and zoom limitations, as the minimum detectable landform width was 20 meters, potentially excluding smaller features. Seasonal variability of the images in the ESRI satellite basemap further complicated the process, with snowpacks identifiable only in summer images, excluding all autumn (September) imagery. On the other hand, more extensive snow cover on certain images obscured some areas, hindering the accurate inventory of RTS and their attributes in these regions. Additionally, visual artifacts (blur, glare, clouds, contrails) in some imagery led to the omission of some cells, though this accounted for less than 0.5% of the total dataset.

Temporal constraints posed another issue, as working with a single satellite image captured at a specific time could mean that some features were not visible or detectable under those conditions, leading to potential underrepresentation of RTS features. The rapid evolution of RTS added difficulty, with two RTSs of a single morphology potentially merging into a complex morphology, creating challenges in morphology classification. Similar challenges were reported in the literature (Huang et al., 2020; Rodenhizer et al., 2024). Additionally, updates to the ESRI satellite basemap during the mapping effort sometimes introduced inconsistencies across different stages of our workflow, e.g. between the initial mapping of RTS as points, the subsequent addition of attributes, and the later correction loop (Fig. 2). To alleviate some of these challenges, we effectively used the ESRI Wayback time series to verify uncertain landforms or attributes.

Visual identification also had several challenges. Stabilized RTSs were difficult to recognize. Challenges were also faced when classifying partially stabilized RTS. The limitations concerning distinguishing slowly stabilizing slumps from stabilized slumps using optical data were also reported in the literature (Bernhard et al., 2020). The sediment accumulation as a secondary indicator for coastal thermo-erosion was found debatable due to its temporary nature. Some landforms, such as curved riverbanks, wave-cut lakeshores, active layer detachments (ALDs), and first-stage thermokarst mound (baydzherakh) development, could have been easily misclassified as RTS, leading to false positives.



Subjectivity was another significant limitation. Atypical for this area, Yedoma RTSs in the northern Gydan region differed significantly in appearance from other RTSs. Yedoma RTS's visual characteristics were not properly addressed in the initial visual identification protocol, leading to potential misidentifications.

4.2 Accuracy

Human subjectivity, even if mapping is conducted by experienced researchers, can influence the results and contribute to dataset uncertainties. For RTS mapping, this has been demonstrated before in a mapping exercise with multiple operators with varying degrees of experience (Nitze et al., 2024). Our subjectivity assessment using a subset of 120 RTS samples revealed that 3–16.6% were classified as non-RTS, with an average false positive rate of approximately 8.5% and a median of 4.1%. Consequently, the accuracy of our dataset based on this experiment averages around 0.91.

The degree of classification similarity among the five co-authors, compared to the original dataset, exhibited a clear trend influenced by spatial organization, morphology, and two concurrent processes—coastal thermo-erosion and lateral thermo-erosion—which were generally the most subjective. Spatial organization emerged as the most subjective parameter, with classifications showing the alignment in only half of the 120 sample points on average (Fig. 16a).

To further quantify classification variability, we calculated Jensen-Shannon distances (Fig. 16b), a metric for measuring similarity between probability distributions. This value ranges from 0.0, indicating identical distributions, to 1.0, representing completely distinct distributions. The results confirmed the overall trend of morphology, coastal thermo-erosion, and lateral thermo-erosion being the most subjective parameters, except for spatial organization, which showed minor differences in probability distributions. Coastal thermo-erosion exhibited the highest variation in classification probability distributions, likely due to two distinct hotspots observed in the heatmap.

Overall, the probability distributions of most classified parameters were either highly or moderately similar to those in the original dataset. This suggests a generally consistent perception of RTS classification among the co-authors in the experiment. RTS location accuracy was estimated for the area around the Vaskiny Dachi research station in central Yamal and central Gydan Peninsulas, with helicopter surveys conducted in 2020 and 2023 (see Appendix B). RTS location accuracy assessments for all areas revealed very high precision compared to the ground truth, confirming the reliability of the dataset (Table 2). A relatively low recall, even after applying mapping style adjustments, indicates an approximate 50% underestimation of small RTSs in the study area (Table 2) primarily due to the reasons described in the Data Limitations section (see Sect. 4.1). Please, note that in this context, precision specifically refers to the metric used in the F1-score calculation and should not be mistaken for measurement precision, as no actual measurements were conducted.




Figure 16 Classification subjectivity assessment: (a) Heatmap of the proportion of similar classifications by five co-authors compared to the classification in the dataset.; (b) Heatmap of Jensen-Shannon distances explaining deviation of classifications by five co-authors compared to the classification in the dataset.



Table 2 Average results of RTS location accuracy assessment for all three sets of ground truth field data: central Yamal and central Gydan (2020 and 2023). The adjusted value represents the accuracy measure calculated by comparing our dataset to the ground truth datasets adapted to the RTS mapping protocol applied for manual collection. The unadjusted value represents the accuracy measure calculated by comparing our dataset to the original ground truth datasets.

Average results for all three sets of ground truth field data		
	Adjusted to the mapping style	Unadjusted to the mapping style
Precision	0.96	0.96
Recall	0.44	0.38
F1 score	0.60	0.54

The relatively low F1 scores observed in our study can be attributed primarily to high underestimation (i.e., low recall) when compared to field data. Manual mapping of RTS using remote sensing data is often regarded as the most accurate approach (Swanson and Nolan, 2018; Segal et al., 2016a, b; Young et al., 2022; Luo et al., 2022). Efforts to enhance accuracy, particularly in terms of precision, have been made by incorporating multi-year datasets (Huang et al., 2021) and conducting multiple rounds of expert review (Segal et al., 2016b; Young et al., 2022). To ensure the reliability of manual mapping, Young et al. (2022) employed aerial field survey data for visual validation; however, their study did not report the initial recall of manual RTS mapping against field observations.

To the best of our knowledge, there are no existing studies that quantitatively assess the recall uncertainty of RTS manual mapping using remote sensing compared to field data, particularly over large spatial extents. Lewkowicz and Way (2019) attempted to estimate recall accuracy for manual RTS mapping in Banks Island, Canada (70000 km²), but their evaluation was based on a comparison with another remote sensing dataset rather than ground-based field observations. This limitation is largely due to the challenges associated with field data collection in remote study areas. Moreover, since field data provides only a single snapshot in time, some RTS classified as false positives based on remote sensing data may be true RTS that were simply not captured in the field dataset.

Despite these uncertainties, manually mapped RTS datasets serve as validation sources for automated deep-learning-based mapping algorithms (Nitze et al., 2021; Yang et al., 2023; Xia et al., 2022; Huang et al., 2021). Notably, relatively high F1 scores ($F1 > \sim 0.7$) for automated RTS mapping have been reported, but these assessments were primarily conducted against internal training datasets covering limited spatial extents and derived from manual mapping rather than field data (Huang et al., 2020; Nitze et al., 2021; Witharana et al., 2022; Yang et al., 2023).



Our findings demonstrate that manual mapping using remote sensing data cannot be considered a definitive ground truth and is associated with a certain degree of inaccuracy, particularly concerning recall. Our accuracy assessment highlights the overall subjectivity in defining RTS morphology and spatial organization. These parameters critically influence what is visually identified as RTS in satellite imagery. This subjectivity aligns with previous RTS mapping experiments, where "mapping style" and the scientific background of domain experts were found to impact RTS delineation (Nitze et al., 2024). Our results demonstrate that, despite standardized instructions, both morphology and spatial organization remain the most subjective parameters in RTS classification.

4.3 Data applicability

The collected data on RTSs holds significant potential for future applications and research across various disciplines. It can serve as a foundation for a more detailed characterization of the permafrost region. The spatial distribution and clustering of RTSs in West Siberia, combined with cryostratigraphic and geomorphological analyses, can help unravel driving processes and improve our understanding of these dynamic landforms.

This dataset can also guide further research efforts, such as field surveys aimed at monitoring cryogenic processes as well as studies to uncover the ground ice origin. In addition, it provides a valuable reference for ground-truthing in machine learning applications, enabling more accurate automated remote sensing classifications and predictive modeling.

The dataset is particularly relevant to ecologists, biogeochemists, geomorphologists, climatologists, permafrost scientists, hazard researchers, and remote sensing specialists. This data can also be useful in the context of managing permafrost-related risks and planning sustainable development in vulnerable regions.

5. Data availability

The dataset is available at Nesterova et al., 2025 (<https://doi.pangaea.de/10.1594/PANGAEA.974406>).

6. Conclusions

In this study, we present the first large-scale manual RTS mapping effort with accuracy assessments based on field data. We present a comprehensive, manually mapped dataset of 6168 current retrogressive thaw slumps (RTS) for a large region in the West Siberian Arctic. Each RTS in the dataset was classified according to its morphology, spatial organization, terrain position, and concurrent permafrost relief-forming processes. Accuracy assessments with independent field data and expert knowledge indicate a high accuracy of the dataset while also highlighting some subjectivity in the classifications. Due to resolution limitations in the satellite image basemaps used for mapping, the dataset may underestimate the occurrence of small RTS in the region, resulting in an overall conservative estimate. Despite these constraints, our new RTS inventory offers valuable



insights for a wide range of research fields aiming at further investigations of RTS formation and dynamics, permafrost-climate interactions, permafrost-ecosystem feedbacks, and ground ice distribution in West Siberia.

Appendices

Appendix A




Figure A1. Density maps of RTS points counted per 30 × 30 km hexagonal grid cell located at the (a) river, (b) gully, and (c) slope. Projection: WGS 84 UTM Zone 43. Basemap: OSM Standard.



Appendix B

Table B Results of RTS location accuracy assessment for all three sets of ground truth field data: central Yamal Vaskiny Dachi research station and central Gydan Helicopter Survey (2020 and 2023). The adjusted value represents the accuracy measure calculated by comparing our dataset to the ground truth datasets adapted to the RTS mapping protocol applied for manual collection. The unadjusted value represents the accuracy measure calculated by comparing our dataset to the original ground truth datasets.

Vaskiny Dachi research station, central Yamal		
	Adjusted to the mapping style	Unadjusted to the mapping style
Precision	0.88	0.88
Recall	0.44	0.37
F-1 score	0.59	0.52
Gydan Helicopter Survey 2020		
	Adjusted to the mapping style	Unadjusted to the mapping style
Precision	1	1
Recall	0.3	0.2
F-1 score	0.46	0.33
Gydan Helicopter Survey 2023		
Precision		1
Recall		0.58
F-1 score		0.73



Author contribution

NN: conceptualization, resources (data collection and correction), investigation, and writing (original draft preparation). IT: conceptualization, resources (data collection and correction), and writing (review and editing). ML: conceptualization, supervision, and writing (review and editing). AKh: resources (field data collection) and writing (review and editing). AK: writing (review and editing). IN: writing (review and editing). GG: supervision, and writing (review and editing).

Supplement

The supplement of the decision-tree schemes and the collection of screenshots of different RTSs to help classify RTSs in West Siberia is available online at Zenodo (Nesterova and Tarasevich, 2025, <https://doi.org/10.5281/zenodo.15063753>).

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgments

NN was funded by a DAAD fellowship (“STIBET-I”). IT, ML, and AKh were funded by the state assignment of the Ministry of Science and Higher Education of the Russian Federation (grant no. FWRZ-2021-0012). IN and GG were funded by the German Federal Ministry for Economic Affairs and Climate Action (BMWK) project ML4EARTH, European Space Agency (ESA) CCI+ Permafrost, as well as NSF (NSF Opp: #1927872 and #2052107) and google.org “Permafrost Discovery Gateway”. AK was funded by the Lomonosov Moscow State University state assignment “The cryosphere evolution under climate change and anthropogenic impact” (#121051100164-0).

References

- Aastrup, P., Ackrén, M., Allard, M., Archambault, P., Arendt, K., Barrette, C., Bélanger, S., Bell, T., Berteaux, D., and Bjella, K.: AMAP 2017. Adaptation actions for a changing Arctic: Perspectives from the Baffin Bay/Davis Strait Region, 2018.
- ESRI World Imagery Wayback: <https://livingatlas.arcgis.com/wayback/>, last access: 1 October 2024.
- Babkina, E. A., Leibman, M. O., Dvornikov, Yu. A., Fakashchuk, N. Yu., Khairullin, R. R., and Khomutov, A. V.: Activation of Cryogenic Processes in Central Yamal as a Result of Regional and Local Change in Climate and Thermal State of Permafrost, Russian Meteorology and Hydrology, 44, 283–290, <https://doi.org/10.3103/S1068373919040083>, 2019.



- 480 Badu, Y. B.: Ice content of cryogenic strata (permafrost interval) in gas-bearing structures, Northern Yamal, Earth's
 481 Cryosphere, 19, 9–18, 2015.
- 482 Baulin, V. V. and Danilova, N. S.: West Siberia (In Russian), in: Fundamental of Geocryology. Regional and historical
 483 geogryology of the world, Moscow State University, Moscow, 252–271, 1998.
- 484 Baulin, V. V., Belopukhova, E. B., Dubikov, G. I., and Shmelev, L. M.: Geocryological conditions of western Siberia Lowland,
 485 Nauka, Moscow, Russia, 1967.
- 486 Bernhard, P., Zwieback, S., Leinss, S., and Hajnsek, I.: Mapping Retrogressive Thaw Slumps Using Single-Pass TanDEM-X
 487 Observations, IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing, 13, 3263–3280,
 488 <https://doi.org/10.1109/JSTARS.2020.3000648>, 2020.
- 489 Bernhard, P., Zwieback, S., Bergner, N., and Hajnsek, I.: Assessing volumetric change distributions and scaling relations of
 490 retrogressive thaw slumps across the Arctic, The Cryosphere, 16, 1–15, <https://doi.org/10.5194/tc-16-1-2022>, 2022.
- 491 Biskaborn, B. K., Smith, S. L., Noetzli, J., Matthes, H., Vieira, G., Streletskiy, D. A., Schoeneich, P., Romanovsky, V. E.,
 492 Lewkowicz, A. G., Abramov, A., Allard, M., Boike, J., Cable, W. L., Christiansen, H. H., Delaloye, R., Diekmann, B.,
 493 Drozdov, D., Etzelmüller, B., Grosse, G., Guglielmin, M., Ingeman-Nielsen, T., Isaksen, K., Ishikawa, M., Johansson, M.,
 494 Johannsson, H., Joo, A., Kaverin, D., Kholodov, A., Konstantinov, P., Kröger, T., Lambiel, C., Lanckman, J.-P., Luo, D.,
 495 Malkova, G., Meiklejohn, I., Moskalenko, N., Oliva, M., Phillips, M., Ramos, M., Sannel, A. B. K., Sergeev, D., Seybold, C.,
 496 Skryabin, P., Vasiliev, A., Wu, Q., Yoshikawa, K., Zheleznyak, M., and Lantuit, H.: Permafrost is warming at a global scale,
 497 Nature Communications, 10, 264, <https://doi.org/10.1038/s41467-018-08240-4>, 2019.
- 498 Brown, J., Hinkel, K. M., and Nelson, F. E.: The circumpolar active layer monitoring (calm) program: Research designs and
 499 initial results 1, Polar Geography, 24, 166–258, <https://doi.org/10.1080/10889370009377698>, 2000.
- 500 Cassidy, A. E., Christen, A., and Henry, G. H. R.: Impacts of active retrogressive thaw slumps on vegetation, soil, and net
 501 ecosystem exchange of carbon dioxide in the Canadian High Arctic, Arctic Science, 3, 179–202, [https://doi.org/10.1139/as-](https://doi.org/10.1139/as-2016-0034)
 502 [2016-0034](https://doi.org/10.1139/as-2016-0034), 2017.
- 503 Dixon, P. M.: Ripley's K function, in: Encyclopedia of Environmetrics, vol. 3, John Wiley & Sons, Ltd, Chichester, Chichester,
 504 1796–1803, 2002.
- 505 Günther, F., Overduin, P. P., Sandakov, A., Grosse, G., and Grigoriev, M. N.: Thermo-erosion along the yedoma coast of the
 506 Buor Khaya peninsula, Laptev Sea, east Siberia, Proceedings of the Tenth International Conference on Permafrost, Volume 1:
 507 International Contributions, 137–142, 2012.
- 508 Harris, S. A. and Permafrost Subcommittee, Associate Committee on Geotechnical Research, National Research Council of
 509 Canada (Eds.): Glossary of permafrost and related ground-ice terms, Ottawa, Ontario, Canada, 156 pp., 1988.



- 510 Huang, L., Luo, J., Lin, Z., Niu, F., and Liu, L.: Using deep learning to map retrogressive thaw slumps in the Beiluhe region
 511 (Tibetan Plateau) from CubeSat images, *Remote Sensing of Environment*, 237, 111534,
 512 <https://doi.org/10.1016/j.rse.2019.111534>, 2020.
- 513 Huang, L., Liu, L., Luo, J., Lin, Z., and Niu, F.: Automatically quantifying evolution of retrogressive thaw slumps in Beiluhe
 514 (Tibetan Plateau) from multi-temporal CubeSat images, *International Journal of Applied Earth Observation and*
 515 *Geoinformation*, 102, 102399, <https://doi.org/10.1016/j.jag.2021.102399>, 2021.
- 516 Huang, L., Willis, M. J., Li, G., Lantz, T. C., Schaefer, K., Wig, E., Cao, G., and Tiampo, K. F.: Identifying active retrogressive
 517 thaw slumps from ArcticDEM, *ISPRS Journal of Photogrammetry and Remote Sensing*, 205, 301–316,
 518 <https://doi.org/10.1016/j.isprsjprs.2023.10.008>, 2023.
- 519 Kerfoot, D. E.: The geomorphology and permafrost conditions of Garry Island, N.W.T., Text, 1969.
- 520 Khomutov, A., Leibman, M., Dvornikov, Yu., Gubarkov, A., Mullanurov, D., and Khairullin, R.: Activation of Cryogenic
 521 Earth Flows and Formation of Thermocirques on Central Yamal as a Result of Climate Fluctuations, in: *Advancing Culture of*
 522 *Living with Landslides. Proceedings of World Landslide Forum 4*, May 29 – June 2, 2017, Ljubljana, Slovenia. Vol.5,
 523 *Landslides in Different Environments*, Ljubljana, 209–216, 2017.
- 524 Khomutov, A. V., Babkina, E. A., Khairullin, R. R., and Dvornikov, Yu. A.: Factors of thermal denudation activation and
 525 thermicirques activity on central Yamal in 2010–2018 (In Russian), *Arctic and Antarctic Research*, 70, 222–237, 2024.
- 526 Lantuit, H. and Pollard, W. H.: Temporal stereophotogrammetric analysis of retrogressive thaw slumps on Herschel Island,
 527 Yukon Territory, *Natural Hazards and Earth System Sciences*, 5, 413–423, <https://doi.org/10.5194/nhess-5-413-2005>, 2005.
- 528 Lantz, T. C., Kokelj, S. V., Gergel, S. E., and Henry, G. H. R.: Relative impacts of disturbance and temperature: persistent
 529 changes in microenvironment and vegetation in retrogressive thaw slumps, *Global Change Biology*, 15, 1664–1675,
 530 <https://doi.org/10.1111/j.1365-2486.2009.01917.x>, 2009.
- 531 Leibman, M. O. and Kizyakov, A. I.: Cryogenic landslides of the Yamal and Yugorsky Peninsulas (In Russian), *Earth*
 532 *Cryosphere Institute SB RAS*, Moscow, 2007.
- 533 Leibman, M. O., Khomutov, A. V., Gubarkov, A. A., Dvornikov, Y. A., and Mullanurov, D. R.: The research station "Vaskiny
 534 Dachi", Central Yamal, West Siberia, Russia—a review of 25 years of permafrost studies, *Fennia-International Journal of*
 535 *Geography*, 193, 3–30, 2015.
- 536 Lewkowicz, A. G. and Way, R. G.: Extremes of summer climate trigger thousands of thermokarst landslides in a High Arctic
 537 environment, *Nature Communications*, 10, 1329, <https://doi.org/10.1038/s41467-019-09314-7>, 2019.
- 538 Luo, D., Wu, Q., Jin, H., Marchenko, S. S., Lü, L., and Gao, S.: Recent changes in the active layer thickness across the northern
 539 hemisphere, *Environmental Earth Sciences*, 75, 555, <https://doi.org/10.1007/s12665-015-5229-2>, 2016.



- 540 Luo, J., Niu, F., Lin, Z., Liu, M., Yin, G., and Gao, Z.: Inventory and Frequency of Retrogressive Thaw Slumps in Permafrost
 541 Region of the Qinghai–Tibet Plateau, Geophysical Research Letters, 49, e2022GL099829,
 542 <https://doi.org/10.1029/2022GL099829>, 2022.
- 543 Mackay, J. R.: Segregated epigenetic ice and slumps in permafrost, Mackenzie Delta area, NWT, Geographical Bulletin, 8,
 544 59–80, 1966.
- 545 Miner, K. R., Turetsky, M. R., Malina, E., Bartsch, A., Tamminen, J., McGuire, A. D., Fix, A., Sweeney, C., Elder, C. D., and
 546 Miller, C. E.: Permafrost carbon emissions in a changing Arctic, Nature Reviews Earth & Environment, 3, 55–67,
 547 <https://doi.org/10.1038/s43017-021-00230-3>, 2022.
- 548 Nesterova, N. and Tarasevich, I.: Decision tree for the classification of retrogressive thaw slumps in West Siberia. Zenodo,
 549 <https://doi.org/10.5281/zenodo.15063753>, 2025.
- 550 Nesterova, N., Leibman, M., Kizyakov, A., Lantuit, H., Tarasevich, I., Nitze, I., Veremeeva, A., and Grosse, G.: Review
 551 article: Retrogressive thaw slump characteristics and terminology, The Cryosphere, 18, 4787–4810, [https://doi.org/10.5194/tc-](https://doi.org/10.5194/tc-18-4787-2024)
 552 [18-4787-2024](https://doi.org/10.5194/tc-18-4787-2024), 2024.
- 553 Nesterova, N., Tarasevich, I., Leibman, M., Khomutov, A. V., Kizyakov, A., Nitze, I., and Grosse, G.: Manually mapped
 554 retrogressive thaw slumps in West Siberia, version 0.1 [dataset]. PANGAEA,
 555 <https://doi.pangaea.de/10.1594/PANGAEA.974406>, 2025.
- 556 Nesterova, N. B., Khomutov, A. V., Leibman, M. O., Safonov, T. A., and Belova, N. G.: The inventory of retrogressive thaw
 557 slumps (thermocirques) in the north of West Siberia based on 2016–2018 satellite imagery mosaic, Earth’s Cryosphere, 25,
 558 <https://doi.org/10.15372/KZ20210604>, 2021.
- 559 Nitze, I., Grosse, G., Jones, B. M., Romanovsky, V. E., and Boike, J.: Remote sensing quantifies widespread abundance of
 560 permafrost region disturbances across the Arctic and Subarctic, Nature Communications, 9, [https://doi.org/10.1038/s41467-](https://doi.org/10.1038/s41467-018-07663-3)
 561 [018-07663-3](https://doi.org/10.1038/s41467-018-07663-3), 2018.
- 562 Nitze, I., Heidler, K., Barth, S., and Grosse, G.: Developing and Testing a Deep Learning Approach for Mapping Retrogressive
 563 Thaw Slumps, Remote Sensing, 13, 4294, 2021.
- 564 Nitze, I., Van der Sluijs, J., Barth, S., Bernhard, P., Huang, L., Kizyakov, A., Lara, M. J., Nesterova, N., Runge, A., Veremeeva,
 565 A., Ward Jones, M., Witharana, C., Xia, Z., and Liljedahl, A. K.: A Labeling Intercomparison of Retrogressive Thaw Slumps
 566 by a Diverse Group of Domain Experts, Permafrost and Periglacial Processes, 36, 83–92, <https://doi.org/10.1002/ppp.2249>,
 567 2024a.



- 568 Nitze, I., Heidler, K., Nesterova, N., Küpper, J., Schütt, E., Hölzer, T., Barth, S., Lara, M. J., Liljedahl, A., and Grosse, G.:
 569 DARTS: Multi-year database of AI detected retrogressive thaw slumps (RTS) and active layer detachment slides (ALD) in
 570 hotspots of the circum-arctic permafrost region-v1, 2024b.
- 571 Obu, J.: How Much of the Earth's Surface is Underlain by Permafrost?, *Journal of Geophysical Research: Earth Surface*, 126,
 572 e2021JF006123, <https://doi.org/10.1029/2021JF006123>, 2021.
- 573 Pollard, W. H. and French, H. M.: A first approximation of the volume of ground ice, Richards Island, Pleistocene Mackenzie
 574 Delta, Northwest Territories, Canada, *Canadian Geotechnical Journal*, 17, 509–516, 1980.
- 575 Rodenhizer, H., Yang, Y., Fiske, G., Potter, S., Windholz, T., Mullen, A., Watts, J. D., and Rogers, B. M.: A Comparison of
 576 Satellite Imagery Sources for Automated Detection of Retrogressive Thaw Slumps, *Remote Sensing*, 16,
 577 <https://doi.org/10.3390/rs16132361>, 2024.
- 578 Runge, A., Nitze, I., and Grosse, G.: Remote sensing annual dynamics of rapid permafrost thaw disturbances with LandTrendr,
 579 *Remote Sensing of Environment*, 268, 112752, <https://doi.org/10.1016/j.rse.2021.112752>, 2022.
- 580 Schuur, E. A. G., McGuire, A. D., Schädel, C., Grosse, G., Harden, J. W., Hayes, D. J., Hugelius, G., Koven, C. D., Kuhry,
 581 P., Lawrence, D. M., Natali, S. M., Olefeldt, D., Romanovsky, V. E., Schaefer, K., Turetsky, M. R., Treat, C. C., and Vonk, J.
 582 E.: Climate change and the permafrost carbon feedback, *Nature*, 520, 171–179, <https://doi.org/10.1038/nature14338>, 2015.
- 583 Segal, R. A., Lantz, T. C., and Kokelj, S. V.: Acceleration of thaw slump activity in glaciated landscapes of the Western
 584 Canadian Arctic, *Environmental Research Letters*, 11, 34025, 2016a.
- 585 Segal, R. A., Kokelj, S. V., Lantz, T. C., Pierce, K. L., Durkee, K., Gervais, S., Mahon, E., Snijders, M., Buysse, J., and
 586 Schwarz, S.: NWT Open Report 2016-023 Mapping of Terrain Affected by Retrogressive Thaw Slumping in Northwestern
 587 Canada, NWT Open Report 2016-023, 5 p., 2016b.
- 588 Smith, S. L., O'Neill, H. B., Isaksen, K., Noetzli, J., and Romanovsky, V. E.: The changing thermal state of permafrost, *Nature*
 589 *Reviews Earth & Environment*, 3, 10–23, <https://doi.org/10.1038/s43017-021-00240-1>, 2022.
- 590 Streletskaya, I., Gusev, E., Vasiliev, A., Oblogov, G., and Molodkov, A.: Pleistocene-Holocene palaeoenvironmental records
 591 from permafrost sequences at the Kara Sea coast (NW Siberia, Russia), *Geography, environment, sustainability*, 6, 60–76,
 592 2013.
- 593 Streletskaya, I. D., Vasiliev, A. A., and Vanstein, B. G.: Erosion of Sediment and Organic Carbon from the Kara Sea Coast,
 594 *Arctic, Antarctic, and Alpine Research*, 41, 79–87, <https://doi.org/10.1657/1523-0430-41.1.79>, 2009.
- 595 Swanson, D. K. and Nolan, M.: Growth of retrogressive thaw slumps in the Noatak Valley, Alaska, 2010–2016, measured by
 596 airborne photogrammetry, *Remote Sensing*, 10, <https://doi.org/10.3390/rs10070983>, 2018.



- 597 Thienpont, J. R., Ruehland, K. M., Pisaric, M. F. J., Kokelj, S. V., Kimpe, L. E., Blais, J. M., and Smol, J. P.: Biological
 598 responses to permafrost thaw slumping in Canadian Arctic lakes, *Freshwater Biology*, 58, 337–353, 2013.
- 599 Vasiliev, A. A., Drozdov, D. S., Gravis, A. G., Malkova, G. V., Nyland, K. E., and Streletskiy, D. A.: Permafrost degradation
 600 in the Western Russian Arctic, *Environmental Research Letters*, 15, 045001, <https://doi.org/10.1088/1748-9326/ab6f12>, 2020.
- 601 Ward Jones, M. K., Pollard, W. H., and Jones, B. M.: Rapid initialization of retrogressive thaw slumps in the Canadian high
 602 Arctic and their response to climate and terrain factors, *Environmental Research Letters*, 14, [https://doi.org/10.1088/1748-](https://doi.org/10.1088/1748-9326/ab12fd)
 603 [9326/ab12fd](https://doi.org/10.1088/1748-9326/ab12fd), 2019.
- 604 Witharana, C., Udawalpola, M. R., Liljedahl, A. K., Jones, M. K. W., Jones, B. M., Hasan, A., Joshi, D., and Manos, E.:
 605 Automated Detection of Retrogressive Thaw Slumps in the High Arctic Using High-Resolution Satellite Imagery, *Remote*
 606 *Sensing*, 14, 4132, <https://doi.org/10.3390/rs14174132>, 2022.
- 607 Xia, Z., Huang, L., Fan, C., Jia, S., Lin, Z., Liu, L., Luo, J., Niu, F., and Zhang, T.: Retrogressive thaw slumps along the
 608 Qinghai–Tibet Engineering Corridor: a comprehensive inventory and their distribution characteristics, *Earth Syst. Sci. Data*,
 609 14, 3875–3887, <https://doi.org/10.5194/essd-14-3875-2022>, 2022.
- 610 Yang, Y., Rogers, B. M., Fiske, G., Watts, J., Potter, S., Windholz, T., Mullen, A., Nitze, I., and Natali, S. M.: Mapping
 611 retrogressive thaw slumps using deep neural networks, *Remote Sensing of Environment*, 288, 113495,
 612 <https://doi.org/10.1016/j.rse.2023.113495>, 2023.
- 613 Young, J. M., Alvarez, A., van der Sluijs, J., Kokelj, S. V., Rudy, A., McPhee, A., Stoker, B. J., Margold, M., and Froese, D.:
 614 Recent Intensification (2004–2020) of Permafrost Mass-Wasting in the Central Mackenzie Valley Foothills Is a Legacy of
 615 Past Forest Fire Disturbances, *Geophysical Research Letters*, 49, e2022GL100559, <https://doi.org/10.1029/2022GL100559>,
 616 2022.
- 617