

Mapping the global distribution of lead and its isotopes in seawater with explainable machine learning

Arianna Olivelli^{1,2,*}, Rossella Arcucci¹, Mark Rehkämper¹, Tina van de Flierdt¹

¹ Grantham Institute for Climate Change and the Environment, Imperial College London, South Kensington Campus, London SW7 2AZ, United Kingdom

² Department of Earth Science and Engineering, Imperial College London, South Kensington Campus, London SW7 2AZ, United Kingdom

* Author now at the Flanders Marine Institute (VLIZ), Jacobsenstraat 1, 8400, Ostend, Belgium

Correspondence to: Arianna Olivelli (arianna.olivelli@vliz.be)

Abstract. Lead (Pb) and its isotopes are powerful tools to study the pathways of Pb pollution from land to sea and, simultaneously, investigate biogeochemical processes in the ocean. However, the scarcity and sparsity of *in situ* measurements of Pb concentrations and isotope compositions do not allow for a comprehensive understanding of Pb pollution pathways and biogeochemical cycling on a global scale. Here, we present three machine learning models developed to map seawater Pb concentrations and isotope compositions leveraging the global GEOTRACES dataset as well as historical data. The models use climatologies of oceanographic and atmospheric variables as features from which to predict Pb concentrations, ²⁰⁶Pb/²⁰⁷Pb, and ²⁰⁸Pb/²⁰⁷Pb. Using Shapley Additive Values (SHAP), we found that seawater temperature, atmospheric dust and black carbon, and salinity are the most important features for predicting Pb concentrations. Dissolved oxygen concentration, salinity, temperature, and atmospheric dust are the most important features for predicting ²⁰⁶Pb/²⁰⁷Pb, while atmospheric black carbon and dust, seawater temperature, and surface chlorophyll-a for ²⁰⁸Pb/²⁰⁷Pb. In line with observations, our model outputs show that (i) the surface Indian Ocean has the highest levels of pollution, (ii) pollution from previous decades is sinking in the North Atlantic and Pacific Ocean, and (iii) waters characterised by highly anthropogenic Pb isotope fingerprints are spreading from the Southern Ocean throughout the Southern Hemisphere at intermediate depths. By analysing the uncertainty associated with our maps, we identified the Southern Ocean as the key area to prioritise in future sampling campaigns. Our datasets, models and their outputs, in the form of Pb concentration, ²⁰⁶Pb/²⁰⁷Pb, and ²⁰⁸Pb/²⁰⁷Pb climatologies, are made freely available to the community at Olivelli et al. (2024a, <https://doi.org/10.5281/zenodo.14261154>) and https://github.com/OlivelliAri/Pb-ML_GEOTRACES.

Deleted: ,

Deleted:

Formatted: Font: Not Italic

Deleted: a.olivelli21@imperial.ac.uk

Deleted: a

Deleted: tool

Deleted: Our

Deleted: a

Deleted: fingerprint

1 Introduction

45 The low concentrations at which lead (Pb) is present in the ocean (on the order of parts per trillion) and the ubiquity of Pb contamination during sampling and sample processing did not allow for successful measurements of seawater Pb concentration and isotope composition until the late 1970s (Schaule and Patterson, 1981, 1983). Indeed, since the early 1900s, human activities have caused an increase in Pb concentrations in the environment, both on land and in the ocean, which altered the background levels and the biogeochemical cycling of Pb (Boyle et al., 2014; Nriagu, 1979; Nriagu and Pacyna, 1988; Pacyna and Pacyna, 2001).

50 This increase in Pb concentrations has been accompanied by a shift in the Pb isotope compositions of seawater, reflecting the predominance of anthropogenically-sourced over naturally-sourced Pb (Boyle et al., 2014). In fact, the relative abundances of stable Pb isotopes (^{204}Pb , ^{206}Pb , ^{207}Pb , and ^{208}Pb), expressed as ratios such as $^{206}\text{Pb}/^{207}\text{Pb}$ and $^{208}\text{Pb}/^{207}\text{Pb}$, can be used to identify and quantify the contributions of the different sources of Pb (Reuer and Weiss, 2002).

55 In the years and decades that followed the first successful measurements of Pb and its isotopes in seawater, sampling efforts mainly concentrated in the North Atlantic Ocean (Alleman et al., 1999; Boyle et al., 1986; Helmers et al., 1991; Pohl et al., 1993; Shen and Boyle, 1988; Sherrell et al., 1992; Véron et al., 1994, 1998, 1999). This basin, surrounded by the early-developed economies of North America and Western Europe, initially saw a sharp increase in pollution followed by a decrease thanks to environmental policies that phased out and banned the use of leaded gasoline since the 1970s (Bridgestock et al., 2016; Weiss et al., 2003).

60 A major breakthrough for the understanding of the marine biogeochemistry and pollution of Pb on a global scale came with the international marine geochemistry programme GEOTRACES, which has run since 2006 and included Pb concentrations and isotope compositions as key parameters of interest to measure on all its cruise sections (GEOTRACES Planning Group, 2006). However, despite these great efforts, the majority of the world's ocean remains unsampled for Pb concentrations and isotope compositions, especially in the Southern Hemisphere (Fig. 1).

65

Deleted: ¶

Deleted: until 1963 (Tatsumoto & Patterson, 1963)

Deleted: 1976

Deleted: &

Deleted:)

Deleted: &

Deleted: &

Deleted: &

Deleted: &




Figure 1. Geographical distribution of all samples in the Pb concentration (A), $^{206}\text{Pb}/^{207}\text{Pb}$ (C), and $^{208}\text{Pb}/^{207}\text{Pb}$ (E) datasets. Panels A, C, and E include all samples from the top 100 m. Panels B, D, and F represent the frequency distribution of sampling depths for panels A, C, and E, respectively.

80 In this context, modelling studies represent a powerful tool to expand our knowledge about the distribution and biogeochemical cycle of Pb, complementary to *in situ* observations. While deterministic models require a good understanding of the processes at play in the biogeochemical of Pb, machine learning (ML) models do not require such prior knowledge as they are data-driven (Glover et al., 2011). Therefore, the latter can be regarded as a computationally efficient way to investigate the

distribution of Pb and its isotopes in the global ocean without the need to explicitly parametrise causal relationships derived
85 from fundamental knowledge of the biogeochemical cycle of Pb.

In recent years, ML algorithms have been employed to investigate several topics within chemical oceanography, including the distribution and cycling of tracers, such as iron (Huang et al., 2022), zinc (Roshan et al., 2018), copper (Roshan et al., 2020), and barium (Mete et al., 2023), iodide (Sherwen et al., 2019), and nitrogen isotopes (Rafter et al., 2019). A thorough overview of the use and potential of ML in ocean sciences is provided in Sonnewald et al. (2021).

90 In this study, we leverage the growing GEOTRACES dataset, in combination with climatologies of several oceanographic and atmospheric variables, to map the distribution of Pb and its isotopes in the global ocean using the ML regression algorithm XGBoost. With this approach, we aim to (i) produce three-dimensional global maps of Pb concentrations and isotope compositions ($^{206}\text{Pb}/^{207}\text{Pb}$ and $^{208}\text{Pb}/^{207}\text{Pb}$) to use as a benchmark for future levels of pollution, and to (ii) identify areas where to concentrate sampling efforts in the upcoming years and decades by analysing the uncertainty of our model outputs. We
95 make the resulting data products available to the community, in the form of mean global climatologies, and envisage updating the models' outputs as new observations become available.

2 Materials and Methods

Regression ML models require training on existing observations of a target variable (to be predicted by the model) alongside a number of features (i.e. ancillary variables) in order to make accurate predictions. Therefore, three different datasets were
100 compiled to develop three separate ML models to map the global distribution of Pb concentrations, $^{206}\text{Pb}/^{207}\text{Pb}$, and $^{208}\text{Pb}/^{207}\text{Pb}$, respectively. Each dataset includes information on 14 geographical, oceanographic and atmospheric features (Table 1) and a target variable (either Pb concentration, $^{206}\text{Pb}/^{207}\text{Pb}$, or $^{208}\text{Pb}/^{207}\text{Pb}$). The features included in the datasets were chosen due to their proven predictive power in other ML studies focusing on ocean chemistry (Huang et al., 2022; Mete et al., 2023; Roshan et al., 2018, 2020; Sherwen et al., 2019) and their likely connection to the biogeochemical cycle of Pb.

105 The three datasets and models were built following the same procedure, explained in sections 2.1 and 2.2. All data preparation, modelling and analyses were done in Python 3.11.5 in a Linux operating system.

110

Table 1. List of features used for dataset compilation and model development for Pb concentration, $^{206}\text{Pb}/^{207}\text{Pb}$, and $^{208}\text{Pb}/^{207}\text{Pb}$.

Feature name	Data source	Spatial resolution	Temporal resolution
Geographical features			
Latitude [$^{\circ}\text{N}$]*	Sampling information		
Longitude [$^{\circ}\text{E}$]*	Sampling information		
Depth [m]	Sampling information		
Distance from bottom depth [m]	Derived from sampling information		
Oceanographic features			
Temperature [$^{\circ}\text{C}$]	World Ocean Atlas 2018	$1^{\circ}\times 1^{\circ}$	Climatology, 01/2005 - 12/2017
Salinity	World Ocean Atlas 2018	$1^{\circ}\times 1^{\circ}$	Climatology, 01/2005 - 12/2017
Dissolved nitrate [$\mu\text{mol/kg}$]	World Ocean Atlas 2018	$1^{\circ}\times 1^{\circ}$	Climatology, all available data (1955 - 2017)
Dissolved phosphate [$\mu\text{mol/kg}$]	World Ocean Atlas 2018	$1^{\circ}\times 1^{\circ}$	Climatology, all available data (1955 - 2017)
Dissolved silicate [$\mu\text{mol/kg}$]	World Ocean Atlas 2018	$1^{\circ}\times 1^{\circ}$	Climatology, all available data (1955 - 2017)
Density anomaly (σ_{θ}) [kg/m^3]	World Ocean Atlas 2018	$1^{\circ}\times 1^{\circ}$	Climatology, 01/2005 - 12/2017
Dissolved oxygen [$\mu\text{mol/kg}$]	World Ocean Atlas 2018	$1^{\circ}\times 1^{\circ}$	Climatology, all available data (1955 - 2017)
Apparent oxygen utilization [$\mu\text{mol/kg}$]	World Ocean Atlas 2018	$1^{\circ}\times 1^{\circ}$	Climatology, all available data (1955 - 2017)
Mixed layer depth [m]	World Ocean Atlas 2018	$1^{\circ}\times 1^{\circ}$	Climatology, 01/2005 - 12/2017
Chlorophyll-a [mg/m^3]	CMEMS ^a	$0.25^{\circ}\times 0.25^{\circ}$	Monthly, 01/2005 - 12/2017 (transformed to climatology)
Atmospheric features			
Black carbon AOD ^b	CAMS ^c	$0.75^{\circ}\times 0.75^{\circ}$	Monthly, 01/2005 - 12/2017 (transformed to climatology)
Dust AOD ^b	CAMS ^c	$0.75^{\circ}\times 0.75^{\circ}$	Monthly, 01/2005 - 12/2017 (transformed to climatology)

* Variables used for dataset compilation and for an initial version of the models, but dropped for the final models (see more in SI note 1)

^a Copernicus Marine Environment Monitoring Service

^b Aerosol Optical Depth

^c Copernicus Atmosphere Monitoring Service

2.1 Data sources

2.1.1 Pb concentrations and isotope compositions

Observations of seawater dissolved and total dissolvable Pb concentrations, $^{206}\text{Pb}/^{207}\text{Pb}$, and $^{208}\text{Pb}/^{207}\text{Pb}$ included in the GEOTRACES Intermediate Data Product 2021v2 (IDP2021v2; GEOTRACES Intermediate Data Product Group, 2023) form the basis of this study. Only GEOTRACES IDP2021v2 data with quality control flag equal to 1, equivalent to good data, were included in our Pb concentrations, $^{206}\text{Pb}/^{207}\text{Pb}$, and $^{208}\text{Pb}/^{207}\text{Pb}$ datasets. Among these good data points, 8 samples from the southern sector of the Indian Ocean and the Southern Ocean were not included in the Pb concentrations dataset as they had unusually high values compared to those collected at surrounding locations. Additionally, one sample from the Arctic Ocean was not included in our dataset as it had a negative Pb concentration (-0.05 pmol/kg).

Feature name	Data source
Geographical features	
Latitude [$^{\circ}\text{N}$]*	Sampling information
Longitude [$^{\circ}\text{E}$]*	Sampling information
Depth [m]	Sampling information
Distance from bottom depth [m]	Derived from sampling information
Oceanographic features	
Temperature [$^{\circ}\text{C}$]	World Ocean Atlas 2018
Salinity	World Ocean Atlas 2018
Dissolved nitrate [$\mu\text{mol/kg}$]	World Ocean Atlas 2018
Dissolved phosphate [$\mu\text{mol/kg}$]	World Ocean Atlas 2018
Dissolved silicate [$\mu\text{mol/kg}$]	World Ocean Atlas 2018
Density anomaly (σ_{θ}) [kg/m^3]	World Ocean Atlas 2018
Dissolved oxygen [$\mu\text{mol/kg}$]	World Ocean Atlas 2018
Apparent oxygen utilization [$\mu\text{mol/kg}$]	World Ocean Atlas 2018
Mixed layer depth [m]	World Ocean Atlas 2018
Chlorophyll-a [mg/m^3]	CMEMS ^a
Atmospheric features	
Black carbon AOD ^b	CAMS ^c
Dust AOD ^b	CAMS ^c

* Variables used for dataset compilation and for an initial version of the models, but dropped for the final models (see more in SI note 1)

^a Copernicus Marine Environment Monitoring Service

^b Aerosol Optical Depth

^c Copernicus Atmosphere Monitoring Service

Deleted:

Formatted: Font: Bold

Formatted: Space After: 12 pt, Line spacing: 1.5 lines

120 Additional data were included in our dataset to increase the number of observations available; these encompass published
(Chen et al., 2023b; Chien et al., 2017; Lanning et al., 2023; Olivelli et al., 2023, 2024b) and unpublished studies (from the
MAGIC Laboratories at Imperial College London). As previous studies have shown that dissolved and total dissolvable Pb
concentrations and isotope compositions are comparable in the open ocean (Bridgestock et al., 2018; Olivelli et al., 2023,
2024b; Schlosser et al., 2019), data for dissolved and total dissolvable Pb concentrations, $^{206}\text{Pb}/^{207}\text{Pb}$, and $^{208}\text{Pb}/^{207}\text{Pb}$ were
125 pooled together for the purpose of this work.

Deleted: 2023a

Formatted: Font: Symbol

Formatted: Font: Symbol

2.1.2 Oceanographic and atmospheric features

Global gridded climatologies of seawater temperature, salinity, density anomaly (σ_θ), nitrate, phosphate, silicate, oxygen,
apparent oxygen utilization (AOU), and mixed layer depth (MLD) were obtained from the World Ocean Atlas 2018 product
(WOA18; Garcia, 2019). The WOA18 has a $1^\circ \times 1^\circ$ spatial resolution and 102 depth levels (spaced every 5 m between 0 and
130 100 m, every 25 m between 100 and 500 m, every 50 m between 500 and 2000 m, and every 100 m between 2000 and 5500
m). Chlorophyll-a concentrations in the surface ocean were obtained from the Copernicus Marine Service Global Ocean
Biogeochemistry Hindcast product ($0.25^\circ \times 0.25^\circ$ spatial resolution) as monthly averages for the period between January 2005
and December 2017, and averaged over the entire period to create a climatology.

Formatted: Font: Symbol

Deleted: ,

Dust aerosol optical depth (AOD) and black carbon AOD were included as most Pb enters the ocean via atmospheric transport.
135 Dust was found to be an important source of natural Pb to the surface ocean in several regions, including the Atlantic Ocean
(Bridgestock et al., 2016; Olivelli et al., 2023) and Red Sea (Benaltabet et al., 2020). Black carbon, on the other hand, was
chosen as an indicator of Pb emissions from industrial and high-temperature activities, which are known sources of
anthropogenic Pb to the environment (Nriagu and Pacyna, 1988; Pacyna and Pacyna, 2001). Data for the years between January
2005 and December 2017 were obtained from CAMS global reanalysis (ECMWF Atmospheric Composition Reanalysis 4;
140 EAC4) monthly averaged fields, with a $0.75^\circ \times 0.75^\circ$ spatial resolution, and were averaged over the whole period to obtain a
climatology.

Deleted: , southern Indian Ocean (Lee et al., 2015),

Formatted: Placeholder Text

Formatted: Font colour: Text 1

Formatted: Font colour: Text 1

Deleted: &

Deleted: &

Formatted: Font colour: Text 1

Deleted: °x

2.1.3 Data handling and dataset preparation

The WOA18 grid was chosen as a reference for all feature and target variables in this study. Therefore, Pb concentrations, $^{206}\text{Pb}/^{207}\text{Pb}$, $^{208}\text{Pb}/^{207}\text{Pb}$, chlorophyll-a, dust AOD, and black carbon AOD values, which were either obtained as point observations or gridded at higher resolution, were re-gridded according to it. Lead data were binned to the $1^\circ \times 1^\circ$ grid cell the samples were collected in and assigned to the depth level closest to the depth at which the samples were collected. All samples collected at depth larger than 5500 m were discarded, as 5500 m is the deepest depth level considered in the WOA18. Distance from bottom depth was calculated as the difference between the bottom depth value reported for the sampling station and the WOA18 depth level to which the observation was assigned. Chlorophyll-a, dust AOD, and black carbon AOD were re-gridded by assigning each $1^\circ \times 1^\circ$ grid cell the feature value of the respective product cell located the closest to the centre of the WOA18-gridded cell.

All samples in each $1^\circ \times 1^\circ$ column were attributed the same dust AOD, black carbon AOD, surface chlorophyll-a, and mixed layer depth values, regardless of their depth. This was done to assess whether characteristics of the surface ocean have an important impact throughout the water column. These features represent atmospheric inputs (dust AOD and black carbon AOD), surface primary production as a proxy for sinking particulate matter in the water column (surface chlorophyll-a), and the thickness of the ocean layer that is directly influenced by the atmosphere (mixed layer depth).

In total, the Pb concentration dataset includes 9920 observations, the $^{206}\text{Pb}/^{207}\text{Pb}$ dataset 2014, and the $^{208}\text{Pb}/^{207}\text{Pb}$ dataset 2010 (Fig. 1). In the WOA18 grid, 9920 observations cover $\sim 0.3\%$ of the total gridded ocean volume, while 2014 and 2010 cover $\sim 0.06\%$.

In the initial phase of model development, latitude and longitude were included as features and deconvoluted in a three-dimensional coordinate system to allow for continuity (Gregor et al., 2017; Huang et al., 2022). However, when including coordinates as features, the global maps of Pb concentration and isotope compositions showed unrealistic spatial artefacts (see SI Note 1). For this reason, coordinates were excluded as features for further model development and are not included in the results presented here.

Formatted: Font: Symbol

Formatted: Font: Symbol

Formatted: Font: Symbol

2.2 Model development and validation

The non-linear regression algorithm XGBoost (eXtreme Gradient Boosting; Chen & Guestrin, 2016) was chosen to develop three separate models for Pb concentration, $^{206}\text{Pb}/^{207}\text{Pb}$ and $^{208}\text{Pb}/^{207}\text{Pb}$. Indeed, evidence shows that tree-based models consistently outperform deep learning models on tabular-style datasets (Grinsztajn et al., 2022). Moreover, in recent years, XGBoost has proven to perform well on a variety of applications, spanning from finance to medicine, and including Earth sciences (Biass et al., 2022; Xie et al., 2024; Ye et al., 2023). XGBoost is an ensemble algorithm that consists of individual decision trees built in a sequential manner. In brief, each successive tree aims to reduce the errors perpetrated by its predecessor. The final prediction made by the model equals to the weighted sum of the predictions made by all trees in the ensemble. XGBoost was preferred to the Random Forest algorithm as the sequential building of trees in XGBoost allows the algorithm to focus on poorly understood fields, and because Random Forest is more prone to overfitting than XGBoost. Moreover, the Random Forest algorithm was found to perform worse than XGBoost for the Pb concentration and $^{208}\text{Pb}/^{207}\text{Pb}$ models, and comparably for the $^{206}\text{Pb}/^{207}\text{Pb}$ model (SI Note 2).

In addition to performing well on a variety of applications, tree-based models also offer a higher level of interpretability compared to deep learning and linear models (Lundberg et al., 2020). Among the different approaches that have been used to explain predictions of XGBoost models, Shapley Additive exPlanations (SHAP) have stood out in recent years as a unified approach to explain local predictions and gain a global understanding of a model's structure (Lundberg et al., 2020; Lundberg and Lee, 2017; Molnar, 2022).

The SHAP approach is based on game theory (i.e. features are considered as players that contribute to a prediction) and allows to interpret the importance of each feature on a given prediction by calculating the marginal contribution of each feature across all possible permutations. For each sample, a positive (negative) SHAP value indicates that a specific feature contributes towards increasing (decreasing) the final predicted value. Additionally, features with a larger mean absolute SHAP value are considered to contribute more to a given prediction compared to features with a smaller mean absolute SHAP value.

In recent years, several studies in Earth sciences have used SHAP values for interpreting predictions made by tree-based and deep learning models, providing insightful findings on atmospheric pollution (Hou et al., 2022; Qin et al., 2022; Stimberg et al., 2021), ocean dynamics (Clare et al., 2022), and vegetation vulnerability (Biass et al., 2022). Here, we used the

Deleted: &

Deleted: &

TreeExplainer method from the 'shap' library in Python (Lundberg et al., 2020; Lundberg and Lee, 2017) to compute SHAP values, and to evaluate the contribution of each feature on predicted Pb concentrations, $^{206}\text{Pb}/^{207}\text{Pb}$, and $^{208}\text{Pb}/^{207}\text{Pb}$. Interactions between features, which are defined as the additional combined feature effects after accounting for the individual feature effects, were calculated using SHAP interaction values.

The first step of model development consisted of withholding a cruise transect in the Atlantic Ocean from the Pb concentration, $^{206}\text{Pb}/^{207}\text{Pb}$, and $^{208}\text{Pb}/^{207}\text{Pb}$ datasets. This group of withheld samples, identified as "geographic test set" (Fig. S1), included 216 samples for the Pb concentration model and 26 samples for the Pb isotope ratios models, and was used to mimic areas of the ocean where *in situ* samples have never been collected and assess the ability of the model to generalise to those. The remainder of each of the three datasets was split into a training test, consisting of 80% of the data, and a randomly selected test set, consisting of 20% of the data.

To identify the best model architectures, hyperparameters were tuned using 5-fold cross-validation on the training set using the grid search method. Cross-validation is a technique that allows to evaluate the performance of a machine learning model during the hyperparameter tuning phase. It consists of splitting the training dataset into a specified number of subsets (k , in this case equal to 5), and using $k-1$ subsets for training and one subset for testing. The training and testing process is repeated k times (i.e. k -fold cross-validation), and the performance of each hyperparameter configuration is calculated by averaging the scores obtained for each fold. Hyperparameters are a set of parameters that control the learning process of a ML model and are tuned to obtain the most optimal model performance. The grid search method consists of specifying a set of possible values for each hyperparameter and subsequently training and testing the models built with each unique combination of hyperparameter values. The hyperparameters tuned for our Pb concentration, $^{206}\text{Pb}/^{207}\text{Pb}$, and $^{208}\text{Pb}/^{207}\text{Pb}$ include: the number of decision trees built and boosted (' $n_estimators$ '), the rate at which the model learns information (' $learning_rate$ '), the maximum number of split nodes in a tree (' max_depth '), the minimum number of samples a node must represent in order to be split further (' min_child_weight '), the percentage of features used to construct each tree (' $colsample_bytree$ '), and the L1 and L2 regularization terms (' reg_alpha ' and ' reg_lambda ', respectively). The two regularization terms, L1 and L2, were only tuned for the Pb isotope ratio models to reduce the computational costs of hyperparameter tuning for the Pb concentration

model, which was built using a much larger dataset than the $^{206}\text{Pb}/^{207}\text{Pb}$ and $^{208}\text{Pb}/^{207}\text{Pb}$ models. As the target variables of the three models (Pb concentrations, $^{206}\text{Pb}/^{207}\text{Pb}$, and $^{208}\text{Pb}/^{207}\text{Pb}$) were not uniformly distributed, a least-square loss function was chosen as the preferred method for model optimization, as squaring the error penalises the model more when the size of the error increases.

Different performance metrics were used to assess model performance during cross validation and testing on the geographic and random test sets. These include root mean square error (RMSE), mean average percentage error (MAPE), and the coefficient of determination (R^2), and are calculated as follows:

$$RMSE = \sqrt{\frac{1}{n} \sum_{i=1}^n (y_o - y_p)^2}$$

$$MAPE = 100 \frac{1}{n} \sum_{i=1}^n \left| \frac{y_o - y_p}{y_o} \right|$$

$$R^2 = 1 - \frac{\sum_{i=1}^n (y_o - y_p)^2}{\sum_{i=1}^n (y_o - \bar{y})^2}$$

where n is the number of samples, y_o and y_p the observed and predicted value, respectively, and \bar{y} is the mean of the observed values.

Root mean square error was chosen as the primary evaluation metric for the two isotope ratio models. Indeed, RMSE is reported in the same unit as the target variables (in this case unitless) and makes it more intuitive to interpret the results. On the other hand, MAPE was chosen as the preferred metric on which to evaluate the performance of Pb concentration models. This was done as Pb concentrations in the dataset vary between 0.10 and 95.00 pmol/kg, with an average value of 17.33 ± 13.05 (1SD, $n = 9920$). While MAPE might be harder to interpret, as it is reported as a percentage rather than an absolute value with the same unit of measurement as the target variable, minimising the percentage error ensures that basins and locations where Pb concentrations are low are not overlooked by the model.

3 Results and discussion

3.1 Model development and performance

During hyperparameter tuning, we assessed model configurations by changing hyperparameter grid spacing and increasing
245 granularity to get as close as possible to the best combination achievable. In the final phase of the tuning, 1280 different
configurations were assessed for Pb concentrations, and 11520 both for ²⁰⁶Pb/²⁰⁷Pb and ²⁰⁸Pb/²⁰⁷Pb. The hyperparameter space
explored and the best hyperparameters identified for each of the three models are reported in Table 2. The variations in MAPE
and RMSE values for different combinations of hyperparameters are visible in Fig. S2.

The mean MAPE values of the Pb concentration models calculated on the cross-validation, random test set and geographic test
250 set were 22.7 ± 2.2 %, 20.7 ± 1.5 %, and 19.5 ± 4.1 % (n = 1280, 2SD), respectively. The ten best hyperparameter configurations
for the Pb concentration model shared the same learning rate (0.01) and number of columns subsampled by each tree (0.7).
The maximum difference in MAPE values between these ten models was 0.05 % on the cross-validation set, 0.17 % on the
random test set, and 0.82 % on the geographic test set.

Table 2. Hyperparameter space explored for the XGBoost regression models. Bold values identify the combination of
255 hyperparameters that returned the best model performance.

Hyperparameter	Pb concentration					²⁰⁶ Pb/ ²⁰⁷ Pb					²⁰⁸ Pb/ ²⁰⁷ Pb				
learning_rate	0.01	0.05	0.1	0.3		0.01	0.05	0.1			0.01	0.05	0.1		
n_estimators	1200	1300	1400	1500	1600	1100	1200	1300	1400	1500	700	800	900	1000	1100
max_depth	12	14	16	18		4	6	8	10		16	18	20	22	
colsample_bytree	0.5	0.7	0.9	1		0.5	0.7	0.9	1		0.5	0.7	0.9	1	
min_child_weight	6	8	10	12		6	8	10	12		12	14	16	18	
reg_alpha						0	0.1	1			0	0.1	1		
reg_lambda						0	0.1	1	10		6	8	10	12	

The ²⁰⁶Pb/²⁰⁷Pb models had mean RMSE values of 0.008 ± 0.004 (cross-validation), 0.007 ± 0.003 (random test set), and 0.006
± 0.002 (geographic test set; n = 11520, 2SD). The 20 best performing hyperparameter configurations shared the same learning
rate (0.01), L1 regularization (0) and number of columns subsampled by each tree (0.7). These models have a maximum
260 difference in RMSE values of 0.000024 on the cross-validation set, 0.000194 on the random test set, and 0.000298 on the
geographic test set. Lastly, the ²⁰⁸Pb/²⁰⁷Pb models had mean RMSE values of 0.007 ± 0.003 (cross-validation), 0.007 ± 0.003

(random test set), and 0.008 ± 0.003 (geographic test set; $n = 11520$, 2SD). The best five hyperparameter configurations have the same learning rate (0.01), number of columns subsampled by each tree (0.5), and L1 regularization (0). The maximum difference in RMSE values between these models was 0.000003 on the cross-validation set, 0.000046 on the random test set, and 0.000069 on the geographic test set.

Overall, given the very comparable performances of the best hyperparameter configurations on cross-validation, random test set and geographic test set for each of the three models described above, we built the final Pb concentration, $^{206}\text{Pb}/^{207}\text{Pb}$, and $^{208}\text{Pb}/^{207}\text{Pb}$ models using the hyperparameter configuration that performed best on cross-validation.

Table 3. Performance of the best Pb concentration, $^{206}\text{Pb}/^{207}\text{Pb}$, and $^{208}\text{Pb}/^{207}\text{Pb}$ models.

Target variable	Random test set				Geographic test set (Atlantic Ocean)			
	<i>n</i>	<i>R</i> ²	RMSE	MAPE [%]	<i>n</i>	<i>R</i> ²	RMSE	MAPE [%]
[Pb]	1941	0.87	4.84 pmol/kg	20.2	216	0.82	5.06 pmol/kg	18.3
$^{206}\text{Pb}/^{207}\text{Pb}$	398	0.80	0.006	0.3	26	0.77	0.005	0.3
$^{208}\text{Pb}/^{207}\text{Pb}$	397	0.71	0.006	0.1	26	0.52	0.006	0.2

The predictive performance of the final models was assessed on both the random and geographic test sets and is reported in Table 3. For all three models, the agreement between observed and predicted values (R^2 score) is higher for the random test set than for the geographic test set. However, MAPE and RMSE are not always better for the random test set than for the geographic one, indicating that the model is able to generalise well without drastically reducing its performance. Indeed, the Pb concentration model has a lower MAPE for the geographic test set (18.3 %) than for the random one (20.2 %), and the $^{206}\text{Pb}/^{207}\text{Pb}$ model has a lower RMSE for the geographic (0.005) than the random test set (0.006).

Model performance on the random test set was also assessed by splitting the data between ocean basins to identify which areas are better reproduced by the models (Fig. 2). The Pb concentration model achieves the highest R^2 on the Pacific Ocean (0.93) and the lowest MAPE on the Atlantic Ocean (14 %). The model fitting of Indian Ocean observations is overall very good ($R^2 = 0.84$), however the MAPE is relatively high at 33 %, mostly due to a poor performance on data between 25 and 65 pmol/kg. Lastly, the two high-latitude basins have the worst R^2 (Arctic Ocean: 0.49; Southern Ocean: 0.58) and high MAPE values (Arctic Ocean: 37 %; Southern Ocean: 29 %), due to the presence of outliers as well as a high density of low concentration

values (i.e. MAPE is higher for a given difference between modelled and actual value when observation values are closer to zero).




Figure 2. Basin-wise model performances assessed on all samples from each basin in the random test set. Top row: Pb concentration; Middle row: $^{206}\text{Pb}/^{207}\text{Pb}$; Bottom row: $^{208}\text{Pb}/^{207}\text{Pb}$.

The Pb isotope ratio models perform the worst on the Arctic Ocean ($^{206}\text{Pb}/^{207}\text{Pb}$: $R^2 = 0.52$, $\text{RMSE} = 0.010$; $^{208}\text{Pb}/^{207}\text{Pb}$: $R^2 = 0.55$, $\text{RMSE} = 0.011$), and show discrepant performances on the Atlantic, Pacific, Indian and Southern Ocean. The $^{206}\text{Pb}/^{207}\text{Pb}$ model performs best on the two basins with the lowest number of samples in the random test, namely the Southern ($R^2 = 0.93$, $\text{RMSE} = 0.004$) and Indian Ocean ($R^2 = 0.87$, $\text{RMSE} = 0.004$). Samples from the Pacific Ocean are well reproduced ($R^2 =$

0.80, RMSE = 0.004), while the performance on the Atlantic Ocean is poorer ($R^2 = 0.59$, RMSE = 0.006). Lastly, for the $^{208}\text{Pb}/^{207}\text{Pb}$ model, the Indian Ocean is the basin with the best performance ($R^2 = 0.89$, RMSE = 0.003), followed by the Atlantic ($R^2 = 0.74$, RMSE = 0.004) and the Pacific ($R^2 = 0.57$, RMSE 0.005).

3.2 Model explanation

SHAP values were calculated for all samples in the Pb concentration, $^{206}\text{Pb}/^{207}\text{Pb}$, and $^{208}\text{Pb}/^{207}\text{Pb}$ datasets to maximise the interpretability of the models. In the next subsections, the four most important features for each model, as well as their interaction terms, are explained.

Table 4. SHAP importance values for all features included in the Pb concentration, $^{206}\text{Pb}/^{207}\text{Pb}$, and $^{208}\text{Pb}/^{207}\text{Pb}$ models.

Feature Name	Pb concentration		$^{206}\text{Pb}/^{207}\text{Pb}$		$^{208}\text{Pb}/^{207}\text{Pb}$	
	Importance	Rank	Importance	Rank	Importance	Rank
Apparent oxygen utilization [$\mu\text{mol/kg}$]	0.22	14	0.00033	14	0.00025	14
Black Carbon AOD	2.01	3	0.00141	6	0.00185	1
Bottom distance [m]	0.49	11	0.00074	10	0.00091	5
Chlorophyll-a [mg/m^3]	0.63	9	0.00070	11	0.00097	4
Density [kg/m^3]	0.76	5	0.00144	5	0.00064	10
Depth [m]	0.68	8	0.00086	7	0.00073	8
Dust AOD	2.84	2	0.00154	4	0.00114	2
Mixed layer depth [m]	0.71	7	0.00074	9	0.00079	6
Nitrate [$\mu\text{mol/kg}$]	0.41	13	0.00034	13	0.00048	13
Oxygen [$\mu\text{mol/kg}$]	0.46	12	0.00191	1	0.00078	7
Phosphate [$\mu\text{mol/kg}$]	0.56	10	0.00064	12	0.00049	12
Salinity	0.99	4	0.00170	2	0.00057	11
Silicate [$\mu\text{mol/kg}$]	0.74	6	0.00083	8	0.00069	9
Temperature [$^{\circ}\text{C}$]	3.19	1	0.00155	3	0.00102	3

3.2.1. Pb concentrations

Temperature, dust AOD, black carbon AOD, and salinity are the most important features for the Pb concentration model (Table 4, Fig. 3). For temperature values above 5.8 $^{\circ}\text{C}$, SHAP values are consistently positive, with warmer waters having a larger impact on predicted Pb concentrations (Fig. 3A). Black carbon AOD has the largest interaction with temperature, showing positive SHAP values in areas where both features are high. Taken together, these results suggest that surface and intermediate waters at low and mid latitudes (where black carbon AOD is highest) are associated with higher predictions of Pb concentrations. An exception to this trend is observed for samples with low temperatures and extremely low black carbon

Formatted: Font: Symbol

AOD, which have unexpectedly positive SHAP values for temperature (Fig. 3A). Analogously, SHAP values for black carbon AOD are positive for the lowest black carbon AOD values (< 0.001 ; Fig. 3C). The positive impact of extremely low black carbon concentrations in the atmosphere cannot be explained by a physico-chemical process, as one would expect lower values of black carbon AOD to lead to lower Pb concentrations in the ocean. However, the positive SHAP values for temperature and black carbon AOD can be explained to the presence of samples with high Pb concentrations in the Southern Ocean (Fig. 1A), where black carbon AOD is lowest, and seawater temperatures are below 5.0°C (Fig. S3 and S4). Further evidence supporting the positive impact of the lowest black carbon AOD values on Pb concentrations in the Southern Ocean can be found in the interaction between black carbon AOD and salinity observed for salinity SHAP values (Fig. 3D). Indeed, this interaction indicates positive SHAP values for extremely low black carbon AOD and salinities around 34.0 ± 0.2 (Fig. S5), which are characteristics of the higher latitudes of the Southern Hemisphere.

Formatted: Font: Symbol




Figure 3. SHAP values for temperature (A), dust AOD (B), black carbon AOD (C), and salinity (D) for the Pb concentration model. The colour of the dots represents the value of the feature that has the highest interaction (reported on the right-hand-side of each plot). Y-axis values differ among the panels.

SHAP values are consistently positive for dust AOD values above 0.003 (Fig. 3B), and they show an interesting interaction with distance from bottom depth, which is especially visible for larger values of dust AOD (> 0.05). Indeed, as all grid-cells at the same location were assigned the same dust AOD value throughout the water column, the interaction between dust AOD and distance from bottom depth shows that samples more distant from the seafloor (i.e. closer to the surface) have higher SHAP values than their deeper counterparts (Fig. 3B). This finding agrees with observations of higher Pb concentrations at the surface ocean due to atmospheric deposition (Duce et al., 1991; Patterson and Settle, 1987), therefore suggesting that the model has learned from the data the key process of atmospheric Pb sourcing to the ocean.

Deleted: &

3.2.2. Pb isotope compositions

Seawater oxygen concentration, salinity, temperature, and dust AOD are the four most important features for the $^{206}\text{Pb}/^{207}\text{Pb}$ model (Table 4; Fig. 4). Oxygen concentration values below $220 \mu\text{mol/kg}$ have SHAP values between -0.004 and 0 , and a sharp transition to positive SHAP values can be observed at oxygen concentrations of $\sim 240 \mu\text{mol/kg}$ (Fig. 4A). The strong interaction between oxygen and nitrate concentrations suggests that areas of high dissolved oxygen and low nitrate concentrations, such as the Arctic and the North Atlantic Ocean (Fig. S6 and S7), have a large positive impact on predicted $^{206}\text{Pb}/^{207}\text{Pb}$.

Formatted: Font: Symbol

Formatted: Font: Symbol


SHAP values for salinity show an increasing trend as the latter varies between 32.5 and 37.0 (Fig. 4B). This is especially visible for surface and intermediate waters with densities (σ_θ) between 25 and 35 kg/m^3 , indicating that the model learned well from observations, which show higher Pb concentrations at shallower depths and a general decrease in Pb concentration values from tropical and subtropical to polar latitudes (Fig. 1A). SHAP values for temperature show a decreasing trend as waters get warmer (Fig. 4C). Interestingly, and expectedly from a theoretical perspective, this trend is reversed for density, which is the fifth most important feature for the model and shows increasing SHAP values with as waters become denser (Fig. S8). Both temperature and density also show a clear interaction with salinity, which can be explained by the fact that temperature and salinity are the two conservative parameters that contribute to seawater density. Overall, the SHAP values of and interactions between these three variables agree with the general view of Pb isotope ratios as tracers of water mass movements and ocean ventilation (Bridgestock et al., 2018; Frank, 2002; Henderson and Maier-Reimer, 2002).

Deleted: (σ

Deleted: km

Deleted: &

350 Dust AOD shows a more variable distribution of SHAP values, **which is better visualised on a logarithmic scale (Fig. 4D).**
Strongly positive SHAP values (up to 0.009) **are visible** for dust AODs lower than 0.004, followed by a rapid decrease to
strongly negative SHAP values (up to -0.007) for dust AODs between 0.004 and 0.007. As dust AOD increases **further**, SHAP
values **first** show a **decreasing trend for** dust AODs **up to 0.12**, **whilst they** increase afterwards (Fig. 4D).



355 **Figure 4. SHAP values for oxygen (A), salinity (B), temperature (C), and dust AOD (D – on a logarithmic scale) for the**
 $^{206}\text{Pb}/^{207}\text{Pb}$ model. The colour of the dots represents the value of the feature that has the highest interaction (reported
on the right-hand-side of each plot). Y-axis values differ among the panels.

Perhaps surprisingly, given the generally observed correlation between the $^{206}\text{Pb}/^{207}\text{Pb}$ and $^{208}\text{Pb}/^{207}\text{Pb}$ isotope ratios (Boyle et al., 2012; GEOTRACES Intermediate Data Product Group, 2023), the most important features for $^{208}\text{Pb}/^{207}\text{Pb}$ (Fig. 5) do not
360 match closely those of $^{206}\text{Pb}/^{207}\text{Pb}$ (Fig. 4). In fact, the four most important features for the $^{208}\text{Pb}/^{207}\text{Pb}$ model include black carbon AOD, dust AOD, seawater temperature, and surface chlorophyll-a concentration (Table 4; Fig. 5). Black carbon AOD has mostly positive SHAP values for feature values above 0.0045 and negative otherwise (Fig. 5A). It has the strongest interaction with surface chlorophyll-a, and vice versa. Moreover, SHAP values for chlorophyll-a are mostly negative for the lowest chlorophyll-a concentrations, and range between 0.003 and -0.002 for chlorophyll-a concentrations above 0.13 mg/m^3

Moved (insertion) [1]

Deleted: with strongly

Moved up [1]: 4D).

Deleted: (Fig.

Deleted: flat

Deleted: , followed by a slight decrease between

Deleted: of 0.05 and

Deleted: and an

Deleted: 4B). The interaction between dust AOD and silicate highlights that for dust AOD values above 0.007, SHAP values tend to be higher where silicate concentrations are lower, such as in the surface layer (besides in the Southern Ocean and northern North Pacific), and the intermediate and deep Arctic and North Atlantic Ocean (Fig. S9 and S10).

(Fig. 5D). The interaction between these two features, however, does not show a clear pattern and cannot be interpreted intuitively, as both high and low chlorophyll-a concentrations have high and low SHAP values for black carbon AOD, and the other way round (Fig. 5A and 5D). SHAP values of dust AOD (Fig. 5B) show a similar pattern to the $^{206}\text{Pb}/^{207}\text{Pb}$ model, with lower dust AOD values having both the highest and lowest SHAP values. The interaction between dust AOD and black carbon AOD suggests that when both features are low, dust AOD has a strong positive impact on predicted $^{208}\text{Pb}/^{207}\text{Pb}$. Contrarily, when dust AOD is low and black carbon AOD higher than 0.004, dust AOD contributes to reducing predicted $^{208}\text{Pb}/^{207}\text{Pb}$ values (Fig. 5B). Lastly, SHAP values for temperature also show a similar decreasing trend to those of the $^{206}\text{Pb}/^{207}\text{Pb}$ model and have a strong interaction with dissolved nitrate concentrations (Fig. 5C). This interaction indicates that warmer waters with low nitrate content, such as those in tropical and subtropical areas (Fig. S4 and S7), are associated with a reduction in predicted $^{208}\text{Pb}/^{207}\text{Pb}$ values.

3.2.3. SHAP model explanations and current understanding of marine Pb cycling

SHAP explanations for the Pb concentration, $^{206}\text{Pb}/^{207}\text{Pb}$, and $^{208}\text{Pb}/^{207}\text{Pb}$ models align well the current understanding of marine biogeochemical cycling of Pb, with respect to the oceanographic predictors identified as most important features for the three models. This indicates that the models are able to identify the key oceanographic patterns and processes that drive the distribution of Pb and its isotopes on a global scale. However, some regional explanations of dust and black carbon AODs for the Pb concentration and isotope composition models appear to contradict the theoretical intuition that dust and black carbon are sources of Pb to the ocean, with Pb isotope compositions broadly reflecting those of natural and anthropogenic land sources, respectively. In particular, the high SHAP values obtained for extremely low black carbon AODs in the Pb concentration model disagree with the hypothesis that low black carbon concentrations in the atmosphere should result in lower Pb concentrations in the ocean. Analogously, the highly positive SHAP values obtained for extremely low values of dust AOD for the $^{206}\text{Pb}/^{207}\text{Pb}$ and $^{208}\text{Pb}/^{207}\text{Pb}$ models, and the consistently positive SHAP values obtained for relatively high black carbon AODs for the $^{208}\text{Pb}/^{207}\text{Pb}$ model do not match the intuition that dust should contribute to shifting the Pb isotope composition of seawater toward more natural values and black carbon toward more anthropogenic ones.




Figure 5. SHAP values for black carbon AOD (A), dust AOD (B – on a logarithmic scale), temperature (C), and surface chlorophyll-a (D) for the $^{208}\text{Pb}/^{207}\text{Pb}$ model. The colour of the dots represents the value of the feature that has the highest interaction (reported on the right-hand-side of each plot). Y-axis values differ among the panels.

405 Possible explanations for the partial mismatch between expected and obtained local SHAP values are provided in the following, but these will need validation through new campaigns at sea and further studies of the Pb isotope compositions of dust and black carbon sources to the atmosphere, [such as the one by Nizam et al. \(2020\)](#). A first explanation is that dust and black carbon are not the only sources of Pb to the global ocean and this might have important implications on a regional scale. An example is the Southern Ocean, where ice melting and runoff could contribute significantly to the marine Pb budget of the area. Another possibility is that global climatologies of dust AOD and black carbon AOD do not capture the regional and temporal variability of the Pb isotope compositions of natural and anthropogenic Pb sources to the environment. Moreover, black carbon is not only emitted due to fossil fuel consumption, but also from forest fires and coal combustion. [In detail, radiocarbon measurements show that the relative contributions from biomass and fossil fuel combustion to the composition of Asian black carbon can vary on a regional scale, whereby fossil fuel is the predominant source of black carbon to the Tibetan Plateau whilst biomass burning predominates for western India and the Maldives \(Gustafsson et al., 2009; Li et al., 2016\). As](#)

Deleted: .

Deleted: , which might have substantially different Pb isotope signatures compared to other industrial and urban sources.

420 such, regional variations in the relative contributions of different atmospheric black carbon sources are expected to be associated with variations in the Pb isotope signatures associated with these black carbon emissions. Therefore, while the global climatologies of dust AOD and black carbon AOD used as model predictors are currently the best available proxies for atmospheric sources of Pb, they might not fully encompass the complexity of Pb sources to the environment, resulting in locally counterintuitive SHAP explanations for the three models.

3.3. Global maps of Pb concentrations and isotope compositions in seawater

425 Global reconstructions of Pb concentrations, ²⁰⁶Pb/²⁰⁷Pb, and ²⁰⁸Pb/²⁰⁷Pb were made for each grid cell in the WOA18 using the same structure and ancillary features used for model development (Sect. 2.1).

Figure 6 shows the global distribution of Pb concentration, ²⁰⁶Pb/²⁰⁷Pb, and ²⁰⁸Pb/²⁰⁷Pb at four depth levels (10 m, 1000 m, 2500 m, and 4000 m), arbitrarily chosen to represent surface, intermediate, deep, and bottom waters, respectively. These maps provide the first global-scale estimates of Pb concentrations and isotopes and can be used to further understand the dynamics and mechanisms that govern the large-scale distribution of Pb.

3.3.1. Pb concentrations

Lead concentrations vary between 2.05 and 77.65 pmol/kg. Generally, concentrations are higher at the surface and show a decreasing trend with depth (Fig. 6A). The ~~main~~ exception to this trend is ~~visible at intermediate and deep depths in the North Atlantic Ocean, where mapped Pb concentrations at 1000 m and 2500 m are higher than at 10 m (Fig. 6A). However, subsurface~~ maxima in Pb concentrations are also visible between 70 and 500 m in the North Pacific, in line with recent observations (Chan et al., 2024; Jiang et al., 2025; Wu et al., 2010; Zheng et al., 2019; Zurbrick et al., 2017). Both the general trend and the ~~exceptions of the North Atlantic and, to a lesser extent, the North Pacific~~ are in line with the expected distribution of Pb concentrations. Indeed, the majority of Pb is sourced to the ocean via atmospheric wet deposition and then sinks towards the bottom aided by particulate matter (Duce et al., 1991; Patterson and Settle, 1987). Therefore, high Pb concentrations are expected for the surface ocean, with values decreasing with depth. However, extensive historical pollution from leaded gasoline in North America and Europe (Kelly et al., 2009) led to high Pb concentrations in North Atlantic Ocean surface waters between the 1950s and 1980s. Lead concentrations eventually decreased due to environmental policies that phased out and

Formatted: Justified


Deleted: only

Deleted: .

Deleted: exception

Deleted: &

banned leaded gasoline, and Pb pollution in surface waters has decreased steadily since 1975 (Boyle et al., 2014; Bridgestock et al., 2016; Kelly et al., 2009). Therefore, the relatively high Pb concentrations in intermediate- and deep-water reflect the sinking of elemental Pb pollution from earlier decades (Fig. 6A), in agreement with observations (Fig. 1A; Noble et al., 2015).



Deleted: at

Deleted: depths reflects

Deleted: 1

455 **Figure 6. Global maps of reconstructed Pb concentrations (A) and $^{206}\text{Pb}/^{207}\text{Pb}$ (B), and $^{208}\text{Pb}/^{207}\text{Pb}$ (C). The four panels represent different depth levels (10 m, 1000 m, 2500 m, 4000 m), with white patches corresponding to seafloor. The filled circles with white edges represent true data values for comparison with the modelled values in the background.**

460 Mapped surface water Pb concentrations are highest in the Indian Ocean (average $[\text{Pb}] = 32.0 \pm 14.2$ pmol/kg, 1SD) and in the North Pacific (33.6 ± 11.3 pmol/kg, 1SD), in line with recent observations from the two basins (Chen et al., 2023b; Echegoyen et al., 2014; Lanning et al., 2023). The North Pacific also shows high mapped Pb concentrations at depths up to 1000 m (Fig. 6A), with subsurface maxima of $[\text{Pb}]$ up to 70 pmol/kg. These features can be reconciled with pollution from previous decades (Lanning et al., 2023; Wu et al., 2010), like in the North Atlantic, and subduction of mode and intermediate waters from their areas of formation in the north-west Pacific (Jiang et al., 2021).

465 Relatively high Pb concentrations are furthermore found at the surface level near South and Central America, Western Africa, and Southeast Asia. Except for samples collected close to the coast in the South Atlantic Ocean, these areas have never been sampled before and the high concentrations predicted by the model cannot be directly verified. However, the distribution of predictor features shows that these are all areas with high seawater temperatures, and relatively high black carbon and dust AODs (Fig. S4, S3 and S9). Therefore, following the model's interpretation with SHAP values (Sect. 3.2.1), these features must be strongly contributing to the high predictions made by the model.

470 Lastly, the polar regions are characterised by the lowest Pb concentrations throughout the water column, with average Pb concentrations of 9.0 ± 3.1 pmol/kg (1SD) for the Arctic Ocean and 11.9 ± 3.6 pmol/kg (1SD) for the Southern Ocean. Additionally, both polar oceans have the least variable Pb concentrations with depth (Fig. S11), which denotes them as the areas of the global ocean that are least affected by anthropogenic pollution.

3.3.2. Pb isotope compositions

475 Global maps of $^{206}\text{Pb}/^{207}\text{Pb}$ and $^{208}\text{Pb}/^{207}\text{Pb}$ ratios show similar patterns at the four different depth levels analysed (Fig 6B, C). Globally, $^{206}\text{Pb}/^{207}\text{Pb}$ ratios vary between 1.144 and 1.200, and $^{208}\text{Pb}/^{207}\text{Pb}$ between 2.420 and 2.479. Generally, lower $^{206}\text{Pb}/^{207}\text{Pb}$ and $^{208}\text{Pb}/^{207}\text{Pb}$ values are associated with Pb sourced from anthropogenic activities, while higher values reflect natural sources of Pb (Bollhöfer and Rosman, 2000, 2001). However, the exact isotope composition of natural and

Deleted: surface

Deleted: 2023a

Deleted: depth

Deleted: which

Deleted: &

anthropogenic Pb varies between sources and basins, and there is no single reference value that is valid on a global scale due to the short residence time of Pb in seawater and the heterogeneity of global Pb sources. Recent work has, furthermore, shown that reversible scavenging can affect and modify the advected Pb isotope signature in areas of high suspended particulate matter (Lanning et al., 2023; Olivelli et al., 2024b). However, historically Pb isotopes have been regarded as tracers of water mass movements and ventilation in the North Atlantic Ocean (e.g. Véron et al., 1998, 1999). We therefore additionally assessed the distribution of $^{206}\text{Pb}/^{207}\text{Pb}$ and $^{208}\text{Pb}/^{207}\text{Pb}$ across different ranges of potential density anomaly (σ_θ ; Fig. 6 and Fig. S12).

The median $^{206}\text{Pb}/^{207}\text{Pb}$ value is highest for the Arctic (1.186) and lowest for the Indian Ocean (1.165; Fig 7). Median values for the North and South Pacific, South Atlantic and Southern Ocean are very similar, varying between 1.168 and 1.171, while the North Atlantic Ocean has a higher median $^{206}\text{Pb}/^{207}\text{Pb}$ of 1.180. Additionally, the Arctic Ocean shows the smallest difference between the first and third quartile, indicating that Pb isotope compositions are rather constant throughout the geographical domain and the water column, while the Southern Ocean has the most variable distribution (Fig. 6).

In the surface layer (10 m), the lowest $^{206}\text{Pb}/^{207}\text{Pb}$ and $^{208}\text{Pb}/^{207}\text{Pb}$ values are observed in the northern Indian Ocean, while the Arctic Ocean has the highest $^{206}\text{Pb}/^{207}\text{Pb}$ values and coastal areas around North America, the Arctic Ocean and the eastern Pacific Ocean have the highest $^{208}\text{Pb}/^{207}\text{Pb}$ values (Fig. 6B, C). Additionally, the distribution of both isotope ratios shows the presence of low values ($^{206}\text{Pb}/^{207}\text{Pb}$: 1.148 – 1.160; $^{208}\text{Pb}/^{207}\text{Pb}$: 2.420 – 2.437) at 1000 m depth, with a core at the location of the Subantarctic Front ($\sim 48^\circ\text{S}$), expanding horizontally to all basins in the Southern Hemisphere and vertically to the deep layer (2500 m). Low $^{206}\text{Pb}/^{207}\text{Pb}$ and $^{208}\text{Pb}/^{207}\text{Pb}$ values at intermediate depths have been observed in the Indian (Lee et al., 2015), South Atlantic (Olivelli et al., 2024b), and Southern Ocean off the coast of Tasmania (GEOTRACES cruise GS01; unpublished data from the MAGIC group at Imperial College London). Historically, anthropogenic emissions from Australia, New Zealand, Chile and South Africa record the lowest Pb isotope ratios and are mostly associated with the use of Broken Hill-type leaded gasoline (with $^{206}\text{Pb}/^{207}\text{Pb}$ and $^{208}\text{Pb}/^{207}\text{Pb}$ as low as 1.060 and 2.328, respectively; Bollhöfer & Rosman, 2000). Our maps, therefore, suggest that Antarctic Intermediate Water (AAIW), which sinks to depth at latitudes between 50 and 60 $^\circ\text{S}$ across the Southern Ocean, in the eastern South Pacific and near the Drake Passage, has the most anthropogenic signature of all non-surface water masses considered (Fig. 6B, C). Further evidence for the highly anthropogenic signature of

Formatted: Font colour: Text 1

Formatted: Font colour: Text 1

Deleted: (σ_θ ;

Formatted: Font colour: Text 1

Deleted: $^\circ$


Deleted: $^\circ$

510 AAIW can be found by analysing the distribution of Pb isotope compositions across density layers. Indeed, AAIW, which has a potential density anomaly between 27.0 – 27.4 kg/m³, corresponds to the layer with the lowest median ²⁰⁶Pb/²⁰⁷Pb in the South Pacific, South Atlantic, and Indian Ocean (Fig. 7). As leaded gasoline was phased out in the abovementioned countries, and their neighbours, before the time of sampling on GEOTRACES cruises, the widespread anthropogenic signature observed at 1000 m in the Southern Hemisphere either reflects persisting pollution from previous decades that was deposited in the Southern Ocean and is transported northward or the remobilisation of previously deposited Pb on land and subsequent atmospheric deposition in the formation areas of AAIW.

In the North Pacific, North Atlantic and Southern Ocean, the median ²⁰⁶Pb/²⁰⁷Pb and ²⁰⁸Pb/²⁰⁷Pb values increase with depth and potential density ranges (Fig 6B, C and Fig. 7). Contrarily, in the South Atlantic, Indian, and, to a lesser extent, the South Pacific Ocean, a decrease in ²⁰⁶Pb/²⁰⁷Pb values can be observed between surface and intermediate water masses, generally followed by an increase as water masses become denser (Fig. 7). The mapped distributions of Pb isotope compositions, taken together with that of Pb concentrations, therefore indicate that the deepest and densest water masses are least affected by anthropogenic pollution on a global scale, as they have the lowest Pb concentrations and highest ²⁰⁶Pb/²⁰⁷Pb and ²⁰⁸Pb/²⁰⁷Pb ratios. On the other hand, surface and intermediate waters are most affected by pollution, which is discernible (i) in the Northern Hemisphere from Pb concentration data and to some extent from Pb isotope compositions, although natural and anthropogenic sources affecting those basins have a relatively small difference in Pb isotope fingerprints, and (ii) in the Southern Hemisphere, and the northern Indian Ocean, from the low ratios of ²⁰⁶Pb/²⁰⁷Pb and ²⁰⁸Pb/²⁰⁷Pb.

3.4. Uncertainty quantification

Three different sources of uncertainty were identified for our model outputs and are described, and where possible quantified, in this section. The first source of uncertainty relates to the observed values of Pb concentrations and isotope compositions, the second to the ancillary variables used as features in the models, and the third to the random split of the data into a training and a test set.







Figure 7. Violin plots of $^{206}\text{Pb}/^{207}\text{Pb}$ distributions in the different ocean basin (top left panel) and within each basin at different potential density ranges. The white dashed line in each violin represents the median value, while the dotted lines represent the lower and upper quartiles (Q1 and Q3, respectively).

All observations of Pb concentrations, $^{206}\text{Pb}/^{207}\text{Pb}$, and $^{208}\text{Pb}/^{207}\text{Pb}$ ratios are characterised by a fundamental level of uncertainty associated with measuring Pb and its isotopes in seawater. While the GEOTRACES programme ensures data quality of the highest standard through a thorough process of intercalibration, it cannot eliminate uncertainty that is intrinsic to sample processing and analysis. In the case of Pb concentrations, individual sample uncertainties, arising from replicate measurements
545 generally range between 0.1% and 80%, and on average are around 10 to 25 % (relative 1SD; e.g. Boyle et al., 2020; Echegoyen et al., 2014). Inter-laboratory uncertainties arising from the use of different sample processing methodologies and different instrumentation and assessed on intercalibration samples, however, typically show a better reproducibility. In fact, the relative standard deviation of the North Atlantic GEOTRACES and SAFe reference samples is between 3.5% and 9.4% (1SD; GEOTRACES, 2009, 2013). For Pb isotope compositions, individual sample uncertainties, assessed either as replicate
550 measurements of the sample itself or extrapolated from replicates measurements of NIST 981 standard reference material is generally around 0.2 – 0.5‰ for both $^{206}\text{Pb}/^{207}\text{Pb}$ and $^{208}\text{Pb}/^{207}\text{Pb}$ (e.g. Boyle et al., 2020; Bridgestock et al., 2016). Inter-laboratory calibration exercises of Pb isotope composition measurements of seawater have returned uncertainties ranging between 0.1‰ and 3.5‰ for both isotope ratios considered (Boyle et al., 2012). So, if we consider the MAPE as a reflection of the uncertainty associated with the models' predictions, predicted values of Pb concentration have an uncertainty (20.2%
555 random test set, 18.3% geographic test set; Table 3) that falls within the average values for replicate measurements (10 – 25%). Similarly, the uncertainty associated with predicted $^{206}\text{Pb}/^{207}\text{Pb}$ and $^{208}\text{Pb}/^{207}\text{Pb}$ values (MAPE: 1.0 – 3.0‰; Table 3) falls within the range observed for inter-laboratory calibration.

Pooling together dissolved and total dissolvable Pb concentration and isotope composition data creates an additional layer of uncertainty. Our choice is backed by the evidence that in the open ocean, dissolved and total dissolvable Pb data are comparable
560 (Bridgestock et al., 2018; Olivelli et al., 2023, 2024b; Schlosser et al., 2019). However, this might not be the case in estuaries and coastal areas, where boundary exchange and isotopic re-equilibration are believed to play an important role in determining the observed Pb concentrations and isotope compositions (Chen et al., 2016, 2023a). Therefore, new observations, particularly
from coastal areas and other sparsely sampled locations, would provide valuable data for the validation of our modelling results. The uncertainty associated with the models' features cannot be quantified directly, and contributes to the models'
565 MAPE values, but is nevertheless expected to be small and not create systematic bias. Indeed, the data products used in this

Deleted: 2023b)

Deleted: ¶



Moved down [2]: study, namely WOA18 for oceanographic variables, CMEMS's Global Ocean Biogeochemistry Hindcast product for chlorophyll-a, and CAMS's ECMWF Atmospheric Composition Reanalysis 4 for black carbon AOD and dust AOD, are all subjected to stringent quality control. [\[1\]](#)

Figure 8. Coefficient of variation for reconstructed Pb concentration (A), $^{206}\text{Pb}/^{207}\text{Pb}$ (B), and $^{208}\text{Pb}/^{207}\text{Pb}$ (C) values. As for Figure 4, the different panels represent different depth levels (10 m, 1000 m, 2500 m, 4000 m) and the white patches correspond to seafloor locations. Please note that the colour scales differ for the three panels.

study, namely WOA18 for oceanographic variables, CMEMS's Global Ocean Biogeochemistry Hindcast product for chlorophyll-a, and CAMS's ECMWF Atmospheric Composition Reanalysis 4 for black carbon AOD and dust AOD, are all subjected to stringent quality control.

Moved (insertion) [2]

580 Lastly, to quantify the uncertainty arising from the random splitting of the data into a training and test set, the best Pb concentration, $^{206}\text{Pb}/^{207}\text{Pb}$, and $^{208}\text{Pb}/^{207}\text{Pb}$ model architectures were initiated with 100 different random train-test splits while keeping the hyperparameters constant. The means and standard deviations of the model ensemble predictions for the global maps of Pb concentrations and isotope compositions were used to calculate the coefficients of variation for the mapped Pb distributions (Fig. 8). The coefficient of variation was calculated as the ratio between the average value of each cell across the 585 100 different ensemble members and its associated standard deviation. For both Pb concentrations and isotope compositions, the coefficients of variation are highest in the Arctic and Southern Ocean, especially in the surface and intermediate layer (Fig. 8). Both the Arctic and Southern Ocean are characterised by the lowest Pb concentrations observed, therefore any variability in the outputs of the ensemble would cause a larger coefficient of variation as the mean concentrations are low. However, the spatial variability in observed Pb concentrations in the Southern Ocean is much larger than in the Arctic Ocean. Indeed, high 590 concentrations are found at the surface and in the water column near the Antarctic Peninsula, likely caused by glacial meltwater inputs, while much lower values are visible in the Indian and Australian sectors. Therefore, the different splits between training and test set lead to much more variable model outputs for the Southern Ocean than for its Arctic counterpart. The high latitudes are also the areas with the largest coefficients of variation for $^{206}\text{Pb}/^{207}\text{Pb}$ and $^{208}\text{Pb}/^{207}\text{Pb}$, arguably because of the significant presence of both anthropogenic and natural Pb. Based on the variability of the outputs of the three model ensembles, it can be 595 argued that at-sea sampling efforts should focus on the Southern Ocean to obtain a clearer understanding of the distribution and cycling of Pb and its isotopes in that area, and to reduce the uncertainty associated with our modelled outputs.

4 Model and data applications

More than 20 years ago, Henderson & Maier-Reimer (2002) used a general circulation model to investigate the natural marine cycle of Pb and its isotopes. As these authors did not consider inputs from anthropogenic activities, we here present the first 600 study that provides global maps of the distribution of Pb concentrations, and $^{206}\text{Pb}/^{207}\text{Pb}$ and $^{208}\text{Pb}/^{207}\text{Pb}$ ratios.

In addition to providing a detailed demonstration of the applicability of machine learning to marine (isotope) geochemistry, the new models and their outputs can be used to (i) benchmark future levels of pollution, (ii) inform sampling strategies and campaigns, and (iii) compare outputs from different models, either process-based or data-driven. Lead concentration and isotope composition data from our 12-year climatologies provide useful information on the levels of marine pollution, both in terms of sources and spatial extent. This insight can be used to assess the impact of human activities and any potential environmental policies implemented in the future. Both the mapped distributions and their associated uncertainties can be used by the community to support any process-oriented or large-scale sampling campaigns. In particular, the large uncertainty associated with Pb concentrations in the Southern Ocean and the widespread distribution of low $^{206}\text{Pb}/^{207}\text{Pb}$ and $^{208}\text{Pb}/^{207}\text{Pb}$ values at intermediate depth throughout the Southern Hemisphere suggest that these areas should be prioritised in the future. Lastly, the $1^\circ \times 1^\circ$ data products developed will be a useful and robust comparison for any global and regional modelling studies that attempt to parametrise and reproduce the processes driving the biogeochemical cycle of Pb, which still require future work.

Formatted: Font: Symbol

5 Data and code availability

All scripts generated to build the Pb concentration, $^{206}\text{Pb}/^{207}\text{Pb}$, and $^{208}\text{Pb}/^{207}\text{Pb}$ datasets and the corresponding models are available at https://github.com/OlivelliAri/Pb-ML_GEOTRACES. The gridded climatologies of Pb concentrations, $^{206}\text{Pb}/^{207}\text{Pb}$, and $^{208}\text{Pb}/^{207}\text{Pb}$, as well as the results of the model ensemble for uncertainty calculations, are available in Olivelli et al. (2024a, <https://doi.org/10.5281/zenodo.14261154>).

Deleted: <https://doi.org/10.5281/zenodo.14261154>

Deleted: .

Formatted: Font colour: Black

6 Conclusion

We successfully generated global climatologies of Pb concentrations, $^{206}\text{Pb}/^{207}\text{Pb}$, and $^{208}\text{Pb}/^{207}\text{Pb}$ ratios using the non-linear regression algorithm XGBoost, trained and tested on high-quality Pb data collected as part of the GEOTRACES programme. We found that Pb concentrations are highest in the surface layer of the Indian and North Pacific Ocean and at intermediate depths in the North Atlantic, and lowest in the Arctic Ocean. The $^{206}\text{Pb}/^{207}\text{Pb}$ and $^{208}\text{Pb}/^{207}\text{Pb}$ ratios are lowest in the surface Indian Ocean and at intermediate depths in the Southern Hemisphere, and highest throughout the water column in the Arctic Ocean. Taken together, the distributions of Pb concentrations, $^{206}\text{Pb}/^{207}\text{Pb}$, and $^{208}\text{Pb}/^{207}\text{Pb}$ ratios indicate that (i) the surface

Indian Ocean has the highest levels of pollution, (ii) pollution from previous decades is sinking in the North Atlantic and Pacific Ocean, and (iii) that a highly anthropogenic fingerprint originating in the Southern Ocean is spreading at intermediate depths throughout the Southern Hemisphere with AAIW. The lower model performance and higher uncertainty associated with reconstructions of the Southern Ocean suggest that this is a key area to prioritise future sampling campaigns to expand our understanding of biogeochemical processes that drive the distribution and cycling of Pb and its isotopes in the ocean. More broadly, our approach demonstrates the applicability of machine learning to marine (isotope) geochemistry, even when data are scarce and very sparse, as for Pb isotope compositions. Lastly, our model output will provide a useful benchmark for future levels of pollutions and a valuable comparison for process-based models that aim to improve understanding of the biogeochemical processes governing the marine cycling of Pb.

Deleted: is

Acknowledgements

All members of the Data Learning group at Imperial College London are thanked for the useful discussions and suggestions throughout this study. Rob Middag and Kyas Seyitmuhammedov are thanked for the information they shared regarding Pb concentration samples collected near the Antarctic Peninsula on GEOTRACES cruise GPpr08. Francois van Shalkwyk is thanked for his support with computational resources.

Funding

AO was supported by the Natural Environment Research Council (NERC) (NE/S007415/1). The International GEOTRACES Programme is possible in part thanks to the support from the U.S. National Science Foundation (Grant OCE-2140395) to the Scientific Committee on Oceanic Research (SCOR). For the purpose of open access, the author has applied a Creative Commons Attribution (CC BY) license to any Author Accepted Manuscript version arising.

Deleted: Arianna Olivelli

Author contribution statement

AO devised the project, curated the data, developed the models, led the interpretation and wrote the initial draft of this manuscript with support from RA, MR and TvdF. All authors contributed to the final version of the manuscript.

Deleted: Arianna Olivelli

Deleted: Rossella Arcucci

Deleted: Mark Rehkämper

Deleted: Tina van de Fliert

Competing Interests

The authors declare that they have no conflict of interest.

References

660 Alleman, L. Y., Véron, A. J., Church, T. M., Flegal, A. R., and Hamelin, B.: Invasion of the abyssal North Atlantic
by modern anthropogenic lead. *Geophys Res Lett*, 26, 1477–1480. <https://doi.org/10.1029/1999GL900287>, 1999.

Benaltabet, T., Lapid, G., and Torfstein, A.: Seawater Pb concentration and isotopic composition response to daily
time scale dust storms in the Gulf of Aqaba, Red Sea. *Mar Chem*, 227, 103895, 2020.
<https://doi.org/10.1016/j.marchem.2020.103895>.

665 Biass, S., Jenkins, S. F., Aeberhard, W. H., Delmelle, P., and Wilson, T.: Insights into the vulnerability of
vegetation to tephra fallouts from interpretable machine learning and big Earth observation data. *Natural Hazards*
and Earth System Sciences, 22, 2829–2855. <https://doi.org/10.5194/nhess-22-2829-2022>, 2022.

Bollhöfer, A. and Rosman, K. J. R.: Isotopic source signatures for atmospheric lead: The Southern Hemisphere.
Geochim Cosmochim Acta, 64, 3251–3262, 2000.

670 Bollhöfer, A. and Rosman, K. J. R.: Isotopic source signatures for atmospheric lead: The Northern Hemisphere.
Geochim Cosmochim Acta, 65, 1727–1740, 2001.

Boyle, E. A., Chapnick, S. D., Shen, G. T., and Bacon, M. P.: Temporal variability of lead in the western North
Atlantic. *J Geophys Res Oceans*, 91, 8573–8593. <https://doi.org/10.1029/jc091ic07p08573>, 1986.

Boyle, E. A., John, S., Abouchami, W., Adkins, J. F., Echegoyen-Sanz, Y., Ellwood, M., Flegal, A. R., Fornace,
K., Gallon, C., Galer, S., Gault-Ringold, M., Lacan, F., Radic, A., Rehkämper, M., Rouxel, O., Sohrin, Y., Stirling,
C., Thompson, C., Vance, D., Xue, Z., and Zhao, Y.: GEOTRACES IC1 (BATS) contamination-prone trace
element isotopes Cd, Fe, Pb, Zn, Cu, and Mo intercalibration. *Limnol Oceanogr Methods*, 10, 653–665.
<https://doi.org/10.4319/lom.2012.10.653>, 2012.

675 Boyle, E. A., Lee, J. M., Noble, A. E., Moos, S., Carrasco, G., Zhao, N., Kayser, R., Zhang, J., Gamo, T., Obata,
H., and Norisuye, K.: Anthropogenic lead emissions in the ocean: The evolving global experiment. *Oceanography*,
27, 69–75, 2014.

Boyle, E. A., Zurbick, C., Lee, J. M., Till, R., Till, C. P., Zhang, J., and Flegal, A. R.: Lead and lead isotopes in
the U.S. GEOTRACES East Pacific zonal transect (GEOTRACES GP16). *Mar Chem*, 227,
<https://doi.org/10.1016/j.marchem.2020.103892>, 2020.

680 Bridgestock, L., Van De Flierdt, T., Rehkämper, M., Paul, M., Middag, R., Milne, A., Lohan, M. C., Baker, A. R.,
Chance, R., Khondoker, R., Strekopytov, S., Humphreys-Williams, E., Achterberg, E. P., Rijkenberg, M. J. A.,

Formatted: Indent: Left: 1.69 cm, First line: 0 cm, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Deleted: &...nd Hamelin, B. (1999). ... [1]

Deleted: . *Geophysical Research Letters*

Formatted: Font: Not Italic

Deleted: (10),... 1477–1480. ... [2]

Deleted: &...nd Torfstein, A. (2020). ... [3]

Deleted: . *Marine Chemistry*

Formatted: Font: Not Italic

Deleted: .

Deleted: &...nd Wilson, T. (2022). ... [4]

Deleted: .

Formatted: Font: Not Italic

Deleted: (9),... 2829–2855. ... [5]

Formatted: Font: Not Italic

Deleted: ., &... and Rosman, K. J. R. (2000),...: Isotopic source signatures for atmospheric lead: The Southern Hemisphere. *Geochimica et Cosmochimica* ... [6]

Deleted: (19),

Formatted: ... [7]

Deleted: ., &... and Rosman, K. J. R. (2001),...: Isotopi ... [8]

Deleted: (11),

Formatted: ... [9]

Deleted: &...nd Bacon, M. P. (1986). ... [10]

Deleted: . *Journal of Geophysical Research:*

Deleted: (C7),... 8573–8593. ... [12]

Formatted: ... [11]

Deleted:ue, Z., and Zhao, Y. (2012). ... [13]

Deleted: . *Limnology and Oceanography:*

Deleted: (9),... 653–665. ... [15]

Formatted: ... [14]

Deleted: &...nd Norisuye, K. (2014),...: Anthropogen ... [16]

Formatted: ... [17]

Deleted: (1),... 69–75. ... [18]

Deleted: &...nd Flegal, A. R. (2020). ... [19]

Deleted: . *Marine Chemistry*

Deleted: .

Formatted: Font: Not Italic

Moved down [3]: Bridgestock, L., Rehkämper, M., van de

Deleted: & Achterberg, E. P. (2018). The distribution ... [20]

Gerringa, L. J. A., and De Baar, H. J. W.: Return of naturally sourced Pb to Atlantic surface waters, *Nat Commun.*, 7, <https://doi.org/10.1038/ncomms12921>, 2016.

Bridgestock, L., Rehkämper, M., van de Flierdt, T., Paul, M., Milne, A., Lohan, M. C., and Achterberg, E. P.: The distribution of lead concentrations and isotope compositions in the eastern Tropical Atlantic Ocean, *Geochim Cosmochim Acta*, 225, 36–51, <https://doi.org/10.1016/j.gca.2018.01.018>, 2018.

Chan, C. Y., Zheng, L., and Sohrin, Y.: The behaviour of aluminium, manganese, iron, cobalt, and lead in the subarctic Pacific Ocean: boundary scavenging and temporal changes, *J Oceanogr*, 80, 99–115, <https://doi.org/10.1007/s10872-023-00710-8>, 2024.

Chen, M., Boyle, E. A., Lee, J. M., Nurhati, I., Zurbrück, C., Switzer, A. D., and Carrasco, G.: Lead isotope exchange between dissolved and fluvial particulate matter: A laboratory study from the Johor River estuary, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 374, <https://doi.org/10.1098/rsta.2016.0054>, 2016.

Chen, M., Carrasco, G., Zhao, N., Wang, X., Lee, J. N., Tanzil, J. T. I., Annammala, K. V., Poh, S. C., Lauro, F. M., Ziegler, A. D., Duangnamon, D., and Boyle, E. A.: Boundary exchange completes the marine Pb cycle jigsaw, *Proc Natl Acad Sci U S A*, 120, <https://doi.org/10.1073/pnas.2213163120>, 2023a.

Chen, M., Boyle, E. A., Jiang, S., Liu, Q., Zhang, J., Wang, X., and Zhou, K.: Dissolved Lead (Pb) Concentrations and Pb Isotope Ratios Along the East China Sea and Kuroshio Transect—Evidence for Isopycnal Transport and Particle Exchange, *J Geophys Res Oceans*, 128, <https://doi.org/10.1029/2022JC019423>, 2023b.

Chen, T., and Guestrin, C.: XGBoost: A scalable tree boosting system, in: *Proceedings of the ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, 785–794, <https://doi.org/10.1145/2939672.2939785>, 2016.

Chien, C. Te, Ho, T. Y., Sanborn, M. E., Yin, Q. Z., and Paytan, A.: Lead concentrations and isotopic compositions in the Western Philippine Sea, *Mar Chem*, 189, 10–16, <https://doi.org/10.1016/j.marchem.2016.12.007>, 2017.

Clare, M. C. A., Sonnewald, M., Lguensat, R., Deshayes, J., and Balaji, V.: Explainable Artificial Intelligence for Bayesian Neural Networks: Toward Trustworthy Predictions of Ocean Dynamics, *J Adv Model Earth Syst*, 14, <https://doi.org/10.1029/2022MS003162>, 2022.

Duce, R. A., Liss, P. S., Merrill, J. T., Atlas, E. L., Buat-Menard, P., Hicks, B. B., Miller, J. M., Prospero, J. M., Arimoto, R., Church, T. M., Ellis, W., Galloway, J. N., Hansen, L., Jickells, T. D., Knap, A. H., Reinhardt, K. H., Schneider, B., Soudine, A., Tokos, J. J., Tsunogai, S., Wollast, R., and Zhou, M.: The atmospheric input of trace species to the world ocean, *Global Biogeochem Cycles*, 5, 193–259, <https://doi.org/10.1029/91GB01778>, 1991.

Echegoyen, Y., Boyle, E. A., Lee, J. M., Gamou, T., Obata, H., and Norisuye, K.: Recent distribution of lead in the Indian Ocean reflects the impact of regional emissions, *Proc Natl Acad Sci U S A*, 111, 15328–15331, <https://doi.org/10.1073/pnas.1417370111>, 2014.

- Deleted:** &...nd De Baar, H. J. W. (2016)....: Return of naturally sourced Pb to Atlantic surface waters. *Nature Communications*... [21]
- Deleted:** .
- Formatted:** Font: Not Italic
- Moved (insertion) [3]**
- Moved down [4]:** Chen, M., Boyle, E. A., Jiang, S., Liu, Q.,
- Deleted:** & Zhou, K. (2023a). Dissolved Lead (Pb) [22]
- Deleted:** .
- Deleted:** (2081).
- Formatted:** Font: Not Italic
- Formatted:** Font: Not Italic
- Deleted:** &...nd Boyle, E. A. (2023b)....: Boundary ex [23]
- Deleted:** (6).
- Formatted:** Font: Not Italic
- Moved (insertion) [4]**
- Deleted:** ., &... and Guestrin, C. (2016)....: XGBoost: [25]
- Formatted:** Font: Not Italic
- Formatted:** [24]
- Deleted:** 13-17-August-2016, ...85–794. [26]
- Deleted:** &...nd Paytan, A. (2017). [27]
- Deleted:** . *Marine Chemistry*
- Deleted:** .
- Formatted:** Font: Not Italic
- Deleted:** &...nd Balaji, V. (2022). [28]
- Deleted:** . *Journal of Advances in Modeling*
- Deleted:** *Systems*
- Deleted:** (11).
- Formatted:** Font: Not Italic
- Formatted:** Font: Not Italic
- Deleted:**sunogai, S., Wollast, R., and Zhou, M. ([29]
- Deleted:** .
- Deleted:** *Biogeochemical*
- Deleted:** (3),... 193–259. [31]
- Formatted:** Font: Not Italic
- Formatted:** [30]
- Deleted:** &...nd Norisuye, K. (2014). [32]
- Deleted:** . *Proceedings of the National Academy of Sci* [33]
- Deleted:** (43),... 15328–15331. [34]
- Formatted:** Font: Not Italic

Frank, M.: Radiogenic isotopes: Tracers of past ocean circulation and erosional input, *Reviews of Geophysics*, **40**, 1-1-1-38, <https://doi.org/10.1029/2000RG000094>, 2002.

GEOTRACES: Summary of Consensus Values for samples collected on the 2009 GEOTRACES Intercomparison Cruise, 2009.

GEOTRACES: Dissolved Lead - Consensus values (± 1 std. dev.) for North Atlantic GEOTRACES Reference Samples as of May 2013, 2013.

GEOTRACES Intermediate Data Product Group: The GEOTRACES Intermediate Data Product 2021v2 (IDP2021v2), 2023.

GEOTRACES Planning Group: GEOTRACES Science Plan, Baltimore, Maryland, 2006.

Glover, D., Jenkins, W., and Doney, S.: Modeling Methods for Marine Science, Cambridge University Press, 2011.

Gregor, L., Kok, S., and Monteiro, P. M. S.: Empirical methods for the estimation of Southern Ocean CO₂: Support vector and random forest regression, *Biogeosciences*, **14**, 5551–5569, <https://doi.org/10.5194/bg-14-5551-2017>, 2017.

Grinsztajn, L., Oyallon, E., and Varoquaux, G.: Why do tree-based models still outperform deep learning on typical tabular data?, in: 36th Conference on Neural Information Processing Systems (NeurIPS 2022) Track on Datasets and Benchmarks, 2022.

Gustafsson, O., Kruså, M., Zencak, Z., Sheesley, R. J., Granat, L., Engström, E., Praveen, P. S., Rao, P. S. P., Leck, C., and Rodhe, H.: Brown Clouds over South Asia: Biomass or Fossil Fuel Combustion?, *Science* (1979), **323**, 492–495, <https://doi.org/10.1126/science.1165884>, 2009.

Helmers, E., Mart, L., and Schrems, O.: Lead in Atlantic surface waters as a tracer for atmospheric input, *Journal of Fresenius J Anal Chem*, Springer-Verlag, 580–584 pp., 1991.

Henderson, G. M., and Maier-Reimer, E.: Advection and removal of 210Pb and stable Pb isotopes in the oceans: A general circulation model study, *Geochim Cosmochim. Acta*, **66**, 257–272, [https://doi.org/10.1016/S0016-7037\(01\)00779-7](https://doi.org/10.1016/S0016-7037(01)00779-7), 2002.

Hou, L., Dai, Q., Song, C., Liu, B., Guo, F., Dai, T., Li, L., Liu, B., Bi, X., Zhang, Y., and Feng, Y.: Revealing Drivers of Haze Pollution by Explainable Machine Learning, *Environ Sci Technol Lett*, **9**, 112–119, <https://doi.org/10.1021/acs.estlett.1c00865>, 2022.

Huang, Y., Tagliabue, A., and Cassar, N.: Data-Driven Modeling of Dissolved Iron in the Global Ocean, *Front Mar Sci*, **9**, <https://doi.org/10.3389/fmars.2022.837183>, 2022.

Jiang, S., Zhang, J., Zhou, H., Xue, Y., and Zheng, W.: Concentration of dissolved lead in the upper Northwestern Pacific Ocean, *Chem Geol*, **577**, <https://doi.org/10.1016/j.chemgeo.2021.120275>, 2021.

Jiang, S., Lanning, N., Boyle, E., Fitzsimmons, J., Ramezani, J., Wang, A. G., and Zhang, J.: Meridional Central Pacific Ocean Depth Section for Pb and Pb Isotopes (GEOTRACES GP15, 152°W, 56°N to 20°S) Including Shipboard Aerosols, *J Geophys Res Oceans*, **130**, <https://doi.org/10.1029/2024JC021674>, 2025.

Deleted: . (2002)....: Radiogenic isotopes: Tracers of p... [35]
Formatted ... [36]
Deleted: (1),
Deleted: .
Deleted: . (2009).
Formatted ... [37]
Deleted: . <https://www.geotraces.org/wp->... [38]
Deleted: . (2013).
Formatted ... [39]
Deleted: .
Deleted: . (2023).
Deleted: .). In *NERC EDS British Oceanographic Data*... [41]
Deleted: . (2006).
Deleted: . <http://www.jhu.edu/scor/>
Formatted ... [42]
Deleted: &...nd Doney, S. (2011).
Deleted: .
Formatted ... [44]
Deleted: &...nd Monteiro, P. M. S. (2017).
Deleted: .
Deleted: (23),... 5551–5569.
Formatted ... [46]
Deleted: &...nd Varoquaux, G. (2022).
Deleted: ?
Formatted ... [49]
Deleted: &...nd Schrems, O. (1991)....: Lead in Atlan... [51]
Formatted ... [52]
Formatted ... [50]
Deleted: (Vol. 340).
Deleted: ., &... and Maier-Reimer, E. (2002).
Deleted: . *Geochimica et Cosmochimica*
Deleted: (2),... 257–272.
Formatted ... [54]
Deleted: &...nd Feng, Y. (2022).
Deleted: . *Environmental Science and Technology Letters*
Deleted: (2),... 112–119.
Formatted ... [57]
Deleted: &...nd Cassar, N. (2022)....: Data-Driven M... [59]
Deleted: .
Formatted ... [60]

020 Kelly, A. E., Reuer, M. K., Goodkin, N. F., and Boyle, E. A.: Lead concentrations and isotopes in corals and water
near Bermuda, 1780-2000, *Earth Planet Sci Lett*, 283, 93–100, <https://doi.org/10.1016/j.epsl.2009.03.045>, 2009.

Lanning, N. T., Jiang, S., Amaral, V. J., Mateos, K., Steffen, J. M., Lam, P. J., Boyle, E. A., and Fitzsimmons, J.
N.: Isotopes illustrate vertical transport of anthropogenic Pb by reversible scavenging within Pacific Ocean particle
veils, *Proc Natl Acad Sci U S A*, 120, <https://doi.org/10.1073/pnas.2219688120>, 2023.

025 Lee, J. M., Boyle, E. A., Gamo, T., Obata, H., Norisuye, K., and Echegoyen, Y.: Impact of anthropogenic Pb and
ocean circulation on the recent distribution of Pb isotopes in the Indian Ocean, *Geochim Cosmochim Acta*, 170,
126–144, <https://doi.org/10.1016/j.gca.2015.08.013>, 2015.

Li, C., Bosch, C., Kang, S., Andersson, A., Chen, P., Zhang, Q., Cong, Z., Chen, B., Qin, D., and Gustafsson, Ö.: Sources
of black carbon to the Himalayan-Tibetan Plateau glaciers, *Nat Commun*, 7, <https://doi.org/10.1038/ncomms12574>,
2016.

030 Lundberg, S. M. and Lee, S.-I.: A Unified Approach to Interpreting Model Predictions, in: 31st Conference on Neural
Information Processing Systems (NIPS 2017), Long Beach, CA, USA, 2017.

Lundberg, S. M., Erion, G., Chen, H., DeGrave, A., Prutkin, J. M., Nair, B., Katz, R., Himmelfarb, J., Bansal, N.,
and Lee, S. I.: From local explanations to global understanding with explainable AI for trees, *Nat Mach Intell*, 2,
56–67, <https://doi.org/10.1038/s42256-019-0138-9>, 2020.

035 Mete, Ö. Z., Subhas, A. V., Kim, H. H., Dunlea, A. G., Whitmore, L. M., Shiller, A. M., Gilbert, M., Leavitt, W.
D., and Horner, T. J.: Barium in seawater: dissolved distribution, relationship to silicon, and barite saturation state
determined using machine learning, *Earth Syst Sci Data*, 15, <https://doi.org/10.5194/essd-15-4023-2023>, 2023.

Molnar, C.: *Interpretable Machine Learning: A Guide for Making Black Box Models Explainable*, 2nd ed., 2022.

040 Nizam, S., Sen, I. S., Vinoj, V., Galy, V., Selby, D., Azam, M. F., Pandey, S. K., Creaser, R. A., Agarwal, A. K., Singh,
A. P., and Bizimis, M.: Biomass-Derived Provenance Dominates Glacial Surface Organic Carbon in the Western
Himalaya, *Environ Sci Technol*, 54, 8612–8621, <https://doi.org/10.1021/acs.est.0c02710>, 2020.

Noble, A. E., Echegoyen-Sanz, Y., Boyle, E. A., Ohnemus, D. C., Lam, P. J., Kayser, R., Reuer, M., Wu, J., and
Smethie, W.: Dynamic variability of dissolved Pb and Pb isotope composition from the U.S. North Atlantic
GEOTRACES transect, *Deep Sea Res 2 Top Stud Oceanogr*, 116, 208–225,
045 <https://doi.org/10.1016/j.dsr2.2014.11.011>, 2015.

Nriagu, J. O.: Global Inventory of Natural and Anthropogenic Emissions of Trace Metals to the Atmosphere,
Nature, 279, 409–411, 1979.

Nriagu, J. O. and Pacyna, J. M.: Quantitative assessment of worldwide contamination of air, water and soils by
trace metals, *Nature*, 333, 134–139, 1988.

050 Olivelli, A., Murphy, K., Bridgestock, L., Wilson, D. J., Rijkenberg, M., Middag, R., Weiss, D. J., van de Flierdt,
T., and Rehkämper, M.: Decline of anthropogenic lead in South Atlantic Ocean surface waters from 1990 to 2011:

Deleted: &...nd Boyle, E. A. (2009). ... [62]

Formatted ... [61]

Deleted: .

Deleted: and Planetary Science Letters

Deleted: (1–4),... 93–100. ... [65]

Formatted ... [63]

Formatted ... [64]

Deleted: &

Deleted: . (2023).

Deleted: . *Proceedings of the National Academy of Sci* ... [66]

Deleted: (23).

Formatted ... [67]

Deleted: &...nd Echegoyen, Y. (2015). ... [68]

Deleted: . *Geochimica et Cosmochimica*

Formatted ... [69]

Deleted: .

Moved (insertion) [5]

Formatted ... [70]

Deleted: &

Deleted: . (2020),...: From local explanations to global ... [71]

Deleted: (1).

Formatted ... [72]

Deleted: .

Moved up [5]: Lundberg, S. M

Deleted: ., & Lee, S.-I. (2017). A Unified Approach to ... [73]

Deleted: &...nd Horner, T. J. (2023). ... [74]

Deleted: .

Deleted: *System Science*

Deleted: (9).

Formatted ... [75]

Formatted ... [76]

Deleted: . (2022).

Deleted: (... 2nd ed.). ... [78]

Formatted ... [77]

Deleted: &

Formatted ... [79]

Deleted: . (2015).

Deleted: .

Deleted: -

Deleted: *Research Part II: Topical Studies in Oceanography*

Deleted: .

Formatted ... [80]

Formatted ... [81]

Formatted ... [82]

Deleted: . (1979),...: Global Inventory of Natural and ... [83]

Formatted ... [84]

Formatted ... [85]

Deleted: ., &... and Pacyna, J. M. (1988). ... [86]

Deleted: .

Formatted ... [87]

Deleted: Olivelli, A., Arcucci, R., Rehkämper, M., & v ... [88]

Deleted: &...nd Rehkämper, M. (2023). ... [89]

	New constraints from concentration and isotope data, <i>Mar Pollut Bull</i> , 189, https://doi.org/10.1016/j.marpolbul.2023.114798 , 2023.	Deleted: . <i>Marine Pollution Bulletin</i> Deleted: . Formatted: ... [90] Formatted: ... [91] Deleted: &...nd Van De Flierdt, T. (2024b). ... [92] Deleted: . Deleted: and <i>Planetary Science Letters</i> , 646...lanet Sc... [94] Formatted: ... [93] Deleted: ., &... and Pacyna, E. G. (2001). ... [95] Deleted: . Deleted: (4),... 269–298. ... [97] Formatted: ... [96] Deleted: ., &... and Settle, D. M. (1987). ... [98] Deleted: . <i>Marine Chemistry</i> Deleted: (2–4),... 137–162. ... [100] Formatted: ... [99] Deleted: &...nd Schulz-Baldes, M. (1993). ... [101] Deleted: . Formatted: ... [102] Deleted: &...nd Deng, C. (2022). ... [103] Deleted: . <i>Atmospheric Chemistry and Physics</i> Deleted: (24), Formatted: ... [104] Deleted: . Deleted: &...nd Devries, T. (2019). ... [105] Deleted: . Deleted: (13),... 2617–2633. ... [107] Formatted: ... [106] Deleted: ., &... and Weiss, D. J. (2002)...: Anthrope... [108] Deleted: (1801), Formatted: ... [109] Formatted: ... [110] Deleted: . Deleted: & Deleted: . (2020). Formatted: ... [111] Deleted: . Deleted: <i>Biogeochemical</i> Deleted: (11). Formatted: ... [112] Formatted: ... [113] Moved up [6]: Roshan, S., DeVries, T., Deleted: Wu, J., & Chen, G. (2018). The Internal Cyc... [114] Deleted: . Deleted: and <i>Planetary Science Letters</i> Formatted: ... [115] Formatted: ... [116]
155	Olivelli, A., Arcucci, R., Rehkämper, M., and van de Flierdt, T.: Data for: Mapping the global distribution of lead and its isotopes in seawater with explainable machine learning [Data set], 2024a.	
160	Olivelli, A., Paul, M., Xu, H., Kreissig, K., Coles, B. J., Moore, R. E. T., Bridgestock, L., Rijkenberg, M., Middag, R., Lohan, M. C., Weiss, D. J., Rehkämper, M., and Van De Flierdt, T.: Vertical transport of anthropogenic lead by reversible scavenging in the South Atlantic Ocean, <i>Earth Planet Sci Lett</i> , 118980, https://doi.org/10.1016/j.epsl.2024.118980 , 2024b.	
165	Pacyna, J. M. and Pacyna, E. G.: An assessment of global and regional emissions of trace metals to the atmosphere from anthropogenic sources worldwide, <i>Environmental Reviews</i> , 9, 269–298, https://doi.org/10.1139/er-9-4-269 , 2001.	
170	Patterson, C. C. and Settle, D. M.: Review of data on eolian fluxes of industrial and natural lead to the lands and seas in remote regions on a global scale, <i>Mar Chem</i> , 22, 137–162, https://doi.org/10.1016/0304-4203(87)90005-3 , 1987.	
175	Pohl, C., Kattner, G., and Schulz-Baldes, M.: Cadmium, copper, lead and zinc on transects through Arctic and Eastern Atlantic surface and deep waters, <i>Journal of Marine Systems</i> , 4, 17–29, 1993.	
180	Qin, X., Zhou, S., Li, H., Wang, G., Chen, C., Liu, C., Wang, X., Huo, J., Lin, Y., Chen, J., Fu, Q., Duan, Y., Huang, K., and Deng, C.: Enhanced natural releases of mercury in response to the reduction in anthropogenic emissions during the COVID-19 lockdown by explainable machine learning, <i>Atmos Chem Phys</i> , 22, 15851–15865, https://doi.org/10.5194/acp-22-15851-2022 , 2022.	
	Rafter, P., Bagnell, A., Marconi, D., and Devries, T.: Global trends in marine nitrate N isotopes from observations and a neural network-based climatology, <i>Biogeosciences</i> , 16, 2617–2633, https://doi.org/10.5194/bg-16-2617-2019 , 2019.	
	Reuer, M. K. and Weiss, D. J.: Anthropogenic lead dynamics in the terrestrial and marine environment, <i>Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences</i> , 360, 2889–2904, https://doi.org/10.1098/rsta.2002.1095 , 2002.	
	Roshan, S., DeVries, T., Wu, J., and Chen, G.: The Internal Cycling of Zinc in the Ocean, <i>Global Biogeochem Cycles</i> , 32, 1833–1849, https://doi.org/10.1029/2018GB006045 , 2018.	
	Roshan, S., DeVries, T., and Wu, J.: Constraining the Global Ocean Cu Cycle With a Data-Assimilated Diagnostic Model, <i>Global Biogeochem Cycles</i> , 34, https://doi.org/10.1029/2020GB006741 , 2020.	
	Schaule, B. K. and Patterson, C. C.: Lead concentrations in the northeast Pacific: evidence for global anthropogenic perturbations, <i>Earth Planet Sci Lett</i> , 54, 97–116, 1981.	

Schaule, B. K. and Patterson, C. C.: Perturbations of the Natural Lead Depth Profile in the Sargasso Sea by Industrial Lead, in: *Trace Metals in Sea Water*, Springer US, Boston, MA, 487–503, https://doi.org/10.1007/978-1-4757-6864-0_29, 1983.

Schlosser, C., Karstensen, J., and Woodward, E. M. S.: Distribution of dissolved and leachable particulate Pb in the water column along the GEOTRACES section GA10 in the South Atlantic, *Deep Sea Res. I Oceanogr Res Pap.*, 148, 132–142, <https://doi.org/10.1016/j.dsr.2019.05.001>, 2019.

Shen, G. T. and Boyle, E. A.: Thermocline ventilation of anthropogenic lead in the western North Atlantic, *J Geophys Res Oceans*, 93, 15715–15732, <https://doi.org/10.1029/JC093iC12p15715>, 1988.

Sherrell, R. M., Boyle, E. A., and Hamelin, B.: Isotopic equilibration between dissolved and suspended particulate lead in the Atlantic Ocean: evidence from 210Pb and stable Pb isotopes, *J Geophys Res*, 97, 11257–11268, <https://doi.org/10.1029/92jc00759>, 1992.

Sherwen, T., Chance, R. J., Tinell, L., Ellis, D., Evans, M. J., and Carpenter, L. J.: A machine-learning-based global sea-surface iodide distribution, *Earth Syst Sci Data*, 11, 1239–1262, <https://doi.org/10.5194/essd-11-1239-2019>, 2019.

Sonnenwald, M., Lguensat, R., Jones, D. C., Dueben, P. D., Brajard, J., and Balaji, V.: Bridging observations, theory and numerical simulation of the ocean using machine learning, *Environmental Research Letters*, 16, <https://doi.org/10.1088/1748-9326/ac0eb0>, 2021.

Stirnberg, R., Cermak, J., Kotthaus, S., Haefelin, M., Andersen, H., Fuchs, J., Kim, M., Petit, J. E., and Favez, O.: Meteorology-driven variability of air pollution (PM1) revealed with explainable machine learning, *Atmos Chem Phys*, 21, 3919–3948, <https://doi.org/10.5194/acp-21-3919-2021>, 2021.

Véron, A. J., Church, T. M., Patterson, C. C., and Flegal, A. R.: Use of stable lead isotopes to characterize the sources of anthropogenic lead in North Atlantic surface waters, *Geochim Cosmochim Acta*, 58, 3199–3206, 1994.

Véron, A. J., Church, T. M., and Flegal, A. R.: Lead Isotopes in the Western North Atlantic: Transient Tracers of Pollutant Lead Inputs, *Environ Res*, 78, 104–111, 1998.

Véron, A. J., Church, T. M., Rivera-Duarte, I., and Flegal, A. R.: Stable lead isotopic ratios trace thermohaline circulation in the subarctic North Atlantic, *Deep-Sea Research II*, 46, 919–935, 1999.

Weiss, D., Boyle, A. B., Wu, J., Chavagnac, V., Michel, A., and Reuer, M. K.: Spatial and temporal evolution of lead isotope ratios in the North Atlantic Ocean between 1981 and 1989, *J Geophys Res Oceans*, 108, <https://doi.org/10.1029/2000jc000762>, 2003.

Wu, J., Rember, R., Jin, M., Boyle, E. A., and Flegal, A. R.: Isotopic evidence for the source of lead in the North Pacific abyssal water, *Geochim Cosmochim Acta*, 74, 4629–4638, <https://doi.org/10.1016/j.gca.2010.05.017>, 2010.

Formatted	... [117]
Deleted: &...nd Woodward, E. M. S. (2019).	... [118]
Deleted: .	
Formatted	... [119]
Deleted: -	
Formatted	... [120]
Deleted: Research Part I: Oceanographic Research Papers	
Formatted	... [121]
Deleted: .	
Deleted: ., &... and Boyle, E. A. (1988).... Thermoc	... [122]
Deleted: (C12),... 15715–15732.	... [124]
Formatted	... [123]
Deleted: &...nd Hamelin, B. (1992).	... [125]
Deleted: . Journal of Geophysical Research	
Deleted: (C7),... 11257–11268.	... [127]
Formatted	... [126]
Deleted: &...nd Carpenter, L. J. (2019).	... [128]
Deleted: .	
Deleted: System Science	
Deleted: (3),... 1239–1262.	... [131]
Formatted	... [129]
Formatted	... [130]
Deleted: &...nd Balaji, V. (2021).	... [132]
Deleted: .	
Deleted: (7).	
Formatted	... [133]
Deleted: &	
Deleted: . (2021).... Meteorology-driven variability	... [134]
Deleted: (5),... 3919–3948.	... [136]
Formatted	... [135]
Moved down [7]: Véron, A. J., Church, T. M.,	
Deleted: Tatsumoto, M., & Patterson, C. C. (1963).	... [137]
Deleted: & ...atterson, C. C., and Flegal, A. R. (1998)	... [138]
Deleted: Patterson, C. C., & Flegal, A. R. (1994).	
Deleted: *. Geochimica et Cosmochimica	
Formatted	... [139]
Moved (insertion) [7]	
Deleted: &...nd Flegal, A. R. (1999).	... [141]
Formatted	... [140]
Deleted: .	
Formatted	... [142]
Deleted: &...nd Reuer, M. K. (2003).	... [143]
Deleted: . Journal of Geophysical Research:	
Deleted: (10).	
Formatted	... [144]
Deleted: &...nd Flegal, A. R. (2010).	... [145]
Deleted: . Geochimica et Cosmochimica	
Deleted: (16),... 4629–4638.	... [147]
Formatted	... [146]

405

Xie, J., Liu, X., Jasechko, S., Berghuijs, W. R., Wang, K., Liu, C., Reichstein, M., Jung, M., ~~and~~ Koirala, S.: Majority of global river flow sustained by groundwater, *Nat Geosci*, 17, 770–777, <https://doi.org/10.1038/s41561-024-01483-5>, 2024.

410

Ye, M., Zhu, L., Li, X., Ke, Y., Huang, Y., Chen, B., Yu, H., Li, H., ~~and~~ Feng, H.: Estimation of the soil arsenic concentration using a geographically weighted XGBoost model based on hyperspectral data, *Science of the Total Environment*, 858, <https://doi.org/10.1016/j.scitotenv.2022.159798>, 2023.

415

~~Zheng, L., Minami, T., Konagaya, W., Chan, C. Y., Tsujisaka, M., Takano, S., Norisuye, K., and Sohrin, Y.: Distinct basin-scale-distributions of aluminum, manganese, cobalt, and lead in the North Pacific Ocean, *Geochim Cosmochim Acta*, 254, 102–121, <https://doi.org/10.1016/j.gca.2019.03.038>, 2019.~~
~~Zurbrück, C. M., Gallon, C., and Flegel, A. R.: Historic and Industrial Lead within the Northwest Pacific Ocean Evidenced by Lead Isotopes in Seawater, *Environ Sci Technol*, 51, 1203–1212, <https://doi.org/10.1021/acs.est.6b04666>, 2017.~~

Deleted: &

Deleted: . (2024).

Deleted: . *Nature Geoscience*

Deleted: (8),

Deleted: .

Formatted: Font: Not Italic

Deleted: &

Deleted: . (2023).

Deleted: .

Formatted: Font: Not Italic

Deleted: .

Formatted: Font: Not Italic

Deleted:

Page 31: [1] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
Page 31: [1] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
Page 31: [2] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
Page 31: [2] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
Page 31: [3] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
Page 31: [3] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
Page 31: [4] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
Page 31: [4] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
Page 31: [5] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
Page 31: [5] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
Page 31: [6] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
Page 31: [6] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
Page 31: [6] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
Page 31: [7] Formatted	Olivelli, Arianna	16/04/2025 10:22:00

▼.....▲
Page 31: [8] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼.....▲
Page 31: [8] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼.....▲
Page 31: [9] Formatted Olivelli, Arianna 16/04/2025 10:22:00
Font: Not Italic

▲.....▼
Page 31: [9] Formatted Olivelli, Arianna 16/04/2025 10:22:00
Font: Not Italic

▲.....▼
Page 31: [10] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼.....▲
Page 31: [10] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼.....▲
Page 31: [11] Formatted Olivelli, Arianna 16/04/2025 10:22:00
Font: Not Italic

▲.....▼
Page 31: [11] Formatted Olivelli, Arianna 16/04/2025 10:22:00
Font: Not Italic

▲.....▼
Page 31: [12] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼.....▲
Page 31: [12] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼.....▲
Page 31: [13] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼.....▲
Page 31: [13] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼.....▲
Page 31: [14] Formatted Olivelli, Arianna 16/04/2025 10:22:00
Font: Not Italic

▲.....▼
Page 31: [14] Formatted Olivelli, Arianna 16/04/2025 10:22:00

Page 31: [15] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 31: [16] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 31: [16] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 31: [16] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 31: [17] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Font: Not Italic

Page 31: [17] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Font: Not Italic

Page 31: [18] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 31: [18] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 31: [19] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 31: [19] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 31: [20] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [21] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [21] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [21] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [23] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [23] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [23] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [24] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Indent: Left: 1.69 cm, First line: 0 cm, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Page 32: [25] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [25] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [25] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [26] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [26] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [27] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [27] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [28] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [28] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [29] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [30] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Font: Not Italic

Page 32: [30] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Font: Not Italic

Page 32: [31] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [31] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [31] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [32] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [32] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [32] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [33] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [34] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [34] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 32: [34] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 33: [35] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 33: [35] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 33: [35] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 33: [36] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Font: Not Italic

Page 33: [36] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Font: Not Italic

Page 33: [37] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Font: Not Italic

Font: Not Italic

▲	▼	Page 33: [40] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
---	---	-----------------------	-------------------	---------------------

▼	▲	Page 33: [41] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
---	---	-----------------------	-------------------	---------------------

▼	▲	Page 33: [42] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
---	---	-------------------------	-------------------	---------------------

Font: Not Italic

▲	▼	Page 33: [43] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
---	---	-----------------------	-------------------	---------------------

▼	▲	Page 33: [43] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
---	---	-----------------------	-------------------	---------------------

▼	▲	Page 33: [44] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
---	---	-------------------------	-------------------	---------------------

Font: Not Italic

▲	▼	Page 33: [45] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
---	---	-----------------------	-------------------	---------------------

▼	▲	Page 33: [45] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
---	---	-----------------------	-------------------	---------------------

▼	▲	Page 33: [46] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
---	---	-------------------------	-------------------	---------------------

Font: Not Italic

▲	▼	Page 33: [46] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
---	---	-------------------------	-------------------	---------------------

Font: Not Italic

▲	▼	Page 33: [47] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
---	---	-----------------------	-------------------	---------------------

▼	▲	Page 33: [47] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
---	---	-----------------------	-------------------	---------------------

▼	▲	Page 33: [48] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
---	---	-----------------------	-------------------	---------------------

▼	▲	Page 33: [48] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
---	---	-----------------------	-------------------	---------------------

Page 33: [50] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Indent: Left: 1.69 cm, First line: 0 cm, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Page 33: [51] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 33: [51] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 33: [51] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 33: [52] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Font: Not Italic

Page 33: [53] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 33: [53] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 33: [54] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Font: Not Italic

Page 33: [54] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Font: Not Italic

Page 33: [55] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 33: [55] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 33: [56] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 33: [56] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 33: [57] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Page 33: [58] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 33: [59] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 33: [59] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 33: [59] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 33: [60] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Font: Not Italic

Page 34: [61] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Indent: Left: 1.69 cm, First line: 0 cm, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Page 34: [62] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 34: [62] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 34: [63] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Font: Not Italic

Page 34: [64] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Font: Not Italic

Page 34: [65] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 34: [65] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 34: [66] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

Page 34: [67] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

▼ ▲

Page 34: [68] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

▲ ▼

Page 34: [69] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Font: Not Italic

▲ ▼

Page 34: [69] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Font: Not Italic

▲ ▼

Page 34: [70] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Indent: Left: 1.69 cm, First line: 0 cm, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

▲ ▼

Page 34: [71] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

▲ ▼

Page 34: [71] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

▲ ▼

Page 34: [72] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Font: Not Italic

▲ ▼

Page 34: [73] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

▲ ▼

Page 34: [74] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

▲ ▼

Page 34: [74] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
-----------------------	-------------------	---------------------

▲ ▼

Page 34: [75] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Font: Not Italic

▲ ▼

Page 34: [76] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Font: Not Italic

▲ ▼

Page 34: [76] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Font: Not Italic

▲ ▼

Page 34: [77] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Font: Not Italic

▼ ▲
Page 34: [79] Formatted Olivelli, Arianna 16/04/2025 10:22:00

Indent: Left: 1.69 cm, First line: 0 cm, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

▲
Page 34: [80] Formatted Olivelli, Arianna 16/04/2025 10:22:00

Font: Not Italic

▲
Page 34: [81] Formatted Olivelli, Arianna 16/04/2025 10:22:00

Font: Not Italic

▲
Page 34: [82] Formatted Olivelli, Arianna 16/04/2025 10:22:00

Font: Not Italic

▲
Page 34: [83] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼ ▲
Page 34: [83] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼ ▲
Page 34: [84] Formatted Olivelli, Arianna 16/04/2025 10:22:00

Font: Not Italic

▲
Page 34: [85] Formatted Olivelli, Arianna 16/04/2025 10:22:00

Font: Not Italic

▲
Page 34: [86] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼ ▲
Page 34: [86] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼ ▲
Page 34: [87] Formatted Olivelli, Arianna 16/04/2025 10:22:00

Font: Not Italic

▲
Page 34: [87] Formatted Olivelli, Arianna 16/04/2025 10:22:00

Font: Not Italic

▲
Page 34: [88] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼ ▲
Page 34: [88] Deleted Olivelli, Arianna 16/04/2025 10:22:00

Page 35: [99] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
-------------------------	-------------------	---------------------

Font: Not Italic

Page 35: [100] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
------------------------	-------------------	---------------------

Page 35: [100] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
------------------------	-------------------	---------------------

Page 35: [101] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
------------------------	-------------------	---------------------

Page 35: [101] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
------------------------	-------------------	---------------------

Page 35: [102] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
--------------------------	-------------------	---------------------

Font: Not Italic

Page 35: [102] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
--------------------------	-------------------	---------------------

Font: Not Italic

Page 35: [103] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
------------------------	-------------------	---------------------

Page 35: [103] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
------------------------	-------------------	---------------------

Page 35: [104] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
--------------------------	-------------------	---------------------

Font: Not Italic

Page 35: [105] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
------------------------	-------------------	---------------------

Page 35: [105] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
------------------------	-------------------	---------------------

Page 35: [106] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
--------------------------	-------------------	---------------------

Font: Not Italic

Page 35: [106] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
--------------------------	-------------------	---------------------

Font: Not Italic

▼ ▲ Page 35: [108] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼ ▲ Page 35: [108] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼ ▲ Page 35: [108] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼ ▲ Page 35: [109] Formatted Olivelli, Arianna 16/04/2025 10:22:00

Font: Not Italic

▲ Page 35: [110] Formatted Olivelli, Arianna 16/04/2025 10:22:00

Font: Not Italic

▲ Page 35: [111] Formatted Olivelli, Arianna 16/04/2025 10:22:00

Indent: Left: 1.69 cm, First line: 0 cm, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

▲ Page 35: [112] Formatted Olivelli, Arianna 16/04/2025 10:22:00

Font: Not Italic

▲ Page 35: [113] Formatted Olivelli, Arianna 16/04/2025 10:22:00

Font: Not Italic

▲ Page 35: [113] Formatted Olivelli, Arianna 16/04/2025 10:22:00

Font: Not Italic

▲ Page 35: [114] Deleted Olivelli, Arianna 16/04/2025 10:22:00

✖ Page 35: [114] Deleted Olivelli, Arianna 16/04/2025 10:22:00

✖ Page 35: [114] Deleted Olivelli, Arianna 16/04/2025 10:22:00

✖ Page 35: [115] Formatted Olivelli, Arianna 16/04/2025 10:22:00

Font: Not Italic

▲ Page 35: [116] Formatted Olivelli, Arianna 16/04/2025 10:22:00

Font: Not Italic

▼ ▲

Page 36: [118] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
------------------------	-------------------	---------------------

▼ ▲

Page 36: [119] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
--------------------------	-------------------	---------------------

Font: Not Italic

▲

Page 36: [120] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
--------------------------	-------------------	---------------------

Font: Not Italic

▲

Page 36: [121] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
--------------------------	-------------------	---------------------

Font: Not Italic

▲

Page 36: [122] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
------------------------	-------------------	---------------------

▼ ▲

Page 36: [122] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
------------------------	-------------------	---------------------

▼ ▲

Page 36: [122] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
------------------------	-------------------	---------------------

▼ ▲

Page 36: [123] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
--------------------------	-------------------	---------------------

Font: Not Italic

▲

Page 36: [123] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
--------------------------	-------------------	---------------------

Font: Not Italic

▲

Page 36: [124] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
------------------------	-------------------	---------------------

▼ ▲

Page 36: [124] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
------------------------	-------------------	---------------------

▼ ▲

Page 36: [125] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
------------------------	-------------------	---------------------

▼ ▲

Page 36: [125] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
------------------------	-------------------	---------------------

▼ ▲

Page 36: [126] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
------------------------	-------------------	---------------------

▼

▲

Page 36: [127] Deleted

Olivelli, Arianna

16/04/2025 10:22:00

▼

▲

Page 36: [128] Deleted

Olivelli, Arianna

16/04/2025 10:22:00

▼

▲

Page 36: [128] Deleted

Olivelli, Arianna

16/04/2025 10:22:00

▼

▲

Page 36: [129] Formatted

Olivelli, Arianna

16/04/2025 10:22:00

Font: Not Italic

▼

▲

Page 36: [130] Formatted

Olivelli, Arianna

16/04/2025 10:22:00

Font: Not Italic

▼

▲

Page 36: [130] Formatted

Olivelli, Arianna

16/04/2025 10:22:00

Font: Not Italic

▼

▲

Page 36: [131] Deleted

Olivelli, Arianna

16/04/2025 10:22:00

▼

▲

Page 36: [131] Deleted

Olivelli, Arianna

16/04/2025 10:22:00

▼

▲

Page 36: [132] Deleted

Olivelli, Arianna

16/04/2025 10:22:00

▼

▲

Page 36: [132] Deleted

Olivelli, Arianna

16/04/2025 10:22:00

▼

▲

Page 36: [133] Formatted

Olivelli, Arianna

16/04/2025 10:22:00

Font: Not Italic

▼

▲

Page 36: [133] Formatted

Olivelli, Arianna

16/04/2025 10:22:00

Font: Not Italic

▼

▲

Page 36: [134] Deleted

Olivelli, Arianna

16/04/2025 10:22:00

▼

▲

Page 36: [134] Deleted

Olivelli, Arianna

16/04/2025 10:22:00

▼ ▲ Page 36: [136] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼ ▲ Page 36: [137] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼ ▲ Page 36: [138] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼ ▲ Page 36: [138] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼ ▲ Page 36: [139] Formatted Olivelli, Arianna 16/04/2025 10:22:00

Font: Not Italic

▼ ▲ Page 36: [139] Formatted Olivelli, Arianna 16/04/2025 10:22:00

Font: Not Italic

▼ ▲ Page 36: [140] Formatted Olivelli, Arianna 16/04/2025 10:22:00

Indent: Left: 1.69 cm, First line: 0 cm, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

▼ ▲ Page 36: [141] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼ ▲ Page 36: [141] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼ ▲ Page 36: [142] Formatted Olivelli, Arianna 16/04/2025 10:22:00

Font: Not Italic

▼ ▲ Page 36: [142] Formatted Olivelli, Arianna 16/04/2025 10:22:00

Font: Not Italic

▼ ▲ Page 36: [143] Deleted Olivelli, Arianna 16/04/2025 10:22:00

▼ ▲ Page 36: [143] Deleted Olivelli, Arianna 16/04/2025 10:22:00

Page 36: [145] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
------------------------	-------------------	---------------------

▼.....▲.....◀.....

Page 36: [145] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
------------------------	-------------------	---------------------

▼.....▲.....◀.....

Page 36: [146] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
--------------------------	-------------------	---------------------

Font: Not Italic

Page 36: [146] Formatted	Olivelli, Arianna	16/04/2025 10:22:00
--------------------------	-------------------	---------------------

Font: Not Italic

Page 36: [147] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
------------------------	-------------------	---------------------

▼.....▲.....◀.....

Page 36: [147] Deleted	Olivelli, Arianna	16/04/2025 10:22:00
------------------------	-------------------	---------------------

▼.....▲.....◀.....