

# Indicators of Global Climate Change 2024: annual update of key indicators of the state of the climate system and human influence

Piers M. Forster<sup>1</sup>, Chris Smith<sup>2,3</sup>, Tristram Walsh<sup>4</sup>, William F. Lamb<sup>1, 5</sup>, Robin Lamboll<sup>6</sup>, Christophe Cassou<sup>7, 8</sup>, Mathias Hauser<sup>9</sup>, Zeke Hausfather<sup>10, 11</sup>, June-Yi Lee<sup>12, 13</sup>, Matthew D. Palmer<sup>14, 15</sup>, Karina von Schuckmann<sup>16</sup>, Aimée B.A. Slangen<sup>17</sup>, Sophie Szopa<sup>18</sup>, Blair Trewin<sup>19</sup>, Jeongeun Yun<sup>12</sup>, Nathan P. Gillett<sup>20</sup>, Stuart Jenkins<sup>4</sup>, H. Damon Matthews<sup>21</sup>, Krishnan Raghavan<sup>22</sup>, Aurélien Ribes<sup>23</sup>, Joeri Rogelj<sup>3, 6, 24</sup>, Debbie Rosen<sup>1</sup>, Xuebin Zhang<sup>25</sup>, Myles Allen<sup>4, 26</sup>, Lara Aleluia Reis<sup>27, 28</sup>, Robbie M. Andrew<sup>29</sup>, Richard A. Betts<sup>14, 30</sup>, Alex Borger<sup>31</sup>, Jiddu A. Broersma<sup>31</sup>, Samantha N. Burgess<sup>32</sup>, Lijing Cheng<sup>33</sup>, Pierre Friedlingstein<sup>7, 30</sup>, Catia M. Domingues<sup>34, 35</sup>, Marco Gambarini<sup>27, 28</sup>, Thomas Gasser<sup>3</sup>, Johannes Gütschow<sup>36</sup>, Masayoshi Ishii<sup>37</sup>, Christopher Kadow<sup>38</sup>, John Kennedy<sup>39</sup>, Rachel E. Killick<sup>14</sup>, Paul B. Krummel<sup>40</sup>, Aurélien Liné<sup>8, 16, 41</sup>, Didier P. Monselesan<sup>42</sup>, Colin Morice<sup>14</sup>, Jens Mühle<sup>43</sup>, Vaishali Naik<sup>44</sup>, Glen P. Peters<sup>29</sup>, Anna Pirani<sup>45</sup>, Julia Pongratz<sup>46, 47</sup>, Jan. C. Minx<sup>1, 5</sup>, Matthew Rigby<sup>48</sup>, Robert Rohde<sup>10</sup>, Abhishek Savita<sup>49, 50</sup>, Sonia I. Seneviratne<sup>9</sup>, Peter Thorne<sup>51</sup>, Christopher Wells<sup>1</sup>, Luke M. Western<sup>48</sup>, Guido R. van der Werf<sup>52</sup>, Susan E. Wijffels<sup>42, 53</sup>, Valérie Masson-Delmotte<sup>18</sup>, Panmao Zhai<sup>54</sup>

1. Priestley Centre for Climate Futures, University of Leeds, Leeds, LS2 9JT, UK
2. Department of Water and Climate, Vrije Universiteit Brussel, Brussels, Belgium
3. International Institute for Applied Systems Analysis (IIASA), Vienna, Austria
4. Environmental Change Institute, University of Oxford, Oxford, UK
5. Potsdam Institute for Climate Impact Research (PIK), Member of the Leibniz Association, Potsdam, Germany
6. Centre for Environmental Policy, Imperial College London, London, UK
7. Laboratoire de Météorologie Dynamique/Institut Pierre-Simon Laplace, CNRS, Ecole Normale Supérieure/Université PSL, Paris, France
8. CECI, Université de Toulouse, CERFACS, CNRS, Toulouse, France
9. Institute for Atmospheric and Climate Science, Department of Environmental Systems Science, ETH Zurich, Zurich, Switzerland
10. Berkeley Earth, Berkeley, CA, USA
11. Stripe Inc., South San Francisco, CA, USA
12. Research Center for Climate Sciences, Pusan National University, Busan, Republic of Korea
13. Center for Climate Physics, Institute for Basic Science, Busan, Republic of Korea
14. Met Office Hadley Centre, Exeter, UK
15. School of Earth Sciences, University of Bristol, Bristol, UK
16. Mercator Ocean international, Toulouse, France

34 17. NIOZ Royal Netherlands Institute for Sea Research, Department of Estuarine and Delta Systems, Yerseke, the  
 35 Netherlands  
 36 18. Institut Pierre Simon Laplace, Laboratoire des Sciences du Climat et de l'Environnement (UMR 8212 CEA-CNRS-  
 37 UVSQ), Université Paris-Saclay, Gif-sur-Yvette, France  
 38 19. Bureau of Meteorology, Melbourne, Australia  
 39 20. Canadian Centre for Climate Modelling and Analysis, Environment and Climate Change Canada, Victoria, BC,  
 40 Canada  
 41 21. Concordia University, Montreal, Canada  
 42 22. Indian Institute of Tropical Meteorology, Pune, India  
 43 23. CNRM, Université de Toulouse, Météo France, CNRS, Toulouse, France  
 44 24. Grantham Institute for Climate Change and Environment, Imperial College London, United Kingdom  
 45 25. Pacific Climate Impacts Consortium, University of Victoria, Victoria, Canada  
 46 26. Atmospheric, Oceanic and Planetary Physics, Department of Physics, University of Oxford, UK  
 47 27. CMCC Foundation, Euro-Mediterranean Center on Climate Change, Lecce, Italy  
 48 28. RFF-CMCC, European Institute on Economics and the Environment, Milan, Italy  
 49 29. CICERO Center for International Climate Research, Oslo, Norway  
 50 30. Global Systems Institute, Science and Economy, University of Exeter, Exeter, UK  
 51 31. Climate Change Tracker, Data for Action Foundation, Amsterdam, the Netherlands  
 52 32. European Centre for Medium-Range Weather Forecasts, ECWMF, Reading, United Kingdom  
 53 33. State Key Laboratory of Earth System Numerical Modeling and Application, Institute of Atmospheric Physics,  
 54 Chinese Academy of Sciences, Beijing, China  
 55 34. Marine Physics and Ocean Climate, National Oceanography Centre, Southampton, UK  
 56 35. Environmental Business Unit, CSIRO, Hobart, Australia  
 57 36. Climate Resource, Melbourne, Australia  
 58 37. Meteorological Research Institute, Tsukuba, Japan  
 59 38. German Climate Computing Center, Hamburg, Germany (DKRZ)  
 60 39. No affiliation, Verdun, France  
 61 40. CSIRO Environment, Aspendale, Australia  
 62 41. Institut de Mécanique des Fluides de Toulouse, Université de Toulouse, INP, CNRS, Toulouse, France  
 63 42. CSIRO, Environment Research Unit, Climate Intelligence, Climate variability and hazards, Hobart, Tasmania,  
 64 Australia  
 65 43. Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA, USA  
 66 44. NOAA Geophysical Fluid Dynamics Laboratory, Princeton, NJ, USA  
 67 45. Euro-Mediterranean Center on Climate Change (CMCC), Venice, Italy; Università Cà Foscari, Venice, Italy  
 68 46. Ludwig-Maximilians-Universität München, München, Germany  
 69 47. Max Planck Institute for Meteorology, Hamburg, Germany

- 70 48. School of Chemistry, University of Bristol, Bristol, United Kingdom  
71 49. Centre for Atmospheric Sciences, Indian Institute of Technology Delhi, Delhi, India  
72 50. Department of Atmospheric Sciences, Rosenstiel School of Marine, Atmospheric and Earth Science, Miami, USA  
73 51. ICARUS Climate Research Centre, Maynooth University, Maynooth, Ireland  
74 52. Wageningen University and Research, Wageningen, The Netherlands  
75 53. Physical Oceanography, Woods Hole Oceanographic Institution, Woods Hole, Massachusetts, USA  
76 54. Chinese Academy of Meteorological Sciences, Beijing, China  
77

78 *Correspondence to:* Piers. M. Forster ([p.m.forster@leeds.ac.uk](mailto:p.m.forster@leeds.ac.uk))

79  
80 **Abstract.**

81 In a rapidly changing climate, evidence-based decision-making benefits from up-to-date and timely information. Here  
82 we compile monitoring datasets (published here, <https://doi.org/10.5281/zenodo.15639576>, Smith et al., 2025a) to  
83 produce updated estimates for key indicators of the state of the climate system: net emissions of greenhouse gases and  
84 short-lived climate forcers, greenhouse gas concentrations, radiative forcing, the Earth's energy imbalance, surface  
85 temperature changes, warming attributed to human activities, the remaining carbon budget, and estimates of global  
86 temperature extremes. This year, we additionally include indicators for sea-level rise and land precipitation change.  
87 We follow methods as closely as possible to those used in the IPCC Sixth Assessment Report (AR6) Working Group  
88 One report.  
89

90 The indicators show that human activities are increasing the Earth's energy imbalance and driving faster sea-level rise  
91 compared to the AR6 assessment. For the 2015–2024 decade average, observed warming relative to 1850–1900 was  
92 1.24 [1.11 to 1.35] °C, of which 1.22 [1.0 to 1.5] °C was human-induced. The 2024 observed [best estimate of](#) global  
93 surface temperature (1.52 °C) is well above the best estimate of human-caused warming (1.36°C). However, the 2024  
94 observed warming can still be regarded as a typical year, considering the human induced warming level and the state  
95 of internal variability associated with the phase of El Niño and Atlantic variability. Human-induced warming has been  
96 increasing at a rate that is unprecedented in the instrumental record, reaching 0.27 [0.2 - 0.4] °C per decade over 2015–  
97 2024. This high rate of warming is caused by a combination of greenhouse gas emissions being at an all-time high of  
98 53.6 ± 5.2 GtCO<sub>2e</sub> per year over the last decade (2014–2023), as well as reductions in the strength of aerosol cooling.  
99 Despite this, there is evidence that the rate of increase in CO<sub>2</sub> emissions over the last decade has slowed compared to  
100 the 2000s, and depending on societal choices, a continued series of these annual updates over the critical 2020s decade  
101 could track decreases or increases in the rate of the climatic changes presented here.

Deleted: <https://doi.org/10.5281/zenodo.15327155>

Deleted: (WGI)

Deleted: record in

Deleted: best estimate

106 **1 Introduction**

107 IPCC AR6 provided an assessment of human influence on key indicators of the state of climate grounded in available  
108 data at the time of publication. The preparation for the next IPCC report, the Seventh Assessment Report (AR7), has  
109 started and the assessment is due in around 5 years. Given the speed of recent change, and the need for updated climate  
110 knowledge to inform evidence-based decision-making, the Indicators of Global Climate Change (IGCC) was initiated  
111 to provide policymakers with annual updates of the latest scientific understanding on the state of selected critical  
112 indicators of the climate system and where possible of the quantified human influence upon these.

113  
114 IGCC compliments other annual updates, most notably, the BAMS State of the Climate Report (Dunn et al., 2024)  
115 and the WMO State of the Global Climate (WMO, 2024). The main difference is that this work goes beyond the  
116 observations to make process level estimates of effective radiative forcing and attributed human-induced response  
117 using methods rigorously assessed in AR6.


118  
119 This third annual update follows broadly the format of last year (Forster et al., 2024), which extended indicators  
120 through 2023. The work focuses on indicators related to heating of the climate system, building from greenhouse gas  
121 emissions towards estimates of human-induced warming and the remaining carbon budget for 1.5 °C and other policy-  
122 relevant temperature thresholds. Fig. 1 presents an overview of the aspects assessed and their interlinkages from cause  
123 (emissions) through effect (changes in physical indicators) to Climatic Impact-Drivers. It also provides a visual  
124 roadmap as to the structure of remaining sections in this paper to guide the reader.

Deleted:

Deleted: ing

Deleted: thresholds.

Deleted: .



**Figure 1 The flow chart of data production from emissions to human induced warming, the remaining carbon budget, and changes to Climatic Impact-Drivers, illustrating both the rationale and workflow within the paper production.**

The update is based on methodologies assessed by the IPCC Sixth Assessment Report (AR6) of the physical science basis of climate change (Working Group One (WGI) report; IPCC, 2021a) as well as Chap. 2 of the WGIII report (Dhakal et al., 2022) and is aligned with the efforts initiated in AR6 to implement FAIR (Findable, Accessible, Interoperable, Reusable) principles for reproducibility and reusability (Pirani et al., 2022; Iturbide et al., 2022). IPCC reports make a much wider assessment of the science and methodologies – we do not attempt to reproduce the comprehensive nature of these IPCC assessments here. We also do not consider adopting fundamentally different approaches to AR6. Rather, our aim is to rigorously track both climate system change and evolving methodological improvements between IPCC report cycles, thereby increasing transparency and consistency in between successive reports.

This annual update is organised as follows: greenhouse gas (GHG) emissions (Sect. 2), greenhouse gas concentrations (Sect. 3) and emissions of short-lived climate forcers (Sect. 4) are used to develop updated estimates of effective radiative forcing (Sect. 5). The Earth energy imbalance (Sect. 6) and observations of global surface temperature change (Sect. 7) are key global indicators of a warming world. The contributions to global surface temperature change from human and natural influences are formally attributed in Sect. 8, which tracks the level and rate of human-induced warming. Sect. 9 updates the remaining carbon budget for policy-relevant temperature thresholds. Sect. 10 gives an example of global-scale indicators associated with climate extremes of maximum land surface temperatures and Sect. 11 shows land-surface precipitation trends traceable to AR6, a new addition to this year's update. Sect. 12 presents updated estimates of global mean sea-level rise, also a new addition. Code and data availability are given in Sect. 13, and conclusions are presented in Sect. 14. Data are available at <https://doi.org/10.5281/zenodo.15639576> (Smith et al., 2025a).

Deleted: c

Deleted: <https://doi.org/10.5281/zenodo.15327155>

## **2 Greenhouse gas emissions**

Historic GHG emissions from human activity were assessed in both AR6 WGI and WGIII. Chapter 5 of WGI assessed CO<sub>2</sub> and CH<sub>4</sub> emissions in the context of the carbon cycle (Canadell et al., 2021). Chapter 2 of WGIII, published one year later (Dhakal et al., 2022), assessed the sectoral sources of emissions and gave the most up-to-date understanding of the current level of emissions. This section bases its methods and data on those employed in this WGIII chapter.

### **2.1 Methods of estimating greenhouse gas emissions changes**

Like in AR6 WGIII, net GHG emissions in this paper refer to releases of GHGs from anthropogenic sources minus removals by anthropogenic sinks, for the set of GHGs outlined in the United Nations Framework Convention on Climate Change (UNFCCC). These include: CO<sub>2</sub> emissions from fossil fuels and industry (CO<sub>2</sub>-FFI); net CO<sub>2</sub>

emissions from land use, land-use change and forestry (CO<sub>2</sub>-LULUCF); CH<sub>4</sub> emissions; N<sub>2</sub>O emissions; and fluorinated gas (F-gas) emissions comprising hydrofluorocarbons (HFCs), perfluorocarbons (PFCs), sulphur hexafluoride (SF<sub>6</sub>) and nitrogen trifluoride (NF<sub>3</sub>) - hereafter the “UNFCCC F-gases”.

Despite an extensive literature on GHG emissions, there remains important differences in reporting conventions and system boundaries between assessments. These differences relate to three underlying issues: (1) emissions data sets vary in their coverage of sources and sectors; (2) there are different approaches to determining the ‘anthropogenic’ component of LULUCF emissions and removals; and (3) the Paris Agreement does not cover all relevant sources of emissions (Lamb et al., 2025).

Concerning the first issue, there are several possible emissions datasets to draw from, each with varying coverage and update schedules. Emissions data are gathered by countries and submitted to the UNFCCC in the form of national inventory reports and common reporting tables. However, these “national inventories” are generally incomplete and are not kept up to date for all countries. Emissions reporting therefore often relies on “third-party” datasets compiled by research organisations, including: the Global Carbon Budget (GCB; Friedlingstein et al., 2024); the Emissions Database for Global Atmospheric Research (EDGAR; Crippa et al., 2023); the Potsdam Real-time Integrated Model for probabilistic Assessment of emissions Paths (PRIMAP-hist; Gütschow et al., 2016; Gütschow et al., 2025)<sup>1</sup>; the Community Emissions Data System (CEDS; Hoesly et al., 2018; Hoesly and Smith, 2024); and the Global Fire Emissions Database (GFED; van der Werf et al., 2017). As detailed below, for various reasons not all these datasets were employed in this update.

Concerning the second issue, there are varying conventions used to quantify CO<sub>2</sub>-LULUCF fluxes. These include the use of bookkeeping models and aggregated national inventory reporting (Pongratz et al., 2021), which differ in terms of their applied system boundaries and definitions, and in particular how they treat “indirect anthropogenic effects” such as the influence of increased atmospheric CO<sub>2</sub> on vegetation growth. As such, the CO<sub>2</sub>-LULUCF emissions estimates generated using bookkeeping models versus national inventories are not directly comparable and differ by about 7.5GtCO<sub>2</sub>yr<sup>-1</sup> (2013-2022 average), but there are now methods to “translate” between these two approaches (Friedlingstein et al., 2022; Grassi et al., 2023; Schwingshackl et al., 2022). Assessments also differ with respect to biomass fire emissions and to what extent components of these are treated as anthropogenic (Lamb et al., 2025).

Finally, two categories of emissions are not directly covered by the Paris Agreement but might be considered depending on the objectives of an assessment. These include the Ozone Depleting Substances (hereafter the “ODS F-gases”) comprising halons, chlorofluorocarbons (CFCs) and hydrochlorofluorocarbons (HCFCs). The ODS F-gases

---

<sup>1</sup> PRIMAP is a synthetic dataset that includes two time-series: PRIMAP Hist-TP, which is compiled from other underlying products such as EDGAR; and PRIMAP Hist-CR, which prioritises data from national inventories but gap-fills these where necessary.

were initially controlled under the Montreal Protocol and its amendments and are therefore not included in national inventories submitted to the UNFCCC, nor in many third-party emissions datasets - in contrast to the UNFCCC F-gases. Another important omission is the cement carbonation sink. To date this has also been excluded from national reporting under the UNFCCC, but plans for a new chapter covering these removals in the IPCC Task Force on National Greenhouse Gas Inventory Guidelines indicates a pathway for its eventual inclusion (IPCC, 2025).

The IPCC AR6 WGIII addressed these issues as follows. Total net GHG emissions were calculated as the sum of CO<sub>2</sub>-FFI, CH<sub>4</sub>, N<sub>2</sub>O and UNFCCC F-gases from EDGAR (version 6, with a fast-track methodology applied for the final year of data - 2019), and net CO<sub>2</sub>-LULUCF emissions from the GCB (the 2020 version; Friedlingstein et al., 2020). Net CO<sub>2</sub>-LULUCF emissions followed the GCB convention and were derived from the average of three bookkeeping models (Hansis et al., 2015; Houghton and Nassikas, 2017; Gasser et al., 2020). “Indirect anthropogenic effects” on the terrestrial carbon fluxes were therefore excluded from totals (i.e., they were treated as part of the natural land sink). Further, the GCB methodology (and thus reporting in IPCC AR6 WGIII) includes CO<sub>2</sub> emissions from deforestation and forest degradation fires, but excludes those from wildfires, which are classified as natural even if climate change affects their intensity and frequency. Similarly, the EDGAR dataset used in AR6 includes some non-CO<sub>2</sub> biomass fire emissions in the agricultural sector, but otherwise excludes those from wildfires. Sources not covered by inventories or the Paris Agreement (ODS F-gases and cement carbonation) were also excluded. Together these choices ensured consistency with the Integrated Assessment Model (IAM) benchmarks reported in WGIII and were closely focused on direct anthropogenic emissions under the UNFCCC, reflecting the importance of human-driven technology and policy options in shaping the future climate response.

The analysis presented here continues to provide an “WGIII update” estimate that tracks the same system boundary and compilation of GHGs as in AR6 WGIII, albeit with some differences in the selected data sources. As in previous years, we use GCB data for CO<sub>2</sub>-FFI. We also continue to use GCB for CO<sub>2</sub>-LULUCF, which has now been updated to use the average of four (rather than three) bookkeeping models (BLUE by Hansis et al., 2015; H&C by Houghton and Castanho, 2023; OSCAR by Gasser et al., 2020; LUCE by Qin et al., 2024). We use PRIMAP Hist-TP data for CH<sub>4</sub> and N<sub>2</sub>O, and inversions of atmospheric concentrations tracked by NOAA and AGAGE with best-estimate lifetimes for UNFCCC F-gas emissions based on analysis in the subsequent Section 3 (Lan et al., 2025; Dutton et al., 2024; Prinn et al., 2018). We follow the same approach for estimating uncertainties and CO<sub>2</sub>-equivalent emissions as in AR6, as described in the Supplement.

In addition to the WGIII update, we provide two further estimates that provide clarity and comparison to other assessment approaches. This reflects the fact that other decision criteria for tracking emissions are possible. First, in cases where assessments prioritise calculating the best estimate of fluxes to the atmosphere, it would be important to include ODS F-gases, cement carbonation and all non-CO<sub>2</sub> biomass fire emissions, including those from wildfires. Indeed, these are included in this article in subsequent assessments of concentration change (including compounds



234 formed in the atmosphere as ozone), effective radiative forcing, human-induced warming, carbon budgets and climate  
235 impacts, in line with the WGI assessment. We therefore provide an “IPCC update + additional sources and sinks”  
236 estimate that shows the change implied by including these three components in the global total. Second, the IPCC  
237 AR7 report outline foresees the tracking of “inventory-aligned” emissions that are consistent with national reporting.  
238 Full alignment between emissions inventories and WGIII emissions consistent with IAM benchmarks is essential for  
239 an accurate assessment and stocktake of the Nationally Determined Contributions (NDCs) and pathways to net-zero  
240 emissions (Grassi et al., 2021; Gidden et al., 2023; Allen et al., 2025). We therefore provide an “Inventory-aligned”  
241 estimate that follows the inventory approach to accounting for LULUCF emissions, while also integrating the latest  
242 national inventory data from the Common Reporting Tables. The data sources associated with these additional  
243 estimates are detailed in Table S1 in the Supplement.

244  
245 We expect to see differences between the three estimates, most notably between the “WGIII update” and “Inventory-  
246 aligned” estimates. As discussed above, these differ conceptually in their treatment of the LULUCF sector. However,  
247 national inventory reporting can also differ from third-party datasets in terms of underlying methods: in some  
248 countries, investments into statistical infrastructures have enabled the use of more precise emissions factors in  
249 inventories to estimate fluxes according to local or national conditions, while in others this may not be the case. In  
250 contrast, third-party datasets often use globally consistent emissions factors. Notably, the PRIMAP Hist-CR dataset,  
251 which is here used to represent national inventories, has significantly lower total CH<sub>4</sub> emissions relative to other  
252 datasets reported here, as well as the global atmospheric inversion estimates evaluated in this paper. A substantive  
253 body of literature has found that, on average, national inventories tend to underestimate CH<sub>4</sub> compared to inversions  
254 (Deng et al., 2022; Tibrewal et al., 2024; Janardanan et al., 2024; Scarpelli et al., 2022).

255 **2.2 Updated greenhouse gas emissions**

256 Updated GHG emission estimates following the WGIII assessment are presented in Fig. 2 and Table 1. Total global  
257 GHG emissions were  $55.4 \pm 5.1$  GtCO<sub>2</sub>e in 2023. Of this total, CO<sub>2</sub>-FFI contributed  $37.8 \pm 3.0$  GtCO<sub>2</sub>, CO<sub>2</sub>-LULUCF  
258 contributed  $3.6 \pm 2.5$  GtCO<sub>2</sub>, CH<sub>4</sub> contributed  $9.2 \pm 2.7$  GtCO<sub>2</sub>e, N<sub>2</sub>O contributed  $2.9 \pm 1.7$  GtCO<sub>2</sub>e and F-gas  
259 emissions contributed  $1.9 \pm 0.6$  GtCO<sub>2</sub>e.

260  
261 [Note the recent history of emissions in these datasets are continually revised, so there are small differences between](#)  
262 [each annual update in emission estimates over the recent past.](#) Initial projections for 2024 indicate that CO<sub>2</sub> emissions  
263 from fossil fuels and industry increased to  $38.2 \pm 3.0$ , and CO<sub>2</sub> emissions from land-use change increased to  
264  $4.2 \pm 2.8$  GtCO<sub>2</sub> (Friedlingstein et al., 2024; Deng et al., 2024). The significant increase in land-use change emissions  
265 is connected to high emissions from tropical deforestation and degradation fires in the aftermath of the El Niño with  
266 droughts in South America continuing since 2023. Synchronous large fires occurred in North America, where the  
267 record-breaking Canadian fires of 2023 were followed by another well above average year in 2024, but are attributable  
268 to climate variability and climate change, and not anthropogenic land-use change (Friedlingstein et al., 2024).

Deleted: 1

Deleted: 6

Deleted: 5

272  
 273 Average annual GHG emissions for the decade 2014–2023 were  $53.6 \pm 5.2$  GtCO<sub>2</sub>e. Average decadal GHG emissions  
 274 have increased steadily since the 1970s across all major groups of GHGs, driven primarily by increasing CO<sub>2</sub>  
 275 emissions from fossil fuel and industry but also rising emissions of CH<sub>4</sub> and N<sub>2</sub>O. Emissions of UNFCCC F-gases  
 276 have grown more rapidly than other GHG, but from low levels. Both the magnitude and trend of CO<sub>2</sub> emissions from  
 277 land-use change remain highly uncertain, with the latest data indicating an average net flux between 4–5 GtCO<sub>2</sub> yr<sup>-1</sup>  
 278 for the past few decades.

279  
 280 The fossil fuel share of global GHG emissions was approximately 70% in 2023 (GWP100 weighted), based on the  
 281 EDGAR v9 dataset (Crippa et al., 2023) and net land-use CO<sub>2</sub> emissions from the Global Carbon Budget  
 282 (Friedlingstein et al. 2024). The remaining share of non-fossil fuel emissions are mostly from land-use change,  
 283 agriculture, cement production, waste and F-gas emissions.

284  
 285 Different emissions assessment approaches are shown in Fig. 3. Increasing the scope of the WGIII update to include  
 286 ODS F-gases, cement carbonation, and CH<sub>4</sub> and N<sub>2</sub>O from biomass burning results in emissions of  $56.9 \pm 5.2$  GtCO<sub>2</sub>e  
 287 yr<sup>-1</sup> in 2023, or a total change of  $+1.6$  GtCO<sub>2</sub>e yr<sup>-1</sup>. ODS F-gas emissions have declined substantially since the 1990s  
 288 under the Montreal Protocol and its amendments, reaching  $1.3$  GtCO<sub>2</sub>e yr<sup>-1</sup> in 2023, with a stalling rate of reduction  
 289 in the past decade. The cement carbonation sink has steadily increased alongside cement production to reach  $-0.8$   
 290 GtCO<sub>2</sub>e yr<sup>-1</sup> in 2023. Biomass fire emissions have a more variable trend and 2023 was a relatively extreme year at  $1$   
 291 GtCO<sub>2</sub>e yr<sup>-1</sup>, compared to an average of  $0.7$  GtCO<sub>2</sub>e yr<sup>-1</sup> in the preceding decade.

292  
 293 Emissions according to national inventories were  $47.1 \pm 4.7$  GtCO<sub>2</sub>e yr<sup>-1</sup> in 2023, or  $8.3$  GtCO<sub>2</sub>e yr<sup>-1</sup> lower than the  
 294 WGIII update (Fig. 3). The main reason is due to diverging estimates of net LULUCF emissions, which according to  
 295 inventory accounts were on average  $7.5$  GtCO<sub>2</sub> lower over the past decade (2014–2023). Additional differences result  
 296 from a lower estimate of Energy, Industrial Process, Agriculture and Waste emissions in inventories ( $-1.8$  GtCO<sub>2</sub>e  
 297 yr<sup>-1</sup>), particularly for CH<sub>4</sub> ( $-0.7$  GtCO<sub>2</sub>e yr<sup>-1</sup>).

298  
 299 Emerging literature, published after AR6, suggests that increases in atmospheric CH<sub>4</sub> concentrations may also be  
 300 driven by methane emissions from wetland changes resulting from climate change and variability (e.g. Basu et al.,  
 301 2022; Hardy et al., 2023; Peng et al., 2022; Nisbet et al., 2023; Zhang et al., 2023). There is also a possible effect from  
 302 CO<sub>2</sub> fertilisation (Feron et al., 2024; Hu et al., 2023). The latest global methane budget estimates indirect  
 303 anthropogenic CH<sub>4</sub> fluxes from wetlands and freshwater bodies of approximately  $2.4$  GtCO<sub>2</sub>e yr<sup>-1</sup> (Saunois et al.,  
 304 2024). Such emissions are not captured in the WGIII estimate here as they are not a direct emission from human  
 305 activity, but rather a feedback induced by a changing climate, yet they will contribute to GHG concentration rise,  
 306 forcing and energy budget changes discussed in the next sections. They will become more important to properly

Deleted: 6

Deleted: 5


Deleted: 4

Deleted: 7

Deleted: 9

Deleted: 5

313 account for in future years. Note that these indirect CH<sub>4</sub> emissions are not used to determine the effective radiative  
314 forcing in Sect. 5.  
315



316  
317 **Figure 2** Annual global anthropogenic GHG emissions by source, 1970–2023. Refer to Sect. 2.1 and Table S1 for a list of  
318 datasets. Datasets with an asterisk (\*) indicate the sources used to compile global total greenhouse gas emissions following  
319 the WGIII assessment in (a). CO<sub>2</sub>-equivalent emissions in (a) and (f) are calculated using GWP100 from the AR6 WGI  
320 Chap. 7 (Forster et al., 2021). F-gas emissions in (a) comprise only UNFCCC F-gas emissions (see Sect. 2.1 for a list of  
321 species). F-gas emissions in (f) refer to UNFCCC F-gases, except for “CIP v2024.04 [ODS F-gases]”. Some of the major  
322 depicted differences between datasets (e.g. between GCB v2024 and Grassi NGHGI v2024 in panel c) are due to varying  
323 system boundaries, rather than underlying uncertainties in activity levels or emissions factors.

325 **Table 1 Global anthropogenic greenhouse gas emissions by source and decade following the WGIII assessment. All numbers**  
326 **refer to decadal averages, except for annual estimates in 2023 and 2024. CO<sub>2</sub>-equivalent emissions are calculated using**  
327 **GWP100 from AR6 WGI Chap. 7 (Forster et al., 2021). Projections of non-CO<sub>2</sub> GHG emissions in 2024 remain unavailable**  
328 **at the time of publication. Uncertainties are ±8 % for CO<sub>2</sub>-FFI, ±70 % for CO<sub>2</sub>-LULUCF, ±30 % for CH<sub>4</sub> and F-gases, and**  
329 **±60 % for N<sub>2</sub>O, corresponding to a 90 % confidence interval. “GHG” in row one is the sum of the other rows.**


Units: GtCO <sub>2</sub> e	1970- 1979	1980- 1989	1990- 1999	2000- 2009	2010- 2019	2014- 2023	2023	2024 (projectio n)
GHG	30.9±4.5	34.6±4.6	39.3±5.1	45.1±5.1	52.9±5.4	53.6±5.2	55.4±5.1	
CO <sub>2</sub> - FFI	17.3±1.4	20.3±1.6	23.6±1.9	28.9±2.3	35.4±2.8	36.3±2.9	37.8±3.0	38.2±3.0
CO <sub>2</sub> - LULUCF	5.2±3.7	5.1±3.6	5.7±4.0	5.2±3.6	4.9±3.4	4.1±2.9	3.6±2.5	4.2±2.8
CH <sub>4</sub>	6.3±1.9	6.7±2	7.2±2.2	7.7±2.3	8.4±2.5	8.7±2.6	9.2±2.7	
N <sub>2</sub> O	1.9±1.1	2.2±1.3	2.3±1.4	2.5±1.5	2.7±1.6	2.8±1.7	2.9±1.7	
UNFCCC F-gases	0.2±0.01	0.4±0.1	0.5±0.2	0.8±0.3	1.4±0.4	1.6±0.5	1.9±0.6	

Deleted: 1

Deleted: 6

Deleted: 5

330  
331



332 **Figure 3 Annual global anthropogenic greenhouse gas emissions by assessment convention in 2023. Refer to Table 1 for a**  
333 **list of underlying datasets. Differences between conventions are primarily due to differences in system boundaries (Lamb**  
334 **et al., 2025). Uncertainties are ±8 % for CO<sub>2</sub>-FFI, ±70 % for CO<sub>2</sub>-LULUCF, ±30 % for CH<sub>4</sub> and F-gases, and ±60 % for**  
335 **N<sub>2</sub>O, corresponding to a 90 % confidence interval.**  
336


340 **3 Well-mixed greenhouse gas concentrations**

341 As in Forster et al. (2024), we report best-estimate global mean concentrations for 52 well-mixed GHGs. These  
342 concentrations are updated to 2024. CO<sub>2</sub> mixing ratios were taken from the NOAA Global Monitoring Laboratory  
343 (GML) and are updated here through 2024 (Lan et al., 2025a). As in Forster et al. (2023, 2024), CO<sub>2</sub> is reported on  
344 the WMO-CO<sub>2</sub>-X2019 scale, which differs from the WMO-CO<sub>2</sub>-X2007 scale used in AR6 with WMO-CO<sub>2</sub>-X2019  
345 being around 0.18 ppm higher than WMO-CO<sub>2</sub>-X2007 in recent years. For consistency with WMO-CO<sub>2</sub>-X2019, the  
346 AR6 CO<sub>2</sub> concentrations that make up the 1750 to 1978 period in the IGCC dataset (before recent NOAA updates)  
347 have been converted to the WMO-CO<sub>2</sub>-X2019 scale. Other GHG records were compiled from NOAA and AGAGE  
348 global networks or extrapolated from literature. An average of NOAA and AGAGE data, updated through 2024, were  
349 used for N<sub>2</sub>O, CH<sub>4</sub>, CFC-11, CFC-12, CCl<sub>4</sub>, HCFC-22, HFC-134a, and HFC-125 (Lan et al., 2025; Dutton et al., 2024;  
350 Prinn et al., 2018), which, along with CO<sub>2</sub>, account for over 97% of the ERF from well-mixed GHGs. Several other  
351 species also use means from the NOAA and AGAGE networks, where the NOAA data is updated to 2024 from the  
352 values given in the BAMS State of the Climate Report (Dunn et al., 2024) and AGAGE data up until 2022 is available;  
353 for 2023 and 2024, an offset to the NOAA data was applied which was equal to the mean difference between the  
354 NOAA and AGAGE datasets over the recent past. In cases where no updated information is available, global estimates  
355 were extrapolated from Vimont et al. (2022), Western et al. (2023, 2024), or other literature and scaled to be consistent  
356 with those reported in AR6. Some extrapolations are based on data from the mid-2010s (Droste et al., 2020; Laube et  
357 al., 2014; Simmonds et al., 2017; Vollmer et al., 2018), but have an imperceptible effect on the total ERF assessed in  
358 Sect. 5, and are included to maintain consistency with AR6. Mixing ratio uncertainties for 2024 are assumed to be like  
359 2019, and we adopt the same uncertainties as assessed in AR6 WGI.

360  
361 Fig. 4 shows recent GHG concentrations and their changes. Table S2 in the Supplement shows specific updated  
362 concentrations for all the GHGs considered. The global surface mean concentrations of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O in 2024  
363 were 422.8 [±0.4] parts per million (ppm), 1929.7 [±3.3] parts per billion (ppb) and 337.9 [±0.4] ppb, respectively.  
364 Concentrations of all three major GHGs have increased since 2019, with CO<sub>2</sub> increasing by 12.7 ppm, CH<sub>4</sub> by 63.3  
365 ppb, and N<sub>2</sub>O by 5.8 ppb. Increases since 2019 are consistent with those from the CSIRO network (Francey et al.,  
366 1999), which are 13.0 ppm, 61.9 ppb, and 6.0 ppb for CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O, respectively. With few exceptions,  
367 concentrations of ozone-depleting substances, such as CFC-11 and CFC-12, continue to decline, while those of  
368 replacement compounds (HFCs) have increased. HFC-134a, for example, has increased 25% since 2019 from 107.6  
369 to 134.7 parts per trillion (ppt). Aggregated across all gases, PFCs have increased from 109.7 to an estimated 117.4  
370 ppt CF<sub>4</sub>-eq from 2019 to 2024, HFCs from 237 to 321.2 ppt HFC-134a-eq, while ozone depleting gases have declined  
371 from 1032 to 995.4 ppt CFC-12-eq. Mixing ratio equivalents are determined by the radiative efficiencies of each GHG  
372 from Hodnebrog et al. (2020).

Deleted: 1

Deleted: 6



**Figure 4 Atmospheric concentrations of a set of well mixed greenhouse gases over 2000-2024. The grey shaded region represents continuing changes since AR6. Note the different vertical scales.**

Ozone and other non-methane SLCFs are not well-mixed in the atmosphere and are thus discussed separately (in Section 4). For this reason, the warming impact of ozone, the third most important GHG (in terms of current

381 contribution to warming) is not included in the contribution of well-mixed GHGs to observed warming, consistently  
382 with AR6. Note that change in methane concentration affects ozone, but this indirect effect is not accounted for in the  
383 estimate of the warming due to the evolution of well mixed GHG concentrations.  
384

Deleted: in

#### 385 4 Non-methane short-lived climate forcers

386 Chapter 6 of WGI assessed emissions in the context of understanding the climate and air quality impacts of SLCFs  
387 (Szopa et al., 2021). Methane is a SLCF but also a well mixed GHG and is discussed in Sections 2 and 3. Trends in  
388 SLCFs emissions are spatially heterogeneous (Szopa et al., 2021), with strong shifts in the locations of reductions and  
389 increases over the decade 2010–2019 (Hodnebrog et al., 2024). Concentrations of non-methane SLCFs are  
390 heterogeneously distributed in the atmosphere and the observation networks are too sparse to report globally averaged  
391 concentrations. Typically, a combination of satellite data, where available, and global models and reanalysis are relied  
392 upon for producing global-scale distributions. In the case of models, production of near-real time information relies  
393 upon the availability of near-real time updates to SLCF emissions which are still challenging. Little information,  
394 whether from observations from local monitoring networks, satellite data or from global model reanalysis, is released  
395 in near real time.

396  
397 In addition to GHG emissions, we provide an update of anthropogenic emissions of non-methane SLCFs (SO<sub>2</sub>, black  
398 carbon (BC), organic carbon (OC), NO<sub>x</sub>, volatile organic compounds (VOCs), CO and NH<sub>3</sub>). Data are presented in  
399 Table 2 and the evolution of SLCF emission estimates from the AR6 to this study is presented in Sect. S4 of the  
400 Supplement. Consistency between emission trends and concentrations is considered whenever feasible. HFCs,  
401 whatever their lifetimes, were considered in Sect. 2.2.

402  
403 Sectoral emissions of SLCFs are derived from two sources: CEDS, which was used in the AR6 and in CMIP6 to  
404 assess historical evolution of atmospheric composition and that has been updated since then, and the Copernicus  
405 Atmosphere Monitoring Service (CAMS). The most recent release of the CEDS anthropogenic emissions dataset  
406 (Hoesly et al., 2025) covers the 1750-2023 period (Hoesly et al., 2018; Hoesly and Smith, 2024). Since 2023,  
407 CAMS has released regular updates of their global emission dataset (Soulie et al., 2023). For the year 2024, we  
408 apply, for each compound, the trend in emission from the CAMS dataset to the 2023 CEDS emission. The CAMS  
409 dataset is essentially based on the EDGARv6/v7 emissions as well as on CEDS, so CEDS and CAMS are not  
410 entirely independent. The temporal extension is based on evolution of drivers of emissions (energy consumption,  
411 production rates) and trends in technologies that affect the emissions factors (e.g. fleet renewal and abatement  
412 systems) (Denier van der Gon et al., 2023).

413  
414 The CAMS v6.2 emission dataset (ECCAD, 2025) indicates a decrease in global anthropogenic emissions of the  
415 primary SLCFs (NO<sub>x</sub>, CO, NMVOCs, SO<sub>2</sub>, BC and OC) since the COVID hiatus in emissions, except for NH<sub>3</sub>,

417 whose emissions are steadily increasing. SLCF emissions from biomass burning are taken from GFED (van der  
 418 Werf et al., 2017) with small fires (GFED4.1s) updated to 2024 (following AR6 WGIII (Dhakal et al., 2022)).  
 419 Estimates from GFED for 2017 to 2024 are provisional and will be updated with GFED5 in future datasets which  
 420 will provide substantially higher emissions for most species. The estimate of global carbon emissions due to  
 421 wildfires in 2024 is slightly lower than in 2023 (both were higher than average fire years). These lower overall  
 422 carbon emissions in 2024 hide an increase in CO<sub>2</sub> emissions (accompanied by an increase in NO<sub>x</sub> emissions) but a  
 423 decrease in CH<sub>4</sub> and CO emissions accompanied by a decrease in carbonaceous aerosols and NMVOC emissions.  
 424  
 425 The decrease of global NO<sub>x</sub> emissions, despite very heterogeneous regional trends (Szopa et al., 2021), is confirmed  
 426 by global NO<sub>2</sub> satellite observations from OMI (tropospheric NO<sub>2</sub> column from OMI visualised through the  
 427 Giovanni system, Acker and Leptoukh, 2007). The trends in global CO concentration are less clear. Surface data  
 428 from MOPITT and AIRS show a slight increase over the last three years. CO does not result solely from CO  
 429 emissions but also from VOC including methane oxidation which can explain differences in trends between  
 430 emissions and concentrations.  
 431  
 432 Overall, the trends in emissions were similar (see Supplement Sect. S4) over the 2020-23 period in the most recent  
 433 CEDS dataset to our previous estimate (Forster et al., 2024) but with a lower post COVID rebound for NO<sub>x</sub> and SO<sub>2</sub>.  
 434 Regarding SO<sub>2</sub>, the CEDS datasets (v2024\_04\_01 used in Forster et al., 2024 and v2025\_03\_18 used here) account  
 435 for the introduction of strict fuel sulphur controls brought in by the International Maritime Organization in January  
 436 2020. Total SO<sub>2</sub> emissions in 2019 were 80.9 TgSO<sub>2</sub> (Table 2). The SO<sub>2</sub> emissions from international shipping  
 437 declined by 8.4 TgSO<sub>2</sub> from 10.4 TgSO<sub>2</sub> in 2019 to 2.0 TgSO<sub>2</sub> in 2020, which is close to the expected 8.5 TgSO<sub>2</sub>  
 438 reduction estimated by the International Maritime Organization. This decrease was estimated at 7.4 TgSO<sub>2</sub> in the  
 439 previous CEDS version used in Forster et al. (2024). More generally, the reduction pace of the global SO<sub>2</sub> emission  
 440 over the last ten years corresponds to that of the first ten years of the SSP scenarios assuming strong air pollution  
 441 control (SSP1 and SSP5).  
 442  
 443 Using our combined estimate of GFED, and CEDS (with a 2024 extrapolation based on CAMS), emissions of all  
 444 SLCFs were reduced in 2022 relative to 2019, but rebounded in 2023 and then slightly decreased in 2024 (relative to  
 445 2023) for all compounds except NO<sub>x</sub> whose increase is partly driven by increased emissions from biomass burning  
 446 (Table 2 and Supplement Sect. S4). 2023 was a record year for emissions of organic carbon (driven again by a very  
 447 active biomass burning season) and ammonia (driven by a steady background increase in agricultural sources, plus a  
 448 contribution from biomass burning). Fires can be worsened by climate change, because of increased fire prone weather  
 449 conditions (Burton et al., 2024). Strictly speaking, such fires could sometimes be considered as feedbacks and not be  
 450 included in anthropogenic forcings. However, we choose to include fires in our tracking, as historical biomass burning  
 451 emissions inventories have previously been consistently treated as an anthropogenic forcing (for example in CMIP6),  
 452 though this assumption may need to be revisited in the future (see also discussion in Sect. 5). This differs from the  
 453 treatment of accounting for CO<sub>2</sub> and CH<sub>4</sub> emissions at present (Sect. 2.2), where we do not include natural emissions

Commented [1]: "should partly"?

Deleted: should be



in the inventories. As described in Sect. 5, this treatment of all biomass burning emissions as a forcing has implications for several categories of anthropogenic radiative forcing.

**Table 2 Emissions of the major SLCFs in 1750, 2019, and 2024 from a combination of CEDS and GFED. Emissions of SO<sub>2</sub>+SO<sub>4</sub> use SO<sub>2</sub> molecular weights. Emissions of NO<sub>x</sub> use NO<sub>2</sub> molecular weights. VOCs are for the total mass. [Note that estimates for previous years have been revised and updated. WGI 2019 estimates from Smith et al. \(2021\).](#)**

Compound	SLCF emissions (Tg yr <sup>-1</sup> )				
	1750	2019 (WGI for ERF estimates)	2019 (updated)	2023 (updated)	2024 (updated)
Sulphur dioxide (SO <sub>2</sub> ) + sulfate (SO <sub>4</sub> <sup>2-</sup> )	2.8	83.7	80.9	72.7	71.2
Black carbon (BC)	2.1	7.8	7.3	7.6	7.5
Organic carbon (OC)	15.5	29.8	33.0	41.0	36.1
Ammonia (NH <sub>3</sub> )	6.6	64.9	66.3	72.7	70.6
Oxides of nitrogen (NO <sub>x</sub> )	19.4	135.3	133.6	128.4	130.4
Volatile organic compounds (VOCs)	60.9	209.1	204.8	224.1	212.7
Carbon monoxide (CO)	348.4	855.0	816.1	896.0	845.3

Uncertainties associated with these emission estimates are difficult to quantify. From the non-biomass-burning sectors they are estimated to be smallest for SO<sub>2</sub> (±14 %), largest for black carbon (BC) (a factor of 2) and intermediate for other species (Smith et al., 2011; Bond et al., 2013; Hoesly et al., 2018). Relative uncertainties are also likely to increase both backwards in time (Hoesly et al., 2018) and again in the most recent years. Future updates of CEDS are expected to include uncertainties (Hoesly et al., 2018).

Deleted: 4.2

Deleted: 5

Deleted: 34.2

Deleted: 7.6

Deleted: 41.7

Deleted: 17.3

Deleted: 3.8

477 **5 Effective radiative forcing (ERF)**

478 ERFs were principally assessed in Chap. 7 of AR6 WGI (Forster et al., 2021), which focussed on assessing ERF from  
479 changes in atmospheric concentrations; it also supported estimates of ERF in Chap. 6 that attributed forcing to specific  
480 precursor emissions (Szopa et al., 2021) and generated the time history of ERF shown in AR6 WGI Fig. 2.10 and  
481 discussed in Chap. 2 (Gulev et al., 2021).

482  
483 The ERF calculation follows the methodology used in AR6 WGI (Smith et al., 2021) as updated by Forster et al.  
484 (2024) and described in the Supplement Sect. S5). One methodological update is incorporated in IGCC 2024 for the  
485 ERF from land use surface reflection and irrigation (Supplement Sect. S5.4). For each category of forcing, a 100,000-  
486 member probabilistic Monte Carlo ensemble is sampled to span the assessed uncertainty range in each forcing.  
487 Uncertainties account for systematic, structured random and random components. All uncertainties are reported as  
488 5 %–95 % ranges and provided in square brackets. The methods are all detailed in the Supplement, Sect. S5.

489  
490 The summary results for the anthropogenic constituents of ERF and solar irradiance in 2024 relative to 1750 are shown  
491 in Fig. 5a. In Table 3 these are summarised alongside the equivalent ERFs from AR6 (1750–2019) and last year’s  
492 Climate Indicators update (1750–2023). Fig. 5b shows the time evolution of ERF from 1750 to 2024.

493  
494 **Table 3 Contributions to anthropogenic effective radiative forcing (ERF) for 1750–2024 assessed in this section. Data is for**  
495 **single year estimates unless specified. All values are in watts per square metre ( $\text{W m}^{-2}$ ), and 5 %–95 % ranges are in square**  
496 **brackets. As a comparison, the equivalent assessments from AR6 (1750–2019) and last year’s Climate Indicators (1750–**  
497 **2023) are shown. Solar ERF is included and unchanged from AR6, based on the most recent solar cycle (2009–2019), thus**  
498 **differing from the single-year estimate in Fig. 5a. Volcanic ERF is excluded due to the sporadic nature of eruptions.**

Forcer	1750-2019 [ $\text{W m}^{-2}$ ] (AR6)	1750-2023 [ $\text{W m}^{-2}$ ] (Forster et al., 2024)	1750-2024 [ $\text{W m}^{-2}$ ]	Reason for change since last year
CO <sub>2</sub>	2.16 [1.90 to 2.41]	2.28 [2.01 to 2.56]	2.33 [2.05 to 2.61]	Increases in GHG concentrations resulting from increases in emissions
CH <sub>4</sub>	0.54 [0.43 to 0.65]	0.56 [0.45 to 0.68]	0.57 [0.45 to 0.68]	
N <sub>2</sub> O	0.21 [0.18 to 0.24]	0.22 [0.19 to 0.26]	0.23 [0.20 to 0.26]	
Halogenated GHGs	0.41 [0.33 to 0.49]	0.41 [0.33 to 0.49]	0.41 [0.33 to 0.49]	
Ozone	0.47 [0.24 to 0.71]	0.51 [0.25 to 0.76]	0.50 [0.25 to 0.75]	
Stratospheric water vapour	0.05 [0.00 to 0.10]	0.05 [0.00 to 0.10]	0.05 [0.00 to 0.11]	


|

499

Aerosol-radiation interactions	-0.22 [-0.47 to +0.04]	-0.26 [-0.50 to -0.03]	-0.22 [-0.44 to +0.01]	Decrease in most aerosol and aerosol precursor emissions (Table 2)
Aerosol-cloud interactions	-0.84 [-1.45 to -0.25]	-0.91 [-1.80 to -0.27]	-0.85 [-1.65 to -0.25]	
Land use (surface albedo changes and effects of irrigation)	-0.20 [-0.30 to -0.10]	-0.20 [-0.31 to -0.10]	-0.19 [-0.30 to -0.05]	Separation of albedo and irrigation components; updated data source and methodology
Light-absorbing particles on snow and ice	0.08 [0.00 to 0.18]	0.08 [0.00 to 0.17]	0.08 [0.00 to 0.19]	
Contrails and contrail-induced cirrus	0.06 [0.02 to 0.10]	0.05 [0.02 to 0.09]	0.06 [0.02 to 0.10]	
Total anthropogenic	2.72 [1.96 to 3.48]	2.79 [1.78 to 3.61]	2.97 [2.05 to 3.77]	Increasing positive GHG forcing and decreasing negative aerosol forcing
Solar irradiance	0.01 [-0.06 to 0.08]	0.01 [-0.06 to 0.08]	0.01 [-0.06 to 0.08]	

Deleted: 6

500






Figure 5 Effective radiative forcing (ERF) from 1750–2024. (a) 1750–2024 change in ERF, showing best estimates (bars) and 5 %–95 % uncertainty ranges (lines) from major anthropogenic components to ERF, total anthropogenic ERF and solar forcing. Note that solar forcing in 2024 is a single-year estimate and hence differs from Table 3. (b) Time evolution of ERF from 1750 to 2024. Best estimates from major anthropogenic categories are shown along with solar and volcanic forcing (thin coloured lines), total (thin black line), and anthropogenic total (thick black line). The 5 %–95 % uncertainty in the anthropogenic forcing is shown by grey shading.

Total anthropogenic ERF has increased to 2.97 [2.05 to 3.77]  $\text{W m}^{-2}$  in 2024 relative to 1750, compared to 2.72 [1.96 to 3.48]  $\text{W m}^{-2}$  for 2019 relative to 1750 in AR6. The ERF has increased considerably from the 2023 estimate of 2.79 [1.79 to 3.61]  $\text{W m}^{-2}$ . 2023 was a year associated with high biomass burning aerosol which resulted in a stronger negative aerosol forcing than recent trends. Biomass burning was also high in 2024, but lower than 2023 levels. Sulphur emissions from shipping have declined since 2020, weakening the aerosol ERF and adding around +0.1  $\text{W m}^{-2}$  over 2020 to 2024 (see Sect. 7.2 and Supplement Sects. S5 and S7). The approach of including all biomass burning aerosols is consistent with reporting ERF based on concentration increase of GHGs independent of whether  $\text{CO}_2$  and  $\text{CH}_4$  are caused by anthropogenic emissions or a smaller part is caused by any feedbacks such as from biomass burning fires or wetlands. Changes in mineral dust and sea salt are not easily relatable to human activity and are not included in the ERF of aerosols.

The ERF from well-mixed GHGs is 3.54 [3.22 to 3.85]  $\text{W m}^{-2}$  for 1750–2024, of which 2.33  $\text{W m}^{-2}$  is from  $\text{CO}_2$ , 0.57  $\text{W m}^{-2}$  from  $\text{CH}_4$ , 0.23  $\text{W m}^{-2}$  from  $\text{N}_2\text{O}$  and 0.41  $\text{W m}^{-2}$  from halogenated gases. This is an increase of around 7% from 3.32 [3.03 to 3.61]  $\text{W m}^{-2}$  for 1750–2019 in AR6. ERFs from  $\text{CO}_2$ ,  $\text{CH}_4$  and  $\text{N}_2\text{O}$  have all increased since the AR6 WG1 assessment for 1750–2019, owing to increases in atmospheric concentrations.

Deleted: 6

526  
527 The total aerosol ERF (sum of the ERF from aerosol-radiation interactions (ERF<sub>ari</sub>)  
528 and aerosol-cloud interactions (ERF<sub>aci</sub>)) for 1750–2024 is  $-1.07$  [ $-1.90$  to  $-0.43$ ]  $\text{W m}^{-2}$   
529 compared to  $-1.18$  [ $-2.10$  to  $-0.49$ ]  $\text{W m}^{-2}$  for 1750–2023 (Forster et al., 2024) and  $-1.06$   
530 [ $-1.71$  to  $-0.41$ ]  $\text{W m}^{-2}$  assessed for 1750–2019 in AR6 WGI. Attributing year-to-year trends to aerosol forcing  
531 is problematic due to the variability in biomass burning emissions. Increasing biomass burning emissions since AR6  
532 have been mostly offset by a decrease in emissions from energy and industrial sectors, leading to best estimates of  
533 ERF<sub>ari</sub> and ERF<sub>aci</sub> that are virtually unchanged from the 1750–2019 AR6 assessment to the 1750–2024 determination  
534 here (Table 3).

535  
536 Ozone ERF is determined to be  $0.50$  [ $0.25$  to  $0.75$ ]  $\text{W m}^{-2}$  for 1750–2024, slightly higher than the AR6 assessment of  
537  $0.47$  [ $0.24$  to  $0.71$ ]  $\text{W m}^{-2}$  for 1750–2019. This is due to the increase in emissions of some of its precursors (CO,  
538 VOC, CH<sub>4</sub>), but this result is highly uncertain since consolidated ozone trends are not yet released. Stratospheric water  
539 vapour from methane oxidation is unchanged (to two decimal places) since AR6. ERF from light-absorbing particles  
540 on snow and ice being  $0.08$  [ $0.00$  to  $0.19$ ]  $\text{W m}^{-2}$  for 1750–2024, like AR6. We determine from provisional data that  
541 aviation activity in 2024 has returned to pre-COVID levels (IATA, 2024). Therefore, ERF from contrails and contrail-  
542 induced cirrus is the same as in AR6, at  $0.06$  [ $0.02$  to  $0.10$ ]  $\text{W m}^{-2}$  in 2024. The methodology to determine land-use  
543 ERF has been updated (Sect. S5.4) but this forcing has a similar best estimate to 2023 and AR6, with a wider  
544 uncertainty range that accounts for the separate assessment of irrigation forcing.

545  
546 The headline assessment of solar ERF has not been re-assessed, at  $0.01$  [ $-0.06$  to  
547  $+0.08$ ]  $\text{W m}^{-2}$  from pre-industrial to the 2009–2019 solar cycle mean (Table 3). Separate to the assessment of solar  
548 forcing over complete solar cycles, we provide a single-year solar ERF for 2024 of  $+0.09$  [ $+0.01$  to  $+0.17$ ]  $\text{W m}^{-2}$   
549 (Fig. 5a). This is higher than the single-year estimate of solar ERF for 2019 (a solar  
550 minimum) of  $-0.02$  [ $-0.08$  to  $0.06$ ]  $\text{W m}^{-2}$ .

551  
552 Volcanic ERF is included in the overall time series (Fig. 5b) but following IPCC convention we do not provide a  
553 single-year estimate for 2024 given the sporadic nature of volcanoes. Alongside the time series of stratospheric aerosol  
554 optical depth derived from proxies and satellite products, for 2022–2024 we include the stratospheric water vapour  
555 contribution from the Hunga Tonga-Hunga Ha’apai (HTHH) eruption derived from Microwave Limb Sounder (MLS)  
556 data. We estimate a net positive (positive forcing from stratospheric water vapour more than outweighing negative  
557 forcing from stratospheric aerosols) forcing from HTHH through 2024 (Supplementary Material Sect. S5), though  
558 note that other studies find the net HTHH forcing to be negative (Gupta et al., 2025) or close to zero (Schoeberl et al.,  
559 2024).

560

Deleted: .6.2


## 6 Earth energy imbalance (EEI)

EEI, assessed in Chap. 7 of AR6 WGI (Forster et al., 2021), provides a measure of accumulated surplus energy (heating) in the climate system, and is hence an essential indicator to monitor the current and future status of global warming. It represents the difference between the radiative forcing acting to warm the climate, and Earth's radiative response, which acts to oppose this warming. Under stable climate conditions, i.e., in the absence of anthropogenic climate forcing, this difference would be balanced over interdecadal time scales. Since at least 1970 there has been a persistent imbalance in the energy flows that has led to excess energy being absorbed by the climate system (Forster et al., 2021). On annual and longer timescales, the global Earth heat inventory changes associated with EEI are dominated by the changes in global ocean heat content (OHC), which accounts for about 90 % of global heating since the 1970s (Forster et al., 2021). This planetary heating results in changes in all components of the Earth system such as sea-level rise, ocean warming, ice loss, rises in temperature and water vapor in the atmosphere, changes in ocean and atmospheric circulation, continental warming and permafrost thawing (e.g. Cheng et al., 2022; von Schuckmann et al., 2023a), with adverse impacts for ecosystems and human systems (Douville et al., 2021; IPCC, 2022).

On decadal timescales, changes in global surface temperatures (Sect. 5) can become decoupled from EEI by ocean heat rearrangement processes (e.g. Palmer and McNeall, 2014; Allison et al., 2020). Therefore, the increase in the Earth heat inventory arguably provides a more robust indicator of the rate of global change on interannual-to-decadal timescales (Cheng et al., 2019; Forster et al., 2021; von Schuckmann et al., 2023a). AR6 WGI found increased confidence in the assessment of change in the Earth heat inventory compared to previous IPCC reports due to observational advances and joint closure of the energy and global sea level budgets (Forster et al., 2021; Fox-Kemper et al., 2021).

AR6 estimated that EEI increased from  $0.50 [0.32\text{--}0.69] \text{ W m}^{-2}$  during the period 1971–2006 to  $0.79 [0.52\text{--}1.06] \text{ W m}^{-2}$  during the period 2006–2018 (Forster et al., 2021). The contributions to increases in the Earth heat inventory throughout 1971–2018 remained stable: 91 % for the full-depth ocean, 5 % for the land, 3 % for the cryosphere and about 1 % for the atmosphere (Forster et al., 2021). Two recent studies demonstrated independently and consistently that since 1960, the rate of warming of the world ocean is increasing at a relatively consistent pace of  $0.15 \pm 0.05 \text{ W m}^{-2}$  per decade (Minière et al., 2023; Storto and Yang, 2024; Merchant et al., 2025), while the rate of warming for the land, cryosphere, and atmosphere has been increasing at rate of  $0.013 \pm 0.003 \text{ W m}^{-2}$  per decade (Minière et al., 2023). The increase in EEI over the last several decades (Fig. 6) has also been reported by Cheng et al. (2019), von Schuckmann et al. (2020, 2023a), Loeb et al. (2021), Hakuba et al. (2021), Kramer et al. (2021), Raghuraman et al. (2021) and Minère et al. (2023). The observed increase in EEI over the most recent period (i.e. past 2 decades) are helping to drive exceptionally warm conditions (Sect. 7; Minobe et al., 2025). The increase in has been linked to rising concentrations of well-mixed GHGs and recent reductions in aerosol emissions (Sect. 5; Raghuraman et al., 2021; Kramer et al., 2021; Hansen et al., 2023), and to an increase in absorbed solar radiation associated with

597 decreased reflection by clouds and sea-ice and a decrease in outgoing longwave radiation (OLR) due to increases in  
 598 trace gases and water vapor (Loeb et al., 2021; Goesling et al., 2025; [Allan and Merchant, 2025](#)).  
 599  
 600 We carry out an update to the AR6 estimate of changes in the Earth heat inventory based on updated observational  
 601 time series for the period 1971–2020 (Table 4 and Fig. 6). Time series of heating associated with loss of ice and  
 602 warming of the atmosphere and continental land surface are obtained from the recent Global Climate Observing  
 603 System (GCOS) initiative (von Schuckmann et al., 2023b; Adusumilli et al., 2022; Cuesta-Valero et al., 2023;  
 604 Vanderkelen and Thiery, 2022; Nitzbon et al., 2022; Kirchengast et al., 2022). We use the original AR6 time series  
 605 ensemble OHC time series for the period 1971–2018 and then an updated five-member ensemble for the period 2019–  
 606 2024. We “splice” the two sets of time series by adding an offset as needed to ensure that the 2018 values are identical.  
 607 The AR6 heating rates and uncertainties for the ocean below 2000 m are assumed to be constant throughout the period.  
 608 The time evolution of the Earth heat inventory is determined as a simple summation of time series of atmospheric  
 609 heating; continental land heating; heating of the cryosphere; and heating of the ocean over three depth layers: 0–700,  
 610 700–2000 and below 2000 m (Fig. 6a). While von Schuckmann et al. (2023a) have also quantified heating of  
 611 permafrost and inland lakes and reservoirs, these additional terms are small and not included here for consistency with  
 612 AR6 (Forster et al., 2021).  
 613



615 Figure 6 (a) Observed changes in the Earth heat inventory for the period 1971–2020, with component contributions as  
616 indicated in the figure legend. (b) Estimates of the Earth energy imbalance for the IPCC AR6 assessment periods, for  
617 consecutive 20-year periods and the most recent decade. Shaded regions indicate the *very likely* range (90 % to 100 %  
618 probability). Data use and approach are based on the AR6 methods and further described in Supplement Sect. S6. For the  
619 IPCC AR6 periods our assessment closely matches that in AR6. Note the periods in our assessment overlap with different  
620 IPCC AR6 periods.

621 In our updated analysis, we find successive increases in EEI for each 20-year period since 1975, with an estimated  
622 value of 0.43 [0.03 to 0.83] W m<sup>-2</sup> during 1975–1994 that more than doubled to 0.89 [0.7 to 1.09] W m<sup>-2</sup> during  
623 2005–2024 (Fig. 6b). In addition, there is some evidence that the warming signal is propagating into the deeper ocean  
624 over time, as seen by a robust increase of ocean warming in the 700–2000m depth layer since the 1990s (von  
625 Schuckmann et al., 2020; 2023; Cheng et al., 2019, 2022). The model simulations qualitatively agree with the  
626 observational evidence (e.g. Gleckler et al., 2016; Cheng et al., 2019), further suggesting that more than half of the  
627 OHC increase since the late 1800s occurs after the 1990s.

628  
629 The update of the AR6 assessment periods to end in 2024 results in systematic increases of EEI: 0.68 W m<sup>-2</sup> during  
630 1977–2024 compared to 0.57 W m<sup>-2</sup> during 1971–2018; and 0.99 W m<sup>-2</sup> during 2012–2024 compared to 0.79 W m<sup>-2</sup>  
631 2006–2018 (Table 4). The trend and interannual variability of EEI can largely be explained by a combination of  
632 surface temperature changes and radiative forcing (Hodnebrog et al., 2024). However, there was a jump in 2023 and  
633 2024 which is still being investigated (see Sect. 7.2), but which is also discussed in the light of recent exceptional  
634 extreme climate conditions (Minobe et al., 2025).

635  
636 Table 4 Estimates of the Earth energy imbalance (EEI) for AR6 and the present study.

Time Period	Earth energy imbalance (W m <sup>-2</sup> ). Square brackets [show 90% confidence intervals].	
	IPCC AR6	This Study
1971-2018	0.57 [0.43 to 0.72]	0.57 [0.43 to 0.72]
1971-2006	0.50 [0.32 to 0.69]	0.50 [0.31 to 0.68]
2006-2018	0.79 [0.52 to 1.06]	0.79 [0.52 to 1.07]
1977-2024	-	0.68 [0.52 to 0.85]



2012-2024	-	0.99 [0.70 to 1.28]
-----------	---	---------------------

**7 Observed surface temperature change**

**7.1 Change since 1850-1900**

AR6 WGI Chap. 2 assessed the 2001–2020 globally averaged surface temperature change above an 1850–1900 baseline to be 0.99 [0.84 to 1.10] °C and 1.09 [0.95 to 1.20] °C for 2011–2020 (Gulev et al., 2021). Updated estimates to 2013–2022 of 1.15 [1.00–1.25] °C were given in AR6 SYR (Lee et al., 2023), matching the estimate in Forster et al. (2023).

There are choices around the methods used to aggregate surface temperatures into a global average, how to correct for systematic errors in measurements, methods of infilling missing data, and whether surface measurements or atmospheric temperatures just above the surface are used. These choices, and others, affect temperature change estimates and contribute to their uncertainty (AR6 WGI Chap. 2, Cross Chap. Box 2.3, Gulev et al., 2021). The methods chosen here closely follow AR6 WGI and are presented in the Supplement Sect. S7. Confidence intervals are taken from AR6 as only one of the employed datasets regularly updates ensembles (see Supplement Sect. S7).

Based on the updates available as of March 2025, the change in global surface temperature from 1850–1900 to 2015–2024 is presented in Fig. 7. These data, using the same underlying datasets (with some version changes: see Supplement Sect. S7) and methodology as AR6, estimate 1.24 [1.11–1.35] °C of warming, an increase of 0.15 °C within four years from the 2011–2020 value reported in AR6 WGI (Table 5), or 0.14 °C from the 2011–2020 value in the most recent dataset version. The decade 2015–2024 was 0.31 °C warmer than the previous decade (2005–2014). These changes, although amplified somewhat by the exceptionally warm years in 2023 and 2024, are broadly consistent with typical warming rates over the last few decades, which were assessed in AR6 as 0.76 °C over the 1980–2020 period (using ordinary-least-square linear trends) or 0.019 °C per year (Gulev et al., 2021). They are also broadly consistent with projected warming rates from 2001–2020 to 2021–2040 reported in AR6, which have a very likely range between 0.016 °C per year and 0.036 °C per year under SSP2-4.5 (Lee et al., 2021, their Table 4.5), and with human-induced warming rates discussed in Sect. 8.4.

Land temperatures have increased by 1.79 [1.56–2.03] °C from 1850-1900 to 2015-2024, and ocean temperatures by 1.02 [0.81-1.13] °C over the same period, implying that most land areas have already experienced more than 1.5 °C of warming from the 1850–1900 period. As was the case for the periods reported in AR6, the ratio of observed land

Deleted: IPCC

668 to ocean warming is in the vicinity of 1.75, somewhat higher than the ratio of 1.5 [1.4–1.7] projected by the end of the  
669 century in CMIP6 models (AR6, their Table 4.2 and Section 4.5.1.1.1). The additional observed warming since 2020  
670 in the most recent dataset versions (0.21 °C for land, 0.13 °C for ocean) has a ratio within the CMIP6 projections  
671 range.

672  
673 **Table 5 Estimates of global surface temperature change from 1850–1900 [*very likely* (90 %–100 % probability) ranges] for**  
674 **IPCC AR6 and the present study.**

Time period	Temperature change from 1850-1900 (°C)	
	IPCC AR6 (as reported)	This study
Global, most recent 10 years	1.09 [0.95 to 1.20] (to 2011-2020)	1.24 [1.11 to 1.35] (to 2015-2024)
Global, most recent 20 years	0.99 [0.84 to 1.10] (to 2001-2020)	1.09 [0.93 to 1.20] (to 2005-2024)
Land, most recent 10 years	1.59 [1.34 to 1.83] (to 2011-2020)	1.79 [1.56 to 2.03] (to 2015-2024)
Ocean, most recent 10 years	0.88 [0.68 to 1.01] (to 2011-2020)	1.02 [0.81 to 1.13] (to 2015-2024)




Figure 7 Annual (thin line) and decadal (thick line) means of global surface temperature (expressed as a change from the 1850–1900 reference period). Temperatures are based on an average of four datasets following AR6, see Supplement Sect. S7 for details.

## 7.2 2023-2024 global mean temperature -anomalies

At the time, 2023 set a new global annual-mean surface temperature change record, with a best estimate of 1.44 °C, beating 2016 by 0.16 °C. 2024 surpassed this, reaching 1.52  $\pm$  0.13 °C; 2024, becoming the first calendar year since preindustrial more likely than not exceeding 1.5 °C (Fig. 7). The assessed uncertainty range is based on that in AR6 WGI (Gulev et al., 2021). All four individual datasets are well inside the range (ranging from 1.46 to 1.56 oC). Natural drivers and internal variability are expected to modulate human-caused warming at interannual-to-decadal timescales. 2024 is assessed to be 0.16 °C warmer than the updated human-induced value (Table 6) while 2022 was 0.06 °C colder. These values are not inconsistent with AR6, which estimated the effect of internal variability in any single year be  $\pm$  0.25 °C based on CMIP6 models, nor with the lower estimated ranges ( $\pm$  0.17 °C) when calculated from observational products (Trewin, 2022).

Deleted: a best estimate of

692 The probability of seeing an observed temperature of 1.52 °C in 2024 considering a human-induced warming equal to  
 693 1.36 °C is about 1 chance out of 6 (Fig. 8a). The methodology to calculate this probability consists in comparing the  
 694 GSAT observed anomaly to those expected from CMIP6 models following the framework adopted in AR6 in Chapter  
 695 3 (Eyring et al., 2021) for decadal trends and adapted here for interannual time scale issues. The same probability but  
 696 conditional to the fact that 2024 followed an El Niño year and that the Atlantic Multidecadal Variability (AMV) was  
 697 in a positive phase (Supplement Sect. S7), rises to 1 chance out of 2. 2024 can therefore be treated as a “normal” year,  
 698 i.e. very much expected at the actual human-caused global warming level when the internal modes of variability are  
 699 taken into account and when assessed from a very large number of simulations from large ensembles. Based on the  
 700 same calculation, we estimate that a year as warm as 2023 would occur once in 4 years at human-induced warming  
 701 equal to 1.31 °C (Fig. 8b). It drops to 1-in-14 [10-20, CI 5-95%] year event, i.e. a rare-to-exceptional event, when  
 702 considering that 2023 followed a La Nina year and despite persistent positive AMV. [Note that the probability of the](#)  
 703 [large jump in global temperatures was increased by the fact that the El Niño followed an extended La Niña over 2020-](#)  
 704 [2022 \(Raghuraman et al., 2024\).](#) Within such a framework, 2022, that was colder than human-induced warming, could  
 705 be interpreted as a normal/expected year considering that 2021 was a La Nina year and AMV positive (Fig. 8c). [These](#)  
 706 [results show that human induced warming combined with particular modes of natural variability shifts the odds of](#)  
 707 [global surface temperatures passing 1.5 oC, making it more likely. Sect. 8 has a fuller discussion of human induced](#)  
 708 [warming.](#)  
 709  
 710




Figure 8 a) Gray histograms of global surface air temperature (GSAT) interannual anomalies estimated from 15 CMIP6 models extracted from all available SSP scenarios (~700 members) at anthropogenic global warming levels (ANT\_GWL) corresponding to a) 2024, b) 2023, c) 2022. The red vertical bar stands for the observational consolidated GSAT annual anomalies (Sect. 7.1). The return period of the observed annual GSAT event estimated from the CMIP6 distribution is provided (upper-corner). Associated [5-95%] likely range is assessed through bootstrapping. Interannual anomalies are obtained following Trewin (2022) method over 10-yr sliding windows. Only models providing large-ensembles (n members >5) and having at least one member whose interannual variance of GSAT is compatible with observational estimates, are selected. Colored histograms stand for the same distribution but conditional to the combined phase of El Niño Southern Oscillation (ENSO) and Atlantic Multidecadal Variability (AMV). SST Anomalies for the modes of variability are calculated from the residual of SST obtained after removing the modelled forced response estimated as model ensemble mean. A year is considered as an El Niño/La Niña year if the (October-December) Oceanic Niño Index (ONI) index of the previous year is greater/lower than one standard deviation. A year is considered as an AMV+ year if the annual North Atlantic average SST is greater than one standard deviation. Light pink represents years when ONI and AMV are concomitantly positive and light-blue when ONI is negative.

The increase in global temperature between 2022 and 2023 and in particular in global sea surface temperature is exceptional based on model estimates accounting for projected known human and natural forcings plus internal variability (Rantanen and Laaksonen, 2024; Terhaar et al., 2025; Cattiaux et al., 2024). The La Niña-to-El Niño sequence is of key importance and has been likely reinforced by enhanced energy uptake due to multi-year persistence in the preceding La Niña. The temporal synchronicity between the modes of variability in all basins is hypothesized to have played a role in the jump (Minobe et al., 2025) with the North Atlantic being record warm (Guinaldo et al., 2025) and the austral sea ice extent being record low (Purish and Doddridge, 2023).

Possible specific causes beyond internal variability, many of which are already accounted for in the estimated human-induced warming level, have been postulated e.g.: International Maritime Organization rules on shipping fuel sulphur content that came into force in January 2021; the eruption of Hunga Tonga Hunga Ha'apai in January 2022 and other subsequent smaller volcanic activity; and a faster-than-expected onset of Solar Cycle 25 (see Supplement Section S7 for details and references). A key diagnostic of these changes including both external forcing and internal variability was the exceptional magnitude of the net energy increase into the Earth system from mid-2022 to mid-2023, driven in large part by the reduced reflectance and greater absorption of solar radiation (Hodnebrog et al., 2024; Goessling et al., 2024; Minobe et al. 2025; [Allan and Merchant, 2025](#)), which may be influenced by cloud feedbacks (Tselioudis et al., 2024) as well as surface reflectance and atmospheric composition change (see also Sect. 6).

Our analysis, detailed in Supplement Sect. S7, makes use of estimates of variability and radiative forcing contributions and their uncertainty based on Sect 5. and the published literature. It shows that the increase in 2023 and 2024 compared to previous years could be explained by a combination of factors. In summary, our analyses show that, although the relative weight between the physical processes in explaining the high surface temperatures remain to be

750 better quantified, the 2023 and 2024 observed temperatures are not inconsistent with the level of human induced  
751 warming assessed next, in Sect. 8.

## 753 **8 Human contribution to surface temperature change**

754 Human-induced warming, also known as anthropogenic warming, refers to the component of observed global surface  
755 temperature increase attributable to both the direct and indirect effects of human activities, which are typically grouped  
756 as follows: well-mixed GHGs (consisting of CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O and F-gases) and other human forcings (consisting of  
757 aerosol–radiation interaction, aerosol–cloud interaction, black carbon on snow, contrails, ozone, stratospheric H<sub>2</sub>O  
758 and land use) (Eyring et al., 2021). The remaining contributors to total warming are natural: consisting of both natural  
759 forcings (such as solar and volcanic activity) and internal variability of the climate system (such as variability related  
760 to El Niño/La Nina events).

761  
762 An assessment of human-induced warming was provided in two reports within the IPCC's Sixth Assessment cycle:  
763 first in SR1.5 in 2018 [Chap. 1 Sect. 1.2.1.3 and Fig. 1.2 (Allen et al., 2018), summarised in the Summary for  
764 Policymakers (SPM) Sect. A.1 and Fig. SPM.1 (IPCC, 2018)] and second in AR6 in 2021 [WGI Chap. 3 Sect. 3.3.1.1.2  
765 and Fig. 3.8 (Eyring et al., 2021), summarised in the WGI Summary for Policymakers (SPM) Sect. A.1.3 and Fig.  
766 SPM.2 (IPCC, 2021b)], and quoted again without any updates in SYR [Sect. 2.1.1 and Fig. 2.1 (IPCC,2023a) and  
767 SYR Summary for Policymakers (SPM) Sect. A.1.2. (IPCC 2023b)].

### 768 **8.1 Warming period definitions in the IPCC Sixth Assessment cycle**

769 Temperature increases are defined relative to a baseline; IPCC assessments typically use the 1850–1900 average  
770 temperature as a proxy for the climate in pre-industrial times, referred to as the period before 1750, even though a  
771 small amount of warming likely occurred over 1750-1850 (see AR6 WGI Cross Chapter Box 1.2). Temperatures in  
772 the IPCC were reported as either GMST or GSAT, see Supplement Sect. 8.1 for details.

773  
774 Tracking progress towards the long-term global goal to limit warming, in line with the Paris Agreement, requires the  
775 assessment of both what the current level of global surface temperatures are and whether a level of global warming,  
776 such as 1.5 °C, is being reached. Definitions for these were not specified in the Paris Agreement, and several ways of  
777 tracking levels of global warming are in use; here we focus on those adopted within AR6. When determining whether  
778 warming thresholds have been passed, both AR6 and SR1.5 adopted definitions that depend on future warming; in  
779 practice, levels of current warming were therefore reported in AR6 and SR1.5 using additional definitions that  
780 circumvented the need to wait for observations of the future climate, as described next. AR6 defined crossing-time for  
781 a level of global warming as the midpoint of the first 20-year period during which the average *observed* warming for  
782 that period exceeds that level of warming (see AR6 WGI Chapter 2 Box 2.3) (the level of warming for a given year  
783 defined in this way is therefore not known until 10 years after that year). AR6 therefore reported current levels of both

784 *observed* and *human-induced* warming as their averages over just the most recent 10 years (which gives warming that  
 785 lags by only 5 years instead of 10 years) (see AR6 WGI Chapter 3 their Sect. 3.3.1.1.2); we refer to this definition as  
 786 the “AR6 decade-average” warming. SR1.5 defined the level of warming in a given year as the average *human-induced*  
 787 warming, in GMST, of a 30-year period centred on that year; when the given year is the *current* year, SR1.5 specified  
 788 that the future 15 years (required for the mean) are revealed by extrapolating the multidecadal trend (see SR1.5 Chapter  
 789 1, their Sect. 1.2.1); we refer to this definition as the “SR1.5 trend-based” warming. If the multidecadal trend is  
 790 interpreted as being linear (which it has been very close to over recent decades), this definition of current warming is  
 791 equivalent to the end-point of the trend line through the most recent 15 years of human-induced warming, and therefore  
 792 provides a definition of warming for the current year that depends only on historical warming. This interpretation  
 793 produces results that in recent years have been identical (or extremely close) to the current annual mean value of  
 794 human-induced warming (see results in Sect. 8.2, and Supplement Sect. S8.3), so in practice the attribution assessment  
 795 in SR1.5 was based not on the trend-based definition, but on the simple annual-year attributed warming; we refer to  
 796 this definition as the “SR1.5 annual-mean” warming. A diagram of these three definitions is given in Supplement Fig.  
 797 S11.  
 798

## 799 **8.2 Updated assessment approach of human-induced warming to date**

800 This paper provides an update of the AR6 WGI and SR1.5 human-induced warming assessments including, for  
 801 completeness, all three definitions (AR6 decade-average, SR1.5 trend-based [single year](#), and SR1.5 annual-mean  
 802 [single year](#)). The 2024 updates in this paper follow the same methods and process as the 2022 and 2023 updates  
 803 provided in Forster et al. (2023, 2024). Global mean surface temperature (GMST) is adopted as the definition of global  
 804 surface temperature (see Supplement Sect. S8.1). The three attribution methods used in AR6 are retained: the Global  
 805 Warming Index (GWI) (building on Haustein et al., 2017), regularised optimal fingerprinting (ROF) (as in Gillett et  
 806 al., 2021) and kriging for climate change (KCC) (Ribes et al., 2021). Details of each method, their different uses in  
 807 SR1.5 and AR6, and any methodological changes, are provided in Supplement Sect. S8.2; method-specific results are  
 808 also provided in Supplement Sect. S8.3. The overall estimate of attributed global warming for each definition (decade-  
 809 average, trend-based, and annual-mean), is based on a multi-method assessment of the three attribution methods (GWI,  
 810 KCC, ROF); the best estimate is given as the 0.01 °C-precision mean of the 50th percentiles from each method, and  
 811 the *likely* range is given as the smallest 0.1 °C-precision range that envelops the 5th to 95th percentile ranges of each  
 812 method. This assessment approach is identical to last year’s update (Forster et al. (2024)); it is directly traceable to  
 813 and fully consistent with the assessment approach in AR6, though it has been lightly extended in ways that are  
 814 explained in Supplement Sect. S8.4.

815

816 Results are summarised in Table 6 and Fig. 9. Method-specific contributions to the assessment results, along with time

817 series, are given in the Supplement, Sect. S8.3. Where results reported in GSAT differ from those reported in GMST

818 (see Supplement Sect. S8.1), the additional GSAT results are given in Supplement Sect. S8.3.

819






Figure 9 Updated assessed contributions to observed warming relative to 1850–1900; see AR6 WGI SPM.2. Results for all time periods in this figure are calculated using updated datasets and methods. The 2015–2024 average and 2024 results are this year’s updated assessments for AR6 and SR1.5, respectively. Panel (a) shows updated observed global warming from Sect. 7, expressed as total global mean surface temperature (GMST), due to both anthropogenic and natural influences. Whiskers give the “very likely” range. Panels (b) and (c) show updated assessed contributions to warming, expressed as global mean surface temperature (GMST), from natural forcings and total human-induced forcings, which in turn consist of contributions from well-mixed GHGs and other human forcings. Whiskers give the “likely” range. Changes to warming levels since the IPCC sixth assessment cycle are depicted in Supplement Fig. S10.

Table 6 Updates to assessments in the IPCC 6th assessment cycle of warming attributable to multiple influences. Estimates of warming attributable to multiple influences, in °C, relative to the 1850–1900 baseline period. Results are given as best estimates, with the likely range in brackets, and reported as global mean surface temperature (GMST). Results from the IPCC 6th assessment cycle, for both AR6 and SR1.5, are quoted in columns labelled (i) and are compared with repeat calculations in columns labelled (ii) for the same period using the updated methods and datasets to see how methodological and dataset updates alone would change previous assessments. Assessments for the updated periods are reported in columns labelled (iii). \* Updated GMST observations, quoted from Sect. 7 of this update, are marked with an asterisk, with “very likely” ranges given in brackets. \*\* In AR6 WGI, best-estimate values were not provided for warming attributable to well-mixed GHGs, other human forcings and natural forcings (though they did receive a “likely” range); for comparison, best estimates (marked with two asterisks) have been retrospectively calculated in an identical way to the best estimate that AR6 provided for anthropogenic warming (see discussion in Supplement Sect. S8.4.1). \*\*\* The SR1.5 assessment drew only on GWI rounded to 0.1°C precision, whereas the repeat and updated calculations use the updated multi-method assessment approach.

Estimates of warming attributable to multiple influences, in °C, relative to the 1850–1900 baseline period Results are given as best estimates, with the likely range in brackets, and reported as Global Mean Surface Temperature (GMST).						
Definition →	(a) IPCC AR6 Attributable Warming Update			(b) IPCC SR1.5 Attributable Warming Update		
	Value for decade (average of previous 10-year period)			Value for single year (30-year mean centred on current year)		
Period →	(i) 2010–2019 Quoted from AR6 Chapter 3 Sect. 3.3.1.1.2	(ii) 2010–2019 Repeat calculation using the updated methods and datasets	(iii) 2015–2024 Updated value using updated methods and datasets	(i) 2017 Quoted from SR1.5 Chapter 1 Sect. 1.2.1.3	(ii) 2017 Repeat calculation using the updated methods and datasets	(iii) 2024 Updated value using updated methods and datasets
Component ↓	Table 3.1 for attributed warming, and Cross-chapter Box 2.3 Table 1 for observed warming					
Observed	1.06 [0.92 to 1.17]	1.07 [0.89 to 1.22] *	1.24 [1.11 to 1.35] *	-	-	1.52 [1.39 to 1.65]
Anthropogenic	1.07 [0.8 to 1.3]	1.09 [0.9 to 1.3]	1.22 [1.0 to 1.5]	1.0 [0.8 to 1.2] ***	1.13 [0.9 to 1.3]	1.36 [1.1 to 1.7]
Well-mixed GHGs	1.40** [1.0 to 2.0]	1.40 [1.0 to 1.9]	1.51 [1.0 to 2.0]	N/A	1.45 [1.0 to 1.9]	1.62 [1.1 to 2.1]
Other human forcings	-0.32** [-0.8 to 0.0]	-0.30 [-0.8 to 0.1]	-0.28 [-0.8 to 0.1]	N/A	-0.31 [-0.8 to 0.1]	-0.26 [-0.8 to 0.1]

Deleted:

Deleted: 1

Deleted: 0

<b>Natural forcings</b>	0.03** [-0.1 to 0.1]	0.05 [-0.1 to 0.2]	0.05 [-0.1 to 0.2]	N/A	0.05 [-0.1 to 0.2]	0.04 [-0.1 to 0.2]
-------------------------	----------------------	--------------------	--------------------	-----	--------------------	--------------------

845  
846 The repeat calculations for attributable warming in 2010–2019 exhibit good correspondence with the results in AR6  
847 WGI for the same period (see also Supplement, Sect. S8). The repeat calculation for the level of attributable  
848 anthropogenic warming in 2017 is about 0.1 °C larger than the estimate provided in SR1.5 for the same period,  
849 resulting from changes in methods and observational data (see AR6 WGI Chapter 2 Box 2.3). The updated results for  
850 warming contributions in 2024 are higher than in 2017 due also to 7 additional years of increasing anthropogenic  
851 forcing. Note also that the SR1.5 assessment only used the GWI method, whereas these annual updates apply the full  
852 AR6 multi-method assessment (see Supplement Sect. S8.4 for details and rationale).

853  
854 In this 2025 update, we assess the 2015–2024 decade average human induced-warming at 1.22 [1.0 to 1.5] °C, which  
855 is 0.15°C above the AR6 assessment for 2010–2019. The single year average human-induced warming is assessed to  
856 be 1.36 [1.1 to 1.7] °C in 2024 relative to 1850–1900. In general, these forced warming levels have evolved steadily  
857 and predictably in line with the current warming rate within uncertainty. The uncertainty range for the single-year  
858 level of anthropogenic warming already included 1.5 °C in previous years’ assessments, and for the first time this year  
859 also lies at the edge of the uncertainty range for the (lagged) decade mean definition. The single-year anthropogenic  
860 warming best estimate is well below the observed best estimate for 2024 (1.52 °C, see Sect. 7), but note that the best  
861 estimate and lower uncertainty for observed warming lies within the uncertainty for single-year anthropogenic  
862 warming from each of the three attribution methods (see Supplement Table S5), whereas the upper uncertainty range  
863 of observed warming lies above the range for anthropogenic warming for the two attribution methods that fully exclude  
864 internal variability.

865  
866 The best estimates for decade-average and single-year human-induced warming are 0.04 °C and 0.05 °C respectively  
867 above the value estimated in the previous update for the year 2023 (Forster et al., 2024), but should not be interpreted  
868 as a substantive increase in the rate of forced anthropogenic warming, as the rate increase is well within uncertainty  
869 ranges (Sect. 8.3).


870  
871 AR6 found that, averaged for the 2010–2019 period, essentially all observed global surface temperature change was  
872 human-induced, with solar and volcanic drivers and internal climate variability making a negligible contribution. This  
873 conclusion remains the same for the 2015–2024 period. Generally, whatever methodology is used, on a global scale,  
874 the best estimate of the current level of human-induced warming is (within uncertainty) similar to the observed global  
875 surface temperature change (Table 6).

876

Deleted: small  
Deleted: ies

879 **8.3 Rate of human-induced global warming**

880 Estimates of the human-induced warming rate follow the same methodology as in the previous year’s update (a rolling  
881 10-year linear trend in attributed anthropogenic warming). A full description of the approach can be found in the  
882 Supplement Sect. S8.5. The rate of increase in attributed anthropogenic warming over time is distinct from the rate of  
883 increase in the observed global surface temperature, which is also affected by internal variability such as El Niño and  
884 natural forcings such as volcanic activity (see discussions in Sect. 7.2). The rate of anthropogenic warming we estimate  
885 here is driven by the rate of change of anthropogenic ERF (Sect. 5), with variations in the climate forcing trend over  
886 time correlating with variations in the rate of attributed warming (Fig. 10).  
887



888  
889 **Figure 10** Rates of (a) attributable warming (global mean surface temperature (GMST)) and (b) effective radiative forcing. The attributable warming rate time-series are calculated using the Global Warming Index method with full ensemble uncertainty. The observed GMST rates included for reference are also calculated with uncertainty from the HadCRUT5 ensemble, and, for consistency with the attributed warming rates, do not include standard regression error, which, for observed warming, would increase the size of the error bars. The effective radiative forcing rates are calculated using a representative 1000-member ensemble of the forcings provided in Sect. 5 of this paper. The depicted rates are the decadal rates, with the end year of the decade in question being the value given on the time axis.

897  
898 Estimates for the trend derived from the three warming attribution methodologies are presented in Table 7, with results  
899 for individual attribution methods detailed in the Supplement Table S6. The GWI (based on observed warming and  
900 forcing) and KCC (based on CMIP simulations) methodologies report results that are in close agreement, while  
901 estimates derived with the ROF method (also based on CMIP simulations) are more strongly influenced by residual  
902 internal variability that remains in the anthropogenic warming signal due to the limitations in size of the [available](#)

CMIP ensemble. The median result is presented at 0.01 °C/decade precision for the overall multi-method rate of warming assessment.

An overall best estimate attributed rate of human-induced warming of 0.27 °C/decade is found for the decade 2015–2024. This increased rate relative to the 0.2 °C/decade AR6 assessment is broken down in the following way: (i) 0.03 °C/decade from changing the rounding precision (updating the AR6 2010–2019 warming rate assessment from 0.2 to 0.23 °C/decade), (ii) 0.03 °C/decade is due to methodological and dataset updates (updating the 2010–2019 warming rate from 0.23 °C/decade to 0.26 °C/decade; including the effect of adding 5 additional observed years to the attribution over the entire historical period), and (iii) 0.01 °C/decade due to a real increase in rate for the 2015–2024 period since the 2010–2019 period (updating 0.26 °C/decade for 2010–2019 to 0.27 °C/decade for 2015–2024), consistent with increased GHG emissions over the last decade. The spread of rates across the three attribution methods remains similar to their spread in AR6, and previous updates of this work, and hence does not support a decrease in the headline uncertainty range. However, as previous assessments suggested, we update the uncertainty range for the rate of human-induced warming from [0.1–0.3] °C/decade in AR6 to [0.2–0.4] °C/decade to better reflect the closer agreement of the 5% floors and the larger spread in the 95% ceilings of the three methods, and higher rate from the ROF method. The rate of human-induced warming for the 2015–2024 decade is concluded to be 0.27 °C/decade with a range of [0.2–0.4] °C/decade. This agrees with the decadal trend in observed warming of 0.26 °C per decade (also calculated as a linear trend through 10-year periods - see Sect. 7.1). It is important to note, however, that internal variability leads to the decadal rates of observed warming being far less stable than for anthropogenic warming, and the very close correspondence between the two this year is somewhat incidental (see Fig. 10).

**Table 7 Updates to the IPCC AR6 rate of human-induced warming.** Results for each method are given in the Supplement Table S6; assessment results are given as a best estimate with *likely* range in brackets. Results from AR6 WGI (Ch.3 Sect. 3.3.1.1.2 Table 3.1) are quoted in column (i), and compared with a repeat calculation using the updated methods and datasets in column (ii), and finally updated for the 2015–2024 period in column (iii). The AR6 assessment result was identical to the SR1.5 assessment result, though the latter was based on a different set of studies and timeframes. \* Note that for clarity and ease of comparison with this year’s updated assessment, the assessed rate in column (i) both quotes the assessment from AR6 and retrospectively applies the median approach adopted in this paper. The observed rates are calculated using the multi-dataset observed temperature dataset from Sect. 7; no ensemble is available for this, hence the absence of an uncertainty range.

Deleted: in

Estimates of anthropogenic warming rate, in °C per decade			
Results are given as best estimates, with brackets giving the <i>likely</i> range for the assessments, and 5-95% uncertainty for the individual methods			
Definition ➡	<b>IPCC AR6 Anthropogenic Warming Rate Update</b> <i>Linear trend in anthropogenic warming over the trailing 10-year period</i>		
Period ➡	<b>(i) 2010-2019</b> <i>Quoted from AR6 Chapter 3 Sect. 3.3.1.1.2 Table 3.1</i>	<b>(ii) 2010-2019</b> <i>Repeat calculation using the updated methods and datasets</i>	<b>(iii) 2015-2024</b> <i>Updated value using updated methods and datasets</i>
<b>Anthropogenic Warming Rate Assessment</b>	Quoted from AR6: 0.2 [0.1 to 0.3]	0.26 [0.2 to 0.4]	0.27 [0.2 to 0.4]

	Using the median approach: 0.23 [0.1 to 0.3] *		
Observed		0.37	0.26

934

935

936   **9 Remaining Carbon Budget**

937   AR5 (IPCC, 2013) assessed that long-term global surface temperature increase caused by CO<sub>2</sub> emissions is close to  
938   linearly proportional to the total amount of cumulative CO<sub>2</sub> emissions (Collins et al., 2013). The most recent AR6  
939   report reaffirmed this assessment and highlights that this near-linear relationship also holds between cumulative CO<sub>2</sub>  
940   emissions and maximum global surface temperature increase caused by CO<sub>2</sub> (Canadell et al., 2021). This near-linear  
941   relationship implies that for keeping global warming below a specified temperature level, one can estimate the total  
942   amount of CO<sub>2</sub> that can ever be emitted. When expressed relative to a recent reference period, this is referred to as the  
943   remaining carbon budget (Rogelj et al., 2018).

944

945   AR6 assessed the remaining carbon budget (RCB) in Chap. 5 of its WGI report (Canadell et al., 2021) for warming  
946   limits ranging from 1.3 to 2.4 °C relative to the 1850-1900 period (see Table 5.8 in Canadell et al., 2021). A selection  
947   of these (1.5, 1.7, and 2 °C) were also reported in its Summary for Policymakers (Table SPM.2, IPCC, 2021b). These  
948   RCB values are updated in this section using the same method as last year (Forster et al., 2024). Data for four warming  
949   limits (1.5, 1.6, 1.7 and 2 °C) are included in Table 8 while figures for more values are included in the Supplement  
950   Sect. S9.

951

952   The RCB is estimated by application of the WGI AR6 method described in Rogelj et al. (2019), which involves the  
953   combination of the assessment of five factors: (i) the amount of human-induced warming for the most recent decade  
954   (given in Sect. 8), (ii) the transient climate response to cumulative emissions of CO<sub>2</sub> (TCRE), which quantifies the  
955   linear proportionality between cumulative CO<sub>2</sub> emissions and CO<sub>2</sub>-induced warming (iii) the zero emissions  
956   commitment (ZEC), representing the expected amount of additional (at present unrealized) warming caused by past  
957   CO<sub>2</sub> emissions (iv) the temperature contribution of future non-CO<sub>2</sub> emissions and (v) an adjustment term for Earth  
958   system feedbacks that are otherwise not captured through the other factors. AR6 WGI reassessed all five terms  
959   (Canadell et al., 2021). Lamboll et al. (2023) further considered the temperature contribution of non-CO<sub>2</sub> emissions  
960   and integrated different uncertainties, while Rogelj and Lamboll (2024) clarified the reductions in non-CO<sub>2</sub> emissions  
961   that are assumed in the RCB estimation.

962

963   The RCB for 1.5, 1.6, 1.7 and 2 °C warming levels is re-assessed based on the most recent available data. Estimated  
964   RCBs are reported in Table 8. They are expressed relative to the start of 2025 for estimates based on the 2015–2024

human-induced warming update (Sect. 8). Based on the variation in non-CO<sub>2</sub> emissions across the scenarios in AR6 WGIII scenario database, the estimated RCB values can be higher or lower by around 200 GtCO<sub>2</sub> depending on how successful non-CO<sub>2</sub> emissions reductions are (Lamboll et al., 2023; Rogelj and Lamboll, 2024). Notably, RCB estimates consider the subset of non-CO<sub>2</sub> emission scenarios in the AR6 WGIII database that are aligned with a global transition to net zero CO<sub>2</sub> emissions (Lamboll et al., 2023; Rogelj and Lamboll, 2024). These estimates assume median reductions in non-CO<sub>2</sub> emissions between 2020–2050 of CH<sub>4</sub> (about 50 %), N<sub>2</sub>O (about 20 %) and SO<sub>2</sub> (about 80 %) (see Supplement, Sect. S9 and Table S7 and (Rogelj and Lamboll, 2024)). If these non-CO<sub>2</sub> GHG emission reductions are not achieved, the RCB for all temperature targets would be smaller than the values reported here in Table 8 (see Lamboll et al., 2023, Rogelj and Lamboll, 2024).

Compared to RCB values reported in AR6, our estimates here are smaller owing to several factors. First, AR6 budgets were expressed from 2020 onwards, and approximately 200 GtCO<sub>2</sub> have been emitted between 2020 and 2024. Second, we use updated physical models of non-CO<sub>2</sub> forcing which lead to an increased estimate of the importance of aerosols that are expected to decline with time in low emissions pathways (Rogelj et al., 2014; Rogelj and Lamboll, 2024). This decreased negative forcing from aerosols is expected to cause additional net non-CO<sub>2</sub> warming because more non-CO<sub>2</sub> GHG warming is being unmasked and this decreases the RCB (Lamboll et al., 2023) by slightly over 100 GtCO<sub>2</sub>. There was also a small reduction in the budget (about 10 GtCO<sub>2</sub>) from using the newer AR6 scenario set. Finally, the updated warming estimate reported in Sect. 8 is slightly increased due to the high observed temperatures in the last few years, which resulted in a further reduction of the budget by around 40 GtCO<sub>2</sub>, relative to values reported in last year’s assessment (Forster et al. 2024). This gives a total reduction in RCB values estimated from the beginning of 2025 of ~370 GtCO<sub>2</sub> compared to the values from 2020 reported in AR6.

**Table 8 Updated estimates of the remaining carbon budget for 1.5, 1.6, 1.7 and 2.0 °C, for five levels of likelihood, considering only uncertainty in TCRE. Estimates are expressed relative to the start of 2024. The probability includes only the uncertainty in how the Earth immediately responds to CO<sub>2</sub> emissions (TCRE), not long-term committed warming or uncertainty in the climate response to other non-CO<sub>2</sub> emissions. All values are rounded to the nearest 10 GtCO<sub>2</sub>. Additional values can be found in the Supplement Tables S7 and S8.**

Temperature (°C)	Estimated remaining carbon budgets from the beginning of 2025 (GtCO <sub>2</sub> )				
Avoidance probability:	17%	33%	50%	67%	83%
1.5	320	200	130	80	30
1.6	620	420	310	240	160
1.7	910	640	490	390	290

2.0	1790	1310	1050	870	690
-----	------	------	------	-----	-----

This year's update of the 1.5 °C budget uses the historical warming level for the 2015-2024 period of 1.24 °C, with 0.11 °C future contribution of non-CO<sub>2</sub> warming. Assuming a median TCRE estimate of 0.45 °C per 1000 GtCO<sub>2</sub> this gives around 340 GtCO<sub>2</sub> from the midpoint of the period, from which we subtract around 210 GtCO<sub>2</sub> (204 GtCO<sub>2</sub> that were already emitted from the middle until the end of the 2015-2024 period, and 7 GtCO<sub>2</sub> that represents the median estimate of the impact of Earth systems feedbacks such as permafrost feedback that would otherwise not be covered). The same method is used to calculate budgets for the other warming levels.

The values in Table 8 are all greater than zero, implying that we have not yet emitted the amount of CO<sub>2</sub> that would commit us to these levels of warming. However, including the uncertainty in ZEC (as in the Supplement Table S8), non-CO<sub>2</sub> emission and forcing uncertainty, and underrepresented Earth-system feedbacks results in negative RCB estimates for limiting warming to low temperature limits with high likelihood. A negative RCB for a specific temperature limit would mean that the world is already committed to this amount of warming, and that net negative emissions would therefore be required to return to the temperature limit after a period of overshoot. The assumption behind such a calculation is that we can treat the warming impact of positive and negative net emissions as approximately symmetric. While the claim of symmetry is likely valid for small emissions values, some model studies have shown that it holds less well for reversal of larger emissions (Canadell et al., 2021, Zickfeld et al., 2021, Vakilifard et al., 2022, Pelz et al., 2025) As such, larger exceedances of the RCB for a particular temperature target would decrease the likelihood that the temperature target could still be achieved by an equivalent amount of net negative emissions.


Note that the 50 % RCB estimate of 130 GtCO<sub>2</sub> would be exhausted in a little more than 3 years if global CO<sub>2</sub> emissions remain at 2024 levels (42 GtCO<sub>2</sub>/yr, see Table 1). This is not expected to correspond exactly to the time that 1.5 °C global warming level is reached due to uncertainty associated with committed warming from past CO<sub>2</sub> emissions (the ZEC) as well as ongoing warming and cooling contributions from non-CO<sub>2</sub> emissions. For comparison, our estimate of 2024 anthropogenic warming (1.36 °C) and the recent rate of increase (0.27 °C/decade) would suggest that continued emissions at current levels would cause human-induced global warming to reach 1.5°C in approximately 5 years.

#### 10 Indicator of climate and weather extremes: land average maximum temperatures

Changes in climate and weather extremes are among the most visible effects of human-induced climate change. Within AR6 WGI, a full chapter was dedicated to the assessment of past and projected changes in extremes on continents (Seneviratne et al., 2021), and the chapter on ocean, cryosphere and sea level changes also provided assessments on changes in marine heatwaves (Fox-Kemper et al., 2021). Global indicators related to climate extremes include averaged changes in climate extremes, for example, the mean increase of annual minimum and maximum temperatures

on land (AR6 WGI Chap. 11, Fig. 11.2, Seneviratne et al., 2021) or the area affected by certain types of extremes (AR6 WGI Chap. 11, Box 11.1, Fig. 1, Seneviratne et al., 2021; Sippel et al., 2015).

The presented climate indicator for changes in temperature extremes consists of land average maximum temperatures for any single day in a year (TXx) (excluding Antarctica). Fig. 11 updates the land mean TXx shown in Forster et al. (2023, 2024), originally based on Fig. 11.2 from Seneviratne et al. (2021). Three datasets are analyzed: HadEX3 (Dunn et al., 2020), Berkeley Earth Surface Temperature (building off Rohde et al., 2013), and the fifth-generation ECMWF atmospheric reanalysis of the global climate (ERA5; Hersbach et al., 2020). HadEX3 is static and has not received any updates. Berkeley Earth has been extended and updated compared to Forster et al. (2024), resulting in TXx differences for most years (less than 0.1°C), and now includes data for 2023. Of the three datasets, only ERA5 covers the whole of 2024 at the present time. TXx is calculated by averaging the annual maximum temperature over all available land grid points (excluding Antarctica) and then converted to anomalies with respect to a base period of 1961–1990. To express the TXx as anomalies with respect to 1850–1900, we add an offset of 0.51 °C to all three datasets. See Supplement Sect. S10 for details on the data selection, averaging and offset computation.



**Figure 11** Time series of observed temperature anomalies for land average annual maximum temperature (TXx) for ERA5 (1950–2024), Berkeley Earth (1955–2023) and HadEX3 (1961–2018), with respect to 1850–1900. The datasets have different spatial coverage and are not coverage-matched. All anomalies are calculated relative to 1961–1990, and an offset of 0.51 °C is added to obtain TXx values relative to 1850–1900. Note that while the HadEX3 numbers are the same as shown in Seneviratne et al. (2021) Fig. 11.2, these numbers were not specifically assessed.

Our climate has warmed rapidly in the last few decades (Sect. 7), which also manifests in changes in the occurrence and intensity of climate and weather extremes. From about 1980 onwards, all datasets point to a strong TXx increase,



1047 which coincides with the transition from global dimming, associated with aerosol increases, to brightening, associated  
1048 with aerosol decreases (Wild et al., 2005, Sect. 4). The ERA5 based TXx warming estimate w.r.t. 1850–1900 for 2024  
1049 is at 2.35 °C; an increase of 0.05 °C compared to 2023, and thus even warmer than the previous record in 2023. On  
1050 longer time scales, land average TXx has warmed 0.49 °C in the past 10 years (comparing the decades 2015–2024 to  
1051 2005–2014) and 1.90 °C with respect to pre-industrial conditions (Table 9). Since the offset relative to our pre-  
1052 industrial baseline period is calculated over 1961–1990, temperature anomalies align by construction over this period  
1053 but can diverge afterwards.

1054  
1055 **Table 9 Anomalies of land average annual maximum temperature (TXx) for recent decades based on HadEX3, Berkeley**  
1056 **Earth, and ERA5, with respect to 1850–1900. All anomalies are calculated relative to 1961–1990, and an offset of 0.51 °C is**  
1057 **added to obtain TXx values relative to 1850–1900.**


	HadEX3	Berkeley Earth	ERA5
2000–2009	1.23	1.18	1.21
2005–2014	1.37	1.31	1.4
2009–2018	1.52	1.41	1.54
2011–2020	-	1.45	1.63
2013–2022	-	1.52	1.72
2014–2023	-	1.6	1.81
2015–2024	-	-	1.9

1058  
  
1059 **11 Global land precipitation**  
  
1060 Anthropogenic radiative forcings modify the Earth’s energy budget and subsequently drive substantial and widespread  
1061 changes in the global water cycle including precipitation, evaporation, atmospheric moisture, and runoff (Forster et  
1062 al., 2021, Douville et al., 2021; Gulev et al. 2021). AR6 Chapter 8 assessed that human-caused climate change has  
1063 driven detectable changes in the global water cycle since the mid-20th century with high confidence, including an  
1064 overall increase in atmospheric moisture (7% per 1 °C of warming), precipitation intensity (1-3% per 1 °C of warming)  
1065 and increased terrestrial evapotranspiration (Douville et al., 2021).

1066  
1067 In AR6, global land precipitation was highlighted as one of the large-scale indicators of climate change rather than  
1068 global precipitation since land precipitation has greater societal relevance and in situ precipitation records over land  
1069 extend back to the early to mid-20th century quasi-globally except Antarctica and parts of Africa and South America  
1070 (Gulev et al., 2021; Lee et al., 2021; Douville et al., 2021). AR6 assessed that global land precipitation has likely  
1071 increased since the middle of the 20th century with a faster increase since the 1980s with large interannual variability

and regional heterogeneity. The observed Northern Hemispheric land summer monsoon precipitation experienced a significant decline during 1901-2014, which has been attributed to the dominant influence of anthropogenic aerosols (Cao et al., 2022). Here, we include an update of global land precipitation change since AR6 (i.e., from 2020 to 2024).

Figure 12 shows annual global land precipitation anomaly relative to 1991-2020, following the current WMO climatology reference, obtained from GPCP V2020 (Schamm et al., 2014), CRU TS 4.08 (Harris et al., 2020), GPCP V.2.3 (Adler et al., 2018), and GHCN V4 (Menne et al., 2018) observed datasets. [The change of reference period to 1991-2020 affects the perceived evolution compared to Figure 2.15c in AR6 WGI which had a 1981-2010 reference period.](#) There is little consistency among datasets due to differences in input data, completeness of records, period of covered, and the gridding procedures applied (Sun et al., 2018; Nogueira, 2020). While the globally averaged land surface specific humidity has continuously increased (Dunn et al., 2024), global land precipitation has exhibited considerable interannual to interdecadal variability (Fig. 12). There was a positive anomaly in global land precipitation in 2024 but a negative anomaly in 2023. The former was contributed to by above-normal precipitation over the Asian and Australian monsoon region, likely associated with La Nina conditions, but was offset by dry conditions over South America and the southern part of Africa. The latter was driven by below normal precipitation over South Asia, Maritime Continents, the southern part of North America and the northern part of South America, due to El Niño conditions, with a corresponding increase in precipitation over the ocean (Adler and Gu, 2024).



**Figure 12** Time series of annual global land precipitation ( $\text{mm yr}^{-1}$ ) from 1891 to date relative to a 1991-2020 climatology obtained from GPCP V2020, CRU TS 4.08, GPCP V2.3, and GHCN V4 (note that different products commence at distinct times). Annual global land precipitation for each observed data is estimated following the AR6 method except the period of climatology and updated from 2020 to 2024. In AR6, the reference period of the climatology was from 1981 to 2010.

1096

1097 **12 Global mean sea-level rise**

1098 Global mean sea-level rise (GMSLR) is included in this annual update of AR6 for the first time. GMSLR is primarily  
1099 driven by: (i) thermal expansion as the ocean warms; and (ii) increases in ocean mass associated with the addition of  
1100 water or ice from land-based reservoirs, including glaciers and ice sheets (Fox-Kemper et al., 2021). Most of these  
1101 processes are directly linked to changes in the global Earth energy inventory (Sect. 6). Sea-level rise can have large  
1102 consequences for coastal ecosystems, safety and management, as it increases the baseline for sea-level extremes  
1103 arising from short-term phenomena such as storm surges, waves and tides.

1104

1105 Observed GMSLR was assessed in IPCC AR6 WG1, in Chapter 2 (their Section 2.3.3.3, Gulev et al., 2021) and  
1106 Chapter 9 (their Section 9.6.1 and Cross-Chapter Box 9.1, Fox-Kemper et al., 2021) on the basis of tide gauge  
1107 reconstructions (up to 1993) and satellite altimeter observations (1993-2018). The assessment of GMSLR from tide  
1108 gauge reconstructions used the ensemble approach presented by Palmer et al. (2021), which quantifies an ensemble  
1109 and its uncertainties by combining an estimate of the structural uncertainty (informed by the ensemble spread) with  
1110 an estimate of the internal uncertainty across the ensemble (i.e. the parametric uncertainty of each of the members in  
1111 the ensemble). The members included in the tide gauge ensemble, which informed the total sea-level change estimate  
1112 for the period 1901-1992, were reconstructions from Church and White (2011), Dangendorf et al (2019), Frederikse  
1113 et al. (2020) and Hay et al. (2015). For the satellite period, from 1993 to 2018, AR6 used the estimate of the WCRP  
1114 Global Sea Level Budget Group (2018), which was constructed from satellite-based GMSLR time series from six  
1115 groups (AVISO/CNES, CSIRO, NASA/GSFC, NOAA, SL\_cci/ESA and University of Colorado). Based on this  
1116 information, AR6 concluded that GMSLR increased by 0.20 [0.15 to 0.25] m over the period 1901 to 2018, with a  
1117 rate of 1.73 [1.28 to 2.17] mm yr<sup>-1</sup> (*high confidence*). Periods closer to the present showed an accelerating GMSLR,  
1118 with a rate of 2.3 [1.6 to 3.1] mm yr<sup>-1</sup> over the period 1971–2018 increasing to 3.7 [3.2 to 4.2] mm yr<sup>-1</sup> over the period  
1119 2006–2018 (*high confidence*).

1120

1121 Here, we extend the AR6 GMSLR time series, which ended in 2018, closer to the present day. We use the same tide  
1122 gauge-based ensemble estimate as in AR6 for the period up to 1993. We do note that two new reconstructions have  
1123 been published recently, both providing rates in line with the AR6 assessment rates given above. The new GMSLR  
1124 reconstruction by Dangendorf et al. (2024) uses a Kalman-smoother and adjusted estimates of the contributions of  
1125 glacial isostatic adjustment, barystatic and sterodynamic changes to sea-level change and finds a trend of  $1.50 \pm 0.20$   
1126 mm/yr for the period 1900-2021. The new reconstruction by Wang et al. (2024) uses an updated vertical land motion  
1127 correction and considers barystatic fingerprints and sterodynamic patterns from CMIP6 models and finds a trend of  
1128  $1.6 \pm 0.2$  mm/yr over 1900-2019.

1129

1130 The satellite record now provides observations up to the end of 2024, for three out of the six satellite data products  
1131 used for the WCRP estimate used in AR6. The three records available to the end of 2024 are from NASA (2025),  
1132 NOAA (2025) and AVISO (2025). All data was downloaded on 19 February 2025. We use the global mean time series  
1133 based on the reference missions, with seasonal signals removed and corrected for glacial isostatic adjustment. We first  
1134 compute annual averages and then an ensemble average time series, which is spliced to the AR6 GMSLR record  
1135 ending in 2018. For consistency, we retain the uncertainties from the six-member WCRP ensemble and propagate  
1136 them over the period 2019-2024. We note that reprocessing of the altimetry record is periodically required to account  
1137 for new insights on instrument drift, retracking and geophysical corrections to the altimetry missions. This  
1138 reprocessing may lead to small differences in the satellite altimeter record and the associated assessment of GMSLR  
1139 in future iterations of IGCC.

1140  
1141 Over the period 2019 to 2024 global mean sea level has increased by 26.1 [19.8 to 32.4] mm. When combining the  
1142 AR6 estimate up to 2018 with the satellite time series for 2019-2024, we find a total GMSLR of 228.0 [176.4 to 279.6]  
1143 mm for the period 1901-2024, which translates to an average rate of 1.85 [1.43 to 2.27] mm yr<sup>-1</sup> (Table 10, Fig. 13).  
1144 The rate increase associated with extending the time series by just 6 years, as well as the increasing rates over  
1145 consecutive 20-yr periods (Fig. 13b), indicate a continuing acceleration of GMSLR. This is in line with the  
1146 assessments of AR6 (Fox-Kemper et al., 2021), SROCC (Oppenheimer et al., 2019) and AR5 (Church et al., 2013)  
1147 that sea-level change has been accelerating over the course of the 20th and early 21st centuries, and consistent with  
1148 the observed acceleration in some components of the Earth heat inventory (see Sect. 6).

1149  
1150 **Table 10** Observed global mean sea-level rise (GMSLR) as presented in IPCC AR6, table 9.5 (Fox-Kemper et al., 2021)  
1151 compared with the extended time series in this study. Values are expressed as the total change ( $\Delta$ ) in the annual mean  
1152 over each period (mm) along with the equivalent rate calculated as the total change divided by the number of years (mm  
1153 yr<sup>-1</sup>). Uncertainties represent the *very likely* range.

Observed GMSLR		IPCC AR6	This study
Start year		End year 2018	End year 2024
1901	$\Delta$ (mm)	201.9 [150.3 to 253.5]	228.0 [176.4 to 279.6]
	mm yr <sup>-1</sup>	1.73 [1.28 to 2.17]	1.85 [1.43 to 2.27]
1971	$\Delta$ (mm)	109.6 [72.8 to 146.4]	135.8 [99.0 to 172.5]
	mm yr <sup>-1</sup>	2.33 [1.55 to 3.12]	2.56 [1.87 to 3.26]
1993	$\Delta$ (mm)	81.2 [72.1 to 90.2]	107.3 [98.2 to 116.4]
	mm yr <sup>-1</sup>	3.25 [2.88 to 3.61]	3.46 [3.17 to 3.75]
2006	$\Delta$ (mm)	44.3 [38.6 to 50.0]	70.4 [64.7 to 76.1]

Deleted: 7

Deleted: 2

Deleted: 7

Deleted: 2

	mm yr <sup>-1</sup>	3.69 [3.21 to 4.17]	3.91 [3.59 to 4.23]
--	---------------------	---------------------	---------------------




Figure 13 (a) Global mean sea-level rise time series 1901-2024 (mm). The GMSLR ensemble from AR6 in black, w.r.t. the period 1995-2014; the updated satellite altimetry ensemble in red, w.r.t. the AR6 ensemble in 2018. Individual time series are shown in dashed lines. (b) GMSLR rates (mm yr<sup>-1</sup>) for different periods. Uncertainties in a) show the likely range and in b) the very likely range, computed relative to 1901, including estimates of both structural uncertainty and parametric uncertainty (Palmer et al., 2021).

### 13 Code, data availability and visualisations

We publish a set of selected key indicators of global climate change via Climate Change Tracker (<https://climatechangetracker.org/>, Climate Change Tracker, 2025), a platform which aims to provide reliable, user-friendly, high-quality interactive dashboards, visualisations, data, and easily accessible insights of this paper.

With Climate Change Tracker we aim to reach a wider public audience, including policymakers involved in UNFCCC negotiations, and decision makers working in climate change mitigation and adaptation. Climate Change Tracker plans to update significant indicators multiple times throughout the year, providing an up-to-date picture of the indicators of climate change. Within the dashboards, all data is traceable to the underlying sources. [Fig. 14 presents a screenshot of the IGCC dashboard.](#)




Figure 14 Screenshot of the IGCC dashboard, (<https://climatechangetracker.org/>, Climate Change Tracker, 2025).

Formatted: No underline, Font colour: Auto

Formatted: Heading 4

The carbon budget calculation is available from <https://github.com/Rlamboll/AR6CarbonBudgetCalc/tree/v1.0.3> (Lamboll and Rogelj, 2025). The code and data used to produce other indicators are available in repositories under <https://github.com/ClimateIndicator/data/tree/v2025.06.11> (Smith et al., 2025b). All data are available from <https://doi.org/10.5281/zenodo.15639576> (Smith et al., 2025a). Data are provided under the CC-BY 4.0 License.

Deleted: <https://github.com/ClimateIndicator/data/releases/tag/v2025.04.30b>

Deleted: 10.5281/zenodo.15327155

HadEX3 [3.0.4] data were obtained from <https://catalogue.ceda.ac.uk/uuid/115d5e4ebf7148ec941423ec86fa9f26> (Dunn et al., 2023) on 5 April 2023 and are © British Crown Copyright, Met Office, 2022, provided under an Open Government Licence; <http://www.nationalarchives.gov.uk/doc/open-government-licence/version/2/> (last access: 2 June 2023).

#### 14 Discussion and conclusions

The third year of the Indicators of Global Climate Change (IGCC) initiative has built on previous years' efforts to provide a comprehensive update of the climate change indicators required to estimate the human-induced warming

and the remaining carbon budget. Table 11 and Fig. 15 present a summary of the headline indicators from each section compared to those given in the AR6 assessment. Table 11 also summarises methodological updates.

**Table 11 Summary of headline results and methodological updates from the Indicators of Global Climate Change (IGCC) initiative.**

Climate Indicator	AR6 2021 assessment	This 2024 assessment	Explanation of changes	Methodological updates since AR6
GHG emissions AR6 WGIII Chapter 2: Dhakal et al. (2022); see also Minx et al. (2021)	2010-2019 average: 55.9 ± 6 GtCO <sub>2</sub> e	2010-2019 average: 52.9 ± 5.4 GtCO <sub>2</sub> e 2014-2023 average: 53.6 ± 5.2 GtCO <sub>2</sub> e	Average emissions in the past decade grew at a slower rate than in the previous decade. The change from AR6 is due to a systematic downward revision in CO <sub>2</sub> -LULUCF and CH <sub>4</sub> estimates. Real-world emissions have slightly increased.	CO <sub>2</sub> -LULUCF emissions revised down. CO <sub>2</sub> GCB Fossil Fuel and Industry emissions used instead of EDGAR. PRIMAP-hist TP used in place of EDGAR for CH <sub>4</sub> and N <sub>2</sub> O emissions, atmospheric measurements taken for F-gas emissions. These changes reduce estimates by around 3 GtCO <sub>2</sub> e (Sect. 2).
GHG concentrations AR6 WGI Chapter 2: Gulev et al. (2021)	2019: CO <sub>2</sub> , 410.1 [± 0.36] ppm CH <sub>4</sub> , 1866.3 [± 3.2] ppb N <sub>2</sub> O, 332.1 [± 0.7] ppb	2024: CO <sub>2</sub> , 422.8 [±0.4] ppm CH <sub>4</sub> , 1929.7 [±3.3] ppb N <sub>2</sub> O, 337.9 [±0.4] ppb	Increases caused by continued GHG anthropogenic emissions	Updates based on NOAA data and AGAGE (Sect. 3)
Effective radiative forcing change since 1750 AR6 WGI Chapter 7: Forster et al. (2021)	2019: 2.72 [1.96 to 3.48] W m <sup>-2</sup>	2024: 2.97 [2.05 to 3.77] W m <sup>-2</sup>	Trend since 2019 is caused by increases in GHG concentrations and reductions in aerosol precursors.	Follows AR6 with minor update to aerosol precursor treatment and emissions dataset that revises 2019 ERF estimate relative to 1750 downwards (more negative) by 0.09 W m <sup>-2</sup> . Added this year is a new method to estimate the ERF from land use surface reflection and irrigation to avoid scaling with cumulative emissions. This

Deleted: 4

Deleted: 8

Deleted: 3

Deleted: 6

				does not materially affect the ERF. (Sect. 5)
Earth's energy imbalance  AR6 WGI Chapter 7: Forster et al. (2021)	2006-2018 average:  0.79 [0.52 to 1.06] W m <sup>-2</sup>	2012-2024 average:  0.99 [0.70 to 1.28] W m <sup>-2</sup>	A 25% increase in energy imbalance estimated based on increased rate of ocean heating.	Ocean heat content timeseries extended from 2018 to 2024 using all of the 5 AR6 datasets. Other heat inventory terms updated following von Schuckmann et al. (2023a). Ocean heat content uncertainty is used as a proxy for total uncertainty. Further details in Sect. 6.
Global mean surface temperature change since 1850-1900  AR6 WGI Chapter 2: Gulev et al. (2021)	2011-2020 average:  1.09 [0.95 to 1.20] °C	2015-2024 average:  1.24 [1.11 to 1.35] °C	An increase of 0.15 °C within four years, indicating a high decadal rate of change which may in part be internal variability.	Methods match four datasets used in AR6. Individual datasets have updated historical data, but these changes are not materially affecting results. (Sect. 7).
Human induced global warming since preindustrial  AR6 WGI Chapter 3: Eyring et al. (2021)  SR1.5 Chapter 1	2010-2019 decade average:  1.07 [0.8 to 1.3] °C  2017 single year: 1.0 [0.8 to 1.2] °C	2015-2024 decade average:  1.22 [1.0 to 1.5] °C  2024 single year: 1.36 [1.1 to 1.7] °C	An increase of 0.15 °C within five years, indicating a high decadal rate of change (broadly consistent with warming projections). The decadal warming rate increased slightly between 2019 and 2024. One of the three AR6 methods is diverging.	The three methods for the basis of the AR6 assessment are retained, but each has new input data (Sect. 8)

Deleted: 4


Deleted: 3



Remaining carbon budget for 50% likelihood of limiting global warming to 1.5 °C  AR6 WGI Chapter 5: Canadell et al. (2021)	From the start of 2020:  500 GtCO <sub>2</sub>	From the start of 2025:  130 GtCO <sub>2</sub>	The 1.5 °C budget is becoming very small. The RCB can exhaust before the 1.5 °C threshold is reached due to having to allow for future non-CO <sub>2</sub> warming.	Emulator and scenario change has reduced budget since 2020 by 100 GtCO <sub>2</sub> (Sect. 9)
Land average maximum temperature change compared to pre-industrial.  AR6 WGI Chapter 11: Seneviratne et al., 2021	2009-2018 average:  1.55 °C	2015-2024 average:  1.90 °C	Rising at a substantially faster rate compared to global mean surface temperature	HadEX3 data used in AR6 replaced with ERA reanalysis data employed in this report which is more updatable going forward. Adds 0.01 °C to estimate (Sect. 10)
Global land precipitation compared to preindustrial (Douville et al., 2021)	Likely increased since the middle of the 20th century with a faster increase since the 1980s with large interannual variability	Large interannual variability associated with El Niño dominates the record in recent years, making long-term trend less clear	2023 exhibited a negative anomaly relative to preindustrial due to El Niño conditions	The four datasets used in AR6 have been extended (Sect. 11)
Global mean sea-level rise since 1901  (Gulev et al., 2021; Fox-Kemper et al., 2021)	1901 to 2018 change  201.9 [150.3 to 253.5] mm  at a rate of  1.73 [1.28 to 2.17] mm yr <sup>-1</sup>	1901 to 2024 change  228.0 [176.4 to 279.6] mm  at a rate of  1.85 [1.43 to 2.27] mm yr <sup>-1</sup>	Sea-level rise continues to accelerate.	AR6 data extended with three of the six datasets from AR6, using latest satellite data (Sect. 12).

Deleted: 7

Deleted: 2



210

211 Figure 15. Infographic for the best estimate of headline indicators assessed in this paper.

Deleted: 4

212 Last year (2024) witnessed global surface temperatures more likely than not exceeding 1.5 °C above preindustrial  
213 levels which has widely been reported in the press. Sects. 7 and 8. show that such high levels of global temperature  
214 anomalies are typical of what we expect from current best estimates of human induced warming, modulated by internal  
215 climate variability.

1216

1218 The overview of key indicators of the state of global climate indeed highlights the multiple fingerprints of the 2023-  
1219 2024 El-Nino event regarding peak global surface temperature (Section 7.2), regional dry anomalies in land  
1220 precipitation (Section 11), and their implications for reduced land carbon sinks and the record growth rate of  
1221 atmospheric CO<sub>2</sub> concentrations in 2024 (Section 3).

1222 The overall increase in land maximum temperatures (Section 10), closely related to global warming levels, drives  
1223 increasing trends in potential evapotranspiration, decreasing trends in soil moisture (Seo et al., 2025), contributing to  
1224 the increased rate of global mean sea-level rise (Section 12).

1225 Methane and biomass emissions had a strong component of change related to climate feedbacks (Sects. 2 and 3). Such  
1226 changes will become increasingly important over this century, even if the direct human influence declines. This year,  
1227 we explored different inventory choices in Sect. 2. In future years a more consistent approach to attribution of  
1228 atmospheric emissions, concentration change and radiative forcing should be developed, so it can be assessed in AR7.

1229 It is hoped that this update can support the science community in its collection and provision of reliable and timely  
1230 global climate data. In future years we are particularly interested in improving SLCF updating methods to get a more  
1231 accurate estimate of short-term ERF changes. The work also highlights the importance of high-quality metadata to  
1232 document changes in methodological approaches over time. This year we have extended the datasets with land  
1233 precipitation and global mean sea\_level rise. In future years we hope to improve the robustness of the indicators  
1234 presented here and could update other AR6 assessments. Parallel efforts could explore how we might update indicators  
1235 of regional climate extremes and their attribution, which are particularly relevant for supporting actions on adaptation  
1236 and loss and damage.

1237

1238 Generally, scientists and scientific organisations have an important role as “watchdogs” to critically inform evidence-  
1239 based decision-making. This annual update traced to IPCC methods can provide a reliable, timely source of  
1240 trustworthy information, [IGCC and the complimentary updates of the State of the Climate \(BAMS\) and State of Global](#)  
1241 [Climate \(WMO\) reports](#), very much rely on continued support for high quality, global monitoring networks of  
1242 atmospheric and climate data, and also on open data sources that are regularly updated and easily accessed.

1243

1244 This is a critical decade: human-induced global warming rates are at their highest historical level, and 1.5 °C global  
1245 warming might be expected to be reached or exceeded in around 5 years in the absence of cooling from major volcanic  
1246 eruptions (Sects. 8 and 9). Yet this is also the decade when global GHG emissions could be expected to peak and  
1247 begin to substantially decline. The indicators of global climate change presented here show that the Earth's energy  
1248 imbalance has increased to around 1.0 W m<sup>-2</sup>, averaged over the last 12 years (Sect. 5), which represents a 25%  
1249 increase on the value assessed for 2006-2018 by AR6. This also has implications for the committed response of slow  
1250 components in the climate system (glaciers, deep ocean, ice sheets) and committed long-term sea-level rise (through  
1251 ocean thermal expansion and land-based ice melt/loss), to be addressed further in future updates. However, rapid and

Deleted: .  
Deleted: It  
Deleted: ies

1255 stringent GHG emission decreases such as those committed to at COP28 could halve warming rates over the next 20  
1256 years (McKenna et al., 2021). Table 1 shows that global GHG emissions are at a long-term high, yet there are signs  
1257 that their rate of increase has slowed. Depending on the societal choices made in this critical decade, a continued series  
1258 of these annual updates could track an improving trend for some of the indicators herein discussed.

1259 **Supplement**

1260 The supplement related to this article is available online.

1261 **Author contributions**

1262 PMF, CS, MA, PF, JR and AP developed the concept of an annual update in discussions with the wider IPCC  
1263 community over many years. CS led the work of the data repositories. VMD, PZ, SS, CS, SIS, VN, AP, NPG, GPP,  
1264 BT, MDP, KvS, JR, PF, MA, JCM, XZ, RAB, CB, CC, SB and PT provided important IPCC and UNFCCC framing.  
1265 PMF coordinated the production of the manuscript with support from DR. WFL led Sect. 2 with contributions from  
1266 PF, GPP, JG, JP, JCM and RA. CS led Sect. 3 with inputs from JM, PK, LW, PMF and MR. SS led Sect. 4 with inputs  
1267 from VK, CS, GvdW, LAR and MG. CS led Sect. 5 with contributions from CW, TG, SS, VN and GvdW. KvS and  
1268 MDP led Sect. 6 with contributions from LC, MI, JR, REK, AS, CMD, DPM and SEW. BT, CC and ZH led Sect. 7  
1269 with contributions from PT, CM, CK, JK, RR, RV, AL and LC. TW led Sect. 8 with contributions and calculations  
1270 from AR, NPG, SJ, CS and MA. RL led Sect. 9 with contributions from JR and HDM. Sect. 10 was led by MH, with  
1271 contributions from SIS, and XZ. JYL, JEY, and RK led Sect. 11 with contributions from VMD, PT, and KvS. AS led  
1272 Sect. 12 with contributions from MDP. All authors either edited or commented on the manuscript. DR, AB and JAB  
1273 coordinated the data visualisation effort.

1274 **Competing interests**

1275 The contact author has declared that none of the authors has any competing interests.

1276 **Disclaimer**

1277 Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and  
1278 institutional affiliations.

1279 **Acknowledgements**

1280 This research has been supported by the European Union's Horizon Europe research and innovation programme under  
1281 Grant Agreement Nos. 820829, 101081395, 101081661 and 821003, the H2020 European Research Council (grant  
1282 no. 951542), the Natural Environment Research Council (NE/X00452X/1) and the Engineering and Physical Research  
1283 Council (EP/V000772/1). Matthew Palmer, Colin Morice, Rachel Killick and Richard Betts were supported by the  
1284 Met Office Hadley Centre Climate Programme funded by DSIT. Peter Thorne was supported by Co-Centre award

number 22/CC/11103. The Co-Centre award is managed by Research Ireland Northern Ireland's Department of Agriculture, Environment and Rural Affairs (DAERA) and UK Research and Innovation (UKRI), and supported via UK's International Science Partnerships Fund (ISPF), and the Irish Government's Shared Island initiative. Analyses and visualizations for concentrations of Short Lived Climate Forcers used in this paper were produced with the Giovanni online data system, developed and maintained by the NASA GES DISC (as available in February 2025). June-Yi Lee and Jung-Eun Yun were supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (No. RS-2024-00416848). Aimée Slangen was supported by the research programme ENW-Vidi (DARSea, project number VI.Vidi.2023.058) funded by the Dutch Research Council (NWO). We thank Xin Lan for assistance with compiling the GHG concentration data.

**References**

Acker, J. G. and Leptoukh, G.: Online analysis enhances use of NASA Earth science data, *EoS Transactions*, 88, 14–17, <https://doi.org/10.1029/2007EO020003>, 2007.

Adler, R. F. and Gu, G.: Global precipitation for the year 2023 and how it relates to longer term variations and trends, *Atmosphere*, 15, 535, 2024.

Adler, R. F., Sapiiano, M. R. P., Huffman, J., Wang, J.-J., Gu G., Bolvin, D., Chiu, L., Schneider, U., Becker, A., Nelkin, E., Xie, P., Ferrarok R., and Shin, D.-B.: The global precipitation climatology project (GPCP) monthly analysis (new version 2.3) and a review of 2017 global precipitation. *Atmosphere*, 9, 138, <https://doi.org/10.3390/atmos9040138>, 2018.

Allan ,R.P. and Merchant, C.J.: Reconciling Earth's growing energy imbalance with ocean warming. *Environ. Res. Lett*, 20 04402, <https://doi.org/10.1088/1748-9326/adb448>, 2025.

Allen, M. R., O. P. Dube, W. Solecki, F. Aragón-Durand, W. Cramer, S. Humphreys, M. Kainuma, J. Kala, N. Mahowald, Y. Mulugetta, R. Perez, M. Wairiu, and K. Zickfeld, 2018: Framing and Context. In: *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty* [Masson-Delmotte, V., P. Zhai, H.-O. Pörtner, D. Roberts, J. Skea, P.R. Shukla, A. Pirani, W. Moufouma-Okia, C. Péan, R. Pidcock, S. Connors, J.B.R. Matthews, Y. Chen, X. Zhou, M.I. Gomis, E. Lonnoy, T. Maycock, M. Tignor, and T. Waterfield (eds.)], Cambridge University Press, Cambridge, UK and New York, NY, USA, 49-92, <https://doi.org/10.1017/9781009157940.003>, 2018.

Allen, M. R., Frame, D. J., Friedlingstein, P., Gillett, N. P., Grassi, G., Gregory, J. M., Hare, W., House, J., Huntingford, C., Jenkins, S., Jones, C. D., Knutti, R., Lowe, J. A., Matthews, H. D., Meinshausen, M., Meinshausen, N., Peters, G. P., Plattner, G.-K., Raper, S., Rogelj, J., Stott, P. A., Solomon, S., Stocker, T. F., Weaver, A. J., and Zickfeld, K.: Geological Net Zero and the need for disaggregated accounting for carbon sinks, *Nature*, 638, 343–350, <https://doi.org/10.1038/s41586-024-08326-8>, 2025.

1318 Allison, L. C., Palmer, M. D., Allan, R. P., Hermanson, L., Liu, C., and Smith, D. M.: Observations of planetary  
 1319 heating since the 1980s from multiple independent datasets, *Environ. Res. Commun.*, 2, 101001,  
 1320 <https://doi.org/10.1088/2515-7620/abbb39>, 2020.

1321 AVISO: Mean sea-level product, [https://www.aviso.altimetry.fr/en/data/products/ocean-indicators-products/mean-](https://www.aviso.altimetry.fr/en/data/products/ocean-indicators-products/mean-sea-level/data-access.html)  
 1322 [sea-level/data-access.html](https://www.aviso.altimetry.fr/en/data/products/ocean-indicators-products/mean-sea-level/data-access.html), [data set], accessed 19 February 2025, 2025.

1323 Barnes, C., Boulanger, Y., Keeping, T., Gachon, P., Gillett, N., Haas, O., Wang, X., Roberge, F., Kew, S., Heinrich,  
 1324 D., Singh, R., Vahlberg, M., Van Aalst, M., Otto, F., Kimutai, J., Boucher, J., Kasoar, M., Zachariah, M., and Krikken,  
 1325 F.: Climate change more than doubled the likelihood of extreme fire weather conditions in Eastern Canada, Imperial  
 1326 College London, <https://doi.org/10.25561/105981>, 2023.

1327 Basu, S., Lan, X., Dlugokencky, E., Michel, S., Schwietzke, S., Miller, J. B., Bruhwiler, L., Oh, Y., Tans, P. P.,  
 1328 Apadula, F., Gatti, L. V., Jordan, A., Necki, J., Sasakawa, M., Morimoto, S., Di Iorio, T., Lee, H., Arduini, J., and  
 1329 Manca, G.: Estimating emissions of methane consistent with atmospheric measurements of methane and  $\delta^{13}\text{C}$  of  
 1330 methane, *Atmos. Chem. Phys.*, 22, 15351–15377, <https://doi.org/10.5194/acp-22-15351-2022>, 2022.

1331 Bellouin, N., Davies, W., Shine, K. P., Quaas, J., Mülmenstädt, J., Forster, P. M., Smith, C., Lee, L., Regayre, L.,  
 1332 Brasseur, G., Sudarchikova, N., Bouarar, I., Boucher, O., and Myhre, G.: Radiative forcing of climate change from  
 1333 the Copernicus reanalysis of atmospheric composition, *Earth Syst. Sci. Data*, 12, 1649–1677,  
 1334 <https://doi.org/10.5194/essd-12-1649-2020>, 2020.

1335 Betts, R. A., Belcher, S. E., Hermanson, L., Klein Tank, A., Lowe, J. A., Jones, C. D., Morice, C. P., Rayner, N. A.,  
 1336 Scaife, A. A., and Stott, P. A.: Approaching 1.5 °C: how will we know we’ve reached this crucial warming mark?,  
 1337 *Nature*, 624, 33–35, <https://doi.org/10.1038/d41586-023-03775-z>, 2023.

1338 Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S.,  
 1339 Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman,  
 1340 C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z.,  
 1341 Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black  
 1342 carbon in the climate system: A scientific assessment, *J. Geophys. Res.-Atmos.*, 118, 5380–5552,  
 1343 <https://doi.org/10.1002/jgrd.50171>, 2013.

1344 Bun, R., Marland, G., Oda, T., See, L., Puliafito, E., Nahorski, Z., Jonas, M., Kovalyshyn, V., Ialongo, I., Yashchun,  
 1345 O., and Romanchuk, Z.: Tracking unaccounted greenhouse gas emissions due to the war in Ukraine since 2022,  
 1346 *Science of The Total Environment*, 914, 169879, <https://doi.org/10.1016/j.scitotenv.2024.169879>, 2024.

1347 Burton, C., Lampe, S., Kelley, D. I., Thiery, W., Hantson, S., Christidis, N., Gudmundsson, L., Forrest, M., Burke,  
 1348 E., Chang, J., Huang, H., Ito, A., Kou-Giesbrecht, S., Lasslop, G., Li, W., Nieradzik, L., Li, F., Chen, Y., Randerson,  
 1349 J., Reyer, C. P. O., and Mengel, M.: Global burned area increasingly explained by climate change, *Nat. Clim. Chang.*,  
 1350 14, 1186–1192, <https://doi.org/10.1038/s41558-024-02140-w>, 2024.

1351 Canadell, J. G., P. M. S. Monteiro, M. H. Costa, L. Cotrim da Cunha, P. M. Cox, A. V. Eliseev, S. Henson, M. Ishii, S.  
 1352 Jaccard, C. Koven, A. Lohila, P. K. Patra, S. Piao, J. Rogelj, S. Syampungani, S. Zaehle, and K. Zickfeld: Global

Carbon and other Biogeochemical Cycles and Feedbacks. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 673–816, <https://doi.org/10.1017/9781009157896.007>, 2021.

Cao, J., Wang, H., Wang, B., Zhao, H., Wang, C., and Zhu, X.: Higher sensitivity of Northern Hemisphere monsoon to anthropogenic aerosol than greenhouse gases, *Geophys. Res. Lett.*, 49, e2022GL100270, <https://doi.org/10.1029/2022GL100270>, 2022.

Cattiaux, J., Ribes, A., and Cariou, E.: How Extreme Were Daily Global Temperatures in 2023 and Early 2024?, *Geophysical Research Letters*, 51, e2024GL110531, <https://doi.org/10.1029/2024GL110531>, 2024.

Cheng, L., Abraham, J., Hausfather, Z., and Trenberth, K. E.: How fast are the oceans warming?, *Science*, 363, 128–129, <https://doi.org/10.1126/science.aav7619>, 2019.

Cheng, L., Von Schuckmann, K., Abraham, J. P., Trenberth, K. E., Mann, M. E., Zanna, L., England, M. H., Zika, J. D., Fasullo, J. T., Yu, Y., Pan, Y., Zhu, J., Newsom, E. R., Bronselaer, B., and Lin, X.: Past and future ocean warming, *Nat. Rev. Earth. Environ.*, 3, 776–794, <https://doi.org/10.1038/s43017-022-00345-1>, 2022.

Church, J. A., White, N. J., Konikow, L. F., Domingues, C. M., Cogley, J. G., Rignot, E., Gregory, J. M., Van Den Broeke, M. R., Monaghan, A. J., and Velicogna, I.: Revisiting the Earth’s sea-level and energy budgets from 1961 to 2008: SEA-LEVEL AND ENERGY BUDGETS, *Geophys. Res. Lett.*, 38, n/a-n/a, <https://doi.org/10.1029/2011GL048794>, 2011.

Climate Change Tracker, <https://climatechangetracker.org/igcc>, accessed 20.05.2025, 2025.

Collins, M., Knutti, R., Arblaster, J., Dufresne, J.-L., Fichefet, T., Friedlingstein, P., Gao, X., Gutowski, W.J., Johns, T., Krinner, G., Shongwe, M., Tebaldi, C., Weaver, A.J. & Wehner, M.: Long-term Climate Change: Projections, Commitments and Irreversibility. In: V.B. Stocker T.F., .D. Qin, G.K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia & P.M. Midgley (eds.). *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, United Kingdom and New York, NY, USA, Cambridge University Press. pp. 1029–1136, 2013.

Crippa, M., Guizzardi, D., Schaaf, E., Monforti-Ferrario, F., Quadrelli, R., Riskez Martin, A., Rossi, S., Vignati, E., Muntean, M., Brandao De Melo, J., Oom, D., Pagani, F., Banja, M., Taghavi-Moharamli, P., Köykkä, J., Grassi, G., Branco, A., and San-Miguel, J.: GHG emissions of all world countries – 2023, Publications Office of the European Union, <https://doi.org/doi/10.2760/953322>, 2023.

Cuesta-Valero, F. J., Beltrami, H., García-García, A., Krinner, G., Langer, M., MacDougall, A., Nitzbon, J., Peng, J., von Schuckmann, K., Seneviratne, S., Thiery, W., Vanderkelen, I., Wu, T.: GCOS EHI 1960-2020 Continental Heat Content (Version 2), World Data Center for Climate (WDCC) at DKRZ, [https://doi.org/10.26050/WDCC/GCOS\\_EHI\\_1960-2020\\_CoHC\\_v2](https://doi.org/10.26050/WDCC/GCOS_EHI_1960-2020_CoHC_v2), 2023.

1388 Dangendorf, S., Hay, C., Calafat, F. M., Marcos, M., Piecuch, C. G., Berk, K., and Jensen, J.: Persistent acceleration  
 1389 in global sea-level rise since the 1960s, *Nat. Clim. Chang.*, 9, 705–710, <https://doi.org/10.1038/s41558-019-0531-8>,  
 1390 2019.

1391 Dangendorf, S., Sun, Q., Wahl, T., Thompson, P., Mitrovica, J. X., and Hamlington, B.: Probabilistic reconstruction  
 1392 of sea-level changes and their causes since 1900, *Earth Syst. Sci. Data*, 16, 3471–3494, [https://doi.org/10.5194/essd-](https://doi.org/10.5194/essd-16-3471-2024)  
 1393 [16-3471-2024](https://doi.org/10.5194/essd-16-3471-2024), 2024.

1394 Deng, Z., Ciais, P., Tzompa-Sosa, Z. A., Saunio, M., Qiu, C., Tan, C., Sun, T., Ke, P., Cui, Y., Tanaka, K., Lin, X.,  
 1395 Thompson, R. L., Tian, H., Yao, Y., Huang, Y., Lauerwald, R., Jain, A. K., Xu, X., Bastos, A., Sitch, S., Palmer, P.  
 1396 I., Lauvaux, T., d’Aspremont, A., Giron, C., Benoit, A., Poulter, B., Chang, J., Petrescu, A. M. R., Davis, S. J., Liu,  
 1397 Z., Grassi, G., Albergel, C., Tubiello, F. N., Perugini, L., Peters, W., and Chevallier, F.: Comparing national  
 1398 greenhouse gas budgets reported in UNFCCC inventories against atmospheric inversions, *Earth System Science Data*,  
 1399 14, 1639–1675, <https://doi.org/10.5194/essd-14-1639-2022>, 2022.

1400 Deng, Z., Zhu, B., Davis, S. J., Ciais, P., Guan, D., Gong, P., and Liu, Z.: Global carbon emissions and decarbonization  
 1401 in 2024, *Nat Rev Earth Environ*, 6, 231–233, <https://doi.org/10.1038/s43017-025-00658-x>, 2025.

1402 Denier van der Gon, H., Gauss, M., Granier, C., Arellano, S., Benedictow, A., Darras, S., Dellaert, S., Guevara, M.,  
 1403 Jalkanen, J.-P., Krueger, K., Kuenen, J., Liaskoni, M., Liousse, C., Markova, J., Prieto Perez, A., Quack, B., Simpson,  
 1404 D., Sindelarova, K., and Soulie, A.: Documentation of CAMS emission inventory products,  
 1405 <https://doi.org/10.24380/Q2SI-TI6I>, 2023.

1406 Dhakal, S., J. C. Minx, F. L. Toth, A. Abdel-Aziz, M. J. Figueroa Meza, K. Hubacek, I. G. C. Jonckheere, Yong-Gun  
 1407 Kim, G. F. Nemet, S. Pachauri, X. C. Tan, T. Wiedmann: Emissions Trends and Drivers. In IPCC, 2022: Climate  
 1408 Change 2022: Mitigation of Climate Change. Contribution of Working Group III to the Sixth Assessment Report of  
 1409 the Intergovernmental Panel on Climate Change [P.R. Shukla, J. Skea, R. Slade, A. Al Khourdajie, R. van Diemen,  
 1410 D. McCollum, M. Pathak, S. Some, P. Vyas, R. Fradera, M. Belkacemi, A. Hasija, G. Lisboa, S. Luz, J. Malley,  
 1411 (eds.)]. Cambridge University Press, Cambridge, UK and New York, NY, USA,  
 1412 <https://doi.org/10.1017/9781009157926.004>, 2022.

1413 Douville, H., K. Raghavan, J. Renwick, R.P. Allan, P.A. Arias, M. Barlow, R. Cerezo-Mota, A. Cherchi, T.Y. Gan, J.  
 1414 Gergis, D. Jiang, A. Khan, W. Pokam Mba, D. Rosenfeld, J. Tierney, and O. Zolina: Water Cycle Changes. In Climate  
 1415 Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the  
 1416 Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S.  
 1417 Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K.  
 1418 Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United  
 1419 Kingdom and New York, NY, USA, pp. 1055–1210, <https://doi.org/10.1017/9781009157896.010>, 2021.

1420 Droste, E. S., Adcock, K. E., Ashfold, M. J., Chou, C., Fleming, Z., Fraser, P. J., Gooch, L. J., Hind, A. J., Langenfelds,  
 1421 R. L., Leedham Elvidge, E. C., Mohd Hanif, N., O’Doherty, S., Oram, D. E., Ou-Yang, C.-F., Panagi, M., Reeves, C.  
 1422 E., Sturges, W. T., and Laube, J. C.: Trends and emissions of six perfluorocarbons in the Northern Hemisphere and  
 1423 Southern Hemisphere, *Atmos. Chem. Phys.*, 20, 4787–4807, <https://doi.org/10.5194/acp-20-4787-2020>, 2020.



1424 Dunn, R. J. H., Alexander, L. V., Donat, M. G., Zhang, X., Bador, M., Herold, N., Lippmann, T., Allan, R., Aguilar,  
1425 E., Barry, A. A., Brunet, M., Caesar, J., Chagnaud, G., Cheng, V., Cinco, T., Durre, I., Guzman, R., Htay, T. M., Wan  
1426 Ibadullah, W. M., Bin Ibrahim, M. K. I., Khoshkam, M., Kruger, A., Kubota, H., Leng, T. W., Lim, G., Li-Sha, L.,  
1427 Marengo, J., Mbatha, S., McGree, S., Menne, M., Milagros Skansi, M., Ngwenya, S., Nkrumah, F., Oonariya, C.,  
1428 Pabon-Caicedo, J. D., Panthou, G., Pham, C., Rahimzadeh, F., Ramos, A., Salgado, E., Salinger, J., Sané, Y.,  
1429 Sopaheluwakan, A., Srivastava, A., Sun, Y., Timbal, B., Trachow, N., Trewin, B., Schrier, G., Vazquez-Aguirre, J.,  
1430 Vasquez, R., Villarroel, C., Vincent, L., Vischel, T., Vose, R., and Bin Hj Yussof, M. N.: Development of an updated  
1431 global land in situ-based data set of temperature and precipitation extremes: HadEX3, *J. Geophys. Res.-Atmos.*, 125,  
1432 e2019JD032263, <https://doi.org/10.1029/2019JD032263>, 2020.

1433 Dunn, R. J. H., Donat, M. G., and Alexander, L. V.: Comparing extremes indices in recent observational and reanalysis  
1434 products, *Front. Clim.*, 4, 98905, <https://doi.org/10.3389/fclim.2022.989505>, 2022.

1435 Dunn, R.J.H., Alexander, L., Donat, M., Zhang, X., Bador, M., Herold, N., Lippmann, T., Allan, R.J., Aguilar, E.,  
1436 Aziz, A., Brunet, M., Caesar, J., Chagnaud, G., Cheng, V., Cinco, T., Durre, I., de Guzman, R., Htay, T.M., Wan  
1437 Ibadullah, W.M., Bin Ibrahim, M.K.I., Khoshkam, M., Kruge, A., Kubota, H., Leng, T.W., Lim, G., Li-Sha, L.,  
1438 Marengo, J., Mbatha, S., McGree, S., Menne, M., de los Milagros Skansi, M., Ngwenya, S., Nkrumah, F., Oonariya,  
1439 C., Pabon-Caicedo, J.D., Panthou, G., Pham, C., Rahimzadeh, F., Ramos, A., Salgado, E., Salinger, J., Sane, Y.,  
1440 Sopaheluwakan, A., Srivastava, A., Sun, Y., Trimbal, B., Trachow, N., Trewin, B., van der Schrier, G., Vazquez-  
1441 Aguirre, J., Vasquez, R., Villarroel, C., Vincent, L., Vischel, T., Vose, R., Bin Hj Yussof, and M.N.A.: HadEX3:  
1442 Global land-surface climate extremes indices v3.0.4 (1901-2018), NERC EDS Centre for Environmental Data  
1443 Analysis [data set], <https://dx.doi.org/10.5285/115d5e4ebf7148ec941423ec86fa9f26>, 2023.

1444 Dunn, R. J. H., Blannin, J., Gobron, N., Miller, J. B. and Willett, K. M. eds: Global climate [in “State of the Climate  
1445 in 2023”]. *Bull. Amer. Meteor. Soc.*, 105, S12-S155, <https://doi.org/10.1175/BAMS-D-24-0116.1>, 2024.

1446 Dutton, G.S., B. D. Hall, S.A. Montzka, J. D. Nance, S. D. Clingan, K. M. Petersen, Combined Atmospheric  
1447 Chlorofluorocarbon-12 Dry Air Mole Fractions from the NOAA GML Halocarbons Sampling Network, 1977-2024,  
1448 Version: 2024-03-07, <https://doi.org/10.15138/PJ63-H440>, 2024.

1449 ECCAD: CAMS database version 6.2 (v6.2), <https://permalink.acris-data.fr/CAMS-GLOB-ANT>, [data set], accessed  
1450 20 April 2025, 2025.

1451 Eyring, V., N. P. Gillett, K.M. Achuta Rao, R. Barimalala, M. Barreiro Parrillo, N. Bellouin, C. Cassou, P. J. Durack,  
1452 Y. Kosaka, S. McGregor, S. Min, O. Morgenstern, and Y. Sun: Human Influence on the Climate System. In *Climate  
1453 Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the  
1454 Intergovernmental Panel on Climate Change*[Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S.  
1455 Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K.  
1456 Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United  
1457 Kingdom and New York, NY, USA, pp. 423–552, <http://doi:10.1017/9781009157896.005>, 2021.

1458 Feron, S., Malhotra, A., Bansal, S., Fluet-Chouinard, E., McNicol, G., Knox, S. H., Delwiche, K. B., Cordero, R. R.,  
1459 Ouyang, Z., Zhang, Z., Poulter, B., and Jackson, R. B.: Recent increases in annual, seasonal, and extreme methane  
1460 fluxes driven by changes in climate and vegetation in boreal and temperate wetland ecosystems, *Global Change*  
1461 *Biology*, 30, e17131, <https://doi.org/10.1111/gcb.17131>, 2024.

1462 Forster, P. M., Forster, H. I., Evans, M. J., Gidden, M. J., Jones, C. D., Keller, C. A., Lamboll, R. D., Le Quéré, C.,  
1463 Rogelj, J., Rosen, D., Schleussner, C. F., Richardson, T. B., Smith, C. J. and Turnock, S. T.: Current and future global  
1464 climate impacts resulting from COVID-19, *Nature Clim. Chang.*, 10, 913–919, [https://doi.org/10.1038/s41558-020-](https://doi.org/10.1038/s41558-020-0883-0)  
1465 [0883-0](https://doi.org/10.1038/s41558-020-0883-0), 2020.

1466 Forster, P., T. Storelvmo, K. Armour, W. Collins, J.-L. Dufresne, D. Frame, D.J. Lunt, T. Mauritsen, M.D. Palmer,  
1467 M. Watanabe, M. Wild, and H. Zhang, 2021: The Earth's Energy Budget, Climate Feedbacks, and Climate Sensitivity.  
1468 In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment*  
1469 *Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors,  
1470 C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews,  
1471 T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United  
1472 Kingdom and New York, NY, USA, pp. 923–1054, <https://doi.org/10.1017/9781009157896.009>, 2021.

1473 Forster, P., Smith, C., Walsh, T., Lamb, W., Lamboll, R., Hauser, M., Ribes, A., Rosen, D., Gillett, N., Palmer, M.,  
1474 Rogelj, J., von Schuckmann, K., Seneviratne, S., Trewin, B., Zhang, X., Allen, M., Andrew, R., Birt, A., Borger, A.,  
1475 Boyer, T., Broersma, J., Cheng, L., Dentener, F., Friedlingstein, P., Gutiérrez, J., Gütschow, J., Hall, B., Ishii, M.,  
1476 Jenkins, S., Lan, X., Lee, J.-Y., Morice, C., Kadow, C., Kennedy, J., Killick, R., Minx, J., Naik, V., Peters, G., Pirani,  
1477 A., Pongratz, J., Schleussner, C.-F., Szopa, S., Thorne, P., Rohde, R., Rojas Corradi, M., Schumacher, D., Vose, R.,  
1478 Zickfeld, K., Masson-Delmotte, V., and Zhai, P.: Indicators of Global Climate Change 2022: annual update of large-  
1479 scale indicators of the state of the climate system and human influence, *Earth System Science Data*, 15, 2295–2327,  
1480 <https://doi.org/10.5194/essd-15-2295-2023>, 2023.

1481 Forster, P. M., Smith, C., Walsh, T., Lamb, W. F., Lamboll, R., Hall, B., Hauser, M., Ribes, A., Rosen, D., Gillett, N.  
1482 P., Palmer, M. D., Rogelj, J., Von Schuckmann, K., Trewin, B., Allen, M., Andrew, R., Betts, R. A., Borger, A.,  
1483 Boyer, T., Broersma, J. A., Buontempo, C., Burgess, S., Cagnazzo, C., Cheng, L., Friedlingstein, P., Gettelman, A.,  
1484 Gütschow, J., Ishii, M., Jenkins, S., Lan, X., Morice, C., Mühle, J., Kadow, C., Kennedy, J., Killick, R. E., Krummel,  
1485 P. B., Minx, J. C., Myhre, G., Naik, V., Peters, G. P., Pirani, A., Pongratz, J., Schleussner, C.-F., Seneviratne, S. I.,  
1486 Szopa, S., Thorne, P., Kovilakam, M. V. M., Majamäki, E., Jalkanen, J.-P., Van Marle, M., Hoesly, R. M., Rohde, R.,  
1487 Schumacher, D., Van Der Werf, G., Vose, R., Zickfeld, K., Zhang, X., Masson-Delmotte, V., and Zhai, P.: Indicators  
1488 of Global Climate Change 2023: annual update of key indicators of the state of the climate system and human  
1489 influence, *Earth Syst. Sci. Data*, 16, 2625–2658, <https://doi.org/10.5194/essd-16-2625-2024>, 2024.

1490 Fox-Kemper, B., Fox-Kemper, B., H. T. Hewitt, C. Xiao, G. Aðalgeirsdóttir, S.S. Drijfhout, T. L. Edwards, N. R.  
1491 Golledge, M. Hemer, R. E. Kopp, G. Krinner, A. Mix, D. Notz, S. Nowicki, I. S. Nurhati, L. Ruiz, J.-B. Sallée, A. B.  
1492 A. Slangen, and Y. Yu: Ocean, Cryosphere and Sea Level Change. In *Climate Change 2021: The Physical Science*  
1493 *Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate*

1494 Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb,  
1495 M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu,  
1496 and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1211–  
1497 1362, <https://doi.org/10.1017/9781009157896.011>, 2021.

1498 Francey, R.J., L.P. Steele, R.L. Langenfelds and B.C. Pak, High precision long-term monitoring of radiatively-active  
1499 trace gases at surface sites and from ships and aircraft in the Southern Hemisphere atmosphere, *J. Atmos. Science*, 56,  
1500 279–285 [https://doi.org/10.1175/1520-0469\(1999\)056<0279:HPLTMO>2.0.CO;2](https://doi.org/10.1175/1520-0469(1999)056<0279:HPLTMO>2.0.CO;2), 1999.

1501 Frederikse, T., Landerer, F., Caron, L., Adhikari, S., Parkes, D., Humphrey, V. W., Dangendorf, S., Hogarth, P.,  
1502 Zanna, L., Cheng, L., and Wu, Y.-H.: The causes of sea-level rise since 1900, *Nature*, 584, 393–397,  
1503 <https://doi.org/10.1038/s41586-020-2591-3>, 2020.

1504 Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Olsen, A., Peters, G. P., Peters, W.,  
1505 Pongratz, J., Sitch, S., Le Quéré, C., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S., Aragão, L. E. O. C., Armeth,  
1506 A., Arora, V., Bates, N. R., Becker, M., Benoit-Cattin, A., Bittig, H. C., Bopp, L., Bultan, S., Chandra, N., Chevallier,  
1507 F., Chini, L. P., Evans, W., Florentie, L., Forster, P. M., Gasser, T., Gehlen, M., Gilfillan, D., Gkritzalis, T., Gregor,  
1508 L., Gruber, N., Harris, I., Hartung, K., Haverd, V., Houghton, R. A., Ilyina, T., Jain, A. K., Joetzjer, E., Kadono, K.,  
1509 Kato, E., Kitidis, V., Korsbakken, J. I., Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Liu, Z., Lombardozzi,  
1510 D., Marland, G., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., Niwa, Y., O’Brien, K., Ono, T., Palmer,  
1511 P. I., Pierrot, D., Poulter, B., Resplandy, L., Robertson, E., Rödenbeck, C., Schwinger, J., Séférian, R., Skjelvan, I.,  
1512 Smith, A. J. P., Sutton, A. J., Tanhua, T., Tans, P. P., Tian, H., Tilbrook, B., van der Werf, G., Vuichard, N., Walker,  
1513 A. P., Wanninkhof, R., Watson, A. J., Willis, D., Wiltshire, A. J., Yuan, W., Yue, X., and Zaehle, S.: Global carbon  
1514 budget 2020, *Earth Syst. Sci. Data*, 12, 3269–3340, <https://doi.org/10.5194/essd-12-3269-2020>, 2020.

1515 Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Gregor, L., Hauck, J., Le Quéré, C., Luijkx, I. T.,  
1516 Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R.  
1517 B., Alin, S. R., Alkama, R., Armeth, A., Arora, V. K., Bates, N. R., Becker, M., Bellouin, N., Bittig, H. C., Bopp, L.,  
1518 Chevallier, F., Chini, L. P., Cronin, M., Evans, W., Falk, S., Feely, R. A., Gasser, T., Gehlen, M., Gkritzalis, T.,  
1519 Gloege, L., Grassi, G., Gruber, N., Gürses, Ö., Harris, I., Hefner, M., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina,  
1520 T., Jain, A. K., Jersild, A., Kadono, K., Kato, E., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I.,  
1521 Landschützer, P., Lefèvre, N., Lindsay, K., Liu, J., Liu, Z., Marland, G., Mayot, N., McGrath, M. J., Metzl, N.,  
1522 Monacci, N. M., Munro, D. R., Nakaoka, S.-I., Niwa, Y., O’Brien, K., Ono, T., Palmer, P. I., Pan, N., Pierrot, D.,  
1523 Pocock, K., Poulter, B., Resplandy, L., Robertson, E., Rödenbeck, C., Rodriguez, C., Rosan, T. M., Schwinger, J.,  
1524 Séférian, R., Shutler, J. D., Skjelvan, I., Steinhoff, T., Sun, Q., Sutton, A. J., Sweeney, C., Takao, S., Tanhua, T., Tans,  
1525 P. P., Tian, X., Tian, H., Tilbrook, B., Tsujino, H., Tubiello, F., van der Werf, G. R., Walker, A. P., Wanninkhof, R.,  
1526 Whitehead, C., Willstrand Wranne, A., et al.: Global Carbon Budget 2022, *Earth Syst. Sci. Data*, 14, 4811–4900,  
1527 <https://doi.org/10.5194/essd-14-4811-2022>, 2022.

1528 Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Bakker, D. C. E., Hauck, J., Landschützer, P., Le  
 1529 Quéré, C., Luijkx, I. T., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P.,  
 1530 Jackson, R. B., Alin, S. R., Anthoni, P., Barbero, L., Bates, N. R., Becker, M., Bellouin, N., Decharme, B., Bopp, L.,  
 1531 Brasika, I. B. M., Cadule, P., Chamberlain, M. A., Chandra, N., Chau, T.-T.-T., Chevallier, F., Chini, L. P., Cronin,  
 1532 M., Dou, X., Enyo, K., Evans, W., Falk, S., Feely, R. A., Feng, L., Ford, D. J., Gasser, T., Ghattas, J., Gkritzalis, T.,  
 1533 Grassi, G., Gregor, L., Gruber, N., Gürses, Ö., Harris, I., Hefner, M., Heinke, J., Houghton, R. A., Hurtt, G. C., Iida,  
 1534 Y., Ilyina, T., Jacobson, A. R., Jain, A., Jarníková, T., Jersild, A., Jiang, F., Jin, Z., Joos, F., Kato, E., Keeling, R. F.,  
 1535 Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Körtzinger, A., Lan, X., Lefèvre, N., Li, H., Liu, J.,  
 1536 Liu, Z., Ma, L., Marland, G., Mayot, N., McGuire, P. C., McKinley, G. A., Meyer, G., Morgan, E. J., Munro, D. R.,  
 1537 Nakaoka, S.-I., Niwa, Y., O'Brien, K. M., Olsen, A., Omar, A. M., Ono, T., Paulsen, M., Pierrot, D., Pocock, K.,  
 1538 Poulter, B., Powis, C. M., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Rosan, T. M., Schwinger, J.,  
 1539 Séférian, R., et al.: Global Carbon Budget 2023, *Earth System Science Data*, 15, 5301–5369,  
 1540 <https://doi.org/10.5194/essd-15-5301-2023>, 2023.  
 1541  
 1542 Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Landschützer, P., Le Quéré, C., Li, H.,  
 1543 Luijkx, I. T., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P.,  
 1544 Jackson, R. B., Alin, S. R., Arneeth, A., Arora, V., Bates, N. R., Becker, M., Bellouin, N., Berghoff, C. F., Bittig, H.,  
 1545 C., Bopp, L., Cadule, P., Campbell, K., Chamberlain, M. A., Chandra, N., Chevallier, F., Chini, L. P., Colligan, T.,  
 1546 Decayeux, J., Djéutchouang, L. M., Dou, X., Duran Rojas, C., Enyo, K., Evans, W., Fay, A. R., Feely, R. A., Ford, D.  
 1547 J., Foster, A., Gasser, T., Gehlen, M., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, Ö., Harris, I., Hefner,  
 1548 M., Heinke, J., Hurtt, G. C., Iida, Y., Ilyina, T., Jacobson, A. R., Jain, A. K., Jarníková, T., Jersild, A., Jiang, F., Jin,  
 1549 Z., Kato, E., Keeling, R. F., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Lan, X., Lauvset, S. K., Lefèvre, N.,  
 1550 Liu, Z., Liu, J., Ma, L., Maksyutov, S., Marland, G., Mayot, N., McGuire, P. C., Metzl, N., Monacchi, N. M., Morgan,  
 1551 E. J., Nakaoka, S.-I., Neill, C., Niwa, Y., Nützel, T., Olivier, L., Ono, T., Palmer, P. I., Pierrot, D., Qin, Z., Resplandy,  
 1552 L., Roobaert, A., Rosan, T. M., Rödenbeck, C., Schwinger, J., Smallman, T. L., Smith, S. M., Sospedra-Alfonso, R.,  
 1553 Steinhoff, T., Sun, Q., Sutton, A. J., Séférian, R., Takao, S., Tatebe, H., Tian, H., Tilbrook, B., Torres, O., Tourigny,  
 1554 E., Tsujino, H., Tubiello, F., van der Werf, G., Wanninkhof, R., Wang, X., Yang, D., Yang, X., Yu, Z., Yuan, W.,  
 1555 Yue, X., Zaehle, S., Zeng, N., and Zeng, J.: Global Carbon Budget 2024, *Earth Syst. Sci. Data*, 17, 965–1039,  
 1556 <https://doi.org/10.5194/essd-17-965-2025>, 2025.  
 1557 Gasser, T., Crepin, L., Quilcaille, Y., Houghton, R. A., Ciais, P., and Obersteiner, M.: Historical CO<sub>2</sub> emissions from  
 1558 land use and land cover change and their uncertainty, *Biogeosciences*, 17, 4075–4101, [https://doi.org/10.5194/bg-17-](https://doi.org/10.5194/bg-17-4075-2020)  
 1559 [4075-2020](https://doi.org/10.5194/bg-17-4075-2020), 2020.  
 1560 Gidden, M. J., Gasser, T., Grassi, G., Forsell, N., Janssens, I., Lamb, W. F., Minx, J., Nicholls, Z., Steinhauser, J., and  
 1561 Riahi, K.: Aligning climate scenarios to emissions inventories shifts global benchmarks, *Nature*, 624, 102–108,  
 1562 <https://doi.org/10.1038/s41586-023-06724-y>, 2023.

1563 Gillett, N.P., Kirchmeier-Young, M., Ribes, A., Shiogama, H., Hegerl, G.C., Knutti, R., Gastineau, G., John, J.G., Li,  
 1564 L., Nazarenko, L., Rosenbloom, N., Seland, Ø., Wu, T., Yukimoto, S., and Ziehn, T.: Constraining human  
 1565 contributions to observed warming since the pre-industrial period, *Nat. Clim. Chang.*, 11, 207–212,  
 1566 <https://doi.org/10.1038/s41558-020-00965-9>, 2021.

1567 Gleckler, P. J., Durack, P. J., Stouffer, R. J., Johnson, G. C., and Forest, C. E.: Industrial-era global ocean heat uptake  
 1568 doubles in recent decades, *Nat. Clim. Chang.*, 6, 394–398, <https://doi.org/10.1038/nclimate2915>, 2016.

1569 Goessling, H. F., Rackow, T., and Jung, T.: Recent global temperature surge intensified by record-low planetary  
 1570 albedo, *Science*, 387, 68–73, <https://doi.org/10.1126/science.adq7280>, 2025.

1571 Grassi, G., Stehfest, E., Rogelj, J., Van Vuuren, D., Cescatti, A., House, J., Nabuurs, G.-J., Rossi, S., Alkama, R.,  
 1572 Viñas, R. A., Calvin, K., Ceccherini, G., Federici, S., Fujimori, S., Gusti, M., Hasegawa, T., Havlik, P., Humpeöder,  
 1573 F., Korosuo, A., Perugini, L., Tubiello, F. N., and Popp, A.: Critical adjustment of land mitigation pathways for  
 1574 assessing countries' climate progress, *Nat. Clim. Chang.*, 11, 425–434, <https://doi.org/10.1038/s41558-021-01033-6>,  
 1575 2021.

1576 Grassi, G., Schwingshackl, C., Gasser, T., Houghton, R. A., Sitch, S., Canadell, J. G., Cescatti, A., Ciais, P., Federici,  
 1577 S., Friedlingstein, P., Kurz, W. A., Sanz Sanchez, M. J., Abad Viñas, R., Alkama, R., Bultan, S., Ceccherini, G., Falk,  
 1578 S., Kato, E., Kennedy, D., Knauer, J., Korosuo, A., Melo, J., McGrath, M. J., Nabel, J. E. M. S., Poulter, B.,  
 1579 Romanovskaya, A. A., Rossi, S., Tian, H., Walker, A. P., Yuan, W., Yue, X., and Pongratz, J.: Harmonising the land-  
 1580 use flux estimates of global models and national inventories for 2000–2020, *Earth Syst. Sci. Data*, 15, 1093–1114,  
 1581 <https://doi.org/10.5194/essd-15-1093-2023>, 2023.

1582 Guinaldo, T., Cassou, C., Sallée, J.-B., and Liné, A.: Internal variability effect doped by climate change drove the  
 1583 2023 marine heat extreme in the North Atlantic, *Commun Earth Environ*, 6, 291, <https://doi.org/10.1038/s43247-025-02197-1>,  
 1584 2025.

1585 Gulev, S. K., P. W. Thorne, J. Ahn, F. J. Dentener, C. M. Domingues, S. Gerland, D. Gong, D. S. Kaufman, H. C.  
 1586 Nnamchi, J. Quaas, J.A. Rivera, S. Sathyendranath, S.L. Smith, B. Trewin, K. von Schuckmann, and R. S. Vose:  
 1587 Changing State of the Climate System. In *Climate Change 2021: The Physical Science Basis. Contribution of Working*  
 1588 *Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*[Masson-Delmotte, V., P.  
 1589 Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell,  
 1590 E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge  
 1591 University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 287–422,  
 1592 <https://doi.org/10.1017/9781009157896.004>, 2021.

1593 Gupta, A. K., Mittal, T., Fauria, K. E., Bennartz, R., and Kok, J. F.: The January 2022 Hunga eruption cooled the  
 1594 southern hemisphere in 2022 and 2023, *Commun Earth Environ*, 6, 240, [https://doi.org/10.1038/s43247-025-02181-](https://doi.org/10.1038/s43247-025-02181-9)  
 1595 [9](https://doi.org/10.1038/s43247-025-02181-9), 2025.

1596 Gütschow, J., Jeffery, M. L., Gieseke, R., Gebel, R., Stevens, D., Krapp, M., and Rocha, M.: The PRIMAP-hist  
 1597 national historical emissions time series, *Earth Syst. Sci. Data*, 8, 571–603, <https://doi.org/10.5194/essd-8-571-2016>,  
 1598 2016.

1599 Gütschow, J., and Busch, D., and Pflüger, M.: The PRIMAP-hist national historical emissions time series v2.6.1  
1600 (1750-2023) (2.6.1), Zenodo [data set] <https://doi.org/10.5281/zenodo.15016289>, 2025.

1601 Hardy, A., Palmer, P. I., and Oakes, G.: Satellite data reveal how Sudd wetland dynamics are linked with globally-  
1602 significant methane emissions, *Environ. Res. Lett.*, 18, 074044, <https://doi.org/10.1088/1748-9326/ace272>, 2023.

1603 Hay, C. C., Morrow, E., Kopp, R. E., and Mitrovica, J. X.: Probabilistic reanalysis of twentieth-century sea-level rise,  
1604 *Nature*, 517, 481–484, <https://doi.org/10.1038/nature14093>, 2015.

1605 Hakuba, M. Z., Frederikse, T., and Landerer, F. W.: Earth's energy imbalance from the ocean perspective (2005–  
1606 2019), *Geophys Res Lett*, 48, e2021GL093624, <https://doi.org/10.1029/2021GL093624>, 2021.

1607 Hansen, J. E., Sato, M., Simons, L., Nazarenko, L. S., Sangha, I., Kharecha, P., Zachos, J. C., von Schuckmann, K.,  
1608 Loeb, N. G., Osman, M. B., Jin, Q., Tselioudis, G., Jeong, E., Lacis, A., Ruedy, R., Russell, G., Cao, J., and Li, J.:  
1609 Global warming in the pipeline, *Oxford Open Climate Change*, 3, kgad008, <https://doi.org/10.1093/oxfclm/kgad008>,  
1610 2023.

1611 Hansis, E., Davis, S. J., and Pongratz, J.: Relevance of methodological choices for accounting of land use change  
1612 carbon fluxes, *Global Biogeochem. Cy.*, 29, 1230–1246, <https://doi.org/10.1002/2014GB004997>, 2015.

1613 Haustein, K., Allen, M. R., Forster, P. M., Otto, F. E. L., Mitchell, D. M., Matthews, H. D., and Frame, D. J.: A real-  
1614 time Global Warming Index, *Sci Rep*, 7, 15417, <https://doi.org/10.1038/s41598-017-14828-5>, 2017.

1615 Harris, I., Osborn, T. J., Jones, P., and Lister, D.: Version 4 of the CRU TS monthly high-resolution gridded  
1616 multivariate climate dataset, *Scientific data*, 7, 109, <https://doi.org/10.1038/s41597-020-045303>, 2020

1617 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R.,  
1618 Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J.,  
1619 Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes,  
1620 M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P.,  
1621 Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global  
1622 reanalysis, *Q. J. R. Meteorol. Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.

1623 Hodnebrog, Ø., Aamaas, B., Fuglestad, J. S., Marston, G., Myhre, G., Nielsen, C. J., Sandstad, M., Shine, K. P., and  
1624 Wallington, T. J.: Updated Global Warming Potentials and Radiative Efficiencies of Halocarbons and Other Weak  
1625 Atmospheric Absorbers, *Rev. Geophys.*, 58, e2019RG000691, <https://doi.org/10.1029/2019RG000691>, 2020.

1626 Hodnebrog, Ø., Myhre, G., Jouan, C., Andrews, T., Forster, P. M., Jia, H., Loeb, N. G., Olivieri, D. J. L., Paynter, D.,  
1627 Quaas, J., Raghuraman, S. P., and Schulz, M.: Recent reductions in aerosol emissions have increased Earth's energy  
1628 imbalance, *Communications Earth & Environment*, 5, 166, <https://doi.org/10.1038/s43247-024-01324-8>, 2024.

1629 Hoesly, R., Smith, S. J., Ahsan, H., Prime, N., O'Rourke, P., Crippa, M., Klimont, Z., Guizzardi, D., Feng, L., Harkins,  
1630 C., MCDONALD, B., and Wang, S.: CEDS v\_2025\_03\_18 Aggregate Data (v\_2025\_03\_18),  
1631 <https://doi.org/10.5281/ZENODO.15059443>, 2025.

1632 Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres,  
1633 R. J., Bolt, R. M., Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J.-I., Li, M., Liu, L., Lu, Z., Moura, M. C. P.,

1634 O'Rourke, P. R., and Zhang, Q.: Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from  
 1635 the Community Emissions Data System (CEDS), *Geosci. Model. Dev.*, 11, 369–408, [https://doi.org/10.5194/gmd-11-](https://doi.org/10.5194/gmd-11-369-2018)  
 1636 [369-2018](https://doi.org/10.5194/gmd-11-369-2018), 2018.

1637 Hoesly, R., & Smith, S., CEDS v\_2024\_04\_01 Release Emission Data (v\_2024\_04\_01) [Data set], Zenodo.  
 1638 <https://doi.org/10.5281/zenodo.10904361>, 2024.

1639 Houghton, R. A., and Nassikas, A. A.: Global and regional fluxes of carbon from land use and land cover change  
 1640 1850–2015, *Global Biogeochem. Cy.*, 31, 456–472, <https://doi.org/10.1002/2016GB005546>, 2017.

1641 Houghton, R. A. and Castanho, A.: Annual emissions of carbon from land use, land-use change, and forestry from  
 1642 1850 to 2020, *Earth System Science Data*, 15, 2025–2054, <https://doi.org/10.5194/essd-15-2025-2023>, 2023.

1643 Hu, Y., Yue, X., Tian, C., Zhou, H., Fu, W., Zhao, X., Zhao, Y., and Chen, Y.: Identifying the main drivers of the  
 1644 spatiotemporal variations in wetland methane emissions during 2001–2020, *Frontiers in Environmental Science*, 11,  
 1645 <https://doi.org/10.3389/fenvs.2023.1275742>, 2023.

1646 Huang, B., Yin, X., Boyer, T., Liu, C., Menne, M., Rao, Y. D., Smith, T., Vose, R., and Zhang, H.-M.: Extended  
 1647 Reconstructed Sea Surface Temperature, Version 6 (ERSSTv6). Part I: An Artificial Neural Network Approach,  
 1648 *Journal of Climate*, 38, 1105–1121, <https://doi.org/10.1175/JCLI-D-23-0707.1>, 2025.

1649 IATA: Air Passenger Monthly Analysis March 2024, [https://www.iata.org/en/iata-repository/publications/economic-](https://www.iata.org/en/iata-repository/publications/economic-reports/air-passenger-market-analysis-march-2024/)  
 1650 [reports/air-passenger-market-analysis-march-2024/](https://www.iata.org/en/iata-repository/publications/economic-reports/air-passenger-market-analysis-march-2024/), accessed 20.05.2024, 2024.

1651 IEA: CO2 Emissions in 2023. <https://www.iea.org/reports/co2-emissions-in-2023>, accessed 20.04.2024, 2024.

1652 IPCC: Sixty-second Session of the IPCC (IPCC-62), Fifteenth Session of the IPCC Working Group I (WGI-15),  
 1653 Thirteenth Session of the IPCC Working Group II (WGII-13), and Fifteenth Session of the IPCC Working Group III  
 1654 (WGIII-15), <https://www.ipcc.ch/meeting-doc/ipcc-62/>, accessed 20 April 2025, 2025.

1655 IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment  
 1656 Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K.  
 1657 Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge,  
 1658 United Kingdom and New York, NY, USA, 1535 pp, <https://doi:10.1017/CBO9781107415324>, 2013.

1659 IPCC: Summary for Policymakers. In: Global Warming of 1.5°C. An IPCC Special Report on the impacts of global  
 1660 warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of  
 1661 strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate  
 1662 poverty [Masson-Delmotte, V., P. Zhai, H.-O. Pörtner, D. Roberts, J. Skea, P.R. Shukla, A. Pirani, W. Moufouma-  
 1663 Okia, C. Péan, R. Pidcock, S. Connors, J.B.R. Matthews, Y. Chen, X. Zhou, M.I. Gomis, E. Lonnoy, T. Maycock, M.  
 1664 Tignor, and T. Waterfield (eds.)]. Cambridge University Press, Cambridge, UK and New York, NY, USA, pp. 3-24,  
 1665 <https://doi.org/10.1017/9781009157940.001>, 2018.

1666 IPCC: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment  
 1667 Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, United Kingdom  
 1668 and New York, NY, USA, <https://doi.org/10.1017/9781009157896>, 2021a.

1669 IPCC: Summary for Policymakers, in: Climate Change 2021: The Physical Science Basis.  
 1670 Contribution of Working Group I to the Sixth Assessment Report of the  
 1671 Intergovernmental Panel on Climate Change, edited by: Masson-Delmotte, V., Zhai, P.,  
 1672 Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M.  
 1673 I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T.,  
 1674 Yelekçi, O., Yu, R., and Zhou, B., Cambridge University Press, Cambridge, United  
 1675 Kingdom and New York, NY, USA, pp.3–32 <https://doi.org/10.1017/9781009157896.001>, 2021b.

1676 IPCC: Climate Change 2022: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Sixth  
 1677 Assessment Report of the Intergovernmental Panel on Climate Change [H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S.  
 1678 Poloczanska, K. Mintenbeck, A. Alegría, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, B. Rama (eds.)].  
 1679 Cambridge University Press. Cambridge University Press, Cambridge, UK and New York, NY, USA, 3056 pp.,  
 1680 <https://doi.org/10.1017/9781009325844>, 2022.

1681 IPCC, 2023: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth  
 1682 Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and J. Romero  
 1683 (eds.)]. IPCC, Geneva, Switzerland., Intergovernmental Panel on Climate Change (IPCC),  
 1684 <https://doi.org/10.59327/IPCC/AR6-9789291691647>, 2023a.

1685 IPCC, 2023: Climate Change 2023: Summary for Policy Makers. Contribution of Working Groups I, II and III to the  
 1686 Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and J.  
 1687 Romero (eds.)]. IPCC, Geneva, Switzerland., Intergovernmental Panel on Climate Change (IPCC),  
 1688 <https://doi.org/10.59327/IPCC/AR6-9789291691647>, 2023b.

1689 Iturbide, M., Fernández, J., Gutiérrez, J. M., Pirani, A., Huard, D., Al Khourdajie, A., Baño-Medina, J., Bedia, J.,  
 1690 Casanueva, A., Cimadevilla, E., Cofiño, A. S., De Felice, M., Díez-Sierra, J., García-Díez, M., Goldie, J., Herrera, D.  
 1691 A., Herrera, S., Manzanar, R., Milovac, J., Radhakrishnan, A., San-Martín, D., Spinuso, A., Thyng, K. M., Trenham,  
 1692 C., and Yelekçi, Ö.: Implementation of FAIR principles in the IPCC: the WGI AR6 Atlas repository, Sci Data, 9, 629,  
 1693 <https://doi.org/10.1038/s41597-022-01739-y>, 2022.

1694 Janardanan, R., Maksyutov, S., Wang, F., Nayagam, L., Sahu, S. K., Mangaraj, P., Saunio, M., Lan, X., and  
 1695 Matsunaga, T.: Country-level methane emissions and their sectoral trends during 2009–2020 estimated by high-  
 1696 resolution inversion of GOSAT and surface observations, Environ. Res. Lett., 19, 034007,  
 1697 <https://doi.org/10.1088/1748-9326/ad2436>, 2024.

1698 Jenkins, S., Povey, A., Gettelman, A., Grainger, R., Stier, P., and Allen, M.: Is Anthropogenic Global Warming  
 1699 Accelerating?, Journal of Climate, 35, 7873–7890, <https://doi.org/10.1175/JCLI-D-22-0081.1>, 2022.



1700 Jenkins, S., Smith, C., Allen, M., and Grainger, R.: Tonga eruption increases chance of temporary surface temperature  
 1701 anomaly above 1.5 °C, *Nature Clim. Chang.*, 13, 127–129, <https://doi.org/10.1038/s41558-022-01568-2>, 2023.

1702 Kirchengast, G., Gorfer, M., Mayer, M., Steiner, A. K., and Haimberger, L.: GCOS EHI 1960-2020 Atmospheric Heat  
 1703 Content, [https://doi.org/10.26050/WDCC/GCOS\\_EHI\\_1960-2020\\_AHC](https://doi.org/10.26050/WDCC/GCOS_EHI_1960-2020_AHC), 2022.

1704 Kramer, R. J., He, H., Soden, B. J., Oreopoulos, L., Myhre, G., Forster, P. M., and Smith, C. J., Observational evidence  
 1705 of increasing global radiative forcing, *Geophys. Res. Lett.*, 48, e2020GL091585,  
 1706 <https://doi.org/10.1029/2020GL091585>, 2021.

1707 Lamb, W., Andrew, R., Jones, M., Nicholls, Z., Peters, G., Smith, C., Saunio, M., Grassi, G., Pongratz, J., Smith, S.,  
 1708 Tubiello, F., Crippa, M., Gidden, M., Friedlingstein, P., Minx, J., and Forster, P.: Differences in anthropogenic  
 1709 greenhouse gas emissions estimates explained, <https://doi.org/10.5194/essd-2025-188>, 24 April 2025.

1710 Lamboll, R. D., Jones, C. D., Skeie, R. B., Fiedler, S., Samset, B. H., Gillett, N. P., Rogelj, J., and Forster, P. M.:  
 1711 Modifying emissions scenario projections to account for the effects of COVID-19: protocol for CovidMIP,  
 1712 *Geoscientific Model Development*, 14, 3683–3695, <https://doi.org/10.5194/gmd-14-3683-2021>, 2021.

1713 Lamboll, R. D. and Rogelj, J.: Code for estimation of remaining carbon budget in IPCC AR6 WGI, Zenodo [code],  
 1714 <https://doi.org/10.5281/zenodo.6373365>, 2022.

1715 Lamboll, R. and Rogelj, J.: Carbon Budget Calculator, 2025, Github [code],  
 1716 <https://github.com/RLamboll/AR6CarbonBudgetCalc/tree/v1.0.1>, last access: 25 April 2025, 2025.

1717 Lamboll, R. D., Nicholls, Z. R. J., Smith, C. J., Kikstra, J. S., Byers, E., and Rogelj, J.: Assessing the size and  
 1718 uncertainty of remaining carbon budgets, *Nature Climate Change*, 13, 1360–1367, <https://doi.org/10.1038/s41558-023-01848-5>, 2023.

1720 Lan, X., Tans, P. and Thoning, K.W.: Trends in globally-averaged CO<sub>2</sub> determined from NOAA Global Monitoring  
 1721 Laboratory measurements, Version Monday, 14-Apr-2025 09:08:57 MDT <https://doi.org/10.15138/9N0H-ZH07>,  
 1722 2025.

1723 Lan, X., Thoning, K. W., and Dlugokencky, E.J.: Trends in globally-averaged CH<sub>4</sub> N<sub>2</sub>O, and SF<sub>6</sub> determined from  
 1724 NOAA Global Monitoring Laboratory measurements, Version 2023-04, <https://doi.org/10.15138/P8XG-AA10>,  
 1725 2023b.

1726 Laube, J., Newland, M., Hogan, C., Brenninkmeijer, A.M., Fraser, P.J., Martinerie, P., Oram, D.E., Reeves, C.E.,  
 1727 Röckmann, T., Schwander, J., Witrant, E., Sturges, W.T.: Newly detected ozone-depleting substances in the  
 1728 atmosphere. *Nature Geosci.*, 7, 266–269, <https://doi.org/10.1038/ngeo2109>, 2014.

1729 Lee, J.-Y., J. Marotzke, G. Bala, L. Cao, S. Corti, J.P. Dunne, F. Engelbrecht, E. Fischer, J.C. Fyfe, C. Jones, A.  
 1730 Maycock, J. Mutemi, O. Ndiaye, S. Panickal, and T. Zhou: Future Global Climate: Scenario-Based Projections and  
 1731 Near-Term Information. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to  
 1732 the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*[Masson-Delmotte, V., P. Zhai, A.  
 1733 Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E.

1734 Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)). Cambridge  
 1735 University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 553–  
 1736 672, <https://doi.org/10.1017/9781009157896.006>, 2021.

1737 Lee, H., K. Calvin, D. Dasgupta, G. Krinner, A. Mukherji, P. Thorne, C. Trisos, J. Romero, P. Aldunce, K. Barrett,  
 1738 G. Blanco, W.W.L. Cheung, S.L. Connors, F. Denton, A. Diongue-Niang, D. Dodman, M. Garschagen, O. Geden, B.  
 1739 Hayward, C. Jones, F. Jotzo, T. Krug, R. Lasco, J.-Y. Lee, V. Masson-Delmotte, M. Meinshausen, K. Mintenbeck, A.  
 1740 Mokssit, F.E.L. Otto, M. Pathak, A. Pirani, E. Poloczanska, H.-O. Pörtner, A. Revi, D.C. Roberts, J. Roy, A.C. Ruane,  
 1741 J. Skea, P.R. Shukla, R. Slade, A. Slangen, Y. Sokona, A.A. Sörensson, M. Tignor, D. van Vuuren, Y.-M. Wei, H.  
 1742 Winkler, P. Zhai, and Z. Zommers: Synthesis Report of the IPCC Sixth Assessment Report (AR6): Summary for  
 1743 Policymakers. Intergovernmental Panel on Climate Change [accepted], available at  
 1744 <https://www.ipcc.ch/report/ar6/syr/>, 2023.

1745 Liu, Z., Deng, Z., Davis, S. J., and Ciais, P.: Global carbon emissions in 2023, *Nature Reviews Earth & Environment*,  
 1746 5, 253–254, <https://doi.org/10.1038/s43017-024-00532-2>, 2024.

1747 Loeb, N. G., Johnson, G. C., Thorsen, T. J., Lyman, J. M., Rose, F. G., Kato, S.: Satellite and ocean data reveal marked  
 1748 increase in Earth's heating rate. *Geophys. Res. Lett.*, 48, e2021GL093047, <https://doi.org/10.1029/2021GL093047>,  
 1749 2021.

1750 van Marle, M. J. E., Kloster, S., Magi, B. I., Marlon, J. R., Daniau, A.-L., Field, R. D., Arneeth, A., Forrest, M.,  
 1751 Hantson, S., Kehrwald, N. M., Knorr, W., Lasslop, G., Li, F., Mangeon, S., Yue, C., Kaiser, J. W., and van der Werf,  
 1752 G. R.: Historic global biomass burning emissions for CMIP6 (BB4CMIP) based on merging satellite observations  
 1753 with proxies and fire models (1750–2015), *Geosci. Model Dev.*, 10, 3329–3357, [https://doi.org/10.5194/gmd-10-](https://doi.org/10.5194/gmd-10-3329-2017)  
 1754 [3329-2017](https://doi.org/10.5194/gmd-10-3329-2017), 2017.

1755 McKenna, C. M., Maycock, A. C., Forster, P. M., Smith, C. J., and Tokarska, K. B.: Stringent mitigation substantially  
 1756 reduces risk of unprecedented near-term warming rates, *Nature Climate Change*, 11, 126–131,  
 1757 <https://doi.org/10.1038/s41558-020-00957-9>, 2021.

1758 Menne, M. J., Williams, C. N., Gleason, B. E., Rennie, J. J., and Lawrimore, J. H.: The global historical climatology  
 1759 network monthly temperature dataset, version 4, *J. Climate*, 31, 9835–9854, [https://doi.org/10.1175/JCLI-D-18-](https://doi.org/10.1175/JCLI-D-18-0094.1)  
 1760 [0094.1](https://doi.org/10.1175/JCLI-D-18-0094.1), 2018.

1761 [Merchant, C. J., Allan, R.P., Embury, O.: Quantifying the acceleration of multidecadal global sea surface warming](https://doi.org/10.1088/1748-9326/adaa8a)  
 1762 [driven by Earth's energy imbalance, \*Environ. Res. Lett.\*, 20, 024037, https://doi.org/10.1088/1748-9326/adaa8a, 2025.](https://doi.org/10.1088/1748-9326/adaa8a)

1763 Minière, A., von Schuckmann, K., Sallée, J.-B., and Vogt, L.: Robust acceleration of Earth system heating observed  
 1764 over the past six decades, *Scientific Reports*, 13, 22975, <https://doi.org/10.1038/s41598-023-49353-1>, 2023.

1765 Minobe, S., Behrens, E., Findell, K. L., Loeb, N. G., Meyssignac, B., and Sutton, R.: Global and regional drivers for  
 1766 exceptional climate extremes in 2023-2024: beyond the new normal, *npj Clim Atmos Sci*, 8, 138,  
 1767 <https://doi.org/10.1038/s41612-025-00996-z>, 2025.

Minx, J. C., Lamb, W. F., Andrew, R. M., Canadell, J. G., Crippa, M., Döbbeling, N., Forster, P. M., Guizzardi, D., Olivier, J., Peters, G. P., Pongratz, J., Reisinger, A., Rigby, M., Saunio, M., Smith, S. J., Solazzo, E., and Tian, H.: A comprehensive and synthetic dataset for global, regional, and national greenhouse gas emissions by sector 1970–2018 with an extension to 2019, *Earth Syst. Sci. Data*, 13, 5213–5252, <https://doi.org/10.5194/essd-13-5213-2021>, 2021.

NASA: Satellite sea level observations, [data set], <https://sealevel.nasa.gov/understanding-sea-level/key-indicators/global-mean-sea-level/>, accessed 19 February 2025, 2025.

Nickolay A. Krotkov, Lok N. Lamsal, Sergey V. Marchenko, Edward A. Celarier, Eric J. Bucsela, William H. Swartz, Joanna Joiner and the OMI core team, OMI/Aura NO<sub>2</sub> Cloud-Screened Total and Tropospheric Column L3 Global Gridded 0.25 degree x 0.25 degree V3, NASA Goddard Space Flight Center, Goddard Earth Sciences Data and Information Services Center (GES DISC), Accessed: [Data Access 22 April 2024], <https://doi.org/10.5067/Aura/OMI/DATA3007>, 2019.

Nisbet, E. G., Manning, M. R., Dlugokencky, E. J., Michel, S. E., Lan, X., Roeckmann, T., Gon, H. A. D. V. D., Palmer, P., Oh, Y., Fisher, R., Lowry, D., France, J. L., and White, J. W. C.: Atmospheric methane: Comparison between methane’s record in 2006–2022 and during glacial terminations, Preprints, <https://doi.org/10.22541/essoar.167689502.25042797/v1>, 2023.

Nitzbon, J., Krinner, G., Langer, M.: GCOS EHI 1960–2020 Permafrost Heat Content, World Data Center for Climate (WDCC) at DKRZ, [https://doi.org/10.26050/WDCC/GCOS\\_EHI\\_1960-2020\\_PHC](https://doi.org/10.26050/WDCC/GCOS_EHI_1960-2020_PHC), 2022.

NOAA: Global sea level timeseries, [https://www.star.nesdis.noaa.gov/socd/lisa/SeaLevelRise/LSA\\_SLR\\_timeseries.php](https://www.star.nesdis.noaa.gov/socd/lisa/SeaLevelRise/LSA_SLR_timeseries.php), [data set], accessed 19 February 2025, 2025.

Palmer, M. D. and McNeall, D. J.: Internal variability of Earth’s energy budget simulated by CMIP5 climate models, *Environ. Res. Lett.*, 9, 034016, <https://doi.org/10.1088/1748-9326/9/3/034016>, 2014.

Palmer, M. D., Domingues, C. M., Slangen, A. B. A., and Boeira Dias, F.: An ensemble approach to quantify global mean sea-level rise over the 20th century from tide gauge reconstructions, *Environ. Res. Lett.*, 16, 044043, <https://doi.org/10.1088/1748-9326/abdaec>, 2021.

Peng, S., Lin, X., Thompson, R. L., Xi, Y., Liu, G., Hauglustaine, D., Lan, X., Poulter, B., Ramonet, M., Saunio, M., Yin, Y., Zhang, Z., Zheng, B., and Ciais, P.: Wetland emission and atmospheric sink changes explain methane growth in 2020, *Nature*, 612, 477–482, <https://doi.org/10.1038/s41586-022-05447-w>, 2022.

Pelz, S., Ganti, G., Lamboll, R., Grant, L., Smith, C., Pachauri, S., Rogelj, J., Riahi, K., Thiery, W., and Gidden, M. J.: Using net-zero carbon debt to track climate overshoot responsibility, *Proc. Natl. Acad. Sci. U.S.A.*, 122, e2409316122, <https://doi.org/10.1073/pnas.2409316122>, 2025.

Pirani, A., Alegria, A., Khourdajie, A. A., Gunawan, W., Gutiérrez, J. M., Holsman, K., Huard, D., Juckes, M., Kawamiya, M., Klutse, N., Krey, V., Matthews, R., Milward, A., Pascoe, C., Van Der Shrier, G., Spinuso, A., Stockhause, M., and Xiaoshi Xing: The implementation of FAIR data principles in the IPCC AR6 assessment process, <https://doi.org/10.5281/ZENODO.6504469>, 2022.

1803 Pongratz, J., Schwingshackl, C., Bultan, S., Obermeier, W., Havemann, F., and Guo, S.: Land Use Effects on Climate:  
 1804 Current State, Recent Progress, and Emerging Topics, *Curr. Clim. Change Rep.*, 7, 99–120,  
 1805 <https://doi.org/10.1007/s40641-021-00178-y>, 2021.

1806 Prinn, R. G., Weiss, R. F., Arduini, J., Arnold, T., DeWitt, H. L., Fraser, P. J., Ganesan, A. L., Gasore, J., Harth, C.  
 1807 M., Hermansen, O., Kim, J., Krummel, P. B., Li, S., Loh, Z. M., Lunder, C. R., Maione, M., Manning, A. J., Miller,  
 1808 B. R., Mitrevski, B., Mühle, J., O'Doherty, S., Park, S., Reimann, S., Rigby, M., Saito, T., Salameh, P. K., Schmidt,  
 1809 R., Simmonds, P. G., Steele, L. P., Vollmer, M. K., Wang, R. H., Yao, B., Yokouchi, Y., Young, D., and Zhou, L.:  
 1810 History of chemically and radiatively important atmospheric gases from the Advanced Global Atmospheric Gases  
 1811 Experiment (AGAGE), *Earth Syst. Sci. Data*, 10, 985–1018, <https://doi.org/10.5194/essd-10-985-2018>, 2018.

1812 Purich, A. and Doddridge, E. W.: Record low Antarctic sea ice coverage indicates a new sea ice state, *Commun Earth*  
 1813 *Environ.*, 4, 314, <https://doi.org/10.1038/s43247-023-00961-9>, 2023.

1814 Quaas, J., Jia, H., Smith, C., Albright, A. L., Aas, W., Bellouin, N., Boucher, O., Doutriaux-Boucher, M., Forster, P.  
 1815 M., Grosvenor, D., Jenkins, S., Klimont, Z., Loeb, N. G., Ma, X., Naik, V., Paulot, F., Stier, P., Wild, M., Myhre, G.,  
 1816 and Schulz, M.: Robust evidence for reversal of the trend in aerosol effective climate forcing, *Atmos. Chem. Phys.*,  
 1817 22, 12221–12239, <https://doi.org/10.5194/acp-22-12221-2022>, 2022.

1818 Qin, Z., Zhu, Y., Canadell, J. G., Chen, M., Li, T., Mishra, U., and Yuan, W.: Global spatially explicit carbon emissions  
 1819 from land-use change over the past six decades (1961–2020), *One Earth*, 7, 835–847,  
 1820 <https://doi.org/10.1016/j.oneear.2024.04.002>, 2024.

1821 Raghuraman, S. P., Paynter, D. and Ramaswamy, V.: Anthropogenic forcing and response yield observed positive  
 1822 trend in Earth's energy imbalance, *Nat. Commun.* 12, 4577, <https://doi.org/10.1038/s41467-021-24544-4>, 2021.

1823 Raghuraman, S. P., Soden, B., Clement, A., Vecchi, G., Menemenlis, S., and Yang, W.: The 2023 global warming  
 1824 spike was driven by the El Niño–Southern Oscillation, *Atmos. Chem. Phys.*, 24, 11275–11283,  
 1825 <https://doi.org/10.5194/acp-24-11275-2024>, 2024.

1826 Ribes, A., Qasmi, S., and Gillett, N. P.: Making climate projections conditional on historical observations, *Sci. Adv.*,  
 1827 7, eabc0671, <https://doi.org/10.1126/sciadv.abc0671>, 2021.

1828 Rogelj, J., D. Shindell, K. Jiang, S. Ffita, P. Forster, V. Ginzburg, C. Handa, H. Kheshgi, S. Kobayashi, E. Kriegler,  
 1829 L. Mundaca, R. Séférián, and M. V. Vilarinho: Mitigation Pathways Compatible with 1.5°C in the Context of  
 1830 Sustainable Development. In: *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming*  
 1831 *of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of*  
 1832 *strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate*  
 1833 *poverty* [Masson-Delmotte, V., P. Zhai, H.-O. Pörtner, D. Roberts, J. Skea, P.R. Shukla, A. Pirani, W. Moufouma-  
 1834 Okia, C. Péan, R. Pidcock, S. Connors, J. B. R. Matthews, Y. Chen, X. Zhou, M. I. Gomis, E. Lonnoy, T. Maycock,  
 1835 M. Tignor, and T. Waterfield (eds.)]. Cambridge University Press, Cambridge, UK and New York, NY, USA, pp. 93-  
 1836 174, <https://doi.org/10.1017/9781009157940.004>, 2018.

1837 Rogelj, J., Forster, P. M., Kriegler, E., Smith, C. J., and Séférian, R.: Estimating and tracking the remaining carbon  
1838 budget for stringent climate targets, *Nature*, 571, 335–342, <https://doi.org/10.1038/s41586-019-1368-z>, 2019.

1839 Rogelj, J., Lamboll, R.D.: Substantial reductions in non-CO2 greenhouse gas emissions reductions implied by IPCC  
1840 estimates of the remaining carbon budget. *Communications Earth Environ* 5, 35. [https://doi.org/10.1038/s43247-023-](https://doi.org/10.1038/s43247-023-01168-8)  
1841 [01168-8](https://doi.org/10.1038/s43247-023-01168-8), 2024.

1842 Rogelj, J., Rao, S., McCollum, D. L., Pachauri, S., Klimont, Z., Krey, V., and Riahi, K: Air-pollution emission ranges  
1843 consistent with the representative concentration pathways, *Nature Clim. Chang.*, 4 (6), 446–450,  
1844 <https://doi.org/10.1038/nclimate2178>, 2014.

1845 Rohde, R., Muller, R., Jacobsen, R., Perlmutter, S., Rosenfeld, A. et al.: Berkeley Earth Temperature Averaging  
1846 Process, *Geoinfor. Geostat.: An Overview* 1:2., <http://dx.doi.org/10.4172/gigs.1000103>, 2013.

1847 Saunio, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A., Dlugokencky,  
1848 E. J., Houweling, S., Patra, P. K., Ciais, P., Arora, V. K., Bastviken, D., Bergamaschi, P., Blake, D. R., Brailsford, G.,  
1849 Bruhwiler, L., Carlson, K. M., Carrol, M., Castaldi, S., Chandra, N., Crevoisier, C., Crill, P. M., Covey, K., Curry, C.  
1850 L., Etiopie, G., Frankenberg, C., Gedney, N., Hegglin, M. I., Höglund-Isaksson, L., Hugelius, G., Ishizawa, M., Ito,  
1851 A., Janssens-Maenhout, G., Jensen, K. M., Joos, F., Kleinen, T., Krummel, P. B., Langenfelds, R. L., Laruelle, G. G.,  
1852 Liu, L., Machida, T., Maksyutov, S., McDonald, K. C., McNorton, J., Miller, P. A., Melton, J. R., Morino, I., Müller,  
1853 J., Murguía-Flores, F., Naik, V., Niwa, Y., Noce, S., O'Doherty, S., Parker, R. J., Peng, C., Peng, S., Peters, G. P.,  
1854 Prigent, C., Prinn, R., Ramonet, M., Regnier, P., Riley, W. J., Rosentreter, J. A., Segers, A., Simpson, I. J., Shi, H.,  
1855 Smith, S. J., Steele, L. P., Thornton, B. F., Tian, H., Tohjima, Y., Tubiello, F. N., Tsuruta, A., Viovy, N., Voulgarakis,  
1856 A., Weber, T. S., Van Weele, M., Van Der Werf, G. R., Weiss, R. F., Worthy, D., Wunch, D., Yin, Y., Yoshida, Y.,  
1857 Zhang, W., Zhang, Z., Zhao, Y., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: The Global Methane Budget 2000–  
1858 2017, *Earth Syst. Sci. Data*, 12, 1561–1623, <https://doi.org/10.5194/essd-12-1561-2020>, 2020.

1859 Sato, K., Sato, K., Savita, A., Schweiger, A., Shepherd, A., Seneviratne, S. I., Simons, L., Slater, D. A., Slater, T.,  
1860 Steiner, A. K., Suga, T., Szekely, T., Thiery, W., Timmermans, M.-L., Vanderkelen, I., Wjiffels, S. E., Wu, T., and  
1861 Zemp, M.: GCOS EHI 1960-2020 Earth Heat Inventory Ocean Heat Content (Version 2),  
1862 [https://doi.org/10.26050/WDCC/GCOS\\_EHI\\_1960-2020\\_OHC\\_v2](https://doi.org/10.26050/WDCC/GCOS_EHI_1960-2020_OHC_v2), 2023b.

1863 Scarpelli, T. R., Jacob, D. J., Grossman, S., Lu, X., Qu, Z., Sulprizio, M. P., Zhang, Y., Reuland, F., Gordon, D., and  
1864 Worden, J. R.: Updated Global Fuel Exploitation Inventory (GFEL) for methane emissions from the oil, gas, and coal  
1865 sectors: evaluation with inversions of atmospheric methane observations, *Atmos. Chem. Phys.*, 22, 3235–3249,  
1866 <https://doi.org/10.5194/acp-22-3235-2022>, 2022.

1867 Schamm, K., Ziese, M., Becker A., Finger, P., Meyer-Christoffer, A., Schneider, U., Schroder, M., and Stender, P.:  
1868 Global gridded precipitation over land: a description of the new GPCC First Guess Daily product, *Earth Syst. Sci.*  
1869 *Data*, 6, 49-60. <https://doi.org/10.5194/essd-6-49-2014>, 2014.

1870 Schmidt, G.: Climate models can't explain 2023's huge heat anomaly — we could be in uncharted territory, *Nature*,  
1871 627, 467–467, <https://doi.org/10.1038/d41586-024-00816-z>, 2024.

1872 von Schuckmann, K., Cheng, L., Palmer, M. D., Hansen, J., Tassone, C., Aich, V., Adusumilli, S., Beltrami, H., Boyer,  
1873 T., Cuesta-Valero, F. J., Desbruyères, D., Domingues, C., García-García, A., Gentine, P., Gilson, J., Gorfer, M.,  
1874 Haimberger, L., Ishii, M., Johnson, G. C., Killick, R., King, B. A., Kirchengast, G., Kolodziejczyk, N., Lyman, J.,  
1875 Marzeion, B., Mayer, M., Monier, M., Monselesan, D. P., Purkey, S., Roemmich, D., Schweiger, A., Seneviratne, S.  
1876 I., Shepherd, A., Slater, D. A., Steiner, A. K., Straneo, F., Timmermans, M.-L., and Wijffels, S. E.: Heat stored in the  
1877 Earth system: where does the energy go?, *Earth Syst. Sci. Data*, 12, 2013–2041, [https://doi.org/10.5194/essd-12-2013-](https://doi.org/10.5194/essd-12-2013-2020)  
1878 [2020](https://doi.org/10.5194/essd-12-2013-2020), 2020.

1879 von Schuckmann, K., Minière, A., Gues, F., Cuesta-Valero, F. J., Kirchengast, G., Adusumilli, S., Straneo, F., Ablain,  
1880 M., Allan, R. P., Barker, P. M., Beltrami, H., Blazquez, A., Boyer, T., Cheng, L., Church, J., Desbruyeres, D., Dolman,  
1881 H., Domingues, C. M., García-García, A., Giglio, D., Gilson, J. E., Gorfer, M., Haimberger, L., Hakuba, M. Z.,  
1882 Hendricks, S., Hosoda, S., Johnson, G. C., Killick, R., King, B., Kolodziejczyk, N., Korosov, A., Krinner, G., Kuusela,  
1883 M., Landerer, F. W., Langer, M., Lavergne, T., Lawrence, I., Li, Y., Lyman, J., Marti, F., Marzeion, B., Mayer, M.,  
1884 MacDougall, A. H., McDougall, T., Monselesan, D. P., Nitzbon, J., Otsuka, I., Peng, J., Purkey, S., Roemmich, D.,  
1885 Sato, K., Sato, K., Savita, A., Schweiger, A., Shepherd, A., Seneviratne, S. I., Simons, L., Slater, D. A., Slater, T.,  
1886 Steiner, A. K., Suga, T., Szekely, T., Thiery, W., Timmermans, M.-L., Vanderkelen, I., Wijffels, S. E., Wu, T., and  
1887 Zemp, M.: Heat stored in the Earth system 1960–2020: where does the energy go?, *Earth System Science Data*, 15,  
1888 1675–1709, <https://doi.org/10.5194/essd-15-1675-2023>, 2023a.

1889 ▼

1890 Schoeberl, M. R., Wang, Y., Taha, G., Zawada, D. J., Ueyama, R., and Dessler, A.: Evolution of the Climate Forcing  
1891 During the Two Years After the Hunga Tonga-Hunga Ha’apai Eruption, *JGR Atmospheres*, 129, e2024JD041296,  
1892 <https://doi.org/10.1029/2024JD041296>, 2024.

1893 Schwingshackl, C., Obermeier, W. A., Bultan, S., Grassi, G., Canadell, J. G., Friedlingstein, P., Gasser, T., Houghton,  
1894 R. A., Kurz, W. A., Sitch, S., and Pongratz, J.: Differences in land-based mitigation estimates reconciled by separating  
1895 natural and land-use CO<sub>2</sub> fluxes at the country level, *One Earth*, 5, 1367–1376,  
1896 <https://doi.org/10.1016/j.oneear.2022.11.009>, 2022.

1897 Seneviratne, S. I., X. Zhang, M. Adnan, W. Badi, C. Dereczynski, A. Di Luca, S. Ghosh, I. Iskandar, J. Kossin, S.  
1898 Lewis, F. Otto, I. Pinto, M. Satoh, S. M. Vicente-Serrano, M. Wehner, and B. Zhou: Weather and Climate Extreme  
1899 Events in a Changing Climate. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group*  
1900 *I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai,  
1901 A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E.  
1902 Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge  
1903 University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1513–1766,  
1904 doi:10.1017/9781009157896.013.1513–1766, <https://doi.org/10.1017/9781009157896.013>, 2021.

1905 Seo, K.-W., Ryu, D., Jeon, T., Youm, K., Kim, J.-S., Oh, E. H., Chen, J., Famiglietti, J. S., and Wilson, C. R.: Abrupt  
1906 sea level rise and Earth’s gradual pole shift reveal permanent hydrological regime changes in the 21st century, *Science*,  
1907 387, 1408–1413, <https://doi.org/10.1126/science.adq6529>, 2025.

**Deleted:** von Schuckmann, K., Minière, A., Gues, F., Cuesta-Valero, F. J., Kirchengast, G., Adusumilli, S., Straneo, F., Ablain, M., Allan, R. P., Barker, P. M., Beltrami, H., Blazquez, A., Boyer, T., Cheng, L., Church, J., Desbruyeres, D., Dolman, H., Domingues, C. M., García-García, A., Giglio, D., Gilson, J. E., Gorfer, M., Haimberger, L., Hakuba, M. Z., Hendricks, S., Hosoda, S., Johnson, G. C., Killick, R., King, B., Kolodziejczyk, N., Korosov, A., Krinner, G., Kuusela, M., Landerer, F. W., Langer, M., Lavergne, T., Lawrence, I., Li, Y., Lyman, J., Marti, F., Marzeion, B., Mayer, M., MacDougall, A. H., McDougall, T., Monselesan, D. P., Nitzbon, J., Otsuka, I., Peng, J., Purkey, S., Roemmich, D., ...

1921 Sherwin, E. D., Rutherford, J. S., Zhang, Z., Chen, Y., Wetherley, E. B., Yakovlev, P. V., Berman, E. S. F., Jones, B.  
 1922 B., Cusworth, D. H., Thorpe, A. K., Ayasse, A. K., Duren, R. M., and Brandt, A. R.: US oil and gas system emissions  
 1923 from nearly one million aerial site measurements, *Nature*, 627, 328–334, [https://doi.org/10.1038/s41586-024-07117-](https://doi.org/10.1038/s41586-024-07117-5)  
 1924 [5](https://doi.org/10.1038/s41586-024-07117-5), 2024.

1925 Simmonds, P. G., Rigby, M., McCulloch, A., O'Doherty, S., Young, D., Mühle, J., Krummel, P. B., Steele, P., Fraser,  
 1926 P. J., Manning, A. J., Weiss, R. F., Salameh, P. K., Harth, C. M., Wang, R. H. J., and Prinn, R. G.: Changing trends  
 1927 and emissions of hydrochlorofluorocarbons (HCFCs) and their hydrofluorocarbon (HFCs) replacements, *Atmos.*  
 1928 *Chem. Phys.*, 17, 4641–4655, <https://doi.org/10.5194/acp-17-4641-2017>, 2017.

1929 Sippel, S., Zscheischler, J., Heimann, M., Otto, F. E. L., Peters, J., and Mahecha, M. D.: Quantifying changes in  
 1930 climate variability and extremes: Pitfalls and their overcoming, *Geophys. Res. Lett.*, 42, 9990–9998,  
 1931 <https://doi.org/10.1002/2015GL066307>, 2015.

1932 Smith, C., Walsh, T., Gillett, N., Hall, B., Hauser, M., Krummel, P., Lamb, W., Lamboll, R., Muhle, J., Palmer, M.,  
 1933 Ribes, A., Seneviratne, S., Trewin, B., von Schuckmann, K., and Forster, P.: ClimateIndicator/data: Indicators of  
 1934 Global Climate Change 2024 submission (v2025.06.11), [Data set], <https://doi.org/10.5281/zenodo.15639576>, 2025a.

1935 Smith, C., Walsh, T., Gillett, N., Hall, B., Hauser, M., Krummel, P., Lamb, W., Lamboll, R., X., Muhle, J., Palmer,  
 1936 M., Ribes, A., Seneviratne, S., Trewin, B., von Schuckmann, K., and Forster, P.: Data repository for Indicators of  
 1937 Global Climate Change, Github [code], <https://github.com/ClimateIndicator/data/tree/v2025.06.11>, last access: 11  
 1938 June, 2025, 2025b

1939 Smith, C., Nicholls, Z. R. J., Armour, K., Collins, W., Forster, P., Meinshausen, M., Palmer, M. D., and Watanabe,  
 1940 M.: The Earth's Energy Budget, Climate Feedbacks, and Climate Sensitivity Supplementary Material, in: *Climate*  
 1941 *Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the*  
 1942 *Intergovernmental Panel on Climate Change*, edited by: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L.,  
 1943 Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews,  
 1944 J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., 2021.

1945 [Smith, C., Forster, P., Palmer, M., Collins, B., Leach, N., Watanabe, M., Berger, S., Hall, B., Zelinka, M., Lunt, D.,](https://doi.org/10.5281/zenodo.5211358)  
 1946 [Cain, M., Harris, G., and Ringer, M.: IPCC WGI AR6 Chapter 7 \(v.1.0\). Zenodo.](https://doi.org/10.5281/zenodo.5211358)  
 1947 <https://doi.org/10.5281/zenodo.5211358>, 2021.

1948 Smith, S. J., van Aardenne, J., Klimont, Z., Andres, R. J., Volke, A., and Delgado Arias, S.: Anthropogenic sulfur  
 1949 dioxide emissions: 1850–2005, *Atmos. Chem. and Phys.*, 11, 1101–1116, <https://doi.org/10.5194/acp-11-1101-2011>,  
 1950 2011.

1951 Soulie, A., C. Granier, S. Darras, N. Zilbermann, T. Doumbia, M. Guevara, J.-P. Jalkanen, S. Keita, C. Liousse, M.  
 1952 Crippa, D. Guizzardi, R. Hoesly, S. J. Smith Global Anthropogenic Emissions (CAMSGLOBANT) for the  
 1953 Copernicus Atmosphere Monitoring Service Simulations of Air Quality Forecasts and Reanalyses Earth Syst. Sci.  
 1954 Data, 2023.

Deleted: v2025.05.02

Deleted: <https://doi.org/10.5281/zenodo.15327155>

Deleted: <https://github.com/ClimateIndicator/data/tree/v2025.05.02>

Deleted: 2 May

1960 Storto, A. and Yang, C.: Acceleration of the ocean warming from 1961 to 2022 unveiled by large-ensemble reanalyses,  
 1961 Nature Communications, 15, 545, <https://doi.org/10.1038/s41467-024-44749-7>, 2024.

1962 Szopa, S., V. Naik, B. Adhikary, P. Artaxo, T. Bernsten, W.D. Collins, S. Fuzzi, L. Gallardo, A. Kiendler-Scharr, Z.  
 1963 Klimont, H. Liao, N. Unger, and P. Zanis: Short-Lived Climate Forcers. In Climate Change 2021: The Physical  
 1964 Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on  
 1965 Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L.  
 1966 Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi,  
 1967 R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp.  
 1968 817–922, <https://doi.org/10.1017/9781009157896.008>, 2021.

1969 <https://doi.org/10.1038/s41586-025-08674-z>, 2025.

1970 Terhaar, J., Burger, F.A., Vogt, L. et al.: Record sea surface temperature jump in 2023–2024 unlikely but not  
 1971 unexpected, Nature 639, 942–946, <https://doi.org/10.1038/s41586-025-08674-z>, 2025.

1972 Trewin, B.: Assessing Internal Variability of Global Mean Surface Temperature From Observational Data and  
 1973 Implications for Reaching Key Thresholds, JGR Atmospheres, 127, e2022JD036747,  
 1974 <https://doi.org/10.1029/2022JD036747>, 2022.

1975 Tibrewal, K., Ciais, P., Saunio, M., Martinez, A., Lin, X., Thanwerdas, J., Deng, Z., Chevallier, F., Giron, C.,  
 1976 Albergel, C., Tanaka, K., Patra, P., Tsuruta, A., Zheng, B., Belikov, D., Niwa, Y., Janardanan, R., Maksyutov, S.,  
 1977 Segers, A., Tzompa-Sosa, Z. A., Bousquet, P., and Sciare, J.: Assessment of methane emissions from oil, gas and coal  
 1978 sectors across inventories and atmospheric inversions, Communications Earth & Environment, 5, 26,  
 1979 <https://doi.org/10.1038/s43247-023-01190-w>, 2024.

1980 Vakilifard, N., Williams, R. G., Holden, P. B., Turner, K., Edwards, N. R., and Beerling, D. J.: Impact of negative and  
 1981 positive CO<sub>2</sub> emissions on global warming metrics using an ensemble of Earth system model simulations,  
 1982 Biogeosciences, 19, 4249–4265, <https://doi.org/10.5194/bg-19-4249-2022>, 2022.

1983 Vanderkelen, I. and Thiery, W.: GCOS EHI 1960-2020 Inland Water Heat Content,  
 1984 [https://doi.org/10.26050/WDCC/GCOS\\_EHI\\_1960-2020\\_IWHC](https://doi.org/10.26050/WDCC/GCOS_EHI_1960-2020_IWHC), 2022.

1985 Vimont, I. J. B. D. Hall, G. Dutton, S. A. Montzka, J. Mühle, M. Crotwell, K. Petersen, S. Clingan, and D. Nance, [in  
 1986 “State of the Climate in 2022”]. Bull. Amer. Meteor. Soc., 104, 9, S76–S78, [https://doi.org/10.1175/BAMS-D-23-](https://doi.org/10.1175/BAMS-D-23-0090.1)  
 1987 [0090.1](https://doi.org/10.1175/BAMS-D-23-0090.1), 2022.

1988 Vollmer, M. K., Young, D., Trudinger, C. M., Mühle, J., Henne, S., Rigby, M., Park, S., Li, S., Guillevis, M.,  
 1989 Mitrevski, B., Harth, C. M., Miller, B. R., Reimann, S., Yao, B., Steele, L. P., Wyss, S. A., Lunder, C. R., Arduini, J.,  
 1990 McCulloch, A., Wu, S., Rhee, T. S., Wang, R. H. J., Salameh, P. K., Hermansen, O., Hill, M., Langenfelds, R. L., Ivy,  
 1991 D., O'Doherty, S., Krummel, P. B., Maione, M., Etheridge, D. M., Zhou, L., Fraser, P. J., Prinn, R. G., Weiss, R. F.,  
 1992 and Simmonds, P. G.: Atmospheric histories and emissions of chlorofluorocarbons CFC-13 (CClF<sub>3</sub>), ΣCFC-114  
 1993 (C<sub>2</sub>Cl<sub>2</sub>F<sub>4</sub>), and CFC-115 (C<sub>2</sub>ClF<sub>5</sub>), Atmos. Chem. Phys., 18, 979–1002, <https://doi.org/10.5194/acp-18-979-2018>,  
 1994 2018.

**Deleted:** Terhaar, J., Vogt, L., and Foukal, N. P.: Atlantic overturning inferred from air-sea heat fluxes indicates no decline since the 1960s, Nat Commun, 16, 222, <https://doi.org/10.1038/s41467-024-55297-5>, 2025.



1999 Watson-Parris, D., Christensen, M. W., Laurenson, A., Clewley, D., Gryspeerdt, E., and Stier, P.: Shipping regulations  
2000 lead to large reduction in cloud perturbations, *Proc. Natl. Acad. Sci. U.S.A.*, 119, e2206885119,  
2001 <https://doi.org/10.1073/pnas.2206885119>, 2022.

2002 WCRP Global Sea Level Budget Group: Global sea-level budget 1993–present, *Earth Syst. Sci. Data*, 10, 1551–1590,  
2003 <https://doi.org/10.5194/essd-10-1551-2018>, 2018.

2004 Western, L. M., Vollmer, M. K., Krummel, P. B., Adcock, K. E., Fraser, P. J., Harth, C. M., Langenfelds, R. L.,  
2005 Montzka, S. A., Mühle, J., O'Doherty, S., Oram, D. E., Reimann, S., Rigby, M., Vimont, I., Weiss, R. F., Young, D.,  
2006 and Laube, J. C.: Global increase of ozone-depleting chlorofluorocarbons from 2010 to 2020, *Nat. Geosci.*, 16, 309–  
2007 313, <https://doi.org/10.1038/s41561-023-01147-w>, 2023.

2008 Western, L. M., Daniel, J. S., Vollmer, M. K., Clingan, S., Crotwell, M., Fraser, P. J., Ganesan, A. L., Hall, B., Harth,  
2009 C. M., Krummel, P. B., Mühle, J., O'Doherty, S., Salameh, P. K., Stanley, K. M., Reimann, S., Vimont, I., Young,  
2010 D., Rigby, M., Weiss, R. F., Prinn, R. G., and Montzka, S. A.: A decrease in radiative forcing and equivalent effective  
2011 chlorine from hydrochlorofluorocarbons, *Nat. Clim. Chang.*, 14, 805–807, [https://doi.org/10.1038/s41558-024-02038-](https://doi.org/10.1038/s41558-024-02038-7)  
2012 [7](https://doi.org/10.1038/s41558-024-02038-7), 2024.

2013 van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van Marle,  
2014 M. J. E., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during  
2015 1997–2016, *Earth System Science Data*, 9, 697–720, <https://doi.org/10.5194/essd-9-697-2017>, 2017.

2016 Wild, M., Gilgen, H., Roesch, A., Ohmura, A., Long, C. N., Dutton, E. G., Forgan, B., Kallis, A., Russak, V., and  
2017 Tsvetkov, A.: From Dimming to Brightening: Decadal Changes in Solar Radiation at Earth's Surface, *Science*, 308,  
2018 847–850, <https://doi.org/10.1126/science.1103215>, 2005.

2019 World Meteorological Organization (WMO). State of the Global Climate 2024. WMO-No. 1368. 2025. ISBN: 978-  
2020 92-63-11368-5.

2021 Xu, Q., Wei, S., Li, Z., and Li, Q.: A New Evaluation of Observed Changes in Diurnal Temperature Range,  
2022 *Geophysical Research Letters*, 52, e2024GL113406, <https://doi.org/10.1029/2024GL113406>, 2025.

2023 Zickfeld, K., Azevedo, D., Mathesius, S., and Matthews, H. D.: Asymmetry in the climate–carbon cycle response to  
2024 positive and negative CO<sub>2</sub> emissions, *Nat. Clim. Chang.*, 11, 613–617, <https://doi.org/10.1038/s41558-021-01061-2>,  
2025 2021.

2026 Zhang, Z., Poulter, B., Feldman, A.F., Ying, Q., Ciais, P., Peng, S. and Xin, L.: Recent intensification of wetland  
2027 methane feedback, *Nat. Clim. Chang.* 13, 430–433, <https://doi.org/10.1038/s41558-023-01629-0>, 2023.

2028